# On Sequential Locally Repairable Codes

Wentu Song, Kai Cai and Chau Yuen

#### Abstract

We consider the locally repairable codes (LRC), aiming at sequential recovering multiple erasures. We define the (n, k, r, t)-SLRC (Sequential Locally Repairable Codes) as an [n, k] linear code where any  $t'(\leq t)$  erasures can be sequentially recovered, each one by r ( $2 \leq r < k$ ) other code symbols. Sequential recovering means that the erased symbols are recovered one by one, and *an already recovered symbol can be used* to recover the remaining erased symbols. This important recovering method, in contrast with the vastly studied parallel recovering, is currently far from understanding, say, lacking codes constructed for arbitrary  $t \geq 3$  erasures and bounds to evaluate the performance of such codes.

We first derive a tight upper bound on the code rate of (n, k, r, t)-SLRC for t = 3 and  $r \ge 2$ . We then propose two constructions of binary (n, k, r, t)-SLRCs for general  $r, t \ge 2$  (Existing constructions are dealing with  $t \le 7$  erasures). The first construction generalizes the method of direct product construction. The second construction is based on the resolvable configurations and yields SLRCs for any  $r \ge 2$  and odd  $t \ge 3$ . For both constructions, the rates are optimal for  $t \in \{2, 3\}$  and are higher than most of the existing LRC families for arbitrary  $t \ge 4$ .

#### **Index Terms**

Distributed storage, locally repairable codes, parallel recovery, sequential recovery.

#### I. INTRODUCTION

To avoid the inefficiency of straightforward replication of data, various coding techniques are introduced to the distributed storage system (DSS), among which the linear locally repairable codes, also known as locally recoverable codes (LRC) [3], [4], attracted much attention recently. Roughly speaking, a linear LRC with locality r is an [n, k] linear code such that the value of each coordinate (code symbol) can be computed from the values of at most r other coordinates.

In a DSS system where a LRC C is used, the information stored in each storage node corresponds to one coordinate of C. Hence, each *single* node failure (erasure) can be recovered by a set of at most r other nodes. However, it is very common that two or more storage nodes fail in the system. This problem, which has become a central focus for the LRC society, are recently investigated by many authors (e.g. [6]-[21]). Basically, when multiple erasures occur, the recovering performance can be heavily depends on the recovering strategy in use, say, recovering the erasures simultaneously or one by one. The two strategies were first distinguished as *parallel approach* and *sequential approach* in [17]. Comparing with the parallel approach, the sequential approach recovery erasures one by one and hence the already fixed erasure nodes can be used in the next round of recovering. Potentially, for the same LRC, using the sequential approach can fix more erasures than using the parallel approach, and hence the sequential approach is a better candidate than the parallel approach in practice. However, due to technique difficulties, this more important approach remains far from understood, say, lacking of both code constructions and bounds to evaluate the code performance. In contrast with the vastly studied parallel approach [6]-[16], existing work on the sequential approach up to date are limited to dealing with  $t \leq 7$  erasures. For example, the case of t = 2 are considered in [17], where the authors derived upper bounds on the code rate as well as minimum distance and also constructed a family of distance-optimal codes based on Turán graphs. For the code rate, they proved that:

$$\frac{k}{n} \le \frac{r}{r+2}.\tag{1}$$

The original version of this work [18], firstly considered the case of t = 3 and gave both constructions and code rate bounds for  $t \in \{2, 3\}$  (in a more generalized manner of functional recovering). Of great relevance to the present work are the results recently obtained in [20] and [21], where the authors derived a lower bound on code length n of *binary code* for t = 3 and an upper bound on code rate of *binary code* for t = 4. A couple of optimal or high rate constructions were provided in these two papers, say, rate-optimal codes for  $t \in \{2, 3, 4\}$ , and high rate codes for r = 2 and  $t \in \{5, 6, 7\}$ . Here, we note that, by using orthogonal Latin squares, the authors in [20] gave an interesting construction of sequential locally recoverable codes for *any odd*  $t \ge 3$  with rate  $k/n = 1/(1 + \frac{t-r}{r} + \frac{1}{r^2})$ . Obviously, the SLRCs can deal with any t erasures and having high code rate and are highly desired in both theory and practices.

W. Song and C. Yuen are with Singapore University of Technology and Design, Singapore (e-mails: {wentu\_song, yuenchau}@sutd.edu.sg). Kai Cai is with the Department of Mathematics, University of Hong Kong. (e-mail: kcai@hku.hk).

# A. Our Contribution

In practice, high rate LRCs are desired since they mean low storage overhead. In this work, we are interested in the high rate LRCs for sequential recovering any  $t \ge 3$  erasures, by defining the (n, k, r, t)-SLRC (Sequential Locally Repairable Code) as an [n, k] linear code in which any t' ( $t' \le t$ ) erased code symbols can be sequentially recovered, *each one* by at most r ( $2 \le r < k$ ) other symbols. Our first contribution is an upper bound on the code rate for (n, k, r, t)-SLRC with t = 3 and any  $k > r \ge 2$ . The bound is derived by using a graph theoretical method, say, we associate each (n, k, r, t)-SLRC with a set of directed acyclic graphs, called repair graphs, and then obtain the bound by studying the structural properties of the so-called *minimal repair graph*. The sprit of this method lies in [23], [24]. For general  $t \ge 5$ , deriving an achievable, explicit upper bound of the rate of (n, k, r, t)-SLRC seems very challenging, and we give some discussions and conjectures on this issue.

Then we construct two families of binary (n, k, r, t)-SLRC. The first family, which contains the product of m copies of the binary [r + 1, r] single-parity code [10] as a special case, is for any positive integers  $r (\geq 2)$  and t, and has rate

$$\frac{k}{n} = \frac{1}{\sum_{s=0}^{t} \frac{1}{r^{|\operatorname{supp}_m(s)|}}},$$

where m is any given positive integer satisfying  $t \le 2^m - 1$  and  $\operatorname{supp}_m(s)$  is the support of the m-digit binary representation<sup>1</sup> of s. The second family is constructed for any  $r \ge 2$  and any odd integer  $t \ge 3$  and is based on *resolvable configurations*. This family has code rate

$$\frac{k}{n} = \left(1 + \frac{t-1}{r} + \left\lceil \frac{1}{r^2} \right\rceil\right)^{-1},$$

which is the same with the Latin square-based code constructed in [20]. For  $t \in \{2, 3\}$ , the code rates of these two constructions are optimal.

A basic and important fact revealed by our study is: the sequential approach can have much better performance than parallel approach, e.g., for the direct product of m copies of the binary [r + 1, r] single-parity code, it can recover m erasures with locality r by the parallel approach [10], but  $2^m - 1$  erasures with the same locality by the sequential approach.

#### B. Related Work

Except that mentioned previously, most existing work focus on [n, k] linear LRCs with parallel approach. In [7], the authors defined and constructed the  $(r, t + 1)_a$  code, for which each code symbol i is contained in a punctured code (local code) with length  $\leq r + t$  and minimum distance  $\geq t + 1$ . Clearly, for such codes, any t erased code symbols can be recovered in parallel by at most tr other code symbols, among which, each erased symbol can be recovered by at most r symbols. The code rate of this family satisfies [14]

$$\frac{k}{n} \le \frac{r}{r+t}.$$
(2)

Another family is the codes with locality r and availability t [8], [9], for which, each code symbol has t disjoint recovering sets of size at most r. An upper bound on the code rate of such codes is proved in [10]:

$$\frac{k}{n} \le \frac{1}{\prod_{j=1}^{t} (1 + \frac{1}{jr})}.$$
(3)

Unfortunately, for  $t \ge 3$ , the tightness of bound (3) is not known and most of existing construction have rate  $\le \frac{r}{r+t}$  (e.g., see [12], [15], [16]). Constructions with rate  $> \frac{r}{r+t}$  are proposed only for some very special values, e.g.,  $(n, k, r, t) = (2^{r+1} - 1, 2^r - 1, r, r+1)$  [16]. The third family of parallel recovery LRC is proposed in [6], in which, for any set  $E \subseteq [n]$  of erasures of size at most t and any  $i \in E$ , the *i*th code symbol has a recovering set of size at most r contained in  $[n] \setminus E$ . The fourth family, called codes with cooperative local repair, is proposed in [13] and defined by a stronger condition: each subset of t code symbols can be *cooperatively* recovered from at most r other code symbols. For this family, an upper bound of the code rate with exactly the same form as (2) is derived [13]. By far, constructing LRCs with high code rate (e.g.,  $\frac{k}{n} > \frac{r}{r+t}$ ) is still an interesting open problem, both for parallel recovery and for sequential recovery.

#### C. Organization

The rest of this paper is organized as follows. In Section II, we define the (n, k, r, t)-SLRC and then present some basic and useful facts. In section III, we first investigate the (minimal) repair graphs of the SLRC and then prove the upper bound on the code rate of (n, k, r, t)-SLRC for  $t \in \{2, 3\}$ . Before constructing the first family of SLRC in Section V, we first study an example in Section IV. Then, the second family of SLRC is constructed in Section VI. Finally, the paper is concluded in Section VII.

<sup>1</sup>The *m*-digit binary representation of any positive integer  $s \leq 2^m - 1$  is the binary vector  $(\lambda_m, \lambda_{m-1}, \dots, \lambda_1) \in \mathbb{Z}_2^m$  such that  $s = \sum_{j=1}^m \lambda_j 2^{j-1}$ .

## D. Notations

For any positive integer n,  $[n] := \{1, 2, \dots, n\}$ . For any set A, |A| is the size (the number of elements) of A. If  $B \subseteq A$  and |B| = t, then B is called a t-subset of A. For any real number x,  $\lceil x \rceil$  is the smallest integer greater than or equal to x. If C is an [n, k] linear code and  $A \subseteq [n]$ , then  $C|_A$  denotes the punctured code by puncturing coordinates in  $\overline{A} := [n] \setminus A$ . For any codeword  $x = (x_1, x_2, \dots, x_n) \in C$ ,  $\operatorname{supp}(x) := \{i \in [n]; x_i \neq 0\}$  is the support of x.

# II. PRELIMINARY

## A. Sequential Locally repairable code (SLRC)

Let C be an [n, k] linear code over the finite field  $\mathbb{F}$  and  $i \in [n]$ . A subset  $R \subseteq [n] \setminus \{i\}$  is called a *recovering set* of i if there exists an  $a_j \in \mathbb{F} \setminus \{0\}$  for each  $j \in R$  such that  $x_i = \sum_{j \in R} a_j x_j$  for all  $x = (x_1, x_2, \dots, x_n) \in C$ . Equivalently, there exists a codeword y in the dual code  $C^{\perp}$  such that  $\sup(y) = R \cup \{i\}$ .

Definition 1 (Sequential Locally Repairable Code): For any  $E \subseteq [n]$ , C is said to be (E, r)-recoverable if E can be sequentially indexed, say  $E = \{i_1, i_2, \dots, i_{|E|}\}$ , such that each  $i_{\ell} \in E$  has a recovering set  $R_{\ell} \subseteq \overline{E} \cup \{i_1, \dots, i_{\ell-1}\}$  of size  $|R_{\ell}| \leq r$ , where  $\overline{E} := [n] \setminus E$ ; C is called an (n, k, r, t)-sequential locally repairable code (SLRC) (or simply (r, t)-SLRC) if C is (E, r)-recoverable for each  $E \subseteq [n]$  of size  $|E| \leq t$ , where r is called the locality of C.

As a special case of Definition 1, if for each  $E \subseteq [n]$  of size  $|E| \leq t$  and each  $i \in E$ , *i* has a recovering set  $R \subseteq \overline{E}$  of size  $|R| \leq r$ , then C is called an (n, k, r, t)-parallel locally repairable code (PLRC). This special case is first considered in [6].

By the definition, we can have  $r \le k$  for any (n, k, r, t)-SLRC. Throughout this paper, we assume that a recovering set R has size  $2 \le |R| \le r < k$ . The following equivalent form of Definition 1 will be frequently used in our paper.

Lemma 2: C is an (n, k, r, t)-SLRC if and only if for any nonempty  $E \subseteq [n]$  of size  $|E| \leq t$ , there exists an  $i \in E$  such that i has a recovering set  $R \subseteq [n] \setminus E$ .

*Proof:* Let C be an (n, k, r, t)-SLRC and  $\emptyset \neq E \subseteq [n]$  of size  $|E| \leq t$ . Then by Definition 1, E can be sequentially indexed as  $E = \{i_1, i_2, \dots, i_{|E|}\}$  such that  $i_1$  has a recovering set  $R_1 \subseteq [n] \setminus E$ .

Conversely, for any  $E \subseteq [n]$  of size  $|E| \leq t$ , by assumption, one can find an  $i_1 \in E$  such that  $i_1$  has a recovering set  $R_1 \subseteq [n] \setminus E$ . Further, since  $|E \setminus \{i_1\}| < |E| \leq t$ , then by assumption, there exists an  $i_2 \in E \setminus \{i_1\}$  such that  $i_2$  has a recovering set  $R_2 \subseteq [n] \setminus (E \setminus \{i_1\}) = \overline{E} \cup \{i_1\}$ . Similarly, we can find an  $i_3 \in E \setminus \{i_1, i_2\}$  such that  $i_3$  has a recovering set  $R_3 \subseteq \overline{E} \cup \{i_1, i_2\}$ , and so on. Then E can be sequentially indexed as  $E = \{i_1, i_2, \dots, i_{|E|}\}$  such that each  $i_\ell \in E$  has a recovering set  $R_\ell \subseteq \overline{E} \cup \{i_1, \dots, i_{\ell-1}\}$ . So by definition 1, C is an (n, k, r, t)-SLRC.

The following lemma gives a sufficient condition of (r, t)-SLRC, which reflects the difference between the sequential recovery and the parallel recovery.

Lemma 3: Suppose  $[n] = A \cup B$  and  $A \cap B = \emptyset$ . Suppose  $t_1, t_2 \ge 0$  and C is an [n, k] linear code such that

(1) For any nonempty  $E \subseteq A$  of size  $|E| \leq t_1$ , there exists an  $i \in E$  such that i has a recovering set  $R \subseteq A \setminus E$ ;

(2) For any nonempty  $E \subseteq A$  of size  $|E| \leq t_1 + t_2 + 1$ , there exists an  $i \in E$  such that i has a recovering set  $R \subseteq [n] \setminus E$ ;

(3) For any nonempty  $E \subseteq B$  of size  $|E| \leq t_2$ , there exists an  $i \in E$  such that i has a recovering set  $R \subseteq B \setminus E$ ;

(4) For any nonempty  $E \subseteq B$  of size  $|E| \leq t_1 + t_2 + 1$ , there exists an  $i \in E$  such that i has a recovering set  $R \subseteq [n] \setminus E$ . Then C is an (r, t)-SLRC with  $t = t_1 + t_2 + 1$ .

*Proof:* We prove, by Lemma 2, that for any nonempty  $E \subseteq [n]$  of size  $|E| \leq t_1 + t_2 + 1$ , there exists an  $i \in E$  such that i has a recovering set  $R \subseteq [n] \setminus E$ . Obviously, it holds when  $E \subseteq A$  or  $E \subseteq B$  (by condition (2) or (4)). So we assume  $E \cap A \neq \emptyset$  and  $E \cap B \neq \emptyset$ . Consider the following two cases.

Case 1:  $0 < |E \cap A| \le t_1$ . By condition (1), there exists an  $i \in E$  such that i has a recovering set  $R \subseteq A \setminus E \subseteq [n] \setminus E$ .

Case 2:  $|E \cap A| > t_1$ . Since  $|E| \le t_1 + t_2 + 1$  and  $A \cap B = \emptyset$ , then  $0 < |E \cap B| \le t_2$ . By condition (3), there exists an  $i \in E$  such that i has a recovering set  $R \subseteq B \setminus E \subseteq [n] \setminus E$ .

The proof is completed by combining the above cases.

#### B. Repair Graph and Minimal Repair Graph

Let  $G = (\mathcal{V}, \mathcal{E})$  be a directed, acyclic graph, where  $\mathcal{V}$  is the vertex set and  $\mathcal{E}$  is the (directed) edge set. A directed edge e from vertex u to v is denoted by an ordered pair e = (u, v), where u is called the *tail* of e and v the *head* of e. Moreover, u is called an *in-neighbor* of v and v an *out-neighbor* of u. For each  $v \in \mathcal{V}$ , let  $\ln(v)$  and  $\operatorname{Out}(v)$  denote the set of in-neighbors and out-neighbors of v respectively. If  $\ln(v) = \emptyset$ , we call v a *source*; otherwise, v is called an *inner vertex*. Denote by S(G) the set of all sources of G. For any  $E \subseteq \mathcal{V}$ , let

$$\operatorname{Out}(E) = \bigcup_{v \in E} \operatorname{Out}(v) \backslash E.$$
(4)

By (4), we have  $E \cap \text{Out}(E) = \emptyset$ . For any  $v \in \mathcal{V}$ , denote

$$\operatorname{Out}^{2}(v) = \bigcup_{u \in \operatorname{Out}(v)} \operatorname{Out}(u) \setminus \operatorname{Out}(v)$$
(5)

4

i.e.,  $\operatorname{Out}^2(v)$  is the set of all  $w \in \mathcal{V}$  such that w is an out-neighbor of some  $u \in \operatorname{Out}(v)$  but not an out-neighbor of v.

As an example, consider the graph depicted in Fig. 1, where vertices are indexed by  $\{1, 2, \dots, 16\}$ . Then  $Out(3) = \{9, 10\}$ ,  $Out(4) = \{10, 11\}$  and  $Out^2(3) = \{13, 15, 16\}$ . Let  $E = \{3, 4\}$ . Then  $Out(E) = \{9, 10, 11\}$ .




Fig 1. An example repair graph with n = 16, r = 2 and |S(G)| = 8.

Definition 4 (Repair Graph): Let C be an (n, k, r, t)-SLRC and  $G = (\mathcal{V}, \mathcal{E})$  be a directed, acyclic graph such that  $\mathcal{V} = [n]$ . G is called a *repair graph* of C if for all inner vertex  $i \in \mathcal{V}$ ,  $\ln(i)$  is a recovering set of i.

Obviously, an (n, k, r, t)-SLRC may have many repair graphs. If C is an (n, k, r, t)-SLRC, we usually use  $\{G_{\lambda}; \lambda \in \Lambda\}$  to denote the set of all repair graphs of C, where  $\Lambda$  is some proper index set. It should be noted that the repair graph defined here has subtle differences with the recovering graph defined in [10], e.g., it must be acyclic and an (n, k, r, t)-SLRC may have many repair graphs such that for each  $i \in [n]$ , at most one recovery set of i is considered in each repair graph. The key ingredient of our technique is the so-called *minimal repair graph* as defined follows.

Let C be an (n, k, r, t)-SLRC and  $\{G_{\lambda}; \lambda \in \Lambda\}$  be the set of all repair graphs of C. Recall that for each  $\lambda \in \Lambda$ ,  $S(G_{\lambda})$  is the set of all sources of  $G_{\lambda}$ . Denote

$$\delta^* \triangleq \min\{|\mathbf{S}(G_\lambda)|; \lambda \in \Lambda\}.$$
(6)

Definition 5 (Minimal Repair Graph): A repair graph  $G_{\lambda_0}, \lambda_0 \in \Lambda$ , is called a minimal repair graph of C if  $|S(G_{\lambda_0})| = \delta^*$ . Remark 6: It is easy to see that any (n, k, r, t)-SLRC has at least one minimal repair graph by noticing that the set  $\{|S(G_{\lambda})|; \lambda \in \Lambda\} \subseteq [n]$  is finite.

# III. AN UPPER BOUND ON THE CODE RATE

Before proposing the main result of this section, we need first investigate properties of the minimal repair graphs of (n, k, r, t)-SLRC.

## A. Properties of the Minimal Repair Graph

In this subsection, we always assume that C is an (n, k, r, t)-SLRC and  $G_{\lambda_0} = (\mathcal{V}, \mathcal{E})$  is a minimal repair graph of C. The following two results are of fundamental.

Lemma 7:

$$(n - \delta^*)r \ge |\mathcal{E}|.\tag{7}$$

*Proof:* By the definition,  $G_{\lambda_0}$  has  $n - \delta^*$  inner vertices and each of them has at most r in-neighbors, and hence the result follows.

Lemma 8:

$$k < \delta^*. \tag{8}$$

*Proof:* According to Definition 4, for each  $j \in [n]$ , the *j*th code symbol of C is a linear combination of the code symbols in  $\operatorname{In}(j)$ . In other words, the code symbols of  $\operatorname{In}(j)$  spans the code symbols of  $\{j\} \cup \operatorname{In}(j)$ . Moreover, since  $G_{\lambda_0}$  is acyclic, then inductively, the code symbols of  $S(G_{\lambda_0})$  spans C, which proves  $k \leq |S(G_{\lambda_0})| = \delta^*$ .

The following is a key lemma to investigate the structure of  $G_{\lambda_0}$ .

Lemma 9: For any  $E \subseteq [n]$  of size  $|E| \leq t$ ,

$$|\operatorname{Out}(E)| \ge |E \cap \mathbf{S}(G_{\lambda_0})|. \tag{9}$$

*Proof:* Suppose, on the contrary, there exists an  $E = \{i_1, i_2, \dots, i_{t'}\} \subseteq [n]$  such that  $|E| = t' \leq t$  and  $|\operatorname{Out}(E)| < |E \cap S(G_{\lambda_0})|$ . By definition of (n, k, r, t)-SLRC, we can let  $R_\ell \subseteq \overline{E} \cup \{i_1, \dots, i_{\ell-1}\}$  be a recovering set of  $i_\ell$  for each  $\ell \in [t']$ . We can construct a graph  $C_\ell$  from  $C_\ell$  by delating and adding edges as follows: First for each  $i \in E \sqcup \operatorname{Out}(E)$  and

We can construct a graph  $G_{\lambda_1}$  from  $G_{\lambda_0}$  by deleting and adding edges as follows: First, for each  $i \in E \cup \text{Out}(E)$  and  $j \in \text{In}(i)$ , delete (j, i) if it is an edge of  $G_{\lambda_0}$ , and denote the resulted graph as  $G_{\lambda'_1}$ ; Second, for each  $i_{\ell} \in E$  and each  $j \in R_{\ell}$ ,

add a directed edge from j to  $i_{\ell}$ , and let the resulted graph be  $G_{\lambda_1}$ . Clearly,  $G_{\lambda'_1}$  is acyclic because  $G_{\lambda_0}$  is acyclic. Moreover, since  $R_{\ell} \subseteq \overline{E} \cup \{i_1, \dots, i_{\ell-1}\}$  for each  $\ell \in [t']$ , then by construction,  $G_{\lambda_1}$  is also acyclic.

We declare that  $G_{\lambda_1}$  is a repair graph of C and  $|S(G_{\lambda_1})| < |S(G_{\lambda_0})|$ , which contradicts to the minimality of  $G_{\lambda_0}$ .

In fact, by construction,  $S(G_{\lambda_1}) = (S(G_{\lambda_0}) \setminus E) \cup Out(E)$ . Then for each inner node *i* of  $G_{\lambda_1}$ , we have the following two cases:

Case 1:  $i \in E$ . Then  $i = i_{\ell}$  for some  $\ell \in [t']$  and by the construction of  $G_{\lambda_1}$ ,  $\operatorname{In}(i) = R_{\ell}$  is a recovering set of i. Case 2: i is an inner vertex of  $G_{\lambda_0}$  and  $i \notin \operatorname{Out}(E)$ . Then considering  $G_{\lambda_0}$ ,  $\operatorname{In}(i) \subseteq \overline{E} = [n] \setminus E$  is a recovering set of i.

So In(i) is always a recovering set of *i*. Hence,  $G_{\lambda_1}$  is a repair graph of C.

On the other hand, note that by definition,  $S(G_{\lambda_0}) \cap Out(E) = \emptyset$  and  $E \cap Out(E) = \emptyset$ . So if we assume that  $|Out(E)| < |E \cap S(G_{\lambda_0})|$ , then we have

$$|\mathbf{S}(G_{\lambda_1})| = |(\mathbf{S}(G_{\lambda_0}) \setminus E) \cup \operatorname{Out}(E)|$$
  
=  $|(\mathbf{S}(G_{\lambda_0}) \setminus E)| + |\operatorname{Out}(E)|$   
=  $|(\mathbf{S}(G_{\lambda_0})| - |E \cap \mathbf{S}(G_{\lambda_0})| + |\operatorname{Out}(E)|$   
<  $|\mathbf{S}(G_{\lambda_0})|,$  (10)

which completes the proof.

The following example illustrates the construction of  $G_{\lambda_1}$  in the proof of Lemma 9.

*Example 10:* Consider the graph in Fig. 1, which we denote as  $G_{\lambda_0}$  here. Suppose it is a repair graph of a (r = 2, t = 3)-SLRC. We can see that  $\{2,3\}$  is a recovering set of 9,  $\{3,4\}$  is a recovering set of 10, and etc.

Let  $E = \{2,3,9\}$  and assume the recovering sets of 2, 3 and 9 are  $\{1,10\}, \{12,13\}$  and  $\{11,14\}$ , respectively. Then we can construct a graph  $G_{\lambda_1}$  as follows. Since  $Out(E) = \{10\}$ , thus, in the first step, we delete edges (2,9), (3,9), (3,10) and (4,10); and in the second step, we add edges (1,2), (10,2), (12,3), (13,3), (11,9) and (14,9). The resulted graph  $G_{\lambda_1}$  is shown in Fig. 2. We can see that  $|S(G_{\lambda_1})| = |(S(G_{\lambda_0}) \setminus E) \cup Out(E)| = |\{1,4,5,6,7,8,10\}| = 7 < 8 = |S(G_{\lambda_0})|$ . So the graph in Fig. 1 is not a minimal repair graph.




Fig 2. Construction of  $G_{\lambda_1}$  from the graph in Fig. 1.

The following two corollaries give some explicit structural properties of the minimal repair graphs of (n, k, r, t)-SLRC. Corollary 11: If  $t \ge 3$ , for any  $v \in S(G_{\lambda_0})$ , the following hold:

- 1)  $|Out(v)| \ge 1$ .
- 2) If  $\operatorname{Out}(v) = \{v'\}$ , then  $\operatorname{Out}^2(v) = \operatorname{Out}(v') \neq \emptyset$ .
- 3) If  $\operatorname{Out}(v) = \{v_1\}$  and  $\operatorname{Out}(v_1) = \{v_2\}$ , then  $\operatorname{Out}(v_2) \neq \emptyset$ .
- 4) If  $\operatorname{Out}(v) = \{v_1\}$  and  $\operatorname{Out}(v_1) = \{v_2\}$ , then  $|\operatorname{Out}(u)| \ge 2$  for any source  $u \in \operatorname{In}(v_2)$ .
- 5) If v, w are two distinct sources and |Out(v)| = |Out(w)| = 1, then  $Out(v) \neq Out(w)$ .

*Proof:* We can prove all claims by contradiction.

1) Suppose otherwise  $|\operatorname{Out}(v)| = 0$ . Let  $E = \{v\}$ . Then,  $|\operatorname{Out}(E)| = |\operatorname{Out}(v)| = 0 < 1 = |\{v\}| = |E \cap S(G_{\lambda_0})|$ , which contradicts to Lemma 9.

2) Since  $G_{\lambda_0}$  is acyclic and  $\operatorname{Out}(v) = \{v'\}$ , then from (5),  $\operatorname{Out}^2(v) = \operatorname{Out}(v')$ . If  $\operatorname{Out}(v') = \emptyset$ , then by letting  $E = \{v, v'\}$ , we have  $|\operatorname{Out}(E)| = |\emptyset| = 0 < 1 = |\{v\}| = |E \cap S(G_{\lambda_0})|$ , which contradicts to Lemma 9.

3) If  $\operatorname{Out}(v_2) = \emptyset$ , then by letting  $E = \{v, v_1, v_2\}$ , we have  $|\operatorname{Out}(E)| = |\emptyset| = 0 < 1 = |\{v\}| = |E \cap S(G_{\lambda_0})|$ , which contradicts to Lemma 9.

4) By assumption, we can see that  $u \neq v$ . Suppose otherwise |Out(u)| = 1. Then  $Out(u) = \{v_2\}$  since  $u \in In(v_2)$ . Let  $E = \{v, v_1, u\}$ . We have  $|Out(E)| = |\{v_2\}| = 1 < 2 = |\{v, u\}| = |E \cap S(G_{\lambda_0})|$ , which contradicts to Lemma 9.

5) Suppose otherwise  $Out(v) = Out(w) = \{v_1\}$ . Let  $E = \{v, w\}$ . Then we have  $|Out(E)| = |\{v_1\}| = 1 < 2 = |\{v, w\}| = |E \cap S(G_{\lambda_0})|$ , which contradicts to Lemma 9.

We give in the below an example and a counterexample of minimal repair graph that can be verified by Corollary 11. *Example 12:* Consider the repair graph  $G_{\lambda_0}$  in Fig. 3, where the vertex set is  $\mathcal{V} = \{1, 2, \dots, 15\}$ . We can check that  $|\text{Out}(E)| \ge |E \cap S(G_{\lambda_0})|$  for each  $E \subseteq [n]$  of size  $|E| \le t$ . Corresponding to items 1)–5) of Corollary 11, we can check:

- 1) For every  $v \in S(G_{\lambda_0})$ ,  $|Out(v)| \ge 1$ .
- 2) For v = 5 and v' = 10, we have  $Out(v) = \{v'\}$  and  $Out^2(v) = Out(v') = \{12, 13\}$ .
- 3) For v = 1,  $v_1 = 8$  and  $v_2 = 11$ , we have  $Out(v) = \{v_1\}$ ,  $Out(v_1) = \{v_2\}$  and  $Out(v_2) = \{14\}$ .
- 4) For v = 1,  $v_1 = 8$  and  $v_2 = 11$ , we have  $u = 6 \in In(v_2)$  is a source and  $|Out(u)| = |\{10, 11\}| \ge 2$ .
- 5) For v = 1 and w = 5, we have |Out(v)| = |Out(w)| = 1 and  $Out(v) = \{8\} \neq Out(w) = \{10\}$ .




Fig 3. An example repair graph  $G_{\lambda_0} = (\mathcal{V}, \mathcal{E})$ , where  $\mathcal{V} = \{1, 2, \cdots, 15\}$ .

*Example 13:* Any one of the following five observations, which violates the corresponding five cases of Lemma 11, can show that the graph in Fig. 1 is not a minimal repair graph.

1) For the source v = 1, we have  $Out(1) = \emptyset$ .

2) For the source v = 2, we have  $Out(2) = \{9\}$  and  $Out(9) = \emptyset$ .

3) For the source v = 5, we have  $Out(5) = \{11\}$ ,  $Out(11) = \{13\}$  and  $Out(13) = \emptyset$ .

4) For the source v = 6, we have  $Out(6) = \{12\}$ ,  $Out(12) = \{14\}$  and there is another source  $u = 8 \in In(14)$  such that  $Out(8) = \{14\}$ .

5) For the two sources v = 6 and w = 7, we have  $Out(6) = Out(7) = \{12\}$ .

*Remark 14:* In Corollary 11, claim 1) holds for all  $t \ge 1$ , since the contradiction is derived from a subset E of size 1. And claims 2), 5) hold for all  $t \ge 2$  since the contradictions are derived from subsets of size 2.

Corollary 15: Suppose  $t \ge 3$  and  $v \in S(G_{\lambda_0})$  such that  $Out(v) = \{v_1, v_2\}$ . Then the following hold:

1)  $\operatorname{Out}(v_1) \neq \emptyset$  or  $\operatorname{Out}(v_2) \neq \emptyset$ .

2) If  $\{v_1\} = \operatorname{Out}(u)$  for some source u, then  $\operatorname{Out}(v_2) \neq \emptyset$ .

3) If  $\{v_1\} = \operatorname{Out}(u)$  for some source u, then  $|\operatorname{Out}(w)| \ge 2$  for any source  $w \in \operatorname{In}(v_2)$ .

*Proof:* All the claims can be proved by assuming the converse and choosing a proper E as in the proof of Lemma 11 and then derive a contradiction.

1) Suppose otherwise  $\operatorname{Out}(v_1) = \operatorname{Out}(v_2) = \emptyset$ . We let  $E = \{v, v_1, v_2\}$  and have  $|\operatorname{Out}(E)| = |\emptyset| = 0 < 1 = |\{v\}| = |E \cap S(G_{\lambda_0})|$ , which contradicts to Lemma 9.

2) Suppose otherwise  $Out(v_2) = \emptyset$ . Similarly, we can get a contradiction by letting  $E = \{u, v, v_2\}$ .

3) Suppose otherwise there exist a source w such that  $Out(w) = \{v_2\}$ . A contradiction can be obtained by letting  $E = \{u, v, w\}$ .

We give in the below an example and a counterexample of minimal repair graph that can be verified by Corollary 15.

*Example 16:* Again consider the repair graph  $G_{\lambda_0}$  in Fig. 3. Let v = 3,  $v_1 = 8$  and  $v_2 = 9$ . Then  $v \in S(G_{\lambda_0})$  and  $Out(v) = \{v_1, v_2\}$ . Corresponding to items 1)-3) of Corollary 15, we can check:

1)  $\operatorname{Out}(v_1) = \operatorname{Out}(v_2) = \{11\} \neq \emptyset.$ 

2) For u = 1, we have  $\{v_1\} = Out(u)$  and  $Out(v_2) = \{11\} \neq \emptyset$ .

3) For w = 4, we can see that  $w \in In(v_2)$  is a source and  $|Out(w)| = \{9, 10, 12\}| \ge 2$ .

*Example 17:* Let G be a repair graph as shown in Fig. 4. Then any one of the following three observations, which violates the corresponding three cases of Corollary 15, can show that G is not a minimal repair graph.

1) There exists a source v = 5 such that  $Out(v) = \{9, 10\}$  and  $Out(9) = Out(10) = \emptyset$ .

2) There exists a source v = 2 such that  $Out(2) = \{7, 8\}$ , and a source u = 1 such that  $Out(1) = \{7\}$  and  $Out(8) = \emptyset$ .

3) There exist three sources v = 2, u = 1 and w = 3 such that  $Out(2) = \{7, 8\}$ ,  $Out(1) = \{7\}$  and  $Out(3) = \{8\}$ .

# B. Upper Bound on the Code Rate for (n, k, r, 3)-SLRC

In this subsection, we assume C is an (n, k, r, 3)-SLRC and  $G_{\lambda_0} = (\mathcal{V}, \mathcal{E})$  is a minimal repair graph of C. Recall that  $S(G_{\lambda_0})$  is the set of all sources of  $G_{\lambda_0}$ . We divide  $S(G_{\lambda_0})$  into four subsets as follows.

$$A = \{ v \in \mathbf{S}(G_{\lambda_0}); |\operatorname{Out}(v)| \ge 3 \},$$
(11)

$$B = \{ v \in \mathbf{S}(G_{\lambda_0}); |\operatorname{Out}(v)| = 2 \},$$
(12)




Fig 4. An example repair graph with n = 12 and r = 2.

$$C_1 = \{ v \in \mathbf{S}(G_{\lambda_0}); |\mathsf{Out}(v)| = 1 \text{ and } |\mathsf{Out}^2(v)| = 1 \}$$
(13)

and

$$C_2 = \{ v \in \mathbf{S}(G_{\lambda_0}); |\operatorname{Out}(v)| = 1 \text{ and } |\operatorname{Out}^2(v)| \ge 2 \}.$$
(14)

Clearly,  $A, B, C_1$  and  $C_2$  are mutually disjoint. Moreover, by 1), 2) of Corollary 11,  $S(G_{\lambda_0}) = A \cup B \cup C_1 \cup C_2$ . Hence,

$$\delta^* = |\mathbf{S}(G_{\lambda_0})| = |A| + |B| + |C_1| + |C_2|.$$
(15)

We define three types of edges of  $G_{\lambda_0}$ , denoted by red edge, green edge and blue edge respectively, as follows.

Firstly, an edge is called a *red edge* if its tail is a source. For each  $v \in S(G_{\lambda_0})$ , let  $\mathcal{E}_{red}(v)$  be the set of all red edges whose tail is v and denote

$$\mathcal{E}_{\mathrm{red}} = \bigcup_{v \in \mathbf{S}(G_{\lambda_0})} \mathcal{E}_{\mathrm{red}}(v)$$

Then  $\mathcal{E}_{red}$  is the set of all red edges. Clearly,  $|\mathcal{E}_{red}(v)| = |Out(v)|$  and  $\mathcal{E}_{red}(w) \cap \mathcal{E}_{red}(v) = \emptyset$  for any source  $w \neq v$ . So by (11)–(14), we have

$$|\mathcal{E}_{\text{red}}| = \sum_{v \in \mathcal{S}(G_{\lambda_0})} |\text{Out}(v)| \ge 3|A| + 2|B| + |C_1| + |C_2|.$$
(16)

Secondly, an edge is called a *green edge* if its tail is the unique out-neighbor of some source in  $C_1 \cup C_2$ . For each  $v \in C_1 \cup C_2$ , let  $\mathcal{E}_{green}(v)$  be the set of all green edges whose tail is the unique out-neighbor of v. Clearly,  $|\mathcal{E}_{green}(v)| = |\operatorname{Out}^2(v)|$ . Let

$$\mathcal{E}_{\text{green}} = \bigcup_{v \in C_1 \cup C_2} \mathcal{E}_{\text{green}}(v)$$

be the set of all green edges. Note that if  $v \neq w \in C_1 \cup C_2$ , then by 5) of Corollary 11,  $v' \neq w'$ , where v'(resp. w') is the unique out-neighbor of v(resp. w). So  $\mathcal{E}_{\text{green}}(v) \cap \mathcal{E}_{\text{green}}(w) = \emptyset$ . Hence, by (13) and (14),

$$|\mathcal{E}_{\text{green}}| = \sum_{v \in C_1 \cup C_2} |\text{Out}^2(v)| \ge |C_1| + 2|C_2|.$$
(17)

Thirdly, suppose  $e \in \mathcal{E}$  is not a green edge and  $v \in B \cup C_1$ . *e* is called a *blue edge belonging to* v if one of the following two conditions hold:

(a)  $v \in B$  and the tail of e belongs to Out(v).

(b)  $v \in C_1$  and the tail of e belongs to  $Out^2(v)$ .

Let  $\mathcal{E}_{blue}(v)$  be the set of all blue edges belonging to v and let

$$\mathcal{E}_{\text{blue}} = \bigcup_{v \in B \cup C_1} \mathcal{E}_{\text{blue}}(v)$$

be the set of all blue edges. Then we have the following lemma.

Lemma 18: The number of blue edges is lower bounded by

$$|\mathcal{E}_{\text{blue}}| \ge \frac{|B| + |C_1|}{r}.$$
(18)

*Proof:* It is sufficient to prove : i) For each  $v \in B \cup C_1$ , there exists at least one blue edge belonging to v; and ii) Each blue edge belongs to at most r different  $v \in B \cup C_1$ . To prove these two statements, we will use the definition of red edge, green edge and blue edge repeatedly.

We first prove i) by considering the cases of  $v \in B$  and  $v \in C_1$ .

Let  $v \in B$ , and we look for a blue edge belonging to v. In this case, by (12), we can assume  $Out(v) = \{v_1, v_2\}$  (see Fig. 5(a)). Then, by 1) of Corollary 15,  $Out(v_1) \neq \emptyset$  or  $Out(v_2) \neq \emptyset$ . Without loss of generality, assume  $Out(v_1) \neq \emptyset$  and  $v_3 \in Out(v_1)$ . Consider  $(v_1, v_3)$ . If it is not a green edge, then by definition, it is a blue edge belonging to v. So we assume

that  $(v_1, v_3)$  is a green edge. Then by definition,  $\{v_1\} = \text{Out}(u)$  for some  $u \in C_1 \cup C_2$ . By 2) of Corollary 15,  $\text{Out}(v_2) \neq \emptyset$ and we can let  $v_4 \in \text{Out}(v_2)$ , as illustrated in Fig. 5(a). Consider  $(v_2, v_4)$ . By 3) of Corollary 15,  $|\text{Out}(w)| \ge 2$  for any source  $w \in \text{In}(v_2)$ , which implies  $(v_2, v_4)$  is not a green edge. So  $(v_2, v_4)$  is a blue edge belonging to v. Hence, for each  $v \in B$ , we can always find a blue edge belonging to v.




Fig 5. Two local graphs.

Now, let  $v \in C_1$  and we look for a blue edge belonging to v. By (13), we can assume  $Out(v) = \{v_1\}$  and  $Out^2(v) = \{v_2\}$  (see Fig.5(b)). By 3) of Corollary 11, we have  $Out(v_2) \neq \emptyset$ . Let  $v_3 \in Out(v_2)$ . Note that by 4) of Corollary 11,  $|Out(u)| \ge 2$  for any source  $u \in In(v_2)$  (see Fig. 5(b) as illustration), which implies  $(v_2, v_3)$  is not a green edge. So  $(v_2, v_3)$  is a blue edge belonging to v. Hence, for each  $v \in C_1$ , we can always find a blue edge belonging to v.

By the above discussion, statement i) holds.

Let (u', u'') be a blue edge and S be the set of all  $v \in B \cup C_1$  such that (u', u'') belongs to v. To prove statement ii), we prove that there is an *injection*, namely  $\varphi$ , from S to  $\operatorname{In}(u')$ . Then ii) follows from the fact that  $\operatorname{In}(u')$  has size at most r. The injection of  $\varphi(v)$  can be constructed as follows: If  $v \in B$ , simply let  $\varphi(v) = v$ . If  $v \in C_1$ , let  $\varphi(v) = v'$ , where  $\{v'\} = \operatorname{Out}(v)$ . It is easy to see that  $\varphi(v)$  is an injection (noticing 5) of Corollary 11), which completes the proof of statement ii).

*Example 19:* Consider the repair graph in Fig. 3. We have  $A = \{2, 4, 7\}$ ,  $B = \{3, 6\}$ ,  $C_1 = \{1\}$  and  $C_2 = \{5\}$ , and the edges with tails from 1 to 7 are red edges, as illustrated in Fig. 6.

Moreover, one can check that  $\mathcal{E}_{\text{green}}(1) = \{(8, 11)\}$  and  $\mathcal{E}_{\text{green}}(5) = \{(10, 12), (10, 13)\}$ . As for blue edges, since  $1 \in C_1$  and  $11 \in \text{Out}^2(1)$ , then  $(11, 14) \in \mathcal{E}_{\text{blue}}(1)$ ; Since  $11 \in \text{Out}(6)$  and  $6 \in B$ , then  $(11, 14) \in \mathcal{E}_{\text{blue}}(6)$ ; Since  $3 \in B$  and  $9 \in \text{Out}(3)$ , then  $(9, 11) \in \mathcal{E}_{\text{blue}}(3)$ . One can check that  $\mathcal{E}_{\text{blue}}(1) = \mathcal{E}_{\text{blue}}(6) = \{(11, 14)\}$  and  $\mathcal{E}_{\text{blue}}(3) = \{(9, 11)\}$ . The green edges and blue edges are also illustrated in Fig. 6.




Fig 6. Illustration of red edge, green edge and blue edge of minimal repair graph.

Now, we can propose our main theorem of this section. *Theorem 20:* For (n, k, r, 3)-SLRC, we have <sup>2</sup>

$$\frac{k}{n} \le \left(\frac{r}{r+1}\right)^2. \tag{19}$$

<sup>2</sup>In the original version [18] of this paper, bound (19) was presented equivalently in terms of the code length as  $n \ge k + \left\lfloor \frac{2k + \lceil \frac{k}{r} \rceil}{r} \right\rfloor$ .

*Proof:* By definition, we can easily see that  $\mathcal{E}_{red}$ ,  $\mathcal{E}_{green}$  and  $\mathcal{E}_{blue}$  are mutually disjoint. Then by (15)-(18), we have

$$\begin{split} |\mathcal{E}| &\geq |\mathcal{E}_{\rm red}| + |\mathcal{E}_{\rm green}| + |\mathcal{E}_{\rm blue}| \\ &\geq (3|A| + 2|B| + |C_1| + |C_2|) \\ &+ (|C_1| + 2|C_2|) + \frac{|B| + |C_1|}{r} \\ &= 2(|A| + |B| + |C_1| + |C_2|) \\ &+ (|A| + |C_2| + \frac{|B| + |C_1|}{r}) \\ &= 2\delta^* + \frac{r|A| + r|C_2| + |B| + |C_1|}{r} \\ &\geq 2\delta^* + \frac{|A| + |C_2| + |B| + |C_1|}{r} \\ &\geq 2\delta^* + \frac{\delta^*}{r}. \end{split}$$

That is,  $|\mathcal{E}| \ge 2\delta^* + \frac{\delta^*}{r}$ . Combining this with Lemma 7, we have

$$(n-\delta^*)r \ge |\mathcal{E}| \ge 2\delta^* + \frac{\delta^*}{r}.$$

So

$$(n-\delta^*)r \ge 2\delta^* + \frac{\delta^*}{r}$$

Solving n from the above equation, we have

$$n \ge \delta^* + \frac{2\delta^* + \frac{\delta^*}{r}}{r}.$$
(20)

By Lemma 8,  $\delta^* \ge k$ . So (20) implies that

$$a \ge k + \frac{2k + \frac{\kappa}{r}}{r}$$
$$= k \left( 1 + \frac{2}{r} + \frac{1}{r^2} \right)$$
$$= k \left( \frac{r+1}{r} \right)^2.$$

Hence,

 $\frac{k}{n} \le \left(\frac{r}{r+1}\right)^2,$ 

r

which proves the theorem.

We will later construct two families of (n, k, r, 3)-SLRCs achieving (19) and hence show the tightness of this bound.

## C. Code Rate for (n, k, r, 2)-SLRC

In this subsection, we give a new proof of the bound (1) for the (n, k, r, 2)-SLRC using the similar techniques as in Subsection B. Assume that C is an (n, k, r, 2)-SLRC and  $G_{\lambda_0} = (\mathcal{V}, \mathcal{E})$  is a minimal repair graph of C.

*Proof of Bound* (1): By Remark 14 and 1) of Corollary 11, each source of  $G_{\lambda_0}$  has at least one out-neighbor. Let A be the set of sources that has only one out-neighbor and let  $\mathcal{E}_{red}$  be the set of all edges e, called red edges, such that the tail of e is a source. Then the number of red edges is

$$|\mathcal{E}_{\text{red}}| \ge |A| + 2|\mathbf{S}(G_{\lambda_0}) \setminus A| = 2\delta^* - |A|.$$
<sup>(21)</sup>

For each  $v \in A$ , let v' be the unique out-neighbor of v and  $\mathcal{E}_{green}(v)$  be the set of all edges whose tail is v'. By Remark 14 and 2) of Corollary 11,  $\operatorname{Out}^2(v) = \operatorname{Out}(v') \neq \emptyset$ . So  $|\mathcal{E}_{green}(v)| = |\operatorname{Out}(v')| \ge 1$ . Let  $\mathcal{E}_{green}$  be the set of all green edges. For any two different  $v_1, v_2 \in A$ , let  $v'_1, v'_2$  be the unique out-neighbor of  $v_1, v_2$ , respectively. By Remark 14 and 5) of Corollary 11,  $v'_1 \neq v'_2$ . So  $\mathcal{E}_{green}(v_1) \cap \mathcal{E}_{green}(v_2) = \emptyset$ . Hence,

$$|\mathcal{E}_{\text{green}}| = \left| \bigcup_{v \in A} \mathcal{E}_{\text{green}}(v) \right| = \sum_{v \in A} |\mathcal{E}_{\text{green}}(v)| \ge |A|.$$
(22)

Clearly,  $\mathcal{E}_{red} \cap \mathcal{E}_{green} = \emptyset$ . Then by (21) and (22),

$$|\mathcal{E}| \ge |\mathcal{E}_{\text{red}}| + |\mathcal{E}_{\text{green}}| \ge 2\delta^*.$$

On the other hand, by Lemma 7,

$$(n-\delta^*)r \ge |\mathcal{E}|.$$

So  $(n - \delta^*)r \ge 2\delta^*$ , which implies  $n \ge \delta^* + \frac{2\delta^*}{r} \ge k + \frac{2k}{r}$  (Lemma 8). Hence,  $\frac{k}{n} \le \frac{r}{r+2}$ , which completes the proof.




Fig 7. (a) The index set of  $\mathbb{Z}_3^3$ , where  $(i_3, i_2, i_1)$  is simply written as  $i_3i_2i_1$ ; (b) The recovering sets  $R_1, \dots, R_7$  of  $\alpha_1, \dots, \alpha_7$ , where  $R_1 = \{(000), (002)\}$ ,  $R_2 = \{(010), (011)\}$ ,  $R_3 = \{(020), (021)\}$ ,  $R_4 = \{(011), (211)\}$ ,  $R_5 = \{\alpha_2, (212)\}$ ,  $R_6 = \{(021), (221)\}$  and  $R_7 = \{\alpha_3, (222)\}$  drawn in heavy lines.

# IV. AN EXAMPLE OF SLRC

In order to have a better understanding of the binary (r, t)-SLRC constructed in the next section, we give an example in this section. Let r = 2, m = 3 and C be the product code of m copies of the binary [r + 1, r] single parity check code. Then C has length  $n = (r + 1)^3 = 27$  and dimension  $k = r^3 = 8$ . It is convenient to use  $\mathbb{Z}_3^3 = \{(i_3, i_2, i_1); i_1, i_2, i_3 \in \mathbb{Z}_3\}$  instead of [n] as the index set of the coordinates of C, and let  $\mathbb{Z}_2^3 = \{(i_3, i_2, i_1); i_1, i_2, i_3 \in \mathbb{Z}_3\}$  be the information set of C. Here,  $\mathbb{Z}_3 = \{0, 1, 2\}$  and  $\mathbb{Z}_2 = \{0, 1\}$  are simply viewed as two sets and  $\mathbb{Z}_2 \subseteq \mathbb{Z}_3$  (no algebraic meaning is considered here).

The index set  $\mathbb{Z}_3^3$  is depicted in Fig. 7(a). By definition, each code symbol (coordinate) of  $\mathcal{C}$  can be recovered by all the other symbols on the same (red, green or blue) line. Hence, each code symbol of  $\mathcal{C}$  has m = 3 disjoint recovering sets (red, green and blue) of size r = 2, and  $\mathcal{C}$  can recover any 3 erasures by parallel recovery. However, we can prove (*see details in the next section*) that it can recover any  $t = 2^m - 1 = 7$  erasures by sequential recovery. For example, consider an erasure of 7 code symbols, say,  $E = \{\alpha_1, \dots, \alpha_7\}$ , where  $\alpha_1 = (001)$ ,  $\alpha_2 = (012)$ ,  $\alpha_3 = (022)$ ,  $\alpha_4 = (111)$ ,  $\alpha_5 = (112)$ ,  $\alpha_6 = (121)$  and  $\alpha_7 = (122)$ , as illustrated in Fig. 7(b). We can select a sequence of recovering sets  $R_1 = \{(000), (002)\}$ ,  $R_2 = \{(010), (011)\}$ ,  $R_3 = \{(020), (021)\}$ ,  $R_4 = \{(011), (211)\}$ ,  $R_5 = \{\alpha_2, (212)\}$ ,  $R_6 = \{(021), (221)\}$  and  $R_7 = \{\alpha_3, (222)\}$  (see Fig. 7(b)). It is easy to check that  $R_1, \dots, R_7$  sequentially repair  $\{\alpha_1, \dots, \alpha_7\}$ .

In general, by puncturing C properly, we can obtain (r, t)-SLRC for any  $t \in \{1, 2, \dots, 6\}$ . As an example, we construct an (r, 5)-SLRC as follows. For each  $j \in \mathbb{Z}_2 = \{0, 1\}$ , let

$$A_j = \{(j, i_2, i_1); i_2, i_1 \in \mathbb{Z}_3\}$$

and let

$$A = A_0 \cup A_1$$

$$B = \{(2, i_2, i_1); i_2 \in \mathbb{Z}_2 \text{ and } i_1 \in \mathbb{Z}_3\} \\ = \{(200), (201), (202), (210), (211), (212)\} \}$$

Let  $\Omega = A \cup B$ , as depicted in Fig. 8. Then, the punctured code  $\mathcal{C}|_{\Omega}$  is an (n', k, r, 5)-SLRC, with  $n' = |\Omega| = 24$ . In fact, one can see that the following items hold.

- i) For any nonempty  $E \subseteq A$  of size  $|E| \le t_1 = 3$ , there exists an  $\alpha \in E$  such that  $\alpha$  has a recovering set  $R \subseteq A \setminus E$ .
- ii) For any nonempty  $E \subseteq A$  of size  $|E| \le t = 5$ , there exists an  $\alpha \in E$  such that  $\alpha$  has a recovering set  $R \subseteq \Omega \setminus E$ .
- iii) For any nonempty  $E \subseteq B$  of size  $|E| \le t_2 = 1$ , there exists an  $\alpha \in E$  such that  $\alpha$  has a recovering set  $R \subseteq B \setminus E$ .
- iv) For any nonempty  $E \subseteq B$  of size  $|E| \le t = 5$ , there exists an  $\alpha \in E$  such that  $\alpha$  has a recovering set  $R \subseteq \Omega \setminus E$ .

11

In the above, items i), iii), iv) can be easily verified. For example, one can see that the punctured codes  $C|_{A_0}$  and  $C|_{A_1}$  are both (r, 3)-SLRC and  $C|_B$  is a (r, 1)-SLRC, hence i) and iii) hold. From Fig. 8, one can see that each  $(2, i_2, i_1) \in B$  has a recovery set (red line)  $R = \{(0, i_2, i_1), (1, i_2, i_1)\} \subseteq A$ , hence iv) holds. To prove ii), we consider the following two cases:

- 1)  $E \subseteq A_0$  or  $E \subseteq A_1$ . Without loss of generality, assume  $E \subseteq A_1$ . If  $E \subseteq \{(120), (121), (122)\}$ , then each  $(1, 2, i) \in E$  has a recovery set (green line)  $R = \{(1, 0, i), (1, 1, i)\} \subseteq \Omega \setminus E$ ; Otherwise, there exists a  $(1, i_2, i_1) \in E \cap \{(100), (101), (102), (110), (111), (112)\}$  which has a recovery set (red line)  $R = \{(0, i_2, i_1), (2, i_2, i_1)\} \subseteq \Omega \setminus E$ .
- 2)  $E \cap A_0 \neq \emptyset$  and  $E \cap A_1 \neq \emptyset$ . Since  $|E| \leq 5$ , then  $|E \cap A_0| \leq 3$  or  $|E \cap A_1| \leq 3$ . Note that both  $C|_{A_0}$  and  $C|_{A_1}$  are (r, 3)-SLRCs, by Lemma 2, there exists an  $\alpha \in E$  and  $j \in \{0, 1\}$  such that  $\alpha$  has a recovering set  $R \subseteq A_j \setminus E \subseteq A \setminus E \subseteq \Omega \setminus E$ .

Then, by Lemma 3,  $C|_{\Omega}$  is an (n', k, r, 5)-SLRC. The generalization of this example as well as the formal proof will be given in the next section.




Fig 8. Graphical illustration of a subset of  $\mathbb{Z}_3^3$ .

# V. CONSTRUCTION OF (n, k, r, t)-SLRC

In this section, we construct a family of binary (n, k, r, t)-SLRC for any positive integers  $r (\geq 2)$  and t. It will be shown that the code rate of this family is greater than  $\frac{r}{r+t}$ , and in particular, for  $t \in \{2, 3\}$ , it achieves the bounds (1) and (19), respectively.

We first need introduce some notations. For any positive integers r and m, where  $r \ge 2$ , let  $\mathbb{Z}_r = \{0, 1, \dots, r-1\}$  and  $\mathbb{Z}_r^m = \{(i_m, i_{m-1}, \dots, i_1); i_m, i_{m-1}, \dots, i_1 \in \mathbb{Z}_r\}$ . Here,  $\mathbb{Z}_r$  is simply viewed as a set (without any algebraic meaning). So  $\mathbb{Z}_r \subseteq \mathbb{Z}_{r+1} = \{0, 1, \dots, r-1, r\}$ . We will use  $\alpha, \beta, \gamma$ , etc, to denote elements (points) of  $\mathbb{Z}_{r+1}^m$ . Note that by the notation, for each  $\alpha = (i_m, i_{m-1}, \dots, i_1) \in \mathbb{Z}_r$  and  $\ell \in [m]$ ,  $i_\ell$  is the  $\ell$ th coordinate of  $\alpha$  from the right.

For each  $\alpha = (i_m, i_{m-1}, \dots, i_1) \in \mathbb{Z}_{r+1}^m$ , we let

$$\mathbf{U}^{(m)}(\alpha) = \{\ell \in [m]; i_{\ell} = r\},\tag{23}$$

and

$$\mathbf{T}^{(m)}(\alpha) = \{\ell \in [m]; i_{\ell} \in \mathbb{Z}_r\}.$$
(24)

Further, we let

$$\mathcal{L}^{(m)}(\alpha) = \{ (j_m, j_{m-1}, \cdots, j_1) \in \mathbb{Z}_r^m; j_\ell = i_\ell, \forall \ell \in \mathbf{T}^{(m)}(\alpha) \}.$$
(25)

Clearly,  $\mathbf{U}^{(m)}(\alpha) \cap \mathbf{T}^{(m)}(\alpha) = \emptyset$  and  $\mathbf{U}^{(m)}(\alpha) \cup \mathbf{T}^{(m)}(\alpha) = [m]$ . Moreover, for each  $\alpha \in \mathbb{Z}_{r+1}^m \setminus \mathbb{Z}_r^m$ ,  $\mathbf{U}^{(m)}(\alpha) \neq \emptyset$  and  $\mathcal{L}^{(m)}(\alpha) \neq \emptyset$ . In particular, if  $\alpha = (r, r, \cdots, r)$ , then  $\mathbf{U}^{(m)}(\alpha) = [m]$  and  $\mathcal{L}^{(m)}(\alpha) = \mathbb{Z}_r^m$ .

As an example, let r = 2, m = 6 and  $\alpha = (1, 0, 2, 1, 1, 2) \in \mathbb{Z}_3^6$ . Then  $\mathbf{U}^{(m)}(\alpha) = \{4, 1\}, \mathbf{T}^{(m)}(\alpha) = \{6, 5, 3, 2\}$  and  $\mathcal{L}^{(m)}(\alpha) = \{(1, 0, i_4, 1, 1, i_1); i_4, i_1 \in \mathbb{Z}_2\} = \{(1, 0, 0, 1, 1, 0), (1, 0, 0, 1, 1, 1), (1, 0, 1, 1, 1, 0), (1, 0, 1, 1, 1, 1)\}.$ 

Note that for any integer s such that  $0 \le s \le 2^m - 1$ , s has a unique *m*-digit binary representation, say  $(\lambda_m \lambda_{m-1} \cdots \lambda_1)$ . That is,  $(\lambda_m, \lambda_{m-1}, \cdots, \lambda_1) \in \{0, 1\}^m$  and  $s = \sum_{\ell=1}^m \lambda_\ell 2^{\ell-1}$ . Denote by  $\operatorname{supp}_m(s)$  the support of  $(\lambda_m, \lambda_{m-1}, \cdots, \lambda_1)$ . Let

$$\Gamma_s^{(m)} = \{ \alpha \in \mathbb{Z}_{r+1}^m; \mathbf{U}^{(m)}(\alpha) = \operatorname{supp}_m(s) \}$$
(26)

and

$$\Omega_s^{(m)} = \bigcup_{\ell=0}^s \Gamma_\ell^{(m)}.$$
(27)

For example, suppose r = 2, m = 6 and s = 22. Then (010110) is the unique 6-digit binary representation of s and

For example, suppose r = 2, m = 0 and s = 22. Then (010110) is the unique o digit charge representation of r = 1, m = 0 and s = 22. Then (010110) is the unique o digit charge representation of r = 1,  $supp_m(s) = \{5, 3, 2\}$ . From (26), we have  $\Gamma_{22}^{(6)} = \{(i_6, 2, i_4, 2, 2, i_1); i_6, i_4, i_1 \in \mathbb{Z}_2\}$ . Clearly,  $\Gamma_0^{(m)}, \Gamma_1^{(m)}, \cdots, \Gamma_{2m-1}^{(m)}$  are mutually disjoint and  $|\Gamma_s^{(m)}| = r^{m-|supp_m(s)|}$ , for  $s = 0, 1, \cdots, 2^m - 1$ . In particular,  $|\Gamma_0^{(m)}| = r^m$  and  $|\Gamma_{2m-1}^{(m)}| = 1$ . Moreover, by definition, we have  $\Omega_0^{(m)} = \Gamma_0^{(m)} = \mathbb{Z}_r^m$  and  $\Omega_{2m-1}^{(m)} = \mathbb{Z}_{r+1}^m$ . For any positive integers  $r \ (\geq 2)$  and t, we can always pick an integer m such that  $t \le 2^m - 1$  and, using the above notations, define a matrix  $H_t^{(m)} = (h_{\alpha,\beta})$  satisfying the following two properties.

(1) The rows of  $H_t^{(m)}$  are indexed by  $\Omega_t^{(m)} \setminus \Omega_0^{(m)}$  and the columns of  $H_t^{(m)}$  are indexed by  $\Omega_t^{(m)}$ ; (2) For each  $\alpha \in \Omega_t^{(m)} \setminus \Omega_0^{(m)}$  and  $\beta \in \Omega_t^{(m)}$ ,

$$h_{\alpha,\beta} = \begin{cases} 1, & \text{if } \beta \in \mathcal{L}^{(m)}(\alpha) \cup \{\alpha\};\\ 0, & \text{Otherwise.} \end{cases}$$
(28)

It should be noted that,  $H_t^{(m)} = (h_{\alpha,\beta})$  is an  $h \times n$  binary matrix, where  $h = \left|\Omega_t^{(m)} \setminus \Omega_0^{(m)}\right|$ ,  $n = \left|\Omega_t^{(m)}\right|$ . The sub-matrix of  $H_t^{(m)}$ , formed by the columns indexed by  $\Omega_t^{(m)} \setminus \Omega_0^{(m)}$ , is a permutation matrix. Hence, rank  $\left(H_t^{(m)}\right) = h$ .

Theorem 21: Let  $\mathcal{C}_t^{(m)}$  be the binary code that has a parity check matrix  $H_t^{(m)}$ . Then  $\mathcal{C}_t^{(m)}$  is an (n, k, r, t)-SLRC with

$$n = r^m \sum_{s=0}^t \frac{1}{r^{|\text{supp}_m(s)|}}$$
(29)

and

$$k = r^m. ag{30}$$

Hence, the code rate of  $C_t^{(m)}$  is

$$\frac{k}{n} = \frac{1}{\sum_{s=0}^{t} \frac{1}{r^{|\text{supp}_m(s)|}}},\tag{31}$$

where  $r (\geq 2)$  and t are any positive integers and m is any integer satisfying  $t \leq 2^m - 1$ .

Remark 22: We have some remarks about the construction.

- 1) The example codes given in the last section are just  $C_t^{(m)}$  for r = 2, m = 3 and t = 7,5 respectively. In general, for  $t = 2^m 1$ , it is easy to check that  $\Omega_{2^m-1}^{(m)} = \mathbb{Z}_{r+1}^m$  and  $C_{2^m-1}^{(m)}$  is the product of m copies of the [r+1,r] binary code. If  $t < 2^m 1$ , then  $C_t^{(m)}$  is the punctured code of  $C_{2^m-1}^{(m)}$  with respect to  $\Omega_t^{(m)}$ .
- 2) For  $t \in \{2,3\}$ , we can let m = 2 and from (31), the code rates of our construction are  $\frac{r}{r+2}$  and  $\left(\frac{r}{r+1}\right)^2$  respectively, which are optimal according to (1) and (19). For  $t \ge 4$ , by (31), the code rate of  $\mathcal{C}_t^{(m)}$  is higher than  $\frac{r}{r+t}$  for all  $r \ge 2$ .
- 3) It was shown in [10] that  $\mathcal{C}_{2^m-1}^{(m)}$  has locality r and availability m, which implies that it can recover m erasures with locality r using the parallel approach. In contrast, by Theorem 21, it can recover  $t = 2^m - 1$  erasures with the same locality when using the sequential approach, which is a significant advantage of the product code for the sequential recovery. In particular, the product of two copies of the [r+1, r] binary code is not optimal (in rate) among codes with locality r and availability t = 2 [12], but optimal among (r, t = 3)-SLRCs.

In the rest of this section, we will prove Theorem 21. To prove that  $C_t^{(m)}$  is an (r, t)-SLRC, we will prove a more general claim, say, for any binary linear code C, if C has a parity check matrix H which contains all rows of  $H_t^{(m)}$  (not necessarily  $H = H_t^{(m)}$ ), then C is an (r, t)-SLRC. We first make some clarifications on the construction by two simple remarks.

*Remark 23:* Let C be a binary linear code. If the code symbols of C are indexed by  $\Omega_t^{(m)}$ , then, by construction of  $H_t^{(m)}$ , C has a parity check matrix which contains all rows of  $H_t^{(m)}$  if and only if for each  $\alpha \in \Omega_t^{(m)} \setminus \Omega_0^{(m)}$ ,

$$x_{\alpha} = \sum_{\beta \in \mathcal{L}^{(m)}(\alpha)} x_{\beta}.$$
(32)

If the code symbols of  $\mathcal{C}$  are indexed by S, where  $S \neq \Omega_t^{(m)}$ , then  $\mathcal{C}$  has a parity check matrix which contains all rows of  $H_t^{(m)}$  if and only if there is a bijection  $\psi: \Omega_t^{(m)} \to S$  such that for each  $\alpha \in \Omega_t^{(m)} \setminus \Omega_0^{(m)}$ ,

$$x_{\psi(\alpha)} = \sum_{\beta \in \mathcal{L}^{(m)}(\alpha)} x_{\psi(\beta)}.$$
(33)

*Remark 24:* Since  $1 \le t \le 2^m - 1$ , we can find a  $m_0 \in [m]$  such that  $2^{m_0-1} - 1 < t \le 2^{m_0} - 1$ . Let  $t_1 = 2^{m_0-1} - 1$  and  $t_2 = t - t_1 - 1$ . Then  $0 \le t_2 \le t_1 \le 2^{m-1} - 1$  and  $\Omega_t^{(m)}$  can be partitioned into two disjoint nonempty subsets

$$A = \Omega_{t_1}^{(m)} = \bigcup_{s=0}^{t_1} \Gamma_s^{(m)}$$

and

$$B = \Omega_t^{(m)} \backslash A = \bigcup_{s=t_1+1}^t \Gamma_s^{(m)}.$$

Moreover, noticing that  $supp_m(s) \subseteq \{1, 2, \dots, m_0 - 1\}$  for  $0 \leq s \leq t_1$ , then A can be partitioned into r mutually disjoint nonempty subsets, according to the values of the  $m_0$ th coordinate (from the right) of its elements, as follows.

$$A_j = \{(i_m, i_{m-1}, \cdots, i_1) \in A; \ i_{m_0} = j\}, \ \forall j \in \mathbb{Z}_r.$$

In the following, if there is no other specification, we always assume that C is a binary linear code and has a parity check matrix which contains all rows of  $H_t^{(m)}$ . Without loss of generality, we assume that the code symbols of C are indexed by  $\Omega_t^{(m)}$ . To prove Theorem 21, we need the following three lemmas.

- Lemma 25: Suppose m > 1. With notations in Remark 24, the following hold.
- 1) For each  $j \in \mathbb{Z}_r$ , the punctured code  $\mathcal{C}|_{A_j}$  has a parity check matrix which contains all rows of  $H_{t_1}^{(m-1)}$ .
- 2) If  $t_2 \ge 1$ , the punctured code  $\mathcal{C}|_B$  has a parity check matrix which contains all rows of  $H_{t_2}^{(m-1)}$ .

*Proof:* For each  $j \in \mathbb{Z}_{r+1} = \{0, 1, \cdots, r\}$ , let

$$\psi_j: \mathbb{Z}_{r+1}^{m-1} \to \mathbb{Z}_{r+1}^m$$

be such that  $\psi_j(\alpha) = (i_{m-1}, \dots, i_{m_0}, j, i_{m_0-1}, \dots, i_1)$  for each  $\alpha \in (i_{m-1}, \dots, i_{m_0}, i_{m_0-1}, \dots, i_1) \in \mathbb{Z}_{r+1}^{m-1}$ . That is,  $\psi_j(\alpha)$  is obtained by inserting j as a coordinate between the  $(m_0-1)$ th and  $m_0$ th coordinate (from the right) of  $\alpha$ .

1) For each  $j \in \mathbb{Z}_r$ , it is a mechanical work to check that  $\psi_j$  induces a bijection between  $\Omega_{t_1}^{(m-1)}$  and  $A_j$  such that for each  $\alpha \in \Omega_{t_1}^{(m-1)} \setminus \Omega_0^{(m-1)}$ ,

$$\mathcal{L}^{(m)}(\psi_j(\alpha)) = \{\psi_j(\beta); \beta \in \mathcal{L}^{(m-1)}(\alpha)\}$$

Since C has a parity check matrix containing all rows of  $H_t^{(m)}$ , then by (32), we have

$$x_{\psi_j(\alpha)} = \sum_{\substack{\beta' \in \mathcal{L}^{(m)}(\psi_j(\alpha))\\}} x_{\beta'}$$
$$= \sum_{\substack{\beta \in \mathcal{L}^{(m-1)}(\alpha)}} x_{\psi_j(\beta)}.$$

Hence, by Remark 23,  $C|_{A_j}$  has a parity check matrix which contains all rows of  $H_{t_1}^{(m-1)}$ . 2) Recall that  $t_1 = 2^{m_0-1} - 1$ . Then for each  $s \in \{t_1 + 1, t_1 + 2, \dots, t\}$ , we have

$$\operatorname{supp}_m(s) = \operatorname{supp}_{m-1}(s') \cup \{m_0\},\$$

where  $s' = s - 2^{m_0-1} \in \{0, 1, \dots, t_2\}$ . So similar to 1), we can check that  $\psi_r$  induces a bijection between  $\Omega_{t_2}^{(m-1)}$  and  $B = \Omega_t^{(m)} \setminus \Omega_{t_1}^{(m)}$  such that for each  $\alpha \in \Omega_{t_2}^{(m-1)} \setminus \Omega_0^{(m-1)}$ ,

$$x_{\psi_r(\alpha)} = \sum_{\beta \in \mathcal{L}^{(m-1)}(\alpha)} x_{\psi_r(\beta)}.$$

Hence, by Remark 23,  $C|_B$  has a parity check matrix which contains all rows of  $H_{t_2}^{(m-1)}$ .

For each  $\alpha = (i_m, i_{m-1}, \dots, i_1) \in \mathbb{Z}_{r+1}^m$  and  $\ell \in [m]$ , let

$$L_{\alpha}^{(\ell)} = \{ (i_m, \cdots, i_{\ell+1}, i'_{\ell}, i_{\ell-1}, \cdots, i_1); i'_{\ell} \in \mathbb{Z}_{r+1} \}.$$
(34)

That is,  $L_{\alpha}^{(\ell)}$  consists of  $\alpha$  as well as the points in  $\mathbb{Z}_{r+1}^m$  which differs from  $\alpha$  only at the  $\ell$ th coordinate (from the right). Lemma 26: For each  $\alpha \in \Omega_t^{(m)}$  and  $\ell \in [m]$ , if  $L_{\alpha}^{(\ell)} \subseteq \Omega_t^{(m)}$ , then  $R = L_{\alpha}^{(\ell)} \setminus \{\alpha\}$  is a recovering set of  $\alpha$ .

Proof: Let

$$\alpha = (i_m, \cdots, i_{\ell+1}, i_\ell, i_{\ell-1}, \cdots, i_1)$$

Then by (34),

$$L_{\alpha}^{(\ell)} = \{\alpha_0, \alpha_1, \cdots, \alpha_r\},\$$

where  $\alpha_j = (i_m, \cdots, i_{\ell+1}, j, i_{\ell-1}, \cdots, i_1)$  for each  $j \in \mathbb{Z}_{r+1}$  and  $\alpha = \alpha_{i_\ell}$ . From (25), it is easy to see that

$$\mathcal{L}^{(m)}(\alpha_r) = \bigcup_{j=0}^{r-1} \mathcal{L}^{(m)}(\alpha_j)$$
(35)

and for distinct  $j_1, j_2 \in \mathbb{Z}_r$ ,

$$\mathcal{L}^{(m)}(\alpha_{j_1}) \cap \mathcal{L}^{(m)}(\alpha_{j_1}) = \emptyset.$$
(36)

So combining (32), (35) and (36), we have

$$x_{\alpha_r} = \sum_{\beta \in \mathcal{L}^{(m)}(\alpha_r)} x_{\beta}$$
$$= \sum_{j=0}^{r-1} \left( \sum_{\beta \in \mathcal{L}^{(m)}(\alpha_j)} x_{\beta} \right)$$
$$= \sum_{j=0}^{r-1} x_{\alpha_j}$$

which is equivalent to (noticing that C is a binary code)

$$x_{\alpha} = \sum_{\beta \in L_{\alpha}^{(\ell)} \setminus \{\alpha\}} x_{\beta}$$

Note that from (34),  $L_{\alpha}^{(\ell)}$  has size r + 1. So  $R = L_{\alpha}^{(\ell)} \setminus \{\alpha\}$  has size r, hence is a recovering set of  $\alpha$ . Lemma 27: For any nonempty  $E \subseteq \Omega_t^{(m)} \setminus \Omega_0^{(m)}$ , there exists an  $\alpha \in E$  which has a recovering set  $R \subseteq \Omega_t^{(m)} \setminus E$ . Proof: Let s be the smallest number such that  $E \cap \Gamma_s^{(m)} \neq \emptyset$ . Since  $E \subseteq \Omega_t^{(m)} \setminus \Omega_0^{(m)}$ , then  $s \ge 1$  and  $\operatorname{supp}_m(s) \neq \emptyset$ . Hence, we can always find a  $\ell \in \operatorname{supp}_m(s)$  and a s' < s such that

$$\operatorname{supp}_m(s) = \operatorname{supp}_m(s') \cup \{\ell\}. \tag{37}$$

Pick  $\alpha \in E \cap \Gamma_s^{(m)}$ . Then by (26),  $\mathbf{U}^{(m)}(\alpha) = \operatorname{supp}_m(s)$ . Further, by (34) and (37),  $\mathbf{U}^{(m)}(\beta) = \operatorname{supp}_m(s')$  for each  $\beta \in L_{\alpha}^{(\ell)} \setminus \{\alpha\}$ . Then again by (26), we have

$$L_{\alpha}^{(\ell)} \setminus \{\alpha\} \subseteq \Gamma_{s'}^{(m)}. \tag{38}$$

Since s' < s and s is the smallest number such that  $E \cap \Gamma_s^{(m)} \neq \emptyset$ , then  $E \cap \Gamma_{s'}^{(m)} = \emptyset$ . Hence,

$$L_{\alpha}^{(\ell)} \setminus \{\alpha\} \subseteq \Gamma_{s'}^{(m)} \setminus E \subseteq \Omega_{s'}^{(m)} \setminus E \subseteq \Omega_t^{(m)} \setminus E,$$

and by Lemma 26,  $R = L_{\alpha}^{(\ell)} \setminus \{\alpha\}$  is a recovering set of  $\alpha$ .

Now, we can prove Theorem 21.

*Proof of Theorem 21:* By the construction, it is easy to see that the code length of  $\mathcal{C}_t^{(m)}$  is

$$\begin{split} n &= \left| \Omega_t^{(m)} \right| \\ &= \sum_{s=0}^t \left| \Gamma_s^{(m)} \right| \\ &= \sum_{s=0}^t r^{m-|\mathrm{supp}_m(s)|} \\ &= r^m \sum_{s=0}^t \frac{1}{r^{|\mathrm{supp}_m(s)|}}, \\ &k &= \left| \Omega_0^{(m)} \right| = r^m. \end{split}$$

and the dimension of  $C_t^{(m)}$  is

$$\frac{k}{n} = \frac{1}{\sum_{s=0}^{t} \frac{1}{r^{|\operatorname{supp}_m(s)|}}}$$

We then need to prove that  $\mathcal{C}_t^{(m)}$  is a (r, t)-SLRC. It is sufficient to prove that for any binary linear code  $\mathcal{C}$ , if  $\mathcal{C}$  has a parity

check matrix containing all rows of  $H_t^{(m)}$ , then C is an (r, t)-SLRC. We will prove this by induction on m. First, for m = 1, since  $1 \le t \le 2^m - 1$ , we have t = 1. By (26) and (27),  $\Gamma_0^{(1)} = \mathbb{Z}_r$ ,  $\Gamma_1^{(1)} = \{r\}$  and  $\Omega_1^{(1)} = \mathbb{Z}_{r+1}$ . So

$$H_1^{(1)} = (1, 1, \cdots, 1)_{1 \times (r+1)}.$$

Clearly, the binary linear code C with parity check matrix containing  $H_1^{(1)}$  is a (r, 1)-SLRC. Now, suppose m > 1 and the induction assumption holds for all m' < m and  $t' \le 2^{m'} - 1$ . We consider m and  $t \le 2^m - 1$ . Using the same notations as in Remark 24, we have the following four claims.

- For any nonempty  $E \subseteq A$  of size  $|E| \leq t_1$ , there exists an  $\alpha \in E$  such that  $\alpha$  has a recovering set  $R \subseteq A \setminus E$ . i)
- For any nonempty  $E \subseteq A$  of size  $|E| \leq t$ , there exists an  $\alpha \in E$  such that  $\alpha$  has a recovering set  $R \subseteq \Omega_t^{(m)} \setminus E$ . ii)
- For any nonempty  $E \subseteq B$  of size  $|E| \leq t_2$ , there exists an  $\alpha \in E$  such that  $\alpha$  has a recovering set  $R \subseteq B \setminus E$ . iii)
- For any nonempty  $E \subseteq B$  of size  $|E| \leq t$ , there exists an  $\alpha \in E$  such that  $\alpha$  has a recovering set  $R \subseteq \Omega_t^{(m)} \setminus E$ . iv)

We will prove them one by one as follows.

i): Since  $E \subseteq A$  and, by Remark 24,  $A = \bigcup_{j=0}^{r-1} A_j$ , then  $E \cap A_{j_0} \neq \emptyset$  for some  $j_0 \in \mathbb{Z}_r$ . By 1) of Lemma 25,  $\mathcal{C}|_{A_{j_0}}$  has a parity check matrix containing all rows of  $H_{t_1}^{(m-1)}$ . So by induction assumption,  $\mathcal{C}|_{A_{j_0}}$  is an  $(r, t_1)$ -SLRC. Moreover, since  $|E \cap A_{j_0}| \le |E| \le t_1$ , hence, by Lemma 2, there exists an  $\alpha \in E \cap A_{j_0}$  such that  $\alpha$  has a recovering set  $R \subseteq A_{j_0} \setminus E \subseteq A \setminus E$ .

ii): According to Remark 24,  $\{A_j; j \in \mathbb{Z}_r\}$  is a partition of A. We can consider the following two cases.

Case 1: There are  $j_1, j_2 \in \mathbb{Z}_r$ ,  $j_1 \neq j_2$ , such that  $E \cap A_{j_1} \neq \emptyset$  and  $E \cap A_{j_2} \neq \emptyset$ . According to Remark 24,  $t \leq 2^{m_0} - 1 =$  $2t_1 + 1$ . Then either  $|E \cap A_{j_1}| \le t_1$  or  $|E \cap A_{j_2}| \le t_1$ . Without loss of generality, assume  $|E \cap A_{j_1}| \le t_1$ . Similar to the proof

of 1),  $C|_{A_{j_1}}$  is an  $(r, t_1)$ -SLRC and there exists an  $\alpha \in E \cap A_{j_1}$  such that  $\alpha$  has a recovering set  $R \subseteq A_{j_1} \setminus E \subseteq A \setminus E$ . Case 2:  $E \subseteq A_{j_1}$  for some  $j_1 \in \mathbb{Z}_r$ . In this case, if  $E \cap \Omega_0^{(m)} = \emptyset$ . Then the expected  $\alpha$  exists by Lemma 27. So we assume  $E \cap \Omega_0^{(m)} \neq \emptyset$ . Pick an  $\alpha \in E \cap \Omega_0^{(m)}$ . Recall that  $t_1 = 2^{m_0-1} - 1$ . By (34), we can check that  $L_{\alpha}^{(m_0)} \subseteq \Omega_0^{(m)} \bigcup \Gamma_{t_1+1}^{(m)}$  and  $L_{\alpha}^{(m_0)} \cap A_{j_1} = \{\alpha\}$ . Since  $E \subseteq A_{j_1}$ , then  $R = L_{\alpha}^{(m_0)} \setminus \{\alpha\} \subseteq \Omega_{t_1+1}^{(m)} \setminus E \subseteq \Omega_t^{(m)} \setminus E$ . By Lemma 26, R is a recovering set of  $\alpha$ . iii): If  $t_2 = 0$ , the claim is naturally true. Assume  $t_2 \ge 1$ . By 2) of Lemma 25,  $C|_B$  has a parity check matrix containing all norms of  $U_{\alpha}^{(m-1)}$ . rows of  $H_{t_2}^{(m-1)}$ . So by induction assumption,  $\mathcal{C}|_B$  is an  $(r, t_2)$ -SLRC. Hence, by Lemma 2, there exists an  $\alpha \in E$  such that

 $\alpha$  has a recovering set  $R \subseteq B \setminus E$ . iv): In this case, by the definition of B, we have  $E \cap \Omega_0^{(m)} = \emptyset$ . Hence, by Lemma 27, there exists an  $\alpha \in E$  such that  $\alpha$ 

has a recovering set  $R \subseteq \Omega_t^{(m)} \setminus E$ .

Combining i)-iv) and by Lemma 3, the result follows.

# VI. CONSTRUCTION FROM RESOLVABLE CONFIGURATIONS

In [20], by using t-3 mutually orthogonal latin squares (MOLS) of order r, the authors construct a family of binary (r,t)-SLRC with  $k = r^2$  and code rate  $\frac{k}{n} = 1/(1 + \frac{t-1}{r} + \frac{1}{r^2})$  for odd t. A limitation of this construction is  $t \le r+2$ , since, a necessary condition of existing  $\ell$  MOLS of order r (r > 1) is  $\ell \le r-1$  [25]. In this section, by using the resolvable configurations, we give a new family of binary (r, t)-SLRC achieving the same rate  $\frac{k}{r} = 1/(1 + \frac{t-1}{r} + \frac{1}{r^2})$  for any r and any odd  $t \ge 3$  (not limited by  $t \le r+2$ ). First, we introduce a definition [25], [26].

Definition 28: Let X be a set of k elements, called points, and A be a collection of subsets of X, called lines. The pair  $(X, \mathcal{A})$  is called a  $(k_{t-1}, b_r)$  configuration if the following three conditions hold.

- (1) Each line contains r points;
- (2) Each point belongs to t-1 lines;

(3) Every pair of distinct points belong to at most one line;

Clearly, condition (3) is equivalent to the following condition.

(3') Every pair of distinct lines have at most one point in common;

The configuration  $(X, \mathcal{A})$  is called *resolvable*, if further

(4) All lines in A can be partitioned into t-1 parallel classes, where a parallel class is a set of lines that partition X.

For any  $(k_{t-1}, b_r)$  resolvable configuration  $(X, \mathcal{A})$ , one can see that r|k and each parallel class contains  $s = \frac{k}{r}$  lines. So,  $b = \frac{k}{r}(t-1) = s(t-1)$  in such a case. As usual, the incidence matrix of a  $(k_{t-1}, b_r)$  configuration  $(X, \mathcal{A})$ , where  $X = \{x_1, \dots, x_k\}$  and  $\mathcal{A} = \{A_1, \dots, A_b\}$ , is defined as a  $b \times k$  binary matrix  $M = (m_{i,j})$  such that

$$m_{i,j} = \begin{cases} 1, & \text{if } x_j \in A_i; \\ 0, & \text{otherwise.} \end{cases}$$

Clearly, any configuration is uniquely determined by its incidence matrix.

*Example 29:* We can check that the following matrix determines a  $(k_{t-1}, b_r)$  resolvable configuration  $(X, \mathcal{A})$  with k = 9, t-1 = 4, b = 12 and r = 3. Clearly,  $\{A_1, A_2, A_3\}$ ,  $\{A_4, A_5, A_6\}$ ,  $\{A_7, A_8, A_9\}$  and  $\{A_{10}, A_{11}, A_{12}\}$  are four parallel classes of  $(X, \mathcal{A})$  and any pair of lines in different parallel classes have one point in common.

$$M = \left(\begin{array}{cccccccccccccc} 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 \end{array}\right)$$

Resolvable configurations was recently used for constructing codes whose information symbols have locality r and availability t by Su [27]. The author also constructed some resolvable configurations in the paper, for example, the  $(k_{t-1}, b_r)$  resolvable configurations with  $k = r^m$  and  $t - 1 \le \frac{r^{m-1}}{r-1}$ , where  $m \ge 2$  and r is a prime power. The following construction, using the free  $\mathbb{Z}_r$ -module [28], not only generalize the result of [27], but also enable us to construct  $(k_{t-1}, b_r)$  resolvable configuration for any  $r, t \ge 2$  (r need not be a prime power), and further (r, t)-SLRCs for any  $r \ge 2$  and odd integer  $t \ge 3$ .

Lemma 30: For any  $r, t \ge 2$ , there exists a  $(k_{t-1}, b_r)$  resolvable configuration with  $k = r^m$ , where m is an arbitrary integer such that  $m \ge \log_2 t$ .

*Proof:* Consider the free  $\mathbb{Z}_r$ -module  $X = \mathbb{Z}_r^m$ , where  $\mathbb{Z}_r$  is the ring of integers modulo r. For any  $\alpha \in \mathbb{Z}_r^m$ , we use  $\alpha(j)$  to denote the *j*th coordinate of  $\alpha$ . For example, if  $\alpha = (i_1, i_2, \cdots, i_m)$ , then  $\alpha(j) = i_j$ .

For each nonempty  $S \subseteq [m]$ , let  $\alpha_S \in \mathbb{Z}_r^m$  be such that  $\alpha_S(j) = 1$  for  $j \in S$  and  $\alpha_S(j) = 0$  otherwise. Let

$$A_{S,0} \triangleq \{i \cdot \alpha_S; i \in \mathbb{Z}_r\}$$

Clearly,  $A_{S,0}$  is a submodule of  $\mathbb{Z}_r^m$  with r elements and  $A_{S,0} \cap A_{S',0} = (0, 0, \dots, 0)$  for any two distinct nonempty subsets S and S' of [m]. Let

$$\mathcal{A}_S = \{A_{S,\ell}, \ \ell = 0, 1, \cdots, r^{m-1} - 1\}$$

be the collection of all cosets of  $A_{S,0}$ . Then  $\alpha_1 - \alpha_2 \in A_{S,0}$  for any  $\ell \in \{0, 1, \dots, r^{m-1} - 1\}$  and any  $\alpha_1, \alpha_2 \in A_{S,\ell}$ .

Note that  $m \ge \log_2 t$  (i.e.,  $t-1 \le 2^m - 1$ ) and [m] has  $2^m - 1$  nonempty subsets. We can always pick t-1 nonempty subsets of [m], say  $S_1, S_2, \dots, S_{t-1}$ . Let

$$\mathcal{A} = \bigcup_{i=1}^{t-1} \mathcal{A}_{S_i}$$

We claim that  $(X = \mathbb{Z}_r^m, \mathcal{A})$  is a  $(k_{t-1}, b_r)$  resolvable configuration, which can be seen as follows.

Firstly, noticing that for each nonempty  $S \subseteq [m]$ ,  $\mathcal{A}_S$  is a partition of X, then Conditions (1), (2), (4) of Definition 28 hold. Secondly, if S, S' are two distinct nonempty subsets of [m] and  $\ell, \ell' \in \{0, 1, \dots, 2^m - 1\}$ , then we have  $|A_{S,\ell} \cap A_{S',\ell'}| \leq 1$ . Since, if otherwise, suppose  $\alpha_1, \alpha_2 \in A_{S,\ell} \cap A_{S',\ell'}$ , then  $\alpha_1 - \alpha_2 \in A_{S,0} \cap A_{S',0} = (0, 0, \dots, 0)$ . Hence, we have  $\alpha_1 = \alpha_2$ , i.e.  $|A_{S,\ell} \cap A_{S',\ell'}| \leq 1$ . Moreover, since for each nonempty  $S \subseteq [m]$ ,  $\mathcal{A}_S$  is a partition of X, so Condition (3) of Definition 28 holds, which completes the proof.

In the rest of this section, we always assume that  $(X, \mathcal{A})$  is a  $(k_{t-1}, b_r)$  resolvable configuration and  $\mathcal{A} = \{A_1, \dots, A_b\}$ . Firstly, we need a lemma for the property of the resolvable configuration  $(X, \mathcal{A})$  with odd t.

Lemma 31: Let E be a t-subset of X and t be an odd integer. Then there exists an  $A_j \in \mathcal{A}$  such that  $|E \cap A_j| = 1$ .

*Proof:* Consider a parallel class of  $(X, \mathcal{A})$ . Since it is a partition of X and |E| = t is odd, there exists some  $A_{j_1}$  in the class such that  $|E \cap A_{j_1}|$  is odd. If  $|E \cap A_{j_1}| = 1$ , then we have done. So suppose  $E = \{i_1, \dots, i_t\}$  and  $\{i_1, i_2, i_3\} \subseteq E \cap A_{j_1}$ . Since each point belongs to t - 1 lines, we can assume  $i_1$  belongs to lines  $A_{j_1}, A_{j_2}, \dots, A_{j_{t-1}}$ , where  $A_{j_1}, A_{j_2}, \dots, A_{j_{t-1}}$  belong to different parallel classes. Moreover, since every pair of distinct points belong to at most one line, then  $i_2, i_3 \notin A_{j_\ell}, \forall \ell \in \{2, \dots, t-1\}$  and each point  $i_\ell, \ell \in \{4, \dots, t\}$ , belongs to at most one line in  $\{A_{j_2}, \dots, A_{j_{t-1}}\}$ . Hence, there exists a line  $A_j \in \{A_{j_2}, \dots, A_{j_{t-1}}\}$  that contains no point in  $\{i_2, \dots, i_t\}$ . That is to say,  $E \cap A_j = \{i_1\}$ , which completes the proof.

From now on, we let X = [k] and  $A_1 = \{A_1, \dots, A_s\}$ ,  $A_2 = \{A_{s+1}, \dots, A_{2s}\}$ ,  $\dots$ ;  $A_{t-1} = \{A_{(t-2)s+1}, \dots, A_b\}$  be the t-1 parallel classes of  $(X, \mathcal{A})$ . We further partition [s] into  $\lceil \frac{s}{r} \rceil$  nonempty subsets, say  $B_1, \dots, B_{\lceil \frac{s}{r} \rceil}$ , such that  $|B_i| \leq r$  for all  $i \in \{1, \dots, \lceil \frac{s}{r} \rceil\}$ . Such a partition plays a subtle role in our construction, as will become clear later. Now, let  $W = (w_{i,j})$  be a  $\lceil \frac{s}{r} \rceil \times b$  matrix defined by

$$w_{i,j} = \begin{cases} 1, & \text{if } j \in B_i \\ 0, & \text{otherwise.} \end{cases}$$

Let M be the incidence matrix of  $(X, \mathcal{A})$  and

$$H = \begin{pmatrix} M & I_b & O_{b \times \lceil \frac{s}{r} \rceil} \\ O_{\lceil \frac{s}{r} \rceil \times k} & W & I_{\lceil \frac{s}{r} \rceil} \end{pmatrix}$$
(39)

where  $I_{\ell}$  denotes the  $\ell \times \ell$  identity matrix and  $O_{\ell \times \ell'}$  denotes the  $\ell \times \ell'$  all-zero matrix for any positive integers  $\ell$  and  $\ell'$ . Clearly, H has  $b + \lceil \frac{s}{r} \rceil$  rows,  $n = k + b + \lceil \frac{s}{r} \rceil$  columns and rank  $b + \lceil \frac{s}{r} \rceil$ .

As an example, consider the resolvable configuration  $(X, \mathcal{A})$  in Example 29. We have  $s = \frac{k}{r} = 3$  and  $\lceil \frac{s}{r} \rceil = 1$ . So we can construct  $W = (1, 1, 1, 0, \dots, 0)_{1 \times 12}$  and  $I_{\lceil \frac{s}{r} \rceil} = (1)_{1 \times 1}$ , and according to (39), further construct a matrix H as in (40).

Let C be a binary linear code with parity check matrix H as (40). Then from the first 12 rows of H, we can see that the coordinate 1 has 4 disjoint recovering sets, i.e.,  $\{2,3,10\}$ ,  $\{4,7,13\}$ ,  $\{6,8,16\}$  and  $\{5,9,19\}$ , and the coordinate 10 has a recovering set  $\{1,2,3\} \subseteq \{1,\cdots,9\}$ . Moreover, from the last row of H, we can see that  $\{11,12,22\}$  is a recovering set of 10 and  $\{10,11,12\}$  is a recovering set of 22. In general, we have the following lemma.

Lemma 32: Let C be an [n, k] binary linear code with parity check matrix H as in (40). Then, the following hold.

1) Each  $i \in [k]$  has t-1 disjoint recovering sets, i.e.,  $A_{j_{\ell}} \cup \{k+j_{\ell}\} \setminus \{i\}$ , where  $A_{j_{\ell}}$ ,  $\ell = 1, \dots, t-1$ , are lines containing i.

2) Each  $i \in \{k+1, \dots, k+b\}$  has a recovering set  $R \subseteq [k]$ .

3) Each  $i \in \{k+1, \dots, k+s\}$  has a recovering set  $R \subseteq \{k+1, \dots, k+s\} \cup \{k+b+1, \dots, n\} \setminus \{i\}$ .

4) Each  $i \in \{k+b+1, \dots, n\}$  has a recovering set  $R \subseteq \{k+1, \dots, k+s\}$ .

*Proof:* 1) and 2) are obtained by considering the first b rows of H; 3) and 4) are obtained by considering the last  $\lceil \frac{s}{r} \rceil$  rows of H.

Theorem 33: If t is odd, then the binary linear code C with parity check matrix H as in (39) is an (n, k, r, t)-SLRC with rate

$$\frac{k}{n} = \left(1 + \frac{t-1}{r} + \left\lceil \frac{1}{r^2} \right\rceil\right)^{-1}$$

*Proof:* By the construction, C has block length

$$n = k + b + \left\lceil \frac{s}{r} \right\rceil$$
$$= k \left( 1 + \frac{t-1}{r} + \left\lceil \frac{1}{r^2} \right\rceil \right).$$

and dimension  $n - (b + \lfloor \frac{s}{r} \rfloor) = k$ . So the code rate is

$$\frac{k}{n} = \left(1 + \frac{t-1}{r} + \left\lceil \frac{1}{r^2} \right\rceil\right)^{-1}$$

We now prove, according to Lemma 2, that for any  $E \subseteq [n]$  with  $|E| \leq t$ , there exists an  $i \in E$  such that i has a recovering set  $R \subseteq [n] \setminus E$ . Consider the following cases.

Case 1:  $E \cap [k] = \emptyset$ . Then we have  $E \subseteq \{k + 1, \dots, n\}$ . If  $E \cap \{k + 1, \dots, k + b\} \neq \emptyset$ , then by 2) of Lemma 32, each  $i \in E \cap \{k + 1, \dots, k + b\}$  has a recovering set  $R \subseteq [k] \subseteq [n] \setminus E$ ; Otherwise,  $E \subseteq \{k + b + 1, \dots, n\}$ , then by 4) of Lemma 32, each  $i \in E$  has a recovering set  $R \subseteq \{k + 1, \dots, k + s\} \subseteq \{k + 1, \dots, k + b\} \subseteq [n] \setminus E$ .

Case 2:  $E \cap [k] \neq \emptyset$ . Pick an  $i_1 \in E \cap [k]$ . Let

$$R_{\ell} = A_{j_{\ell}} \cup \{k + j_{\ell}\} \setminus \{i_1\}, \ \ell = 1, \cdots, t - 1,$$
(41)

where  $A_{j_1} \cdots A_{j_{t-1}}$  are the t-1 lines containing  $i_1$ . By 1) of Lemma 32,  $R_1, \cdots, R_{t-1}$  are t-1 disjoint recovering sets of  $i_1$ . If  $R_{\ell} \subseteq [n] \setminus E$  for some  $\ell \in \{1, \dots, t-1\}$ , then we are done. So we assume  $E \cap R_{\ell} \neq \emptyset$  for all  $\ell \in \{1, \dots, t-1\}$ . Since all  $R_{\ell}$ s are disjoint, so |E| = t,  $|E \cap R_{\ell}| = 1$ ,  $\ell = 1, \dots, t-1$ , and  $E \subseteq \{i_1\} \bigcup (\bigcup_{\ell=1}^{t-1} R_{\ell})$ . We have the following three subcases:

Case 2.1:  $E \cap R_{\ell} \subseteq [k], \forall \ell \in \{1, \dots, t-1\}$ . Then  $E \subseteq [k]$ . Since |E| = t is odd, by Lemma 31,  $|E \cap A_i| = 1$  for some  $A_i \in \mathcal{A}$ . Let  $E \cap A_i = \{i_2\}$ . By 1) of Lemma 32,  $R = A_i \cup \{k+i\} \setminus \{i_2\} \subseteq [n] \setminus E$  is a recovering set of  $i_2$ .

Case 2.2:  $E \cap R_{\ell_1} \subseteq [k]$  and  $E \cap R_{\ell_2} \not\subseteq [k]$  for some  $\{\ell_1, \ell_2\} \subseteq \{1, \dots, t-1\}$ . Without loss of generality, assume  $i_2 \in E \cap R_1 \subseteq [k]$  and  $E \cap R_2 \not\subseteq [k]$ . Then according to (41), we have  $E \cap R_2 = \{k + j_2\}$ . By 1) of Lemma 32, we can let  $R'_1, \dots, R'_{t-1}$  are t-1 disjoint recovering sets of  $i_2$ , where  $R'_1 = A_{j_1} \cup \{k + j_1\} \setminus \{i_2\}$  and  $R'_\ell = A_{j'_\ell} \cup \{k + j'_\ell\} \setminus \{i_2\}$ ,  $\ell = 2, \dots, t-1$ , such that  $A_{j_1}$  together with  $A_{j'_2}, \dots, A_{j'_{t-1}}$  are the t-1 lines containing  $i_2$  (see Fig. 9). Note that  $A_{j_1}$  is the only line containing both  $i_1$  and  $i_2$ , one can see that  $\{i_1, i_2, k + j_2\} \cap R'_\ell = \emptyset, \forall \ell \in \{2, \dots, t-1\}$ . Since |E| = t, then there exists an  $\ell_0 \in \{2, \dots, t-1\}$  such that  $E \cap R'_{\ell_0} = \emptyset$ . Hence,  $R'_{\ell_0} \subseteq [n] \setminus E$  is a recovering sets of  $i_2$ .

Case 2.3:  $E \cap R_{\ell} \not\subseteq [k]$  for all  $\ell \in \{1, \dots, t-1\}$ . Then we have  $E \cap R_{\ell} = \{k+j_{\ell}\}$ . Note that  $A_{j_1}, \dots, A_{j_{t-1}}$  belong to distinct parallel classes (since all of them contain  $i_1$ ). Without loss of generality, we assume  $A_{j_{\ell}} \in \mathcal{A}_{\ell}, \ell \in \{1, \dots, t-1\}$ . Then  $j_1 \leq s$  and  $s < j_{\ell} \leq b, \ell = 2, \dots, t-1$ . By 3) of Lemma 32,  $k+j_1$  has a recovering set  $R \subseteq \{k+1, \dots, k+s\} \cup \{k+b+1, \dots, n\} \setminus \{j_1\} \subseteq [n] \setminus E$ .

By the above discussion, for any  $E \subseteq [n]$  of size  $|E| \leq t$ , there exists an  $i \in E$  such that i has a recovering set  $R \subseteq [n] \setminus E$ . So by Lemma 2, C is an (n, k, r, t)-SLRC.




Fig 9. Illustration of points and recovering sets:  $R_{\ell} = A_{j_{\ell}} \cup \{k+j_{\ell}\} \setminus \{i_1\}, \ell = 1, \cdots, t-1$ , are t-1 recovering sets of  $i_1$ ;  $R_1$  and  $R'_{\ell} = A_{j'_{\ell}} \cup \{k+j'_{\ell}\} \setminus \{i_2\}, \ell = 2, \cdots, t-1$ , are t-1 recovering sets of  $i_2$ .

## VII. CONCLUSIONS AND FUTURE WORK

In this paper, we investigated sequential locally repairable codes (SLRC) by proposing an upper bound on the code rate of (n, k, r, t)-SLRC for t = 3, and constructed two families of (n, k, r, t)-SLRC for  $r, t \ge 2$  (for the second family, t is odd). Both of our constructions have code rate  $> \frac{r}{r+t}$  and are optimal for  $t \in \{2, 3\}$  with respect to the proposed bound.

It is still an open problem to determine the optimal code rate of (n, k, r, t)-SLRCs for general t, i.e.,  $t \ge 5$ . Here, we conjecture that an achievable upper bound of the code rate of (n, k, r, t)-SLRCs has the following form:

$$\frac{k}{n} \le \left(1 + \sum_{i=1}^{m} \frac{a_i}{r^i}\right)^{-1},\tag{42}$$

where  $m = \lceil \log_r k \rceil$ , all  $a_i \ge 0$  are integers such that  $\sum_{i=1}^m a_i = t$ . This conjecture can be verified for  $t \in \{1, 2, 3, 4\}$ , for which the values of the *m*-tuple  $(a_1, \dots, a_m)$ , denoted by  $\alpha_t$  for each *t*, are listed in the following table, where, the cases of t = 2, 3 are due to [17] and this work, respectively. The case of t = 4 (for binary code) is recently due to Balaji et al [21].

t	$a_1$	$a_2$	$a_3$	•••	$a_m$
1	1	0	0	• • •	0
2	2	0	0		0
3	2	1	0		0
4	2	2	0	• • •	0

Table 1. The known values of  $\alpha_t = (a_1, a_2, \cdots, a_m)$ .

It is very hard to give the explicit values of  $\alpha_t$  for general  $t \ge 5$ , even a LP-based or recursive formulation of  $\alpha_t$  seems difficult. Further, we conjecture that  $\alpha_5 = (2, 2, 1, 0, \dots, 0)$  and  $\alpha_6 = (2, 3, 1, 0, \dots, 0)$ . We would like to take the above problems our future work.

#### REFERENCES

- A. G. Dimakis, B. Godfrey, Y.Wu, M. J. Wainwright, and K. Ramchandran, "Network coding for distributed storage systems," *IEEE Trans. Inf. Theory*, vol. 56, no. 9, pp. 4539-4551, Sep. 2010.
- [2] D. S. Papailiopoulos and A. G. Dimakis, "Locally repairable codes," in Proc. IEEE Int. Symp. Inf. Theory (ISIT), Cambridge, USA, Jul. 2012, pp. 2771-2775.
- [3] P. Gopalan, C. Huang, H. Simitci, and S. Yekhanin, "On the locality of codeword symbols," IEEE Trans. Inf. Theory, vol. 58, no. 11, pp. 6925-6934, Nov. 2012.
- [4] F. Oggier and A. Datta, "Self-repairing homomorphic codes for distributed storage systems," in INFOCOM, 2011 Proceedings IEEE, pp. 1215-1223, IEEE, 2011.
- [5] D. S. Papailiopoulos, J. Luo, A. G. Dimakis, C. Huang, and J. Li, "Simple regenerating codes: Network coding for cloud storage," in INFOCOM, 2012 Proceedings IEEE, pp. 2801-2805, IEEE, 2012.
- [6] L. Pamies-Juarez, H. D. L. Hollmann, and F. Oggier, "Locally repairable codes with multiple repair alternatives," in Proc. IEEE Int. Symp. Inf. Theory (ISIT), Istanbul, Turkey, Jul. 2013, pp. 892-896.
- [7] N. Prakash, G. M. Kamath, V. Lalitha, and P. V. Kumar, "Optimal linear codes with a local-error-correction property," in Proc. IEEE Int. Symp. Inf. Theory (ISIT), Cambridge, USA, Jul. 2012, pp. 2776-2780.
- [8] A. Wang and Z. Zhang, "Repair locality with multiple erasure tolerance," IEEE Trans. Inf. Theory, vol. 60, no. 11, pp. 6979-6987, Nov. 2014.
- [9] A. S. Rawat, D. S. Papailiopoulos, A. G. Dimakis, and S. Vishwanath, "Locality and Availability in Distributed Storage," in Proc. IEEE Int. Symp. Inf. Theory (ISIT), Honolulu, HI, USA, June. 2014, pp. 681-685.
- [10] I. Tamo, A. Barg, "Bounds on locally recoverable codes with multiple recovering sets," in Proc. IEEE Int. Symp. Inform. Theory (ISIT), Honolulu, HI, USA, June. 2014, pp. 691-695.
- [11] I. Tamo, A. Barg, "A family of optimal locally recoverable codes," IEEE Trans. Inf. Theory, vol. 60, no. 80, pp. 4661-4676, Aug. 2014.
- [12] A. Wang, Z. Zhang, and M. Liu, "Achieving Arbitrary Locality and Availability in Binary Codes," in Proc. IEEE Int. Symp. Inform. Theory (ISIT), pp. 1866-1870, 2015.
- [13] A. S. Rawat, A. Mazumdar, and S. Vishwanath, "On cooperative local repair in distributed storage," in Proc. 48th Annu. Conf. Inf. Sci. Syst., Mar. 2014, pp. 1-5.
- [14] W. Song, S. H. Dau, C. Yuen, and T. J. Li, "Optimal locally repairable linear codes," IEEE J. Sel. Areas Commun., vol. 32, no. 5, pp. 1019-1036, May 2014.
- [15] P. Huang, E. Yaakobi, H. Uchikawa, and P. H. Siegel, "Linear locally repairable codes with availability," in Proc. IEEE Int. Symp. Inform. Theory (ISIT), pp. 1871-1875, 2015.
- [16] A. Wang, Z. Zhang, and D. Lin, "Two Classes of (r, t)-Locally Repairable Codes," in Proc. IEEE Int. Symp. Inf. Theory (ISIT), Barcelona, Spain, Jul. 2016, pp. 445 449.
- [17] N. Prakash, V. Lalitha, and P. V. Kumar, "Codes with locality for two erasures," in Proc. IEEE Int. Symp. Inform. Theory (ISIT), Honolulu, HI, USA, June. 2014, pp. 1962-1966.
- [18] W. Song and C. Yuen, "Locally Repairable Codes with Functional Repair and Multiple Erasure Tolerance," http://arxiv.org/abs/1507.02796, 2015.
- [19] W. Song and C. Yuen, "Binary locally repairable codes sequential repair for multiple erasures," in *Proc. Globecom* 2016, also available at: http://arxiv.org/abs/1511.06034, 2015.
- [20] S. B. Balaji, K. P. Prasanth, and P. V. Kumar, "Binary codes with locality for multiple erasures having short block length," in Proc. IEEE Int. Symp. Inf. Theory (ISIT), Barcelona, Spain, Jul. 2016, pp. 655 - 659.
- [21] S. B. Balaji, K. P. Prasanth, and P. V. Kumar, "Binary Codes with Locality for Four Erasures," Available: https://arxiv.org/abs/1607.02817
- [22] A. Wang and Z. Zhang, "An Integer Programming-Based Bound for Locally Repairable Codes," *IEEE Trans. Inf. Theory*, vol. 61, no. 10, pp. 5280-5294, Oct. 2015.
- [23] C. Fragouli and E. Soljanin,"Information flow decomposition for network coding," IEEE Trans. Inf. Theory, vol. 52, no. 3, pp. 829-848, Mar. 2006.
- [24] W. Song, K. Cai, R. Feng and C. Yuen, "The Complexity of Network Coding With Two Unit-Rate Multicast Sessions," *IEEE Trans. Inf. Theory*, vol. 59, no. 9, pp. 5692-5707, Sept. 2013.
- [25] C. J. Colbourn and J. H. Denitz, Handbook of Combinatorial Designs, 2nd ed. Chapman and Hall/CRC, 2006.
- [26] T. Pisanski and B. Servatius, Configurations from a graphical viewpoint, Birkhauser, Basel, 2013.
- [27] Y.S. Su, "Design of Membership Matrices for (r, t)-Availability in Distributed Storage," in Proc. IEEE Int. Symp. Inf. Theory (ISIT), Barcelona, Spain, Jul. 2016, pp. 998 - 1002.
- [28] W.C. Brown, Matrices over Commutative Rings, Dekker, New York, 1993.