

MATRICES WITH GAUSSIAN NOISE: OPTIMAL ESTIMATES FOR SINGULAR SUBSPACE PERTURBATION

SEAN O'ROURKE, VAN VU, AND KE WANG

ABSTRACT. The Davis–Kahan–Wedin $\sin \Theta$ theorem describes how the singular subspaces of a matrix change when subjected to a small perturbation. This classic result is sharp in the worst case scenario. In this paper, we prove a stochastic version of the Davis–Kahan–Wedin $\sin \Theta$ theorem when the perturbation is a Gaussian random matrix. Under certain structural assumptions, we obtain an optimal bound that significantly improves upon the classic Davis–Kahan–Wedin $\sin \Theta$ theorem. One of our key tools is a new perturbation bound for the singular values, which may be of independent interest.

1. INTRODUCTION

Consider an $N \times n$ (data) matrix A . In practice, it is common that we only have access to a corrupted (noisy) version \tilde{A} given by

$$\tilde{A} := A + E, \tag{1}$$

where E represents the noise matrix. As a result, one must use \tilde{A} as input for all calculations and algorithms intended for A . A question of fundamental interest is to estimate the impact of the noise E on the output; see for instance [31, 32, 33, 37, 40, 41, 55, 57, 60, 65, 72] and references therein.

In modern studies, noise is often assumed to be random (e.g., Gaussian) and the data matrix A possesses certain structural properties. For example, in a vast number of studies, researchers assume that A has low rank [18, 19, 20, 63], and our main results focus on this case.

Assume that the $N \times n$ data matrix A has rank $r \geq 1$. We will often think of r as a constant (or a parameter very small compared to the dimensions N and n such as $r \leq \log n$ or $r \leq n^\epsilon$). The singular value decomposition (SVD) of A takes the form $A = U\Sigma V^T$, where $\Sigma = \text{diag}(\sigma_1, \dots, \sigma_r)$ is a diagonal matrix containing the non-zero singular values $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r > 0$ of A ; the columns of the matrices $U = (u_1, \dots, u_r)$ and $V = (v_1, \dots, v_r)$ are the orthonormal left and right singular vectors of A , respectively. In other words, u_i and v_i are the left and right singular vectors corresponding to σ_i . It follows that $U^T U = V^T V = I_r$, where I_r is the $r \times r$ identity matrix. For convenience, we will take $\sigma_{r+i} = 0$ for all $i \geq 1$.

Recall that \tilde{A} is given in (1). Denote the SVD of \tilde{A} similarly by $\tilde{A} = \tilde{U}\tilde{\Sigma}\tilde{V}^T$, where the diagonal entries of $\tilde{\Sigma}$ are the singular values $\tilde{\sigma}_1 \geq \tilde{\sigma}_2 \geq \dots \geq \tilde{\sigma}_{\min\{N,n\}} \geq$

S. O'Rourke has been supported in part by NSF grant DMS-1810500.

V. Vu is partially supported by NSF grant DMS-1902825.

K. Wang is partially supported by Hong Kong RGC grant GRF 16308219, GRF 16304222 and ECS 26304920.

0, and the columns of \tilde{U} and \tilde{V} are the orthonormal left and right singular vectors, denoted by \tilde{u}_i and \tilde{v}_i , respectively.

Let Π_s denotes the orthogonal projection onto the subspace spanned by the s leading singular vectors of A (either left or right). The matrix $\Pi_s A$ is the best rank s approximation of A [34, Section 2.4] and plays an important role in applications in almost every fields of science involving large data sets. Given the noise issue, it is thus of fundamental interest to bound the difference between Π_s and its “noisy” counterpart $\tilde{\Pi}_s$ (the projection onto the subspace formed by the leading s singular vectors of \tilde{A}). The goal of this paper is to bound this difference.

As our main result is bit technical, let us first consider a toy case. Assume we want to compute the first (left) singular vector u_1 of the matrix A . If we only have access to the noisy matrix \tilde{A} , we can only compute \tilde{u}_1 . The famous Davis–Kahan–Wedin $\sin \Theta$ theorem [25, 70] provides a bound on the difference between \tilde{u}_1 and u_1 . Two parameters appear in this bound: the gap (or separation) δ_1 between the largest singular values of A given by

$$\delta_1 := \sigma_1 - \sigma_2$$

and the spectral norm of E defined by

$$\|E\| := \max_{\|u\|=1} \|Eu\|,$$

where $\|u\|$ denotes the Euclidean norm of the vector u .

Theorem 1 (Davis–Kahan–Wedin $\sin \Theta$ theorem). *One has*

$$\sin \angle(u_1, \tilde{u}_1) \leq 2 \frac{\|E\|}{\delta_1},$$

where $\angle(u_1, \tilde{u}_1)$ is the acute angle between u_1 and \tilde{u}_1 , taken in $[0, \pi/2]$. The same bound holds for $\sin \angle(v_1, \tilde{v}_1)$.

Theorem 1 follows as a simple corollary of the Davis–Kahan–Wedin $\sin \Theta$ theorem; see Theorem 4 from [51], which also contains an example explaining the necessity of the appearance of the gap δ_1 .

In [51], the current authors considered random noise and improved Theorem 1 by showing that a stronger bound

$$\sin \angle(u_1, \tilde{u}_1) \lesssim \frac{C(r)}{\delta_1} + \frac{\|E\|}{\sigma_1} + \frac{\|E\|^2}{\sigma_1 \delta_1} \quad (2)$$

holds with high probability (the probability space is generated by the randomness of the noise matrix E ; see [51] for details). Here, $C(r)$ is a parameter depending polynomially on r . To see how this improves upon the Davis–Kahan–Wedin $\sin \Theta$ theorem, let us mention that in most settings, the norm of the random matrix E is polynomial in $N + n$. Thus, in the setting where r is significantly smaller than the dimensions N, n , the first term $C(r)/\delta_1$ improves upon the term $\|E\|/\delta_1$ as it replaces a polynomial in $N + n$ by a polynomial in r . The second term $\|E\|/\sigma_1$ represents the signal-to-noise ratio; notice that the denominator is the singular value σ_1 which is usually much larger than the gap δ_1 between σ_1 and σ_2 . Finally, the third term $\frac{\|E\|^2}{\sigma_1 \delta_1}$ improves upon the term $\|E\|/\delta_1$ by the a factor involving the noise-to-signal ratio $\|E\|/\sigma_1$. This theorem generalized Theorem 8 of [66], where the second author considered Bernoulli random matrices.

The first term on the right-hand side of (2) was already conjectured in [66]. As noted above, the second term represents the signal-to-noise ratio. Both terms are necessary (see below for further details). Indeed, if the gap δ_1 is too small, there is a chance that the two leading singular values (and thus the corresponding singular vectors) get “swapped.” Moreover, if the signal-to-noise ratio is low, then the data matrix is overwhelmed by noise, and the singular vectors behave more like random vectors. In random matrix theory, this phenomenon is known as the BBP phase transition, named after Baik, Ben Arous, and P ech e [5]; see [5, 8] and references therein for further details.

It is conjectured that the third term on the right-hand side of (2) is not necessary, under some mild assumptions. In this paper, we introduce a new method of analyzing random perturbations and confirm this conjecture (up to lower-order corrections). As the precise result is a bit technical, let us first state a simpler, but easier to read, version of our main result. The asymptotic notation is used under the assumption that the dimensions n and N tend to infinity.

Theorem 2 (Perturbation with Gaussian noise; simplified asymptotic version). *Let A and E be $N \times n$ real matrices, where A is deterministic and the entries of E are jointly independent standard Gaussian random variables. If A has rank $r \geq 1$, then, with probability $1 - o(1)$,*

$$\sin \angle(u_1, \tilde{u}_1) \lesssim \frac{C(r, \log(N+n))}{\delta_1} + \frac{\|E\|}{\sigma_1}, \quad (3)$$

where $C(s, t)$ is a positive parameter which grows at most polynomially in s and t .

Theorem 2 shows that the third term on the right-hand side of (2) is not necessary and verifies the conjecture discussed above. In Section 2, we state the more detailed version of Theorem 2 and its generalizations. In this paper, we focus on the case when the entries of E are jointly independent Gaussian random variables. This assumption simplifies parts of the (already technical) proofs, but the method does extend to other distributions. For instance, the results can be extended to the case when E has independent and identically distributed sub-gaussian entries. To relax the Gaussian assumption, one needs to establish a variant of the isotropic local law presented in Lemma 9 below. Similar results have been established for more general entry distributions (see the discussion regarding the isotropic local laws in Section 3.2 for references). We plan to discuss the non-Gaussian case in a separate paper.

One can also consider the case when the entries of E are not necessarily identically distributed (but are still jointly independent). For example, consider the case when the entries of $E = (E_{ij})$ are jointly independent normal random variables, where E_{ij} has mean zero and variance σ_{ij}^2 . Depending on the values of the variances σ_{ij}^2 , the resolvent of E may no longer satisfy an isotropic law and instead may satisfy an anisotropic law [43]. It is unclear if our methods can be adapted to deal with the anisotropic case. Similarly, while we expect our results to also hold when there is a small dependence between the entries of E , our methods heavily rely on the joint independence of the entries.

It is worth noting that when $\delta_1 = \sigma_1$ (i.e., the rank-1 case) or when the spectral gap is of similar magnitude to the signal strength, $\delta_1 \asymp \sigma_1$, the right-hand side of (3) is proportional to $\|E\|/\sigma_1$. Theorem 3 below implies that, up to lower order terms and constants, this bound is optimal. Nevertheless, based on our bounds, it

is apparent that $\sin \angle(u_1, \tilde{u}_1)$ can be reasonably bounded in most cases, even when $\delta_1 \ll \sigma_1$. Such relaxation offers improved outcomes in numerous applications, which will be elaborated upon in a forthcoming paper.

We remark that the bound in (3) does not depend on the condition number $\kappa = \sigma_1/\sigma_r$ of A . Instead, we have modeled our main results after the Davis–Kahan–Wedin $\sin \Theta$ theorem (Theorem 1), which instead relies on the gap δ_1 to quantify how sensitive the matrix is to perturbations.

Before concluding this section, we discuss the optimality of Theorem 2. Numerical simulations show that, up to the particular form that $C(s, t)$ takes, the first term on the right-hand side of (3) is necessary; see Figure 1. The second term on the right-hand side of (3) represents the signal-to-noise ratio; this term is also necessary as can be seen from the following lower bound.




FIGURE 1. A plot of the cumulative distribution function F of $\sin \angle(u_1, \tilde{u}_1)$, where $F(x) = \mathbb{P}(\sin \angle(u_1, \tilde{u}_1) \leq x)$ for $0 \leq x \leq 1$. We take $N = n = 1000$ and $A = \text{diag}(300, 300 - \delta, 0, \dots, 0)$ with rank $r = 2$ where the spectral gap $\delta = \delta_1$ is chosen to be 20, 10, 5, and 2. E is a Gaussian matrix as in Theorem 2. Each curve is generated from 400 samples.

Theorem 3 (Lower bound). *Let A and E be $N \times n$ real matrices, where A is deterministic with rank r satisfying $1 \leq r \leq \frac{1}{2} \max(N, n)$ and the entries of E are independent and identically distributed sub-gaussian¹ random variables with mean zero and unit variance. Then*

$$\max\{\sin \angle(u_1, \tilde{u}_1), \sin \angle(v_1, \tilde{v}_1)\} \geq \frac{1}{8\sqrt{2}} \frac{\frac{\|E\|}{\sigma_1}}{1 + (1 + \sqrt{2}) \frac{\|E\|}{\sigma_1}} \quad (4)$$

with probability at least $1 - C \exp(-c \max(N, n))$, where $C, c > 0$ are constants depending only on the sub-gaussian norm of the entries of E .

Notice that if $\frac{\|E\|}{\sigma_1} \leq 1$, then the right-hand side of (4) simplifies to $c \frac{\|E\|}{\sigma_1}$ for some constant c . The proof of Theorem 3 can be found in Appendix A. This result can be extended to more general random matrices with independent entries and matrix ensembles satisfying rotational invariance. Due to space limitation, we do not pursue such generalizations here.

¹A random variable ξ is sub-gaussian if there is a positive constant c such that $\mathbb{P}(|\xi| \geq t) \leq 2 \exp(-ct^2)$ for all $t \geq 0$; the largest constant $c > 0$ for which this property holds is called the sub-gaussian norm of ξ . Standard normal random variables are sub-gaussian.

ACKNOWLEDGEMENTS

The authors are grateful to the anonymous referees for valuable feedback and corrections.

2. MAIN RESULTS

We state all of our main results in non-asymptotic forms (without using any asymptotic notation) and state all constants explicitly so that the results can be applied to matrices of any dimension.

2.1. Individual singular vector bounds. We first state the technical version of Theorem 2 with precise dependences between the parameters.

Theorem 4 (Perturbation with Gaussian noise). *Let A and E be $N \times n$ real matrices, where A is deterministic with rank $r \geq 1$ and the entries of E are jointly independent standard Gaussian random variables. Let K be an arbitrary positive constant, and denote $\eta := 54r\sqrt{(K+8)\log(N+n)}$. If $\sigma_1 \geq 4(\sqrt{N} + \sqrt{n}) + 140\eta r$ and $\delta_1 = \sigma_1 - \sigma_2 \geq 100\eta r$, then with probability at least $1 - 15(N+n)^{-K}$*

$$\sin \angle(u_1, \tilde{u}_1) \leq 54^2 \sqrt{K+8} \frac{r \sqrt{\log(N+n)}}{\delta_1} + 2 \frac{\|E\|}{\sigma_1} \quad (5)$$

whenever $\frac{(\sqrt{N}+\sqrt{n})^2}{\log(N+n)} > 64(K+9)$. The same bound holds for $\sin \angle(v_1, \tilde{v}_1)$.

Remark 5. Note that the constants in this theorem (such as 54 and 100) and our theorems below are chosen for convenience. In order to keep the proof presentable, we have not tried to optimize these values. However, the values can be significantly improved by tracking the constants throughout the proof. For example, if one replaces the assumptions $\sigma_1 \geq 4(\sqrt{N} + \sqrt{n}) + 140\eta r$ and $\frac{(\sqrt{N}+\sqrt{n})^2}{\log(N+n)} > 64(K+9)$ by $\sigma_1 \geq 11(\sqrt{N} + \sqrt{n}) + 150\eta r$ and $\frac{(\sqrt{N}+\sqrt{n})^2}{\log(N+n)} > 200(K+9)$, then the constant 54^2 appearing on the right-hand side of (5) can be replaced by 180.

Theorem 4 improves upon (2) and Theorem 1 when the rank r is sufficiently small and $\delta_1 \ll \|E\|$. As discussed in the previous section, the bound given in (5) is optimal, up to the choice of constants and the particular polynomial dependence on r and $\log(N+n)$.

2.2. Singular subspace bounds. The results stated so far have focused on the singular vectors corresponding to the largest singular value. More generally, we will consider the singular subspaces spanned by the first j ($1 \leq j \leq r$) singular vectors. Define

$$\begin{aligned} U_j &:= \text{Span}\{u_1, \dots, u_j\}, & V_j &:= \text{Span}\{v_1, \dots, v_j\}, \\ \tilde{U}_j &:= \text{Span}\{\tilde{u}_1, \dots, \tilde{u}_j\}, & \tilde{V}_j &:= \text{Span}\{\tilde{v}_1, \dots, \tilde{v}_j\}. \end{aligned} \quad (6)$$

Even more generally, for any $1 \leq k \leq s \leq r$, let us denote

$$U_{k,s} := \text{Span}\{u_k, \dots, u_s\}, \quad \tilde{U}_{k,s} := \text{Span}\{\tilde{u}_k, \dots, \tilde{u}_s\}$$

and analogously for $V_{k,s}$ and $\tilde{V}_{k,s}$.

Recall that if U and V are two subspaces of the same dimension, then the largest principal angle $\angle(U, V)$ between them is given by

$$\sin \angle(U, V) := \max_{u \in U; u \neq 0} \min_{v \in V; v \neq 0} \sin \angle(u, v) = \|P_U - P_V\| = \|P_{U^\perp} P_V\|, \quad (7)$$

where P_W is the orthogonal projection matrix onto subspace W . We define the gaps (or separations) between the singular values of A by

$$\delta_i = \sigma_i - \sigma_{i+1}$$

for $1 \leq i \leq r$, where we use the convention that $\sigma_{r+1} = 0$.

Theorem 2 can be generalized to the following.

Theorem 6 (Singular subspace bounds; simplified asymptotic version). *Let A and E be $N \times n$ real matrices, where A is deterministic with rank $r \geq 1$ and the entries of E are jointly independent standard Gaussian random variables. For any $1 \leq r_0 \leq r$, if $\min_{\sigma_l \neq \sigma_j, 1 \leq l, j \leq r_0} |\sigma_l - \sigma_j| \geq C(r, \log(N+n))$, then, with probability $1 - o(1)$,*

$$\sin \angle(U_{r_0}, \tilde{U}_{r_0}) \lesssim \frac{C(r, \log(N+n))}{\delta_{r_0}} + \frac{\|E\|}{\sigma_{r_0}},$$

where $C(s, t)$ is a positive parameter that grows at most polynomially in s and t .

Here, the minimum $\min_{\sigma_l \neq \sigma_j, 1 \leq l, j \leq r_0} |\sigma_l - \sigma_j|$ is over all distinct singular values $\sigma_l \neq \sigma_j$, $j, l \leq r_0$. In particular, this includes the case when some of the singular values of A may be repeated. Since the minimum is only over distinct singular values, even if some singular values occur with multiplicity, it is always the case that $\min_{\sigma_l \neq \sigma_j, 1 \leq l, j \leq r_0} |\sigma_l - \sigma_j| > 0$. The technical version of Theorem 6 is given below.

Theorem 7 (Singular subspace bounds). *Let A and E be $N \times n$ real matrices, where A is deterministic and the entries of E are jointly independent standard Gaussian random variables. Assume A has rank $r \geq 1$. Let $K > 0$ be any constant and denote $\eta := 54r \sqrt{(K+8) \log(N+n)}$. Assume $\frac{(\sqrt{N} + \sqrt{n})^2}{\log(N+n)} > 64(K+9)$. For any $1 \leq r_0 \leq r$, if $\sigma_{r_0} \geq 4(\sqrt{N} + \sqrt{n}) + 140\eta r$, $\delta_{r_0} \geq 100\eta r$ and $\min_{\sigma_l \neq \sigma_j, 1 \leq l, j \leq r_0} |\sigma_l - \sigma_j| \geq 100\eta r$, then*

$$\sin \angle(U_{r_0}, \tilde{U}_{r_0}) \leq 21\sqrt{2}\eta \sqrt{\sum_{j=1}^{r_0} \frac{1}{(\sigma_j - \sigma_{r_0+1})^2}} + 2 \frac{\|E\|}{\sigma_{r_0}} \quad (8)$$

with probability at least $1 - 15(N+n)^{-K}$. The same conclusion also holds for $\sin \angle(V_{r_0}, \tilde{V}_{r_0})$.

In addition, for any $1 < k \leq s \leq r_0$, if $\min\{\delta_{k-1}, \delta_s\} \geq 100\eta r$, then

$$\begin{aligned} \sin \angle(U_{k,s}, \tilde{U}_{k,s}) &\leq 21\sqrt{2}\eta \left(\sqrt{\sum_{j=1}^{k-1} \frac{1}{(\sigma_j - \sigma_k)^2}} + \sqrt{\sum_{j=k}^s \frac{1}{(\sigma_j - \sigma_{s+1})^2}} \right) \\ &\quad + 2 \left(\frac{\|E\|}{\sigma_{k-1}} + \frac{\|E\|}{\sigma_s} \right) \end{aligned}$$

with probability at least $1 - 15(N+n)^{-K}$. The same conclusion also holds for $\sin \angle(V_{k,s}, \tilde{V}_{k,s})$.

The choice of constants, such as 100 and 140, in Theorem 7 is for convenience; an inspection of the proof will reveal exactly how much these constants can be optimized (see Remark 5).

The key to proving our main results is a precise prediction for the location of the singular values of $A + E$. In order to obtain optimal control of the singular values, one cannot simply compare $\tilde{\sigma}_j$ to σ_j (or more conveniently $\tilde{\sigma}_j^2$ to σ_j^2). Instead, we

compare σ_j^2 to $\tilde{\sigma}_j^2 + \varepsilon$, where ε is a random correction term (depending only on the matrix E). This random correction term allows us to obtain a more precise prediction for the singular values. The precise result is given in Theorem 12, which can be found in Section 3.3, after appropriate notations have been introduced.

2.3. Comparison to other results in the literature. Many classical results compare the singular vectors (alternatively, eigenvectors) of $A + E$ to those of A . The study of eigenvector perturbations dates back to at least Rayleigh [54] and Schrödinger [58]. More recently, these results include the Davis–Kahan–Wedin $\sin \Theta$ theorem [25, 70]. For the singular values (alternatively, eigenvalues), there are many classical results, including Weyl’s bound [13]. In contrast to this work, all of these classical results focus on the case when A and E are deterministic. We refer the reader to the classical texts [13, 38, 59] for further details and generalizations.

The case when E is random has only been studied more recently. As discussed above, our main results in this paper improve upon the works [51, 66]. A number of similar results have focused on the case when E has Gaussian entries. For example, in [44], Koltchinskii and Xia derive concentration bounds for linear forms involving the singular vectors and this was later extended to tensors by Xia and Zhou in [71]. The non-asymptotic distribution of the singular vectors, up to rotation, is studied by R. Wang in [69]. A perturbative expansion of the coordinates of the eigenvectors is given in [9]. Allez and Bouchaud studied the eigenvector dynamics of $A + E$ when both A and E are real symmetric matrices and the entries of E are constructed from a family of independent real Brownian motions [2].

In the random matrix theory literature, there are a number of perturbation results; in contrast to this work, many of these results focus on the case when $\|A\|$ and $\|E\|$ are proportional. The works of Benaych–Georges and Nadakuditi [11, 12] have influenced this paper (and we discuss these works more below). The results in [11, 12] establish the almost sure convergence of the projection of the outlier singular vectors (resp. eigenvectors) onto the r -dimensional singular vector subspace (resp. eigenspace) of A , assuming r , the rank of A , is fixed and the dimensions N, n tend to infinity. The limiting distribution of such projections is explicitly given in [7]. In these papers, the norm of A and E must be comparable. We make no such assumption here. (In fact, in applications, the intensity of the noise is expected to be much smaller than the key signals.) Several related results for eigenvectors of random matrices are also discussed in the survey [50]. A different yet closely related type of perturbation comes from the spiked covariance model (see [5, 6, 16, 17, 49] and many references therein).

Another class of results in the literature is motivated by applications. Motivated by statistical machine learning, Abbe, Fan, Wang, and Zhong [1] provide entry-wise bounds between the eigenvectors of a random matrix and those of its expectation. With similar motivations, the geometry of the singular subspaces are studied using the two-to-infinity subordinate vector norm on matrices in [22]. In the real symmetric case, when the matrix A is incoherent and has low rank, ℓ^∞ -norm bounds for the eigenvectors are given in [33]. Similar entrywise-type behaviors for the eigenvectors are studied in [76, 77]. Both deviation and fluctuation results for the eigenvectors are presented in [21] based on statistical motivations. Applications of principal components analysis have also motivated a number of similar works, including [75] and references therein.

The stochastic block model has been extensively studied in recent years, especially in connection with spectral algorithms, which often take advantage of eigenvector perturbation results. For example, motivated by the stochastic block model, Eldridge, Belkin, and Wang [26] investigated random perturbations of real symmetric matrices. In particular, their results focus on the eigenvalues and ℓ^∞ -norm bounds for the eigenvectors, which improve upon classical bounds. Additionally, we highlight the works [24, 35, 47, 48, 55, 61, 67, 73, 74] concerning the stochastic block model, which are perhaps the most relevant to this paper.

The list of works discussed above is far from complete and represents only a small fraction of the literature.

2.4. Outline and notation. The paper is organized as follows. Section 3 establishes the preliminary tools needed for the proofs of our main results. We introduce a key lemma, Lemma 9, that shows that the resolvent of the random noise can be well approximated by a diagonal matrix. The proof of Lemma 9 is deferred to Section 6. Section 3 also contains a brief overview of our proof techniques. In Section 4, we prove Theorem 7 using Lemma 9. Section 5 contains the proof of Theorem 12, which describes the precise location of the perturbed singular values of \tilde{A} . The proof of Theorem 3 is presented in Appendix A. In Appendix B, we collect the proofs of Propositions 8, 10, 11, (21) and Lemma 13.

Without loss of generality, we always assume $N \leq n$, for if not, one can simply apply the results to the transposes of the matrices. We introduce the following notation. For a finite set S , $|S|$ denotes the cardinality of S . Recall that $\|M\|$ denotes the spectral norm of the matrix M , and let $\|M\|_F$ denote the Frobenius norm. For a vector x , $\|x\|$ will be its Euclidean norm. The matrix I_n is the $n \times n$ identity matrix. We will often simply write I when the size can be deduced from context. For integers $m_2 \geq m_1 \geq 1$, we let $\llbracket m_1, m_2 \rrbracket := \{m_1, \dots, m_2\}$ denote the discrete interval. The distance between a point $z \in \mathbb{C}$ and a set $\mathcal{G} \subset \mathbb{C}$ is $\text{dist}(z, \mathcal{G}) := \inf_{w \in \mathcal{G}} |z - w|$. For two sets $\mathcal{F}, \mathcal{G} \subset \mathbb{C}$, the distance between them is $\text{dist}(\mathcal{F}, \mathcal{G}) := \inf_{z \in \mathcal{F}, w \in \mathcal{G}} |z - w|$. For two random elements x and y , we write $x \sim y$ if x and y have the same distribution. The function $\log(\cdot)$ will always denote the natural logarithm.

3. BASIC TOOLS AND AN OVERVIEW OF THE PROOF

In this section, we develop the preliminary tools needed to establish our main results. Section 3.5 contains a brief overview of our main proof techniques.

3.1. Linear algebra. We first apply a linearization trick, which allows us to consider the eigenvalues and eigenvectors of a symmetric matrix instead of studying the singular values and vectors of a non-symmetric matrix.

Consider the $(N + n) \times (N + n)$ matrices

$$\mathcal{A} := \begin{pmatrix} 0 & A \\ A^T & 0 \end{pmatrix} \quad \text{and} \quad \mathcal{E} := \begin{pmatrix} 0 & E \\ E^T & 0 \end{pmatrix}$$

in block form. Define

$$\tilde{\mathcal{A}} := \mathcal{A} + \mathcal{E}.$$

The non-zero eigenvalues of \mathcal{A} are given by $\pm\sigma_1, \dots, \pm\sigma_r$. Indeed, $\mathcal{A}(u_j^T, v_j^T)^T = \sigma_j(u_j^T, v_j^T)^T$ and $\mathcal{A}(u_j^T, -v_j^T)^T = -\sigma_j(u_j^T, -v_j^T)^T$. Denote these eigenvalues by $\lambda_j = \sigma_j$ and $\lambda_{j+r} = -\sigma_j$ for $1 \leq j \leq r$. Then $\mathbf{u}_j := \frac{1}{\sqrt{2}}(u_j^T, v_j^T)^T$ and $\mathbf{u}_{j+r} :=$

$\frac{1}{\sqrt{2}}(u_j^T, -v_j^T)^T$ for $1 \leq j \leq r$ are their corresponding orthonormal eigenvectors. The spectral decomposition of \mathcal{A} is given by

$$\mathcal{A} = \mathcal{U}\mathcal{D}\mathcal{U}^T, \quad (9)$$

where $\mathcal{U} := (\mathbf{u}_1, \dots, \mathbf{u}_{2r})$ and $\mathcal{D} := \text{diag}(\lambda_1, \dots, \lambda_{2r})$. It follows that $\mathcal{U}^T\mathcal{U} = I_{2r}$. Similarly, the non-zero eigenvalues of $\tilde{\mathcal{A}}$ are denoted by $\tilde{\lambda}_j = \tilde{\sigma}_j$ and $\tilde{\lambda}_{j+\min\{N,n\}} = -\tilde{\sigma}_j$ for $1 \leq j \leq \min\{N,n\}$. The eigenvector corresponding to $\tilde{\lambda}_j$ is denoted by $\tilde{\mathbf{u}}_j$ and is formed by the right and left singular vectors of \tilde{A} .

For $J \subset \llbracket 1, 2r \rrbracket$, we introduce the notation \mathcal{U}_J to denote the $(N+n) \times |J|$ matrix formed from \mathcal{U} by removing the columns containing \mathbf{u}_i for $i \notin J$. Similarly, \mathcal{D}_J will denote the $|J| \times |J|$ matrix formed from \mathcal{D} by removing the rows and columns containing λ_i for $i \notin J$. Let $I := \llbracket 1, 2r \rrbracket \setminus J$. In this way, we can decompose A as

$$\mathcal{A} = \mathcal{U}\mathcal{D}\mathcal{U}^T = \mathcal{U}_J\mathcal{D}_J\mathcal{U}_J^T + \mathcal{U}_I\mathcal{D}_I\mathcal{U}_I^T. \quad (10)$$

With a slight abuse of notation, we also denote the subspace

$$\mathcal{U}_J := \text{Span}\{\mathbf{u}_k : k \in J\}.$$

Let P_J be the orthogonal projection onto the subspace \mathcal{U}_J . Clearly, $P_J = \mathcal{U}_J\mathcal{U}_J^T$. Analogous notations $\tilde{\mathcal{U}}_J, \tilde{P}_J, \tilde{\mathcal{D}}_J$ are also defined for $\tilde{\mathcal{A}}$.

For the remainder of the paper, it suffices to derive results on the eigenspaces of \tilde{A} by noting the following linear algebra fact.

Proposition 8. *Let $U, \tilde{U} \subset \mathbb{R}^N$ and $V, \tilde{V} \subset \mathbb{R}^n$ be subspaces of the same dimension p . Let W and \tilde{W} be subspaces in \mathbb{R}^{N+n} obtained by concatenating vectors from U, V and \tilde{U}, \tilde{V} respectively, i.e. $W = \{w \in \mathbb{R}^{N+n} : w = (u^T, v^T)^T, u \in U, v \in V\}$ and $\tilde{W} = \{\tilde{w} \in \mathbb{R}^{N+n} : \tilde{w} = (\tilde{u}^T, \tilde{v}^T)^T, \tilde{u} \in \tilde{U}, \tilde{v} \in \tilde{V}\}$. Then*

$$\max\{\sin \angle(U, \tilde{U}), \sin \angle(V, \tilde{V})\} = \sin \angle(W, \tilde{W}).$$

The proof of Proposition 8 is deferred to Appendix B.1. In particular, as a special case of Proposition 8 one has

$$\max\{\sin \angle(U_{r_0}, \tilde{U}_{r_0}), \sin \angle(V_{r_0}, \tilde{V}_{r_0})\} = \sin \angle(\mathcal{U}_I, \tilde{\mathcal{U}}_I) \quad (11)$$

for the index set $I := \llbracket 1, r_0 \rrbracket \cup \llbracket r+1, r+r_0 \rrbracket$ and

$$\max\{\sin \angle(U_{k,s}, \tilde{U}_{k,s}), \sin \angle(V_{k,s}, \tilde{V}_{k,s})\} = \sin \angle(\mathcal{U}_l, \tilde{\mathcal{U}}_l) \quad (12)$$

where $l := \llbracket k, s \rrbracket \cup \llbracket r+k, r+s \rrbracket$, $1 \leq r_0 \leq r$, and $1 \leq k \leq s \leq r_0$.

We also recall the Weinstein-Aronszajn identity (also called Sylvester's determinant identity), see page 271 of [53]: if B is an $n \times k$ matrix and C is a $k \times n$ matrix then

$$\det(I_n - BC) = \det(I_k - CB). \quad (13)$$

When $k < n$, (13) allows us to reduce an $n \times n$ determinant to a smaller $k \times k$ determinant.

Weyl's inequality (see [13, Corollary III.2.6]) states that if B and C are $n \times n$ real symmetric matrices with eigenvalues $\lambda_1(B) \geq \dots \geq \lambda_n(B)$ and $\lambda_1(C) \geq \dots \geq \lambda_n(C)$, then

$$\max_{1 \leq j \leq n} |\lambda_j(B) - \lambda_j(C)| \leq \|B - C\|. \quad (14)$$

The resolvent identity

$$B^{-1} - C^{-1} = B^{-1}(C - B)C^{-1} \quad (15)$$

holds for invertible matrices B and C .

3.2. Resolvent. For $z \in \mathbb{C}$ with $|z| > \|\mathcal{E}\|$, we define the resolvent of \mathcal{E} as

$$G(z) := (zI - \mathcal{E})^{-1}.$$

Often we will drop the identity matrix and simply write $(z - \mathcal{E})^{-1}$ for this matrix. We use $G_{ij}(z)$ to denote the (i, j) -entry of $G(z)$.

The resolvent $G(z)$ is a heavily studied object in random matrix theory. For example, one can study the eigenvalues of \mathcal{E} (and hence the singular values of E) by analyzing the trace, $\text{tr} G(z)$, since the trace is a meromorphic function with poles precisely at the eigenvalues. In addition, the matrix $G(z)$ encodes the behavior of the eigenvectors of \mathcal{E} (and hence the singular vectors of E). For many matrix models (see [15, 27, 28, 30, 36, 42, 43, 45, 46] and references therein), the resolvent $G(z)$ can be approximated by a diagonal matrix. Here, we consider a random diagonal matrix

$$\Phi(z) := \begin{pmatrix} \frac{1}{\phi_1(z)} I_N & 0 \\ 0 & \frac{1}{\phi_2(z)} I_n \end{pmatrix}, \quad (16)$$

where

$$\phi_1(z) := z - \sum_{t \in \llbracket N+1, N+n \rrbracket} G_{tt}(z), \quad \phi_2(z) := z - \sum_{s \in \llbracket 1, N \rrbracket} G_{ss}(z). \quad (17)$$

Under the assumptions of Theorem 7, where $\eta = 54r\sqrt{(K+8)\log(N+n)}$, define a set in the complex plane in the neighborhood of any $\sigma \in \mathbb{R}$ by

$$S_\sigma := \{w \in \mathbb{C} : |\text{Im}(w)| \leq 20\eta r, \sigma - 20\eta r \leq \text{Re}(w) \leq \frac{8}{7}\sigma + 20\eta r\}. \quad (18)$$

In the remainder of this paper, we define an index

$$i_0 := \min\{j \in \llbracket 1, r_0 \rrbracket : \sigma_j \leq n^2\}. \quad (19)$$

Hence, for any the index $l < i_0$, $\sigma_l > n^2$. Note that i_0 may not exist; in this case, $\sigma_j > n^2$ for all $1 \leq j \leq r_0$.

Lemma 9. *Under the assumptions of Theorem 7 and the additional assumption that i_0 defined in (19) exists, one has*

$$\max_{j \in \llbracket i_0, r_0 \rrbracket} \max_{z \in S_{\sigma_j}} |z|^2 \|\mathcal{U}^\top (G(z) - \Phi(z)) \mathcal{U}\| \leq 54r\sqrt{(K+8)\log(N+n)}$$

with probability at least $1 - 10(N+n)^{-(K+1)}$.

Lemma 9 is similar to many isotropic laws for random matrices; see, for instance, [15, 36, 42, 43] and references therein. Roughly speaking, Lemma 9 quantitatively controls how close $G(z)$ is to the diagonal matrix $\Phi(z)$ and will be a fundamental tool in our proofs. Unfortunately, we are not aware of any results in the literature that imply Lemma 9 as stated due to the block structure that \mathcal{E} takes and the particular spectral domain S_σ we are interested in. We present the proof of this lemma in Section 6. The assumption $\sigma_j \leq n^2$ is purely technical: it is used in the proof for carrying out a volume argument on a bounded set. The cutoff n^2 is chosen for convenience. An inspection of the proof reveals the conclusion of Lemma 9 holds for all σ_j 's satisfying $\sigma_j \leq n^c$ for a fixed $c > 1$ by adjusting the tail probability accordingly.

Finally, we collect some basic facts about the matrix $\Phi(z)$. By setting

$$\mathcal{I}^u := \begin{pmatrix} I_N & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad \mathcal{I}^d := \begin{pmatrix} 0 & 0 \\ 0 & I_n \end{pmatrix},$$

one can rewrite (17) as

$$\phi_1(z) = z - \text{tr} \mathcal{I}^d G(z), \quad \phi_2(z) = z - \text{tr} \mathcal{I}^u G(z). \quad (20)$$

It can be derived by elementary linear algebra (see Appendix B.4 for the proof) that

$$\phi_1(z) = \phi_2(z) - \frac{1}{z}(n - N). \quad (21)$$

From the definition of \mathcal{U} in (9), it is easy to verify that

$$\mathcal{U}^T \Phi(z) \mathcal{U} = \begin{pmatrix} \alpha(z) I_r & \beta(z) I_r \\ \beta(z) I_r & \alpha(z) I_r \end{pmatrix}, \quad (22)$$

where we denote

$$\alpha(z) := \frac{1}{2} \left(\frac{1}{\phi_1(z)} + \frac{1}{\phi_2(z)} \right) \quad \text{and} \quad \beta(z) := \frac{1}{2} \left(\frac{1}{\phi_1(z)} - \frac{1}{\phi_2(z)} \right)$$

for notational brevity. It follows that

$$\|\mathcal{U}^T \Phi(z) \mathcal{U}\| = \max \left\{ \frac{1}{|\phi_1(z)|}, \frac{1}{|\phi_2(z)|} \right\}. \quad (23)$$

Sometimes, we drop the z -dependence of $\alpha(z), \beta(z), \phi_1(z), \phi_2(z)$ and simply write $\alpha, \beta, \phi_1, \phi_2$ when the context is clear.

The following technical results can be derived via basic linear algebra and the proofs are deferred to Appendix B.

Proposition 10. *For $1 \leq r_0 < r$, denote the index sets $I := \llbracket 1, r_0 \rrbracket \cup \llbracket r+1, r+r_0 \rrbracket$ and $J := \llbracket 1, 2r \rrbracket \setminus I$. For any $x \in \mathbb{R}$ satisfying $|x| > \|\mathcal{E}\|$, the singular values of $I_{2r-2r_0} - \mathcal{U}_J^T \Phi(x) \mathcal{U}_J \mathcal{D}_J$ are given by*

$$\left| \sqrt{1 + \beta(x)^2 \sigma_t^2} \pm |\alpha(x)| \sigma_t \right|$$

for $r_0 + 1 \leq t \leq r$.

Proposition 11. *For any $z \in \mathbb{C}$ satisfying $|z| > \|\mathcal{E}\|$, the matrix*

$$\mathcal{D}^{-1} - \mathcal{U}^T \Phi(z) \mathcal{U}$$

is invertible with probability 1, and one has

$$\left\| (\mathcal{D}^{-1} - \mathcal{U}^T \Phi(z) \mathcal{U})^{-1} \right\| = \max_{1 \leq l \leq r} \frac{\sigma_l}{|\sigma_l^2 - \phi_1 \phi_2|} \mathcal{Q}^{1/2},$$

where

$$\mathcal{Q} := |\phi_1 \phi_2|^2 + \frac{1}{2} \sigma_l^2 (|\phi_1|^2 + |\phi_2|^2) + \frac{1}{2} \sigma_l [4|\phi_1 \phi_2|^2 |\phi_1 + \bar{\phi}_2|^2 + \sigma_l^2 (|\phi_1|^2 - |\phi_2|^2)^2]^{1/2}.$$

3.3. Singular value locations. In this subsection, we introduce results that describe the precise locations of the singular values of $\mathcal{A} + \mathcal{E}$, and they will play a key role in the proof of Theorem 7.

To this end, consider the random function

$$\varphi(z) := \phi_1(z)\phi_2(z), \quad (24)$$

where $\phi_1(z)$ and $\phi_2(z)$ are defined in (17), and recall the set S_σ for any $\sigma \in \mathbb{R}$ defined in (18).

Theorem 12 (Singular value locations). *Under the assumptions of Theorem 7 and the additional assumption that i_0 defined in (19) exists, for any $i_0 \leq j \leq r_0$, if σ_j has multiplicity α_j and is denoted by $\sigma_j = \sigma_{j+1} = \dots = \sigma_{j+\alpha_j-1}$, then $\tilde{\sigma}_j, \tilde{\sigma}_{j+1}, \dots, \tilde{\sigma}_{j+\alpha_j-1}$ are in the set S_{σ_j} specified in (18), and*

$$|\varphi(\tilde{\sigma}_{j+s}) - \sigma_{j+s}^2| \leq \frac{115}{4}\eta r \left(\tilde{\sigma}_{j+s} + \frac{8}{7}\sigma_{j+s} \right) \quad \text{for } 0 \leq s \leq \alpha_j - 1 \quad (25)$$

with probability at least $1 - 10(N+n)^{-K}$.

Since $\varphi(z) = z^2 + \varepsilon(z)$, where $\varepsilon(z)$ is a random term depending on z and the resolvent $G(z)$, Theorem 12 allows us to approximate the (squared) singular values of \tilde{A} with those of A , up to the random $\varepsilon(z)$ correction term. While this random correction term may seem odd at first, it allows for the much sharper bound appearing on the right-hand side of (25), which in many cases is a significant improvement over classic, deterministic bounds (such as Weyl's inequality). Intuitively, since the singular values of \tilde{A} are random, it makes sense that one cannot only use deterministic values to accurately predict their locations. One of the key differences between the techniques in this paper and those in [51] is that we take into account the precise behavior of this random correction term.

The singular values considered in Theorem 12 are no larger than n^2 . If a singular value σ_l is sufficiently large, the effect of the noise \mathcal{E} is negligible compared to the strong signal and consequently, the location of $\tilde{\sigma}_l$ is very close to σ_l . The next lemma provides the perturbed singular value locations for large singular values. We defer its proof to Appendix B.5.

Lemma 13. *Under the assumptions of Theorem 7,*

$$\max_{l \in [1, r_0]: \sigma_l > \frac{1}{2}n^2} |\tilde{\sigma}_l - \sigma_l| \leq \eta r$$

with probability at least $1 - (N+n)^{-2r^4(K+8)}$.

3.4. Additional tools. We now present a few additional tools we will need in the proofs. The first lemma captures the tail behavior for sub-exponential random variables. A random variable X with mean $\mu = \mathbb{E}X$ is *sub-exponential* with non-negative parameters (ν^2, α) if

$$\mathbb{E} \left(e^{\lambda(X-\mu)} \right) \leq e^{\frac{\nu^2\lambda^2}{2}} \quad \text{for all } |\lambda| < \frac{1}{\alpha}.$$

It is easy to verify that for a standard Gaussian random variable Z , Z^2 is sub-exponential with parameters $(4, 4)$, i.e.

$$\mathbb{E} \left(e^{\lambda(Z^2-1)} \right) \leq e^{\frac{4\lambda^2}{2}}, \quad \text{for all } |\lambda| < \frac{1}{4}. \quad (26)$$

Lemma 14 (Bernstein's inequality for sub-exponential random variables; see Proposition 2.9 in [68]). *Suppose that X is sub-exponential with parameters (ν^2, α) . Then*

$$\mathbb{P}(|X - \mathbb{E}X| \geq t) \leq 2 \exp\left(-\frac{1}{2} \min\left\{\frac{t^2}{\nu^2}, \frac{t}{\alpha}\right\}\right).$$

The following lemma provides a non-asymptotic bound for the spectral norm of Gaussian matrices.

Lemma 15 (Spectral norm bound; see (2.3) from [56]). *Let E be an $N \times n$ matrix whose entries are independent standard Gaussian random variables. Then*

$$\|E\| \leq 2(\sqrt{N} + \sqrt{n})$$

with probability at least $1 - 2e^{-(\sqrt{N} + \sqrt{n})^2/2}$. More generally,

$$\mathbb{P}(\|E\| \leq \sqrt{N} + \sqrt{n} + t) \geq 1 - 2e^{-t^2/2}.$$

The next lemma bounds the operator norms of the resolvent and is used frequently in the proof. Let $\mathcal{E}^{(k)}$ be the minor of \mathcal{E} with the k th row and column replaced by zeros and $G^{(k)}(z)$ the resolvent of $\mathcal{E}^{(k)}$ (see the precise definition at the beginning of Section 6). The proof is given in Appendix C.

Lemma 16. *On the event where $\|E\| \leq 2(\sqrt{N} + \sqrt{n})$,*

$$\|G(z)\| \leq \frac{2}{|z|}, \quad \|G^{(k)}(z)\| \leq \frac{2}{|z|}$$

and

$$\frac{7}{8}|z| \leq |\phi_i(z)| \leq \frac{9}{8}|z| \quad \text{for } i = 1, 2 \tag{27}$$

for any $z \in \mathbb{C}$ with $|z| \geq 4(\sqrt{N} + \sqrt{n})$ and for any $k \in \llbracket 1, N + n \rrbracket$.

The following lemma suggests that when $|z|$ is large, the resolvent $G(z)$ can be well approximated by a simple matrix. We defer its proof to Appendix C.

Lemma 17. *On the event where $\|E\| \leq 2(\sqrt{N} + \sqrt{n})$,*

$$\left\| G(z) - \frac{1}{z} I_{N+n} - \frac{\mathcal{E}}{z^2} \right\| \leq \frac{2\|\mathcal{E}\|^2}{|z|^3}$$

for any $z \in \mathbb{C}$ with $|z| \geq 4(\sqrt{N} + \sqrt{n})$.

The next lemma bounds the operator norm of the random matrix $\mathcal{U}^T \mathcal{E} \mathcal{U}$ of size $2r \times 2r$ where \mathcal{U} is defined in (9). Its proof can be found in Appendix C.

Lemma 18. *Let K be an arbitrary positive constant. With probability at least $1 - 2(N + n)^{-K}$, we have*

$$\|\mathcal{U}^T \mathcal{E} \mathcal{U}\| \leq 2\sqrt{r} + \sqrt{2K \log(N + n)}.$$

3.5. Overview of the proofs. We now provide a brief overview of the proofs of our main results. For simplicity, we focus on the proof of Theorem 4, although the proof of Theorem 7 is similar.

As noted above, Proposition 8 allows us to focus on the eigenvectors of \mathcal{A} and $\tilde{\mathcal{A}}$. In this case, we have

$$\sin^2 \angle(\mathbf{u}_1, \tilde{\mathbf{u}}_1) = 1 - |\mathbf{u}_1^T \tilde{\mathbf{u}}_1|^2 = \sum_{j \neq 1} |\mathbf{u}_1^T \tilde{\mathbf{u}}_j|^2.$$

Thus, our task reduces to controlling the inner products $\mathbf{u}_1^T \tilde{\mathbf{u}}_j$ for $j \neq 1$. For simplicity, let us focus on the case when $j = 2$. We start with the eigenvector-eigenvalue equation

$$\tilde{\mathcal{A}}\tilde{\mathbf{u}}_2 = \tilde{\lambda}_2\tilde{\mathbf{u}}_2.$$

Rearranging this equation and using that $\tilde{\mathcal{A}} = \mathcal{A} + \mathcal{E}$, we arrive at

$$\tilde{\mathbf{u}}_2 = G(\tilde{\lambda}_2)\mathcal{A}\tilde{\mathbf{u}}_2, \quad (28)$$

provided $|\tilde{\lambda}_2| > \|\mathcal{E}\|$. Multiplying (28) by \mathbf{u}_1^T , we find

$$\mathbf{u}_1^T \tilde{\mathbf{u}}_2 = \mathbf{u}_1^T G(\tilde{\lambda}_2)\mathcal{U}\mathcal{D}\mathcal{U}^T \tilde{\mathbf{u}}_2, \quad (29)$$

where we used the spectral decomposition given in (9). Using Lemma 9, the resolvent matrix on the right-hand side of (29) can be approximated by the matrix $\Phi(\tilde{\lambda}_2)$. Accordingly, we split the right-hand side of (29) into two terms

$$\mathbf{u}_1^T \tilde{\mathbf{u}}_2 = \mathbf{u}_1^T \Phi(\tilde{\lambda}_2)\mathcal{U}\mathcal{D}\mathcal{U}^T \tilde{\mathbf{u}}_2 + \mathbf{u}_1^T \left(G(\tilde{\lambda}_2) - \Phi(\tilde{\lambda}_2) \right) \mathcal{U} \cdot \mathcal{D}\mathcal{U}^T \tilde{\mathbf{u}}_2, \quad (30)$$

and Lemma 9 facilitates precise control over the second term appearing on the right-hand side of (30). In addition, we see that the term $\mathcal{U}^T \tilde{\mathbf{u}}_2$ appearing in the first term on the right-hand side of (30) contains a copy of the inner product $\mathbf{u}_1 \cdot \tilde{\mathbf{u}}_2$ as one of its entries. This will allow us to turn (29) into a recursive equation, with the inner product appearing on both sides of the equation. More precisely, from the spectral decomposition

$$\mathcal{A} = \mathcal{U}\mathcal{D}\mathcal{U}^T = \lambda_1 \mathbf{u}_1 \mathbf{u}_1^T + \lambda_{r+1} \mathbf{u}_{r+1} \mathbf{u}_{r+1}^T + \mathcal{U}_0 \mathcal{D}_0 \mathcal{U}_0^T,$$

where \mathcal{U}_0 is obtained from \mathcal{U} by removing the columns \mathbf{u}_1 and \mathbf{u}_{r+1} and \mathcal{D}_0 from \mathcal{D} by removing the rows/columns containing λ_1 and λ_{r+1} . Hence, from (22), we arrive at

$$\mathbf{u}_1^T \Phi(\tilde{\lambda}_2)\mathcal{U}\mathcal{D}\mathcal{U}^T \tilde{\mathbf{u}}_2 = \lambda_1 \alpha(\tilde{\lambda}_2) \mathbf{u}_1^T \tilde{\mathbf{u}}_2 + \lambda_{r+1} \beta(\tilde{\lambda}_2) \mathbf{u}_{r+1}^T \tilde{\mathbf{u}}_2.$$

Plugging the above equation back into (30) and rearranging the terms, we see that

$$\left(1 - \lambda_1 \alpha(\tilde{\lambda}_2) \right) \mathbf{u}_1^T \tilde{\mathbf{u}}_2 = \lambda_{r+1} \beta(\tilde{\lambda}_2) \mathbf{u}_{r+1}^T \tilde{\mathbf{u}}_2 + \mathbf{u}_1^T \left(G(\tilde{\lambda}_2) - \Phi(\tilde{\lambda}_2) \right) \mathcal{U} \cdot \mathcal{D}\mathcal{U}^T \tilde{\mathbf{u}}_2. \quad (31)$$

Understanding the location of $\tilde{\lambda}_2$ is key to analyzing the left-hand side of (31), and Theorem 12 will allow us to accurately estimate

$$1 - \lambda_1 \alpha(\tilde{\lambda}_2) = 1 - \sigma_1 \alpha(\tilde{\sigma}_2) \approx 1 - \sigma_1 / \sigma_2;$$

this estimate explains the appearance of the spectral gap δ_1 in the results.

Actually, the term $\mathbf{u}_{r+1}^T \tilde{\mathbf{u}}_2$ on the right-hand side of (31) also encompasses information regarding $\mathbf{u}_1^T \tilde{\mathbf{u}}_2$ by the definition of the $\tilde{\mathbf{u}}_j$'s and should not be overlooked. Besides, it is also wasteful to estimate each inner product $|\mathbf{u}_1^T \tilde{\mathbf{u}}_j|$ separately. Instead, when proving Theorem 7, we will group the inner products $|\mathbf{u}_1^T \tilde{\mathbf{u}}_j|$ into two categories: when $j \in \llbracket r_0 + 1, r \rrbracket \cup \llbracket r + r_0 + 1, 2r \rrbracket$ and when $j > 2r$. The arguments for the first category are similar to the methods discussed above (see Lemma 20). The bound for the second category is much simpler and results in the signal-to-noise ratio term appearing in Theorem 7 (see Lemma 19).

4. PROOF OF THEOREM 7

In this section, we prove Theorem 7 using Theorem 12 and Lemma 9. In view of (11), it suffices to bound $\sin \angle(\mathcal{U}_I, \tilde{\mathcal{U}}_I)$ for $I := \llbracket 1, r_0 \rrbracket \cup \llbracket r+1, r+r_0 \rrbracket$. Assume the null space of \mathcal{A} is spanned by the orthonormal basis $\{\mathbf{u}_{2r+1}, \dots, \mathbf{u}_{N+n}\}$; in general, there will be many choices for this orthonormal basis and any choice will suffice for our purposes.

The main idea of the proof is to divide the bound on $\sin \angle(\mathcal{U}_I, \tilde{\mathcal{U}}_I)$ into two parts. The first part involves projections of the vectors $\mathbf{u}_{r_0+1}, \dots, \mathbf{u}_r, \mathbf{u}_{r+r_0+1}, \dots, \mathbf{u}_{2r}$ while the second part involves projections of the vectors $\mathbf{u}_{2r+1}, \dots, \mathbf{u}_{n+N}$. The latter terms can be controlled using the noise-to-signal ratio $\|E\|/\sigma_{r_0}$. The main argument is the estimate of the first term, which reflects that when the signal is stronger than the noise, the action of \mathcal{E} is essentially on the $2r$ -dimensional subspace spanned by the eigenvectors of \mathcal{A} corresponding to the non-trivial eigenvalues. We use the resolvent $G(z)$ to extract information about the perturbed eigenvectors with the aid of Lemma 9 and Theorem 12.

If $r_0 = r$, then $\delta_r = \sigma_r$ and the conclusion follows from the Davis–Kahan–Wedin sine Θ theorem [25, 70], which implies that

$$\max\{\sin \angle(U_r, \tilde{U}_r), \sin \angle(V_r, \tilde{V}_r)\} \leq 2 \frac{\|E\|}{\sigma_r}.$$

Thus we assume $1 \leq r_0 \leq r-1$ throughout the proof. For the remainder of this section, we fix the index sets

$$I := \llbracket 1, r_0 \rrbracket \cup \llbracket r+1, r+r_0 \rrbracket$$

and

$$J := \llbracket 1, 2r \rrbracket \setminus I = \llbracket r_0+1, r \rrbracket \cup \llbracket r+r_0+1, 2r \rrbracket.$$

Lemma 19. *Under the assumptions of Theorem 7,*

$$\sin \angle(\mathcal{U}_I, \tilde{\mathcal{U}}_I) := \|P_I - \tilde{P}_I\| \leq \|P_J \tilde{P}_I\| + 2 \frac{\|E\|}{\sigma_{r_0}}$$

with probability 1.

Proof. We assume $\sigma_{r_0} > 2\|E\|$ as the bound is trivial otherwise. In view of [13, Exercise VII.1.11],

$$\begin{aligned} \|P_I - \tilde{P}_I\| &= \|P_{I^c} \tilde{P}_I\| \\ &\leq \|P_J \tilde{P}_I\| + \|P_{\llbracket 2r+1, N+n \rrbracket} \tilde{P}_I\|. \end{aligned}$$

We aim to bound

$$\|P_{\llbracket 2r+1, N+n \rrbracket} \tilde{P}_I\| \leq \|\mathcal{U}_{\llbracket 2r+1, N+n \rrbracket}^T \tilde{\mathcal{U}}_I\|.$$

From the spectral decomposition of $\tilde{\mathcal{A}}$, we have

$$(\mathcal{A} + \mathcal{E})\tilde{\mathcal{U}}_I = \tilde{\mathcal{U}}_I \tilde{\mathcal{D}}_I.$$

Multiplying by $\mathcal{U}_{\llbracket 2r+1, N+n \rrbracket}^T$ on the left of the equation above, we further have

$$\mathcal{U}_{\llbracket 2r+1, N+n \rrbracket}^T \mathcal{E} \tilde{\mathcal{U}}_I = \mathcal{U}_{\llbracket 2r+1, N+n \rrbracket}^T \tilde{\mathcal{U}}_I \tilde{\mathcal{D}}_I.$$

As $\sigma_{r_0} > 2\|E\|$ by supposition, Weyl's inequality (14) implies that

$$\tilde{\sigma}_i \geq \sigma_i - \|E\| \geq \frac{1}{2} \sigma_i \tag{32}$$

for $1 \leq i \leq r_0$. Hence, $\tilde{\mathcal{D}}_I$ is invertible since $|\tilde{\lambda}_i| = \tilde{\sigma}_i \geq \sigma_i - \|E\| > 0$ for $i \in \llbracket 1, r_0 \rrbracket$ and $|\tilde{\lambda}_i| = \tilde{\sigma}_{i-r} > 0$ for $i \in \llbracket r+1, r+r_0 \rrbracket$. It follows that

$$\|\mathcal{U}_{\llbracket 2r+1, N+n \rrbracket}^T \tilde{\mathcal{U}}_I\| = \|\mathcal{U}_{\llbracket 2r+1, N+n \rrbracket}^T \mathcal{E} \tilde{\mathcal{U}}_I \tilde{\mathcal{D}}_I^{-1}\| \leq \|\mathcal{E}\| \|\tilde{\mathcal{D}}_I^{-1}\| = \frac{\|E\|}{\tilde{\sigma}_{r_0}}.$$

Thus by another application of (32), we get

$$\|P_{\llbracket 2r+1, N+n \rrbracket} \tilde{P}_I\| \leq \frac{\|E\|}{\tilde{\sigma}_{r_0}} \leq 2 \frac{\|E\|}{\sigma_{r_0}}. \quad (33)$$

as desired. \square

It remains to bound $\|P_J \tilde{P}_I\|$, which is the content of the following lemma.

Lemma 20. *Under the assumptions of Theorem 7, we have*

$$\|P_J \tilde{P}_I\| \leq 21\sqrt{2}\eta \sqrt{\sum_{j=1}^{r_0} \frac{1}{(\sigma_j - \sigma_{r_0+1})^2}}$$

with probability at least $1 - 15(N+n)^{-K}$.

Proof. In the following, we work on the event where $\|E\| \leq 2(\sqrt{N} + \sqrt{n})$; Lemma 15 shows this event holds with probability at least $1 - 2e^{-(\sqrt{N} + \sqrt{n})^2/2} \geq 1 - 2(N+n)^{-32(K+2)}$ since $(\sqrt{N} + \sqrt{n})^2 > 64(K+2) \log(N+n)$ by assumption. We start with the bound

$$\|P_J \tilde{P}_I\| \leq \|\mathcal{U}_J^T \tilde{\mathcal{U}}_I\| \leq \|\mathcal{U}_J^T \tilde{\mathcal{U}}_I\|_F = \sqrt{\sum_{i \in I} \|\mathcal{U}_J^T \tilde{\mathbf{u}}_i\|^2}. \quad (34)$$

Now we estimate $\|\mathcal{U}_J^T \tilde{\mathbf{u}}_i\|$ for each $i \in I$. We split the index set I into two disjoint sets:

$$\mathcal{I}_s := \{i \in I : |\lambda_i| \leq n^2\} \quad \text{and} \quad \mathcal{I}_b := \{i \in I : |\lambda_i| > n^2\}.$$

Note that \mathcal{I}_s or \mathcal{I}_b could be the empty set.

Case 1: estimate $\|\mathcal{U}_J^T \tilde{\mathbf{u}}_i\|$ for $i \in \mathcal{I}_s$. We first obtain an identity for the eigenvector $\tilde{\mathbf{u}}_i$. By Weyl's inequality, $|\tilde{\lambda}_i| \geq \tilde{\sigma}_{r_0} \geq \sigma_{r_0} - \|E\| > \|E\| = \|\mathcal{E}\|$ by supposition on σ_{r_0} and thus $G(\tilde{\lambda}_i)$ and $\Phi(\tilde{\lambda}_i)$ are well-defined. As $(\mathcal{A} + \mathcal{E})\tilde{\mathbf{u}}_i = \tilde{\lambda}_i \tilde{\mathbf{u}}_i$, we solve for $\tilde{\mathbf{u}}_i$ to obtain

$$\tilde{\mathbf{u}}_i = (\tilde{\lambda}_i I - \mathcal{E})^{-1} \mathcal{A} \tilde{\mathbf{u}}_i = G(\tilde{\lambda}_i) \mathcal{A} \tilde{\mathbf{u}}_i.$$

Rewrite the above equation

$$\tilde{\mathbf{u}}_i = \Phi(\tilde{\lambda}_i) \mathcal{A} \tilde{\mathbf{u}}_i + (G(\tilde{\lambda}_i) - \Phi(\tilde{\lambda}_i)) \mathcal{A} \tilde{\mathbf{u}}_i$$

and multiply on the left by \mathcal{U}_J^T to get

$$\mathcal{U}_J^T \tilde{\mathbf{u}}_i = \mathcal{U}_J^T \Phi(\tilde{\lambda}_i) \mathcal{A} \tilde{\mathbf{u}}_i + \mathcal{U}_J^T (G(\tilde{\lambda}_i) - \Phi(\tilde{\lambda}_i)) \mathcal{A} \tilde{\mathbf{u}}_i. \quad (35)$$

Plugging in (10), we further have

$$\mathcal{U}_J^T \tilde{\mathbf{u}}_i = \mathcal{U}_J^T \Phi(\tilde{\lambda}_i) \mathcal{U}_J \mathcal{D}_J \mathcal{U}_J^T \tilde{\mathbf{u}}_i + \mathcal{U}_J^T (G(\tilde{\lambda}_i) - \Phi(\tilde{\lambda}_i)) \mathcal{U} \mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i,$$

where we used $\mathcal{U}_J^T \Phi(\tilde{\lambda}_i) \mathcal{U}_I = 0$. Hence,

$$\left(I_{2r-2r_0} - \mathcal{U}_J^T \Phi(\tilde{\lambda}_i) \mathcal{U}_J \mathcal{D}_J \right) \mathcal{U}_J^T \tilde{\mathbf{u}}_i = \mathcal{U}_J^T (G(\tilde{\lambda}_i) - \Phi(\tilde{\lambda}_i)) \mathcal{U} \mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i. \quad (36)$$

We are now in position to bound $\|\mathcal{U}_J^T \tilde{\mathbf{u}}_i\|$. This can be achieved by obtaining an upper bound for the right-hand side of (36) and estimating the smallest singular value of the matrix

$$I_{2r-2r_0} - \mathcal{U}_J^T \Phi(\tilde{\lambda}_i) \mathcal{U}_J \mathcal{D}_J \quad (37)$$

on the left-hand side of (36). We establish these estimates in the following two steps. Recall the index i_0 from (19). We will work on the event $\mathbf{E} := \cap_{i=i_0}^{r_0} \mathbf{E}_i$ where

$$\mathbf{E}_i := \left\{ \tilde{\sigma}_i \in S_{\sigma_i}, \|\mathcal{U}^T (G(\tilde{\sigma}_i) - \Phi(\tilde{\sigma}_i)) \mathcal{U}\| \leq \frac{\eta}{\tilde{\sigma}_i^2}, |\phi_1(\tilde{\sigma}_i) \phi_2(\tilde{\sigma}_i) - \sigma_i^2| \leq \frac{115}{4} \eta r (\tilde{\sigma}_i + \frac{8}{7} \sigma_i) \right\}. \quad (38)$$

By Theorem 12 and Lemma 9, the event \mathbf{E} holds with probability at least $1 - 12(N+n)^{-K}$.

Step 1. Upper bound for the right-hand side of (36). Recall $I = \llbracket 1, r_0 \rrbracket \cup \llbracket r+1, r+r_0 \rrbracket$. We first consider the case when $i \in \llbracket 1, r_0 \rrbracket$ and $\tilde{\lambda}_i = \tilde{\sigma}_i$. Note that $\mathcal{U}_J^T (G(\tilde{\sigma}_i) - \Phi(\tilde{\sigma}_i)) \mathcal{U}$ is a sub-matrix of $\mathcal{U}^T (G(\tilde{\sigma}_i) - \Phi(\tilde{\sigma}_i)) \mathcal{U}$. Thus, using (38) and the fact that the spectral norm of any sub-matrix is bounded by the spectral norm of the full matrix, we deduce that

$$\|\mathcal{U}_J^T (G(\tilde{\sigma}_i) - \Phi(\tilde{\sigma}_i)) \mathcal{U} \mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i\| \leq \frac{\eta}{\tilde{\sigma}_i^2} \|\mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i\|.$$

Observe from $(\mathcal{A} + \mathcal{E}) \tilde{\mathbf{u}}_i = \tilde{\sigma}_i \tilde{\mathbf{u}}_i$ that $\mathcal{U} \mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i = (\tilde{\sigma}_i I - \mathcal{E}) \tilde{\mathbf{u}}_i$. Multiplying \mathcal{U}^T on both sides, we get the bound

$$\|\mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i\| \leq \|E\| + \tilde{\sigma}_i \leq 2\sigma_i \quad (39)$$

using the assumption $\|E\| \leq \frac{1}{2} \sigma_i$ and Weyl's inequality. Hence,

$$\left\| \mathcal{U}_J^T \left(G(\tilde{\lambda}_i) - \Phi(\tilde{\lambda}_i) \right) \mathcal{U} \mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i \right\| \leq 2 \frac{\eta \sigma_i}{\tilde{\sigma}_i^2}. \quad (40)$$

For the case when $i \in \llbracket r+1, r+r_0 \rrbracket$, $\tilde{\lambda}_i = -\tilde{\sigma}_{i-r}$. Observe that

$$G(-\tilde{\sigma}_{i-r}) = (-\tilde{\sigma}_{i-r} - \mathcal{E})^{-1} = -(\tilde{\sigma}_{i-r} + \mathcal{E})^{-1} \sim -(\tilde{\sigma}_{i-r} - \mathcal{E})^{-1} = -G(\tilde{\sigma}_{i-r})$$

because the distribution of \mathcal{E} is symmetric. Hence

$$\Phi(-\tilde{\sigma}_{i-r}) \sim -\Phi(\tilde{\sigma}_{i-r})$$

by the definition (16). Repeating the arguments from the previous case, we see that

$$\left\| \mathcal{U}_J^T \left(G(\tilde{\lambda}_i) - \Phi(\tilde{\lambda}_i) \right) \mathcal{U} \mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i \right\| \leq 2 \frac{\eta \sigma_{i-r}}{\tilde{\sigma}_{i-r}^2}. \quad (41)$$

Step 2. Lower bound for the smallest singular value of the matrix (37). By Proposition 10, the singular values of $I_{2r-2r_0} - \mathcal{U}_J^T \Phi(\tilde{\lambda}_i) \mathcal{U}_J \mathcal{D}_J$ are given by

$$\left| \sqrt{1 + \beta(\tilde{\lambda}_i)^2 \sigma_t^2} \pm |\alpha(\tilde{\lambda}_i)| \sigma_t \right| \quad (42)$$

for $r_0 + 1 \leq t \leq r$.

In order to bound the singular values, we first estimate $\phi_1(\tilde{\sigma}_i) \phi_2(\tilde{\sigma}_i)$, $\phi_1(\tilde{\sigma}_i)$ and $\phi_2(\tilde{\sigma}_i)$ for $i_0 \leq i \leq r_0$. Since $\tilde{\sigma}_i \in S_{\sigma_i}$ by (38) where S_{σ_i} is defined in (18), we have

$$\tilde{\sigma}_i \geq \sigma_i - 20\eta r \geq 4(\sqrt{N} + \sqrt{n})$$

and

$$\frac{9}{7}\sigma_i \geq \frac{8}{7}\sigma_i + 20\eta r \geq \tilde{\sigma}_i \geq \sigma_i - 20\eta r \geq \frac{6}{7}\sigma_i$$

by the supposition $\sigma_i \geq 4(\sqrt{N} + \sqrt{n}) + 140\eta r$.

Continuing from (38), we have

$$|\phi_1(\tilde{\sigma}_i)\phi_2(\tilde{\sigma}_i) - \sigma_i^2| \leq \frac{1955}{28}\eta r\sigma_i \quad (43)$$

and consequently, $\frac{393}{784}\sigma_i^2 \leq \phi_1(\tilde{\sigma}_i)\phi_2(\tilde{\sigma}_i) \leq \frac{1175}{784}\sigma_i^2$ from the supposition that $\sigma_i \geq 140\eta r$. Observe from (27) that

$$\frac{7}{8}\tilde{\sigma}_i \leq \phi_1(\tilde{\sigma}_i) \leq \frac{9}{8}\tilde{\sigma}_i. \quad (44)$$

Thus

$$\frac{3}{4}\sigma_i \leq \phi_1(\tilde{\sigma}_i) \leq \frac{81}{56}\sigma_i.$$

The same bound also holds from $\phi_2(\tilde{\sigma}_i)$. Using these estimates, we crudely bound

$$0 < \alpha(\tilde{\sigma}_i) = \frac{1}{2} \left(\frac{1}{\phi_1(\tilde{\sigma}_i)} + \frac{1}{\phi_2(\tilde{\sigma}_i)} \right) \leq \frac{4}{3\sigma_i},$$

and with (21)

$$\beta(\tilde{\sigma}_i) = \frac{1}{2} \left(\frac{1}{\phi_1(\tilde{\sigma}_i)} - \frac{1}{\phi_2(\tilde{\sigma}_i)} \right) = \frac{\phi_2(\tilde{\sigma}_i) - \phi_1(\tilde{\sigma}_i)}{2\phi_1(\tilde{\sigma}_i)\phi_2(\tilde{\sigma}_i)} = \frac{n - N}{\tilde{\sigma}_i} \frac{1}{2\phi_1(\tilde{\sigma}_i)\phi_2(\tilde{\sigma}_i)} \leq \frac{49}{786\sigma_i}$$

by noting that $\tilde{\sigma}_i, \sigma_i \geq 4(\sqrt{N} + \sqrt{n})$.

We are ready to bound the singular values of $I_{2r-r_0} - \mathcal{U}_J^\top \Phi(\tilde{\sigma}_i) \mathcal{U}_J \mathcal{D}_J$. We start with the case when $i \in \llbracket 1, r_0 \rrbracket$ and $\tilde{\lambda}_i = \tilde{\sigma}_i$. In view of (42), the goal is to bound

$$\begin{aligned} \min_{r_0+1 \leq t \leq r} \left| \sqrt{1 + \beta(\tilde{\sigma}_i)^2 \sigma_t^2} \pm |\alpha(\tilde{\sigma}_i)| \sigma_t \right| &= \min_{r_0+1 \leq t \leq r} \left| \sqrt{1 + \beta(\tilde{\sigma}_i)^2 \sigma_t^2} - \alpha(\tilde{\sigma}_i) \sigma_t \right| \\ &= \min_{r_0+1 \leq t \leq r} \left| \frac{1 - (\alpha(\tilde{\sigma}_i)^2 - \beta(\tilde{\sigma}_i)^2) \sigma_t^2}{\sqrt{1 + \beta(\tilde{\sigma}_i)^2 \sigma_t^2} + \alpha(\tilde{\sigma}_i) \sigma_t} \right| \\ &= \min_{r_0+1 \leq t \leq r} \frac{\left| 1 - \frac{\sigma_t^2}{\phi_1(\tilde{\sigma}_i)\phi_2(\tilde{\sigma}_i)} \right|}{\sqrt{1 + \beta(\tilde{\sigma}_i)^2 \sigma_t^2} + \alpha(\tilde{\sigma}_i) \sigma_t}. \end{aligned}$$

The upper bounds of $\alpha(\tilde{\sigma}_i)$ and $\beta(\tilde{\sigma}_i)$ obtained above, together with $\sigma_t/\sigma_i \leq 1$, yield that

$$\sqrt{1 + \beta(\tilde{\sigma}_i)^2 \sigma_t^2} + \alpha(\tilde{\sigma}_i) \sigma_t \leq \sqrt{1 + \left(\frac{49}{786} \right)^2 \frac{\sigma_t^2}{\sigma_i^2}} + \frac{4\sigma_t}{3\sigma_i} \leq 2.5$$

for any $r_0 + 1 \leq t \leq r$. Hence,

$$\begin{aligned} \min_{r_0+1 \leq t \leq r} \left| \sqrt{1 + \beta(\tilde{\sigma}_i)^2 \sigma_t^2} \pm \alpha(\tilde{\sigma}_i) \sigma_t \right| &\geq \frac{1}{2.5} \min_{r_0+1 \leq t \leq r} \left| \frac{\phi_1(\tilde{\sigma}_i)\phi_2(\tilde{\sigma}_i) - \sigma_t^2}{\phi_1(\tilde{\sigma}_i)\phi_2(\tilde{\sigma}_i)} \right| \\ &\geq \frac{1}{2.5} \frac{\sigma_i^2 - \frac{1955}{28}\eta r\sigma_i - \sigma_{r_0+1}^2}{\phi_1(\tilde{\sigma}_i)\phi_2(\tilde{\sigma}_i)}. \end{aligned}$$

In the last inequality above, we invoked (43) which implies

$$\phi_1(\tilde{\sigma}_i)\phi_2(\tilde{\sigma}_i) - \sigma_t^2 \geq \sigma_i^2 - \frac{1955}{28}\eta r\sigma_i - \sigma_{r_0+1}^2 > 0$$

by the supposition $\sigma_i - \sigma_{r_0+1} \geq \delta_{r_0} \geq 100\eta r$. Applying this supposition again, together with (44), we further deduce that

$$\begin{aligned} \min_{r_0+1 \leq t \leq r} \left| \sqrt{1 + \beta(\tilde{\sigma}_i)^2 \sigma_t^2} \pm \alpha(\tilde{\sigma}_i) \sigma_t \right| &\geq \frac{1}{\tilde{\sigma}_i^2} \frac{1}{2.5} \left(\frac{8}{9} \right)^2 \left(\sigma_i^2 - \sigma_{r_0+1}^2 - \frac{1955}{28} \eta r \sigma_i \right) \\ &\geq \frac{\sigma_i^2 - \sigma_{r_0+1}^2}{\tilde{\sigma}_i^2} \frac{1}{2.5} \left(\frac{8}{9} \right)^2 \left(1 - \frac{1955}{28} \frac{1}{100} \right) \\ &\geq 0.0953 \frac{\sigma_i^2 - \sigma_{r_0+1}^2}{\tilde{\sigma}_i^2}. \end{aligned} \quad (45)$$

For the case when $i \in \llbracket r+1, r+r_0 \rrbracket$ and $\tilde{\lambda}_i = -\tilde{\sigma}_{i-r}$. Use the observation that $\alpha(\tilde{\lambda}_i) \sim -\alpha(\tilde{\sigma}_{i-r})$ and $\beta(\tilde{\lambda}_i) \sim -\beta(\tilde{\sigma}_{i-r})$ from the definitions (20). A simple modification of the proof for the first case shows that

$$\min_{r_0+1 \leq t \leq r} \left| \sqrt{1 + \beta(\tilde{\lambda}_i)^2 \sigma_t^2} \pm |\alpha(\tilde{\lambda}_i)| \sigma_t \right| \geq 0.0953 \frac{\sigma_{i-r}^2 - \sigma_{r_0+1}^2}{\tilde{\sigma}_{i-r}^2}. \quad (46)$$

Step 3. Combining the bounds above. With the estimates deduced in the previous two steps, we are in a position to bound $\|\mathcal{U}_J^T \tilde{\mathbf{u}}_i\|$. For $i \in \llbracket 1, r_0 \rrbracket \cap \mathcal{I}_s$, plugging (40) and (45) and into (36), we find that

$$\|\mathcal{U}_J^T \tilde{\mathbf{u}}_i\| \leq \frac{2\eta\sigma_i}{0.0953(\sigma_i^2 - \sigma_{r_0+1}^2)} \leq \frac{21\eta}{\sigma_i - \sigma_{r_0+1}}, \quad (47)$$

and for $i \in \llbracket r+1, r+r_0 \rrbracket \cap \mathcal{I}_s$, using (41) and (46), we get

$$\|\mathcal{U}_J^T \tilde{\mathbf{u}}_i\| \leq \frac{21\eta}{\sigma_{i-r} - \sigma_{r_0+1}}. \quad (48)$$

Case 2: estimate $\|\mathcal{U}_J^T \tilde{\mathbf{u}}_i\|$ for $i \in \mathcal{I}_b$. By Weyl's inequality, $|\tilde{\lambda}_i| \geq n^2 - \|\mathcal{E}\| \geq 4(\sqrt{N} + \sqrt{n})$ for every $i \in \mathcal{I}_b$. Hence, we apply Lemma 17 to get

$$\left\| G(\tilde{\lambda}_i) - \Psi(\tilde{\lambda}_i) \right\| \leq \frac{2\|\mathcal{E}\|^2}{|\tilde{\lambda}_i|^3} \quad (49)$$

where $\Psi(z) := \frac{1}{z} I_{N+n} + \frac{1}{z^2} \mathcal{E}$. Repeating the arguments as in the beginning of Case 1, we obtain the following equation similar to (35):

$$\mathcal{U}_J^T \tilde{\mathbf{u}}_i = \mathcal{U}_J^T \Psi(\tilde{\lambda}_i) \mathcal{A} \tilde{\mathbf{u}}_i + \mathcal{U}_J^T \left(G(\tilde{\lambda}_i) - \Psi(\tilde{\lambda}_i) \right) \mathcal{A} \tilde{\mathbf{u}}_i.$$

Plugging in (10) and using the facts $\mathcal{U}_J^T \mathcal{U}_I = 0$ and $\mathcal{U}_J^T \mathcal{U}_J = I$, we further get

$$\mathcal{U}_J^T \tilde{\mathbf{u}}_i = \frac{1}{\tilde{\lambda}_i} \mathcal{D}_J \mathcal{U}_J^T \tilde{\mathbf{u}}_i + \frac{1}{\tilde{\lambda}_i^2} \mathcal{U}_J^T \mathcal{E} \mathcal{U} \mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i + \mathcal{U}_J^T \left(G(\tilde{\lambda}_i) - \Psi(\tilde{\lambda}_i) \right) \mathcal{U} \mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i,$$

which, by rearranging the terms, is reduced to

$$(\tilde{\lambda}_i I - \mathcal{D}_J) \mathcal{U}_J^T \tilde{\mathbf{u}}_i = \frac{1}{\tilde{\lambda}_i} \mathcal{U}_J^T \mathcal{E} \mathcal{U} \mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i + \tilde{\lambda}_i \mathcal{U}_J^T \left(G(\tilde{\lambda}_i) - \Psi(\tilde{\lambda}_i) \right) \mathcal{U} \mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i.$$

Hence,

$$\min_{j \in J} |\tilde{\lambda}_i - \lambda_j| \cdot \|\mathcal{U}_J^T \tilde{\mathbf{u}}_i\| \leq \frac{1}{|\tilde{\lambda}_i|} \|\mathcal{U}^T \mathcal{E} \mathcal{U}\| \cdot \|\mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i\| + |\tilde{\lambda}_i| \|G(\tilde{\lambda}_i) - \Psi(\tilde{\lambda}_i)\| \cdot \|\mathcal{D} \mathcal{U}^T \tilde{\mathbf{u}}_i\|.$$

Note that $\|\mathcal{D}\mathcal{U}^T\tilde{\mathbf{u}}_i\| \leq \|\mathcal{E}\| + |\tilde{\lambda}_i| \leq 2|\tilde{\lambda}_i|$ as in (39). Inserting (49) into the above inequality, we arrive at

$$\min_{j \in \mathcal{J}} |\tilde{\lambda}_i - \lambda_j| \cdot \|\mathcal{U}_j^T \tilde{\mathbf{u}}_i\| \leq 2\|\mathcal{U}^T \mathcal{E}\mathcal{U}\| + 4 \frac{\|E\|^2}{|\tilde{\lambda}_i|}. \quad (50)$$

For the remaining arguments, we work on the event

$$\mathbf{F} := \left\{ \|\mathcal{U}^T \mathcal{E}\mathcal{U}\| \leq 2\sqrt{r} + \sqrt{10(K+8)\log(N+n)} \right\} \cap \left\{ \max_{i \in \llbracket 1, r_0 \rrbracket; \sigma_i > \frac{1}{2}n^2} |\tilde{\sigma}_i - \sigma_i| \leq \eta r \right\}. \quad (51)$$

By Lemma 13 and Lemma 18, the event \mathbf{F} holds with probability at least

$$1 - 2(N+n)^{-5(K+8)} - (N+n)^{-2r^4(K+8)} \geq 1 - 3(N+n)^{-2(K+8)}.$$

We continue the estimation of $\|\mathcal{U}_j^T \tilde{\mathbf{u}}_i\|$ from (50). Note from (51) that

$$\|\mathcal{U}^T \mathcal{E}\mathcal{U}\| \leq 2\sqrt{r} + \sqrt{10(K+8)\log(N+n)} \leq \eta.$$

Also, $\|E\|^2/|\tilde{\lambda}_i| \leq 8(2\sqrt{n})^2/n^2 \leq \eta$ where we used the crude bound $|\tilde{\lambda}_i| \geq \frac{1}{2}n^2$ by Weyl's inequality. It follows that

$$\min_{j \in \mathcal{J}} |\tilde{\lambda}_i - \lambda_j| \cdot \|\mathcal{U}_j^T \tilde{\mathbf{u}}_i\| \leq 6\eta. \quad (52)$$

To bound the left-hand side of (52), we first consider $i \in \llbracket 1, r_0 \rrbracket \cap \mathcal{I}_b$. Then

$$\min_{j \in \mathcal{J}} |\tilde{\lambda}_i - \lambda_j| = \min_{r_0+1 \leq j \leq r} |\tilde{\sigma}_i - \sigma_j| = \tilde{\sigma}_i - \sigma_{r_0+1}$$

by (51) and the supposition $\delta_{r_0} = \sigma_{r_0} - \sigma_{r_0+1} \geq 100\eta r$. Next, applying $\delta_{r_0} \geq 100\eta r$ again, we get

$$\min_{j \in \mathcal{J}} |\tilde{\lambda}_i - \lambda_j| = \sigma_i - \sigma_{r_0+1} + (\tilde{\sigma}_i - \sigma_i) \geq 0.99(\sigma_i - \sigma_{r_0+1}).$$

It follows from (52) that

$$\|\mathcal{U}_j^T \tilde{\mathbf{u}}_i\| \leq \frac{7\eta}{\sigma_i - \sigma_{r_0+1}} \quad (53)$$

for every $i \in \llbracket 1, r_0 \rrbracket \cap \mathcal{I}_b$. Finally, for $i \in \llbracket r+1, r+r_0 \rrbracket \cap \mathcal{I}_b$, analogous arguments yield that

$$\|\mathcal{U}_j^T \tilde{\mathbf{u}}_i\| \leq \frac{7\eta}{\sigma_{i-r} - \sigma_{r_0+1}}. \quad (54)$$

The proof is now complete by inserting (47), (48), (53) and (54) into (34). \square

Combining Lemma 19 and Lemma 20 gives

$$\sin \angle(\mathcal{U}_I, \tilde{\mathcal{U}}_I) \leq 21\sqrt{2}\eta \sqrt{\sum_{i=1}^{r_0} \frac{1}{(\sigma_i - \sigma_{r_0+1})^2}} + 2 \frac{\|E\|}{\sigma_{r_0}}. \quad (55)$$

The proof of (8) now follows by (11).

Finally, the bound for $\sin \angle(U_{\llbracket k, s \rrbracket}, \tilde{U}_{\llbracket k, s \rrbracket})$ and $\sin \angle(V_{\llbracket k, s \rrbracket}, \tilde{V}_{\llbracket k, s \rrbracket})$ can be derived by an analogous procedure, and we briefly sketch the details below. First, recall from (12) that it suffices to bound $\sin \angle(\mathcal{U}_l, \tilde{\mathcal{U}}_l)$ where $l := \llbracket k, s \rrbracket \cup \llbracket r+k, r+s \rrbracket$.

Denote

$$\mathbf{J} := \llbracket 1, 2r \rrbracket \setminus l = \mathbf{J}_1 \cup \mathbf{J}_2,$$

where

$$J_1 := \llbracket 1, k-1 \rrbracket \cup \llbracket r+1, r+k-1 \rrbracket \quad \text{and} \quad J_2 := \llbracket s+1, r \rrbracket \cup \llbracket r+s+1, 2r \rrbracket.$$

Observe that

$$\begin{aligned} \sin \angle(\mathcal{U}_1, \tilde{\mathcal{U}}_1) &= \|P_1 - \tilde{P}_1\| = \|P_{J_1} \tilde{P}_1\| \\ &\leq \|P_{J_1} \tilde{P}_1\| + \|P_{J_2} \tilde{P}_1\| + \|P_{\llbracket 2r+1, N+n \rrbracket} \tilde{P}_1\|. \end{aligned}$$

Using the same method as in Lemma 19, one can bound

$$\|P_{\llbracket 2r+1, N+n \rrbracket} \tilde{P}_1\| \leq 2 \frac{\|E\|}{\sigma_s}.$$

Similarly, following the arguments in the proof of Lemma 20, one sees that

$$\|P_{J_2} \tilde{P}_1\| \leq 21\sqrt{2}\eta \sqrt{\sum_{i=k}^s \frac{1}{(\sigma_i - \sigma_{s+1})^2}}.$$

To conclude the result, we only need to bound

$$\begin{aligned} \|P_{J_1} \tilde{P}_1\| &= \|(P_{J_1} - \tilde{P}_{J_1}) \tilde{P}_1\| \\ &\leq \|P_{J_1} - \tilde{P}_{J_1}\| = \sin \angle(\mathcal{U}_{J_1}, \tilde{\mathcal{U}}_{J_1}) \leq 21\sqrt{2}\eta \sqrt{\sum_{i=1}^{k-1} \frac{1}{(\sigma_i - \sigma_k)^2}} + 2 \frac{\|E\|}{\sigma_{k-1}} \end{aligned}$$

by applying (55). Hence, we obtain

$$\begin{aligned} \sin \angle(\mathcal{U}_1, \tilde{\mathcal{U}}_1) &\leq 21\sqrt{2}\eta \sqrt{\sum_{i=1}^{k-1} \frac{1}{(\sigma_i - \sigma_k)^2}} + 21\sqrt{2}\eta \sqrt{\sum_{i=k}^s \frac{1}{(\sigma_i - \sigma_{s+1})^2}} + 2 \frac{\|E\|}{\sigma_{k-1}} + 2 \frac{\|E\|}{\sigma_s}. \end{aligned}$$

The proof is complete by (12).

5. PROOF OF THEOREM 12

This section is devoted to the proof of Theorem 12. Throughout the proof, we work on the event where $\|\mathcal{E}\| = \|E\| \leq 2(\sqrt{N} + \sqrt{n})$; recall that Lemma 15 provides with probability at least $1 - 2e^{-(\sqrt{N} + \sqrt{n})^2/2} \geq 1 - 2(N+n)^{-32(K+2)}$ that this event holds. For convenience, denote

$$M := 4(\sqrt{N} + \sqrt{n}).$$

Note that the assumptions of Theorem 12 guarantees that for any $z \in S_{\sigma_j}$ ($i_0 \leq j \leq r_0$), $|z| \geq \operatorname{Re}(z) \geq \sigma_j - 20\eta r > M$.

Our next lemma provides a way to locate the eigenvalues of a perturbed real symmetric matrix. Similar results have been applied in the random matrix theory literature to study eigenvalues for both symmetric and non-symmetric random matrices [4, 10, 11, 23, 62].

Lemma 21 (Eigenvalue location criterion). *Assume \mathcal{A} has rank $2r$ with the spectral decomposition $\mathcal{A} = \mathcal{U}\mathcal{D}\mathcal{U}^T$, where \mathcal{U} is an $(N+n) \times 2r$ matrix satisfying $\mathcal{U}^T\mathcal{U} = I_{2r}$ and \mathcal{D} is a $2r \times 2r$ diagonal matrix with non-zero $\lambda_1, \dots, \lambda_{2r}$ on the diagonal. Then the eigenvalues of $\mathcal{A} + \mathcal{E}$ outside of $[-\|\mathcal{E}\|, \|\mathcal{E}\|]$ are the zeros of the function*

$$z \mapsto \det(\mathcal{D}^{-1} - \mathcal{U}^T G(z) \mathcal{U}).$$

Moreover, the algebraic multiplicity of each eigenvalue matches the corresponding multiplicity of each zero.

Proof. The eigenvalues of $\mathcal{A} + \mathcal{E}$ are the zeros of the polynomial $\det(zI - \mathcal{A} - \mathcal{E})$. For $|z| > \|\mathcal{E}\|$,

$$\begin{aligned} \det(zI - \mathcal{A} - \mathcal{E}) &= \det(zI - \mathcal{E}) \det(I - G(z)\mathcal{A}) \\ &= \det(zI - \mathcal{E}) \det(I - G(z)\mathcal{U}\mathcal{D}\mathcal{U}^\top) \\ &= \det(zI - \mathcal{E}) \det(I - \mathcal{U}^\top G(z)\mathcal{U}\mathcal{D}) \\ &= \det(zI - \mathcal{E}) \det(\mathcal{D}) \det(\mathcal{D}^{-1} - \mathcal{U}^\top G(z)\mathcal{U}). \end{aligned}$$

by the Weinstein-Aronszajn identity (13). Since $\det(zI - \mathcal{E}) \neq 0$ for $|z| > \|\mathcal{E}\|$, the claim follows. \square

Define the functions

$$f(z) := \det(\mathcal{D}^{-1} - \mathcal{U}^\top G(z)\mathcal{U}), \quad g(z) := \det(\mathcal{D}^{-1} - \mathcal{U}^\top \Phi(z)\mathcal{U}),$$

where $\Phi(z)$ is given in (16). Observe that, by Lemma 16, $1/\phi_1(z)$, $1/\phi_2(z)$ and thus $\Phi(z)$ are well-defined for any $|z| > M$. Therefore, f and g are both complex analytic in the region $\{z \in \mathbb{C} : |z| > M\}$.

An easy computation, together with (22), yields that

$$g(z) = \prod_{l=1}^r \left(\frac{1}{\phi_1(z)\phi_2(z)} - \frac{1}{\sigma_l^2} \right)$$

and thus the zeros of $g(z)$ are the values $z \in \mathbb{C}$ which satisfy the equations $\phi_1(z)\phi_2(z) = \sigma_l^2$.

Recall from (24) and (20) that

$$\varphi(z) = \phi_1(z)\phi_2(z) = (z - \operatorname{tr} \mathcal{I}^d G(z))(z - \operatorname{tr} \mathcal{I}^u G(z)).$$

The next lemma establishes several properties of φ in the complex plane and on the real line.

Lemma 22. *The function φ satisfies the following properties.*

(i) For $z, w \in \mathbb{C}$ with $|z|, |w|, |z+w| > M$,

$$\frac{9}{16}|z^2 - w^2| \leq |\varphi(z) - \varphi(w)| \leq \frac{23}{16}|z^2 - w^2|. \quad (56)$$

(ii) (Monotone) φ is real-valued and strictly increasing on $[M, \infty)$.

(iii) (Crude bounds) $0 < \varphi(z) < z^2$ for any $z \in [M, \infty)$.

Proof. Since $\varphi(z) = z^2 - z \operatorname{tr} G(z) + \operatorname{tr} \mathcal{I}^u G(z) \operatorname{tr} \mathcal{I}^d G(z)$, we first have

$$\begin{aligned} \varphi(z) - \varphi(w) &= (z^2 - w^2) - (z \operatorname{tr} G(z) - w \operatorname{tr} G(w)) \\ &\quad + (\operatorname{tr} \mathcal{I}^u G(z) \operatorname{tr} \mathcal{I}^d G(z) - \operatorname{tr} \mathcal{I}^u G(w) \operatorname{tr} \mathcal{I}^d G(w)). \end{aligned} \quad (57)$$

To establish (56), observe that

$$\begin{aligned} z \operatorname{tr} G(z) - w \operatorname{tr} G(w) &= (z \operatorname{tr} G(z) - w \operatorname{tr} G(z)) + w(\operatorname{tr} G(z) - \operatorname{tr} G(w)) \\ &= (z - w) \operatorname{tr} G(z) + w(w - z) \operatorname{tr} G(z)G(w) \end{aligned}$$

by the resolvent identity (15). For $|z|, |w|, |z + w| > M$, using Lemma 16, we get

$$\begin{aligned}
|z \operatorname{tr} G(z) - w \operatorname{tr} G(w)| &= |z - w|(|\operatorname{tr} G(z)| + |w| |\operatorname{tr} G(z)G(w)|) \\
&\leq |z - w|((N + n)\|G(z)\| + |w|(N + n)\|G(z)G(w)\|) \\
&\leq |z - w|(N + n) \left(\frac{2}{|z|} + |w| \frac{4}{|z||w|} \right) \\
&= |z - w|(N + n) \frac{6}{|z|} \\
&\leq \frac{3}{8}|z^2 - w^2|. \tag{58}
\end{aligned}$$

Analogously, one can split

$$\begin{aligned}
&|\operatorname{tr} \mathcal{I}^u G(z) \operatorname{tr} \mathcal{I}^d G(z) - \operatorname{tr} \mathcal{I}^u G(w) \operatorname{tr} \mathcal{I}^d G(w)| \\
&\leq |\operatorname{tr} \mathcal{I}^u G(z) \operatorname{tr} \mathcal{I}^d G(z) - \operatorname{tr} \mathcal{I}^u G(w) \operatorname{tr} \mathcal{I}^d G(z)| \\
&\quad + |\operatorname{tr} \mathcal{I}^u G(w) \operatorname{tr} \mathcal{I}^d G(z) - \operatorname{tr} \mathcal{I}^u G(w) \operatorname{tr} \mathcal{I}^d G(w)|. \tag{59}
\end{aligned}$$

For the first term on the right-hand side of (59), by (15) and Lemma 16, we have

$$\begin{aligned}
|\operatorname{tr} \mathcal{I}^u G(z) \operatorname{tr} \mathcal{I}^d G(z) - \operatorname{tr} \mathcal{I}^u G(w) \operatorname{tr} \mathcal{I}^d G(z)| &= |\operatorname{tr} \mathcal{I}^d G(z)| |\operatorname{tr} \mathcal{I}^u G(z) - \operatorname{tr} \mathcal{I}^u G(w)| \\
&= |\operatorname{tr} \mathcal{I}^d G(z)| |(z - w) \operatorname{tr} \mathcal{I}^u G(z)G(w)| \\
&\leq \frac{1}{8}|z||z - w| |\operatorname{tr} \mathcal{I}^u G(z)G(w)| \\
&\leq \frac{1}{8}|z||z - w|(N + n)\|G(z)\| \|G(w)\| \\
&\leq \frac{1}{2} \frac{N + n}{|w|} |z - w| \\
&\leq \frac{1}{32}|z^2 - w^2|,
\end{aligned}$$

where in the last inequality we used $\frac{N+n}{|w|} \leq \frac{1}{16}|z + w|$ since $|w|, |z + w| > M$. The second term on the right-hand side of (59) can be estimated likewise, and we conclude that

$$|\operatorname{tr} \mathcal{I}^u G(z) \operatorname{tr} \mathcal{I}^d G(z) - \operatorname{tr} \mathcal{I}^u G(w) \operatorname{tr} \mathcal{I}^d G(w)| \leq \frac{1}{16}|z^2 - w^2|.$$

Combining the above bound and (58) with (57) yields (56).

To prove property (ii), it suffices to show that $\phi_1(z), \phi_2(z)$ are both positive and strictly increasing on $[M, \infty)$. By (20) and the bound in (95), $\phi_1(z) > 0$ for $z \geq M$. Using the expression in (93), we observe that

$$\phi_1(z) = z - \frac{1}{2} \sum_{i=1}^N \left(\frac{1}{z - \eta_i} + \frac{1}{z + \eta_i} \right) - \frac{1}{z}(n - N), \tag{60}$$

where η_1, \dots, η_N are the singular values of E . Thus, $\phi_1(z)$ is a strictly increasing function since $z \geq M \geq |\eta_i|$ for all i . A similar argument shows that $\phi_2(z)$ is also positive and strictly increasing on $[M, \infty)$. Thus, $\varphi(z) = \phi_1(z)\phi_2(z)$ is the product of two positive strictly increasing functions and so is strictly increasing. In addition, it follows from (60) that $0 < \phi_1(z) < z$ (and similarly $0 < \phi_2(z) < z$) for $z \geq M$. Thus, property (iii) follows immediately. \square

Fix an index $j \in \llbracket 1, r_0 \rrbracket$. Since $\varphi(M) < M^2$ and $\lim_{z \rightarrow \infty} \varphi(z) = \infty$, it follows from the previous lemma that there exists a unique positive real number $z_j > M$ such that $\varphi(z_j) = \sigma_j^2$. Similarly, if $\sigma_l > M$ for $\sigma_l \neq \sigma_j$, then there exists a unique positive real number z_l with $\varphi(z_l) = \sigma_l^2$ so that $z_j > z_l$ if $l > j$ and $z_j < z_l$ if $l < j$.

For the next result, we define the half space

$$H_j := \{z \in \mathbb{C} : \operatorname{Re}(z) \geq z_j - 20\eta r\}.$$

Proposition 23. *Under the assumptions of Theorem 7, for every $z \in H_j$,*

$$|z| \geq \sigma_j \geq M.$$

In particular,

$$\sigma_j < z_j < \frac{8}{7}\sigma_j. \quad (61)$$

Proof. In view of Lemma 22, it follows that $\sigma_j^2 = \varphi(z_j) < z_j^2$. Thus, as $\sigma_j \geq M + 140\eta r$ by assumption,

$$|z| \geq \operatorname{Re}(z) \geq z_j - 20\eta r \geq \sigma_j - 20\eta r \geq M + 100\eta r \geq M$$

for any $z \in H_j$.

Since $\sigma_j^2 = \varphi(z_j) = (z_j - \operatorname{tr} \mathcal{I}^d G(z_j))(z_j - \operatorname{tr} \mathcal{I}^u G(z_j))$,

$$z_j^2 - \sigma_j^2 = z_j \operatorname{tr} G(z_j) - \operatorname{tr} \mathcal{I}^u G(z_j) \operatorname{tr} \mathcal{I}^d G(z_j) \leq 2(N + n), \quad (62)$$

where we invoked Lemma 16 and the fact that $\operatorname{tr} \mathcal{I}^u G(z_j) \operatorname{tr} \mathcal{I}^d G(z_j) \geq 0$ (see (93) and (94)) in the last inequality. Hence, using that $z_j \geq M$ and $\sigma_j \geq M$, we conclude that

$$z_j - \sigma_j \leq \frac{N + n}{M} \leq \frac{1}{4}(\sqrt{N} + \sqrt{n}) \leq \frac{1}{16}\sigma_j,$$

where the last inequality follows from the assumption that $n + N \geq 32$. This proves (61) (where we use a slightly worse but simpler constant). \square

If σ_j is sufficiently large, we get a finer estimate for z_j than what is given in (61).

Proposition 24. *If $\sigma_j > \frac{1}{2}n^2$, then $|z_j - \sigma_j| \leq \frac{8}{n}$.*

Proof. Following the computation in (62), together with Lemma 16, we get

$$\begin{aligned} |z_j - \sigma_j|(z_j + \sigma_j) &= |z_j^2 - \sigma_j^2| \leq z_j |\operatorname{tr} G(z_j)| + \operatorname{tr} \mathcal{I}^u G(z_j) \operatorname{tr} \mathcal{I}^d G(z_j) \\ &\leq z_j(N + n) \|G(z_j)\| + Nn \|G(z_j)\|^2 \\ &\leq 2(N + n) + 4 \frac{Nn}{z_j^2} \\ &\leq 2(N + n) + 4 \frac{Nn}{\sigma_j^2}, \end{aligned}$$

where we used (61) in the last inequality. Since $z_j + \sigma_j \geq 2\sigma_j$, we further get

$$|z_j - \sigma_j| \leq \frac{N + n}{\sigma_j} + 2 \frac{Nn}{\sigma_j^3},$$

and the conclusion follows from $\sigma_j > \frac{1}{2}n^2$ and the supposition $N \leq n$. \square

We now complete the proof of Theorem 12. Let j be a fixed index in $\llbracket i_0, r_0 \rrbracket$. We will work in the set $H_j \cap S_{\sigma_j}$, where S_{σ_j} is specified in (18). By Corollary 2.14 in [39], it follows that

$$\frac{|f(z) - g(z)|}{|g(z)|} \leq (1 + \varepsilon(z))^{2r} - 1, \quad (63)$$

where

$$\varepsilon(z) := \left\| (\mathcal{D}^{-1} - \mathcal{U}^T \Phi(z) \mathcal{U})^{-1} \right\| \left\| \mathcal{U}^T (G(z) - \Phi(z)) \mathcal{U} \right\|.$$

The next result facilitates the comparison of the numbers of zeros of f and g inside a region and will be used repeatedly in the later arguments.

Lemma 25. *For any circle $\mathcal{C} \subset \mathbb{C}$, if $\varepsilon(z) \leq \frac{1}{4r}$ for all $z \in \mathcal{C}$, then the number of zeros of f inside \mathcal{C} is the same as the number of zeros of g inside \mathcal{C} .*

Proof. Continuing from (63), we find that

$$\frac{|f(z) - g(z)|}{|g(z)|} \leq \left(1 + \frac{1}{4r} \right)^{2r} - 1 \leq e^{1/2} - 1 < 1 \quad (64)$$

for each $z \in \mathcal{C}$. Therefore, by Rouché's theorem, we conclude that the numbers of zeros of f and g inside \mathcal{C} are the same. \square

We first bound $\varepsilon(z)$ for $z \in S_{\sigma_j}$. By Proposition 11,

$$\left\| (\mathcal{D}^{-1} - \mathcal{U}^T \Phi(z) \mathcal{U})^{-1} \right\| = \max_{1 \leq l \leq r} \frac{\sigma_l}{|\sigma_l^2 - \phi_1 \phi_2|} \mathcal{Q}^{1/2},$$

where

$$\mathcal{Q} := |\phi_1 \phi_2|^2 + \frac{1}{2} \sigma_l^2 (|\phi_1|^2 + |\phi_2|^2) + \frac{1}{2} \sigma_l [4|\phi_1 \phi_2|^2 |\phi_1 + \bar{\phi}_2|^2 + \sigma_l^2 (|\phi_1|^2 - |\phi_2|^2)^2]^{1/2}.$$

Using (27) from Lemma 16, for $z \in S_{\sigma_j}$, we get

$$\begin{aligned} \mathcal{Q} &\leq \left(\frac{9}{8} \right)^4 |z|^4 + \left(\frac{9}{8} \right)^2 \sigma_l^2 |z|^2 + \left(\frac{9}{8} \right)^2 \sigma_l |z|^2 \sqrt{\sigma_l^2 + 4 \left(\frac{9}{8} \right)^2 |z|^2} \\ &\leq \frac{9}{4} |z|^4 + \frac{27}{4} |z|^3 \sigma_l + \frac{9}{2} |z|^2 \sigma_l^2 \\ &\leq \frac{9}{4} |z|^2 \left(|z| + \frac{3}{2} \sigma_l \right)^2, \end{aligned}$$

and thus

$$\left\| (\mathcal{D}^{-1} - \mathcal{U}^T \Phi(z) \mathcal{U})^{-1} \right\| \leq \frac{3}{2} |z| \max_{1 \leq l \leq r} \frac{\sigma_l (|z| + \frac{3}{2} \sigma_l)}{|\sigma_l^2 - \phi_1(z) \phi_2(z)|}.$$

Combining this bound with the bound in Lemma 9, we obtain with probability at least $1 - 10(N+n)^{-(K+1)}$

$$\varepsilon(z) \leq \max_{1 \leq l \leq r} \frac{3}{2} \frac{\eta}{|z|} \frac{\sigma_l (|z| + \frac{3}{2} \sigma_l)}{|\sigma_l^2 - \phi_1(z) \phi_2(z)|} \quad (65)$$

for all $z \in S_{\sigma_j}$.

We now restrict ourselves to values of z contained on a circle \mathcal{C}_j in $H_j \cap S_{\sigma_j}$. Here we take \mathcal{C}_j to be the circle of radius $20\eta r$ centered at z_j (which by definition is contained in $H_j \cap S_{\sigma_j}$).

The goal is to show $\varepsilon(z)$ is small for all $z \in \mathcal{C}_j$. Continuing from (65), it suffices to show

$$\frac{3}{2} \frac{\eta}{|z|} \frac{\sigma_l(|z| + \frac{3}{2}\sigma_l)}{|\sigma_l^2 - \phi_1(z)\phi_2(z)|}$$

is small for all $1 \leq l \leq r$. We split the discussion into two cases: $l = j$ and $l \neq j$.

Case 1. When $l = j$, in view of (56), for $z \in \mathcal{C}_j$, we have

$$\begin{aligned} |\sigma_j^2 - \phi_1(z)\phi_2(z)| &= |\phi_1(z_j)\phi_2(z_j) - \phi_1(z)\phi_2(z)| \\ &\geq \frac{9}{16}|z_j^2 - z^2| = \frac{45}{4}\eta r|z_j + z| \geq \frac{585}{28}\eta r\sigma_j, \end{aligned}$$

where, in the last inequality, we used $|z + z_j| = |2z_j + z - z_j| \geq 2z_j - 20\eta r \geq \frac{13}{7}\sigma_j$ by (61) and the assumption $\sigma_j \geq 140\eta r$. Similarly, using (56) and Proposition 23,

$$|\sigma_j^2 - \phi_1(z)\phi_2(z)| \leq \frac{23}{16}|z_j^2 - z^2| \leq \frac{1955}{28}\eta r\sigma_j. \quad (66)$$

Thus, to bound $\varepsilon(z)$, we apply the bound

$$\frac{6}{7}\sigma_j \leq z_j - 20\eta r \leq |z| \leq z_j + 20\eta r \leq \frac{9}{7}\sigma_j \quad (67)$$

that follows from (61) and $\sigma_j \geq 140\eta r$ to obtain

$$\frac{3}{2} \frac{\eta}{|z|} \frac{\sigma_j(|z| + \frac{3}{2}\sigma_j)}{|\sigma_j^2 - \phi_1(z)\phi_2(z)|} \leq \frac{0.13}{r} < \frac{1}{4r}.$$

Case 2. When $l \neq j$, note that

$$\begin{aligned} |\sigma_l^2 - \phi_1(z)\phi_2(z)| &\geq |\sigma_l^2 - \sigma_j^2| - |\sigma_j^2 - \phi_1(z)\phi_2(z)| \\ &\geq |\sigma_j - \sigma_l|(\sigma_j + \sigma_l) - \frac{1955}{28}\eta r\sigma_j \\ &\geq \frac{169}{560}|\sigma_j - \sigma_l|(\sigma_j + \sigma_l) \end{aligned}$$

by (66) and the assumption that $|\sigma_j - \sigma_l| \geq 100\eta r$. Hence, using (67), we get

$$\frac{3}{2} \frac{\eta}{|z|} \frac{\sigma_l(|z| + \frac{3}{2}\sigma_l)}{|\sigma_l^2 - \phi_1(z)\phi_2(z)|} \leq \frac{3}{2} \frac{\eta}{\frac{6}{7}\sigma_j} \frac{560\sigma_l(\frac{9}{7}\sigma_j + \frac{3}{2}\sigma_l)}{169|\sigma_j - \sigma_l|(\sigma_j + \sigma_l)} \leq 9 \frac{\eta\sigma_l}{\sigma_j|\sigma_j - \sigma_l|}.$$

If $\sigma_l \leq 1.8\sigma_j$, then

$$\frac{3}{2} \frac{\eta}{|z|} \frac{\sigma_l(|z| + \frac{3}{2}\sigma_l)}{|\sigma_l^2 - \phi_1(z)\phi_2(z)|} \leq 9 \frac{1.8\eta}{100\eta r} < \frac{1}{4r}.$$

If $\sigma_l \geq 1.8\sigma_j$, then $\sigma_l - \sigma_j \geq \frac{4}{9}\sigma_l$ and

$$\frac{3}{2} \frac{\eta}{|z|} \frac{\sigma_l(|z| + \frac{3}{2}\sigma_l)}{|\sigma_l^2 - \phi_1(z)\phi_2(z)|} \leq 9 \frac{\eta}{140\eta r \frac{4}{9}} < \frac{1}{4r}.$$

Thus, we conclude that

$$\varepsilon(z) \leq \frac{1}{4r} \quad (68)$$

for all $z \in \mathcal{C}_j$. By Lemma 25, the number of zeros of f inside \mathcal{C}_j is the same as the number of zeros of g inside \mathcal{C}_j . Since g has α_j zeros inside \mathcal{C}_j , it follows that there exists exactly α_j values of $l_1(j) \leq \dots \leq l_{\alpha_j}(j)$ such that

$$|\tilde{\sigma}_{l_s(j)} - z_j| < 20\eta r \quad \text{for } 1 \leq s \leq \alpha_j. \quad (69)$$

In particular, this means that for $1 \leq s \leq \alpha_j$

$$|\varphi(\tilde{\sigma}_{l_s(j)}) - \varphi(z_j)| \leq \frac{23}{16} |\tilde{\sigma}_{l_s(j)}^2 - z_j^2| \leq \frac{115}{4} \eta r (\tilde{\sigma}_{l_s(j)} + z_j) \leq \frac{115}{4} \eta r \left(\tilde{\sigma}_{l_s(j)} + \frac{8}{7} \sigma_j \right)$$

by Lemma 22 and (61). Since $\varphi(z_j) = \sigma_j^2$, we conclude that

$$|\varphi(\tilde{\sigma}_{l_s(j)}) - \sigma_j^2| \leq \frac{115}{4} \eta r \left(\tilde{\sigma}_{l_s} + \frac{8}{7} \sigma_j \right). \quad (70)$$

It remains to show that $l_s(j) = j + s - 1$ for $1 \leq s \leq \alpha_j$ and $i_0 \leq j \leq r_0$. We will do so by proving the following claims hold with probability at least $1 - 30(N+n)^{-(K+1)}$ (see Figure 2 for an illustration):



FIGURE 2. Distinct circles \mathcal{C}_j with centers z_j on the real line and radius $20\eta r$ for $i_0 \leq j \leq r_0$.

Claim 1. Distinct circles do not intersect.

Claim 2. $\mathcal{A} + \mathcal{E}$ has exactly $i_0 - 1$ eigenvalues larger than $z_{i_0} + 20\eta r$.

Claim 3. No eigenvalues of $\mathcal{A} + \mathcal{E}$ lie between distinct circles.

For the moment, let us assume these claims are true. Note that $\tilde{\sigma}_{i_0}$ has to lie inside one of the \mathcal{C}_j 's ($i_0 \leq j \leq r_0$) because it is the largest eigenvalue of $\mathcal{A} + \mathcal{E}$ that is no larger than $z_{i_0} + 20\eta r$ (due to *Claim 2*) and thus it satisfies $\tilde{\sigma}_{i_0} > z_{r_0} - 20\eta r$ by (69). Since the number of zeros of $g(z)$ located inside \mathcal{C}_j 's for all $i_0 \leq j \leq r_0$, which is $r_0 - i_0 + 1$, is the same as that of $f(z)$ inside \mathcal{C}_j 's ($i_0 \leq j \leq r_0$), we have $\tilde{\sigma}_{i_0}, \dots, \tilde{\sigma}_{r_0}$ lie inside \mathcal{C}_j 's ($i_0 \leq j \leq r_0$). The conclusion follows by the fact that the number of zeros of $g(z)$ in each \mathcal{C}_j is the same as that of $f(z)$.

We start with the proof of the *Claim 1*. For $\sigma_l \neq \sigma_j$, by Lemma 22,

$$|z_l^2 - z_j^2| \geq \frac{16}{23} |\varphi(z_l) - \varphi(z_j)| = \frac{16}{23} |\sigma_l^2 - \sigma_j^2| \geq \frac{16}{23} 100\eta r (\sigma_l + \sigma_j).$$

Since $|z_l^2 - z_j^2| = (z_l + z_j)|z_l - z_j| \leq \frac{8}{7} (\sigma_l + \sigma_j) |z_l - z_j|$ by Proposition 23, we have

$$|z_l - z_j| \geq \frac{1400}{23} \eta r, \quad (71)$$

and thus

$$\text{dist}(\mathcal{C}_j, \mathcal{C}_l) \geq |z_l - z_j| - 40\eta r > 20\eta r.$$

Next, we prove *Claim 2*. We split the proof into two cases: $i_0 = 1$ and $i_0 > 1$.

Case 1: $i_0 = 1$. We prove that no eigenvalues of $\mathcal{A} + \mathcal{E}$ are larger than $z_1 + 20\eta r$. We now take \mathcal{C}_0 to be any circle with radius $20\eta r$ centered at a point $z_0 > z_1 + 20\eta r$ on the real line inside the region $H_1 \cap \hat{S}_{\sigma_1}$ such that $\text{dist}(z_1, \mathcal{C}_0) \geq 20\eta r$. Here

$$\hat{S}_{\sigma_1} := \{w \in \mathbb{C} : |\text{Im}(w)| \leq 20\eta r, 4(\sqrt{N} + \sqrt{n}) + 120\eta r \leq \text{Re}(w) \leq \frac{3}{2}\sigma_1 + 20\eta r\} \quad (72)$$

is a slight modification of the set S_σ in (18). Note that $\tilde{\sigma}_1 \in \hat{S}_{\sigma_1}$: the upper bound $\tilde{\sigma}_1 \leq \frac{3}{2}\sigma_1$ follows from the Weyl's inequality and the supposition $\|\mathcal{E}\| \leq \frac{1}{2}\sigma_1$; the lower bound is because it is the largest eigenvalue and $\tilde{\sigma}_1 \geq z_j - 20\eta r \geq \sigma_j - 20\eta r$ from (69). An inspection of the proof of Lemma 9 reveals that the conclusion of Lemma 9 also holds on the set \hat{S}_{σ_1} . Hence, the bound (65) also holds for $z \in \hat{S}_{\sigma_1}$. We show

$$\varepsilon(z) < \frac{1}{4r}$$

for all $z \in \mathcal{C}_0$. The proof is similar to the proof of (68) and we sketch it here. For any $z \in \mathcal{C}_0$, from $|z - z_0| = 20\eta r$, $|z - z_1| \geq 20\eta r$ and $z_0 - z_1 > 40\eta r$, we obtain $|z| \leq z_0 + 20\eta r$ and

$$|z| \geq z_0 - 20\eta r \geq z_1 + 20\eta r > \sigma_1 + 20\eta r > \sigma_1.$$

Again, by Lemma 22, we see for any $1 \leq l \leq r$,

$$\begin{aligned} |\sigma_l^2 - \phi_1(z)\phi_2(z)| &= |\varphi(z_l) - \varphi(z)| \\ &\geq \frac{9}{16}|z_l^2 - z^2| \\ &\geq \frac{9}{16}(z_l + \operatorname{Re}(z))(\operatorname{Re}(z) - z_l) \\ &\geq \frac{9}{16}(\sigma_l + z_0 - 20\eta r)(z_0 - z_l - 20\eta r) \\ &\geq \frac{9}{16}(\sigma_l + z_0 - 20\eta r)20\eta r. \end{aligned} \quad (73)$$

Plugging these estimates back into (65), we see

$$\varepsilon(z) \leq \max_{1 \leq l \leq r} \frac{3}{2} \frac{\eta \sigma_l}{\sigma_1} \frac{z_0 + 20\eta r + \frac{3}{2}\sigma_l}{\frac{9}{16}(\sigma_l + z_0 - 20\eta r)20\eta r} < \frac{1}{4r},$$

where we used the bound $z_0 + 20\eta r + \frac{3}{2}\sigma_l \leq \frac{3}{2}(\sigma_l + z_0 - 20\eta r)$ in the last inequality.

By Lemma 25, f has the same number of zeros inside \mathcal{C}_0 as g . As g has no zeros inside \mathcal{C}_0^2 , $\mathcal{A} + \mathcal{E}$ has no eigenvalues inside \mathcal{C}_0 . Since the circle \mathcal{C}_0 was arbitrarily chosen inside this region, we conclude that $\mathcal{A} + \mathcal{E}$ has no eigenvalues larger than $z_1 + 20\eta r$.

Case 2: $i_0 > 1$. We work on the event

$$\max_{l \in [1, r_0]; \sigma_l > n^2/2} |\tilde{\sigma}_l - \sigma_l| \leq \eta r. \quad (74)$$

By Lemma 13, this event holds with probability at least $1 - (N + n)^{-2r^4(K+8)}$.

Note that $\sigma_{i_0-1} > n^2$. Combining (74), Proposition 24 and $z_{i_0-1} - z_{i_0} \geq \frac{1400}{23}\eta r > 60\eta r$ from (71), we get

$$\tilde{\sigma}_{i_0-1} \geq \sigma_{i_0-1} - \eta r \geq z_{i_0-1} - \frac{8}{n} - \eta r \geq z_{i_0} + 59\eta r - \frac{8}{n} > z_{i_0} + 20\eta r.$$

Hence, $\mathcal{A} + \mathcal{E}$ has at least $i_0 - 1$ eigenvalues larger than $z_{i_0} + 20\eta r$.

We first consider $\sigma_{i_0} > \frac{1}{2}n^2$. It follows from (74) and Proposition 24 that

$$\tilde{\sigma}_{i_0} \leq \sigma_{i_0} + \eta r \leq z_{i_0} + \frac{8}{n} + \eta r \leq z_{i_0} + 20\eta r.$$

This shows that $\mathcal{A} + \mathcal{E}$ has exactly $i_0 - 1$ eigenvalues larger than $z_{i_0} + 20\eta r$.

²This follows from Lemma 22 and the fact that $\operatorname{Im}(\varphi(z)) \neq 0$ whenever $\operatorname{Im} z \neq 0$ for all $|z| > M$.

Now consider $\sigma_{i_0} \leq \frac{1}{2}n^2$. By Weyl's inequality, $\tilde{\sigma}_{i_0} \leq \sigma_{i_0} + \|E\| < \frac{3}{2}\sigma_{i_0}$. If $\frac{3}{2}\sigma_{i_0} \leq z_{i_0} + 20\eta r$, the proof is already done. Now we assume $\frac{3}{2}\sigma_{i_0} > z_{i_0} + 20\eta r$. If $\frac{3}{2}\sigma_{i_0} - (z_{i_0} + 20\eta r) < 40\eta r$, following (61), we have $\frac{8}{7}\sigma_{i_0} > z_{i_0} > \frac{3}{2}\sigma_{i_0} - 60\eta r$ and thus $\sigma_{i_0} < 168\eta r$. From the assumption $\sigma_{i_0} \geq 4(\sqrt{N} + \sqrt{n}) + 140\eta r$, we further have $4(\sqrt{N} + \sqrt{n}) \leq 28\eta r$ and hence $\|E\| \leq 2(\sqrt{N} + \sqrt{n}) < 14\eta r$. It follows from Weyl's inequality and (61) that $\tilde{\sigma}_{i_0} \leq \sigma_{i_0} + \|E\| \leq z_{i_0} + 14\eta r < z_{i_0} + 20\eta r$.

It remains to consider the case when $\frac{3}{2}\sigma_{i_0} - (z_{i_0} + 20\eta r) \geq 40\eta r$. To prove $\tilde{\sigma}_{i_0} \leq z_{i_0} + 20\eta r$, we show that f has no zeros on the interval $(z_{i_0} + 20\eta r, \frac{3}{2}\sigma_{i_0})$. The proof is similar to the proof of *Case 1* when $i_0 = 1$. We only mention the differences. Define $\hat{S}_{\sigma_{i_0}}$ as in (72) and the bound (65) also holds for $z \in \hat{S}_{\sigma_{i_0}}$. The goal is to show $\varepsilon(z) < 1/4r$ for all $z \in \mathcal{C}_0$, where \mathcal{C}_0 is any circle with radius $20\eta r$ centered at a point $z_0 \in (z_{i_0} + 20\eta r, \frac{3}{2}\sigma_{i_0})$ inside the region $H_{i_0} \cap \hat{S}_{\sigma_{i_0}}$ such that $\text{dist}(z_0, z_{i_0} + 20\eta r) \geq 20\eta r$ and $\text{dist}(z_0, \frac{3}{2}\sigma_{i_0}) \geq 20\eta r$. If so, by Lemma 25, f has the same number of zeros inside \mathcal{C}_0 as g . Note that g has no zeros inside \mathcal{C}_0 since $\text{Im}(\varphi(z)) \neq 0$ whenever $\text{Im} z \neq 0$ for all $|z| > M$ and $z_{i_0-1} \geq \sigma_{i_0-1} - \frac{8}{n} > n^2 - \frac{8}{n} > \frac{3}{2}\sigma_{i_0}$ by Proposition 24. Since \mathcal{C}_0 was arbitrarily chosen, $\mathcal{A} + \mathcal{E}$ has no eigenvalues on $(z_{i_0} + 20\eta r, \frac{3}{2}\sigma_{i_0})$.

It remains to bound $\varepsilon(z)$ from (65). The same arguments as those in *Case 1* yield that

$$\max_{i_0 \leq l \leq r} \frac{3}{2} \frac{\eta}{|z|} \frac{\sigma_l(|z| + \frac{3}{2}\sigma_l)}{|\sigma_l^2 - \phi_1(z)\phi_2(z)|} < \frac{1}{4r}$$

for any $z \in \mathcal{C}_0$. We only need to control

$$\max_{1 \leq l \leq i_0-1} \frac{3}{2} \frac{\eta}{|z|} \frac{\sigma_l(|z| + \frac{3}{2}\sigma_l)}{|\sigma_l^2 - \phi_1(z)\phi_2(z)|}.$$

For any $z \in \mathcal{C}_0$, $|z| \geq z_0 - 20\eta r \geq z_{i_0} \geq \sigma_{i_0}$ and $|z| \leq z_0 + 20\eta r \leq \frac{3}{2}\sigma_{i_0} + 40\eta r$. For any $1 \leq l \leq i_0 - 1$, using similar computation from (73), we get

$$|\sigma_l^2 - \phi_1(z)\phi_2(z)| \geq \frac{9}{16}(\sigma_l + z_0 - 20\eta r)(z_l - z_0 - 20\eta r).$$

Note that $z_0 \geq z_{i_0} + 40\eta r \geq \sigma_{i_0} + 40\eta r$. Hence, $\sigma_l + z_0 - 20\eta r \geq \sigma_l + \sigma_{i_0} + 20\eta r$. From $\sigma_l \geq n^2$, we see $\sigma_{i_0} \leq \frac{1}{2}n^2 \leq \frac{1}{2}\sigma_l$. This, together with (61) and $z_0 \leq \frac{3}{2}\sigma_{i_0} - 20\eta r$, implies that

$$z_l - z_0 - 20\eta r \geq \sigma_l - \frac{3}{2}\sigma_{i_0} \geq \sigma_l - \frac{3}{4}\sigma_l = \frac{1}{4}\sigma_l.$$

Hence,

$$\begin{aligned} \max_{1 \leq l \leq i_0-1} \frac{3}{2} \frac{\eta}{|z|} \frac{\sigma_l(|z| + \frac{3}{2}\sigma_l)}{|\sigma_l^2 - \phi_1(z)\phi_2(z)|} &\leq \frac{3}{2} \frac{\eta}{\sigma_{i_0}} \max_{1 \leq l \leq i_0-1} \frac{\sigma_l(\frac{3}{2}\sigma_{i_0} + 40\eta r + \frac{3}{2}\sigma_l)}{\frac{9}{16}(\sigma_l + \sigma_{i_0} + 20\eta r)\frac{1}{4}\sigma_l} \\ &\leq \frac{16\eta}{\sigma_{i_0}} \\ &< \frac{1}{4r} \end{aligned}$$

using the assumption $\sigma_{i_0} > 140\eta r$. Therefore, $\varepsilon(z) < 1/4r$ for all $z \in \mathcal{C}_0$.

The proof of *Claim 3* is similar to the previous argument. Let $\mathcal{C}_{j_1}, \mathcal{C}_{j_2}$ be two distinct adjacent circles. Note that from the proof of *Claim 1*, $\text{dist}(\mathcal{C}_{j_1}, \mathcal{C}_{j_2}) > 20\eta r$. We show that $\mathcal{A} + \mathcal{E}$ has no eigenvalues lying on the real line between \mathcal{C}_{j_1} and \mathcal{C}_{j_2} . Take any point x on the real line between the two circles so that \mathcal{C}_x , the circle

centered at x with radius $10\eta r$, is inside the region $H_{j_1} \cap S_{\sigma_{j_1}}$ or $H_{j_2} \cap S_{\sigma_{j_2}}$, where $\text{dist}(x, \mathcal{C}_{j_1}) > 10\eta r$ and $\text{dist}(x, \mathcal{C}_{j_2}) > 10\eta r$. Then using similar calculations as in the proof of *Claim 2*, it suffices to show that $\varepsilon(z) < \frac{1}{4r}$. The remaining arguments are similar to those in the proof of *Claim 2*; we omit the details.

6. PROOF OF LEMMA 9

In this section, we present the proof of Lemma 9. We first show that $G(z)$ is close to $\Phi(z)$ for any fixed $z \in \mathbb{C}$ satisfying $|z| \geq 4(\sqrt{N} + \sqrt{n})$. Then we extend this result to any $z \in S_{\sigma_j}$ by a net argument. Throughout the proof, we will sometimes write G instead of $G(z)$ for convenience. The proof presented here takes advantage of the fact that the entries of E are jointly independent standard Gaussian random variables; this assumption greatly simplifies the forthcoming calculations, although the method can be extended to other distributions. In particular, when the entries are non-Gaussian, one also needs to estimate the off-diagonal entries of G . This can be accomplished by modifying some of the techniques from [15].

We begin with the following notation.

Definition 26 (Minors). For $I \subset \llbracket 1, N+n \rrbracket$, we define $\mathcal{E}^{(I)}$ by

$$\mathcal{E}_{st}^{(I)} := \begin{cases} \mathcal{E}_{st} & \text{if } s, t \notin I, \\ 0 & \text{otherwise.} \end{cases}$$

We define the resolvent of $\mathcal{E}^{(I)}$ by

$$G_{st}^{(I)}(z) := \begin{cases} (z - \mathcal{E}^{(I)})_{st}^{-1} & \text{if } s, t \notin I, \\ 0 & \text{otherwise,} \end{cases}$$

whenever the inverse is defined. We use the summation notation

$$\sum_s^{(I)} := \sum_{s \in \llbracket 1, N+n \rrbracket : s \notin I}.$$

When $I = \{a\}$, we abbreviate $(\{a\})$ by (a) in the above definitions.

Lemma 27 (Resolvent identities). *For any $k \in \llbracket 1, N+n \rrbracket$ and for $|z| > \|\mathcal{E}\|$*

$$G_{kk}(z) = \frac{1}{z - \mathcal{E}_{kk} - \sum_{s,t}^{(k)} \mathcal{E}_{sk} G_{st}^{(k)}(z) \mathcal{E}_{tk}}.$$

Moreover, for $i \neq j$ and any $|z| > \|\mathcal{E}\|$,

$$G_{ij}(z) = -G_{ii}(z) \sum_k^{(i)} \mathcal{E}_{ik} G_{kj}^{(i)}(z).$$

Proof. The formula for the diagonal entries follows from the Schur complement (see [3, Theorem A.4]). The off-diagonal entries can be computed in a similar way (see [27, Lemma 4.2] or [29, Lemma 6.10]). \square

Next, we show that the resolvent matrix $G(z) = (z - \mathcal{E})^{-1}$ is well approximated by the diagonal matrix $\Phi(z)$ for any fixed $z \in \mathbb{C}$ with sufficiently large modulus.

Lemma 28. *Let $K > 0$ be any constant and assume $(\sqrt{N} + \sqrt{n})^2 \geq 64(K + 2) \log(N+n)$. For any $z \in \mathbb{C}$ with $|z| \geq 4(\sqrt{N} + \sqrt{n})$,*

$$\|\mathcal{U}^T (G(z) - \Phi(z)) \mathcal{U}\| \leq 48r \frac{\sqrt{(K+1) \log(N+n)}}{|z|^2}$$

with probability at least $1 - (N + n)^{-K}$.

Proof. By the rotational invariance of E , it suffices to assume that U is the matrix with columns e_1, \dots, e_r , where e_1, \dots, e_N is the canonical basis in \mathbb{R}^N , and the columns of V are given by f_1, \dots, f_r , where f_1, \dots, f_n is the canonical basis in \mathbb{R}^n . We use the shorthand notations $\bar{i} = i - r$ and $\bar{j} = j - r$ if $r + 1 \leq i, j \leq 2r$. Thus,

$$2(\mathcal{U}^T G \mathcal{U})_{ij} = \begin{cases} G_{ij} + G_{i,N+j} + G_{N+i,j} + G_{N+i,N+j} & \text{if } i, j \in \llbracket 1, r \rrbracket; \\ G_{i\bar{j}} - G_{i,N+\bar{j}} + G_{N+i,\bar{j}} - G_{N+i,N+\bar{j}} & \text{if } i \in \llbracket 1, r \rrbracket, j \in \llbracket r+1, 2r \rrbracket; \\ G_{\bar{i}j} + G_{\bar{i},N+j} - G_{N+\bar{i},j} - G_{N+\bar{i},N+j} & \text{if } i \in \llbracket r+1, 2r \rrbracket, j \in \llbracket 1, r \rrbracket; \\ G_{\bar{i}\bar{j}} - G_{\bar{i},N+\bar{j}} - G_{N+\bar{i},\bar{j}} + G_{N+\bar{i},N+\bar{j}} & \text{if } i, j \in \llbracket r+1, 2r \rrbracket. \end{cases}$$

Fix $z \in \mathbb{C}$ with $|z| \geq 4(\sqrt{N} + \sqrt{n})$. Denote the set

$$\mathcal{S}_r := \{(i, j) : i, j \in \llbracket 1, r \rrbracket \cup \llbracket N+1, N+r \rrbracket\}.$$

Since $\|\mathcal{M}\| \leq 2r \|\mathcal{M}\|_{\max} := 2r \max_{1 \leq i, j \leq 2r} |\mathcal{M}_{ij}|$ for any $2r \times 2r$ matrix \mathcal{M} , we first get

$$\|\mathcal{U}^T (G(z) - \Phi(z)) \mathcal{U}\| \leq 2r \max_{1 \leq i, j \leq 2r} |(\mathcal{U}^T G(z) \mathcal{U})_{ij} - (\mathcal{U}^T \Phi(z) \mathcal{U})_{ij}|. \quad (75)$$

In order to prove Lemma 28, we claim that it suffices to show that

$$\max_{(i,j) \in \mathcal{S}_r} |G_{ij}(z) - \Phi_{ij}(z)| \leq 12 \frac{\sqrt{(K+1) \log(N+n)}}{|z|^2}. \quad (76)$$

with probability at least $1 - 10(N+n)^{-K}$.

Let us assume for the moment that (76) holds. For any $(i, j) \in \llbracket 1, r \rrbracket$, in view of (22) and the definition of $\Phi(z)$ in (16), we find that

$$\begin{aligned} & 2r |(\mathcal{U}^T G(z) \mathcal{U})_{ij} - (\mathcal{U}^T \Phi(z) \mathcal{U})_{ij}| \\ &= r \left| G_{ij} + G_{i,N+j} + G_{N+i,j} + G_{N+i,N+j} - \frac{1}{\phi_1(z)} \delta_{ij} - \frac{1}{\phi_2(z)} \delta_{ij} \right| \\ &\leq r |G_{ij}(z) - \Phi_{ij}(z)| + r |G_{i,N+j}(z) - \Phi_{i,N+j}(z)| \\ &\quad + r |G_{N+i,j}(z) - \Phi_{N+i,j}(z)| + r |G_{N+i,N+j}(z) - \Phi_{N+i,N+j}(z)| \\ &\leq 48r \frac{\sqrt{(K+1) \log(N+n)}}{|z|^2}. \end{aligned}$$

Similar discussion applies to every $(i, j) \in \llbracket 1, 2r \rrbracket$ and the details are omitted. Hence,

$$2r \max_{1 \leq i, j \leq 2r} |(\mathcal{U}^T G(z) \mathcal{U})_{ij} - (\mathcal{U}^T \Phi(z) \mathcal{U})_{ij}| \leq 48r \frac{\sqrt{(K+1) \log(N+n)}}{|z|^2}$$

and the conclusion of Lemma 28 follows from (75).

Now we turn to the proof of (76). For the remainder of the proof, we work on the event where $\|\mathcal{E}\| = \|E\| \leq 2(\sqrt{N} + \sqrt{n})$; recall that Lemma 15 shows this event holds with probability at least $1 - 2e^{-(\sqrt{N} + \sqrt{n})^2/2} \geq 1 - 2(N+n)^{-32(K+2)}$ since $\frac{(\sqrt{N} + \sqrt{n})^2}{\log(N+n)} > 64(K+2)$. We start by controlling the diagonal entries of $G(z)$. By Lemma 27, for $k \in \llbracket 1, N+n \rrbracket$,

$$G_{kk}(z) = \frac{1}{z - \mathcal{E}_{kk} - \sum_{s,t} \binom{k}{s,t} \mathcal{E}_{sk} G_{st}^{(k)}(z) \mathcal{E}_{tk}},$$

and thus, for $1 \leq k \leq r$, by the block definition of \mathcal{E} and the expression of $\phi_1(z)$ in (17) and (20), we have

$$\begin{aligned} \left| G_{kk}(z) - \frac{1}{\phi_1(z)} \right| &= \left| \frac{1}{z - \sum_{1 \leq i, j \leq n}^{(k)} E_{ki} G_{N+i, N+j}^{(k)}(z) E_{kj}} - \frac{1}{\phi_1(z)} \right| \\ &\leq \frac{|\sum_{1 \leq i, j \leq n}^{(k)} E_{ki} G_{N+i, N+j}^{(k)}(z) E_{kj} - \sum_{t \in \llbracket N+1, N+n \rrbracket} G_{tt}(z)|}{|z - \sum_{1 \leq i, j \leq n}^{(k)} E_{ki} G_{N+i, N+j}^{(k)}(z) E_{kj}| |z - \text{tr } \mathcal{I}^d G(z)|}. \end{aligned} \quad (77)$$

We now turn to bounding the right-hand side of (77). We start with obtaining an upper bound for the numerator of this term. By the triangle inequality,

$$\mathbb{P} \left(\left| \sum_{1 \leq i, j \leq n}^{(k)} E_{ki} G_{N+i, N+j}^{(k)}(z) E_{kj} - \sum_{t \in \llbracket N+1, N+n \rrbracket} G_{tt}^{(k)}(z) \right| \geq t \right) \quad (78)$$

$$\leq \mathbb{P} \left(\left| \sum_{1 \leq i, j \leq n}^{(k)} E_{ki} \text{Re } G_{N+i, N+j}^{(k)}(z) E_{kj} - \sum_{t \in \llbracket N+1, N+n \rrbracket} \text{Re } G_{tt}^{(k)}(z) \right| \geq t/2 \right) \quad (79)$$

$$+ \mathbb{P} \left(\left| \sum_{1 \leq i, j \leq n}^{(k)} E_{ki} \text{Im } G_{N+i, N+j}^{(k)}(z) E_{kj} - \sum_{t \in \llbracket N+1, N+n \rrbracket} \text{Im } G_{tt}^{(k)}(z) \right| \geq t/2 \right).$$

Since the k -th row of E is independent of $G^{(k)}$, we condition on $G^{(k)}$ in the following estimates. We start with the term in (79). For notational convenience, denote $\mathbf{X} := \left(\text{Re } G_{N+i, N+j}^{(k)}(z) \right)_{i, j \neq k}$ and \mathbf{g}^T the k th row of E with the k th entry removed.

By assumption, \mathbf{g} is a standard Gaussian vector in \mathbb{R}^{n-1} . Rewrite

$$\sum_{1 \leq i, j \leq n}^{(k)} E_{ki} \text{Re } G_{N+i, N+j}^{(k)}(z) E_{kj} = \mathbf{g}^T \mathbf{X} \mathbf{g}$$

and $\sum_{t \in \llbracket N+1, N+n \rrbracket} \text{Re } G_{tt}^{(k)}(z) = \mathbb{E} \mathbf{g}^T \mathbf{X} \mathbf{g}$. Assume the singular value decomposition of \mathbf{X} is given by $\mathbf{X} = O_1 \Sigma O_2$ where $\Sigma = \text{diag}(s_1, \dots, s_{n-1})$ and O_1, O_2 are orthogonal matrices. Due to the rotation invariance property of Gaussian vectors, $\mathbf{g}^T \mathbf{X} \mathbf{g} \sim \mathbf{g}^T \Sigma \mathbf{g}$ and $\mathbb{E} \mathbf{g}^T \mathbf{X} \mathbf{g} = \mathbb{E} \mathbf{g}^T \Sigma \mathbf{g} = \sum_{i=1}^{n-1} s_i$. To bound (79), it is equivalent to bound $\mathbb{P}(|\mathbf{g}^T \Sigma \mathbf{g} - \mathbb{E} \mathbf{g}^T \Sigma \mathbf{g}| \geq t/2)$. In order to apply Lemma 14, we can verify that $\mathbf{g}^T \Sigma \mathbf{g} = \sum_{i=1}^{n-1} s_i g_i^2$ is sub-exponential with parameters $(4\|\mathbf{X}\|_F^2, 4\|\mathbf{X}\|)$. Indeed, by independence and (26),

$$\mathbb{E} e^{\lambda(\mathbf{g}^T \Sigma \mathbf{g} - \mathbb{E} \mathbf{g}^T \Sigma \mathbf{g})} = \mathbb{E} e^{\sum_{i=1}^{n-1} \lambda(s_i - 1) g_i^2} = \prod_{i=1}^{n-1} \mathbb{E} e^{\lambda s_i (g_i^2 - 1)} \leq \prod_{i=1}^{n-1} e^{\frac{4s_i^2 \lambda^2}{2}} = e^{\frac{4\|\mathbf{X}\|_F^2 \lambda^2}{2}}$$

for all $|\lambda| < \frac{1}{4 \max_i s_i} = \frac{1}{4\|\mathbf{X}\|}$. It follows from Lemma 14 that

$$\left| \sum_{1 \leq i, j \leq n}^{(k)} E_{ki} \text{Re } G_{N+i, N+j}^{(k)}(z) E_{kj} - \sum_{t \in \llbracket N+1, N+n \rrbracket} \text{Re } G_{tt}^{(k)}(z) \right| \leq t \quad (80)$$

with probability at least $1 - 2 \exp(-\frac{1}{32} \min\{t^2|z|^2/n, 2t|z|\})$ by combining the following estimates

$$\|\mathbf{X}\| \leq \left\| \left(G_{N+i, N+j}^{(k)}(z) \right)_{1 \leq i, j \leq n} \right\| \leq \|G^{(k)}(z)\| \leq \frac{2}{|z|}$$

and

$$\|\mathbf{X}\|_F^2 \leq \left\| \left(G_{N+i, N+j}^{(k)}(z) \right)_{1 \leq i, j \leq n} \right\|_F^2 \leq n \left\| \left(G_{N+i, N+j}^{(k)}(z) \right)_{1 \leq i, j \leq n} \right\|^2 \leq \frac{4n}{|z|^2}$$

due to Lemma 16. Since $|z| \geq 4(\sqrt{N} + \sqrt{n})$, by selecting $t = \sqrt{2(K+1) \log(N+n)}$ in (80), we find that

$$\left| \sum_{1 \leq i, j \leq n}^{(k)} E_{ki} \operatorname{Re} G_{N+i, N+j}^{(k)}(z) E_{kj} - \sum_{t \in \llbracket N+1, N+n \rrbracket} \operatorname{Re} G_{tt}^{(k)}(z) \right| \leq \sqrt{2(K+1) \log(N+n)}$$

with probability at least $1 - 2(N+n)^{-(K+1)}$. To be more specific, to get this probability bound, we need the following discussion. When $t|z| \leq 2n$ and thus $\min\{t^2|z|^2/n, 2t|z|\} = t^2|z|^2/n$, the probability bound in (80) is at least $1 - 2(N+n)^{-(K+1)}$ since $t|z| \geq 4\sqrt{2(K+1) \log(N+n)}(\sqrt{N} + \sqrt{n})$. When $t|z| > 4n$ and $\min\{t^2|z|^2/n, 2t|z|\} = 2t|z| > 2n$, we obviously still have the probability bound in (80) is at least $1 - 2 \exp(-n/8) > 1 - 2(N+n)^{-2(K+2)}$ by the suppositions $(\sqrt{N} + \sqrt{n})^2 > 64(K+2) \log(N+n)$ and $n \geq N$.

Likewise, one also has

$$\left| \sum_{1 \leq i, j \leq n}^{(k)} E_{ki} \operatorname{Im} G_{N+i, N+j}^{(k)}(z) E_{kj} - \sum_{t \in \llbracket N+1, N+n \rrbracket} \operatorname{Im} G_{tt}^{(k)}(z) \right| \leq \sqrt{2(K+1) \log(N+n)}$$

with probability at least $1 - 2(N+n)^{-(K+1)}$. Inserting the above estimates back into (78), we find that

$$\left| \sum_{1 \leq i, j \leq n}^{(k)} E_{ki} G_{N+i, N+j}^{(k)}(z) E_{kj} - \sum_{t \in \llbracket N+1, N+n \rrbracket} G_{tt}^{(k)}(z) \right| \leq 2\sqrt{2(K+1) \log(N+n)} \quad (81)$$

with probability at least $1 - 4(N+n)^{-(K+1)}$.

Next, we show that the difference between $\sum_{t \in \llbracket N+1, N+n \rrbracket} G_{tt}^{(k)}(z)$ and $\sum_{t \in \llbracket N+1, N+n \rrbracket} G_{tt}(z)$ is quite small. Thus, combining (81), we get an upper bound for the term in the numerator of the right-hand side of (77). Rewrite

$$\left| \sum_{t \in \llbracket N+1, N+n \rrbracket} G_{tt}^{(k)}(z) - \sum_{t \in \llbracket N+1, N+n \rrbracket} G_{tt}(z) \right| = \left| \operatorname{tr} \mathcal{I}^d(G^{(k)}(z) - G(z)) \right|.$$

By the resolvent identity (15), the above term is written as

$$\left| \operatorname{tr} \mathcal{I}^d(G^{(k)}(z) - G(z)) \right| = \left| \operatorname{tr} \mathcal{I}^d G(z) (\mathcal{E} - \mathcal{E}^{(k)}) G^{(k)}(z) \right|.$$

Since $\mathcal{E} - \mathcal{E}^{(k)}$ has at most rank 2, we conclude that

$$\begin{aligned} \left| \operatorname{tr} \mathcal{I}^d G(z) (\mathcal{E} - \mathcal{E}^{(k)}) G^{(k)}(z) \right| &\leq 2 \|G(z) (\mathcal{E} - \mathcal{E}^{(k)}) G^{(k)}(z)\| \\ &\leq 2 \|G(z)\| (\|\mathcal{E}\| + \|\mathcal{E}^{(k)}\|) \|G^{(k)}(z)\| \\ &\leq \frac{16 \|E\|}{|z|^2} \\ &\leq \frac{8}{|z|} \end{aligned}$$

since $|z| \geq 4(\sqrt{N} + \sqrt{n})$ and $\|E\| \leq 2(\sqrt{N} + \sqrt{n})$. Thus, we have shown that

$$\left| \sum_{t \in \llbracket N+1, N+n \rrbracket} G_{tt}^{(k)}(z) - \sum_{t \in \llbracket N+1, N+n \rrbracket} G_{tt}(z) \right| \leq \frac{8}{|z|} \leq \frac{2}{\sqrt{N} + \sqrt{n}}.$$

Returning to (81), we see that

$$\left| \sum_{1 \leq i, j \leq n}^{(k)} E_{ki} G_{N+i, N+j}^{(k)}(z) E_{kj} - \sum_{t \in \llbracket N+1, N+n \rrbracket} G_{tt}(z) \right| \leq 4\sqrt{(K+1) \log(N+n)} \quad (82)$$

with probability at least $1 - 4(N+n)^{-(K+1)}$.

To finish the estimate of (77), we provide a lower bound for the denominator of the right-hand side of (77). Note from (95) that

$$\left| \sum_{t \in \llbracket N+1, N+n \rrbracket} G_{tt}(z) \right| = |\operatorname{tr} \mathcal{I}^d G(z)| \leq \frac{|z|}{8}.$$

Combined with (82) and the fact that

$$4\sqrt{(K+2) \log(N+n)} \leq \frac{1}{2}(\sqrt{N} + \sqrt{n}) \leq \frac{|z|}{8}$$

since $(\sqrt{N} + \sqrt{n})^2 \geq 64(K+2) \log(N+n)$ by supposition and $|z| \geq 4(\sqrt{N} + \sqrt{n})$, we arrive at

$$\left| \sum_{1 \leq i, j \leq n}^{(k)} E_{ki} G_{N+i, N+j}^{(k)}(z) E_{kj} \right| \leq \frac{|z|}{4}.$$

Hence, by triangle inequality,

$$\left| z - \sum_{1 \leq i, j \leq n}^{(k)} E_{ki} G_{N+i, N+j}^{(k)}(z) E_{kj} \right| \geq \frac{21}{32} |z|^2.$$

Together with (82), inserting the above estimates into (77) yields

$$\left| G_{kk}(z) - \frac{1}{\phi_1(z)} \right| \leq \frac{128}{21} \frac{\sqrt{(K+1) \log(N+n)}}{|z|^2}$$

with probability at least $1 - 4(N+n)^{-(K+1)}$ for $1 \leq k \leq r$. Analogously, one also has

$$\left| G_{kk}(z) - \frac{1}{\phi_2(z)} \right| \leq \frac{128}{21} \frac{\sqrt{(K+1) \log(N+n)}}{|z|^2}$$

with probability at least $1 - 4(N+n)^{-(K+1)}$ for $N+1 \leq k \leq N+r$. Union bounding over all $k \in \llbracket 1, r \rrbracket \cup \llbracket N+1, N+r \rrbracket$ completes the proof for the diagonal entries.

For the off-diagonal entries, by Lemma 27, we have

$$|G_{ij}(z)| \leq \frac{2}{|z|} \left| \sum_k^{(i)} \mathcal{E}_{ik} G_{kj}^{(i)}(z) \right|$$

on the event where $\|E\| \leq 2(\sqrt{N} + \sqrt{n})$. Since the i -th row of \mathcal{E} is independent of $G^{(i)}$, conditioning on $G^{(i)}$, $\sum_k^{(i)} \mathcal{E}_{ik} G_{kj}^{(i)}(z)$ still has a Gaussian distribution. In particular, the real part $\sum_k^{(i)} \mathcal{E}_{ik} \operatorname{Re} G_{kj}^{(i)}(z)$ has a Gaussian distribution with mean 0 and variance $\sum_k^{(i)} \left(\operatorname{Re} G_{kj}^{(i)} \right)^2 \leq \sum_k^{(i)} \overline{G_{kj}^{(i)}} G_{kj}^{(i)} = ((G^{(i)})^* G^{(i)})_{jj}$. Likewise, the imaginary part $\sum_k^{(i)} \mathcal{E}_{ik} \operatorname{Im} G_{kj}^{(i)}(z)$ also has a Gaussian distribution with mean 0 and variance at most $((G^{(i)})^* G^{(i)})_{jj}$. Using the tail bounds for the Gaussian distribution [64, Proposition 2.1.2], we get

$$\begin{aligned} \left| \sum_k^{(i)} \mathcal{E}_{ik} G_{kj}^{(i)}(z) \right| &\leq \left| \sum_k^{(i)} \mathcal{E}_{ik} \operatorname{Re} G_{kj}^{(i)}(z) \right| + \left| \sum_k^{(i)} \mathcal{E}_{ik} \operatorname{Im} G_{kj}^{(i)}(z) \right| \\ &\leq 2\sqrt{2(K+1)\log(N+n)} \sqrt{((G^{(i)})^* G^{(i)})_{jj}} \end{aligned}$$

with probability at least $1 - 0.5(N+n)^{-(K+1)}$. By bounding $((G^{(i)})^* G^{(i)})_{jj} \leq \|G^{(i)}\|^2 \leq (2/|z|)^2$, we conclude that

$$|G_{ij}(z)| \leq \frac{8\sqrt{2(K+1)\log(N+n)}}{|z|^2}$$

with probability at least $1 - 0.5(N+n)^{-(K+1)}$. Union bounding over $i, j \in \llbracket 1, r \rrbracket \cup \llbracket N+1, N+r \rrbracket$ completes the proof. \square

We conclude this section with the proof of Lemma 9.

Proof of Lemma 9. Fix an index $j \in \llbracket i_0, r_0 \rrbracket$. By Lemma 28, for any $z \in \mathbb{C}$ with $|z| \geq 4(\sqrt{N} + \sqrt{n})$,

$$\|\mathcal{U}^T (G(z) - \Phi(z)) \mathcal{U}\| \leq 48r \frac{\sqrt{(K+8)\log(N+n)}}{|z|^2}$$

with probability at least $1 - (N+n)^{-(K+7)}$. Note that every $z \in S_{\sigma_j}$ satisfies $|z| \geq 4(\sqrt{N} + \sqrt{n})$ by assumption. Let \mathcal{N} be a 1-net of S_{σ_j} . Since $\sigma_j \leq n^2$, a simple volume argument (see for instance [52, Lemma 3.3]) shows \mathcal{N} can be chosen so that $|\mathcal{N}| \leq 20[(20\eta r)^2 + (8n^2/7 + 20\eta r)^2]$. By the union bound, the supposition $(\sqrt{N} + \sqrt{n})^2 \geq 64(K+9)\log(N+n)$ and Lemma 28,

$$\max_{z \in \mathcal{N}} |z|^2 \|\mathcal{U}^T (G(z) - \Phi(z)) \mathcal{U}\| \leq 48r \sqrt{(K+8)\log(N+n)} \quad (83)$$

with probability at least $1 - 10(N+n)^{-(K+1)}$. We now wish to extend this bound to all $z \in S_{\sigma_j}$.

Define the functions

$$f(z) := z^2 \mathcal{U}^T G(z) \mathcal{U}, \quad g(z) := z^2 \mathcal{U}^T \Phi(z) \mathcal{U}.$$

In order to complete the proof, it suffices to show that f and g are 8-Lipschitz in S_{σ_j} . In other words, we want to show that $\|f(z) - f(w)\| \leq 8|z - w|$ and $\|g(z) - g(w)\| \leq 8|z - w|$ for all $z, w \in S_{\sigma_j}$. Indeed, in view of (83), if $z \in S_{\sigma_j}$, then there exists $w \in \mathcal{N}$ so that $|z - w| \leq 1$, and hence

$$\begin{aligned} |z|^2 \|\mathcal{U}^T G(z)\mathcal{U} - \mathcal{U}^T \Phi(z)\mathcal{U}\| &\leq \|f(z) - f(w)\| + \|f(w) - g(w)\| + \|g(w) - g(z)\| \\ &\leq 16 + |w|^2 \|\mathcal{U}^T G(w)\mathcal{U} - \mathcal{U}^T \Phi(w)\mathcal{U}\| \\ &\leq 16 + 48r\sqrt{(K+8)\log(N+n)} \\ &\leq 54r\sqrt{(K+8)\log(N+n)}, \end{aligned}$$

where we used the Lipschitz continuity of f and g in the second inequality.

It remains to show that f and g are 8-Lipschitz in S_{σ_j} . To do so we will only work on the event where $\|E\| \leq 2(\sqrt{N} + \sqrt{n})$; the probability of this event is at least $1 - 2e^{-(\sqrt{N} + \sqrt{n})^2/2} \geq 1 - 2(N+n)^{-32(K+9)}$ by Lemma 15 and the supposition $(\sqrt{N} + \sqrt{n})^2 \geq 64\log(N+n)(K+9)$. Let $z, w \in S_{\sigma_j}$, and assume without loss of generality that $|z| \geq |w| \geq 4(\sqrt{N} + \sqrt{n})$. Then

$$\begin{aligned} \|f(z) - f(w)\| &\leq \|z^2\mathcal{U}^T G(z)\mathcal{U} - zw\mathcal{U}^T G(z)\mathcal{U}\| + \|zw\mathcal{U}^T G(z)\mathcal{U} - w^2\mathcal{U}^T G(z)\mathcal{U}\| \\ &\quad + \|w^2\mathcal{U}^T G(z)\mathcal{U} - w^2\mathcal{U}^T G(w)\mathcal{U}\| \\ &\leq |z|\|G(z)\||z - w| + |w|\|z - w\|\|G(z)\| + |w|^2|z - w|\|G(z)\|\|G(w)\| \\ &\leq 8|z - w|, \end{aligned}$$

where we used the resolvent identity (15), Lemma 16, and the fact that $\frac{|w|}{|z|} \leq 1$. This shows that f is 8-Lipschitz in S_{σ_j} .

The proof for g is similar. First, for $|z| \geq |w| \geq 4(\sqrt{N} + \sqrt{n})$, by the triangle inequality, we have

$$\begin{aligned} \|g(z) - g(w)\| &\leq \|z^2\mathcal{U}^T \Phi(z)\mathcal{U} - zw\mathcal{U}^T \Phi(z)\mathcal{U}\| + \|zw\mathcal{U}^T \Phi(z)\mathcal{U} - w^2\mathcal{U}^T \Phi(z)\mathcal{U}\| \\ &\quad + \|w^2\mathcal{U}^T \Phi(z)\mathcal{U} - w^2\mathcal{U}^T \Phi(w)\mathcal{U}\| \\ &\leq |z|\|z - w\|\|\mathcal{U}^T \Phi(z)\mathcal{U}\| + |w|\|z - w\|\|\mathcal{U}^T \Phi(z)\mathcal{U}\| + |w|^2\|\mathcal{U}^T (\Phi(z) - \Phi(w))\mathcal{U}\|. \end{aligned} \tag{84}$$

Using the explicit expression in (22), we find that

$$\|\mathcal{U}^T (\Phi(z) - \Phi(w))\mathcal{U}\| = \max \left\{ \frac{|\phi_1(z) - \phi_1(w)|}{|\phi_1(z)\phi_1(w)|}, \frac{|\phi_2(z) - \phi_2(w)|}{|\phi_2(z)\phi_2(w)|} \right\}.$$

By (20) and the resolvent identity (15),

$$\begin{aligned} |\phi_1(z) - \phi_1(w)| &= |z - w - \text{tr } \mathcal{I}^d(G(z) - G(w))| \\ &= |z - w - (z - w) \text{tr } \mathcal{I}^d G(z)G(w)| \\ &\leq |z - w| (1 + (N+n)\|G(z)\|\|G(w)\|) \\ &\leq |z - w| \left(1 + \frac{4(N+n)}{|z||w|} \right) \\ &\leq \frac{5}{4}|z - w|, \end{aligned}$$

where we used Lemma 16 and the facts that $\frac{N+n}{|w|} \leq \frac{|w|}{16}$ and $\frac{|w|}{|z|} \leq 1$. The same upper bound also holds for $|\phi_2(z) - \phi_2(w)|$. Combining these estimates with (27), we have

$$\|\mathcal{U}^T(\Phi(z) - \Phi(w))\mathcal{U}\| \leq \frac{80}{49} \frac{|z-w|}{|z||w|}.$$

Notice that $\|\mathcal{U}^T\Phi(z)\mathcal{U}\| \leq \frac{8}{7|z|}$ for any $|z| \geq 4(\sqrt{N} + \sqrt{n})$, which can be verified using (23) and the bounds in (27). Inserting these bounds into (84) yields that g is 8-Lipschitz in S_{σ_j} ; we omit the details.

The proof is complete by taking a union bound over $j \in \llbracket i_0, r_0 \rrbracket$. \square

APPENDIX A. PROOF OF THEOREM 3

This section is devoted to the proof of Theorem 3. Unless otherwise specified, we let C and c be positive constants, which may change from one occurrence to the next, depending only on the sub-gaussian moment of the entries of E . Without loss of generality, we assume $N \leq n$, for if not, one can simply replace A and E by their transposes.

Let $P_1 \in \mathbb{R}^{N \times N}$ be the orthogonal projection matrix onto the subspace $\text{Span}\{u_1, \dots, u_r\}^\perp$ and $P_2 \in \mathbb{R}^{n \times n}$ be that onto $\text{Span}\{v_1, \dots, v_r\}^\perp$.

We first prove that

$$\begin{aligned} \tilde{\sigma}_1 \sin \angle(u_1, \tilde{u}_1) + \sqrt{2}\|E\| \sin \angle(v_1, \tilde{v}_1) &\geq \|P_1 E v_1\|, \\ \tilde{\sigma}_1 \sin \angle(v_1, \tilde{v}_1) + \sqrt{2}\|E\| \sin \angle(u_1, \tilde{u}_1) &\geq \|P_2 E^T u_1\|. \end{aligned} \quad (85)$$

To start, observe that

$$\begin{aligned} \sin^2 \angle(u_1, \tilde{u}_1) &= 1 - \langle u_1, \tilde{u}_1 \rangle^2 = \|\tilde{u}_1\|^2 - \langle u_1, \tilde{u}_1 \rangle^2 \\ &= \sum_{i=2}^r \langle u_i, \tilde{u}_1 \rangle^2 + \|P_1 \tilde{u}_1\|^2 \geq \|P_1 \tilde{u}_1\|^2. \end{aligned} \quad (86)$$

By multiplying by P_1 on all sides of the equation $\tilde{A}\tilde{v}_1 = A\tilde{v}_1 + E\tilde{v}_1 = \tilde{\sigma}_1\tilde{u}_1$, we get $P_1 E \tilde{v}_1 = \tilde{\sigma}_1 P_1 \tilde{u}_1$. Continuing from (86), we have

$$\tilde{\sigma}_1 \sin \angle(u_1, \tilde{u}_1) \geq \|P_1 E \tilde{v}_1\|. \quad (87)$$

Likewise, we also have

$$\tilde{\sigma}_1 \sin \angle(v_1, \tilde{v}_1) \geq \|P_2 E^T \tilde{u}_1\|. \quad (88)$$

Let α denote the angle between u_1 and \tilde{u}_1 (taken in $[0, \pi]$) and β denote the angle between v_1 and \tilde{v}_1 (taken in $[0, \pi]$). By possibly multiplying \tilde{u}_1, \tilde{v}_1 by -1 , it suffices to consider one of the following cases: either (i) $\alpha, \beta \in [0, \frac{\pi}{2}]$; or (ii) $\alpha \in [\frac{\pi}{2}, \pi]$ and $\beta \in [0, \frac{\pi}{2}]$.

If $\alpha, \beta \in [0, \frac{\pi}{2}]$, by simple trigonometric identities,

$$\|u_1 - \tilde{u}_1\| = 2 \sin(\alpha/2) = \frac{\sin \alpha}{\cos(\alpha/2)} \leq \sqrt{2} \sin \alpha = \sqrt{2} \sin \angle(u_1, \tilde{u}_1). \quad (89)$$

Similarly, we have

$$\|v_1 - \tilde{v}_1\| \leq \sqrt{2} \sin \angle(v_1, \tilde{v}_1).$$

From (87) and the triangle inequality, we obtain

$$\begin{aligned}\tilde{\sigma}_1 \sin \angle(u_1, \tilde{u}_1) &\geq \|P_1 E \tilde{v}_1\| \\ &\geq \|P_1 E v_1\| - \|P_1 E(\tilde{v}_1 - v_1)\| \\ &\geq \|P_1 E v_1\| - \|E\| \|\tilde{v}_1 - v_1\| \\ &\geq \|P_1 E v_1\| - \sqrt{2} \|E\| \sin \angle(v_1, \tilde{v}_1).\end{aligned}$$

Repeating the same argument, we obtain

$$\tilde{\sigma}_1 \sin \angle(v_1, \tilde{v}_1) \geq \|P_2 E^T u_1\| - \sqrt{2} \|E\| \sin \angle(u_1, \tilde{u}_1).$$

In the second case, if $\alpha \in [\frac{\pi}{2}, \pi]$ and $\beta \in [0, \frac{\pi}{2}]$, we still have $\|v_1 - \tilde{v}_1\| \leq \sqrt{2} \sin \angle(v_1, \tilde{v}_1)$, and hence

$$\tilde{\sigma}_1 \sin \angle(u_1, \tilde{u}_1) \geq \|P_1 E v_1\| - \sqrt{2} \|E\| \sin \angle(v_1, \tilde{v}_1).$$

Since $\theta := \pi - \alpha \in [0, \frac{\pi}{2}]$, using the same estimate as in (89), we see that

$$\|u_1 + \tilde{u}_1\| = 2 \sin(\theta/2) = \frac{\sin \theta}{\cos(\theta/2)} \leq \sqrt{2} \sin \theta = \sqrt{2} \sin(\pi - \alpha) = \sqrt{2} \sin \angle(u_1, \tilde{u}_1).$$

Thus, from (88), we conclude that

$$\begin{aligned}\tilde{\sigma}_1 \sin \angle(v_1, \tilde{v}_1) &\geq \|P_2 E^T \tilde{u}_1\| \\ &\geq \|P_2 E^T u_1\| - \|P_2 E^T(\tilde{u}_1 + u_1)\| \\ &\geq \|P_2 E^T u_1\| - \|E\| \|\tilde{u}_1 + u_1\| \\ &\geq \|P_2 E^T u_1\| - \sqrt{2} \|E\| \sin \angle(u_1, \tilde{u}_1).\end{aligned}$$

Rearranging the terms yields (85).

We now turn to the proof of Theorem 3. It follows immediately from (85) that

$$\max\{\sin \angle(u_1, \tilde{u}_1), \sin \angle(v_1, \tilde{v}_1)\} \geq \frac{\max\{\|P_1 E v_1\|, \|P_2 E^T u_1\|\}}{\tilde{\sigma}_1 + \sqrt{2} \|E\|} \geq \frac{\max\{\|P_1 E v_1\|, \|P_2 E^T u_1\|\}}{\sigma_1 + (1 + \sqrt{2}) \|E\|}, \quad (90)$$

where in the last inequality we used $\tilde{\sigma}_1 \leq \sigma_1 + \|E\|$ from the classical Weyl's inequality. Furthermore, using [50, Lemma 11.8], one has

$$\mathbb{P}(|\|P_2 E^T u_1\|^2 - (n - r)| > t) \leq C \exp\left(-\min\left\{\frac{t^2}{n - r}, t\right\}\right)$$

for any $t \geq 0$. Thus, since $N \leq n$,

$$\max\{\|P_1 E v_1\|, \|P_2 E^T u_1\|\} \geq \|P_2 E^T u_1\| \geq \frac{1}{2} \sqrt{n - r}$$

with probability at least $1 - C \exp(-c(n - r))$. Plugging into (90), we get

$$\max\{\sin \angle(u_1, \tilde{u}_1), \sin \angle(v_1, \tilde{v}_1)\} \geq \frac{1}{2} \frac{\sqrt{n - r}}{\sigma_1 + (1 + \sqrt{2}) \|E\|} = \frac{\|E\|}{2\sigma_1} \frac{\frac{\sqrt{n - r}}{\|E\|}}{1 + (1 + \sqrt{2}) \frac{\|E\|}{\sigma_1}}.$$

Applying [56, Proposition 2.4], we have $\|E\| \leq 2(\sqrt{N} + \sqrt{n})$ with probability at least $1 - C \exp(-c(\sqrt{N} + \sqrt{n})^2)$. It follows that $\frac{\sqrt{n - r}}{\|E\|} \geq \frac{1}{4\sqrt{2}}$, where we used that

$r \leq n/2$. Finally, we conclude that

$$\max\{\sin \angle(u_1, \tilde{u}_1), \sin \angle(v_1, \tilde{v}_1)\} \geq \frac{1}{8\sqrt{2}} \frac{\frac{\|E\|}{\sigma_1}}{1 + (1 + \sqrt{2}) \frac{\|E\|}{\sigma_1}}$$

with probability at least $1 - C \exp(-c(n-r))$. The claim now follows from the fact that $r \leq n/2$.

APPENDIX B. PROOFS OF PROPOSITIONS 8, 10, 11, (21) AND LEMMA 13

In this section, we collect the proofs of Propositions 8, 10, 11, (21) and Lemma 13 from Section 3. We continue to use the notation introduced in the previous sections.

B.1. Proof of Proposition 8. Let $0 \leq \alpha_1 \leq \dots \leq \alpha_p \leq \pi/2$ be the principal angles between U and \tilde{U} . Then

$$\cos \alpha_p = \cos \angle(U, \tilde{U}).$$

Let $\{u_1, \dots, u_p\}$ (resp. $\{\tilde{u}_1, \dots, \tilde{u}_p\}$) be an orthonormal basis for U (resp. for \tilde{U}). Denote the matrices $\mathbf{U} := (u_1, \dots, u_p)$ and $\mathbf{V} := (v_1, \dots, v_p)$. [14, Theorem 1] provides the connection between the principle angles and the singular values of $\mathbf{U}^T \tilde{\mathbf{U}}$. That is, consider the SVD of $\mathbf{U}^T \tilde{\mathbf{U}}$ given by

$$\mathbf{U}^T \tilde{\mathbf{U}} = Y_1 C_1 Z_1^T,$$

where C_1 is a diagonal matrix composed of the singular values and Y_1, Z_1 are $p \times p$ orthogonal matrices. Then the diagonal entries of C_1 are exactly $\cos \alpha_1, \dots, \cos \alpha_p$.

Analogously, let $0 \leq \beta_1 \leq \dots \leq \beta_p \leq \pi/2$ be the principal angles between V and \tilde{V} . Then

$$\cos \beta_p = \cos \angle(V, \tilde{V}).$$

For the orthonormal basis $\{v_1, \dots, v_p\}$ for V and $\{\tilde{v}_1, \dots, \tilde{v}_p\}$ for \tilde{V} , denote $\mathbf{V} := (v_1, \dots, v_p)$ and $\tilde{\mathbf{V}} := (\tilde{v}_1, \dots, \tilde{v}_p)$. Consider the SVD

$$\mathbf{V}^T \tilde{\mathbf{V}} = Y_2 C_2 Z_2^T,$$

where C_2 is a diagonal matrix composed of the singular values and Y_2, Z_2 are orthogonal matrices. Then the diagonal entries of C_2 are $\cos \beta_1, \dots, \cos \beta_p$.

Next, define $\mathbf{w}_i := (u_i^T, v_i^T)^T$ and $\mathbf{w}_{i+p} := (u_i^T, -v_i^T)^T$ for $1 \leq i \leq p$. Define $\tilde{\mathbf{w}}_i$'s ($1 \leq i \leq 2p$) analogously. It is easy to verify that $\frac{1}{\sqrt{2}}\{\mathbf{w}_1, \dots, \mathbf{w}_{2p}\}$ forms an orthonormal basis of W and $\frac{1}{\sqrt{2}}\{\tilde{\mathbf{w}}_1, \dots, \tilde{\mathbf{w}}_{2p}\}$ forms an orthonormal basis of \tilde{W} . Denote the matrices $\mathbf{W} := \frac{1}{\sqrt{2}}(\mathbf{w}_1, \dots, \mathbf{w}_{2p})$ and $\tilde{\mathbf{W}} := \frac{1}{\sqrt{2}}(\tilde{\mathbf{w}}_1, \dots, \tilde{\mathbf{w}}_{2p})$. Let $0 \leq \gamma_1 \leq \dots \leq \gamma_{2p} \leq \pi/2$ be the principal angles between W and \tilde{W} . Then $\cos \gamma_1, \dots, \cos \gamma_{2p}$ are the singular values of $\mathbf{W}^T \tilde{\mathbf{W}}$ by [14, Theorem 1]. In particular, the smallest singular value of $\mathbf{W}^T \tilde{\mathbf{W}}$ is $\cos \gamma_{2p} = \cos \angle(W, \tilde{W})$.

Observe that

$$\mathbf{W}^T \tilde{\mathbf{W}} = \frac{1}{2} \begin{pmatrix} U & U \\ V & -V \end{pmatrix}^T \begin{pmatrix} \tilde{U} & \tilde{U} \\ \tilde{V} & -\tilde{V} \end{pmatrix} = \frac{1}{2} \begin{pmatrix} U^T \tilde{U} + V^T \tilde{V} & U^T \tilde{U} - V^T \tilde{V} \\ U^T \tilde{U} - V^T \tilde{V} & U^T \tilde{U} + V^T \tilde{V} \end{pmatrix}.$$

Using the SVDs of $U^T \tilde{U}$ and $V^T \tilde{V}$ and simple computations, we further get

$$\mathbf{W}^T \tilde{\mathbf{W}} = \frac{1}{\sqrt{2}} \begin{pmatrix} Y_1 & -Y_2 \\ Y_1 & Y_2 \end{pmatrix} \begin{pmatrix} C_1 & 0 \\ 0 & C_2 \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} Z_1^T & Z_1^T \\ -Z_2^T & Z_2^T \end{pmatrix} =: YCZ^T.$$

Since Y_1, Y_2, Z_1, Z_2 are orthogonal matrices, it is easy to verify that both Y and Z are $2p \times 2p$ orthogonal matrices. Hence, YCZ^T is the SVD of $\mathbf{W}^T \tilde{\mathbf{W}}$. By [14, Theorem 1], the singular values $\cos \gamma_1, \dots, \cos \gamma_{2p}$ of $\mathbf{W}^T \tilde{\mathbf{W}}$ are exactly

$$\cos \alpha_1, \dots, \cos \alpha_p, \cos \beta_1, \dots, \cos \beta_p.$$

Hence,

$$\cos \angle(W, \tilde{W}) = \cos \gamma_{2p} = \min\{\cos \alpha_p, \cos \beta_p\} = \min\{\cos \angle(U, \tilde{U}), \cos \angle(V, \tilde{V})\},$$

which is equivalent to

$$\sin \angle(W, \tilde{W}) = \max\{\sin \angle(U, \tilde{U}), \sin \angle(V, \tilde{V})\}.$$

This completes the proof.

B.2. Proof of Propositions 10. Note that $\Phi(x)$ is well-defined when $|x| > \|\mathcal{E}\|$. From the expression of $\mathcal{U}^T \Phi(x) \mathcal{U}$ in (22), using the block structure of matrices, we rewrite

$$I_{2r-2r_0} - \mathcal{U}_J^T \Phi(x) \mathcal{U}_J \mathcal{D}_J = \begin{pmatrix} I_{r-r_0} - \alpha \mathcal{D}_{[[r_0+1, r]]} & -\beta \mathcal{D}_{[[r+r_0+1, 2r]]} \\ -\beta \mathcal{D}_{[[r_0+1, r]]} & I_{r-r_0} - \alpha \mathcal{D}_{[[r+r_0+1, 2r]]} \end{pmatrix}.$$

Note that $\mathcal{D}_{[[r+r_0+1, 2r]]} = -\mathcal{D}_{[[r_0+1, r]]}$ by the definition of \mathcal{D} . We compute the eigenvalues of $(I_{2r-2r_0} - \mathcal{U}_J^T \Phi(\tilde{\sigma}_j) \mathcal{U}_J \mathcal{D}_J)(I_{2r-2r_0} - \mathcal{U}_J^T \Phi(\tilde{\sigma}_j) \mathcal{U}_J \mathcal{D}_J)^T$, which, after simplification, has the following format

$$\begin{pmatrix} (I - \alpha \mathcal{D}_{[[r_0+1, r]]})^2 + \beta^2 \mathcal{D}_{[[r_0+1, r]]}^2 & 2\alpha\beta \mathcal{D}_{[[r_0+1, r]]}^2 \\ 2\alpha\beta \mathcal{D}_{[[r_0+1, r]]}^2 & (I + \alpha \mathcal{D}_{[[r_0+1, r]]})^2 + \beta^2 \mathcal{D}_{[[r_0+1, r]]}^2 \end{pmatrix}. \quad (91)$$

Note that each block of (91) is a diagonal matrix. Using basic linear algebra and a simple computation, we further obtain the eigenvalues of the above matrix are given by

$$1 + \sigma_t^2(\alpha^2 + \beta^2) \pm 2|\alpha|\sigma_t \sqrt{1 + \beta^2 \sigma_t^2} = \left(\sqrt{1 + \beta^2 \sigma_t^2} \pm |\alpha|\sigma_t \right)^2$$

for $r_0 + 1 \leq t \leq r$. Taking the square roots yields the singular values of $I_{2r-2r_0} - \mathcal{U}_J^T \Phi(x) \mathcal{U}_J \mathcal{D}_J$ and completes the proof.

B.3. Proof of Propositions 11. Note that $\Phi(z)$ is well-defined when $|z| > \|\mathcal{E}\|$. We first compute the eigenvalues of

$$(\mathcal{D}^{-1} - \mathcal{U}^T \Phi(z) \mathcal{U}) (\mathcal{D}^{-1} - \mathcal{U}^T \Phi(z) \mathcal{U})^*, \quad (92)$$

where we remind the readers that \mathcal{D} and \mathcal{U} are defined in Section 3.1. In particular, $\mathcal{D} = \text{diag}(\text{diag}(D), -\text{diag}(D))$. Recall the definition of $\mathcal{U}^T \Phi(z) \mathcal{U}$ in (22). Note that

$$\mathcal{D}^{-1} - \mathcal{U}^T \Phi(z) \mathcal{U} = \begin{pmatrix} D^{-1} - \alpha I_r & -\beta I_r \\ -\beta I_r & -D^{-1} - \alpha I_r \end{pmatrix}$$

where each block is a diagonal matrix. Elementary (yet tedious) calculations yield that the eigenvalues of (92) are

$$|\alpha|^2 + |\beta|^2 + \frac{1}{\sigma_t^2} \pm \left((2|\alpha|^2 + 2|\beta|^2 + \bar{\alpha}^2 - \bar{\beta}^2 + \alpha^2 - \beta^2) \frac{1}{\sigma_t^2} + 2|\alpha|^2 |\beta|^2 + \bar{\alpha}^2 \beta^2 + \alpha^2 \bar{\beta}^2 \right)^{1/2}.$$

With further simplification after plugging in the expressions of α, β , we denote these eigenvalues by

$$\gamma_{\pm} := \frac{1}{\sigma_l^2} + \frac{1}{2} \left(\frac{1}{|\phi_1|^2} + \frac{1}{|\phi_2|^2} \right) \pm \frac{1}{2} \left[\frac{4}{\sigma_l^2} \left| \frac{1}{\phi_1} + \frac{1}{\bar{\phi}_2} \right|^2 + \left(\frac{1}{|\phi_1|^2} - \frac{1}{|\phi_2|^2} \right)^2 \right]^{1/2}$$

for $1 \leq l \leq r$. Since the entries of E have continuous distribution, $\mathcal{D}^{-1} - \mathcal{U}^T \Phi(z) \mathcal{U}$ is invertible with probability 1. Consequently,

$$\left\| (\mathcal{D}^{-1} - \mathcal{U}^T \Phi(z) \mathcal{U})^{-1} \right\|^2 = \max_{1 \leq l \leq r} \frac{1}{\gamma_{l-}} = \max_{1 \leq l \leq r} \frac{\sigma_l^2}{|\sigma_l^2 - \phi_1 \phi_2|^2} \mathcal{Q},$$

where

$$\mathcal{Q} := |\phi_1 \phi_2|^2 + \frac{1}{2} \sigma_l^2 (|\phi_1|^2 + |\phi_2|^2) + \frac{1}{2} \sigma_l [4|\phi_1 \phi_2|^2 |\phi_1 + \bar{\phi}_2|^2 + \sigma_l^2 (|\phi_1|^2 - |\phi_2|^2)^2]^{1/2}.$$

The conclusion of Propositions 11 follows by taking a square root.

B.4. Proof of (21). The singular value decomposition of E is given by $E = X \text{diag}(\eta_1, \dots, \eta_N) Y^T$, where the columns of X (resp. Y) are $\mathbf{x}_1, \dots, \mathbf{x}_N \in \mathbb{R}^N$ (resp. $\mathbf{y}_1, \dots, \mathbf{y}_N \in \mathbb{R}^n$). Clearly, some of the η_i 's may be zero. However, if $n > N$, then E trivially maps an $(n - N)$ -dimensional space to zero; we assume this subspace has orthonormal basis $\mathbf{h}_1, \dots, \mathbf{h}_{n-N} \in \mathbb{R}^n$.

The spectral decomposition of \mathcal{E} is then given by

$$\mathcal{E} = \mathcal{W} \text{diag}(\eta_1, \dots, \eta_N, -\eta_1, \dots, -\eta_N) \mathcal{W}^T,$$

where the columns of \mathcal{W} are the orthonormal vectors $\mathbf{w}_i = \frac{1}{\sqrt{2}} (\mathbf{x}_i^T, \mathbf{y}_i^T)^T \in \mathbb{R}^{N+n}$ and $\mathbf{w}_{N+i} = \frac{1}{\sqrt{2}} (\mathbf{x}_i^T, -\mathbf{y}_i^T)^T \in \mathbb{R}^{N+n}$ for $1 \leq i \leq N$. Likewise, \mathcal{E} trivially maps an $(n - N)$ -dimensional space to zero; this subspace is spanned by the orthonormal vectors $\mathbf{w}_{2N+j} = (0, \mathbf{h}_j^T)^T \in \mathbb{R}^{N+n}$ for $j = 1, \dots, n - N$. Thus, the spectral decomposition of the resolvent $G(z)$ can be expressed as

$$G(z) = (z - \mathcal{E})^{-1} = \sum_{i=1}^N \frac{\mathbf{w}_i \mathbf{w}_i^T}{z - \eta_i} + \sum_{i=1}^N \frac{\mathbf{w}_{N+i} \mathbf{w}_{N+i}^T}{z + \eta_i} + \frac{1}{z} \sum_{j=1}^{n-N} \mathbf{w}_{2N+j} \mathbf{w}_{2N+j}^T.$$

It follows that

$$\text{tr} \mathcal{I}^d G(z) = \frac{1}{2} \sum_{i=1}^N \left(\frac{1}{z - \eta_i} + \frac{1}{z + \eta_i} \right) + \frac{1}{z} (n - N), \quad (93)$$

$$\text{tr} \mathcal{I}^u G(z) = \frac{1}{2} \sum_{i=1}^N \left(\frac{1}{z - \eta_i} - \frac{1}{z + \eta_i} \right), \quad (94)$$

and thus, from (20), we see that (21) holds.

B.5. Proof of Lemma 13. We work on the event that $\|E\| \leq 2(\sqrt{N} + \sqrt{n})$, which holds with probability at least $1 - 2e^{-(\sqrt{N} + \sqrt{n})^2/2}$ by Lemma 15.

If $r^2 > \frac{1}{27}(\sqrt{N} + \sqrt{n})$, then the conclusion follows directly from the Weyl's inequality and the supposition on n, N :

$$\max_{1 \leq l \leq r} |\tilde{\sigma}_l - \sigma_l| \leq \|E\| \leq 2(\sqrt{N} + \sqrt{n}) \leq \eta r = 54r^2 \sqrt{(K+8) \log(N+n)}.$$

If $r^2 \leq \frac{1}{27}(\sqrt{N} + \sqrt{n})$, then $\eta r < \frac{1}{4}n^2 < \frac{1}{2}\sigma_l$ for any $\sigma_l > \frac{1}{2}n^2$. We apply [51, Theorem 23] by noting that E is $(2, \frac{1}{2}, 2)$ -concentrated in the Definition 11 from [51] to get

$$\tilde{\sigma}_l \geq \sigma_l - \eta r$$

uniformly for any $1 \leq l \leq r$ with probability at least $1 - 9^{r+1} \exp(-(\eta r)^2/32)$. Conditioning on this event, with the assumption that $\sigma_l > \frac{1}{2}n^2$ which implies $\tilde{\sigma}_l > \frac{1}{2}\sigma_l$, [51, Theorem 23] yields the lower bound

$$\tilde{\sigma}_l \leq \sigma_l + t\sqrt{r} + 128\frac{\sqrt{r}}{n} + 2^9\frac{r}{n^{5/2}}$$

with probability at least $1 - 4 \cdot 9^{2r} \exp(-rt^2/32)$ for any $t > 0$. Note that $128\frac{\sqrt{r}}{n} + 2^9\frac{r}{n^{5/2}} \leq (3/100)\eta r$ by the supposition on n . The conclusion follows by taking $t\sqrt{r} = \frac{1}{2}\eta r$, a union bound over $l \leq r_0$ and simplifying the probability bounds using $(\eta r)^2 > 54^2 r^4$.

APPENDIX C. PROOFS OF LEMMA 16, 17, AND 18

In this section, we collect the proofs of Lemma 16, 17, and 18.

C.1. Proof of Lemma 16. By writing $G(z) := (z - \mathcal{E})^{-1}$ as a Neumann series, we find

$$\|G(z)\| \leq \frac{1}{|z|} \sum_{k=0}^{\infty} \left(\frac{\|\mathcal{E}\|}{|z|} \right)^k.$$

Since $\|E\| = \|\mathcal{E}\|$ and $\frac{\|\mathcal{E}\|}{|z|} \leq 1/2$ by assumption, the claim for $\|G(z)\|$ follows. Since $\|\mathcal{E}^{(k)}\| \leq \|\mathcal{E}\|^k$, the same proof also applies to $G^{(k)}(z)$. The bounds in (27) follow from (20), the bound

$$\max\{|\operatorname{tr} \mathcal{I}^u G(z)|, |\operatorname{tr} \mathcal{I}^d G(z)|\} \leq (N+n)\|G(z)\| \leq \frac{2(N+n)}{|z|} \leq \frac{|z|}{8}, \quad (95)$$

and the triangle inequality.

C.2. Proof of Lemma 17. By writing $G(z) := (z - \mathcal{E})^{-1}$ as a Neumann series, we see that

$$\left\| G(z) - \frac{1}{z} I_{N+n} - \frac{\mathcal{E}}{z^2} \right\| \leq \frac{1}{|z|} \sum_{k=2}^{\infty} \left(\frac{\|\mathcal{E}\|}{|z|} \right)^k = \frac{\|\mathcal{E}\|^2}{|z|^2(|z| - \|\mathcal{E}\|)} \leq \frac{2\|\mathcal{E}\|^2}{|z|^3}$$

since $\frac{\|\mathcal{E}\|}{|z|} \leq 1/2$ and $|z| - \|\mathcal{E}\| \geq \frac{1}{2}|z|$ by assumption.

C.3. Proof of Lemma 18. By the rotational invariance of E , it suffices to assume that U is the matrix with columns e_1, \dots, e_r , where e_1, \dots, e_N is the canonical basis in \mathbb{R}^N , and the columns of V are given by f_1, \dots, f_r , where f_1, \dots, f_n is the canonical basis in \mathbb{R}^n . By the definition of \mathcal{U} in (9), it is easy to verify that

$$2(\mathcal{U}^T \mathcal{E} \mathcal{U})_{ij} = \begin{cases} E_{ij} + E_{ji} & \text{for } 1 \leq i, j \leq r; \\ -E_{i,j-r} + E_{j-r,i} & \text{for } 1 \leq i \leq r, r+1 \leq j \leq 2r; \\ E_{i-r,j} - E_{j,i-r} & \text{for } r+1 \leq i \leq 2r, 1 \leq j \leq r; \\ -E_{i-r,j-r} - E_{j-r,i-r} & \text{for } r+1 \leq i, j \leq 2r. \end{cases}$$

Denote E_r the $r \times r$ upper-left corner of E . We rewrite

$$\mathcal{U}^T \mathcal{E} \mathcal{U} = \frac{1}{2} \begin{pmatrix} E_r + E_r^T & -E_r + E_r^T \\ E_r - E_r^T & -E_r - E_r^T \end{pmatrix} = \frac{1}{2} \begin{pmatrix} I_r & I_r \\ I_r & -I_r \end{pmatrix} \begin{pmatrix} E_r & 0 \\ 0 & E_r^T \end{pmatrix} \begin{pmatrix} I_r & -I_r \\ I_r & I_r \end{pmatrix}.$$

By elementary computation and the fact that $E_r E_r^T$ and $E_r^T E_r$ share the same non-trivial eigenvalues, we get

$$\|\mathcal{U}^T \mathcal{E} \mathcal{U}\| \leq \frac{1}{2} \left\| \begin{pmatrix} I_r & I_r \\ I_r & -I_r \end{pmatrix} \right\| \left\| \begin{pmatrix} E_r & 0 \\ 0 & E_r^T \end{pmatrix} \right\| \left\| \begin{pmatrix} I_r & -I_r \\ I_r & I_r \end{pmatrix} \right\| = \|E_r\|.$$

Hence, by invoking Lemma 15, we have

$$\|\mathcal{U}^T \mathcal{E} \mathcal{U}\| \leq \|E_r\| \leq 2\sqrt{r} + \sqrt{2K \log(N+n)}$$

with probability at least $1 - 2(N+n)^{-K}$.

REFERENCES

- [1] E. Abbe, J. Fan, K. Wang, and Y. Zhong. Entrywise eigenvector analysis of random matrices with low expected rank. *Ann. Statist.*, 48(3):1452–1474, 2020.
- [2] R. Allez and J.-P. Bouchaud. Eigenvector dynamics under free addition. *Random Matrices Theory Appl.*, 3(3):1450010, 17, 2014.
- [3] Z. Bai and J. W. Silverstein. *Spectral analysis of large dimensional random matrices*. Springer Series in Statistics. Springer, New York, second edition, 2010.
- [4] Z. Bai and J.-f. Yao. Central limit theorems for eigenvalues in a spiked population model. *Ann. Inst. Henri Poincaré Probab. Stat.*, 44(3):447–474, 2008.
- [5] J. Baik, G. Ben Arous, and S. Péché. Phase transition of the largest eigenvalue for nonnull complex sample covariance matrices. *Ann. Probab.*, 33(5):1643–1697, 2005.
- [6] Z. Bao, X. Ding, J. Wang, and K. Wang. Statistical inference for principal components of spiked covariance matrices. *Ann. Statist.*, 50(2):1144–1169, 2022.
- [7] Z. Bao, X. Ding, and K. Wang. Singular vector and singular subspace distribution for the matrix denoising model. *The Annals of Statistics*, 49(1):370–392, 2021.
- [8] Z. Bao and D. Wang. Eigenvector distribution in the critical regime of BBP transition. *Probab. Theory Related Fields*, 182(1-2):399–479, 2022.
- [9] F. Benaych-Georges, N. Enriquez, and A. Michail. Eigenvectors of a matrix under random perturbation. *Random Matrices Theory Appl.*, 10(2):Paper No. 2150023, 17, 2021.
- [10] F. Benaych-Georges, A. Guionnet, and M. Maida. Fluctuations of the extreme eigenvalues of finite rank deformations of random matrices. *Electron. J. Probab.*, 16:no. 60, 1621–1662, 2011.
- [11] F. Benaych-Georges and R. R. Nadakuditi. The eigenvalues and eigenvectors of finite, low rank perturbations of large random matrices. *Adv. Math.*, 227(1):494–521, 2011.
- [12] F. Benaych-Georges and R. R. Nadakuditi. The singular values and vectors of low rank perturbations of large rectangular random matrices. *J. Multivariate Anal.*, 111:120–135, 2012.
- [13] R. Bhatia. *Matrix analysis*, volume 169 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1997.
- [14] A. Björck and G. H. Golub. Numerical methods for computing angles between linear subspaces. *Math. Comp.*, 27:579–594, 1973.
- [15] A. Bloemendal, L. Erdős, A. Knowles, H.-T. Yau, and J. Yin. Isotropic local laws for sample covariance and generalized Wigner matrices. *Electron. J. Probab.*, 19:no. 33, 53, 2014.
- [16] A. Bloemendal, A. Knowles, H.-T. Yau, and J. Yin. On the principal components of sample covariance matrices. *Probability theory and related fields*, 164(1):459–552, 2016.
- [17] T. Cai, Z. Ma, and Y. Wu. Optimal estimation and rank detection for sparse spiked covariance matrices. *Probab. Theory Related Fields*, 161(3-4):781–815, 2015.
- [18] E. J. Candès and Y. Plan. Matrix completion with noise. *Proceedings of the IEEE*, 98:925–936, 2010.
- [19] E. J. Candès and B. Recht. Exact matrix completion via convex optimization. *Found. Comput. Math.*, 9(6):717–772, 2009.

- [20] E. J. Candès, J. Romberg, and T. Tao. Robust uncertainty principles: exact signal reconstruction from highly incomplete frequency information. *IEEE Trans. Inform. Theory*, 52(2):489–509, 2006.
- [21] J. Cape, M. Tang, and C. E. Priebe. Signal-plus-noise matrix models: eigenvector deviations and fluctuations. *Biometrika*, 106(1):243–250, 2019.
- [22] J. Cape, M. Tang, and C. E. Priebe. The two-to-infinity norm and singular subspace geometry with applications to high-dimensional statistics. *Ann. Statist.*, 47(5):2405–2439, 2019.
- [23] M. Capitaine, C. Donati-Martin, and D. Féral. Central limit theorems for eigenvalues of deformations of Wigner matrices. *Ann. Inst. Henri Poincaré Probab. Stat.*, 48(1):107–133, 2012.
- [24] P. Chin, A. Rao, and V. Vu. Stochastic block model and community detection in sparse graphs: A spectral algorithm with optimal rate of recovery. In P. Grünwald, E. Hazan, and S. Kale, editors, *Proceedings of The 28th Conference on Learning Theory*, volume 40 of *Proceedings of Machine Learning Research*, pages 391–423, Paris, France, 03–06 Jul 2015. PMLR.
- [25] C. Davis and W. M. Kahan. The rotation of eigenvectors by a perturbation. III. *SIAM J. Numer. Anal.*, 7:1–46, 1970.
- [26] J. Eldridge, M. Belkin, and Y. Wang. Unperturbed: spectral analysis beyond Davis-Kahan. In *Algorithmic learning theory 2018*, volume 83 of *Proc. Mach. Learn. Res. (PMLR)*, page 38. Proceedings of Machine Learning Research PMLR, [place of publication not identified], 2018.
- [27] L. Erdős, H.-T. Yau, and J. Yin. Bulk universality for generalized Wigner matrices. *Probab. Theory Related Fields*, 154(1-2):341–407, 2012.
- [28] L. Erdős, H.-T. Yau, and J. Yin. Rigidity of eigenvalues of generalized Wigner matrices. *Adv. Math.*, 229(3):1435–1515, 2012.
- [29] L. Erdős, A. Knowles, H.-T. Yau, and J. Yin. Spectral statistics of erdős-rényi graphs ii: Eigenvalue spacing and the extreme eigenvalues. *Communications in Mathematical Physics*, 314(3):587–640, 2012.
- [30] L. Erdős, A. Knowles, H.-T. Yau, and J. Yin. Spectral statistics of erdős-rényi graphs I: Local semicircle law. *Ann. Probab.*, 41(3B):2279–2375, 2013.
- [31] J. Fan and X. Han. Estimation of the false discovery proportion with unknown dependence. *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 79(4):1143–1164, 2017.
- [32] J. Fan, Y. Liao, and M. Mincheva. Large covariance estimation by thresholding principal orthogonal complements. *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 75(4):603–680, 2013. With 33 discussions by 57 authors and a reply by Fan, Liao and Mincheva.
- [33] J. Fan, W. Wang, and Y. Zhong. An ℓ_∞ eigenvector perturbation bound and its application to robust covariance estimation. *J. Mach. Learn. Res.*, 18:Paper No. 207, 42, 2017.
- [34] G. H. Golub and C. F. Van Loan. *Matrix computations*. JHU press, 2013.
- [35] O. Guédon and R. Vershynin. Community detection in sparse networks via Grothendieck’s inequality. *Probab. Theory Related Fields*, 165(3-4):1025–1049, 2016.
- [36] Y. He, A. Knowles, and R. Rosenthal. Isotropic self-consistent equations for mean-field random matrices. *Probab. Theory Related Fields*, 171(1-2):203–249, 2018.
- [37] N. J. Higham. A survey of componentwise perturbation theory in numerical linear algebra. In *Mathematics of Computation 1943–1993: a half-century of computational mathematics (Vancouver, BC, 1993)*, volume 48 of *Proc. Sympos. Appl. Math.*, pages 49–77. Amer. Math. Soc., Providence, RI, 1994.
- [38] R. A. Horn and C. R. Johnson. *Matrix analysis*. Cambridge University Press, Cambridge, second edition, 2013.
- [39] I. C. F. Ipsen and R. Rehman. Perturbation bounds for determinants and characteristic polynomials. *SIAM J. Matrix Anal. Appl.*, 30(2):762–776, 2008.
- [40] I. T. Jolliffe. *Principal component analysis*. Springer Series in Statistics. Springer-Verlag, New York, 1986.
- [41] R. Kannan and S. Vempala. Spectral algorithms. *Found. Trends Theor. Comput. Sci.*, 4(3-4):front matter, 157–288 (2009), 2008.
- [42] A. Knowles and J. Yin. The isotropic semicircle law and deformation of Wigner matrices. *Comm. Pure Appl. Math.*, 66(11):1663–1750, 2013.
- [43] A. Knowles and J. Yin. Anisotropic local laws for random matrices. *Probab. Theory Related Fields*, 169(1-2):257–352, 2017.

- [44] V. Koltchinskii and D. Xia. Perturbation of linear forms of singular vectors under Gaussian noise. In *High dimensional probability VII*, volume 71 of *Progr. Probab.*, pages 397–423. Springer, [Cham], 2016.
- [45] J. O. Lee and K. Schnelli. Local law and Tracy-Widom limit for sparse random matrices. *Probab. Theory Related Fields*, 171(1-2):543–616, 2018.
- [46] J. O. Lee, K. Schnelli, B. Stetler, and H.-T. Yau. Bulk universality for deformed Wigner matrices. *Ann. Probab.*, 44(3):2349–2425, 2016.
- [47] M. Lelarge, L. Massoulié, and J. Xu. Reconstruction in the labelled stochastic block model. *IEEE Trans. Network Sci. Eng.*, 2(4):152–163, 2015.
- [48] F. McSherry. Spectral partitioning of random graphs. In *42nd IEEE Symposium on Foundations of Computer Science (Las Vegas, NV, 2001)*, pages 529–537. IEEE Computer Soc., Los Alamitos, CA, 2001.
- [49] B. Nadler. Finite sample approximation results for principal component analysis: a matrix perturbation approach. *Ann. Statist.*, 36(6):2791–2817, 2008.
- [50] S. O’Rourke, V. Vu, and K. Wang. Eigenvectors of random matrices: A survey. *Journal of Combinatorial Theory, Series A*, 144:361 – 442, 2016. Fifty Years of the Journal of Combinatorial Theory.
- [51] S. O’Rourke, V. Vu, and K. Wang. Random perturbation of low rank matrices: improving classical bounds. *Linear Algebra Appl.*, 540:26–59, 2018.
- [52] S. O’Rourke and N. Williams. Pairing between zeros and critical points of random polynomials with independent roots. *Trans. Amer. Math. Soc.*, 371(4):2343–2381, 2019.
- [53] C. Pozrikidis. *An introduction to grids, graphs, and networks*. Oxford University Press, Oxford, 2014.
- [54] J. W. S. Rayleigh, Baron. *The Theory of Sound*. Dover Publications, New York, N. Y., 1945. 2d ed.
- [55] K. Rohe, S. Chatterjee, and B. Yu. Spectral clustering and the high-dimensional stochastic blockmodel. *Ann. Statist.*, 39(4):1878–1915, 2011.
- [56] M. Rudelson and R. Vershynin. Non-asymptotic theory of random matrices: extreme singular values. In *Proceedings of the International Congress of Mathematicians. Volume III*, pages 1576–1602. Hindustan Book Agency, New Delhi, 2010.
- [57] P. Sarkar and P. J. Bickel. Role of normalization in spectral clustering for stochastic block-models. *Ann. Statist.*, 43(3):962–990, 2015.
- [58] E. Schrödinger. Quantisierung als eigenwertproblem. *Annalen der Physik*, 384(4):361–376, 1926.
- [59] G. W. Stewart and J. G. Sun. *Matrix perturbation theory*. Computer Science and Scientific Computing. Academic Press, Inc., Boston, MA, 1990.
- [60] M. Tang, A. Athreya, D. L. Sussman, V. Lyzinski, Y. Park, and C. E. Priebe. A semiparametric two-sample hypothesis testing problem for random graphs. *J. Comput. Graph. Statist.*, 26(2):344–354, 2017.
- [61] M. Tang, J. Cape, and C. E. Priebe. Asymptotically efficient estimators for stochastic block-models: the naive MLE, the rank-constrained MLE, and the spectral estimator. *Bernoulli*, 28(2):1049–1073, 2022.
- [62] T. Tao. Outliers in the spectrum of iid matrices with bounded rank perturbations. *Probab. Theory Related Fields*, 155(1-2):231–263, 2013.
- [63] C. Tomasi and T. Kanade. Shape and motion from image streams: a factorization method. *Proceedings of the National Academy of Sciences*, 90(21):9795–9802, 1993.
- [64] R. Vershynin. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge university press, 2018.
- [65] U. von Luxburg. A tutorial on spectral clustering. *Stat. Comput.*, 17(4):395–416, 2007.
- [66] V. Vu. Singular vectors under random perturbation. *Random Structures Algorithms*, 39(4):526–538, 2011.
- [67] V. Vu. A simple SVD algorithm for finding hidden partitions. *Combin. Probab. Comput.*, 27(1):124–140, 2018.
- [68] M. J. Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press, 2019.
- [69] R. Wang. Singular vector perturbation under Gaussian noise. *SIAM J. Matrix Anal. Appl.*, 36(1):158–177, 2015.

- [70] P.-A. Wedin. Perturbation bounds in connection with singular value decomposition. *Nordisk Tidskr. Informationsbehandling (BIT)*, 12:99–111, 1972.
- [71] D. Xia and F. Zhou. The sup-norm perturbation of HOSVD and low rank tensor denoising. *J. Mach. Learn. Res.*, 20:Paper No. 61, 42, 2019.
- [72] Y. Yu, T. Wang, and R. J. Samworth. A useful variant of the Davis-Kahan theorem for statisticians. *Biometrika*, 102(2):315–323, 2015.
- [73] S.-Y. Yun and A. Proutiere. Accurate community detection in the stochastic block model via spectral algorithms. Available at arXiv:1412.7335, 2014.
- [74] S.-Y. Yun and A. Proutiere. Optimal cluster recovery in the labeled stochastic block model. In D. Lee, M. Sugiyama, U. Luxburg, I. Guyon, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 29. Curran Associates, Inc., 2016.
- [75] X. Zhong, C. Su, and Z. Fan. Empirical Bayes PCA in high dimensions. *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 84(3):853–878, 2022.
- [76] Y. Zhong. Eigenvector under random perturbation: A nonasymptotic rayleigh-schrödinger theory. Available at arXiv:1702.00139, 2017.
- [77] Y. Zhong and N. Boumal. Near-optimal bounds for phase synchronization. *SIAM J. Optim.*, 28(2):989–1016, 2018.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COLORADO AT BOULDER, BOULDER, CO 80309
Email address: sean.d.orourke@colorado.edu

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, PO BOX 208283, NEW HAVEN, CT 06520-8283, USA
Email address: van.vu@yale.edu

DEPARTMENT OF MATHEMATICS, HONG KONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, HONG KONG
Email address: kewang@ust.hk