A QUANTITATIVE IMPROVEMENT FOR ROTH'S THEOREM ON ARITHMETIC PROGRESSIONS

THOMAS F. BLOOM

ABSTRACT. We improve the quantitative estimate for Roth's theorem on threeterm arithmetic progressions, showing that if $A \subset \{1, \ldots, N\}$ contains no nontrivial three-term arithmetic progressions then $|A| \ll N(\log \log N)^4 / \log N$. By the same method we also improve the bounds in the analogous problem over $\mathbb{F}_q[t]$ and for the problem of finding long arithmetic progressions in a sumset.

1. INTRODUCTION

In this paper we prove the following quantitative improvement for Roth's theorem on arithmetic progressions.

Theorem 1.1. If $A \subset \{1, ..., N\}$ contains no non-trivial three-term arithmetic progressions then

$$|A| \ll \frac{(\log \log N)^4}{\log N} N.$$

This problem has a long history, the first significant quantitative bound being given by Roth [1953]. Let R(N) denote the size of the largest subset of $\{1, \ldots, N\}$ which contains no non-trivial three-term arithmetic progressions. For comparison the table below summarises the history of upper bounds for R(N).

Roth [1953]	$N/\log\log N$
Szemerédi [1990] and Heath-Brown [1987]	$N/(\log N)^c$ for some $c > 0$
Bourgain [1999]	$(\log \log N)^{1/2} N / (\log N)^{1/2}$
Bourgain [2008]	$(\log \log N)^2 N / (\log N)^{2/3}$
Sanders [2012]	$N/(\log N)^{3/4-o(1)}$
Sanders [2011]	$(\log \log N)^6 N / \log N$

The claimed bound of $(\log \log N)^5 N / \log N$ in Sanders [2011] is due to a calculation error, and the method there in fact delivers $(\log \log N)^6 N / \log N$ as in the table above. The best known lower bound has the shape $R(N) \gg N \exp(-c\sqrt{\log N})$ for some absolute constant c > 0, due to Behrend [1946], and so Theorem 1.1 still leaves much to be desired.

Not only does the method of this paper deliver a quantitative improvement but it manages to do so without the powerful combinatorial tools used in Sanders [2011], and instead operates almost entirely in 'frequency space', exploiting a new lemma concerning structural properties of the large Fourier spectrum inspired by a recent breakthrough in Roth's theorem in \mathbb{F}_p^n by Bateman and Katz [2012]. In

The author was supported by an EPSRC doctoral training grant.

particular, we hope that the methods in this paper could be used along with more combinatorial techniques to yield further quantitative progress.

The significant new ingredient in the proof of Theorem 1.1 is a new lemma about the structural properties of the set of large Fourier coefficients of a given set, which under certain circumstances offers a quantitative improvement over a related wellknown lemma of Chang [2002]. As a further demonstration of the utility of this lemma we outline how it can be combined with the technique of Sanders [2008] to prove the following quantitative improvement to the problem of finding long arithmetic progressions in a sumset.

Theorem 1.2. If $A, B \subset \{1, ..., N\}$ with both $c_1 \alpha N \leq |A| \leq |B| \leq c_2 \alpha N$ then A + B contains an arithmetic progression of length at least

$$\exp\left(cf(\alpha)\sqrt{\log N}\right),$$

where $f(\alpha) = \alpha^{1/2} / \log(1/\alpha)$ and the constant c > 0 depends only on c_1 and c_2 .

By contrast, Sanders [2008] used Chang's lemma to prove a similar result with $f(\alpha) = \alpha$, a result which was first proved using a different method by Green [2002]. This was subsequently improved by Croot, Laba, and Sisask [2013] to $f(\alpha) = \alpha^{1/2} (\log(1/\alpha))^{-3/2}$. The proof of Theorem 1.2 can be easily adapted to give a similar bound when the sizes of A and B are not comparable but in this situation the method of Croot, Laba, and Sisask [2013] delivers superior bounds.

We shall present our method in a general setting, which will allow us to prove more general versions of Theorem 1.1 in both the integers and their finite-characteristic analogue $\mathbb{F}_q[t]$. In general, we will be concerned with counting non-trivial solutions to equations of the shape

(1)
$$c_1 x_1 + c_2 x_2 + c_3 x_3 = 0,$$

where $c_1 + c_2 + c_3 = 0$. A trivial solution is one where $x_1 = x_2 = x_3$. Applied to any such equation our method yields the following result over the integers.

Theorem 1.3. If $c_1, c_2, c_3 \in \mathbb{Z} \setminus \{0\}$ are such that $c_1 + c_2 + c_3 = 0$ and $A \subset \{1, \ldots, N\}$ contains no non-trivial solutions to (1) then

$$|A| \ll_{\mathbf{c}} \frac{(\log \log N)^4}{\log N} N.$$

In particular, Theorem 1.1 follows by considering the coefficients $\mathbf{c} = (1, 1, -2)$. The polynomial ring $\mathbb{F}_q[t]$ is, in many respects, a finite-characteristic analogue of the integers, so we should expect a result of similar strength to hold. In Bloom [2012] the method of Sanders [2011] was adapted to give a quantitative result for $\mathbb{F}_q[t]$. In this paper we shall similarly prove an analogue of Theorem 1.3 for $\mathbb{F}_q[t]$. As in Bloom [2012] the finite-characteristic property leads to a slight improvement in this case (due to the preponderance of subgroups).

Theorem 1.4. Let $A \subset \mathbb{F}_q[t]_{\deg \leq n}$. If $c_1, c_2, c_3 \in \mathbb{F}_q[t] \setminus \{0\}$ are such that $c_1 + c_2 + c_3 = 0$ and A contains no non-trivial solutions to (1) then

$$|A| \ll_{\mathbf{c}} \frac{(\log n)^2}{n} q^n.$$

We remark that if $c_1, c_2, c_3 \in \mathbb{F}_q \setminus \{0\}$ then the problem is much simpler and better bounds are available. In particular, Liu and Spencer [2009] proved that in this case we have the upper bound $|A| \ll_{\mathbf{c}} q^n/n$; when $c_1, c_2, c_3 \in \mathbb{F}_p \setminus \{0\}$ this bound was first provided by Meshulam [1995]. It is also likely that the work of Bateman and Katz [2012] in \mathbb{F}_p^n could be adapted to deliver a bound of $|A| \ll_{\mathbf{c}} q^n/n^{1+\epsilon}$ for some absolute constant $\epsilon > 0$, when $c_1, c_2, c_3 \in \mathbb{F}_q \setminus \{0\}$.

The core of our argument is a qualitatively stronger alternative to a well-known lemma of Chang [2002] concerning the additive structure of the large spectrum of the Fourier transform in finite groups. More particularly, if G is a finite abelian group and $A \subset G$ with density $\alpha = |A| / |G|$ then we define the large spectrum $\Delta_{\eta}(A)$ as the set of characters $\gamma \in \widehat{G}$ such that $|\widehat{A}(\gamma)| \geq \eta |A|$. It follows immediately from Parseval's theorem that $|\Delta_{\eta}(A)| \leq \eta^{-2} \alpha^{-1}$, which is the best possible bound on the cardinality. Chang's lemma shows that a smaller set can be found which additively 'controls' the spectrum. In particular, we say that Δ is *d*-covered if there exists Λ of size $|\Lambda| \leq d$ such that

$$\Delta \subset \left\{ \sum_{\lambda \in \Lambda} \epsilon_{\lambda} \lambda : \epsilon_{\lambda} \in \{-1, 0, 1\} \right\}.$$

In this language, Chang's lemma may be stated as follows.

Theorem 1.5 (Chang [2002]). If $A \subset G$ with density $\alpha = |A| / |G|$ then $\Delta_{\eta}(A)$ is *d*-covered for some

$$d \ll \eta^{-2} \log(1/\alpha).$$

This important structural lemma has found many applications since, and in particular has been instrumental in the recent advances in Roth's theorem. By contrast, in this paper our method does not directly use Chang's lemma, but rather the following theorem which is, for some applications, much stronger.

Theorem 1.6. If $A \subset G$ with density $\alpha = |A| / |G|$ then there exists $\Delta' \subset \Delta_{\eta}(A)$ of size $|\Delta'| \gg \eta |\Delta_{\eta}(A)|$ which is d-covered for some

$$d \ll \eta^{-1} \log(1/\alpha).$$

In particular, we can save a factor of η in the dimension while only losing a factor of η on the size of the set considered. We prove Theorem 1.6, or rather, a more general version, by considering the additive energy of large spectra, building on work by Shkredov [2008] and Bateman and Katz [2012].

For our applications we shall discuss the ideas leading to Theorem 1.6 in a general setting that also applies to covering 'relative' to a given set, which is needed for the density increment strategy used for Roth's theorem.

2. NOTATION AND DEFINITIONS

We fix some finite abelian group G with dual group \widehat{G} . Let N = |G|. For convenience we shall use the counting measure on both G and \widehat{G} . In particular all L^p norms on both G and \widehat{G} are defined with respect to the counting measure, so that if $f: G \to \mathbb{C}$ and $p \ge 1$ then

$$||f||_p = \left(\sum_x |f(x)|^p\right)^{1/p}$$
 and $||f||_{\infty} = \sup_{x \in G} |f(x)|$,

and similarly for $\omega : \widehat{G} \to \mathbb{C}$. For any function $f : G \to \mathbb{C}$ we define the Fourier transform $\widehat{f} : \widehat{G} \to \mathbb{C}$ by

$$\widehat{f}(\gamma) = \sum_{x} f(x) \gamma(x).$$

For all functions $f, g: G \to \mathbb{C}$ we have Parseval's identity

$$\langle f,g \rangle = \sum_{x} f(x)\overline{g(x)} = N^{-1} \sum_{\gamma} \widehat{f}(\gamma)\overline{\widehat{g}(\gamma)}.$$

If $B \subset G$ and $\Gamma \subset \widehat{G}$ then for any $\epsilon \in [0, 2]$ we say that B has ϵ -control of Γ if for all $x \in B$ and $\gamma \in \Gamma$ we have

$$|1 - \gamma(x)| \le \epsilon.$$

This condition will be important in several places; in general, the hypothesis of control presents no serious difficulties as we will just define the sets we are working with precisely so that they have the required control. The details will vary on the choice of the group G, and hence for most of the paper we simply present the necessary control hypothesis and take care of how to ensure it in our applications in the final sections of the paper.

For any $\eta \in [0,1]$ and $f: G \to \mathbb{C}$ we define the spectrum

$$\Delta_{\eta}(f) = \left\{ \gamma \in \widehat{G} : \left| \widehat{f}(\gamma) \right| \ge \eta \left\| f \right\|_{1} \right\}$$

and the level spectrum

$$\tilde{\Delta}_{\eta}(f) = \left\{ \gamma \in \widehat{G} : \eta \left\| f \right\|_{1} \le \left| \widehat{f}(\gamma) \right| < 2\eta \left\| f \right\|_{1} \right\}.$$

For $0 < \delta \leq 1$ we shall use the convenient shorthand $\mathcal{L}(\delta)$ to denote $2 + \lceil \log(1/\delta) \rceil$. Finally, it will often be convenient for certain sets to renormalise the counting measure to be compact; these sets will always be denoted by B (possibly with some subscripts or superscripts) and then for any $A \subset G$ we define $\beta(A) = |A \cap B| / |B|$. In general, if we speak of $A \subset B$ having relative density α then this means $\beta(A) = \alpha$.

We will frequently abuse notation by conflating a set and its characteristic function; thus, for example, if $\Gamma \subset \widehat{G}$ then $\Gamma(\gamma) = 1$ if $\gamma \in \Gamma$ and 0 otherwise.

For any $\Gamma \subset \widehat{G}$ and $\omega : \widehat{G} \to \mathbb{R}_+$ and integer $m \ge 1$ we define the additive energy as

$$E_{2m}(\omega,\Gamma) = \sum_{\gamma_1,\dots,\gamma'_m} \omega(\gamma_1)\cdots\omega(\gamma'_m)\Gamma\left(\sum_{i=1}^m \gamma_i - \sum_{j=1}^m \gamma'_j\right).$$

Similarly, we define the restricted energy as

$$E_{t_1,t_2}^{\sharp}(\omega,\Gamma) = \sum_{\substack{\Delta_1 \in \binom{\hat{G}}{t_1}, \Delta_2 \in \binom{\hat{G}}{t_2} \\ \Delta_1 \cap \Delta_2 = \emptyset}} \prod_{\gamma \in \Delta_1 \cup \Delta_2} \omega(\gamma) \Gamma\left(\sum_{\gamma \in \Delta_1} \gamma - \sum_{\gamma' \in \Delta_2} \gamma'\right).$$

We write E_{2m}^{\sharp} for $E_{m,m}^{\sharp}$ and for any ω and Γ we define $E_0(\omega, \Gamma) = E_0^{\sharp}(\omega, \Gamma) = 1$. Observe that E and E^{\sharp} differ, not only in the restriction on repeating elements, but also in that the former is sensitive to permutations of the γ_i . We say that S is d-covered by Γ if there exists $\Lambda \subset \widehat{G}$ of size $|\Lambda| \leq d$ such that

$$S \subset \Gamma - \Gamma + \langle \Lambda \rangle_{\mathcal{S}}$$

where

$$\langle \Lambda \rangle = \left\{ \sum_{\lambda \in \Lambda} \epsilon_{\lambda} \lambda : \epsilon_{\lambda} \in \{-1, 0, 1\} \right\}$$

and $\langle \emptyset \rangle = \{0\}$.

We say that Δ is Γ -dissociated if for all $k \geq 1$ and $\lambda \in \widehat{G}$ there are at most 2^k many pairs Δ_1, Δ_2 of disjoint subsets of Δ such that $|\Delta_1 \cup \Delta_2| = k$ and

$$\sum_{\gamma \in \Delta_1} \gamma - \sum_{\gamma' \in \Delta_2} \gamma' \in \Gamma + \lambda.$$

Finally, we say that S has Γ -dimension of d if d is the size of the largest Γ -dissociated subset of S. We observe that the dimension is always at least 1 since any singleton set is trivially Γ -dissociated for all $\Gamma \subset \widehat{G}$. Furthermore, if Δ is Γ -dissociated then it is also Γ' -dissociated for any translate Γ' of Γ .

3. Additive energy

In this section we discuss the relationship between the dimension of a set and its additive energy. If a set has a very large dimension then almost all of the set is dissociated, and hence one would expect few additive relations between its elements, so it should have small additive energy. The following lemma verifies this intuition.

Lemma 3.1. If $S \subset \widehat{G}$ has Γ -dimension |S| - k then for all $m \geq t_1, t_2 \geq 0$

1

$$E^{\sharp}_{t_1, t_2}(S, \Gamma) \le 4^{k+m}.$$

Proof. Let $S = S_0 \sqcup S_1$ where S_0 is Γ -dissociated and $|S_1| = k$. By separating the contribution from the subsets of S_1 we obtain the estimate

$$E_{t_1,t_2}^{\sharp}(S,\Gamma) = \sum_{\substack{\Delta_1 \in \binom{S}{t_1}, \Delta_2 \in \binom{S}{t_2} \\ \Delta_1 \cap \Delta_2 = \emptyset}} \Gamma\left(\sum_{\gamma \in \Delta_1} \gamma - \sum_{\gamma' \in \Delta_2} \gamma'\right)$$
$$\leq \sum_{\substack{0 \le r_1 \le t_1 \\ 0 \le r_2 \le t_2}} \binom{k}{r_1} \binom{k}{r_2} \sup_{\substack{\lambda \\ \lambda_1 \in \binom{S_0}{t_1 - r_1}, \Delta_2 \in \binom{S_0}{t_2 - r_2}}} \Gamma\left(\sum_{\gamma \in \Delta_1} \gamma - \sum_{\gamma' \in \Delta_2} \gamma' + \lambda\right)$$

Since S_0 is Γ -dissociated, however, the inner summand is bounded above by $2^{t_1+t_2}$ and the lemma follows.

For the main result of this section we need to convert a conclusion about the restricted additive energy to the full additive energy, for which the following lemma will suffice.

Lemma 3.2. For any $\Gamma \subset \widehat{G}$, weight function $\omega : \widehat{G} \to \mathbb{R}_+$ and integer $m \geq 2$ we have

$$E_{2m}(\omega,\Gamma) \le 2^{4m} (m!)^2 \|\omega\|_2^{2m} \sum_{0 \le t_1, t_2 \le m} \frac{\|\omega\|_2^{-t_1-t_2}}{((m-t_1)!(m-t_2)!)^{1/2}} \sup_{\lambda} E_{t_1, t_2}^{\sharp}(\omega, \Gamma+\lambda).$$

Proof. We divide the range of summation of E_{2m} according to the size of the subsets of $\{\gamma_1, \ldots, \gamma_m\}$ and $\{\gamma'_1, \ldots, \gamma'_m\}$ consisting of elements that each occur with multiplicity 1. This leads to the upper bound

(2)
$$E_{2m}(\omega,\Gamma) \leq \sum_{0 \leq l_1, l_2 \leq m} G_{m-l_1}(\omega) G_{m-l_2}(\omega) \binom{m}{l_1} \binom{m}{l_2} \sup_{\lambda} F_{\lambda}(l_1, l_2)$$

where

$$F_{\lambda}(l_1, l_2) = \sum_{\substack{\gamma_1, \dots, \gamma'_{l_2} \\ \gamma_i \neq \gamma_j \ \gamma'_i \neq \gamma'_j \ i \neq j}} \omega(\gamma_1) \cdots \omega(\gamma'_{l_2}) \Gamma\left(\sum_{i=1}^{l_1} \gamma_i - \sum_{j=1}^{l_2} \gamma'_j - \lambda\right)$$

and

(3)
$$G_k(\omega) = \sum_{\Delta}^* \prod_{\gamma \in \Delta} \omega(\gamma) \le \frac{k!}{(\lfloor k/2 \rfloor)!} \left(\sum_{\gamma} \omega(\gamma)^2 \right)^{k/2},$$

the first sum being restricted to those ordered k-tuples $\Delta \in \widehat{G}^k$ where each element occurs with multiplicity at least 2. The sum $F_{\lambda}(l_1, l_2)$ is almost a renormalised version of the restricted energy E_{l_1, l_2}^{\sharp} except that it lacks the restriction $\gamma_i \neq \gamma'_j$ for all $1 \leq i \leq l_1$ and $1 \leq j \leq l_2$. To introduce this we partition $F_{\lambda}(l_1, l_2)$ according to the number of common elements between the γ_i and γ'_i ; thus

(4)
$$F_{\lambda}(l_1, l_2) \leq \sum_{i=0}^{\min(l_1, l_2)} {l_1 \choose i} {l_2 \choose i} i! \|\omega\|_2^{2i} (l_1 - i)! (l_2 - i)! E_{l_1 - i, l_2 - i}^{\sharp} (\omega, \Gamma + \lambda).$$

Combining (2), (3) and (4) and simplifying the expression implies that $E_{2m}(\omega, \Gamma)$ is at most

$$(m!)^{2} \|\omega\|_{2}^{2m} \sum_{0 \le l_{1}, l_{2} \le m} \sum_{i=0}^{\min(l_{1}, l_{2})} \frac{\|\omega\|_{2}^{2i-l_{1}-l_{2}}}{i!(\lfloor (m-l_{1})/2 \rfloor)!(\lfloor (m-l_{2})/2 \rfloor)!} \sup_{\lambda} E_{l_{1}-i, l_{2}-i}^{\sharp}(\omega, \Gamma+\lambda)$$

Relabelling $t_1 = l_1 - i$ and $t_2 = l_2 - i$ this is at most

$$(m!)^2 \|\omega\|_2^{2m} \sum_{0 \le t_1, t_2 \le m} \|\omega\|_2^{-t_1 - t_2} \sup_{\lambda} E_{t_1, t_2}^{\sharp}(\omega, \Gamma + \lambda) f(m, t_1, t_2)$$

where

$$f(m, t_1, t_2) = \sum_{i \ge \max(t_1, t_2)}^{m - \max(t_1, t_2)} \frac{1}{i! (\lfloor (m - t_1 - i)/2 \rfloor)! (\lfloor (m - t_2 - i)/2 \rfloor)!}$$

Finally, a tedious calculation using the elementary inequality $n!/(\lfloor n/2 \rfloor!)^2 \leq 2(n+1)^{1/2}2^n$, valid for all $n \geq 0$, shows that the inner sum is at most $2^{4m}((m-t_1)!(m-t_2)!)^{-1/2}$ and the lemma follows.

The final technical lemma of this section provides a relationship between covering and dimension that will be important in the proof of Theorem 3.1.

Lemma 3.3. Let $\Gamma \subset \widehat{G}$ be a symmetric set. If $\Delta \subset \widehat{G}$ has Γ -dimension r then there is a partition $\widehat{G} = \Lambda_0 \sqcup \Lambda_1$ where Λ_0 is 2r-covered by Γ and for all $\gamma \in \Lambda_1$ the set $\Delta \cup \{\gamma\}$ has Γ -dimension r + 1. *Proof.* By hypothesis we can decompose Δ as $\Delta_0 \sqcup \Delta_1$ where Δ_0 is Γ -dissociated and $|\Delta_0| = r$. Let Δ' be the set of all $\gamma \in \widehat{G}$ such that $\Delta_0 \cup \{\gamma\}$ is not Γ -dissociated. We claim that if we let $\Lambda_0 = \Delta' \cup \Delta_0$ and $\Lambda_1 = \widehat{G} \setminus \Lambda_0$ then this is a suitable decomposition.

Firstly, let $\gamma \in \Lambda_1$. By construction the set $\Delta_0 \cup \{\gamma\}$ is Γ -dissociated, and hence $\Delta \cup \{\gamma\}$ has Γ -dimension at least r + 1 by definition, since $|\Delta_0 \cup \{\gamma\}| = r + 1$. It remains to show that Λ_0 is 2*r*-covered by Γ ; for this, it suffices to show that

$$\Delta_0 \cup \Delta' \subset \Gamma - \Gamma + \langle \Delta_0 \rangle + \langle \Delta_0 \rangle.$$

This is obvious for Δ_0 . Let $\gamma \in \Delta'$. By construction $\Delta_0 \cup \{\gamma\}$ is not Γ -dissociated, and hence there exists $k \geq 1$ and $\lambda \in \widehat{G}$ such that there are more than 2^k many triples $(\epsilon, \Delta'_1, \Delta'_2)$ such that $\epsilon \in \{-1, 0, 1\}$, the sets Δ'_1 and Δ'_2 are disjoint subsets of Δ_0 with $|\Delta'_1 \cup \Delta'_2| + |\epsilon| = k$, and

$$\epsilon \gamma + \sum_{\gamma_1' \in \Delta_1'} \gamma_1' - \sum_{\gamma_2' \in \Delta_2'} \gamma_2' \in \Gamma + \lambda.$$

If there exists at least one such triple with $\epsilon = 0$ and at least one with $\epsilon \neq 0$ then it is easy to check that this implies that $\gamma \in \Gamma - \Gamma + \langle \Delta_0 \rangle - \langle \Delta_0 \rangle$ as required. If $\epsilon \equiv 0$ for all such triples then this contradicts the Γ -dissociativity of Δ_0 . Hence we can assume that $\epsilon \in \{-1, 1\}$ for all such triples; this is clearly impossible for k = 1, and for k > 1 we observe that by the pigeonhole principle there are strictly more than 2^{k-1} such triples with identical ϵ . This, however, is another contradiction to the Γ dissociativity of Δ_0 , considering the translate $\Gamma + \lambda - \epsilon \gamma$. Thus $\gamma \in \Gamma - \Gamma + \langle \Delta_0 \rangle - \langle \Delta_0 \rangle$ as required, and the proof is complete. \Box

The following theorem is crucial, and uses random sampling to prove a hereditary version of our earlier intuition: namely, if a set is such that every large subset is not efficiently covered then we must have particularly small additive energy. The argument is a variant on that used in [Bateman and Katz, 2012, Section 5]. There, however, they only wish to bound the 8-fold additive energy, whereas for our purposes we shall need to deal with the 2m-fold additive energy where $m \to \infty$ as $N \to \infty$ (for our applications we shall in fact take $m \approx (\log \log N)^{1+o(1)}$), and hence we have taken care to make the dependence on m explicit.

We treat the constants in this argument, as in the rest of this paper, quite crudely; it is certainly possible to improve them, but such improvements would have a negligible effect on the main results.

Theorem 3.1. Let $\Gamma \subset \widehat{G}$ be a symmetric set and $\omega : \widehat{G} \to \mathbb{R}_+$. Let $m \geq 2$ and $d \geq n \geq 2$ be such that $m \leq d/4$ and $\|\omega\|_2 \leq m^{1/2}d^{-1} \|\omega\|_1$. Then either there is a finite set $\Delta \subset \widehat{G}$ such that

$$\sum_{\gamma \in \Delta} \omega(\gamma) \geq \frac{n}{d} \, \|\omega\|_1$$

and Δ is 2*d*-covered by Γ , or

$$E_{2m}(\omega,\Gamma) \le 2^{13m+6n} m^{2m} d^{-2m} \|\omega\|_1^{2m}.$$

Proof. Without loss of generality we may suppose that $\|\omega\|_1 = 1$. We first observe that either we are in the first case, or every subset $\Delta \subset \widehat{G}$ which is 2*d*-covered by Γ satisfies $\sum_{\gamma \in \Delta} \omega(\gamma) \leq nd^{-1}$, which we shall assume henceforth.

Let $S \subset \widehat{G}$ be a random set of size at most d chosen by selecting d elements of \widehat{G} at random, where we choose $\gamma \in \widehat{G}$ with probability $\omega(\gamma)$. We claim that for $k \geq 0$ the set S has Γ -dimension d - k with probability at most $n^k/k!$.

For suppose we have selected $d' \leq d$ elements of S, say S', and suppose that S' has Γ -dimension r. By Lemma 3.3 we can partition $\widehat{G} = \Lambda_0 \sqcup \Lambda_1$ such that Λ_0 is 2*d*-covered by Γ and for all $\gamma \in \Lambda_1$ the set $S' \cup \{\gamma\}$ has Γ -dimension at least r + 1. Thus, in our model,

$$\mathbb{P}(\dim(S' \cup \{\gamma\}) \le \dim(S')) \le \sum_{\gamma \in \Lambda_0} \omega(\gamma) \le \frac{n}{d},$$

since Λ_0 is 2*d*-covered by Γ . From this estimate, combined with the trivial observations that the empty set has Γ -dimension 0 and that Γ -dimension is non-decreasing, it follows that the probability that S has Γ -dimension d - k is at most the probability that k events with probability at most n/d occur in d independent trials, which is at most

$$\binom{d}{k} n^k d^{-k} \le n^k / k!$$

as required. By Lemma 3.1 it follows that for all $\lambda \in \widehat{G}$ and integers $t_1, t_2 \leq m$ we have

$$\mathbb{E}E^{\sharp}_{t_1,t_2}(S,\Gamma+\lambda) \le 4^m \sum_{k=0}^{\infty} \frac{(4n)^k}{k!} = 4^m e^{4n}.$$

Let $1 \leq k \leq 2m$. For any distinct $\gamma_1, \ldots, \gamma_k \in \widehat{G}$ the probability that $\gamma_1, \ldots, \gamma_k \in S$ is at least

$$k! \binom{d}{k} \omega(\gamma_1) \cdots \omega(\gamma_k) \left(1 - \sum_{i=1}^k \omega(\gamma_i) \right)^{d-k}$$

Since $k \leq d/2$ we have $k!\binom{d}{k} \geq (d/2)^k$. Furthermore, by the Cauchy-Schwarz inequality

$$\sum_{i=1}^{k} \omega(\gamma_i) \le (2m)^{1/2} \|\omega\|_2 \le 2md^{-1} \le 1/2,$$

so that the second factor is at least

$$\exp\left(-d\sum_{i=1}^k \omega(\gamma_i)\right) \ge e^{-2m}.$$

It follows that the probability that $\gamma_1, \ldots, \gamma_k \in S$ is at least $2^{-5m} d^k \omega(\gamma_1) \cdots \omega(\gamma_k)$. By linearity of expectation, assuming $t_1 + t_2 \leq m$,

$$\mathbb{E}E^{\sharp}_{t_1,t_2}(S,\Gamma+\lambda) \ge 2^{-5m} d^{t_1+t_2} E^{\sharp}_{t_1,t_2}(\omega,\Gamma+\lambda),$$

and so, for all $\lambda \in \widehat{G}$ and $0 \leq t_1, t_2 \leq m$,

$$E_{t_1,t_2}^{\sharp}(\omega,\Gamma+\lambda) \le 2^{7m} e^{4n} d^{-t_1-t_2}.$$

From Lemma 3.2 it follows that

$$E_{2m}(\omega,\Gamma) \le 2^{11m} e^{4n} m! \|\omega\|_2^{2m} \left(\sum_{0 \le t \le m} \frac{(m!)^{1/2} (\|\omega\|_2^{-1} d^{-1})^t}{((m-t)!)^{1/2}} \right)^2.$$

For brevity let $r = \|\omega\|_2^{-2} d^{-2} \ge m^{-1}$. By the Cauchy-Schwarz inequality the inner factor is at most

$$(m+1)\sum_{0\le t\le m}\frac{m!}{(m-t)!}r^t\le (m+1)\sum_{0\le t\le m}(erm)^t\le (m+1)^2(erm)^m,$$

say. In particular

$$E_{2m}(\omega,\Gamma) \le 2^{13m} e^{4n} m^{2m} \|\omega\|_2^{2m} r^m = 2^{13m} e^{4n} m^{2m} d^{-2m},$$

and the lemma follows.

4. STRUCTURE IN SPECTRA

The results in the previous section are extremely general, and can be used to deduce facts about the dimensions of arbitrary sets from lower bounds on their additive energy. We shall use them to derive a structural result about the large spectrum of a function $f: G \to \mathbb{C}$, and for this we require a lower bound on the additive energy of such spectra. A suitable lower bound was provided by Shkredov [2008], who showed that if $A \subset G$ with density $\alpha = |A|/N$ and $\Delta \subset \Delta_{\eta}(A)$ then $E_{2m}(\Delta, \{0\}) \gg \eta^{2m} \alpha |\Delta|^{2m}$. A simpler proof of this fact was given in Shkredov [2009]; the following is a simple generalisation of this proof.

Lemma 4.1. Let $\epsilon \in [0,1]$. For any $B \subset G$ and $\eta \in [0,1]$ let $f : B \to \mathbb{C}$ and $\omega : \widehat{G} \to \mathbb{R}_+$ be supported on \widehat{G} . Then for all integers $m \ge 1$

$$E_{2m}(\omega, \Delta_{\epsilon}(B)) \ge \|\omega\|_{1}^{2m} \left(\left(\eta \frac{\|f\|_{1}}{\|f\|_{2m/(2m-1)} |B|^{1/2m}} \right)^{2m} - \epsilon \right).$$

To recover the bound of Shkredov we set B = G and let $\epsilon \to 0$.

Proof. Let χ be defined by letting $\widehat{\chi}(\gamma) = \overline{c_{\gamma}}\omega(\gamma)$, where $c_{\gamma}\widehat{f}(\gamma) = |\widehat{f}(\gamma)|$. By construction whenever $\omega(\gamma) \neq 0$ we have $|\widehat{f}(\gamma)| \geq \eta ||f||_1$, whence

$$\sum_{x} f(x)\overline{\chi(x)} = N^{-1} \sum_{\gamma} \widehat{f}(\gamma)\overline{\widehat{\chi}(\gamma)} = N^{-1} \sum_{\gamma} \omega(\gamma) \left| \widehat{f}(\gamma) \right| \ge N^{-1} \eta \left\| f \right\|_{1} \left\| \omega \right\|_{1}.$$

By Hölder's inequality, however,

$$\left(\sum_{x} f(x)\overline{\chi(x)}\right)^{2m} \le \left(\sum_{x} |f(x)|^{2m/(2m-1)}\right)^{2m-1} \left(\sum_{x} B(x) |\chi(x)|^{2m}\right).$$
mains to note that, by the triangle inequality

It remains to note that, by the triangle inequality,

$$\sum_{x} B(x) |\chi(x)|^{2m} = N^{-2m} \sum_{x} B(x) \left| \sum_{\gamma} c_{\gamma} \omega(\gamma) \gamma(x) \right|^{2m}$$
$$\leq N^{-2m} \sum_{\gamma_{1}, \dots, \gamma'_{m}} \omega(\gamma_{1}) \cdots \omega(\gamma'_{m}) \left| \widehat{B}(\gamma_{1} + \dots + \gamma_{m} - \gamma'_{1} - \dots - \gamma'_{m}) \right|.$$

It follows that

$$\eta^{2m} \|f\|_{1}^{2m} \|\omega\|_{1}^{2m} \leq \|f\|_{2m/(2m-1)}^{2m} \sum_{\gamma_{1},\dots,\gamma'_{m}} \omega(\gamma_{1})\cdots\omega(\gamma'_{m}) \left| \widehat{B}(\gamma_{1}+\cdots-\gamma'_{m}) \right|$$
$$\leq \|f\|_{2m/(2m-1)}^{2m} \left(|B| E_{2m}(\omega,\Delta_{\epsilon}(B)) + \epsilon |B| \|\omega\|_{1}^{2m} \right),$$

and the proof is complete.

Finally, we can prove the technical heart of our argument, the aforementioned alternative to Chang's lemma. Again, for our application we need a fairly general statement, but at first glance the reader should take B = G and any $\epsilon > 0$, so that $\Delta_{\epsilon}(B) = \{0\}$.

Theorem 4.1. Suppose that $f : B \to \mathbb{C}$ and let $\alpha = \|f\|_1 / \|f\|_{\infty} |B|$. Let $\omega : \widehat{G} \to \mathbb{R}_+$ be supported on $\Delta_{\eta}(f)$ and let $0 \le \epsilon \le \exp(-8\mathcal{L}(\eta)\mathcal{L}(\alpha))$. There is a set $\Delta' \subset \Delta_{\eta}(f)$ such that

$$\sum_{\gamma \in \Delta'} \omega(\gamma) \ge 2^{-12} \eta \, \|\omega\|_1$$

and Δ' is $2^{14}\mathcal{L}(\alpha)\eta^{-1}$ -covered by $\Delta_{\epsilon}(B)$.

Proof. Without loss of generality we may suppose that $\|\omega\|_1 = 1$. Suppose first that $\|\omega\|_2 \ge 2^{-12} \mathcal{L}(\alpha)^{-1/2} \eta$ and let Δ' be a random set selected by including $\gamma \in \widehat{G}$ independently with probability $2^{13} \eta^{-1} \mathcal{L}(\alpha) \omega(\gamma)$. Then, if Δ' is this randomly chosen set we have, by Chernoff's inequality, that $|\Delta'| \le 2^{14} \eta^{-1} \mathcal{L}(\alpha)$ with probability at least 7/8, say, and

$$\mathbb{E}\sum_{\gamma\in\Delta'}\omega(\gamma)\geq 2^{13}\eta^{-1}\mathcal{L}(\alpha)\left\|\omega\right\|_{2}^{2}\geq 2^{-11}\eta,$$

and hence by Markov's inequality we have $\sum_{\gamma \in \Delta'} \omega(\gamma) \geq 2^{-12} \eta$ with probability at least 1/2, and the lemma follows.

Otherwise, we let $n = m = \mathcal{L}(\alpha)$ and $d = \lfloor 2^{12} \eta^{-1} m \rfloor$ and apply Lemmata 4.1 and 3.1. By the above we can suppose that $\|\omega\|_2 \leq 2^{-12} \mathcal{L}(\alpha)^{-1/2} \eta \leq m^{1/2} d^{-1}$, as is necessary for the application of Lemma 3.1. Lemma 4.1 implies that

$$E_{2m}(\omega; \Delta_{\epsilon}(B)) \ge \left(\eta \frac{\|f\|_{1}}{\|f\|_{2m/(2m-1)} |B|^{1/2m}}\right)^{2m} - \epsilon.$$

We have the trivial bound $||f||_{2m/(2m-1)} \leq ||f||_{\infty}^{1/2m} ||f||_{1}^{1-1/2m}$, and hence if $\epsilon \leq \eta^{2m} \alpha/2$ then

$$E_{2m}(\omega; \Delta_{\epsilon}(B)) \ge \eta^{2m} \alpha/2$$

By Lemma 3.1 either there is a set Δ' such that

$$\sum_{\gamma \in \Delta'} \omega(\gamma) \ge \frac{m}{d}$$

and Δ' is 2*d*-covered by $\Delta_{\epsilon}(B)$, or

$$\eta^{2m} \alpha \le 2^{19m+1} m^{2m} d^{-2m}.$$

In particular,

$$d \le 2^{10} m \eta^{-1} \alpha^{-1/2m},$$

which contradicts our initial choice of d and m, and the proof is complete.

Theorem 1.6 is a special case of this; to obtain it one simply sets B = G, lets $\epsilon \to 0$ and takes ω as the characteristic function of $\Delta_{\eta}(f)$.

5. The density increment

In this section we show how to use the structural result Theorem 4.1 to obtain the usual density increment lemma which can be iterated to yield the main theorems. The density increment strategy originated with Roth [1953] but has seen various technical simplifications since which we incorporate here. Roughly speaking, the idea is to show that if A does not contain the expected number of solutions to (1) then it has large Fourier coefficients, and this information can be translated into finding some group-like $B \subset G$ such that $|A \cap B| / |B|$ is larger than the expected |A|/N, and the argument can then be iterated until it halts due to the trivial bound $|A \cap B| \leq |B|$.

The first lemma is the standard conversion of L^2 -information on the balanced function of a set to a density increment, a technique first exploited for Roth's theorem by Heath-Brown [1987] and Szemerédi [1990].

Lemma 5.1. Let $f: B \to [0,1]$ and $\mathbf{f} = f - \alpha B$, where $\alpha = \|f\|_1 / |B|$. Suppose that

$$\sum_{\gamma \in \Gamma} \left| \widehat{\mathbf{f}}(\gamma) \right|^2 \ge \nu \alpha \left\| f \right\|_1 N.$$

Then if B' is a symmetric set such that for every $\gamma \in \Gamma$ we have

$$\left|\widehat{B'}(\gamma)\right| \ge 2^{-1} \left|B\right|,$$

and furthermore $|(2B'+B)\backslash B| \leq 2^{-4}\nu\alpha |B|$ then

$$\|f * B'\|_{\infty} \ge (1 + 2^{-3}\nu)\alpha |B'|.$$

Proof. By hypothesis we have

$$\sum_{\gamma} \left| \widehat{\mathbf{f}}(\gamma) \right|^2 \left| \widehat{B'}(\gamma) \right|^2 \ge 2^{-2} \nu \alpha \left\| f \right\|_1 \left| B' \right|^2 N.$$

In particular,

$$\|\mathbf{f} * B'\|_{2}^{2} = N^{-1} \sum_{\gamma} \left| \widehat{\mathbf{f}}(\gamma) \right|^{2} \left| \widehat{B'}(\gamma) \right|^{2} \ge 2^{-2} \nu \alpha \left\| f \right\|_{1} \left| B' \right|^{2}.$$

Expanding out the L^2 norm we obtain that

$$\sum_{x} |f * B'(x)|^{2} + \alpha^{2} ||B * B'||_{2}^{2} - 2\alpha \langle B * B', f * B' \rangle \ge 2^{-2} \nu \alpha ||f||_{1} |B'|^{2}.$$

By hypothesis

$$\begin{split} \left| \langle B * B', f * B' \rangle - \left\| f \right\|_1 \left| B' \right|^2 \right| &\leq \left| B' \right|^2 \sup_{x, y \in B'} \sum_{z \notin B} f(z + x - y) \\ &\leq \left| B' \right|^2 \left| (2B' + B) \backslash B \right| \\ &\leq 2^{-4} \nu \left\| f \right\|_1 \left| B' \right|^2. \end{split}$$

It follows that

$$\|f * B'\|_{2}^{2} \ge (1 + 2^{-3}\nu)\alpha \|f\|_{1} |B'|^{2}.$$

The left hand side is at most $\|f\|_1 \|B'\| \|f * B'\|_{\infty}$ and the lemma follows.

The second lemma shows that if a set has small dimension then control on only a few elements gives control on the whole set.

Lemma 5.2. If $\Delta \subset \widehat{G}$ is d-covered by Γ then there is $\Lambda \subset \widehat{G}$ of size at most d such that if B has $(4d)^{-1}$ -control of Λ and $\Gamma(1/8)$ then for every $\gamma \in \Delta$ we have

$$\left|\widehat{B}(\gamma)\right| \ge 2^{-1} \left|B\right|$$

Proof. By hypothesis there is some $\Lambda \subset \widehat{G}$ such that $|\Lambda| \leq d$ and

$$\Delta \subset \Gamma - \Gamma + \langle \Lambda \rangle.$$

Let $\gamma \in \Delta$, so that there exists some $\gamma_0, \gamma_1 \in \Gamma$ and $\epsilon \in \{-1, 0, 1\}^{\Lambda}$ such that

$$\gamma = \gamma_0 - \gamma_1 + \sum_{\lambda \in \Lambda} \epsilon_\lambda \lambda.$$

By the triangle inequality, for every $x \in B$ we have

$$|1 - \gamma(x)| \le |1 - \gamma_0(x)| + |1 - \gamma_1(x)| + d \sup_{\lambda \in \Lambda} |1 - \lambda(x)| \le 1/2.$$

It follows that

$$\left| \widehat{B}(\gamma) - |B| \right| \le \sum_{x \in B} |1 - \gamma(x)| \le 2^{-1} |B|$$

and the conclusion follows.

Finally, recalling Theorem 4.1 we see that to make use of Lemma 5.2 we will need to have good control on the spectrum of a given set. The following lemma gives a more useful criterion to ensure this.

Lemma 5.3. If c > 0 and $|(B + B') \setminus B| \le c\epsilon |B|$ then B' has 2c-control of $\Delta_{\epsilon}(B)$. *Proof.* Choose γ such that $|\widehat{B}(\gamma)| \ge \epsilon |B|$ and $x \in B'$. Then

$$|1 - \gamma(x)| |B| \le \epsilon^{-1} \left| \sum_{y \in B} \gamma(y) - \sum_{y \in B+x} \gamma(y) \right|$$
$$\le 2\epsilon^{-1} |(B + B') \setminus B|$$
$$\le 2c |B|.$$

We now combine all our tools thus far to prove the following efficient density increment theorem.

Theorem 5.1. Let $B, B' \subset G$ be any sets. Let $A \subset B$ with density $\alpha = |A| / |B|$ and $f : B' \to [-1, 1]$ with density $\tau = \|f\|_1 / |B|$. Let $\mathbf{A}(x) = A(x) - \alpha B(x)$ and suppose that

$$\sum_{\gamma} \left| \widehat{f}(\gamma) \right| \left| \widehat{\mathbf{A}}(\gamma) \right|^2 = \nu \alpha \left\| f \right\|_1 \left| A \right| N_{\gamma}$$

Then there is a finite set $\Lambda \subset \widehat{G}$ of size

$$d = |\Lambda| \le 2^{16} \mathcal{L}(\tau) (\nu \alpha)^{-1}$$

such that if a symmetric set $B'' \subset G$ has $(4d)^{-1}$ -control of Λ ,

$$\left| (2B'' + B) \backslash B \right| \le 2^{-17} \nu \alpha \left| B \right|,$$

and

$$|(B''+B')\backslash B'| \le 2^{-4} \exp(-2^4 \mathcal{L}(\tau) \mathcal{L}(\nu\alpha)) |B'|$$

then there exists x such that

$$(A - x) \cap B''| \ge (1 + 2^{-16}\nu)\alpha |B''|.$$

Proof. Let $\Delta = \Delta_{\eta}(f)$ where $\eta = \nu \alpha/2$. In particular,

$$\sum_{\gamma \notin \Delta} \left| \widehat{f}(\gamma) \right| \left| \widehat{\mathbf{A}}(\gamma) \right|^2 \le 2^{-1} \nu \alpha \left\| f \right\|_1 \left\| \mathbf{A} \right\|_2^2 N \le 2^{-1} \nu \alpha \left\| f \right\|_1 \left| A \right| N.$$

In particular,

$$\sum_{\gamma \in \Delta} \left| \widehat{f}(\gamma) \right| \left| \widehat{\mathbf{A}}(\gamma) \right|^2 \ge 2^{-1} \nu \alpha \left\| f \right\|_1 |A| N.$$

We now perform a dyadic decomposition of Δ into $\Delta_i = \tilde{\Delta}_{2^i\eta}(f)$, and apply Theorem 4.1 to each Δ_i with the weight function

$$\omega(\gamma) = \left| \widehat{f}(\gamma) \right| \left| \widehat{\mathbf{A}}(\gamma) \right|^2$$

and

$$\epsilon = \exp(-2^4 \mathcal{L}(\nu \alpha) \mathcal{L}(\tau)) \le \exp(-8\mathcal{L}(\eta) \mathcal{L}(\tau)).$$

Then, if Δ'_i is the set provided by Theorem 4.1, we have

$$2^{-12} \sum_{\gamma \in \Delta_{i}} \left| \widehat{f}(\gamma) \right| \left| \widehat{\mathbf{A}}(\gamma) \right|^{2} \leq (2^{i}\eta)^{-1} \sum_{\gamma \in \Delta_{i}^{\prime}} \left| \widehat{f}(\gamma) \right| \left| \widehat{\mathbf{A}}(\gamma) \right|^{2}$$
$$\leq 2 \left\| f \right\|_{1} \sum_{\gamma \in \Delta_{i}^{\prime}} \left| \widehat{\mathbf{A}}(\gamma) \right|^{2}.$$

Summing over all $i \ge 0$ implies that

$$\sum_{\gamma \in \Delta'} \left| \widehat{\mathbf{A}}(\gamma) \right|^2 = \sum_{i} \sum_{\gamma \in \Delta'_i} \left| \widehat{\mathbf{A}}(\gamma) \right|^2 \ge 2^{-13} \nu \alpha \left| A \right| N,$$

where $\Delta' = \bigcup_{i \ge 0} \Delta'_i$. Furthermore, since each Δ'_i is $2^{14} \mathcal{L}(\tau) (2^i \eta)^{-1}$ -covered by $\Delta_{\epsilon}(B')$ it follows that Δ' is *d*-covered by $\Delta_{\epsilon}(B')$ where

$$d \le 2^{14} \mathcal{L}(\tau) \sum_{i} (2^{i} \eta)^{-1} \le 2^{16} \mathcal{L}(\tau) (\nu \alpha)^{-1}.$$

Since B'' satisfies $|(B'' + B') \setminus B'| \leq 2^{-4} \epsilon |B'|$ it follows from Lemma 5.3 that B'' has 2^{-3} -control of $\Delta_{\epsilon}(B')$. Hence by Lemma 5.2 there is a Λ of size $d \leq 2^{16} \mathcal{L}(\tau)(\nu \alpha)^{-1}$ such that if B'' has $(4d)^{-1}$ -control of Λ then for every $\gamma \in \Delta'$ we have

$$\left|\widehat{B''}(\gamma)\right| \ge 2^{-1} \left|B''\right|.$$

Since we also have $|(2B'' + B) \setminus B| \le 2^{-17} \nu \alpha |B|$ it follows from Lemma 5.1 that

$$|(A - x) \cap B''| \ge (1 + 2^{-16}\nu)\alpha |B''|$$

as required.

We now at last recall our purpose, which is to study the number of solutions to (1) in a given set A. This count is given by $\langle (c_1 \cdot A) * (c_2 \cdot A), (-c_3 \cdot A) \rangle$. It is now straightforward to use Parseval's theorem and Theorem 5.1 to prove the following general density increment theorem, which shows that either this count is large or some dilation of A has increased density on some structured subset.

Theorem 5.2. Suppose that $B' \subset B \subset G$ and we have $A_1 \subset B'$ and $A_2, A_3 \subset B$ each with relative densities α_i . Let $\alpha = 2^{-1} \min(2^{-5}, \alpha_1, \alpha_2, \alpha_3)$ and suppose that

$$|(B'+B)\backslash B| \le 2^{-2}\alpha |B|$$

Then either

(5)
$$\langle A_1 * A_2, A_3 \rangle \ge 2^{-2} \alpha_1 \alpha_2 \alpha_3 |B| |B'|$$

or there is a finite set $\Lambda \subset \widehat{G}$ of size

$$d = |\Lambda| \le 2^{19} \mathcal{L}(\alpha) \alpha^{-1}$$

such that if a symmetric set B'' has $(4d)^{-1}$ -control of Λ ,

$$|(2B'' + B) \setminus B| \le 2^{-19} \alpha |B|,$$

and

$$|(B'' + B') \setminus B'| \le 2^{-4} \exp(-2^5 \mathcal{L}(\alpha)^2)) |B'|$$

then there exists $x \in G$ and $i \in \{2,3\}$ such that

$$|(A_i - x) \cap B''| \ge (1 + 2^{-18})\alpha_i |B''|.$$

Proof. Let $\mathbf{A}_i = A_i - \alpha_i B$ for i = 2, 3 and $\mathbf{A}_1 = A_1 - \alpha_1 B'$. We have

$$\sum_{x} A_1 * \mathbf{A_2}(x) \mathbf{A_3}(x) = \sum_{x} A_1 * A_2(x) A_3(x) - \alpha_2 \sum_{x \in B} A_3 * A_1^-(x)$$
$$-\alpha_3 \sum_{x \in B} A_1 * A_2(x) + \alpha_2 \alpha_3 \sum_{x \in B} B * A_1(x).$$

Observe that if A' is supported on B' and A is supported on B then

$$\left| \sum_{x \in B} A' * A(x) - |A'| |A| \right| \le |A'| |(B' + B) \setminus B| \le 2^{-2} |A'| |A|.$$

In particular,

$$\sum_{x} A_1 * \mathbf{A_2}(x) \mathbf{A_3}(x) \le \sum_{x} A_1 * A_2(x) A_3(x) - 2^{-1} \alpha_1 \alpha_2 \alpha_3 |B| |B'|.$$

By the triangle inequality either (5) is true or

$$\sum_{\gamma} \left| \widehat{A_1}(\gamma) \right| \left| \widehat{A_2}(\gamma) \right| \left| \widehat{A_3}(\gamma) \right| \ge 2^{-2} \alpha_1 \alpha_2 \alpha_3 \left| B' \right| \left| B \right| N.$$

Applying the Cauchy-Schwarz inequality it follows that

$$\left(\sum_{\gamma} \left|\widehat{A}_{1}(\gamma)\right| \left|\widehat{\mathbf{A}}_{2}(\gamma)\right|^{2}\right) \left(\sum_{\gamma} \left|\widehat{A}_{1}(\gamma)\right| \left|\widehat{\mathbf{A}}_{3}(\gamma)\right|^{2}\right) \geq 2^{-4} \left|A_{1}\right|^{2} \alpha_{2}^{2} \alpha_{3}^{2} \left|B\right|^{2} N^{2}.$$

It follows that for some $i \in \{2, 3\}$ we have

$$\sum_{\gamma} \left| \widehat{A_1}(\gamma) \right| \left| \widehat{\mathbf{A}_i}(\gamma) \right|^2 \ge 2^{-2} \left| A_1 \right| \alpha_i^2 \left| B \right| N$$

and the conclusion then follows from Theorem 5.1.

6. POLYNOMIAL RINGS

For the first demonstration of this method we choose $G = \mathbb{F}_q[t]/(p(t))$, where p(t) is some prime of degree n, so that $N = q^n$, and q is some odd prime power. We denote this group by \mathfrak{A}_N . This group is rich enough to mirror the behaviour of the integers, while it has the significant technical advantage of finite characteristic.

We fix some coefficients $c_1, c_2, c_3 \in \mathbb{F}_q[t] \setminus \{0\}$ such that $c_1 + c_2 + c_3 = 0$, and some finite set $A \subset \mathfrak{A}_N$. We suppose that N is large enough so that these coefficients are coprime to p(t), and hence each acts faithfully on \mathfrak{A}_N . We wish to study the function that counts solutions to $c_1x_1 + c_2x_2 + c_3x_3 = 0$ with $x_i \in A$, denoted by

(6)
$$\Upsilon_{\mathbf{c}}(A) = \langle (c_1 \cdot A) * (c_2 \cdot A), (-c_3 \cdot A) \rangle.$$

We note that $\Upsilon_{\mathbf{c}}(A)$ is invariant under dilations and translations of A.

We recall that both \mathfrak{A}_N and $\widehat{\mathfrak{A}_N}$ are \mathbb{F}_q -vector spaces of dimension n. Furthermore, any $a \in \mathfrak{A}_N$ acts on $\widehat{\mathfrak{A}_N}$ by letting $(a\gamma)(x) = \gamma(ax)$. The Bohr space of $\Gamma \subset \widehat{\mathfrak{A}_N}$ is defined as

$$B(\Gamma) = \{ x \in \mathfrak{A}_N : \gamma(x) = 1 \text{ for all } \gamma \in \Gamma \}.$$

If B is a Bohr space then the frequency set of B is

$$[B] = \{ \gamma \in \widehat{\mathfrak{A}_N} : \gamma(x) = 1 \text{ for all } x \in B \},\$$

and we define the rank of B as the \mathbb{F}_q -dimension of [B].

Lemma 6.1. Every Bohr space is an \mathbb{F}_q -subspace of \mathfrak{A}_N , and conversely, every \mathbb{F}_q -subspace of \mathfrak{A}_N is a Bohr space. Furthermore if B is a Bohr space then

$$|B| = q^{-\mathrm{rk}(B)}N$$

Proof. We first claim that if V is a \mathbb{F}_q -subspace of \mathfrak{A}_N then |V||[V]| = N. Since V is closed under addition it is easy to check that if $\gamma \notin [V]$ then $\widehat{V}(\gamma) = 0$, and if $\gamma \in [V]$ then $\widehat{V}(\gamma) = |V|$, and hence by Parseval's identity,

$$|V|N = \sum_{\gamma} \left| \widehat{V}(\gamma) \right|^2 = \sum_{\gamma \in [V]} \left| \widehat{V}(\gamma) \right|^2 = |[V]| \left| V \right|^2$$

which proves our initial claim. The rest of the lemma follows easily.

Corollary 6.1. If B is a Bohr space and $c \in \mathfrak{A}_N \setminus \{0\}$ then $c \cdot B$ is also a Bohr space and

$$\operatorname{rk}(c \cdot B) = \operatorname{rk}(B).$$

Since Bohr spaces are closed under addition our density increment tool, Theorem 5.2, takes the following particularly simple form in this setting.

Theorem 6.1. Suppose that $B \subset \mathfrak{A}_N$ is a Bohr space and we have $A_1, A_2, A_3 \subset B$ with relative densities $\alpha_i = |A_i| / |B|$. Let $\alpha = \min(\alpha_1, \alpha_2, \alpha_3)$. Then either

$$\langle A_1 * A_2, A_3 \rangle \gg \alpha_1 \alpha_2 \alpha_3 | B$$

or there is a Bohr space $B' \subset B$ of rank

$$\operatorname{rk}(B') \leq \operatorname{rk}(B) + O(\mathcal{L}(\alpha)\alpha^{-1})$$

such that there exists $x \in \mathfrak{A}_N$ and $i \in \{2, 3\}$ such that

$$|(A_i - x) \cap B'|) \ge (1 + \Omega(1))\alpha_i |B'|.$$

We may then iterate this density increment in the usual fashion to obtain a lower bound on $\Upsilon_{\mathbf{c}}(A)$, from which Theorem 1.4 follows easily.

Theorem 6.2. Suppose that $\mathbf{c} \in (\mathbb{F}_q[t] \setminus \{0\})^3$ is such that $c_1 + c_2 + c_3 = 0$ and N is sufficiently large, depending only on \mathbf{c} . For any $A \subset \mathfrak{A}_N$

$$\Upsilon_{\mathbf{c}}(A) \ge \exp_q \left(-O_{\mathbf{c}}(\mathcal{L}(\alpha)^2 \alpha^{-1}) \right) N^2.$$

Proof. Let $\ell = \max_{1 \le i \le 3} \deg c_i$. Let K be maximal such that there exists a sequence of Bohr spaces

$$\mathfrak{A}_N=B_0\supset B_1\supset\cdots\supset B_K,$$

and a sequence of sets $A_i \subset B_i$ with density $\alpha_i = |A_i| / |B_i|$ such that, for $0 \leq i < K$, there exists Λ_{i+1} of size $O(\mathcal{L}(\alpha)\alpha_i^{-1})$ and $a_i \in \mathfrak{A}_N \setminus \{0\}$ such that there exist Γ_i with $B_i = B(\Gamma_i)$ and

(8)
$$\Gamma_{i+1} \subset (a_i \cdot \{1, \dots, t^{3\ell}\} \cdot \Gamma_i) \cup \Lambda_{i+1}$$

and for some absolute constant c > 0 we have

$$\alpha_{i+1} \ge (1+c)\,\alpha_i,$$

and furthermore $\Upsilon_{\mathbf{c}}(A_{i+1}) \leq \Upsilon_{\mathbf{c}}(A_i)$. In particular,

$$1 \ge \alpha_K \ge (1+c)^K \alpha,$$

and hence $K \ll \mathcal{L}(\alpha)$. Furthermore it follows from (8) and induction that for $1 \leq J \leq K$, if $\Lambda_0 = \{0\}$, there exist $a'_j \in \mathfrak{A}_N \setminus \{0\}$ for $0 \leq j < J$ such that

$$\Gamma_J \subset \Lambda_J \cup \bigcup_{j=0}^{J-1} a'_j \cdot \{1, \dots, t^{3\ell(J-j)}\} \cdot \Lambda_j,$$

and so in particular

(9)
$$\operatorname{rk}(B_K) \ll_{\ell} K \sum_{j=0}^{K} |\Lambda_j| \ll_{\ell} K \mathcal{L}(\alpha) \alpha^{-1} \sum_{i=0}^{K-1} (1+c)^{-i} \ll_{\ell} \mathcal{L}(\alpha)^2 \alpha^{-1}.$$

We will demonstrate that

(10)
$$\Upsilon_{\mathbf{c}}(A_K) \gg \alpha^3 |B_K|^2$$

and the theorem follows from (7) and (9).

Let $B' = B(\{1, \ldots, t^{3\ell}\} \cdot \Gamma_K)$. We observe that $a \cdot B' \subset B_K$ for all $a \in \mathbb{F}_q[t]$ with deg $a \leq 3\ell$ and if B'' is a Bohr space of the shape $a \cdot B'$ and $a \neq 0$ then

$$[B''] \subset a^{-1} \cdot \{1, \dots, t^{3\ell}\} \cdot \Gamma_K$$

Consider $B'' = c_1 c_2 c_3 \cdot B'$ and $B^{(i)} = c_i^{-1} \cdot B''$ for $1 \le i \le 3$. Since $B^{(i)} \subset B_K$ for $1 \le i \le 3$

$$\sum_{x \in B_K} (A_K * \beta^{(1)} + A_K * \beta^{(2)} + A_K * \beta^{(3)})(x) = 3 |A_K|,$$

and so for some $x \in B_K$

$$A_K * \beta^{(1)}(x) + A_K * \beta^{(2)}(x) + A_K * \beta^{(3)}(x) \ge 3\alpha_K$$

If for some $1 \leq i \leq 3$ we have $A_K * \beta^{(i)}(x) \geq \alpha_K(1+c)$ then this contradicts the maximality of K, letting $B_{K+1} = B^{(i)}$ and $A_{K+1} = A_K - x$. Otherwise, for $1 \leq i \leq 3$,

$$A_K * \beta^{(i)}(x) \ge (1 - 4c)\alpha_K.$$

If we let $A'_i = c_i \cdot ((A_K - x) \cap B'')$ then we have satisfied the hypotheses required to apply Theorem 6.1. It follows that either (10) holds or we may choose B_{K+1} to be the sub-Bohr set of B_K with control of Λ and A_{K+1} to be a translation of some A'_i . Since

$$|\Lambda| = O(\alpha_K^{-1} \mathcal{L}(\alpha))$$

and

$$\alpha_{K+1} \ge (1 + \Omega(1))(1 - 4c)\alpha_K \ge (1 + c)\alpha_K$$

if we choose c sufficiently small, this contradicts the maximality of K and thus concludes the proof.

7. Integers

We now prove Theorem 1.3 by applying these methods to the case $G = \mathbb{Z}_N$, where N is some large prime. Analogously to the case $G = \mathfrak{A}_N$ we fix some coefficients $c_1, c_2, c_3 \in \mathbb{Z} \setminus \{0\}$, all coprime to N, such that $c_1 + c_2 + c_3 = 0$ and for $A \subset \mathbb{Z}_N$ define $\Upsilon_{\mathbf{c}}(A)$ as in (6). For $\Gamma \subset \widehat{\mathbb{Z}}_N$ and $\rho : \Gamma \to [0, 2]$ we define the Bohr set $B_{\rho}(\Gamma)$ as

$$\{n \in \mathbb{Z}_N : |\gamma(n) - 1| < \rho(\gamma)\}.$$

We call Γ the frequency set of B and ρ the width, and define the rank of B to be the size of Γ . In fact, when we speak of a Bohr set we implicitly refer to the triple $(\Gamma, \rho, B_{\rho}(\Gamma))$, since the set $B_{\rho}(\Gamma)$ does not uniquely determine the frequency set or the width. Furthermore, if $\rho : \Gamma \to [0, 2]$ and $\rho' : \Gamma' \to [0, 2]$ then we define $\rho \wedge \rho' : \Gamma \cup \Gamma' \to [0, 2]$ by

$$(\rho \land \rho')(\gamma) = \begin{cases} \rho(\gamma) & \text{if } \gamma \in \Gamma \backslash \Gamma' \\ \rho'(\gamma) & \text{if } \gamma \in \Gamma' \backslash \Gamma, \text{ and} \\ \min(\rho(\gamma), \rho'(\gamma)) & \text{if } \gamma \in \Gamma \cap \Gamma'. \end{cases}$$

We no longer have the convenient property of Bohr sets being closed under addition, but Bourgain [1999] observed that certain Bohr sets have a weak version of this property suitable for our applications. A Bohr set $B_{\rho}(\Gamma)$ of rank d is regular if for all $|\kappa| \leq 2^{-6}d^{-1}$ we have

$$\left| \left| B_{\rho(1+\kappa)}(\Gamma) \right| - \left| B_{\rho}(\Gamma) \right| \right| \le 2^{6} d \left| \kappa \right| \left| B_{\rho}(\Gamma) \right|.$$

If $B = B_{\rho}(\Gamma)$ then we write $B(\lambda)$ for the Bohr set $B_{\lambda\rho}(\Gamma)$. For further discussion of Bohr sets and proofs of the following basic lemmas see, for example, [Tao and Vu, 2006, Chapter 4].

Lemma 7.1. For any Bohr set B there exists $\lambda \in [1/2, 1]$ such that $B(\lambda)$ is regular. **Lemma 7.2.** If $\rho' : \Gamma' \to [0, 2]$ then

$$|B_{\rho \wedge \rho'}(\Gamma \cup \Gamma')| \ge \prod_{\gamma \in \Gamma'} (\rho'(\gamma)/4) |B_{\rho}(\Gamma)|.$$

Furthermore, if B is a Bohr set of rank d then $|B(\lambda)| \ge \lambda^{O(d)} |B|$.

The following lemma follows easily from the observation that if $c \in \mathbb{Z}_N \setminus \{0\}$ then $c \cdot B_{\rho}(\Gamma) = B_{\rho}(c \cdot \Gamma)$.

Lemma 7.3. For any Bohr set B and $c \in \mathbb{Z}_N^*$ the set $c \cdot B$ is also a Bohr set of rank $\operatorname{rk}(B)$, and furthermore for any $\lambda > 0$

$$c \cdot (B(\lambda)) = (c \cdot B)(\lambda).$$

The notion of regularity allows us to exert the required amount of additive control, and our density increment tool, Theorem 5.2, becomes the following.

Theorem 7.1. There exists an absolute constant c > 0 such that the following holds. Let $B \subset \mathbb{Z}_N$ be a regular Bohr set of rank d where $d \leq \exp(c\mathcal{L}(\alpha)^2)$. Let $A_1, A_2 \subset B$ and $A_3 \subset B(\delta)$, each with relative densities α_i . Let $\alpha = \min(c, \alpha_1, \alpha_2, \alpha_3)$ and suppose that $B(\delta)$ is also regular and $cd^{-1}\alpha/4 \leq \delta \leq cd^{-1}\alpha$. Then either

(11)
$$\langle A_1 * A_2, A_3 \rangle \gg \alpha_1 \alpha_2 \alpha_3 |B| |B(\delta)|$$

or there is a regular Bohr set B' of rank

$$\operatorname{rk}(B') \le d + O(\mathcal{L}(\alpha)\alpha^{-1})$$

and size

(12) $|B'| \ge \exp(-O(\mathcal{L}(\alpha)^2(d + \mathcal{L}(\alpha)\alpha^{-1})))|B|$

such that there exists $x \in \mathbb{Z}_N$ and $i \in \{1, 2\}$ with

 $|(A_i - x) \cap B'| \ge (1 + c)\alpha_i |B'|.$

Proof. We first observe that since B is regular we have that

(13)
$$|(B(\delta) + B) \setminus B| \ll d\delta |B| \le 2^{-2} \alpha |B|,$$

provided δ is sufficiently small. The hypotheses of Theorem 5.2 are now met, so that either (11) holds or there is a finite set $\Lambda \subset \widehat{G}$ of size $l = O(\mathcal{L}(\alpha)\alpha^{-1})$ such that if a symmetric set B' has $(4l)^{-1}$ -control of Λ ,

(14)
$$|(2B'+B)\backslash B| \le 2^{-19}\alpha |B|,$$

and

(15)
$$|(B' + B(\delta)) \setminus B(\delta)| \le 2^{-4} \exp(-2^5 \mathcal{L}(\alpha)^2)) |B(\delta)|$$

then there exists $x \in \mathbb{Z}_N$ and $i \in \{1, 2\}$ such that

$$|(A_i - x) \cap B'| \ge (1 + 2^{-18})\alpha_i |B'|.$$

Let $\rho': \Lambda \to [0,2]$ be defined as $\rho'(\lambda) = 1/4l$ for all $\lambda \in \Lambda$. If $B = B_{\rho}(\Gamma)$ then let $B^* = B_{\delta\rho\wedge\rho'}(\Gamma\cup\Lambda)$, and let $B' = B^*(\delta')$ for some δ' to be chosen later, but chosen such that B' is regular. Clearly, B' is a regular Bohr set of rank $d + O(\mathcal{L}(\alpha)\alpha^{-1})$ with $(4l)^{-1}$ -control of Λ by choice of ρ' . Provided $\delta' \leq 1/2$ we have that $2B' \subset B(\delta)$, and hence (14) is satisfied, arguing as for (13). Furthermore, since $B(\delta)$ is regular and $B' \subset B(\delta\delta')$ it follows that

$$\left| (B' + B(\delta)) \setminus B(\delta) \right| \ll d\delta' \left| B(\delta) \right|,$$

and hence (15) is satisfied for some $\delta' \gg \exp(-O(\mathcal{L}(\alpha)^2)))d^{-1}$. Finally, by Lemma 7.2 we have

$$|B'| \ge (\delta')^{O(l+d)} |B^*| \ge l^{-O(l)} \delta^d(\delta')^{O(l+d)} |B|,$$

and (12) follows from our bounds on δ , δ' and l, and the proof is complete.

This density increment lemma may then be iterated in the standard fashion to yield a lower bound for $\Upsilon_{\mathbf{c}}(A)$.

Theorem 7.2. Let $\mathbf{c} \in (\mathbb{Z} \setminus \{0\})^3$ be such that $c_1 + c_2 + c_3 = 0$. For any prime N sufficiently large, depending only on \mathbf{c} , and any $A \subset \mathbb{Z}_N$ we have

$$\Upsilon_{\mathbf{c}}(A) \ge \exp\left(-O_{\mathbf{c}}\left(\mathcal{L}(\alpha)^4 \alpha^{-1}\right)\right) N^2.$$

Theorem 1.3 follows by embedding $\{1, \ldots, N\}$ into a suitable subinterval I of $\mathbb{Z}_{N'}$, where $N' \ll_{\mathbf{c}} N$ is some prime large enough such that if $x_1, x_2, x_3 \in I$ is a solution to $c_1x_1 + c_2x_2 + c_3x_3 = 0$ in $\mathbb{Z}_{N'}$ then it is also a solution in $\{1, \ldots, N\}$.

Proof. Let K be maximal such that there exists a sequence of regular Bohr sets

$$\mathbb{Z}_N = B_0 \supset B_1 \supset \cdots \supset B_K,$$

each with frequency set Γ_i , rank d_i , width ρ_i and a sequence of sets $A_i \subset B_i$ with density $\alpha_i = |A_i| / |B_i|$ such that, for $0 \le i < K$ we have

$$d_{i+1} \le d_i + O(\mathcal{L}(\alpha)\alpha_i^{-1}) \le \exp(O(\mathcal{L}(\alpha)^2)),$$

(16)
$$|B_{i+1}| \ge \exp(-O(\mathcal{L}(\alpha)^2(d_i + \mathcal{L}(\alpha)\alpha_i^{-1}))) |B_i|,$$
$$\alpha_{i+1} \ge (1+c)\alpha_i,$$

and furthermore $\Upsilon_{\mathbf{c}}(A_{i+1}) \leq \Upsilon_{\mathbf{c}}(A_i)$. In particular, this will follow provided A_{i+1} is a subset of a dilation of a translation of A_i . We observe that

$$1 \ge \alpha_K \ge (1+c)^K \alpha_s$$

and hence $K \ll \mathcal{L}(\alpha)$. Furthermore, for $0 \leq j \leq K$ we have the estimate $d_j \ll$ $\mathcal{L}(\alpha)\alpha^{-1}$, and so

$$|B_K| \ge \exp\left(-O_{\mathbf{c}}\left(\mathcal{L}(\alpha)^2 \sum_{i=0}^K d_i\right)\right) N \ge \exp(-O(\mathcal{L}(\alpha)^4 \alpha^{-1}))N.$$

We will demonstrate that

$$\Upsilon_{\mathbf{c}}(A_K) \gg \exp(-O(d_K \mathcal{L}(\alpha)^2)) |B_K|^2$$

and the theorem follows.

1

For brevity, let $B' = B_K(\rho'), B'' = B'(\rho') = B_K(\rho'\rho'')$, and $B''' = B''(\rho''') =$ $B_K(\rho'\rho''\rho''')$, where ρ', ρ'' and ρ''' will be chosen later, but chosen such that all Bohr sets considered are regular. Let

$$B^{(1)} = c_2 c_3 \cdot B'', \ B^{(2)} = c_1 c_3 \cdot B'', \ \text{and} \ B^{(3)} = c_2 c_3 \cdot B'''$$

so that in particular if $\rho'' \ll 1/c_1c_2c_3$ we have $B^{(i)} \subset B'$ for $1 \leq i \leq 3$, and hence by the regularity of B_K

$$\left|\sum_{x\in B_K} A_K * \beta^{(i)}(x) - |A_K|\right| \le |(B'+B_K)\backslash B_K| \ll d\rho' |B_K|.$$

In particular for any $\epsilon > 0$ we may choose $\rho' \gg_{\epsilon} d^{-1} \alpha_K$ such that

$$\sum_{x \in B_K} (A_K * \beta^{(1)} + A_K * \beta^{(2)} + A_K * \beta^{(3)})(x) \ge (3 - \epsilon) |A_K|,$$

and hence, for some $x \in B$,

$$A_K * \beta^{(1)}(x) + A_K * \beta^{(2)}(x) + A_K * \beta^{(3)}(x) \ge (3 - \epsilon)\alpha_K.$$

If $A_K * \beta^{(i)}(x) \ge \alpha(1+c)$ for some $1 \le i \le 3$ then this contradicts the maximality of K, letting $B_{K+1} = B^{(i)}$ and $A_{K+1} = (A_K - x) \cap B_{K+1}$. In particular note that $B^{(i)}$ also has rank d_K and

$$\left|B^{(i)}\right| \ge |B^{\prime\prime\prime}| \ge \exp(-O(d_K \mathcal{L}(\rho^{\prime} \rho^{\prime\prime} \rho^{\prime\prime\prime}))) |B_K|,$$

and we shall see that our choices for the ρ parameters give the lower bound (16). Otherwise, for $1 \leq i \leq 3$, if we choose ϵ sufficiently small depending on c, then for $1 \leq i \leq 3$

$$A_K * \beta^{(i)}(x) \ge (1-c)\alpha_K.$$

Let $A_i = c_i \cdot (A' - x) \cap c_i \cdot B^{(i)}$, and observe that

$$c_3 \cdot B^{(3)} = c_1 c_2 c_3 \cdot B^{\prime \prime \prime} \subset (c_1 c_2 c_3 \cdot B^{\prime \prime})(\rho^{\prime \prime \prime}),$$

and hence there exists $\rho''' \gg d_K^{-1} \alpha_K$ such that we are in a position to apply Theorem 7.1. In particular, either

$$\Upsilon_{\mathbf{c}}(A_K) \gg \alpha_K^3 |B''| |B'''| \gg \exp(-O(d_K \mathcal{L}(\alpha)^2)) |B_K|^2$$

or there is a regular Bohr set B^{\sharp} of rank

$$d^{\sharp} \le d + O(\mathcal{L}(\alpha)\alpha_K^{-1})$$

and size

$$|B^{\sharp}| \geq \exp(-O(\mathcal{L}(\alpha)^{2}(d+\mathcal{L}(\alpha)\alpha_{K}^{-1})))|B''| \geq \exp(-O(\mathcal{L}(\alpha)^{2}(d+\mathcal{L}(\alpha)\alpha_{K}^{-1})))|B_{K}|,$$

and $i \in \{1, 2\}$ such that

$$\left| (A_i - x) \cap B^{\sharp} \right| \ge (1 + \Omega(1))(1 - c)\alpha_K \left| B^{\sharp} \right| \ge (1 + c)\alpha_K \left| B^{\sharp} \right|$$

if we choose c sufficiently small, which contradicts the maximality of K. This concludes the proof.

8. ARITHMETIC PROGRESSIONS IN SUMSETS

In this section we sketch how our improved structural result for spectra may be combined with the methods of Sanders [2008] to prove Theorem 1.2. The difference lies in an improvement of the iteration lemma, Lemma 6.4 of Sanders [2008]. In particular, Theorem 1.2 follows immediately from the proof of Sanders [2008], replacing the use of its Lemma 6.4 by the following quantitatively superior version.

Lemma 8.1. Let $B \subset \mathbb{Z}_N$ be a regular Bohr set and $A_1, A_2 \subset B$ with densities α_1 and α_2 respectively. For any $\sigma \in (0, 1]$ either

- (1) there is a regular Bohr set $B' = B(\rho)$ such that $A_1 + A_2$ contains at least a proportion 1σ of B' and $\delta' \gg \alpha^2 d^{-1}$, or
- (2) there is a regular Bohr set $B'' \subset B$ of rank

$$\operatorname{rk}(B'') \le \operatorname{rk}(B) + O(\alpha^{-1}\mathcal{L}(\sigma))$$

and width

$$\rho(B'') \gg \rho(B)(\operatorname{rk}(B))^{-1} \exp(-O(\mathcal{L}(\sigma)\mathcal{L}(\alpha)))$$

such that for some absolute constant c > 0

$$||A_1 * \beta''||_{\infty} ||A_2 * \beta''||_{\infty} \ge \alpha_1 \alpha_2 (1+c).$$

Proof. Let $\mathbf{A}_i = A_i - \alpha B$, and $B' = B(\rho)$ for some suitable $\rho \gg \alpha^4 d^{-1}$ such that B' is regular. Arguing as in the proof of Lemma 6.4 in Sanders [2008] we see that either the first case holds or there is some $S \subset B'$ and $1 \le i \le 2$ such that

$$\sum_{\gamma \in \widehat{\mathbb{Z}_N}} \left| \widehat{\mathbf{A}_i}(\gamma) \right|^2 \left| \widehat{S}(\gamma) \right| \gg \alpha_i \left| A_i \right| \left| S \right| N.$$

It follows from Theorem 5.1 that if we choose our Bohr set $B'' \subset B'$ such that it has $\Omega(l^{-1})$ -control of Λ , where Λ is a set of size $l \ll \alpha_i^{-1} \mathcal{L}(\sigma)$ and

$$|(B'' + B') \setminus B'| \ll \exp(-\mathcal{L}(\sigma)\mathcal{L}(\alpha_i)) |B'|$$

then $||A_i * \beta''|| \ge (1+c)\alpha_i$. The proof is then completed as in the proof of Lemma 6.4 in Sanders [2008].

Acknowledgements

The author would like to thank Julia Wolf and Trevor Wooley for their advice and helpful comments on a preliminary draft of this paper, and Kevin Henriot for pointing out several flaws in an earlier version of the argument.

References

- Bateman, M. and N. H. Katz "New bounds on cap sets" J. Amer. Math. Soc. 25 (2012): 585–613.
- Behrend, F. A. "On sets of integers which contain no three terms in arithmetical progression" Proc. Nat. Acad. Sci. U. S. A. 32 (1946): 331–332.
- Bloom, T. F. "Translation invariant equations and the method of Sanders" Bull. Lond. Math. Soc. 44 (2012): 1050–1067.
- Bourgain, J. "On triples in arithmetic progression" *Geom. Funct. Anal.* 9 (1999): 968–984.
- Bourgain, J. "Roth's theorem on progressions revisited" J. Anal. Math. 104 (2008): 155–192.
- Chang, M.-C. "A polynomial bound in Freiman's theorem" *Duke Math. J.* 113 (2002): 399–419.
- Croot, E., Laba, I. and O. Sisask "Arithmetic progressions in sumsets and L^{p} -almost-periodicity" Combin. Probab. Comput. 22 (2013): 351–365.+
- Green, B. "Arithmetic progressions in sumsets" *Geom. Funct. Anal.* 12 (2002): 584–597.
- Heath-Brown, D. R. "Integer sets containing no arithmetic progressions" J. London Math. Soc. 35 (1987): 385–394.
- Liu, Y.-R. and C. V. Spencer "A generalization of Roth's theorem in function fields" Int. J. Number Theory 5 (2009): 1149–1154.
- Meshulam, R. "On subsets of finite abelian groups with no 3-term arithmetic progressions" J. Combin. Theory Ser. A 71 (1995): 168–172.
- Roth, K. F. "On certain sets of integers" J. London Math. Soc. 28 (1953): 104-109.
- Sanders, T. "Additive structures in sumsets" Math. Proc. Cambridge Philos. Soc. 144 (2008): 289–316.
- Sanders, T. "On Roth's theorem on progressions" Ann. of Math. 174 (2011): 619–636.
- Sanders, T. "On certain other sets of integers" J. Anal. Math. 116 (2012): 53-82.
- Shkredov, I. D. "On sets of large trigonometric sums" *Izv. Math.* 72 (2008): 149–168.
- Shkredov, I. D. "On sumsets of dissociated sets" Online J. Anal. Comb. 4 (2009).
- Szemerédi, E. "Integer sets containing no arithmetic progressions" Acta Math. Hungar. 56 (1990): 155–158.
- Tao, T., and V. Vu. Additive Combinatorics, 1st ed. Cambridge University Press, 2006.

THOMAS F. BLOOM

THOMAS BLOOM, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRISTOL, UNIVERSITY WALK, CLIFTON, BRISTOL BS8 1TW, UNITED KINGDOM *E-mail address:* matfb@bristol.ac.uk