

DIAGRAMS FOR CONTACT 5-MANIFOLDS

FAN DING, HANSJÖRG GEIGES, AND OTTO VAN KOERT

ABSTRACT. According to Giroux, contact manifolds can be described as open books whose pages are Stein manifolds. For 5-dimensional contact manifolds the pages are Stein surfaces, which permit a description via Kirby diagrams. We introduce handle moves on such diagrams that do not change the corresponding contact manifold. As an application, we derive classification results for subcritically Stein fillable contact 5-manifolds and characterise the standard contact structure on the 5-sphere in terms of such fillings. This characterisation is discussed in the context of the Andrews–Curtis conjecture concerning presentations of the trivial group. We further illustrate the use of such diagrams by a covering theorem for simply connected spin 5-manifolds and a new existence proof for contact structures on simply connected 5-manifolds.

1. INTRODUCTION

The aim of this paper is to develop a diagrammatic language for 5-dimensional contact manifolds. This is motivated by (but does not depend on) the deep result of Giroux [18], which says that any closed contact manifold (in any odd dimension) admits an open book decomposition adapted to the contact structure, where the pages are Stein manifolds and the monodromy is a symplectic diffeomorphism. In the case of a 5-dimensional contact manifold, the pages are Stein surfaces. These permit a description via Kirby diagrams, where the attaching circles for the 2-handles are Legendrian knots in the standard contact structure on the boundary $\#_k S^1 \times S^2$ of the 1-handlebody. Provided the monodromy is given as a product of Dehn twists along Lagrangian spheres corresponding to the 2-handles, this too can be encoded in the Kirby diagram.

Since Legendrian knots are faithfully represented by their front projection in the 2-plane, we obtain a description of 5-manifolds in terms of 2-dimensional diagrams.

The combination of stabilisations of the open book with handle slides in the Stein page leads to a couple of moves on such diagrams that do not change the contact 5-manifold. These moves are introduced in Section 4.1, after a brief discussion of open books and their monodromy in Sections 2 and 3. We then present two simple applications of these moves to diagrams without 1-handles. In Section 4.2 we describe an integer family of contact structures on $S^2 \times S^3$ and $S^2 \tilde{\times} S^3$, the non-trivial S^3 -bundle over S^2 . We also give a diagrammatic proof of the diffeomorphism

$$S^2 \tilde{\times} S^3 \# S^2 \tilde{\times} S^3 \cong S^2 \times S^3 \# S^2 \tilde{\times} S^3$$

and its contact analogue. In Section 4.3 we classify contact 5-manifolds that admit Stein fillings made up of a single 0-handle and 2-handles only.

In Section 5 we turn our attention to general subcritically Stein fillable contact 5-manifolds. These can be described by open books with trivial monodromy, whose diagrammatic representations are particularly tractable. We show how to implement the Tietze moves on presentations of the fundamental group as diagrammatic

moves. As an application, we prove that subcritically fillable contact 5-manifolds are classified by their fundamental group, up to connected sums with $S^2 \times S^3$ and $S^2 \tilde{\times} S^3$ (with their standard contact structures), see Theorem 5.3. There is a corresponding classification of the subcritical Stein fillings (Corollary 5.4).

Roughly speaking, these results can be phrased as saying that 6-dimensional compact subcritical Stein manifolds are determined by topological data. This contrasts sharply with the general situation for Stein manifolds. In any even dimension ≥ 8 there are countably many pairwise distinct Stein manifolds of finite type, all diffeomorphic to Euclidean space; this was proved by McLean [30], building on earlier work of Seidel and Smith [34].

In Section 6 we specialise to subcritical Stein fillings of the 5-sphere and give a characterisation of the standard contact structure on S^5 in terms of such fillings. We discuss the relevance of this result in the context of the Andrews–Curtis conjecture concerning presentations of the trivial group.

Finally, in Section 7 we exhibit diagrams for some special simply connected 5-manifolds. These diagrams are then used to show that every 5-dimensional simply connected spin manifold is a double branched cover of the 5-sphere. A further application is a new proof that every simply connected 5-manifold admits a contact structure in each homotopy class of almost contact structures.

2. OPEN BOOK DECOMPOSITIONS

In this section we review the basic aspects of the Giroux correspondence between contact structures and open books. We describe three essential operations on open books: Dehn–Seidel twist, stabilisation, and open book connected sum. In the last part of this section we recall how to compute the homology of open books.

2.1. Open books. Recall that a compact Stein manifold is a compact complex manifold Σ admitting a strictly plurisubharmonic function $f: \Sigma \rightarrow \mathbb{R}$ that is constant on the boundary $\partial\Sigma$ and has no critical points there. Then the exact 2-form $i\partial\bar{\partial}f$ defines a symplectic structure on Σ compatible with the complex structure. Notice that compact Stein manifolds are in particular of **finite type**, i.e. they have finite handlebody decompositions.

According to a fundamental theorem of Giroux [18], any cooriented contact structure on a closed manifold is supported by an open book decomposition of that manifold, where the pages are compact Stein manifolds and the monodromy is symplectic. For the purposes of the present article it suffices to understand how one finds a contact structure adapted to a given open book decomposition. Here we briefly recall this construction.

Let $(\Sigma, \omega = d\lambda)$ be a compact Stein manifold, and let ψ be a symplectomorphism of (Σ, ω) equal to the identity near $\partial\Sigma$. By a lemma of Giroux, cf. [16, Lemma 7.3.4], we may assume without loss of generality that the symplectomorphism is exact, that is, $\psi^*\lambda = \lambda + dh$ for some smooth function $h: \Sigma \rightarrow \mathbb{R}^+$.

The mapping torus

$$A := \Sigma \times \mathbb{R} / (x, \varphi) \sim (\psi(x), \varphi - h(x))$$

carries the contact form $\lambda + d\varphi$. Since ψ equals the identity near $\partial\Sigma$, we can glue A to $B := \partial\Sigma \times D^2$ along their common boundary $\partial\Sigma \times S^1$. In terms of polar coordinates (r, φ) on D^2 , one can define a contact form on B by the ansatz

$$h_1(r) \lambda|_{\partial\Sigma} + h_2(r) d\varphi.$$

With the functions h_1 and h_2 chosen as in Figure 1, this will indeed be a contact form that glues smoothly with $\lambda + d\varphi$ on A , resulting in a contact form α on $M := A \cup_{\partial} B$. This description of the manifold M is called an **open book decomposition**. The codimension 2 submanifold $\partial\Sigma \times \{0\} \subset B \subset M$ is called the **binding** of the open book. Up to diffeomorphism, A can be identified with

$$\Sigma \times [0, 2\pi]/(x, 2\pi) \sim (\psi(x), 0).$$

In terms of this description, the **pages** of the open book are the codimension 1 submanifolds

$$\Sigma \times \{\varphi\} \cup \partial\Sigma \times \{re^{i\varphi} \in D^2 : r \in [0, 1]\},$$

which are diffeomorphic copies of Σ . The map ψ is called the **monodromy** of the open book. For more details see [16, Sections 4.4.2 and 7.3].




FIGURE 1. The functions h_1 and h_2 .

It is not too difficult to see that the resulting contact manifold $(M, \ker \alpha)$ is determined, up to contactomorphism, by Σ and ψ . In fact, it is enough to know the symplectomorphism type of the completion of Σ in the sense of [13], see [18, Proposition 9]. For a compact Stein manifold, the completion is simply the corresponding open Stein manifold. We are therefore justified in denoting this contact manifold $(M, \ker \alpha)$ by $\text{Open}(\Sigma, \psi)$ and call it a **contact open book**. When M is 5-dimensional, the pages Σ of the open book are Stein surfaces, which allow a description in terms of Kirby diagrams. This description of contact 5-manifolds will form the basis of our discussion.

2.2. Dehn–Seidel twists. Let $L \cong S^n$ be a Lagrangian sphere in a compact Stein manifold $(\Sigma, d\lambda)$ of real dimension $2n$. By the Weinstein neighbourhood theorem [37] there is a neighbourhood of L symplectomorphic to the cotangent bundle T^*S^n with its canonical symplectic structure $d\lambda_{\text{can}}$, which is defined as follows. Using Cartesian coordinates $(\mathbf{q}, \mathbf{p}) \in \mathbb{R}^{n+1} \times \mathbb{R}^{n+1}$, we can describe the cotangent bundle $T^*S^n \subset \mathbb{R}^{2n+2}$ by the equations

$$\mathbf{q} \cdot \mathbf{q} = 1 \text{ and } \mathbf{q} \cdot \mathbf{p} = 0;$$

the canonical 1-form is given by $\lambda_{\text{can}} = \mathbf{p} d\mathbf{q}$.

For each $k \in \mathbb{Z}$, one can define a so-called k -fold Dehn twist

$$\tau_k : (T^*S^n, d\lambda_{\text{can}}) \longrightarrow (T^*S^n, d\lambda_{\text{can}})$$

as follows. First consider the normalised geodesic flow σ_t on $T^*S^n \setminus S^n$ given by

$$\sigma_t(\mathbf{q}, \mathbf{p}) = \begin{pmatrix} \cos t & |\mathbf{p}|^{-1} \sin t \\ -|\mathbf{p}| \sin t & \cos t \end{pmatrix} \begin{pmatrix} \mathbf{q} \\ \mathbf{p} \end{pmatrix}.$$

Then set

$$\tau_k(\mathbf{q}, \mathbf{p}) = \sigma_{g_k(|\mathbf{p}|)}(\mathbf{q}, \mathbf{p}),$$

where $r \mapsto g_k(r)$ is a smooth function that interpolates monotonically between $k\pi$ near $r = 0$ and 0 for large r . For $\mathbf{p} = 0$ we read this as $\tau_k(\mathbf{q}, 0) = ((-1)^k \mathbf{q}, 0)$. Then τ_k is an exact symplectomorphism of $(T^*S^n, d\lambda_{\text{can}})$, see [27], equal to the identity for $|\mathbf{p}|$ large. This allows us to regard τ_k as a symplectomorphism of Σ . Viewed this way, τ_k is called a k -fold **Dehn twist** along $L \subset \Sigma$. The map τ_1 is called a right-handed Dehn (or Dehn–Seidel) twist [32, Section 6], cf. [33]; for $n = 1$ this coincides with the classical notion of a Dehn twist.

2.3. Stabilisations. Suppose we are given a contact manifold $\text{Open}(\Sigma, \psi)$ and a properly embedded Lagrangian disc $L \subset \Sigma$ with Legendrian boundary $\partial L \subset \partial \Sigma$. We can construct a Stein manifold Σ' by attaching an n -handle to Σ along ∂L . This new Stein (and hence symplectic) manifold contains a Lagrangian sphere L' , given as the union of L and the core of the n -handle. Let $\tau_{L'}$ be a right-handed Dehn twist along L' . The contact manifold $\text{Open}(\Sigma', \psi \circ \tau_{L'})$ is called a **right-handed stabilisation** of $\text{Open}(\Sigma, \psi)$ along L .

Giroux has announced the following result.

Proposition 2.1 (Giroux). *A right-handed stabilisation of $\text{Open}(\Sigma, \psi)$ does not change its contactomorphism type.* \square

For a detailed proof see [26]; here is the main idea. The Legendrian sphere $\partial L \subset \partial \Sigma$ in the binding of the open book is an isotropic sphere in the ambient contact manifold $\text{Open}(\Sigma, \psi)$. So the attaching of a handle to each page along ∂L may be seen as a contact surgery (in the sense of [11, 38], cf. [16]) along this isotropic sphere, where the necessary trivialisation of the symplectic normal bundle of ∂L in $\text{Open}(\Sigma, \psi)$ is provided by the trivialisation of the normal bundle of the binding $\partial \Sigma$ in the open book. Performing a right-handed Dehn twist along the Lagrangian sphere L' in the new page Σ' may be regarded as a Legendrian surgery on $L' \subset \text{Open}(\Sigma', \psi)$, see Theorem 7.2 below. This second surgery can be seen to cancel the first, even on the level of symplectic handle attachments.

2.4. Book connected sum. A connected sum operation for open books has been described in [28]. Let $\text{Open}(\Sigma_i, \psi_i)$, $i = 1, 2$, be two open books of dimension $2n + 1$. We write $\partial \Sigma_i$ for the binding. One can form the connected sum of these two open books by cutting out discs D_i^{2n+1} embedded in such a way that the pair $(D_i^{2n+1}, D_i^{2n+1} \cap \partial \Sigma_i)$ is diffeomorphic to the standard disc pair (D^{2n+1}, D^{2n-1}) . Then the connected sum

$$\text{Open}(\Sigma_1, \psi_1) \# \text{Open}(\Sigma_2, \psi_2)$$

is diffeomorphic to

$$\text{Open}(\Sigma_1 \natural \Sigma_2, \psi_1 \natural \psi_2),$$

where $\Sigma_1 \natural \Sigma_2$ denotes the boundary connected sum of the pages, and $\psi_1 \natural \psi_2$ is the obvious map on this boundary connected sum that restricts to ψ_i on Σ_i .

This construction is compatible with the contact structures on these open books if $\Sigma_1 \natural \Sigma_2$ is interpreted as the boundary connected sum of Stein or symplectic manifolds as in [11, 38], see [24].

2.5. Homology of open books. The homology of an open book $M = A \cup_{\partial} B$ can easily be computed in terms of the homology of the page Σ and the action of the monodromy ψ on homology. This will turn out to be especially useful in the discussion of simply connected 5-manifolds, whose diffeomorphism type is determined by homological data.

We only consider this 5-dimensional case, and we assume for simplicity that Σ is composed of one 0-handle and only 2-handles. With Q denoting the intersection form on $H_2(\Sigma)$, one finds that $H_1(\partial\Sigma) \cong \text{coker } Q$, cf. [5, pp. 427–8]. This is the essential part of the homology of $B \simeq \partial\Sigma$.

We briefly recall the argument for this statement about $H_1(\partial\Sigma)$, since we need explicit information about the homological generators in the proof of Proposition 7.1 below. The homology $H_2(\Sigma)$ is isomorphic to \mathbb{Z}^m , with m denoting the number of 2-handles, freely generated by the surfaces obtained by gluing a Seifert surface of each attaching circle in $S^3 = \partial D^4$ with the core disc of the corresponding 2-handle. The relative homology group $H_2(\Sigma, \partial\Sigma)$ is likewise isomorphic to \mathbb{Z}^m ; here the generators are meridional discs of the attaching circles whose boundary lies on the boundary of the 2-handle. The homology $H_1(\partial\Sigma)$ is generated by the meridians of the attaching circles, i.e. the images of the generators of $H_2(\Sigma, \partial\Sigma)$ under the boundary homomorphism. In terms of the described generators, the homomorphism $H_2(\Sigma) \rightarrow H_2(\Sigma, \partial\Sigma)$ is given by the intersection form Q . The result $H_1(\partial\Sigma) \cong \text{coker } Q$ now follows from the homology exact sequence of the pair $(\Sigma, \partial\Sigma)$.

The homology of the Σ -bundle A over S^1 can be computed using the Wang sequence [31, Lemma 8.4]. The relevant part of this sequence looks as follows.

$$\dots \longrightarrow H_3(A) \longrightarrow H_2(\Sigma) \xrightarrow{\psi_* - \text{id}} H_2(\Sigma) \longrightarrow H_2(A) \longrightarrow H_1(\Sigma) \longrightarrow \dots$$

This information on A and B can be combined to obtain the homology of M via the Mayer–Vietoris sequence of the decomposition $M = A \cup_{\partial} B$.

3. MONODROMY

The symplectomorphism group of a given Stein manifold Σ is not known, in general. Therefore we restrict our attention to symplectomorphisms that can be written as compositions of Dehn twists along Lagrangian spheres $L \subset \Sigma$.

3.1. Action of Dehn twists on homology. Let Σ be a symplectic manifold of dimension $2n$ and $L \subset \Sigma$ a Lagrangian sphere. Write $[L] \in H_n(\Sigma)$ for the homology class represented by L . As before, we denote the intersection form on $H_n(\Sigma)$ by Q . With this notation, the homomorphism on homology induced by a right-handed Dehn twist τ_L along L is given by

$$\begin{aligned} (\tau_L)_*: H_n(\Sigma) &\longrightarrow H_n(\Sigma) \\ u &\longmapsto u + (-1)^{1+n(n+1)/2} Q([L], u) \cdot [L]. \end{aligned}$$

This can be seen as follows. First suppose u is a homology class represented by a closed, oriented submanifold of Σ . This submanifold can be isotoped to intersect L transversely in a finite number of points. Think of a neighbourhood of L in Σ as T^*L ; the submanifold representing u may then be assumed to intersect T^*L in a finite number of fibres. On the level of homology, a Dehn twist along L adds $\pm[L]$ to u for each of these intersections. The dependence of the sign in $\pm[L]$ on the

sign of the transverse intersection can be computed explicitly in the local model; we leave this to the reader.

The same argument applies for any singular cycle u , as can be verified by using the intersection theory developed in [35, §73] for pairs of singular chains of complementary dimensions in a given manifold.

3.2. Monodromy and Stein filling. For the most part we shall be concerned with open books that have trivial monodromy. The following proposition tells us that this is a reasonably interesting class of manifolds to consider. Recall that a contact manifold (M, ξ) is said to be **Stein fillable** if it arises as the boundary of a compact Stein manifold, with ξ given as the complex tangencies on the boundary. As is well known, a Stein manifold of real dimension $2n + 2$ has a handle decomposition with handles up to index $n + 1$ only. The filling is called **subcritical** if there are no handles of index $n + 1$.

Proposition 3.1. *A contact structure ξ on a closed manifold M is supported by an open book with trivial monodromy if and only if (M, ξ) is subcritically Stein fillable.*

Proof. Suppose $(M, \xi) = \text{Open}(\Sigma, \text{id})$. Then

$$M = \Sigma \times S^1 \cup_{\partial} \partial\Sigma \times D^2 = \partial(\Sigma \times D^2).$$

If J is the complex structure on Σ and $f: \Sigma \rightarrow \mathbb{R}$ a strictly plurisubharmonic function, then $\Sigma \times D^2 \ni (p, x, y) \mapsto f(p) + x^2 + y^2$ defines a strictly plurisubharmonic function on $\Sigma \times D^2$ with respect to the obvious complex structure (J, i) , and this function has no critical points of maximal index.

The converse is due to Cieliebak [6], who showed that any subcritical Stein manifold is equivalent to a product $\Sigma \times \mathbb{C}$. \square

According to a result of Loi–Piergallini and Giroux [18], cf. [17], the 3-dimensional Stein fillable contact manifolds are characterised as those contact manifolds that admit a supporting open book whose monodromy is a composition of right-handed Dehn twists. In higher dimensions, this condition on the monodromy is still sufficient for Stein fillability.

Proposition 3.2. *If a contact manifold is supported by an open book whose monodromy is a composition of right-handed Dehn twists along Lagrangian spheres on the pages, then it is Stein fillable.* \square

This appears to be a folklore theorem; a proof can be found in [26], see also [1]. The main idea is that a Lagrangian sphere on a page can be made Legendrian with respect to the contact structure on the open book; a right-handed Dehn twist then corresponds to a Legendrian surgery, which preserves Stein fillability.

In [4] left-handed Dehn twists are used to construct algebraically overtwisted contact manifolds, i.e. contact manifolds with vanishing contact homology, which are conjecturally not strongly fillable (and hence not Stein fillable).

4. DIAGRAMS FOR 5-MANIFOLDS

We now want to give diagrammatic representations of 5-dimensional contact open books. Here the page is a Stein surface, which can be described by a Kirby diagram. According to a result of Eliashberg [11], worked out further by Gompf [20], any compact Stein surface can be obtained from the 4-disc in \mathbb{C}^2 by attaching a finite number of 1- and 2-handles, where each 2-handle is attached along a Legendrian

knot (in the standard contact structure on the relevant boundary 3-manifold), with framing -1 relative to the contact framing of that Legendrian knot. Conversely (this is the easier part), this recipe always produces a Stein surface.

The boundary 3-manifold obtained by attaching k 1-handles to the 4-disc is the connected sum $\#_k(S^1 \times S^2)$ with its unique tight contact structure; the attaching circles for the 2-handles form a Legendrian link in this contact manifold. In other words, the Kirby diagram of a Stein surface consists of a finite collection of pairs of attaching balls for the 1-handles in $\mathbb{R}^3 \subset S^3$ with its standard contact structure $\xi_{\text{st}} = \ker(dz + x dy)$, and the front projection to the yz -plane of a Legendrian link. For more information see [20] and [21].

In order to obtain a representation of the contact manifold $\text{Open}(\Sigma, \psi)$, one also needs to encode the monodromy ψ in the diagram. This is possible in special cases. For instance, if one of the Legendrian knots in the Kirby diagram bounds in an obvious way a Lagrangian disc in the Stein surface Σ , the union of this disc with the core disc of the 2-handle is a Lagrangian sphere, and we can speak of a Dehn twist along this sphere. Beware that different such Dehn twists will not, in general, commute with each other. In cases where the order of the Dehn twists is inessential, we simply write the relevant Dehn twist next to the Legendrian knot representing the corresponding Lagrangian sphere. In particular, stabilisations can be so encoded, provided one understands the monodromy of the given diagram.

By Section 2.4, the diagram for the connected sum of two contact open books is simply given by drawing the attaching balls and Legendrian attaching circles in a single diagram, separated by a hyperplane.

4.1. Handle moves. We illustrate the use of (de-)stabilisations in the following two handle moves on a diagram of $\text{Open}(\Sigma, \psi)$. These moves do not change the contactomorphism type of $\text{Open}(\Sigma, \psi)$.

4.1.1. Move I. Assume we are given a Kirby diagram for Σ that includes a Legendrian knot K with the property that the monodromy ψ equals the identity on the handle attached along K . Add a standard Legendrian unknot K_0 with Thurston–Bennequin invariant $\text{tb}(K_0) = -1$ to the diagram, unlinked with the given Legendrian link and not passing over 1-handles. This K_0 bounds an obvious Lagrangian disc, so a right-handed Dehn twist τ along the corresponding 2-sphere (see the following remark) amounts to a stabilisation of $\text{Open}(\Sigma, \psi)$.

Remark 4.1. It is not accidental that the knot K_0 we use for the stabilisation is chosen to have $\text{tb}(K_0) = -1$. Write L' for the Lagrangian 2-sphere obtained by gluing the Lagrangian disc L bounded by K_0 to the core disc of the handle attached along K_0 . A neighbourhood of L' looks like T^*S^2 , so L' has self-intersection -2 . If we push the core disc along its boundary in the direction of the surgery framing, we obtain a disjoint disc. Hence, pushing L in that direction along its boundary leads to a disc having intersection -2 with L . If L is topologically isotopic to a disc in $\partial\Sigma$ (as in the case of K_0), this means that the linking number of K_0 with its push-off in the direction of the surgery framing equals -2 . Since the surgery framing is obtained from the contact framing by adding a negative twist, we conclude $\text{tb}(K_0) = -1$.

Beware that $\text{tb}(K_0)$ may take other values, with K_0 regarded as a knot in (S^3, ξ_{st}) , if the Lagrangian disc bounded by K_0 goes over 2-handles.

Move I is now performed in three steps, see Figure 2:

- (i) Stabilise $\text{Open}(\Sigma, \psi)$ by adding (K_0, τ) to the diagram as described.
- (ii) Perform a handle slide of K over K_0 ; this is possible via a Legendrian isotopy, see [8, Proposition 1].
- (iii) Destabilise by removing (K_0, τ) from the picture.




FIGURE 2. Move I.

Remark 4.2. A Legendrian isotopy of a Legendrian submanifold in a contact manifold $(M, \xi = \ker \alpha)$ extends to a contact isotopy ϕ_t of (M, ξ) , cf. [16, Theorem 2.6.2], and thence to an \mathbb{R} -invariant symplectic isotopy Φ_t of the symplectisation $(\mathbb{R} \times M, d(e^s \alpha))$. Write $\phi_t^* \alpha = e^{h_t} \alpha$ with a smooth function $h_t: M \rightarrow \mathbb{R}$ and set $\Phi_t(s, x) = (s - h_t(x), \phi_t(x))$. Then $\Phi_t^*(e^s \alpha) = e^s \alpha$, which implies that Φ_t is a Hamiltonian isotopy. By cutting off the corresponding Hamiltonian function, this isotopy may be assumed to coincide with Φ_t on $\{r\} \times M$ and to be stationary outside $(0, R) \times M$ for $R > r > 0$ sufficiently large.

Hence, if (M, ξ) has a strong symplectic filling (W, ω) , this cut off symplectic isotopy extends to an isotopy of the symplectic completion of (W, ω) in the sense of [13]. So the time-1 map Φ_1 may be regarded as a symplectomorphism of the symplectic completion of (W, ω) . Alternatively, we may view Φ_1 as a symplectomorphism of the filling $(W, \omega) \cup ([0, r] \times M, d(e^s \alpha))$ of (M, ξ) onto its image, which on the boundary induces the contactomorphism ϕ_1 .

It will be in this sense that we interpret step (ii) of move I as a symplectomorphism of the filling.

The effect of this move I is to replace K by its double Legendrian stabilisation $S_+ S_- K$, i.e. a Legendrian knot which has two additional zigzags (one up, one down). Notice that the contact framing of $S_+ S_- K$ differs from that of K by two negative (i.e. left-handed) twists; if K is homologically trivial this means $\text{tb}(S_+ S_- K) = \text{tb}(K) - 2$. The rotation number rot does not change under this move.

One of the potential uses of move I is the following. As shown by Fuchs and Tabachnikov [14, Theorem 4.4], cf. [10], any topological isotopy of Legendrian knots can be turned into a Legendrian isotopy of suitable Legendrian stabilisations. Thus, after a repeated application of move I, two topologically isotopic Legendrian knots with the same rotation number will become Legendrian isotopic.

4.1.2. *Move II.* Our second move gives us even greater flexibility, for it allows us to change crossings of Legendrian knots K, K' (or a self-crossing of K) in the Kirby diagram for Σ , at the cost of replacing K by its fourfold Legendrian stabilisation $S_+^2 S_-^2 K$. The move is a combination of move I with a further (de-)stabilisation and handle slide; a pictorial description is given in Figure 3.




FIGURE 3. Move II.

The first two steps in Figure 3 can be performed even if the monodromy along K is not the identity, but the destabilisation may not be possible. Move II without the final destabilisation still allows us to change a given Kirby diagram into one where the Legendrian link is topologically a link of unknots.

4.2. **Diagrams for $S^2 \times S^3$ and $S^2 \tilde{\times} S^3$.** We now use moves I and II from the preceding section to give a characterisation of the contact manifolds $\text{Open}(\Sigma, \text{id})$ where Σ has a Kirby diagram consisting of a single knot K .

Since $\pi_1(\text{SO}(4)) = \mathbb{Z}_2$, there are exactly two S^3 -bundles over S^2 up to bundle isomorphism, the trivial and the non-trivial one. Their total spaces are non-diffeomorphic and are denoted by $S^2 \times S^3$ and $S^2 \tilde{\times} S^3$, respectively.

If M is a closed simply connected 5-manifold with $H_2(M; \mathbb{Z}) \cong \mathbb{Z}$, then by Barden's classification [3] one has

$$M \cong \begin{cases} S^2 \times S^3 & \text{if } w_2(M) = 0, \\ S^2 \tilde{\times} S^3 & \text{if } w_2(M) \neq 0. \end{cases}$$

Proposition 4.3. *Let K be an oriented Legendrian knot in (S^3, ξ_{st}) . Then the contact manifold $(M, \xi) = \text{Open}(\Sigma, \text{id})$, with Σ the Stein surface given by the Kirby diagram consisting of K only, is one of the following:*

- $(S^2 \times S^3, \xi_{|\text{rot}(K)|})$ if $\text{rot}(K)$ is even,
- $(S^2 \tilde{\times} S^3, \xi_{|\text{rot}(K)|})$ if $\text{rot}(K)$ is odd.

As the notation suggests, the contact structure $\xi_{|\text{rot}(K)|}$ depends, up to diffeomorphism, only on $|\text{rot}(K)|$.

Proof. Using move II we can untie any Legendrian knot without changing its rotation number, so we may assume that K is an unknot. Then Σ is a 2-disc bundle over S^2 . The zero section S^2 of this bundle, oriented in such a way that the disc $D^2 \subset S^2$ bounded by K in S^3 is oriented consistently with K , represents the positive generator of $H_2(\Sigma)$. Since the monodromy is the identity, we have $M \cong \partial(\Sigma \times D^2)$. This means that M is the boundary of a 4-disc bundle over S^2 , i.e. an S^3 -bundle.

Next we determine the first Chern class of ξ . The cohomology exact sequence for the pair $(\Sigma \times D^2, \partial(\Sigma \times D^2))$ shows that the inclusion $i: \partial(\Sigma \times D^2) \rightarrow \Sigma \times D^2$ induces an isomorphism $i^*: H^2(\Sigma \times D^2) \rightarrow H^2(\partial(\Sigma \times D^2))$. The contact structure ξ is given by the complex tangencies of $\partial(\Sigma \times D^2)$ (after the smoothing of corners), and the complementary complex line bundle in $T(\Sigma \times D^2)|_{\partial(\Sigma \times D^2)}$ is trivial, so we have $c_1(\xi) = i^*c_1(\Sigma \times D^2)$. The class $c_1(\Sigma \times D^2)$ can be naturally identified with $c_1(\Sigma)$, which by [20, Proposition 2.3] equals $\text{rot}(K)h$, where h is the positive generator of $H^2(\Sigma)$. Thus, under the mentioned identifications we have $c_1(\xi) = \text{rot}(K)h$.

The second Stiefel–Whitney class $w_2(M)$ equals the mod 2 reduction of $c_1(\xi)$. So the diffeomorphism type of M is as claimed.

It remains to show that the contact structure is determined by the absolute value of its first Chern class. Thus, let K_1 and K_2 be two oriented Legendrian unknots with $\text{rot}(K_1) = \pm \text{rot}(K_2)$. The orientation of these knots has no bearing on the resulting contact manifold, only on the designation of one of the generators of H_2 as the positive one. So we may orient K_1 and K_2 such that $\text{rot}(K_1) = \text{rot}(K_2)$. The parity condition $\text{tb}(K_i) + \text{rot}(K_i) \equiv 1 \pmod{2}$, cf. [16], allows us to achieve in addition that $\text{tb}(K_1) = \text{tb}(K_2)$ after a repeated application of move I to one of the two knots. From the classification of Legendrian unknots by Eliashberg and Fraser [12] it follows that K_1 and K_2 are then Legendrian isotopic, so the corresponding 5-dimensional contact manifolds are contactomorphic. \square

Remark 4.4. Both $S^2 \times S^3$ and $S^2 \tilde{\times} S^3$ admit orientation-preserving diffeomorphisms that act as minus the identity on H^2 [3, Theorem 2.2], and orientation-reversing diffeomorphisms that act as the identity on H^2 [16, p. 399]. This explains why the orientation of M and the sign of the rotation number are irrelevant for the diffeomorphism classification.


From Barden’s classification it follows that $S^2 \tilde{\times} S^3 \# S^2 \tilde{\times} S^3$ is diffeomorphic to $S^2 \times S^3 \# S^2 \tilde{\times} S^3$. This can also be seen by diagram moves, which proves a little more.

Proposition 4.5. *With the notation from the preceding proposition, we have a contactomorphism*

$$(S^2 \tilde{\times} S^3, \xi_{2m+1}) \# (S^2 \tilde{\times} S^3, \xi_1) \cong (S^2 \times S^3, \xi_{2n}) \# (S^2 \tilde{\times} S^3, \xi_1)$$

for any $m, n \in \mathbb{N}_0$.

Proof. The proof consists of sliding the 2-handle corresponding to ξ_{2m+1} over the one corresponding to ξ_1 ; Figure 4 shows this for $m = 0$, $n = 1$. From (i) to (ii) one performs a handle slide; from (ii) one gets to (iii) by using move II to change crossings and the reverse of move I to perform double Legendrian destabilisations. Observe that the rotation numbers add under a handle slide. If we perform a handle subtraction instead of a handle addition (i.e. a handle slide with the orientation of the shark on the right-hand side reversed), we obtain $n = 0$. \square


 FIGURE 4. The connected sum $S^2 \tilde{\times} S^3 \# S^2 \tilde{\times} S^3$.

4.3. Subcritical fillings without 1-handles. We are now in a position to classify 5-dimensional contact manifolds that admit subcritical Stein fillings without 1-handles. The first Chern class $c_1(\xi)$ of a contact structure on a simply connected 5-manifold determines the homotopy class of the underlying reduction of the structure group to $U(2) \times 1$. When there exists a subcritical Stein filling, it actually determines the contact structure.

Theorem 4.6. *Let (M_i, ξ_i) , $i = 1, 2$, be two simply connected contact 5-manifolds that admit subcritical Stein fillings without 1-handles. If there is an isomorphism $\phi: H^2(M_1) \rightarrow H^2(M_2)$ such that $\phi(c_1(\xi_1)) = c_1(\xi_2)$, then (M_1, ξ_1) and (M_2, ξ_2) are contactomorphic.*

Proof. The assumption on the existence of subcritical Stein fillings implies that the (M_i, ξ_i) can be realised as contact open books whose monodromy is the identity. With the help of moves I and II, the corresponding diagrams (which, by assumption, contain no 1-handles) can be turned into a collection of unlinked Legendrian unknots. If one so wishes, one can arrange the rotation number of at most one of the knots to be odd by the argument in the preceding proposition. In particular, we see that the M_i are diffeomorphic to a connected sum of copies of $S^2 \times S^3$, possibly with one additional summand $S^2 \tilde{\times} S^3$.

Fixing an ordering of the unknots in the respective diagram, and an orientation for each unknot, amounts to fixing an identification of $H^2(M_i)$ with \mathbb{Z}^k , where $k \in \mathbb{N}$ is the number of unknots in either diagram. Then ϕ may be regarded as an element of $GL(k, \mathbb{Z})$. By elementary row and column operations (over the integers), this matrix can be converted to the identity matrix. So it suffices to show that these elementary operations can be effected by a change in the respective diagram. The condition $\phi(c_1(\xi_1)) = c_1(\xi_2)$ implies (by Proposition 4.3 and its proof) that between the two new diagrams there is a bijective pairing of Legendrian unknots with the same rotation number, which amounts to the claimed contactomorphism.

Here are the diagrammatic realisations of the elementary row and column operations, the former being changes in the diagram for M_2 , the latter in that of M_1 .

1. Add a row/column to another one: this amounts to a handle slide, see Figure 5.
2. Multiply a row/column by -1 : a change of sign of one of the generators of \mathbb{Z}^k simply amounts to changing the orientation on the corresponding unknot. This reverses the sign of the rotation number of that unknot. \square

Remark 4.7. M.-L. Yau [39] has shown that the contact homology of a subcritically Stein fillable contact manifold (M, ξ) is determined by the homology of the manifold, provided $c_1(\xi)$ evaluates to zero on $\pi_2(M)$ (i.e. homology classes represented by maps $S^2 \rightarrow M$). In particular, if we write the subcritical Stein filling of a 5-dimensional contact manifold as $\Sigma \times D^2$, then $HC_2(M, \xi) \cong H_2(\Sigma) \cong H_2(M)$. In view of this result, the theorem above says that contact homology is a complete




FIGURE 5. Sliding the handles to get the desired map on cohomology.

invariant for 5-dimensional contact manifolds (M, ξ) that admit a subcritical Stein filling without 1-handles and with $c_1(\xi) = 0$.

5. DIAGRAMS FOR SUBCRITICALLY FILLABLE CONTACT 5-MANIFOLDS

In this section we prove a result that goes some way towards classifying subcritically fillable contact 5-manifolds and their Stein fillings. By realising Tietze moves on group presentations via handle moves in Kirby diagrams, we show that subcritically fillable contact 5-manifolds are determined, up to a connected sum with copies of $S^2 \times S^3$ and $S^2 \tilde{\times} S^3$, by their fundamental group; see Theorem 5.3 for the precise formulation. The corresponding classification of 6-dimensional subcritical Stein manifolds up to symplectomorphism is formulated in Corollary 5.4.

5.1. Tietze moves and Legendrian isotopies. Let $\langle g_1, \dots, g_k | r_1, \dots, r_l \rangle$ be a finite presentation of a group G . We can then realise this group G as the fundamental group of a Stein surface by associating a 1-handle with each of the generators g_1, \dots, g_k , and an oriented Legendrian attaching circle with each of the relations r_1, \dots, r_l . Our conventions for the translation from a group presentation to a Kirby diagram in standard form as in [20] are as follows.

- (i) The 1-handles are represented by horizontal pairs of attaching balls.
- (ii) A Legendrian curve going over the 1-handle corresponding to a generator g in the way shown in Figure 6 is read as the letter g in the word represented by that curve.
- (iii) The relations, which are words in the generators, are translated into a curve by reading the word from left to right.

5.1.1. Equivalent diagrams. Since we are dealing with subcritical Stein fillings, we may restrict our attention to open books having trivial monodromy. So we are free to use the moves introduced in Section 4.1; these do not change the contact manifold or its filling.



FIGURE 6. Attaching circle going over a 1-handle.

- (i) Move I: replace a Legendrian attaching knot K by its double Legendrian stabilisation S_+S_-K .
- (ii) Move II: change crossings of the Legendrian attaching circles at the price of adding Legendrian stabilisations.

In particular, the crossings of the Legendrian attaching circles are ultimately of no importance.

In order to realise all Tietze moves on group presentations — these moves will be described presently — we need to allow one further change in the diagram:

- (iii) Add an unlinked standard Legendrian unknot K_0 with $\text{tb}(K_0) = -1$ to the diagram.

On the level of contact 5-manifolds, this third move corresponds to taking the connected sum with $(S^2 \times S^3, \xi_0)$, see Proposition 4.3.

We remark that the particular points where the attaching circles go over the 1-handles are irrelevant. This is illustrated in Figure 7 (without loss of generality, strand 2 is assumed to be Legendrian stabilised).




FIGURE 7. Changing the order of attaching points.

5.1.2. *Tietze moves.* According to the Tietze theorem [7, pp. 43–4], any two finite presentations of a given group are related by a sequence of the following two moves.

T 1. Add or remove a relation s that is a so-called consequence of the other relations:

$$\langle g_1, \dots, g_k | r_1, \dots, r_l \rangle \rightsquigarrow \langle g_1, \dots, g_k | r_1, \dots, r_l, s \rangle$$

For the relation s to be a **consequence** of the relations r_1, \dots, r_l means that s (which is an element of the free group generated by g_1, \dots, g_k) is contained in every normal subgroup that contains r_1, \dots, r_l .

This move, read from left to right, can be written as a sequence of the following submoves.

- (i) Double a relation or add an inverse of a relation:

$$\langle g_1, \dots, g_k | r_1, \dots, r_l \rangle \rightsquigarrow \langle g_1, \dots, g_k | r_1, r_1^{\pm 1}, \dots, r_l \rangle$$

- (ii) Conjugate a relation by a generator:

$$r \rightsquigarrow grg^{-1} \text{ or } g^{-1}rg$$

- (iii) Replace one relation by its product with another relation:

$$r_1, r_2 \rightsquigarrow r_1, r_1 r_2 \text{ or } r_1, r_2 r_1$$

T 2. Add or remove a generator g and a relation $g = w$ expressing g as a word w in the other generators and their inverses:

$$\langle g_1, \dots, g_k | r_1, \dots, r_l \rangle \rightsquigarrow \langle g_1, \dots, g_k, g | r_1, \dots, r_l, gw^{-1} \rangle$$

5.1.3. *Realising the Tietze moves.* We now want to show that these moves correspond to Legendrian isotopies of the attaching circles and handle cancellations, modulo the equivalences described in Section 5.1.1.

T 1. (i) A relation represented by a Legendrian attaching circle K can be doubled by adding the Legendrian push-off of K to the diagram. Giving this push-off the opposite orientation amounts to adding the inverse relation. By [8, Proposition 2], this Legendrian push-off is Legendrian isotopic to a meridian of K . With the help of crossing changes we can disentangle this meridian from the rest of the diagram. In other words, we can double a relation by first adding a standard Legendrian unknot to the diagram, i.e. by taking a connected sum with $(S^2 \times S^3, \xi_0)$, and then turning this into a push-off of K with the help of moves I (and its inverse) and II.

T 1. (ii) The Legendrian knot representing grg^{-1} as shown in Figure 8 is Legendrian isotopic to that representing r : first move the cusp formed by the strands 1 and 4 over the 1-handle g , then perform a first Reidemeister move.

In Figure 8 we have indicated a base point that one needs to fix in order to set up a one-to-one correspondence between loops at this base point and words in the generators. For the Tietze move, however, one only needs to check that attaching a handle along r has the same effect as attaching it along grg^{-1} , and for that the described Legendrian isotopy is not required to fix the base point

T 1. (iii) This corresponds to a handle slide or second Kirby move in the sense of [8].

T 2. In Figure 9 we have indicated the relation $g = w$, where the word w is supposed to be given by a curve that may go over the 1-handles corresponding to the generators g_1, \dots, g_k . We need to show that the generator g and this relation can be cancelled by diagram moves.

Slide the right-hand attaching ball of the 1-handle g along the Legendrian curve representing the word w^{-1} ; this can be done via a contact isotopy. In the process, we “accumulate” cusps. See Figure 10 for an example.





 FIGURE 8. The Tietze move **T 1** (ii).

 FIGURE 9. The relation $g = w$.


FIGURE 10. Sliding the attaching ball of a 1-handle.

Positive and negative Legendrian stabilisations can be removed in pairs using the light bulb trick [8, Figure 21]. So ultimately we obtain a diagram as shown in Figure 11, cf. [9, Section 5].




FIGURE 11. A cancelling handle pair.

Then the 1-handle g and the 2-handle attached along the Legendrian circle going once over the 1-handle form a cancelling pair. *Topologically* this follows because the attaching circle of the 2-handle intersects the belt sphere of the 1-handle in exactly one point, see [21, Proposition 4.2.9]. In fact, this diagram with a cancelling handle pair also represents the standard 4-disc *symplectically*, see [26]. In the given dimension, this is also a consequence of the deep result of Gromov [22, p. 311], cf. [29, Theorem 9.4.2], that any strong symplectic filling of (S^3, ξ_{st}) not containing homologically non-trivial 2-spheres (in particular a filling known to be topologically a disc) is *symplectomorphic* to the 4-disc.

5.2. Classification of subcritically fillable contact 5-manifolds. As a first step towards showing how the fundamental group determines subcritically fillable contact 5-manifolds, the following lemma tells us that two diagrams representing isomorphic fundamental groups can be turned into two diagrams giving identical group presentations.

Lemma 5.1. *Let (M, ξ) and (M', ξ') be two subcritically Stein fillable contact 5-manifolds with $\pi_1(M) \cong \pi_1(M')$. Then there are $k, k' \in \mathbb{N}_0$ and diagrams D, D' for*

$$(M, \xi) \# k(S^2 \times S^3, \xi_0) \quad \text{and} \quad (M', \xi') \# k'(S^2 \times S^3, \xi_0),$$

respectively, such that the presentations of $\pi_1(M), \pi_1(M')$ determined by D, D' are identical.

Proof. Write $(M, \xi) = \text{Open}(\Sigma, \text{id})$, so that the subcritical Stein filling is given by $\Sigma \times D^2$, and similarly for M' . The theorem of Seifert and van Kampen, applied to the decomposition

$$M = \Sigma \times S^1 \cup_{\partial} \partial \Sigma \times D^2,$$

shows that $\pi_1(M) \cong \pi_1(\Sigma)$. So any choice of Kirby diagram for Σ and Σ' gives rise to two presentations of the same group.

Now invoke Tietze's theorem and our implementation of the Tietze moves. Since we merely want to turn both group presentations into identical ones by appropriate changes in the diagrams, rather than converting one presentation to the other, we need the Tietze move **T 1** only going from left to right, i.e. we do not care about accumulating redundant relations. As we saw in Section 5.1.3, the move **T 1** (i) amounts to a connected sum with $(S^2 \times S^3, \xi_0)$; the other moves do not change the contact manifold. \square

This lemma does not say anything about the contact framings of the Legendrian knots in the diagrams D, D' . In particular, we cannot expect that $M \# k(S^2 \times S^3)$ and $M' \# k'(S^2 \times S^3)$ are diffeomorphic, in general. The most simple example would be $M = S^2 \times S^3$ and $M' = S^2 \tilde{\times} S^3$, where we could take $k = k' = 0$.

The following lemma is the key to showing that a summand $(S^2 \tilde{\times} S^3, \xi_1)$ will give us complete control over the contactomorphism type of the resulting manifold.

Lemma 5.2. *Let Σ be a Stein surface containing at least one 2-handle attached along a Legendrian knot K . Let Σ_{\pm} be the Stein surface where this one attaching circle is replaced by its Legendrian stabilisation $S_{\pm}K$. Then*

$$\text{Open}(\Sigma, \text{id}) \# (S^2 \tilde{\times} S^3, \xi_1) \cong \text{Open}(\Sigma_{\pm}, \text{id}) \# (S^2 \tilde{\times} S^3, \xi_1).$$

Proof. The contact manifold $\text{Open}(\Sigma, \text{id}) \# (S^2 \tilde{\times} S^3, \xi_1)$ is represented by a diagram for Σ with one additional shark. After a handle slide of K over the shark (with one or the other orientation), and by applying moves I and II, we obtain a diagram with K replaced by $S_{\pm}K$, see Figure 12. In (i) we have added a shark to the given diagram, i.e. formed the connected sum with $(S^2 \tilde{\times} S^3, \xi_1)$. We then perform a handle slide over the shark to obtain (ii). From there one gets to (iii) and (iv) as in Figure 4. \square




FIGURE 12. Handle slide turning K into $S_{\pm}K$.

Here is our main classification result.

Theorem 5.3. *Let (M, ξ) and (M', ξ') be two subcritically Stein fillable contact 5-manifolds with $\pi_1(M) \cong \pi_1(M')$. Then there are $k, k' \in \mathbb{N}_0$ with $k - k' = \text{rank } H_2(M') - \text{rank } H_2(M)$ such that*

$$(M, \xi) \# k(S^2 \times S^3, \xi_0) \# (S^2 \tilde{\times} S^3, \xi_1) \cong (M', \xi') \# k'(S^2 \times S^3, \xi_0) \# (S^2 \tilde{\times} S^3, \xi_1).$$

Proof. By Lemma 5.1 there are $k_0, k'_0 \in \mathbb{N}_0$ such that $(M, \xi) \# k_0(S^2 \times S^3, \xi_0)$ and $(M', \xi') \# k'_0(S^2 \times S^3, \xi_0)$ are represented by diagrams that give identical presentations of $\pi_1(M) \cong \pi_1(M')$. Notice that the two diagrams may contain Legendrian knots that do not go over the 1-handles and hence do not contribute to the fundamental group or its presentation.

With move II we can change the crossings of the Legendrian attaching circles, so we may assume that the two diagrams are topologically identical, up to a finite number of unlinked unknots. These unlinked unknots correspond to summands $S^2 \times S^3$ or $S^2 \tilde{\times} S^3$ with one of the standard contact structures described in Proposition 4.3.

By the theorem of Fuchs and Tabachnikov [14, Theorem 4.4], this topological isotopy can be realised as a Legendrian isotopy of suitable stabilisations of the Legendrian knots. As shown in the preceding lemma, a summand $(S^2 \tilde{\times} S^3, \xi_1)$ enables us to perform such Legendrian stabilisations.

Finally, Proposition 4.5 allows us, up to contactomorphism, to turn all but one summand $(S^2 \tilde{\times} S^3, \xi_1)$ into $(S^2 \times S^3, \xi_0)$. \square

It is possible to combine this argument with that for Theorem 4.6 in order to formulate a more precise classification result that involves conditions on the first Chern class, but we have opted for the more transparent formulation of the statement.

The moves in the proof of Theorem 5.3 all extend to symplectomorphisms of the filling, so the following corollary is immediate. Here we write $(S^2 \times D^4, \omega_0)$ for the standard filling of $(S^2 \times S^3, \xi_0)$, and $(S^2 \tilde{\times} D^4, \omega_1)$ for that of $(S^2 \tilde{\times} S^3, \xi_1)$. ‘Symplectomorphism’ is to be understood in the sense of Remark 4.2.

Corollary 5.4. *Let W and W' be two compact subcritical Stein manifolds of dimension 6 with $\pi_1(W) \cong \pi_1(W')$. Then there are $k, k' \in \mathbb{N}_0$ with $k - k' = \text{rank } H_2(W') - \text{rank } H_2(W)$ such that the boundary connected sums*

$$W \natural k(S^2 \times D^4, \omega_0) \natural (S^2 \tilde{\times} D^4, \omega_1)$$

and

$$W' \natural k'(S^2 \times D^4, \omega_0) \natural (S^2 \tilde{\times} D^4, \omega_1)$$

are symplectomorphic. \square

Remark 5.5. It seems feasible to prove Theorem 5.3 by working directly with the subcritical filling. The fundamental group of the 6-dimensional subcritical Stein manifold $\Sigma \times D^2$, which equals that of M , has a presentation with generators given by the 1-handles and relations given by the attaching circles of the 2-handles. In the 6-dimensional Stein manifold we now have an h -principle for the attaching maps.

6. SUBCRITICAL FILLINGS OF S^5

The purpose of the present section is to show how the so-called Andrews–Curtis moves on balanced group presentations can be realised as moves in our diagrams for contact 5-manifolds. As an application we prove that any subcritically Stein fillable contact structure on the 5-sphere is the standard one, provided the filling gives rise to an Andrews–Curtis trivial presentation of the trivial group.

Let ξ be a contact structure on S^5 admitting a subcritical Stein filling W . From the homology long exact sequence of the pair (W, S^5) we deduce that W has the homology of a point. By the result of Cieliebak [6] we can write $W = \Sigma \times D^2$.

Hence $\pi_1(W) = \pi_1(\Sigma) = \pi_1(S^5) = \{1\}$, cf. the proof of Lemma 5.1. From the h -cobordism theorem it follows that W is diffeomorphic to D^6 .

Since the homology of W can also be computed from a cellular decomposition, it follows that a handlebody decomposition of W with a single 0-handle must contain an equal number of 1- and 2-handles. This gives rise to a **balanced presentation** of the trivial group $\pi_1(W)$, i.e. a presentation containing as many relations as generators.

If $\partial\Sigma$ were the standard 3-sphere, we could conclude immediately that ξ is the standard contact structure on S^5 . Unfortunately, the boundary of a contractible 4-manifold Σ (even with a finite handlebody decomposition as described) will, in general, be some complicated homology 3-sphere, as pointed out in [19].

6.1. Andrews–Curtis moves. In the attempt to prove that ξ is the standard contact structure on S^5 , one can try to show that the balanced presentation of the trivial group $\pi_1(W)$ can be converted to the empty presentation via balanced presentations, and then to implement these moves as transformations of the handlebody W . As we shall see in the next section, this topological implementation does not pose any difficulties. The algebraic part of this strategy, however, remains unresolved and forms the content of the Andrews–Curtis conjecture. There are various forms of this conjecture; the following is the weaker of the two versions given in [2].

Conjecture 6.1. *Any balanced presentation of the trivial group can be reduced to the empty presentation (via balanced presentations) by the following transformations:*

- AC 1.** *Replace a relation by its inverse.*
- AC 2.** *Conjugate a relation by a generator.*
- AC 3.** *Replace one relation by its product with another relation.*
- AC 4.** *Add or remove a generator g together with the relation g .*

A balanced presentation of the trivial group is called **Andrews–Curtis trivial** if it can be reduced to the empty presentation using Andrews–Curtis moves. In other words, the Andrews–Curtis conjecture can be rephrased as saying that any balanced presentation of the trivial group is Andrews–Curtis trivial.

This conjecture is still unresolved, but potential counterexamples have been suggested in [19].

6.2. Realising the Andrews–Curtis moves. The move **AC 1** amounts to reversing the orientation of the Legendrian attaching circle representing the relation in question; the moves **AC 2** and **AC 3** are the Tietze moves **T 1** (ii) and **T 1** (iii), respectively; the fourth Andrews–Curtis move is a special case of the second Tietze move. So all the Andrews–Curtis moves can be realised in our diagrams as in Section 5.1.3.

Some versions of the Andrews–Curtis conjecture also allow a generator to be replaced by its product with another generator, or by its inverse. These moves, too, can be realised in our diagrams as follows.

Multiplying a generator by another one amounts to a 1-handle slide [21, Figure 5.2]. Such a 1-handle slide can also be performed in a diagram with Legendrian attaching circles.

In order to replace a generator g by its inverse, one needs to flip the two attaching balls of the 1-handle corresponding to g . This can be done after sliding the Legendrian curves that go over this 1-handle halfway around the attaching balls, cf. [20, Figure 16].

6.3. A characterisation of the standard S^5 . With the Andrews–Curtis moves at our disposal, we can now give a characterisation of the standard contact structure on S^5 in terms of subcritical Stein fillings.

Proposition 6.2. *Let ξ be a contact structure on S^5 with a subcritical Stein filling W . If W admits a plurisubharmonic Morse function that induces a handlebody decomposition of W giving rise to an Andrews–Curtis trivial presentation of the trivial group $\pi_1(W)$, then ξ is diffeomorphic to the standard contact structure on S^5 .*

Proof. The subcritical Stein manifold W corresponds to a description of (S^5, ξ) as a contact open book $\text{Open}(\Sigma, \text{id})$ with a handlebody decomposition of the page Σ giving rise to an Andrews–Curtis trivial presentation of the trivial group. By realising the Andrews–Curtis moves that transform this presentation to the empty one, we transform the open book to the one described by the empty diagram, which represents the standard contact structure on S^5 . \square

Depending on one’s predisposition, one may regard this result as evidence for the conjecture that among the contact structures on S^5 only the standard one admits a subcritical Stein filling, or as a potential means for disproving the Andrews–Curtis conjecture. Indeed, one way to read the proposition is that any exotic (i.e. non-standard) contact structure on S^5 with a subcritical Stein filling W gives rise to a balanced presentation of the trivial group $\pi_1(W)$ that is not Andrews–Curtis trivial. Unfortunately, the result of M.-L. Yau cited at the end of Section 4.3 implies that cylindrical contact homology is not sensitive enough to detect such examples.

Although the general belief appears to be that the Andrews–Curtis conjecture is false, we should admit in all fairness that this suggested strategy for disproving it is not the most promising one. It seems more likely that methods such as those indicated in Remark 5.5 allow one to give a direct proof that the standard contact structure on S^5 is the only one admitting a subcritical Stein filling.

7. DIAGRAMS FOR SIMPLY CONNECTED 5-MANIFOLDS

In this section we exhibit diagrams for contact structures on some simply connected 5-manifolds. As a corollary, we obtain a branched covering description of 5-dimensional simply connected spin manifolds, and a new proof that every simply connected 5-manifold admits a contact structure in each homotopy class of almost contact structures.

7.1. Barden’s classification. Barden [3] has given a complete classification of simply connected 5-manifolds. Here we are only interested in those manifolds that potentially carry contact structures. If a 5-manifold M admits a contact structure, then its structure group reduces to $U(2) \times 1$; such a reduction is called an **almost contact structure**. Necessary and sufficient for the existence of an almost contact structure is the vanishing of the third integral Stiefel–Whitney class $W_3(M)$, see [16, Proposition 8.1.1].

According to Barden's classification, any simply connected 5-manifold M with $W_3(M) = 0$ decomposes as the connected sum of finitely many manifolds from the following list of examples:

- (i) manifolds M_k , $k \in \mathbb{N}$, characterised by $H_2(M_k) \cong \mathbb{Z}_k \oplus \mathbb{Z}_k$,
- (ii) $S^2 \times S^3$,
- (iii) $S^2 \tilde{\times} S^3$.

The manifold M_1 is the 5-sphere; the manifolds M_k are prime if and only if k is a prime power p^j , $j \geq 1$. One obtains a unique prime decomposition of a given M if one requires that only the M_{p^j} and at most one summand $S^2 \tilde{\times} S^3$ are used, cf. Proposition 4.5. All the prime manifolds in this list, with the exception of $S^2 \tilde{\times} S^3$, are spin manifolds, i.e. their second Stiefel–Whitney class vanishes.

7.2. Diagrams for the M_k . Let (N_k, η_k) , $k \in \mathbb{N}$, be the contact 5-manifold represented by the diagram depicted in Figure 13. Write Σ_k for the page of the open book represented by this diagram, so that $(N_k, \eta_k) = \text{Open}(\Sigma_k, (\tau_{K_1} \circ \tau_{K_2})^2)$.




FIGURE 13. Diagram for M_k .

Proposition 7.1. *For each $k \in \mathbb{N}$ the manifold N_k is diffeomorphic to M_k .*

One ingredient in the argument will be the following folklore theorem, which is proved in [26]; cf. [17] for a proof in the 3-dimensional case.

Theorem 7.2. *Let $L \subset \text{Open}(\Sigma, \psi)$ be a Legendrian sphere in a contact open book that sits on a page of the open book as a Lagrangian submanifold. Then the contact manifold obtained by Legendrian surgery along L is contactomorphic to $\text{Open}(\Sigma, \psi \circ \tau_L)$, where τ_L denotes the right-handed Dehn twist along $L \subset \Sigma$. \square*

Proof of Proposition 7.1. We construct a Stein filling W_k of (N_k, η_k) by starting with the subcritical Stein manifold $\Sigma_k \times D^2$, which is a filling for $\text{Open}(\Sigma_k, \text{id})$, and attaching a 3-handle for each Dehn twist. The first two 3-handles corresponding to τ_{K_1} and τ_{K_2} cancel the 2-handles of Σ_k , so we obtain a 6-dimensional disc D^6 .

The further two 3-handles corresponding to the iterated application of τ_{K_1} and τ_{K_2} then yields the Stein filling W_k . This implies $H_3(W_k) \cong \mathbb{Z}^2$, and the skew-symmetric intersection form Q_{W_k} on $H_3(W_k)$ has to look like

$$\begin{pmatrix} 0 & l_k \\ -l_k & 0 \end{pmatrix}$$

for some $l_k \in \mathbb{Z}$. By the argument mentioned in Section 2.5 we conclude

$$H_2(N_k) \cong H_2(\partial W_k) \cong \text{coker } Q_{W_k} \cong \mathbb{Z}_{l_k} \oplus \mathbb{Z}_{l_k}.$$

The manifold N_k is obviously simply connected, and it satisfies $W_3(N_k) = 0$ because it carries a contact structure. So in order to establish that N_k is diffeomorphic to M_k it suffices to show, by Barden's classification, that $|l_k| = k$.

Decompose the open book N_k as $N_k = A_k \cup_{\partial} B_k$ as in Section 2.1, with A_k the mapping torus of Σ_k , and $B_k = \partial\Sigma_k \times D^2 \simeq \partial\Sigma_k$. We can then compute the homology of N_k by the procedure outlined in Section 2.5. In particular, we consider the Mayer–Vietoris sequence

$$\begin{aligned} H_3(N_k) \longrightarrow H_2(A_k \cap B_k) \longrightarrow H_2(A_k) \oplus H_2(B_k) \longrightarrow \\ H_2(N_k) \longrightarrow H_1(A_k \cap B_k) \longrightarrow H_1(A_k) \oplus H_1(B_k). \end{aligned}$$

In terms of the standard generators of $H_2(\Sigma_k)$, the intersection form Q_{Σ_k} is given by

$$\begin{pmatrix} -2 & k \\ k & -2 \end{pmatrix}.$$

Thus, as shown in Section 3.1, the action of the two Dehn twists on $H_2(\Sigma_k)$ is given by

$$(\tau_{K_1})_* = \begin{pmatrix} -1 & k \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad (\tau_{K_2})_* = \begin{pmatrix} 1 & 0 \\ k & -1 \end{pmatrix},$$

hence

$$(\tau_{K_1} \circ \tau_{K_2})_*^2 = \begin{pmatrix} k^4 - 3k^2 + 1 & 2k - k^3 \\ -2k + k^3 & -k^2 + 1 \end{pmatrix}.$$

From the Wang sequence of the mapping torus A_k we then have

$$H_2(A_k) \cong \text{coker} \begin{pmatrix} k^4 - 3k^2 & 2k - k^3 \\ -2k + k^3 & -k^2 \end{pmatrix}.$$

Given a homomorphism $\phi: \mathbb{Z}^m \rightarrow \mathbb{Z}^m$ with $\det \phi \neq 0$, a simple algebraic consideration shows that $|\text{coker } \phi| = |\det \phi|$. This allows us to conclude that

$$|H_2(A_k)| = k^2 |k^2 - 4| \quad \text{for } k \neq 2.$$

For $k = 2$ we obtain $H_2(A_2) \cong \mathbb{Z} \oplus \mathbb{Z}_4$.

From the Künneth theorem we have

$$H_1(A_k \cap B_k) \cong H_0(\partial\Sigma_k) \oplus H_1(\partial\Sigma_k).$$

From the Wang sequence one sees that $H_1(A_k)$ is generated by the class of $\{p\} \times S^1$, where p is any point of $\partial\Sigma_k$. Combining these two pieces of information, one deduces that the homomorphism

$$H_1(A_k \cap B_k) \longrightarrow H_1(A_k) \oplus H_1(B_k)$$

in the Mayer–Vietoris sequence is an isomorphism.

Similarly, we have

$$H_2(A_k \cap B_k) \cong H_1(\partial\Sigma_k) \oplus H_2(\partial\Sigma_k),$$

and the second summand $H_2(\partial\Sigma_k)$ maps isomorphically into the second summand of $H_2(A_k) \oplus H_2(B_k)$ in the Mayer–Vietoris sequence.

So that sequence tells us that $H_2(N_k)$ is a quotient of $H_2(A_k)$, and hence at most of rank 1. Combining this with the information that $H_2(N_k)$ is isomorphic to $\mathbb{Z}_{l_k} \oplus \mathbb{Z}_{l_k}$, we conclude $l_k \neq 0$. Since $H_1(N_k) = 0$, we obtain $H^2(N_k) = 0$ from the

universal coefficient theorem, whence $H_3(N_k) = 0$ by Poincaré duality. Thus, the Mayer–Vietoris sequence reduces to

$$0 \longrightarrow H_1(\partial\Sigma_k) \longrightarrow H_2(A_k) \longrightarrow H_2(N_k) \longrightarrow 0.$$

Recall that $H_1(\partial\Sigma_k) \cong \text{coker } Q_{\Sigma_k}$. For $k \neq 2$ this is a finite group of order $|k^2 - 4|$. It follows that $H_2(N_k)$ is a finite group of order k^2 , and hence $|l_k| = k$, as we wanted to show.

For $k = 2$ the short exact sequence becomes

$$0 \longrightarrow \mathbb{Z} \oplus \mathbb{Z}_2 \longrightarrow \mathbb{Z} \oplus \mathbb{Z}_4 \longrightarrow H_2(N_k) \longrightarrow 0.$$

Recall that $H_1(\partial\Sigma_2)$ is generated by meridional loops u_1, u_2 around K_1, K_2 , respectively. The homomorphism $H_1(\partial\Sigma_2) \rightarrow H_2(A_2)$ is given by sending u_i to the class represented by the torus $u_i \times S^1 \subset \partial\Sigma_2 \times S^1 \subset A_2$. Now cut this torus along a meridian $u_i \times *$ and insert two meridional discs in Σ_2 (of opposite orientation); this gives us a 2-sphere in A_2 representing the same homology class. Now flow one of the meridional discs and the cylindrical part of that 2-sphere along a vector field on the mapping torus that is transverse to the fibres and whose time-1 map, say, gives the monodromy map from a fibre to itself.

The geometric intersection number of the meridional disc D_i to u_i with the spherical generator S_j of $H_2(\Sigma_2)$ corresponding to K_j (made up of a Seifert disc for K_j and the core disc of the 2-handle) equals the Kronecker δ_{ij} . The self-intersection number of the S_i is -2 , and the intersection number between S_1 and S_2 equals $k = 2$. Hence, with the observation from Section 3.1 we infer that the monodromy $(\tau_{K_1} \circ \tau_{K_2})^2$ acts on the D_i as follows:

$$\begin{array}{lll} D_1 & \mapsto & D_1 \\ & \mapsto & D_1 + S_1 + 2S_2 \\ D_2 & \mapsto & D_2 + S_2 \\ & \mapsto & D_2 + 2S_1 + 4S_2 \end{array} \quad \begin{array}{lll} & \mapsto & D_1 + S_1 \\ & \mapsto & D_1 + 4S_1 + 2S_2, \\ & \mapsto & D_2 + 2S_1 + S_2 \\ & \mapsto & D_2 + 6S_1 + 4S_2. \end{array}$$

The homology group $H_2(A_2) \cong \mathbb{Z} \oplus \mathbb{Z}_4$ is generated by the classes of S_1 and S_2 , subject to the relation $4(S_1 + S_2) = 0$. So the effect of the monodromy homomorphism can also be written as

$$\begin{array}{ll} D_1 & \mapsto D_1 - 2S_2, \\ D_2 & \mapsto D_2 + 2S_1. \end{array}$$

It follows that the 2-torus $u_1 \times S^1$ (resp. $u_2 \times S^1$) is homologically equivalent to $-2S_2$ (resp. $2S_1$) in A_2 . Thus, in terms of suitable generators, the homomorphism $\mathbb{Z} \oplus \mathbb{Z}_2 \rightarrow \mathbb{Z} \oplus \mathbb{Z}_4$ in the Mayer–Vietoris sequence for N_2 is multiplication by 2, hence $H_2(N_2) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$, i.e. $|l_2| = 2$. \square

7.3. Spin manifolds as branched covers. Here is an amusing corollary of the above example.

Corollary 7.3. *Any closed, simply connected 5-dimensional spin manifold is a double branched cover of the 5-sphere.*

Proof. By Barden’s classification, any closed, simply connected 5-dimensional spin manifold M can be decomposed as

$$M \cong \#_m S^2 \times S^3 \# M_{k_1} \# \cdots \# M_{k_n}.$$

This classification of *spin* manifolds had actually been achieved earlier by Smale [36]. By the preceding section, each M_{k_i} is diffeomorphic to an open book $\text{Open}(\Sigma_{k_i}, \psi_i^2)$.

The manifold $S^2 \times S^3$ can likewise be written as an open book with quadratic monodromy. Namely, as page take the cotangent unit disc bundle DT^*S^2 of S^2 . With Proposition 4.3 the manifold $\text{Open}(DT^*S^2, \text{id})$ can be shown to be diffeomorphic to $S^2 \times S^3$. Let τ be a right-handed Dehn twist along the zero section of DT^*S^2 . Since τ^2 is isotopic, relative to the boundary, to the identity map, it follows that $S^2 \times S^3$ is diffeomorphic to $\text{Open}(DT^*S^2, \tau^2)$, see [27] and [23, p. 36]. So M can be written as

$$M \cong \text{Open}(\natural_m DT^*S^2 \natural_{\Sigma_{k_1}} \natural \cdots \natural_{\Sigma_{k_n}} \natural_m \tau^2 \natural_{\psi_1^2} \natural \cdots \natural_{\psi_n^2}).$$

This implies that M is the double branched cover of

$$\text{Open}(\natural_m DT^*S^2 \natural_{\Sigma_{k_1}} \natural \cdots \natural_{\Sigma_{k_n}} \natural_m \tau \natural_{\psi_1} \natural \cdots \natural_{\psi_n}),$$

branched along the binding of the open book. That last open book, however, is a right-handed stabilisation of $\text{Open}(D^4, \text{id})$, which is diffeomorphic to the 5-sphere. \square

7.4. Existence of contact structures. The following theorem was first proved by the second author [15], using contact surgery. The second proof was given by the third author [25], using open book decompositions. Here we give a diagrammatic proof.

Theorem 7.4. *Every closed, oriented, simply connected 5-manifold admits a contact structure in each homotopy class of almost contact structures.*

Proof. Homotopy classes of almost contact structures on oriented, simply connected 5-manifolds are classified by the first Chern class, cf. [16, Proposition 8.1.1]. So each M_k in Barden's classification admits a unique almost contact structure up to homotopy (for either of its orientations). The result now follows from Section 7.2 and Proposition 4.3. \square

Acknowledgements. F. D. is partially supported by grant no. 10631060 of the National Natural Science Foundation of China.

This project was initiated during the conference “Symplectic Topology, Contact Topology, and Applications” at Hokkaido University, organised by O. v. K. (then a post-doc at Hokkaido University) and funded by JSPS grant 19-07804.

Some parts of this research were done while H. G. and O. v. K. attended the workshop “Symplectic Techniques in Conservative Dynamics” at the Lorentz Center, Leiden, organised by V. Ginzburg, F. Pasquotto, B. Rink and R. Vandervorst. We thank the organisers for this opportunity to meet, and the Lorentz Center for the excellent working environment.

The final writing was done while F. D. stayed at the Universität zu Köln, supported by a DAAD – K. C. Wong fellowship, grant no. A/09/99005.

REFERENCES

- [1] S. Akbulut and M. F. Arıkan, Lefschetz fibrations on compact Stein manifolds, [arXiv: 1003.2200v3](#).
- [2] J. J. Andrews and M. L. Curtis, Free groups and handlebodies, *Proc. Amer. Math. Soc.* **16** (1965), 192–195.
- [3] D. Barden, Simply connected five-manifolds, *Ann. of Math. (2)* **82** (1965), 365–385.
- [4] F. Bourgeois and O. van Koert, Contact homology of left-handed stabilizations and plumbing of open books, *Commun. Contemp. Math.* **12** (2010), 223–263.

- [5] G. E. Bredon, *Topology and Geometry*, Grad. Texts in Math., Vol. 139 (Springer-Verlag, Berlin, 1993).
- [6] K. Cieliebak, Subcritical Stein manifolds are split, [arXiv: math.DG/0204351](#).
- [7] R. H. Crowell and R. H. Fox, *Introduction to Knot Theory*, Grad. Texts in Math., Vol. 57 (Springer-Verlag, Berlin, 1977).
- [8] F. Ding and H. Geiges, Handle moves in contact surgery diagrams, *J. Topol.* **2** (2009), 105–122.
- [9] F. Ding and H. Geiges, The diffeotopy group of $S^1 \times S^2$ via contact topology, *Compos. Math.* **146** (2010), 1096–1112.
- [10] F. Ding and H. Geiges, Neighbourhoods and isotopies of knots in contact 3-manifolds, [arXiv: 1012.1721](#).
- [11] Ya. Eliashberg, Topological characterization of Stein manifolds of dimension > 2 , *Internat. J. Math.* **1** (1990), 29–46.
- [12] Ya. Eliashberg and M. Fraser, Topologically trivial Legendrian knots, *J. Symplectic Geom.* **7** (2009), 77–127.
- [13] Ya. Eliashberg and M. Gromov, Convex symplectic manifolds, in *Several Complex Variables and Complex Geometry (Santa Cruz, CA, 1989)*, Proc. Sympos. Pure Math., Vol. 52, Part 2 (American Mathematical Society, Providence, RI, 1991), pp. 135–162.
- [14] D. Fuchs and S. Tabachnikov, Invariants of Legendrian and transverse knots in the standard contact space, *Topology* **36** (1997), 1025–1053.
- [15] H. Geiges, Contact structures on 1-connected 5-manifolds, *Mathematika* **38** (1991), 303–311.
- [16] H. Geiges, *An Introduction to Contact Topology*, Cambridge Stud. Adv. Math., Vol. 109 (Cambridge University Press, 2008).
- [17] H. Geiges, Contact structures and geometric topology, [arXiv: 1004.3172](#).
- [18] E. Giroux, Géométrie de contact: de la dimension trois vers les dimensions supérieures, in *Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002)*, (Higher Ed. Press, Beijing, 2002), pp. 405–414.
- [19] R. E. Gompf, Killing the Akbulut-Kirby 4-sphere, with relevance to the Andrews–Curtis and Schoenflies problems, *Topology* **30** (1991), 97–115.
- [20] R. E. Gompf, Handlebody construction of Stein surfaces, *Ann. of Math. (2)* **148** (1998), 619–693.
- [21] R. E. Gompf and A. I. Stipsicz, *4-Manifolds and Kirby Calculus*, Grad. Stud. Math., Vol. 20 (American Mathematical Society, Providence, RI, 1999).
- [22] M. Gromov, Pseudoholomorphic curves in symplectic manifolds, *Invent. Math.* **82** (1985), 307–347.
- [23] F. Hirzebruch and K. H. Mayer, *O(n)-Mannigfaltigkeiten, exotische Sphären und Singularitäten*, Lecture Notes in Math., Vol. 57 (Springer-Verlag, Berlin, 1968).
- [24] O. van Koert, Open books for contact five-manifolds and applications of contact homology, Inaugural-Dissertation, Universität zu Köln (2005).
- [25] O. van Koert, Open books on contact five-manifolds, *Ann. Inst. Fourier (Grenoble)* **58** (2008), 139–157.
- [26] O. van Koert, Lecture notes on stabilization of contact open books, [arXiv: 1012.4359](#).
- [27] O. van Koert and K. Niederkrüger, Open book decompositions for contact structures on Brieskorn manifolds, *Proc. Amer. Math. Soc.* **133** (2005), 3679–3686.
- [28] O. Manzoli Neto, S. Massago and O. Saeki, Open book structures on $(n - 1)$ -connected $(2n + 1)$ -manifolds, *J. Math. Sci. Univ. Tokyo* **13** (2006), 439–523.
- [29] D. McDuff and D. Salamon, *J-Holomorphic Curves and Symplectic Topology*, Amer. Math. Soc. Colloq. Publ., Vol. 52 (American Mathematical Society, Providence, RI, 2004).
- [30] M. McLean, Lefschetz fibrations and symplectic homology, *Geom. Topol.* **13** (2009), 1877–1944.
- [31] J. Milnor, *Singular Points of Complex Hypersurfaces*, Ann. of Math. Stud, Vol. 61 (Princeton University Press, 1968).
- [32] P. Seidel, Lagrangian two-spheres can be symplectically knotted, *J. Differential Geom.* **52** (1999), 145–171.
- [33] P. Seidel, Lectures on four-dimensional Dehn twists, in *Symplectic 4-Manifolds and Algebraic Surfaces*, Lecture Notes in Math., Vol. 1938 (Springer-Verlag, Berlin, 2008), pp. 231–267.
- [34] P. Seidel and I. Smith, The symplectic topology of Ramanujam’s surface, *Comment. Math. Helv.* **80** (2005), 859–881.

- [35] H. Seifert and W. Threlfall, *Lehrbuch der Topologie* (Teubner, Leipzig, 1934).
- [36] S. Smale, On the structure of 5-manifolds, *Ann. of Math. (2)* **75** (1962), 38–46.
- [37] A. Weinstein, Symplectic manifolds and their Lagrangian submanifolds, *Advances in Math.* **6** (1971), 329–346.
- [38] A. Weinstein, Contact surgery and symplectic handlebodies, *Hokkaido Math. J.* **20** (1991), 241–251.
- [39] M.-L. Yau, Cylindrical contact homology of subcritical Stein-fillable contact manifolds, *Geom. Topol.* **8** (2004), 1243–1280.

DEPARTMENT OF MATHEMATICS, PEKING UNIVERSITY, BEIJING 100871, P. R. CHINA
E-mail address: `dingfan@math.pku.edu.cn`

MATHEMATISCHES INSTITUT, UNIVERSITÄT ZU KÖLN, WEYERTAL 86–90, 50931 KÖLN, GERMANY
E-mail address: `geiges@math.uni-koeln.de`

DEPARTMENT OF MATHEMATICAL SCIENCES, SEOUL NATIONAL UNIVERSITY, SAN56-1 SHILLIM-DONG KWANAK-GU, SEOUL 151-747, KOREA
E-mail address: `okoert@snu.ac.kr`