

EXPONENTIAL MOMENTS OF SELF-INTERSECTION LOCAL TIMES OF STABLE RANDOM WALKS IN SUBCRITICAL DIMENSIONS

FABIENNE CASTELL, CLÉMENT LAURENT, AND CLOTHILDE MÉLOT

ABSTRACT. Let $(X_t, t \geq 0)$ be an α -stable random walk with values in \mathbb{Z}^d . Let $l_t(x) = \int_0^t \delta_x(X_s) ds$ be its local time. For $p > 1$, not necessarily integer, $I_t = \sum_x l_t^p(x)$ is the so-called p -fold self-intersection local time of the random walk. When $p(d - \alpha) < d$, we derive precise logarithmic asymptotics of the probability $\mathbb{P}[I_t \geq r_t]$ for all scales $r_t \gg \mathbb{E}[I_t]$. Our result extends previous works by Chen, Li & Rosen 2005, Becker & König 2010, and Laurent 2012.

1. INTRODUCTION

Intersections of random paths have been an extensively studied topic, not only due to its fundamental importance in the theory of stochastic processes, but also because they play a central role in various physical models as the "polaron problem" in quantum field theory [18], the parabolic Anderson model describing diffusion in random potential ([24, 32, 8]), or random polymer models ([17, 19, 20]). Therefore, various mathematical objects have been introduced to quantify these intersections: the range of a random walk, the volume of the Wiener sausage, or the self-intersection local times, which is the main object of the present paper.

1.1. Self-intersection local times. Let $(X_t, t \geq 0)$ be a continuous time random walk on \mathbb{Z}^d , with jump rate 1, and generator A :

$$Af(x) = \sum_{y \in \mathbb{Z}^d} \mu(y - x)(f(y) - f(x)),$$

where μ denotes the law of the increments of the random walk. Throughout the paper, we assume that μ is in the domain of attraction of a stable law of index $\alpha \in (0; 2]$ and that μ is symmetric. More precise assumptions on μ will be given later. For any $x \in \mathbb{Z}^d$, let $l_t(x) = \int_0^t \delta_x(X_s) ds$ be the time spent by the random walk on site x up to time t . For a positive real number $p \geq 1$, consider the ℓ_p norm of l_t

$$N_p(l_t) = \left(\sum_{x \in \mathbb{Z}^d} l_t(x)^p \right)^{1/p}$$

Note that for $p = 1$, $N_1(l_t) = t$, while for $p = 0$, it is equal to the number of distinct sites visited by the random walk up to time t . Moreover, since $l_t(x) \leq t$ for all $x \in \mathbb{Z}^d$, $N_p(l_t) \leq t$. If p is an integer,

$$I_t := N_p^p(l_t) = \int_0^t dt_1 \cdots \int_0^t dt_p \mathbb{1}_{X_{t_1} = \cdots = X_{t_p}},$$

is the so called p -fold intersection local time of the random walk, which measures the time spent by the random walk on sites visited at least p times. To begin our study of the large time behavior

2000 *Mathematics Subject Classification.* 60F10; 60J55.

Key words and phrases. Self-intersection local times; Random walk in random scenery; large deviations; stable random walk

This research was supported by the french ANR project MEMEMO2.

of $N_p(l_t)$, let us have a look at its typical behavior, which depends on the recurrence/transience property of the random walk. We state the results in terms of I_t :

- **Transient case** ($d > \alpha$): in this case, the range of the random walk is of order t and the random walk spends a time of order 1 on each visited site, so that I_t is of order t . In the introduction of [23], Kesten & Spitzer proved that I_t/t converges a.s. to some deterministic constant.
- **Recurrent case** ($d < \alpha$, i.e $d = 1, \alpha > 1$): the range of the random walk is now of order $t^{1/\alpha}$; the random walk spends a time of order $t^{1-1/\alpha}$ on each visited site, so that I_t is of order $t^{p-(p-1)/\alpha}$. More precisely, $I_t/t^{p-(p-1)/\alpha}$ converges in distribution to the L_p -norm of the stable limiting process local time (see lemme 6 in [23], or lemma 14 in [15]).
- **Critical case** ($d = \alpha$; i.e $d = \alpha = 1, d = \alpha = 2$): the range of the random walk is now of order $t/\log(t)$; the random walk spends a time of order $\log(t)$ on each visited site, so that I_t is of order $t \log(t)^{p-1}$. More precisely, $I_t/(t \log(t)^{p-1})$ converges a.s. to some explicit deterministic constant (see [11] or lemme 4 in [10]).

To summarize, we get

$$\mathbb{E}(I_t) \asymp \begin{cases} t^{p-(p-1)/\alpha} & \text{if } d < \alpha, \\ t \log(t)^{p-1} & \text{if } d = \alpha, \\ t & \text{if } d > \alpha. \end{cases} \quad (1)$$

1.2. Main result. In this paper we are interested in the large deviations asymptotics of $N_p(l_t)$ when the walk produces an excess of self-intersections. More precisely, we give exact logarithmic asymptotics of $\mathbb{P}[I_t \geq r_t]$ for scales $t^p \gg r_t \gg \mathbb{E}(I_t)$. This problem has received a lot of attention during the last decade, and we refer the reader to the recent monograph [12], or to the survey paper [25] for an up-to-date picture of known results and of the various technics used in order to prove them. Let us stress that on a heuristic level, there is a phase transition in the optimal behavior of the random walk to produce many self-intersections:

- In the supercritical case $d > \alpha q$ (where q is the conjugate exponent of p), this optimal behavior is to stay confined in a ball of radius of order 1 during a time period of order $r_t^{1/p} \ll t$. This confinement happens with probability of order $\exp(-r_t^{1/p})$. Rough logarithmic asymptotics in the supercritical case were first proved in [2] (for $p = \alpha = 2$) and later refined in precise logarithmic asymptotics in [1, 26].
- In the subcritical case $d < \alpha q$, the optimal behavior for the random walk is to remain up to time t in a ball whose volume is of order $t^q r_t^{-q/p} \gg 1$. This confinement happens with probability of order $\exp(-t^{1-(\alpha q)/d} r_t^{(\alpha q)/(pd)})$. Precise logarithmic asymptotics were first proved in $d = 1$ in [14] for $\alpha = 2$, and later extended in [15] for $\alpha > 1$. Very recently, the case $d \geq 2, \alpha = 2$ was treated in [7] (with the restriction $d < 2/(p-1) < 2q$), and [27].
- In the critical case $d = \alpha q$, any confinement strategy in a ball of volume $R^d \in [1; t^q r_t^{-q/p}]$ during a time period of order $r_t^{1/p} R^{d/q}$ has a probability of order $\exp(-r_t^{1/p})$, so that the intuitive picture is not clear. Nevertheless, precise logarithmic asymptotics were established in [9] for $\alpha = 2$, and later extended in [26] to other values of α .

The main result of this paper is to prove logarithmic asymptotics of $\mathbb{P}[I_t \geq r_t]$ in the subcritical case, for any value of $\alpha \in (0, 2]$, thus extending [15] to $d \geq 2$ and $\alpha < 1$, and [7, 27] to $\alpha < 2$. To state our result, we have to introduce some notations and make precise assumptions on μ . For $p > 1$, $L^p(\mathbb{R}^d)$ is the usual Sobolev space of p -integrable functions w.r.t Lebesgue measure. $\|\cdot\|_p$ is the norm on $L^p(\mathbb{R}^d)$. For any function g in $L^2(\mathbb{R}^d)$, we denote by $\mathcal{F}(g)$ its Fourier transform:

$$\mathcal{F}(g)(\omega) = \int_{\mathbb{R}^d} g(x) \exp(-2i\pi \langle x, \omega \rangle) dx, \omega \in \mathbb{R}^d. \quad (2)$$

Throughout the paper, we assume that

(H1): μ is symmetric.

(H2): μ is in the normal domain of attraction of a strictly (symmetric) stable law S_α with index $\alpha \in (0, 2]$ with characteristic function given by $\exp(-|t|^\alpha)$. This means that there exists a constant c such that $(\frac{c}{t^{1/\alpha}}X_{st}, s \in [0; 1])$ converges in distribution when $t \rightarrow +\infty$, in the space $D([0, 1])$ of càdlàg functions, endowed with J_1 -topology. Without loss of generality, we will assume that $c = 2\pi$. This choice is made in order to get nice statements using (2) as a definition of Fourier transform.

(H3): μ is non-arithmetic, i.e $\{\omega \in \mathbb{R}^d, \text{ such that } |\sum_{x \in \mathbb{Z}^d} \exp(2i\pi\langle \omega, x \rangle) \mu(x)| = 1\} = \mathbb{Z}^d$

(H4): Let $p_t(x, y)$ be the transition probabilities of $(X_t, t \geq 0)$. There exists a constant κ such that for any $t > 0$, and any $x, y \in \mathbb{Z}^d$,

$$p_t(x, y) \leq \kappa \left(\frac{1}{t^{d/\alpha}} \wedge \frac{t}{|x - y|^{d+\alpha}} \right), \quad (3)$$

where $x \wedge y = \min(x, y)$. When $\alpha = 2$, this is true by standard Gaussian estimates. When $\alpha < 2$, assumption **(H1)** and the following condition on μ are shown to imply (3) in [6]: there are constants c_1, c_2 , such that for any $x \in \mathbb{Z}^d, x \neq 0$,

$$\frac{c_1}{|x|^{d+\alpha}} \leq \mu(x) \leq \frac{c_2}{|x|^{d+\alpha}}.$$

We next define

$$\rho_{\alpha, d, p} = \left(\frac{\alpha q}{\alpha q - d} \right) \left(\frac{\alpha q - d}{d} \right)^{d/\alpha q} \inf_{g: \mathbb{R}^d \rightarrow \mathbb{R}} \left\{ \|g\|_2^{2(1-d/\alpha q)} \left(\int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2 d\omega \right)^{d/\alpha q}, \|g\|_{2p} = 1 \right\}. \quad (4)$$

$$\chi_{\alpha, d, p} := \inf_{g: \mathbb{R}^d \rightarrow \mathbb{R}} \left\{ \int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2 d\omega, \|g\|_{2p} = \|g\|_2 = 1 \right\}. \quad (5)$$

Note that the expression in the infimum appearing in (4) is invariant under the transformation $g \mapsto g_\lambda = \lambda^{d/(2p)} g(\lambda \cdot)$, so that one can reduce this infimum to functions g satisfying $\|g\|_{2p} = \|g\|_2 = 1$. Therefore,

$$\rho_{\alpha, d, p} = \frac{\alpha q}{\alpha q - d} \left(\frac{\alpha q - d}{d} \chi_{\alpha, d, p} \right)^{d/(\alpha q)}.$$

It is proved in [16] that these constants are not degenerate when $d < \alpha q$. We have then the following result.

Theorem 1. *Assume **(H1-4)**, and that $p > 1$, $\alpha \in (0; 2]$ and $d < \alpha q$, where q is the conjugate exponent of p ($1/p + 1/q = 1$). Let $(\beta_t, t \geq 0)$ be such that for large t ,*

- $1 \ll \beta_t^\alpha \ll t$ when $d < \alpha$,

- $1 \ll \beta_t^d \ll \frac{t}{\log(t)}$ when $d = \alpha$,

- $1 \ll \beta_t^d \ll t$ when for $d > \alpha$.

Then for any $\theta \geq 0$,

$$\lim_{t \rightarrow +\infty} \frac{\beta_t^\alpha}{t} \log \mathbb{E} \left[\exp \left(\theta \beta_t^{d/q - \alpha} N_p(t) \right) \right] = \left(\frac{\theta}{\rho_{\alpha, d, p}} \right)^{\frac{\alpha q}{\alpha q - d}} \quad (6)$$

and

$$\lim_{t \rightarrow +\infty} \frac{\beta_t^\alpha}{t} \log \mathbb{P} \left(I_t \geq \theta t^p \beta_t^{-d(p-1)} \right) = -\theta^{\alpha/(d(p-1))} \chi_{\alpha, d, p}. \quad (7)$$

Remark 2. Denoting $r_t = t^p \beta_t^{-d(p-1)}$, we stress the fact that the conditions on β_t appearing in theorem 1 are equivalent to $t^p \gg r_t \gg \mathbb{E}(I_t)$. Moreover, using the fact that the function $\theta \mapsto |\theta|^{(\alpha q)/(\alpha q - d)}$ is essentially smooth, (7) is a straightforward consequence of (6) and the Gärtner-Ellis theorem for positive random variables large deviations principle (see Theorem 1.2.3 in [12]), and is equivalent to a full large deviations principle for $t^{-p} \beta_t^{d(p-1)} I_t$ (see Theorem 1.2.1 in [12]).

1.3. Comments on the proof. Literature remarks. To understand the main difficulty in obtaining Theorem 1, let us introduce for any $x \in \mathbb{R}^d$,

$$L_t(x) = \frac{\beta_t^d}{t} l_t(\lfloor \beta_t x \rfloor).$$

Note that L_t is an element of the set $\mathcal{M}_1(\mathbb{R}^d)$ of probability measures on \mathbb{R}^d . Following Donsker-Varadhan's results, it is easy to check that L_t satisfies a weak large deviations principles on $\mathcal{M}_1(\mathbb{R}^d)$, endowed with the weak convergence defined by duality with compactly supported continuous functions (see section 5 Theorem 11). The speed of this large deviations principle is t/β_t^α , while its rate function is defined by

$$\mathcal{I}(\mu) = \begin{cases} \int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2 d\omega & \text{if } d\mu = g^2 dx, \\ +\infty & \text{otherwise.} \end{cases}$$

Moreover $\|L_t\|_p^p = \beta_t^{d(p-1)} t^{-p} I_t$. Applying the contraction principle to $\mu \in \mathcal{M}^1(\mathbb{R}^d) \mapsto \left\| \frac{d\mu}{dx} \right\|_p$ would lead to Theorem 1. The main problem here is that the latter function is lower semicontinuous but not continuous in weak topology. We can therefore deduce the lower bounds in Theorem 1 by contraction, but this is no more the case for the upper bounds. To circumvent this problem, various strategies have been proposed. One can first try to smoothen L_t . Given a regularizing kernel ϕ_ϵ , the map $\mu \in \mathcal{M}^1(\mathbb{R}^d) \mapsto \left\| \phi_\epsilon \star \frac{d\mu}{dx} \right\|_p$ is now continuous in weak topology, so that we just have to prove that $\|\phi_\epsilon \star L_t\|_p$ and $\|L_t\|_p$ are exponentially close. Typically, this can be done in "small" dimension using the regularity of local times. Roughly speaking, this is the way followed by Bass, Chen & Rosen in a series of works starting with the paper of Chen & Li [14] about the large deviations of the self-intersections ($p = 2$) of Brownian motion or random walk in $d = 1$. Later, they studied the cases $d = 1, \alpha > d$ in [15], $p = d = \alpha = 2$ in [4], $p = 2$ and $d \geq \alpha > 2d/3$ in [5]. In $d = 1$, they worked directly on the self-intersections of the limiting stable process, and used the existence and the regularity of their local times, thus reducing their result to $\alpha > d$. In $d \geq 2$, they first proved large deviations results for intersections of p -independent processes or random walks using a regularisation procedure in [13, 16], and then transferred these results to large deviations for self-intersections. Here the assumption $p = 2$ is crucial since this transfer is done via the bisection method introduced by Varadhan in [31], which does not work for $p \neq 2$. Becker and König [7] proved Theorem 1 for $\alpha = 2, p < 2q/p (< 2q)$ and $\beta_t \ll (t/\log(t))^{d/(d+2)}$, using an upper bound for the joint density of the local times of a Markov chain obtained in [3]. Unfortunately, this upper bound is not precise enough to obtain the result in its full extent. We propose here to use a method of proof introduced in [9] in the critical case, and successfully extended in [26] to the supercritical case, and to the subcritical case for $\alpha = 2$ in [27]. One of the main tool in the proof is the Dynkin's isomorphism theorem, according to which the law of the local times of a symmetric recurrent Markov process stopped at an independent exponential time, is related to the law of the square of a Gaussian process whose covariance function is the Green kernel of the stopped Markov process. This allows to control the exponential moments of $N_p(l_t)$ by the exponential moments of $N_{2p, R\beta_t}(Z)^2$ where

- $N_{2p, R\beta_t}(f)$ is the ℓ_{2p} -norm of a real valued function f defined on $\mathbb{T}_{R\beta_t}$ the discrete torus of radius $R\beta_t$;

- $(Z_x, x \in \mathbb{T}_{R\beta_t})$ is a centered Gaussian process independent of $(X_t, t \geq 0)$, whose covariance function is given by the Green kernel $G_{R\beta_t, \lambda_t}(x, y)$ of $(X_t, t \geq 0)$ projected on $\mathbb{T}_{R\beta_t}$, stopped at an independent exponential time with parameter $\lambda_t = a\beta_t^{-\alpha}$.

Loosely speaking, this leads to a non asymptotic upper bound

$$\mathbb{P}(N_p(l_t) \geq \theta^{1/p} t \beta_t^{-d/q}) \leq \exp(at\beta_t^{-\alpha}(1 + o(1))) \mathbb{P}(N_{p, R\beta_t}(Z^2) \geq 2\theta^{1/p} t \beta_t^{-d/q}),$$

where the first term in the right hand side comes from the stopping at an exponential time. As soon as the median of $N_{p, R\beta_t}(Z^2)$ is negligible with respect to $t\beta_t^{-d/q}$ (which is equivalent to the conditions on β_t), concentration inequalities for norms of Gaussian processes yields

$$\mathbb{P}(N_p(l_t) \geq \theta^{1/p} t \beta_t^{-d/q}) \leq \exp(at\beta_t^{-\alpha}(1 + o(1))) \exp(-\theta^{1/p} t \beta_t^{-d/q} \sigma^{-2}),$$

where σ^2 is the maximal variance of the process Z viewed as an element of $l_{2p}(\mathbb{T}_{R\beta_t})$:

$$\sigma^2 = \sup_{f: \mathbb{T}_{R\beta_t} \rightarrow \mathbb{R}} \{ \langle f, G_{R\beta_t, \lambda_t} f \rangle_{R\beta_t}; N_{(2p)', R\beta_t}(f) = 1 \},$$

where $\langle \cdot, \cdot \rangle_{R\beta_t}$ is the scalar product on $l^2(\mathbb{T}_{R\beta_t})$, and $G_{R\beta_t, \lambda_t} f(x) = \sum_{y \in \mathbb{T}_{R\beta_t}} G_{R\beta_t, \lambda_t}(x, y) f(y)$ for any $f \in l^2(\mathbb{T}_{R\beta_t})$ and any $x \in \mathbb{T}_{R\beta_t}$. Since $G_{R\beta_t, \lambda_t} = (\lambda_t \text{Id} - A_{R\beta_t})^{-1}$ (where $A_{R\beta_t}$ is the generator of the random walk on the torus), we get

$$\mathbb{P}(N_p(l_t) \geq \theta^{1/p} t \beta_t^{-d/q}) \leq \exp(-t\beta_t^{-\alpha}(-a + \theta^{1/p} \beta_t^{\alpha-d/q} \rho(a, R, t) + o(1))), \quad (8)$$

with

$$\rho(a, R, t) = \inf_{h: \mathbb{T}_{R\beta_t} \rightarrow \mathbb{R}} \{ a\beta_t^{-\alpha} N_{2, R\beta_t}^2(h) - \langle h, A_{R\beta_t} h \rangle_{R\beta_t}, N_{2p, R\beta_t}(h) = 1 \}. \quad (9)$$

Up to this point, the proof is the same in the critical case, the supercritical case, or the subcritical case. We just have to use the correct scalings. It remains now to study the limit of the upper bound (8) when t goes to infinity. This is where the proofs differ. In the critical case and the supercritical case, the limiting constant is still given by a variational formula involving functions defined on the grid \mathbb{Z}^d (see [9, 26]), while on the subcritical case, the limiting constant is given by a variational formula involving functions defined on \mathbb{R}^d . Therefore, we have to interpolate the minimisers in $\rho(a, R, t)$. When $\alpha = 2$, the operator A is local, and this interpolation is done via linear interpolation (see [27]). This proof does not work anymore when $\alpha < 2$, and we use here interpolation via Fourier transform.

The paper is organized as follows. Section 2 is a reminder of classical results in Fourier analysis. Section 3 is devoted to the upper bound in (6). It relies on Proposition 10 giving the asymptotic behavior of $\rho(a, R, t)$, and Lemma 9 about estimates of $G_{R\beta_t, \lambda_t}(0, 0)$ whose proofs are presented in Section 4. Finally, we prove the lower bound in (6) in Section 5.

2. PRELIMINARIES RESULTS ON FOURIER TRANSFORMS

We gather in this section notations and well known results about Fourier transform used throughout the paper.

2.1. Fourier transform on the discrete torus. Let R be an integer and let $u : \mathbb{T}_R \mapsto \mathbb{C}$ be a function defined on the discrete d -dimensional torus \mathbb{T}_R of radius R . We denote by $F_R(u) : \mathbb{T}_R \mapsto \mathbb{C}$ its Fourier transform:

$$\forall n \in \mathbb{T}_R, F_R(u)(n) = \sum_{k \in \mathbb{T}_R} u(k) \exp\left(-2i\pi \frac{\langle k, n \rangle}{R}\right).$$

We have an inversion formula and a Parseval formula:

Proposition 3. For $u : \mathbb{T}_R \mapsto \mathbb{C}$,

$$\forall k \in \mathbb{T}_R, u(k) = \frac{1}{R^d} \sum_{n \in \mathbb{T}_R} F_R(u)(n) \exp\left(2i\pi \frac{\langle k, n \rangle}{R}\right)$$

$$\text{and } \sum_{n \in \mathbb{T}_R} |F_R(u)(n)|^2 = R^d \sum_{k \in \mathbb{T}_R} |u(k)|^2.$$

2.2. Fourier transform on \mathbb{Z}^d . For a sequence $u \in l_2(\mathbb{Z}^d)$, we denote by $F(u)$ its Fourier transform. $F(u)$ is the 1-periodic function defined by

$$\forall \omega \in \mathbb{R}^d, F(u)(\omega) = \sum_{z \in \mathbb{Z}^d} u(z) \exp(-2i\pi \langle z, \omega \rangle).$$

There is an inversion and Parseval's formulas:

Proposition 4.

$$u(z) = \int_{[-1/2, 1/2]^d} F(u)(\omega) \exp(2i\pi \langle z, \omega \rangle) d\omega \text{ and } \|F(u)\|_{2, [-1/2, 1/2]^d}^2 = N_2^2(u)$$

These two types of Fourier transforms are linked through periodization: if $\forall x \in \mathbb{T}_R$, $g_R(x) = \sum_{z \in \mathbb{Z}^d} g(x + Rz)$ is the periodised version of g , then

$$\forall x \in \mathbb{T}_R, F_R(g_R)(x) = F(g)\left(\frac{x}{R}\right). \quad (10)$$

2.3. Fourier transform on the torus. For g_R a R -periodic function on \mathbb{R}^d , we consider its Fourier coefficients:

$$\forall n \in \mathbb{Z}^d, \mathcal{F}_R(g_R)(n) = \frac{1}{R^d} \int_{[0, R]^d} g_R(x) \exp\left(-2i\pi \frac{\langle n, x \rangle}{R}\right) dx.$$

Denoting by $\|\cdot\|_{2,R}$ the norm in $L^2([0, R]^d)$, we get the following inversion and Parseval's formulas:

Proposition 5.

$$\forall x \in \mathbb{R}^d, g_R(x) = \sum_{n \in \mathbb{Z}^d} \mathcal{F}_R(g_R)(n) \exp\left(2i\pi \frac{\langle n, x \rangle}{R}\right) \text{ and } N_2^2(\mathcal{F}_R(g_R)) = \frac{1}{R^d} \|g_R\|_{2,R}^2.$$

2.4. Fourier transform on \mathbb{R}^d . When $g : \mathbb{R}^d \mapsto \mathbb{R}$ is an element of $L^2(\mathbb{R}^d)$, we define its Fourier transform $\mathcal{F}(g) : \mathbb{R}^d \mapsto \mathbb{C}$ by equation (2). With this definition, Parseval's identity and inversion formula read

$$\|g\|_2 = \|\mathcal{F}(g)\|_2, \quad g(x) = \int_{\mathbb{R}^d} \mathcal{F}(g)(\omega) \exp(2i\pi \langle x, \omega \rangle) d\omega, \quad x \in \mathbb{R}^d.$$

To end this section, we remind the reader the following Young inequalities:

Proposition 6. let $p > 2$ and q be its conjugate exponent. For any $u : \mathbb{T}_R \mapsto \mathbb{C}$,

$$N_{p,R}(u) \leq R^{-d/q} N_{q,R}(F_R(u)).$$

For any R -periodic function $g_R : \mathbb{R}^d \mapsto \mathbb{C}$,

$$\|g_R\|_{p,R} \leq R^{d/p} N_q(\mathcal{F}_R(g_R)).$$

These inequalities are straightforward applications of the Riesz-Thorin interpolation theorem (see Theorem 1.3.4 in [22] for example) and Parseval's formula.

3. EXPONENTIAL MOMENTS UPPER BOUND

This section is devoted to the proof of the upper bound in (6).

Let $a > 0$ and τ be an exponential time of parameter $\lambda_t = a\beta_t^{-\alpha}$ independent of $(X_t, t \geq 0)$. Let $(X_s^{R\beta_t}, s \geq 0)$ be the projection of $(X_t, t \geq 0)$ on the discrete torus $\mathbb{T}_{R\beta_t}$:

$$X_s^{R\beta_t} = X_s \bmod(R\beta_t).$$

Let $l_{R\beta_t, t}(x) = \int_0^t \delta_x(X_s^{R\beta_t}) ds$ be its local time at site x up to time t .

$$\begin{aligned} N_p^p(l_t) &= \sum_{x \in \mathbb{Z}^d} l_t^p(x) = \sum_{x \in \mathbb{T}_{R\beta_t}} \sum_{k \in \mathbb{Z}^d} l_t^p(x + kR\beta_t) \\ &\leq \sum_{x \in \mathbb{T}_{R\beta_t}} \left(\sum_{k \in \mathbb{Z}^d} l_t(x + kR\beta_t) \right)^p = \sum_{x \in \mathbb{T}_{R\beta_t}} l_{p, R\beta_t}^p(x) = N_{p, R\beta_t}^p(l_{R\beta_t, t}). \end{aligned}$$

Then using the fact that $\tau \sim \mathcal{E}(\lambda_t)$ is independent of $(X_s, s \geq 0)$ we get for all $\theta > 0$:

$$\mathbb{E} \left[\exp \left(\theta \beta_t^{d/q-\alpha} N_p(l_t) \right) \right] \exp(-t\lambda_t) \leq \mathbb{E} \left[\exp \left(\theta \beta_t^{d/q-\alpha} N_{p, R\beta_t}(l_{R\beta_t, t}) \right); \tau \geq t \right] \quad (11)$$

$$\leq \mathbb{E} \left[\exp \left(\theta \beta_t^{d/q-\alpha} N_{p, R\beta_t}(l_{R\beta_t, \tau}) \right) \right] \quad (12)$$

We are now going to use the following version of Dynkyn's isomorphism theorem:

Theorem 7. (*Eisenbaum, see for instance corollary 8.1.2 page 364 in [30]*).

Let $\tau \sim \mathcal{E}(\lambda_t)$ independent of $(X_s, s \geq 0)$, and let $(Z_x, x \in \mathbb{T}_{R\beta_t})$ be a centered Gaussian process with covariance matrix $G_{R\beta_t, \lambda_t}(x, y) = E_x \left[\int_0^\tau \delta_y(X_s^{R\beta_t}) ds \right]$ independent of τ and of the random walk $(X_s, s \geq 0)$. For $s \neq 0$, consider the process $S_x := l_{R\beta_t, \tau}(x) + \frac{1}{2}(Z_x + s)^2$. Then for all measurable and bounded function $F : \mathbb{R}^{\mathbb{T}_{R\beta_t}} \mapsto \mathbb{R}$:

$$\mathbb{E} [F((S_x; x \in \mathbb{T}_{R\beta_t}))] = \mathbb{E} \left[F \left(\left(\frac{1}{2}(Z_x + s)^2; x \in \mathbb{T}_{R\beta_t} \right) \left(1 + \frac{Z_0}{s} \right) \right) \right].$$

This theorem allows to compare the tail behavior of $N_{p, R\beta_t}(l_{R\beta_t, \tau})$ with the tail behavior of $N_{2p, R\beta_t}(Z)$. Indeed, using that

$$S_x^p \geq l_{R\beta_t, \tau}^p(x) + \left(\frac{1}{2}(Z_x + s)^2 \right)^p,$$

and the independence of $(Z_x, x \in \mathbb{T}_{R\beta_t})$ with the random walk $(X_s, s \geq 0)$ and the exponential time τ , we get for all $\epsilon > 0$, and all $y > 0$,

$$\begin{aligned} &\mathbb{P} \left(N_{p, R\beta_t}^p(l_{R\beta_t, \tau}) \geq y^p t^p \beta_t^{d(1-p)} \right) \mathbb{P} \left(\frac{1}{2^p} N_{2p, R\beta_t}^{2p}(Z + s) \geq t^p \beta_t^{d(1-p)} \epsilon^p \right) \\ &\leq \mathbb{P} \left(N_{p, R\beta_t}^p(l_{R\beta_t, \tau}) + \frac{1}{2^p} N_{2p, R\beta_t}^{2p}(Z + s) \geq t^p \beta_t^{d(1-p)} (y^p + \epsilon^p) \right) \\ &\leq \mathbb{P} \left(N_{p, R\beta_t}^p(S) \geq t^p \beta_t^{d(1-p)} (y^p + \epsilon^p) \right) \\ &= \mathbb{E} \left[\left(1 + \frac{Z_0}{s} \right); \frac{1}{2^p} N_{2p, R\beta_t}^{2p}(Z + s) \geq t^p \beta_t^{d(1-p)} (y^p + \epsilon^p) \right] \end{aligned} \quad (13)$$

where the last equality comes from Theorem 7. Moreover by Hölder's inequality, for all $\epsilon > 0$,

$$\begin{aligned} & \mathbb{E} \left[\left(1 + \frac{Z_0}{s} \right); \frac{1}{2^p} N_{2p, R\beta_t}^{2p}(Z + s) \geq t^p \beta_t^{d(1-p)} (y^p + \epsilon^p) \right] \\ & \leq \mathbb{E} \left[\left| 1 + \frac{Z_0}{s} \right|^{1+1/\epsilon} \right]^{\epsilon/(1+\epsilon)} \mathbb{P} \left(N_{2p, R\beta_t}^{2p}(Z + s) \geq 2^p t^p \beta_t^{d(1-p)} (y^p + \epsilon^p) \right)^{1/(1+\epsilon)}. \end{aligned} \quad (14)$$

Combining (13) and (14), we obtain that for all $a, \epsilon > 0$,

$$\begin{aligned} & \mathbb{P} \left(N_{p, R\beta_t}(l_{R\beta_t, \tau}) \geq yt\beta_t^{-d/q} \right) \\ & \leq \mathbb{E} \left[\left| 1 + \frac{Z_0}{s} \right|^{1+1/\epsilon} \right]^{\frac{\epsilon}{1+\epsilon}} \frac{\mathbb{P} \left(N_{2p, R\beta_t}(Z + s) \geq \sqrt{2t} \beta_t^{-d/(2q)} (y^p + \epsilon^p)^{1/(2p)} \right)^{\frac{1}{1+\epsilon}}}{\mathbb{P}(N_{2p, R\beta_t}(Z + s) \geq \sqrt{2t\epsilon} \beta_t^{-d/(2q)})}. \end{aligned} \quad (15)$$

Then using the fact that $\text{Var}(Z_0) = G_{R\beta_t, \lambda_t}(0, 0)$, and Markov's inequality, we obtain that for all $\gamma > 0$,

$$\begin{aligned} & \mathbb{P} \left(N_{p, R\beta_t}(l_{R\beta_t, \tau}) \geq yt\beta_t^{-d/q} \right) \\ & \leq C(\epsilon) \left(1 + \frac{\sqrt{G_{R\beta_t, \lambda_t}(0, 0)}}{s} \right) \frac{\mathbb{P} \left(N_{2p, R\beta_t}(Z + s) \geq \sqrt{2t} \beta_t^{-d/(2q)} (y^p + \epsilon^p)^{1/(2p)} \right)^{1/(1+\epsilon)}}{\mathbb{P}(N_{2p, R\beta_t}(Z + s) \geq \sqrt{2t\epsilon} \beta_t^{-d/(2q)})}, \quad (16) \\ & \leq C(\epsilon) \left(1 + \frac{\sqrt{G_{R\beta_t, \lambda_t}(0, 0)}}{s} \right) \frac{\mathbb{E} \left[\exp \left(\frac{\gamma}{2} \beta_t^{d/q - \alpha} N_{2p, R\beta_t}^2(Z + s) \right) \right]^{1/(1+\epsilon)}}{\mathbb{P}(N_{2p, R\beta_t}(Z + s) \geq \sqrt{2t\epsilon} \beta_t^{-d/(2q)})} \exp \left(-\gamma t \beta_t^{-\alpha} \frac{(y^p + \epsilon^p)^{1/p}}{1 + \epsilon} \right). \end{aligned} \quad (17)$$

Therefore,

$$\begin{aligned} & \mathbb{E} \left[\exp \left(\theta \beta_t^{d/q - \alpha} N_{p, R\beta_t}(l_{R\beta_t, \tau}) \right) \right] \\ & = 1 + \int_0^{+\infty} \mathbb{P} \left(N_{p, R\beta_t}(l_{R\beta_t, \tau}) \geq yt\beta_t^{-d/q} \right) \theta t \beta_t^{-\alpha} \exp(\theta t y \beta_t^{-\alpha}) dy \\ & \leq 1 + C(\epsilon) \left(1 + \frac{\sqrt{G_{R\beta_t, \lambda_t}(0, 0)}}{s} \right) \frac{\mathbb{E} \left[\exp \left(\frac{\gamma}{2} \beta_t^{d/q - \alpha} N_{2p, R\beta_t}^2(Z + s) \right) \right]^{1/(1+\epsilon)}}{\mathbb{P}(N_{2p, R\beta_t}(Z + s) \geq \sqrt{2t\epsilon} \beta_t^{-d/(2q)})} \\ & \quad \int_0^{+\infty} \theta t \beta_t^{-\alpha} e^{-t\beta_t^{-\alpha} y (\frac{\gamma}{1+\epsilon} - \theta)} dy. \end{aligned}$$

Note that the integral is finite if and only if $\theta < \frac{\gamma}{1+\epsilon}$. In this case,

$$\begin{aligned} & \mathbb{E} \left[\exp \left(\theta \beta_t^{d/q - \alpha} N_{p, R\beta_t}(l_{R\beta_t, \tau}) \right) \right] \\ & \leq 1 + C(\epsilon, \theta, \gamma) \left(1 + \frac{\sqrt{G_{R\beta_t, \lambda_t}(0, 0)}}{s} \right) \frac{\mathbb{E} \left[\exp \left(\frac{\gamma}{2} \beta_t^{d/q - \alpha} N_{2p, R\beta_t}^2(Z + s) \right) \right]^{1/(1+\epsilon)}}{\mathbb{P}(N_{2p, R\beta_t}(Z + s) \geq \sqrt{2t\epsilon} \beta_t^{-d/(2q)})}. \end{aligned}$$

Choosing $s = \frac{\epsilon\sqrt{2t}}{R^{\frac{d}{2p}}\beta_t^{d/2}}$, using triangle inequality and the fact that $N_{2p,R\beta_t}(s) = s(R\beta_t)^{\frac{d}{2p}}$, we have:

$$\begin{aligned} & \mathbb{E} \left[\exp \left(\theta \beta_t^{d/q-\alpha} N_{p,R\beta_t}(l_{R\beta_t,\tau}) \right) \right] \\ & \leq 1 + C(\epsilon, \theta, \gamma) \left(1 + \frac{R^{d/(2p)} \sqrt{\beta_t^d G_{R\beta_t,\lambda_t}(0,0)}}{\epsilon\sqrt{2t}} \right) e^{\gamma t \epsilon \beta_t^{-\alpha}} \frac{\mathbb{E} \left[\exp \left(\frac{\gamma(1+\epsilon)}{2} \beta_t^{d/q-\alpha} N_{2p,R\beta_t}^2(Z) \right) \right]^{1/(1+\epsilon)}}{\mathbb{P} \left(N_{2p,R\beta_t}(Z) \geq \sqrt{2t\epsilon} \beta_t^{-d/(2q)} (1 + \sqrt{\epsilon}) \right)} \end{aligned} \quad (18)$$

Combining (12) and (18), we have thus proved that for all $\theta > 0, \epsilon > 0, \gamma > 0$ such that $\theta < \frac{\gamma}{1+\epsilon}$,

$$\begin{aligned} & \mathbb{E} \left[\exp \left(\theta \beta_t^{d/q-\alpha} N_p(l_t) \right) \right] e^{-at\beta_t^{-\alpha}} \\ & \leq \left[1 + C(\epsilon, \theta, \gamma) \left(1 + \frac{R^{d/(2p)} \sqrt{\beta_t^d G_{R\beta_t,\lambda_t}(0,0)}}{\epsilon\sqrt{2t}} \right) e^{\gamma t \epsilon \beta_t^{-\alpha}} \frac{\mathbb{E} \left[\exp \left(\frac{\gamma(1+\epsilon)}{2} \beta_t^{d/q-\alpha} N_{2p,R\beta_t}^2(Z) \right) \right]^{1/(1+\epsilon)}}{\mathbb{P} \left(N_{2p,R\beta_t}(Z) \geq \sqrt{2t\epsilon} \beta_t^{-d/(2q)} (1 + \sqrt{\epsilon}) \right)} \right]. \end{aligned} \quad (19)$$

We now use concentration inequalities for norms of Gaussian processes.

Lemma 8. *Large deviations for $N_{2p,R}(Z)$.*

Let τ and $(Z_x, x \in \mathbb{T}_{R\beta_t})$ be defined as in Theorem 7, and $\rho(a, R, t)$ be defined by (9). Under assumptions of theorem 1,

- (1) For all $a > 0, R > 0, t > 0$, $\rho(a, R, t) \leq aR^{d/q}\beta_t^{d/q-\alpha}$.
- (2) $\forall a, \epsilon, R, t > 0$,

$$\mathbb{P} \left[N_{2p,R\beta_t}(Z) \geq \sqrt{t\epsilon} \beta_t^{-d/(2q)} \right] \geq \frac{\beta_t^{d/(2q)}}{\sqrt{2\pi t \epsilon \rho(a, R, t)}} \left(1 - \frac{\beta_t^{d/q}}{t\epsilon \rho(a, R, t)} \right) \exp \left(-\frac{1}{2} t \beta_t^{-d/q} \epsilon \rho(a, R, t) \right).$$

- (3) $\forall a, \gamma, \epsilon, R, t > 0$ such that $\gamma(1+\epsilon)^2 < \rho(a, R, t)\beta_t^{\alpha-d/q}$

$$\mathbb{E} \left[\exp \left(\frac{\gamma(1+\epsilon)}{2} \beta_t^{d/q-\alpha} N_{2p,R\beta_t}^2(Z) \right) \right] \leq \frac{2}{\sqrt{1 - \frac{\gamma(1+\epsilon)^2 \beta_t^{d/q-\alpha}}{\rho(a, R, t)}}}.$$

Proof. (1) It suffices to take $f = (R\beta_t)^{-d/2p}$ in (9) to obtain the result.

- (2) By Hölder's inequality, for any f such that $N_{(2p)',R\beta_t}(f) = 1$,

$$\mathbb{P} \left[N_{2p,R\beta_t}(Z) \geq \sqrt{t\epsilon} \beta_t^{-d/(2q)} \right] \geq \mathbb{P} \left[\sum_{x \in \mathbb{T}_{R\beta_t}} f_x Z_x \geq \sqrt{t\epsilon} \beta_t^{-d/(2q)} \right].$$

Since $\sum_{x \in \mathbb{T}_{R\beta_t}} f_x Z_x$ is a real centered Gaussian variable with variance

$$\sigma_{a,R,t}^2(f) = \sum_{x,y \in \mathbb{T}_{R\beta_t}} G_{R\beta_t,\lambda_t}(x,y) f_x f_y,$$

we have:

$$\begin{aligned} \mathbb{P} \left[\|Z\|_{2p, R\beta_t} \geq \sqrt{t\epsilon} \beta_t^{-d/(2q)} \right] &\geq \frac{\sigma_{a, R, t}(f) \beta_t^{d/(2q)}}{\sqrt{2\pi t\epsilon}} \left(1 - \frac{\sigma_{a, R, t}^2(f) \beta_t^{d/q}}{t\epsilon} \right)_+ \exp \left(-\frac{t\beta_t^{-d/q} \epsilon}{2\sigma_{a, R, t}^2(f)} \right) \\ &\geq \frac{\sigma_{a, R, t}(f) \beta_t^{d/(2q)}}{\sqrt{2\pi t\epsilon}} \left(1 - \frac{\rho_1(a, R, t) \beta_t^{d/q}}{t\epsilon} \right)_+ \exp \left(-\frac{t\beta_t^{-d/q} \epsilon}{2\sigma_{a, R, t}^2(f)} \right), \end{aligned}$$

where $\rho_1(a, R, t) = \sup \left\{ \sigma_{a, R, t}^2(f), N_{(2p)', R\beta_t}(f) = 1 \right\}$. Taking the supremum over f we obtain that $\forall a, R, t, \epsilon > 0$,

$$\begin{aligned} &\mathbb{P} \left[N_{2p, R\beta_t}(Z) \geq \sqrt{t\epsilon} \beta_t^{-d/(2q)} \right] \\ &\geq \frac{\sqrt{\rho_1(a, R, t)} \beta_t^{d/(2q)}}{\sqrt{2\pi t\epsilon}} \left(1 - \frac{\rho_1(a, R, t) \beta_t^{d/q}}{t\epsilon} \right)_+ \exp \left(-\frac{t\beta_t^{-d/q} \epsilon}{2\rho_1(a, R, t)} \right). \end{aligned}$$

Then it suffices to prove that $\rho_1(a, R, t) = \frac{1}{\rho(a, R, t)}$ to have the result.

On one hand, by Hölder inequality,

$$\langle f, G_{R\beta_t, \lambda_t} f \rangle_{R\beta_t} \leq N_{2p, R\beta_t}(G_{R\beta_t, \lambda_t} f), \quad \forall f \text{ such that } N_{(2p)', R\beta_t}(f) = 1.$$

Since $G_{R\beta_t, \lambda_t}^{-1} = \lambda_t - A_{R\beta_t}$,

$$\begin{aligned} \langle f, G_{R\beta_t, \lambda_t} f \rangle_{R\beta_t} &= \langle G_{R\beta_t, \lambda_t}^{-1} G_{R\beta_t, \lambda_t} f, G_{R\beta_t, \lambda_t} f \rangle_{R\beta_t} \\ &= \lambda_t N_{2p, R\beta_t}^2(G_{R\beta_t, \lambda_t} f) - \langle A_{R\beta_t} G_{R\beta_t, \lambda_t} f, G_{R\beta_t, \lambda_t} f \rangle \\ &\geq \rho(a, R, t) N_{2p, R\beta_t}^2(G_{R\beta_t, \lambda_t} f). \end{aligned}$$

Therefore, for all f such that $N_{(2p)', R\beta_t}(f) = 1$, $\langle f, G_{R\beta_t, \lambda_t} f \rangle_{R\beta_t} \leq \frac{\langle f, G_{R\beta_t, \lambda_t} f \rangle_{R\beta_t}}{\rho(a, R, t)}$. Then, taking the supremum over f , $\rho_1(a, R, t) \leq 1/\rho(a, R, t)$.

On the other hand, let f_0 achieving the infimum in the definition of $\rho(a, R, t)$.

$$\begin{aligned} \rho_1(a, R, t) &= \sup_{N_{(2p)', R\beta_t}(f)=1} \{ \langle f, G_{R\beta_t, \lambda_t} f \rangle_{R\beta_t} \} \\ &\geq \frac{\langle G_{R\beta_t, \lambda_t}^{-1} f_0, f_0 \rangle_{R\beta_t}}{N_{(2p)', R\beta_t}^2(G_{R\beta_t, \lambda_t}^{-1} f_0)} = \frac{\rho(a, R, t)}{N_{(2p)', R\beta_t}^2(G_{R\beta_t, \lambda_t}^{-1} f_0)}. \end{aligned}$$

Furthermore, using the Lagrange multipliers method, we know that

$N_{(2p)', R\beta_t}(G_{R\beta_t, \lambda_t}^{-1} f_0) = \rho(a, R, t)$. Hence $\rho_1(a, R, t) \geq 1/\rho(a, R, t)$, and then $\rho_1(a, R, t) = 1/\rho(a, R, t)$.

(3) Let M be a median of $N_{2p, R\beta_t}(Z)$. We can easily see that

$$\begin{aligned} &\mathbb{E} \left[\exp \left(\frac{\gamma(1+\epsilon)}{2} \beta_t^{d/q-\alpha} N_{2p, R\beta_t}^2(Z) \right) \right] \\ &\leq \mathbb{E} \left[\exp \left(\frac{\gamma(1+\epsilon)^2}{2} \beta_t^{d/q-\alpha} |N_{2p, R\beta_t} - M|^2 \right) \right] \exp \left(\frac{\gamma(1+\epsilon)^2}{2\epsilon} \beta_t^{d/q-\alpha} M^2 \right) \end{aligned} \quad (20)$$

Let us now prove that under our assumptions we have $\beta_t^{d/q-\alpha} M^2 = o(t\beta_t^{-\alpha})$ which is equivalent to $M^2 = o(t\beta_t^{-d/q})$. Since $M = \left(\text{median} \left(\sum_{x \in \mathbb{T}_{R\beta_t}} Z_x^{2p} \right) \right)^{1/2p}$ and that for $X \geq$

0, $\text{median}(X) \leq 2\mathbb{E}[X]$, we get:

$$\begin{aligned} M^2 &= \left(\text{median} \left(\sum_{x \in \mathbb{T}_{R\beta_t}} Z_x^{2p} \right) \right)^{1/p} \leq \left(2\mathbb{E} \left[\sum_{x \in \mathbb{T}_{R\beta_t}} Z_x^{2p} \right] \right)^{1/p} \\ &\leq 2^{1/p} \left(\sum_{x \in \mathbb{T}_{R\beta_t}} G_{R\beta_t, \lambda_t}(0, 0)^p \mathbb{E}[Y^{2p}] \right)^{1/p}, \text{ where } Y \sim \mathcal{N}(0, 1) \\ &\leq C(p)(R\beta_t)^{d/p} G_{R\beta_t, \lambda_t}(0, 0). \end{aligned}$$

The asymptotic behavior of $G_{R\beta_t, \lambda_t}(0, 0)$ is given by the following lemma whose proof is postponed in section 4.

Lemma 9. *Behavior of $G_{R\alpha_t, \lambda_t}(0, 0)$.*

Assume **(H4)**, and that $\lambda_t = a\beta_t^{-\alpha}$ and $\beta_t \gg 1$. Then for any $a, R > 0$,

- (a) for $d < \alpha$, $G_{R\beta_t, \lambda_t}(0, 0) = O(\beta_t^{\alpha-d})$.
- (b) for $d = \alpha$, $G_{R\beta_t, \lambda_t}(0, 0) = O(\log \beta_t)$.
- (c) for $d > \alpha$, $G_{R\beta_t, \lambda_t}(0, 0) = O(1)$.

Hence, for $d < \alpha$, $M^2 = O(\beta_t^{\alpha-d/q}) = o(t\beta_t^{-d/q})$ as soon as $\beta_t^\alpha \ll t$. For $d = \alpha$, $M^2 = O(\beta_t^{d/p} \log \beta_t) = o(t\beta_t^{-d/q})$ as soon as $\beta_t^d \ll \frac{t}{\log(t)}$. For $d > \alpha$, $M^2 = O(\beta_t^{d/p}) = o(t\beta_t^{-d/q})$ as soon as $\beta_t^d \ll t$.

Let us now work on the expectation in (20).

$$\begin{aligned} &\mathbb{E} \left[\exp \left(\frac{\gamma(1+\epsilon)^2}{2} \beta_t^{d/q-\alpha} |N_{2p, R\beta_t} - M|^2 \right) \right] \\ &= 1 + \int_0^{+\infty} \frac{\gamma(1+\epsilon)^2}{2} \beta_t^{d/q-\alpha} \exp \left(\frac{\gamma(1+\epsilon)^2}{2} \beta_t^{d/q-\alpha} y \right) \mathbb{P}(|N_{2p, R\beta_t}(Z) - M|^2 > y) dy \end{aligned}$$

Using concentration inequalities for norms of Gaussian processes (see for instance lemma 3.1 in [29]), for all $y > 0$,

$$\mathbb{P}(|N_{2p, R\beta_t}(Z) - M| \geq \sqrt{y}) \leq 2\mathbb{P} \left(Y \geq \sqrt{y\rho(a, R, t)} \right), \text{ where } Y \sim \mathcal{N}(0, 1).$$

Hence,

$$\begin{aligned} &\mathbb{E} \left[\exp \left(\frac{\gamma(1+\epsilon)^2}{2} \beta_t^{d/q-\alpha} |N_{2p, R\beta_t} - M|^2 \right) \right] \\ &\leq 1 + 2 \int_0^{+\infty} \gamma(1+\epsilon)^2 \beta_t^{d/q-\alpha} \exp \left(\frac{\gamma(1+\epsilon)^2}{2} \beta_t^{d/q-\alpha} y \right) \mathbb{P} \left(Y \geq \sqrt{y\rho(a, R, t)} \right) dy \\ &= 2\mathbb{E} \left[\exp \left(\frac{\gamma(1+\epsilon)^2}{2\rho(a, R, t)} \beta_t^{d/q-\alpha} Y^2 \right) \right] - 1 \\ &\leq \frac{2}{\sqrt{1 - \frac{\gamma(1+\epsilon)^2 \beta_t^{d/q-\alpha}}{\rho(a, R, t)}}}, \text{ if } \gamma(1+\epsilon)^2 < \rho(a, R, t) \beta_t^{\alpha-d/q}. \end{aligned}$$

□

We now end the proof of the upper bound in (6). To begin with, we state the following lemma about the asymptotic behavior of $\rho(a, R, t)$. Its proof will be given in Section 4.

Proposition 10. *For positive real numbers a, R , let us define*

$$\rho(a, R) = \inf_{\substack{g: \mathbb{R}^d \rightarrow \mathbb{R} \\ R\text{-periodic}}} \left\{ a \|g\|_{2,R}^2 + R^d \sum_{z \in \mathbb{Z}^d} |\mathcal{F}_R(g)(z)|^2 \left| \frac{z}{R} \right|^\alpha, \|g\|_{2p,R} = 1 \right\}. \quad (21)$$

$$\rho(a) = \inf_{g: \mathbb{R}^d \rightarrow \mathbb{R}} \left\{ a \|g\|_2^2 + \int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2, \|g\|_{2p} = 1 \right\}. \quad (22)$$

Then, under assumptions of Theorem 1,

$$\liminf_{t \rightarrow +\infty} \beta_t^{\alpha-d/q} \rho(a, R, t) \geq \rho(a, R), \quad \text{and} \quad \liminf_{R \rightarrow +\infty} \rho(a, R) \geq \rho(a).$$

Moreover, $\rho(a) = a^{1-d/(\alpha q)} \rho_{\alpha,d,p}$, where $\rho_{\alpha,d,p}$ is defined by (4).

Let us fix $\theta, a, R, \gamma, \epsilon$ such that $\theta(1+\epsilon) < \gamma < (1+\epsilon)^{-2} \rho(a, R) \leq \liminf_{t \rightarrow +\infty} \beta_t^{\alpha-d/q} \rho(a, R, t)$. By lemma 9,

$$\limsup_{t \rightarrow +\infty} \frac{t}{\beta_t^\alpha} \log \left(\frac{\beta_t^d G_{R\beta_t, \lambda_t}(0, 0)}{t} \right) \leq 0.$$

By 3 of lemma 8,

$$\limsup_{t \rightarrow +\infty} \frac{\beta_t^\alpha}{t} \log \mathbb{E} \left[\exp \left(\frac{\gamma(1+\epsilon)}{2} \beta_t^{d/q-\alpha} N_{2p, R\beta_t}^2(Z) \right) \right] \leq 0, \quad (23)$$

and by 1 and 2 of lemma 8,

$$\liminf_{t \rightarrow +\infty} \frac{\beta_t^\alpha}{t} \log \mathbb{P} \left(N_{2p, R\beta_t}(Z) \geq \sqrt{2t\epsilon} \beta_t^{-d/(2q)} (1 + \sqrt{\epsilon}) \right) \geq -a\epsilon(1 + \sqrt{\epsilon})^2 R^{d/q}. \quad (24)$$

Therefore, it follows from (19) that for $\theta(1+\epsilon) < \gamma < (1+\epsilon)^{-2} \rho(a, R)$,

$$\limsup_{t \rightarrow +\infty} \frac{\beta_t^\alpha}{t} \log \mathbb{E} \left[\exp \left(\theta \beta_t^{d/q-\alpha} N_p(l_t) \right) \right] \leq a + \epsilon\gamma + a\epsilon(1 + \sqrt{\epsilon})^2 R^{d/q}. \quad (25)$$

Letting ϵ go to 0, we obtain that for $\theta < \gamma < \rho(a, R)$

$$\limsup_{t \rightarrow +\infty} \frac{\beta_t^\alpha}{t} \log \mathbb{E} \left[\exp \left(\theta \beta_t^{d/q-\alpha} N_p(l_t) \right) \right] \leq a. \quad (26)$$

Letting R go to $+\infty$, the same inequality is true for $\theta < \rho(a)$. Thus, for all $\theta > 0$,

$$\begin{aligned} \limsup_{t \rightarrow +\infty} \frac{\beta_t^\alpha}{t} \log \mathbb{E} \left[\exp \left(\theta \beta_t^{d/q-\alpha} N_p(l_t) \right) \right] &\leq \inf \{ a, \rho(a) > \theta \} \\ &= \inf \left\{ a > 0, a^{1-\frac{d}{\alpha q}} \rho_{\alpha,d,p} > \theta \right\} = \left(\frac{\theta}{\rho_{\alpha,d,p}} \right)^{\frac{\alpha q}{\alpha q - d}} \end{aligned}$$

which is the desired result.

4. ASYMPTOTIC BEHAVIOR OF $\rho(a, R, t)$ AND $G_{R\beta_t, \lambda_t}(0, 0)$.

The aim of this section is to prove Proposition 10 and Lemma 9.

4.1. **Proof of Lemma 9.** . Let p_s and $p_s^{R\beta_t}$ the transition probabilities of X_s and $X_s^{R\beta_t}$.

$$\begin{aligned} G_{R\beta_t, \lambda_t}(0, 0) &= \int_0^{+\infty} \exp(-\lambda_t s) p_s^{R\beta_t}(0, 0) ds \\ &\leq 1 + \int_1^{+\infty} \exp(-\lambda_t s) \sum_{z \in \mathbb{Z}^d} p_s(0, zR\beta_t) ds. \end{aligned}$$

By **(H4)**, there exists C such that

$$\forall s > 0, \forall z \in \mathbb{Z}^d, p_s(0, zR\beta_t) \leq C \left(s^{-d/\alpha} \wedge \frac{s}{|zR\beta_t|^{d+\alpha}} \right).$$

Then,

$$\begin{aligned} G_{R\beta_t, \lambda_t}(0, 0) &\leq 1 + C \int_1^{+\infty} \exp(-\lambda_t s) \left(s^{-d/\alpha} + \sum_{0 < |z| \leq \frac{s^{1/\alpha}}{R\beta_t}} s^{-d/\alpha} + \sum_{|z| > \frac{s^{1/\alpha}}{R\beta_t}} \frac{s}{|zR\beta_t|^{d+\alpha}} \right) ds \\ &\leq 1 + C \left(\int_1^{\beta_t^\alpha} s^{-d/\alpha} ds + \int_{\beta_t^\alpha}^{+\infty} \frac{\exp(-\lambda_t s)}{\beta_t^d} ds \right) + C \int_1^{+\infty} \frac{\exp(-\lambda_t s)}{(R\beta_t)^d} ds \\ &\leq 1 + C \int_1^{\beta_t^\alpha} s^{-d/\alpha} ds + C \frac{\exp(-\lambda_t \beta_t^\alpha)}{\lambda_t \beta_t^d} + C \frac{\exp(-\lambda_t)}{\lambda_t (R\beta_t)^d}. \end{aligned}$$

Remember that $\lambda_t = a\beta_t^{-\alpha}$. Then for a constant C depending on a and R , we have,

$$G_{R\beta_t, \lambda_t}(0, 0) \leq 1 + C \int_1^{\beta_t^\alpha} s^{-d/\alpha} ds + O\left(\beta_t^{\alpha-d}\right).$$

This inequality yields the result in the three cases $\alpha < d$, $\alpha = d$ and $\alpha > d$.

4.2. **Proof of Proposition 10. Asymptotic behavior of $\rho(a, R, t)$ when $t \rightarrow +\infty$.**

We want to prove that $\liminf_{t \rightarrow +\infty} \rho(a, R, t) \geq \rho(a, R)$.

4.2.1. *Expression of $\rho(a, R, t)$ in terms of Fourier transform.* Let us rewrite $\rho(a, R, t)$ defined by (9) in terms of Fourier transform. Let h be the function achieving the infimum in the definition of

$\rho(a, R, t)$.

$$\begin{aligned}
& \langle h, A_{R\beta_t} h \rangle_{R\beta_t} \\
&= \sum_{x \in \mathbb{T}_{R\beta_t}} h(x) \sum_{y \in \mathbb{T}_{R\beta_t}} \mu_{R\beta_t}(y-x)(h(y) - h(x)) \\
&= -N_{2,R\beta_t}^2(h) + \sum_{x \in \mathbb{T}_{R\beta_t}} \sum_{y \in \mathbb{T}_{R\beta_t}} h(x)\mu_{R\beta_t}(y-x)h(y) \\
&= -\frac{1}{(R\beta_t)^d} N_{2,R\beta_t}^2(F_{R\beta_t}(h)) \\
&\quad + \sum_{x \in \mathbb{T}_{R\beta_t}} \sum_{y \in \mathbb{T}_{R\beta_t}} h(x)h(y) \sum_{z \in \mathbb{T}_{R\beta_t}} \frac{1}{(R\beta_t)^d} F_{R\beta_t}(\mu_{R\beta_t})(z) \exp\left(2i\pi \frac{\langle z, y-x \rangle}{R\beta_t}\right) \\
&= -\frac{1}{(R\beta_t)^d} N_{2,R\beta_t}^2(F_{R\beta_t}(h)) + \frac{1}{(R\beta_t)^d} \sum_{z \in \mathbb{T}_{R\beta_t}} |F_{R\beta_t}(h)(z)|^2 F_{R\beta_t}(\mu_{R\beta_t})(z) \\
&= \frac{1}{(R\beta_t)^d} \sum_{z \in \mathbb{T}_{R\beta_t}} |F_{R\beta_t}(h)(z)|^2 \left(F(\mu)\left(\frac{z}{R\beta_t}\right) - 1\right).
\end{aligned}$$

We have thus shown that

$$\beta_t^{\alpha-d/q} \rho(a, R, t) = a\beta_t^{-\frac{d}{q}} N_{2,R\beta_t}^2(h) + \frac{\beta_t^{\alpha-\frac{d}{q}}}{(R\beta_t)^d} \sum_{z \in \mathbb{T}_{R\beta_t}} |F_{R\beta_t}(h)(z)|^2 (1 - F(\mu)\left(\frac{z}{R\beta_t}\right)). \quad (27)$$

We call $a\beta_t^{-\frac{d}{q}} N_{2,R\beta_t}^2(h)$ the 2-norm part and the second term in the right-hand side of (27) the gradient part. We set for $x \in \mathbb{R}^d$,

$$g_R(x) = \beta_t^{d/(2p)} \sum_{k \in \mathbb{T}_{R\beta_t}} h(k) \varphi(\beta_t x - k),$$

where φ is the $R\beta_t$ -periodic function from \mathbb{R}^d to \mathbb{C} , whose Fourier's coefficients are

$$\mathcal{F}_{R\beta_t}(\varphi)(n) = \begin{cases} \frac{1}{(R\beta_t)^d} \frac{\sqrt{1-F(\mu)\left(\frac{n}{R\beta_t}\right)}}{\left|\frac{n}{R\beta_t}\right|^{\alpha/2}} \mathbb{I}_{\left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}\right]^d}(n) & \text{if } n \neq 0, \\ \frac{1}{(R\beta_t)^d} & \text{otherwise,} \end{cases}$$

where for any $a, b \in \mathbb{R}$, $\llbracket a, b \rrbracket^d = [a, b]^d \cap \mathbb{Z}^d$.

Note that μ being a symmetric probability measure, $F(\mu)$ is real valued and $|F(\mu)| \leq 1$, so that $\mathcal{F}_{R\beta_t}(\varphi)(n)$ is well-defined. By definition, g_R is a R -periodic function defined on \mathbb{R}^d and for all $z \in \mathbb{Z}^d$,

$$\mathcal{F}_R(g_R)(z) = \beta_t^{d/2p} F_{R\beta_t}(h)(z) \mathcal{F}_{R\beta_t}(\varphi)(z).$$

g_R is our candidate to achieve the infimum in the definition (21) of $\rho(a, R)$.

4.2.2. *Gradient part.* The function g_R was built to preserve the gradient part. Indeed,

$$\begin{aligned}
& \frac{\beta_t^{\alpha-\frac{d}{q}}}{(R\beta_t)^d} \sum_{z \in \mathbb{T}_{R\beta_t}} |F_{R\beta_t}(h)(z)|^2 \left(1 - F(\mu)\left(\frac{z}{R\beta_t}\right)\right) \\
&= \frac{\beta_t^{\alpha-\frac{d}{q}}}{(R\beta_t)^d} \sum_{z \in \left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}\right]^d} |F_{R\beta_t}(h)(z)|^2 \mathcal{F}_{R\beta_t}(\varphi)(z)^2 (R\beta_t)^{2d} \left|\frac{z}{R\beta_t}\right|^\alpha \\
&= R^d \sum_{z \in \mathbb{Z}^d} |\mathcal{F}_R(g_R)(z)|^2 \left|\frac{z}{R}\right|^\alpha. \tag{28}
\end{aligned}$$

4.2.3. *2-norm part.* We work now on the 2-norm part. By Parseval's equality,

$$\begin{aligned}
\|g_R\|_{2,R}^2 &= R^d N_2^2(\mathcal{F}_R(g_R)) = R^d \sum_{z \in \left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}\right]^d} \beta_t^{d/p} |F_{R\beta_t}(h)(z)|^2 \left|\mathcal{F}_{R\beta_t}(\varphi)\left(\frac{z}{R\beta_t}\right)\right|^2 \\
&= \frac{R^d \beta_t^{d/p}}{(R\beta_t)^{2d}} N_{2,R\beta_t}^2(F_{R\beta_t}(h)) + \frac{R^d \beta_t^{d/p}}{(R\beta_t)^{2d}} \sum_{z \in \left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}\right]^d, z \neq 0} |F_{R\beta_t}(h)(z)|^2 \left[\frac{1 - F(\mu)\left(\frac{z}{R\beta_t}\right)}{\left|\frac{z}{R\beta_t}\right|^\alpha} - 1 \right] \\
&= \beta_t^{-d/q} N_{2,R\beta_t}^2(h) + \frac{R^d \beta_t^{d/p}}{(R\beta_t)^{2d}} \sum_{z \in \left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}\right]^d, z \neq 0} |F_{R\beta_t}(h)(z)|^2 \left[\frac{1 - F(\mu)\left(\frac{z}{R\beta_t}\right)}{\left|\frac{z}{R\beta_t}\right|^\alpha} - 1 \right].
\end{aligned}$$

Under assumptions **(H1)** and **(H2)** $F(\mu)(u) = 1 - |u|^\alpha + o(|u|^\alpha)$ (see for instance Feller [21], Chapter XVI.5). So, for any $\epsilon > 0$, one can find $\delta > 0$ such that for all $|z| < \delta$, we have $\left|\frac{1 - F(\mu)(z)}{|z|^\alpha} - 1\right| \leq \epsilon$. Then,

$$\begin{aligned}
& \frac{R^d \beta_t^{d/p}}{(R\beta_t)^{2d}} \sum_{z \in \left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}\right]^d, z \neq 0} |F_{R\beta_t}(h)(z)|^2 \left[\frac{1 - F(\mu)\left(\frac{z}{R\beta_t}\right)}{\left|\frac{z}{R\beta_t}\right|^\alpha} - 1 \right] \\
&\leq \frac{R^d \beta_t^{d/p}}{(R\beta_t)^{2d}} \left(\epsilon \sum_{z/\left|\frac{z}{R\beta_t}\right| < \delta} |F_{R\beta_t}(h)(z)|^2 + \delta^{-\alpha} \sum_{z/\delta \leq \left|\frac{z}{R\beta_t}\right| \leq \frac{1}{2}} |F_{R\beta_t}(h)(z)|^2 \left|1 - F(\mu)\left(\frac{z}{R\beta_t}\right)\right| \right) \\
&\leq \epsilon \beta_t^{-d/q} N_{2,R\beta_t}^2(h) + \delta^{-\alpha} R^d \beta_t^{d/p} \sum_{z \in \mathbb{Z}^d} |F_{R\beta_t}(h)(z) \mathcal{F}_{R\beta_t}(\varphi)(z)|^2 \left|\frac{z}{R\beta_t}\right|^\alpha \\
&= \epsilon \beta_t^{-d/q} N_{2,R\beta_t}^2(h) + \delta^{-\alpha} R^d \beta_t^{-\alpha} \sum_{z \in \mathbb{Z}^d} |\mathcal{F}_R(g_R)(z)|^2 \left|\frac{z}{R}\right|^\alpha.
\end{aligned}$$

Moreover, by (28) and (27) it is easy to see that

$$R^d \sum_{z \in \mathbb{Z}^d} |\mathcal{F}_R(g_R)(z)|^2 \left|\frac{z}{R}\right|^\alpha \leq \beta_t^{\alpha-d/q} \rho(a, R, t). \tag{29}$$

Hence, $\forall \epsilon > 0, \exists \delta > 0$ such that

$$\|g_R\|_{2,R}^2 \leq (1 + \epsilon) \beta_t^{-d/q} N_{2,R\beta_t}^2(h) + (\delta \beta_t)^{-\alpha} \left(\beta_t^{\alpha-d/q} \rho(a, R, t) \right). \tag{30}$$

4.2.4. *2p-norm part.* We work now on the most difficult part, that is the $2p$ -norm. Using Fourier inversion formula,

$$g_R(x) = \beta_t^{d/(2p)} \sum_{n \in \mathbb{Z}^d} F_{R\beta_t}(h)(n) \mathcal{F}_{R\beta_t}(\varphi)(n) \exp\left(2i\pi \frac{\langle x, n \rangle}{R}\right).$$

$$\|g_R\|_{2p,R} \geq \frac{\beta_t^{d/(2p)}}{(R\beta_t)^d} \left\| \sum_{n \in \left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}\right]^d} F_{R\beta_t}(h)(n) \exp\left(2i\pi \frac{\langle \cdot, n \rangle}{R}\right) \right\|_{2p,R}$$

$$- \beta_t^{d/(2p)} \left\| \sum_{n \in \left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}\right]^d} F_{R\beta_t}(h)(n) \left(\mathcal{F}_{R\beta_t}(\varphi)(n) - \frac{1}{(R\beta_t)^d} \right) \exp\left(2i\pi \frac{\langle \cdot, n \rangle}{R}\right) \right\|_{2p,R} \quad (31)$$

We want to prove that the first term is close to the $2p$ -norm of h , and that the second term is negligible. We first work on the second term in (31). Let $\delta_t \xrightarrow[t \rightarrow +\infty]{} 0$ to be chosen later. We first use inversion formula of Proposition 5, Young inequality of Proposition 6 and we again cut the sum in two parts. Denoting by $(2p)'$ the conjugate exponent of $2p > 2$, we are led to

$$\beta_t^{d/(2p)} \left\| \sum_{n \in \left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}\right]^d} F_{R\beta_t}(h)(n) \left(\mathcal{F}_{R\beta_t}(\varphi)(n) - \frac{1}{(R\beta_t)^d} \right) \exp\left(2i\pi \frac{\langle \cdot, n \rangle}{R}\right) \right\|_{2p,R}$$

$$\leq (R\beta_t)^{d/(2p)} N_{(2p)', R\beta_t} \left(F_{R\beta_t}(h)(\cdot) \mathbf{1}_{\left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}\right]^d}(\cdot) \left(\mathcal{F}_{R\beta_t}(\varphi)(\cdot) - \frac{1}{(R\beta_t)^d} \right) \right) \leq T_1 + T_2,$$

$$\text{with } T_1 = (R\beta_t)^{d/(2p)} \left(\sum_{n/|\frac{n}{R\beta_t}| \leq \delta_t} \left| F_{R\beta_t}(h)(n) \left(\mathcal{F}_{R\beta_t}(\varphi)(n) - \frac{1}{(R\beta_t)^d} \right) \right|^{(2p)'} \right)^{1/(2p)'}$$

$$T_2 = (R\beta_t)^{d/(2p)} \left(\sum_{n/|\frac{n}{R\beta_t}| > \delta_t} \left| F_{R\beta_t}(h)(n) \left(\mathcal{F}_{R\beta_t}(\varphi)(n) - \frac{1}{(R\beta_t)^d} \right) \right|^{(2p)'} \right)^{1/(2p)'}$$

Let us focus on the first term T_1 . Set $\epsilon(u) = |u|^{-\alpha/2} \sqrt{1 - F(\mu)(u)} - 1$. By our assumptions on μ , $\lim_{u \rightarrow 0} \epsilon(u) = 0$. Using Hölder inequality ($(2p)' < 2$) and Parseval's equality,

$$T_1 \leq \frac{(R\beta_t)^{d/(2p)}}{(R\beta_t)^d} \left(\sum_{n/0 < |\frac{n}{R\beta_t}| \leq \delta_t} \left| F_{R\beta_t}(h)(n) \left(\frac{\sqrt{1 - F(\mu)\left(\frac{n}{R\beta_t}\right)}}{\left|\frac{n}{R\beta_t}\right|^{\alpha/2}} - 1 \right) \right|^{(2p)'} \right)^{1/(2p)'}$$

$$\leq \frac{(R\beta_t)^{d/(2p)}}{(R\beta_t)^d} N_{2, R\beta_t}(F_{R\beta_t}(h)) \left(\sum_{n/0 < |\frac{n}{R\beta_t}| \leq \delta_t} \left| \frac{\sqrt{1 - F(\mu)\left(\frac{n}{R\beta_t}\right)}}{\left|\frac{n}{R\beta_t}\right|^{\alpha/2}} - 1 \right|^{2q} \right)^{1/(2q)}$$

$$= \frac{(R\beta_t)^{d/(2p)}}{(R\beta_t)^{d/2}} N_{2, R\beta_t}(h) \left(\sum_{n/|\frac{n}{R\beta_t}| \leq \delta_t} \left| \epsilon\left(\frac{n}{R\beta_t}\right) \right|^{2q} \right)^{1/(2q)}$$

$$\leq \frac{1}{\sqrt{a}} \left((\beta_t \delta_t)^{d/(2q)} \sup_{|u| \leq \delta_t} |\epsilon(u)| \right) \sqrt{\beta_t^{\alpha-d/q} \rho(a, R, t)},$$

by (27). To be negligible compared to $\sqrt{\beta_t^{\alpha-d/q} \rho(a, R, t)}$, we have to choose δ_t such that

$$(\beta_t \delta_t)^{d/(2q)} \sup_{|u| \leq \delta_t} |\epsilon(u)| \xrightarrow{t \rightarrow +\infty} 0. \quad (32)$$

Let us turn to the second term T_2 . By assumption **(H3)**, there exists $C > 0$ such that for $|x| \leq \frac{1}{2}$, $|x|^{\alpha/2} \leq C \sqrt{1 - F(\mu)(x)}$. Then, using the fact that $\left| \frac{n}{R\beta_t} \right| \leq \frac{1}{2}$ and Hölder inequality,

$$\begin{aligned} T_2 &\leq C \frac{(R\beta_t)^{d/(2p)}}{(R\beta_t)^d} \left(\sum_{n/\frac{1}{2} \geq \left| \frac{n}{R\beta_t} \right| > \delta_t} \left| F_{R\beta_t}(h)(n) \frac{\sqrt{1 - F(\mu)\left(\frac{n}{R\beta_t}\right)}}{\left| \frac{n}{R\beta_t} \right|^{\alpha/2}} \right|^{(2p)'} \right)^{1/(2p)'} \\ &\leq C \frac{(R\beta_t)^{d/(2p)}}{(R\beta_t)^d} N_{2, R\beta_t} \left(F_{R\beta_t}(h)(\cdot) \sqrt{1 - F(\mu)\left(\frac{\cdot}{R\beta_t}\right)} \right) (R\beta_t)^{\alpha/2} \left(\sum_{n/|n| > R\beta_t \delta_t} |n|^{-\alpha q} \right)^{1/(2q)} \\ &\leq C (R\beta_t)^{\alpha/2 - d/(2q) - d/2} N_{2, R\beta_t} \left(F_{R\beta_t}(h)(\cdot) \sqrt{1 - F(\mu)\left(\frac{\cdot}{R\beta_t}\right)} \right) \left((R\beta_t \delta_t)^{d - \alpha q} \right)^{1/(2q)}. \end{aligned}$$

Using (27), we have therefore

$$T_2 \leq C \delta_t^{\frac{1}{2}(d/q - \alpha)} \sqrt{\rho(a, R, t)}.$$

To be negligible compared to $\sqrt{\beta_t^{\alpha-d/q} \rho(a, R, t)}$, it is enough to choose δ_t such that

$$\beta_t \delta_t \rightarrow +\infty. \quad (33)$$

Note that conditions (32) and (33) are compatible. Indeed, set $\eta(x) = x \sup_{|u| \leq x} |\epsilon(u)|^{2q/d}$ for $x \in \mathbb{R}^+$. η is an increasing function, and (32) and (33) are equivalent to $\beta_t^{-1} \ll \delta_t \ll \eta^{-1}(\beta_t^{-1})$. Such a δ_t can be found as soon as $\lim_{x \rightarrow 0} \eta^{-1}(x)/x = +\infty$, which is the case since $\lim_{x \rightarrow 0} x/\eta(x) = +\infty$. We have now succeeded to control the second term in (31) by proving that one can find u_t , $\lim_{t \rightarrow +\infty} u_t = 0$ such that

$$\beta_t^{d/(2p)} \left\| \sum_{n \in \left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2} \right]^d} F_{R\beta_t}(h)(n) \left(\mathcal{F}_{R\beta_t}(\varphi)(n) - \frac{1}{(R\beta_t)^d} \right) e^{2i\pi \frac{\langle \cdot, n \rangle}{R}} \right\|_{2p, R} \leq u_t \sqrt{\beta_t^{\alpha-d/q} \rho(a, R, t)}. \quad (34)$$

It remains to control the first term of the sum in (31). Performing the change of variable $x \rightarrow x\beta_t$,

$$\frac{\beta_t^{d/(2p)}}{(R\beta_t)^d} \left\| \sum_{n \in \left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2} \right]^d} F_{R\beta_t}(h)(n) e^{2i\pi \frac{\langle \cdot, n \rangle}{R}} \right\|_{2p, R} = \left\| \frac{1}{(R\beta_t)^d} \sum_{n \in \left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2} \right]^d} F_{R\beta_t}(h)(n) e^{2i\pi \frac{\langle \cdot, n \rangle}{R\beta_t}} \right\|_{2p, R\beta_t}.$$

For $x \in \mathbb{R}^d$, let

$$f(x) := \frac{1}{(R\beta_t)^d} \sum_{n \in \left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2} \right]^d} F_{R\beta_t}(h)(n) \exp\left(2i\pi \frac{\langle x, n \rangle}{R\beta_t}\right).$$

Note that $f(x) = h(x)$, $\forall x \in \mathbb{T}_{R\beta_t}$. We define an approximation of f by:

$$\forall x \in \mathbb{R}^d, \bar{f}(x) = \sum_{k \in \left[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2} \right]^d} \bar{f}_k \mathbb{1}_{Q_k}(x)$$

where $Q_k = [k - \frac{1}{2}; k + \frac{1}{2}]^d$ is the unit cube centered on k , and $\bar{f}_k = \frac{1}{|Q_k|} \int_{Q_k} f(x) dx$ is the mean of f on Q_k . For any $x \in \mathbb{R}^d$, we denote by $\lfloor x \rfloor$ the unique $k \in \mathbb{Z}^d$ such that $x \in Q_k$. Note that $\forall x \in Q_k, \lfloor x \rfloor = k$ and $f(\lfloor x \rfloor) = h(k)$. Introducing the $R\beta_t$ -periodic functions, $f(\lfloor \cdot \rfloor)$ and $\bar{f}(\cdot)$, we have that:

$$\|f\|_{2p, R\beta_t} \geq \|f(\lfloor \cdot \rfloor)\|_{2p, R\beta_t} - \|\bar{f} - f(\lfloor \cdot \rfloor)\|_{2p, R\beta_t} - \|f - \bar{f}\|_{2p, R\beta_t}. \quad (35)$$

The first term is exactly the $2p$ -norm of h . Let us consider the second one. For $x \in \mathbb{R}^d$,

$$\begin{aligned} \bar{f}(x) - f(\lfloor x \rfloor) &= \frac{1}{(R\beta_t)^d} \sum_{n \in \lfloor [-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d} F_{R\beta_t}(h)(n) \underbrace{\int_{Q_0} e^{2i\pi \frac{\langle n, y \rangle}{R\beta_t}} dy}_{C\left(\frac{n}{R\beta_t}\right)} \sum_{k \in \lfloor [-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d} e^{2i\pi \frac{\langle n, k \rangle}{R\beta_t}} \mathbb{1}_{Q_k}(x) \\ &\quad - \frac{1}{(R\beta_t)^d} \sum_{n \in \lfloor [-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d} F_{R\beta_t}(h)(n) e^{2i\pi \frac{\langle n, \lfloor x \rfloor \rangle}{R\beta_t}} \\ &= \frac{1}{(R\beta_t)^d} \sum_{n \in \lfloor [-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d} F_{R\beta_t}(h)(n) \left(C\left(\frac{n}{R\beta_t}\right) - 1 \right) \exp\left(2i\pi \frac{\langle n, \lfloor x \rfloor \rangle}{R\beta_t}\right) \end{aligned}$$

Therefore, using inversion formula of Proposition 3, Young inequality of Proposition 6 and Hölder inequality:

$$\begin{aligned} &\|\bar{f} - f(\lfloor \cdot \rfloor)\|_{2p, R\beta_t} \\ &\leq (R\beta_t)^{-d/(2p)'} N_{(2p)'} \left(F_{R\beta_t}(h)(\cdot) \mathbb{1}_{\lfloor [-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d}(\cdot) \left(C\left(\frac{\cdot}{R\beta_t}\right) - 1 \right) \right) \\ &\leq (R\beta_t)^{-d/(2p)'} N_{2, R\beta_t} \left(F_{R\beta_t}(h)(\cdot) \sqrt{1 - F(\mu)\left(\frac{\cdot}{R\beta_t}\right)} \right) N_{2q} \left(\mathbb{1}_{\lfloor [-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d}(\cdot) \frac{C\left(\frac{\cdot}{R\beta_t}\right) - 1}{\sqrt{1 - F(\mu)\left(\frac{\cdot}{R\beta_t}\right)}} \right). \end{aligned}$$

Let us have a look at $N_{2q} \left(\mathbb{1}_{\lfloor [-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d}(\cdot) \frac{C\left(\frac{\cdot}{R\beta_t}\right) - 1}{\sqrt{1 - F(\mu)\left(\frac{\cdot}{R\beta_t}\right)}} \right)$. A simple computation shows that

$$C(x) = \prod_{j=1}^d e^{i\pi x_j} \frac{\sin(\pi x_j)}{\pi x_j}, \quad C(x) \underset{0}{\sim} 1 + O(|x|).$$

Combining this with the fact that $1 - F(\mu)(x) \underset{0}{\sim} |x|^\alpha$, we have that $\frac{C(x) - 1}{\sqrt{1 - F(\mu)(x)}} = O(|x|^{1-\alpha/2})$.

Moreover, by **(H3)**, $F(\mu)(x) = 1 \Leftrightarrow x \in \mathbb{Z}^d$. Then the function $x \rightarrow \frac{C(x) - 1}{\sqrt{1 - F(\mu)(x)}}$ is uniformly

bounded for $|x| \leq \frac{1}{2}$ and $N_{2q} \left(\mathbb{1}_{\lfloor [-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d}(\cdot) \frac{C\left(\frac{\cdot}{R\beta_t}\right) - 1}{\sqrt{1 - F(\mu)\left(\frac{\cdot}{R\beta_t}\right)}} \right) = O((R\beta_t)^{d/(2q)})$. Therefore, using (27),

$$\|\bar{f} - f(\lfloor \cdot \rfloor)\|_{2p, R\beta_t} \leq C \sqrt{\beta_t^{-\alpha+d/q}} \sqrt{\beta_t^{\alpha-d/q} \rho(a, R, t)}. \quad (36)$$

We turn now to the third term of (35). By Poincaré's inequality, as the unit cube is a Lipschitz's domain,

$$\begin{aligned} \|f - \bar{f}\|_{2p, R\beta_t}^{2p} &= \int_{[-R\beta_t/2, R\beta_t/2]^d} |f(x) - \bar{f}(x)|^{2p} dx = \sum_{k \in \mathbb{I}_{[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d}} \int_{Q_k} |f(x) - \bar{f}_k|^{2p} dx \\ &\leq C \sum_{k \in \mathbb{I}_{[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d}} \|\nabla f\|_{2p, Q_k}^{2p} = C \|\nabla f\|_{2p, R\beta_t}^{2p}, \end{aligned}$$

where C depends only on d, p and $|Q_0|$. Hence, denoting by ψ_j the j -th coordinate function $\psi_j(n) = n_j$, and using inversion formula of Proposition 5, Young's and Hölder's inequalities,

$$\begin{aligned} \|f - \bar{f}\|_{2p, R\beta_t} &\leq \frac{C}{(R\beta_t)^d} \sum_{j=1}^d \left\| \sum_{n \in \mathbb{I}_{[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d}} F_{R\beta_t}(h)(n) \frac{2i\pi\psi_j(n)}{R\beta_t} \exp\left(2i\pi \frac{\langle n, \cdot \rangle}{R\beta_t}\right) \right\|_{2p, R\beta_t} \\ &\leq C \frac{(R\beta_t)^{d/(2p)}}{(R\beta_t)^d} \sum_{j=1}^d N_{(2p)'} \left(F_{R\beta_t}(h)(\cdot) \frac{2i\psi_j(\cdot)}{R\beta_t} \mathbb{I}_{\mathbb{I}_{[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d}(\cdot)} \right) \\ &\leq C \frac{(R\beta_t)^{d/(2p)}}{(R\beta_t)^d} N_{(2p)'} \left(F_{R\beta_t}(h)(\cdot) \frac{|\cdot|}{R\beta_t} \mathbb{I}_{\mathbb{I}_{[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d}(\cdot)} \right) \\ &\leq C \frac{(R\beta_t)^{d/(2p)}}{(R\beta_t)^d} N_{2, R\beta_t} \left(F_{R\beta_t}(h)(\cdot) \sqrt{1 - F(\mu)\left(\frac{\cdot}{R\beta_t}\right)} \right) \\ &\quad N_{2q} \left(\frac{|\cdot|}{R\beta_t} \left(1 - F(\mu)\left(\frac{\cdot}{R\beta_t}\right)\right)^{-1/2} \mathbb{I}_{\mathbb{I}_{[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d}(\cdot)} \right) \end{aligned}$$

As previously, $1 - F(\mu)(x) \sim |x|^\alpha$ implies that $\frac{|x|}{\sqrt{1 - F(\mu)(x)}} \sim |x|^{1-\alpha/2}$, which is bounded around 0 since $\alpha \in]0, 2]$. By **(H3)**, we deduce that the function $x \rightarrow \frac{|x|}{\sqrt{1 - F(\mu)(x)}}$ is uniformly bounded for $|x| \leq 1/2$. Therefore, $N_{2q} \left(\frac{|\cdot|}{R\beta_t} \left(1 - F(\mu)\left(\frac{\cdot}{R\beta_t}\right)\right)^{-1/2} \mathbb{I}_{\mathbb{I}_{[-\frac{R\beta_t}{2}, \frac{R\beta_t}{2}]^d}(\cdot)} \right) = O((R\beta_t)^{d/(2q)})$, and using (27),

$$\|f - \bar{f}\|_{2p, R\beta_t} \leq C \sqrt{\beta_t^{-\alpha+d/q}} \sqrt{\beta_t^{\alpha-d/q} \rho(a, R, t)}. \quad (37)$$

Combining (35), (36), (37) and the fact that $N_{2p, R\beta_t}(h) = 1$, we have shown that

$$\|f\|_{2p, R\beta_t} \geq 1 - C \sqrt{\beta_t^{-\alpha+d/q}} \sqrt{\beta_t^{\alpha-d/q} \rho(a, R, t)}. \quad (38)$$

Putting (31), (34), (38) together, as $\alpha - d/q > 0$, we have controlled the $2p$ -norm of g_R by:

$$\|g_R\|_{2p, R} \geq 1 - u_t \sqrt{\beta_t^{\alpha-d/q} \rho(a, R, t)}, \quad (39)$$

for some positive function $(u_t, t \geq 0)$ satisfying $\lim_{t \rightarrow +\infty} u_t = 0$. Finally, $\forall R > 0, \forall \epsilon > 0, \exists \delta > 0$ such that

$$\begin{aligned} & \beta_t^{\alpha-d/q} \rho(a, R, t) \\ &= a \beta_t^{-\frac{d}{q}} N_{2, R\beta_t}^2(h) - \beta_t^{\alpha-\frac{d}{q}} \langle h, A_{R\beta_t} h \rangle_{R\beta_t} \\ &\geq \frac{1}{1+\epsilon} \left(a \|g_R\|_{2, R}^2 + R^d \sum_{z \in \mathbb{T}_{R\beta_t}} |\mathcal{F}_R(g_R)(z)|^2 \left| \frac{z}{R} \right|^\alpha \right) - \frac{a}{1+\epsilon} (\delta \beta_t)^{-\alpha} \left(\beta_t^{\alpha-d/q} \rho(a, R, t) \right) \\ &\geq \frac{\|g_R\|_{2p, R}^2}{1+\epsilon} \rho(a, R) - \frac{a}{1+\epsilon} (\delta \beta_t)^{-\alpha} \left(\beta_t^{\alpha-d/q} \rho(a, R, t) \right) \\ &\geq \frac{1 - u_t \sqrt{\beta_t^{\alpha-d/q} \rho(a, R, t)}}{1+\epsilon} \rho(a, R) - \frac{a}{1+\epsilon} (\delta \beta_t)^{-\alpha} \left(\beta_t^{\alpha-d/q} \rho(a, R, t) \right). \end{aligned}$$

We can assume that $\liminf_{t \rightarrow +\infty} \beta_t^{\alpha-d/q} \rho(a, R, t) < +\infty$. Otherwise, there is nothing to prove. Letting first $t \rightarrow +\infty$, then $\epsilon \rightarrow 0$ in the above inequality, we obtain that for all $R > 0$,

$$\liminf_{t \rightarrow +\infty} \beta_t^{\alpha-d/q} \rho(a, R, t) \geq \rho(a, R).$$

4.3. Proof of Proposition 10. Asymptotic behavior of $\rho(a, R)$ when $R \rightarrow +\infty$.

It remains now to prove that for any $a > 0$,

$$I := \liminf_{R \rightarrow +\infty} \rho(a, R) \geq \rho(a). \quad (40)$$

This statement was already proved in [7, 27] for $\alpha = 2$. We show it for $\alpha < 2$.

We assume that $I < +\infty$, otherwise there is nothing to prove. Let then $\epsilon > 0$. For any $R_0 > 0$, let $R \geq R_0$ and let g_R be such that $\|g_R\|_{2p, R} = 1$ and

$$a \|g_R\|_{2, R}^2 + R^d \sum_{z \in \mathbb{Z}^d} \left| \frac{z}{R} \right|^\alpha |\mathcal{F}_R(g_R)(z)|^2 \leq I + \epsilon \quad (41)$$

Note that any translation of order θ of g_R also satisfies (41). Indeed, setting $g_{R, \theta} = g_R(\theta + \cdot)$, and using that g_R is periodic, we get $\|g_{R, \theta}\|_{2p, R} = \|g_R\|_{2p, R}$, $\|g_{R, \theta}\|_{2, R} = \|g_R\|_{2, R}$, $\mathcal{F}_R(g_R(\theta + \cdot))(n) = \exp\left(2i\pi \frac{\langle n, \theta \rangle}{R}\right) \mathcal{F}_R(g_R)(n)$, and $\sum_{z \in \mathbb{Z}^d} \left| \frac{z}{R} \right|^\alpha |\mathcal{F}_R(g_R)(z)|^2 = \sum_{z \in \mathbb{Z}^d} \left| \frac{z}{R} \right|^\alpha |\mathcal{F}_R(g_{R, \theta})(z)|^2$.

Let $E_R = \left([0; \sqrt{R}] \cup [R - \sqrt{R}; R]\right)^d$ and $Q_R = [0, R]^d$.

$$\inf_{\theta \in Q_R} \int_{E_R} |g_{R, \theta}(x)|^{2p} dx \leq \frac{1}{R^d} \int_{Q_R} \int_{E_R} |g_{R, \theta}(x)|^{2p} dx d\theta = (2R^{-1/2})^d.$$

Therefore one can find θ such that $\int_{E_R} |g_{R, \theta}(x)|^{2p} dx \leq 3^d R^{-d/2}$, and we can assume without loss of generality that g_R also satisfies

$$\int_{E_R} |g_R(x)|^{2p} dx \leq 3^d R^{-d/2}. \quad (42)$$

Let ϕ_R be the function from \mathbb{R} to \mathbb{R} , with compact support in $[0; R]$, equal to 1 on $[0, R] \setminus ([0; \sqrt{R}] \cup [R - \sqrt{R}; R])$, and which is linear on $[0; \sqrt{R}]$ and $[R - \sqrt{R}; R]$. Let $\psi_R : \mathbb{R}^d \mapsto [0, 1]$ be defined by $\psi_R(x) = \prod_{i=1}^d \phi_R(x_i)$. Note that ψ_R has compact support in Q_R , is equal to 1 on $Q_R \setminus E_R$, and that for all $x \in \mathbb{R}^d$, $\|\nabla \psi_R(x)\| \leq \sqrt{d}/R$.

Let take $g(x) = g_R(x)\psi_R(x)$. g is our candidate to realize the infimum defining $\rho(a)$. Note that

$$\|g\|_2^2 = \int_{\mathbb{R}^d} g_R^2(x)\psi_R^2(x) dx \leq \int_{Q_R} g_R^2(x) dx. \quad (43)$$

$$\|g\|_{2p}^{2p} = \int_{\mathbb{R}^d} |g_R(x)|^{2p} \psi_R^{2p}(x) dx \geq \int_{Q_R \setminus E_R} |g_R(x)|^{2p} dx \geq 1 - 3^d R^{-d/2}, \quad (44)$$

where the last inequality comes from (42).

Let us now estimate $\int_{\mathbb{R}^d} |\mathcal{F}(g)(\omega)|^2 |\omega|^\alpha d\omega$. To begin with, note that

$$\int_{\mathbb{R}^d} |\mathcal{F}(g)(\omega)|^2 |\omega|^\alpha d\omega = c_{\alpha,d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(g(x) - g(y))^2}{|x - y|^{d+\alpha}} dx dy, \quad (45)$$

for some constant $c_{\alpha,d} \in]0; +\infty[$. Indeed, using Parseval's identity,

$$\begin{aligned} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(g(x) - g(y))^2}{|x - y|^{d+\alpha}} dx dy &= \int_{\mathbb{R}^d} \frac{1}{|z|^{d+\alpha}} \left(\int_{\mathbb{R}^d} (g(z+y) - g(y))^2 dy \right) dz \\ &= \int_{\mathbb{R}^d} \frac{1}{|z|^{d+\alpha}} \left(\int_{\mathbb{R}^d} |e^{-2\pi i \langle z; \omega \rangle} - 1|^2 |\mathcal{F}(g)(\omega)|^2 d\omega \right) dz \\ &= \int_{\mathbb{R}^d} |\mathcal{F}(g)(\omega)|^2 \left(\int_{\mathbb{R}^d} \frac{4 \sin^2(\pi \langle z; \omega \rangle)}{|z|^{d+\alpha}} dz \right) d\omega \end{aligned}$$

Let $H(\omega) = \int_{\mathbb{R}^d} \frac{4 \sin^2(\pi \langle z; \omega \rangle)}{|z|^{d+\alpha}} dz$. For any orthogonal matrix O of \mathbb{R}^d , $H(O\omega) = H(\omega)$, so that

$$H(\omega) = H(|\omega| e_1) = |\omega|^\alpha \int_{\mathbb{R}^d} \frac{4 \sin^2(\pi z_1)}{|z|^{d+\alpha}} dz := |\omega|^\alpha c_{\alpha,d},$$

where $c_{\alpha,d} \in]0; \infty[$ for $\alpha \in]0; 2[$. This gives (45). Similarly, one obtains that

$$R^d \sum_{z \in \mathbb{Z}^d} \left| \frac{z}{R} \right|^\alpha |\mathcal{F}_R(g_R)(z)|^2 = c_{\alpha,d} \int_{Q_R} \int_{\mathbb{R}^d} \frac{(g_R(x) - g_R(x+y))^2}{|y|^{d+\alpha}} dy dx, \quad (46)$$

for the same constant $c_{\alpha,d}$ as in (45). Now, $\forall \delta > 0$,

$$\begin{aligned} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(g(x) - g(y))^2}{|x - y|^{d+\alpha}} dx dy &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{[(g_R(x) - g_R(y))\psi_R(x) + g_R(y)(\psi_R(x) - \psi_R(y))]^2}{|x - y|^{d+\alpha}} dx dy \\ &\leq (1 + \delta) \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(g_R(x) - g_R(y))^2}{|x - y|^{d+\alpha}} \psi_R^2(x) dx dy + \frac{1 + \delta}{\delta} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} g_R^2(y) \frac{(\psi_R(x) - \psi_R(y))^2}{|x - y|^{d+\alpha}} dx dy \\ &\leq (1 + \delta) \int_{Q_R} \int_{\mathbb{R}^d} \frac{(g_R(x) - g_R(x+z))^2}{|z|^{d+\alpha}} dz dx + \frac{1 + \delta}{\delta} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} g_R^2(y) \frac{(\psi_R(x) - \psi_R(y))^2}{|x - y|^{d+\alpha}} dx dy. \end{aligned}$$

But, $\forall x, y \in \mathbb{R}^d$, $|\psi_R(x) - \psi_R(y)| \leq \min(\sqrt{\frac{d}{R}} |x - y|, 1)(\mathbb{1}_{Q_R}(x) + \mathbb{1}_{Q_R}(y))$. This leads to

$$\begin{aligned} &\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} g_R^2(y) \frac{(\psi_R(x) - \psi_R(y))^2}{|x - y|^{d+\alpha}} dx dy \\ &\leq \int_{Q_R} g_R^2(y) \left(\int_{\mathbb{R}^d} \frac{\min(\sqrt{\frac{d}{R}} |z|, 1)^2}{|z|^{d+\alpha}} dz \right) dy + \int_{Q_R} \int_{\mathbb{R}^d} g_R^2(x+z) \frac{\min(\sqrt{\frac{d}{R}} |z|, 1)^2}{|z|^{d+\alpha}} dz dx \\ &= 2 \left(\int_{Q_R} g_R^2(y) dy \right) \left(\int_{\mathbb{R}^d} \frac{\min(\sqrt{\frac{d}{R}} |z|, 1)^2}{|z|^{d+\alpha}} dz \right). \end{aligned}$$

It is easy to check that there exists a constant C (only depending on α and d) such that $\int_{\mathbb{R}^d} \frac{\min(\sqrt{\frac{d}{R}}|z|, 1)^2}{|z|^{d+\alpha}} dz \leq CR^{-\alpha/2}$. It follows then from (41), (45), (46) that $\exists C$ such that $\forall \delta > 0$,

$$\int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2 d\omega \leq (1 + \delta)R^d \sum_{z \in \mathbb{Z}^d} \left| \frac{z}{R} \right|^\alpha |\mathcal{F}_R(g_R)(z)|^2 + C \frac{1 + \delta}{\delta} \frac{I + \epsilon}{a} R^{-\alpha/2}. \quad (47)$$

Combining (43), (44), (47), we get that $\exists C$ such that $\forall \epsilon > 0, \forall R_0 > 0, \exists R \geq R_0$ such that $\forall \delta > 0$,

$$\begin{aligned} & \inf \left\{ a \|f\|_2^2 + \int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(f)(\omega)|^2 d\omega; \|f\|_{2p} = 1 \right\} \\ & \leq \frac{a \|g\|_2^2 + \int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2 d\omega}{\|g\|_{2p}^2} \\ & \leq (1 + \delta) \frac{a \|g_R\|_{2,R}^2 + R^d \sum_{z \in \mathbb{Z}^d} \left| \frac{z}{R} \right|^\alpha |\mathcal{F}_R(g_R)(z)|^2}{1 - 3^d R^{-d/2}} + C \frac{1 + \delta}{\delta} \frac{I + \epsilon}{a} \frac{R^{-\alpha/2}}{1 - 3^d R^{-d/2}} \\ & \leq (1 + \delta) \frac{I + \epsilon}{1 - 3^d R^{-d/2}} + C \frac{1 + \delta}{\delta} \frac{I + \epsilon}{a} \frac{R^{-\alpha/2}}{1 - 3^d R^{-d/2}} \end{aligned}$$

This ends the proof of (40) by first letting R_0 go to ∞ , and then δ and ϵ go to 0.

To see that $\rho(a) = a^{1-d/(\alpha q)} \rho_{\alpha, d, p}$, use the transformation $g \mapsto g_\lambda = \lambda^{d/(2p)} g(\lambda \cdot)$ ($\lambda > 0$), and optimize over λ . This ends the proof of Proposition 10.

5. EXPONENTIAL MOMENTS LOWER BOUND.

The aim of this section is to prove the lower bound in (6). This part is inspired by the proof of Theorem 1.3 in [14]. Let us assume for a while the following theorem:

Theorem 11. *For any continuous function f on \mathbb{R}^d with compact support,*

$$\begin{aligned} & \liminf_{t \rightarrow +\infty} \frac{\beta_t^\alpha}{t} \log \mathbb{E} \exp \left(\beta_t^{-\alpha} \int_0^t f \left(\frac{X_s}{\beta_t} \right) ds \right) \\ & \geq \sup \left\{ \int_{\mathbb{R}^d} f(x) g(x)^2 dx - \int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2 d\omega, \|g\|_2 = 1, g \geq 0 \right\}. \end{aligned}$$

It follows from $I_t = \beta_t^d \int_{\mathbb{R}^d} l_t(\lfloor \beta_t x \rfloor)^p dx$ that

$$\frac{\beta_t^\alpha}{t} \log \mathbb{E} \exp \left(\theta \beta_t^{d/q - \alpha} N_p(l_t) \right) = \frac{\beta_t^\alpha}{t} \log \mathbb{E} \exp \left(\theta \beta_t^{d-\alpha} \left(\int_{\mathbb{R}^d} l_t(\lfloor \beta_t x \rfloor)^p dx \right)^{1/p} \right). \quad (48)$$

Let f be a continuous function with compact support such that $\|f\|_q = 1$. By Hölder inequality we have that:

$$\begin{aligned} \left(\int_{\mathbb{R}^d} l_t(\lfloor \beta_t x \rfloor)^p dx \right)^{1/p} & \geq \int_{\mathbb{R}^d} f(x) l_t(\lfloor \beta_t x \rfloor) dx \\ & = \beta_t^{-d} \int_{\mathbb{R}^d} f \left(\frac{x}{\beta_t} \right) l_t(\lfloor x \rfloor) dx \\ & = \beta_t^{-d} \sum_{x \in \mathbb{Z}^d} l_t(x) \int_{[x, x+1]^d} f \left(\frac{y}{\beta_t} \right) dy. \end{aligned}$$

As f is uniformly continuous and $\beta_t \rightarrow +\infty$,

$$\begin{aligned} \left(\int_{\mathbb{R}^d} l_t(\lfloor \beta_t x \rfloor)^p dx \right)^{1/p} &\geq \beta_t^{-d} \sum_{x \in \mathbb{Z}^d} l_t(x) \left(f\left(\frac{x}{\beta_t}\right) + o(1) \right) \\ &= \beta_t^{-d} \left(\int_0^t f\left(\frac{X_s}{\beta_t}\right) ds + o(1) \right). \end{aligned} \quad (49)$$

Combining (48) and (49),

$$\frac{\beta_t^\alpha}{t} \log \mathbb{E} \exp \left(\theta \beta_t^{d/q-\alpha} N_p(l_t) \right) \geq \frac{\beta_t^\alpha}{t} \log \mathbb{E} \exp \left(\beta_t^{-\alpha} \left(\int_0^t \theta f\left(\frac{X_s}{\beta_t}\right) ds + o(1) \right) \right)$$

Then taking the limit over t and using Theorem 11,

$$\begin{aligned} \liminf_{t \rightarrow +\infty} \frac{\beta_t^\alpha}{t} \log \mathbb{E} \exp \left(\theta \beta_t^{d/q-\alpha} N_p(l_t) \right) \\ \geq \sup \left\{ \theta \int_{\mathbb{R}^d} f(x) g(x)^2 dx - \int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2 d\omega, \|g\|_2 = 1, g \geq 0 \right\}. \end{aligned}$$

We take the supremum over all functions f with compact support such that $\|f\|_q = 1$ and exchange the two supremum. This immediately leads to

$$\liminf_{t \rightarrow +\infty} \frac{\beta_t^\alpha}{t} \log \mathbb{E} \exp \left(\theta \beta_t^{d/q-\alpha} N_p(l_t) \right) \geq \sup \left\{ \theta \|g\|_{2p}^2 - \int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2 d\omega, \|g\|_2 = 1, g \geq 0 \right\}. \quad (50)$$

Note that by (45), $\int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2 d\omega \geq \int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(|g|)(\omega)|^2 d\omega$, so that we can remove the constraint $g \geq 0$ in the supremum in (50). All that remains to show now is

$$\sup \left\{ \theta \|g\|_{2p}^2 - \int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2 d\omega, \|g\|_2 = 1 \right\} = \left(\frac{\theta}{\rho_{\alpha,d,p}} \right)^{\frac{\alpha q}{\alpha q - d}}. \quad (51)$$

Using the transformation $g \mapsto \lambda^{d/2} g(\lambda)$ ($\lambda > 0$) and optimizing over λ yields

$$\begin{aligned} \sup \left\{ \theta \|g\|_{2p}^2 - \int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2 d\omega, \|g\|_2 = 1 \right\} \\ = \frac{\alpha q - d}{\alpha q} \left(\frac{d}{\alpha q} \right)^{\frac{d}{\alpha q - d}} \theta^{\frac{\alpha q}{\alpha q - d}} \sup \left\{ \|g\|_{2p}^{2\frac{\alpha q}{\alpha q - d}} \left(\int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2 d\omega \right)^{-\frac{d}{\alpha q - d}}, \|g\|_2 = 1 \right\}. \end{aligned}$$

The expression in the supremum above is invariant under the transformation $g \mapsto \lambda^{d/2} g(\lambda)$. We can therefore freely add the constraint $\|g\|_{2p} = 1$. This gives (51) and ends the proof of the lower bound in (6).

It remains to prove Theorem 11.

Proof of Theorem 11.

We first split the time interval $[0, t]$ into interval of length $\lfloor \beta_t^\alpha \rfloor$. We set $\gamma_t = \lfloor \frac{t}{\beta_t^\alpha} \rfloor$.

$$\mathbb{E} \exp \left(\beta_t^{-\alpha} \int_0^t f\left(\frac{X_s}{\beta_t}\right) ds \right) \geq \exp(-2\|f\|_\infty) \mathbb{E} \exp \left(\beta_t^{-\alpha} \int_{\lfloor \beta_t^\alpha \rfloor}^{\lfloor \beta_t^\alpha \rfloor \gamma_t} f\left(\frac{X_s}{\beta_t}\right) ds \right).$$

Let us introduce the two following operators. For any ξ in $\ell^2(\mathbb{Z}^d)$ and for any $x \in \mathbb{Z}^d$:

$$\begin{aligned}\Pi_t \xi(x) &= \mathbb{E}_x \left[\exp \left(\beta_t^{-\alpha} \int_0^{\lfloor \beta_t^\alpha \rfloor} f \left(\frac{X_s}{\beta_t} \right) ds \right) \xi(X_{\lfloor \beta_t^\alpha \rfloor}) \right] \\ T_t \xi(x) &= \mathbb{E}_x \left[\exp \left(\beta_t^{-\alpha} \int_0^1 f \left(\frac{X_s}{\beta_t} \right) ds \right) \xi(X_1) \right]\end{aligned}$$

($X_t, t \geq 0$) being symmetric, T_t is self-adjoint and the Markov property implies that $T_t^{\lfloor \beta_t^\alpha \rfloor} = \Pi_t$. It follows that Π_t is also self-adjoint. Now, let us introduce a non negative function $g \in \mathcal{C}_c^\infty(\mathbb{R}^d)$, the set of infinitely differentiable function with compact support, such that $\|g\|_2 = 1$. We assume that the support of g is included in $[-M, M]^d$ and define ξ_t by $\xi_t(x) = \beta_t^{-d/2} g\left(\frac{x}{\beta_t}\right)$.

$$\begin{aligned}& \mathbb{E} \left[\exp \left(\beta_t^{-\alpha} \int_{\lfloor \beta_t^\alpha \rfloor}^{\lfloor \beta_t^\alpha \rfloor \gamma_t} f \left(\frac{X_s}{\beta_t} \right) ds \right) \right] \\ &= \sum_{x \in \mathbb{Z}^d} \mathbb{P}(X_{\lfloor \beta_t^\alpha \rfloor} = x) \mathbb{E}_x \left[\exp \left(\beta_t^{-\alpha} \int_0^{\lfloor \beta_t^\alpha \rfloor (\gamma_t - 1)} f \left(\frac{X_s}{\beta_t} \right) ds \right) \right] \\ &\geq \frac{1}{\sup_{x \in \mathbb{Z}^d} |\xi_t(x)|^2} \sum_{x \in \mathbb{Z}^d} \mathbb{P}(X_{\lfloor \beta_t^\alpha \rfloor} = x) \xi_t(x) \mathbb{E}_x \left[\exp \left(\beta_t^{-\alpha} \int_0^{\lfloor \beta_t^\alpha \rfloor (\gamma_t - 1)} f \left(\frac{X_s}{\beta_t} \right) ds \right) \xi_t(X_{\lfloor \beta_t^\alpha \rfloor (\gamma_t - 1)}) \right] \\ &= \frac{\beta_t^d}{\sup_{x \in \mathbb{Z}^d} |g(x)|^2} \sum_{x \in \mathbb{Z}^d} \mathbb{P}(X_{\lfloor \beta_t^\alpha \rfloor} = x) \xi_t(x) \Pi_t^{\gamma_t - 1} \xi_t(x)\end{aligned}$$

According to the local limit theorem (Remark page 661 of Le Gall and Rosen [28]),

$$\lim_{t \rightarrow +\infty} \sup_{x \in \mathbb{Z}^d} \left| \beta_t^d \mathbb{P}(X_{\lfloor \beta_t^\alpha \rfloor} = x) - p_1 \left(\frac{x}{\beta_t} \right) \right| = 0,$$

where p_1 is the transition density of the limit process. Since ξ_t is supported by $[-M\beta_t, M\beta_t]^d$, we only sum over this box. Moreover, there exists $\delta > 0$ such that $p_1(x) > \delta$ for all $x \in [-M, M]^d$. ξ_t being non negative, we get for any $\delta > 0$ and any t sufficiently large:

$$\mathbb{E} \left[\exp \left(\beta_t^{-\alpha} \int_{\lfloor \beta_t^\alpha \rfloor}^{\lfloor \beta_t^\alpha \rfloor \gamma_t} f \left(\frac{X_s}{\beta_t} \right) ds \right) \right] \geq \frac{\delta}{2 \sup_{x \in \mathbb{Z}^d} |g(x)|^2} \sum_{x \in \mathbb{Z}^d} \xi_t(x) \Pi_t^{\gamma_t - 1} \xi_t(x) = C \langle \xi_t, \Pi_t^{\gamma_t - 1} \xi_t \rangle.$$

Using the spectral representation of the operator Π_t , there exists a probability measure μ_{ξ_t} such that $\langle \xi_t, \Pi_t \xi_t \rangle = \int_0^{+\infty} \lambda d\mu_{\xi_t}$. It follows then from Jensen's inequality that

$$\langle \xi_t, \Pi_t^{\gamma_t - 1} \xi_t \rangle = \int_0^{+\infty} \lambda^{\gamma_t - 1} d\mu_{\xi_t} \geq \left(\int_0^{+\infty} \lambda d\mu_{\xi_t} \right)^{\gamma_t - 1} = \langle \xi_t, \Pi_t \xi_t \rangle^{\gamma_t - 1}.$$

We have thus proved:

$$\liminf_{t \rightarrow +\infty} \frac{\beta_t^\alpha}{t} \log \mathbb{E} \left[\exp \left(\beta_t^{-\alpha} \int_0^t f \left(\frac{X_s}{\beta_t} \right) ds \right) \right] \geq \liminf_{t \rightarrow +\infty} \log \langle \xi_t, \Pi_t \xi_t \rangle.$$

Now, remember that $(X_t, t \geq 0)$ is in the domain of a stable process $(U_t, t \geq 0)$, i.e: $\frac{1}{t^{1/\alpha}}(X_{ts}, s \in [0; 1]) \rightarrow (U_s, s \in [0; 1])$ in the Skorokhod's J1 topology.

$$\begin{aligned} \langle \xi_t, \Pi_t \xi_t \rangle &= \sum_{x \in \mathbb{Z}^d} \xi_t(x) \mathbb{E}_x \left[\exp \left(\beta_t^{-\alpha} \int_0^{\lfloor \beta_t^\alpha \rfloor} f \left(\frac{X_s}{\beta_t} \right) ds \right) \xi_t(X_{\lfloor \beta_t^\alpha \rfloor}) \right] \\ &= \beta_t^{-d} \sum_{x \in \mathbb{Z}^d} g \left(\frac{x}{\beta_t} \right) \mathbb{E}_0 \left[\exp \left(\int_0^1 f \left(\frac{X_{s \lfloor \beta_t^\alpha \rfloor} + x}{\beta_t} \right) ds \right) g \left(\frac{X_{\lfloor \beta_t^\alpha \rfloor} + x}{\beta_t} \right) \right] \\ &\xrightarrow{t \rightarrow +\infty} \int_{\mathbb{R}^d} g(x) \mathbb{E}_0 \left[\exp \left(\int_0^1 f(U_s + x) ds \right) g(U_1 + x) \right] dx \\ &= \int_{\mathbb{R}^d} g(x) \mathbb{E}_x \left[\exp \left(\int_0^1 f(U_s) ds \right) g(U_1) \right] dx. \end{aligned}$$

Let us justify the convergence over t above. Let F be the functional, defined for any function j by:

$$F(j) = \exp \left(\int_0^1 f \circ j(s) ds \right) g \circ j(1).$$

We have to prove that

$$\int_{\mathbb{R}^d} g(x) \mathbb{E} [F((U_s + x, s \in [0, 1]))] dx - \beta_t^{-d} \sum_{x \in \mathbb{Z}^d} g \left(\frac{x}{\beta_t} \right) \mathbb{E} \left[F \left(\frac{1}{\beta_t} (X_{s \lfloor \beta_t^\alpha \rfloor} + x, s \in [0, 1]) \right) \right] \xrightarrow{t \rightarrow +\infty} 0.$$

Introducing $\int_{\mathbb{R}^d} g(x) \mathbb{E} \left[F \left(\frac{1}{\beta_t} X_{s \lfloor \beta_t^\alpha \rfloor} + x, s \in [0, 1] \right) \right] dx$ in the previous sum, we can apply the dominate convergence to obtain:

$$\int_{\mathbb{R}^d} g(x) \mathbb{E} [F((U_s + x, s \in [0, 1]))] dx - \int_{\mathbb{R}^d} g(x) \mathbb{E} \left[F \left(\frac{1}{\beta_t} X_{s \lfloor \beta_t^\alpha \rfloor} + x, s \in [0, 1] \right) \right] dx \xrightarrow{t \rightarrow +\infty} 0.$$

Indeed, on one hand F is continuous in the Skorokhod's J1 topology because f and g are continuous, and on the other hand F is bounded since f and g are bounded. Moreover we can see that

$$\int_{\mathbb{R}^d} g(x) \mathbb{E} \left[F \left(\frac{X_{s \lfloor \beta_t^\alpha \rfloor} + x, s \in [0, 1] \right) \right] dx - \beta_t^{-d} \sum_{y \in \mathbb{Z}^d} g \left(\frac{y}{\beta_t} \right) \mathbb{E} \left[F \left(\frac{X_{s \lfloor \beta_t^\alpha \rfloor} + y, s \in [0, 1] \right) \right] \xrightarrow{t \rightarrow +\infty} 0.$$

Indeed, f and g being continuous with compact support, the function $x \in \mathbb{R}^d \mapsto F \left(\frac{X_{s \lfloor \beta_t^\alpha \rfloor} + x, s \in [0, 1] \right)$ is uniformly continuous, and its modulus of continuity does not depend of t .

At this point, we have proved that

$$\begin{aligned} \liminf_{t \rightarrow +\infty} \frac{\beta_t^\alpha}{t} \log \mathbb{E} \left[\exp \left(\beta_t^{-\alpha} \int_0^t f \left(\frac{X_s}{\beta_t} \right) ds \right) \right] &\geq \log \int_{\mathbb{R}^d} g(x) \mathbb{E}_x \left[\exp \left(\int_0^1 f(U_s) ds \right) g(U_1) \right] dx. \\ &= \log \langle g, S_1 g \rangle \end{aligned}$$

where S_t is the semigroup of operators defined on $L^2(\mathbb{R}^d)$ by:

$$S_t h(x) = E_x \left[\exp \left(\int_0^t f(U_s) ds \right) h(U_t) \right].$$

This semigroup is self-adjoint due to the symmetry of the process $(U_t, t \geq 0)$. Thus, its infinitesimal operator \mathcal{A} is also self-adjoint. Therefore, using the spectral representation of \mathcal{A} and Jensen's

inequality, we obtain:

$$\begin{aligned} \langle g, S_1 g \rangle &= \langle g, \exp(\mathcal{A})g \rangle = \int_{\mathbb{R}} \exp(\lambda) d\mu_g(\lambda) \\ &\geq \exp \int_{\mathbb{R}} \lambda d\mu_g(\lambda) = \exp \langle g, \mathcal{A}g \rangle = \exp \left(\int_{\mathbb{R}^d} f(x)g^2(x)dx - \int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}g(\omega)|^2 d\omega \right). \end{aligned}$$

Hence,

$$\liminf_{t \rightarrow +\infty} \frac{\beta_t^\alpha}{t} \log \mathbb{E} \left[\exp \left(\beta_t^{-\alpha} \int_0^t f \left(\frac{X_s}{\beta_t} \right) ds \right) \right] \geq \int_{\mathbb{R}^d} f(x)g(x)^2 dx - \int_{\mathbb{R}^d} |\omega|^\alpha |\mathcal{F}(g)(\omega)|^2 d\omega.$$

for any non negative function $g \in C_c^\infty(\mathbb{R}^d)$. We now take the supremum over non negative functions $g \in C_c^\infty(\mathbb{R}^d)$. Using representation (45) and boundedness of f , one can easily see that this is the same as taking the supremum over non negative functions in $L^2(\mathbb{R}^d)$, and this ends the proof of Theorem 11. \square

REFERENCES

- [1] Asselah, A. *Large deviation principle for self-intersection local times for random walk in \mathbb{Z}^d with $d \geq 5$* . ALEA Lat. Am. J. Probab. Math. Stat. 6 (2009), 281-322.
- [2] Asselah, A.; Castell, F. *Random walk in random scenery and self-intersection local times in dimensions $d \geq 5$* . Probab. Theory Related Fields 138 (2007), no. 1-2, 1-32.
- [3] Brydges, D.; van der Hofstad, R.; König, W. *Joint density for the local times of continuous-time Markov chains*. Ann. Probab. 35 (2007), no. 4, 1307-1332.
- [4] Bass, R.; Chen, Xia; Rosen, J. *Moderate deviations and laws of the iterated logarithm for the renormalized self-intersection local times of planar random walks*. Electron. J. Probab. 11 (2006), no. 37, 993-1030 (electronic).
- [5] Bass, R.; Chen, Xia; Rosen, J. *Large deviations for renormalized self-intersection local times of stable processes*. Ann. Probab. 33, (2005), no. 3, 984-1013.
- [6] Bass, R.; Levin, D. A. *Transition probabilities for symmetric jump processes*. Trans. Amer. Math. Soc. 354 (2002), no. 7, 2933-2953 (electronic).
- [7] Becker, M.; König, W. *Self-intersection local times of random walks: Exponential moments in subcritical dimensions* To appear in Probab. Theory Relat. Fields
- [8] Carmona, René A.; Molchanov, S. A. *Parabolic Anderson problem and intermittency*. Mem. Amer. Math. Soc. 108 (1994), no. 518, viii+125 pp
- [9] Castell, F. *Large deviations for intersection local times in critical dimension*. Ann. Probab. 38 (2010), no. 2, 927-953.
- [10] Castell, F.; Guillotin-Plantard, N.; Pène, F. *Limit theorems for one and two-dimensional random walk in random scenery*. Preprint arXiv:1103.4453 [math.PR]. To appear in Ann. Inst. Henri Poincaré-PR.
- [11] Černý, Jiří. *Moments and distribution of the local time of a two-dimensional random walk*. Stochastic Process. Appl. 117 (2007), no. 2, 262-270.
- [12] Chen, Xia. *Random walk intersections. Large deviations and related topics*. Volume 157 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2010.
- [13] Chen, Xia. *Exponential asymptotics and law of the iterated logarithm for intersection local times of random walks*. Ann. Probab. 32 (2004), no. 4, 3248-3300.
- [14] Chen, Xia; Li, Wenbo V. *Large and moderate deviations for intersection local times*. Probab. Theory Related Fields 128 (2004), no. 2, 213-254.
- [15] Chen, Xia; Li, Wenbo V.; Rosen, Jay. *Large deviations for local times of stable processes and stable random walks in 1 dimension*. Electron. J. Probab. 10 (2005), no. 16, 577-608.
- [16] Chen, Xia; Rosen, Jay. *Exponential asymptotics for intersection local times of stable processes and random walks*. Ann. Inst. H. Poincaré Probab. Statist. 41 (2005), no. 5, 901-928.
- [17] C. Domb and G.S. Joyce. *Cluster expansion for a polymer chain*. J. Phys, 5, 956-976, 1972.
- [18] Donsker, M. D.; Varadhan, S. R. S. *The polaron problem and large deviations*. New stochastic methods in physics. Phys. Rep. 77 (1981), no. 3, 235-237.
- [19] S. F. Edwards. *The statistical mechanics of polymers with excluded volume*. Proc. Phys. Soc. 85, 613-624, 1965.
- [20] de Gennes, P. G. *Scaling concepts in Polymer Physics*. Cornell University Press, Ithaca (1979).
- [21] Feller W. *An introduction to probability theory and its applications. Vol II*. Second Wiley series in probability and mathematical statistics, Second edition, John Wiley & Sons, New York (1971), xxiv+669 pages.

- [22] Grafakos, L. *Classical Fourier analysis*, Graduate Texts in Mathematics 249 , Second edition, Springer, New York (2008), xvi+489 pages.
- [23] Kesten, H.; Spitzer, F. *A limit theorem related to a new class of self-similar processes*. Z. Wahrsch. Verw. Gebiete 50 (1979), no. 1, 5-25.
- [24] Khanin, K. M.; Mazel, A. E.; Shlosman, S. B.; Sinai, Ya. G. *Loop condensation effects in the behavior of random walks*. The Dynkin Festschrift, 167-184, Progr. Probab., **34**, Birkhäuser Boston, Boston, MA, 1994.
- [25] König, W. *Upper tails of self-intersection local times of random walks*. Survey of proof techniques. Actes rencontr. CIRM **2:1**, 15-24 (2010).
- [26] Laurent, C. *Large deviations for self-intersection local times of stable random walks*. Stochastic Process. Appl. 120 (2010), no. 11, 2190-2211.
- [27] Laurent, C. *Large deviations for self-intersection local times in subcritical dimensions*. Electron. J. Probab. 17 (2012), no. 21, 1-20.
- [28] Le Gall, J-F., Rosen, Jay. *The range of stable random walks*, Ann. Probab. 19 (1991), no 2, 650-705.
- [29] Ledoux, M.; Talagrand, M. *Probability in Banach spaces*. Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol 23, Springer-Verlag, Berlin (1991), xii+480 pages.
- [30] Marcus, M. B.; Rosen, Jay. *Markov processes, Gaussian processes, and local times*. Cambridge Studies in Advanced Mathematics 100, Cambridge University Press, Cambridge (2006), x+620 pages.
- [31] Varadhan, S.R.S. *Appendix to Euclidean quantum field theory, by K. Symanzik*. Local Quantum Theory (R. Jost, ed). Academic New York (1969).
- [32] Zel'dovich, Ya. B.; Molchanov, S. A.; Ruzmaikin, A. A.; Sokoloff, D. D. *Intermittency, diffusion and generation in a nonstationary random medium*. Mathematical physics reviews, **Vol. 7**, 3-110, Soviet Sci. Rev. Sect. C Math. Phys. Rev., 7, Harwood Academic Publ., Chur, 198

LATP, UMR CNRS 7353. CENTRE DE MATHÉMATIQUES ET INFORMATIQUE. AIX-MARSEILLE UNIVERSITÉ.
39, RUE JOLIOT CURIE. 13 453 MARSEILLE CEDEX 13. FRANCE.
E-mail address: Fabienne.Castell@cmi.univ-mrs.fr

LATP, UMR CNRS 7353. CENTRE DE MATHÉMATIQUES ET INFORMATIQUE. AIX-MARSEILLE UNIVERSITÉ.
39, RUE JOLIOT CURIE. 13 453 MARSEILLE CEDEX 13. FRANCE.
E-mail address: Clement.Laurent@cmi.univ-mrs.fr

LATP, UMR CNRS 7353. CENTRE DE MATHÉMATIQUES ET INFORMATIQUE. AIX-MARSEILLE UNIVERSITÉ.
39, RUE JOLIOT CURIE. 13 453 MARSEILLE CEDEX 13. FRANCE.
E-mail address: Clothilde.Melot@cmi.univ-mrs.fr