

Fine structure of one-dimensional discrete point system

V. A. Malyshev

Abstract

We consider the system of N points on the segment of the real line with the nearest-neighbor Coulomb repulsive interaction and external force F . For the fixed points of such systems (fixed configurations) we study the asymptotics (in N and l) of finite differences of order l . Classical theory of finite differences is extensively used.

Assume that the system

$$0 \leq x_1 < x_2 < \dots < x_N < 1$$

of N different points on the segment $[0, 1] \in \mathcal{R}$ is given. If this system is a random system, there exist a lot of ways to characterize its structure, for example, as a random process of increments $\Delta_i = x_{i+1} - x_i$. If there is no any randomness, then there is no conventional way to characterize its organization. One of the possible ways is to consider the system of finite differences of several orders $l = 1, 2, \dots$

$$\nabla x_i = \Delta_i = x_{i+1} - x_i, \nabla^2 x_i = x_{i+2} - 2x_{i+1} + x_i, \dots, \nabla^l x_i, \dots$$

as the natural local characteristics of this system of points (or numbers), even if we would not know the metrics of the space where they are embedded and even would not know the space itself. This corresponds to the situation in the analysis where the existence of derivatives of sufficiently large order indicates the «quality» of the function. In the discrete case there is no existence problem (for example, on a circle the difference of arbitrary order are defined), but a possible substitute for the existence can be the decay rate of finite differences in l (for sufficiently large N). To go further, everything depends on the way how this systems of points is defined. Most popular way is the discretization of smooth functions. Already for a long time discrete differences have been used in numerical mathematics in connection with approximation, interpolation, solution of differential equations, etc. However, in these disciplines one does not need differences of high orders. For arbitrary orders there exists classical science - the theory of finite differences, see [1, 2, 3]), worked out already at Newton's time (the Newton series, divided differences, etc.), In section 2 we give necessary definitions and results from this science.

We are interested in the case when there is no natural smooth function such that the point system is its discretization. Such systems appear, for example, as the fixed points of natural dynamical systems in physics.

We suggest a new insight on the system of differences as the indicators of the deviation scale from the ideal system. We call the system **ideal system** if all distances between neighbors are equal. Then all differences of the order greater than 1 are zero. In physics this corresponds to the ideal crystal, and many papers were devoted to the proof that the system becomes ideal in the (thermodynamic) limit $N \rightarrow \infty$, see [7, 8, 9, 10, 11]. We are interested in the cases when the system is (in some sense) close to ideal. Here the thermodynamic limit is counter-productive - our problem is finer.

The differences of given order l depend on N and on i . We are interested in the asymptotics or, at least, in the bounds from above for given l , uniform in i . Let us say that the differences of order l are defined on the scale $\kappa(N, l)$ (not seen on the scales higher than $\kappa(N, l)$), if the number $\kappa(N, l)$ is the asymptotics for the differences of order l uniform in i (bound from above, for given l , uniform in i). We call **fine structure** of the point system the set of numbers $\kappa(N, l)$.

For example, in physics the scale of order 1 corresponds to macro-scale, and the scale N^{-1} corresponds to the micro-scale.

Formulation of the problem Often, the system of point is not given explicitly, but as a configuration, yielding minimum to some potential, or as a fixed point of some dynamics. We study here concrete system of points on $[0, 1]$, defined by the system of equations

$$f(x_i - x_{i-1}) - f(x_{i+1} - x_i) + F(x_i) = 0, i = 2, \dots, N - 1 \quad (1)$$

and for any end point x_1 and x_N there is an alternative - either $x_1 = 0$ ($x_N = 1$) and moreover correspondingly

$$-f(x_2 - x_1) + F(x_1) \leq 0 \quad (f(x_N - x_{N-1}) + F(x_N) \geq 0)$$

or, correspondingly,

$$-f(x_2 - x_1) + F(x_1) = 0 \quad (f(x_N - x_{N-1}) + F(x_N) = 0)$$

This system is interpreted as a fixed configuration, when x_i are subjected to the interaction forces $f(x_i - x_{i-1})$ and $-f(x_{i+1} - x_i)$ with the left and right neighbor correspondingly (we consider the Coulomb repulsive interaction $f(r) = r^{-2}$). Moreover, there is external force $F(x)$. It is clear that if $F(x) \equiv 0$, then $x_1=0, x_N = 1$, and all $x_{i+1} - x_i$ are equal - the ideal case.

In this paper, which is a natural continuation of the papers [5, 6], we consider three examples of the external force F - constant, linear and power functions. Only in the first case we get exact asymptotics, which is based on the techniques, developed in combinatorics for Stirling numbers of the second kind. In two other cases we get only bounds from above, which seem to be close to exact, as our estimates seem to be «on the edge» of exact estimates. Proof, in the linear case, are based again on the combinatorics and inductive procedure in l . In the power case we use a different inductive construction, however each step of this construction contains one more inductive procedure, similar to the one used in linear case.

It was proved in [5, 6], that for any monotone external force $F(x)$ the fixed configuration exists, is unique and such that $x_1 = 0, x_N = L$, and for $i = 2, \dots, N - 1$ the equations (1). Moreover, it was proved there that the first differences have asymptotics (uniform in i)

$$\Delta_i \sim \frac{L}{N-1}$$

The central fact - this asymptotics does not depend on F . In [5] the second term of the asymptotic expansion was obtained

$$(x_{i+1} - x_i) - \frac{L}{N-1} \sim \frac{F}{2} N^{-3} \left(\frac{N}{2} - i \right), i = 2, \dots, N - 1$$

for constant external force F . Formal calculation shows that the main term of the asymptotics for the second difference should be of the order N^{-3} , and does not depend on i , and moreover using the results of [5], one can show that the higher terms of the asymptotic expansion do not change this conclusion. One can say that only second difference shows up the macro force F . Here we study the asymptotics of all higher differences. Denote

$$\Delta_1 = x_2 - x_1 = \Delta, \Delta_i = x_{i+1} - x_i = \Delta(1 + \delta_i), i = 1, \dots, N - 1$$

Thus $\delta_1 = 0$ by definition, and in [5, 6] it was proved that $\Delta \sim \frac{1}{N}$. All proofs below are based on the study of the following system of equations for the unknowns $\delta_i, i = 2, \dots, N - 1$

$$f(\Delta(1 + \delta_i)) - f(\Delta) = \sum_{i=2}^i F(x_i), x_i = (i-1)\Delta + \Delta \sum_{i=2}^i \delta_i$$

which can be obtained by summing up the equations (1) from 2 to i . Then we have

$$\delta_i = (1 + Q_i)^{-\frac{1}{2}} - 1 = \sum_{m=1}^{\infty} \delta_{i,m}, \delta_{i,m} = a_m Q_i^m \quad (2)$$

where

$$Q_i = \Delta^2 \sum_{i=2}^i F(x_i)$$

$$(-1)^m a_m = \frac{1 \cdot 3 \dots (2m-1)}{2^m m!} = \frac{(2m)!}{(2^m m!)^2} \sim \frac{1}{\sqrt{\pi m}}, |a_m| \leq 1 \quad (3)$$

We see that for non-constant function F , this system is strongly non-linear and the equations are intertwined.

1 Finite differences

Let the function $g_i = g(i), i \in Z$, be given on the lattice Z . Call

$$\nabla g(i) = \nabla^+ g(i) = g(i+1) - g(i), (\nabla^- g)(i) = g(i) - g(i-1) \quad (4)$$

its right and left finite difference (discrete derivative). If the function is defined on the part of the lattice, for example only for $i = 1, \dots, N$, then we consider only such differences which are defined. For example, the difference $\nabla^l g(i)$ is defined iff $i+l \leq N$.

Note that

$$\nabla^n = (S-1)^n = \sum_{k=0}^n C_n^k (-1)^k S^{n-k} = \sum_{k=0}^n C_n^k (-1)^{n-k} S^k \quad (5)$$

where S is the shift operator

$$(Sf)(i) = f(i+1)$$

It follows in particular

$$\nabla x_i = \Delta_i = x_{i+1} - x_i, \nabla^2 x_i = x_{i+2} - 2x_{i+1} + x_i = \nabla \Delta_i = \Delta \nabla \delta_i = \Delta(\delta_{i+1} - \delta_i) \quad (6)$$

Thus the derivatives commute with S and the following (Leibniz) formulas hold

$$\nabla(gf)(i) = f(i+1)(\nabla g)(i) + g(i)(\nabla f)(i) = (Sf)(\nabla g) + g(\nabla f) = (Sg)(\nabla f) + f(\nabla g) \quad (7)$$

$$\nabla(f_1 \dots f_n) = (\nabla f_1)S(f_2 \dots f_n) + f_1 \nabla(f_2 \dots f_n) = \dots = \sum_{k=1}^n f_1 \dots f_{k-1} (\nabla f_k) S(f_{k+1} \dots f_n) \quad (8)$$

In the continuous case the differentiation of the product formula (with arbitrary l and n) is

$$(f_1 \dots f_n)^{(l)} = \sum_Q \frac{l!}{\prod_{k=1}^n l_k!} f_1^{(l_1)} \dots f_n^{(l_n)} \quad (9)$$

where the sum is over all arrays $Q = (l_1, \dots, l_n)$ of non-negative integers, satisfying the condition $l_1 + \dots + l_n = l$, and index (l) indicates the derivative of the order l . In the discrete case we will need the analog of the formula (9)

$$\begin{aligned} \nabla^l(f_1 f_2) &= \sum_{k=0}^l C_l^k (\nabla^k f_1) (\nabla^{l-k} S^k f_2) \\ \nabla^l(f_1 \dots f_n) &= \sum_Q \frac{l!}{\prod_{k=1}^n l_k!} (\nabla^{l_1} f_1) (\nabla^{l_2} S^{l_1} f_1) (\nabla^{l_3} S^{l_1+l_2} f_1) \dots (\nabla^{l_n} S^{l_1+\dots+l_{n-1}} f_n) \end{aligned} \quad (10)$$

Both formulas are easily proved by induction, the first one is the formula (48) in [16]. However, shift operators and their powers will not play role for us. If one denotes

$$\gamma_k(q) = \max_{i=1, \dots, l-q} |S^i \nabla^q f_k|$$

then in two last sections we will need the inequality

$$\nabla^l(f_1 \dots f_n) \leq \sum_Q \frac{l!}{\prod_{k=1}^n l_k!} \gamma_1(l_1) \dots \gamma_n(l_n) \quad (11)$$

Note also that if the function $f(i)$ does not depend on i , then its differentiation gives zero. It is also easy to show that

$$\nabla^l i^n = 0, l > n \quad (12)$$

Higher differences of the power function The numbers

$$S(n, l, i) = \frac{1}{l!} \nabla^l i^n$$

are sometimes called generalized Stirling numbers of the second kind [14]. A particular case is the ordinary Stirling numbers of the second kind

$$\left\{ \begin{matrix} n \\ l \end{matrix} \right\} = S(n, l) = S(n, l, 0) = \frac{1}{l!} \nabla^l 0^n = \frac{1}{l!} \sum_{k=0}^l (-1)^{l-k} C_l^k k^n$$

where the value $\nabla^l i^n$ for $i = 0$ is denoted by $\nabla^l 0^n$. It is well-known that

$$\left\{ \begin{matrix} n \\ l \end{matrix} \right\} = 0, n < l, \left\{ \begin{matrix} n \\ l \end{matrix} \right\} = 1, n = l, \quad (13)$$

Only some asymptotics for $S(n, l, i)$ are known:

1. Riordan asymptotics for $n \rightarrow \infty, l = \text{const}$

$$S(n, l) \sim \frac{l^n}{l!}$$

2. for $l \rightarrow \infty, n = l + k$ with bounded k , for any i one has [14]

$$S(n, l, i) \sim C_{l+k}^k \left(i + \frac{l}{2}\right)^k \quad (14)$$

the particular case is the result of [12] for $S(n, l)$;

3. if $n \rightarrow \infty, l \rightarrow \infty$ so that $\kappa = \frac{k}{l}$ is bounded away from zero and infinity. This is the result of Good [13] for $i = 0$, which was generalized in [14] for i 's with not too fast growth.

Discretization of smooth functions The following example demonstrates what one can expect in case of analytic function discretization. If some function $g(x)$ is defined on the circle S_1 , then it is equivalent to the function on all \mathbb{R} , periodic with period 1. Example is $g(x) = \sin 2\pi x$. Put

$$g_i = \sin \frac{2\pi i}{N}, i = 1, \dots, N \quad (15)$$

$$g_{i,n} = \nabla^n g_i, \Delta = \frac{1}{N}$$

Then

$$\begin{aligned} g_{i,1} &= \nabla g_i = g_{i+1} - g_i = \int_{x_i}^{x_i+\Delta} g^{(1)}(y_1) dy_1, \\ g_{i,2} &= \nabla g_{i,1} = g_{i+1,1} - g_{i,1} = \int_{x_{i+1}}^{x_{i+1}+\Delta} g^{(1)}(y) dy - \int_{x_i}^{x_i+\Delta} g^{(1)}(y) dy = \\ &= \int_{x_i}^{x_i+\Delta} (g^{(1)}(y_1 + \Delta) - g^{(1)}(y_1)) dy_1 = \int_{x_i}^{x_i+\Delta} dy_1 \int_{y_1}^{y_1+\Delta} g^{(2)}(y_2) dy_2, \end{aligned}$$

and so on

$$g_{i,n+1} = \nabla^{n+1} g_i = g_{i+1,n} - g_{i,n} = \int_{x_i}^{x_i+\Delta} dy_1 \left(\int_{y_1}^{y_1+\Delta} dy_2 \dots \left(\int_{y_n}^{y_n+\Delta} dy_{n+1} g^{(n+1)}(y_{n+1}) \right) \right)$$

That is why, if $|g^{(n)}(x)| \leq C^{n+1}$ for some constant $C = C(g) > 0$, then

$$|g_{i,n}| \leq C^{n+1} \Delta^n$$

2 Constant external force

Theorem 1 *Let $F(x) = F$ be constant, then as $N \rightarrow \infty$ and $1 \leq l = o(N)$ uniformly in $i < N - l$*

$$\nabla^{l+1} x_i \sim (-1)^l \sqrt{2} \left(\frac{F}{e} l \Delta^2\right)^l \Delta$$

Proof. Due to (6) it is sufficient to prove that

$$\nabla^l \delta_i \sim \sqrt{2} \left(\frac{F}{e} l \Delta^2\right)^l$$

We have from (2)

$$\nabla^l \delta_{i+1} = \sum_{m=1}^{\infty} \nabla^l \delta_{i+1,m}, \delta_{i+1,m} = a_m i^m \Delta^{2m} F^m \quad (16)$$

By (12) or (13) for $l > m$

$$\nabla^l \delta_{i,m} = 0$$

Lemma 2 *For $l = m$*

$$\nabla^l \delta_{i+1,m} = a_l l! (F \Delta^2)^l$$

Note that by (5)

$$\nabla^l i^{l+p} = \sum_{k=0}^l C_l^k (-1)^{l-k} (i+k)^{l+p} \quad (17)$$

Then lemma follows from the chain of equalities

$$\nabla^m i^m = \sum_{k=0}^m C_m^k (-1)^{m-k} (i+k)^m = \sum_{k=0}^m C_m^k (-1)^{m-k} k^m = m! \left\{ \begin{matrix} m \\ m \end{matrix} \right\} = m!$$

where the second equality has two explanations: either using (13) or because any differentiation lowers by 1 the degree of polynomial of i , that is why the second expression does not depend on i .

To prove the theorem we will show that the sum of the series (16) for $m > l$ is asymptotically less than the main term with $l = m$.

Further on $m = l + p, p > 0$. We consider two cases. Firstly let $p > l$, then from (17) it follows that

$$|(\nabla^+)^l \delta_{i+1,l+p}| \leq |a_{l+p}| (F \Delta^2)^{l+p} 2^l N^{l+p} = |a_{l+p}| 2^l F^{2l} \Delta^{2l} (F \Delta)^{p-l}$$

Take minimal l_0 , such that for any $l \geq l_0$

$$l! > 2^l F^l$$

Then for $l \geq l_0$

$$|a_{l+p}| 2^l F^{2l} \Delta^{2l} (F \Delta)^{p-l} \leq |a_l| (F \Delta^2)^l (F \Delta)^{p-l} l! \leq (F \Delta)^{p-l} |\nabla^l \delta_{i+1,l}|$$

that means that the sum over $p > l$ of the terms with $l \geq l_0$ is (asymptotically) majorized by the "main" term. If $l < l_0$ then the corresponding term does not exceed

$$C(l_0, F) \Delta^{p-l} |\nabla^l \delta_{i+1,l}|$$

for some constant $C(l_0, F) > 0$.

The case $1 \leq p \leq l$ is more complicated. One can easily show the known, see for example [14], equality

$$\nabla^l i^{l+p} = \sum_{q=0}^p C_{l+p}^q i^q (\nabla^l 0^{l+p-q})$$

For $l \rightarrow \infty$ we will need the fact that finite differences $\nabla^l i^n$ (and Stirling numbers) have simple combinatorial interpretation [14]. Remind that $\nabla^l 0^n$ is equal to the number of ways to place n different objects in l different cells so that no cell is empty. Similarly let we have $l + i$ different cells, l of them being marked out. Then $\nabla^l i^n$ is the number of ways to place n different objects in these cells so that no of the marked out l cells is empty. It follows that $\nabla^l i^n$ are non-negative (if $i \geq 0$) and increase with n .

Lemma 3

$$\nabla^l i^{n+1} \leq (l + i + nl) \nabla^l i^n$$

Proof. We have

$$\nabla^l i^n \leq \nabla^l i^{n+1} = (l + i) \nabla^l i^n + l \nabla^{l-1} i^n$$

In fact, the inequality is evident, and the equality can be explained as follows. Fix one element (for example the last one in some fixed enumeration). The configuration of the other elements can be of two types: 1) such that all l marked out cells were nonempty. Then the fixed element can be placed in $l + i$ ways, 2) exactly one of l marked out cells is empty, then the fixed element should be placed in this free cell.

Moreover it is clear that

$$\nabla^{l-1} i^n \leq n \nabla^l i^n$$

Lemma is proved.

As $l + i \leq N$ and $nl \leq 2l^2 = o(\Delta^2)$, we get from the lemma

$$\left| \frac{\nabla^l \delta_{i+1, n+1}}{\nabla^l \delta_{i+1, n}} \right| \leq \left| \frac{a_{l+1}}{a_l} \right| (l + i + nl) F \Delta^2 = o(\Delta)$$

and that the sum of the terms with $1 \leq p < l$ is asymptotically less than $\nabla^l \delta_{i+1, n}$. The theorem is proved.

Remark. The asymptotics for the case when l is of order N , is related to the unsolved combinatorial problem of finding the maximum of unimodal sequence of Stirling numbers, see [15], proposition 3.30. But, for example, the estimates for $l \sim \epsilon N$

$$(1 - C\epsilon) < \frac{|\nabla^l \delta_i|}{\sqrt{2} \left(\frac{F}{\epsilon} l \Delta^2\right)^l} < (1 + C\epsilon)$$

for some constant $C > 0$ and sufficiently small $\epsilon > 0$ follow from the proof above.

3 Linear external force

Theorem 4 Let $F(x) = \alpha x, \alpha \geq 1$. Then uniformly in $2 \leq l \leq N - i$

$$|\nabla^l x_i| \leq \Delta (C\Delta)^l l!$$

Proof. By (6) it is sufficient to prove that

$$|\nabla^{l-1} \delta_i| \leq (C\Delta)^l l!$$

Similarly to the previous section

$$\nabla^{l-1} \delta_i = \sum_{m=1}^{\infty} \nabla^{l-1} \delta_{i, m}, \delta_{i, m} = a_m \Delta^{2m} \alpha^m \left(\sum_{k=2}^i x_k \right)^m \quad (18)$$

The estimation method of each summand will depend on the pair m, l .

Small m, l For $l = 2, m = 1$

$$\nabla \delta_{i, 1} = a_1 \Delta^2 \alpha \nabla \sum_{k=2}^i x_k = a_1 \Delta^2 \alpha x_{i+1} \quad (19)$$

For $l = 3, m = 1$

$$\nabla^2 \delta_{i, 1} = a_2 \Delta^2 \alpha \nabla^2 \sum_{k=2}^i x_k = a_2 \Delta^2 \alpha \nabla x_{i+1} \sim a_2 \alpha \Delta^3 \quad (20)$$

For $l = 3, m = 2$

$$\nabla^2 \delta_{i, 2} = a_2 \Delta^4 \alpha^2 \nabla^2 \left(\sum_{k=2}^i x_k \right)^2 = O(\Delta^4) \quad (21)$$

Case $2 \leq l \leq m$ We “honestly” differentiate once, and for the rest $l - 2$ differentiations we use the following evident bound (which holds for any $f(i)$)

$$|\nabla^{l-2} f(i)| \leq 2^{l-2} \max |f(i)| \quad (22)$$

Namely, denoting $\psi_i = \sum_{k=2}^i x_k$, we have

$$\begin{aligned} |(\nabla^+)^{l-1} \delta_{i,m}| &\leq \Delta^{2m} \alpha^m |(\nabla^+)^{l-2} [x_{i+1}(\psi_{i+1}^{m-1} + \psi_{i+1}^{m-2} \psi_i + \dots + \psi_i^{m-1})]| \leq \\ &\leq \Delta^{2m} \alpha^m 2^{l-2} m(i+1)^{m-1} \leq \alpha^m \Delta^{m+1} 2^{l-2} m, \end{aligned}$$

and

$$\sum_{m \geq l \geq 2} |(\nabla^+)^{l-1} \delta_{i,m}| \leq 2^{l-2} \sum_{m \geq l \geq 2} m \alpha^m \Delta^{m+1} \leq l(2\alpha)^l \Delta^{l+1}$$

We proved even more, namely that the formulas (19) and (20) give asymptotics for $l = 2$ and $l = 3$ correspondingly.

Case $m < l \leq N$ Here the bounds are essentially more complicated. Denoting

$$\gamma_k = \max_{i:i+k \leq N} |\nabla^k x_i|$$

we use (for $l \geq 3$) the following inductive hypothesis

$$\gamma_k \leq \Delta (C\Delta)^k k!, \quad k = 2, \dots, l-1 \quad (23)$$

Consider the first summand (that is with $m = 1$) in (18), which is equal to

$$a_1 \alpha \Delta^2 \nabla^{l-2} x_{i+1}$$

and where the result follows directly from the inductive hypothesis.

For $m \geq 2$, using the formulas (10), the modulus of the expression

$$\nabla^{l-1} \left(\sum_{k=2}^i x_k \right)^m$$

after $l - 1$ differentiations can be estimated as

$$|\nabla^{l-1} \left(\sum_{k=2}^i x_k \right)^m| \leq \sum_Q C(m, l-1 | q_0, \dots, q_{l-1}) (\max_i \sum_{k=2}^i x_k)^{q_0} (\max_i x_{i+1})^{q_1} (\max_i |\nabla x_{i+1}|)^{q_2} \prod_{k \geq 3} \gamma_{k-1}^{q_k} \quad (24)$$

where \sum_Q is the sum over finite ordered arrays $Q = q_0, q_1, \dots, q_l$ of non-negative integers such that

$$q_0 + \sum_{k \geq 1} q_k = m, \quad \sum_{k \geq 1} k q_k = l - 1$$

Such arrays will be called admissible. Their meaning is that exactly q_0 factors $\sum_{k=2}^i x_k$ are not differentiated at all, q_1 factors are differentiated exactly once after what they become equal to x_{i+1} , etc., q_{l-1} factors are differentiated exactly $l - 1$ times. Enumerate m factors in $(\sum_{k=2}^i x_k)^m$ from 1 to m . Any of the subsequent differentiations is applied to one of these factors, giving different summands in the formula (12). Moreover, as we take maximum in i , one may not take into account the shift operators in the formula (10).

For given Q consider finite enumerated arrays $\alpha = \{A_k, k = 0, \dots, l-1\}$ of subsets of the set $\{1, \dots, m\}$, such that $|A_k| = q_k$ and

$$\{1, \dots, m\} = \cup A_k$$

The set A_k contain those and only those elements which have been differentiated exactly k times.

Moreover,, the sequence of $l - 1$ differentiations can be subdivided onto groups $B_{k,p}, p = 1, \dots, q_k$, so that

$$\{1, \dots, l-1\} = \cup_{k,p} B_{k,p}, \quad |B_{k,p}| = k$$

The meaning of the set $B_{k,p}$ is that each differentiation from $B_{k,p}$ is applied to the p -th element of the set A_k .

The constants $C(m, l-1 | q_0, \dots, q_{l-1})$ for admissible arrays Q are equal to the number of such partitions, and thus are equal to

$$\begin{aligned} & (C_m^{q_0} C_{m-q_0}^{q_1} C_{m-q_0-q_1}^{q_2} \dots) (C_{l-1}^{q_1} (q_1!)) (C_{l-1-q_1}^{2q_2} \frac{(2q_2)!}{2^{q_2}}) \dots (C_{l-1-q_1-2q_2 \dots}^{kq_k} \frac{(kq_k)!}{(k!)^{q_k}}) \dots = \\ & = \frac{m!}{\prod_{k=0}^{l-1} q_k!} \frac{(l-1)!}{\prod_{k=1}^{l-1} (k!)^{q_k}} \end{aligned}$$

Let us estimate from above the moduli of the summands in (18), using (24). In this we use inductive hypothesis (23) to estimate γ_k , and evident bounds for $|\sum x_i|, |x_i|, |\nabla x_{i+1}|$. As a result we get

$$|\nabla^{l-1} \delta_{i,m}| \leq \alpha^m \Delta^{2m} N^{q_0} \Delta^{\sum_{k \geq 3} q_k} \prod_{k \geq 2} ((C\Delta)^{k-1} (k-1)!)^{q_k} \frac{m!}{\prod_{k=0}^{l-1} q_k!} \frac{(l-1)!}{\prod_{k=1}^{l-1} (k!)^{q_k}}$$

Finally, the power of Δ will be

$$\begin{aligned} 2m - q_0 + m - q_0 - q_1 - q_2 + \sum_{k \geq 2} (k-1)q_k &= 2m - q_0 + m - q_0 - q_1 - q_2 + l - 1 - q_1 - m + q_0 + q_1 = \\ &= 2m + l - 1 - q_0 - q_1 - q_2 \geq m + l - 1 \end{aligned}$$

and the constant C will have power

$$l - 1 - q_1 - m + q_0 + q_1 = l - 1 - m + q_0$$

The final estimate is

$$\begin{aligned} & \alpha^m \Delta^{m+l-1} C^{l-1-m+q_0} \frac{m!}{\prod_{k=0}^{l-1} q_k!} (l-1)! \prod_{k \geq 2} \frac{((k-1)!)^{q_k}}{(k!)^{q_k}} \leq \\ & \leq \alpha^m (C\Delta)^{l-1} (l-1)! \Delta^2 m^2 2^{-\frac{m}{2}} C^{-m+q_0} \frac{1}{\prod_{k=0}^{l-1} q_k!} \prod_{k \geq 2} \frac{1}{k^{q_k}} \end{aligned}$$

In fact, as for any $k \geq 2$

$$k! \leq k^k 2^{-\frac{k}{2}}$$

then

$$\Delta^m m! \leq \Delta^m m^m 2^{-\frac{m}{2}} \leq \Delta^2 m^2 2^{-\frac{m}{2}}$$

Now we have only to do summation over all Q

$$\sum_Q C^{q_0} \frac{1}{\prod_{k=0}^{l-1} q_k!} \prod_{k \geq 2} \frac{1}{k^{q_k}} \leq \left(\sum_{q_0} \frac{C^{q_0}}{q_0!} \right) \prod_{k=1}^{l-1} \sum_{q_k=0}^m \frac{k^{-q_k}}{q_k!} \leq e^C e^{\sum_i^{l-1} k^{-1}} \leq e^{C+\ln(l-1)} = e^C (l-1)$$

after what we have the estimate of any summand (18) for given m

$$(C\Delta)^l l! \Delta^2 m^2 C^{-m} e^C 2^{-\frac{m}{2}} \alpha^m$$

Summation in m gives

$$(C\Delta)^l l! \Delta^2 \sum_{m=1}^{l-1} m^2 C^{-m} e^C 2^{-\frac{m}{2}} \alpha^m$$

Note that for $C > \frac{\alpha}{\sqrt{2}}$

$$\sum_{m=1}^{l-1} m^2 C^{-m} 2^{-\frac{m}{2}} e^C \alpha^m$$

is uniformly bounded in l (and in N). Taking into account (18), we see that the theorem has been proved with the constant

$$C = 2\alpha$$

Remark. The case $\alpha < 1$ can be considered similarly - only the constant C may change. We saw also that for $l = 2$ the asymptotics depends on i , but for $l = 3$ it does not.

4 Power external force

Theorem 5 Let $F(x) = \alpha x^n$, $\alpha \geq 1$. Then for any i and all $2 \leq l \leq N - i$

$$|\nabla^l x_i| \leq \Delta(\Delta C)^l l!, C = 2\alpha n! e^6$$

Proof. Similarly to above one has to estimate

$$\nabla^{l-1} \delta_i = \sum_{m=1}^{\infty} a_m \Delta^{2m} \alpha^m \nabla^{l-1} \left(\sum_{k=2}^i x_k^n \right)^m \quad (25)$$

Case $l = 2, m = 1$

$$\nabla \delta_{i,1} = a_1 \Delta^2 \alpha \nabla \sum_{k=2}^i x_k^n = a_1 \Delta^2 \alpha x_{i+1}^n \quad (26)$$

Case $2 \leq l \leq m$ We “honestly” differentiate once (using formula (8)), and for the rest $l - 2$ differentiations we use the estimate (22). Namely, denoting $\psi_i = \sum_{k=2}^i x_k^n$, we have

$$\begin{aligned} |\nabla^{l-1} \delta_{i,m}| &\leq \Delta^{2m} \alpha^m |\nabla^{l-2} [\psi_{i+1}^{m-1} + \psi_{i+1}^{m-2} \psi_i + \dots + \psi_i^{m-1}]| \leq \\ &\leq \Delta^{2m} \alpha^m 2^{l-2} m(i+1)^{m-1} \leq \alpha^m \Delta^{m+1} 2^{l-2} m, \end{aligned}$$

and

$$2^{l-2} \sum_{m \geq l \geq 2} m \alpha^m \Delta^{m+1} \leq l(2\alpha)^l \Delta^{l+1}$$

From this and from (26) the result of the theorem follows for $l = 2$.

First inductive procedure One has to estimate

$$\sum_{m=1}^{l-1} \nabla^{l-1} \delta_{i,m} = \sum_{m=1}^{l-1} a_m \alpha^m \Delta^{2m} \nabla^{l-1} \left(\sum_{k=2}^i x_k^n \right)^m \quad (27)$$

For this we use the inductive construction very similar to the one used for linear external force. Together with this (for $l \geq 3$) we use the inductive assumption

$$\gamma_k = \max_{i:i+k \leq N} |\nabla^k x_i| \leq \Delta(C\Delta)^k k!, k = 2, \dots, l-1 \quad (28)$$

to get the bound for $|\nabla^{l-1} x_i^n|$. Similarly to the previous section we have

$$|\nabla^{l-1} x_{i+1}^n| \leq \sum_Q C(n, l-1 | q_0, \dots, q_l) (\max_i x_{i+1})^{q_0} (\max_i |\nabla x_{i+1}|)^{q_1} \prod_{k \geq 2}^{l-1} \gamma_k^{q_k} \quad (29)$$

where

$$\sum_{k \geq 0} q_k = n, \sum_{k \geq 1} k q_k = l-1$$

The constants $C(n, l-1 | q_0, \dots, q_l)$ for the admissible arrays Q are the same as in the previous section

$$\frac{n!}{\prod_{k=0}^{l-1} q_k!} \frac{(l-1)!}{\prod_{k=1}^{l-1} (k!)^{q_k}}$$

Using the inductive assumption, the expression (29) can be estimated from above as

$$\begin{aligned} &\Delta \sum_{k \geq 1}^{q_k} \prod_{k \geq 2}^{l-1} ((C\Delta)^k k!)^{q_k} \frac{n!}{\prod_{k=0}^{l-1} q_k!} \frac{(l-1)!}{\prod_{k=1}^{l-1} (k!)^{q_k}} = \\ &= \Delta^{l-1 + \sum_{k \geq 2} q_k} C^{l-1-q_1} \frac{n!}{\prod_{k=0}^{l-1} q_k!} (l-1)! \leq (C\Delta)^{l-1} (l-1)! n! \Delta^{\sum_{k \geq 2} q_k} C^{-q_1} \frac{1}{\prod_{k=0}^{l-1} q_k!} \end{aligned}$$

We have to do summation over all Q

$$\sum_Q \Delta^{\sum_{k \geq 2} q_k} C^{-q_1} \frac{1}{\prod_{k=0}^{l-1} q_k!} \leq \sum_{q_0} \frac{1}{q_0!} \sum_{q_1} \frac{C^{-q_1}}{q_1!} \prod_{k=2}^{l-1} \sum_{q_k=0}^{l-1} \frac{\Delta^{q_k}}{q_k!} \leq e^{1+c^{-1}+(l-3)\Delta}$$

Finally we get

$$|\nabla^{l-1} x_{i+1}^n| \leq (C\Delta)^{l-1} (l-1)! n! e^{2+C^{-1}} \quad (30)$$

Case 1 = $m < l \leq N$ Similarly, we estimate the summand with $m = 1$, using

$$\begin{aligned} |\nabla^{l-1} \delta_{i,1}| &\leq a_1 \alpha \Delta^2 |\nabla^{l-1} (\sum_{k=2}^i x_k^n)| = a_1 \alpha \Delta^2 |\nabla^{l-2} x_{i+1}^n| \leq \\ &\leq a_1 \alpha \Delta^2 \sum_Q C(n, l-2 | q_0, \dots, q_{l-2}) (\max_i x_{i+1}^n)^{q_0} (\max_i |\nabla x_{i+1}^n|)^{q_1} \prod_{k \geq 2}^{l-2} (\gamma_k)^{q_k} \end{aligned} \quad (31)$$

where

$$\sum_{k \geq 0} q_k = n, \quad \sum_{k \geq 1} k q_k = l-2$$

The constants $C(n, l-2 | q_0, \dots, q_{l-2})$ for the admissible Q 's are the same as before

$$\frac{n!}{\prod_{k=0}^{l-2} q_k!} \frac{(l-2)!}{\prod_{k=1}^{l-2} (k!)^{q_k}}$$

Using the inductive assumption (30), the modulus of the expression (31) in the sum can be estimated as follows

$$\begin{aligned} &\alpha \Delta^2 \Delta^{\sum_{k \geq 1} q_k} \prod_{k \geq 2}^{l-2} ((C\Delta)^k k!)^{q_k} \frac{n!}{\prod_{k=0}^{l-2} q_k!} \frac{(l-2)!}{\prod_{k=1}^{l-2} (k!)^{q_k}} = \\ &= \alpha \Delta^{2+l-2+\sum_{k \geq 2} q_k} C^{l-2-q_1} \frac{n!}{\prod_{k=0}^{l-2} q_k!} (l-2)! \leq \alpha (C\Delta)^l l! n! \frac{1}{l(l-1)} \Delta^{\sum_{k \geq 2} q_k} C^{-2-q_1} \frac{1}{\prod_{k=0}^{l-2} q_k!} \end{aligned}$$

Again, we have to do summation over all Q

$$\sum_Q \Delta^{\sum_{k \geq 2} q_k} C^{-q_1} \frac{1}{\prod_{k=0}^{l-2} q_k!} \leq \sum_{q_0} \frac{1}{q_0!} \sum_{q_1} \frac{\Delta^{q_1} C^{-q_1}}{q_1!} \prod_{k=2}^{l-2} \sum_{q_k=0}^{l-2} \frac{\Delta^{q_k}}{q_k!} \leq e^{1+C^{-1}+(l-4)\Delta}$$

As the result we get

$$\alpha (C\Delta)^l l! n! \frac{1}{l(l-1)} \frac{e^{2+C^{-1}}}{C^2}$$

Second inductive procedure Now, using the estimate (30), we consider the case $2 \leq m < l \leq N$, that is estimate for $m \geq 2$ the expression

$$a_m \alpha^m \Delta^{2m} \nabla^{l-1} (\sum_{k=2}^i x_k^n)^m$$

Denote now

$$\beta_k = \max_i |\nabla^k x_{i+1}^n|$$

It can be written, after $l-1$ differentiations, as

$$a_m \alpha^m \Delta^{2m} \sum_Q C(m, l-1 | q_0, \dots, q_{l-1}) (\max_i \sum_{k=2}^i x_k^n)^{q_0} (\max_i x_{i+1}^n)^{q_1} \prod_{k \geq 2}^{l-1} \beta_k^{q_k} \quad (32)$$

Finally we have

$$|\nabla^{l-1} \delta_{i,m}| = |a_m \Delta^{2m} \alpha^m \nabla^{l-1} (\sum_{k=2}^i x_k^n)^m| \leq$$

$$\leq |\alpha^m \Delta^{2m} N^{q_0} \Delta^{\sum_{k \geq 3} q_k} \prod_{k \geq 2} ((C\Delta)^{(k-1)} (k-1)! n! e^{2+C^{-1}})^{q_k} \frac{m!}{\prod_{k=0}^{l-1} q_k!} \frac{(l-1)!}{\prod_{k=1}^{l-1} (k!)^{q_k}}|$$

Now it will be more convenient to consider the factors separately.

The power of Δ is

$$2m - q_0 + \sum_{k \geq 3} q_k + l - q_1 - m + q_0 + q_1 = m + l + \sum_{k \geq 3} q_k$$

the power of C is

$$l - q_1 - m + q_0 + q_1 = l - m + q_0$$

The remaining constants are

$$(n! e^{2+C^{-1}})^{m-q_0-q_1}$$

Besides the factor $(C\Delta)^l l!$ we have the factor

$$m! l^{-1} \Delta^m \leq l^{-1}$$

The summation gives

$$\sum_Q \left[\sum_{q_0=0}^{l-1} \frac{1}{q_0!} \right] \left[\sum_{q_1=0}^{l-1} \frac{1}{q_1!} \right] \prod_{k=2}^{l-1} \left[\sum_{q_k=0}^{l-1} \frac{1}{q_k!} \frac{1}{k^{q_k}} \Delta^{q_k} \right] \leq e^3$$

Finally we get the constants

$$C^{-m+q_0} (\alpha n! e^{2+C^{-1}})^{m-q_0-q_1} e^3$$

which proves the theorem with

$$C = 2\alpha n! e^6$$

References

- [1] Ch. Jordan. Calculus of finite differences. 2nd edition, 1950. NY.
- [2] L. Milne-Thomson. The calculus of finite differences. 1933. London.
- [3] A. O. Gelfond. Calculus of finite differences. 1959. Moscow.
- [4] Ph. Flajolet, R. Sedgewick. Mellin transforms and asymptotics: finite differences and Rice's integrals. Theor. Comp. Science, 1995, v. 144, 101-124.
- [5] V. A. Malyshev. Fixed points for one-dimensional particle system with strong interaction. Moscow Math. Journal, 2012, v. 12, No. 1.
- [6] V. A. Malyshev. Critical states of multi-particle systems with strong interaction on a circle. Problems of information transmission, 2011, v. 47, No. 2, 117-127.
- [7] W. Ventevogel. On the configuration of a one-dimensional system of interacting particles with minimum potential energy per particle. Physica A, 1978, 92, No. 3-4, pp. 343-361.-
- [8] M. Duneau, A. Katz. Structural stability of classical lattices in one-dimension. Annales de l'I.H.P., section A, 1984, 41, No. 3, 269-290.
- [9] Ch. Radin. Existence of ground state configurations. Math. Physics Electronic J., 2004, v. 10.
- [10] Ch. Radin. Crystals and Quasicrystals: a lattice gas model. Physics Letters, 1986, 114A, No. 7, 381-383.
- [11] Ch. Radin, L. Schulman. Periodicity of classical ground states. Phys. Rev. Letters, 1983, 51, No. 8, 621-622.
- [12] L. C. Hsu. Note on the asymptotic expansion of the n-th difference of zero. Annals of Math. Stat., 1948, v. 19, 273-277.
- [13] I. Good. An asymptotic formula for the differences of the power at zero. Annals Math. Stat., 1961, 32, 1, 249-256.
- [14] Yu. Medvedev, G. Ivchenko. Asymptotic representations of finite differences of a power function at an arbitrary point. Theory of Probability and applications, 1965, v. 10, No. 1, 151-156.
- [15] M. Aigner. Combinatorial theory. 1979, Springer, NY. (Russian translation Moscow, 1982).
- [16] M. Spiegel. Calculus of finite differences and difference equations. 1971. McGraw-Hill.