STABILITY OF THE SLOW MANIFOLD IN THE PRIMITIVE EQUATIONS

R. TEMAM AND D. WIROSOETISNO

ABSTRACT. We show that, under reasonably mild hypotheses, the solution of the forced-dissipative rotating primitive equations of the ocean loses most of its fast, inertia-gravity, component in the small Rossby number limit as $t \rightarrow \infty$. At leading order, the solution approaches what is known as "geostrophic balance" even under ageostrophic, slowly time-dependent forcing. Higher-order results can be obtained if one further assumes that the forcing is time-independent and sufficiently smooth. If the forcing lies in some Gevrey space, the solution will be exponentially close to a finite-dimensional "slow manifold" after some time.

1. INTRODUCTION

One of the most basic models in geophysical fluid dynamics is the primitive equations, understood here to be the hydrostatic approximation to the rotating compressible Navier–Stokes equations, which is believed to describe the large-scale dynamics of the atmosphere and the ocean to a very good accuracy. An important feature of such large-scale dynamics is that it largely consists of slow motions in which the pressure gradient is nearly balanced by the Coriolis force, a state known as geostrophic balance. Various physical explanations have been given, some supported by numerical simulations, to describe how this comes about, but to our knowledge no rigorous mathematical proof has been proposed. (For a review of the geophysical background, see, e.g., [7].) One aim of this article is to prove that, in the limit of strong rotation and stratification, the solution of the primitive equations will approach geostrophic balance as $t \to \infty$, in the sense that the ageostrophic energy will be of the order of the Rossby number.

As illustrated by the simple one-dimensional model (4.3), here the basic mechanism for balance is the viscous damping of rapid oscillations, leaving the slow dynamics mostly unchanged. Separation of timescale, characterised by a small parameter ε , is therefore crucial for our result; this is obtained by considering the limit of strong rotation and stratification, or in other words, small Rossby number with Burger number of order one. We note that there are other physical mechanisms through which a balanced state may be reached. Working in an unbounded domain, an important example is the radiation of inertia–gravity waves to infinity in what is known as the classical geostrophic adjustment problem (see [11, §7.3] and further developments in [27]).

²⁰⁰⁰ Mathematics Subject Classification. Primary: 35B40, 37L25, 76U05.

Key words and phrases. Slow manifold, exponential asymptotics, primitive equations.

This research was partially supported by the National Science Foundation under grant NSF-DMS-0604235, by the Research Fund of Indiana University, and by a grant from the Nuffield Foundation.

Attempts to extend geostrophic balance to higher orders, and the closely related problem of eliminating rapid oscillations in numerical solutions (e.g., [3, 20, 16, 34]), led naturally to the concept of *slow manifold* [19], which has since become important in the study of rotating fluids (and more generally of systems with multiple timescales). We refer the reader to [21] for a thorough review, but for our purposes here, a slow manifold means a manifold in phase space on which the normal velocity is small; if the normal velocity is zero, we have an exact slow manifold. In the geophysical literature, there have been many papers proposing various formal asymptotic methods to construct slow manifolds (e.g., [38, 37]). A number of numerical studies closely related to the stability of slow manifolds have also been done (e.g., [10, 26]).

It was realised early on [19, 36] that in general no exact slow manifold exists and any construction is generally asymptotic in nature. For finite-dimensional systems, this can often be proved using considerations of exponential asymptotics (see, e.g., [15]). More recently, it has been shown explicitly [33] in an infinite-dimensional rotating fluid model that exponentially weak fast oscillations are generated spontaneously by vortical motion, implying that slow manifolds could at best be exponentially accurate (meaning the normal velocity on it be exponentially small). Theorem 2 shows, given the hypotheses, that an exponential accuracy can indeed be achieved for the primitive equations, albeit with a weaker dependence on ε .

From a more mathematical perspective, our exponentially slow manifold (see Lemma 2), which is also presented in [31] in a slightly different form, is obtained using a technique adapted from that first proposed in [22]. It involves truncating the PDE to a finite-dimensional system whose size depends on ε and applying a classical estimate from perturbation theory to the finite system. By carefully balancing the truncation size and the estimates on the finite system, one obtains a finite-dimensional exponentially accurate slow manifold. This estimate is local in time and only requires that the (instantaneous) variables and the forcing be in some Sobolev space H^s ; it (although not the long-time asymptotic result below) can thus be obtained for the inviscid equations as well. If our solution is also Gevrey (which is true for the primitive equations given Gevrey forcing), the ignored high modes are exponentially small, so the "total error" (i.e. normal velocity on the slow manifold) is also exponentially small.

Gevrey regularity of the solution is therefore crucial in obtaining exponential estimates. As with the Navier–Stokes equations [9], in the absence of boundaries and with Gevrey forcing, one can prove that the strong solution of the primitive equations also has Gevrey regularity [25]. For the present article, we need uniform bounds on the norms, which have been proved recently [23] following the global regularity results of [6, 13, 14]. Since our result also assumes strong rotation, however, one could have used an earlier work [2] which proved global regularity under a sufficiently strong rotation and then used [25] to obtain Gevrey regularity.

While our earlier paper [31] is concerned with a finite-time estimate on pointwise accuracy ("predictability"), in this article our aim is to obtain long-time asymptotic estimates (on "balance"). In this regard, the main problem for both the leading-order (Theorem 1) and higher-order (Theorem 2) estimates are the same: to bound the energy transfer, through the nonlinear term, from the slow to fast modes at the same order as the fast modes themselves. For this, one needs to handle not only *exact* fast–fast–slow resonances, whose absence has long been known in the

geophysical literature (cf. e.g., [4, 8, 17, 35] for discussions of related models), but also *near* resonances. A key part in our approach is an estimate involving near resonances in the primitive equations (cf. Lemma 1). Another method based on algebraic geometry to handle related near resonances can be found in [1].

Taken together with [31], the results here may be regarded as an extension of the single-frequency exponential estimates obtained in [22] to the ocean primitive equations, which have an infinite number of frequencies. Alternately, one may view Theorem 2 as an extension to exponential order of the leading-order results of [2] for a closely related model. Finally, our results here put a strong constraint on the nature of the global attractor [12] in the strong rotation limit: the attractor will have to lie within an exponentially thin neighbourhood of the slow manifold.

The rest of this article is arranged as follows. We begin in the next section by describing the ocean primitive equations (henceforth OPE) and recalling the known regularity results. In Section 3, we write the OPE in terms of fast-slow variables and in Fourier modes, followed by computing explicitly the operator corresponding to the nonlinear terms and describing its properties. In Section 4, we state and prove our leading-order estimate, that the solution of the OPE will be close to geostrophic balance as $t \to \infty$. In the last section, we state and prove our exponential-order estimate.

2. The Primitive Equations

We start by recalling the basic settings of the ocean primitive equations [18], and then recast the system in a form suitable for our aim in this article.

2.1. Setup. We consider the primitive equations for the ocean, scaled as in [24]

(2.1)
$$\partial_t \boldsymbol{v} + \frac{1}{\varepsilon} \left[\boldsymbol{v}^{\perp} + \nabla_2 p \right] + \boldsymbol{u} \cdot \nabla \boldsymbol{v} = \mu \Delta \boldsymbol{v} + f_{\boldsymbol{v}}$$
$$\frac{\partial_t \rho - \frac{1}{\varepsilon} u^3 + \boldsymbol{u} \cdot \nabla \rho = \mu \Delta \rho + f_{\rho},$$
$$\nabla \cdot \boldsymbol{u} = \nabla \cdot \boldsymbol{v} + \partial_z u^3 = 0,$$
$$\rho = -\partial_z p.$$

Here $\boldsymbol{u} = (u^1, u^2, u^3)$ and $\boldsymbol{v} = (u^1, u^2, 0)$ are the three- and two-dimensional fluid velocity, with $\boldsymbol{v}^{\perp} := (-u^2, u^1, 0)$. The variable ρ can be interpreted in two ways: One can take it to be the departure from a stably-stratified profile (with the usual Boussinesq approximation), with the full density of the fluid given by

(2.2)
$$\rho_{\text{full}}(x, y, z, t) = \rho_0 - \varepsilon^{-1} z \rho_1 + \rho(x, y, z, t),$$

for some positive constants ρ_0 and ρ_1 . Alternately, one can think of it to be, e.g., salinity or temperature that contributes linearly to the density. The pressure p is determined by the hydrostatic relation $\partial_z p = -\rho$ and the incompressibility condition $\nabla \cdot \boldsymbol{u} = 0$, and is not (directly) a function of ρ . We write $\nabla := (\partial_x, \partial_y, \partial_z)$, $\nabla_2 := (\partial_x, \partial_y, 0), \Delta := \partial_x^2 + \partial_y^2 + \partial_z^2$ and $\Delta_2 := \partial_x^2 + \partial_y^2$. The parameter ε is related to the Rossby and Froude numbers; in this paper we shall be concerned with the limit $\varepsilon \to 0$. In general the viscosity coefficients for \boldsymbol{v} and ρ are different; we have set them both to μ for clarity of presentation (the general case does not introduce any more essential difficulty). The variables (\boldsymbol{v}, ρ) evidently depend on the parameters ε and μ as well as on (\boldsymbol{x}, t) , but we shall not write this dependence explicitly. We work in three spatial dimensions, $\boldsymbol{x} := (x, y, z) = (x^1, x^2, x^3) \in [0, L_1] \times [0, L_2] \times [-L_3/2, L_3/2] =: \mathcal{M}$, with periodic boundary conditions assumed; we write $|\mathcal{M}| := L_1 L_2 L_3$. Moreover, following the practice in numerical simulations of stratified turbulence (see, e.g., [4]), we impose the following symmetry on the dependent variables:

(2.3)
$$\begin{aligned} & \boldsymbol{v}(x,y,-z) = \boldsymbol{v}(x,y,z), & p(x,y,-z) = p(x,y,z), \\ & u^3(x,y,-z) = -u^3(x,y,z), & \rho(x,y,-z) = -\rho(x,y,z); \end{aligned}$$

we say that v and p are even in z, while u^3 and ρ are odd in z. For this symmetry to persist, f_v must be even and f_ρ odd in z. Since u^3 and ρ are also periodic in z, we have $u^3(x, y, -L_3/2) = u^3(x, y, L_3/2) = 0$ and $\rho(x, y, -L_3/2) = \rho(x, y, L_3/2) = 0$; similarly, $\partial_z u^1 = 0$, $\partial_z u^2 = 0$ and $\partial_z p = 0$ on $z = 0, \pm L_3/2$ if they are sufficiently smooth (as will be assumed below). One may consider the symmetry conditions (2.3) as a way to impose the boundary conditions $u^3 = 0$, $\rho = 0$, $\partial_z u^1 = 0$, $\partial_z u^2 = 0$ and $\partial_z p = 0$ on z = 0 and $z = L_3/2$ in the effective domain $[0, L_1] \times [0, L_2] \times [0, L_3/2]$. All variables and the forcing are assumed to have zero mean in \mathcal{M} ; the symmetry conditions above ensure that this also holds for their products that appear below. It can be verified that the symmetry (2.3) is preserved by the OPE (2.1); that is, if it holds at t = 0, it continues to hold for t > 0.

2.2. Determining the pressure and vertical velocity. Since $u^3 = 0$ at z = 0, we can use (2.1c) to write

(2.4)
$$u^3(x,y,z) = -\int_0^z \nabla \cdot \boldsymbol{v}(x,y,z') \, \mathrm{d}z'.$$

Similarly, the pressure p can be written in terms of the density ρ as follows (cf. [28]). Let $p(x, y, z) = \langle p(x, y) \rangle + \delta p(x, y, z)$ where $\langle \cdot \rangle$ denotes z-average and where

(2.5)
$$\delta p(x, y, z) = -\int_{z_0}^{z} \rho(x, y, z') \, \mathrm{d}z'$$

with $z_0(x, y)$ chosen such that $\langle \delta p \rangle = 0$; this is most conveniently done using Fourier series (see below). Using the fact that

(2.6)
$$\int_{-L_3/2}^{L_3/2} \nabla \cdot \boldsymbol{v} \, \mathrm{d}z = -\int_{-L_3/2}^{L_3/2} \partial_z u^3 \, \mathrm{d}z = u^3(\cdot, -L_3/2) - u^3(\cdot, L_3/2) = 0,$$

and taking 2d divergence of the momentum equation (2.1a), we find

(2.7)
$$\frac{1}{\varepsilon} \left[\nabla \cdot \langle \boldsymbol{v}^{\perp} \rangle + \Delta_2 \langle \boldsymbol{p} \rangle \right] + \nabla \cdot \langle \boldsymbol{u} \cdot \nabla \boldsymbol{v} \rangle = \mu \Delta \nabla \cdot \langle \boldsymbol{v} \rangle + \nabla \cdot \langle \boldsymbol{f}_{\boldsymbol{v}} \rangle.$$

Here we have used the fact that z-integration commutes with horizontal differential operators. We can now solve for the average pressure $\langle p \rangle$,

(2.8)
$$\langle p \rangle = \Delta_2^{-1} \left[-\nabla \cdot \langle \boldsymbol{v}^{\perp} \rangle + \varepsilon \left(-\nabla \cdot \langle \boldsymbol{u} \cdot \nabla \boldsymbol{v} \rangle + \mu \Delta \nabla \cdot \langle \boldsymbol{v} \rangle + \nabla \cdot \langle f_{\boldsymbol{v}} \rangle \right) \right]$$

where Δ_2^{-1} is uniquely defined to have zero *xy*-average. With this, the momentum equation now reads

(2.9)
$$\partial_t \boldsymbol{v} + \frac{1}{\varepsilon} \left[\boldsymbol{v}^{\perp} - \nabla \Delta_2^{-1} \nabla \cdot \langle \boldsymbol{v}^{\perp} \rangle + \nabla_2 \delta \boldsymbol{p} \right] + \boldsymbol{u} \cdot \nabla \boldsymbol{v} - \nabla \Delta_2^{-1} \nabla \cdot \langle \boldsymbol{u} \cdot \nabla \boldsymbol{v} \rangle$$
$$= \mu \Delta \left(\boldsymbol{v} - \nabla \Delta_2^{-1} \nabla \cdot \langle \boldsymbol{v} \rangle \right) + f_{\boldsymbol{v}} - \nabla \Delta_2^{-1} \nabla \cdot \langle f_{\boldsymbol{v}} \rangle$$

2.3. Canonical form and regularity results. Besides the usual $L^{p}(\mathcal{M})$ and $H^{s}(\mathcal{M})$, with $p \in [1, \infty]$ and $s \geq 0$, we shall also need the Gevrey space $G^{\sigma}(\mathcal{M})$, defined as follows. For $\sigma \geq 0$, we say that $u \in G^{\sigma}(\mathcal{M})$ if

(2.10)
$$|e^{\sigma(-\Delta)^{1/2}}u|_{L^2} =: |u|_{G^{\sigma}} < \infty.$$

Let us denote our state variable $W = (\mathbf{v}, \rho)^{\mathrm{T}}$. We write $W \in L^{p}(\mathcal{M})$ if $\mathbf{v} \in L^{p}(\mathcal{M})^{2}$, $\rho \in L^{p}(\mathcal{M}), (\mathbf{v}, \rho)$ has zero average over \mathcal{M} and (\mathbf{v}, ρ) satisfies the symmetry (2.3), in the distribution sense as appropriate; analogous notations are used for $W \in H^{s}(\mathcal{M})$ and $W \in G^{\sigma}(\mathcal{M})$, and for the forcing f (which has to preserve the symmetries of W).

With u^3 given by (2.4) and δp by (2.5), we can write the OPE (2.1b) and (2.9) in the compact form

(2.11)
$$\partial_t W + \frac{1}{\varepsilon} LW + B(W, W) + AW = f.$$

The operators L, B and A are defined by

$$LW = \left(\boldsymbol{v}^{\perp} - \nabla\Delta_{2}^{-1}\nabla\cdot\langle\boldsymbol{v}^{\perp}\rangle + \nabla_{2}\delta p, -u^{3}\right)^{\mathrm{T}}$$

(2.12)
$$B(W, \hat{W}) = \left(\boldsymbol{u}\cdot\nabla\hat{\boldsymbol{v}} - \nabla\Delta_{2}^{-1}\nabla\cdot\langle\boldsymbol{u}\cdot\nabla\hat{\boldsymbol{v}}\rangle, \boldsymbol{u}\cdot\nabla\hat{\rho}\right)^{\mathrm{T}}$$
$$AW = -\left(\mu\Delta(\boldsymbol{v}-\nabla\Delta_{2}^{-1}\nabla\cdot\langle\boldsymbol{v}\rangle), \mu\Delta\rho\right)^{\mathrm{T}},$$

and the force f is given by

(2.13)
$$f = (f_{\boldsymbol{v}} - \nabla \Delta_2^{-1} \nabla \cdot \langle f_{\boldsymbol{v}} \rangle, f_{\rho})^{\mathrm{T}}.$$

The following properties are known (see, e.g., [25]). The operator L is antisymmetric: for any $W \in L^2(\mathcal{M})$

 $(2.14) \quad (LW, W)_{L^2} = 0;$

B conserves energy: for any $W \in H^1(\mathcal{M})$ and $\hat{W} \in H^1(\mathcal{M})$,

 $(2.15) \quad (W, B(\hat{W}, W))_{L^2} = 0;$

and A is coercive: for any $W \in H^2(\mathcal{M})$,

 $(2.16) \quad (AW,W)_{L^2} = \mu \, |\nabla W|_{L^2}^2.$

We shall need the following regularity results for the OPE (here K_s and M_{σ} are continuous increasing functions of their arguments):

Theorem 0. Let $W_0 \in H^1$ and $f \in L^{\infty}(\mathbb{R}_+; L^2)$. Then for all $t \ge 0$ there exists a solution $W(t) \in H^1$ of (2.11) with $W(0) = W_0$ and

 $(2.17) \quad |W(t)|_{H^1} \le K_0(|W_0|_{H^1}, ||f||_0)$

where, here and henceforth, $||f||_s := \operatorname{ess\,sup}_{t\geq 0} |f(t)|_{H^s}$ for $s \geq 0$. Moreover, there exists a time $T_1(|W_0|_{H^1}, ||f||_0)$ such that for $t \geq T_1$,

 $(2.18) |W(t)|_{H^1} \le K_1(||f||_0).$

Similarly, if $f \in L^{\infty}(\mathbb{R}_+; H^{s-1})$, there exists a time $T_s(|W_0|_{H^1}, ||f||_{s-1})$ such that (2.19) $|W(t)|_{H^s} \leq K_s(||f||_{s-1})$

for $t \geq T_s$. Finally, fixing $\sigma > 0$, if also $\nabla f \in L^{\infty}(\mathbb{R}_+; G^{\sigma})$, there exists a time $T_{\sigma}(|W_0|_{H^1}, |\nabla f|_{G^{\sigma}})$ such that, for $t \geq T_{\sigma}$

(2.20) $|\nabla^2 W(t)|_{G^{\sigma}} \leq M_{\sigma}(|\nabla f|_{G^{\sigma}}).$

The proof of (2.17)–(2.18) can be found in [12]; the higher-order results (2.19) can be found in [23]. Both these works followed [6] and [13]. The result (2.20) follows from [25] and using (2.19) for s = 2.

Since we are concerned with the limit of small ε , however, one might also be able to obtain (2.17) and (2.19) following the method used in [2] for the Boussinesq (non-hydrostatic) model. One could then proceed to obtain (2.20) as above.

3. Normal Modes

In this section, we decompose the solution W into its slow and fast components, expand them in Fourier modes, and state a lemma that will be used in sections 4 and 5 below.

3.1. Fast and slow variables. The Ertel potential vorticity

(3.1)
$$q_E = \nabla^{\perp} \cdot \boldsymbol{v} - \partial_z \rho + \varepsilon [(\partial_z \boldsymbol{v}) \cdot \nabla^{\perp} \rho - \partial_z \rho (\nabla^{\perp} \cdot \boldsymbol{v})],$$

where $\nabla^{\perp} := (-\partial_y, \partial_x, 0)$, plays a central role in geophysical fluid dynamics since it is a material invariant in the absence of forcing and viscosity. In this paper, however, it is easier to work with the *linearised* potential vorticity (henceforth simply called *potential vorticity*)

$$(3.2) \quad q := \nabla^{\perp} \cdot \boldsymbol{v} - \partial_z \rho$$

From (2.1), its evolution equation is

(3.3)
$$\partial_t q + \nabla^{\perp} \cdot (\boldsymbol{u} \cdot \nabla \boldsymbol{v}) - \partial_z (\boldsymbol{u} \cdot \nabla \rho) = \mu \Delta q + f_q$$

where $f_q := \nabla^{\perp} \cdot f_v - \partial_z f_{\rho}$. Let $\psi^0 := \Delta^{-1}q$, uniquely defined by requiring that ψ^0 has zero integral over \mathcal{M} , and let

(3.4)
$$W^0 := \begin{pmatrix} \boldsymbol{v}^0 \\ \rho^0 \end{pmatrix} := \begin{pmatrix} \nabla^{\perp} \psi^0 \\ -\partial_z \psi^0 \end{pmatrix}.$$

We note a mild abuse of notation on v^0 and ∇^{\perp} : $W^0 = (-\partial_y \psi^0, \partial_x \psi^0, -\partial_z \psi^0)^{\mathrm{T}}$.

A little computation shows that W^0 lies in the kernel of the antisymmetric operator L, that is, $LW^0 = 0$. Conversely, if LW = 0, then $W = (\nabla^{\perp}\Psi, -\partial_z\Psi)^{\mathrm{T}}$ for some Ψ : Since $u^3 = 0$, we have $\nabla \cdot \boldsymbol{v} = 0$, so $\boldsymbol{v} = \nabla^{\perp}\Psi + \boldsymbol{V}$ for some $\Psi(x, y, z)$ and $\boldsymbol{V}(z)$. Now

(3.5)
$$0 = \boldsymbol{v}^{\perp} - \nabla_2 \Delta_2^{-1} \nabla \cdot \langle v^{\perp} \rangle + \nabla_2 \delta p$$
$$= -\nabla_2 \Psi + \boldsymbol{V}^{\perp} + \nabla_2 \Delta_2^{-1} \Delta_2 \langle \Psi \rangle + \nabla_2 \delta p.$$

Since all other terms are horizontal gradients and V does not depend on (x, y), we must have V = 0. Writing $\Psi(x, y, z) = \tilde{\Psi}(x, y, z) + \langle \Psi(x, y) \rangle$ where $\tilde{\Psi}(x, y, z)$ has zero z-average, the terms that do not depend on z cancel and we are left with

$$(3.6) \quad -\nabla_2 \Psi + \nabla_2 \delta p = 0.$$

So $\delta p(x, y, z) = \tilde{\Psi}(x, y, z) + \Phi(z)$; but since $\langle \delta p \rangle = 0$, $\Phi = 0$ and thus $\rho = -\partial_z \Psi$ by (2.5). Therefore the null space of L is completely characterised by (3.4),

(3.7)
$$\ker L = \{ W^0 : W^0 = (\nabla^{\perp} \psi^0, -\partial_z \psi^0)^{\mathrm{T}} \}.$$

With $\psi^0 = \Delta^{-1} (\nabla^{\perp} \cdot \boldsymbol{v} - \partial_z \rho)$ as above, this also defines a projection $W \mapsto W^0$. We call W^0 our *slow variable*. Letting B^0 be the projection of B to ker L,

(3.8)
$$B^{0}(W,\hat{W}) := \begin{pmatrix} \nabla^{\perp} \Delta^{-1} [\nabla^{\perp} \cdot (\boldsymbol{u} \cdot \nabla \hat{\boldsymbol{v}}) - \partial_{z} (\boldsymbol{u} \cdot \nabla \hat{\rho})] \\ -\partial_{z} \Delta^{-1} [\nabla^{\perp} \cdot (\boldsymbol{u} \cdot \nabla \hat{\boldsymbol{v}}) - \partial_{z} (\boldsymbol{u} \cdot \nabla \hat{\rho})] \end{pmatrix},$$

we find that W^0 satisfies

(3.9) $\partial_t W^0 + B^0(W, W) + AW^0 = f^0$

where $f^0 = (\nabla^{\perp} \Delta^{-1} f_q, -\partial_z \Delta^{-1} f_q)^{\mathrm{T}}$ is the slow forcing. Now let

(3.10)
$$W^{\varepsilon} = \begin{pmatrix} \boldsymbol{v}^{\varepsilon} \\ \rho^{\varepsilon} \end{pmatrix} := W - W^{0} = \begin{pmatrix} \boldsymbol{v} - \boldsymbol{v}^{0} \\ \rho - \rho^{0} \end{pmatrix}.$$

It will be seen below in Fourier representation that W^{ε} is a linear combination of eigenfunctions of L with imaginary eigenvalues whose moduli are bounded from below; we thus call W^{ε} our *fast variable*. Since $\nabla \cdot \boldsymbol{v}^0 = 0$, the vertical velocity u^3 is a purely fast variable. In analogy with (3.9), we have

(3.11)
$$\partial_t W^{\varepsilon} + \frac{1}{\varepsilon} L W^{\varepsilon} + B^{\varepsilon}(W, W) + A W^{\varepsilon} = f^{\varepsilon}$$

where $B^{\varepsilon}(W, \hat{W}) := B(W, \hat{W}) - B^0(W, \hat{W})$ and $f^{\varepsilon} := f - f^0$.

The fast variable has no potential vorticity, as can be seen by computing $\nabla^{\perp} \cdot \boldsymbol{v}^{\varepsilon} - \partial_z \rho^{\varepsilon} = q - \nabla^{\perp} \cdot \nabla^{\perp} \psi^0 - \partial_{zz} \psi^0 = 0$. Since the slow variable is completely determined by the potential vorticity, this implies that the fast and slow variables are orthogonal in $L^2(\mathcal{M})$,

(3.12)
$$(W^0, W^{\varepsilon})_{L^2} = (\boldsymbol{v}^0, \boldsymbol{v}^{\varepsilon})_{L^2} + (\rho^0, \rho^{\varepsilon})_{L^2} = (\nabla^{\perp} \psi^0, \boldsymbol{v}^{\varepsilon})_{L^2} - (\partial_z \psi^0, \rho^{\varepsilon})_{L^2} = (\psi_0, -\nabla^{\perp} \cdot \boldsymbol{v}^{\varepsilon} + \partial_z \rho^{\varepsilon})_{L^2} = 0.$$

Of central interest in this paper is the "fast energy"

 $(3.13) \quad \frac{1}{2} |W^{\varepsilon}|_{L^2}^2 = \frac{1}{2} \left(|\boldsymbol{v}^{\varepsilon}|_{L^2}^2 + |\rho^{\varepsilon}|_{L^2}^2 \right).$

Its time derivative can be computed as follows. Using (3.12), we have after integrating by parts

$$(3.14) \quad (W^{\varepsilon}, \partial_t W)_{L^2} = (W^{\varepsilon}, \partial_t W^0)_{L^2} + (W^{\varepsilon}, \partial_t W^{\varepsilon})_{L^2} = \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} |W^{\varepsilon}|_{L^2}^2.$$

Now (2.15) implies that

$$(3.15) \quad (W^{\varepsilon}, B(W, W))_{L^2} = (W^{\varepsilon}, B(W, W^0 + W^{\varepsilon}))_{L^2} = (W^{\varepsilon}, B(W, W^0))_{L^2}.$$

Putting these together with (2.14) and (2.16), we find

(3.16)
$$\frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} |W^{\varepsilon}|_{L^{2}}^{2} + \mu |\nabla W^{\varepsilon}|_{L^{2}}^{2} = -(W^{\varepsilon}, B(W, W^{0}))_{L^{2}} + (W^{\varepsilon}, f^{\varepsilon})_{L^{2}}.$$

3.2. Fourier expansion. Thanks to the regularity results in Theorem 0, our solution W(t) is smooth and we can thus expand it in Fourier series,

(3.17) $\boldsymbol{v}(\boldsymbol{x},t) = \sum_{\boldsymbol{k}} \boldsymbol{v}_{\boldsymbol{k}}(t) e^{i\boldsymbol{k}\cdot\boldsymbol{x}}$ and $\rho(\boldsymbol{x},t) = \sum_{\boldsymbol{k}} \rho_{\boldsymbol{k}}(t) e^{i\boldsymbol{k}\cdot\boldsymbol{x}}.$

Here $\mathbf{k} = (k_1, k_2, k_3) \in \mathbb{Z}_L$ where $\mathbb{Z}_L = \mathbb{R}^3 / \mathcal{M} = \{(2\pi l_1 / L_1, 2\pi l_2 / L_2, 2\pi l_3 / L_3) : (l_1, l_2, l_3) \in \mathbb{Z}^3\}$; any wavevector \mathbf{k} is henceforth understood to live in \mathbb{Z}_L . We also denote $\mathbf{k}' := (k_1, k_2, 0)$ and write $\mathbf{k}' \wedge \mathbf{j}' := k_1 j_2 - k_2 j_1$. Since our variables have zero average over \mathcal{M} , $\mathbf{v}_{\mathbf{k}} = 0$ when $\mathbf{k} = 0$; moreover, since ρ is odd in $z, \rho_{\mathbf{k}} = 0$

whenever $k_3 = 0$. Thus $W_{\mathbf{k}} := (\mathbf{v}_{\mathbf{k}}, \rho_{\mathbf{k}}) = 0$ when $\mathbf{k} = 0$, which allows us to write the H^s norm simply as

(3.18) $|W|_{H^s}^2 = \sum_{k} |k|^{2s} |W_k|^2$

and (see (2.10) for the definition of G^{σ})

(3.19) $|W|_{G^{\sigma}}^2 = \sum_{k} e^{2\sigma |k|} |W_{k}|^2.$

The antisymmetric operator L is diagonal in Fourier space, meaning that $L_{kl} = 0$ when $k \neq l$; we shall thus write $L_k := L_{kk}$. When $k_3 \neq 0$, we have

(3.20)
$$L_{\mathbf{k}} = \begin{pmatrix} 0 & -1 & -k_1/k_3 \\ 1 & 0 & -k_2/k_3 \\ k_1/k_3 & k_2/k_3 & 0 \end{pmatrix}$$

For $\mathbf{k}' \neq 0$, its eigenvalues are $\omega_{\mathbf{k}}^0 = 0$ and $i\omega_{\mathbf{k}}^{\pm} = \pm i|\mathbf{k}|/k_3$, where $|\mathbf{k}| := (k_1^2 + k_2^2 + k_3^2)^{1/2}$, with eigenvectors

(3.21)
$$X_{\boldsymbol{k}}^{0} = \frac{1}{|\boldsymbol{k}|} \begin{pmatrix} k_{2} \\ -k_{1} \\ k_{3} \end{pmatrix}$$
 and $X_{\boldsymbol{k}}^{\pm} = \frac{1}{\sqrt{2}|\boldsymbol{k}'||\boldsymbol{k}|} \begin{pmatrix} -k_{2}k_{3} \pm ik_{1}|\boldsymbol{k}| \\ k_{1}k_{3} \pm ik_{2}|\boldsymbol{k}| \\ |\boldsymbol{k}'|^{2} \end{pmatrix}$.

When $\mathbf{k}' = 0$, we have $\omega_{\mathbf{k}}^0 = 0$ and $i\omega_{\mathbf{k}}^{\pm} = \pm i$ as eigenvalues with eigenvectors

(3.22)
$$X_{\boldsymbol{k}}^{0} = \begin{pmatrix} 0\\ 0\\ \operatorname{sgn} k_{3} \end{pmatrix}$$
 and $X_{\boldsymbol{k}}^{\pm} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1\\ \mp i\\ 0 \end{pmatrix}$.

For k fixed, these eigenvectors are orthonormal under the inner product \cdot in \mathbb{C}^3 .

When $k_3 = 0$, the fact that $\rho_{\mathbf{k}} = 0$ and $\mathbf{k} \cdot \mathbf{v}_{\mathbf{k}} = 0$ implies that the space is onedimensional for each \mathbf{k} (in fact, it is known that the vertically-averaged dynamics is that of the rotating 2d Navier–Stokes equations). Since projecting to the $k_3 = 0$ subspace is equivalent to taking vertical average, we compute

(3.23)
$$\langle LW \rangle = (\langle \boldsymbol{v}^{\perp} \rangle - \nabla_2 \Delta_2^{-1} \nabla \cdot \langle \boldsymbol{v}^{\perp} \rangle, 0)^{\mathrm{T}}$$

where we have used $\langle u^3 \rangle = 0$ (since u^3 is odd) and $\langle \delta p \rangle = 0$ (by definition). Reasoning as in (3.5)–(3.6) above, we find that $\langle LW \rangle = 0$, that is, the verticallyaveraged ($k_3 = 0$) component is completely slow. In this case we can thus write

(3.24)
$$\omega_{\mathbf{k}}^{0} = 0$$
 and $X_{\mathbf{k}}^{0} = \frac{1}{|\mathbf{k}'|} \begin{pmatrix} k_{2} \\ -k_{1} \\ 0 \end{pmatrix}$,

which can be included in the generic case $\mathbf{k}' \neq 0$ in computations. Since the $k_3 = 0$ component is completely slow, $\langle W^{\varepsilon} \rangle = 0$, there is no need to fix $X_{\mathbf{k}}^{\pm}$.

We note that since $k_3 \neq 0$, $|\omega_{\mathbf{k}}^{\pm}| \geq 1$, viz.,

(3.25)
$$\inf |\omega_{\mathbf{k}}^{\pm}|^2 = \inf_{k_3 \neq 0} \left\{ \frac{k_1^2 + k_2^2 + k_3^2}{k_3^2}, 1 \right\} = 1$$

In what follows, it is convenient to use $\{X_{k}^{0}, X_{k}^{\pm}\}$ as basis. We can now write

(3.26)
$$\begin{aligned} W^{0}(\boldsymbol{x},t) &:= \sum_{\boldsymbol{k}} w_{\boldsymbol{k}}^{0}(t) X_{\boldsymbol{k}}^{0} \mathrm{e}^{i\boldsymbol{k}\cdot\boldsymbol{x}} \\ W^{\varepsilon}(\boldsymbol{x},t) &:= \sum_{\boldsymbol{k}}^{s} w_{\boldsymbol{k}}^{s}(t) X_{\boldsymbol{k}}^{s} \mathrm{e}^{-\mathrm{i}\omega_{\boldsymbol{k}}^{s}t/\varepsilon} \mathrm{e}^{\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}}, \end{aligned}$$

where $s \in \{-1, +1\}$, which we write as $\{-, +\}$ when it appears as a label. The Fourier coefficients $w_{\mathbf{k}}^0$ and $w_{\mathbf{k}}^{\pm}$ are complex numbers that depend on t only, with $w_0^0 = 0$ and $w_{(k_1,k_2,0)}^{\pm} = 0$. With $\alpha \in \{-1, 0, +1\}$, they can be computed using

(3.27)
$$w_{\boldsymbol{k}}^{\alpha}(t) = \frac{1}{|\mathcal{M}|} \int_{\mathcal{M}} W(\boldsymbol{x}, t) \cdot X_{\boldsymbol{k}}^{\alpha} e^{\mathrm{i}\omega_{\boldsymbol{k}}^{\alpha}t/\varepsilon - \mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}} \,\mathrm{d}\boldsymbol{x}.$$

The following relations hold:

(3.28)
$$|W^0|_{L^2}^2 = \sum_{k} |w_k^0|^2$$
 and $|W^{\varepsilon}|_{L^2}^2 = \sum_{k}^{s} |w_k^s|^2$.

In addition, the fact that (v^0, ρ^0) is real implies

(3.29)
$$w_{-\mathbf{k}}^0 = -\overline{w_{\mathbf{k}}^0}$$
 and $w_{(k_1,k_2,-k_3)}^0 = w_{(k_1,k_2,k_3)}^0$

where overbars denote complex conjugation. Similarly, since $(v^{\varepsilon}, \rho^{\varepsilon})$ is real,

(3.30)
$$w_{-\mathbf{k}}^{\pm} = w_{\mathbf{k}}^{\pm}$$
 and $w_{(k_1,k_2,-k_3)}^{\pm} = -w_{(k_1,k_2,k_3)}^{\pm}$

when $\mathbf{k}' \neq 0$ and, when $\mathbf{k}' = 0$,

$$(3.31) \quad w_{(0,0,-k_3)}^{\pm} = w_{(0,0,k_3)}^{\mp}$$

We shall see below that, the linear oscillations having been factored out, the variable w_{k}^{s} is slow at leading order. Similarly to W, we write the forcing f as

(3.32)
$$\begin{aligned} f^0(\boldsymbol{x},t) &:= \sum_{\boldsymbol{k}} f^0_{\boldsymbol{k}}(t) X^0_{\boldsymbol{k}} \mathrm{e}^{\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}} \\ f^{\varepsilon}(\boldsymbol{x},t) &:= \sum_{\boldsymbol{k}}^s f^s_{\boldsymbol{k}}(t) X^s_{\boldsymbol{k}} \mathrm{e}^{\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}}, \end{aligned}$$

where, unlike in (3.26), there is no factor of $e^{-i\omega_{k}^{s}t/\varepsilon}$ in the definition of f^{ε} . As noted above, f must satisfy the same symmetries as W, so the above properties of $w_{\mathbf{k}}^{\alpha}$ also hold for $f_{\mathbf{k}}^{\alpha}$; we note in particular that $f_{\mathbf{k}}^{\pm} = 0$ when $k_3 = 0$. For later convenience, we define the operator ∂_t^* by

(3.33)
$$\partial_t^* W := \mathrm{e}^{-tL/\varepsilon} \partial_t \, \mathrm{e}^{tL/\varepsilon} W.$$

From (2.11), we find

$$(3.34) \quad \partial_t^* W + B(W, W) + AW = f,$$

which is $\partial_t W$ with the large antisymmetric term removed.

Now the nonlinear term on the rhs of (3.16) can be written as

(3.35)
$$(W^{\varepsilon}, B(W^{0} + W^{\varepsilon}, W^{0}))_{L^{2}} = (W^{\varepsilon}, B(W^{0}, W^{0}))_{L^{2}} + (W^{\varepsilon}, B(W^{\varepsilon}, W^{0}))_{L^{2}} = (W^{\varepsilon}, B(W^{0}, W^{0}))_{L^{2}} - (W^{0}, B(W^{\varepsilon}, W^{\varepsilon}))_{L^{2}},$$

where the identity $(W^0, B(W^{\varepsilon}, W^{\varepsilon}))_{L^2} = -(W^{\varepsilon}, B(W^{\varepsilon}, W^0))_{L^2}$ had been obtained from (2.15).

First, let

$$(W^{\varepsilon}, B(W^{0}, W^{0}))_{L^{2}} = |\mathcal{M}| \sum_{\boldsymbol{jkl}}^{s} w_{\boldsymbol{j}}^{0} w_{\boldsymbol{k}}^{0} \overline{w_{\boldsymbol{l}}^{s}} \operatorname{i}(X_{\boldsymbol{j}}^{0} \cdot \boldsymbol{k}') (X_{\boldsymbol{k}}^{0} \cdot X_{\boldsymbol{l}}^{s}) \, \delta_{\boldsymbol{j}+\boldsymbol{k}-\boldsymbol{l}} \operatorname{e}^{\mathrm{i}\omega_{\boldsymbol{l}}^{s}t/\varepsilon}$$

$$(3.36)$$

$$= \sum_{\boldsymbol{jkl}}^{s} w_{\boldsymbol{j}}^{0} w_{\boldsymbol{k}}^{0} \overline{w_{\boldsymbol{l}}^{s}} B_{\boldsymbol{jkl}}^{00s} \operatorname{e}^{\mathrm{i}\omega_{\boldsymbol{l}}^{s}t/\varepsilon}$$

where $\delta_{j+k-l} = 1$ when j + k = l and 0 otherwise, and where $(3.37) \quad B^{00s}_{\boldsymbol{j}\boldsymbol{k}\boldsymbol{l}} := \mathrm{i} \left| \mathcal{M} \right| \delta_{\boldsymbol{j}+\boldsymbol{k}-\boldsymbol{l}}(X^{0}_{\boldsymbol{j}} \cdot \boldsymbol{k}')(X^{0}_{\boldsymbol{k}} \cdot X^{s}_{\boldsymbol{l}}).$

It is easy to verify from (3.37) that $B_{jkl}^{00s} = 0$ when $|j'| |k'| l_3 = 0$, so we consider the other cases. For the first factor, we have

(3.38)
$$X_{\boldsymbol{j}}^{0} \cdot \boldsymbol{k}' = \frac{\boldsymbol{k}' \wedge \boldsymbol{j}'}{|\boldsymbol{j}|}.$$

For the second factor, we have

(3.39)

$$X_{\boldsymbol{k}}^{0} \cdot X_{\boldsymbol{l}}^{s} = \frac{k_{2} - \mathrm{i}sk_{1}}{\sqrt{2}|\boldsymbol{k}|} \qquad \text{when } \boldsymbol{l}' = 0, \text{ and}$$

$$X_{\boldsymbol{k}}^{0} \cdot X_{\boldsymbol{l}}^{s} = \frac{k_{3}|\boldsymbol{l}'|^{2} - (\boldsymbol{k}' \cdot \boldsymbol{l}')l_{3} - \mathrm{i}s(\boldsymbol{l}' \wedge \boldsymbol{k}')|\boldsymbol{l}|}{\sqrt{2}|\boldsymbol{k}||\boldsymbol{l}||\boldsymbol{l}'|} \qquad \text{when } \boldsymbol{l}' \neq 0.$$

From these, we have the bound

$$(3.40) |B_{\boldsymbol{j}\boldsymbol{k}\boldsymbol{l}}^{00s}| \leq \frac{3|\mathcal{M}|}{\sqrt{2}} \frac{|\boldsymbol{k}'||\boldsymbol{j}'|}{|\boldsymbol{j}|}.$$
Next, we consider
$$(W^{0}, B(W^{\varepsilon}, W^{\varepsilon}))_{L^{2}}$$

$$(3.41) = |\mathcal{M}| \sum_{\boldsymbol{j}\boldsymbol{k}\boldsymbol{l}}^{rs} w_{\boldsymbol{j}}^{r} w_{\boldsymbol{k}}^{s} \overline{w_{\boldsymbol{l}}^{0}} \operatorname{i} (\mathsf{V}X_{\boldsymbol{j}}^{r} \cdot \boldsymbol{k}) (X_{\boldsymbol{k}}^{s} \cdot X_{\boldsymbol{l}}^{0}) \,\delta_{\boldsymbol{j}+\boldsymbol{k}-\boldsymbol{l}} \,\mathrm{e}^{-\mathrm{i}(\omega_{\boldsymbol{j}}^{r}+\omega_{\boldsymbol{k}}^{s})t/\varepsilon}$$

$$= \sum_{\boldsymbol{j}\boldsymbol{k}\boldsymbol{l}}^{rs} w_{\boldsymbol{j}}^{r} w_{\boldsymbol{k}}^{s} \overline{w_{\boldsymbol{l}}^{0}} \, B_{\boldsymbol{j}\boldsymbol{k}\boldsymbol{l}}^{rs0} \,\mathrm{e}^{-\mathrm{i}(\omega_{\boldsymbol{j}}^{r}+\omega_{\boldsymbol{k}}^{s})t/\varepsilon}$$

where

(3.42)
$$B_{\boldsymbol{j}\boldsymbol{k}\boldsymbol{l}}^{rs0} := \mathrm{i} |\mathcal{M}| \, \delta_{\boldsymbol{j}+\boldsymbol{k}-\boldsymbol{l}} \, (\mathsf{V}X_{\boldsymbol{j}}^r \cdot \boldsymbol{k}) (X_{\boldsymbol{k}}^s \cdot X_{\boldsymbol{l}}^0)$$

and where the operator V, which produces an incompressible velocity vector out of X_{j}^{r} , is defined by

and

(3.43)
$$\begin{array}{l} \mathsf{V}X_{\bm{j}}^{r} = X_{\bm{j}}^{r} & \text{when } j_{3}|\bm{j}'| = 0, \\ \mathsf{V}X_{\bm{j}}^{r} = \frac{1}{\sqrt{2}|\bm{j}||\bm{j}'|} \begin{pmatrix} -j_{2}j_{3} + \mathrm{i}rj_{1}|\bm{j}| \\ j_{1}j_{3} + \mathrm{i}rj_{2}|\bm{j}| \\ -\mathrm{i}r|\bm{j}'|^{2}|\bm{j}|/j_{3} \end{pmatrix} & \text{when } j_{3}|\bm{j}'| \neq 0. \end{array}$$

Thus, we have $VX_{j}^{r} \cdot \mathbf{k} = 0$ when $j_{3} = 0$,

(3.44)
$$\forall X_{\boldsymbol{j}}^r \cdot \boldsymbol{k} = (k_1 - \mathrm{i} r k_2)/\sqrt{2}$$

when j' = 0, and

(3.45)
$$\forall X_{\boldsymbol{j}}^{r} \cdot \boldsymbol{k} = \frac{j_{3}(\boldsymbol{j}' \wedge \boldsymbol{k}') + \mathrm{i}r|\boldsymbol{j}|(\boldsymbol{j}' \cdot \boldsymbol{k}') - \mathrm{i}r|\boldsymbol{j}'|^{2}|\boldsymbol{j}| k_{3}/j_{3}}{\sqrt{2}|\boldsymbol{j}||\boldsymbol{j}'|}$$

in the generic case $j_3|\boldsymbol{j}'| \neq 0$. In all cases, we have the bound (3.46) $|\nabla X_{\boldsymbol{j}}^r \cdot \boldsymbol{k}| \leq |\mathcal{M}| (\sqrt{2} |\boldsymbol{k}'| + |\boldsymbol{j}'| |k_3|/|j_3|).$ Next, $X_{\boldsymbol{k}}^s \cdot X_{\boldsymbol{l}}^0 = 0$ when $k_3 = 0$ or $\boldsymbol{k}' = \boldsymbol{l}' = 0$, and

$$\begin{aligned} X_{\boldsymbol{k}}^{s} \cdot X_{\boldsymbol{l}}^{0} &= \frac{l_{2} + \mathrm{i} s l_{1}}{\sqrt{2} |\boldsymbol{l}|} & \text{when } \boldsymbol{l}' \neq 0 \text{ and } \boldsymbol{k}' = 0, \\ (3.47) \quad X_{\boldsymbol{k}}^{s} \cdot X_{\boldsymbol{l}}^{0} &= \mathrm{sgn} \, l_{3} \frac{|\boldsymbol{k}'|}{\sqrt{2} |\boldsymbol{k}|} & \text{when } \boldsymbol{l}' = 0 \text{ and } \boldsymbol{k}' \neq 0, \\ X_{\boldsymbol{k}}^{s} \cdot X_{\boldsymbol{l}}^{0} &= \frac{-(\boldsymbol{k}' \cdot \boldsymbol{l}') k_{3} + \mathrm{i} s(\boldsymbol{k}' \wedge \boldsymbol{l}') |\boldsymbol{k}| + |\boldsymbol{k}'|^{2} l_{3}}{\sqrt{2} |\boldsymbol{k}| |\boldsymbol{k}'| |\boldsymbol{l}|} & \text{when } |\boldsymbol{k}'| \, |\boldsymbol{l}'| \, k_{3} \neq 0. \end{aligned}$$

These give us the bound

(3.48) $|X_{\boldsymbol{k}}^s \cdot X_{\boldsymbol{l}}^0| \le \sqrt{5/2}$ in all cases and, together with (3.46), when $j_3 \ne 0$,

(3.49) $|B_{jkl}^{rs0}| \le \sqrt{5} |\mathcal{M}| (|\mathbf{k}'| + |\mathbf{j}'| |k_3|/|j_3|).$

When $j_3k_3 = 0$ or $\boldsymbol{l} = 0$, we have $B_{\boldsymbol{j}\boldsymbol{k}\boldsymbol{l}}^{rs0} = 0$.

3.3. Fast–Fast–Slow Resonances. We first write (3.41) as

$$(3.50) \quad (W^0, B(W^{\varepsilon}, W^{\varepsilon}))_{L^2} = \frac{1}{2} \sum_{\boldsymbol{jkl}}^{rs} w_{\boldsymbol{j}}^r w_{\boldsymbol{k}}^s \overline{w_{\boldsymbol{l}}^0} \left(B_{\boldsymbol{jkl}}^{rs0} + B_{\boldsymbol{kjl}}^{sr0} \right) \mathrm{e}^{-\mathrm{i}(\omega_{\boldsymbol{j}}^r + \omega_{\boldsymbol{k}}^s)t/\varepsilon}.$$

It has long been known in the geophysical community that many rotating fluid models "have no fast–fast–slow resonances" (see, e.g., [35] for the shallow-water equations and [4] for the Boussinesq equations). In our notation, the absence of *exact* fast–fast–slow resonances means that $B_{jkl}^{rs0} + B_{kjl}^{sr0} = 0$ whenever $\omega_j^r + \omega_k^s = 0$; the significance of this will be apparent below [see the development following (4.16)]. For our purpose, however, we also need to consider *near* resonances, i.e. those cases when $|\omega_j^r + \omega_k^s|$ is small but nonzero. The following "no-resonance" lemma contains the estimate we need:

Lemma 1. For any $\boldsymbol{j}, \boldsymbol{k}, \boldsymbol{l} \in \mathbb{Z}_L$ with $\boldsymbol{l} \neq 0$,

(3.51)
$$|B_{jkl}^{rs0} + B_{kjl}^{sr0}| \le c_{nr} |\mathcal{M}| \left(\frac{|j||k|}{|l|} + |j_3| + |k_3|\right) |\omega_j^r + \omega_k^s|$$

where c_{nr} is an absolute constant.

We note that $B_{jkl}^{rs0} = B_{kjl}^{sr0} = 0$ when l = 0 by (3.41), so this case is trivial. We defer the proof to Appendix A.

4. Leading-Order Estimates

In this section, we discuss the leading-order case of our general problem. This is done separately due to its geophysical interest and since it requires qualitatively weaker hypotheses. As before, $W(t) = W^0(t) + W^{\varepsilon}(t)$ is the solution of the OPE (2.11) with initial conditions $W(0) = W_0$, and $K_{\rm g}(\cdot)$ is a continuous and increasing function of its argument.

Theorem 1. Suppose that the initial data $W_0 \in H^1(\mathcal{M})$ and that the forcing $f \in L^{\infty}(\mathbb{R}_+; H^2) \cap W^{1,\infty}(\mathbb{R}_+; L^2)$, with

(4.1) $||f||_{g} := \operatorname{ess\,sup}_{t>0} (|f(t)|_{H^2} + |\partial_t f(t)|_{L^2}).$

Then there exist $T_{g} = T_{g}(|W_{0}|_{H^{1}}, ||f||_{g}, \varepsilon)$ and $K_{g} = K_{g}(||f||_{g})$, such that for $t \ge T_{g}$, (4.2) $|W^{\varepsilon}(t)|_{L^{2}} \le \sqrt{\varepsilon} K_{g}(||f||_{g})$.

In geophysical parlance, our result states that, for given initial data and forcing, the solution of the OPE will become geostrophically balanced (in the sense that the ageostrophic component W^{ε} is of order $\sqrt{\varepsilon}$) after some time. We note that the forcing may be time-dependent (although $||f||_{\rm g}$ cannot depend on ε) and need not be geostrophic; this will not be the case when we consider higher-order balance later. Also, in contrast to the higher-order result in the next section, no restriction on ε is necessary in this case.

The linear mechanism of this "geostrophic decay" may be appreciated by modelling (3.11), without the nonlinear term, by the following ODE

(4.3)
$$\frac{\mathrm{d}x}{\mathrm{d}t} + \frac{\mathrm{i}}{\varepsilon}x + \mu x = f$$

where $\mu > 0$ is a constant and f = f(t) is given independently of ε . The skewhermitian term ix/ε causes oscillations of x whose frequency grows as $\varepsilon \to 0$. In this limit, the forcing becomes less effective since f varies slowly by hypothesis while the damping remains unchanged, so x will eventually decay to the order of the "net forcing" $\sqrt{\varepsilon}f$. More concretely, let $z(t) = e^{it/\varepsilon}x(t)$ and write (4.3) as

(4.4)
$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\mathrm{e}^{\mu t/2} z \right) + \frac{\mu}{2} \mathrm{e}^{\mu t/2} z = \mathrm{e}^{\mu t/2 - \mathrm{i}t/\varepsilon} f,$$

from which it follows that

(4.5)
$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\mathrm{e}^{\mu t/2} |z|^2 \right) + \mu \mathrm{e}^{\mu t/2} |z|^2 = 2 \mathrm{e}^{\mu t/2} \mathrm{Re} \left(\mathrm{e}^{-\mathrm{i}t/\varepsilon} \bar{z} f \right)$$

Integrating, we find

(4.6)
$$e^{\mu t/2} |z(t)| - |z(0)|^2 + \mu \int_0^t e^{\mu \tau/2} |z(\tau)|^2 d\tau = 2 \int_0^t e^{\mu \tau/2} \operatorname{Re}\left(e^{-i\tau/\varepsilon} \bar{z}f\right) d\tau$$
$$= 2\varepsilon \left[e^{\mu \tau/2} \operatorname{Re}\left(ie^{-it/\varepsilon} \bar{z}f\right)\right]_0^t - 2\varepsilon \int_0^t \operatorname{Re}\left[ie^{-i\tau/\varepsilon} \partial_\tau (e^{\mu \tau/2} \bar{z}f)\right] d\tau,$$

where the second equality is obtained by integration by parts. Since $\partial_t f$ is bounded independently of ε , the integral can be bounded using (4.4) and the integral on the left-hand side. This leaves us with

(4.7)
$$|z(t)|^2 \le e^{-\mu t/2} c_1(|f|) |z(0)|^2 + \frac{\varepsilon}{\mu} (1 - e^{-\mu t/2}) K(|f|, |\partial_t f|, \mu).$$

Most of the work in the proof below is devoted to handling the nonlinear term, where particular properties of the OPE come into play. A PDE application of this principle can be found in [29].

4.1. **Proof of Theorem 1.** In this proof, we omit the subscript in the inner product $(\cdot, \cdot)_{L^2}$ when the meaning is unambiguous; similarly, $|\cdot| \equiv |\cdot|_{L^2}$. We start by writing (3.16) as

(4.8)
$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t} |W^{\varepsilon}|^2 + 2\mu |\nabla W^{\varepsilon}|^2 \\ &= -2(W^{\varepsilon}, B(W^0, W^0)) - 2(W^{\varepsilon}, B(W^{\varepsilon}, W^0)) + 2(W^{\varepsilon}, f^{\varepsilon}) \\ &= -2(W^{\varepsilon}, B(W^0, W^0)) + 2(W^0, B(W^{\varepsilon}, W^{\varepsilon})) + 2(W^{\varepsilon}, f^{\varepsilon}). \end{aligned}$$

Using the Poincaré inequality, $|W^{\varepsilon}|^2 \leq c_p |\nabla W^{\varepsilon}|^2$, and multiplying the left-hand side by $2e^{\nu t}$ where $\nu := \mu c_p$, we have

(4.9)
$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\mathrm{e}^{\nu t} |W^{\varepsilon}|^{2} \right) + \mu \mathrm{e}^{\nu t} |\nabla W^{\varepsilon}|^{2} \leq \mathrm{e}^{\nu t} \left(\frac{\mathrm{d}}{\mathrm{d}t} |W^{\varepsilon}|^{2} + \mu |\nabla W^{\varepsilon}|^{2} + \mu |\nabla W^{\varepsilon}|^{2} \right).$$

With this, (4.8) becomes

(4.10)
$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\mathrm{e}^{\nu t} |W^{\varepsilon}|^{2} \right) + \mu \mathrm{e}^{\nu t} |\nabla W^{\varepsilon}|^{2}$$
$$\leq 2 \, \mathrm{e}^{\nu t} \left(W^{\varepsilon}, f^{\varepsilon} \right) - 2 \, \mathrm{e}^{\nu t} \left(W^{\varepsilon}, B(W^{0}, W^{0}) \right) + 2 \, \mathrm{e}^{\nu t} \left(W^{0}, B(W^{\varepsilon}, W^{\varepsilon}) \right).$$

We now integrate this inequality from 0 to t. On the left-hand side we have

(4.11)
$$\int_0^t \left\{ \frac{\mathrm{d}}{\mathrm{d}\tau} \left(\mathrm{e}^{\nu\tau} |W^{\varepsilon}|^2 \right) + \mu \mathrm{e}^{\nu\tau} |\nabla W^{\varepsilon}|^2 \right\} \mathrm{d}\tau \\ = \mathrm{e}^{\nu t} |W^{\varepsilon}(t)|^2 - |W^{\varepsilon}(0)|^2 + \mu \int_0^t \mathrm{e}^{\nu\tau} |\nabla W^{\varepsilon}|^2 \,\mathrm{d}\tau.$$

Using the expansion (3.26) of W^{ε} , we integrate the right-hand side by parts to bring out a factor of ε ; that is, we integrate the rapidly oscillating exponential $e^{i\omega_k^s t/\varepsilon}$ and leave everything else. For the force term, we have

(4.12)

$$\int_{0}^{t} e^{\nu\tau} (W^{\varepsilon}, f^{\varepsilon}) d\tau = |\mathcal{M}| \sum_{\boldsymbol{k}}^{s} \int_{0}^{t} e^{\nu\tau + \mathrm{i}\omega_{\boldsymbol{k}}^{s}\tau/\varepsilon} \overline{w_{\boldsymbol{k}}^{s}} f_{\boldsymbol{k}}^{s} d\tau$$

$$= \varepsilon |\mathcal{M}| \sum_{\boldsymbol{k}}^{s'} \frac{1}{\mathrm{i}\omega_{\boldsymbol{k}}^{s}} [\overline{w_{\boldsymbol{k}}^{s}(t)} f_{\boldsymbol{k}}^{s}(t) e^{\nu t + \mathrm{i}\omega_{\boldsymbol{k}}^{s}t/\varepsilon} - \overline{w_{\boldsymbol{k}}^{s}(0)} f_{\boldsymbol{k}}^{s}(0)]$$

$$- \varepsilon |\mathcal{M}| \int_{0}^{t} \sum_{\boldsymbol{k}}^{s'} \frac{\mathrm{e}^{\mathrm{i}\omega_{\boldsymbol{k}}^{s}\tau/\varepsilon}}{\mathrm{i}\omega_{\boldsymbol{k}}^{s}} \frac{\mathrm{d}}{\mathrm{d}\tau} (\overline{w_{\boldsymbol{k}}^{s}} f_{\boldsymbol{k}}^{s} e^{\nu\tau}) d\tau.$$

Here the prime on \sum' indicates that terms for which $\omega_{\mathbf{k}}^s = 0$ are omitted since then $w_{\mathbf{k}}^s = 0$. Introducing the integration operator I_{ω} defined by

(4.13)
$$\mathsf{I}_{\omega}W^{\varepsilon}(\boldsymbol{x},t) := \sum_{\boldsymbol{k}}^{s'} \frac{\mathrm{i}}{\omega_{\boldsymbol{k}}^{s}} w_{\boldsymbol{k}}^{s}(t) X_{\boldsymbol{k}}^{s} \mathrm{e}^{-\mathrm{i}\omega_{\boldsymbol{k}}^{s}t/\varepsilon} \mathrm{e}^{\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}},$$

which is well-defined since $|\omega_{\boldsymbol{k}}^{s}| \geq 1$, we can write this as

(4.14)
$$\int_{0}^{t} e^{\nu\tau} (W^{\varepsilon}, f^{\varepsilon}) d\tau = \varepsilon e^{\nu t} (\mathsf{I}_{\omega} W^{\varepsilon}(t), f^{\varepsilon}(t)) - \varepsilon (\mathsf{I}_{\omega} W^{\varepsilon}(0), f^{\varepsilon}(0)) - \varepsilon \int_{0}^{t} e^{\nu\tau} \{ \nu (\mathsf{I}_{\omega} W^{\varepsilon}, f^{\varepsilon}) + (\mathsf{I}_{\omega} \partial_{\tau}^{*} W^{\varepsilon}, f^{\varepsilon}) + (\mathsf{I}_{\omega} W^{\varepsilon}, \partial_{\tau} f^{\varepsilon}) \} d\tau.$$

Similarly, integrating the next term by parts we find

(4.15)

$$\int_{0}^{t} e^{\nu\tau} (W^{\varepsilon}, B(W^{0}, W^{0})) d\tau$$

$$= \varepsilon e^{\nu t} (\mathbf{I}_{\omega} W^{\varepsilon}, B(W^{0}, W^{0}))(t) - \varepsilon (\mathbf{I}_{\omega} W^{\varepsilon}, B(W^{0}, W^{0}))(0)$$

$$- \varepsilon \int_{0}^{\tau} e^{\nu\tau} \{ \nu (\mathbf{I}_{\omega} W^{\varepsilon}, B(W^{0}, W^{0})) + (\mathbf{I}_{\omega} \partial_{\tau}^{*} W^{\varepsilon}, B(W^{0}, W^{0}))$$

$$+ (\mathbf{I}_{\omega} W^{\varepsilon}, B(\partial_{\tau} W^{0}, W^{0})) + (\mathbf{I}_{\omega} W^{\varepsilon}, B(W^{0}, \partial_{\tau} W^{0})) \} d\tau$$

Next, we consider

$$\begin{aligned} \int_{0}^{t} \mathrm{e}^{\nu\tau} \left(W^{0}, B(W^{\varepsilon}, W^{\varepsilon}) \right) \, \mathrm{d}\tau \\ &= \int_{0}^{t} \frac{1}{2} \sum_{\boldsymbol{jkl}}^{rs} \mathrm{e}^{-\mathrm{i}(\omega_{\boldsymbol{j}}^{r} + \omega_{\boldsymbol{k}}^{s})\tau/\varepsilon} \left(B_{\boldsymbol{jkl}}^{rs0} + B_{\boldsymbol{kjl}}^{sr0} \right) w_{\boldsymbol{j}}^{r} w_{\boldsymbol{k}}^{s} \overline{w_{\boldsymbol{l}}^{0}} \, \mathrm{e}^{\nu\tau} \, \mathrm{d}\tau \\ (4.16) &= \frac{\varepsilon \mathrm{i}}{2} \sum_{\boldsymbol{jkl}}^{rs'} \frac{B_{\boldsymbol{jkl}}^{rs0} + B_{\boldsymbol{kjl}}^{sr0}}{\omega_{\boldsymbol{j}}^{r} + \omega_{\boldsymbol{k}}^{s}} \left[w_{\boldsymbol{j}}^{r}(t) w_{\boldsymbol{k}}^{s}(t) \overline{w_{\boldsymbol{l}}^{0}(t)} \mathrm{e}^{\nu t - \mathrm{i}(\omega_{\boldsymbol{j}}^{r} + \omega_{\boldsymbol{k}}^{s})t/\varepsilon} \right. \\ &- w_{\boldsymbol{j}}^{r}(0) w_{\boldsymbol{k}}^{s}(0) \overline{w_{\boldsymbol{l}}^{0}(0)} \right] \\ &- \frac{\varepsilon \mathrm{i}}{2} \int_{0}^{t} \sum_{\boldsymbol{jkl}}^{rs'} \frac{B_{\boldsymbol{jkl}}^{rs0} + B_{\boldsymbol{kjl}}^{sr0}}{\omega_{\boldsymbol{j}}^{r} + \omega_{\boldsymbol{k}}^{s}} \, \mathrm{e}^{-\mathrm{i}(\omega_{\boldsymbol{j}}^{r} + \omega_{\boldsymbol{k}}^{s})\tau/\varepsilon} \frac{\mathrm{d}}{\mathrm{d}\tau} \left[w_{\boldsymbol{j}}^{r} w_{\boldsymbol{k}}^{s} \overline{w_{\boldsymbol{l}}^{0}} \mathrm{e}^{\nu\tau} \right] \, \mathrm{d}\tau. \end{aligned}$$

Here the prime on \sum' indicates that exactly resonant terms, for which $\omega_{j}^{r} + \omega_{k}^{s} = 0$ and $B_{jkl}^{rs0} + B_{kjl}^{sr0} = 0$, are excluded. Using the bilinear operator B_{ω} , defined for any W^{ε} , \hat{W}^{ε} and \tilde{W}^{0} by

$$(4.17) \quad (\tilde{W}^0, B_{\omega}(W^{\varepsilon}, \hat{W}^{\varepsilon})) := \frac{\mathrm{i}}{2} \sum_{\boldsymbol{jkl}}^{rs} \frac{B_{\boldsymbol{jkl}}^{rs0} + B_{\boldsymbol{kjl}}^{sr0}}{\omega_{\boldsymbol{j}}^r + \omega_{\boldsymbol{l}}^s} w_{\boldsymbol{j}}^r \hat{w}_{\boldsymbol{k}}^s \overline{w}_{\boldsymbol{l}}^0 \mathrm{e}^{-\mathrm{i}(\omega_{\boldsymbol{j}}^r + \omega_{\boldsymbol{k}}^s)t/\varepsilon},$$

we can write (4.16) in the more compact form

(4.18)

$$\int_{0}^{t} e^{\nu\tau} (W^{0}, B(W^{\varepsilon}, W^{\varepsilon})) d\tau$$

$$= \varepsilon e^{\nu t} (W^{0}, B_{\omega}(W^{\varepsilon}, W^{\varepsilon}))(t) - \varepsilon (W^{0}, B_{\omega}(W^{\varepsilon}, W^{\varepsilon}))(0)$$

$$- \varepsilon \int_{0}^{t} e^{\nu\tau} \left\{ \nu (W^{0}, B_{\omega}(W^{\varepsilon}, W^{\varepsilon})) + (\partial_{\tau} W^{0}, B_{\omega}(W^{\varepsilon}, W^{\varepsilon})) + (W^{0}, \partial_{\tau}^{*} B_{\omega}(W^{\varepsilon}, W^{\varepsilon})) + (W^{0}, \partial_{\tau}^{*} B_{\omega}(W^{\varepsilon}, W^{\varepsilon})) \right\} d\tau$$

Putting these together, (4.10) integrates to

$$e^{\nu t} |W^{\varepsilon}(t)|^{2} - |W^{\varepsilon}(0)|^{2} + \mu \int_{0}^{t} e^{\nu \tau} |\nabla W^{\varepsilon}|^{2} d\tau$$

$$\leq 2\varepsilon e^{\nu t} \left(\mathbf{I}_{\omega} W^{\varepsilon}, f^{\varepsilon} \right)(t) - 2\varepsilon \left(\mathbf{I}_{\omega} W^{\varepsilon}, f^{\varepsilon} \right)(0)$$

$$(4.19) \qquad - 2\varepsilon e^{\nu t} \left(\mathbf{I}_{\omega} W^{\varepsilon}, B(W^{0}, W^{0}) \right)(t) + 2\varepsilon \left(\mathbf{I}_{\omega} W^{\varepsilon}, B(W^{0}, W^{0}) \right)(0)$$

$$+ 2\varepsilon e^{\nu t} \left(W^{0}, B_{\omega} (W^{\varepsilon}, W^{\varepsilon}) \right)(t) - 2\varepsilon \left(W^{0}, B_{\omega} (W^{\varepsilon}, W^{\varepsilon}) \right)(0)$$

$$+ 2\varepsilon \int_{0}^{t} e^{\nu \tau} \left\{ I_{0}(\tau) - I_{1}(\tau) + I_{2}(\tau) \right\} d\tau.$$

Here the integrands are

$$(4.20) \quad I_0 := \nu(\mathsf{I}_{\omega}W^{\varepsilon}, f^{\varepsilon}) + (\mathsf{I}_{\omega}W^{\varepsilon}, \partial_{\tau}f^{\varepsilon}) + (\mathsf{I}_{\omega}\partial_{\tau}^*W^{\varepsilon}, f^{\varepsilon}),$$

$$(4.21) \quad I_1 := \nu(\mathsf{I}_{\omega}W^{\varepsilon}, B(W^0, W^0)) + (\mathsf{I}_{\omega}\partial_{\tau}^*W^{\varepsilon}, B(W^0, W^0)) + (\mathsf{I}_{\omega}W^{\varepsilon}, B(W^0, \partial_{\tau}W^0)) + (\mathsf{I}_{\omega}W^{\varepsilon}, B(W^0, \partial_{\tau}W^0)),$$

14

and

(4.22)
$$I_2 := \nu \left(W^0, B_\omega(W^\varepsilon, W^\varepsilon) \right) + \left(\partial_\tau W^0, B_\omega(W^\varepsilon, W^\varepsilon) \right) \\ + \left(W^0, \partial_\tau^* B_\omega(W^\varepsilon, W^\varepsilon) \right).$$

We now bound the right-hand side of (4.19). On the second line, we have

(4.23)
$$\begin{aligned} \left| \mathrm{e}^{\nu t} \left(\mathsf{I}_{\omega} W^{\varepsilon}(t), f^{\varepsilon}(t) \right) - \left(\mathsf{I}_{\omega} W^{\varepsilon}(0), f^{\varepsilon}(0) \right) \right| \\ &\leq \mathrm{e}^{\nu t} \left| W^{\varepsilon}(t) \right| \left| f^{\varepsilon}(t) \right| + \left| W^{\varepsilon}(0) \right| \left| f^{\varepsilon}(0) \right|, \end{aligned}$$

where we have used the fact that, thanks to (3.25),

(4.24)
$$|\nabla^{\alpha}|_{\omega}W^{\varepsilon}| \leq |\nabla^{\alpha}W^{\varepsilon}|, \quad \text{for } \alpha = 0, 1, 2, \cdots$$

To bound the next line, we use the estimate

(4.25)
$$\begin{aligned} |(\tilde{W}, B(W^{0}, \tilde{W}))| &\leq C \, |\tilde{W}|_{L^{6}} \, |W^{0}|_{L^{3}} \, |\nabla \tilde{W}|_{L^{2}} \\ &\leq C \, |\nabla \tilde{W}| \, |W^{0}|^{1/2} \, |\nabla W^{0}|^{1/2} \, |\nabla \hat{W}| \end{aligned}$$

(note that the first argument of B is W^0) to obtain

(4.26)
$$\begin{aligned} \left| e^{\nu t} \left(\mathsf{I}_{\omega} W^{\varepsilon}, B(W^{0}, W^{0}) \right)(t) - \left(\mathsf{I}_{\omega} W^{\varepsilon}, B(W^{0}, W^{0})(0) \right) \\ & \leq e^{\nu t} \left| \nabla W^{\varepsilon}(t) \right| |W^{0}(t)|^{1/2} |\nabla W^{0}(t)|^{3/2} \\ & + \left| \nabla W^{\varepsilon}(0) \right| |W^{0}(0)|^{1/2} |\nabla W^{0}(0)|^{3/2} \end{aligned}$$

In (4.25) and in the rest of this proof, C and c denote generic constants which may not be the same each time the symbol is used; such constants may depend on \mathcal{M} but not on any other parameter. Numbered constants may also depend on μ .

but not on any other parameter. Numbered constants may also depend on μ . We now derive a bound involving B_{ω} . Since $B_{jkl}^{rs0} + B_{kjl}^{sr0} = 0$ in the case of exact resonance, we assume that $\omega_j^r + \omega_k^s \neq 0$. Then (3.51) implies

(4.27)
$$\frac{\left|B_{\boldsymbol{j}\boldsymbol{k}\boldsymbol{l}}^{rs0} + B_{\boldsymbol{j}\boldsymbol{j}\boldsymbol{l}}^{sr0}\right|}{\left|\omega_{\boldsymbol{j}}^{r} + \omega_{\boldsymbol{k}}^{s}\right|} \le C \left|\boldsymbol{j}\right| \left|\boldsymbol{k}\right|.$$

With this, we have for any W^{ε} , \hat{W}^{ε} and \tilde{W}^{0} ,

$$(4.28) \qquad \left| (\tilde{W}^{0}, B_{\omega}(W^{\varepsilon}, \hat{W}^{\varepsilon})) \right| \leq \frac{1}{2} \sum_{\boldsymbol{j}\boldsymbol{k}\boldsymbol{l}}^{rs} \left| \frac{B_{\boldsymbol{j}\boldsymbol{k}\boldsymbol{l}}^{rs} + B_{\boldsymbol{k}\boldsymbol{j}\boldsymbol{l}}^{s}}{\omega_{\boldsymbol{j}}^{r} + \omega_{\boldsymbol{k}}^{s}} \right| |w_{\boldsymbol{j}}^{r}| |\hat{w}_{\boldsymbol{k}}^{s}| |\tilde{w}_{\boldsymbol{l}}^{0}| \leq C \sum_{\boldsymbol{j}+\boldsymbol{k}=\boldsymbol{l}}^{rs} |\boldsymbol{j}| |\boldsymbol{k}| |w_{\boldsymbol{j}}^{r}| |\hat{w}_{\boldsymbol{k}}^{s}| |\tilde{w}_{\boldsymbol{l}}^{0}| \leq \int_{\mathcal{M}} \theta(\boldsymbol{x}) \,\xi(\boldsymbol{x}) \,\zeta(\boldsymbol{x}) \,\mathrm{d}\boldsymbol{x}^{3} \leq C |\nabla W^{\varepsilon}|_{L^{p}} |\nabla \hat{W}^{\varepsilon}|_{L^{q}} |\tilde{W}^{0}|_{L^{m}},$$

with 1/p + 1/q + 1/m = 1 and where on the penultimate line

(4.29)
$$\theta(\boldsymbol{x}) := \sum_{\boldsymbol{j}}^{r} |\boldsymbol{j}| |w_{\boldsymbol{j}}^{r}| e^{i\boldsymbol{j}\cdot\boldsymbol{x}}, \xi(\boldsymbol{x}) := \sum_{\boldsymbol{k}}^{s} |\boldsymbol{k}| |\hat{w}_{\boldsymbol{k}}^{s}| e^{i\boldsymbol{k}\cdot\boldsymbol{x}} \text{ and } \zeta(\boldsymbol{x}) := \sum_{\boldsymbol{l}} |\tilde{w}_{\boldsymbol{l}}^{0}| e^{i\boldsymbol{l}\cdot\boldsymbol{x}}.$$

Using (4.28) with p = q = 2 and $m = \infty$, plus the embedding $H^2 \subset L^{\infty}$, we have the bound

(4.30)
$$\frac{\left| \mathrm{e}^{\nu t} \left(W^{0}, B_{\omega}(W^{\varepsilon}, W^{\varepsilon}) \right)(t) - \left(W^{0}, B_{\omega}(W^{\varepsilon}, W^{\varepsilon}) \right)(0) \right|}{\leq C \left(\mathrm{e}^{\nu t} |\nabla W^{\varepsilon}(t)|^{2} |\nabla^{2} W^{0}(t)| + |\nabla W^{\varepsilon}(0)|^{2} |\nabla^{2} W^{0}(0)| \right).$$

To bound the integrand in (4.19), we need estimates on $\partial_t W^0$ and $\partial_t^* W^{\varepsilon}$ in addition to those already obtained. Using the bound

 $\begin{array}{ll} (4.31) & |B^0(W,W)|_{L^2} \leq C \, |\nabla W|_{L^4}^2 \leq C \, |\nabla W|_{H^{3/4}}^2 \leq C \, |\nabla^2 W|^{3/2} |\nabla W|^{1/2}, \\ \text{we find from (3.9)} \end{array}$

(4.32)
$$\begin{aligned} |\partial_t W^0|_{L^2} &\leq C \, |\nabla W|^2_{H^{3/4}} + \mu \, |\nabla^2 W^0| + |f| \\ &\leq C \, |\nabla^2 W|^{3/2} |\nabla W|^{1/2} + \mu \, |\nabla^2 W^0| + |f|. \end{aligned}$$

Similarly, we find from (3.34)

(4.33)
$$\begin{aligned} |\partial_t^* W^{\varepsilon}|_{L^2} &\leq C \, |\nabla W|_{H^{3/4}}^2 + \mu \, |\nabla^2 W^{\varepsilon}| + |f| \\ &\leq C \, |\nabla^2 W|^{3/2} |\nabla W|^{1/2} + \mu \, |\nabla^2 W^{\varepsilon}| + |f|. \end{aligned}$$

Now using the bound

 $\begin{aligned} (4.34) \quad |\nabla B(W,W)|_{L^2} \leq C \, |\nabla^2 W|_{L^{12/5}} |\nabla W|_{L^{12}} \leq C \, |\nabla^2 W|_{H^{1/4}} \\ \text{we find} \end{aligned}$

$$(4.35) \begin{aligned} |\nabla \partial_t W^0|_{L^2} &\leq C \, |\nabla^2 W|^2_{H^{1/4}} + \mu \, |\nabla^3 W^0| + |\nabla f^0| \\ &\leq C \, |\nabla^3 W|^{1/2} |\nabla^2 W|^{3/2} + \mu \, |\nabla^3 W^0| + |\nabla f^0|, \\ |\nabla \partial_t^* W^{\varepsilon}|_{L^2} &\leq C \, |\nabla^2 W|^2_{H^{1/4}} + \mu \, |\nabla^3 W^{\varepsilon}| + |\nabla f^{\varepsilon}| \\ &\leq C \, |\nabla^3 W|^{1/2} |\nabla^2 W|^{3/2} + \mu \, |\nabla^3 W^{\varepsilon}| + |\nabla f^{\varepsilon}|. \end{aligned}$$

The bound for I_0 follows by using (4.33),

(4.36)
$$\begin{aligned} |I_0|_{L^2} &\leq C \left(|\nabla W|^2_{H^{3/4}} + (\mu + c) |\nabla^2 W^{\varepsilon}| + |f^{\varepsilon}| \right) \left(|f^{\varepsilon}| + |\partial_t f^{\varepsilon}| \right) \\ &\leq C \left(|\nabla W|^2_{H^{3/4}} + (\mu + c) |\nabla^2 W| + ||f||_g \right) ||f||_g, \end{aligned}$$

where we have used the fact that $|\nabla^{\alpha}W^{\varepsilon}|^2 \leq |\nabla^{\alpha}W^{\varepsilon}|^2 + |\nabla^{\alpha}W^0|^2 = |\nabla^{\alpha}W|^2$. Next, using (4.28) we bound I_2 as

$$(4.37) \begin{aligned} |I_2|_{L^2} &\leq \mu c \, |W^0|_{L^{\infty}} |\nabla W^{\varepsilon}|^2 + c \, |\partial_{\tau} W^0| \, |\nabla W^{\varepsilon}|_{L^4}^2 \\ &+ c \, |W^0|_{L^{\infty}} |\nabla W^{\varepsilon}| \, |\nabla \partial_t^* W^{\varepsilon}| \\ &\leq \mu c \, |\nabla^2 W^0| \, |\nabla W^{\varepsilon}|^2 + c \left(|\nabla W|_{H^{3/4}}^2 + \mu \, |\nabla^2 W^0| + |f^0| \right) |\nabla W^{\varepsilon}|_{H^{3/4}}^2 \\ &+ c \, |\nabla^2 W^0| \, |\nabla W^{\varepsilon}| \left(|\nabla^2 W|_{H^{1/4}}^2 + \mu \, |\nabla^3 W^{\varepsilon}| + |\nabla f^{\varepsilon}| \right) \\ &\leq c \, |\nabla W| \, |\nabla^2 W| \, |\nabla^2 W|_{H^{1/4}}^2 + \mu c \, |\nabla^3 W| \, |\nabla^2 W| \, |\nabla W| \\ &+ |\nabla^2 W|^{3/2} |\nabla W|^{1/2} \, \|f\|_g \end{aligned}$$

where interpolation inequalities have been used for the last step. The bound for I_1 is majorised by that for I_2 .

Putting everything together, we have from (4.19)

$$(4.38) e^{\nu t} |W^{\varepsilon}(t)|^{2} - |W^{\varepsilon}(0)|^{2} \\ \leq \varepsilon c_{2} e^{\nu t} |\nabla^{2}W(t)| (|\nabla W(t)|^{2} + ||f||_{g}) + \varepsilon c_{2} |\nabla^{2}W_{0}| (|\nabla W_{0}|^{2} + ||f||_{g}) \\ + \varepsilon c_{3} \int_{0}^{t} e^{\nu \tau} \{ |W|_{H^{1}} |W|_{H^{2}} |W|_{H^{9/4}}^{2} + \mu |W|_{H^{3}} |W|_{H^{2}} |W|_{H^{1}} \\ + (|W|_{H^{2}}^{3/2} |W|_{H^{1}}^{1/2} + (\mu + c) |W|_{H^{2}} + ||f||_{g}) ||f||_{g} \} d\tau$$

Now by (2.18) and (2.19), we can find $K_*(||f||_g)$ and $T_*(|\nabla W_0|, ||f||_g)$ such that, for $t \ge T_*$,

(4.39)
$$c |\nabla^s W(t)|^2 + (\mu + c') |\nabla^s W(t)| + ||f||_g \leq K,$$

for $s \in \{0, 1, 2, 3\}$. Let $t' := t - T_*$ and relabel t in (4.38) as t'. We can then bound the integral in (4.38) as

(4.40)
$$\int_0^{t'} \mathrm{e}^{\nu \tau} \{\cdots\} \mathrm{d}\tau \le \frac{\mathrm{e}^{\nu t'} - 1}{\nu} c_4 K_* (\|f\|_{\mathrm{g}})^2.$$

Bounding the remaining terms in (4.38) similarly, we find

(4.41)
$$|W^{\varepsilon}(t)|^{2} \leq e^{-\nu(t-T_{*})} |W^{\varepsilon}(T_{*})|^{2} + \varepsilon c_{5} \left(K_{*}^{2} + K_{*}^{3/2}\right) \\ \leq e^{-\nu(t-T_{*})} |W(T_{*})|^{2} + \varepsilon c_{5} \left(K_{*}^{2} + K_{*}^{3/2}\right).$$

This proves the theorem, with $K_{\rm g}(\|f\|_{\rm g})^2 = 2 c_5 \left(K_*^2 + K_*^{3/2}\right)$ and $T_{\rm g}(|\nabla W_0|, \|f\|_{\rm g}, \varepsilon) = T_* - \log \left[\varepsilon c_5 \left(K_* + K_*^{1/2}\right)\right]/\nu$.

5. Higher-Order Estimates

When $\partial_t f = 0$ in the very simple model (4.3), we can obtain a better estimate on x' = x - U where $U = \varepsilon f/(\varepsilon \mu + i)$ than on x, namely that $x'(t) \to 0$ as $t \to \infty$; here U is the (exact, higher-order) slow manifold. The situation is more complicated when f is time-dependent, or when x is coupled to a slow variable ywith the evolution equations having nonlinear terms. In this case, it is not generally possible to find U (explicit examples are known where no such U exists), and thus $x'(t) \not\to 0$ as $t \to \infty$ for any $U(y, f; \varepsilon)$. Nevertheless, it is often possible to find a U^* that gives an exponentially small bound on x'(t) for large t. We shall do this for the primitive equations.

More concretely, in this section we show that, with reasonable regularity assumptions on the forcing f, the leading-order estimate on the fast variable W^{ε} in the previous section can be sharpened to an exponential-order estimate on $W^{\varepsilon} - U^*(W^0, f; \varepsilon)$, where U^* is computed below. As in [31], we make use of the Gevrey regularity of the solution and work with a finite-dimensional truncation of the system, whose description now follows.

Given a fixed $\kappa > 0$, we define the low-mode truncation of W by

(5.1)
$$W^{<}(\boldsymbol{x},t) = (\mathsf{P}^{<}W)(\boldsymbol{x},t) := \sum_{|\boldsymbol{k}| < \kappa}^{\alpha} w_{\boldsymbol{k}}^{\alpha} X_{\boldsymbol{k}}^{\alpha} \mathrm{e}^{-\mathrm{i}\omega_{\boldsymbol{k}}^{\alpha}t/\varepsilon} \mathrm{e}^{\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}}$$

where the sum is taken over $\alpha \in \{0, \pm 1\}$ and $\mathbf{k} \in \mathbb{Z}_L$ with $|\mathbf{k}| < \kappa$. We also define the high-mode part of W by $W^> := W - W^<$. The low- and high-mode parts of the slow and fast variables, $W^{0<}$, $W^{0>}$, $W^{\varepsilon<}$ and $W^{\varepsilon>}$, are defined in the obvious manner, i.e. $W^{0<}$ with $\alpha = 0$ in (5.1) and $W^{\varepsilon<}$ with $\alpha \in \{\pm 1\}$. It is clear from (5.1) and (3.18) that the projection $\mathsf{P}^<$ is orthogonal in H^s , so $\mathsf{P}^<$ commutes with both A and L in (2.11). We denote $\mathsf{P}^< B$ by $B^<$.

It follows from the definition that the low-mode part $W^{<}$ satisfies a "reverse Poincaré" inequality, i.e. for any $s \ge 0$,

(5.2) $|\nabla W^{<}|_{H^{s}} \leq \kappa |W^{<}|_{H^{s}}.$

If $W \in G^{\sigma}(\mathcal{M})$, the exponential decay of its Fourier coefficients implies that $W^{>}$ is exponentially small, that is, for any $s \geq 0$,

(5.3)
$$|W^{>}|_{H^s} \leq C_s \,\kappa^s \,\mathrm{e}^{-\sigma\kappa} |W|_{G^{\sigma}}$$
.

The first inequality evidently also applies to the slow and fast parts separately, i.e. with $W^{<}$ replaced by $W^{0<}$ or $W^{\varepsilon<}$; as for (5.3), it also holds when $W^{>}$ on the lhs is replaced by $W^{0>}$ or $W^{\varepsilon>}$.

We recall that the global regularity results of Theorem 0 imply that, with Gevrey forcing, any solution $W \in H^1(\mathcal{M})$ will be in $G^{\sigma}(\mathcal{M})$ after a short time. As in [31] and following [22], the central idea here is to split W^{ε} into its low- and high-mode parts. The high-mode part $W^{\varepsilon>}$ is exponentially small by (5.3). We then compute $U^*(W^{0<}, f^{<}; \varepsilon)$ such that $W^{\varepsilon<} - U^*$ becomes exponentially small after some time.

Following historical precedent in the geophysical literature, it is natural to present our results in two parts, first locally in time and second globally. (Here "local in time" is used in a sense similar to "local truncation error" in numerical analysis, giving a bound on the time derivative of some "error".) The following lemma states that, in a suitable finite-dimensional space, we can find a "slow manifold" $W^{\varepsilon <} = U^*(W^{0<}, f^{<}; \varepsilon)$ on which the normal velocity of $W^{\varepsilon <}$ is at most exponentially small:

Lemma 2. Let s > 3/2 and $\eta > 0$ be fixed. Given $W^0 \in H^s(\mathcal{M})$ and $f \in H^s(\mathcal{M})$ with $\partial_t f = 0$, there exists $\varepsilon_{**}(|W^0|_{H^s}, |f|_{H^s}, \eta)$ such that for $\varepsilon \leq \varepsilon_{**}$ one can find $\kappa(\varepsilon)$ and $U^*(W^{0<}, f^{<}; \varepsilon)$ that makes the remainder function

(5.4)
$$\mathcal{R}^{*}(W^{0<}, f^{<}; \varepsilon) := \mathsf{P}^{<}[(\mathsf{D}U^{*})\mathcal{G}^{*}] + \frac{1}{\varepsilon}LU^{*} + B^{\varepsilon<}(W^{0<} + U^{*}, W^{0<} + U^{*}) + AU^{*} - f^{\varepsilon<}$$

exponentially small in ε ,

(5.5)
$$|\mathcal{R}^*(W^{0<}, f^{<}; \varepsilon)|_{H^s} \le c_r \left[(|W^{0<}|_{H^s} + \eta)^2 + |f|_{H^s} \right] \exp(-\eta/\varepsilon^{1/4});$$

here $\mathsf{D}U^*$ is the derivative of U^* with respect to $W^{0<}$ and

(5.6)
$$\mathcal{G}^* := -B^{0<}(W^{0<} + U^*, W^{0<} + U^*) - AW^{0<} + f^{0<}.$$

Remarks.

1. The bounds may depend on s, μ and \mathcal{M} as well as on η , but only the latter is indicated explicitly here and in the proof below.

2. Given κ fixed, U^* lives in the same space as $W^{\varepsilon <}$, that is, $(W^0, U^*)_{L^2} = 0$ and $\mathsf{P}^{\leq} U^* = U^*$.

3. In the leading-order case of §4, the slow manifold is $U^0 = 0$ and the local error estimate is incorporated directly into the proof of Theorem 1; we therefore did not put these into a separate lemma.

4. Unlike formal constructions in the geophysical literature (see, e.g., [5, 37]), our slow manifold is not defined for all possible W^0 and ε . Instead, given that $|W^0|_{G^{\sigma}} \leq R$, we can define U^* for all $\varepsilon \leq \varepsilon_{**}(R, \sigma)$; generally, the larger the set of W^0 over which U^* is to be defined, the smaller ε will have to be.

5. In what follows, we will often write $U^*(W^0, f; \varepsilon)$ for $U^*(\mathsf{P}^{\leq}W^0, \mathsf{P}^{\leq}f; \varepsilon)$; this should not cause any confusion.

Using the Lemma and a technique similar to that used to prove Theorem 1, we can bound the "net forcing" on $W' = W^{\varepsilon <} - U^*$ by \mathcal{R}^* . The dissipation term AW' then ensures that W' eventually decays to an exponentially small size. This gives us our global result:

Theorem 2. Let $W_0 \in H^1(\mathcal{M})$ and $\nabla f \in G^{\sigma}(\mathcal{M})$ be given with $\partial_t f = 0$. Then there exist $\varepsilon_*(f;\sigma)$ and $T_*(|\nabla W_0|, |\nabla f|_{G^{\sigma}})$ such that for $\varepsilon \leq \varepsilon_*$ and $t \geq T_*$, we can approximate the fast variable $W^{\varepsilon}(t)$ by a function $U^*(W^0(t), f; \varepsilon)$ of the slow variable $W^0(t)$ up to an exponential accuracy,

(5.7)
$$|W^{\varepsilon}(t) - U^*(W^0(t), f; \varepsilon)|_{L^2} \le K_*(|\nabla f|_{G^{\sigma}}, \sigma) \exp(-\sigma/\varepsilon^{1/4}).$$

As in Theorem 1, here K_* is a continuous increasing function of its arguments; $W(t) = W^0(t) + W^{\varepsilon}(t)$ is the solution of (2.11) with initial condition $W(0) = W_0$. As before, the bounds depend on μ and \mathcal{M} , but these are not indicated explicitly.

Remarks.

6. With very minor changes in the proof of Theorem 2 below, one could also show that, if $f \in H^{n+1}$ and $\partial_t f = 0$, then $|W^{\varepsilon}(t) - U^n(W^0(t), f; \varepsilon)|_{L^2}$ is bounded as $\varepsilon^{n/4}$ for sufficiently large n and possibly something better for smaller n.

7. Recalling remark 4 above, our slow manifold is only defined for ε sufficiently small for a given $|W^{0<}|$ (or equivalently, for $|W^{0<}|$ sufficiently small for a given ε). The results of Theorem 0 tell us that W(t) will be inside a ball in $G^{\sigma}(\mathcal{M})$ after a sufficiently large t; we use (twice) the radius of this absorbing ball to fix the restriction on ε . Thus our approach sheds no light on the analogous problem in the inviscid case, which has no absorbing set.

8. As proved in [12, 13, 23], assuming sufficiently smooth forcing, the primitive equations admit a finite-dimensional global attactor. Theorem 2 states that, for $\varepsilon \leq \varepsilon_*(|f|_{G^{\sigma}})$, the solution will enter, and remain in, an exponentially thin neighbourhood of $U^*(W^{0<}, f^{<}; \varepsilon)$ in $L^2(\mathcal{M})$ after some time. It follows that the global attractor must then be contained in this exponentially thin neighbourhood as well.

9. The dynamics on this attractor is generally thought to be chaotic [30]. Thus our present results do not qualitatively affect the finite-time predictability estimate of [31].

10. When $\partial_t f \neq 0$, the slaving relation U^* would have a non-local dependence on t. Quasi-periodic forcing, however, can be handled by introducing an auxiliary variable $\boldsymbol{\theta} = (\theta_1, \dots, \theta_n)$, where n is the number of independent frequencies of f. The slaving relation U^* would then depend on $\boldsymbol{\theta}$ as well as on $W^{0<}$.

11. Bounds of this type are only available for the fast variable W^{ε} ; no special bounds exist for the slow variable W^{0} except in special cases, such as when the forcing f is completely fast, $(W^{0}, f)_{L^{2}} = 0$.

We next present the proofs of Lemma 2 and Theorem 2. The first one follows closely that in [31] which used a slightly different notation; we redo it here for notational coherence and since some estimates in it are needed in the proof of Theorem 2. As before, we write $(\cdot, \cdot) \equiv (\cdot, \cdot)_{L^2}$ and $|\cdot| \equiv |\cdot|_{L^2}$ when there is no ambiguity.

5.1. **Proof of Lemma 2.** As usual, we use c to denote a generic constant which may not be the same each time it appears. Constants may depend on s and the domain \mathcal{M} (and also on μ for non-generic ones), but dependence on η is indicated explicitly. Since s > 3/2, $H^s(\mathcal{M})$ is a Banach algebra, so if u and $v \in H^s$,

 $(5.8) \quad |uv|_s \le c \, |u|_s |v|_s$

where here and henceforth $|\cdot|_s := |\cdot|_{H^s}$. Let us take $\varepsilon \leq 1$ and κ as given for now; restrictions on ε will be stated as we go along and κ will be fixed in (5.22) below.

We construct the function U^* iteratively as follows. First, let

(5.9)
$$\frac{1}{\varepsilon}LU^1 = -B^{\varepsilon <}(W^{0<}, W^{0<}) + f^{\varepsilon <},$$

where $U^1 \in \text{range } L$ for uniqueness; similarly, $U^n \in \text{range } L$ in what follows. For $n = 1, 2, \cdots$, let

(5.10)
$$\frac{1}{\varepsilon}LU^{n+1} = -\mathsf{P}^{<}[(\mathsf{D}U^{n})\mathcal{G}^{n}] - B^{\varepsilon<}(W^{0<} + U^{n}, W^{0<} + U^{n}) - AU^{n} + f^{\varepsilon<},$$

where $\mathsf{D}U^n$ is the Fréchet derivative of U^n with respect to $W^{0<}$ (regarded as living in an appropriate Hilbert space) and

(5.11)
$$\mathcal{G}^n := -B^{0<}(W^{0<} + U^n, W^{0<} + U^n) - AW^{0<} + f^{0<}.$$

We note that the right-hand sides of (5.9) and (5.10) do not lie in ker L, so U^1 and U^{n+1} are well defined. Moreover, U^n lives in the same space as $W^{\varepsilon <}$, that is, $U^n \in \mathsf{P}^{\leq}$ range L; in other words, $(W^0, U^n) = 0$ and $\mathsf{P}^{\leq} U^n = U^n$.

For $\eta > 0$, let $D_{\eta}(W^{0<})$ be the complex η -neighbourhood of $W^{0<}$ in $\mathsf{P}^{\leq}H^{s}(\mathcal{M})$. With $W^{0<}$ defined by (5.1), this is

(5.12)
$$D_{\eta}(W^{0}) = \left\{ \hat{W}^{0} : \hat{W}^{0}(\boldsymbol{x}, t) = \sum_{|\boldsymbol{k}| < \kappa} \hat{w}_{\boldsymbol{k}}^{0} X_{\boldsymbol{k}}^{0} \mathrm{e}^{\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}} \text{ with} \\ \hat{w}_{(k_{1},k_{2},k_{3})}^{0} = \hat{w}_{(k_{1},k_{2},-k_{3})}^{0} \text{ and } \sum_{|\boldsymbol{k}| < \kappa} |\boldsymbol{k}|^{2s} |\hat{w}_{\boldsymbol{k}}^{0} - w_{\boldsymbol{k}}^{0}|^{2} < \eta^{2} \right\}$$

Since $W^0(\boldsymbol{x},t)$ and $X^0_{\boldsymbol{k}}$ are real, $w^0_{\boldsymbol{k}}$ must satisfy (3.29a), but $\hat{w}^0_{\boldsymbol{k}}$ in (5.12) need not satisfy this condition although it must satisfy (3.29b). We can thus regard $D_{\eta}(W^{0<}) \subset \{(w_{\boldsymbol{k}}): 0 < |\boldsymbol{k}| < \kappa \text{ and } w_{(k_1,k_2,-k_3)} = w_{k_1,k_2,k_3}\} \cong \mathbb{C}^m$ for some m. Let $\delta > 0$ be given; it will be fixed below in (5.22). For any function g of $W^{0<}$, let $(5.13) |g(W^{0<})|_{s;n} := \sup_{W \in D_{\eta-n\delta}(W^{0<})} |g(W)|_s$;

this expression is meaningful when $D_{\eta-n\delta}(W^{0<})$ is non-empty, that is, for $n \in \{0, \cdots, \lfloor \eta/\delta \rfloor =: n_*\}$. For future reference, we note that (5.14) $|W^{0<}|_{s:0} \leq |W^{0<}|_s + \eta$.

Our first step is to obtain by induction a couple of uniform bounds (5.25)–(5.26), valid for $n \in \{1, \dots, n_*\}$, which will be useful later. First, for U^1 , we have

(5.15)
$$\frac{1}{\varepsilon} |LU^1|_{s;1} \le |B^{\varepsilon <}(W^{0<}, W^{0<})|_{s;1} + |f^{\varepsilon <}|_s$$

which, using the estimate $|B(W,W)|_s \leq c |\nabla W|_s^2$ and (5.2), implies (5.16) $|U^1|_{s,1} \leq \varepsilon c_0 \left(\kappa^2 |W^{0<}|_{s,1}^2 + |f^{\varepsilon <}|_s\right).$ Next, we derive an iterative estimate for $|U^n|_{s;n}$. Using the fact that $|\cdot|_{s;m} \leq |\cdot|_{s;n}$ whenever $m \geq n$, we have for $n = 1, 2, \cdots$,

(5.17)
$$\frac{1}{\varepsilon} |U^{n+1}|_{s;n+1} \le |(\mathsf{D}U^n)\mathcal{G}^n|_{s;n+1} + |B^{\varepsilon <}(W^{0<} + U^n, W^{0<} + U^n)|_{s;n} + \mu\kappa^2 |W^{0<}|_{s;n} + |f^{\varepsilon <}|_s.$$

The first term on the right-hand side can be bounded by a technique based on Cauchy's integral formula: Let $D_{\eta}(z_0) \subset \mathbb{C}$ be the complex η -neighbourhood of z_0 . For $\varphi : D_{\eta}(z_0) \to \mathbb{C}$ analytic and $\delta \in (0, \eta)$, we can bound $|\varphi'|$ in $D_{\eta-\delta}(z_0)$ by $|\varphi|$ in $D_{\eta}(z_0)$ as

(5.18)
$$|\varphi' \cdot z|_{D_{\eta-\delta}(z_0)} \leq \frac{1}{\delta} |\varphi|_{D_{\eta}(z_0)} |z|_{\mathbb{C}}.$$

Now by (5.9) U^1 is an analytic function of the finite-dimensional variable $W^{0<}$, so assuming that U^n is analytic in $W^{0<}$ we can regard the Fréchet derivative $\mathsf{D}U^n$ as an ordinary derivative. Taking for φ' in (5.18) the derivative of U^n in the direction \mathcal{G}^n (i.e. working on the complex plane containing 0 and \mathcal{G}^n), we have

(5.19)
$$|(\mathsf{D}U^n)\mathcal{G}^n|_{s;n+1} \leq \frac{1}{\delta} |U^n|_{s;n} |\mathcal{G}^n|_{s;n}$$

Using the estimate

$$(5.20) |B^{\varepsilon<}(W^{0<} + U^n, W^{0<} + U^n)|_{s;n} \le c |\nabla(W^{0<} + U^n)|_{s;n}^2 \le c \,\kappa^2 |W^{0<} + U^n|_{s;r}^2$$
 we have

(5.21)
$$|U^{n+1}|_{s;n+1} \leq \frac{\varepsilon c}{\delta} |U^n|_{s;n} \left(c \,\kappa^2 \, |W^{0<} + U^n|_{s;n}^2 + \mu \kappa^2 \, |W^{0<}|_{s;n} + |f^{0<}|_s \right) \\ + \varepsilon \kappa^2 \, c \, |W^{0<} + U^n|_{s;n}^2 + \mu \varepsilon \kappa^2 \, |U^n|_{s;n} + \varepsilon \, |f^{\varepsilon<}|_s \,.$$

To complete the inductive step, let us now set

(5.22) $\delta = \varepsilon^{1/4}$ and $\kappa = \varepsilon^{-1/4}$.

With this, we have from (5.21)

$$(5.23) \quad \begin{aligned} |U^{n+1}|_{s;n+1} &\leq \varepsilon^{1/4} c_1 |U^n|_{s;n} \left(|W^{0<} + U^n|_{s;n}^2 + \mu |W^{0<}|_{s;n} + \varepsilon^{1/2} |f^{0<}|_s \right) \\ &+ \varepsilon^{1/2} c_2 \left(|W^{0<} + U^n|_{s;n}^2 + \mu |U^n|_{s;n} + \varepsilon^{1/2} |f^{\varepsilon<}|_s \right). \end{aligned}$$

We require ε to be such that

 $(5.24) \quad \varepsilon^{1/4} \left(c_0 + c_1 + c_2 \right) \left(|W^{0<}|_{s;0}^2 + \mu \, |W^{0<}|_{s;0} + |f|_s \right) \leq \frac{1}{4} \min\{1, |W^{0<}|_s\}$

and claim that with this we have

 $(5.25) \quad |U^n|_{s;n} \le \varepsilon^{1/4} c_U \left(|W^{0<}|_{s;0}^2 + \mu \, |W^{0<}|_{s;0} + |f^{<}|_s \right)$

with $c_U = 4 (c_0 + c_1 + c_2)$. Now since $\varepsilon \leq 1$, (5.16) implies that it holds for n = 1, so let us suppose that it holds for $m = 0, \dots, n$ for some $n < n_*$. Now (5.24) and (5.25) imply that

(5.26)
$$|U^m|_{s;m} \le |W^{0<}|_s \le |W^{0<}|_{s;0}$$
 and $|U^m|_{s;m} \le 1$

for $m = 0, \dots, n$. Using these in (5.23), we have

(5.27)
$$\begin{aligned} |U^{n+1}|_{s;n+1} &\leq 4 \,\varepsilon^{1/4} \, c_1 \left(|W^{0<}|_{s;0}^2 + \mu \, |W^{0<}|_{s;0} + |f^{<}|_s \right) |U^n|_{s;n} \\ &+ 4 \,\varepsilon^{1/2} \, c_2 \left(|W^{0<}|_{s;0}^2 + \mu \, |W^{0<}|_{s;0} + |f^{<}|_s \right) \\ &\leq \varepsilon^{1/4} c_U \left(|W^{0<}|_{s;0}^2 + \mu \, |W^{0<}|_{s;0} + |f^{<}|_s \right). \end{aligned}$$

This proves (5.25) and (5.26) for $n = 0, \dots, n_*$.

We now turn to the remainder

(5.28) $\mathcal{R}^{0} := B^{\varepsilon <}(W^{0<}, W^{0<}) - f^{\varepsilon <}$ and, for $n = 1, \cdots$, (5.29) $\mathcal{R}^{n} := \mathsf{P}^{<}[(\mathsf{D}U^{n})\mathcal{G}^{n}] + \frac{1}{\varepsilon}LU^{n} + B^{\varepsilon <}(W^{0<} + U^{n}, W^{0<} + U^{n}) + AU^{n} - f^{\varepsilon <}.$

We seek to show that, for $n = 0, \dots, n_*$, it scales as e^{-n} . We first note that by construction $\mathcal{R}^n \notin \ker L$, so $L^{-1}\mathcal{R}^n$ is well-defined. Taking $U^0 = 0$, we have

(5.30)
$$\mathcal{R}^n = \frac{1}{\varepsilon} L \left(U^n - U^{n+1} \right).$$

We then compute

$$\begin{aligned} \mathcal{R}^{n+1} &= \mathsf{P}^{<}[(\mathsf{D}U^{n+1})\,\mathcal{G}^{n+1}] + \frac{1}{\varepsilon}LU^{n+1} \\ &+ B^{\varepsilon <}(W^{0 <} + U^{n+1}, W^{0 <} + U^{n+1}) + AU^{n+1} - f^{\varepsilon <} \\ &= \mathsf{P}^{<}[(\mathsf{D}U^{n+1})(\mathcal{G}^{n} + \delta\mathcal{G}^{n})] + \frac{1}{\varepsilon}LU^{n} - \mathcal{R}^{n} \\ &+ B^{\varepsilon <}(W^{0 <} + U^{n}, W^{0 <} + U^{n}) - \varepsilon \, B^{\varepsilon <}(W^{0 <} + U^{n}, L^{-1}\mathcal{R}^{n}) \\ &- \varepsilon \, B^{\varepsilon <}(L^{-1}\mathcal{R}^{n}, W^{0 <} + U^{n+1}) + AU^{n} - \varepsilon \, AL^{-1}\mathcal{R}^{n} - f^{\varepsilon <} \\ &= \mathsf{P}^{<}[(\mathsf{D}U^{n})\,\delta\mathcal{G}^{n}] - \varepsilon \, L^{-1}\mathsf{P}^{<}[(\mathsf{D}\mathcal{R}^{n})\,\mathcal{G}^{n+1}] - \varepsilon \, AL^{-1}\mathcal{R}^{n} \\ &- \varepsilon \, B^{\varepsilon <}(L^{-1}\mathcal{R}^{n}, W^{0 <} + U^{n+1}) - \varepsilon \, B^{\varepsilon <}(W^{0 <} + U^{n}, L^{-1}\mathcal{R}^{n}), \end{aligned}$$

where we have used (5.30) and where

(5.32)
$$\begin{aligned} \delta \mathcal{G}^n &:= \mathcal{G}^{n+1} - \mathcal{G}^n \\ &= \varepsilon B^{0<} (W^{0<} + U^{n+1}, L^{-1} \mathcal{R}^n) + \varepsilon B^{0<} (L^{-1} \mathcal{R}^n, W^{0<} + U^n). \end{aligned}$$

To obtain a bound on \mathcal{R}^n , we compute using (5.26)

(5.33)
$$\begin{aligned} |\mathcal{G}^{n}|_{s;n} &\leq c \left(|\nabla (W^{0<} + U^{n})|_{s;n}^{2} + \mu |\Delta W^{0<}|_{s;n} + |f^{0<}|_{s} \right) \\ &\leq c \kappa^{2} \left(|W^{0<}|_{s;0}^{2} + \mu |W^{0<}|_{s;0} + |f|_{s} \right), \end{aligned}$$

as well as

(5.34)
$$\begin{aligned} |\delta \mathcal{G}^{n}|_{s;n+1} &\leq \varepsilon \, c \, |\nabla (W^{0<} + U^{n+1})|_{s;n+1} |\nabla L^{-1} \mathcal{R}^{n}|_{s;n+1} \\ &+ \varepsilon \, c \, |\nabla L^{-1} \mathcal{R}^{n}|_{s;n+1} |\nabla (W^{0<} + U^{n})|_{s;n} \\ &\leq \varepsilon \kappa^{2} \, c \, |\mathcal{R}^{n}|_{s;n+1} |W^{0<}|_{s;0} \, . \end{aligned}$$

(Note that we can only estimate $\delta \mathcal{G}^n$ in $D_{\eta-(n+1)\delta}$ and not in $D_{\eta-n\delta}$; similarly, since the definition of \mathcal{R}^n involves $\mathsf{D}U^n$, it can only be estimated in $D_{\eta-(n+1)\delta}$.)

22

We then have

$$\begin{aligned} |\mathcal{R}^{n+1}|_{s;n+2} &\leq |\mathsf{D}U^n|_{s;n+1} |\delta\mathcal{G}^n|_{s;n+1} + \varepsilon |L^{-1}\mathsf{D}\mathcal{R}^n|_{s;n+2} |\mathcal{G}^{n+1}|_{s;n+1} \\ &+ \varepsilon \mu \kappa^2 |\mathcal{R}^n|_{s;n+1} + \varepsilon |\nabla L^{-1}\mathcal{R}^n|_{s;n+1} |\nabla (W^{0<} + U^{n+1})|_{s;n+1} \\ &+ \varepsilon |\nabla (W^{0<} + U^n)|_{s;n} |\nabla L^{-1}\mathcal{R}^n|_{s;n+1} \\ &\leq \frac{1}{\delta} |U^n|_{s;n} \varepsilon \kappa^2 |\mathcal{R}^n|_{s;n+1} |W^{0<}|_{s;0} + c \frac{\varepsilon}{\delta} |\mathcal{R}^n|_{s;n+1} |\mathcal{G}^{n+1}|_{s;n+1} \\ &+ 4 \varepsilon \kappa^2 |\mathcal{R}^n|_{s;n+1} |W^{0<}|_{s;0} + \varepsilon \mu \kappa^2 |\mathcal{R}^n|_{s;n+1} \\ &\leq \varepsilon^{1/4} |\mathcal{R}^n|_{s;n+1} c_e (|W^{0<}|_{s;0}^2 + \mu |W^{0<}|_{s;0} + |f^{<}|_{s} + \mu) \end{aligned}$$

where for the last inequality we have assumed that

(5.36)
$$\varepsilon^{1/4} \le \min\{\mu/|W^{0<}|_{s;0}, \mu c_U/4\}.$$

If we require ε to satisfy, in addition to $\varepsilon \leq 1$, (5.24) and (5.36),

(5.37)
$$\varepsilon^{1/4} c_e(|W^{0<}|_{s;0}^2 + \mu |W^{0<}|_{s;0} + |f^{<}|_s + \mu) \le \frac{1}{e},$$

we have, for $n = 0, 1, \dots, n_* - 1$,

(5.38)
$$|\mathcal{R}^{n+1}|_{s;n+2} \le \frac{1}{\mathrm{e}} |\mathcal{R}^{n}|_{s;n+1}.$$

Along with the estimate

(5.39)
$$|\mathcal{R}^0|_{s;1} \le c_r (|W^{0<}|_{s;0}^2 + |f^{<}|_s),$$

taking $n = n_* - 1$ leads us to

(5.40)
$$\begin{aligned} |\mathcal{R}^{n_*-1}|_{H^s} &\leq |\mathcal{R}^{n_*-1}|_{s;n_*} \leq c_r \left(|W^{0<}|_{s;0}^2 + |f^{<}|_s\right) \exp(-\eta/\varepsilon^{1/4}) \\ &\leq c_r \left[\left(|W^{0<}|_s + \eta\right)^2 + |f^{<}|_s\right] \exp(-\eta/\varepsilon^{1/4}). \end{aligned}$$

The lemma follows by setting $U^* = U^{n_*-1}$ and taking as ε_{**} the largest value that satisfies $\varepsilon \leq 1$, (5.24), (5.36) and (5.37).

For use later in the proof of Theorem 2, we also bound

(5.41)
$$\begin{aligned} \left| \nabla (1 - \mathsf{P}^{<}) [(\mathsf{D}U^{*})\mathcal{G}^{*}] \right|_{L^{2}} &\leq c \, \mathrm{e}^{-\sigma\kappa} |(\mathsf{D}U^{*})\mathcal{G}^{*}|_{2,n_{*}} \\ &\leq c \, \mathrm{e}^{-\sigma\kappa} \, \frac{1}{\delta} \, |U^{*}|_{2,n_{*}-1} |\mathcal{G}^{*}|_{2,n_{*}-1} \\ &\leq c \, \mathrm{e}^{-\sigma\kappa} \, \kappa^{2} \, (|W^{0<}|_{2;0}^{2} + \mu \, |W^{0<}|_{2;0} + |f|_{2})^{2} \end{aligned}$$

where for the last inequality we have used (5.25) and (5.33) with $n = n_* - 1$.

5.2. **Proof of Theorem 2.** We follow the conventions of the proofs of Theorem 1 and Lemma 2 on constants. We will be rather terse in parts of this proof which mirror a development in the proof of Theorem 1.

First, we recall Theorem 0 and consider $t \ge T := \max\{T_2, T_\sigma\}$ so that $|\nabla^2 W(t)| \le K_2$ and $|\nabla^2 W(t)|_{G^{\sigma}} \le M_{\sigma}$. We use Lemma 2 with s = 2 and, collecting the constraints on ε there, require that

(5.42)
$$\begin{aligned} \varepsilon^{1/4} c_U \left((K_2 + \eta)^2 + \mu (K_2 + \eta) + |f|_2 \right) &\leq \frac{1}{4} \min\{1, K_2\}, \\ \varepsilon^{1/4} &\leq \min\{\mu/(K_2 + \eta), \mu c_U/4, 1\}, \\ \varepsilon^{1/4} c_e \left((K_2 + \eta)^2 + \mu (K_2 + \eta) + \mu + |f|_2 \right) &\leq \frac{1}{e}, \end{aligned}$$

where c_e is that in (5.37). (We note that all these constraints are convex in K_2 , so they do not cause problems when $|W^{0<}| < K_2$.) Further constraints on ε will be imposed below. We note the bound (5.40) and

(5.43)
$$|U^*|_{H^2} \le |U^*|_{2;n_*} \le \varepsilon^{1/4} c_U \left((K_2 + \eta)^2 + \mu (K_2 + \eta) + |f|_2 \right)$$

which follows from (5.26)

which follows from (5.26). We fix $\kappa = \varepsilon^{-1/4}$ as in (5.22) and consider the equation of motion for the low modes $W^{<}$,

(5.44)
$$\partial_t W^{<} + \frac{1}{\varepsilon} L W^{<} + B^{<}(W^{<}, W^{<}) + A W^{<} - f^{<} = -B^{<}(W^{>}, W) - B^{<}(W^{<}, W^{>}) =: \hat{\mathcal{H}}.$$

Writing

(5.45) $W^{\varepsilon <} = U^*(W^{0<}, f^{<}; \varepsilon) + W',$

the equation governing the finite-dimensional variable $W^\prime(t)$ is

(5.46)
$$\partial_t W' + \frac{1}{\varepsilon} LW' + B^{\varepsilon <} (W^<, W^<) + AW' \\ = -\partial_t U^* - \frac{1}{\varepsilon} LU^* - AU^* + f^{\varepsilon <} + \hat{\mathcal{H}}^{\varepsilon}.$$

Using (5.4), this can be written as

(5.47)
$$\partial_t W' + \frac{1}{\varepsilon} LW' + B^{\varepsilon <} (W^<, W') + B^{\varepsilon <} (W', W^{0<} + U^*) + AW'$$
$$= -\mathcal{R}^* - (1 - \mathsf{P}^<)[(\mathsf{D}U^*) \mathcal{G}^*] + \hat{\mathcal{H}}^{\varepsilon}$$
$$=: -\mathcal{R}^* + \mathcal{H}^{\varepsilon}.$$

Multiplying by W' in $L^2(\mathcal{M})$, we find

(5.48)
$$\frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} |W'|^2 + (W', B^{\varepsilon <}(W', W^{0 <} + U^*)) + \mu |\nabla W'|^2 = -(W', \mathcal{R}^*) + (W', \mathcal{H}^{\varepsilon})$$

We now write the nonlinear term as

$$(W', B^{\varepsilon <}(W', W^{0<} + U^*)) = (W', B(W', W^{0<} + U^*))$$

(5.49)
$$= (W', B(W', U^*)) + (W', B(W', W^{0<}))$$
$$= (W', B(W', U^*)) - (W^{0<}, B(W', W')).$$

Following the proof of Theorem 1 [cf. (4.10)], we rewrite (5.48) as

(5.50)
$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\mathrm{e}^{\nu t} |W'|^2 \right) + \mu \, \mathrm{e}^{\nu t} \, |\nabla W'|^2 \leq -2 \, \mathrm{e}^{\nu t} \, (W', \mathcal{R}^*) + 2 \, \mathrm{e}^{\nu t} \, (W', \mathcal{H}^\varepsilon) \\ - 2 \, \mathrm{e}^{\nu t} \, (W', B(W', U^*)) + 2 \, \mathrm{e}^{\nu t} \, (W^{0<}, B(W', W')).$$

We bound the first two terms on the right-hand side as

(5.51)
$$2 |(W', \mathcal{R}^*)| \leq \frac{\mu}{6} |\nabla W'|^2 + \frac{c}{\mu} |\mathcal{R}^*|^2,$$
$$2 |(W', \mathcal{H}^\varepsilon)| \leq \frac{\mu}{6} |\nabla W'|^2 + \frac{c}{\mu} |\mathcal{H}^\varepsilon|^2.$$

As for the third term in (5.50), we bound it as

(5.52)
$$2 |(W', B(W', U^*))| \le c |W'|_{L^6} |\nabla W'|_{L^2} |\nabla U^*|_{L^3} \le |\nabla W'|^2 c_1 \varepsilon^{1/4} ((|W^{0<}|_2 + \eta)^2 + \mu (|W^{0<}|_2 + \eta) + |f^{<}|_2)$$

where we have used (5.43) in the last step. We now require ε to be small enough so that

(5.53)
$$\varepsilon^{1/4} c_1 \left((K_2 + \eta)^2 + \mu K_2 + \mu \eta + |f|_2 \right) \le \frac{\mu}{6},$$

which implies that, since $|W^{0<}|_2 \leq K_2$ by hypothesis,

(5.54)
$$2|(W', B(W', U^*))| \le \frac{\mu}{6} |\nabla W'|^2.$$

With these estimates, (5.50) becomes

(5.55)
$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\mathrm{e}^{\nu t} |W'|^2 \right) + \frac{\mu}{2} \, \mathrm{e}^{\nu t} \, |\nabla W'|^2 \le \frac{c}{\mu} \, \mathrm{e}^{\nu t} \left(|\mathcal{R}^*|^2 + |\mathcal{H}^\varepsilon|^2 \right) \\ + 2 \, \mathrm{e}^{\nu t} \left(W^{0<}, B(W', W') \right).$$

Integrating this inequality and multiplying by $e^{-\nu T}$, we find

(5.56)
$$e^{\nu t} |W'(T+t)|^2 - |W'(T)|^2 + \frac{\mu}{2} \int_T^{T+t} e^{\nu(\tau-T)} |\nabla W'|^2 d\tau \leq \int_T^{T+t} e^{\nu(\tau-T)} \left\{ \frac{c}{\mu} (|\mathcal{R}^*|^2 + |\mathcal{H}^\varepsilon|^2) + 2 (W^{0<}, B(W', W')) \right\} d\tau.$$

We then integrate the last term by parts as in (4.18),

(5.57)
$$\int_{T}^{T+t} e^{\nu(\tau-T)} (W^{0<}, B(W', W')) d\tau$$
$$= \varepsilon e^{\nu t} (W^{0<}, B_{\omega}(W', W'))(T+t) - \varepsilon (W^{0<}, B_{\omega}(W', W'))(T)$$
$$- \varepsilon \int_{T}^{T+t} e^{\nu(\tau-T)} \left\{ \nu (W^{0<}, B_{\omega}(W', W')) + (\partial_{\tau} W^{0<}, B_{\omega}(W', W')) + 2 (W^{0<}, B_{\omega}(\partial_{\tau}^{*} W', W')) \right\} d\tau.$$

To bound the terms in the integral, we first need to estimate

(5.58)

$$\begin{aligned} |\nabla B^{\varepsilon <}(W', W^{0 <} + U^{*})|_{L^{2}} &\leq \kappa |B^{\varepsilon <}(W', W^{0 <} + U^{*})|_{L^{2}} \\ &\leq c \kappa |\nabla W'|_{L^{2}} |\nabla (W^{0 <} + U^{*})|_{L^{\infty}} \\ &\leq c \kappa^{2} |\nabla W'| |W^{0 <} + U^{*}|_{H^{2}} \\ &\leq c \kappa^{2} |\nabla W'| |\nabla^{2} W^{0}| \end{aligned}$$

where for the last inequality we have used (5.26). Using this and the bound (5.59) $|\nabla B^{\varepsilon <}(W^<, W')|_{L^2} \leq c \kappa |\nabla W^<|_{L^\infty} |\nabla W'|_{L^2} \leq c \kappa^2 |\nabla^2 W| |\nabla W'|$ for the term $B^{\varepsilon <}(W^<, W')$ in (5.47) gives us (5.60) $|\nabla \partial_t^* W'|_{L^2} \leq c \kappa^2 |\nabla W'| |\nabla^2 W| + \mu \kappa^2 |\nabla W'| + |\nabla \mathcal{R}^*| + |\nabla \mathcal{H}^{\varepsilon}|.$ The worst term in (5.57) can now be bounded as

(5.61)

$$\varepsilon |(W^{0<}, B_{\omega}(\partial_t^*W', W'))| \le \varepsilon c |W^{0<}|_{L^{\infty}} |\nabla W'|_{L^2} |\nabla \partial_t^*W'|_{L^2}$$

$$\le c_2 \varepsilon \kappa^2 K_2^2 |\nabla W'|^2 + c_3 \varepsilon \kappa^2 \mu K_2 |\nabla W'|^2$$

$$+ \frac{\mu}{48} |\nabla W'|^2 + \frac{\varepsilon^2 c K_2^2}{\mu} (|\nabla \mathcal{R}^*|^2 + |\nabla \mathcal{H}^{\varepsilon}|^2)$$

If we now require that ε satisfy

(5.62)
$$\varepsilon^{1/2} c_2 K_2^2 \le \frac{\mu}{48}$$
 and $\varepsilon^{1/2} c_3 K_2 \le \frac{1}{48}$,

we have

(5.63)
$$\varepsilon |(W^{0<}, B_{\omega}(\partial_t^*W', W'))| \leq \frac{\mu}{16} |\nabla W'|^2 + \frac{\varepsilon^2 c K_2^2}{\mu} (|\nabla \mathcal{R}^*|^2 + |\nabla \mathcal{H}^{\varepsilon}|^2).$$

Bounding another term in (5.57) as

(5.64)

$$\varepsilon \left| (\partial_t W^{0<}, B_{\omega}(W', W')) \right| \leq \varepsilon c \left| \partial_t W^0 \right|_{L^{\infty}} |\nabla W'|^2_{L^2}$$

$$\leq \varepsilon c \left| \partial_t W^0 \right|_{H^2} |\nabla W'|^2_{L^2}$$

$$\leq \varepsilon c_4 \left(\kappa K_2^2 + \mu \kappa^2 K_2 + |f|_2 \right) |\nabla W'|^2$$

and requiring that ε also satisfy

(5.65)
$$\varepsilon^{1/2} c_4 \left(K_2^2 + \mu K_2 + |f|_2 \right) \le \frac{\mu}{12},$$

plus a similar estimate for the first (easiest) term in (5.57), we can bound the integral on the r.h.s. as

(5.66)
$$\int_{T}^{T+t} e^{\nu(\tau-T)} \left| (W^{0<}, B(W', W')) \right| d\tau \\ \leq \frac{\mu}{2} \int_{T}^{T+t} e^{\nu(\tau-T)} |\nabla W'|^2 d\tau + \frac{\varepsilon c}{\mu^2} K_2^2 \left(\|\nabla \mathcal{R}^*\|^2 + \|\nabla \mathcal{H}^\varepsilon\|^2 \right) (e^{\nu t} - 1)$$

where $\|\nabla \mathcal{R}^*\| := \sup_{|W^0| \leq K_2} |\nabla \mathcal{R}^*(W^0, f; \varepsilon)|$ and similarly for $\|\nabla \mathcal{H}^{\varepsilon}\|$. Bounding the limit term in (5.57) as

$$(5.67) |(W^{0<}, B_{\omega}(W', W'))| \le C |W^{0<}|_{L^{\infty}} |\nabla W'|_{L^{2}}^{2} \le c K_{2} \kappa^{2} |W'|^{2},$$

(5.56) becomes

(5.68)
$$(1 - \varepsilon^{1/2} c_5 K_2) |W'(T+t)|^2 \\ \leq e^{-\nu t} (1 + \varepsilon^{1/2} c_5 K_2) |W'(T)|^2 + \frac{\varepsilon c}{\mu^2} K_2^2 (\|\nabla \mathcal{R}^*\|^2 + \|\nabla \mathcal{H}^\varepsilon\|^2)$$

To estimate $\|\nabla \mathcal{H}^{\varepsilon}\|$, we use (5.44), (5.3) and (2.20), to obtain

(5.69)
$$\begin{aligned} |\nabla B^{\varepsilon <}(W^{>},W)|_{L^{2}} + |\nabla B^{\varepsilon <}(W^{<},W^{>})|_{L^{2}} &\leq c \,\kappa \, |\nabla W^{>}|_{L^{4}} |\nabla W|_{L^{4}} \\ &\leq c \,\kappa \,\mathrm{e}^{-\sigma\kappa} \, M_{\sigma} \, K_{2} \,. \end{aligned}$$

Now (5.41) implies that

(5.70) $|\nabla(1 - \mathsf{P}^{<})[(\mathsf{D}U^{*})\mathcal{G}^{*}]|_{L^{2}} \leq c \,\mathrm{e}^{-\sigma\kappa} \,\kappa^{2} \left((K_{2} + \eta)^{2} + \mu (K_{2} + \eta) + |f|_{2}\right)^{2};$ this and the previous estimate give us (5.71) $\|\nabla\mathcal{H}^{\varepsilon}\|_{L^{2}} \leq c \,\mathrm{e}^{-\sigma\kappa} \,\kappa^{2} \left[M_{\sigma}K_{2} + \left((K_{2} + \eta)^{2} + \mu (K_{2} + \eta) + |f|_{2}\right)^{2}\right].$

26

Meanwhile, using (5.40) we have

(5.72)
$$\|\nabla \mathcal{R}^*\|_{L^2} \le c \left((K_2 + \eta)^2 + |f|_2 \right) \exp(-\eta/\varepsilon^{1/4})$$

Setting $\eta = \sigma$ and requiring ε to satisfy, in addition to (5.42), (5.62) and (5.65),

(5.73)
$$\varepsilon^{1/2} c_5 K_2 \leq \frac{1}{2}$$
,

we have

(5.74)
$$|W'(T+t)|^2 \le 4 e^{-\nu t} |W'(T)|^2 + \frac{c}{\mu^2} \left[(K_2 + \sigma)^4 + \mu^2 (K_2 + \sigma)^2 + |f|_2^2 + |f|_2^4 + M_\sigma^2 K_2^2 \right] \exp(-2\sigma/\varepsilon^{1/4})$$

Since $|W'(T)| \leq c K_2$ by Theorem 0, by taking t sufficiently large we have

(5.75)
$$|W'(T+t)| \leq \frac{c}{\mu} \Big[(K_2 + \sigma)^4 + \mu^2 (K_2 + \sigma)^2 + |f|_2^2 + |f|_2^4 + M_\sigma^2 K_2^2 \Big] \times \varepsilon^{1/2} \exp(-\sigma/\varepsilon^{1/4}).$$

And since

(5.76) $|W^{\varepsilon} - U^*|^2 \le |W^{\varepsilon}|^2 + |W'|^2 \le c M_{\sigma}^2 \exp(-2\sigma/\varepsilon^{1/4}) + |W'|^2$,

the theorem follows by the same argument used to obtain Theorem 1.

APPENDIX A.

Proof of Lemma 1. Since $B_{jkl}^{rs0} = 0$ when $j_3k_3|l| = 0$, we assume that $j_3k_3|l| \neq 0$ in the rest of this proof. As before, all wavevectors are understood to live in $\mathbb{Z}_L - \{0\}$ and their third component take values in $\{0, \pm 2\pi/L_3, \pm 4\pi/L_3, \cdots\}$.

We start by noting that an *exact* resonance is only possible when j and k lie on the same "resonance cone", that is, when $|\mathbf{j}|/|j_3| = |\mathbf{k}|/|k_3|$, or equivalently, when $|\mathbf{j}'|/|j_3| = |\mathbf{k}'|/|k_3|$. There are only two cases to consider: (a') When $\mathbf{j}' = \mathbf{k}' = 0$, we have $B_{\mathbf{j}\mathbf{k}\mathbf{l}}^{rs0} = B_{\mathbf{k}\mathbf{j}\mathbf{l}}^{sr0} = 0$.

(b') In the generic case $j_3k_3|\mathbf{j}'||\mathbf{k}'| \neq 0$, direct computation using the resonance relation $r|\mathbf{j}|/j_3 + s|\mathbf{k}|/k_3 = 0$ gives $B_{\mathbf{j}\mathbf{k}\mathbf{l}}^{rs0} + B_{\mathbf{k}\mathbf{j}\mathbf{l}}^{sr0} = 0$. This result also follows as the special case $\omega_{j}^{r} + \omega_{k}^{s} = 0$ in (A.9) below.

Now we turn to *near* resonances. There are several cases to consider, and we start with the generic (and hardest) one.

(a) Suppose that $|j'| |k'| \neq 0$ with $l' \neq 0$. We define Ω and θ by

(A.1)
$$2\Omega := \omega_{j}^{r} - \omega_{k}^{s}$$
 and $2\theta\Omega := \omega_{j}^{r} + \omega_{k}^{s}$

(We note that Ω and θ could take either sign. Our concern is obviously with small $|\theta|$, when when ω_{i}^{r} and ω_{k}^{s} are nearly resonant, so we will restrict θ below.) Now this implies that

(A.2)
$$\omega_{j}^{r} = (1+\theta)\Omega$$
 and $\omega_{k}^{s} = (\theta-1)\Omega$.

We first note that

(A.3)
$$|\mathbf{j}'|^2 / |j_3|^2 = (1+\theta)^2 \Omega^2 - 1$$
 and $|\mathbf{k}'|^2 / |k_3|^2 = (1-\theta)^2 \Omega^2 - 1$
and compute

(A.4)
$$|\mathbf{j}'|^2 \frac{k_3}{j_3} - |\mathbf{k}'|^2 \frac{j_3}{k_3} = 4\theta \Omega^2 j_3 k_3 = -\frac{4\theta}{1-\theta^2} rs |\mathbf{j}| |\mathbf{k}|.$$

Direct computation gives us

(A.5)
$$B_{jkl}^{rs0} + B_{kjl}^{sr0} = \frac{i|\mathcal{M}|\delta_{j+k-l}j_{3}k_{3}}{2|j||j'||k||k'||l|} [(P+P')(Q+Q') + (-P+P'')(Q+Q'')] = \frac{i|\mathcal{M}|\delta_{j+k-l}j_{3}k_{3}}{2|j||j'||k||k'||l|} [P(Q'-Q'') + (P'+P'')Q + P'Q' + P''Q'']$$

where

$$P := \mathbf{j}' \wedge \mathbf{k}' \qquad Q := -\mathbf{j}' \cdot \mathbf{k}'$$
(A.6)
$$P' := i\frac{r|\mathbf{j}|}{j_3}(\mathbf{j}' \cdot \mathbf{k}') - i\frac{r|\mathbf{j}|}{j_3}\frac{k_3}{j_3}|\mathbf{j}'|^2 \qquad Q' := -i\frac{s|\mathbf{k}|}{k_3}(\mathbf{j}' \wedge \mathbf{k}') + \frac{j_3}{k_3}|\mathbf{k}'|^2$$

$$P'' := i\frac{s|\mathbf{k}|}{k_3}(\mathbf{j}' \cdot \mathbf{k}') - i\frac{s|\mathbf{k}|}{k_3}\frac{j_3}{k_3}|\mathbf{k}'|^2 \qquad Q'' := i\frac{r|\mathbf{j}|}{j_3}(\mathbf{j}' \wedge \mathbf{k}') + \frac{k_3}{j_3}|\mathbf{j}'|^2.$$

After some computation, we find

.

$$P' + P'' = 2\theta\Omega i \left(\mathbf{j}' \cdot \mathbf{k}' + \frac{2rs}{1 - \theta^2} |\mathbf{j}| |\mathbf{k}| - \frac{|\mathbf{j}'|^2}{2} \frac{k_3}{j_3} - \frac{|\mathbf{k}'|^2}{2} \frac{j_3}{k_3} \right),$$
(A.7)

$$Q' - Q'' = 2\theta\Omega \left(\frac{2rs/\Omega}{1 - \theta^2} |\mathbf{j}| |\mathbf{k}| - i(\mathbf{j}' \wedge \mathbf{k}') \right),$$

$$P'Q' + P''Q'' = 2\theta\Omega \left\{ \frac{2rs |\mathbf{j}| |\mathbf{k}|}{1 - \theta^2} \frac{r|\mathbf{j}| \frac{s|\mathbf{k}|}{j_3}}{k_3} (\mathbf{j}' \wedge \mathbf{k}') + 2irs|\mathbf{j}| |\mathbf{k}| (\mathbf{j}' \cdot \mathbf{k}') + \frac{i}{2} (\mathbf{j}' \cdot \mathbf{k}') \left(|\mathbf{k}'|^2 \frac{j_3}{k_3} + |\mathbf{j}'|^2 \frac{k_3}{j_3} \right) - i |\mathbf{j}'|^2 |\mathbf{k}'|^2 \right\},$$

from which we obtain

(A.8)
$$B_{jkl}^{rs0} + B_{kjl}^{sr0} = 2\theta\Omega \frac{\mathbf{i} |\mathcal{M}| \, \delta_{j+k-l}}{2 |l|} \Big\{ \mathbf{i} \, (\mathbf{j}' \cdot \mathbf{k}') \frac{|\mathbf{j}'|^2 k_3^2 + |\mathbf{k}'|^2 j_3^2}{|\mathbf{j}| \, |\mathbf{j}'| \, |\mathbf{k}|} - 2\mathbf{i} \frac{|\mathbf{j}'| \, |\mathbf{k}'| \, j_3 k_3}{|\mathbf{j}| \, |\mathbf{k}|} \\ - \frac{2\mathbf{i} \, rs \, \theta^2}{1 - \theta^2} \frac{(\mathbf{j}' \cdot \mathbf{k}') j_3 k_3}{|\mathbf{j}'| \, |\mathbf{k}'|} + \frac{2 \, rs / \Omega}{1 - \theta^2} \frac{\mathbf{j}' \wedge \mathbf{k}'}{|\mathbf{j}'| \, |\mathbf{k}'|} \, j_3 k_3 + \frac{2 / \Omega}{1 - \theta^2} \frac{|\mathbf{j}| \, |\mathbf{k}|}{|\mathbf{j}'| \, |\mathbf{k}'|} (\mathbf{j}' \wedge \mathbf{k}') \Big\}$$

Now if we require that $|\theta| \leq \theta_0 < 1$, we have the bound

(A.9)
$$\left| B_{\boldsymbol{j}\boldsymbol{k}\boldsymbol{l}}^{rs0} + B_{\boldsymbol{k}\boldsymbol{j}\boldsymbol{l}}^{sr0} \right| \leq \frac{|\mathcal{M}|}{2} \left(4 + \frac{6}{1 - \theta_0^2} \right) \frac{|\boldsymbol{j}| |\boldsymbol{k}|}{|\boldsymbol{l}|} |\omega_{\boldsymbol{j}}^r + \omega_{\boldsymbol{k}}^s|.$$

To take care of the case $|\theta| > \theta_0$, we note that in this case

(A.10)
$$|\omega_{\boldsymbol{j}}^r + \omega_{\boldsymbol{k}}^s| \ge \theta_0 \left(\frac{|\boldsymbol{j}|}{|j_3|} + \frac{|\boldsymbol{k}|}{|k_3|}\right).$$

We note that since $\theta_0 < 1$ by hypothesis, this inequality holds both when $\omega_j^r \omega_k^s < 0$ and $\omega_{j}^{r}\omega_{k}^{s} > 0$. Using (3.49), we then find

(A.11)
$$|B_{jkl}^{rs0} + B_{kjl}^{sr0}| \le |B_{jkl}^{rs0}| + |B_{kjl}^{sr0}| \le \sqrt{5} |\mathcal{M}| \left(|\mathbf{k}'| + |\mathbf{j}'| + |\mathbf{k}'| \frac{|j_3|}{|k_3|} + |\mathbf{j}'| \frac{|k_3|}{|j_3|} \right).$$

Putting these together, we find after a short computation,

(A.12)
$$\left| B_{\boldsymbol{j}\boldsymbol{k}\boldsymbol{l}}^{rs0} + B_{\boldsymbol{k}\boldsymbol{j}\boldsymbol{l}}^{sr0} \right| \leq \frac{2\sqrt{5}\left|\mathcal{M}\right|}{\theta_0} \left(\left| j_3 \right| + \left| k_3 \right| \right) \left| \omega_{\boldsymbol{j}}^r + \omega_{\boldsymbol{k}}^s \right|.$$

(b) Suppose now that $|\mathbf{j}'| |\mathbf{k}'| \neq 0$ but $\mathbf{l}' = 0$. We find using $\mathbf{j}' + \mathbf{k}' = 0$,

(A.13)
$$B_{jkl}^{rs0} + B_{kjl}^{sr0} = i |\mathcal{M}| \, \delta_{j+k-l} \, \frac{-i \operatorname{sgn} l_3 \, |j'| \, |k'|}{2 \, |j| \, |k|} (j_3 + k_3) (\omega_j^r + \omega_k^s),$$

and thus the bound

(A.14)
$$|B_{jkl}^{rs0} + B_{kjl}^{sr0}| \le \frac{|\mathcal{M}|}{2} (|j_3| + |k_3|) |\omega_j^r + \omega_k^s|.$$

(c) Finally, we consider the case $\mathbf{j}' = 0$ and $\mathbf{k}' \neq 0$ (which obviously implies the case $\mathbf{k}' = 0$ and $\mathbf{j}' \neq 0$). After some computation using $\mathbf{l}' = \mathbf{k}'$, we find

(A.15)
$$B_{jkl}^{rs0} + B_{kjl}^{sr0} = \frac{i |\mathcal{M}| \delta_{j+k-l}}{2 |l| |k|} j_3(k_1 - irk_2) |k'| \left(sr - \frac{|k|}{k_3} \right).$$

But since in this case

(A.16)
$$|\omega_{j}^{r} - \omega_{k}^{s}| = |r \operatorname{sgn} j_{3} - s|\mathbf{k}|/k_{3}| = |rs - |\mathbf{k}|/k_{3}|,$$

we have the bound

(A.17)
$$|B_{jkl}^{rs0} + B_{kjl}^{sr0}| \le \frac{|\mathcal{M}| |j_3| |k'|^2}{\sqrt{2} |k| |l|} |\omega_j^r + \omega_k^s| \le \frac{|\mathcal{M}| |j_3|}{\sqrt{2}} |\omega_j^r + \omega_k^s|,$$

which holds whether or not $l_3 = 0$. We recall that there is nothing to do when j' = k' = 0 since then $B_{jkl}^{rs0} = B_{kjl}^{sr0} = 0$.

The lemma follows upon fixing θ_0 and collecting (A.9), (A.12), (A.14) and (A.17).

References

- A. BABIN, A. MAHALOV, AND B. NICOLAENKO, Global regularity of 3D rotating Navier-Stokes equations for resonant domains, Indiana Univ. Math. J., 48 (1999), pp. 1133–1176.
- [2] —, Fast singular oscillating limits and global regularity for the 3d primitive equations of geophysics, Modél. Math. Anal. Num., 34 (2000), pp. 201–222.
- [3] F. BAER AND J. J. TRIBBIA, On complete filtering of gravity modes through nonlinear initialization, Monthly Weather Review, 105 (1977), pp. 1536–1539.
- [4] P. BARTELLO, Geostrophic adjustment and inverse cascades in rotating stratified turbulence, J. Atmos. Sci., 52 (1995), pp. 4410–4428.
- [5] O. BOKHOVE, Slaving principles, balanced dynamics and the hydrostatic Boussinesq equations, J. Atmos. Sci., 54 (1997), pp. 1662–1674.
- [6] C. CAO AND E. S. TITI, Global well-posedness of the three-dimensional viscous primitive equations of large scale ocean and atmosphere dynamics, Annals Math., 166 (2007), pp. 245– 267.
- [7] R. DALEY, Atmospheric data analysis, Cambridge Univ. Press, 1991.
- [8] P. F. EMBID AND A. J. MAJDA, Averaging over fast gravity waves for geophysical flows with arbitrary potential vorticity, Comm. P.D.E., 21 (1996), pp. 619–658.
- [9] C. FOIAS AND R. TEMAM, Gevrey class regularity for the solutions of the Navier-Stokes equations, J. Funct. Anal., 87 (1989), pp. 359–369.
- [10] R. FORD, M. E. MCINTYRE, AND W. NORTON, Balance and the slow quasi-manifold: some explicit results, J. Atmos. Sci., 57 (2000), pp. 1236–1254.
- [11] A. E. GILL, Atmosphere-ocean dynamics, Academic Press, 1982.
- [12] N. JU, The global attractor for the solutions to the 3D viscous primitive equations, Discrete Cont. Dyn. Systems, Ser. A, 17 (2007), pp. 159–179.
- [13] G. M. KOBELKOV, Existence of a solution 'in the large' for the 3d large-scale ocean dynamics equations, C. R. Acad. Sc. Paris, Sér. I, 343 (2006), pp. 283–286.
- [14] G. M. KOBELKOV, Existence of a solution "in the large" for ocean dynamics equations, J. Math. Fluid Mech., 9 (2007), pp. 588–610.
- [15] M. D. KRUSKAL AND H. SEGUR, Asymptotics beyond all orders in a model of crystal growth, Stud. Appl. Math., 85 (1991), pp. 129–181.

- [16] C. E. LEITH, Nonlinear normal mode initialization and quasi-geostrophic theory, J. Atmos. Sci., 37 (1980), pp. 958–968.
- [17] M.-P. LELONG AND J. J. RILEY, Internal wave-vortical mode interactions in strongly stratified flows, J. Fluid Mech., 232 (1991), pp. 1–19.
- [18] J.-L. LIONS, R. TEMAM, AND S. WANG, New formulations of the primitive equations of the atmosphere and applications, Nonlinearity, 5 (1992), pp. 237–288.
- [19] E. N. LORENZ, On the existence of a slow manifold, J. Atmos. Sci., 43 (1986), pp. 1547–1557.
- [20] B. MACHENHAUER, On the dynamics of gravity oscillations in a shallow water model, with applications to normal mode initialization, Beitr. Phys. Atmos., 50 (1977), pp. 253–271.
- [21] R. S. MACKAY, Slow manifolds, in Energy localisation and transfer, T. Dauxois, A. Litvak-Hinenzon, R. S. MacKay and A. Spanoudaki, eds., World Scientific, 2004, pp. 149–192.
- [22] K. MATTHIES, Time-averaging under fast periodic forcing of parabolic partial differential equations: exponential estimates, J. Diff. Eq., 174 (2001), pp. 133–180.
- [23] M. PETCU, On the three dimensional primitive equations, Adv. Diff. Eq., 11 (2006), pp. 1201– 1226.
- [24] M. PETCU, R. TEMAM, AND D. WIROSOETISNO, Existence and regularity results for the primitive equations in two space dimensions, Comm. Pure Applied Analysis, 3 (2003), pp. 115–131.
- [25] M. PETCU AND D. WIROSOETISNO, Sobolev and Gevrey regularity results for the primitive equations in 3 space dimensions, Applic. Anal., 84 (2005), pp. 769–788.
- [26] L. M. POLVANI, J. C. MCWILLIAMS, M. A. SPALL, AND R. FORD, The coherent structures of shallow-water turbulence: Deformation-radius effects, cyclone/anticyclone asymmetry and gravity-wave generation, Chaos, 4 (1994), pp. 177–186.
- [27] G. M. REZNIK, V. ZEITLIN, AND M. BEN JELLOUL, Nonlinear theory of geostrophic adjustment. Part 1. Rotating shallow water model, J. Fluid Mech., 445 (2001), pp. 93–120.
- [28] R. M. SAMELSON, R. M. TEMAM, C. WANG, AND S. WANG, Surface pressure Poisson equation formulation of the primitive equations: numerical schemes, SIAM J. Numer. Anal., 41 (2003), pp. 1163–1194.
- [29] S. SCHOCHET, Fast singular limits of hyperbolic PDEs, J. Diff. Eq., 114 (1994), pp. 476–512.
- [30] R. TEMAM, Infinite-dimensional dynamical systems in mechanics and physics, 2ed, Springer-Verlag, 1997.
- [31] R. TEMAM AND D. WIROSOETISNO, Exponential approximations for the primitive equations of the ocean, Discr. Cont. Dyn. Sys. B, 7 (2007), pp. 425–440.
- [32] R. TEMAM AND M. ZIANE, Some mathematical problems in geophysical fluid dynamics, in Handbook of mathematical fluid dynamics III, S. Friedlander and D. Serre, eds., Elsevier, 2004, pp. 535–658.
- [33] J. VANNESTE AND I. YAVNEH, Exponentially small inertia-gravity waves and the breakdown of quasi-geostrophic balance, J. Atmos. Sci., 61 (2004), pp. 211–223.
- [34] R. VAUTARD AND B. LEGRAS, Invariant manifolds, quasi-geostrophy and initialization, J. Atmos. Sci., 43 (1986), pp. 565–584.
- [35] T. WARN, Statistical mechanical equilibria of shallow-water equations, Tellus, 38A (1986), pp. 1–11.
- [36] —, Nonlinear balance and quasi-geostrophic sets, Atmos.–Ocean, 35 (1997), pp. 135–145. This paper was written in 1983 but only published (in its original form) in 1997.
- [37] T. WARN, O. BOKHOVE, T. G. SHEPHERD, AND G. K. VALLIS, Rossby number expansions, slaving principles, and balance dynamics, Quart. J. Roy. Met. Soc., 121 (1995), pp. 723–739.
- [38] T. WARN AND R. MÉNARD, Nonlinear balance and gravity-inertial wave saturation in a simple atmospheric model, Tellus, 38A (1986), pp. 285–294.

E-mail address: temam@indiana.edu

URL: http://mypage.iu.edu/~temam

(RT) THE INSTITUTE FOR SCIENTIFIC COMPUTING AND APPLIED MATHEMATICS, INDIANA UNI-VERSITY, RAWLES HALL, BLOOMINGTON, IN 47405–7106, UNITED STATES

E-mail address: djoko.wirosoetisno@durham.ac.uk *URL*: http://www.maths.dur.ac.uk/~dmaOdw

(DW) DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF DURHAM, DURHAM DH1 3LE, UNITED KINGDOM