# An obstacle to a decomposition theorem for near-regular matroids\*

Dillon Mayhew<sup>†</sup> Geoff Whittle<sup>†</sup> Stefan H. M. van Zwam<sup>‡</sup> October 29, 2021

#### Abstract

Seymour's Decomposition Theorem for regular matroids states that any matroid representable over both GF(2) and GF(3) can be obtained from matroids that are graphic, cographic, or isomorphic to  $R_{10}$  by 1-, 2-, and 3-sums. It is hoped that similar characterizations hold for other classes of matroids, notably for the class of near-regular matroids. Suppose that all near-regular matroids can be obtained from matroids that belong to a few basic classes through k-sums. Also suppose that these basic classes are such that, whenever a class contains all graphic matroids, it does not contain all cographic matroids. We show that in that case 3-sums will not suffice.

## 1 Introduction

A regular matroid is a matroid representable over every field. Much is known about this class, the deepest result being Seymour's Decomposition Theorem:

**Theorem 1.1** (Seymour [16]). Let M be a regular matroid. Then M can be obtained from matroids that are graphic, cographic, or equal to  $R_{10}$  through 1-, 2-, and 3-sums.

A class C of matroids is polynomial-time recognizable if there exists an algorithm that decides, for any matroid M, in time  $f(|E(M)|, \tau)$  whether or not  $M \in C$ , where  $\tau$  is the time of one rank evaluation, and f(x,y) a polynomial. Seymour [17] showed that the class of graphic

<sup>\*</sup>Parts of this research have appeared in the third author's PhD thesis [24]. The research of all authors was partially supported by a grant from the Marsden Fund of New Zealand. The first author was also supported by a FRST Science & Technology post-doctoral fellowship. The third author was also supported by NWO, grant 613.000.561.

<sup>†</sup>School of Mathematics, Statistics and Operations Research, Victoria University of Wellington, New Zealand. E-mail: Dillon.Mayhew@msor.vuw.ac.nz,Geoff.Whittle@msor.vuw.ac.nz

<sup>&</sup>lt;sup>‡</sup>Centrum Wiskunde en Informatica, Postbus 94079, 1090 GB Amsterdam, The Netherlands. E-mail: Stefan.van.Zwam@cwi.nl

matroids is polynomial-time recognizable. Also every finite class is polynomial-time recognizable. Using these facts Truemper [18] (see also Schrijver [14, Chapter 20]) showed the following:

**Theorem 1.2.** The class of regular matroids is polynomial-time recognizable.

A near-regular matroid is a matroid representable over every field, except possibly GF(2). Near-regular matroids were introduced by Whittle [19, 20]. The following is one of his results:

**Theorem 1.3** (Whittle [20]). Let M be a matroid. The following are equivalent:

- i. M is representable over GF(3), GF(4), and GF(5);
- ii. M is representable over  $\mathbb{Q}(\alpha)$  by a totally near-unimodular matrix:
- iii. M is near-regular.

In this theorem  $\alpha$  is an indeterminate. A totally near-unimodular matrix is a matrix over  $\mathbb{Q}(\alpha)$  such that the determinant of every square submatrix is either zero or equal to  $(-1)^s \alpha^i (1-\alpha)^j$  for some  $s,i,j\in\mathbb{Z}$ . Whittle [20, 21] wondered if an analogue of Theorem 1.1 would hold for the class of near-regular matroids. The following conjecture was made:

Conjecture 1.4. Let M be a near-regular matroid. Then M can be obtained from matroids that are signed-graphic, their duals, or members of some finite set through 1-, 2-, and 3-sums.

A matroid is *signed-graphic* if it can be represented by a GF(3)-matrix with at most two nonzero entries in each column (see Zaslavsky [22, 23] for more on these matroids). One difference with the regular case is that not every signed-graphic matroid is near-regular.

Several people have made an effort to understand the structure of near-regular matroids. Oxley et al. [7] studied maximum-sized near-regular matroids. Hliněný [5] and Pendavingh [10] have both written software to investigate all 3-connected near-regular matroids up to a certain size. Pagano [9] studied signed-graphic near-regular matroids, and Pendavingh and Van Zwam [11] studied a closely related class of matroids which they call near-regular-graphic.

Despite these efforts, an analogue to Theorem 1.1 is still not in sight. In this paper we record an obstacle we found, that will have to be taken into account in any structure theorem. Our result is the following:

**Theorem 1.5.** Let  $G_1, G_2$  be graphs. There exists an internally 4-connected near-regular matroid M having both  $M(G_1)$  and  $M(G_2)^*$  as a minor.

From this, and the fact that not all cographic matroids are signed-graphic, it follows that Conjecture 1.4 is false. More generally, suppose we want to find a decomposition theorem for near-regular matroids, such that each basic class that contains all graphic matroids, does

not contain all cographic matroids. Theorem 1.5 implies that such a characterization must employ at least 4-sums.

The paper is organized as follows. In Section 2 we give some preliminary definitions. In Section 3 we prove a lemma that shows how generalized parallel connection can preserve representability over a partial field. In Section 4 we prove Theorem 1.5. We conclude in Section 5 with some updated conjectures.

Throughout this paper we assume familiarity with matroid theory as set out in Oxley [8].

## 2 Preliminaries

# 2.1 Connectivity

In addition to the usual definitions of connectivity and separations (see Oxley [8, Chapter 8]) we say a partition (A, B) of the ground set of a matroid is k-separating if  $\operatorname{rk}_M(A) + \operatorname{rk}_M(B) - \operatorname{rk}(M) < k$ . Recall that (A, B) is a k-separation if it is k-separating and  $\min\{|A|, |B|\} \geq k$ .

**Definition 2.1.** A matroid is *internally 4-connected* if it is 3-connected and  $\min\{|X|, |Y|\} = 3$  for every 3-separation (X, Y).

This notion of connectivity is useful in our context. For instance, Theorem 1.1 can be rephrased as follows:

**Theorem 2.2.** Let M be an internally 4-connected regular matroid. Then M is graphic, cographic, or equal to  $R_{10}$ .

Intuitively, separations (X, Y) where both |X| and |Y| are big should give rise to a decomposition into smaller matroids.

**Definition 2.3.** Let M be a matroid, and N a minor of M. Let (X', Y') be a k-separation of N. We say that (X', Y') is *induced* in M if M has a k-separation (X, Y) such that  $X' \subseteq X$  and  $Y' \subseteq Y$ .

At several points we will use the following easy fact:

**Lemma 2.4.** Let M be a matroid, let N be a minor of M, and let (A,B) be a k-separating partition of E(M). Then  $(A \cap E(N), B \cap E(N))$  is k-separating in N.

Note that  $(A \cap E(N), B \cap E(N))$  need not be exactly k-separating.

# 2.2 Partial fields

Our main tool in the proof of Theorem 1.5 is useful outside the scope of this paper. Hence we have stated it in the general framework of partial fields. For that purpose we need a few definitions. More on the theory of partial fields can be found in Semple and Whittle [15] and in Pendavingh and Van Zwam [13, 12].

**Definition 2.5.** A partial field is a pair (R, G), where R is a commutative ring with identity, and G is a subgroup of the group of units of R such that  $-1 \in G$ .

For example, the near-regular partial field is  $(\mathbb{Q}(\alpha), \langle -1, \alpha, 1-\alpha \rangle)$ , where  $\langle S \rangle$  denotes the multiplicative group generated by S. For  $\mathbb{P} = (R, G)$ , we abbreviate  $p \in G \cup \{0\}$  to  $p \in \mathbb{P}$ .

We will adopt the convention that matrices have labelled rows and columns, so an  $X \times Y$  matrix A is a matrix whose rows are labelled by the (ordered) set X and whose columns are labelled by the (ordered) set Y. The identity matrix with rows and columns labelled by X will be denoted by  $I_X$ . We will omit the subscript if it can be deduced from the context.

Let A be an  $X \times Y$  matrix. If  $X' \subseteq X$  and  $Y' \subseteq Y$  then we denote the submatrix of A indexed by X' and Y' by A[X',Y']. If  $Z \subseteq X \cup Y$  then we write  $A[Z] := A[X \cap Z, Y \cap Z]$ . If A is an  $X \times Y$  matrix, where  $X \cap Y = \emptyset$ , then we denote by  $[I \ A]$  the  $X \times (X \cup Y)$  matrix obtained from A by prepending the identity matrix  $I_X$ .

**Definition 2.6.** Let  $\mathbb{P} := (R, G)$  be a partial field, and let A be a matrix with entries in R. Then A is a  $\mathbb{P}$ -matrix if, for every square submatrix A' of A, either  $\det(A') = 0$  or  $\det(A') \in G$ .

**Theorem 2.7.** Let  $\mathbb{P}$  be a partial field, let A be an  $X \times Y$   $\mathbb{P}$ -matrix for disjoint sets X and Y, let  $E := X \cup Y$ , and let  $A' := [I \ A]$ . If  $\mathcal{B} = \{B \subseteq E : |B| = |X|, \det(A'[X, B]) \neq 0\}$ , then  $\mathcal{B}$  is the set of bases of a matroid.

We denote this matroid by M[I A].

#### 2.3 Pivoting

Let A be an  $X \times Y$   $\mathbb{P}$ -matrix. Then X is a basis of  $M[I\ A]$ . We say that X is the displayed basis. Pivoting in the matrix allows us to change the basis that is displayed. Roughly speaking a pivot in A consists of row reduction applied to  $[I\ A]$ , followed by a column exchange. The precise definition is as follows:

**Definition 2.8.** Let A be an  $X \times Y$  matrix over a ring R, and let  $x \in X, y \in Y$  be such that  $A_{xy} \in R^*$ . Then  $A^{xy}$  is the  $(X-x) \cup y \times (Y-y) \cup x$  matrix with entries

$$(A^{xy})_{uv} = \begin{cases} (A_{xy})^{-1} & \text{if } uv = yx\\ (A_{xy})^{-1}A_{xv} & \text{if } u = y, v \neq x\\ -A_{uy}(A_{xy})^{-1} & \text{if } v = x, u \neq y\\ A_{uv} - A_{uy}(A_{xy})^{-1}A_{xv} & \text{otherwise.} \end{cases}$$

We say that  $A^{xy}$  was obtained from A by *pivoting*. Slightly less opaquely, if

$$A = \begin{pmatrix} x & y & Y' \\ x & a & c \\ b & D \end{pmatrix}$$

then

$$A^{xy} = \begin{bmatrix} x & Y' \\ a^{-1} & a^{-1}c \\ -ba^{-1} & D - ba^{-1}c \end{bmatrix}.$$

As Semple and Whittle [15] proved, pivoting maps  $\mathbb{P}\text{-matrices}$  to  $\mathbb{P}\text{-matrices}:$ 

**Proposition 2.9.** Let A be an  $X \times Y$   $\mathbb{P}$ -matrix, and let  $x \in X, y \in Y$  be such that  $A_{xy} \neq 0$ . Then  $A^{xy}$  is a  $\mathbb{P}$ -matrix, and  $M[I \ A] = M[I \ A^{xy}]$ .

Semple and Whittle also showed that pivots can be used to compute determinants of  $\mathbb{P}$ -matrices:

**Lemma 2.10.** Let  $\mathbb{P}$  be a partial field, and let A be an  $X \times Y$   $\mathbb{P}$ -matrix with |X| = |Y|. If  $x \in X, y \in Y$  is such that  $A_{xy} \neq 0$  then

$$\det(A) = (-1)^{x+y} A_{xy} \det(A^{xy} [X - x, Y - y]).$$

# 3 Generalized parallel connection

Recall the generalized parallel connection of two matroids  $M_1$ ,  $M_2$  along a common restriction N, denoted by  $P_N(M_1,M_2)$ . This construction was introduced by Brylawski [1] (see also Oxley [8, Section 12.4]). Brylawski proved that representability over a field can be preserved under generalized parallel connection, provided that the representations of the common minor are identical. Lee [6] generalized Brylawski's result to matroids representable over a field such that all subdeterminants are in a multiplicatively closed set. We generalize Brylawski's result further to matroids representable over a partial field, as follows.

**Theorem 3.1.** Suppose  $A_1$ ,  $A_2$  are  $\mathbb{P}$ -matrices with the following structure:

$$A_1 = \frac{X_1}{X} \begin{bmatrix} Y_1 & Y \\ D_1' & 0 \\ D_1 & D_X \end{bmatrix}, \qquad A_2 = \frac{X}{X_2} \begin{bmatrix} Y & Y_2 \\ D_X & D_2 \\ 0 & D_2' \end{bmatrix},$$

where  $X, Y, X_1, Y_1, X_2, Y_2$  are pairwise disjoint sets. If  $X \cup Y$  is a modular flat of  $M[I \ A_1]$  then

$$A := \begin{array}{ccc} X_1 & Y & Y_2 \\ X_1 & D_1' & 0 & 0 \\ D_1 & D_X & D_2 \\ 0 & 0 & D_2' \end{array} \right]$$

is a  $\mathbb{P}$ -matrix. Moreover, if  $M_1 = M[I \ A_1]$  and  $M_2 = M[I \ A_2]$ , then  $M[I \ A] = P_N(M_1, M_2)$ , where  $N = M[I \ D_X]$ .

The main difficulty is to show that A is a  $\mathbb{P}$ -matrix. To prove this we will use a result known as the modular short-circuit axiom [1, Theorem 3.11]. We use Oxley's formulation [8, Theorem 6.9.9], and refer to that book for a proof.

**Lemma 3.2.** Let M be a matroid and  $X \subseteq E$  nonempty. The following statements are equivalent:

- i. X is a modular flat of M;
- ii. For every circuit C such that  $C X \neq \emptyset$ , there is an element  $x \in X$  such that  $(C X) \cup x$  is dependent.
- iii. For every circuit C, and for every  $e \in C X$ , there are an  $f \in X$  and a circuit C' such that  $e \in C'$  and  $C' \subseteq (C X) \cup f$ .

The following is an extension of Proposition 4.1.2 in [1] to partial fields. Note that Brylawski proves an "if and only if" statement, whereas we only state the "only if" direction.

**Lemma 3.3.** Let  $M = (E, \mathcal{I})$  be a matroid, and X a modular flat of M. Suppose  $B_X$  is a basis for M|X, and  $B \supseteq B_X$  a basis of M. Suppose A is a  $B \times (E - B)$   $\mathbb{P}$ -matrix such that M = M[I A]. Then every column of  $A[B_X, E - (B \cup X)]$  is a  $\mathbb{P}$ -multiple of a column of  $[I A[B_X, X - B]]$ .

Proof of Lemma 3.3. Let  $M, X, B_X, B, A$  be as in the lemma, so

$$A = {B - B_X \atop B_X} \left[ \begin{array}{cc} E - (B \cup X) & X - B \\ D' & 0 \\ D & D_{B_X} \end{array} \right].$$

Let  $v \in E-(B \cup X)$ , and let C be the B-fundamental circuit containing v. If  $C \cap X = \emptyset$  then  $D[B_X, v]$  is an all-zero vector and the result holds, so assume  $B_X \cap C \neq \emptyset$ . By Lemma 3.2(iii) there are an  $x \in X$  and a circuit C' with  $v \in C'$  and  $C' \subseteq (C - X) \cup x$ .

Let  $M' := M/(B - B_X)$ . Then  $C' \cap E(M') = \{v, x\}$  is a circuit of M'. Hence all  $2 \times 2$  subdeterminants of  $[I \ A][B_X, \{v, x\}]$  have to be 0, which implies that  $A[B_X, v]$  is the all-zero vector or parallel to  $[I \ A][B_X, x]$ .

Proof of Theorem 3.1. Let  $A_1$ ,  $A_2$ , A be as in the theorem, and define  $E := X_1 \cup X_2 \cup X \cup Y_1 \cup Y_2 \cup Y$ . Suppose there exists a  $Z \subseteq E$  such that A[Z] is square, yet  $\det(A[Z]) \notin \mathbb{P}$ . Assume  $A_1, A_2, A, Z$  were chosen so that |Z| is as small as possible.

If  $Z \subseteq X_i \cup Y_i \cup X \cup Y$  for some  $i \in \{1,2\}$  then A[Z] is a submatrix of  $A_i$ , a contradiction. Therefore we may assume that Z meets both  $X_1 \cup Y_1$  and  $X_2 \cup Y_2$ . We may also assume that A[Z] contains no row or column with only zero entries, so either there are  $x \in X_1 \cap Z$ ,  $y \in Y_1 \cap Z$  with  $A_{xy} \neq 0$  or  $x \in X \cap Z$ ,  $y \in Y_1 \cap Z$  with  $A_{xy} \neq 0$ .

In the former case, pivoting over xy leaves  $D_X$ ,  $D_2$ , and  $D_2'$  unchanged, yet by Lemma 2.10  $\det(A[Z]) \in \mathbb{P}$  if and only if  $\det(A^{xy}[Z - \{x,y\}]) \in \mathbb{P}$ . This contradicts minimality of |Z|. Therefore  $Z \cap X_1 = \emptyset$ . Similarly,  $Z \cap X_2 = \emptyset$ .

Define  $X':=Z\cap X$ . Now pick some  $y\in Y_1$ . Since  $A[X',Y_1\cup Y]$  is obtained from  $A[X,Y_1\cup Y]$  by deleting rows, it follows from Lemma 3.3, applied to  $M[I\ A_1]$ , that the column A[X',y] is either a unit vector (i.e. a column of an identity matrix) or parallel to A[X',y'] for some  $y'\in Y$ . In the first case, Lemma 2.10 implies again that  $\det(A[Z])\in \mathbb{P}$  if and only if  $\det(A[Z-\{x,y\}])\in \mathbb{P}$ , contradicting minimality of |Z|. In the second case, if  $y'\in Z$  then  $\det(A[Z])=0$ . Otherwise we can replace y by y' without changing  $\det(A[Z])$  (up to possible multiplication with some nonzero  $p\in \mathbb{P}$ ). It follows that  $\det(A[Z])=p'\det(A[Z'])$ , where  $Z'\subseteq X\cup Y\cup Y_2$ , and  $p'\in \mathbb{P}-\{0\}$ . But  $\det(A[Z'])\in \mathbb{P}$ , so also  $\det(A[Z])\in \mathbb{P}$ , a contradiction.

It remains to prove that  $M[I\ A]=P_N(M_1,M_2)$ . Suppose  $\mathbb{P}=(R,G)$ , and let I be a maximal ideal of R. Let  $\varphi:R\to R/I$  be the canonical ring homomorphism. For a square  $\mathbb{P}$ -matrix D we have  $\det(D)=0$  if and only if  $\det(\varphi(D))=0$ . Hence  $M[I\ A]=M[I\ \varphi(A)]$ . But R/I is a field, so the result now follows directly from Brylawski's original theorem.

The special cases  $X = \emptyset$  and  $X = \{p\}$  were previously proven by Semple and Whittle [15].

# 4 The need for 4-sums

The core of the proof of Theorem 1.5 will be a special matroid  $M_{12} := M[I \ A_{12}]$ , where

$$A_{12} = \begin{bmatrix} a & c & f & 4 & 5 & 6 \\ 1 & 0 & 1 & 1 & 1 & 0 \\ 0 & -1 & 1 & 1 & 0 & \alpha \\ 1 & 1 & 0 & 0 & \alpha & -\alpha \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 2 & 0 & 0 & 0 & 1 & -1 \\ 3 & 0 & 0 & 0 & 1 & 1 & 0 \end{bmatrix}.$$
(1)

**Lemma 4.1.** The following hold:

- i.  $M_{12}$  is near-regular;
- ii.  $M_{12}$  is internally 4-connected;
- iii.  $M_{12}$  is self-dual;
- iv.  $M_{12} \setminus \{1, 2, 3, 4, 5, 6\} \cong M(K_4)$ ;
- $v. M_{12}/\{a, b, c, d, e, f\} \cong M(K_4);$
- vi. No triad of  $M_{12} \setminus \{1, 2, 3, 4, 5, 6\}$  is a triad of  $M_{12}$ .

We will omit the proofs, each of which boils down to a finite case check that is easily done on a computer and not too onerous by hand. Specifically, for the first property one can either verify that  $A_{12}$  is totally near-unimodular, or that  $M_{12}$  contains none of the excluded minors for near-regular matroids (see Hall et al. [4]). The latter approach is facilitated by observing that  $M_{12}$  is the signed-graphic matroid associated with the signed graph illustrated in Figure 1. That




Figure 1: Signed-graphic representation of  $M_{12}$ . Negative edges are dashed; positive edges are solid.

graph can also be used to verify (ii), by examining all edge-partitions (A, B) that meet in two or three vertices. The remaining properties are readily extracted from the matrix  $A_{12}$ .

We will use the  $M(K_4)$ -restriction to create the generalized parallel connection of  $M_{12}$  with  $M(K_n)$ . The following is well-known, but we will include the short proof.

**Lemma 4.2.** The matroid  $M(K_n)$  is internally 4-connected.

Proof. Fix an integer n, and suppose (A, B) is a 3-separation of  $M(K_n)$  with  $|A|, |B| \geq 4$ . It follows that  $n \geq 5$ . Assume that  $\mathrm{rk}(A) \geq \mathrm{rk}(B)$ . Note that  $\mathrm{cl}(A)$  and  $\mathrm{cl}(B)$  induce complete subgraphs of  $K_n$ , and that these subgraphs meet in at most three vertices. It follows that, for some vertex v of  $K_n$ , all edges incident with v are in A, or all edges are in B. Assume the former. Then  $\mathrm{cl}(A) = E(K_n)$ , and therefore  $\mathrm{rk}(A) = n - 1$ , and  $\mathrm{rk}(B) = 2$ . But then B is a subset of a triangle of  $K_n$ , a contradiction.

We need to show that in forming the generalized parallel connection we do not introduce unwanted 3-separations. The following lemma takes care of this.

**Lemma 4.3.** Let  $M_1 = M(K_n)$  for some  $n \ge 5$ , and  $M_2$  an internally 4-connected matroid such that there is a set  $X = E(M_1) \cap E(M_2)$  with  $N := M_1 | X = M_2 | X \cong M(K_4)$ . Then  $M := P_N(M_1, M_2)$  is a well-defined matroid. If no triad of N is a triad of  $M_2$  then M is internally 4-connected.

Proof. It is well-known (see [8, Page 236]) that N is a modular flat of  $M_1$ . Hence  $M=P_N(M_1,M_2)$  is well-defined. It remains to prove that M is internally 4-connected. Suppose not. M is obviously connected. Suppose (A,B) is a 2-separation of M. By relabelling we may assume  $|A\cap E(M_1)|\geq |B\cap E(M_1)|$ . By Lemma 2.4 we have that  $(A\cap E(M_1),B\cap E(M_1))$  is 2-separating in  $M_1$  (since  $M_1$  is a restriction of M). But  $M_1$  is 3-connected, so  $|B\cap E(M_1)|\leq 1$ . Similarly we have either  $|A\cap E(M_2)|\leq 1$  or  $|B\cap E(M_2)|\leq 1$ . Since  $|E(M_1)\cap E(M_2)|=6$ , the latter must hold. Hence  $B=\{e,f\}$  for some  $e\in E(M_1)-E(N)$  and  $f\in E(M_2)-E(N)$ . Since  $E(M_1)$  and  $E(M_2)$ 

are flats of M, we have  $\operatorname{rk}_M(\{e,f\}) = 2$ . Moreover  $e \in \operatorname{cl}_M(E(M_1) - e)$  and  $f \in \operatorname{cl}_M(E(M_2) - f)$ , so  $\{e,f\} \subseteq \operatorname{cl}_M(A)$ . But then

$$\operatorname{rk}_{M}(A) + \operatorname{rk}_{M}(B) - \operatorname{rk}(M) = \operatorname{rk}_{M}(B) = 2, \tag{2}$$

contradicting the fact that (A, B) is a 2-separation.

Next suppose that (A, B) is a 3-separation of M with  $|A| \ge 4$  and  $|B| \ge 4$ . By relabelling we may assume  $|A \cap E(M_1)| \ge |B \cap E(M_1)|$ . By Lemma 2.4 again,  $(A \cap E(M_1), B \cap E(M_1))$  is 3-separating in  $M_1$ . Since  $M_1$  is internally 4-connected,  $|B \cap E(M_1)| \le 3$ . Define  $T := B \cap E(M_1)$ .

We will show that  $T \subseteq \operatorname{cl}_M(B-T)$ . Since  $M_1$  has no cocircuits of size less than 4, we have  $T \subseteq \operatorname{cl}_M(A)$ . Therefore

$$\operatorname{rk}_{M}(A \cup T) + \operatorname{rk}_{M}(B - T) - \operatorname{rk}(M) = \operatorname{rk}_{M}(A) + \operatorname{rk}_{M}(B - T) - \operatorname{rk}(M)$$

$$\leq \operatorname{rk}_{M}(A) + \operatorname{rk}_{M}(B) - \operatorname{rk}(M) = 2.$$
(3)

If  $|B-T| \ge 2$  then it follows from 3-connectivity that equality holds in (3), so  $\operatorname{rk}_M(B) = \operatorname{rk}_M(B-T)$ . If |B-T| = 1 then  $\operatorname{rk}_M(B-T) = 1$  and we must have  $\operatorname{rk}_M(B) = 2$ . In that case T is a triangle of  $M_1$  and some element  $e \in E(M_2) - E(M_1)$  is in the closure of T. But no such element e exists, since  $E(M_1)$  is a flat of M.

Note that  $B-T\subseteq E(M_2)$ . Since  $T\subseteq \operatorname{cl}_M(B-T)$  and  $E(M_2)$  is a flat of M, we have that  $T\subseteq E(M_2)$ . Hence  $T\subseteq E(N)$ , and  $B\cap E(M_2)=B$ . Since  $(A\cap E(M_2),B\cap E(M_2))$  is 3-separating and  $|B\cap E(M_2)|=|B|\geq 4$ , we have  $|A\cap E(M_2)|\leq 3$ . But  $|B\cap E(M_1)|\leq 3$ , and therefore  $E(N)-B\subseteq A\cap E(M_2)$ , from which it follows that  $|A\cap E(M_2)|\geq 3$ .

Since no triad of N is a triad of  $M_2$ , we must have that  $A \cap E(M_2)$  is a triangle of  $M_2$ . Hence  $B \cap E(N)$  is a triad of N. Now consider  $(A \cap E(M_1), B \cap E(M_1))$  again. This partition of  $M_1$  must be 3-separating, but  $B \cap E(M_1)$  is not a triangle of  $M_1$ , and  $M_1$  has no 3-element cocircuits. This contradiction completes the proof.

Proof of Theorem 1.5. It suffices to prove the theorem for  $G_1 = G_2 = K_n$ , where  $n \geq 5$ . Label the edges of some  $K_4$ -restriction  $N_1$  of  $G_1$  by  $\{a, b, c, d, e, f\}$ , and define

$$M' := (P_{N_1} (M(G_1), M_{12}))^*. (4)$$

By Theorem 3.1, M' is near-regular, and by Lemma 4.3, M' is internally 4-connected.

Note that we still have  $M'|\{1,2,3,4,5,6\} \cong M(K_4)$ . Label the edges of some  $K_4$ -restriction  $N_2$  of  $G_2$  by  $\{1,2,3,4,5,6\}$ , and define

$$M := P_{N_2}(M(G_2), M'). (5)$$

By Theorem 3.1, M is near-regular, and by Lemma 4.3, M is internally 4-connected. The result follows.

Matroid  $M_{12}$  was found while studying the 3-separations of  $R_{12}$ . The unique 3-separation (X,Y) of  $R_{12}$  with |X| = |Y| = 6 is induced in the class of regular matroids. Pendavingh and Van Zwam had found, using a computer search for blocking sequences, that it is not induced in the class of near-regular matroids.

Unlike  $R_{10}$  and  $R_{12}$  in Seymour's work, the matroid  $M_{12}$  by itself is quite inconspicuous. A natural class of near-regular matroids is the class of near-regular signed-graphic matroids. As indicated earlier,  $M_{12}$  is a member of this class (see Figure 1). The  $K_4$ -restriction is readily identified.  $M_{12}$  is self-dual and has an automorphism group of size 6, generated by (c, e)(d, f)(1, 5)(3, 6) and (a, d)(b, e)(1, 4)(2, 3).

# 5 Conjectures

While Theorem 1.5 is a bit of a setback, we remain hopeful that a satisfactory decomposition theory for near-regular matroids can be found. First of all, the construction in Section 4 employs only graphic matroids. In fact, it seems difficult to extend the  $M(G_1)$ -restriction of the 4-sum to some strictly near-regular matroid. The proof of Theorem 1.5 suggests the following construction:

**Definition 5.1.** Let  $M_1, M_2$  be matroids such that  $E(M_1) \cap E(M_2) = X$ ,  $N := M_1|X = M_2|X \cong M(K_k)$ , and  $M_1$  is graphic. Then the graph k-clique sum of  $M_1$  and  $M_2$  is  $P_N(M_1, M_2) \setminus X$ .

Now we offer the following update of Conjecture 1.4:

Conjecture 5.2. Let M be a near-regular matroid. Then M can be obtained from matroids that are signed-graphic, are the dual of a signed-graphic matroid, or are members of a finite set C, by applying the following operations:

- i. 1-, 2-, and 3-sums;
- ii. Graph k-clique sums and their duals, where  $k \leq 4$ .

Note that the work of Geelen et al. [3], when finished, should imply a decomposition into parts that are bounded-rank perturbations of signed-graphic matroids and their duals. However, the bounds they require on connectivity are huge. Conjecture 5.2 expresses our hope that for near-regular matroids specialized methods will give much more refined results.

As noted in the introduction, Seymour's Decomposition Theorem is not the only ingredient in the proof of Theorem 1.2. Another requirement is that the basic classes can be recognized in polynomial time. The following result suggests that this may not hold for the basic classes of near-regular matroids:

**Theorem 5.3.** Let M be a signed-graphic matroid. Let N be a matroid on E(M) given by a rank oracle. It is not possible to decide if M=N using a polynomial number of rank evaluations.

A matroid is *dyadic* if it is representable over GF(p) for all primes p > 2. Since all signed-graphic matroids are dyadic (which was first observed by Dowling [2]), this in turn implies that dyadic matroids are not polynomial-time recognizable.

A proof of Theorem 5.3, analogous to the proof by Seymour [17] that binary matroids are not polynomial-time recognizable, was found by Jim Geelen and, independently, by the first author. It involves ternary swirls, which have a number of circuit-hyperplanes that is exponential in the rank. To test if the matroid under consideration is really the ternary swirl, all these circuit-hyperplanes have to be examined, since relaxing any one of them again yields a matroid.

However, this family of signed-graphic matroids is not near-regular for all ranks greater than 3. Hence the complexity of recognizing near-regular signed-graphic matroids is still open. The techniques used by Seymour [17] do not seem to extend, but perhaps some new idea can yield a proof of the following conjecture:

Conjecture 5.4. Let C be the class of near-regular signed-graphic matroids. Then C is polynomial-time recognizable.

In fact, we still have some hope for the following:

Conjecture 5.5. The class of near-regular matroids is polynomialtime recognizable.

**Acknowledgements** We thank the anonymous referee for many useful suggestions. The third author thanks Rudi Pendavingh for introducing him to matroid theory in general, and to the problem of decomposing near-regular matroids in particular.

# References

- [1] T. Brylawski, Modular constructions for combinatorial geometries, Trans. Amer. Math. Soc., 203 (1975), pp. 1–44.
- [2] T. A. DOWLING, A class of geometric lattices based on finite groups, J. Combin. Theory Ser. B, 14 (1973), pp. 61–86.
- [3] J. GEELEN, B. GERARDS, AND G. WHITTLE, Towards a matroid-minor structure theory, in Combinatorics, Complexity, and Chance, vol. 34 of Oxford Lecture Ser. Math. Appl., Oxford University Press, 2007, pp. 72–82.
- [4] R. Hall, D. Mayhew, and S. H. M. van Zwam, The excluded minors for near-regular matroids, European J. Combin., (2010). Accepted. Preprint at arXiv:0902.2071v2 [math.CO].
- [5] P. HLINĚNÝ, Using a computer in matroid theory research, Acta Univ. M. Belii Ser. Math., (2004), pp. 27–44.
- [6] J. Lee, The incidence structure of subspaces with well-scaled frames, J. Combin. Theory Ser. B, 50 (1990), pp. 265–287.
- [7] J. OXLEY, D. VERTIGAN, AND G. WHITTLE, On maximumsized near-regular and  $\sqrt[6]{1}$ -matroids, Graphs and Combinatorics, 14 (1998), pp. 163–179.
- [8] J. G. Oxley, Matroid Theory, Oxford University Press, 1992.
- [9] S. R. PAGANO, Separability and Representability of Bias Matroids of Signed Graphs, PhD thesis, State University of New York at Binghamton, 1998.
- [10] R. A. Pendavingh. Personal communication, 2004.
- [11] R. A. Pendavingh and S. H. M. van Zwam, Recognizing near-regular-graphic matroids. In preparation.
- [12] ——, Confinement of matroid representations to subsets of partial fields, J. Combin. Theory Ser. B, 100 (2010), pp. 510–545.
- [13] —, Lifts of matroid representations over partial fields, J. Combin. Theory Ser. B, 100 (2010), pp. 36–67.
- [14] A. Schrijver, Theory of Linear and Integer Programming, John Wiley & Sons, 1986.
- [15] C. SEMPLE AND G. WHITTLE, Partial fields and matroid representation, Adv. in Appl. Math., 17 (1996), pp. 184–208.
- [16] P. D. SEYMOUR, Decomposition of regular matroids, J. Combin. Theory Ser. B, 28 (1980), pp. 305–359.

- [17] P. D. SEYMOUR, *Recognizing graphic matroids*, Combinatorica, 1 (1981), pp. 75–78.
- [18] K. TRUEMPER, On the efficiency of representability tests for matroids, European J. Combin., 3 (1982), pp. 275–291.
- [19] G. Whittle, A characterisation of the matroids representable over GF(3) and the rationals, J. Combin. Theory Ser. B, 65 (1995), pp. 222–261.
- [20] G. WHITTLE, On matroids representable over GF(3) and other fields, Trans. Amer. Math. Soc., 349 (1997), pp. 579–603.
- [21] —, Recent work in matroid representation theory, Discrete Math., 302 (2005), pp. 285–296.
- [22] T. Zaslavsky,  $Signed\ graphs,$  Discrete Appl. Math., 4 (1982), pp. 47–74.
- [23] —, Erratum: "Signed graphs", Discrete Appl. Math., 5 (1983), p. 248.
- [24] S. H. M. VAN ZWAM, Partial Fields in Matroid Theory, PhD thesis, Technische Universiteit Eindhoven, 2009.