The Polyhedron-Hitting Problem^{*}

Ventsislav Chonev Department of Computer Science Oxford University, UK Joël Ouaknine Department of Computer Science Oxford University, UK

James Worrell Department of Computer Science Oxford University, UK

Abstract

We consider polyhedral versions of Kannan and Lipton's Orbit Problem [14, 13]—determining whether a target polyhedron V may be reached from a starting point x under repeated applications of a linear transformation A in an ambient vector space \mathbb{Q}^m . In the context of program verification, very similar reachability questions were also considered and left open by Lee and Yannakakis in [15], and by Braverman in [4]. We present what amounts to a complete characterisation of the decidability landscape for the Polyhedron-Hitting Problem, expressed as a function of the dimension mof the ambient space, together with the dimension of the polyhedral target V: more precisely, for each pair of dimensions, we either establish decidability, or show hardness for longstanding number-theoretic open problems.

1 Introduction

Given a linear transformation A over the vector space \mathbb{Q}^m , together with a starting point x, the **orbit** of x under A is the infinite sequence $\langle x, Ax, A^2x, \ldots, A^jx, \ldots \rangle$. A natural decision problem in discrete linear dynamical systems is whether the orbit of x ever hits a particular target set V.

An early instance of this problem was raised by Harrison in 1969 [12] for the special case in which Vis simply a point in \mathbb{Q}^m . Decidability remained open for over ten years, and was finally settled in a seminal paper of Kannan and Lipton, who moreover gave a polynomial-time decision procedure [14]. In subsequent work [13], Kannan and Lipton noted that the Orbit Problem becomes considerably harder when the target V is replaced by a subspace of \mathbb{Q}^m : indeed, if V has dimension m-1, the problem is equivalent to the *Skolem Problem*, known to be **NP**-hard but whose decidability has remained open for over 80 years [23]. Nevertheless, Kannan and Lipton speculated in [13] that instances of the Orbit Problem with low-dimensional subspaces as target might be decidable. This was finally substantiated in [6], which showed decidability for vector-space targets of dimension at most 3, with polynomial-time complexity for one-dimensional targets, and complexity in **NP**^{**RP**} for two- and three-dimensional targets.

In this paper, we study a natural generalisation of the Orbit Problem, which we call the **Polyhedron-Hitting Problem**, in which the target V is allowed to be an arbitrary (bounded or unbounded) polyhedron.¹ We present what amounts to a complete characterisation of the decidability landscape for this problem, expressed as a function of the dimension m of the ambient space \mathbb{Q}^m , together with the dimension k of the polyhedral target V; more precisely, for each pair of dimensions, we either establish decidability, or show hardness for longstanding number-theoretic open problems. Our results are summarised in Fig. 1. As our algorithms rely on symbolic manipulation of algebraic numbers of unbounded degree and height, all decidable instances lie in **PSPACE**.

A key motivation for studying the Polyhedron-Hitting Problem comes from the area of program verification, and in particular the problem of determining whether a simple while loop with linear (or affine) assignments and guards will terminate or not. Very similar reachability questions were considered and left open by Lee and Yannakakis in [15] for what they termed "real affine transition systems". Similarly, decidability for the special case of the Polyhedron-Hitting Problem in which the polyhedral target consists of a *single* halfspace (rather than an intersection of several halfspaces)

^{*}This research was partially supported by EPSRC.

¹This problem was also considered in [24] under the appellation of *Chamber-Hitting Problem*. However that paper focused on connections with formal language theory rather than on establishing decidability.

	m = 1	m=2	m = 3	m = 4	m = k	$m \ge k+1$
k = 0	Р	Р	Р	Р	Р	Р
k = 1	PSPACE	PSPACE	PSPACE	PSPACE	PSPACE	PSPACE
k = 2		PSPACE	PSPACE	PSPACE	PSPACE	PSPACE
k = 3			PSPACE	S_5	PSPACE	S_5
k = 4				D	D	$D \& S_5$
$k \ge 5$					D	$D \& S_{k+1}$

Figure 1: Upper and lower complexity bounds for instances of the Polyhedron-Hitting Problem in ambient dimension m with a k-dimensional target. The row k = 0 corresponds to Kannan and Lipton's Orbit Problem [14, 13]. Upper complexity bounds are denoted by **P** and **PSPACE**, indicating membership in these classes, whereas lower bounds are denoted by D (indicating reduction from certain Diophantine-approximation problems detailed precisely in Sec. 2.2) and S_d (indicating reduction from Skolem's Problem of order d, defined in Sec. 2.2).

was mentioned as an open problem by Braverman in Sec. 6 of [4].

It should be noted, however, that the problem considered in the present paper differs in one fundamental respect from what is traditionally termed the 'Termination Problem' in the program verification literature (see, e.g., [3]). The latter studies termination of while loops for *all* possible initial starting points (valuations of the variables), rather than for a *fixed* starting point as we consider in this paper. This distinction drastically transforms the nature of the problem at hand.

In [18], the traditional Termination Problem is solved over the integers for while loops under certain restrictions (chiefly, diagonalisability of the associated linear transformation). That paper relies on markedly different techniques from the present one, eschewing Baker's Theorem and relying instead on nonconstructive lower bounds on sums of S-units (which in turn follow from deep results in Diophantine approximation), as well as real algebraic geometry.

The present paper vastly extends our earlier results from [6], in which only vector-space targets were considered. Polyhedra, defined as intersections of (affine) halfspaces, pose substantial new challenges, as evidenced among others by the Diophantine-approximation lower bounds that arise for polyhedral targets of dimension 4 or greater. In addition to classical tools from algebraic and transcendental number theory such as Baker's Theorem, the present paper relies crucially on several tools not invoked in [6] or [18], including techniques from Diophantine approximation, convex geometry, as well as decision procedures for the existential fragment of the first-order theory of the reals.

In terms of future work, either establishing complexity lower bounds, or improving the **PSPACE** membership of the decidable problem instances, stand out as challenging open questions.

2 Polyhedron-Hitting Problem

The focus of this paper is the **Polyhedron-Hitting Problem**: given a square matrix $A \in \mathbb{Q}^{m \times m}$, a vector $x \in \mathbb{Q}^m$ and polyhedron P (represented as the intersection of halfspaces), determine whether there exists a natural number n such that $A^n x \in P$. We will denote by PHP(m, k) the version of the problem in which the ambient space is \mathbb{Q}^m and the target polyhedron has dimension $k \leq m$.

We begin this section with our decidability results for low-dimensional versions of the problem. We define two related problems to which we reduce the Polyhedron-Hitting Problem in order to obtain our complexity upper bounds: the Extended Orbit Problem and the Simultaneous Positivity Problem. Then we proceed to give hardness results for higher-dimensional cases by reducing from Skolem's Problem and from Diophantine approximation.

2.1 Decidability results Our effectiveness result on the Polyhedron-Hitting Problem is the following:

THEOREM 2.1. If $k \leq 2$ or m = k = 3, then PHP(m, k) is in **PSPACE**.

The strategy for PHP(m, k) when $k \leq 2$ is to reduce to the related **Extended Orbit Problem**: given a linear transformation $A \in \mathbb{Q}^{m \times m}$, a vector $x \in \mathbb{Q}^m$, a target \mathbb{Q} -vector space V defined by a basis $\{y_1, \ldots, y_d\} \subseteq \mathbb{Q}^m$ and a constraint matrix $B \in (\mathbb{R} \cap \mathbb{A})^{k \times d}$, determine whether there exists some exponent $n \in \mathbb{N}$ such that $A^n x \in V$ and the coordinates $u = (u_1, \ldots, u_d)^T$ of $A^n x$ with respect to the basis $\{y_1, \ldots, y_d\}$ satisfy $Bu \geq$ 0. This problem essentially specialises the target of PHP to a cone and assumes a particular parametric representation.

We focus first on PHP(m, 1). By Lemma A.2 in Appendix A.1, a one-dimensional polyhedron is of the form

$$P = \{v_1 + \alpha v_2 : \alpha \in I\}$$

where I is one of \mathbb{R} , [0, 1] and $[0, \infty)$. Moreover, this parametric representation is computable in polynomial time from the halfspace description of P. Now suppose we wish to find $n \in \mathbb{N}$ and $u_1, u_2 \in \mathbb{Q}$ such that

$$\begin{bmatrix} A & 0 \\ 0 & 1 \end{bmatrix}^n \begin{bmatrix} x \\ 1 \end{bmatrix} = u_1 \begin{bmatrix} v_1 \\ 1 \end{bmatrix} + u_2 \begin{bmatrix} v_2 \\ 0 \end{bmatrix}$$

The (m + 1)-th component forces any witness to this problem instance to have $u_1 = 1$. Therefore, requiring $u_2 \ge 0$ and $u_1 - u_2 \ge 0$ gives an Extended Orbit instance with a two-dimensional target space which is positive if and only if the segment $\{v_1 + u_2v_2 : u_2 \in [0,1]\}$ intersects the orbit $\{A^nx : n \in \mathbb{N}\}$. Requiring instead only $u_2 \ge 0$ gives the half-line $\{v_1 + u_2v_2 : u_2 \in [0,\infty)\}$, whereas setting no restriction gives the whole line $\{v_1 + u_2v_2 : u_2 \in \mathbb{R}\}$. In all cases, the resulting Extended Orbit instance has target space of dimension two, so by Theorem 3.1 in Section 3, PHP(m, 1) is in **PSPACE**.

Now we move to PHP(m, 2). By Lemma A.1 in Appendix A.1, any two-dimensional polyhedron can be decomposed into a finite union of simple shapes: $P = \bigcup_{i=1}^{s} S_i$ where

$$S_i = \{v_{i_1} + \alpha v_{i_2} + \beta v_{i_3} : \alpha \ge 0 \text{ and } \beta \ge 0 \text{ and } T(\alpha, \beta)\}$$

where the predicate T is either $\alpha + \beta \leq 1$, or $\beta \leq 1$ or true. In fact, it is easy to see from the proof of Lemma A.1 that $s \in 2^{||P||^{O(1)}}$. For each i, the problem of whether there exists n such that $A^n x \in S_i$ reduces to the Extended Orbit Problem with a three-dimensional target. For instance, if the predicate T_i is $\alpha + \beta \leq 1$, that is, S_i is a triangle, then $A^n x \in S_i$ if and only if there exist $u_1, u_2, u_3 \in \mathbb{Q}$ such that $u_2 \geq 0$, $u_3 \geq 0$, $u_1 - u_2 - u_3 \geq 0$ and

$$\begin{bmatrix} A & 0 \\ 0 & 1 \end{bmatrix}^n \begin{bmatrix} x \\ 1 \end{bmatrix} = \begin{bmatrix} v_{i_1} & v_{i_2} & v_{i_3} \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix}$$

As in the reduction from PHP(m, 1), the (m + 1)-th component forces $u_1 = 1$ and allows us to express the constraint $u_2 + u_3 \leq 1$ with a homogeneous inequality. The remaining possible choices of predicate T reduce similarly. By Theorem 3.1 in Section 3, the Extended Orbit Problem with target space of dimension three is in **PSPACE**. Therefore, to solve PHP(m, 2) in **PSPACE**, it suffices to choose nondeterministically a simple two-dimensional target S_i and proceed to solve an Extended Orbit instance.

Finally, consider the Polyhedron-Hitting Problem in the case when the target polyhedron P has dimension m, matching the dimension of the ambient space \mathbb{Q}^m . Consider the halfspace description of P:

$$P = \bigcap_{i=1}^{s} H_i = \bigcap_{i=1}^{s} \{ p \in \mathbb{Q}^m : v_i^T p \ge c_i \}$$

Define the linear recurrence sequences $S_i(n) = v_i^T A^n x$. By the Cayley-Hamilton Theorem, the sequences S_i satisfy a common recurrence equation with characteristic polynomial the minimal polynomial $f_A(x)$ of A. Define also the sequences $S'_i(n) = S_i(n) - c_i$. It is not difficult to show that the latter also satisfy a common recurrence equation, with characteristic polynomial $(x - 1)f_A(x)$. Since f_A has degree at most m, the order of the recurrence equation shared by the sequences $S'_i(n)$ is at most m+1. Moreover $A^n x \in P$ iff $S'_i(n) \ge 0$ for $i = 1, \ldots, s$.

Thus, PHP(m,m) reduces to the **Simultaneous Positivity Problem**: given a family of linear recurrence sequences $S'_i(n)$, i = 1, ..., s, which satisfy a common recurrence relation of order m + 1, does there exist an index n such that $S'_i(n) \ge 0$ for all i? This problem is the focus of Section 4, where we place it in **PSPACE** in the case of LRS over $\mathbb{R} \cap \mathbb{A}$ whose shared recurrence relation is of order at most three, or of order four but with 1 as a characteristic root. This immediately shows that PHP(3,3) is in **PSPACE**, completing the proof of Theorem 2.1.²

2.2 Hardness results Now we proceed to give hardness results for the Polyhedron-Hitting Problem. First, observe that lower-dimensional versions of *PHP* reduce to higher-dimensional ones:

LEMMA 2.1. For all m, k such that $m \ge k$, PHP(m, k)reduces to PHP(m + 1, k) and to PHP(m + 1, k + 1).

Proof. Given $A \in \mathbb{Q}^{m \times m}$, $x \in \mathbb{Q}^m$ and a polyhedron $P \subseteq \mathbb{Q}^m$ with $\dim(P) = k$, we define the polyhedra $P' = \{(t,0) \in \mathbb{Q}^{m+1} : t \in P\}$ and $P'' = \{(t,1) \in \mathbb{Q}^{m+1} : t \in P\}$. Note that $\dim(P') = k$ and $\dim(P'') = k + 1$. Then

$$A^{n}x \in P \quad \Longleftrightarrow \quad \begin{bmatrix} A & 0 \\ 0 & 1 \end{bmatrix}^{n} \begin{bmatrix} x \\ 0 \end{bmatrix} \in P'$$
$$\iff \quad \begin{bmatrix} A & 0 \\ 0 & 1 \end{bmatrix}^{n} \begin{bmatrix} x \\ 1 \end{bmatrix} \in P''$$

which shows both reductions.

²In fact we can solve the problem in greater generality. One can show a **PSPACE** bound in the case of a *simple* shared recurrence with at most four dominant complex roots. This in turn entails membership in **PSPACE** for PHP(4, 4) and PHP(5, 5) in the case of a diagonalisable matrix. We omit this from the present paper for lack of space.

Next, recall that **Skolem's Problem** is the problem of determining, given a linear recurrence sequence S(n) over \mathbb{Q} , whether it has a zero, that is, an index $n \in \mathbb{N}$ such that S(n) = 0. The decidability of Skolem's Problem for sequences of order 5 or greater has been open for decades.

It is easy to show that Skolem's Problem for LRS of order m reduces to PHP(m, m - 1). For a linear recurrence sequence $S(n) = y^T A^n x$, we have S(n) = 0if and only if $A^n x \in P$, where P is the polyhedron $\{t \in \mathbb{Q}^m : y^T t \ge 0 \text{ and } y^T t \le 0\}$. In fact, $P = (span\{y\})^{\perp}$, so dim(P) = m - 1 and this is an instance of PHP(m, m - 1). By Lemma 2.1, it follows that whenever m > k, decidability of PHP(m, k) would imply decidability of Skolem's Problem for LRS of order k + 1.

In fact, we can show that even PHP(4,3) is hard for Skolem's Problem for linear recurrence sequences of order 5.

LEMMA 2.2. Skolem's Problem for LRS of order 5 reduces to PHP(4,3).

Proof. As discussed in reference [19], the only outstanding case of Skolem's Problem of order 5 is when the LRS has five characteristic roots: two pairs of complex conjugates $\lambda_1, \overline{\lambda_1}, \lambda_2, \overline{\lambda_2}$ and a real root ρ , such that $|\lambda_1| = |\lambda_2| > |\rho| > 0$. Therefore, let $S_1(n)$ be such a sequence, given by

$$\mathcal{S}_1(n) = a\lambda_1^n + \overline{a\lambda_1^n} + b\lambda_2^n + \overline{b\lambda_2^n} + c\rho^n$$

Define the order-4 sequence $S_2(n)$ by

$$S_2(n) = \frac{a\lambda_1^n + \overline{a\lambda_1^n} + b\lambda_2^n + \overline{b\lambda_2^n}}{\rho^n}$$

Let A be the 4×4 companion matrix of S_2 , and let x be the vector of initial terms of S_2 , so that

$$A^{n}x = \begin{bmatrix} \mathcal{S}_{2}(n) \\ \mathcal{S}_{2}(n+1) \\ \mathcal{S}_{2}(n+2) \\ \mathcal{S}_{2}(n+3) \end{bmatrix}$$

Then $S_1(n) = 0$ if and only if $S_2(n) = -c$, or equivalently, if there exist u_1, u_2, u_3 such that

$$A^{n}x = \begin{bmatrix} -c \\ 0 \\ 0 \\ 0 \end{bmatrix} + u_{1} \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} + u_{2} \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} + u_{3} \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

which is an instance of PHP(4,3).

Finally, in Section 4 we show that solving PHP(m,k) for $m \ge k \ge 4$ is highly unlikely without

major breakthroughs in analytic number theory. For any real number x, the homogeneous Diophantine approximation type L(x) is a measure of the extent to which x can be well-approximated by rationals. It is defined by:

$$L(x) = \inf \left\{ c \in \mathbb{R} : \exists n, m \in \mathbb{Z}. \left| x - \frac{n}{m} \right| < \frac{c}{m^2} \right\}$$

Much effort has been devoted to the study of the possible values of the approximation type, see for instance [8]. Nonetheless, very little is known about the approximation type of the vast majority of transcendental numbers. In Section 4 we prove that a decision procedure for the Simultaneous Positivity Problem for rational recurrences order at most 4 would entail the computability of $L(\arg \lambda/2\pi)$ for any complex number $\lambda \in \mathbb{Q}(i)$ of absolute value 1.³ Therefore, a decision procedure for PHP(4, 4) is extremely unlikely without significant advances in Diophantine approximation. By Lemma 2.1, the same hardness result holds for PHP(m, k) with $m \geq k \geq 4$. A similar result has been shown in [20] concerning the Positivity Problem for single linear recurrence sequences of order at most 6.

Our results are summarised in tabular form in the figure presented in the Introduction.

3 Extended Orbit Problem

In this section, we give an overview of the *Extended* Orbit Problem. We are given a matrix $A \in \mathbb{Q}^{m \times m}$, an initial point $x \in \mathbb{Q}^m$, and a target cone specified by a set of vectors $\{y_1, \ldots, y_d\} \subseteq \mathbb{Q}^m$ and a constraint matrix $B \in (\mathbb{R} \cap \mathbb{A})^{k \times d}$. The question is whether there exists an exponent $n \in \mathbb{N}$ and coordinates $u = (u_1, \ldots, u_d)^T$ such that $A^n x = \sum_{i=1}^d u_i y_i$ and $Bu \ge 0$. We refer to the space V spanned by y_1, \ldots, y_d , which contains the target cone, as the target space.

Our main decidability result concerning the Extended Orbit Problem is the following:

THEOREM 3.1. The Extended Orbit Problem is in **PTIME** in the case of a one-dimensional target space, and in **PSPACE** in the case of a two- or three-dimensional target space.

Notice that these complexity bounds depend only on the dimension of the target space V, not on the dimension of the ambient space \mathbb{Q}^m .

We now give an overview of the strategy for proving Theorem 3.1, and consign the full proof to Appendix C in the interest of clarity. The decision method

³Recall that a real number x is *computable* if there exists an algorithm which, given any rational $\varepsilon > 0$ as input, computes a rational q such that $|q - x| < \varepsilon$.

constructs a 'Master System' consisting of equations in n and $u = (u_1, \ldots, u_d)$ together with the inequalities $Bu \ge 0$ given as part of the input. The solutions of the Master System are in one-to-one correspondence with the solutions of the problem instance.

When the Master System contains sufficiently many equations, it was shown in reference [5] that a bound N can be derived such that if n > N, then $A^n x \notin V$. Writing ||I|| for the size of the input, we have $N \in ||I||^{O(1)}$ when dim(V) = 1 and $N \in 2^{||I||^{O(1)}}$ when dim $(V) \leq 3$.

With a one-dimensional target, it is sufficient to try all exponents $n \leq N$ to get a polynomial-time algorithm. In the two- and three-dimensional case, the algorithm is a guess-and-check procedure. An exponent $n \leq N$ is nondeterministically chosen as a possible witness. Then $A^n x \in V$ is verified by checking whether the determinant of the matrix with columns $A^n x, y_1, \ldots, y_d$ equals 0. If it does, then $A^n x \in V$, and we proceed to calculate the coefficients u_1, \ldots, u_d witnessing this membership and to verify the inequalities $Bu \geq 0$ which they must satisfy.

In the verification procedure, all numbers are expressed as arithmetic circuits and exponentiation is performed using repeated squaring. Recall that **PosSLP** is the class of problems which reduce in polynomial time to checking whether an arithmetic circuit evaluates to a positive number. The described operations may all be carried out using an oracle for **PosSLP**, so the algorithm gives a complexity upper bound of **NP**^{PosSLP} for the case of a large Master System. The work of Allender et al. [1] places **PosSLP** in the counting hierarchy, which shows the algorithm runs in polynomial space, as Theorem 3.1 claims.

On the other hand, when the Master System contains 'few' equations, we do not have such a bound Nbeyond which membership in V is impossible. These cases are the main focus of Appendix C, where we show how to solve such small Master Systems. The procedure invokes a decision method for the Simultaneous Positivity Problem, which is discussed in Section 4 and is shown to be in **PSPACE** for all orders which arise in the reduction.

4 Simultaneous Positivity

In this section, we consider the **Simultaneous Pos**itivity problem: given linear recurrence sequences S_1, \ldots, S_k over $\mathbb{R} \cap \mathbb{A}$ which satisfy a common recurrence equation, are they ever simultaneously positive, that is, does there exist n such that $S_i(n) \ge 0$ for all $i \in \{1, \ldots, k\}$? Solving this problem is instrumental in our decision procedures for both the Extended Orbit Problem and the Polyhedron-Hitting Problem. The asymptotic behaviour of a linear recurrence sequence S is closely linked to its dominant characteristic roots, that is, the characteristic roots of greatest magnitude. If $\lambda_1, \ldots, \lambda_s$ are the dominant roots, we can write

$$\frac{\mathcal{S}(n)}{|\lambda_1|^n} = P_1(n) \left(\frac{\lambda_1}{|\lambda_1|}\right)^n + \dots + P_s(n) \left(\frac{\lambda_s}{|\lambda_1|}\right)^n + r(n)$$

where r(n) tends to 0 exponentially quickly. We can use the polynomial root-separation bound (A.2) in Appendix A.2 to bound the absolute value of the quotient λ/λ_1 , where λ is a non-dominant characteristic root. Thus we can show:

LEMMA 4.1. Suppose we are given an LRS S as above. Then there exist constants $\varepsilon \in (0, 1)$ and $N \in \mathbb{N}$ such that $N \in 2^{||S||^{O(1)}}$, $\varepsilon^{-1} \in 2^{||S||^{O(1)}}$, and $|r(n)| < (1-\varepsilon)^n$ for all n > N.

4.1 Decidability results In this section we prove the following result:

THEOREM 4.1. The Simultaneous Positivity Problem is in **PSPACE** for sequences over $\mathbb{R} \cap \mathbb{A}$ whose common recurrence equation has order at most 3, or order 4 but with at least one real root.

We will restrict our attention to non-degenerate LRS. As outlined in Appendix A.3, a degenerate sequence can be partitioned into non-degenerate subsequences. Then the Simultaneous Positivity instance is equivalent to the disjunction of all Simultaneous Positivity instances where each degenerate sequence has been replaced by one of its non-degenerate subsequences. In general, this leads to exponentially many non-degenerate problem instances. However, this leaves Theorem 4.1 unaffected, as a non-degenerate problem instance may simply be guessed nondeterministically by a **PSPACE** algorithm.

The assumption of non-degeneracy guarantees that there can be at most one real root among the dominant roots of the sequences. We can assume without loss of generality that any real root of the sequence is positive (otherwise we separately consider the cases of even and odd n).

The algorithm for Simultaneous Positivity is similar to the one for Extended Orbit. We search for witnesses up to some computable bound $N \in 2^{||I||^{O(1)}}$. To this end, we will choose a witness n nondeterministically and then verify $S_j(n) = v_j^T M_j^n w_j \ge 0$ for all j. Recalling that the entries of M_j are algebraic numbers, we can verify this family of inequalities by constructing a sentence τ in the existential first-order theory of the reals which is true if and only if $v_j^T M_j^n w_j \ge 0$ for all j. We specify each real algebraic number with description $(f_{\alpha}, x_0, y_0, R)$ using the first-order formula $\exists z. f_{\alpha}(z) = 0 \land (z - x_0)^2 + y_0^2 \leq R^2$. To ensure that $\|\tau\| \in \|I\|^{O(1)}$, we use repeated squaring to calculate M_j^n . Finally, we check the validity of τ in **PSPACE**, as per Theorem A.2 in Appendix A.4.

We now consider two cases, according to the number of dominant complex roots of the shared recurrence equation.

No dominant complex roots. Suppose the dominant characteristic roots do not include a pair of complex conjugates. Then by the assumption of non-degeneracy, there is one real dominant root $\rho > 0$. Then the *j*-th sequence is given by

$$\frac{\mathcal{S}_j(n)}{\rho^n} = P_j(n) + r_j(n)$$

where r_j is itself a linear recurrence of lower order which converges to 0 exponentially quickly, and $P_i \in$ $(\mathbb{R} \cap \mathbb{A})[x]$. Each polynomial $P_j(n)$ is either identically zero or is ultimately positive or ultimately negative as n tends to infinity. In the latter two cases, there is an effective threshold $N_j \in 2^{||\mathcal{S}_j(n)||^{O(1)}}$ beyond which the sign of \mathcal{S}_j does not change. If some \mathcal{S}_j is ultimately negative, then any witness to the problem instance must be bounded above by N_j . Since N_j is at most exponentially large in the size of the input, we use a guess-and-check procedure and are done. Similarly, for each sequence S_j for which P_j is ultimately positive we can search for witnesses up to the threshold N_i and if none are found, we discard S_j as if it were uniformly positive. Finally, we are left only with sequences S_j for which P_j is identically zero. Then the problem instance is equivalent to Simultaneous Positivity on the sequences r_i . These sequences satisfy a common recurrence equation of lower order, so we proceed inductively.

Two simple dominant complex roots. Suppose now that the dominant roots of the shared recurrence equation include exactly two complex roots $\lambda, \overline{\lambda}$ and possibly a real dominant root $\rho_1 > 0$. Moreover, assume that the roots are all simple, so the *j*-th sequence is given by

 $\mathcal{S}_{i}(n) = a_{i}\lambda^{n} + \overline{a_{i}}\overline{\lambda}^{n} + b_{i}\rho_{1}^{n} + c_{i}\rho_{2}^{n}$

that is,

$$\frac{\mathcal{S}_j(n)}{|\lambda|^n} = 2|a_j|\cos(\alpha_j + n\varphi) + b_j + r_j(n)$$

where $\alpha_j = \arg(a_j)$ and $\varphi = \arg(\lambda)$. Moreover, r_j is a linear recurrence sequence of order at most 2 with real characteristic roots. Observe that for all j, $b_j + r_j(n)$ is either ultimately positive or ultimately negative as n tends to infinity. Furthermore, a threshold beyond which the sign does not change is effectively computable and at most exponential in $||S_j||$. Following the reasoning of the previous case, we see that we can dismiss sequences S_j which have $a_j = 0$.

Assume therefore that $a_j \neq 0$ for all j. By Lemma B.1 in Appendix B, for each sequence S_j there exists an effective threshold $N_j \in 2^{||S_j||^{O(1)}}$ such that for $n > N_j$, $r_j(n)$ is too small to influence the sign of $S_j(n)$. That is, for all $n > N_j$, we have

$$S_j(n) \ge 0 \iff b_j + \cos(\alpha_j + n\varphi) \ge 0$$

Therefore, for $n > N = \max_{j} \{N_j\}$, the problem instance is equivalent to a conjunction of inequalities in n:

$$\forall j. \cos(\alpha_j + n\varphi) \ge -b_j$$

We use guess-and-check to look for witnesses $n \leq N$. If none are found, the problem instance is then decidable in **PSPACE** by Lemma B.2 in Appendix B.

4.2 Hardness We now proceed to show our main hardness result for Simultaneous Positivity and hence for PHP(m,m). Recall that the homogeneous Diophantine approximation type L(x) of a real number x, defined in Section 2.2, is a measure of how well x can be approximated by rationals. Very little progress has been made on calculating the approximation type for the vast majority of transcendental numbers. In this section, we show that a decision procedure for Simultaneous Positivity for LRS with shared recurrence equation of order 4 would entail the computability of the approximation type of all Gaussian rationals:

THEOREM 4.2. Suppose that Simultaneous Positivity is decidable for rational linear recurrence sequences. Then for any $\lambda \in \mathbb{Q}(i)$ on the unit circle, $L(\arg \lambda/2\pi)$ is a computable number.

Suppose we wish to calculate $L(\varphi/2\pi)$, where $\varphi = \arg \lambda$ for some λ of magnitude 1. Consider the following two sequences for some fixed rational number A:

$$S_1(n) = \frac{1}{2} \left((A - in)\lambda^n + (A + in)\overline{\lambda}^n \right)$$
$$S_2(n) = \frac{1}{2} \left((A + in)\lambda^n + (A - in)\overline{\lambda}^n \right)$$

It is straightforward to verify that $S_1(n)$ and $S_2(n)$ are both rational sequences satisfying a common order-4 recurrence with characteristic polynomial $(x - \lambda)^2 (x - \overline{\lambda})^2$. Moreover we have

$$S_1(n) = n \cos(n\varphi - \pi/2) + A \cos(n\varphi)$$

= $A \cos(n\varphi) + n \sin(n\varphi)$

$$S_2(n) = n \cos(n\varphi + \pi/2) + A \cos(n\varphi)$$

= $A \cos(n\varphi) - n \sin(n\varphi)$

Let $w_n = n |\sin(n\varphi)| - A\cos(n\varphi)$. It is clear that $S_1(n) \ge 0$ and $S_2(n) \ge 0$ if and only if $w_n \le 0$. We will show that a Simultaneous Positivity oracle may be used on these sequences for different choices of A to compute arbitrarily good approximations of $L(\varphi/2\pi)$. Throughout this section, write [x] to denote the distance from x to the closest integer multiple of 2π , that is, $[x] = \min\{|x - 2\pi j| : j \in \mathbb{Z}\}.$

Given $\varepsilon \in (0, 1)$, there exists $\delta > 0$ such that for all $x \in [-\delta, \delta]$, the following hold:

(4.1)
$$(1-\varepsilon)|x| \le |\sin x| \le |x|$$

$$(4.2) 1 - \varepsilon \le \cos x$$

Moreover, there exists $N \in \mathbb{N}$ such that $A/N \leq \delta$ and also,

(4.3) if
$$|\sin x| \le A/N$$
, then $|x| \le \delta$.

LEMMA 4.2. Suppose that $n \ge N$ is such that $w_n \le 0$. Then $n[n\varphi] < A/(1-\varepsilon)$.

Proof.

$$|\sin(n\varphi)| \le \frac{A}{n}\cos(n\varphi) \le \frac{A}{N} \quad [\text{as } w_n \le 0, n \ge N]$$

$$\Rightarrow [n\varphi] \le \delta \qquad \qquad [by (4.3)]$$

But from the definition of w_n , inequality (4.1) and $\cos x \leq 1$, we have

$$w_n = n |\sin(n\varphi)| - A\cos(n\varphi) \ge n(1-\varepsilon)[n\varphi] - A$$

Therefore, $n[n\varphi] \leq A/(1-\varepsilon)$.

LEMMA 4.3. Let $n \ge N$ be such that $n[n\varphi] \le A(1-\varepsilon)$. Then $w_n \le 0$.

Proof. Notice that

$$[n\varphi] \le \frac{A(1-\varepsilon)}{n} \le \frac{A}{N} \le \delta$$

so for w_n we have

$$w_n = n |\sin(n\varphi)| - A\cos(n\varphi)$$
 [definition of w_n]

$$\leq n[n\varphi] - A(1-\varepsilon) \qquad \qquad [by (4.1)(4.2)]$$

$$\leq A(1-\varepsilon) - A(1-\varepsilon) = 0$$
 [by premise]

Letting $t = \varphi/2\pi$, we see that

$$2\pi L(t) = \inf_{m \in \mathbb{N}} m[m\varphi]$$

Thus to show computability of L(t) it is enough to show that $\inf_{m \in \mathbb{N}} m[m\varphi]$ is computable. For this in turn it suffices to provide a procedure that, given $a, b \in \mathbb{Q}$ with a < b, computes a threshold $N \in \mathbb{N}$ and either outputs that $\inf_{m \ge N} m[m\varphi] < b$ or $\inf_{m \ge N} m[m\varphi] > a$. (Clearly $\inf_{m < N} m[m\varphi]$ can be computed to any desired precision.)

Given a < b as above, compute ε and A such that

$$a < A(1-\varepsilon) < \frac{A}{1-\varepsilon} < b$$
.

Calculate also the constant N in the statement of Lemmas 4.2 and 4.3 for this choice of ε and A. Then run a Simultaneous Positivity oracle on the N-th tails of the two sequences $S_1(n)$ and $S_2(n)$ to determine whether $w_n \leq 0$ for some $n \geq N$. If the oracle accepts, then $\inf_{m \in \mathbb{N}} m[m\varphi] \leq \frac{A}{1-\varepsilon} < b$ by Lemma 4.2. If the oracle rejects, then $\inf_{m \in \mathbb{N}} m[m\varphi] \geq A(1-\varepsilon) > a$ by Lemma 4.3.

A Preliminaries

A.1 Polyhedra and their representations Here we state some basic properties of polyhedra. For more details we refer the reader to, for example [10, 16, 26]. A halfspace in \mathbb{R}^d is the set of points $x \in \mathbb{R}^d$ satisfying $v^T x \ge c$ for some fixed vector $v \in \mathbb{R}^d$ and real number c. A polyhedron in \mathbb{R}^d is the intersection of finitely many halfspaces:

(A.1)
$$P = \begin{cases} v_1^T x \ge c_1 \\ x \in \mathbb{R}^d : & \vdots \\ v_m^T x \ge c_m \end{cases}$$

We call the set $\{(v_1, c_1), \ldots, (v_m, c_m)\}$ a halfspace description of a polyhedron, or simply an *H*-polyhedron. The problem of determining a minimal subset of the inequalities (A.1) that define the same polyhedron is called the *H*-redundancy removal problem and is solvable in polynomial time by reduction to linear programming. Thus, we may freely assume that there are no redundant constraints in the descriptions of H-polyhedra.

The dimension of a polyhedron P, denoted dim(P), is the dimension of the subspace of \mathbb{R}^d spanned by P. The task of calculating the dimension of an Hpolyhedron, called the *H*-dimension problem, can be done in polynomial time by solving polynomially many linear programs. If dim(P) = d, we call P fulldimensional. The minimal halfspace representation of a full-dimensional polyhedron is unique, up to scaling of the inequalities in (A.1). The *convex cone* of a finite set of vectors v_1, \ldots, v_m is defined as

$$\operatorname{cone}(\{v_1,\ldots,v_m\}) = \{\lambda_1 v_1 + \cdots + \lambda_m v_m : \forall i . \lambda_i \ge 0\}$$

If the vectors v_1, \ldots, v_m are linearly independent, the cone is called *simplicial*. A classical result, due to Carathéodory, states that each finitely generated cone can be written as a finite union of simplicial cones:

THEOREM A.1. (Carathéodory) Let $v_1, \ldots, v_m \in \mathbb{R}^d$. If $v \in cone(v_1, \ldots, v_m)$, then v belongs to the cone generated by a linearly independent subset of $\{v_1, \ldots, v_m\}$.

We use this to prove that any two-dimensional polyhedron decomposes into a finite union of simple twodimensional shapes:

LEMMA A.1. Suppose $P \subseteq \mathbb{R}^d$ is a two-dimensional polyhedron. Then $P = \bigcup_{i=1}^m A_i$, where m is finite and each of A_i is of the form

$$A_i = \{u_i + \alpha v_i + \beta w_i : T_i(\alpha, \beta)\}$$

for vectors $u_i, v_i, w_i \in \mathbb{R}^d$ and predicates $T_i(\alpha, \beta)$ chosen from the following:

- $T_i(\alpha, \beta) \equiv \alpha \geq 0 \land \beta \geq 0$ (A_i is an infinite cone)
- $T_i(\alpha,\beta) \equiv \alpha \ge 0 \land \beta \ge 0 \land \alpha + \beta \le 1$ (A_i is a triangle)
- $T_i(\alpha, \beta) \equiv \alpha \ge 0 \land \beta \ge 0 \land \beta \le 1$ (A_i is an infinite strip)

Furthermore, if we are given a halfspace description of P with length ||P||, the size of the representation of each vector u_i, v_i, w_i is at most $||P||^{O(1)}$.

Proof. Let

$$P = \{x \in \mathbb{R}^d : Ax \ge b\}$$

for some $A \in \mathbb{R}^{m \times d}$, $b \in \mathbb{R}^d$ and define the polygon

$$P' = \{ y \in \mathbb{R}^{d+1} : [A - b] y \ge 0 \}$$

so that $\dim(P') = 3$ and

$$P = \{ x \in \mathbb{R}^d : (x \ 1)^T \in P' \}$$

Notice that P' is specified using only homogeneous inequalities, so there exist vectors $V = \{v_1, \ldots, v_s\}$ such that $P' = \operatorname{cone}(V)$. By scaling if necessary, we can assume the (d + 1)-th component of each v_i is either 0 or 1. Let \mathcal{H} denote the hyperplane in \mathbb{R}^{d+1} where the (d+1)-th coordinate is 1. By Carathéodory's Theorem, P' may be written as the union of finitely many cones generated from linearly independent subsets of V. Let u_i be the projection of v_i to the first d coordinates. Since $\dim(P') = 3$, no more than three elements of V can be linearly independent, so

$$P' = \bigcup_{(i_1, i_2, i_3) \in I} \operatorname{cone}(v_{i_1}, v_{i_2}, v_{i_3})$$

The intersection $\mathcal{H} \cap \operatorname{cone}(v_{i_1}, v_{i_2}, v_{i_3})$ is non-empty if and only if at least one of $v_{i_1}, v_{i_2}, v_{i_3}$ has 1 in the (d+1)th coordinate. Therefore, P is the finite union of shapes A_i with only two degrees of freedom:

$$A_i = \{ \alpha u_{i_1} + \beta u_{i_2} + \gamma u_{i_3} : \alpha, \beta, \gamma \ge 0 \land T_i(\alpha, \beta, \gamma) \}$$

where each predicate T_i is $\alpha = 1$, or $\alpha + \beta = 1$, or $\alpha + \beta + \gamma = 1$. These are precisely the desired three types of parametric shapes. The descriptions of the vectors involved is polynomially large because each vector v_i is the intersection of d of the halfspaces in \mathbb{R}^{d+1} which define P'.

A simpler version of the above result gives a similar parametric form in the case $\dim(P) = 1$:

LEMMA A.2. Suppose $P \subseteq \mathbb{R}^d$ is a one-dimensional polyhedron. Then

$$P = \{v_1 + \alpha v_2 : T(\alpha)\}$$

where the predicate $T(\alpha)$ is one of $\alpha \in \mathbb{R}$, $\alpha \geq 0$ and $\alpha \in [0,1]$. Furthermore, if we are given a halfspace description of P with length ||P||, the size of the representation of v_1, v_2 is at most $||P||^{O(1)}$.

A.2 Algebraic numbers In this section we briefly review relevant notions in algebraic number theory. See, e.g., [7] for more details.

A complex number α is algebraic if there exists a polynomial $p \in \mathbb{Q}[x]$ such that $p(\alpha) = 0$. The set of algebraic numbers, denoted by \mathbb{A} , is a subfield of \mathbb{C} . The minimal polynomial of α , denoted $f_{\alpha}(x)$, is the unique monic polynomial with rational coefficients of least degree which vanishes at α . The degree of $\alpha \in \mathbb{A}$ is defined as the degree of its minimal polynomial and is denoted by n_{α} . The height of α is defined as the maximum absolute value of a numerator or denominator of a coefficient of the minimal polynomial of α , and is denoted by H_{α} . The roots of $f_{\alpha}(x)$ (including α) are called the Galois conjugates of α . An algebraic integer is an algebraic number α such that $f_{\alpha} \in \mathbb{Z}[x]$. The set of algebraic integers, denoted $\mathcal{O}_{\mathbb{A}}$, is a ring under the usual addition and multiplication.

The canonical representation of an algebraic number α is its minimal polynomial $f_{\alpha}(x)$, along with a numerical approximation of $Re(\alpha)$ and $Im(\alpha)$ of sufficient precision to distinguish α from its Galois conjugates. More precisely, we represent α by the tuple

$$(f_{\alpha}, x, y, R) \in \mathbb{Q}[x] \times \mathbb{Q}^3$$

meaning that α is the unique root of f_{α} inside the circle centred at (x, y) in the complex plane with radius R. A bound due to Mignotte [17] states that for roots $\alpha_i \neq \alpha_j$ of a polynomial p(x),

(A.2)
$$|\alpha_i - \alpha_j| > \frac{\sqrt{6}}{n^{(n+1)/2} H^{n-1}}$$

where n and H are the degree and height of p, respectively. Thus, if R is restricted to be less than a quarter of the root separation bound, the representation is well-defined and allows for equality checking. Observe that given f_{α} , the remaining data necessary to describe α is polynomial in the length of the input. It is known how to obtain polynomially many bits of the roots of any $p \in \mathbb{Q}[x]$ in polynomial time [21].

When we say an algebraic number α is given, we assume we have a canonical description of α . We will denote by $\|\alpha\|$ the length of this description, assuming that integers are expressed in binary and rationals are expressed as pairs of integers. Observe that $|\alpha|$ is an exponentially large quantity in $\|\alpha\|$ whereas $\ln |\alpha|$ is polynomially large. Notice also that $1/\ln |\alpha|$ is at most exponentially large in $\|\alpha\|$. For a rational a, $\|a\|$ is just the sum of the lengths of its numerator and denominator written in binary. For a polynomial $p \in$ $\mathbb{Q}[x], \; \|p\|$ will denote $\sum_{i=0}^n \|p_i\|$ where n is the degree of the polynomial and p_i are its coefficients. Using the resultant method, operations may be performed efficiently on algebraic numbers. Specifically, techniques from algebraic number theory [7] yield the following lemma:

LEMMA A.3. Given canonical representations of $\alpha, \beta \in \mathbb{A}$ and a polynomial $p \in \mathbb{Q}[x]$, it is possible to compute canonical descriptions of $\alpha \pm \beta$, $\alpha\beta^{\pm 1}$, $\sqrt{\alpha}$ and $p(\alpha)$, to check the equality $\alpha = \beta$ and α 's membership in $\mathbb{N}, \mathbb{Z}, \mathbb{Q}$, and finally to determine whether α is a root of unity, and if so, to calculate its order and argument. All of these procedures have polynomial running time.

A.3 Linear recurrence sequences We now recall some basic properties of linear recurrence sequences. For more details, we refer the reader to [9, 11]. A *real linear recurrence sequence* (*LRS*) is an infinite sequence $S = \langle S(0), S(1), S(2), \ldots \rangle$ over \mathbb{R} such that there exists a natural number k and real numbers a_1, \ldots, a_k such that $a_k \neq 0$ and S satisfies the linear recurrence equation

(A.3)

$$\mathcal{S}(n+k) = a_1 \mathcal{S}(n+k-1) + a_2 \mathcal{S}(n+k-2) + \dots + a_k \mathcal{S}(n)$$

The recurrence (A.3) is said to have order k. Note that the same sequence can satisfy different recurrence relations, but it satisfies a unique recurrence of minimum order.

The characteristic polynomial of \mathcal{S} is

$$p(x) = x^{k} - a_{1}x^{k-1} - a_{2}x^{k-2} - \dots - a_{k}$$

and its roots are called the *characteristic roots* of the sequence. For real LRS, the set of characteristic roots is closed under complex conjugation. If $\rho_1, \ldots, \rho_l \in \mathbb{R}$ are the real roots of p(x) and $\gamma_1, \overline{\gamma}_1, \ldots, \gamma_m, \overline{\gamma}_m \in \mathbb{C}$ are the complex ones, the sequence is given by

$$\mathcal{S}(n) = \sum_{i=1}^{l} A_i(n)\rho_i^n + \sum_{j=1}^{m} \left(C_j(n)\gamma_j^n + \overline{C_j}(n)\overline{\gamma_j}^n\right)$$

for all $n \geq 0$, where $A_i \in \mathbb{R}[x]$ and $C_j \in \mathbb{C}[x]$ are univariate polynomials whose degrees are at most the multiplicity of the corresponding roots of p(x). The coefficients of A_i, C_i are effectively computable algebraic numbers.

If $M \in \mathbb{R}^{k \times k}$ is a real square matrix and $v, w \in \mathbb{R}^k$ are real column vectors, then it can be shown using the Cayley-Hamilton Theorem that the sequence $S(n) = v^T M^n w$ satisfies a linear recurrence of order k. Conversely, any LRS may be expressed in this way: it is sufficient to take M to be the transposed companion matrix of the characteristic polynomial of S, v to be the vector $(S(k-1), \ldots, S(0))^T$ of initial terms of S in reverse order, and w to be the unit vector $(0, \ldots, 0, 1)^T$. The characteristic roots of the LRS are precisely the eigenvalues of M.

A linear recurrence sequence is called *degenerate* if for some pair of distinct characteristic roots λ_1, λ_2 of its minimum-order recurrence, the ratio λ_1/λ_2 is a root of unity, otherwise the sequence is *non-degenerate*. As pointed out in [9], the study of arbitrary LRS can effectively be reduced to that of non-degenerate LRS by partitioning the original LRS into finitely many non-degenerate subsequences. Specifically, for a given degenerate linear recurrence sequence S with characteristic roots λ_i , let L be the least common multiple of the orders of all ratios λ_i/λ_j which are roots of unity. Then consider the sequences

$$\mathcal{S}^{(j)}(n) = u^T A^{nL+j} v = u^T (A^L)^n (A^j v)$$

where $j \in \{0, \ldots, L-1\}$. Each of these sequences has characteristic roots λ_i^L and is therefore non-degenerate, because $(\lambda_1/\lambda_2)^{Lk} = 1$ implies $\lambda_1^L = \lambda_2^L$. From the crude lower bound $\varphi(r) \geq \sqrt{r/2}$ on Euler's totient function, it follows that if α has degree d and is a primitive r-th root of unity, then $r \leq 2d^2$. Thus, $L \in 2^{||A||^{O(1)}}$, so non-degeneracy can be ensured by considering at most exponentially many subsequences of the original LRS.

A.4 First-order theory of the reals Let x_1, \ldots, x_m be first-order variables ranging over \mathbb{R} , and suppose $\sigma(x_1, \ldots, x_m)$ is a Boolean combination of predicates of the form $g(x_1, \ldots, x_m) \sim 0$, where $g \in \mathbb{Z}[x_1, \ldots, x_m]$ is a polynomial and \sim is > or =. A sentence of the first-order theory of the reals is a formula τ of the form

$$Q_1 x_1 \ldots Q_m x_m \sigma(x_1, \ldots, x_m)$$

where each Q_i is one of the quantifiers \exists and \forall . If all the quantifiers are \exists , then τ is said to be a sentence of the *existential* first-order theory of the reals.

The decidability of the first-order theory of the reals was originally established by Tarski [25]. Many refinements followed over the years, culminating in the analysis of Renegar [22]. We make use of the following result:

THEOREM A.2. Suppose we are given a sentence τ of the form above using only existential quantifiers. The problem of deciding whether τ holds over the reals is in **PSPACE**. Furthermore, if $M \in \mathbb{N}$ is a fixed constant and we restrict the problem to formulae τ where the number of variables is bounded above by M, then the problem is in **PTIME**.

B Technical lemmas

THEOREM B.1. (Baker and Wüstholz [2]) Let $\alpha_1, \ldots, \alpha_m$ be algebraic numbers other than 0 or 1, and let b_1, \ldots, b_m be rational integers. Write

$$\Lambda = b_1 \log \alpha_1 + \dots + b_m \log \alpha_m$$

Let $A_1, \ldots, A_m, B \geq e$ be real numbers such that, for each $j \in \{1, \ldots, m\}$, A_j is an upper bound for the height of α_j , and B is an upper bound for $|b_j|$. Let d be the degree of the extension field $\mathbb{Q}(\alpha_1, \ldots, \alpha_m)$ over \mathbb{Q} . If $\Lambda \neq 0$, then

$$\log |\Lambda| > -(16md)^{2(m+2)} \log(A_1) \dots \log(A_m) \log(B)$$

THEOREM B.2. Suppose $\alpha, \beta, \gamma, A, B, C \in \mathbb{A}$ and the ratios of α, β, γ (where they exist) are not roots of unity. Let $||I|| = ||\alpha|| + ||\beta|| + ||\gamma|| + ||A|| + ||B|| + ||C||$. Then there exist effective bounds $N_1 \in ||I||^{O(1)}$ and $N_2 \in 2^{||I||^{O(1)}}$ such that if $A\alpha^n + B\beta^n = 0$ then $n \leq N_1$, and if $A\alpha^n + B\beta^n + C\gamma^n = 0$ or $A\alpha^n + Bn\beta^{n-1} + C\beta^n = 0$ then $n \leq N_2$. LEMMA B.1. Let $a, \lambda \in \mathbb{A}$ and $C, \chi \in \mathbb{A} \cap \mathbb{R}$ be given where λ is not a root of unity and $|\chi| < |\lambda| = 1$. Let $\alpha = \arg(a)$ and $\varphi = \arg(\lambda)$. Then there exists an effectively computable $N \in \mathbb{N}$ such that for all n > N, $|C + \cos(\alpha + n\varphi)| > |\chi|^n$. Moreover, $N \in 2^{||I||^{O(1)}}$ where $||I|| = ||\lambda|| + ||\chi|| + ||a|| + ||C||$.

Proof. Suppose that $|C| \leq 1$ and let $b = C + i\sqrt{1 - C^2} = e^{i\beta}$, so that $C = \cos(\beta)$. Then b is algebraic with $\deg(b) \in ||I||^{O(1)}$, $H_b \in 2^{||I||^{O(1)}}$. It is clear that

$$C + \cos(\alpha + n\varphi) = 2\cos\frac{\alpha + \beta + n\varphi}{2}\cos\frac{\alpha - \beta + n\varphi}{2}$$

Since λ is not a root of unity, by Lemma B.2, there exists an effective constant $N_1 \in ||I||^{O(1)}$ such that if $ab^{\pm 1}\lambda^n = -1$ then $n \leq N_1$. Therefore, for $n > N_1$, we have $\cos(\alpha \pm \beta + n\varphi) \neq 0$. Let k_n be the unique integer such that $k_n\pi + (\alpha + \beta + n\varphi + \pi)/2 \in [-\pi/2, \pi/2)$. Notice that $|k_n| < 2n$. Then

$$\left|\cos\frac{\alpha+\beta+n\varphi}{2}\right| = \left|\sin\frac{\alpha+\beta+n\varphi+(2k_n+1)\pi}{2}\right|$$
$$\geq \frac{|\alpha+\beta+n\varphi+(2k_n+1)\pi|}{2\pi}$$

by the inequality $|\sin(x)| \ge |x|/\pi$ for $x \in [-\pi/2, \pi/2]$. Note that α , β , φ and π are logarithms of algebraic numbers with degree polynomial in ||I|| and height exponential in ||I||. Then by from Baker's Theorem, there exist effective positive constants $p_1, p_2 \in ||I||^{O(1)}$ such that

$$n > N_1 \Rightarrow \left| \cos \frac{\alpha + \beta + n\varphi}{2} \right| > (p_1 n)^{-p_2}$$

By the same argument with β replaced by $-\beta$, there exist effective positive constants $N_2, p_3, p_4 \in ||I||^{O(1)}$ such that

$$n > N_2 \Rightarrow \left| \cos \frac{\alpha - \beta + n\varphi}{2} \right| > (p_2 n)^{-p_4}$$

However, since χ^n shrinks exponentially with n and $|\chi^{-1}| \in 2^{||I||^{O(1)}}$, it follows that there exists an effective constant $N_3 \in 2^{||I||^{O(1)}}$ such that for all $n > N_3$,

$$(p_1n)^{-p_2}(p_3n)^{-p_4} > |\chi^n|$$

Then for all $n > \max\{N_1, N_2, N_3\}$, we have

$$|C + \cos(\alpha + n\varphi)| > p_1 p_3 n^{-(p_2 + p_4)} > |\chi^n|$$

as desired.

The remaining case |C| > 1 is easy. If C > 1, we have

$$C + \cos(\alpha + n\varphi) > 1 + \cos(\alpha + n\varphi) = \cos(0) + \cos(\alpha + n\varphi)$$

and the lemma follows by the above argument with $\beta = 0$. Similarly when C < -1.

LEMMA B.2. Suppose a_1, \ldots, a_m and λ are all algebraic numbers on the unit circle and λ is not a root of unity. Suppose also $c_1, \ldots, c_m \in \mathbb{R} \cap \mathbb{A}$. Let $\alpha_j = \arg(a_j)$ and $\varphi = \arg(\lambda)$. Then it is decidable whether there exists an integer n such that

$$\cos(\alpha_j + n\varphi) \ge c_j \text{ for all } j = 1, \dots, m$$

Moreover, the decision procedure's running time is $||I||^{O(1)}$ where

$$||I|| = \sum_{j=1}^{m} (||a_j|| + ||c_j||) + ||\lambda||$$

Proof. Inequalities where $c_j \leq -1$ may be discarded, as they are satisfied for all n, whereas the presence of inequalities with $c_j > 1$ immediately makes the problem instance negative. Now assuming $c_j \in (-1, 1]$, each inequality

(B.4)
$$\cos(\alpha_j + n\varphi_1) \ge c_j$$

defines an arc on the unit circle which λ^n must lie within. Specifically, (B.4) holds if and only if λ^n lies on the arc \mathcal{A}_j defined by

$$\mathcal{A}_j = \{ z \in \mathbb{C} : |z| = 1 \text{ and } h(w_1, w_2, z) \le 0 \}$$

where $w_1 = \overline{a_j} \left(c_j - i\sqrt{1 - c_j^2} \right)$ and $w_2 = \overline{a_j} \left(c_j + i\sqrt{1 - c_j^2} \right)$ are the endpoints of the arc, and

$$h(x, y, z) = \begin{vmatrix} Re(x) & Im(x) & 1 \\ Re(y) & Im(y) & 1 \\ Re(z) & Im(z) & 1 \end{vmatrix}$$

is the orientation function.⁴

The endpoints of \mathcal{A}_j are clearly algebraic and may be computed explicitly in polynomial time in ||I||. Then the intersection $\mathcal{A} = \bigcap_j \mathcal{A}_j$ is also computable in polynomial time. Since λ is not a root of unity, the set $\{\lambda^n : n \in \mathbb{N}\}$ is dense on the unit circle. If \mathcal{A} is empty, then the problem instance is negative. If \mathcal{A} is a nontrivial arc on the unit circle, then by density, the problem instance is positive. Finally, \mathcal{A} could be a set of at most two points z_1, z_2 on the unit circle. Then the problem instance is positive if and only if there exists an exponent $n \in \mathbb{N}$ such that $\lambda^n = z_i$ for some *i*. A polynomial bound on *n* then follows from Theorem B.2.

C Extended Orbit Problem

We now give the details of our decision procedure for the Extended Orbit Problem, as promised in Section 3.

C.1 A Master System In [5], we show how to reduce the Orbit Problem (determining whether there exists $n \in \mathbb{N}$ such that $A^n x$ lies in a vector space V) to the matrix power problem: determining whether there exists $n \in \mathbb{N}$ such that A^n lies in the span of $p_1(A), \ldots, p_d(A)$ for given polynomials $p_1, \ldots, p_d \in$ $\mathbb{Q}[x]$. The reduction takes polynomial time, relies on standard linear algebra and is straightforward to extend, mutatis mutandis, in order to include linear inequalities on the coefficients which witness membership of $A^n x$ in the target vector space. Thus, we shall assume that a problem instance of the Extended Orbit Problem is specified by matrices $A \in \mathbb{Q}^{m \times m}, B \in (\mathbb{R} \cap$ $\mathbb{A}^{k \times d}$ and polynomials $p_1, \ldots, p_d \in \mathbb{Q}[x]$ such that $p_1(A), \ldots, p_d(A)$ are linearly independent, and we have to determine whether there exist $n \in \mathbb{N}$ and u = $(u_1,\ldots,u_d) \in \mathbb{Q}^d$ such that

(C.5)
$$A^n = u_1 p_1(A) + \dots + u_d p_d(A) \text{ and } Bu \ge 0$$

We now proceed to show a Master System of equations, which is equivalent to (C.5). Let $f_A(x)$ be the minimal polynomial of A over \mathbb{Q} and let $\alpha_1, \ldots, \alpha_t$ be its roots, that is, the eigenvalues of A. These can be calculated in polynomial time. Throughout this paper, for an eigenvalue α_i we will denote by $mul(\alpha_i)$ the multiplicity of α_i as a root of the minimal polynomial of the matrix.

Fix an exponent n and coefficients u_1, \ldots, u_d and define the polynomials $P(x) = \sum_{i=1}^d u_i p_i(x)$ and $Q(x) = x^n$. It is easy to see that (C.5) is satisfied if and only if

(C.6)
$$Bu \ge 0 \land P^{(j)}(\alpha_i) = Q^{(j)}(\alpha_i)$$

for all $i \in \{1, \ldots, t\}$, $j \in \{0, \ldots, mul(\alpha_i) - 1\}$. Indeed, P - Q is zero at A if and only if $f_A(x)$ divides P - Q, that is, each α_i is a root of P - Q with multiplicity at least $mul(\alpha_i)$. This is equivalent to saying that each α_i is a root of P - Q and its first $mul(\alpha_i) - 1$ derivatives.

Thus, in order to decide whether the problem instance is positive, it is sufficient to solve the system of equations and inequalities (C.6) in the unknowns nand u_1, \ldots, u_d . Each eigenvalue α_i contributes $mul(\alpha_i)$ equations which specify that P(x) - Q(x) and its first $mul(\alpha_i) - 1$ derivatives all vanish at α_i .

For example, if $f_A(x)$ has roots α_1 , α_2 , α_3 with multiplicities $mul(\alpha_i) = i$ and the target space is $span \{p_1(A), p_2(A)\}$ then the system contains six equa-

⁴Recall that h(x, y, z) is positive if the points x, y, z (in that order) are arranged counter-clockwise on the complex plane, negative if they are arranged clockwise, and zero if they are collinear.

tions, in addition to the inequalities $Bu \ge 0$:

$$\begin{array}{rcl}
\alpha_1^n &=& u_1 p_1(\alpha_1) + u_2 p_2(\alpha_1) \\
\alpha_2^n &=& u_1 p_1(\alpha_2) + u_2 p_2(\alpha_2) \\
n\alpha_2^{n-1} &=& u_1 p_1'(\alpha_2) + u_2 p_2'(\alpha_2) \\
\alpha_3^n &=& u_1 p_1(\alpha_3) + u_2 p_2(\alpha_3) \\
n\alpha_3^{n-1} &=& u_1 p_1'(\alpha_3) + u_2 p_2'(\alpha_3) \\
n(n-1)\alpha_3^{n-2} &=& u_1 p_1''(\alpha_3) + u_2 p_2''(\alpha_3)
\end{array}$$

Notice also that we may assume without loss of generality that 0 is not an eigenvalue. Otherwise, its equations in the Master System $0 = u_1 p_1^{(j)}(0) + \cdots + u_d p_d^{(j)}(0)$ either yield a linear dependence on u_1, \ldots, u_d , allowing us to eliminate some u_i and proceed inductively by solving a lower-dimensional Master System, or are trivially satisfied by all u_1, \ldots, u_d and may be dismissed.

C.2 Equivalence classes of \sim Next, we focus on the equivalence relation \sim on the eigenvalues of the input matrix defined by

$$\alpha \sim \beta \iff \alpha/\beta$$
 is a root of unity

The image of an equivalence class of \sim under complex conjugation is also an equivalence class of \sim . If a class is its own image under complex conjugation, then it is called *self-conjugate*. Classes which are not self-conjugate are grouped into *pairs of conjugate classes* which are each other's image under complex conjugation.

If a class C is self-conjugate, then we can write it as

$$\mathcal{C} = \{\alpha\omega_1, \alpha\omega_2, \dots, \alpha\omega_s\}$$

where α is real algebraic and $\omega_1, \ldots, \omega_s$ are roots of unity. This representation is easily computable in polynomial time. Similarly, if two classes C_1, C_2 are each other's image under complex conjugation, they can be written as

$$C_1 = \{\alpha\omega_1, \alpha\omega_2, \dots, \alpha\omega_s\}$$
$$C_2 = \{\overline{\alpha\omega_1}, \overline{\alpha\omega_2}, \dots, \overline{\alpha\omega_s}\}$$

where α is algebraic and $\arg(\alpha)$ is not a rational multiple of 2π . For an equivalence class C of \sim , write Eq(C, j)for the set of *j*-th derivative equations contributed to the Master System by eigenvalues in C. Define also the *multiplicity* of C to be the maximum multiplicity of an eigenvalue in C.

In our work on the Orbit Problem [5], we analysed the equivalence classes of \sim in order to derive a bound on the exponent *n*. We were able to show that if *A* has 'sufficiently many' eigenvalues unrelated by \sim then just the condition $A^n \in span\{p_1(A), \ldots, p_d(A)\}$ on its own is strong enough to bound the exponent, regardless of the linear inequalities $Bu \ge 0$ which the Extended Problem imposes on the coefficients u_1, \ldots, u_d . The following theorem will allow us to focus only on the cases in which \sim has 'few' equivalence classes.

THEOREM C.1. Suppose we are given a problem instance (A, B, p_1, \ldots, p_d) with $d \leq 3$ and let \sim be the relation on the eigenvalues of A defined as above. Write $||I|| = ||A|| + ||p_1|| + \cdots + ||p_d||$. Let R be the sum of the multiplicities of the equivalence classes of \sim . Then if $R \geq d + 1$, then there exists an effectively computable bound $N \in 2^{||I||^{O(1)}}$ such that if $A^n \in$ span({ $p_1(A), \ldots, p_d(A)$ }), then $n \leq N$. Moreover, if d = 1, then $N \in ||I||^{O(1)}$.

C.3 Case analysis on the residue of n Let Lbe the least common multiple of all the orders of the ratios of eigenvalues of A which are roots of unity. Notice that $L \in 2^{\|I\|^{O(1)}}$. In the two- and threedimensional Extended Orbit Problem, we will perform a case analysis on the residue of n modulo L. We will show that for each fixed residue of n, we can either solve the problem instance directly or derive an effective bound N such that any witness n to the problem instance must be bounded above by N. Since L is at most exponentially large, it may be expressed using at most polynomially many bits. Thus, when the relation \sim has too few equivalence classes for Theorem C.1 to apply, our polynomial-space algorithm can guess the residue of $n \mod L$. This greatly simplifies the Master System and either allows us to solve it outright or to reduce it to an instance of the Simultaneous Positivity Problem.

We now consider what happens to the equations in $Eq(\mathcal{C}, j)$ for a fixed residue of n modulo L. Let $\mathcal{C} = \{\alpha \omega_1, \ldots, \alpha \omega_s\}$ and for simplicity consider $Eq(\mathcal{C}, 0)$:

$$(\alpha\omega_1)^n = \sum_{i=1}^d u_i p_i(\alpha\omega_1)$$

...
$$(\alpha\omega_s)^n = \sum_{i=1}^d u_i p_i(\alpha\omega_s)$$

This set of equations is equivalent to

(C.7)
$$\alpha^n = \sum_{i=1}^d u_i \frac{p_i(\alpha \omega_1)}{\omega_1^n} = \dots = \sum_{i=1}^d u_i \frac{p_i(\alpha \omega_s)}{\omega_s^n}$$

For a fixed residue of n modulo L, we see $\omega_1^n, \ldots, \omega_s^n$ are also fixed, so each $p_i(\alpha \omega_j)/\omega_i^n$ is easily computable.

Observe that (C.7) is equivalent to the conjunction of an equation with a linear system:

(C.8)
$$\alpha^n = \sum_{i=1}^d u_i \frac{p_i(\alpha \omega_s)}{\omega_s^n} \text{ and } B'u = 0$$

where B' is an $(s-1) \times k$ matrix over \mathbb{A} defined by

$$B'_{j,i} = \frac{p_i(\alpha\omega_j)}{\omega_j^n} - \frac{p_i(\alpha\omega_{j+1})}{\omega_{j+1}^n}$$

Writing φ_i for $p_i(\alpha \omega_s)/\omega_s^n$ and considering separately the real and imaginary parts of B'u = 0, we see that (C.8) is equivalent to

$$\alpha^n = \varphi_1 u_1 + \dots + \varphi_d u_d$$
 and $B'' u = 0$

where

$$B^{\prime\prime} = \left[\begin{array}{c} Re(B^{\prime}) \\ Im(B^{\prime}) \end{array} \right]$$

is a $2(s-1) \times k$ matrix over $\mathbb{R} \cap \mathbb{A}$. However, u lies in the nullspace of B'' if and only if u is orthogonal to the column space of B''. Thus, if B'' has nonzero column rank, then u_1, \ldots, u_d must have a nontrivial linear dependence $\psi_1 u_1 + \cdots + \psi_d u_d = 0$ for effectively computable $\psi_1, \ldots, \psi_d \in \mathbb{R} \cap \mathbb{A}$. Therefore, we can eliminate some coefficient u_i , replacing all of its occurrences in the Master System (C.6), and proceed inductively to solve a Master System with dimension d-1. Therefore, we can assume that the column rank of B'' is zero, so the constraint B''u = 0 is satisfied by all vectors u.

Thus, for this particular residue of n modulo L, the equations $Eq(\mathcal{C}, 0)$ are equivalent to the single equation $\alpha^n = \varphi_1 u_1 + \cdots + \varphi_d u_d$. Further, if the equivalence class \mathcal{C} is self-conjugate, then $\alpha \in \mathbb{R} \cap \mathbb{A}$, so we may replace each φ_i with its real part and assume $\varphi_i \in \mathbb{R} \cap \mathbb{A}$. Similarly, for j > 0 and a fixed residue of n modulo L, the equations $Eq(\mathcal{C}, j)$ reduce to the equivalent single equation

$$n(n-1)\dots(n-j+1)\alpha^{n-j} = \sum_{i=1}^d u_i \frac{p_i^{(j)}(\alpha\omega_s)}{\omega_s^{n-j}}$$

C.4 One-dimensional case of Extended Orbit In the one-dimensional Extended Orbit Problem, we have to decide whether there exists some $n \in \mathbb{N}$ such that A^n is a non-negative multiple of $p_1(A)$. We show this problem is in **PTIME**.

Begin by observing that if 0 is an eigenvalue of A, then its equations in the Master System are either satisfied for all values of n, or for no values of n. In the former case, they can be discarded, whereas in the latter

case, the problem instance is immediately negative. We will now perform a case analysis on the number of equivalence classes of \sim .

Two or more equivalence classes. When the relation ~ has at least two equivalence classes, by Theorem C.1, there exists a computable bound $N \in ||I||^{O(1)}$ on the exponent *n*. It suffices to try all $n \leq N$, which can be done in polynomial time.

One equivalence class, all roots simple. The second case is when \sim has only one equivalence class and the eigenvalues $\alpha_1, \ldots, \alpha_s$ of A are all simple in the minimal polynomial of A. The Master System is then equivalent to

$$u_1 = \frac{\alpha_1^n}{p_1(\alpha_1)} \ge 0$$
$$\left(\frac{\alpha_i}{\alpha_j}\right)^n = \frac{p_1(\alpha_i)}{p_1(\alpha_j)} \text{ for all } i, j$$

Since all ratios α_i/α_i are roots of unity, each equation $(\alpha_i/\alpha_j)^n = p_1(\alpha_i)/p_1(\alpha_j)$ is either unsatisfiable, making the problem instance immediately negative, or equivalent to some congruence in n. If all equations are satisfiable, then $A^n \in span\{p_1(A)\}$ holds if and only if $n \equiv t_1 \mod t_2$, where t_1, t_2 are effectively computable natural numbers. Moreover, since \sim has only one equivalence class, it must necessarily be self-conjugate, so $\alpha_1 = |\alpha_1|\omega$ for some root of unity ω which can be calculated easily. Since $u_1 = Re(\alpha_1^n)/Re(p_1(\alpha_1))$, we can compute what the sign of $Re(\alpha_1^n)$ should be to ensure $u_1 \geq 0$, that is, whether ω^n must be 1 or -1 for n to be a witness. This leads to another congruence in n which we put in conjunction with $n \equiv t_1 \mod t_2$. The problem instance is positive iff the two congruences have a common solution.

One equivalence class, some repeated roots. As in the previous case, we take the ratios of all pairs of equations $\alpha_i^n = u_1 p_1(\alpha_i)$ and $\alpha_j^n = u_1 p_1(\alpha_j)$, giving

(C.9)
$$\left(\frac{\alpha_i}{\alpha_j}\right)^n = \frac{p_1(\alpha_i)}{p_1(\alpha_j)} \text{ for all } i, j$$

 (\cdot)

Additionally, for each repeated root α_i , we take the ratios of its first and second equation, of its second and third equation, and so on, obtaining

$$\frac{\alpha_i}{n-j} = \frac{p_1^{(j)}(\alpha_i)}{p_1^{(j+1)}(\alpha_i)} \text{ for all } j \in \{0, \dots, mul(\alpha_i) - 1\}$$

If the equations (C.10) point to different values of n, then the problem instance is negative. If they point to the same value of n, but n does not satisfy

the congruence resulting from (C.9), then the problem instance is negative. Otherwise, the problem instance is positive if and only if $u_1 = \alpha_1^n/p_1(\alpha_1)$ is positive. The relation ~ has only one equivalence class, so it must be self-conjugate, so $\alpha_1 = |\alpha_1|\omega$ for some computable root of unity ω . It is easy to check the sign of ω^n , so the decision method is complete.

C.5 Two-dimensional case of Extended Orbit Now suppose we have a problem instance (A, B, p_1, p_2) and we have to determine whether there exist an exponent $n \in \mathbb{N}$ and coefficients $u = (u_1, u_2) \in \mathbb{Q}^2$ such that

$$A^{n} = u_{1}p_{1}(A) + u_{2}p_{2}(A)$$
 and $Bu \ge 0$

We will perform a case analysis on the equivalence classes of \sim . By Theorem C.1, if the sum of the multiplicities of the equivalence classes of \sim is at least 3, then there exists an effective bound $N \in 2^{\|I\|^{O(1)}}$ on *n* such that for n > N, mere membership of A^n in $span(\{p_1(A), p_2(A)\}$ is impossible, regardless of the constraints on the coefficients u_1, u_2 . Then an exponent $n \leq N$ can be chosen nondeterministically and verified using a **PosSLP** oracle. We consider the remaining cases.

One simple equivalence class. Suppose \sim has only one equivalence class and its eigenvalues are all simple in the minimal polynomial of the matrix. We proceed by case analysis on the residue of n, as in Section C.3. For a fixed residue, the Master System reduces to

(C.11)
$$\alpha^n = u_1 \varphi_1 + u_2 \varphi_2 \text{ and } Bu \ge 0$$

where $\varphi_1, \varphi_2 \in \mathbb{R} \cap \mathbb{A}$. Fix the parity of n and therefore assume $\alpha > 0$ by including its sign into φ_1, φ_2 . Now observe that either all values of n satisfy (C.11), or no value of n does. Indeed, if n is a witness with coefficients (u_1, u_2) , then n + 1 and n - 1 are also witnesses, with coefficients $(u_1\alpha, u_2\alpha)$ and $(u_1/\alpha, u_2/\alpha)$, respectively. Therefore, it suffices to try n = 0. This leads to a conjunction of the equation $1 = u_1\varphi_1 + u_2\varphi_2$ with inequalities in u_1, u_2 , which is easy to solve.

Two simple equivalence classes. Suppose that \sim has two equivalence classes and all eigenvalues are simple in the minimal polynomial of the matrix. Proceed by case analysis on the residue of n as before and reduce the Master System to

(C.12)
$$\begin{bmatrix} \alpha^n \\ \beta^n \end{bmatrix} = \begin{bmatrix} \varphi_1 & \varphi_2 \\ \varphi_3 & \varphi_4 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$$
 and $Bu \ge 0$

If the equivalence classes are both self-conjugate, then $\varphi_1, \ldots, \varphi_4, \alpha, \beta$ are all real algebraic, otherwise $\varphi_3 = \overline{\varphi_1}$,

 $\varphi_4 = \overline{\varphi_2}$ and $\alpha = \overline{\beta}$. If the 2 × 2 matrix in (C.12) is invertible, then premultiplying by its inverse yields

$$\left[\begin{array}{c} u_1\\ u_2 \end{array}\right] = \left[\begin{array}{c} \psi_1 & \psi_2\\ \psi_3 & \psi_4 \end{array}\right] \left[\begin{array}{c} \alpha^n\\ \beta^n \end{array}\right] \text{ and } Bu \ge 0$$

where either ψ_1, \ldots, ψ_4 are real, or $\psi_2 = \overline{\psi_1}$ and $\psi_4 = \overline{\psi_3}$. Now observe that u_1, u_2 satisfy a linear recurrence formula with characteristic equation $(x - \alpha)(x - \beta) = 0$. Then Bu is a vector of linear recurrence sequences over $\mathbb{R} \cap \mathbb{A}$. Each sequence $\mathcal{S}_i(n)$ has order at most 2 and is given by

$$\mathcal{S}_i(n) = a_i \alpha^n + b_i \beta^n$$

so they all satisfy the same shared recurrence formula. Further, observe that these sequences are nondegenerate, since α/β is not a root of unity. Therefore, for this particular residue of n, the problem instance reduces to Simultaneous Positivity for sequences of order at most 2. Finally, if the 2×2 matrix in (C.12) is singular, then there is a non-trivial linear combination of the rows which equates to zero. Then the same nontrivial combination of α^n, β^n equals zero. A bound on n follows from Theorem B.2.

One repeated equivalence class. The last remaining case is when there is only one equivalence class of \sim and it contains at least one eigenvalue repeated in the minimal polynomial of A. This reduces to Simultaneous Positivity in the same way as the previous case, but the resulting recurrence sequences have characteristic equation $(x - \alpha)^2 = 0$ and are given by $S_i(n) = (a_i + b_i n)\alpha^n$.

C.6 Three-dimensional case of Extended Orbit Now we consider an instance of the Extended Orbit Problem with a three-dimensional target space. For given (A, B, p_1, p_2, p_3) , we need to determine whether there exist $n \in \mathbb{N}$ and $u = (u_1, u_2, u_3) \in \mathbb{Q}^3$ such that

$$A^n = u_1 p_1(A) + u_2 p_2(A) + u_3 p_3(A)$$
 and $Bu \ge 0$

The strategy is again to show an effective bound N such that if there is a witness (n, u_1, u_2, u_3) to the problem instance, then n < N. By Theorem C.1, we need only bound n in the cases when the multiplicities of the equivalence classes sum to at most 3.

Three simple equivalence classes. If there are exactly three classes, each of multiplicity 1, one must necessarily be self-conjugate whereas the other two can be either self-conjugate or each other's conjugates. Either way, this case is analogous to the case of two simple equivalence classes in the two-dimensional version. After performing a case analysis on the residue of n, we obtain

(C.13)
$$\begin{bmatrix} \alpha^n \\ \beta^n \\ \gamma^n \end{bmatrix} = Tu \text{ and } Bu \ge 0$$

where T is a 3×3 matrix over $\mathbb{R} \cap \mathbb{A}$. If T is invertible, then we multiply both sides of (C.13) by T^{-1} and see that u_1, u_2, u_3 are linear recurrence sequences over $\mathbb{R} \cap \mathbb{A}$ with characteristic roots α, β, γ . Then the lefthand side of each linear inequality $Bu \geq 0$ is also an LRS over $\mathbb{R} \cap \mathbb{A}$ and has order 3. Thus the problem instance reduces to Simultaneous Positivity for order-3 sequences. On the other hand, if T is singular, then a linear combination of its rows is zero, so the same linear combination of $\alpha^n, \beta^n, \gamma^n$ is also zero. Noting that no two of α, β, γ are related by \sim , we obtain a bound on nfrom Theorem B.2.

Two classes, one simple and one repeated. Next, suppose \sim has two equivalence classes, one of multiplicity 1 and the other of multiplicity 2. This is analogous to the previous case. For a fixed residue of nmodulo L, the Master System is equivalent to

(C.14)
$$\begin{bmatrix} \alpha^n \\ n\alpha^{n-1} \\ \beta^n \end{bmatrix} = Tu \text{ and } Bu \ge 0$$

where T is a 3×3 matrix over $\mathbb{R} \cap \mathbb{A}$. Now if T is invertible, then we multiply both sides of (C.14) by T^{-1} and see that each of u_1, u_2, u_3 is a linear recurrence sequence over $\mathbb{R} \cap \mathbb{A}$ with characteristic equation $(x - \alpha)^2(x - \beta) = 0$. Substituting into the homogeneous linear inequalities $Bu \ge 0$, we now have an instance of the Simultaneous Positivity Problem for LRS of order 3 with a repeated characteristic root. If T is singular, then a linear combination of α^n , $n\alpha^{n-1}$ and β^n must equal zero, so a bound on n follows from Theorem B.2, because the ratio of α and β is not a root of unity.

One simple equivalence class. Suppose now that \sim has only one equivalence class and it has multiplicity 1. The situation is analogous to the same case in the two-dimensional version. We have to find n, u_1, u_2, u_3 such that

(C.15)
$$\alpha^n = u_1\varphi_1 + u_2\varphi_2 + u_3\varphi_3 \text{ and } Bu \ge 0$$

Since everything is real, we observe that either all n are witnesses to the problem instance, or none are, so it suffices to consider n = 0, reducing the problem to a conjunction of the linear inequalities $Bu \ge 0$ with the equation $1 = u_1\varphi_1 + u_2\varphi_2 + u_3\varphi_3$.

Two equivalence classes, both simple. Let \sim have two equivalence classes, both of multiplicity 1. For a fixed residue of n modulo L, the Master System is

equivalent to

$$\left[\begin{array}{c} \alpha^n\\ \beta^n \end{array}\right] = Tu \text{ and } Bu \ge 0$$

where T is a 2×3 matrix. All the numbers involved are algebraic. There are two possibilities: either α , β and T are in $\mathbb{R} \cap \mathbb{A}$, or $\alpha = \overline{\beta}$ and the second row of T is the complex conjugate of the first row.

The dimension of the column space of T is 0, 1 or 2. If the dimension of the column space is 0, then the Master System is unsatisfiable, since T maps everything to zero, whereas α^n and β^n cannot be zero. If the dimension of the column space of T is 1, then it is spanned by a single vector (t_1, t_2) . If at least one of t_1, t_2 is zero, then the System is unsatisfiable, because $\alpha, \beta \neq 0$. Otherwise, we can conclude that $(\alpha/\beta)^n =$ t_1/t_2 . Since α/β is not a root of unity, a bound on nwhich is polynomial in ||I|| follows by Theorem B.2.

Assume therefore that the dimension of the column space of T is 2. We consider the real and the complex cases separately. First, suppose T, α, β are real. Each of the inequalities $Bu \ge 0$ specifies that (u_1, u_2, u_3) lies in a halfspace \mathcal{H}_i of \mathbb{R}^3 . The image of each \mathcal{H}_i under T can be the entire plane \mathbb{R}^2 , a half-plane, a line, or a half-line. Each of these images is easy to calculate in polynomial time. If for some i, the image $T\mathcal{H}_i$ is a line or a half-line, with defining vector (t_1, t_2) , then by the same reasoning as above, we see $(\alpha/\beta)^n = t_1/t_2$ and hence obtain a bound on n from Theorem B.2. Otherwise, we can assume that for all i, $T\mathcal{H}_i$ is a halfplane $\{(x,y) : A_ix + B_iy \geq 0\}$ with effectively computable $A_i, B_i \in \mathbb{R} \cap \mathbb{A}$. We have to determine whether there exists $n \in \mathbb{N}$ such that (α^n, β^n) lies in the intersection of these halfplanes. Noting that $A_i \alpha^n + B_i \beta^n$ as a function of n is a linear recurrence sequence over $\mathbb{R} \cap \mathbb{A}$ which has order 2, we see that this is now an instance of the Simultaneous Positivity Problem, so we are done by Theorem 4.1.

Suppose now that α and β are complex conjugates, and the second row of T is the complex conjugate of the first. We may freely assume that $|\alpha| = |\beta| = 1$, since if the inequalities are satisfied by (u_1, u_2, u_3) , then they are also satisfied by $(u_1/|\alpha|^n, u_2/|\alpha|^n, u_3/|\alpha|^n)$. The image under T of each halfspace \mathcal{H}_i is a homogeneous cone in the complex plane. The same is true of the intersection $G = \bigcap_i T \mathcal{H}_i$ of these cones, which may in fact be computed explicitly. We need to determine whether there exists $n \in \mathbb{N}$ such that $\alpha^n \in G$. Notice that $\{\alpha^n : n \in \mathbb{N}\}$ is dense on the unit circle. The intersection of the unit circle with G could be a single point, or an arc.

Representing real and imaginary parts with variables over \mathbb{R} , we construct a sentence τ in the first-order

theory of the reals which states that the intersection of G with the unit circle is a single point. We check the validity of τ , this can be done in polynomial time by Theorem A.2. If τ is false, then G intersects the unit circle in an arc, so by the density of α^n on the unit circle, the Master System is satisfiable. Otherwise, the intersection is a single point $z \in \mathbb{C}$. Moreover, this point is effectively computable - Renegar's algorithm hinges on quantifier elimination, and will produce a quantifier free formula containing exactly the minimal polynomials of Re(z) and Im(z). The procedure is polynomial-time, so $||z|| \in ||I||^{O(1)}$. Now the Master System is satisfiable if and only if there exists $n \in \mathbb{N}$ such that $\alpha^n = z$. As α and z both have descriptions polynomial in the input size and α is not a root of unity, we see there exists a polynomial bound on n from Theorem B.2.

One repeated equivalence class. Finally, suppose \sim has a single equivalence class and its multiplicity is 2. Then for a fixed residue of n modulo L, the Master System is equivalent to

$$\left[\begin{array}{c} \alpha^n\\ n\alpha^{n-1} \end{array}\right] = Tu \text{ and } Bu \ge 0$$

where α and T are both real algebraic. This is now handled analogously to the previous case for a real Tand reduces to Simultaneous Positivity for LRS with characteristic equation $(x - \alpha)^2 = 0$.

References

- Eric Allender, Peter Bürgisser, Johan Kjeldgaard-Pedersen, and Peter Bro Miltersen. On the complexity of numerical analysis. In Proc. 21st ann. IEEE Conf. on Computational Complexity (CCC), pages 331–339, 2006.
- [2] Alan Baker and Gisbert Wüstholz. Logarithmic forms and group varieties. *Jour. Reine Angew. Math.*, 442:19–62, 1993.
- [3] A. M. Ben-Amram and S. Genaim. Ranking functions for linear-constraint loops. *Journal of the ACM (to appear)*, 2014.
- [4] M. Braverman. Termination of integer linear programs. In Proceedings of the 18th International Conference on Computer Aided Verification, CAV, LNCS 4144, pages 372–385. Springer, 2006.
- [5] Ventsislav Chonev, Joël Ouaknine, and James Worrell. On the complexity of the orbit problem. *CoRR*, abs/1303.2981, 2013.
- [6] Ventsislav Chonev, Joël Ouaknine, and James Worrell. The orbit problem in higher dimensions. In STOC, pages 941–950. ACM, 2013.
- [7] H. Cohen. A Course in Computational Algebraic Number Theory. Springer, 1993.

- [8] Thomas W. Cusick and Mary E. Flahive. *The Markoff* and Lagrange Spectra. American Mathematical Society, 1989.
- [9] Graham Everest, Alf van der Poorten, Thomas Ward, and Igor Shparlinski. *Recurrence Sequences*. American Mathematical Society, 2003.
- [10] Branko Grünbaum, Victor Klee, Micha A Perles, and Geoffrey Colin Shephard. *Convex polytopes*. Springer, 1967.
- [11] V. Halava, T. Harju, M. Hirvensalo, and J. Karhumäki. Skolem's problem – on the border between decidability and undecidability. *TUCS Technical Report*, (683), 2005.
- [12] M. Harrison. Lectures on sequential machines. Academic Press, Orlando, 1969.
- [13] R. Kannan and R. Lipton. Polynomial-time algorithm for the orbit problem. *Journal of the ACM*, 33(4):808– 821, 1986.
- [14] Ravindran Kannan and Richard J. Lipton. The orbit problem is decidable. In *Proceedings of the twelfth annual ACM symposium on Theory of computing*, STOC, pages 252–261. ACM, 1980.
- [15] D. Lee and M. Yannakakis. Online minimization of transition systems (extended abstract). In *Proceedings* of the 24th annual ACM symposium on Theory of Computing, STOC, pages 264–274. ACM, 1992.
- [16] Peter McMullen and Geoffrey Colin Shephard. Convex polytopes and the upper bound conjecture, volume 3. CUP Archive, 1971.
- [17] M. Mignotte. Some useful bounds. Computer Algebra, pages 259–263, 1982.
- [18] J. Ouaknine, J. S. Pinto, and J. Worrell. On termination of integer linear loops. In *Proceedings of SODA*. ACM-SIAM, 2015.
- [19] Joël Ouaknine and James Worrell. Decision problems for linear recurrence sequences. In Alain Finkel, Jérôme Leroux, and Igor Potapov, editors, *Reachability Problems*, volume 7550 of *Lecture Notes in Computer Science*, pages 21–28. Springer Berlin Heidelberg, 2012.
- [20] Joël Ouaknine and James Worrell. Positivity problems for low-order linear recurrence sequences. *Proceedings* of SODA, 2014.
- [21] V. Pan. Optimal and nearly optimal algorithms for approximating polynomial zeros. Computers & Mathematics with Applications, 31(12):97 – 138, 1996.
- [22] James Renegar. On the computational complexity and geometry of the first-order theory of the reals. part i: Introduction. preliminaries. the geometry of semialgebraic sets. the decision problem for the existential theory of the reals. *Journal of Symbolic Computation*, 13(3):255 – 299, 1992.
- [23] T. Tao. Structure and randomness: pages from year one of a mathematical blog. American Mathematical Society, 2008.
- [24] S. P. Tarasov and M. N. Vyalyi. Orbits of linear maps and regular languages. In *Proc. Intern. Comp. Sci. Symp. in Russia (CSR)*, volume 6651 of *LNCS*. Springer, 2011.

- [25] Alfred Tarski. A decision method for elementary algebra and geometry. 1951.
- [26] Günter M Ziegler. Lectures on polytopes, volume 152. Springer, 1995.