The Two-Sided Game of Googol and Sample-Based Prophet Inequalities

José Correa, Andrés Cristi, Boris Epstein, and José A. Soto

Universidad de Chile

Abstract

The secretary problem or the game of Googol are classic models for online selection problems that have received significant attention in the last five decades. In this paper we consider a variant of the problem and explore its connections to data-driven online selection. Specifically, we are given n cards with arbitrary non-negative numbers written on both sides. The cards are randomly placed on n consecutive positions on a table, and for each card, the visible side is also selected at random. The player sees the visible side of all cards and wants to select the card with the maximum hidden value. To this end, the player flips the first card, sees its hidden value and decides whether to pick it or drop it and continue with the next card.

We study algorithms for two natural objectives. In the first one, similar to the secretary problem, the player wants to maximize the probability of selecting the maximum hidden value. We show that this can be done with probability at least 0.45292. In the second objective, similar to the prophet inequality, the player wants to maximize the expectation of the selected hidden value. Here we show a guarantee of at least 0.63518 with respect to the expected maximum hidden value.

Our algorithms result from combining three basic strategies. One is to stop whenever we see a value larger than the initial n visible numbers. The second one is to stop the first time the last flipped card's value is the largest of the currently n visible numbers in the table. And the third one is similar to the latter but to stop it additionally requires that the last flipped value is larger than the value on the other side of its card.

We apply our results to the prophet secretary problem with unknown distributions, but with access to a single sample from each distribution. In particular, our guarantee improves upon 1 - 1/e for this problem, which is the currently best known guarantee and only works for the i.i.d. prophet inequality with samples.

1 Introduction

In the classic game of Googol we are given n cards with n different arbitrary positive numbers written on them. The cards are shuffled and spread on a table with the numbers facing down. The cards are flipped one at a time, in a random uniform order, and we have to decide when to stop. The goal is to maximize the probability that the last flipped card has the overall greatest number.

In this paper we study a variant of this problem that we call *The two-sided game of Googol*. Similar to the classic version, we are given n cards that we have to flip in random uniform order. However, here the cards have numbers on both sides, so we have 2n different arbitrary non-negative numbers instead of n, written on each side of each card. The cards are shuffled and spread on a table so that, independently for each card, either side faces up with probability 1/2. We can see all the numbers that landed facing up (while the other side is hidden), and flip one card at a time, revealing the number that was facing down. Again we have to decide when to stop. We study the *secretary* and the *prophet* variants. In the secretary variant, the goal is to maximize the probability of stopping at the maximum number over the numbers that landed facing down. In the prophet variant, the goal is to maximize the ratio between the expectation of the last number revealed before stopping, and the expected maximum of the numbers that landed facing down.

This problem naturally fits within the theory of optimal stopping theory which is concerned with choosing the right time to take a particular action, so as to maximize the expected reward. Two landmark examples within this theory are the secretary problem (or game of Googol) described above, and the prophet inequality. In the latter a gambler faces a finite sequence of non-negative independent random variables X_1, \ldots, X_n with known distributions F_i from which iteratively a prize is drawn. After seeing a prize, the gambler can either accept the prize and the game ends, or reject the prize and the next prize is presented to her. The classical result of Krengel and Sucheston, and Garling [22, 23], states that the gambler can obtain at least half of the expected reward that a prophet can make who knows the realizations of the prizes beforehand. That is, $\sup\{\mathbb{E}[X_t] : t \text{ stopping rule }\} \geq \frac{1}{2}\mathbb{E}\{\sup_{1\leq i\leq n} X_i\}$. Moreover, Krengel and Sucheston also showed that this bound is best possible. Remarkably, Samuel-Cahn [26] showed that the bound of 1/2 can be obtained by a simple threshold rule, which stops as soon as a prize is above a fixed threshold. We refer the reader to the survey by Hill and Kertz [19] for further classic results.

In recent years, motivated by the connections between optimal stopping and posted price mechanisms [17, 4, 8], there has been a regained interest in studying algorithms for variants of the classic prophet inequalities. The recent survey of Lucier [25] is a good starting point to understand the economic perspective of prophet inequalities, while the recent letter of Correa et al. [7] provides a recent overview of results for single choice prophet inequalities. Due to this regained interest, new variants of the prophet inequality setting have been studied, including problems when the gambler has to select multiple items, when the selected set has to satisfy combinatorial constraints, when the market is large, when prior information is inaccurate, among many others (see e.g. [1, 21, 11, 13, 12]).

Particularly relevant to our work in the *prophet secretary* problem. In this version, the random variables are shown to the gambler in uniform random order, as in the secretary problem. The problem was first studied by Esfandiari et al. [14] who found a bound of 1 - 1/e. Later, Eshani et al. [13] showed that the bound of 1 - 1/e can even be achieved using a single threshold. The factor 1 - 1/e was first beaten by Azar et al [3] by a tiny margin, while the current best bound is 0.67 [9]. In terms of upper bounds it was unclear until very recently whether there was a separation between prophet secretary and the i.i.d prophet inequality, where the random variables are identically distributed. For this problem, Hill ad Kertz [18] provided the family of worst possible instances from which [20] proved the largest possible bound one could expect is 0.7451 and Correa et al. [6] proved that this value is actually tight. Very recently, Correa et al. show that no algorithm can achieve an approximation factor better than $\sqrt{3} - 1 \approx 0.71$ for the prophet secretary problem (see the full version of [9]). Interestingly, the tight is still unknown.

On the other hand, some recent work has started to investigate data-driven versions of the prophet inequality since the full distributional knowledge seems quite strong for many applications. In this context, Azar et al. [2] consider a version in which the gambler only has access to one sample from each distribution. They prove a prophet inequality with an approximation guarantee of 1/4 for

this problem and left open whether achieving 1/2 is possible. This question was recently answered on the positive by Wang [27] who uses an elegant approach to prove that just taking the maximum of the samples as threshold leads to the optimal guarantee. If the instance is further specialized to an i.i.d. instance, the best known bound is 1 - 1/e while it is known that one cannot achieve a guarantee better than $\ln(2) \approx 0.69$ [5].

Our problem is closely related to the single-sample version of the secretary problem and the prophet secretary problem, which combine the data-driven approach of Azar et al [2] and the random arrival order of Esfandiari et al. [14]. In these problems we are given n distributions, which are unknown to us, and only have access to a single sample from each. Then n values are drawn, one from each distribution, and presented to us in random order. When we get to see a value, we have to decide whether to keep it and stop, or to drop it and continue. Again in the single-sample secretary problem the goal is to maximize the probability of stopping with the largest value while in the single-sample prophet secretary problem the goal is to maximize the expectation of the value at which we stopped divided by the expectation of the maximum of the values.

It is immediate to observe that an algorithm for the two-sided game of Googol that has a guarantee of α , for either the secretary or prophet variants of the problem, readily implies the same approximation guarantee for the single-sample secretary problem and single-sample prophet secretary problem respectively [27]. Indeed, if we consider an instance of the two-sided game of Googol where the values on card *i* are independent draws from a distribution F_i we exactly obtain the single-sample secretary problems.

1.1 Our results

Most of our results come from analyzing three basic algorithms. The first is the *Open moving window* algorithm, in which we stop the first time the card just flipped is larger than all the currently visible values. The second is the *Closed moving window* algorithm, in which we additionally require that in the last flipped card the value just revealed is larger than the value that was visible before. Finally we consider the *Full window* algorithm in which we simply take the largest value initially visible as threshold (and therefore stop the first time we see a value larger than all values we have seen).

We first study the secretary variant of the two-sided game of Googol in which the goal is to maximize the probability of choosing the maximum hidden value. For this problem we prove that the closed window algorithm gets the maximum value with probability 0.4529. Of course, this value is more than 1/e which is the best possible for the classic secretary problem [10, 24, 15], but it is less than 0.5801 which is the best possible guarantee for the full information secretary problem, i.e., when the full distribution is known [16].

Next, we concentrate in the prophet variant of the two-sided game of Googol in which the goal is to maximize the ratio between the expectation of the chosen value, and that of the maximum hidden value. In this case we start by observing that all three algorithms described above can only give a guarantee of 1/2. Indeed, consider an instance with two cards. Card 1 has values 0 and 1 on each side while card 2 has the numbers ε and ε^2 . Clearly the expectation of the maximum hidden value is $1/2 + O(\varepsilon)$ (the value 1 is hidden with probability 1/2) while the open moving window algorithm gets in expectation $1/4 + O(\varepsilon)$ (to get the value of 1 the algorithm needs that it is hidden and that card 1 is the first card). For the other two algorithms, consider the instance in which card 1 has the values 1 and $1 - \varepsilon$ while card 2 has the values ε and 0. Clearly the expectation of the maximum hidden value is $1 - O(\varepsilon)$, but since now neither of the algorithms can stop when $1 - \varepsilon$ is

Instance		Prophet	Secretary	
		$(\mathbb{E}(ALG) \ge \alpha \mathbb{E}(OPT))$	$(\mathbb{P}(ALG = OPT) \ge \alpha)$	
Independent (Prophet Secretary)	Samples	$\frac{1}{2}$ [27] ≤ 0.635 [*] $\leq \alpha \leq \ln(2)$ [5]	$\frac{1}{e} [10] \le 0.452 [*] \le \alpha$	
	Known distributions	$0.669 \ [9] \le \alpha \le \sqrt{3} - 1 \ [9]$	$\leq 0.580 \ [16]$	
I.I.D.	Samples	$1 - \frac{1}{e} [5] \le 0.635 [*] \le \alpha \le \ln(2) [5]$		
	Known distribution	$0.745[6] \le \alpha \le 0.745$ [18, 20]	$\alpha = 0.580 \; [16]$	

Table 1: Summary of known results for prophet secretary and secretary problems in the singlesample and full information settings. Results marked [*] are proven in this paper.

revealed, both algorithms obtain $1/2 + O(\varepsilon)$. However, by randomizing the choice of the algorithm we significantly improve the approximation ratio.

Our main result in the paper is to show that a simple randomization of the three proposed algorithms achieves a guarantee of 0.635 > 1 - 1/e. Interestingly, our bound surpasses that of Azar et al [3] which until very recently was the best known bound for (the full information) prophet secretary. Furthermore, our bound also beats the bound of 1 - 1/e obtained by Correa et al. [5] for the i.i.d. single-sample prophet inequality. So our bound not only improves upon these known bounds, but also works in a more general setting. The key behind the analysis is a very fine description of the performance of each of the three basic algorithms. Indeed for each algorithm we exactly compute the probability that they obtain any of the 2n possible values.¹ With this performance distribution at hand it remains to set the right probabilities of choosing each so as to maximize the expectation of the obtained value.

To wrap-up the paper we consider a large data-set situation which naturally applies to a slightly restricted version of the single sample prophet secretary problem. The assumption states that, with high probability, the if we rank all 2n values from largest to smallest, the position of maximum minimum value a card lies far down the list (i.e., when ranking the 2n values the first few values correspond to a maximum in its card). This assumption holds for instance in the single-sample i.i.d. prophet inequality, under the large market assumption used by Abolhassan et al. [1], or whenever the underlying distributions of the prophet secretary instance overlap enough. For this variant we design an optimal moving window algorithm and prove that it achieves an approximation ratio of 0.642.

In table 1 we summarize our results and the previous best bounds for the problems considered in this paper.

1.2 Preliminaries and notation

Formal problem statement. We are given 2n different and arbitrary positive numbers, organized into n pairs that we denote as $\{(a_i, b_i)\}_{i=1}^n$, representing the numbers written in both sides of each card. These pairs of numbers are shuffled into sets \mathcal{U} and \mathcal{D} : for each i = 1, ..., n an independent

 $^{^{1}}$ In this respect, our result for the secretary variant can be seen as a warm up for the prophet variant of the problem.

dent unbiased coin is tossed, if the coin lands head, then $a_i = U_i \in \mathcal{U}$ and $b_i = D_i \in \mathcal{D}$, otherwise $b_i = U_i \in \mathcal{U}$ and $a_i = D_i \in \mathcal{D}$. The set $\mathcal{U} = \{U_1, ..., U_n\}$ represents the numbers that landed facing up, and the numbers in $\mathcal{D} = \{D_1, ..., D_n\}$ are those facing down. The numbers in \mathcal{U} are revealed, and a random uniform permutation $\sigma \in \Sigma_n$ is drawn. Then the elements in \mathcal{D} are revealed in n steps: at each step $i \in [n]$, the value $D_{\sigma(i)}$ and the index $\sigma(i)$ are revealed. After this, we must decide whether to stop or to continue to step i + 1.

We study algorithms (stopping rules) for the two variants. Let τ be the step at which we stop. In the secretary variant the objective is to maximize $\mathbb{P}(D_{\sigma(\tau)} = \max \mathcal{D})$, and in the prophet variant the objective is to maximize $\mathbb{E}(D_{\sigma(\tau)})/\mathbb{E}(\max \mathcal{D})$. Note that the latter is equivalent to just maximizing $\mathbb{E}(D_{\sigma(\tau)})$ since the algorithm does not control $\mathbb{E}(\max \mathcal{D})$.

Ranking and couples Our analyses rely of the ranking of the 2n numbers of the instance, so let us denote by $Y_1 > Y_2 > \cdots > Y_{2n}$ the 2n numbers in $\mathcal{U} \cup \mathcal{D}$ ordered from the largest to the smallest. We say that indices *i* and *j* are a couple, and denote it by $i \sim j$ if Y_i and Y_j are written in the two sides of the same card, i.e., if $(Y_i, Y_j) = (a_\ell, b_\ell)$ or $(Y_j, Y_i) = (a_\ell, b_\ell)$ for some $\ell \in [n]$. Let *k* be the smallest index so that *k* is the couple of some k' < k. In particular, $Y_1, Y_2, \ldots, Y_{k-1}$ are all written on different cards. Note that *k* and *k'* only depend on numbers in the cards², i.e., on the numbers $\{(a_i, b_i)\}_{i=1}^n$, and not on the coin tosses or on the permutation σ . We will restrict all our analysis to the numbers Y_1, \ldots, Y_k , which is easily justified by the observation that max \mathcal{D} always lies in the set $\{Y_1, \ldots, Y_k\}$.

Random arrival times. For most of our analyses it will be useful to consider the following reinterpretation of the randomness of both the coins tosses (hidden sides) and the random permutation (flipping order). Consider that each of the 2n numbers arrives in a random uniform time in the interval (-1, 1] as follows. For every index ℓ smaller than its couple ℓ' (i.e. $\ell \sim \ell', Y_{\ell} > Y_{\ell'}$) an independent random arrival time θ_{ℓ} uniform in (-1, 1] is sampled and ℓ' receives opposite arrival time $\theta_{\ell'} = C(\theta_{\ell})$, where

$$C(x) = \begin{cases} x - 1, & \text{if } x > 0, \\ x + 1, & \text{otherwise.} \end{cases}$$

The reinterpretation can the be done as follows. For each $j \in [2n]$, if $\theta_j > 0$, the number Y_j is facing down, i.e., $Y_j \in \mathcal{D}$, and if $\theta_j \leq 0$ then Y_j is facing up, i.e., $Y_j \in \mathcal{U}$. Therefore, we get to see all values whose corresponding arrival time is negative and the order in which we scan the hidden values $Y_j \in \mathcal{D}$ is increasing in θ_j .

Throughout the paper we use the term *value* to refer to the numbers written in the cards Y_1, Y_2, \ldots, Y_{2n} , while we use the term *element* to refer to a side of a particular card, that is an element in $\{U_1, \ldots, U_n, D_1, \ldots, D_n\}$. Of course these sets are the same so the point is that the *i*-th value corresponds to Y_i , whereas the *i*-th element corresponds to the *i*-th number in the list $U_{\sigma(1)}, \ldots, U_{\sigma(n)}, D_{\sigma(1)}, \ldots, D_{\sigma(n)}$.

^{2}This notation was introduced by Wang [27]

2 Basic Algorithms and statement of our results

Once the random coins for all cards and the random uniform permutation are all selected, we are left with the following situation. A list $E = (e_1, \ldots, e_{2n}) = (U_{\sigma(1)}, \ldots, U_{\sigma(n)}, D_{\sigma(1)}, \ldots, D_{\sigma(n)})$ of 2ndifferent positive elements are presented one-by-one to an algorithm. The algorithm must observe and skip the first n elements of the list, since they correspond to the values that landed facing up on the cards. The next n elements correspond to values that landed facing down (in such a way that the s-th and the s + n-th elements form a couple). The algorithm must decide immediately after observing the s-th element whether to select it and stop, or to continue with the next element.

We now present generic moving window algorithms for selecting one of the last n elements of a given list $E = (e_1, \ldots, e_{2n})$. These family of algorithms form the basis of the algorithms employed in this paper, which are described next.

Generic moving window algorithm. For every $s \in \{n+1, ..., 2n\}$, the algorithm first specifies, possibly in a random or implicit way, a window of elements

$$W_s = \{e_r \colon r \in \{\ell_s, \dots, s-1\}\}$$
(1)

scheduled to arrive immediately before s (the window may be empty). Then the algorithm proceeds as follows. It observes the first n elements without selecting them. For every $s \ge n+1$, the algorithm observes the s-th element, e_s , and selects it if e_s is larger than every element in its window³, i.e. if $e_s > \max W_s$. Otherwise, the algorithm rejects it and continues.

We now present three basic moving window algorithms for deciding when to stop. Each of them can be described by a sequence of left extremes $(\ell_s)_{s=n+1}^{2n}$ that will define the elements inside the window W_s .

Open moving window algorithm (ALG_o). This algorithm corresponds to the following strategy: flip the next unflipped card and accept its value x if and only if x is the largest of the n values that are currently visible (i.e., the *i*-th element is accepted if and only if it is larger than the n - i currently unflipped cards, and is also larger than the i - 1 previously flipped cards). In terms of the generic moving window, the left extreme of W_s is $\ell_s = s - n + 1$ for $n + 1 \le s \le 2n$.

Closed moving window algorithm (ALG_c). This algorithm works similarly to the previous one, with a slight difference. In this strategy, the element on the back of the last flipped card, the one that was facing up, is also required to be smaller than the element to be selected. In terms of the generic moving window, the left extreme of W_s is $\ell_s = s - n$ for $n + 1 \le s \le 2n$.

Full window algorithm (ALG_f). This algorithm corresponds to the algorithm presented by Wang [27]. It stops when the revealed element is larger than all element seen so far. Note that this is equivalent to stop with the first element that is larger than all elements that landed facing up. In terms of the generic moving window, the left extreme of W_s is $\ell_s = 1$ for $n + 1 \leq s \leq 2n$.

³If $W_s = \emptyset$ then we say that e_s is trivially selected.

Combining these three algorithms, we can obtain algorithm ALG_r^* , which picks ALG_o and ALG_c with probability (1 - r)/2 and ALG_f with probability r. Our main results are:

Theorem 1. For any instance of the two-sided game of Googol,

$$\mathbb{P}(ALG_c = \max \mathcal{D}) \ge \ln 2 \left(1 - \frac{\ln 2}{2}\right) \approx 0.45292.$$

Theorem 2. There exists a value $r^* \in [0,1]$ such that for any instance of the two-sided game of Googol, $\mathbb{E}(ALG_{r^*}^*) \geq \alpha \mathbb{E}(\max \mathcal{D})$ with $\alpha = \frac{4-5\ln 2}{5-6\ln 2} \approx 0.635184$.

The moving windows can also be seen from the perspective of the random arrival times. For ALG_o, the window for an element that arrives at time $\theta > 0$ can be defined as all elements arriving in $(\theta - 1, \theta)$. For ALG_c, this changes to $[\theta - 1, \theta)$, while for ALG_f it becomes $[-1, \theta)$. Therefore, we can define threshold functions⁴ for ALG_o, ALG_c, and ALG_f as $T_o(\theta) = \max\{Y_j : \theta_j \in (\theta - 1, \theta)\}$ and $T_f(\theta) = \max\{Y_j : \theta_j \in [-1, \theta)\}$, respectively. This motivates the definition of a fourth algorithm, ALG(t), for which the window for an element that arrives at time θ is defined as all elements that arrive in $[\max\{-1, \theta - 1 - t\}, \theta)$.

Our third result shows that for large k, this type of algorithms perform better.

Theorem 3. For $t^* \approx 0.1128904$, and any instance of the two-sided game of Googol, as k goes to infinity, $\mathbb{E}[ALG(t^*)]/\mathbb{E}[\max \mathcal{D}]$ is at least $R_{\infty}(t^*) \approx 0.6426317$.

To wrap up this section, we stablish two basic lemmas that will be useful later on. The first is used to estimate the probability of selecting any element arriving before a given element e_s . It is clear to see that all our algorithms satisfy the conditions.

Lemma 4 (Moving window lemma). Suppose the windows' left extremes $(\ell_s)_{s \in \{n+1,\dots,2n\}}$ specified by a moving window algorithm satisfy

$$\ell_{n+1} \le \ell_{n+2} \le \dots \le \ell_{2n} \le n+1 \tag{2}$$

Let s be an index with $n+1 \leq s$ and let $e_M = \max\{e_r : n+1 \leq r \leq s\}$ be the maximum value arriving between positions n+1 and s. The algorithm selects some element in the set $\{e_r : n+1 \leq r \leq s\}$ if and only if e_M is larger than every element in its window, i.e. $e_M > \max W_M$. In particular, if the window W_s of s is nonempty and $e_J = \max W_s$ then the algorithm does not stop strictly before e_s if and only if $J \leq n$ or $(J \geq n+1 \text{ and } e_J < \max W_J)$.

Proof. Suppose the algorithm selects some e_r with $n + 1 \leq r \leq s$. Note that e_r cannot arrive after e_M because in that case $e_M > e_r$ would be in the window W_r of e_r contradicting that e_r was selected. Then e_r must be e_M or an element arriving before e_M . In any case, by condition (2), $\ell_r \leq \ell_M$. Since e_r is selected, $e_r \geq \max\{e_t : \ell_r \leq t \leq r-1\} \geq \max\{e_t : \ell_M \leq t \leq n\}$. But then, as e_M is the largest element arriving between n + 1 and s, we also conclude that $e_M > \max\{e_t : \ell_M \leq t \leq M - 1\} = \max W_M$. For the converse suppose that $e_M > \max W_M$ then the algorithm will pick e_M unless it has already selected another element e_r arriving before e_M . In any case, the algorithm stops by selecting some element e_r , with $n + 1 \leq r \leq s$.

⁴These thresholds specify the minimum value under which the algorithm stops for each possible arrival time.

Let us show the second statement of the lemma. Suppose that the window of s is nonempty and let J be the index of the maximum element in its window. Let also, M' be the largest element arriving between steps n + 1 and s - 1. Since $e_{M'}$ is also in the window of s, we conclude that either $J \ge n + 1$ and M' = J, or $J \ge n$ and $e_J > e_{M'}$. Suppose $J \ge n + 1$. By the first part of the lemma, not stopping strictly before e_s is equivalent to $e_{M'} < \max W_{M'}$, and since M' = J, $e_J < \max W_J$. If on the other hand $J \le n$ then $e_J > e_{M'}$ and by assumption (2), $\ell_{M'} \le \ell_s \le J \le n + 1 \le M' \le s - 1$. We conclude that e_J is also in the window of M' and therefore $\max W_{M'} \ge e_J > e_{M'}$, which is equivalent to not stopping strictly before e_s .

As a final ingredient for our results we need the following lemma about independent uniform random variables in (-1, 1].

Lemma 5. Let X_0, X_1, \ldots, X_m be i.i.d. random variables distributed uniformly in (-1, 1]. Define the event $E : X_0 > 0$ and, for all $i \in \{1, \ldots, m\}, X_i \in [X_0 - 1, X_0)$. Conditioned on event E, $\{1 - X_0, X_1 + 1 - X_0, X_2 + 1 - X_0, \ldots, X_m + 1 - X_0\}$ is a family of m + 1 i.i.d. random variables distributed uniformly in [0, 1).

Proof. Conditioned on E and on the realization $a \in (0,1]$ of the variable X_0 , the family $\{X_i + 1 - a\}_{i \in \{1,...,m\}}$ is mutually independent. Since this is true for every realization a of X_0 , we can uncondition on a and deduce that the family $\{1 - X_0, X_1 + 1 - X_0, X_2 + 1 - X_0, \ldots, X_m + 1 - X_0\}$ is mutually independent only conditioned on E. Furthermore, since under E, X_0 distributes uniformly in (0, 1] we conclude that $1 - X_0$ is uniform in [0, 1). Finally, under E, for i > 0, X_i distributes uniformly in $[X_0 - 1, X_0)$, and therefore, $X_i + 1 - X_0$ distributes uniformly in [0, 1).

3 Maximizing the probability of picking the maximum

In this section we present a lower bound for the secretary variant of the two-sided game of Googol. Recall that the objective is to maximize the probability of stopping at the maximum value that landed facing down. If we were maximizing the probability of stopping the overall maximum value (the classical secretary problem), the best strategy [24, 10] would be to skip a constant number of elements (roughly 1/e) and then select the first element larger than all previous ones. Since we cannot select any number from the first half, we would be tempted to try the full window algorithm ALG_f that selects the first element in the second half that is larger than all previous ones. Unfortunately, this algorithm does not select any element if the top one, Y_1 , appears in the first half. It turns out that the best algorithm among the three basic algorithms presented is the closed moving window algorithm ALG_c that we analyze below.

Theorem 1. For any instance of the two-sided game of Googol,

$$\mathbb{P}(ALG_c = \max \mathcal{D}) \ge \ln 2\left(1 - \frac{\ln 2}{2}\right) \approx 0.45292.$$

To prove Theorem 1 the interpretation using random arrival times over (-1, 1] will be of use. The idea of the proof is to compute, for a fixed value k, the probability of winning, that is, selecting the largest number in \mathcal{D} . To compute this, we partition the event of winning by identifying which value Y_i is picked and which element Y_j is the largest inside its window. We find out that the worst case is when $k \to \infty$ and conclude the bound. Define the events E_1 : Y_j is the largest element inside Y_i 's window, E_2 : $Y_i = \max \mathcal{D}$, E_3 : the algorithm selects Y_i . Define Q_{ij} as the event of E_1 , E_2 and E_3 happening simultaneously. The following lemma will be of use to compute the probability of winning.

Lemma 6. For $1 \le i \le k - 1$,

$$\mathbb{P}(Q_{ij}) = \begin{cases} \frac{1}{2^j} \frac{1}{i} \left(1 - \frac{1}{j} \right) & i+1 \le j \le k-1, \\ \frac{1}{2^{k-1}} \frac{1}{i} & j=k, \\ 0 & otherwise. \end{cases}$$

Proof. If j < i and E_1 holds, Y_i will not be selected, therefore $\mathbb{P}(Q_{ij}|j < i) = 0$. Also, note that a window of length 1 will contain either Y_k or $Y_{k'}$, where k' is the couple of k, so if j > k, E_1 can not hold and $\mathbb{P}(Q_{ij}|j > k) = 0$. So assume that $i + 1 \le j \le k$.

Observe now that E_1 is equivalent to the event that $C(\theta_1), \ldots, C(\theta_{i-1}), C(\theta_{i+1}), \ldots, C(\theta_{j-1}), \theta_j$ are inside the interval $[\theta_i - 1, \theta_i)$. As this interval has length 1 and all variables are independent and uniformly chosen from [-1, 1), $\mathbb{P}(E_1 | \theta_i > 0, j < k) = 1/2^{j-1}$. The case j = k is slightly different, because $\theta_k = C(\theta_{k'})$. If i = k' then $\theta_k = \theta_i - 1 \in [\theta_i - 1, \theta_i)$, and if $i \neq k'$ then $\theta_k = C(\theta_l)$ for some $l \in \{1, \ldots, j-1\} \setminus \{i\}$. In both cases, we need to impose only j-2 variables in $[\theta_i - 1, \theta_i)$, resulting in $\mathbb{P}(E_1 | \theta_i > 0, j = k) = 1/2^{k-2}$.

Define the auxiliary events $F_1 : \max\{0, C(\theta_1), \ldots, C(\theta_{i-1})\} = 0$ and $F_2 : \max\{0, C(\theta_1), \ldots, C(\theta_{i-1}), C(\theta_{i+1}), \ldots, C(\theta_{j-1}), \theta_j)\} \neq \theta_j$. Note that given E_1 , event E_2 is equivalent to $\theta_1, \ldots, \theta_{i-1} \leq 0 < \theta_i$, which in turn is equivalent to $\theta_i > 0$ and F_1 . From here E_1 and E_2 happening simultaneously is equivalent to E_1, F_1 and $\theta_i > 0$. Also, by Lemma 4, E_3 is equivalent to either $\theta_j \leq 0$, or $\theta_j > 0$ and Y_j is smaller than some element arriving in $[\theta_j - 1, \theta_j)$.

We claim first that for j = k, Q_{ik} is equivalent to E_1 , F_1 and $\theta_i > 0$ (which is equivalent to E_1, E_2). Indeed, suppose that E_1, E_2 hold. If j = k we distinguish 2 cases. If $i \sim k$, then we have that $\theta_k \leq 0$. If $i \not\sim k$ and $\theta_k > 0$ then the couple k' of k, will satisfy that $Y_{k'} > Y_k$ and $\theta_{k'} \in [\theta_j - 1, \theta_j]$. In both cases, E_3 holds.

Now we claim that for $1 \leq j \leq k-1$, Q_{ij} is equivalent to all events E_1, F_1, F_2 and $\theta_i > 0$ happening simultaneously. For that, suppose that E_1 , and E_2 hold. Under that conditioning, E_3 holds in two cases, either $\theta_j \leq 0$, or $\theta_j > 0$ and for some $\ell \in \{1, \ldots, j-1\} \setminus \{i\}, \theta_j - 1 \leq \theta_\ell < \theta_j$, the latter being the same as $C(\theta_\ell) \in [-1, \theta_j - 1) \cup [\theta_j, 1]$. But recall that by E_1 , every $\ell \in \{1, \ldots, j-1\} \setminus \{i\}$ satisfies $C(\theta_\ell) \in [\theta_i - 1, \theta_i)$. Therefore, E_3 holds if and only if $\theta_j \leq 0$ or for some $\ell \in \{1, \ldots, j-1\} \setminus \{i\}, C(\ell) \in [\theta_j, \theta_i)$. This is equivalent to event F_2 , concluding the claim.

We are ready to compute the probability of Q_{ij} . Note that θ_j being or not the maximum of the set $\{0, C(\theta_1), \ldots, C(\theta_{i-1}), C(\theta_{i+1}), \ldots, C(\theta_{j-1}), \theta_j\}$ does not depend on the inner ordering of the set $\{0, C(\theta_1), \ldots, C(\theta_{i-1})\} = 0$. This implies that F_1 and F_2 are independent events. Since F_1 is equivalent to $\max\{1 - \theta_i, C(\theta_1) + 1 - \theta_i, \ldots, C(\theta_{i-1}) + 1 - \theta_i\} = 1 - \theta_i$, from Lemma 5, $\mathbb{P}(F_1|E_1, \theta_i > 0) = 1/i$, and since F_2 is equivalent to $\max\{1 - \theta_i, C(\theta_1) + 1 - \theta_i, \ldots, C(\theta_{i-1}) + 1 - \theta_i, \ldots, C(\theta_{i-1}) + 1 - \theta_i, C(\theta_{i+1}) + 1 - \theta_i, \ldots, C(\theta_{j-1}) + 1 - \theta_i\} \neq \theta_j + 1 - \theta_i$, $\mathbb{P}(F_2|E_1, \theta_i > 0) = 1 - 1/j$. Putting all together and using independence, $\mathbb{P}(Q_{ik}) = \mathbb{P}(E_1, \theta_i > 0, F_1) = \frac{1}{2} \frac{1}{2^{k-2}} \frac{1}{i}$, and for $1 \leq j \leq k - 1$ $\mathbb{P}(Q_{ij}) = \mathbb{P}(E_1, \theta_i > 0, F_1, F_2) = \frac{1}{2} \frac{1}{2^{j-1}} \frac{1}{i} \left(1 - \frac{1}{j}\right)$. This concludes the proof of the lemma.

With Lemma 6, we can compute the probability of winning as:

$$\mathbb{P}(Win) = \sum_{i=1}^{k-1} \sum_{j=i+1}^{k} \mathbb{P}(Q_{ij}) = \sum_{i=1}^{k-1} \left(\sum_{j=i+1}^{k-1} \frac{1}{2^j} \frac{1}{i} \left(1 - \frac{1}{j} \right) + \frac{1}{2^{k-1}} \frac{1}{i} \right) = \frac{H_{k-1}}{2^{k-1}} + \sum_{i=1}^{k-2} \sum_{j=i+1}^{k-1} \frac{1}{2^j} \frac{1}{i} \frac{j-1}{j},$$

where $H_s = \sum_{i=1}^{s} \frac{1}{i}$ is the s-th harmonic number. Note that the probability of winning only depends on k and denote it F(k). The following lemma helps to find the worst case scenario for our algorithm.

Lemma 7. F(k) is non-increasing in k.

Proof. To prove the lemma we note that for $k \geq 2$:

$$F(k+1) - F(k) = \frac{H_k}{2^k} - \frac{H_{k-1}}{2^{k-1}} + \sum_{i=1}^{k-1} \frac{1}{2^k} \frac{1}{i} \left(1 - \frac{1}{k}\right)$$
$$= \frac{H_k}{2^k} + \frac{H_{k-1}}{2^k} \left(1 - \frac{1}{k}\right) - \frac{H_{k-1}}{2^{k-1}} = \frac{1 - H_{k-1}}{k^{2k}} \le 0.$$

Proof of Theorem 1. To prove the main theorem of the section we use the previous results to give a lower bound for the probability of winning. The worst case happens $k \to \infty$:

$$\mathbb{P}(\text{Win}) \ge \min_{k\ge 2} F_k = \lim_{k\to\infty} F_k = \sum_{i=1}^{\infty} \frac{1}{i} \sum_{j=i+1}^{\infty} \frac{1}{2^j} \left(1 - \frac{1}{j}\right)$$
$$= \sum_{i=1}^{\infty} \frac{1}{i} \sum_{j=i+1}^{\infty} \left(\frac{1}{2^j} - \int_0^{1/2} x^{j-1} dx\right) = \sum_{i=1}^{\infty} \frac{1}{i} \left(\frac{1}{2^i} - \int_0^{1/2} \frac{x^i}{1-x} dx\right)$$
$$= \sum_{i=1}^{\infty} \int_0^{1/2} x^{i-1} dx - \int_0^{1/2} \frac{1}{1-x} \int_0^x \sum_{i=1}^{\infty} y^{i-1} dy dx$$
$$= \int_0^{1/2} \frac{1}{1-x} dx - \int_0^{1/2} \frac{1}{1-x} \int_0^x \frac{1}{1-y} dy dx$$
$$= \ln 2 \left(1 - \frac{\ln 2}{2}\right) \approx 0.45292.$$

4 Maximizing the expected value

In this section we present a lower bound for the prophet variant of the two-sided game of Googol, in which the objective is to maximize $\mathbb{E}(D_{\sigma(\tau)})/\mathbb{E}(\max \mathcal{D})$. We analyze a combination of the three basic algorithms presented in Section 2. More precisely, for a fixed $r \in [0, 1]$, we define ALG_r^* as follows. With probability (1 - r)/2 run ALG_o , with probability (1 - r)/2 run ALG_c , and with probability r run ALG_f . This combination allows us to obtain the main result of the section.

Theorem 2. There exists a value $r^* \in [0,1]$ such that for any instance of the two-sided game of Googol, $\mathbb{E}(ALG_{r^*}^*) \geq \alpha \mathbb{E}(\max \mathcal{D})$ with $\alpha = \frac{4-5\ln 2}{5-6\ln 2} \approx 0.635184$.

This ratio is better than the best known guarantee of $1 - 1/e \approx 0.632$ for the i.i.d. prophet secretary problem with samples [5], which is a particular case of our problem.

Recall that none of the 3 basic algorithms can guarantee a ratio better than $\frac{1}{2}$ by itself, even in very simple cases. Indeed, as shown in the introduction, for the instance $\{(1,0), (\varepsilon, \varepsilon^2)\}$ we have that $\mathbb{E}(\max \mathcal{D}) = 1/2 + O(\varepsilon)$ and $\mathbb{E}(ALG_o) = 1/4 + O(\varepsilon)$. Furthermore, if we consider the instance $\{(1, 1 - \varepsilon), (\varepsilon, 0)\}$ then $\mathbb{E}(\max \mathcal{D}) = 1 - O(\varepsilon)$ but $\mathbb{E}(ALG_c) = \mathbb{E}(ALG_f) = 1/2 + O(\varepsilon)$.

The intuition of our result is that the situations where each of the three algorithms perform poorly are very different. As we are able to compute exactly the distribution of the outcome of each algorithm, we can balance their distributions, so that in all cases ALG_r^* performs well. We can state this in terms of the ordered values $Y_1, ..., Y_k$. On the one hand, since ALG_f is very restrictive, it has good chances of stopping at Y_1 when it is in \mathcal{D} . On the other hand, ALG_c has higher probability of stopping at Y_i than ALG_f , for 1 < i < k. But none of the two algorithms can stop at Y_k , whereas ALG_o stops at Y_k with positive probability. This comes at the cost of sacrificing a bit of the better elements, but as showed in the examples, it is very important for the case when k is small.

Roughly speaking, the proof of Theorem 2 goes as follows. For each of the basic algorithms we compute in lemmas 9 to 11 the probability that they stop with the value Y_j for each j < k, and express it as a series truncated in k and an extra term that depends on k. Then in Lemma 12 we prove that in the combined formula, for certain values of r, the extra term can be replaced with the tail of the series, so the formula does not depend on k. Finally, for a specific value of r, we show in lemmas 13 and 14 an approximate stochastic dominance and conclude, i.e., we make use of the following general observation.

Observation 8. If $\min_{j \leq k} \frac{\mathbb{P}(ALG \geq Y_j)}{\mathbb{P}(\max \mathcal{D} \geq Y_j)} \geq \alpha$, then $\mathbb{E}(ALG) \geq \alpha \mathbb{E}(\max \mathcal{D}).^5$

4.1 Distribution of the Maximum

As noted before, $\mathbb{P}(\max \mathcal{D} = Y_i) = 0$ if i > k. For the remaining values, Wang [27] did a simple analysis for the distribution of $\max \mathcal{D}$.

For Y_i to be the maximum of \mathcal{D} we need: (1) $Y_i \in \mathcal{D}$ and (2) $Y_j \in \mathcal{U}$ for $1 \leq j < i$. If $1 \leq i < k$ all these events are independent with probability $\frac{1}{2}$, so we have $\mathbb{P}(\max \mathcal{D} = Y_i) = (\frac{1}{2})^i$. If i = k, the fact that Y_k is facing downwards implies that its couple is facing upwards, so we need one less coin toss for the events to happen simultaneously. Putting all together:

$$\mathbb{P}(\max \mathcal{D} = Y_i) = \begin{cases} \frac{1}{2^i} & i < k\\ \frac{1}{2^{k-1}} & i = k\\ 0 & i > k \end{cases}$$

Therefore we have that, $\mathbb{P}(\max \mathcal{D} \ge Y_i) = 1 - \frac{1}{2^i}$ if $1 \le i < k$ and $\mathbb{P}(\max \mathcal{D} \ge Y_k) = 1$.

4.2 Analysis of the basic algorithms

In this section we precisely derive the distribution of the obtained value of each of the basic algorithms. In the next section we combine these distributions to find an improved randomized algorithm. In what follows, we will denote by k' the couple of k. Note that the identities of k and k' depend only on the instance, and not on the realizations of the random coins or the random permutation.

⁵Recall that max \mathcal{D} can only take values in $\{Y_1, ..., Y_k\}$, so the implication follows immediately.

Lemma 9. For every $1 \le i \le k-1$,

$$\mathbb{P}(ALG_{c} = Y_{i}) = \frac{1}{2^{k-1}} + \sum_{j=i+1}^{k-1} \frac{1}{2^{j}} \left(1 - \frac{1}{j}\right)$$

and

$$\mathbb{P}(\mathrm{ALG}_{\mathsf{c}} \ge Y_k) = \sum_{j=1}^{k-1} \frac{1}{2^j j}.$$

Proof. Intuitively, for each j > i, we condition on that Y_j is the maximum value in the window of Y_i and show that ALG_c stops in Y_i with probability (1 - 1/j). More precisely, fix an index $i \le k - 1$. For $j \le i, j \ne i$, denote by F_j the event that $\theta_i \ge 0$ (or equivalently $Y_i \in \mathcal{D}$) and that j is the smallest index such that $\theta_j \in [\theta_i - 1, \theta_i)$. If j < k then $\mathbb{P}(F_j) = 1/2^j$ because the variables $\{\theta_\ell\}_{\ell \le j}$ are all independent and uniformly distributed in (-1, 1], and for any given $\theta_i \ge 0$, the interval $[\theta_i - 1, \theta_i)$ has length 1. For F_k we have that $\theta_k \in [\theta_i - 1, \theta_i)$ if and only if $\theta_{k'} = C(\theta_k) \notin [\theta_i - 1, \theta_i)$. Since the variables $\{\theta_\ell\}_{\ell \le k, \ell \ne k'}$ are all independent, $\mathbb{P}(F_k) = 1/2^{k-1}$.

Note that if for some $j < i, \theta_j \in [\theta_i - 1, \theta_i)$, then $ALG_c \neq Y_i$, because its threshold $T_c(\theta_i)$ will be at least $Y_j > Y_i$. Also note that either θ_k or $\theta_{k'}$ is in $[\theta_i - 1, \theta_i)$, so the events $(F_j)_{j=i+1}^k$ (that are pairwise disjoint) completely cover the event { $ALG_c = Y_i$ }. Thus we have the identity

$$\mathbb{P}(ALG_{c} = Y_{i}) = \frac{1}{2^{k-1}} \mathbb{P}(ALG_{c} = Y_{i}|F_{k}) + \sum_{j=i+1}^{k-1} \frac{1}{2^{j}} \cdot \mathbb{P}(ALG_{c} = Y_{i}|F_{j}).$$
(3)

Now, for j > i, conditional on F_j , ALG_c = Y_i if and only if ALG_c does not stop before observing Y_i . Using lemmas 4 and 5 we will show that $\mathbb{P}(ALG_c = Y_i|F_k) = 1$ and that $\mathbb{P}(ALG_c = Y_i|F_j) = (1-1/j)$ if i < j < k, and conclude the first formula of the lemma.

Conditional on F_k , the maximum value in the window of Y_i is Y_k , but Y_k is not larger than all elements in its window, because $Y_{k'}$ is in it (it is on the other side of its card). So Lemma 4 implies that $\mathbb{P}(ALG_c = Y_i|F_k) = 1$.

For the case i < j < k, we have that conditional on F_j , Y_j is the maximum value in the window of Y_i , so Lemma 4 implies that ALG_c stops in $(0, \theta_i)$ if and only if $\theta_j \ge 0$ and Y_j is larger than all elements in its window, i.e., $\theta_\ell \notin [\theta_j - 1, \theta_j)$, for all $\ell < j$. This is equivalent to $\max\{0, C(\theta_1), ..., C(\theta_{i-1}), C(\theta_{i+1}), ..., C(\theta_{j-1}), \theta_j\} = \theta_j$. Now, Lemma 5 implies that conditional on F_j , the random variables $\{0 - (\theta_i - 1), C(\theta_1) - (\theta_i - 1), ..., C(\theta_{i-1}) - (\theta_i - 1), C(\theta_{i+1}) - (\theta_i - 1), ..., C(\theta_{j-1}) - (\theta_i - 1), \theta_j - (\theta_i - 1)\}$ are independent and uniformly distributed in (0, 1). Hence, conditional on F_j , ALG_c stops in $(0, \theta_i)$ with probability 1/j, so $\mathbb{P}(ALG_c = Y_i|F_j) = (1 - 1/j)$. This proves the first formula.

For computing $\mathbb{P}(\operatorname{ALG}_c \geq Y_k)$, which is the probability that ALG_c stops, define as E_j the event that j is the smallest index such that $\theta_j > 0$ (or equivalently, that $Y_j = \max \mathcal{D}$). It is clear that for j < k, $\mathbb{P}(E_j) = 1/2^j$, that $\mathbb{P}(E_k) = 1/2^{k-1}$, and that $\mathbb{P}(E_j) = 0$ if j > k. For j < k, conditional on E_j , Lemma 4 implies that ALG_c stops if and only if $\max\{C(\theta_1), C(\theta_2), ..., C(\theta_{j-1}), \theta_j\} = \theta_j$. But this happens with probability 1/j. Conditional on E_k , ALG_c never stops. This concludes the proof of the lemma. Lemma 10. For every $1 \le i \le k-1$,

$$\mathbb{P}(ALG_{o} = Y_{i}) = \frac{1}{2^{k-1}} \left(1 - \frac{\mathbb{1}_{\{i \neq k'\}}}{k-1} \right) + \sum_{j=i+1}^{k-1} \frac{1}{2^{j}} \left(1 - \frac{1}{j} \right)$$

and

$$\mathbb{P}(ALG_{o} \ge Y_{k}) = \frac{1}{2^{k-1}(k-1)} + \sum_{j=1}^{k-1} \frac{1}{2^{j}j}.$$

Proof. Recall that ALG_o cannot select a value Y_j with j > k. Suppose now that we run ALG_o and ALG_c in the same instance and realization. Since ALG_c is more restrictive than ALG_o , if ALG_o stops, it stops earlier than ALG_c . Nevertheless, note that Y_k is the only value that can be accepted by ALG_o but not by ALG_c . Hence, for $1 \le i \le k - 1$, if $ALG_o = Y_i$ then $ALG_c = Y_i$, and if $ALG_c = Y_i$ then either $ALG_o = Y_i$ or $ALG_o = Y_k$. Thus, we can write the following identity for the case $1 \le i \le k - 1$.

$$\mathbb{P}(ALG_{o} = Y_{i}) = \mathbb{P}(ALG_{c} = Y_{i}) - \mathbb{P}(ALG_{c} = Y_{i} \text{ and } ALG_{o} = Y_{k}).$$
(4)

Thus, we just need to compute the negative term in Equation (4). In order to have $ALG_o = Y_k$ we need that $\theta_k > 0$ and that $\theta_j \notin (\theta_k - 1, \theta_k]$, for all $1 \leq j \leq k - 1$, with $j \neq k'$. Note that this is also a sufficient condition, because values that are smaller than Y_k cannot be accepted by ALG_o . Thus, $\mathbb{P}(ALG_o = Y_k) = 1/2^{k-1}$. If i = k', then $\theta_i = \theta_{k'} = \theta_k - 1$, which is negative if θ_k is positive, so $\mathbb{P}(ALG_c = Y_{k'}) = 0$.

Assume now that $i \neq k'$. Conditional on ALG_o = Y_k , we have that ALG_c = Y_i if Y_i arrives after Y_k and all other values in $\{Y_1, ..., Y_k\}$ arrive either before the window of Y_k or after Y_i , i.e., if $\theta_k < \theta_i$ and $\theta_j \in (-1, \theta_k - 1) \cup (\theta_i, 1]$, for all $j \leq k - 1$, with $j \notin \{i, k'\}$. This is equivalent to say that $C(\theta_i) = \min\{0, C(\theta_1), ..., C(\theta_{k'-1}), C(\theta_{k'+1}), ..., C(\theta_{k-1})\}$. Note that Lemma 5 implies that conditional on ALG_o = Y_k , the variables $\{0 - (\theta_k - 1), C(\theta_1) - (\theta_k - 1), ..., C(\theta_{k'-1}) - (\theta_k - 1), C(\theta_{k'-1}) - (\theta_k - 1)\}$ are independent and uniformly distributed in [0, 1]. Then, $\mathbb{P}(ALG_c = Y_i | ALG_o = Y_k) = 1/(k-1)$.

We conclude the first formula by replacing the just computed probability and the formula of Lemma 9 in Equation (4). For the second one, we have that $\mathbb{P}(ALG_o \ge Y_k) = \sum_{i=1}^k \mathbb{P}(ALG_o = Y_i) = \mathbb{P}(ALG_o = Y_k) - (k-2)\frac{1}{2^{k-1}(k-1)} + \sum_{i=1}^{k-1} \mathbb{P}(ALG_c = Y_i) = \frac{1}{2^k} - (k-2)\frac{1}{2^{k-1}(k-1)} + \mathbb{P}(ALG_c \ge Y_k) = \mathbb{P}(ALG_c \ge Y_k) + \frac{1}{2^{k-1}(k-1)}$.

Lemma 11. For every $1 \le i \le k-1$,

$$\mathbb{P}(\text{ALG}_{f} = Y_{i}) = \frac{1}{2^{k-1}} \frac{1}{k-1} + \sum_{j=i+1}^{k-1} \frac{1}{2^{j}(j-1)}$$

and

$$\mathbb{P}(\mathrm{ALG}_{\mathrm{f}} \ge Y_k) = \frac{1}{2}$$

Proof. Let F_j be the event that Y_j is the maximum value (*j* is the minimum index) such that $\theta_j < 0$. Roughly speaking, we condition on each F_j and show that $ALG_f = Y_i$ if Y_i is the value with earliest arrival time from $\{Y_1, ..., Y_{j-1}\}$. Note that the events F_j are pairwise disjoint, and since either θ_k or $\theta_{k'}$ is negative, $\mathbb{P}(F_j) = 0$ if j > k. Thus, the events F_j for $1 \le j \le k$ form a partition of the probability space. Note also that conditional on F_j , by definition ALG_f cannot accept any value smaller than Y_j . Therefore $\mathbb{P}(ALG_f = Y_i) = \sum_{j=i+1}^k \mathbb{P}(ALG_f = Y_i|F_j)\mathbb{P}(F_j).$

For $j \leq k-1$ the arrival times $\theta_1, ..., \theta_j$ are all independent and uniform in (-1, 1], so $\mathbb{P}(F_j) = 1/2^j$. Conditional on F_j we have that $\theta_1, ..., \theta_{j-1}$ are independent and uniform in [0, 1]. Moreover, conditional on F_j , ALG_f simply accepts the element in $\{Y_1, ..., Y_{j-1}\}$ with earliest arrival time, as they are exactly the ones larger than max $\mathcal{U} = Y_j$. Hence, $\mathbb{P}(ALG_f = Y_i|F_j) = 1/(j-1)$.

The case of F_k is slightly different. F_k is equivalent to the event that $\theta_1, ..., \theta_{k-1} > 0$, because $\theta_{k'} > 0$ implies that $\theta_k < 0$, so $\mathbb{P}(F_k) = 1/2^{k-1}$. Again ALG_f simply accepts the first element larger than Y_k , so it accepts Y_i with probability 1/(k-1). This concludes the proof of the first formula.

For computing $\mathbb{P}(ALG_f \geq Y_k)$, note that this is simply the probability that $\theta_1 > 0$, which is 1/2. This is because Y_1 is the overall largest value, so if it arrives before 0, nothing can be accepted, and if it arrives after 0, then Y_1 itself (and possibly other values with earlier arrival times) can be accepted.

4.3 The combined algorithm

We now use the distributions obtained in the last section in order to obtain an improved randomized algorithm. To this end we combine lemmas 9 to 11 to conclude that for $1 \le i \le k-1$,

$$\mathbb{P}(\text{ALG}_{r}^{*} = Y_{i}) = \frac{1-r}{2^{k-1}} + \frac{r-\mathbb{1}_{\{i \neq k'\}} \cdot (1-r)/2}{2^{k-1}(k-1)} + \sum_{j=i+1}^{k-1} \frac{1}{2^{j}} \left((1-r) - \frac{(1-r)}{j} + \frac{r}{j-1} \right) \\
\geq \frac{1-r}{2^{k-1}} + \frac{r-(1-r)/2}{2^{k-1}(k-1)} + \sum_{j=i+1}^{k-1} \frac{1}{2^{j}} \left((1-r) - \frac{(1-r)}{j} + \frac{r}{j-1} \right).$$
(5)

The inequality comes from the fact that $-\mathbb{1}_{\{i \neq k'\}} \geq -1$. Now we use the following lemma to complete the summation in Equation (5) using the extra term.

Lemma 12. For every $\frac{3-4\ln(2)}{5-6\ln(2)} \le r \le 2/3$, and any $k \ge 2$, $\sum_{i>k} \frac{1}{2^j} \left(1-r - \frac{1-r}{j} + \frac{r}{j-1}\right) \le \frac{1-r}{2^{k-1}} + \frac{r-(1-r)/2}{2^{k-1}(k-1)}.$

Proof. Rearranging terms, we obtain the following.

$$\sum_{j\geq k} \frac{1}{2^{j}} \left(1 - r - \frac{1 - r}{j} + \frac{r}{j - 1} \right) = \frac{1 - r}{2^{k - 1}} - \sum_{j\geq k} \frac{1 - r}{2^{j}j} + \sum_{j\geq k-1} \frac{r/2}{2^{j}j}$$
$$= \frac{1 - r}{2^{k - 1}} + \frac{r/2}{2^{k - 1}(k - 1)} - \sum_{j\geq k} \frac{1 - 3r/2}{2^{j}j}$$
$$= \frac{1 - r}{2^{k - 1}} + \frac{r/2}{2^{k - 1}(k - 1)} - \frac{(1 - 3r/2)}{2^{k - 1}(k - 1)} \sum_{j\geq 1} \frac{(k - 1)}{2^{j}(j + k - 1)}.$$
 (6)

Therefore, it is enough to prove that $-(1-3r/2)\sum_{j\geq 1}\frac{(k-1)}{2^{j}(j+k-1)} \leq r-1/2$. Since $r \leq 2/3$, the term (1-3r/2) is positive. Note that for $k\geq 2$, we have $\sum_{j\geq 1}\frac{(k-1)/(j+k-1)}{2^{j}} \geq \sum_{j\geq 1}\frac{1}{2^{j}(j+1)} = 2\ln(2)-1$.

Thus, it would be enough to prove that $-(1-3r/2)(2\ln 2-1) \le r-1/2$. Rearranging the terms in the last expression and multiplying by -1, we obtain the equivalent condition $3-4\ln 2 \le r(5-6\ln 2)$. \Box

In what follows, we apply Lemma 12 to derive a bound on the distribution of the accepted element that does not depend on k. Then, we select a specific value for r that balances these bounds, and allows us to approximate the distribution of max \mathcal{D} . We define now the function

$$a(r) := \sum_{j \ge 2} \frac{1}{2^j} \left((1-r) - \frac{(1-r)}{j} + \frac{r}{j-1} \right) = 1 - \ln 2 + \frac{r(3\ln 2 - 2)}{2}.$$
 (7)

This function appears in the next lemma we prove as the approximation factor of the distribution of max \mathcal{D} .

Lemma 13. For $r^* = \frac{3-4\ln 2}{5-6\ln 2}$ we have that for all $1 \le i \le k-1$,

$$\mathbb{P}(ALG_{r^*}^* = Y_i) \ge \frac{1}{2^i} 2a(r^*) .$$
(8)

Proof. Using Lemma 12 in Equation (5) we can write the following for $r^* \leq r \leq 2/3$.

$$\mathbb{P}(ALG_r^* = Y_i) \ge \sum_{j \ge i+1} \frac{1}{2^j} \left((1-r) - \frac{(1-r)}{j} + \frac{r}{j-1} \right) \,. \tag{9}$$

Note that this immediately gives the desired bound for i = 1 and any r in the interval $[r^*, 2/3]$. We will prove directly that it also holds for i = 2 and will proceed by induction for $i \ge 3$. For i = 2, Equation (9) combined with the definition of a(r) gives that

$$\mathbb{P}(ALG_r^* = Y_2) \ge a(r) - \frac{1}{4} \left((1-r) - \frac{(1-r)}{2} + r \right)$$
$$= a(r) - \frac{1+r}{8}.$$

Therefore, it is enough to prove that $a(r^*) - \frac{1+r^*}{8} \ge a(r^*)/2$, which is equivalent to $4a(r^*) \ge 1 + r^*$. If we replace $a(r^*)$ with the explicit formula in the right hand of Equation (7) and rearrange terms, we obtain the inequality $3 - 4\ln(2) \ge r^*(5 - 6\ln(2))$. Note that this is satisfied with equality by r^* .

For $i \geq 3$ we simply prove that if $r \leq 1/3$, then

$$\sum_{j\geq i+1} \frac{1}{2^j} \left((1-r) - \frac{(1-r)}{j} + \frac{r}{j-1} \right) \ge \frac{1}{2} \sum_{j\geq i} \frac{1}{2^j} \left((1-r) - \frac{(1-r)}{j} + \frac{r}{j-1} \right).$$
(10)

In fact, note that we can change the index in the right-hand side of Equation (10) to get the same range as in the sum of the left-hand side. So when we write the inequality for each term of the two summations, we obtain

$$\begin{aligned} \frac{1}{2^{j}} \left((1-r) - \frac{1-r}{j} + \frac{r}{j-1} \right) &\geq \frac{1}{2^{j}} \left((1-r) - \frac{1-r}{j-1} + \frac{r}{j-2} \right) &\Leftrightarrow \\ \frac{1}{j-1} - \frac{1}{j} &\geq r \left(\frac{1}{j-2} - \frac{1}{j} \right) &\Leftrightarrow \\ \frac{j-2}{2(j-1)} &\geq r \,. \end{aligned}$$

which holds whenever $j \ge 4$ and $r \le 1/3$. Since $r^* = \frac{3-4\ln 2}{5-6\ln 2} \approx 0.270$, we conclude that Equation (10) holds for r^* , and therefore, Equation (8) holds for all $1 \le i \le k-1$.

Lemma 14. For any $k \geq 2$, we have that $\mathbb{P}(ALG_{r^*}^* \geq Y_k) \geq 2a(r^*)$.

Proof. Using lemmas 9 to 11 we get the following inequality.

$$\mathbb{P}(ALG_{r^*}^* \ge Y_k) = (1 - r^*) \left(\frac{1}{2^k(k-1)} + \sum_{j=1}^{k-1} \frac{1}{2^j j} \right) + \frac{r^*}{2}.$$
(11)

Define now the function $G(k) = \frac{1}{2^k(k-1)} + \sum_{j=1}^{k-1} \frac{1}{2^j j}$. We minimize G to obtain a general lower bound for Equation (11). We have that

$$G(k+1) - G(k) = \frac{1}{2^k k} + \frac{1}{2^{k+1} k} - \frac{1}{2^k (k-1)}$$
$$= \frac{2(k-1) + (k-1) - 2k}{2^{k+1} k (k+1)}$$
$$= \frac{k-3}{2^{k+1} k (k+1)}.$$

Hence, $G(2) \ge G(3)$, and $G(k) \le G(k+1)$ for $k \ge 3$, so G is minimized when k = 3. Thus,

$$\mathbb{P}(ALG_{r^*}^* \ge Y_k) \ge (1 - r^*) \left(\frac{1}{16} + \frac{1}{2} + \frac{1}{8}\right) + \frac{r^*}{2}$$
$$= \frac{11 - 3r^*}{16}$$
$$= \frac{55 - 66 \ln 2 - 9 + 12 \ln 2}{16(5 - 6 \ln 2)}$$
$$= \frac{16(4 - 5 \ln 2)}{16(5 - 6 \ln 2)} + \frac{26 \ln 2 - 18}{16(5 - 6 \ln 2)}$$
$$\ge 2a(r^*).$$

The last inequality comes from the fact that $\frac{26\ln 2 - 18}{16(5-6\ln 2)} \approx 0.00162 \geq 0$ and that

$$\begin{aligned} a(r^*) &= 1 - \ln 2 + \frac{r(3\ln 2 - 2)}{2} \\ &= \frac{(2 - 2\ln 2)(5 - 6\ln 2) + (3 - 4\ln 2)(3\ln 2 - 2)}{2(5 - 6\ln 2)} \\ &= \frac{4 - 5\ln 2}{2(5 - 6\ln 2)}. \end{aligned}$$

With the last two lemmas we are ready to prove the main theorem of this section.

Proof of Theorem 2. Summing the lower bound in Lemma 13 it follows that for all $1 \le i \le k-1$,

$$\mathbb{P}(ALG_{r^*}^* \ge Y_i) \ge 2a(r^*) \cdot \left(1 - \frac{1}{2^i}\right)$$
$$= 2a(r^*)\mathbb{P}(\max \mathcal{D} \ge Y_i).$$

From Lemma 14 we get that also $\mathbb{P}(ALG_{r^*}^* \geq Y_k) \geq 2a(r^*) \cdot \mathbb{P}(\max \mathcal{D} \geq Y_k)$. Therefore, as $\max \mathcal{D} \in \{Y_1, ..., Y_k\}$ with probability 1, we conclude that $\mathbb{E}(ALG_{r^*}^*) \geq 2a(r^*) \cdot \mathbb{E}(\max \mathcal{D})$.

5 Large Data Sets (large k)

In this section we consider the case in which k is large and show that for that case one can obtain better guarantees for the prophet two-sided game of Googol. This case appears very often, for instance, in the i.i.d. prophet secretary problem with unknown (continuous) distributions, all permutations of [2n] are equally likely to be the ordering of the 2n values in the cards. The probability that k is at least k_0 equals the probability that the top t elements of the list are written in different cards. Note that for any $\ell = o(\sqrt{n})$,

$$\mathbb{P}(k \ge \ell) = \left(1 - \frac{1}{2n - 1}\right) \cdot \left(1 - \frac{2}{2n - 2}\right) \cdots \left(1 - \frac{\ell - 1}{2n - (\ell - 1)}\right) \ge \left(1 - \frac{\ell}{2n - \ell}\right)^{\ell} \\ \ge 1 - \frac{\ell^2}{2n - \ell} \ge 1 - o(1).$$

The algorithm that we use uses a moving window of length strictly larger than 1, as outlined in Section 2.

Moving window algorithm of length 1 + t (ALG(t)). We first draw *n* uniform variables in [0, 1] and we sort them from smallest to largest⁶ as $0 < \tau_1 < \tau_2 < \cdots < \tau_n < 1$. We interpret τ_j as the arriving time of the *s*-th hidden value that we reveal, and therefore, $\tau_s - 1 = C(\tau_j)$ is the arriving time of the corresponding *s*-th face up value. The algorithm will accept the *s*-th face up value $x = D_{\sigma(s)}$ if and only if *x* is larger than any value arrived in the previous 1 + t time units, i.e. if and only if *x* is larger than all elements arriving in $[\max(\tau_s - 1 - t, -1), \tau_s]$

Observe that ALG(0) and ALG(1) are exactly the algorithms ALG_c and ALG_f defined in previous sections. Below we analyze the performance of ALG(t).

Lemma 15. For every $1 \le i \le k-1$,

$$\mathbb{P}(\text{ALG}(t) = Y_i) = b(k, t) + \sum_{j=i+1}^{k-1} \frac{a(j, t)}{2^j},$$
(12)
where $a(j, t) = \frac{1 - (1 - t)^{j-1}(1 + t)}{j - 1} + (1 - t)^{j-1}(1 + t) - \frac{(1 - t)^j}{j},$
 $b(k, t) = \frac{1}{2^{k-1}} \left(\frac{1 - (1 - t)^{k-1}}{k - 1} + (1 - t)^{k-1} \right),$

and

$$\mathbb{P}(\text{ALG}(t) \ge Y_k) = \frac{1}{2} + \sum_{i=2}^{k-1} \frac{(1-t)^j}{2^j j}.$$

Proof. Fix an index $i \leq k - 1$. Recall that $\theta_i \in [-1, 1)$ denotes the arriving time of Y_i . Let J be the random variable denoting the index of the largest element arriving in the window of Y_i , or equivalently, the smallest index such that $\theta_j \in [\theta_i - 1 - t, \theta_i)$. Since $[\max(\theta_i - 1 - t, -1), \theta_i)$ has length at least 1, it always contains θ_k or $C(\theta_k) = \theta_{k'}$ (or both), we conclude that $J \leq k$.

Note that the algorithm selects Y_i if and only if the next three events occur: (I) $\theta_i > 0$, (II) J > i, as otherwise, $Y_J > Y_i$ would be inside the window of Y_i , and (III) No element is selected

 $^{^{6}}$ We assume that the drawn values are all distinct as this happens with probability 1

before Y_i 's arrival. By Lemma 4, the third event occurs if and only if the maximum element⁷ X arriving in $[0, \theta_i)$ is smaller than some element in its own window. Note that this happens in two cases: either $\theta_J < 0$ (and therefore Y_J is in the window of X), or $\theta_J \ge 0$ and Y_J is smaller than some element arriving in its own window.

In what follows fix some j with $i + 1 \le j \le k - 1$. We will compute the probability of both selecting Y_i and J = j.

Case 1: $0 \le \theta_i \le t$. In order to impose that J = j we need that all j - 2 arrival times ($\theta_l : l \in [j-1] \setminus \{i\}$) are outside the interval $[-1, \theta_i]$ and θ_j is inside. Furthermore, in this case every element arriving in $[0, \theta_j)$ is strictly smaller than Y_j . Therefore in order to impose that no element is selected before Y_i 's arrival we need that $\theta_j < 0$. It follows that

$$\begin{split} \mathbb{P}(\text{ALG}(t) = Y_i, J = j, \theta_i \in [0, t]) &= \frac{1}{2} \int_0^t \mathbb{P}(\theta_j < 0) \prod_{l \in [j-1] \setminus \{i\}} \mathbb{P}(\theta_l > 1 - x) dx \\ &= \frac{1}{2} \int_0^t \frac{1}{2} \left(\frac{1 - x}{2}\right)^{j-2} dx = \frac{1}{2^j} \cdot \frac{1 - (1 - t)^{j-1}}{j - 1}. \end{split}$$

Case 2: $t < \theta_i \leq 1$ and $\theta_j < 0$. We now need that all j - 2 arrival times $\theta_l, l \in [j - 1] \setminus \{i\}$ fall outside $[\theta_i - t - 1, \theta_i]$ and θ_j is inside. Since we are imposing $\theta_j < 0$, we actually require that $\theta_j \in [\theta_i - t - 1, 0]$. Since $\theta_J < 0$ this is enough to guarantee that Y_i will be selected. Therefore,

$$\begin{split} & \mathbb{P}(\mathrm{ALG}(t) = Y_i, J = j, \theta_i \in (t, 1] \land \theta_j < 0) \\ & = \frac{1}{2} \int_t^1 \mathbb{P}(\theta_j \in [x - t - 1, 0]) \prod_{l \in [j - 1] \backslash \{l\}} \mathbb{P}(\theta_l \not\in [x - t - 1, x]) dx \\ & = \frac{1}{2} \int_t^1 \frac{1 + t - x}{2} \left(\frac{1 - t}{2}\right)^{j - 2} dx = \frac{(1 - t)^{j - 1}(1 + t)}{2^{j + 1}}. \end{split}$$

Case 3: $t < \theta_i \le 1$ and $0 \le \theta_j < t$. As before we need that all j-2 arrival times $\theta_l, l \in [j-1] \setminus \{i\}$ fall outside $[\theta_i - t - 1, \theta_i]$ and $\theta_j \in [0, t)$. Since $\theta_j \ge 0$ we also need that at least one of $Y_l, l \in [j-1] \setminus \{j\}$ arrives in Y_j 's window $[-1, \theta_j)$ and since they must be outside Y_i 's window this reduces to the event that not all j-2 arrival times fall in the interval $(\theta_i, 1]$. Therefore,

$$\begin{split} \mathbb{P}(\text{ALG}(t) &= Y_i, J = j, \theta_i \in (t, 1] \land 0 \le \theta_j < t) \\ &= \frac{1}{2} \int_t^1 \mathbb{P}(\theta_j \in [0, t)) \mathbb{P}(\forall l \in [j - 1] \setminus \{i\}, \theta_l \not\in [x - t - 1, x] \text{ and not all in } (x, 1]) dx \\ &= \frac{1}{2} \int_t^1 \frac{t}{2} \left(\left(\frac{1 - t}{2}\right)^{j - 2} - \left(\frac{1 - x}{2}\right)^{j - 2} \right) dx = \frac{t(1 - t)^{j - 1}}{2^j} - \frac{t(1 - t)^{j - 1}}{2^j(j - 1)} \end{split}$$

Case 4: $t < \theta_i \leq 1$ and $t \leq \theta_j < \theta_i$. Once again we need that all j - 2 arrival times $\theta_l, l \in [j-1] \setminus \{i\}$ fall outside $[\theta_i - t - 1, \theta_i]$ and $\theta_j \in [t, \theta_i)$. Since $\theta_j \geq 0$ we also need that at least one of $Y_l, l \in [j-1] \setminus \{j\}$ arrives in Y_j 's window $[\theta_j - 1 - t, \theta_j)$ and since they must be outside Y_i 's window

⁷If no element arrives in $[0, \theta_i)$ then event (III) also occurs

this reduces to the event that not all j-2 arrival times fall in the set $[-1, \theta_j - t - 1) \cup (\theta_i, 1]$. Therefore,

$$\begin{split} \mathbb{P}(\mathrm{ALG}(t) &= Y_i \wedge J = j \wedge \theta_i \in (t, 1] \wedge t \le \theta_j < x) \\ &= \frac{1}{2} \int_t^1 \mathbb{P}(\theta_j \in [t, x)) \mathbb{P}(\forall l \in [j-1] \setminus \{i\}, \theta_l \not\in [x-t-1, x] \text{ and not all in } [-1, \theta_j - t - 1) \cup (x, 1]) dx \\ &= \frac{1}{2} \int_t^1 \int_t^x \left(\left(\frac{1-t}{2}\right)^{j-2} - \left(\frac{1-x+y-t}{2}\right)^{j-2} \right) dy dx \\ &= \frac{1}{2^j} \int_t^1 (1-t)^{j-2} (x-t) - \frac{(1-t)^{j-1} - (1-x)^{j-1}}{j-1} dx = \frac{(1-t)^j}{2^{j+1}} - \frac{(1-t)^j}{2^j j}. \end{split}$$

Putting all cases together we get that for all $1 \le i \le k-1$ and $i+1 \le j \le k-1$,

$$\mathbb{P}(\text{ALG}(t) = Y_i \land J = j) = \frac{1}{2^j} \cdot \left(\frac{1 - (1 - t)^{j-1}(1 + t)}{j - 1} + (1 - t)^{j-1}(1 + t) - \frac{(1 - t)^j}{j}\right).$$
(13)

Now let us consider the case J = k. In this case we only need to impose that all k - 2 arrival times $\theta_l, l \in [k-1] \setminus \{i\}$ fall outside the window of Y_i . No matter who the couple of k is, the previous condition implies that k is inside the window of Y_i . Therefore,

$$\mathbb{P}(ALG(t) = Y_i \wedge J = k) = \frac{1}{2} \int_0^t \left(\frac{1-x}{2}\right)^{k-2} dx + \left(\frac{1-t}{2}\right)^{k-1} \\ = \frac{1}{2^{k-1}} \left(\frac{1-(1-t)^{k-1}}{k-1} + (1-t)^{k-1}\right).$$
(14)

By combining (13) and (14) together we finish the proof of the first equality of this lemma, (12).

We now compute the probability that ALG(t) selects one of the top k values Y_1, \ldots, Y_k . Observe first that the algorithm never selects a value Y_j with j > k. This is because the window of Y_j always contains θ_k or $C(\theta_k) = \theta_{k'}$, and both Y_k and $Y_{k'}$ are larger than Y_j . By this observation, $\mathbb{P}(ALG(t) \ge Y_k)$ equals the probability that the algorithm stops by selecting something.

Let M be the random variable denoting the index of the largest value arriving after 0, i.e. $\theta_M \ge 0$ and $\theta_i < 0$, for all i < M. By Lemma 4, ALG(t) stops by selecting a value if and only if Y_M is larger than every value arriving in its window $[\max(\theta_M - 1 - t, 1), \theta_M))$. Observe that if M = k, this is impossible because the couple of k is always in that interval. Therefore,

$$\mathbb{P}(\text{ALG}(t) \ge Y_k) = \sum_{j=1}^k \mathbb{P}(\text{ALG}(t) \ge Y_k, M = j)$$

= $\sum_{j=1}^k \frac{1}{2} \int_0^1 \mathbb{P}(\forall i \in [j-1], \theta_i \in [-1, x - 1 - t) dx.$
= $\frac{1}{2} + \sum_{j=2}^{k-1} \frac{1}{2} \int_t^1 \left(\frac{x-t}{2}\right)^{j-1} dx = \frac{1}{2} + \sum_{j=2}^{k-1} \frac{(1-t)^j}{2^j j}.$

Algorithm ALG(t) has a very poor performance for k = 2. For example, in the instance $\{(1, 1 - \varepsilon), (\varepsilon, 0)\}$, we have $Y_1 = 1$, $Y_2 = 1 - \varepsilon$ and k = 2. By Lemma 15, $\mathbb{P}(ALG(t) \ge Y_2) = 1/2$. Therefore $\mathbb{E}(ALG(t)) \le 1/2$, while $\mathbb{E}(\max \mathcal{D}) \ge 1 - \varepsilon$.



Figure 1: Auxiliary functions to analyze ALG(t), as k tends to ∞ .

We will now study the behaviour of ALG(t) when k is large. For that, define

$$R_{k}(i,t) = \frac{\mathbb{P}(\text{ALG}(t) \ge Y_{i})}{\mathbb{P}(\max \mathcal{D} \ge Y_{i})} = \frac{1}{1 - (1/2)^{i}} \sum_{l=1}^{i} \sum_{j=l+1}^{k-1} b(k,t) + \frac{a(j,t)}{2^{j}}$$
$$R_{k}(t) = \mathbb{P}(\text{ALG}(t) \text{ stops}) = \frac{1}{2} + \sum_{j=2}^{k-1} \frac{(1-t)^{j}}{2^{j}j}.$$

Consider also their limits as $k \to \infty$, which since $kb(k,t) \to 0$ can be computed as

$$R_{\infty}(i,t) = \frac{1}{1 - (1/2)^{i}} \sum_{l=1}^{i} \sum_{j=l+1}^{\infty} \frac{a(j,t)}{2^{j}}.$$
$$R_{\infty}(t) = \frac{1}{2} + \sum_{j=2}^{\infty} \frac{(1-t)^{j}}{2^{j}j} = \frac{t}{2} + \ln\left(\frac{2}{1+t}\right).$$

Recall that the competitiveness of ALG(t) is at least $\min_{i\geq 1} R_{\infty}(i,t)$, and we are looking for the value t that maximizes this minimum. In Figure 1 we plot a few of these $R_{\infty}(i,t)$, for small values of i, together with function $R_{\infty}(t)$. From the plot we observe that the t^* maximizing the minima of all the curves satisfies $R_{\infty}(1,t^*) = R_{\infty}(t^*)$. By simplifying the sums, we obtain that $t^* \approx 0.1128904$ is the only root of:

$$R_{\infty}(1,t) - R_{\infty}(t) = 2 - (5t)/2 - (3+t)\ln 2 + (4+t)\ln(1+t).$$

To formally prove that $R_{\infty}(t^*) \approx 0.6426317$ is the sought guarantee we need the following lemma.

Lemma 16. Let a(j,t) be defined as in Lemma 15. For all $j \ge 3$, we have $a(j,t^*) \ge a(j+1,t^*)$

Proof. Below, we tabulate a few values of $a(j, t^*)$ with $t^* \approx 0.1128904$, and we observe that $a(j, t^*) \ge a(j+1, t^*)$ for all $j \in \{3, \ldots, 9\}$:

j	3	4	5	6	7	8	9	10
$a(j,t^*)$	0.705194	0.696462	0.65704	0.607906	0.556898	0.50734	0.460684	0.417513

Let us prove that the inequality also holds for $j \ge 10$. By rearranging terms, $a(j,t) \ge a(j+1,t)$ is equivalent to

$$\frac{1}{(1-t)^{j-1}} \left(\frac{1}{j-1} - \frac{1}{j} \right) \ge -t(1+t) - \frac{t(1-t)}{j} - \frac{(1-t)^2}{j+1} + \frac{1+t}{j-1}.$$

But note that the right hand side, evaluated at $t^* \ge 1/9$ is:

$$-t^*(1+t^*) - \frac{t^*(1-t^*)}{j} - \frac{(1-t^*)^2}{j+1} + \frac{1+t^*}{j-1} \le (1+t^*)\left(\frac{1}{j-1} - t^*\right) \le (1+t^*)\left(\frac{1}{j-1} - \frac{1}{9}\right),$$

which is nonpositive for all $j \ge 10$. Then, it is smaller than the left hand side which is positive. \Box

Theorem 3. For $t^* \approx 0.1128904$, and any instance of the two-sided game of Googol, as k goes to infinity, $\mathbb{E}[ALG(t^*)]/\mathbb{E}[\max \mathcal{D}]$ is at least $R_{\infty}(t^*) \approx 0.6426317$.

Proof. We will prove that for all $i \geq 1$, $R_{\infty}(i,t^*) \geq R_{\infty}(t^*)$. Denote for all $i \geq 1$, $P_i = \lim_{k\to\infty} \mathbb{P}(\operatorname{ALG}(t^*) = Y_i) = \sum_{j=i+1}^{\infty} a(j,t^*)/2^j$. By choice of t^* , $R_{\infty}(1,t^*) = R_{\infty}(t^*) \approx 0.6426317$. Furthermore we we can numerically evaluate $R_{\infty}(2,t^*) = \frac{P_1+P_2}{1-1/4} \approx 0.6547 \geq R_{\infty}(t^*)$ and $R_{\infty}(3,t^*) = \frac{P_1+P_2+P_3}{1-1/8} \approx 0.654331 \geq R_{\infty}(t^*)$.

To finish the proof, we will show that for all $i \ge 4$, $R_{\infty}(i, t^*) \ge R_{\infty}(i+1, t^*)$, and therefore, for all $i \ge 4$, $R_{\infty}(i, t^*) \ge \lim_{j\to\infty} R_{\infty}(j, t^*) = R_{\infty}(t^*)$. For this, we will also need the inequality $P_1 \approx 0.3213158 \ge 0.304065 \approx 8P_4$. By Lemma 16, for all $i \ge 2$,

$$P_i = \sum_{j=i+1}^{\infty} \frac{a(j,t^*)}{2^j} \ge \sum_{j=i+1}^{\infty} \frac{a(j+1,t^*)}{2^j} = 2P_{i+1}.$$

Therefore,

$$\sum_{\ell=1}^{i} P_{\ell} \ge 8P_4 + \sum_{\ell=2}^{i} P_{\ell} \ge 2^{i} P_{i+1} + P_{i+1} \sum_{\ell=2}^{i} 2^{i+1-\ell} = P_{i+1}(2^{i+1}-2).$$

We conclude that,

$$R(i+1,t^*) = \frac{P_{i+1} + \sum_{\ell=1}^{i} P_{\ell}}{1 - 1/2^{i+1}} \le \frac{\left(1 + \frac{1}{2^{i+1} - 2}\right)\sum_{\ell=2}^{i} P_{\ell}}{\frac{2^{i+1} - 1}{2^{i+1}}} = \frac{\sum_{\ell=2}^{i} P_{\ell}}{\frac{2^{i+1} - 2}{2^{i+1}}} = R_{\infty}(i,t^*).$$

To conclude this section we observe that the guarantee obtained in the $k = \infty$ case is still useful for moderately high values of k. Indeed, note that for all $j \ge 1$, and all $t \in (0, 1)$,

$$|a(j,t)| \le 2, |b(k,t)| \le \frac{2}{2^{k-1}}.$$

Therefore, for all $1 \le i \le k-1$ and all $k \ge 3$.

$$\left| \mathbb{P}(\text{ALG}(t) = Y_i) - \sum_{j=i+1}^{\infty} \left| \frac{a(j,t)}{2^j} \right| = \left| b(k,t) - \sum_{j=k}^{\infty} \frac{a(j,t)}{2^j} \right| \le \frac{2}{2^{k-1}} + 2\sum_{j=k}^{\infty} \frac{1}{2^j} = \frac{4}{2^{k-1}} = \frac{1}{2^{k-3}}$$

Therefore,

$$|R_k(i,t) - R_{\infty}(i,t)| = \frac{1}{1 - 1/2^k} \left| \sum_{\ell=1}^i \left(\mathbb{P}(\text{ALG}(t) = Y_\ell) - \sum_{j=\ell+1}^\infty \frac{a(j,t)}{2^j} \right) \right| \le \frac{2i}{2^{k-3}} \le \frac{16k}{2^k}$$

We also have that

$$|R_k(t) - R_{\infty}(t)| = \left|\sum_{j=k}^{\infty} \frac{(1-t)^j}{2^j j}\right| \le \frac{1}{2^{k-1}} \le \frac{16k}{2^k}$$

From here,

$$\min(R_k(t^*), \min_{1 \le i \le k-1} R_k(i, t^*)) \ge \min(R_\infty(t^*), \min_{1 \le i \le k-1} R_k(i, t^*)) - \frac{16k}{2^k} = R_\infty(t^*) - \frac{16k}{2^k}.$$

Thus the guarantee of ALG(t) for not so large values of k is already very close to $R_{\infty}(t^*) \approx 0.6426317$. For example, the guarantee of ALG(t) for $k \geq 20$ is at least $R_{\infty}(t^*) - 0.00031$.

Acknowledgments

This work was partially supported by CONICYT under grants CONICYT-PFCHA/Doctorado Nacional/2018-21180347, CONICYT-PFCHA/Magister Nacional/2018-22181138, Conicyt-Fondecyt 1181180, conicyt-fondecyt 1190043 and PIA AFB-170001, and by an Amazon Research Award.

References

- Abolhassani, M., Ehsani, S., Esfandiari, H., Hajiaghayi, M., Kleinberg, R., Lucier, B. Beating 1-1/e for ordered prophets. STOC 2017.
- [2] Azar, P., Kleinberg, R., Weinberg., S.M. Prophet Inequalities With Limited Information. SODA 2014.
- [3] Azar, Y., Chiplunkar, A., Kaplan, H. Prophet secretary: Surpassing the 1-1/e barrier. EC 2018.
- [4] Chawla, S., Hartline, J., Malec, D., Sivan, B. Multi-parameter mechanism design and sequential posted pricing. STOC 2010.
- [5] Correa, J., Duetting, P., Fischer, F., Schewior, K., Prophet inequalities for i.i.d. random variables from an unknown distribution. EC 2019.
- [6] Correa, J., Foncea, P., Hoeksma, R., Oosterwijk, T., Vredeveld, T. Posted price mechanisms for a random stream of customers. EC 2017.
- [7] Correa, J., Foncea, P., Hoeksma, R., Oosterwijk, T., Vredeveld, T. Recent Developments in Prophet Inequalities. ACM SIGecom Exchanges 17(1), 61–70, 2018.

- [8] Correa, J., Foncea, P., Pizarro, D., Verdugo, V. From pricing to prophets, and back!. Operations Research Letters, 47(1), 25–29, 2019.
- [9] Correa, J., Saona, R., Ziliotto, B. Prophet secretary through blind strategies. SODA 2019.
- [10] Dynkin, E.B. The optimum choice of the instant for stopping a Markov process. Soviet Math. Dokl. 4, 627–629, 1963.
- [11] Düetting, P., Feldman, M., Kesselheim, T., Lucier, B. Prophet inequalities made easy: Stochastic optimization by pricing non-stochastic inputs. FOCS 2017.
- [12] Düetting, P., Kesselheim, T. Posted Pricing and Prophet Inequalities with Inaccurate Priors. EC 2019.
- [13] Ehsani, S., Hajiaghayi, M., Kesselheim, T., Singla, S. Prophet secretary for combinatorial auctions and matroids. SODA 2018.
- [14] Esfandiari, H., Hajiaghayi, M., Liaghat, V., Monemizadeh, M. Prophet secretary. ESA 2015.
- [15] Ferguson, T.S. Who solved the secretary problem? Statistical Science 4(3), 282–296, 1989
- [16] Gilbert, J., Mosteller, F. Recognizing the maximum of a sequence. J. Am. Statist. Assoc. 61, 35–73, 1966
- [17] Hajiaghayi, M., Kleinberg, R., Sandholm, T. Automated online mechanism design and prophet inequalities. AAAI 2007.
- [18] Hill, T., Kertz, R. Comparisons of stop rule and supremum expectations of i.i.d. random variables. *The Annals of Probability* 10(2), 336–345, 1982.
- [19] Hill, T., Kertz, R. A survey of prophet inequalities in optimal stopping theory. Contemporary Mathematics 125, 191–207, 1992.
- [20] Kertz, R. Stop rule and supremum expectations of i.i.d. random variables: A complete comparison by conjugate duality. *Journal of Multivariate Analysis* 19, 88–112, 1986.
- [21] Kleinberg, R., Weinberg, S. M. Matroid prophet inequalities. STOC 2012.
- [22] Krengel, U., Sucheston, L. Semiamarts and finite values. Bull. Amer. Math. Soc. 83, 745–747, 1977.
- [23] Krengel, U., Sucheston, L. On semiamarts, amarts, and processes with finite value. Adv. in Probability 4, 197–266, 1978.
- [24] Lindley D.V. Dynamic programming and decision theory. Appl. Statist. 10, 39–51, 1961.
 The optimum choice of the instant for stopping a Markov process. Soviet Math. Dokl. 4 627-629.
- [25] Lucier, B. An economic view of prophet inequalities. ACM SIGecom Exchanges 16(1), 24–47, 2017.
- [26] Samuel-Cahn, E. Comparisons of threshold stop rule and maximum for independent nonnegative random variables. The Annals of Probability 12(4), 1213–1216, 1984.

[27] Wang, J. The prophet inequality can be solved optimally with a single set of samples. ArXiv preprint, 2018.