Embedding spanning trees in random graphs

Michael Krivelevich *

October 31, 2018

Abstract

We prove that if T is a tree on n vertices with maximum degree Δ and the edge probability p(n) satisfies: $np \geq C \max\{\Delta \log n, n^{\epsilon}\}$ for some positive $\epsilon > 0$, then with high probability the random graph G(n,p) contains a copy of T. The obtained bound on the edge probability is shown to be essentially tight for $\Delta = n^{\Theta(1)}$.

1 Introduction

In this paper we consider the problem of embedding a copy of a given spanning tree T on n vertices into the binomial random graph G(n, p).

Embedding problems are one of the most classical subjects in Extremal and Probabilistic Combinatorics. There is a large variety of results about finding given subgraphs, or graphs belonging to a given family, in random graphs. Here we concentrate on embedding large trees in binomial random graphs.

The problem of embedding large or nearly spanning trees in random graphs on n vertices (where by a nearly spanning tree we mean a tree T whose number of vertices is at most (1-c)n for some constant c>0) is a rather well researched subject, especially in the case of trees with bounded maximum degree, see, e.g., [7], [1], [9], [8], [10]. In particular, Alon, Sudakov and the author proved in [2] that for given $\epsilon>0$ and integer d there exists $C=C(d,\epsilon)>0$ such that \mathbf{whp}^1 the random graph G(n,p) with p=C/n has a copy of a tree T on $(1-\epsilon)n$ vertices of maximum degree at most d (in fact [2] proved that such a random graph contains \mathbf{whp} a copy of every such tree); better constant dependence and the resilience version of this result have recently been obtained in [3] and [4], respectively.

In contrast, nearly nothing has been known for the case of embedding *spanning* trees. Even the case of embedding spanning trees of bounded maximum degree appears to be unaddressed, apart from

^{*}School of Mathematical Sciences, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv, 69978, Israel. Email: krivelev@post.tau.ac.il. Research supported in part by a USA-Israel BSF grant and by a grant from the Israel Science Foundation.

¹An event \mathcal{E}_n occurs with high probability, or **whp** for brevity, in the probability space G(n,p) if $\lim_{n\to\infty} Pr[G \sim G(n,p) \in \mathcal{E}_n] = 1$.

some sporadic cases. Of course, the most classical result is about embedding a Hamilton path, or even a Hamilton cycle, in G(n,p); Komlós and Szemerédi [14] and independently Bollobás [5] proved that if $p(n) \geq \frac{\ln n + \ln \ln n + \omega(1)}{n}$, where $\omega(1)$ is any function tending to infinity arbitrarily slowly with n, then **whp** G(n,p) contains a Hamilton cycle. Alon et al. [2] observed that if a tree T has a linear in n number of leaves, then **whp** $G(n,C \ln n/n)$ contains a copy of T for some large enough C > 0; the proof is not that hard and utilizes the embedding result for nearly spanning trees from the same paper. However, no general result for this problem has been obtained, and even the case of the comb (which is the path P_0 of length $\sqrt{n} - 1$ with disjoint paths of length $\sqrt{n} - 1$ attached to each vertex of P_0), interpolating in some sense between the above mentioned solved cases, is open; this natural question has been communicated to us by Jeff Kahn [12].

Here we make a substantial step forward in solving this class of problems. Our main result if the following embedding theorem.

Theorem 1 Let T be a tree on n vertices of maximum degree Δ . Let $0 < \epsilon < 1$ be a constant. If

$$np \ge \frac{40}{\epsilon} \Delta \ln n + n^{\epsilon},$$

then whp a random graph G(n,p) contains a copy of T.

In other words, starting from $\Delta(T) = n^{\epsilon}$, edge probability $p = \frac{C\Delta \ln n}{n}$ is enough to get **whp** a copy of T in G(n, p).

It is not hard to see that the dependence of p on Δ , posted in Theorem 1, is optimal up to a constant factor in the range $np = n^{\Theta(1)}$. In order to state this result formally, for integers $n \geq \Delta \geq 3$ define the tree $T(n, \Delta)$ as follows. Write $n = (\Delta - 1)k - r$, where $0 \leq r \leq \Delta - 2$. Take a path $P = (v_1, \ldots, v_k)$ with k vertices, attach to v_1, \ldots, v_{k-1} vertex disjoint stars with $\Delta - 2$ leaves each, and finally attach to v_k a star with $\Delta - 2 - r$ leaves. The tree $T(n, \Delta)$ has n vertices and is of maximum degree at most Δ . For future reference observe that the k vertices of P dominate the remaining n - k vertices of $T(n, \Delta)$.

Theorem 2 For every $\epsilon > 0$ there exists $\delta > 0$ such that if $n^{\epsilon} \leq \Delta \leq \frac{n}{\ln n}$, then a random graph G(n,p) with $p = \frac{\delta \Delta \ln n}{n}$ whp does not contain a copy of $T(n,\Delta)$.

There is a certain similarity in appearances between the above results and the theorem of Komlós, Sárközy and Szemerédi [13], who proved that for $\delta > 0$ and all large enough n, any graph G on n vertices of minimum degree at least $(1/2+\delta)n$ contains a copy of every tree T on n vertices of maximum degree $\Delta(T) \leq cn/\ln n$, where $c = c(\delta)$ is a small enough constant; they noticed that their condition on $\Delta(T)$ is essentially tight too (actually because of the random graph G(n,p) with $1/2 and the above described tree <math>T(n,\Delta)$, just like in our Theorem 2). The arguments of [13] are naturally very different and do not seem to have much bearing on the situation in (sparse) random graphs.

In order to ease the reader's task we now give a brief description of the proof of Theorem 1. The key definition used is that of a bare path:

Definition 1 A path P in a tree T is called bare if all vertices of P have degree exactly two in T.

In the proof of Theorem 1, we first argue that every tree T on n vertices has a linear in n number of leaves, or a collection of vertex disjoint bare paths of (large) constant length each (Lemma 2.1). The former case is rather easy; similarly to the argument outlined in [2], we first embed the subtree F of T, obtained by deleting from T a linear number of leaves, by a straightforward greedy algorithm (Lemma 2.2). Then we embed the remaining edges between the omitted leaves of T and their fathers (Lemma 2.3); the restriction $np \geq C\Delta \ln n$ is induced by this part. In the complementary case, where the number of leaves of T is relatively small, we first take out a linear number of disjoint constant length bare paths to obtain a subforest F of T; we embed F using the same greedy argument (Lemma 2.2). Then we are left with embedding the remaining bare paths; we do this by reducing the problem to that of finding a factor of cycles (with some extra conditions imposed) in a random graph, and then by invoking a beautiful result of Johansson, Kahn and Vu [11] about factors in random graphs (Lemma 2.4). In both above described cases we need our random edges to come in two independent chunks: the standard trick of representing $G \sim G(n, p)$ as $G = G_1 \cup G_2$, where $G_i \sim G(n, p_i)$ and $1 - p = (1 - p_1)(1 - p_2)$, allows for this readily.

The notation used in the paper is pretty standard. We systematically suppress rounding signs for the sake of clarity of presentation.

The proofs of Theorems 1 and 2 are given in the next section. The last section of the paper is devoted to concluding remarks.

2 Proofs

Lemma 2.1 Let k, l, n > 0 be integers. Let T be a tree on n vertices with at most l leaves. Then T contains a collection of at least $\frac{n-(2l-2)(k+1)}{k+1}$ vertex disjoint bare paths of length k each.

Proof. Define

$$V_1 = \{v \in V(T) : d(v) = 1\},$$

$$V_2 = \{v \in V(T) : d(v) = 2\},$$

$$V_3 = \{v \in V(T) : d(v) \ge 3\}.$$

Clearly V_1 is the set of leaves of T and thus satisfies $|V_1| \leq l$. We have:

$$\begin{split} 2n-2 &= 2|E(T)| = \sum_{v \in V(T)} d(v) &\geq |V_1| + 2|V_2| + 3|V_3| \\ &= 2(|V_1| + |V_2| + |V_3|) + (|V_3| - |V_1|) \\ &= 2n + |V_3| - |V_1| \,, \end{split}$$

implying $|V_3| \le |V_1| - 2 \le l - 2$.

T has $|V_1|+|V_3|-1 \le 2l-3$ internally disjoint paths connecting between the vertices of $V_1 \cup V_3$, with all internal vertices of these paths being of degree two. In each such path, pick a largest collection of vertex disjoint subpaths of length k. This leaves at most k vertices of the path uncovered, so altogether the so formed collection of bare paths of length k in T contains all but at most $(|V_1|+|V_3|)+(|V_1|+|V_3|-1)k < (2l-2)(k+1)$ vertices, implying that the total number of paths of length k in the collection is at least $\frac{n-(2l-2)(k+1)}{k+1}$ as required.

Lemma 2.2 Let 0 < a < 1 be a constant. Let F be a tree on (1-a)n vertices of maximum degree Δ . If $anp \geq 3\Delta + 5 \ln n$, then **whp** a random graph G(n,p) contains a copy of F.

Proof. Choose arbitrarily a root r of F and fix some search order π , say BFS, on F starting from r. Let $\pi = (v_1 = r, \dots, v_m)$ with m = (1 - a)n. We will embed F in $G \sim G(n, p)$ according to π . Let ϕ be the so constructed embedding.

Suppose we are to embed the children of a current vertex v_i , $1 \le i \le m-1$, in G. Let $U_i \subset [n]$ be the set of vertices already used for embedding, clearly $|U_i| < m$. Expose the edges of G from $\phi(v_i)$ to $[n] - U_i$. We need to find at most $d_F(v_i) \le \Delta$ neighbors of $\phi(v_i)$ outside U_i . The probability of this not happening is at most

$$Pr[Bin(n-m,p) < \Delta] \le e^{-\frac{(anp-\Delta)^2}{2anp}} < e^{-\frac{2anp}{9}} \ll \frac{1}{n}$$
.

Taking the union bound over all embedding steps, we conclude that whp G contains a copy of F. \square

Lemma 2.3 Let $0 < d_1, \ldots, d_k$ be integers satisfying: $d_i \le \Delta$, $\sum_{i=1}^k d_i = l$. Let $A = \{a_1, \ldots, a_k\}$, B be disjoint sets of vertices with |B| = l. Let G be a random bipartite graph with sides A and B, where each pair (a,b), $a \in A, b \in B$, is an edge of G with probability p, independently of other pairs. If

$$p \ge \frac{2\Delta \ln l}{l} \,,$$

then whp as $l \to \infty$ the random graph G contains a collection S_1, \ldots, S_k of vertex disjoint stars such that S_i is centered at a_i and has the remaining d_i vertices in B.

Proof. Define an auxiliary (random) bipartite graph G' with sides A' and B, where |A'| = |B| = l. The vertices of A' are partitioned into k pairwise disjoint sets A_1, \ldots, A_k with $|A_i| = d_i$, $1 \le i \le k$. G has an edge between $a' \in A_i$ and $b \in B$ with probability p_i , where $(1 - p_i)^{d_i} = 1 - p$, implying $p_i \ge p/d_i \ge p/\Delta$. The distribution G' induces the distribution of G by the obvious projection: G has an edge between $a \in A$ and $b \in B$ iff G' has some edge between A_i and B. Observe that if G' has a perfect matching M' then G has the desired collection of stars $\{S_i\}$, obtained by projecting A' back into A (the vertices of A_i are projected onto a_i).

By the classical results about random graphs (see, e.g., Section 7.3 of [6]) and the monotonicity of the property of having a perfect matching it is enough to require that all individual edge probabilities in G' are at least $(\ln l + \omega(1))/l$. Recalling that $p_i \geq p/\Delta$, we see that the lemma's assumption $p \geq \frac{2\Delta \ln l}{l}$ guarantees the required condition.

Lemma 2.4 Let $k \ge 3$ be a fixed integer. Let G be distributed as $G((k+1)n_0, p)$. Let $S = \{s_1, \ldots, s_{n_0}\}$, $T = \{t_1, \ldots, t_{n_0}\}$ be disjoint vertex subsets of $[(k+1)n_0]$. If

$$p \ge C \left(\frac{\ln n_0}{n_0^{k-1}}\right)^{1/k} \,,$$

for some large enough constant C = C(k), then whp G contains a family $\{P_i\}_{i=1}^{n_0}$ of vertex disjoint paths, where P_i is a path of length k connecting s_i and t_i .

Proof. Fix a partition of $V(G) - S \cup T$ into vertex disjoint subsets V_1, \ldots, V_{k-1} of cardinality $|V_i| = n_0$ each. Define an auxiliary graph H with vertex set $V(H) = X \cup V_1 \cup \ldots \cup V_{k-1}$, where $X = \{x_1, \ldots, x_{n_0}\}$. For $1 \le i \le k-2$, the edges of H between V_i and V_{i+1} are identical to those of G. For $v \in V_1$ and $x_j \in X$, (v, x_j) is an edge of H iff (v, s_j) is an edge of G. Similarly, for $v \in V_{k-1}$ and $v_j \in X$, $v_j \in G$ is an edge of G. Notice that each relevant pair in V(H) becomes an edge of G independently and with probability G.

Suppose now that H contains a C_k -factor $\{S_1, \ldots, S_{n_0}\}$, where each cycle S_j traverses the sets X, V_1, \ldots, V_{k-1} in this order. Each such cycle S_i translates to a path of length k between s_i and t_i in G, and these paths are pairwise disjoint.

It thus remains to argue that the random graph H contains **whp** the desired collection of cycles. This can be obtained from the result of Johansson, Kahn and Vu [11] through straightforward (but quite tedious) modification of their arguments. (They proved that a random graph G(kn, p) with $p \geq C(k) \left(\frac{\ln n}{n^{k-1}}\right)^{1/k}$ contains **whp** a factor of cycles C_k , we need the factor in a k-partite random graph; moreover, the cycles in the factor are required to traverse the parts in the prescribed order.) \square

Proof of Theorem 1. Set

$$\delta = \frac{\epsilon}{10},$$

$$k = \left[\frac{2}{\epsilon}\right].$$

We consider two cases.

Case 1. T has at least δn leaves.

We represent G as the union $E(G) = E(G_1) \cup E(G_2)$, where G_1 , G_2 are two independent random graphs, both distributed according to G(n, p') with $1 - p = (1 - p')^2$ (and thus $p' \ge p/2$). Let F be a subtree of T obtained by deleting from T an arbitrary set of δn leaves. We first find a copy $\phi(F)$ of F in G_1 – such a copy exists **whp** due to Lemma 2.2. Let now $B = [n] - V(\phi(F))$, and let $A \subset V(\phi(F))$ be the set of images of the fathers of the δn leaves deleted from T to form F. Denote $A = \{a_1, \ldots, a_k\}$, and let $d_i \le \Delta$ be the number of leaves in T connected to the preimage $\phi^{-1}(a_i)$ and left outside F; clearly $\sum_{i=1}^k d_i = \delta n$. In order to complete the embedding of T into G, we need to find in G K vertex disjoint stars S_1, \ldots, S_k , where the star S_i is centered in a_i and has the remaining d_i vertices in B. We

invoke Lemma 2.3 to find such stars **whp** using the (random) edges of G_2 between A and B. Since the edge probability in G_2 is at least p/2, we need to verify that

$$\frac{p}{2} \ge \frac{2\Delta \ln(\delta n)}{\delta n} \ .$$

Recalling that $\delta = \frac{\epsilon}{10}$ and $p \ge \frac{40\Delta \ln n}{\epsilon n}$, we see that this condition is fulfilled indeed.

Case 2. T has less than δn leaves.

We again represent G as the union $E(G) = E(G_1) \cup E(G_2)$ as in the previous case. According to Lemma 2.1, T contains a family of $n_0 = \frac{n - (2\delta n - 2)(k+1)}{k+1} = \Theta(n)$ of vertex disjoint bare paths of length k. Let F be a subforest of T obtained by deleting the internal vertices of such a family of bare paths. We first use the edges of G_1 to find **whp** a copy $\phi(F)$ of F; this is possible again due to Lemma 2.2. It now remains to insert these n_0 bare paths, connecting between prescribed pairs of vertices. We can apply Lemma 2.4 to the edges of G_2 to meet this goal. Since the edge probability in G_2 is at least p/2 and

$$\frac{p}{2} \ge \frac{n^{-1+\epsilon}}{2} \gg \left(\frac{\ln n}{n^{k-1}}\right)^{1/k}$$

(recall $\epsilon > 1/k$), the graph G_2 contains indeed the required collection of paths **whp**. The proof is complete.

Proof of Theorem 2. Set

$$k = \left\lceil \frac{n}{\Delta - 1} \right\rceil .$$

Consider the random graph G(n,p) with $p = \frac{\delta \Delta \ln n}{n}$, the value of $\delta = \delta(\epsilon)$ to be chosen later. Recall that in the tree $T(n,\Delta)$ k vertices of the spine path P dominate the rest of the graph. It thus suffices to show that **whp** G(n,p) has no dominating set of size k. The probability that such a dominating set exists is at most

$$\binom{n}{k} (1 - (1-p)^k)^{n-k} \le \left(\frac{en}{k}\right)^k e^{-(n-k)(1-p)^k}$$

$$\le (3\Delta)^{\frac{n}{\Delta}} e^{-\frac{n}{2}e^{-pk}} \le e^{\frac{2n\ln\Delta}{\Delta} - \frac{n^{1-\delta}}{3}}$$

$$\le e^{2n^{1-\epsilon}\ln n - \frac{n^{1-\delta}}{3}}.$$

Taking $\delta = \epsilon/2$ we see that **whp** the random graph G(n, p) does not contain a dominating set of size k and thus **whp** does not contain a copy of $T(n, \Delta)$.

3 Concluding remarks

We have shown that the (pretty immediate) lower bound on the edge probability $p(n) \ge c\Delta(T) \ln n/n$ for the random graph G(n,p) to contain **whp** a copy of a given spanning tree T of maximum degree Δ is tight up to a constant factor in the range $\Delta = n^{\Theta(1)}$.

The regime $\Delta(T) = n^{o(1)}$ stays largely open. In particular, we were not able to provide a satisfactory solution for the most natural case of embedding spanning trees with bounded maximum degree. Our result only shows that in this case is enough to require $p(n) = n^{-1+o(1)}$; this is probably not the tightest bound possible.

For the case of embedding a bounded degree spanning tree T with cn leaves [2] has shown that it is enough to take $p(n) = C \ln n/n$, where C may depend on c. It is unclear whether such a dependence is necessary. It seems plausible that assuming $p(n) = (1 + o(1)) \ln n/n$ may be enough.

Finally, it would be very interesting to obtain sufficient conditions for embedding spanning trees with given maximum degree applicable to *pseudo-random* graphs.

References

- [1] M. Ajtai, J. Komlós and E. Szemerédi, *The longest path in a random graph*, Combinatorica 1 (1981), 1-12.
- [2] N. Alon, M. Krivelevich and B. Sudakov, *Embedding nearly-spanning bounded degree trees*, Combinatorica 27 (2007), 629–644.
- [3] J. Balogh, B. Csaba, M. Pei and W. Samotij, Large bounded degree trees in expanding graphs, Electr. J. Combinatorics 17 (2010), Publication R6.
- [4] J. Balogh, B. Csaba and W. Samotij, Local resilience of almost spanning trees in random graphs, Random Struct. Algorithms, to appear.
- [5] B. Bollobás, The evolution of sparse graphs, in: *Graph Theory and Combinatorics (Cambridge*, 1983), Academic Press, London, (1984), 35–57.
- [6] B. Bollobás, Random graphs, 2nd ed., Cambridge Studies in Advanced Mathematics, 73, Cambridge University Press, Cambridge, 2001.
- [7] W. Fernandez de la Vega, Long paths in random graphs, Studia Sci. Math. Hungar. 14 (1979), 335-340.
- [8] W. Fernandez de la Vega, Trees in sparse random graphs, J. Combin. Theory Ser. B 45 (1988), 77-85.
- [9] J. Friedman and N. Pippenger, Expanding graphs contain all small trees, Combinatorica 7 (1987), 71–76.
- [10] P. Haxell, Tree embeddings, J. Graph Theory 36 (2001), 121–130.
- [11] A. Johansson, J. Kahn and V. Vu, Factors in random graphs, Random Struct. Algorithms 33 (2008), 1–28.

- [12] J. Kahn, private communication.
- [13] J. Komlós, G. Sárközy and E. Szemerédi, *Spanning trees in dense graphs*, Combin. Prob. Computing 10 (2001), 397–416.
- [14] J. Komlós and E. Szemerédi, Limit distributions for the existence of Hamilton circuits in a random graph, Discrete Math. 43 (1983), 55–63.