DETERMINING ELECTRICAL AND HEAT TRANSFER PARAMETERS USING COUPLED BOUNDARY MEASUREMENTS

KATSIARYNA KRUPCHYK, MATTI LASSAS, AND SAMULI SILTANEN

ABSTRACT. Let $\Omega \subset \mathbb{R}^n$, $n \geq 3$, be a smooth bounded domain and consider a coupled system in Ω consisting of a conductivity equation $\nabla \cdot \gamma(x) \nabla u(t,x) = 0$ and an anisotropic heat equation $\kappa^{-1}(x) \partial_t \psi(t,x) = \nabla \cdot (A(x) \nabla \psi(t,x)) + (\gamma \nabla u(t,x)) \cdot \nabla u(t,x)$, $t \geq 0$. It is shown that the coefficients γ , κ and $A = (a_{jk})$ are uniquely determined from the knowledge of the boundary map $u|_{\partial\Omega} \mapsto \nu \cdot A \nabla \psi|_{\partial\Omega}$, where ν is the unit outer normal to $\partial\Omega$.

The coupled system models the following physical phenomenon. Given a fixed voltage distribution, maintained on the boundary $\partial\Omega$, an electric current distribution appears inside Ω . The current in turn acts as a source of heat inside Ω , and the heat flows out of the body through the boundary. The boundary measurements above then correspond to the map taking a voltage distribution on the boundary to the resulting heat flow through the boundary. The presented mathematical results suggest a new hybrid diffuse imaging modality combining electrical prospecting and heat transfer-based probing.

Keywords: electrical impedance tomography, heat transfer, inverse problem, coupled systems

AMS subject classification: 35K20, 35J25, 35R30, 80A23

1. Introduction

Let us model a physical body by a bounded set $\Omega \subset \mathbb{R}^n$, $n \geq 3$, with smooth boundary $\partial\Omega$, and the following spatially varying quantities: heat capacity c(x), density $\rho(x)$, electric conductivity $\gamma(x)$, and (possibly anisotropic) thermal conductivity $A(x) = (a_{ik}(x))$, each defined for $x \in \overline{\Omega}$.

Consider applying a spatially and temporally variable electrical voltage distribution f(t,x) at the boundary $\partial\Omega$ starting at time t=0. Then, if there are no sinks or sources of current inside Ω , the electric potential u(t,x) inside the body satisfies the conductivity equation

$$\nabla \cdot \gamma \nabla u(t, x) = 0 \quad \text{for } x \in \Omega \text{ and } t \ge 0,$$

$$u(t, \cdot)|_{\partial\Omega} = f(t, \cdot). \tag{1.1}$$

Equation (1.1) is often used as a mathematical model for electrical impedance tomography (EIT), where one measures the current through the boundary caused by a family of static voltage distributions $f(t,x) = \phi(x)$ and recovers $\gamma(x)$ from

such voltage-to-current map $\Lambda_{\gamma}: \phi \mapsto \nu \cdot \nabla u|_{\partial\Omega}$. Here ν is the unit outer normal to $\partial\Omega$. We refer to [39] for an extensive survey of the mathematical developments in EIT. See also [3, 4, 32, 36, 37] for results in the two-dimensional case, and [13, 27, 28, 29, 31, 34, 38] for results in higher dimensional cases. For counterexamples to uniqueness of time-harmonic inverse problems involving very anisotropic and degenerate material parameters, leading to the phenomenon of invisibility, see [10, 11, 12].

Our aim here is somewhat different as we wish to couple *heat conduction* to the problem. Let us denote the electrical power density inside Ω by F:

$$F(t,x) = (\gamma \nabla u(t,x)) \cdot \nabla u(t,x). \tag{1.2}$$

Now F acts as a source of heat inside Ω . Assuming that the body is at a constant (zero) temperature at the time t=0 when the voltage is first applied, and the surface of the body is kept at that temperature at all times, the temperature distribution $\psi(t,x)$ inside Ω satisfies the following heat equation:

$$\kappa^{-1}(x)\partial_t \psi(t,x) = \nabla \cdot (A(x)\nabla \psi(t,x)) + F(t,x) \quad \text{for } x \in \Omega \text{ and } t \ge 0,$$

$$\psi|_{\overline{\mathbb{R}}_+ \times \partial\Omega} = 0, \quad \psi|_{t=0} = 0,$$
 (1.3)

where $\kappa(x) = c(x)^{-1}\rho(x)^{-1}$. The model (1.1), (1.2), (1.3) is based on the physical assumption that the heat transfer is so slow that the quasistatic (DC) model for the electric potential (1.1) is realistic.

Associated to the coupled system (1.1), (1.2), (1.3), we introduce the voltageto-heat flow map $\Sigma_{\gamma,\kappa,A}$ defined by

$$\Sigma_{\gamma,\kappa,A}: f \mapsto \nu \cdot A \nabla \psi|_{\overline{\mathbb{R}}_{+} \times \partial \Omega}. \tag{1.4}$$

The idea is to measure the heat flow through the boundary caused by the heat from the electric current resulting from the applied voltage distribution.

Our main result is Theorem 2.1 below, stating that under certain smoothness assumptions, the coefficients γ , κ , and A are uniquely determined from the knowledge of the voltage-to-heat flow map $\Sigma_{\gamma,\kappa,A}$.

The method of proof of Theorem 2.1 also outlines a constructive reconstruction procedure for recovering conductivity γ from $\Sigma_{\gamma,\kappa,A}$. Namely, it turns out that applying a temporally static voltage distribution $f(t,x) = \phi(x)$ and studying $\Sigma_{\gamma,\kappa,A}f$ at thermal equilibrium $(t \to \infty)$ yields the knowledge of the Dirichlet-to-Neumann map $\Lambda_{\gamma}\phi$ related to the EIT problem. Then one can recover γ using Nachman's reconstruction result [31].

Notice that various hybrid imaging methods have been proposed and analyzed recently. Examples include thermoacoustic and photoacoustic imaging [1, 5, 25], combination of electrical and magnetic probing [26, 33], electrical and acoustic imaging [15] and magnetic and acoustic imaging [30, 2]. Theorem 2.1 suggests a new hybrid imaging method, utilizing two diffuse modes of propagation: electrical prospecting and heat transfer-based probing. We emphasize that the proposed method recovers complementary information about three different physical




FIGURE 1. Schematic illustration of the practical measurement setup motivating the proposed hybrid imaging method. The ideal voltage-to-heat flow map $\Sigma_{\gamma,\kappa,A}$ may be approximated in practice by maintaining fixed voltages at the electrodes while measuring heat flow using the interlaced sensors.

properties. We also note that in many applications where one wants to reconstruct the heat transfer parameters $\kappa(x)$ and A(x), the use of electric boundary sources may be easier than controlling the temperature or the heat flux at the boundary. Concerning inverse problems for the heat equation, we refer to [6, 7, 19, 20, 21, 22, 23, 40].

We remark that in practice one might use a measurement setup shown in Figure 1. However, analysis of such discrete measurements is outside the scope of this paper, and in the mathematical results below we work with the continuum models (1.1), (1.2), (1.3), and (1.4).

This paper is organized as follows. In Section 2 we state our assumptions and results in a mathematically precise form. In Section 3 we give an auxiliary density result for the conductivity equation. Section 4 is devoted to the reconstruction of the conductivity γ . The proof of Theorem 2.1 is completed in Section 5, where we show the identifiability of the heat parameters κ and A. Finally, Appendix A is devoted to the recovery of the boundary values of the matrix A from interior—to—boundary measurements, associated to a suitable elliptic boundary value problem. This result may be of an independent interest.

2. Statement of results

Let $\Omega \subset \mathbb{R}^n$, $n \geq 3$, be a bounded domain with C^{∞} boundary. Let $\gamma \in C^{\infty}(\overline{\Omega})$ be a strictly positive function on $\overline{\Omega}$. Then given $f(t,x) \in C^1(\overline{\mathbb{R}}_+, H^s(\partial\Omega))$, $s \geq 1/2$, on the boundary at time $t \geq 0$, there exists a unique $u \in C^1(\overline{\mathbb{R}}_+, H^{s+1/2}(\Omega))$, which solves the boundary value problem

$$\nabla \cdot \gamma \nabla u(t, x) = 0 \quad \text{for } x \in \Omega \text{ and } t \ge 0,$$

$$u(t, \cdot)|_{\partial \Omega} = f(t, \cdot),$$
 (2.1)

see [16]. We have

$$F(t,x) = (\gamma \nabla u(t,x)) \cdot \nabla u(t,x) \in C^{1}(\overline{\mathbb{R}}_{+}, L^{2}(\Omega)), \tag{2.2}$$

provided that s is taken large enough, say s > (n+1)/2. In what follows we shall always choose the Sobolev index s in this way. Consider the anisotropic heat equation

$$\kappa^{-1}(x)\partial_t \psi(t,x) = \nabla \cdot (A(x)\nabla \psi(t,x)) + F(t,x) \quad \text{for } x \in \Omega \text{ and } t \ge 0,$$

$$\psi|_{\overline{\mathbb{R}}_+ \times \partial\Omega} = 0, \quad \psi|_{t=0} = 0.$$
 (2.3)

Here $A(x) = (a_{jk}(x))$ is a real symmetric $n \times n$ matrix with $a_{j,k}(x) \in C^{\infty}(\overline{\Omega})$, and there exists $C_0 > 0$ such that

$$\sum_{j,k=1}^{n} a_{jk}(x)\xi_{j}\xi_{k} \ge C_{0}|\xi|^{2}, \quad \text{for all } (x,\xi) \in \overline{\Omega} \times \mathbb{R}^{n}.$$
 (2.4)

We shall assume that $0 < \kappa \in C^{\infty}(\overline{\Omega})$. The operator

$$Pv = -\kappa(x)\nabla \cdot (A(x)\nabla v)$$

is formally self-adjoint in $L^2_{\kappa} = L^2(\Omega, \kappa^{-1} dx)$ and we have

$$(Pv, v)_{L^2_x} \ge C_0 ||v||_{L^2}^2, \quad v \in C_0^{\infty}(\Omega), \quad C_0 > 0.$$

We also let P denote the Friedrichs extension of the operator P on $C_0^{\infty}(\Omega)$, so that the domain of the positive self-adjoint operator P is $(H_0^1 \cap H^2)(\Omega)$.

The solution of (2.3) is given by the Duhamel formula

$$\psi(t,x) = \int_0^t (e^{-(t-s)P} \kappa F)(s,x) ds \in C^1(\overline{\mathbb{R}}_+, L^2(\Omega)) \cap C(\overline{\mathbb{R}}_+, (H_0^1 \cap H^2)(\Omega)), \quad (2.5)$$

see [16].

Associated to the coupled system (2.1), (2.2), and (2.3), we consider the voltage-to-heat flow map,

$$\Sigma_{\gamma,\kappa,A}: C^{1}(\overline{\mathbb{R}}_{+}, H^{s}(\partial\Omega)) \to C(\overline{\mathbb{R}}_{+}, H^{1/2}(\partial\Omega)),$$
$$f \mapsto \nu \cdot A\nabla\psi|_{\overline{\mathbb{R}}_{+}\times\partial\Omega},$$

The main result of the paper is as follows.

Theorem 2.1. Assume that $0 < \gamma_j \in C^{\infty}(\overline{\Omega})$, $0 < \kappa_j \in C^{\infty}(\overline{\Omega})$, and A_j are real symmetric $n \times n$ matrices with $C^{\infty}(\overline{\Omega})$ entries, satisfying (2.4), for j = 1, 2. If $\Sigma_{\gamma_1,\kappa_1,A_1} = \Sigma_{\gamma_2,\kappa_2,A_2}$, then $\gamma_1 = \gamma_2$, $\kappa_1 = \kappa_2$ and $A_1 = A_2$.

It turns out that in the course of the proof of Theorem 2.1, we establish a result for the anisotropic heat equation, which may be of independent interest. In order to state the result, consider the inhomogeneous initial boundary value

problem (2.3) for the anisotropic heat equation with an arbitrary source $F \in C^1(\overline{\mathbb{R}}_+, L^2(\Omega))$. Define the map,

$$\Xi_{\kappa,A}: C^{1}(\overline{\mathbb{R}}_{+}, L^{2}(\Omega)) \to C(\overline{\mathbb{R}}_{+}, H^{1/2}(\partial\Omega)),$$
$$F \mapsto \nu \cdot A\nabla \psi^{F}|_{\overline{\mathbb{R}}_{+} \times \partial\Omega},$$

where ψ^F is the solution of (2.3).

Theorem 2.2. Let $\Omega \subset \mathbb{R}^n$, $n \geq 2$, be a bounded domain with C^{∞} boundary. Assume that $0 < \kappa_j \in C^{\infty}(\overline{\Omega})$, and A_j are real symmetric $n \times n$ matrices with $C^{\infty}(\overline{\Omega})$ entries, satisfying (2.4), for j = 1, 2. If $\Xi_{\kappa_1, A_1} = \Xi_{\kappa_2, A_2}$, then $\kappa_1 = \kappa_2$ and $A_1 = A_2$.

3. An auxiliary density result

Let $\Omega \subset \mathbb{R}^n$, $n \geq 3$, be a bounded domain with C^{∞} boundary and let $\gamma \in C^{\infty}(\overline{\Omega})$ be a strictly positive function on $\overline{\Omega}$. We shall need the following density result, which is a quite straightforward consequence of [38].

Proposition 3.1. The set

$$\operatorname{span}\{\gamma \nabla w_1 \cdot \nabla w_2 : w_i \in C^{\infty}(\overline{\Omega}), \nabla \cdot \gamma \nabla w_i = 0, j = 1, 2\}$$

is dense in $L^2(\Omega)$.

Proof. Let $f \in L^2(\Omega)$ be such that

$$\int_{\Omega} f \gamma \nabla w_1 \cdot \nabla w_2 dx = 0, \tag{3.1}$$

for any solution $w_1, w_2 \in C^{\infty}(\overline{\Omega})$ of the conductivity equation

$$\nabla \cdot \gamma \nabla w = 0. \tag{3.2}$$

We have

$$\gamma \nabla w_1 \cdot \nabla w_2 = \frac{1}{2} \nabla \cdot (\gamma \nabla (w_1 w_2)). \tag{3.3}$$

It follows from [38], see also [5], that for any $\rho \in \mathbb{C}^n$ satisfying $\rho \cdot \rho = 0$ and $|\rho| \geq 1$ large enough, the conductivity equation (3.2) has a solution

$$w_{\rho}(x) = e^{i\rho \cdot x} \gamma^{-1/2} (1 + r_{\rho}),$$
 (3.4)

where $r_{\rho} \in C^{\infty}(\overline{\Omega})$ satisfies

$$||r_{\rho}||_{H^{m}(\Omega)} \le \frac{C_{m}}{|\rho|}, \quad m \ge 0.$$
 (3.5)

Here the constant C_m depends on Ω , n, and a finite number of derivatives of γ . Given $\xi \in \mathbb{R}^n$ and R > 0, according to [38], there exist $\rho_1, \rho_2 \in \mathbb{C}^n$ such that $\rho_j \cdot \rho_j = 0$, $\rho_1 + \rho_2 = \xi$ and $|\rho_j| \geq R$, j = 1, 2. For the solutions w_{ρ_1} and w_{ρ_2} of the form (3.4), we get

$$\nabla \cdot (\gamma \nabla (w_{\rho_1} w_{\rho_2})) = \nabla \cdot (\gamma \nabla (\gamma^{-1} e^{i\xi \cdot x} (1 + r_{\rho_1} + r_{\rho_2} + r_{\rho_1} r_{\rho_2})))$$

$$= (\nabla \cdot (\gamma \nabla \gamma^{-1}) + \gamma (\nabla \gamma^{-1}) \cdot i\xi - |\xi|^2) e^{i\xi \cdot x} (1 + r_{\rho_1} + r_{\rho_2} + r_{\rho_1} r_{\rho_2})$$

$$+ (\gamma \nabla \gamma^{-1} + 2i\xi) e^{i\xi \cdot x} \cdot (\nabla r_{\rho_1} + \nabla r_{\rho_2} + \nabla (r_{\rho_1} r_{\rho_2}))$$

$$+ e^{i\xi \cdot x} (\Delta r_{\rho_1} + \Delta r_{\rho_2} + \Delta (r_{\rho_1} r_{\rho_2})).$$

In view of (3.3), we may substitute the latter expression into (3.1) and let $R \to \infty$. Using (3.5), we obtain that

$$\int_{\Omega} (\nabla \cdot (\gamma \nabla \gamma^{-1}) + \gamma (\nabla \gamma^{-1}) \cdot i\xi - |\xi|^2) e^{i\xi \cdot x} f dx = 0 \quad \text{for all } \xi \in \mathbb{R}^n.$$
 (3.6)

Here we shall view γ as a strictly positive C^{∞} function on \mathbb{R}^n , which is equal to a positive constant near infinity. The identity (3.6) is equivalent to

$$\mathcal{F}_{x \to \xi}(\nabla \cdot (\gamma \nabla \gamma^{-1}) \chi_{\Omega} f - \nabla \cdot (\gamma \nabla (\gamma^{-1}) \chi_{\Omega} f) + \Delta(\chi_{\Omega} f)) = 0,$$

where $\mathcal{F}_{x\to\xi}$ denotes the Fourier transformation and χ_{Ω} is the characteristic function of Ω . It follows that $\chi_{\Omega}f$ is a solution of a second order elliptic equation on \mathbb{R}^n with smooth coefficients. Since it is compactly supported, by unique continuation we conclude that $f \equiv 0$ in Ω . This completes the proof.

4. Recovering the conductivity γ from the voltage-to-heat flow map

The purpose of this section is to make the first step in the proof of Theorem 2.1, by establishing the following result. Recall that here $n \geq 3$.

Proposition 4.1. The voltage-to-heat flow map $\Sigma_{\gamma,\kappa,A}$ determines the conductivity γ uniquely.

When proving Proposition 4.1, we let $\alpha \in C^{\infty}(\overline{\mathbb{R}}_+; [0,1])$ be such that

$$\alpha|_{t<1/2} = 0, \quad \alpha|_{t>1} = 1.$$

Then set $f(t,x) = \alpha(t)h(x)$ with $h \in H^s(\partial\Omega)$, s large enough. Using the Duhamel formula (2.5), we shall study the behavior of $\psi(t,x)$ as $t \to +\infty$. The solution u of (2.1) satisfies

$$u(t,x) = \alpha(t)w_0(x),$$

where w_0 solves

$$\nabla \cdot \gamma \nabla w_0(x) = 0 \quad \text{in} \quad \Omega,$$

$$w_0|_{\partial\Omega} = h(x).$$
(4.1)

Thus,

$$F(t,x) = \gamma \alpha^2(t) \nabla w_0 \cdot \nabla w_0,$$

and (2.5) gives

$$\psi(t,x) = \int_0^t \alpha^2(s)e^{-(t-s)P}\gamma\kappa\nabla w_0 \cdot \nabla w_0 ds. \tag{4.2}$$

Lemma 4.2. We have

$$\|\psi(t,\cdot)-P^{-1}(\gamma\kappa\nabla w_0\cdot\nabla w_0)\|_{\mathcal{D}(P)}\to 0, \text{ as } t\to +\infty.$$

Here $\mathcal{D}(P) = (H^2 \cap H_0^1)(\Omega)$ is equipped with the graph norm $\|\psi\|_{\mathcal{D}(P)} = \|\psi\|_{L^2(\Omega)} + \|P\psi\|_{L^2(\Omega)}$.

Proof. We shall first check that

$$\|\psi(t,\cdot) - P^{-1}(\gamma \kappa \nabla w_0 \cdot \nabla w_0)\|_{L^2(\Omega)} \to 0, \text{ as } t \to +\infty.$$
 (4.3)

It follows from (4.2) that for t > 1, we have

$$\psi(t,x) = \int_0^1 \alpha^2(s) e^{-(t-s)P} \gamma \kappa \nabla w_0 \cdot \nabla w_0 ds + \int_0^{t-1} e^{-sP} \gamma \kappa \nabla w_0 \cdot \nabla w_0 ds.$$

Using that

$$\int_0^{+\infty} e^{-tP} dt = P^{-1},$$

in the sense of bounded operators on $L^2(\Omega)$, see [16], we get with L^2 -convergence, as $t \to +\infty$,

$$\int_0^{t-1} e^{-sP} \gamma \kappa \nabla w_0 \cdot \nabla w_0 ds \to P^{-1}(\gamma \kappa \nabla w_0 \cdot \nabla w_0).$$

On the other hand,

$$\begin{split} &\| \int_0^1 \alpha^2(s) e^{-(t-s)P} \gamma \kappa \nabla w_0 \cdot \nabla w_0 ds \|_{L^2(\Omega)} \\ &\leq \int_0^1 \alpha^2(s) \| e^{-(t-s)P} \gamma \kappa \nabla w_0 \cdot \nabla w_0 \|_{L^2(\Omega)} ds \\ &\leq \int_0^1 \alpha^2(s) \| e^{-(t-s)P} \|_{\mathcal{L}(L^2(\Omega), L^2(\Omega))} \| \gamma \kappa \nabla w_0 \cdot \nabla w_0 \|_{L^2(\Omega)} ds \to 0, \text{ as } t \to +\infty, \end{split}$$

since an application of the spectral theorem shows that

$$||e^{-(t-s)P}||_{\mathcal{L}(L^2(\Omega),L^2(\Omega))} \le \sup_{\lambda \in \operatorname{spec}(P)} e^{-\lambda(t-s)} \to 0, \text{ as } t \to +\infty.$$

This establishes (4.3).

Next we shall show that

$$||P\psi(t,\cdot) - \gamma\kappa\nabla w_0 \cdot \nabla w_0||_{L^2(\Omega)} \to 0$$
, as $t \to +\infty$. (4.4)

We have for t > 1,

$$P\psi(t,x) = P \int_0^1 \alpha^2(s) e^{-(t-s)P} \gamma \kappa \nabla w_0 \cdot \nabla w_0 ds + P \int_0^{t-1} e^{-sP} \gamma \kappa \nabla w_0 \cdot \nabla w_0 ds.$$

The formula

$$P \int_0^t e^{-sP} v ds = -e^{-tP} v + v,$$

see [16], implies that

$$P \int_0^{t-1} e^{-sP} \gamma \kappa \nabla w_0 \cdot \nabla w_0 ds = -e^{-(t-1)P} \gamma \kappa \nabla w_0 \cdot \nabla w_0 + \gamma \kappa \nabla w_0 \cdot \nabla w_0,$$

and

$$||e^{-(t-1)P}\gamma\kappa\nabla w_0\cdot\nabla w_0||_{L^2(\Omega)}\to 0$$
, as $t\to +\infty$.

Finally,

$$||P \int_{0}^{1} \alpha^{2}(s) e^{-(t-s)P} \gamma \kappa \nabla w_{0} \cdot \nabla w_{0} ds||_{L^{2}(\Omega)}$$

$$\leq \int_{0}^{1} \alpha^{2}(s) ||P e^{-(t-s)P}||_{\mathcal{L}(L^{2}(\Omega), L^{2}(\Omega))} ||\gamma \kappa \nabla w_{0} \cdot \nabla w_{0}||_{L^{2}(\Omega)} ds \to 0, \text{ as } t \to +\infty,$$

since

$$||Pe^{-(t-s)P}||_{\mathcal{L}(L^2(\Omega),L^2(\Omega))} \le \sup_{\lambda \in \operatorname{spec}(P)} (\lambda e^{-(t-s)\lambda}) \to 0, \text{ as } t \to +\infty.$$

This proves (4.4) and completes the proof of the lemma.

Lemma 4.2 implies that as $t \to +\infty$,

$$\nu \cdot A \nabla \psi|_{\partial\Omega} \to \nu \cdot A \nabla (P^{-1}(\gamma \kappa \nabla w_0 \cdot \nabla w_0))|_{\partial\Omega}$$
 in $H^{1/2}(\partial\Omega)$.

Thus, as $t \to +\infty$, we have by a repeated application of the divergence theorem together with the Cauchy-Schwarz inequality,

$$\int_{\partial\Omega} \Sigma_{\gamma,\kappa,A}(\alpha(t)h(x))dS = \int_{\partial\Omega} \nu \cdot A\nabla \psi dS \to \int_{\partial\Omega} \nu \cdot A\nabla (P^{-1}(\gamma\kappa\nabla w_0 \cdot \nabla w_0))dS$$
$$= \int_{\Omega} \nabla \cdot A\nabla (P^{-1}(\gamma\kappa\nabla w_0 \cdot \nabla w_0))dx = -\int_{\Omega} \gamma\nabla w_0 \cdot \nabla w_0 dx = -\int_{\partial\Omega} \Lambda_{\gamma}(h)hdS.$$

Here

$$\Lambda_{\gamma}: H^{s}(\partial\Omega) \to H^{s-1}(\partial\Omega),$$

 $h \mapsto \gamma \partial_{\nu} w_{0}|_{\partial\Omega},$

is the Dirichlet-to-Neumann map, associated to the problem (4.1). Thus, the knowledge of the voltage-to-heat flow map $\Sigma_{\gamma,\kappa,A}$ determines the Dirichlet-to-Neumann map Λ_{γ} . It follows from [38] that the isotropic conductivity $\gamma \in C^{\infty}(\overline{\Omega})$ is uniquely determined by $\Sigma_{\gamma,\kappa,A}$. This completes the proof of Proposition 4.1.

5. Recovering the heat parameters A and κ

The purpose of this section is to prove the following result.

Proposition 5.1. If
$$\Sigma_{\gamma,\kappa_1,A_1} = \Sigma_{\gamma,\kappa_2,A_2}$$
, then $\kappa_1 = \kappa_2$ and $A_1 = A_2$.

When determining the conductivity γ in the previous section, we were concerned with the power densities F which are supported in the region $t \geq 1/2$ and independent of t near $+\infty$. Here we shall instead concentrate the density F in a small neighborhood of t=0.

Let $\chi \in C^{\infty}(\overline{\mathbb{R}}_+)$ be such that $0 \leq \chi \leq 1$, supp $(\chi) \subset [0,1]$ and

$$\int \chi^2(t)dt = 1.$$

For $\varepsilon > 0$, we define

$$\chi_{\varepsilon}(t) = \varepsilon^{-1/2} \chi(t/\varepsilon), \tag{5.1}$$

so that $\chi^2_{\varepsilon}(t) \to \delta(t)$, the Dirac measure at t = 0, as $\varepsilon \to 0$. Let $h \in H^s(\partial\Omega)$, for s large enough. Then the solution $u = u^{\varepsilon,h}$ of the problem

$$\nabla \cdot \gamma \nabla u(t, x) = 0 \quad \text{in} \quad \Omega,$$

$$u|_{\partial \Omega} = \chi_{\varepsilon}(t)h(x),$$

satisfies $u^{\varepsilon,h}(t,x) = \chi_{\varepsilon}(t)w^h(x)$, where $w^h(x)$ solves

$$\nabla \cdot \gamma \nabla w^h(x) = 0 \quad \text{in} \quad \Omega,$$

$$w^h|_{\partial\Omega} = h(x).$$
(5.2)

Let $h, \widetilde{h} \in H^s(\partial\Omega)$ and

$$F^{\varepsilon,h\pm\widetilde{h}} = \gamma \nabla (u^{\varepsilon,h} \pm u^{\varepsilon,\widetilde{h}}) \cdot \nabla (u^{\varepsilon,h} \pm u^{\varepsilon,\widetilde{h}}) = \gamma \chi_{\varepsilon}^2(t) \nabla (w^h \pm w^{\widetilde{h}}) \cdot \nabla (w^h \pm w^{\widetilde{h}}).$$

Denote by $\psi^{\varepsilon,h\pm\tilde{h},j}$ the solution of (2.3) with $F = F^{\varepsilon,h\pm\tilde{h}}$, $\kappa = \kappa_j$ and $A = A_j$, j = 1, 2. Set

$$\alpha^{(j)}(t,x) = \alpha^{\varepsilon,h,\widetilde{h},j}(t,x) = \frac{1}{4}(\psi^{\varepsilon,h+\widetilde{h},j}(t,x) - \psi^{\varepsilon,h-\widetilde{h},j}(t,x)).$$

Thus, $\alpha^{(j)}$ is a solution of the following inhomogeneous initial boundary value problem,

$$(\partial_{t} + P_{j})\alpha^{(j)} = \frac{1}{4}\kappa_{j}(x)\chi_{\varepsilon}^{2}(t)\gamma$$

$$\left(\nabla(w^{h} + w^{\tilde{h}}) \cdot \nabla(w^{h} + w^{\tilde{h}}) - \nabla(w^{h} - w^{\tilde{h}}) \cdot \nabla(w^{h} - w^{\tilde{h}})\right)$$

$$= \kappa_{j}(x)\chi_{\varepsilon}^{2}(t)\gamma\nabla w^{h} \cdot \nabla w^{\tilde{h}}, \text{ in } \mathbb{R}_{+} \times \Omega,$$

$$\alpha^{(j)}|_{\mathbb{R}_{+} \times \partial \Omega} = 0, \quad \alpha^{(j)}|_{t=0} = 0.$$

$$(5.3)$$

Since

$$P_j = -\kappa_j \nabla \cdot A_j \nabla$$

is a self-adjoint positive operator in $L^2_{\kappa_j}(\Omega)$ with the domain $\mathcal{D}(P_j) = (H^1_0 \cap H^2)(\Omega)$, the spectrum of P_j is discrete, accumulating at $+\infty$, consisting of eigenvalues of finite multiplicity, $0 < \lambda_1^{(j)} \leq \lambda_2^{(j)} \leq \cdots \to \infty$. Associated to the eigenvalues $\lambda_k^{(j)}$ we have the eigenfunctions $\varphi_k^{(j)} \in \mathcal{D}(P_j)$, which form an orthonormal basis in $L^2_{\kappa_j}(\Omega)$. In what follows we shall assume, as we may, that the eigenfunctions $\varphi_k^{(j)}$ are real-valued. Hence,

$$\alpha^{(j)}(t,x) = \sum_{k=1}^{\infty} c_{k,\varepsilon}^{(j)}(t) \varphi_k^{(j)}(x), \quad c_{k,\varepsilon}^{(j)} = (\alpha^{(j)}, \varphi_k^{(j)})_{L_{\kappa_j}^2},$$

with convergence in $C(\overline{\mathbb{R}}_+, \mathcal{D}(P_i))$. Therefore,

$$\nu \cdot A_j \nabla \alpha^{(j)}|_{\partial \Omega} = \sum_{k=1}^{\infty} c_{k,\varepsilon}^{(j)}(t) (\nu \cdot A_j \nabla \varphi_k^{(j)})|_{\partial \Omega}$$

with convergence in $H^{1/2}(\partial\Omega)$, for each fixed $t \geq 1$.

Next we notice that

$$\nu \cdot A_j \nabla \alpha^{(j)}|_{\partial \Omega \times \overline{\mathbb{R}}_+} = \frac{1}{4} \Sigma_{\gamma, \kappa_j, A_j} (\chi_{\varepsilon}(t)(h + \widetilde{h})) - \frac{1}{4} \Sigma_{\gamma, \kappa_j, A_j} (\chi_{\varepsilon}(t)(h - \widetilde{h})).$$

Thus, since $\Sigma_{\gamma,\kappa_1,A_1} = \Sigma_{\gamma,\kappa_2,A_2}$, it follows that for all $t \geq 1$,

$$\sum_{k=1}^{\infty} c_{k,\varepsilon}^{(1)}(t) (\nu \cdot A_1 \nabla \varphi_k^{(1)})|_{\partial \Omega} = \sum_{k=1}^{\infty} c_{k,\varepsilon}^{(2)}(t) (\nu \cdot A_2 \nabla \varphi_k^{(2)})|_{\partial \Omega}.$$
 (5.4)

Here we would like to let $\varepsilon \to 0$. In order to do so, it will be convenient to obtain an explicit representation of the Fourier coefficients $c_{k,\varepsilon}^{(j)}(t)$.

Set $d_k^{(j)} = (\gamma \nabla w^h \cdot \nabla w^{\tilde{h}}, \varphi_k^{(j)})_{L^2}$, where the scalar product is taken in the space $L^2(\Omega, dx)$. It follows from (5.3) that

$$\partial_t c_{k,\varepsilon}^{(j)}(t) + \lambda_k^{(j)} c_{k,\varepsilon}^{(j)}(t) = \chi_{\varepsilon}^2(t) d_k^{(j)},$$
$$c_{k,\varepsilon}^{(j)}(0) = 0.$$

Hence,

$$c_{k,\varepsilon}^{(j)}(t) = e^{-\lambda_k t} d_k^{(j)} \int_0^t e^{\lambda_k^{(j)} s} \chi_{\varepsilon}^2(s) ds, \quad t \ge 0,$$

and $c_{k,\varepsilon}^{(j)}(t)$ is uniformly bounded in k, ε . For $t \ge 1$ and $k = 1, 2, \ldots$, fixed, we get

$$c_{k,\varepsilon}^{(j)}(t) = e^{-\lambda_k^{(j)}t} d_k \int_0^1 \chi^2(s) e^{\lambda_k^{(j)}s\varepsilon} ds \to e^{-\lambda_k^{(j)}t} d_k^{(j)}, \text{ as } \varepsilon \to 0.$$
 (5.5)

Using (5.5), we may let $\varepsilon \to 0$ in (5.4), and conclude that

$$\sum_{k=1}^{\infty} e^{-\lambda_k^{(1)} t} d_k^{(1)} (\nu \cdot A_1 \nabla \varphi_k^{(1)}) |_{\partial \Omega} = \sum_{k=1}^{\infty} e^{-\lambda_k^{(2)} t} d_k^{(2)} (\nu \cdot A_2 \nabla \varphi_k^{(2)}) |_{\partial \Omega}, \quad t \ge 1. \quad (5.6)$$

In what follows we shall have to distinguish the eigenvalues of the operator P_j , j=1,2. In order to do that let us continue to denote by $\lambda_k^{(j)}$ the sequence of distinct eigenvalues of P_j and let $m_k^{(j)}$ denote the multiplicity of $\lambda_k^{(j)}$. Let $\varphi_{k,1}^{(j)}, \ldots, \varphi_{k,m_k^{(j)}}^{(j)}$ be an orthonormal basis of the eigenfunctions, corresponding to the eigenvalue $\lambda_k^{(j)}$. By the uniqueness of the Dirichlet series, see [9], and (5.6), we obtain the following result.

Proposition 5.2. Assume that $\Sigma_{\gamma,\kappa_1,A_1} = \Sigma_{\gamma,\kappa_2,A_2}$. Then for all k = 1, 2, ..., we have

$$\lambda_k^{(1)} = \lambda_k^{(2)},$$

and

$$\sum_{i=1}^{m_k^{(1)}} d_{k,i}^{(1)} (\nu \cdot A_1 \nabla \varphi_{k,i}^{(1)})|_{\partial \Omega} = \sum_{i=1}^{m_k^{(2)}} d_{k,i}^{(2)} (\nu \cdot A_2 \nabla \varphi_{k,i}^{(2)})|_{\partial \Omega}.$$
 (5.7)

Here $d_{k,i}^{(j)} = (\gamma \nabla w^h \cdot \nabla w^{\widetilde{h}}, \varphi_{k,i}^{(j)})_{L^2}$ and $h, \widetilde{h} \in H^s(\partial \Omega)$ are arbitrary functions.

Let us introduce the following linear continuous operators

$$R_k^{(j)}: L^2(\Omega) \to L^2(\partial\Omega),$$

$$R_k^{(j)}(F) = \sum_{i=1}^{m_k^{(j)}} (F, \varphi_{k,i}^{(j)})_{L^2} (\nu \cdot A_j \nabla \varphi_{k,i}^{(j)})|_{\partial \Omega}, \ j = 1, 2, \ k = 1, 2, \dots.$$

Proposition 3.1 together with (5.7) implies that $R_k^{(1)} = R_k^{(2)}$ on a dense subset of $L^2(\Omega)$, and hence, everywhere. On the level of the distribution kernels, we obtain that for all $k = 1, 2, \ldots$,

$$\sum_{i=1}^{m_k^{(1)}} \varphi_{k,i}^{(1)}(x) (\nu \cdot A_1 \nabla \varphi_{k,i}^{(1)})(y) = \sum_{i=1}^{m_k^{(2)}} \varphi_{k,i}^{(2)}(x) (\nu \cdot A_2 \nabla \varphi_{k,i}^{(2)})(y), x \in \Omega, y \in \partial\Omega. \quad (5.8)$$

We shall next need the following result, see [7]. Since the argument is short, for the convenience of the reader, we give it here.

Lemma 5.3. The functions $(\nu \cdot A_j \nabla \varphi_{k,1}^{(j)})|_{\partial\Omega}, \ldots, (\nu \cdot A_j \nabla \varphi_{k,m_k^{(j)}}^{(j)})|_{\partial\Omega}$ are linearly independent, j = 1, 2.

Proof. Assume that there are $c_1, \ldots, c_{m_k^{(j)}} \in \mathbb{R}$ such that

$$\sum_{i=1}^{m_k^{(j)}} c_i(\nu \cdot A_j \nabla \varphi_{k,i}^{(j)})|_{\partial \Omega} = 0.$$

Then $\varphi = \sum_{i=1}^{m_k^{(j)}} c_i \varphi_{k,i}^{(j)}$ satisfies

$$P_j \varphi = \lambda_k^{(j)} \varphi \quad \text{in} \quad \Omega,$$

 $\varphi|_{\partial\Omega} = 0, \quad \nu \cdot A_j \nabla \varphi|_{\partial\Omega} = 0.$

By the unique continuation principle, we get $\varphi = 0$ in Ω . Thus, $c_i = 0$ for $1 \le i \le m_k^{(j)}$. This proves the lemma.

Our next goal is to analyze the consequences of (5.8), and the key step here is the following algebraic result which is similar to [7, Lemma 2.3].

Lemma 5.4. Let $f_i: \Omega \to \mathbb{R}$, $\widetilde{f}_i: \partial \Omega \to \mathbb{R}$, $i = 1, ..., m^{(1)}$, and $g_l: \Omega \to \mathbb{R}$, $\widetilde{g}_l: \partial \Omega \to \mathbb{R}$, $l = 1, ..., m^{(2)}$, be such that

$$\sum_{i=1}^{m^{(1)}} f_i(x)\widetilde{f}_i(y) = \sum_{l=1}^{m^{(2)}} g_l(x)\widetilde{g}_l(y) \quad \text{for all } x \in \Omega, y \in \partial\Omega.$$
 (5.9)

Moreover, assume that the systems $\{f_1, \ldots, f_{m^{(1)}}\}$, $\{\widetilde{f}_1, \ldots, \widetilde{f}_{m^{(1)}}\}$, $\{g_1, \ldots, g_{m^{(2)}}\}$ and $\{\widetilde{g}_1, \ldots, \widetilde{g}_{m^{(2)}}\}$ are all linearly independent. Then $m^{(1)} = m^{(2)}$ and there exists an $m^{(1)} \times m^{(1)}$ invertible matrix T with real entries such that

$$F(x) = TG(x), \quad \widetilde{F}(y) = (T^t)^{-1}\widetilde{G}(y) \quad \textit{for all } x \in \Omega, y \in \partial \Omega.$$

Here we use the notation

$$F(x) = \begin{pmatrix} f_1(x) \\ \vdots \\ f_{m^{(1)}}(x) \end{pmatrix}, \widetilde{F}(y) = \begin{pmatrix} \widetilde{f}_1(y) \\ \vdots \\ \widetilde{f}_{m^{(1)}}(y) \end{pmatrix},$$
$$G(x) = \begin{pmatrix} g_1(x) \\ \vdots \\ g_{m^{(2)}}(x) \end{pmatrix}, \widetilde{G}(y) = \begin{pmatrix} \widetilde{g}_1(y) \\ \vdots \\ \widetilde{g}_{m^{(2)}}(y) \end{pmatrix}.$$

Proof. As \widetilde{f}_1 is not identically zero in $\partial\Omega$, there exists $y_1 \in \partial\Omega$ such that $\widetilde{f}_1(y_1) \neq 0$. Assuming that

$$\det\begin{pmatrix} \widetilde{f}_1(y_1) & \widetilde{f}_2(y_1) \\ \widetilde{f}_1(y) & \widetilde{f}_2(y) \end{pmatrix} = 0 \quad \text{for all } y \in \partial\Omega,$$

we get that $\widetilde{f}_1, \widetilde{f}_2$ are linearly dependent which contradicts the assumptions of the proposition. Thus, there exists $y_2 \in \partial \Omega$ such that

$$\det\begin{pmatrix} \widetilde{f}_1(y_1) & \widetilde{f}_2(y_1) \\ \widetilde{f}_1(y_2) & \widetilde{f}_2(y_2) \end{pmatrix} \neq 0.$$

Continuing in the same way, we find points $y_1, y_2, \ldots, y_{m^{(1)}} \in \partial \Omega$ such that the matrix

$$Q_{\widetilde{f}} = \begin{pmatrix} \widetilde{f}_{1}(y_{1}) & \widetilde{f}_{2}(y_{1}) & \dots & \widetilde{f}_{m^{(1)}}(y_{1}) \\ \widetilde{f}_{1}(y_{2}) & \widetilde{f}_{2}(y_{2}) & \dots & \widetilde{f}_{m^{(1)}}(y_{2}) \\ \vdots & \vdots & \vdots & \vdots \\ \widetilde{f}_{1}(y_{m^{(1)}}) & \widetilde{f}_{2}(y_{m^{(1)}}) & \dots & \widetilde{f}_{m^{(1)}}(y_{m^{(1)}}) \end{pmatrix}$$

is invertible. It follows from (5.9) that $Q_{\tilde{f}}F(x) = Q_{\tilde{g}}G(x)$ for any $x \in \Omega$, where

$$Q_{\widetilde{g}} = \begin{pmatrix} \widetilde{g}_1(y_1) & \dots & \widetilde{g}_{m^{(2)}}(y_1) \\ \vdots & \vdots & \vdots \\ \widetilde{g}_1(y_{m^{(1)}}) & \dots & \widetilde{g}_{m^{(2)}}(y_{m^{(1)}}) \end{pmatrix}.$$

Thus, F(x) = TG(x) for any $x \in \Omega$, where $T = Q_{\widetilde{f}}^{-1}Q_{\widetilde{g}}$, and therefore, $m^{(1)} \leq m^{(2)}$. Similarly, using the fact that $\{\widetilde{g}_1,\ldots,\widetilde{g}_{m^{(2)}}\}$ is linearly independent, we have $G(x) = T_1F(x)$ for any $x \in \Omega$, and $m^{(2)} \leq m^{(1)}$. Hence, $m^{(1)} = m^{(2)}$. Furthermore,

$$F(x) = TT_1F(x)$$
 for all $x \in \Omega$. (5.10)

Since the system $\{f_1,\ldots,f_{m^{(1)}}\}$ is linearly independent, in the same way as above, we see that there are points $x_1,\ldots,x_{m^{(1)}}\in\Omega$ such that the vectors $F(x_1),\ldots,F(x_{m^{(1)}})$ form a basis in $\mathbb{C}^{m^{(1)}}$. Thus, (5.10) implies that $TT_1=I$ and therefore, T is invertible. Similarly, one can see that $\widetilde{F}(y)=\widetilde{T}\widetilde{G}(y)$ for all $y\in\partial\Omega$ with an invertible matrix \widetilde{T} .

It follows from (5.9) that $F(x) \cdot \widetilde{F}(y) = G(x) \cdot \widetilde{G}(y)$ and therefore,

$$(\widetilde{T}^tT-I)G(x)\cdot\widetilde{G}(y)=0\quad\text{for all }x\in\Omega,y\in\partial\Omega.$$

Since there exist points $x_1, \ldots, x_{m^{(1)}} \in \Omega$ and $y_1, \ldots, y_{m^{(1)}} \in \partial \Omega$ such that the vectors $G(x_1), \ldots G(x_{m^{(1)}})$ (respectively, $\widetilde{G}(y_1), \ldots \widetilde{G}(y_{m^{(1)}})$) form a basis in $\mathbb{C}^{m^{(1)}}$, we get $\widetilde{T}^t T = I$. This proves the claim.

If follows from Lemma 5.4 together with (5.8) and Lemma 5.3 that $m_k^{(1)} = m_k^{(2)} =: m_k$, and there exists an $m_k \times m_k$ invertible matrix T such that

$$\begin{pmatrix} \varphi_{k,1}^{(1)}(x) \\ \vdots \\ \varphi_{k,m_k}^{(1)}(x) \end{pmatrix} = T \begin{pmatrix} \varphi_{k,1}^{(2)}(x) \\ \vdots \\ \varphi_{k,m_k}^{(2)}(x) \end{pmatrix}, \quad x \in \Omega,$$
 (5.11)

and

$$\begin{pmatrix} \nu \cdot A_1(y) \nabla \varphi_{k,1}^{(1)}(y) \\ \vdots \\ \nu \cdot A_1(y) \nabla \varphi_{k,m_k}^{(1)}(y) \end{pmatrix} = (T^t)^{-1} \begin{pmatrix} \nu \cdot A_2(y) \nabla \varphi_{k,1}^{(2)}(y) \\ \vdots \\ \nu \cdot A_2(y) \nabla \varphi_{k,m_k}^{(2)}(y) \end{pmatrix}, \quad y \in \partial \Omega. \quad (5.12)$$

Using (5.11) and (5.12), we have

$$T^{t}T\begin{pmatrix} \nu \cdot A_{1}(y)\nabla\varphi_{k,1}^{(2)}(y) \\ \vdots \\ \nu \cdot A_{1}(y)\nabla\varphi_{k,m_{k}}^{(2)}(y) \end{pmatrix} = \begin{pmatrix} \nu \cdot A_{2}(y)\nabla\varphi_{k,1}^{(2)}(y) \\ \vdots \\ \nu \cdot A_{2}(y)\nabla\varphi_{k,m_{k}}^{(2)}(y) \end{pmatrix}, \quad y \in \partial\Omega.$$
 (5.13)

The next step is to show that the matrix T is in fact orthogonal. This will follow once we establish the following result.

Proposition 5.5. If $\Sigma_{\gamma,\kappa_1,A_1} = \Sigma_{\gamma,\kappa_2,A_2}$ then $A_1|_{\partial\Omega} = A_2|_{\partial\Omega}$.

Proof. Consider the following elliptic boundary value problem,

$$P_{j}u = \kappa_{j}\gamma \nabla w^{h} \cdot \nabla w^{\tilde{h}} \quad \text{in} \quad \Omega,$$

$$u|_{\partial\Omega} = 0,$$
 (5.14)

where w^h (respectively, $w^{\tilde{h}}$) is the solution to the problem (5.2) with the boundary source $h \in H^s(\partial\Omega)$ (respectively, $\tilde{h} \in H^s(\partial\Omega)$), $s \geq 1/2$ large enough. We shall now return to the original notation, where each eigenvalue $\lambda_k^{(j)}$ of the operator P_j is repeated according its multiplicity. Since $0 \notin \operatorname{spec}(P_j)$, the problem (5.14) has the unique solution

$$u^{(j)} = \sum_{k=1}^{\infty} \frac{d_k^{(j)}}{\lambda_k^{(j)}} \varphi_k^{(j)}, \quad d_k^{(j)} = (\gamma \nabla w^h \cdot \nabla w^{\tilde{h}}, \varphi_k^{(j)})_{L^2},$$

with convergence in $H^2(\Omega)$. Thus,

$$\nu \cdot A_j \nabla u^{(j)}|_{\partial\Omega} = \sum_{k=1}^{\infty} \frac{d_k^{(j)}}{\lambda_k^{(j)}} \nu \cdot A_j \nabla \varphi_k^{(j)}|_{\partial\Omega},$$

and therefore, it follows from Proposition 5.2 that if $\Sigma_{\gamma,\kappa_1,A_1} = \Sigma_{\gamma,\kappa_2,A_2}$, then

$$\nu \cdot A_1 \nabla u^{(1)}|_{\partial\Omega} = \nu \cdot A_2 \nabla u^{(2)}|_{\partial\Omega}. \tag{5.15}$$

Define the continuous map

$$\Psi^{(j)}: L^2(\Omega) \to L^2(\partial\Omega), \quad F \mapsto \nu \cdot A_j \nabla u^{(F,j)}|_{\partial\Omega},$$

 $u^{(F,j)}$ is a solution to the problem

$$-\nabla \cdot (A_j \nabla u^{(F,j)}) = F \quad \text{in} \quad \Omega,$$
$$u^{(F,j)}|_{\partial \Omega} = 0.$$

It follows from (5.15) together with Proposition 3.1 that $\Psi^{(1)} = \Psi^{(2)}$ on a dense subset of $L^2(\Omega)$, and thus, everywhere. Hence, Proposition A.1 in Appendix A implies that $A_1|_{\partial\Omega} = A_2|_{\partial\Omega}$.

Now going back to equation (5.13), using Lemma 5.3 we obtain that T is an orthogonal matrix. Proposition 5.2 together with (5.11) gives the following result.

Proposition 5.6. Assume that $\Sigma_{\gamma,\kappa_1,A_1} = \Sigma_{\gamma,\kappa_2,A_2}$. Let $\varphi_k^{(1)}$ be an orthonormal basis in $L^2(\Omega, \kappa_1^{-1}dx)$ of the Dirichlet eigenfunctions of the operator P_1 . Then the Dirichlet eigenvalues $\lambda_k^{(1)}$ (respectively, $\lambda_k^{(2)}$) of the operator P_1 (respectively, P_2), counted with multiplicities, satisfy $\lambda_k^{(1)} = \lambda_k^{(2)}$ for all k and there exists an orthonormal basis in $L^2(\Omega, \kappa_2^{-1}dx)$ of the Dirichlet eigenfunctions $\varphi_k^{(2)}$ of the operator P_2 such that $\varphi_k^{(1)} = \varphi_k^{(2)}$ for all k.

We shall next show that Proposition 5.6 yields that $\kappa_1 = \kappa_2$. Indeed, let us write

$$\kappa_1 = \sum_{k=1}^{\infty} c_k \varphi_k^{(1)} = \sum_{k=1}^{\infty} c_k \varphi_k^{(2)}, \quad \kappa_2 = \sum_{k=1}^{\infty} d_k \varphi_k^{(2)},$$

where the Fourier coefficients c_k are given by

$$c_k = \int_{\Omega} \kappa_1 \varphi_k^{(1)} \kappa_1^{-1} dx = \int_{\Omega} \varphi_k^{(2)} dx = \int_{\Omega} \kappa_2 \varphi_k^{(2)} \kappa_2^{-1} dx = d_k.$$

Thus, $\kappa_1 = \kappa_2$. It follows that $P_1 u = P_2 u$, for any $u \in C_0^{\infty}(\Omega)$, and we get $A_1 = A_2$. The proof of Theorem 2.1 is complete.

Theorem 2.2 can be proven by exactly the same arguments presented in this section applied to the problem (2.3) with the right hand sides of the form

$$F(t,x) = \chi_{\varepsilon}^2(t)H(x), \quad H \in L^2(\Omega),$$

where χ_{ε} is given by (5.1).

APPENDIX A. BOUNDARY RECONSTRUCTION

Let $\Omega \subset \mathbb{R}^n$, $n \geq 2$, be a bounded domain with C^{∞} boundary, and $A(x) = (a_{jk}(x))$, $1 \leq j, k \leq n$, be a real symmetric $n \times n$ matrix with $a_{j,k}(x) \in C^{\infty}(\overline{\Omega})$. Assume that there exists $C_0 > 0$ such that

$$\sum_{j,k=1}^{n} a_{jk}(x)\xi_{j}\xi_{k} \ge C_{0}|\xi|^{2} \quad \text{for all } (x,\xi) \in \overline{\Omega} \times \mathbb{R}^{n}.$$

Consider the following elliptic boundary value problem,

$$-\nabla \cdot (A\nabla u) = F \quad \text{in} \quad \Omega,$$

$$u|_{\partial\Omega} = 0. \tag{A.1}$$

For any $F \in L^2(\Omega)$, the problem (A.1) has a unique solution $u = u^F \in H^2(\Omega) \cap H^1_0(\Omega)$ and one can define the map,

$$\Psi: L^2(\Omega) \to L^2(\partial\Omega), \quad \Psi(F) = \nu \cdot A\nabla u^F|_{\partial\Omega},$$
 (A.2)

where ν is the unit outer normal to the boundary $\partial\Omega$. We note that the map Ψ is sometimes used to model boundary measurements for optical tomography with diffusion approximation, [17, 18, 35].

We have the following proposition, which is closely related to the earlier boundary reconstruction results of a Riemannian metric from the Dirichlet—to—Neumann map [24, 29].

Proposition A.1. The knowledge of the map Ψ given by (A.2) determines the values of A on the boundary $\partial\Omega$.

Proof. We shall recover the values of A on the boundary by analyzing the distribution kernel of the map Ψ , obtained by constructing a right parametrix for the boundary value problem (A.1). Let us denote $\mathcal{A} = -\nabla \cdot (A\nabla)$. Since A(x) is a positive definite matrix, smoothly depending on x, we can view $\overline{\Omega}$ as a Riemannian manifold with boundary, equipped with the metric $G = A^{-1}$, $G = (g_{ij})$, $1 \leq i, j \leq n$. To construct a parametrix for (A.1), we shall work locally near a boundary point. Let $x_0 \in \partial \Omega$ and introduce the boundary normal coordinates $y = (y', y_n) \in U$, $y' = (y_1, \dots, y_{n-1})$, centered at x_0 . Here U stands for some open neighborhood of 0 in \mathbb{R}^n . In terms of the boundary normal coordinates, locally near x_0 , the boundary $\partial \Omega$ is defined by $y_n = 0$, and $y_n > 0$ if and only if $x \in \Omega$. In what follows, we shall write again (x', x_n) for the boundary normal coordinates.

In the boundary normal coordinates, the metric G has the form

$$G = \sum_{\alpha,\beta=1}^{n-1} g_{\alpha\beta}(x) dx_{\alpha} dx_{\beta} + (dx_n)^2,$$

see [29], and the principal symbol of the operator \mathcal{A} is given by

$$a_0(x,\xi) = \xi_n^2 + \sum_{\alpha,\beta=1}^{n-1} g^{\alpha\beta}(x)\xi_\alpha\xi_\beta.$$

Therefore, the equation $a_0(x, \xi', \xi_n) = 0$, $\xi' = (\xi_1, \dots, \xi_{n-1})$, has the solutions,

$$\xi_n = \lambda_{\pm}(x, \xi'), \quad \lambda_{\pm}(x, \xi') = \pm i \sqrt{\sum_{\alpha, \beta = 1}^{n-1} g^{\alpha\beta}(x) \xi_{\alpha} \xi_{\beta}}.$$
 (A.3)

We can view \mathcal{A} as a linear continuous map in the space $\mathcal{D}'(U)$. In the boundary normal coordinates, the problem (A.1) has the following form,

$$\mathcal{A}u = F \text{ in } \mathbb{R}^n_+ = \{ x \in \mathbb{R}^n : x_n > 0 \},$$

 $u|_{x_n=0} = 0.$ (A.4)

Let

$$r_0(x,\xi) = a_0(x,\xi)^{-1}(1-\chi(\xi)), \quad x \in U, \quad \xi \in \mathbb{R}^n,$$

where $\chi(\xi) \in C_0^{\infty}(\mathbb{R}^n)$, $\chi(\xi) = 0$ for $|\xi| \ge 1$ and $\chi = 1$ near 0. The operator $Op(r_0)$ is a rough parametrix for the operator \mathcal{A} , which will be sufficient for our

purposes. Here we are using the classical quantization of a symbol $a \in S^k(U \times \mathbb{R}^n)$, which is given by

$$\operatorname{Op}(a)u(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{i(x-y)\cdot\xi} a(x,\xi)u(y) dy d\xi.$$

As usual, we say that $a \in S^k(U \times \mathbb{R}^n)$ if locally uniformly in $x \in U$, we have

$$|\partial_x^{\alpha} \partial_{\xi}^{\beta} a(x,\xi)| \le C_{\alpha\beta} \langle \xi \rangle^{k-|\beta|}, \quad \langle \xi \rangle = \sqrt{1+|\xi|^2}.$$

Let r_+ be the operation of restriction from \mathbb{R}^n to \mathbb{R}^n_+ and let e_+ be the operation of extension by zero from \mathbb{R}^n_+ to \mathbb{R}^n .

We shall construct a right parametrix for the boundary value problem (A.4) in the following form

$$R(F) = r_{+} \operatorname{Op}(r_{0})(e_{+}F) + R_{b}(\psi).$$

Here

$$\psi(x') = -\tau_0 r_+ \operatorname{Op}(r_0)(e_+ F), \quad \tau_0 : u \mapsto u|_{x_n = 0},$$

and R_b will be constructed as a right parametrix for the boundary value problem

$$\mathcal{A}u = 0 \quad \text{in} \quad \mathbb{R}^n_+,
u|_{x_n=0} = \psi(x').$$
(A.5)

In what follows we shall suppress the operator r_+ from the notation, as this will cause no confusion.

When constructing the operator R_b , we shall follow the standard approach in the theory of elliptic boundary value problems, see [8]. To this end, let $\widetilde{\chi} \in C_0^{\infty}(\mathbb{R}^{n-1})$ be such that $\widetilde{\chi} = 1$ for $|\xi'| \leq 1$. Notice that

$$(1 - \widetilde{\chi}(\xi'))\chi(\xi) = 0.$$

Let $\sigma = \sigma(x, \xi')$ be a simple closed C^1 smooth curve in the upper half-plane $\operatorname{Im} \xi_n > 0$, which encircles the root $\lambda_+(x, \xi')$ in the positive sense. In what follows we may and will choose σ so that it is independent of $x \in U$, depending on ξ' only, i.e. $\sigma = \sigma(\xi')$. When $\varphi \in C_0^{\infty}(U \cap \mathbb{R}^{n-1})$, we define the operator

$$(\Pi\varphi)(x) = \frac{1}{i(2\pi)^n} \int_{\mathbb{R}^{n-1}} e^{ix'\cdot\xi'} \widehat{\varphi}(\xi') (1 - \widetilde{\chi}(\xi')) \left(\int_{\xi_n \in \sigma} \frac{e^{ix_n\xi_n}}{a_0(x,\xi)} d\xi_n \right) d\xi', \quad x_n \ge 0,$$
(A.6)

where

$$\widehat{\varphi}(\xi') = \int_{\mathbb{R}^{n-1}} e^{-iy' \cdot \xi'} \varphi(y') dy'$$

is the Fourier transform of φ . By a contour deformation argument in the complex ξ_n -plane, we have

$$\Pi \varphi = \frac{1}{i} \operatorname{Op}((1 - \widetilde{\chi}) a_0^{-1}) (\varphi \otimes \delta_{x_n = 0}), \quad x_n > 0.$$

We get therefore,

$$\mathcal{A}\Pi\varphi = \operatorname{Op}(b)(\varphi \otimes \delta_{x_n=0}), \quad b \in S^{-1}(U \times \mathbb{R}^n), \quad x_n > 0,$$

since the operator \mathcal{A} is local.

We shall take $R_b(\psi) = \Pi \varphi$, for some function φ , defined locally near $0 \in \mathbb{R}^{n-1}$, to be found from the boundary condition, i.e.

$$\tau_0 \Pi \varphi = \psi. \tag{A.7}$$

To this end, we shall now prove that $\tau_0\Pi$ is an elliptic pseudodifferential operator on the boundary and compute its principal symbol. By the residue calculus, using (A.6), we get

$$\tau_0 \Pi = \operatorname{Op}(d_0), \quad d_0(x', \xi') = (1 - \widetilde{\chi}(\xi')) \frac{1}{2\pi i} \int_{\xi_n \in \sigma} \frac{1}{a_0(x', 0, \xi)} d\xi_n$$
$$= (1 - \widetilde{\chi}(\xi')) \frac{1}{\partial_{\xi_n} a_0(x', 0, \xi', \lambda_+)} = (1 - \widetilde{\chi}(\xi')) \frac{1}{2i|\xi'|_A} \in S^{-1}((U \cap \mathbb{R}^{n-1}) \times \mathbb{R}^{n-1}),$$

where

$$|\xi'|_A = \sqrt{\sum_{\alpha,\beta=1}^{n-1} g^{\alpha\beta}(x',0)\xi_\alpha\xi_\beta}.$$

We introduce next a rough parametrix of $\tau_0\Pi$, given by $\operatorname{Op}(\widetilde{d}_0)$, where $\widetilde{d}_0 \in S^1((U \cap \mathbb{R}^{n-1}) \times \mathbb{R}^{n-1})$ is such that

$$\widetilde{d}_0 = 2i|\xi'|_A$$
, for $|\xi'|$ large.

To satisfy (A.7), we choose

$$\varphi = \operatorname{Op}(\widetilde{d}_0)\psi = -\operatorname{Op}(\widetilde{d}_0)(\tau_0\operatorname{Op}(r_0)(e_+F)).$$

This choice of φ completes the construction of a rough parametrix for the boundary value problem (A.5), given by

$$R_b(\psi) = -\Pi \operatorname{Op}(\widetilde{d}_0)(\tau_0 \operatorname{Op}(r_0)(e_+F)).$$

Hence, the parametrix for the problem (A.4) has the form

$$R(F) = \operatorname{Op}(r_0)(e_+F) - \operatorname{\PiOp}(\widetilde{d_0})(\tau_0 \operatorname{Op}(r_0)(e_+F)).$$

In the boundary normal coordinates, the operator Ψ is given by

$$\Psi(F) = \tau_0 \partial_{x_n} u,$$

and therefore, to obtain the claim of the proposition it suffices to analyze the distribution kernel K(x',y) of the operator $\tau_0 \partial_{x_n} R$ given by

$$(\tau_0 \partial_{x_n} R(F))(x') = \int_{\mathbb{R}^n} K(x', y) F(y) dy, \quad x' \in \mathbb{R}^{n-1}, \quad y \in \mathbb{R}^n.$$

Let us first consider the Schwartz kernel of the operator

$$\tau_0 \partial_{x_n} \operatorname{Op}(r_0)(e_+ F) = \frac{1}{(2\pi)^n} \tau_0 \partial_{x_n} \int_{\mathbb{R}^n} e^{ix \cdot \xi} r_0(x, \xi) \widehat{e_+ F}(\xi) d\xi,$$

which is given by

$$K^{(1)}(x',y) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x'-y')\cdot\xi'} e^{-iy_n\xi_n} (i\xi_n r_0(x',0,\xi) + \partial_{x_n} r_0(x',0,\xi)) d\xi.$$

Recall that here $y_n > 0$. Restricting the attention to the region $|\xi'| \ge 1$, by a contour deformation argument to the lower half plane, we find that

$$\int_{\mathbb{R}} e^{-iy_n \xi_n} i \xi_n r_0(x', 0, \xi) d\xi_n = 2\pi i e^{-iy_n \lambda_-} \frac{i \lambda_-}{\partial_{\xi_n} a_0(x', 0, \xi', \lambda_-)} = \pi e^{-y_n |\xi'|_A},$$

and therefore,

$$K^{(1)}(x',y) = \frac{1}{(2\pi)^{n-1}} \int_{\mathbb{R}^{n-1}} e^{i(x'-y')\cdot\xi'} t(x',y_n,\xi') d\xi',$$

where

$$t(x', y_n, \xi') = \frac{e^{-y_n|\xi'|_A}}{2} + e^{-y_n|\xi'|_A} \mathcal{O}\left(\frac{1}{|\xi'|_A}\right), \quad |\xi'| \ge 1.$$

Next, the operator $\tau_0 \partial_{x_n} \Pi$ is a pseudodifferential operator on \mathbb{R}^{n-1} , given by

$$(\tau_0 \partial_{x_n} \Pi) v(x') = \frac{1}{(2\pi)^{n-1}}$$

$$\int_{\xi' \in \mathbb{R}^{n-1}} (1 - \widetilde{\chi}(\xi')) e^{ix' \cdot \xi'} \widehat{v}(\xi') \frac{1}{2\pi i} \int_{\xi_n \in \sigma} \left(\frac{i\xi_n}{a_0(x', 0, \xi)} + \partial_{x_n} \left(\frac{1}{a_0(x', 0, \xi)} \right) \right) d\xi' d\xi_n.$$

The principal symbol of the operator $\tau_0 \partial_{x_n} \Pi$ is therefore

$$\frac{1}{2\pi i} \int_{\xi_n \in \sigma} \frac{i\xi_n}{a_0(x', 0, \xi)} d\xi_n = i \frac{\lambda_+}{\partial_{\xi_n} a_0(x', 0, \xi', \lambda_+)} = \frac{i}{2}, |\xi'| \text{ large enough.}$$

The operator $\operatorname{Op}(\widetilde{d}_0)$ is also a pseudodifferential operator on \mathbb{R}^{n-1} and its principal symbol is given by $2\lambda_+ \in S^1((U \cap \mathbb{R}^{n-1}) \times \mathbb{R}^{n-1})$, $|\xi'|$ large enough. Hence, the principal symbol of the operator $\tau_0 \partial_{x_n} \operatorname{\PiOp}(\widetilde{d}_0)$ is $i\lambda_+ \in S^1((U \cap \mathbb{R}^{n-1}) \times \mathbb{R}^{n-1})$, and therefore, its kernel is given by

$$\begin{split} K_1^{(2)}(x',z') &= \frac{1}{(2\pi)^{n-1}} \int_{\mathbb{R}^{n-1}} e^{i(x'-z')\cdot\xi'} d_1(x',\xi') d\xi', \\ d_1(x',\xi') &= i\lambda_+(x',0,\xi') + p_0(x',\xi'), \quad |\xi'| \text{ large enough,} \end{split}$$

where $p_0 \in S^0((U \cap \mathbb{R}^{n-1}) \times \mathbb{R}^{n-1})$.

Finally, the kernel of the operator $\tau_0 \text{Op}(r_0)$ is given by

$$K_2^{(2)}(z',y) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(z'-y')\cdot\eta'} e^{-iy_n\eta_n} r_0(z',0,\eta) d\eta$$
$$= \frac{1}{(2\pi)^{n-1}} \int_{\mathbb{R}^{n-1}} e^{i(z'-y')\cdot\eta'} d_2(z',y_n,\eta') d\eta',$$
$$d_2(z',y_n,\eta') = \frac{1}{2} \frac{e^{-y_n|\eta'|_A}}{|\eta'|_A}, \quad |\eta'| \text{ large enough.}$$

Here as usual we use the residue calculus, where only the pole in the lower half plane contributes.

Hence, the kernel of the composition $\tau_0 \partial_{x_n} \Pi \operatorname{Op}(\widetilde{d_0}) \tau_0 \operatorname{Op}(r_0)(e_+ F)$ is given by

$$K^{(2)}(x',y) = \int K_1^{(2)}(x',z') K_2^{(2)}(z',y) dz'$$

$$= \frac{1}{(2\pi)^{2(n-1)}} \iiint e^{i(x'-z')\cdot\xi'} e^{i(z'-y')\cdot\eta'} d_1(x',\xi') d_2(z',y_n,\eta') d\xi' d\eta' dz'$$

$$= \frac{1}{(2\pi)^{n-1}} \int e^{i(x'-y')\cdot\eta'} c(x',y_n,\eta') d\eta',$$

where

$$c(x', y_n, \eta') = \frac{1}{(2\pi)^{n-1}} \iint e^{i(x'-z')\cdot(\xi'-\eta')} d_1(x', \xi') d_2(z', y_n, \eta') d\xi' dz'.$$

Here $y_n \ge 0$ occurs as a parameter. Now $d_1(x', \xi') \in S^1((U \cap \mathbb{R}^{n-1}) \times \mathbb{R}^{n-1})$ for large $|\xi'|$ and $d_2(z', y_n, \eta')$ satisfies

$$|\partial_{z'}^{\alpha}\partial_{\eta'}^{\beta}\partial_{y_n}^{\gamma}d_2(z',y_n,\eta')| \leq C_{\alpha,\beta,\gamma}e^{-y_n\langle\eta'\rangle}\langle\eta'\rangle^{-1-|\beta|+|\gamma|},$$

where $\langle \eta' \rangle = \sqrt{1 + |\eta'|_A^2}$ is large enough. Hence, $c(x', y_n, \eta')$, depending on the parameter y_n , is the symbol of the composition of two pseudodifferential operators in the tangential directions. By the standard results on pseudodifferential operators, see [14], it has the following asymptotic expansion,

$$c(x', y_n, \eta') \sim \sum_{|\alpha| \geq 0} \frac{1}{\alpha!} \partial_{\eta'}^{\alpha} d_1(x', \eta') D_{x'}^{\alpha} d_2(x', y_n, \eta'),$$

with the leading term $d_1(x', \eta')d_2(x', y_n, \eta')$. The knowledge of the operator Ψ implies the knowledge of the kernel $K^{(1)}(x', y) - K^{(2)}(x', y)$ for any $x' \in U \cap \mathbb{R}^{n-1}$ and $y \in U \cap \overline{\mathbb{R}^n}$. This implies the knowledge of

$$\frac{e^{-y_n|\xi'|_A}}{2} + e^{-y_n|\xi'|_A} \mathcal{O}\left(\frac{1}{|\xi'|_A}\right) - c(x', y_n, \xi'), \quad \text{for any } |\xi'| \text{ large enough.}$$

The leading term of the latter expression is given by

$$\frac{e^{-y_n|\xi'|_A}}{2} + \frac{e^{-y_n|\xi'|_A}}{2} = e^{-y_n|\xi'|_A}.$$

Varying ξ' , we recover A(x',0). The proof is complete.

Acknowledgements

The research of K.K. was financially supported by the Academy of Finland (project 125599). The research of M.L. and S.S. was financially supported by the Academy of Finland (Center of Excellence programme 213476 and Computational Science Research Programme, project 134868). This project was partially

conducted at the Mathematical Sciences Research Institute, Berkeley, whose hospitality is gratefully acknowledged.

References

- [1] Ammari, H., Bonnetier, E., Capdeboscq, Y., Tanter, M., and Fink, M., Electrical impedance tomography by elastic deformation, SIAM J. Appl. Math. 68 (2008), no. 6, 1557–1573.
- [2] Ammari, H., Capdeboscq, Y., Kang, H., and Kozhemyak, A., Mathematical models and reconstruction methods in magneto-acoustic imaging European Journal of Applied Mathematics, 20 (2009), 303–317.
- [3] Astala, K., Päivärinta, L., Calderón's inverse conductivity problem in the plane, Ann. of Math. (2) **163** (2006), no. 1, 265–299.
- [4] Astala, K., Päivärinta, L., and Lassas, M., Calderón's inverse problem for anisotropic conductivity in the plane, Comm. Partial Differential Equations 30 (2005), no. 1–3, 207–224.
- [5] Bal, G., Uhlmann, G., Inverse Diffusion Theory of Photoacoustics, Inverse Problems, 26 (2010), 085010.
- [6] Bukhgeim, A., Klibanov, M., Uniqueness in the large of a class of multidimensional inverse problems, (Russian) Dokl. Akad. Nauk SSSR, **260** (1981), no. 2, 269–272.
- [7] Canuto, B., Kavian, O., Determining coefficients in a class of heat equations via boundary measurements, SIAM J. Math. Anal. 32 (2001), no. 5, 963–986.
- [8] Chazarain, J., Piriou, A., Introduction to the theory of linear partial differential equations, Translated from the French. Studies in Mathematics and its Applications, 14. North-Holland Publishing Co., Amsterdam-New York, 1982.
- [9] Choulli, M., Une introduction aux problèmes inverses elliptiques et paraboliques, volume 65 of Mathématiques & Applications (Berlin) [Mathematics & Applications], Springer-Verlag, Berlin, 2009.
- [10] Greenleaf, A., Kurylev, Y., Lassas, M., and Uhlmann, G., *Invisibility and inverse problems*, Bull. Amer. Math. Soc. (N.S.) **46** (2009), no. 1, 55–97.
- [11] Greenleaf, A., Kurylev, Y., Lassas, M., and Uhlmann, G., Full-wave invisibility of active devices at all frequencies, Comm. Math. Phys. **275** (2007), no. 3, 749–789.
- [12] Greenleaf, A., Lassas, M., and Uhlmann, G., On nonuniqueness for Calderón's inverse problem, Math. Res. Lett. **10** (2003), no. 5–6, 685–693.
- [13] Greenleaf, A., Lassas, M., and Uhlmann, G., The Calderón problem for conormal potentials. I. Global uniqueness and reconstruction, Comm. Pure Appl. Math. 56 (2003), no. 3, 328–352.
- [14] Grigis, A., Sjöstrand, J., Microlocal analysis for differential operators. An introduction, London Mathematical Society Lecture Note Series, 196. Cambridge University Press, Cambridge, 1994.
- [15] Gebauer, B., Scherzer, O., *Impedance-acoustic tomography*, SIAM J. Appl. Math., **69** (2008), no. 2, 565–576.
- [16] Grubb, G., *Distributions and operators*, Graduate Texts in Mathematics, volume 252, Springer, New York, 2009.
- [17] Haskell, R., Svaasand, L., Tsay, T., Feng, T., McAdams, M., and Tromberg, B., Boundary conditions for the diusion equation in radiative transfer, J. Opt. Soc. Am. A., 11 (1994), 2727–2741.
- [18] Hielscher, A., Jacques, S., Wang, L., and Tittel, F., The influence of boundary conditions on the accuracy of diffusion theory in time-resolved reflectance spectroscopy of biological tissue, Phys. Med. Biol. 40(1995), 1957–1975.
- [19] Imanuvilov, O., Yamamoto, M., Lipschitz stability in inverse parabolic problems by the Carleman estimate, Inverse Problems 14 (1998), no. 5, 1229–1245.

- [20] Isakov, V., Some inverse problems for the diffusion equation, Inverse Problems 15 (1999), no. 1, 3–10.
- [21] Katchalov, A., Kurylev, Y., and Lassas, M., *Inverse boundary spectral problems*, Chapman & Hall/CRC Monographs and Surveys in Pure and Applied Mathematics, 123, 2001.
- [22] Katchalov, A., Kurylev, Y., Lassas, M., and Mandache, N., Equivalence of time-domain inverse problems and boundary spectral problems, Inverse Problems 20 (2004), no. 2, 419–436.
- [23] Klibanov, M., *Inverse problems and Carleman estimates*, Inverse Problems 8 (1992), no. 4, 575–596.
- [24] Kohn, R., Vogelius, M., Determining conductivity by boundary measurements, Comm. Pure Appl. Math. 37 (1984), no. 3, 289–298.
- [25] Kuchment, P., Kunyansky, L., Mathematics of thermoacoustic tomography, European J. Appl. Math. 19 (2008), no. 2, 191–224.
- [26] Kwon, O., Woo, E., Yoon, J., and Seo, J., Magnetic resonance electrical impedance tomography (MREIT): simulation study of J-substitution algorithm, IEEE Trans. Biomed. Eng., 49 (2002), no. 2, 160–167.
- [27] Lassas, M., Uhlmann, G., On determining a Riemannian manifold from the Dirichlet-to-Neumann map. Ann. Sci. École Norm. Sup. (4) **34** (2001), no. 5, 771–787.
- [28] Lassas, M., Taylor, M., and Uhlmann, G., The Dirichlet-to-Neumann map for complete Riemannian manifolds with boundary, Comm. Anal. Geom. 11 (2003), no. 2, 207–221.
- [29] Lee, J., Uhlmann, G., Determining anisotropic real-analytic conductivities by boundary measurements, Comm. Pure Appl. Math., 42 (1989), no. 8, 1097–1112.
- [30] Ma, Q., and He, B., Investigation on magnetoacoustic signal generation with magnetic induction and its application to electrical conductivity reconstruction, Phys. Med. Biol., 52 (2007), 5085–5099.
- [31] Nachman, A., Reconstructions from boundary measurements, Ann. of Math. (2) 128 (1988), no. 3, 531–576.
- [32] Nachman, A., Global uniqueness for a two-dimensional inverse boundary value problem, Ann. of Math. (2) 143 (1996), no. 1, 71–96.
- [33] Nachman, A., Tamasan, A., and Timonov, A., Conductivity imaging with a single measurement of boundary and interior data, Inverse Problems, 23 (2007), 2551–2563.
- [34] Päivärinta, L., Panchenko, A., and Uhlmann, G., Complex geometrical optics solutions for Lipschitz conductivities, Rev. Mat. Iberoamericana 19 (2003), no. 1, 57–72.
- [35] Schweiger, M., Arridge, S., Hiraoka, M, and Delpy, D., The finite element method for the propagation of light in scattering media: Boundary and source conditions, Med Phys, 22 (1995), 1779–1792.
- [36] Sylvester, J., An anisotropic inverse boundary value problem, Comm. Pure Appl. Math. 43 (1990), no. 2, 201–232.
- [37] Sylvester, J., Uhlmann, G., A uniqueness theorem for an inverse boundary value problem in electrical prospection, Comm. Pure Appl. Math. 39 (1986), no. 1, 91–112.
- [38] Sylvester, J., Uhlmann, G., A global uniqueness theorem for an inverse boundary value problem, Ann. of Math. (2), **125** (1987), no. 1, 153–169.
- [39] Uhlmann, G., Electrical impedance tomography and Calderón's problem, Inverse Problems, 25 (2009), 123011.
- [40] Yamamoto, M., Carleman estimates for parabolic equations and applications, Inverse Problems, 25 (2009), 123013.

- K. Krupchyk, Department of Mathematics and Statistics, University of Helsinki,
- P.O. Box 68, FI-00014 Helsinki, Finland

 $E ext{-}mail\ address:$ katya.krupchyk@helsinki.fi

- M. Lassas, Department of Mathematics and Statistics, University of Helsinki,
- P.O. Box 68, FI-00014 Helsinki, Finland

 $E ext{-}mail\ address: matti.lassas@helsinki.fi}$

- S. SILTANEN, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF HELSINKI,
- P.O. Box 68, FI-00014 Helsinki, Finland

 $E ext{-}mail\ address: samuli.siltanen@helsinki.fi}$