

**ON THE EARLY HISTORY OF THE
SINGULAR VALUE DECOMPOSITION**

By

G.W. Stewart

IMA Preprint Series # 952

April 1992

On the Early History
of the
Singular Value Decomposition*

G. W. Stewart[†]

March 1992

ABSTRACT

This paper surveys the contributions of five mathematicians — Eugenio Beltrami (1835–1899), Camille Jordan (1838–1921), James Joseph Sylvester (1814–1897), Erhard Schmidt (1876–1959), and Hermann Weyl (1885–1955) — who were responsible for establishing the existence of the singular value decomposition and developing its theory.

*This report is available by anonymous ftp at `thales.cs.umd.edu` in the directory `pub/reports`.

[†]Department of Computer Science and Institute for Advanced Computer Studies, University of Maryland, College Park, MD 20742. This work was done during while the author was visiting the Institute for Mathematics and Its Applications, University of Minnesota, Minneapolis, MN 55455.

On the Early History of the Singular Value Decomposition

G. W. Stewart

For Gene Golub on his fifteenth birthday

1. Introduction

One of the most fruitful ideas in the theory of matrices is that of a matrix decomposition or canonical form. The theoretical utility of matrix decompositions has long been appreciated. More recently, they have become the mainstay of numerical linear algebra, where they serve as computational platforms from which a variety of problems can be solved.

Of the many useful decompositions, the singular value decomposition — that is, the factorization of a matrix \mathbf{A} into the product $\mathbf{U}\mathbf{\Sigma}\mathbf{V}^H$ of a unitary matrix \mathbf{U} a diagonal matrix $\mathbf{\Sigma}$ and another unitary matrix \mathbf{V}^H — has assumed a special role. There are several reasons. In the first place, the fact that the decomposition is achieved by unitary matrices makes it an ideal vehicle for discussing the geometry of n -space. Second, it is stable; small perturbations in \mathbf{A} correspond to small perturbations in $\mathbf{\Sigma}$, and conversely. Third, the diagonality of $\mathbf{\Sigma}$ makes it easy to determine when \mathbf{A} is near to a rank-degenerate matrix; and when it is, the decomposition provides optimal low rank approximations to \mathbf{A} . Finally, thanks to the pioneering efforts of Gene Golub, there exist efficient, stable algorithms to compute the singular value decomposition.

The purpose of this paper is to survey the contributions of five mathematicians — Eugenio Beltrami (1835–1899), Camille Jordan (1838–1921), James Joseph Sylvester (1814–1897), Erhard Schmidt (1876–1959), and Hermann Weyl (1885–1955) — who were responsible for establishing the existence of the singular value decomposition and developing its theory. Beltrami, Jordan, and Sylvester came to the decomposition through what we should now call linear algebra; Schmidt and Weyl approached it from integral equations. To give this survey context, we will begin with a brief description of the historical background.

It is an intriguing observation that most of the classical matrix decompositions predated the widespread use of matrices: they were cast in terms of determinants, linear systems of equations, and especially bilinear and quadratic forms. Gauss is the father of this development. Writing in 1823 [15, §31], he describes his famous

elimination algorithm as follows.

Specifically, the function Ω [a quadratic function of x, y, z , etc.] can be reduced to the form

$$\frac{u^0 u^0}{\mathcal{A}^0} + \frac{u' u'}{B'} + \frac{u'' u''}{C''} + \frac{u''' u'''}{D'''} + \text{etc.} + M,$$

in which the divisors $\mathcal{A}^0, B', C'', C'''$, etc. are constants and u^0, u', u'', u''' , etc. are linear functions of x, y, z , etc. However, the second function, u' , is independent of x ; the third, u'' , is independent of x and y ; the fourth, u''' is independent of x, y , and z , and so on. The last function $u^{(\pi-1)}$ depends only on the the last of the unknowns x, y, z , etc. Moreover, the coefficients \mathcal{A}^0, B', C'' , etc. multiply x, y, z , etc. in u^0, u', u'' , etc. respectively.

From this we easily see that Gauss's algorithm factors the matrix of the quadratic form $\mathbf{x}^T \mathbf{A} \mathbf{x}$ into the product $\mathbf{R} \mathbf{D}^{-1} \mathbf{R}$, where \mathbf{D} is diagonal and \mathbf{R} is upper triangular with the diagonals of \mathbf{D} on its diagonal. Gauss's functions u^0, u', u'' , etc. are the components of the vector $\mathbf{u} = \mathbf{R} \mathbf{x}$.

Gauss was also able to effectively obtain the inverse of a matrix by a process of *eliminatio indefinita*, in which the system of equations $\mathbf{y} = \mathbf{A} \mathbf{x}$ is transformed into the inverse system $\mathbf{x} = \mathbf{B} \mathbf{y}$. Gauss's skill in manipulating quadratic forms and systems of equations made possible his very general treatment of the theory and practice of least squares.

Other developments followed. Cauchy [6, 1829] established the properties of the eigenvalues and eigenvectors of a symmetric system (including the interlacing property) by considering the corresponding homogeneous system of equations. In 1846, Jacobi [25] gave his famous algorithm for diagonalizing a symmetric matrix, and in a posthumous paper [26, 1857] he obtained the LU decomposition by decomposing a bilinear form in the style of Gauss. Weierstrass [50, 1868] established canonical forms for pairs of bilinear functions — what we should today call the generalized eigenvalue problem. Thus the advent of the singular value decomposition in 1873 is seen as one of a long line of results on canonical forms.

We will use modern matrix notation to describe the early work on the singular value decomposition. Most of it slips as easily into matrix terminology as Gauss's description of his decomposition; and we shall be in no danger of anachronism, provided we take care to use matrix notation only as an expository device, and otherwise stick close to the writer's argument. The greatest danger is that the use of modern notation will trivialize the writer's accomplishments by making them obvious to our eyes. On the other hand, presenting them in the original scalar

form would probably exaggerate the obstacles these people had to overcome, since they were accustomed, as we are not, to grasping sets of equations as a whole,

With a single author, it is usually possible to modernize notation in such a way that it corresponds naturally to what he actually wrote. Here we are dealing with several authors, and uniformity is more important than correspondence with the original. Consequently, throughout paper we will be concerned with the singular value decomposition

$$\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T,$$

where \mathbf{A} is a real matrix of order n ,

$$\mathbf{\Sigma} = \text{diag}(\sigma_1, \sigma_2, \dots, \sigma_n)$$

has nonnegative diagonal elements arranged in descending order of magnitude, and

$$\mathbf{U} = (\mathbf{u}_1 \ \mathbf{u}_2 \ \dots \ \mathbf{u}_n) \quad \text{and} \quad \mathbf{V} = (\mathbf{v}_1 \ \mathbf{v}_2 \ \dots \ \mathbf{v}_n)$$

are orthogonal. The function $\|\cdot\|$ will denote the Frobenius norm defined by

$$\|\mathbf{A}\|^2 = \sum_{i,j} a_{ij}^2 = \sum_i \sigma_i^2.$$

In summarizing the contributions I have followed the principle that if you try to say everything you end up saying nothing. Most of the works treated here are richer than the following sketches would indicate, and the reader is advised to go to the sources for the full story.

2. Beltrami [5, 1873]

Together Beltrami and Jordan are the progenitors of the singular value decomposition, Beltrami by virtue of first publication and Jordan by the completeness and elegance of his treatment. Beltrami's contribution appeared in the *Journal of Mathematics for the Use of the Students of the Italian Universities*, and its purpose was to encourage students to become familiar with bilinear forms.

The Derivation. Beltrami begins with a bilinear form

$$f(\mathbf{x}, \mathbf{y}) = \mathbf{x}^T \mathbf{A} \mathbf{y},$$

where \mathbf{A} is real and of order n . If one makes the substitutions

$$\mathbf{x} = \mathbf{U}\boldsymbol{\xi} \quad \text{and} \quad \mathbf{y} = \mathbf{V}\boldsymbol{\eta},$$

then

$$f(\mathbf{x}, \mathbf{y}) = \boldsymbol{\xi}^T \mathbf{S} \boldsymbol{\eta},$$

where

$$\mathbf{S} = \mathbf{U}^T \mathbf{A} \mathbf{V}. \quad (2.1)$$

Beltrami now observes that if \mathbf{U} and \mathbf{V} are required to be orthogonal then there are $n^2 - n$ degrees of freedom in their choice, and he proposes to use these degrees of freedom to annihilate the off diagonal element of \mathbf{S} .

Assume that \mathbf{S} is diagonal; i.e. $\mathbf{S} = \boldsymbol{\Sigma} = \text{diag}(\sigma_1, \dots, \sigma_n)$. Then it follows from (2.1) and the orthogonality of \mathbf{V} that

$$\mathbf{U}^T \mathbf{A} = \boldsymbol{\Sigma} \mathbf{V}^T. \quad (2.2)$$

Similarly

$$\mathbf{A} \mathbf{V} = \mathbf{U} \boldsymbol{\Sigma}. \quad (2.3)$$

Substituting the value of \mathbf{U} obtained from (2.3) into (2.2), Beltrami obtains the equation

$$\mathbf{U}^T (\mathbf{A} \mathbf{A}^T) = \boldsymbol{\Sigma}^2 \mathbf{U}^T, \quad (2.4)$$

and similarly he obtains

$$(\mathbf{A}^T \mathbf{A}) \mathbf{V} = \mathbf{V} \boldsymbol{\Sigma}^2.$$

Thus the σ_i are the roots of the equations

$$\det(\mathbf{A} \mathbf{A}^T - \sigma^2 I) = 0 \quad (2.5)$$

and

$$\det(\mathbf{A}^T \mathbf{A} - \sigma^2 I) = 0. \quad (2.6)$$

Note that the derivation, as presented by Beltrami, assumes that $\boldsymbol{\Sigma}$, and hence \mathbf{A} , is nonsingular.¹

Beltrami now argues that the two functions (2.5) and (2.6) are identical because they are polynomials of degree n that assume the same values at $\sigma = \sigma_i$ ($i = 1, \dots, n$) and the common value $\det^2(A)$ at $\sigma = 0$, an argument that presupposes that the singular values are distinct and nonzero.

¹However, it is possible to derive the equations without assuming that \mathbf{A} is nonsingular; e.g., $\mathbf{U}^T \mathbf{A} \mathbf{A}^T = \boldsymbol{\Sigma} \mathbf{V}^T \mathbf{A}^T = \boldsymbol{\Sigma}^2 \mathbf{U}^T$, the first equality following on multiplying (2.2) by \mathbf{A}^T , and the second on substituting the transpose of (2.3). Thanks to Ann Greenbaum for pointing this fact out.

Beltrami next states that by a well-known theorem, the roots of (2.5) are real. Moreover, they are positive. To show this he notes that

$$0 < \|\mathbf{x}^T \mathbf{A}\|^2 = \mathbf{x}^T (\mathbf{A} \mathbf{A}^T) \mathbf{x} = \boldsymbol{\xi}^T \boldsymbol{\Sigma}^2 \boldsymbol{\xi}, \quad (2.7)$$

the last equation following from the theory of quadratic forms. This inequality immediately implies that the σ_i^2 are positive.

There is some confusion here. Beltrami appears to be assuming the existence of the vector $\boldsymbol{\xi}$, whose very existence he is trying to establish. The vector required by his argument is an eigenvector of $\mathbf{A} \mathbf{A}^T$ corresponding to σ . The fact that the two vectors turn out to be the same apparently caused Beltrami to leap ahead of himself and use $\boldsymbol{\xi}$ in (2.7).

Beltrami is now ready to give an algorithm to determine the diagonalizing transformation.

1. Find the roots of the equation (2.5).
2. Determine \mathbf{U} from (2.4). Here Beltrami notes that the columns of \mathbf{U} are determined up to factors of ± 1 , which is true only if the σ_i are distinct. He also tacitly assumes that the resulting \mathbf{U} will be orthogonal, which also requires that the σ_i be distinct.
3. Determine \mathbf{V} from (2.2). This step requires that $\boldsymbol{\Sigma}$ be nonsingular.

Discussion. From the foregoing it is clear that Beltrami derived the singular value decomposition for a real, square, nonsingular matrix having distinct singular values. His derivation is the one given in most textbooks, but it lacks the extras needed to handle degeneracies. It may be that in omitting these extras Beltrami was simplifying things for his student audience, but a certain slackness in the exposition suggests that he had not thought the problem through.

3. Jordan [28,29, 1874]

Camille Jordan can fairly be called the codiscoverer of the singular value decomposition. Although he published his derivation a year after Beltrami, it is clear that the work is independent. In fact, the “Mémoire sur les formes bilinéaires” treats three problems, of which the the reduction of a bilinear form to a diagonal form by orthogonal substitutions is the simplest.²

²The other two are to reduce a form by the same substitution of both sets of variables and to reduce a pair of forms by two substitutions, one for each set of variables. Jordan notes that

The Derivation. Jordan starts with the form

$$P = \mathbf{x}^T \mathbf{A} \mathbf{y}$$

and seeks the maximum and minimum of P subject to

$$\|\mathbf{x}\|^2 = \|\mathbf{y}\|^2 = 1. \quad (3.1)$$

The maximum is determined by the equation

$$0 = dP = d\mathbf{x}^T \mathbf{A} \mathbf{y} + \mathbf{x}^T \mathbf{A} d\mathbf{y}, \quad (3.2)$$

which must be satisfied for all $d\mathbf{x}$ and $d\mathbf{y}$ that satisfy

$$d\mathbf{x}^T \mathbf{x} = 0 \quad \text{and} \quad d\mathbf{y}^T \mathbf{y} = 0. \quad (3.3)$$

3 Jordan then asserts that “equation (3.2) will therefore be a combination of the equations (3.3),” from which one obtains³

$$\mathbf{A} \mathbf{y} = \sigma \mathbf{x} \quad (3.4)$$

and

$$\mathbf{x}^T \mathbf{A} = \tau \mathbf{y}^T. \quad (3.5)$$

From (3.4) it follows that the maximum is

$$\mathbf{x}^T (\mathbf{A} \mathbf{y}) = \sigma \mathbf{x}^T \mathbf{x} = \sigma.$$

Similarly the maximum is also τ , so that $\sigma = \tau$.

Jordan now observes that σ is determined by the vanishing of the determinant

$$D = \begin{vmatrix} -\sigma \mathbf{I} & \mathbf{A} \\ \mathbf{A}^T & -\sigma \mathbf{I} \end{vmatrix}$$

of the system (3.4)–(3.5). He shows that this determinant contains only even powers of σ .

the former problem had been considered by Kronecker [31, 1866] in a different form, and the latter by Weierstrass [50, 1868].

³Jordan’s argument is not very clear. Possibly he means to say that for some constants σ and τ we must have $d\mathbf{x}^T \mathbf{A} \mathbf{y} + \mathbf{x}^T \mathbf{A} d\mathbf{y} = \sigma d\mathbf{x}^T \mathbf{x} + \tau d\mathbf{y}^T \mathbf{y}$, from which the subsequent equations follow from the independence of $d\mathbf{x}$ and $d\mathbf{y}$.

Now let σ_1 be a root of the equation $D = 0$, and let the equations (3.4) and (3.5) be satisfied by $\mathbf{x} = \mathbf{u}$ and $\mathbf{y} = \mathbf{v}$, where $\|\mathbf{u}\|^2 = \|\mathbf{v}\|^2 = 1$. (Jordan notes that one can find such a solution, even when it is not unique.) Let

$$\hat{\mathbf{U}} = (\mathbf{u} \ \mathbf{U}_*) \quad \text{and} \quad \hat{\mathbf{V}} = (\mathbf{v} \ \mathbf{V}_*)$$

be orthogonal, and let

$$\mathbf{x} = \hat{\mathbf{U}}\hat{\mathbf{x}} \quad \text{and} \quad \mathbf{y} = \hat{\mathbf{V}}\hat{\mathbf{y}}.$$

With these substitutions, let

$$P = \hat{\mathbf{x}}^T \hat{\mathbf{A}} \hat{\mathbf{y}}.$$

In this system, P attains its maximum⁴ for $\hat{\mathbf{x}} = \hat{\mathbf{y}} = \mathbf{e}_1$, where $\mathbf{e}_1 = (1, 0, \dots, 0)^T$. Moreover, at the maximum we have

$$\hat{\mathbf{A}}\hat{\mathbf{y}} = \sigma_1\hat{\mathbf{x}} \quad \text{and} \quad \hat{\mathbf{x}}^T \hat{\mathbf{A}} = \sigma_1\hat{\mathbf{y}}^T,$$

which implies that

$$\hat{\mathbf{A}} = \begin{pmatrix} \sigma & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_1 \end{pmatrix}.$$

Thus with $\xi_1 = \hat{x}_1$ and $\eta_1 = \hat{y}_1$, P assumes the form

$$\sigma_1 \xi_1 \eta_1 + P_1,$$

where P_1 is independent of ξ_1 and η_1 . Jordan now applies the reduction inductively to P_1 to arrive at the canonical form

$$P = \boldsymbol{\xi}^T \boldsymbol{\Sigma} \boldsymbol{\eta}.$$

Finally, Jordan notes that when the roots of the characteristic equation $D = 0$ are simple, the columns of \mathbf{U} and \mathbf{V} can be calculated directly from (3.1), (3.4), and (3.5).

Discussion. In this paper we see the sure hand of a skilled professional. Jordan proceeds from problem to solution with economy and elegance. His approach of using a partial solution of the problem to reduce it to one of smaller size—*deflation* is the modern term—avoids the degeneracies that complicate Beltrami's approach. Incidentally, the technique of deflation apparently lay fallow until Schur

⁴Jordan nods here, since he has not explicitly selected the largest root σ_1 .

[41, 1917] used it to establish his triangular form of a general matrix. It is now a widely used theoretical and algorithmic tool.

The matrix

$$\begin{pmatrix} \mathbf{0} & \mathbf{A} \\ \mathbf{A}^T & \mathbf{0} \end{pmatrix},$$

from which the determinant D was formed, is also widely used. Its present day popularity is due to Wielandt (see [14, p.113]) and Lanczos [32, 1958], who apparently rediscovered the decomposition independently.

Yet another consequence of Jordan's approach is the variational characterization of the largest singular value as the maximum of a function. This and related characterizations have played an important role in perturbation and localization theorems for singular values (for more see [43, §IV.4]).

4. Sylvester [44,46,45, 1889]

Sylvester wrote a footnote and two papers on the subject of the singular value decomposition. The footnote appears at the end of a paper in *The Messenger of Mathematics* [44] entitled "A New Proof That a General Quadric May Be Reduced to Its Canonical Form (That Is, a Linear Function of Squares) by Means of a Real Orthogonal Substitution." In the paper Sylvester describes an iterative algorithm for reducing a quadratic form to diagonal form. In the footnote he points out that an analogous iteration can be used to diagonalize a bilinear form and says that he has "sent for insertion in the C. R. of the Institute a Note in which I give the rule for effecting this reduction." The rule turns out to be Beltrami's algorithm. In a final *Messenger* paper [45], Sylvester presents both the iterative algorithm and the rule.

The Rule. Here we follow the *Messenger* paper. Sylvester begins with the bilinear form

$$B = \mathbf{x}^T \mathbf{A} \mathbf{y}$$

and considers the quadratic form

$$M = \sum_i \left(\frac{dB}{dy_i} \right)^2$$

(which is $\mathbf{x}^T \mathbf{A} \mathbf{A}^T \mathbf{x}$, a fact tacitly assumed by Sylvester). Let $M = \sum \lambda_i \xi_i^2$ be the canonical form of M . If B has the canonical form $B = \sum \sigma_i \xi_i \eta_i$, then $\sum [\sigma_i \xi_i]^2$ is orthogonally equivalent to $M = \sum \lambda_i \xi_i^2$, which implies that $\lambda_i = \sigma_i^2$ in some order.

To find the substitutions, Sylvester introduces the matrices $\mathbf{M} = \mathbf{A}\mathbf{A}^T$ and $\mathbf{N} = \mathbf{A}^T\mathbf{A}$ and asserts that the substitution for \mathbf{x} is the substitution that diagonalizes \mathbf{M} and substitution for \mathbf{y} is the one that diagonalizes \mathbf{N} . In general, this is true only if the singular values of \mathbf{A} are distinct.

In his *Comptes Rendu* note Sylvester gives the following rule for finding the coefficients of the \mathbf{x} -substitution corresponding to a singular value σ . Strike a row of the matrix $\mathbf{M} - \sigma^2\mathbf{I}$. Then the vector of coefficients is the vector of minors of order $n - 1$ of the reduced matrix normalized so that their sum of squares is one. Coefficients of the \mathbf{y} -substitution may be obtained analogously from $\mathbf{N} - \sigma\mathbf{I}$. This only works if the singular value σ is simple.

Infinitesimal iteration. Sylvester first proposed this method as a technique for showing that a quadratic form could be diagonalized, and he later extended it to bilinear forms. It is already intricate enough for quadratic forms, and we will confine ourselves to a sketch of that case.

Sylvester proceeds inductively, assuming that he can solve a problem of order $n - 1$. Thus for $n = 3$ he can assume the matrix is of the form

$$\mathbf{A} = \begin{pmatrix} a & 0 & f \\ 0 & b & g \\ f & g & c \end{pmatrix},$$

the zeros being introduced by the induction step. His problem is then to get rid of f and g without destroying the zeros previously introduced.

Sylvester proposes to make an “infinitesimal orthogonal substitution” of the form

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 1 & \epsilon & \eta \\ -\epsilon & 1 & \theta \\ -\eta & -\theta & 1 \end{pmatrix} \begin{pmatrix} \xi_1 \\ \xi_2 \\ \xi_3 \end{pmatrix},$$

where the off diagonal quantities are so small that powers higher than the first can be neglected. Then the the (2,1)- and (1,2)-elements of the transformed matrix are

$$(a - b)\epsilon - f\theta - g\eta, \tag{4.1}$$

while the change in $f^2 + g^2$ is given by

$$\frac{1}{2}\delta(f^2 + g^2) = (a - c)f\eta + (b - c)g\theta.$$

If either of $(a - c)f$ or $(b - c)g$ is nonzero, η and θ can be chosen to decrease $f^2 + g^2$. If $(a - b)$ is nonzero, ϵ may then be chosen so that (4.1) is zero; i.e., so that the zero previously introduced is preserved. Sylvester shows how special cases like $a = b$ can be handled by explicitly deflating the problem.

Sylvester now claims that an infinite sequence of these infinitesimal transformations will reduce one of f or g to zero, or will reduce the problem to one of the special cases.

Discussion. These are not easy papers to read. The style is opaque, and Sylvester pontificates without proving, leaving too many details to the reader. The mathematical reasoning harks back to an earlier, less rigorous era.

The fact that Sylvester sent a note to *Comptes Rendu*, the very organ where Jordan announced his results a decade and a half earlier, makes it clear that he was working in ignorance of his predecessors. It also suggests the importance he attached to his discovery, since a note in *Comptes Rendu* was tantamount to laying claim to a new result.

Sylvester was also working in ignorance of the iterative algorithm of Jacobi [25, 1846] for diagonalizing a quadratic form. The generalization of this algorithm to the singular value decomposition is due to Kogbetliantz [30].

It is not clear whether Sylvester intended to ignore second order terms in his iteration or whether he regards the diagonalization as being composed of an (uncountably) infinite number of infinitesimal transformation. Though the preponderance of his statements favor the latter, neither interpretation truly squares with everything he writes. In the first, small, but finite, terms replace the zeros previously introduces, so that a true diagonalization is not achieved. The second has the flavor of some recent algorithms in which discrete transformations are replaced by continuous transformations defined by differential equations (for applications of this approach to the singular value decomposition see [7,9]). But Sylvester does not give enough detail to write down such equations.

5. Schmidt [39, 1907]

Our story now moves from the domain of linear algebra to integral equations, one of the hot topics of the first decades of our century. In his treatment of integral equations with unsymmetric kernels, Erhard Schmidt (of Gram–Schmidt fame) introduced the infinite dimensional analogue of the singular value decomposition. But he went beyond the mere existence of the decomposition by showing how it can be used to obtain optimal, low-rank approximations to an operator. In doing

so he transformed the singular value decomposition from a mathematical curiosity to an important theoretical and computational tool.

Symmetric Kernels. Schmidt's approach is essentially the same as Beltrami's; however, because he worked in infinite dimensional spaces of functions he could not appeal to previous results on quadratic forms. Consequently, the first part of his paper is devoted to symmetric kernels.

Schmidt begins with a kernel $A(s, t)$ that is continuous and symmetric on $[a, b] \times [a, b]$. A continuous, nonvanishing function $\varphi(s)$ satisfying

$$\varphi(s) = \lambda \int_a^b A(s, t) \varphi(t) dt$$

is said to be an eigenfunction of A corresponding to the eigenvalue λ . Note that Schmidt's eigenvalues are the reciprocals of ours.

Schmidt then establishes the following facts.

1. The kernel A has at least one eigenfunction.
2. The eigenvalues and their eigenfunctions are real.
3. Each eigenvalue of A has at most a finite number of linearly independent eigenfunctions.
4. The kernel A has a complete, orthonormal system of eigenfunctions; that is, a sequence $\varphi_1(s), \varphi_2(s), \dots$ of orthonormal eigenfunctions such that every eigenfunction can be expressed as a linear combination of a finite number of the $\varphi_j(s)$.⁵
5. The eigenvalues satisfy

$$\int_a^b \int_a^b (A(s, t))^2 ds dt \geq \sum_i \frac{1}{\lambda_i^2},$$

which implies that the sequence of eigenvalues is unbounded.

Unsymmetric Kernels. Schmidt now allows $A(s, t)$ to be unsymmetric and calls any nonzero pair $u(s)$ and $v(s)$ satisfying

$$u(s) = \lambda \int_a^b A(s, t) v(t) dt$$

⁵This usage of the word "complete" is at variance with today's usage, in which a sequence is complete if its finite linear combinations are dense.

and

$$v(t) = \lambda \int_a^b A(s, t) u(s) ds$$

a pair of adjoint eigenfunctions corresponding to the eigenvalue λ .⁶ He then introduces the symmetric kernels

$$\bar{A}(s, t) = \int_a^b A(s, r) A(t, r) dr$$

and

$$\underline{A}(s, t) = \int_a^b A(r, s) A(r, t) dr$$

and shows that if $u_1(s), u_2(s), \dots$ is a complete orthonormal system for $\bar{A}(s, t)$ corresponding to the eigenvalues $\lambda_1^2, \lambda_2^2, \dots$ then the sequence defined by

$$v_i(t) = \lambda_i \int_a^b A(s, t) u(s) ds, \quad i = 1, 2, \dots$$

is a complete orthonormal system for $\underline{A}(s, t)$. Moreover, for $i = 1, 2, \dots$ the functions $u_i(s)$ and $v_i(s)$ form an adjoint pair for $A(s, t)$.

Schmidt then goes on to consider the expansion of functions in series of eigenfunctions. Specifically, if

$$g(s) = \int_a^b A(s, t) h(t) dt,$$

then

$$g(s) = \sum_i \frac{u_i(s)}{\lambda_i} \int_a^b h(t) v_i(t) dt,$$

and the convergence is absolute and uniform. Finally, he shows that if g and h are continuous then

$$\int_a^b \int_a^b A(s, t) g(s) h(t) ds dt = \sum_i \frac{1}{\lambda_i} \int_a^b g(s) u_i(s) ds \int_a^b h(t) v_i(t) dt, \quad (5.1)$$

an expression which Schmidt says “corresponds to the canonical decomposition of a bilinear form.”

⁶Again the usage differs from ours, but now in two ways. We work with the reciprocal of λ , calling it a singular value, and we distinguish between the singular values of a matrix and its eigenvalues.

The Approximation Theorem. Up to now, our exposition has been cast in the language of integral equations, principally to keep issues of analysis in the foreground. These issues are not as important in what follows, and we will therefore return to matrix notation, taking care, as always, to follow Schmidt's development closely.

The problem Schmidt sets out to solve is that of finding the best approximation to \mathbf{A} of the form

$$\mathbf{A} \cong \sum_{i=1}^k \mathbf{x}_i \mathbf{y}_i^T$$

in the sense that

$$\|\mathbf{A} - \sum_{i=1}^k \mathbf{x}_i \mathbf{y}_i^T\| = \min.$$

In other words, he is looking for the best approximation of rank not greater than k .

Schmidt begins by noting that if

$$\mathbf{A}_k = \sum_{i=1}^k \sigma_i \mathbf{u}_i \mathbf{v}_i^T, \quad (5.2)$$

then

$$\|\mathbf{A} - \mathbf{A}_k\|^2 = \|\mathbf{A}\|^2 - \sum_{i=1}^k \sigma_i^2.$$

Consequently, if it can be shown that for arbitrary \mathbf{x}_i and \mathbf{y}_i

$$\|\mathbf{A} - \sum_{i=1}^k \mathbf{x}_i \mathbf{y}_i^T\| \geq \|\mathbf{A}\|^2 - \sum_{i=1}^k \sigma_i^2, \quad (5.3)$$

then \mathbf{A}_k will be the desired approximation.

Without loss of generality we may assume that the vectors $\mathbf{x}_1, \dots, \mathbf{x}_k$ are orthonormal. For if they are not, we can use Gram-Schmidt orthogonalization to express them as linear combinations of orthonormal vectors, substitute these expressions in $\sum_{i=1}^k \mathbf{x}_i \mathbf{y}_i^T$, and collect terms in the new vectors.

Now

$$\begin{aligned} \|\mathbf{A} - \sum_{i=1}^k \mathbf{x}_i \mathbf{y}_i^T\| &= \text{trace} \left(\left(\mathbf{A} - \sum_{i=1}^k \mathbf{x}_i \mathbf{y}_i^T \right)^T \left(\mathbf{A} - \sum_{i=1}^k \mathbf{x}_i \mathbf{y}_i^T \right) \right) \\ &= \text{trace} \left(\mathbf{A}^T \mathbf{A} + \sum_{i=1}^k (\mathbf{y}_i - \mathbf{A}^T \mathbf{x}_i)(\mathbf{y}_i - \mathbf{A}^T \mathbf{x}_i)^T - \sum_{i=1}^k \mathbf{A}^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{A} \right) \end{aligned}$$

Since $\text{trace}((\mathbf{y}_i - \mathbf{A}^T \mathbf{x}_i)(\mathbf{y}_i - \mathbf{A}^T \mathbf{x}_i)^T) \geq 0$ and $\text{trace}(\mathbf{A} \mathbf{x}_i \mathbf{x}_i^T \mathbf{A}^T) = \|\mathbf{A} \mathbf{x}_i\|^2$, the result will be established if it can be shown that

$$\sum_{i=1}^k \|\mathbf{A} \mathbf{x}_i\|^2 \leq \sum_{i=1}^k \sigma_i^2.$$

Let $\mathbf{V} = (\mathbf{V}_1 \ \mathbf{V}_2)$, where \mathbf{V}_1 has k columns, and let $\mathbf{\Sigma} = \text{diag}(\mathbf{\Sigma}_1, \mathbf{\Sigma}_2)$ be a conformal partition of $\mathbf{\Sigma}$. Then

$$\begin{aligned} \|\mathbf{A} \mathbf{x}_i\|^2 &= \sigma_k^2 + \left(\|\mathbf{\Sigma}_1 \mathbf{V}_1^T \mathbf{x}_i\|^2 - \sigma_k^2 \|\mathbf{V}_1^T \mathbf{x}_i\|^2 \right) \\ &\quad - \left(\sigma_k^2 \|\mathbf{V}_2^T \mathbf{x}_i\|^2 - \|\mathbf{\Sigma}_2 \mathbf{V}_2^T \mathbf{x}_i\|^2 \right) \\ &\quad - \sigma_k^2 (1 - \|\mathbf{V}^T \mathbf{x}_i\|) \end{aligned} \tag{5.4}$$

Now the last two terms in (5.4) are clearly nonnegative. Hence

$$\begin{aligned} \sum_{i=1}^k \|\mathbf{A} \mathbf{x}_i\|^2 &\leq k\sigma_k^2 + \sum_{i=1}^k \left(\|\mathbf{\Sigma}_1 \mathbf{V}_1^T \mathbf{x}_i\|^2 - \sigma_k^2 \|\mathbf{V}_1^T \mathbf{x}_i\|^2 \right) \\ &= k\sigma_k^2 + \sum_{i=1}^k \sum_{j=1}^k (\sigma_j^2 - \sigma_k^2) |\mathbf{v}_j^T \mathbf{x}_i|^2 \\ &= \sum_{j=1}^k \left(\sigma_k^2 + (\sigma_j^2 - \sigma_k^2) \sum_{i=1}^k |\mathbf{v}_j^T \mathbf{x}_i|^2 \right) \\ &\leq \sum_{j=1}^k (\sigma_k^2 + (\sigma_j^2 - \sigma_k^2)) \\ &= \sum_{j=1}^k \sigma_j^2, \end{aligned}$$

which establishes the result.

Discussion. Schmidt's two contributions to the singular value decomposition are its generalization to function spaces and his approximation theorem. Although Schmidt did not refer to earlier work on the decomposition in finite dimensional spaces, the quote following (5.1) suggests that he knew of its existence. Nonetheless, his contribution here is substantial, especially since he had to deal with many of the problems of functional analysis without modern tools.

An important difference in Schmidt's version of the decomposition is the treatment of null-vectors of \mathbf{A} . In his predecessors' treatments they are part of the substitution that reduces the bilinear form $\mathbf{x}^T \mathbf{A} \mathbf{y}$ to its canonical form. For Schmidt

they are not part of the decomposition. The effect of this can be seen in the third term of (5.4), which in the usual approach is zero but in Schmidt's approach can be nonzero.

The crowning glory of Schmidt's work is his approximation theorem, which is nontrivial to conjecture and hard to prove from scratch. Schmidt's proof is certainly not pretty — we will examine the more elegant approach of Weyl in the next section — but it does establish what can properly be termed the fundamental theorem of the singular value decomposition.

6. Weyl [51, 1912]

An important application of the approximation theorem is the determination of the rank of a matrix in the presence of error. If \mathbf{A} is of rank k and $\tilde{\mathbf{A}} = \mathbf{A} + \mathbf{E}$, then the last $n - k$ singular values of $\tilde{\mathbf{A}}$ satisfy

$$\tilde{\sigma}_{k+1}^2 + \cdots + \tilde{\sigma}_n^2 \leq \|\mathbf{E}\|^2, \quad (6.1)$$

so that the defect in rank of \mathbf{A} will be manifest in the size of its trailing singular values.

The inequality (6.1) is actually a perturbation theorem for the zero singular values of a matrix. Weyl's contribution to the theory of the singular value decomposition was to develop a general perturbation theory and use it to give an elegant proof of the approximation theorem. Although Weyl treated integral equations with symmetric kernels, in a footnote on Schmidt's contribution he states, "E. Schmidt's theorem, by the way, treats arbitrary (unsymmetric) kernels; however, our proof can also be applied directly to this more general case." Since here we are concerned with the more general case, we will paraphrase Weyl's development as he might have written it for unsymmetric matrices.

The Location of Singular Values. The heart of Weyl's development is a lemma concerning the singular values of a perturbed matrix. Specifically, if $\mathbf{B}_k = \mathbf{X}\mathbf{Y}^T$, where \mathbf{X} and \mathbf{Y} have k columns (i.e., $\text{rank}(\mathbf{B}_k) \leq k$), then

$$\sigma_1(\mathbf{A} - \mathbf{B}_k) \geq \sigma_{k+1}(\mathbf{A}), \quad (6.2)$$

where $\sigma_i(\cdot)$ denotes the i th singular value of its argument.

The proof is simple. Since \mathbf{Y} has k columns, there is a linear combination

$$\mathbf{v} = \gamma_1 \mathbf{v}_1 + \gamma_2 \mathbf{v}_2 + \cdots + \gamma_{k+1} \mathbf{v}_{k+1}$$

of the first $k + 1$ columns of \mathbf{V} (from the singular value decomposition of \mathbf{A}) such that $\mathbf{Y}^T \mathbf{v} = 0$. Without loss of generality we may assume that $\|\mathbf{v}\| = 1$, or equivalently that $\gamma_1^2 + \cdots + \gamma_{k+1}^2 = 1$. It follows that

$$\begin{aligned} \sigma_1^2(\mathbf{A} - \mathbf{B}) &\geq \mathbf{v}^T(\mathbf{A} - \mathbf{B})^T(\mathbf{A} - \mathbf{B})\mathbf{v} \\ &= \mathbf{v}^T(\mathbf{A}^T\mathbf{A})\mathbf{v} \\ &= \gamma_1^2\sigma_1^2 + \gamma_2^2\sigma_2^2 + \cdots + \gamma_{k+1}^2\sigma_{k+1}^2 \\ &\geq \sigma_{k+1}^2. \end{aligned}$$

Weyl then proves two theorems. The first states that if $\mathbf{A} = \mathbf{A}' + \mathbf{A}''$ then

$$\sigma_{i+j-1} \leq \sigma'_i + \sigma''_j, \quad (6.3)$$

where the σ'_i and σ''_i are the singular values of \mathbf{A}' and \mathbf{A}'' arranged in descending order of magnitude. Weyl begins by establishing (6.3) for $i = j = 1$:

$$\sigma_1 = \mathbf{u}_1^T \mathbf{A} \mathbf{v}_1 = \mathbf{u}_1^T \mathbf{A}' \mathbf{v}_1 + \mathbf{u}_1^T \mathbf{A}'' \mathbf{v}_1 \leq \sigma'_1 + \sigma''_1.$$

To establish the result in general, let \mathbf{A}'_{i-1} and \mathbf{A}''_{j-1} be formed in analogy with (5.2). Then $\sigma_1(\mathbf{A}' - \mathbf{A}'_{i-1}) = \sigma_i(\mathbf{A}')$ and $\sigma_1(\mathbf{A}'' - \mathbf{A}''_{j-1}) = \sigma_j(\mathbf{A}'')$. Moreover $\text{rank}(\mathbf{A}'_{i-1} + \mathbf{A}''_{j-1}) \leq i + j - 2$. From these facts and from (6.2) it follows that

$$\begin{aligned} \sigma'_i + \sigma''_j &= \sigma_1(\mathbf{A}' - \mathbf{A}'_{i-1}) + \sigma_1(\mathbf{A}'' - \mathbf{A}''_{j-1}) \\ &\geq \sigma_1(\mathbf{A} - \mathbf{A}'_{i-1} - \mathbf{A}''_{j-1}) \\ &\geq \sigma_{i+j-1}, \end{aligned}$$

which proves the theorem.

The second theorem is really a corollary of the first. Set $\mathbf{A}' = \mathbf{A} - \mathbf{B}_k$ and $\mathbf{A}'' = \mathbf{B}_k$, where, as above, \mathbf{B}_k has rank k . Since $\sigma_{k+1}(\mathbf{B}_k) = 0$, we have on setting $j = k + 1$ in (6.3)

$$\sigma_i(\mathbf{A} - \mathbf{B}_k) \geq \sigma_{k+i}, \quad i = 1, 2, \dots$$

As a corollary to this result we obtain

$$\|\mathbf{A} - \mathbf{B}_k\|^2 \geq \sigma_{k+1}^2 + \cdots + \sigma_n^2.$$

This inequality is equivalent to (5.3) and thus establishes the approximation theorem.

Discussion. Weyl did not actually write down the development for unsymmetric kernels, and we remind the reader once again of the advisability of consulting original sources. In particular, since symmetric kernels can have negative eigenvalues

as well as positive ones, Weyl wrote down three sequences of inequalities: one for positive eigenvalues, one for negative, and one — corresponding to the inequalities presented here — for the absolute values of the eigenvalues.

Returning to the perturbation problem that opened this section, if in (6.3) we make the identification $\mathbf{A} \leftarrow \tilde{\mathbf{A}}$, $\mathbf{A}' \leftarrow \mathbf{A}$, and $\mathbf{A}'' \leftarrow \mathbf{E}$, then with $j = 1$ we get

$$\tilde{\sigma}_i \leq \sigma_i + \|\mathbf{E}\|_2,$$

where $\|E\|_2 = \sigma_1(E)$. On the other hand, if we make the identifications $\mathbf{A}' \leftarrow \tilde{\mathbf{A}}$ and $\mathbf{A}'' \leftarrow -\mathbf{E}$, then we get

$$\tilde{\sigma}_i \leq \sigma_i - \|\mathbf{E}\|_2.$$

It follows that

$$|\tilde{\sigma}_i - \sigma_i| \leq \|\mathbf{E}\|_2, \quad i = 1, 2, \dots, n.$$

The number $\|\mathbf{E}\|_2$ is called the spectral norm of \mathbf{E} . Thus Weyl's result implies that if the singular values of \mathbf{A} and $\tilde{\mathbf{A}}$ are associated in their natural order, they cannot differ by more than the spectral norm of the perturbation.

7. Envoi

With Weyl's contribution, the theory of the singular value decomposition can be said to have matured. The subsequent history is one of extensions, new discoveries, and applications. What follows is a brief sketch of these developments yet to come.

Extensions. Autonne [2, 1913] extended the decomposition to complex matrices. Eckart and Young [12, 1936], [13, 1939] extended it to rectangular matrices and rediscovered Schmidt's approximation theorem, which is often (and incorrectly) called the Eckart–Young theorem.

Nomenclature.⁷ The term “singular value” seems to have come from the literature on integral equations. A little after the appearance of Schmidt's paper, Bateman [4, 1908] refers to numbers that are essentially the reciprocals of the eigenvalues of the kernel as singular values. Picard [37, 1910] notes that for symmetric kernels Schmidt's eigenvalues are real and in this case (but not in general) he calls them singular values. By 1937, Smithes was referring to singular values of an integral equation in our modern sense of the word. Even at this point, usage had not stabilized. In 1949, Weyl [52] speaks of the “two kinds of eigenvalues of a linear transformation,” and in a 1969 translation of a 1965 Russian treatise on

⁷Parts of this passage were taken from [43, p. 35]

nonselfadjoint operators Gohberg and Krein [16] refer to the “s-numbers” of an operator. For the term “principal component,” see below.

Related Decompositions. Beltrami’s proof of the existence of the singular decomposition shows that it is closely related to the spectral decompositions of $\mathbf{A}^T\mathbf{A}$ and $\mathbf{A}\mathbf{A}^T$. It can also be used to derive the polar decomposition of Autonne [1, 1902], [3, 1915], in which a matrix is factored into the product of a Hermitian matrix and a unitary matrix.

In his investigation of the geometry of n -space, Jordan [27, 1875] introduced canonical bases for pairs of subspaces. This line of development lead to the CS (cosine-sine) decomposition of a partitioned orthogonal matrix introduced implicitly by Davis and Kahan [8, 1970], and explicitly by [42, 1977]. The CS decomposition can in turn be used to derive the generalized singular value decomposition of a matrix, either in the original form introduced by Van Loan [47, 1975] or in the revised version of Paige and Saunders [35, 1981].

Although it is not, strictly speaking, a matrix decomposition, the Moore-Penrose pseudo-inverse [34, 1920], [36, 1955] can be calculated from the singular value decomposition of a matrix as follows. Suppose that the first k singular values of \mathbf{A} are nonzero while the last $n - k$ are zero, and set $\Sigma^\dagger = \text{diag}(\sigma_1^{-1}, \dots, \sigma_k^{-1}, 0, \dots, 0)$. Then the pseudo inverse of \mathbf{A} is given by $\mathbf{A}^\dagger = \mathbf{U}\Sigma^\dagger\mathbf{V}^T$.

Unitarily Invariant Norms. A matrix norm $\|\cdot\|_U$ is unitarily invariant if $\|\mathbf{U}^H\mathbf{A}\mathbf{V}\|_U = \|\mathbf{A}\|_U$ for all unitary matrices U and V . A vector norm $\|\cdot\|_g$ is a symmetric gauge function if $\|\mathbf{P}\mathbf{x}\|_g = \|\mathbf{x}\|_g$ for any permutation matrix and $\|\|\mathbf{x}\|\|_g = \|\mathbf{x}\|_g$. Von Neumann [49, 1937] showed that to any unitarily invariant norm $\|\cdot\|_U$ there corresponds a symmetric gauge function $\|\cdot\|_g$ such that $\|\mathbf{A}\|_U = \|(\sigma_1, \dots, \sigma_n)^T\|_g$; i.e., a unitarily invariant norm is a symmetric gauge function of the singular values of its argument.

Approximation Theorems. Schmidt’s approximation theorem has been generalized in a number of directions. Mirsky [33, 1960] showed that \mathbf{A}_k of (5.2) is a minimizing matrix in any unitarily invariant norm. The case where further restrictions are imposed on the minimizing matrix are treated in [10, 17, 38].

Given matrices \mathbf{A} and \mathbf{B} , The Procrustes problem, which arises in the statistical method of factor analysis, is that of determining a unitary matrix \mathbf{Q} such that $\|\mathbf{A} - \mathbf{B}\mathbf{Q}\|$ is minimized (for the name see [24, 1962]). Green [20, 1952] and Schöneman [40, 1966] showed that if $\mathbf{U}^T\mathbf{A}^T\mathbf{B}\mathbf{V} = \Sigma$ is the singular value decomposition of $\mathbf{A}^T\mathbf{B}$, then the minimizing matrix is $\mathbf{Q} = \mathbf{V}\mathbf{U}^T$. Rao [38, 1980]

considers the more general problem of minimizing $\|\mathbf{PA} - \mathbf{BQ}\|$, where \mathbf{P} and \mathbf{Q} are orthogonal.

Principal Components. An alternative to factor analysis is the principal component analysis of Hotelling [22, 1933]. Specifically, if \mathbf{x}^T is a multivariate random variable with mean zero and common dispersion matrix \mathbf{D} , and $\mathbf{D} = \mathbf{V}\mathbf{\Sigma}\mathbf{V}^T$ is the eigenvalue-eigenvector decomposition of \mathbf{D} , then the components of $\mathbf{x}^T\mathbf{V}$ are uncorrelated with variances σ_i . Hotelling called the transformed variables “the principal components of variance” of \mathbf{x}^T . If the rows of \mathbf{X} consist of independent samples of \mathbf{x}^T , then the expectation of $\mathbf{X}^T\mathbf{X}$ is proportional to $\mathbf{\Sigma}$. It follows that the matrix $\hat{\mathbf{V}}$ obtained from the singular value decomposition of \mathbf{X} is an estimate \mathbf{V} .

Hotelling [23, 1936] also introduced canonical correlations between two sets of random variables that bears the same relation to the generalized singular value decomposition as his principal components bear to the singular value decomposition.

Inequalities Involving Singular Values. Just as Schmidt did not have the last word on approximation theorems, Weyl was not the last to work on inequalities involving singular values. The subject is too voluminous to treat here, and we refer the reader to the excellent survey with references in [21, Ch. 3]. However, mention should be made of a line of research initiated by Weyl [52, 1949] relating the singular values and eigenvalues of a matrix.

Computational Methods The singular value decomposition was introduced into numerical analysis by Golub and Kahan [18, 1965], who proposed a computational algorithm. However, it was Golub [19, 1970] who gave the algorithm that has been the workhorse of the past two decades. Recently, Demmel and Kahan [11, 1990] have proposed an interesting alternative.

8. Acknowledgment

I would like to thank Nick Higham and Hongyuan Zha for reading and commenting on the manuscript.

References

- [1] L. Autonne. Sur les groupes linéaires, réels et orthogonaux. *Bulletin de la Société Mathématique de France*, 30:121–134, 1902.

-
- [2] L. Autonne. Sur les matrices hypohermitiennes et les unitaires. *Comptes Rendus de l'Academie des Sciences, Paris*, 156:858–860, 1913.
 - [3] L. Autonne. Sur les matrices hypohermitiennes et sur les matrices unitaires. *Annales De L'Université de Lyons, Nouvelle Série I*, 38:1–77, 1915.
 - [4] H. Bateman. A formula for the solving function of a certain integral equation of the second kind. *Transactions of the Cambridge Philosophical Society*, 20:179–187, 1908.
 - [5] E. Beltrami. Sulle funzioni bilineari. *Giornale di Matematiche ad Uso degli Studenti Delle Universita*, 11:98–106, 1873. An English translation by D. Boley is available as University of Minnesota, Department of Computer Science, Technical Report 90–37, 1990.
 - [6] A. L. Cauchy. Sur l'équation à l'aide de laquelle on détermine les inégalités séculaires des mouvements des planètes. In *Oeuvres Complètes (II^e Série)*, 1829.
 - [7] M. Chu. A differential equation approach to the singular value decomposition of bidiagonal matrices. *Linear Algebra and Its Applications*, 80:71–79, 1986.
 - [8] C. Davis and W. Kahan. The rotation of eigenvectors by a perturbation. III. *SIAM Journal on Numerical Analysis*, 7:1–46, 1970.
 - [9] P. Deift, J. Demmel, L.-C. Li, and C. Tomei. The bidiagonal singular value decomposition and Hamiltonian mechanics. *SIAM Journal on Numerical Analysis*, 28:1463–1516, 1991.
 - [10] J. Demmel. The smallest perturbation of a submatrix which lowers the rank and constrained total least squares problems. *SIAM Journal on Numerical Analysis*, 24:199–206, 1987.
 - [11] J. Demmel and W. Kahan. Accurate singular values of bidiagonal matrices. *SIAM Journal on Scientific and Statistical Computing*, 11:873–912, 1989.
 - [12] C. Eckart and G. Young. The approximation of one matrix by another of lower rank. *Psychometrika*, 1:211–218, 1936.
 - [13] C. Eckart and G. Young. A principal axis transformation for non-Hermitian matrices. *Bulletin of the American Mathematical Society*, 45:118–121, 1939.

-
- [14] K. Fan and A. J. Hoffman. Some metric inequalities in the space of matrices. *Proceedings of the American Mathematical Society*, 6:111–116, 1955.
 - [15] C. F. Gauss. Theoria combinationis observationum erroribus minimis obnoxiae, pars posterior. In *Werke*, IV, pages 27–53, Königlichen Gesellschaft der Wissenschaften zu Göttingen (1880), 1823.
 - [16] I. C. Gohberg and M. G. Krein. *Introduction to the Theory of Linear Non-selfadjoint Operators*. American Mathematical Society, Providence, Rhode Island, 1969.
 - [17] G. H. Golub, A. Hoffman, and G. W. Stewart. A generalization of the Eckart-Young matrix approximation theorem. *Linear Algebra and Its Applications*, 88/89:317–327, 1987.
 - [18] G. H. Golub and W. Kahan. Calculating the singular values and pseudo-inverse of a matrix. *SIAM Journal on Numerical Analysis*, 2:205–224, 1965.
 - [19] G. H. Golub and C. Reinsch. Singular value decomposition and least squares solution. *Numerische Mathematik*, 14:403–420, 1970. Also in [53, pp.134–151].
 - [20] B. F. Green. The orthogonal approximation of the oblique structure in factor analysis. *Psychometrika*, 17:429–440, 1952.
 - [21] R. A. Horn and C. R. Johnson. *Topics in Matrix Analysis*. Cambridge University Press, Cambridge, 1991.
 - [22] H. Hotelling. Analysis of a complex of statistical variables into principal components. *Journal of Educational Psychology*, 24:417–441 and 498–520, 1933.
 - [23] H. Hotelling. Relation between two sets of variates. *Biometrika*, 28:322–377, 1936.
 - [24] J. R. Hurley and R. B. Cuttall. The procrustes program: direct rotation to test a hypothesized factor structure. *Behavioral Science*, 7:258–262, 1962.
 - [25] C. G. J. Jacobi. Über ein leichtes Verfahren die in der Theorie der Säculärstörungen vorkommenden Gleichungen numerisch aufzulösen. *Journal für die reine und angewandte Mathematik*, 30:51–s94, 1846.

- [26] C. G. J. Jacobi. Über eine elementare Transformation eines in Bezug jedes von zwei Variablen-Systemen linearen und homogenen Ausdrucks. *Journal für die reine und angewandte Mathematik*, 53:265–270, 1857, posthumous.
- [27] C. Jordan. Essai sur la géométrie à n dimensions. *Bulletin de la Société Mathématique*, 3:103–174, 1875.
- [28] C. Jordan. Mémoire sur les formes bilinéaires. *Journal de Mathématiques Pures et Appliquées, Deuxième Série*, 19:35–54, 1874.
- [29] C. Jordan. Sur la réduction des formes bilinéaires. *Comptes Rendus de l'Académie des Sciences, Paris*, 78:614–617, 1874.
- [30] E. G. Kogbetliantz. Solution of linear systems by diagonalization of coefficients matrix. *Quarterly of Applied Mathematics*, 13:123–132, 1955.
- [31] L. Kronecker. Über bilineare Formen. *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin*, 597–613, 1866.
- [32] C. Lanczos. Linear systems in self-adjoint form. *American Mathematical Monthly*, 65:665–679, 1948.
- [33] L. Mirsky. Symmetric gauge functions and unitarily invariant norms. *Quarterly Journal of Mathematics*, 11:50–59, 1960.
- [34] E. H. Moore. On the reciprocal of the general algebraic matrix. *Bulletin of the American Mathematical Society*, 26:394–395, 1920. Abstract.
- [35] C. C. Paige and M. A. Saunders. Toward a generalized singular value decomposition. *SIAM Journal on Numerical Analysis*, 18:398–405, 1981.
- [36] R. Penrose. A generalized inverse for matrices. *Proceedings of the Cambridge Philosophical Society*, 51:406–413, 1955.
- [37] É. Picard. Sur un théorème général relatif aux équations intégrales de première espèce et sur quelques problèmes de physique mathématique. *Rendiconti del Circolo Matematico di Palermo*, 25:79–97, 1910.
- [38] C. R. Rao. Matrix approximations and reduction of dimensionality in multivariate statistical analysis. In P. R. Krishnaiah, editor, *Multivariate Analysis-V*, North-Holland, Amsterdam, 1980.

- [39] E. Schmidt. Zur Theorie der linearen und nichtlinearen Integralgleichungen. I Teil. Entwicklung willkürlichen Funktionen nach System vorgeschriebener. *Mathematische Annalen*, 63:433–476, 1907.
- [40] P. H. Schöneman. A generalized solution of the orthogonal procrustes problem. *Psychometrika*, 31:1–10, 1966.
- [41] J. Schur. Über Potenzreihen, die im Innern des Einheitskreise beschränkt sind. *Journal für die reine und angewandte Mathematik*, 147:205–232, 1917.
- [42] G. W. Stewart. On the perturbation of pseudo-inverses, projections, and linear least squares problems. *SIAM Review*, 19:634–662, 1977.
- [43] G. W. Stewart and J.-G. Sun. *Matrix Perturbation Theory*. Academic Press, Boston, 1990.
- [44] J. J. Sylvester. A new proof that a general quadric may be reduced to its canonical form (that is, a linear function of squares) by means of a real orthogonal substitution. *Messenger of Mathematics*, 19:1–5, 1889.
- [45] J. J. Sylvester. On the reduction of a bilinear quantic of the n^{TH} order to the form of a sum of n products by a double orthogonal substitution. *Messenger of Mathematics*, 19:42–46, 1889.
- [46] J. J. Sylvester. Sur la réduction biorthogonale d’une forme linéo-linéaire à sa forme canonique. *Comptes Rendus de l’Academie des Sciences, Paris*, 108:651–653, 1889.
- [47] C. F. Van Loan. A general matrix eigenvalue algorithm. *SIAM Journal on Numerical Analysis*, 12:819–834, 1975.
- [48] J. von Neumann. *Collected Works*, (A. H. Taub Editor). Pergamon, New York, 1962.
- [49] J. von Neumann. Some matrix-inequalities and metrization of matrix-space. *Tomsk. Univ. Rev.*, 1:286–300, 1937. In [48, v.4, pp.205–219].
- [50] K. Weierstrass. Zur Theorie der bilinearen und quadratischen Formen. *Monatshefte Akademie Wissenschaften Berlin*, 310–38, 1868.
- [51] H. Weyl. Das asymptotische Verteilungsgesetz der Eigenwert linearer partieller Differentialgleichungen (mit einer Anwendung auf der Theorie der Hohlraumstrahlung). *Mathematische Annalen*, 71:441–479, 1912.

- [52] H. Weyl. Inequalities between the two kinds of eigenvalues of a linear transformation. *Proceedings of the National Academy of Sciences*, 35:408–411, 1949.
- [53] J. H. Wilkinson and C. Reinsch. *Handbook for Automatic Computation. Vol. II Linear Algebra*. Springer, New York, 1971.

#	Author/s	Title
866	Gui-Qiang Chen and Tai-Ping Liu,	Zero relaxation and dissipation limits for hyperbolic conservation laws
867	Gui-Qiang Chen and Jian-Guo Liu,	Convergence of second-order schemes for isentropic gas dynamics
868	Aleksander M. Simon and Zbigniew J. Grzywna,	On the Larché-Cahn theory for stress-induced diffusion
869	Jerzy Luczka, Adam Gadowski and Zbigniew J. Grzywna,	Growth driven by diffusion
870	Mitchell Luskin and Tsorng-Whay Pan,	Nonplanar shear flows for nonaligning nematic liquid crystals
871	Mahmoud Affouf,	Unique global solutions of initial-boundary value problems for thermodynamic phase transitions
872	Richard A. Brualdi and Keith L. Chavey,	Rectangular L -matrices
873	Xinfu Chen, Avner Friedman and Bei Hu,	The thermistor problem with zero-one conductivity II
874	Raoul LePage,	Controlling a diffusion toward a large goal and the Kelly principle
875	Raoul LePage,	Controlling for optimum growth with time dependent returns
876	Marc Hallin and Madan L. Puri,	Rank tests for time series analysis a survey
877	V.A. Solonnikov,	Solvability of an evolution problem of thermocapillary convection in an infinite time interval
878	Horia I. Ene and Bogdan Vernescu,	Viscosity dependent behaviour of viscoelastic porous media
879	Kaushik Bhattacharya,	Self-accommodation in martensite
880	D. Lewis, T. Ratiu, J.C. Simo and J.E. Marsden,	The heavy top: a geometric treatment
881	Leonid V. Kalachev,	Some applications of asymptotic methods in semiconductor device modeling
882	David C. Dobson,	Phase reconstruction via nonlinear least-squares
883	Patricio Aviles and Yoshikazu Giga,	Minimal currents, geodesics and relaxation of variational integrals on mappings of bounded variation
884	Patricio Aviles and Yoshikazu Giga,	Partial regularity of least gradient mappings
885	Charles R. Johnson and Michael Lundquist,	Operator matrices with chordal inverse patterns
886	B.J. Bayly,	Infinitely conducting dynamos and other horrible eigenproblems
887	Charles M. Elliott and Stefan Luckhaus,	'A generalised diffusion equation for phase separation of a multi-component mixture with interfacial free energy'
888	Christian Schmeiser and Andreas Unterreiter,	The derivation of analytic device models by asymptotic methods
889	LeRoy B. Beasley and Norman J. Pullman,	Linear operators that strongly preserve the index of imprimitivity
890	Jerry Donato,	The Boltzmann equation with lie and cartan
891	Thomas R. Hoffend Jr., Peter Smereka and Roger J. Anderson,	A method for resolving the laser induced local heating of moving magneto-optical recording media
892	E.G. Kalnins, Willard Miller, Jr. and Sanchita Mukherjee,	Models of q -algebra representations: the group of plane motions
893	T.R. Hoffend Jr. and R.K. Kaul,	Relativistic theory of superpotentials for a nonhomogeneous, spatially isotropic medium
894	Reinhold von Schwerin,	Two metal deposition on a microdisk electrode
895	Vladimir I. Oliker and Nina N. Uraltseva,	Evolution of nonparametric surfaces with speed depending on curvature, III. Some remarks on mean curvature and anisotropic flows
896	Wayne Barrett, Charles R. Johnson, Raphael Loewy and Tamir Shalom,	Rank incrementation via diagonal perturbations
898	Mingxiang Chen, Xu-Yan Chen and Jack K. Hale,	Structural stability for time-periodic one-dimensional parabolic equations
899	Hong-Ming Yin,	Global solutions of Maxwell's equations in an electromagnetic field with the temperature-dependent electrical conductivity
900	Robert Grone, Russell Merris and William Watkins,	Laplacian unimodular equivalence of graphs
901	Miroslav Fiedler,	Structure-ranks of matrices
902	Miroslav Fiedler,	An estimate for the nonstochastic eigenvalues of doubly stochastic matrices
903	Miroslav Fiedler,	Remarks on eigenvalues of Hankel matrices
904	Charles R. Johnson, D.D. Olesky, Michael Tsatsomeros and P. van den Driessche,	Spectra with positive elementary symmetric functions
905	Pierre-Alain Gremaud,	Thermal contraction as a free boundary problem
906	K.L. Cooke, Janos Turi and Gregg Turner,	Stabilization of hybrid systems in the presence of feedback delays
907	Robert P. Gilbert and Yongzhi Xu,	A numerical transmutation approach for underwater sound propagation
908	LeRoy B. Beasley, Richard A. Brualdi and Bryan L. Shader,	Combinatorial orthogonality
909	Richard A. Brualdi and Bryan L. Shader,	Strong hall matrices
910	Håkan Wennerström and David M. Anderson,	Difference versus Gaussian curvature energies; monolayer versus bilayer curvature energies applications to vesicle stability
911	Shmuel Friedland,	Eigenvalues of almost skew symmetric matrices and tournament matrices
912	Avner Friedman, Bei Hu and J.L. Velazquez,	A Free Boundary Problem Modeling Loop Dislocations in Crystals
913	Ezio Venturino,	The Influence of Diseases on Lotka-Volterra Systems
914	Steve Kirkland and Bryan L. Shader,	On Multipartite Tournament Matrices with Constant Team Size

- 915 Richard A. Brualdi and Jennifer J.Q. Massey, More on Structure-Ranks of Matrices
 916 Douglas B. Meade, Qualitative Analysis of an Epidemic Model with Directed Dispersion
 917 Kazuo Murota, Mixed Matrices Irreducibility and Decomposition
 918 Richard A. Brualdi and Jennifer J.Q. Massey, Some Applications of Elementary Linear Algebra in Combinations
 919 Carl D. Meyer, Sensitivity of Markov Chains
 920 Hong-Ming Yin, Weak and Classical Solutions of Some Nonlinear Volterra Integrodifferential Equations
 921 B. Leinkuhler and A. Ruehli, Exploiting Symmetry and Regularity in Waveform Relaxation Convergence Estimation
 922 Xinfu Chen and Charles M. Elliott, Asymptotics for a Parabolic Double Obstacle Problem
 923 Yongzhi Xu and Yi Yan, An Approximate Boundary Integral Method for Acoustic Scattering in Shallow Oceans
 924 Yongzhi Xu and Yi Yan, Source Localization Processing in Perturbed Waveguides
 925 Kenneth L. Cooke and Janos Turi, Stability, Instability in Delay Equations Modeling Human Respiration
 926 F. Bethuel, H. Brezis, B.D. Coleman and F. Hélein, Bifurcation Analysis of Minimizing Harmonic Maps Describing the Equilibrium of Nematic Phases Between Cylinders
 927 Frank W. Elliott, Jr., Signed Random Measures: Stochastic Order and Kolmogorov Consistency Conditions
 928 D.A. Gregory, S.J. Kirkland and B.L. Shader, Pick's Inequality and Tournaments
 929 J.W. Demmel, N.J. Higham and R.S. Schreiber, Block LU Factorization
 930 Victor A. Galaktionov and Juan L. Vazquez, Regional Blow-Up in a Semilinear Heat Equation with Convergence to a Hamilton-Jacobi Equation
 931 Bryan L. Shader, Convertible, Nearly Decomposable and Nearly Reducible Matrices
 932 Dianne P. O'Leary, Iterative Methods for Finding the Stationary Vector for Markov Chains
 933 Nicholas J. Higham, Perturbation theory and backward error for $AX - XB = C$
 934 Z. Strakos and A. Greenbaum, Open questions in the convergence analysis of the lanczos process for the real symmetric eigenvalue problem
 935 Zhaojun Bai, Error analysis of the lanczos algorithm for the nonsymmetric eigenvalue problem
 936 Pierre-Alain Gremaud, On an elliptic-parabolic problem related to phase transitions in shape memory alloys
 937 Bojan Mohar and Neil Robertson, Disjoint essential circuits in toroidal maps
 938 Bojan Mohar, Convex representations of maps on the torous and other flat surfaces
 939 Bojan Mohar and Svatopluk Poljak Eigenvalues in combinatorial optimization
 940 Richard A. Brualdi, Keith L. Chavey and Bryan L. Shader, Conditional sign-solvability
 941 Roger Fosdick and Ying Zhang, The torsion problem for a nonconvex stored energy function
 942 René Ferland and Gaston Giroux, An unbounded mean-field intensity model: Propagation of the convergence of the empirical laws and compactness of the fluctuations
 943 Wei-Ming Ni and Izumi Takagi, Spike-layers in semilinear elliptic singular Perturbation Problems
 944 Henk A. Van der Vorst and Gerard G.L. Sleijpen, The effect of incomplete decomposition preconditioning on the convergence of conjugate gradients
 945 S.P. Hastings and L.A. Peletier, On the decay of turbulent bursts
 946 Apostolos Hadjidimos and Robert J. Plemmons, Analysis of p -cyclic iterations for Markov chains
 947 ÅBjörck, H. Park and L. Eldén, Accurate downdating of least squares solutions
 948 E.G. Kalnins, William Miller, Jr. and G.C. Williams, Recent advances in the use of separation of variables methods in general relativity
 949 G.W. Stewart, On the perturbation of LU , Cholesky and QR factorizations
 950 G.W. Stewart, Gaussian elimination, perturbation theory and Markov chains
 951 G.W. Stewart, On a new way of solving the linear equations that arise in the method of least squares
 952 G.W. Stewart, On the early history of the singular value decomposition
 953 G.W. Stewart, On the perturbation of Markov chains with nearly transient states
 954 Umberto Mosco, Composite media and asymptotic dirichlet forms
 955 Walter F. Mascarenhas, The structure of the eigenvectors of sparse matrices
 956 Walter F. Mascarenhas, A note on Jacobi being more accurate than QR
 957 Raymond H. Chan, James G. Nagy and Robert J. Plemmons, FFT-based preconditioners for Toeplitz-Block least squares problems
 958 Zhaojun Bai, The CSD, GSVD, their applications and computations
 959 D.A. Gregory, S.J. Kirkland and N.J. Pullman, A bound on the exponent of a primitive matrix using Boolean rank
 960 Richard A. Brualdi, Shmuel Friedland and Alex Pothén, Sparse bases, elementary vectors and nonzero minors of compound matrices
 961 J.W. Demmel, Open problems in numerical linear algebra
 962 James W. Demmel and William Gragg, On computing accurate singular values and eigenvalues of acyclic matrices
 963 James W. Demmel, The inherent inaccuracy of implicit tridiagonal QR
 964 J.J.L. Velázquez, Estimates on the $(N - 1)$ -dimensional Hausdorff measure of the blow-up set for a semilinear heat equation