arXiv:1105.1190v1 [math.AP] 5 May 2011

GLOBAL EXPONENTIAL CONVERGENCE TO VARIATIONAL TRAVELING WAVES IN CYLINDERS

C. B. MURATOV* AND M. NOVAGA[†]

Abstract. We prove, under generic assumptions, that the special variational traveling wave that minimizes the exponentially weighted Ginzburg-Landau functional associated with scalar reactiondiffusion equations in infinite cylinders is the long-time attractor for the solutions of the initial value problems with front-like initial data. The convergence to this traveling wave is exponentially fast. The obtained result is mainly a consequence of the gradient flow structure of the considered equation in the exponentially weighted spaces and does not depend on the precise details of the problem. It strengthens our earlier generic propagation and selection result for "pushed" fronts.

 ${\bf Key \ words.} \ reaction-diffusion \ equations, \ front \ propagation, \ nonlinear \ stability, \ front \ selection, \ exponentially \ weighted \ spaces$

AMS subject classifications. 35B40, 35C07, 35K57, 35A15

1. Introduction. One of the most fundamental problems in the theory of reactiondiffusion equations has to do with the long-time asymptotic behavior of solutions of the associated initial value problem on unbounded domains [6, 38, 39]. In its simplest form, it may be formulated for a one-dimensional scalar reaction-diffusion equation

$$u_t = u_{xx} + f(u), \qquad u : \mathbb{R} \times \mathbb{R}^+ \to [0, 1], \tag{1.1}$$

with an unbalanced bistable nonlinearity f(u), i.e., when f is a smooth function which has precisely three non-degenerate zeros in [0, 1], with

$$f(0) = f(1) = 0,$$
 $f'(0) < 0,$ $f'(1) < 0,$ $\int_0^1 f(u) du > 0,$ (1.2)

e.g. $f(u) = u(1-u)(u-\frac{1}{4})$. For such an equation, it was first proved by Kanel' that initial data $u(x,t) = u_0(x)$ with the property that $u_0(x) = 0$ for all x > b, $u_0(x) = 1$ for all x < a, and $u_0(x)$ is monotone decreasing for $x \in (a,b)$, with some $-\infty < a < b < +\infty$, converges uniformly to a (unique up to translations) traveling wave solution, i.e., a solution $u(x,t) = \bar{u}(x-ct)$ of (1.1), with some uniquely determined speed c > 0, connecting monotonically u = 0 at $x = +\infty$ with u = 1 at $x = -\infty$, in a reference frame moving with speed c [14, 15]. In a subsequent work, Fife and McLeod extended this result to a much wider class of initial data and also showed that the convergence is exponentially fast [7]. Qualitatively, the conclusion of these analyses is that the solution of the considered initial value problem with front-like initial data converges exponentially fast to a traveling front invading the "less stable" equilibrium u = 0 by a "more stable" equilibrium u = 1. We note that a similar result was proved for a certain class of monostable nonlinearities [31], but it does *not* hold (in the reference frame moving with constant speed and in the sense of exponential convergence) in the case of the Fisher's equation [17, 35, 4, 16].

^{*}Department of Mathematical Sciences, New Jersey Institute of Technology, Newark, NJ 07102, USA (muratov@njit.edu). The work of this author was supported by NSF via grants DMS-0718027 and DMS-0908279.

[†]Dipartimento di Matematica, Università di Padova, Via Trieste 63, 35121 Padova, Italy (novaga@dm.unipd.it).

In the multi-dimensional setting, these kinds of results were subsequently obtained for initial boundary value problems for equations in infinite cylindrical domains:

$$u_t = \Delta u + f(u, y), \qquad u(x, 0) = u_0(x),$$
(1.3)

where $u: \Sigma \times \mathbb{R}^+ \to \mathbb{R}, \Sigma = \Omega \times \mathbb{R} \subset \mathbb{R}^n, \Omega \subset \mathbb{R}^{n-1}$ is a bounded domain with sufficiently smooth boundary, $f: \mathbb{R} \times \Omega \to \mathbb{R}$ is a nonlinear reaction term, with either Neumann or Dirichlet boundary conditions. By $x = (y, z) \in \Sigma$, we always denote a point with coordinate $y \in \Omega$ on the cylinder cross-section and $z \in \mathbb{R}$ along the cylinder axis. More generally, one can consider either Dirichlet or Neumann boundary conditions on different connected portions of $\partial\Omega$:

$$u\Big|_{\partial\Sigma_{\perp}} = 0, \qquad \nu \cdot \nabla u\Big|_{\partial\Sigma_{0}} = 0, \tag{1.4}$$

where $\partial \Sigma_{\pm} = \partial \Omega_{\pm} \times \mathbb{R}$ and $\partial \Sigma_0 = \partial \Omega_0 \times \mathbb{R}$, allowing for more than one connected component for $\partial \Omega$ (for motivation and further discussion of the boundary conditions, see [24, 25]). Note that transverse advection by a potential flow can also be straightforwardly included in the present treatment, as was done in [24, 25]. For simplicity of presentation, in this paper we do not consider the advection term and concentrate on pure reaction-diffusion problems.

Without loss of generality, we may assume that u = 0 is a trivial solution of (1.3) and consider traveling waves that invade the u = 0 equilibrium, i.e., the solutions of (1.3) and (1.4) in the form $u(x,t) = \bar{u}(y, z - ct)$, for some c > 0, which converge to zero uniformly as $z \to +\infty$. These solutions satisfy the elliptic equation

$$\Delta \bar{u} + c\bar{u}_z + f(\bar{u}, y) = 0, \qquad (1.5)$$

together with the respective boundary conditions in (1.4) (by a solution, we mean a pair (c, \bar{u}) , with $\bar{u} \in C^2(\Sigma) \cap C^1(\overline{\Sigma})$ being a classical solution of (1.5) and (1.4)). We refer to [3, 38, 24] and references therein, for a comprehensive treatment of the subject of traveling waves. In particular, under certain specific assumptions one obtains uniqueness (up to translations) and global exponential convergence to these solutions for the initial value problem with front-like initial data [22, 29, 30] (see the end of Sec. 2 for a more detailed discussion and a comparison with the present results). This property, therefore, indicates the ubiquitous role of the traveling fronts in the behavior of the solutions of (1.3).

Since in general (1.5) may have many solutions, an important question is which of these solutions, if any, can be a long-time limit of the evolution governed by (1.3), for a given class of initial data. As was recently pointed out in [23], in the case of initial data with sufficiently fast exponential decay at $z = +\infty$ the relevant class of traveling wave solutions consists of the so-called variational traveling waves, even for systems of reaction-diffusion equations in which the nonlinearity is a gradient. More recently, we showed that a special class of variational traveling wave solutions that minimize the exponentially weighted Ginzburg-Landau functional (see Sec. 2 for precise definitions and statements) are relevant for the long-time behavior of the initial value problem in the sense of propagation of the leading edge and, in particular, determine the propagation speed for front-like initial data [24]. It is then natural to ask whether these special traveling fronts are also the long-time attractors for the solutions of (1.3) in the moving reference frame. In this paper, we give a positive answer to this question under a few extra non-degeneracy assumptions to those of [24] which hold generically in the considered class of problems. Our paper is organized as follows. In Sec. 2, we introduce the variational formulation for the traveling waves of interest, state the main result and compare it with those available in the literature. In Sec. 3, we list and discuss our assumptions, as well as state a number of auxiliary results used in the paper. In Sec. 4, we perform local stability analysis of the traveling waves of interest in the exponentially weighted Sobolev spaces, and in Sec. 5 we prove convergence to the traveling wave in the large, completing the proof of the main theorem.

Some notation. For every $-\infty \leq a < b \leq +\infty$ and c > 0, the symbol $L^2_c(\Omega \times (a, b))$ denotes the Hilbert space of all functions $u : \Omega \times (a, b) \to \mathbb{R}$ with $||u||^2_{L^2_c(\Omega \times (a, b))} = \int_a^b \int_\Omega e^{cz} u^2(y, z) \, dy \, dz$. Likewise, by $L^2_c(\Sigma)$, $H^1_c(\Sigma)$ and $H^2_c(\Sigma)$, we denote the spaces of functions which are square integrable with the above exponential weight, together with their first and second derivatives, respectively, in Σ . We also use the symbol $C_b(A)$ to denote the space of bounded continuous function on A equipped with the sup-norm. In all statements and proofs the constants are always assumed to implicitly depend on f, Ω and the choice of the boundary conditions. In the proofs the numbers C, M, etc., may change from line to line. We will also use the symbol T_R to denote a translation by R along the z-axis, i.e., $T_R u(\cdot, z) = u(\cdot, z - R)$.

2. Variational formulation and main result.

The fact that (1.5) possesses a variational structure in exponentially weighted Sobolev spaces was, to our knowledge, first pointed out by Heinze [11, 12] (see also [7, 38, 29, 19, 9, 24] in the context of (1.3), and [23, 20, 27, 8] in the context of its extensions). As we recently showed in [24], for scalar reaction-diffusion equations considered here the solution of (1.5) which determines the asymptotic speed of propagation with front-like initial data is a special variational traveling wave which is the minimizer of the the exponentially weighted Ginzburg-Landau functional

$$\Phi_{c}[u] := \int_{\Sigma} e^{cz} \left(\frac{1}{2} |\nabla u|^{2} + V(u, y) \right) \, dx \qquad c > 0,$$
(2.1)

where

$$V(u,y) = -\int_0^u f(s,y)\chi_{[0,1]}(s)\,ds, \qquad \chi_{[0,1]}(s) = \begin{cases} 1, & s \in [0,1]\\ 0, & s \notin [0,1] \end{cases}, \tag{2.2}$$

over all functions lying in the exponentially weighted Sobolev space $H_c^1(\Sigma)$. We point out that such a minimizer can only exist for a specific value of $c = c^{\dagger} > 0$ (see Theorem 2 below). Under quite general assumptions on the potential V, in [24, Theorem 5.8] we proved that the asymptotic speed of propagation of solutions to (1.3) is precisely given by c^{\dagger} , assuming that the initial datum is front-like, i.e., if it stays sufficiently far away from zero as $z \to -\infty$ and decays sufficiently fast to zero as $z \to +\infty$. In this paper, we discuss the local and global stability of such variational traveling waves.

Our main result is contained in the following theorem (for the details of the definitions and hypotheses, see Sec. 3):

THEOREM 1. Assume hypotheses (H1)–(H3) and (N1)–(N2) are satisfied, and let c^{\dagger} , \bar{u} , v be as in Theorem 2. Then there exist $\alpha > 0$ and $\sigma > 0$, such that if $u_0 \in C^0(\overline{\Sigma}) \cap W^{1,\infty}(\Sigma) \cap L^2_{c^{\dagger}}(\Sigma)$ satisfies $0 \le u_0 \le 1$ and

$$\liminf_{z \to -\infty} u_0(\cdot, z) \ge v - \alpha \qquad \text{uniformly in } \Omega \,, \tag{2.3}$$

there exists $R_{\infty} \in \mathbb{R}$, such that if u is the solution of (1.3) and (1.4) with initial datum u_0 , then

$$||\mathbf{T}_{R_{\infty}-c^{\dagger}t}u(\cdot,t)-\bar{u}||_{H^{2}_{a^{\dagger}}(\Sigma)} \le Ce^{-\sigma t}$$

$$(2.4)$$

for every $t \ge t_0$, with arbitrary $t_0 > 0$ and some C > 0 independent of t.

Note that by Proposition 3.1 below we know that $u(\cdot, t)$ is bounded in $W^{2,p}(\Omega \times [M, M+1])$ uniformly in $M \in \mathbb{R}$ and $t \in [t_0, +\infty)$, for all $t_0 > 0$ and $p < \infty$. Since this bound also applies to \bar{u} , from (2.4) we get the following

COROLLARY 2.1. In the statement of Theorem 1, the inequality (2.4) may be replaced with

$$\|\mathbf{T}_{R_{\infty}-c^{\dagger}t}u(\cdot,t)-\bar{u}\|_{C^{1}(\overline{\Omega}\times[z_{0},z_{1}])} \leq Ce^{-\sigma t},$$
(2.5)

for all $z_0 < z_1$, $t \ge t_0 > 0$, and some C > 0 independent of t and z_1 .

Let us point out that the upper bound $u_0 \leq 1$ in Theorem 1 can be replaced with the condition $u_0(\cdot, z) \leq \overline{v}$ for every $z \in \mathbb{R}$, where $\overline{v} \in C^2(\Omega) \cap C^1(\overline{\Omega})$ satisfies

$$\bar{v} > 0, \quad \Delta_y \bar{v} + f(\bar{v}, y) \le 0 \quad \text{for all } y \in \Omega,$$
(2.6)

together with the boundary conditions from (1.4). In this case, the condition $f(1, y) \leq 0$ in assumption (H1) below should be replaced by (2.6), the conditions in (H2) should hold for $0 \leq u \leq \bar{v}$, and the definition of V in (2.2) should be modified accordingly. We note that, in particular, one can choose \bar{v} to be any positive critical point of the energy functional E associated with Φ_c :

$$E[v] := \int_{\Omega} \left(\frac{1}{2} |\nabla_y v|^2 + V(v, y) \right) dy, \qquad v \in H^1(\Omega), \quad v \big|_{\partial \Omega_{\pm}} = 0.$$
(2.7)

To each such \bar{v} one can associate a minimizer of Φ_c in the admissible class of functions that are bounded above by \bar{v} . Then, under the assumption that the initial data approaches \bar{v} uniformly from below as $z \to -\infty$ one can make the conclusion (under generic non-degeneracy assumptions) that the solution of the initial value problem converges exponentially to the corresponding minimizer. Thus, every front-like initial data in a more restricted sense of connecting zero to a critical point \bar{v} of E converges to the minimizer associated with that critical point. More precisely, we have

COROLLARY 2.2. Under hypotheses (H1)-(H3), (N1)-(N2), with the trial function u in hypothesis (H3) satisfying $u \leq \overline{v}$, where $\overline{v} > 0$ is a critical point of E, let \overline{u} be the unique (up to translations) non-trivial minimizer of $\Phi_{c^{\dagger}}$ over functions $u \in H^1_{c^{\dagger}}(\Sigma)$ satisfying (1.4) and $0 \leq u \leq \overline{v}$. Let $u_0 \in C^0(\overline{\Sigma}) \cap W^{1,\infty}(\Sigma) \cap L^2_{c^{\dagger}}(\Sigma)$ satisfy $0 \leq u_0 \leq \overline{v}$ and $u_0(\cdot, z) \to \overline{v}$ uniformly in Ω as $z \to -\infty$. Then the conclusion of Theorem 1 holds.

An important implication of Corollary 2.2 is that \bar{v} selects the attracting variational traveling wave solution in the long time limit. This kind of conclusion was made by us earlier for the propagation speed of the leading edge without the non-degeneracy assumptions of the present paper [24].

We note that the problem of convergence to traveling waves for solutions of (1.3) has been widely considered in the mathematical literature. We refer to [6, 38, 30] and references therein, for a general overview on the subject. Specifically, our result should be compared with [30, Theorem 3.7] by Roquejoffre, where, in particular, convergence to variational traveling waves is proved (in our notation) for initial data that approach

zero from above as $z \to +\infty$ and a non-degenerate local minimizer $\bar{v} > 0$ of E from below as $z \to -\infty$.

Roquejoffre makes a crucial assumption that there exists a variational traveling wave connecting \bar{v} at $z = -\infty$ with zero at $z = +\infty$. In contrast, our results do not require existence of such a traveling wave. Instead, we require that the initial data decay sufficiently rapidly to zero as $z \to +\infty$ and stay approximately above the local minimizer v of E corresponding to the limit at $z = -\infty$ for the special variational traveling wave \bar{u} given by Theorem 2 as $z \to -\infty$. Under this condition the solution of (1.3) is attracted to a translate of \bar{u} on compacts in the moving reference frame (see Theorem 1 for a precise statement). We note that in the class of front-like initial data with sufficiently fast exponential decay considered by us global stability of a traveling wave connecting zero to \bar{v} is a simple consequence of Corollary 2.2. Indeed, if there exists a variational traveling wave u_c connecting zero to \bar{v} , then by Proposition 3.3 we have $u_c = \bar{u}$, where \bar{u} is as in Corollary 2.2 (note that in this case hypotheses (H3) and (N2) are unnecessary). Thus, within the scope of (1.3) and front-like initial data decaying sufficiently fast, our results are applicable to more general initial data than the ones considered in [30] and, most importantly, provide a selection criterion for the limit front in terms of the asymptotic behavior of the initial data as $z \to -\infty$. We also point out that our assumptions concerning the nonlinearity f (see (H1)–(H3) below) are quite general compared to the assumptions usually made in the literature [3, 36, 30]. In particular, these assumptions can be readily verified in practice (for examples see [19, 20, 25]).

3. Preliminaries. Throughout this paper we assume Ω to be a bounded domain (connected open set, not necessarily simply connected) with a boundary of class C^2 . We start by listing the assumptions on the nonlinearity f which we need in Theorem 1. The function $f : [0, 1] \times \overline{\Omega} \to \mathbb{R}$ satisfies:

$$\begin{array}{ll} (\mathbf{H1}) & f(0,y) = 0 & f(1,y) \leq 0 & \text{for all } y \in \Omega, \\ (\mathbf{H2}) & f \in C^{0,\gamma}([0,1] \times \overline{\Omega}) & f_u = \frac{\partial f}{\partial u} \in C^{0,\gamma}([0,1] \times \overline{\Omega}) \text{ for some } \gamma \in (0,1). \end{array}$$

Hypotheses (H1) and (H2) are needed to guarantee, in particular, existence and basic regularity properties of solutions of (1.3). Indeed, from [24, Proposition 5.1] and [21, Chapter 7] we have the following

PROPOSITION 3.1. Under assumptions (H1) and (H2), let $u_0 \in C^0(\overline{\Sigma}) \cap W^{1,\infty}(\Sigma)$. Let also u_0 satisfy the boundary conditions (1.4) and assume $u_0(x) \in [0,1]$ for all $x \in \Sigma$. Then there exists a unique solution (using notation of [5])

$$u \in C_1^2(\Sigma \times (0,\infty)) \cap C^0(\overline{\Sigma} \times [0,+\infty))$$

of (1.3) with boundary conditions (1.4) and initial condition $u(\cdot, 0) = u_0$, which satisfies $0 \le u \le 1$ and $||\nabla u||_{C_b(\overline{\Sigma} \times (0, +\infty))} < \infty$. Moreover, letting $\Sigma_M := \Omega \times [M, M+1]$ for all $M \in \mathbb{R}$, we have

 $\|u(\cdot,t)\|_{W^{2,p}(\Sigma_M)} \le C(t_0,p) \qquad \text{for all } t \ge t_0 > 0, \ p > 1.$ (3.1)

Finally, if $u_0 \in L^2_c(\Sigma)$ for some c > 0, we also have

$$u \in C^{\alpha}((0, +\infty); H_c^2) \cap C^{1,\alpha}((0, +\infty); L_c^2(\Sigma)) \quad \text{for all } \alpha \in (0, 1),$$
(3.2)

and

$$u_t \in C((0, +\infty); H^1_c(\Sigma)).$$
 (3.3)

We now turn to the assumption which is both necessary and sufficient for the existence of the special variational traveling wave solution considered in this paper [19, 24].

(H3) There exist c > 0, satisfying $c^2 + 4\nu_0 > 0$, where

$$\nu_{0} = \min_{\substack{\psi \in H^{1}(\Omega)\\\psi|_{\partial\Omega_{+}}=0}} \frac{\int_{\Omega} (|\nabla_{y}\psi|^{2} - f_{u}(0, y)\psi^{2}) \, dy}{\int_{\Omega} \psi^{2} \, dy},$$
(3.4)

and $u \in H_c^1(\Sigma)$, such that $\Phi_c[u] \leq 0$ and $u \neq 0$.

REMARK 3.2. As was shown in [24], in the case $\nu_0 \ge 0$ the hypothesis (H3) is equivalent to the condition

$$\inf_{\substack{v \in H^1(\Omega) \\ v|_{\partial\Omega_+} = 0}} E[v] < 0.$$
(3.5)

Under the above assumptions, we can state the existence result concerning the variational traveling wave which is the minimizer of Φ_c with a suitably fixed translation.

THEOREM 2. Under hypotheses (H1)-(H3), there exists a unique value of $c^{\dagger} \geq c$, where c is defined by hypothesis (H3), and a unique function $\bar{u} \in C^2(\Sigma) \cap C^1(\overline{\Sigma})$, $\bar{u} \neq 0$, such that (c^{\dagger}, \bar{u}) solve (1.5) and (1.4), and \bar{u} satisfies $||\bar{u}(\cdot, 0)||_{L^{\infty}(\Omega)} =$ $\frac{1}{2} \sup_{z \in \mathbb{R}} ||\bar{u}(\cdot, z)||_{L^{\infty}(\Omega)}$ and minimizes $\Phi_{c^{\dagger}}$ in $H^1_{c^{\dagger}}(\Sigma)$. Moreover $\bar{u} \in H^2_{c^{\dagger}}(\Sigma) \cap W^{1,\infty}(\Sigma)$, $\bar{u}_z \in H^2_{c^{\dagger}}(\Sigma)$, $\bar{u}_z < 0$ in Σ , and

$$\lim_{z \to +\infty} \bar{u}(\cdot, z) = 0, \quad \lim_{z \to -\infty} \bar{u}(\cdot, z) = v \quad \text{in } C^1(\overline{\Omega}), \tag{3.6}$$

where $v: \Omega \to \mathbb{R}$ is a local minimizer of E defined in (2.7), with E[v] < 0.

For the proof see [24, Theorem 3.3] and [20, Proposition 3.3(ii)] (the latter argument also applies to \bar{u}_z by differentiating (1.5) in z).

Let us point out that the minimizer of Theorem 2 is in some sense the "maximal" variational traveling wave solution. More precisely, we have the following result:

PROPOSITION 3.3. Let hypotheses (H1)–(H3) be satisfied, and let (c, u) solve (1.5) and (1.4), with c > 0, $u \in H_c^1(\Sigma)$ and 0 < u < 1. Then, if v, c^{\dagger}, \bar{u} are as in Theorem 2, and

$$\liminf u(\cdot, z) \ge v \qquad \quad uniformly \ in \ \Omega, \tag{3.7}$$

we have $c = c^{\dagger}$ and $u = T_R \bar{u}$, for some $R \in \mathbb{R}$. In particular, the inequality in (3.7) is, in fact, equality.

Proof. First note that we cannot have $c > c^{\dagger}$. Indeed, if this inequality were true, by [20, Proposition 3.5] the pair (c, u) can be taken as a trial function in hypothesis (H3), contradicting the conclusion of Theorem 2 that $c^{\dagger} \ge c$. On the other hand, it is easy to see that $c < c^{\dagger}$ is also impossible. Indeed, arguing as in the proof of [24, Proposition 5.5], for any $c' \in (c, c^{\dagger})$ there exists a non-trivial minimizer $\bar{u}_{c'}$ of $\Phi_{c'}$ in the class of functions in $H^1_{c'}(\Sigma)$ which stay below v and vanish outside $\Sigma_R = \Omega_{\sigma} \times (-R, R)$, with $\Omega_{\sigma} = \{y \in \Omega : \operatorname{dist}(y, \partial \Omega_{\pm}) > \sigma\}$, where R > 0 is large enough and $\sigma > 0$ is small enough.¹ Furthermore, $\bar{u}_{c'}$ is a classical solution of (1.5) with c = c' in Σ_R and $\bar{u}_{c'} \leq \max(0, v - \varepsilon)$ for some $\varepsilon > 0$. Therefore, by (3.7) the function $T_{R'}\bar{u}_{c'} < u$ in Σ for some $R' \in \mathbb{R}$ sufficiently large negative, and by parabolic comparison principle [26] we have $\bar{u}_{c'} < T_{(c-c')t-R'}u$ for all t > 0. However, the latter is impossible, since the right-hand side of this inequality converges to zero in $H_c^1(\Sigma)$. Thus $c = c^{\dagger}$, hence u is a minimizer by [20, Proposition 3.5], and the result follows from [24, Theorem 3.3(v)]. \Box

We note that, in particular, the result in Proposition 3.3 allows to extend the statements about monotonicity and uniqueness of traveling waves established in the classical work of Berestycki and Nirenberg [3] for (1.3) and (1.4) (see also [36, 37]), in the class of variational traveling waves, under only an assumption that the traveling wave approaches a limit from below as $z \to -\infty$, zero from above as $z \to +\infty$, and is sandwiched between these two limits. Indeed, suppose (c, u) is such a traveling wave, with $u(\cdot, z) \to \bar{v}$ as $z \to -\infty$, with $0 < \bar{v} \leq 1$. Then by the argument of [20, Proposition 6.6] v is a critical point of E, and by [24, Proposition 3.5] we have $c^2 + 4\nu_0 > 0$. So by Proposition 3.3 this traveling wave is a non-trivial minimizer of Φ_c in $H_c^1(\Sigma)$ over all positive functions bounded above by \bar{v} , and the result follows from [24, Theorem 3.3]. In particular, we do not require any non-degeneracy assumptions for the limits of $u(\cdot, z)$ as $z \to \pm\infty$, as is done [3]. Thus, we have:

COROLLARY 3.4. Under hypotheses (H1) and (H2), let c > 0, and let $u \in H_c^1(\Sigma)$ be a solution of (1.5) and (1.4), satisfying $u(\cdot, z) \to \overline{v}$ uniformly in Ω as $z \to -\infty$, where $0 < \overline{v} \leq 1$ and $0 < u < \overline{v}$. Then $c^2 + 4\nu_0 > 0$, the value of c is unique, $u_z < 0$, and u is unique up to translations.

We now list two additional technical assumptions (see also [30]), which are generically satisfied and are needed to prove global exponential stability of the minimizers of Φ_c for initial data bounded below by v as $z \to -\infty$.

(N1) For v as in Theorem 2 we have

$$\tilde{\nu}_{0} = \min_{\substack{\psi \in H^{1}(\Omega) \\ \psi|_{\partial\Omega_{+}} = 0}} \frac{\int_{\Omega} (|\nabla_{y}\psi|^{2} - f_{u}(v, y)\psi^{2}) \, dy}{\int_{\Omega} \psi^{2} \, dy} > 0.$$
(3.8)

(N2) For v < 1 as in Theorem 2 there is no solution (c^{\dagger}, \bar{u}) of (1.5) and (1.4), with c^{\dagger} as in Theorem 2, such that $v < \bar{u} < 1$ on Σ .

Conditions (N1) and (N2) are generic in the sense that the set of nonlinearities f such that (N1) or (N2) do not hold is a meager subset of all f's obeying (H1)-(H3), in the natural topology (for similar notions related to perturbations of Ω see [13]). Indeed, condition (N1) is generic, since by the results of [24] we have $\tilde{\nu}_0 \geq 0$, so that (N1) only excludes the degenerate case of $\tilde{\nu}_0 = 0$. Similarly, condition (N2) excludes the non-generic possibility of existence of a traveling front invading v from above with the same speed c^{\dagger} as the front invading zero by v. To see that the only non-trivial alternative would be to have a front invading v with lower speed, consider the following variational problem. Given c > 0 and $h \in H_c^1(\Sigma)$ satisfying (1.4), let

$$\Psi_c^v[h] := \int_{\Sigma} e^{cz} \left(\frac{|\nabla h|^2}{2} + V(v+h,y) - V(v,y) - V'(v,y) h \right) \, dx, \qquad (3.9)$$

¹This choice of Σ_R also corrects a minor inaccuracy in the proof of [24, Proposition 5.5].

where we used the notation $V'(s, y) := \partial V(s, y)/\partial s$. Notice that, if h is a critical point of $\Psi_{c\dagger}^v$, then $\bar{u} = v + \bar{h}$ is a solution of (1.5) and (1.4). We set

$$c_v^{\dagger} := \inf \left\{ c > 0 : \ \Psi_c^v[h] \ge 0 \text{ for all } h \ge 0 \right\}.$$
 (3.10)

Then the following result concerning c_v^{\dagger} holds.

LEMMA 3.5. The functional Ψ_c^v is weakly sequentially lower semicontinuous and coercive in $H_c^1(\Sigma)$ for all $c > c_v^{\dagger}$, and $c_v^{\dagger} \le c^{\dagger}$, where c^{\dagger} is as in Theorem 2. Moreover, under hypothesis (N2) we have $c_v^{\dagger} < c^{\dagger}$.

In other words, under hypothesis (N2) it is only possible to have such a system of stacked waves [38] invading zero, that the front connecting zero with v moves faster than the front invading v from above.

Proof of Lemma 3.5. First of all, reasoning as in [20, Proposition 5.5] and using the fact that $\tilde{\nu}_0 \geq 0$, where $\tilde{\nu}_0$ is defined in (3.8) [24, Theorem 3.3(iv)], one can see that Ψ_c^v is weakly lower semicontinuous in $H_c^1(\Sigma)$ for all c > 0, so the results of [24] apply to Ψ_c^v . Moreover, reasoning as in the proof of [20, Proposition 6.9], we also get that Ψ_c^v is coercive in $H_c^1(\Sigma)$ for all $c > c_v^{\dagger}$.

Let us now prove that $c_v^{\dagger} \leq c^{\dagger}$. Assume by contradiction that there exists $w \geq v$, such that $\Psi_{c^{\dagger}}^v[w-v] < 0$. Slightly perturbing w, we can ensure that w = v for $z \geq z_0$, with $z_0 \in \mathbb{R}$ big enough. Let \bar{u} be the minimizer of $\Phi_{c^{\dagger}}$ given by Theorem 2, and let $\varepsilon \leq -\Psi_{c^{\dagger}}^v[w-v]/2$. Since $\bar{u}(\cdot,z) \to v$ in $H^1(\Omega)$ as $z \to -\infty$, up to a suitable translation we can perturb \bar{u} into a function $\tilde{u} \in H^1_{c^{\dagger}}(\Sigma)$ such that $\tilde{u} = v$ for $z \leq z_0$ and $\Phi_{c^{\dagger}}[\tilde{u}] \leq \varepsilon$. Define $\hat{u} \in H^1_{c^{\dagger}}(\Sigma)$ as

$$\hat{u}(y,z) := \begin{cases} w(y,z) & \text{if } z \leq z_0 \\ \tilde{u}(y,z) & \text{if } z > z_0. \end{cases}$$

Letting $h = w - v \in H^1_{c^{\dagger}}(\Sigma)$ and satisfying (1.4), after an integration by parts and using the Euler-Lagrange equation for E satisfied by v, we get

$$\begin{split} \Phi_{c^{\dagger}}[\hat{u}] &= \int_{-\infty}^{z_0} \int_{\Omega} e^{c^{\dagger} z} \left(\frac{|\nabla(v+h)|^2}{2} + V(v+h,y) \right) \, dy \, dz \\ &+ \int_{z_0}^{+\infty} \int_{\Omega} e^{c^{\dagger} z} \left(\frac{|\nabla \tilde{u}|^2}{2} + V(\tilde{u},y) \right) \, dy \, dz \\ &= \int_{-\infty}^{z_0} \int_{\Omega} e^{c^{\dagger} z} \left(\frac{|\nabla h|^2}{2} + V(v+h,y) - V(v,y) - V'(v,y)h \right) \, dy \, dz \\ &+ \int_{-\infty}^{z_0} \int_{\Omega} e^{c^{\dagger} z} \left(\frac{|\nabla yv|^2}{2} + V(v,y) \right) \, dy \, dz \\ &+ \int_{z_0}^{+\infty} \int_{\Omega} e^{c^{\dagger} z} \left(\frac{|\nabla \tilde{u}|^2}{2} + V(\tilde{u},y) \right) \, dy \, dz \\ &= \Psi_{c^{\dagger}}^v[h] + \Phi_{c^{\dagger}}[\tilde{u}] \leq \frac{\Psi_{c^{\dagger}}^v[h]}{2} < 0 \end{split}$$

which contradicts the minimizing property $\Phi_{c^{\dagger}}[\bar{u}] = 0$ of \bar{u} [20, Proposition 3.2].

To conclude the proof, it remains to prove that $c_v^{\dagger} < c^{\dagger}$ under hypothesis (N2). If $c_v^{\dagger} = c^{\dagger}$, then for every $c \in (0, c^{\dagger})$, there exists a function $h_c \not\equiv 0$, such that $\Psi_c^v[h_c] < 0$. Hence, the analog of hypothesis (H3) holds for Ψ_c^v , and, therefore, there exists a non-trivial minimizer \bar{h} of Ψ_c^v for some $\hat{c} \ge c^{\dagger}$. On the other hand, by the argument of [24, Proposition 5.5], we have $\hat{c} \leq c_v^{\dagger}$. So $\hat{c} = c^{\dagger}$, and since $\hat{u} = v + \bar{h} > v$ is a solution of (1.5) and (1.4) with $c = \hat{c}$, this violates assumption (N2).

Finally, we note that if either (N1) or (N2) are violated, one would not expect exponential stability of \bar{u} in the reference frame moving with speed c^{\dagger} any more. Therefore, in some sense these conditions are also necessary for the results obtained by us.

4. Local stability in $L^2_{c^{\dagger}}(\Sigma)$. In this section we prove stability of the variational traveling wave \bar{u} minimizing $\Phi_{c^{\dagger}}$ in the reference frame moving with speed c^{\dagger} up to perturbations which are small in the $L^2_{c^{\dagger}}$ -norm and stay approximately above v behind the front.

THEOREM 3. Assume hypotheses (H1)-(H3) and (N1)-(N2) hold, and let \bar{u} and c^{\dagger} be as in Theorem 2. Then there exist $\alpha > 0$ and $\sigma > 0$, such that for every u_0 as in Theorem 1 and for every $\omega > 0$ there exists $\varepsilon > 0$, such that if

$$||u_0 - \bar{u}||_{L^2_{c^{\dagger}}(\Sigma)} \le \varepsilon, \tag{4.1}$$

the solution u(x,t) of

$$u_t = \Delta u + c^{\dagger} u_z + f(u, y), \tag{4.2}$$

with boundary conditions in (1.4) and $u(x,0) = u_0(x)$ satisfies

$$||u(\cdot,t) - \mathcal{T}_{R_{\infty}}\bar{u}||_{L^{2}_{c^{\dagger}}(\Sigma)} \le \omega e^{-\sigma t}, \qquad |R_{\infty}| \le \omega,$$
(4.3)

for some $R_{\infty} \in \mathbb{R}$.

We note that our approach differs somewhat from the conventional approach to the studies of front stability [32, 2, 28, 29] in the way we treat translations along the cylinder axis. We track the front position by minimizing the $L_{c^{\dagger}}^2$ -distance between the solution of (4.2) and a translate of \bar{u} . As a consequence, the deviation between the solution and the closest translate of \bar{u} is automatically orthogonal to the null-space of the linearization operator, allowing to readily establish the exponential decay of the $L_{c^{\dagger}}^2$ -distance. Thus, our method is more variational in nature. Let us also point out that, in contrast to the usual approach, our initial data do not need to be close to \bar{u} in L^{∞} in the whole cylinder, they may be significantly larger than \bar{u} at large negative z.

Throughout the rest of this section, hypotheses (H1)–(H3) and (N1)–(N2) are assumed to hold, and c^{\dagger} , \bar{u} , v always refer to the minimizer in Theorem 2. We begin with the following basic lemma concerning the linearization around \bar{u} .

LEMMA 4.1. There exists K > 0, such that

$$\int_{\Sigma} e^{c^{\dagger} z} \left(|\nabla w|^2 - f_u(\bar{u}, y) w^2 \right) dx \ge K \int_{\Sigma} e^{c^{\dagger} z} w^2 dx, \tag{4.4}$$

for all $w \in H^1_{c^{\dagger}}(\Sigma)$ satisfying $\int_{\Sigma} e^{c^{\dagger} z} w \bar{u}_z dx = 0$.

Proof. First of all, observe that by choosing R_1 and R_2 sufficiently large, we have

$$\frac{\int_{-\infty}^{-R_1} \int_{\Omega} e^{c^{\dagger} z} \left(|\nabla w|^2 - f_u(\bar{u}, y) w^2 \right) dy \, dz}{\int_{-\infty}^{-R_1} \int_{\Omega} e^{c^{\dagger} z} w^2 \, dy \, dz} \ge K_1 > 0, \tag{4.5}$$

$$\frac{\int_{R_2}^{+\infty} \int_{\Omega} e^{c^{\dagger} z} \left(|\nabla w|^2 - f_u(\bar{u}, y) w^2 \right) dy \, dz}{\int_{R_2}^{+\infty} \int_{\Omega} e^{c^{\dagger} z} w^2 \, dy \, dz} \ge K_2 > 0, \tag{4.6}$$

for all $w \in H^1_{c^{\dagger}}(\Sigma)$. Indeed, if z is large enough negative, then $\bar{u}(\cdot, z)$ is sufficiently close in $L^{\infty}(\Omega)$ to v. Hence, (4.5) holds in view of (3.8). On the other hand, by the estimate of [20, Lemma 5.1], we have

$$\int_{R_2}^{+\infty} \int_{\Omega} e^{c^{\dagger} z} \left(|\nabla w|^2 - f_u(0, y) w^2 \right) dy dz$$

$$\geq \int_{R_2}^{+\infty} \int_{\Omega} e^{c^{\dagger} z} \left(\frac{c^{\dagger^2}}{4} w^2 + |\nabla_y w|^2 + f_u(0, y) w^2 \right) dy dz$$

$$\geq \left(\frac{c^{\dagger^2}}{4} + \nu_0 \right) \int_{R_2}^{+\infty} \int_{\Omega} e^{c^{\dagger} z} w^2 dy dz.$$

So, by hypothesis (H3) and (3.6), the inequality in (4.6) holds for some $K_2 > 0$ and R_2 large enough.

Let us now show that the inequality in (4.4) holds with K = 0 for all $w \in H_{c^{\dagger}}^{1}(\Sigma)$ and that equality holds if and only if w is a multiple of \bar{u}_{z} (the proof essentially follows the ideas of concentration compactness principle in the case of exponentially weighted Sobolev spaces [18, 33] and relies on the maximum principle). Indeed, denote by H[w]the left-hand side of (4.4) and let (w_n) be a minimizing sequence for H subject to the constraint $||w_n||_{L^2_{c^{\dagger}}(\Sigma)} = 1$. By coercivity of H on the constraint, ensured by hypothesis (H2), we have $w_n \to w_0$ in $H_{c^{\dagger}}^1(\Sigma)$. In fact, $w_0 \neq 0$, since otherwise $w_n \to 0$ in $L^2_{loc}(\Sigma)$, and so $\int_{-\infty}^{-R_1} \int_{\Omega} e^{c^{\dagger} z} w_n^2 dx + \int_{R_2}^{+\infty} \int_{\Omega} e^{c^{\dagger} z} w_n^2 dx \ge 1-\varepsilon$ for any $\varepsilon > 0$ and large enough n. Therefore, by (4.5) and (4.6) we would have $H[w_n] \ge K > 0$. However, this contradicts the fact (first pointed out in [1]) that \bar{u}_z is an eigenfunction associated with zero eigenvalue of the linearization of (1.5) around \bar{u} (related to the translational symmetry in the z-direction [1, 32, 2, 28]), which can be seen by differentiating (1.5) with respect to z and noting that $\bar{u}_z \in H_{c^{\dagger}}^2(\Sigma)$ by Theorem 2. In view of lower semicontinuity of H with respect to the weak convergence in

In view of lower semicontinuity of H with respect to the weak convergence in $H_{c^{\dagger}}^{1}(\Sigma)$, which follows from [20, Proposition 5.5], hypothesis (H3) and Theorem 2, we have $H[w_{0}] \leq \liminf_{n \to \infty} H[w_{n}] \leq ||\bar{u}_{z}||_{L^{2}_{c^{\dagger}}(\Sigma)}^{2} H[\bar{u}_{z}] = 0$. Then, since $w_{0} \neq 0$, the function $\bar{w} = ||w_{0}||_{L^{2}_{c^{\dagger}}(\Sigma)}^{-1}|w_{0}| \geq 0$ is a minimizer of the considered constrained minimization problem. In fact, $H[\bar{w}] = 0$, since otherwise \bar{w} must be orthogonal to \bar{u}_{z} , which is impossible due to the fact that $\bar{u}_{z} < 0$ by Theorem 2. So $H[w] \geq 0$ for all $w \in H_{c^{\dagger}}^{1}(\Sigma)$. Moreover, H[w] = 0 implies that w is a multiple of \bar{u}_{z} (compare also with [2, 28, 30]). If not, there exists a minimizer w' which is orthogonal to \bar{u}_{z} in $L^{2}_{c^{\dagger}}(\Sigma)$ and, therefore, changes sign. But |w'| is also a minimizer, hence both w' and |w'| satisfy the linearized version of (1.5) in the classical sense, thanks to hypothesis (H2) and Theorem 2. So by strong maximum principle |w'| = 0, leading to a contradiction.

To complete the proof of the lemma, suppose, to the contrary of its statement, there exists a sequence (w_n) with the properties that $||w_n||_{L^2_{c^{\dagger}}(\Sigma)} = 1$, $\int_{\Sigma} e^{c^{\dagger} z} \bar{u}_z w_n dx = 0$ and $H[w_n] \to 0$ as $n \to \infty$. Hence w_n is a minimizing sequence and converges to a non-trivial multiple of \bar{u}_z weakly in $H^1_{c^{\dagger}}(\Sigma)$. But this contradicts the orthogonality of w_n to \bar{u}_z , which is preserved in the limit as $n \to \infty$. \square

Let us note that one may naturally think that the result of Lemma 4.1 may be used to show that the minimizer \bar{u} is, in fact, a strict minimizer of $\Phi_{c^{\dagger}}$ on a suitable subset of $H^1_{c^{\dagger}}(\Sigma)$. This, however, proves difficult, since the functional $\Phi_c[u]$ is not *a priori* twice continuously differentiable in $H^1_c(\Sigma)$. We will get back to this question after Proposition 4.4 below.

Our next result shows that, if a solution to (4.2) with initial datum satisfying (2.3) with α sufficiently small is close enough to a suitable translate of \bar{u} in $L^2_{c^{\dagger}}(\Sigma)$, then it is also close in L^{∞} on some growing portion of Σ , provided that ε is small enough. More precisely, let $R : [0, \infty) \to \mathbb{R}$. For a given $\delta > 0$, we define $z_{\delta} : [0, \infty) \to \mathbb{R}$ as

$$z_{\delta}(t) := \sup\left\{z \in \mathbb{R} : \|u(\cdot, z, t) - \bar{u}(\cdot, z - R(t))\|_{L^{\infty}(\Omega)} > \delta\right\} \qquad \forall t \ge 0.$$

$$(4.7)$$

Then, the following result holds true.

PROPOSITION 4.2. There exists b > 0, such that for every $\delta > 0$ sufficiently small there exist $\alpha = \alpha(\delta) > 0$, $a = a(\delta) > 0$, $\overline{z}_0 = \overline{z}_0(\delta, u_0) \in \mathbb{R}$ and $\eta = \eta(\delta, u_0) > 0$, such that for every $z_0 \leq \overline{z}_0$ there exists $\varepsilon = \varepsilon(\delta, z_0) > 0$ such that for all T > 0

$$z_{\delta}(t) \le z_0 + a - bt \qquad \forall t \in [0, T], \tag{4.8}$$

whenever

$$|R(t)| \le \delta \quad \text{and} \quad \|u(\cdot, t) - \mathcal{T}_{R(t)}\bar{u}\|_{L^2_{c^{\dagger}}(\Sigma)} \le \eta \qquad \forall t \in [0, T],$$

$$(4.9)$$

where u_0 , u, α , ε are as in Theorem 3.

Proof. By (4.9) and the uniform Lipschitz continuity of $u(\cdot, t)$ in Σ , reasoning as in the proof of [20, Proposition 3.3(iii)] we have the following L^{∞} -estimate:

$$||u(\cdot,t) - \mathbf{T}_{R(t)}\bar{u}||_{C_{b}(\bar{\Omega}\times[z_{0},+\infty))}^{n+2} \le C\eta^{2}e^{-c^{\dagger}z_{0}},$$
(4.10)

for any $z_0 \in \mathbb{R}$, any $t \in [0, T]$ and some C > 0 depending on $||\nabla u||_{C_b(\overline{\Sigma} \times (0, +\infty))}$ (see Proposition 3.1). On the other hand, by Theorem 2 for any $\alpha > 0$ there exists $\overline{z}_0 \in \mathbb{R}$, such that

$$||\bar{u}(\cdot, z - R(t)) - v||_{C^0(\overline{\Omega})} \le \alpha \qquad \forall z \le \bar{z}_0, \ \forall t \in [0, T].$$

$$(4.11)$$

Recalling (2.3) and possibly reducing \bar{z}_0 , we can also assume that

$$u_0(\cdot, z) \ge v - 2\alpha \qquad \forall z \le \bar{z}_0. \tag{4.12}$$

Now, choosing $\eta > 0$ sufficiently small, the right-hand side of (4.10) can be bounded by α^{n+2} at $z_0 = \bar{z}_0$, so we have

$$||u(\cdot,t) - \mathcal{T}_{R(t)}\bar{u}||_{C_b(\overline{\Omega} \times [\bar{z}_0, +\infty))} \le \alpha \qquad \forall t \in [0,T].$$

$$(4.13)$$

Therefore,

$$||u(\cdot, z, t) - \bar{u}(\cdot, z - R(t))||_{C^0(\overline{\Omega})} \le \delta \qquad \forall z \ge \bar{z}_0, \ \forall t \in [0, T],$$
(4.14)

as long as $\alpha \leq \delta$, so that $z_{\delta}(t) \leq \overline{z}_0$ for all $t \in [0, T]$.

It remains to show that the inequality in (4.14) also holds for $z \in [z_0 + a - bt, \bar{z}_0]$, for some positive a and b, for small enough α and ε . We proceed by constructing explicit upper and lower barriers for (4.2) in $\Omega \times (-\infty, \bar{z}_0] \times [0, T]$.

Subsolution. First, consider the case of $\partial \Omega_{\pm} = \emptyset$, i.e., pure Neumann boundary conditions in (1.4). Then, it is straightforward to verify that by hypotheses (H1)–(H2) and (N1) the function $v_{\delta}^- = v - C\delta\tilde{\psi}_0$, where $\tilde{\psi}_0 > 0$ is an eigenfunction associated with $\tilde{\nu}_0$ in (3.8) and $C = ||\tilde{\psi}_0||_{C^0(\overline{\Omega})}^{-1}$, is the desired subsolution, provided that δ is sufficiently small.

The construction is more delicate in the presence of Dirichlet boundary conditions, since we do not wish to put any restrictions on the derivative of the initial data near the Dirichlet portion of the boundary. So, let us now assume that $\partial \Omega_{\pm} \neq \emptyset$, implying, in particular, that v < 1 in $\overline{\Omega}$. We construct a subsolution in the form of a non-negative local minimizer of E that lies sufficiently close to and below v, and vanishes identically within some small distance to $\partial \Omega_{\pm}$.

We proceed in the usual way by introducing the modified energy \tilde{E} , given by (2.7) in which V is replaced by $\tilde{V}(u, y) = -\int_0^u \tilde{f}(s, y)ds$, where \tilde{f} is obtained from f by the odd extension for u < 0 and the C^1 linear extrapolation for $|u - v| > \delta$, for some fixed $0 < \delta \ll 1$ and each $y \in \Omega$. We note that $\tilde{f}(u, y) = f(u, y)$ whenever $|u - v| \le \delta$ and $u \ge 0$. Now, by hypotheses (H2) and (N1) the energy \tilde{E} is strictly convex for all functions vanishing on $\partial \Omega_{\pm}$ and, hence, admits a unique minimizer $v_{\delta}^- \in H^1(\Omega)$ in the class of functions vanishing outside $\Omega_{\sigma} = \{y \in \Omega : \operatorname{dist}(y, \partial \Omega_{\pm}) > \sigma\}$, with $\sigma > 0$ sufficiently small. Moreover, we have $|v_{\delta}^- - v| = O(\sigma)$ in Ω_{σ} . Indeed, testing \tilde{E} with $\tilde{v} = \max(0, v - C\sigma)$ for C > 0 so large that $\tilde{v} \equiv 0$ in $\Omega \setminus \Omega_{\sigma}$ and using coercivity of \tilde{E} and the fact that v satisfies the Euler-Lagrange equation for \tilde{E} in the whole of Ω , we obtain that $||v_{\delta}^- - v||_{L^2(\Omega)} = O(\sigma)$. Therefore, by elliptic regularity theory [10] and possibly reducing σ , we have $||v_{\delta}^- - v||_{L^{\infty}(\Omega)} = O(\sigma) \le \delta$, and so v_{δ}^- satisfies the Euler-Lagrange equation for the original energy E whenever $v_{\delta}^- > 0$.

In fact, $v_{\delta}^- \geq 0$ in Ω and is strictly positive in Ω_{σ} . Indeed, by its definition the function $\tilde{V}(u, \cdot)$ is even, whenever $|u-v| \leq \delta$. Hence, if v_{δ}^- is a minimizer satisfying the latter inequality, so is $|v_{\delta}^-|$. But by uniqueness the two must be equal. On the other hand, this implies that v_{δ}^- is a critical point of the original energy E. Therefore, by strong maximum principle we have $v_{\delta}^- > 0$ in Ω_{σ} . Similarly, we must have $v_{\delta}^- < v$ in Ω , since $\bar{v} = v + a \tilde{\psi}_0$ is a strict supersolution for any $0 < a \ll 1$ and, therefore, cannot touch v_{δ}^- from above. Thus, we constructed a function v_{δ}^- which is a non-negative subsolution of the Euler-Lagrange equation for E, and $0 \leq v_{\delta}^- \leq v$. In particular, by construction

$$v - \delta \le v_{\delta}^{-} \le \max(0, v - 2\alpha), \tag{4.15}$$

in Ω , for α sufficiently small, depending only on δ . Finally, extending this function to $\Sigma \times [0,T]$ by defining $u^-(y,z,t) := v_{\delta}^-(y)$, we obtain a subsolution on the desired domain.

Supersolution. Let $\sigma > 0$ be sufficiently small. By the same type of argument as in the construction of v_{δ}^- above, there exists a local minimizer v_{δ}^+ of E, such that $v_{\delta}^+(y) = \sigma$ for all $y \in \partial \Omega_{\pm}$, and we have $v + \beta \leq v_{\delta}^+ \leq v + \frac{1}{4}\delta$, for some $\beta > 0$.

Now, let $c \in (c_v^{\dagger}, c^{\dagger})$, and consider $\Psi_c^{v_{\delta}^{\dagger}}$ defined in (3.9) with v_{δ}^{+} in place of v. Then, by an extension of the argument of Lemma 3.5 it is not difficult to see that there exists a minimizer \bar{h} of $\Psi_c^{v_{\delta}^{\dagger}}$ in the set

$$X := \left\{ h \in H^1_c(\Sigma) : 0 \le h \le 1 - v^+_{\delta}, \ h = 1 - v^+_{\delta} \text{ in } \Omega \times (-\infty, 0], \\ h(y, z) = (1 - v^+_{\delta}(y))\eta(z) \text{ for } (y, z) \in \partial\Omega_{\pm} \times \mathbb{R} \right\},$$

where $\eta \in C^{\infty}(\mathbb{R})$ is a cutoff function with the property that $\eta(z) = 1$ for all z < 0 and $\eta(z) = 0$ for all z > 1. Indeed, semicontinuity and coercivity of $\Psi_c^{v_{\delta}^+}$ only depend on the behavior of the functional for large values of z. Since v_{δ}^+ is still a local minimizer of

E, the functional $\Psi_c^{v_{\delta}^{\dagger}}$ is lower semicontinuous by [20, Proposition 5.5]. Furthermore, by hypothesis (H2) and Taylor formula

$$\begin{split} \Psi_c^{v_{\delta}^+}[h] &= \Psi_c^v[h] + \int_{\Sigma} \int_0^h e^{cz} \Big(f_u(v+s,y) - f_u(v_{\delta}^+ + s,y) \Big) (h-s) ds \, dx \\ &\geq \Psi_c^v[h] - C ||v_{\delta}^+ - v||_{L^{\infty}(\Sigma)}^{\gamma} \int_{\Sigma} \int_0^h e^{cz} (h-s) ds \, dx \\ &\geq \Psi_c^v[h] - C \delta^{\gamma} ||h||_{L^2(\Sigma)}^2, \end{split}$$

for some C > 0, implying coercivity of $\Psi_c^{v_{\delta}^+}$ for small enough δ by the argument of [20, Proposition 6.9]. So the minimizer \bar{h} of $\Psi_c^{v_{\delta}^+}$ exists, and $\bar{h}(\cdot, z) \to 0$ uniformly in Ω , as $z \to +\infty$ (indeed, the convergence is exponential by [20, Proposition 3.3(iii)]). Therefore, there exists a > 0 such that $\bar{h}(\cdot, z) \leq \frac{1}{4}\delta$ for all $z \geq a$.

We finally let $u^+(y, z, t) := v_{\delta}^+(y) + \bar{h}(y, z - z_0 + bt)$, with $b := c^{\dagger} - c > 0$, which is a supersolution for (4.2) on $\Sigma \times [0, T]$. Notice that

$$u^+(\cdot, 0) = 1 \qquad \text{on } \Omega \times (-\infty, z_0], \tag{4.16}$$

and

$$u^+(\cdot, z, t) \le v + \frac{\delta}{2}$$
 on Ω $\forall t \ge 0 \ \forall z \ge z_0 + a - bt.$ (4.17)

Comparison. From (4.12) and (4.15) for α small enough we have

$$u^{-}(\cdot, 0) \le u_0$$
 on $\Omega \times (-\infty, \bar{z}_0]$.

Also, by (4.11), (4.13) and (4.15) for η small enough we have

$$u^{-}(\cdot, \bar{z}_0, t) \le u(\cdot, \bar{z}_0, t) \quad \text{on } \Omega \quad \forall t \in [0, T].$$
(4.18)

Therefore, by parabolic comparison principle [26] we obtain

$$u^- \le u$$
 on $\Omega \times (-\infty, \bar{z}_0] \times [0, T]$. (4.19)

In particular, by (4.15) and the fact that by Theorem 2 we have $\bar{u}(\cdot, z) < v$ for every $z \in \mathbb{R}$, it follows that

$$u(\cdot, z, t) \ge \bar{u}(\cdot, z - R(t)) - \delta \qquad \forall z \le \bar{z}_0, \ \forall t \ge 0.$$

$$(4.20)$$

On the other hand, in view of (4.16), the fact that $u^+ \ge v + \beta$, (4.10) with η replaced by ε at t = 0 due to (4.1), and the fact that $|R(0)| \le \delta$, for every z_0 it is possible to choose ε small enough, so that

$$u_0 \le u^+(\cdot, 0) \qquad \text{on } \Sigma. \tag{4.21}$$

Then, by parabolic comparison principle we have

$$u \le u^+$$
 on $\Sigma \times [0, +\infty),$ (4.22)

and, possibly reducing \bar{z}_0 to ensure that $\bar{u}(\cdot, \bar{z}_0 + a + \delta) \ge v - \frac{1}{2}\delta$, in view of monotonicity of $\bar{u}(\cdot, z)$ by Theorem 2, for every $z_0 \le \bar{z}_0$ we obtain

$$u(\cdot, z, t) \le \bar{u}(\cdot, z - R(t)) + \delta \qquad \forall z \in [z_0 + a - bt, \bar{z}_0 + a] \quad \forall t \ge 0.$$

$$(4.23)$$

Finally, combining (4.20) with (4.23) and (4.14), we get (4.8).

We now prove a technical lemma that will be useful in the proof of Proposition 4.4.

LEMMA 4.3. There exist $0 < C_1 < C_2$, such that for all $|R| \leq 1$ we have

$$C_1|R| \le ||\mathbf{T}_R \bar{u} - \bar{u}||_{L^2_{c^{\dagger}}(\Sigma)} \le C_2|R|.$$
(4.24)

Furthermore,

$$||\mathbf{T}_R \bar{u} - \bar{u}||_{L^2_{c^{\dagger}}(\Sigma)} \ge C_1,$$
 (4.25)

for all $|R| \geq 1$.

Proof. Let us first prove the upper bound. Notice that, thanks to Theorem 2, the functions \bar{u} belongs to $H^1_{c^{\dagger}}(\Sigma)$, hence in particular the map $\eta \mapsto T_{\eta} \bar{u}$ defines a differentiable curve in $L^2_{c^{\dagger}}(\Sigma)$. A direct computation then gives

$$\begin{aligned} ||\mathbf{T}_{R}\bar{u} - \bar{u}||_{L^{2}_{c^{\dagger}}(\Sigma)} &= \left\| \int_{0}^{R} \mathbf{T}_{\eta}\bar{u}_{z} \, d\eta \right\|_{L^{2}_{c^{\dagger}}(\Sigma)} \leq \int_{-|R|}^{|R|} ||\mathbf{T}_{\eta}\bar{u}_{z}||_{L^{2}_{c^{\dagger}}(\Sigma)} \, d\eta \\ &\leq \left(\int_{-|R|}^{|R|} e^{\frac{c^{\dagger}}{2}\eta} \, d\eta \right) ||\bar{u}_{z}||_{L^{2}_{c^{\dagger}}(\Sigma)} \leq C|R|, \end{aligned}$$

where we used the identity $||T_{\eta}\bar{u}_z||_{L^2_{c^{\dagger}}(\Sigma)} = e^{\frac{c^{\dagger}\eta}{2}}||\bar{u}_z||_{L^2_{c^{\dagger}}(\Sigma)}.$

To obtain the lower bound in (4.24), we observe that for any $\Sigma_0 \subseteq \Sigma$ compact, we have

$$||\mathbf{T}_{R}\bar{u} - \bar{u}||_{L^{2}_{c^{\dagger}}(\Sigma)}^{2} \geq \int_{\Sigma_{0}} e^{c^{\dagger}z} (\bar{u}(y, z - R) - \bar{u}(y, z))^{2} dx$$
$$= R^{2} \int_{\Sigma_{0}} e^{c^{\dagger}z} \bar{u}_{z}^{2}(y, z - \tilde{R}(y, z)) dx, \qquad (4.26)$$

for some $0 < |\tilde{R}(y,z)| < |R|$. The lower bound then follows from the fact that $\bar{u}_z < 0$ in Σ and, hence, $|u_z(y, z - \tilde{R}(y, z))|$ is bounded away from zero in Σ_0 , as long as $|R| \leq 1$. Finally, to get (4.25) we observe that $||T_R \bar{u} - \bar{u}||^2_{L^2_{c^{\dagger}}(\Sigma)}$ is monotonically increasing in |R|. \square

We now look for a suitable translation of \bar{u} which serves as the best approximation, in some sense, to the solution of (4.2). For a given $u \in H^1_{c^{\dagger}}(\Sigma)$ and $R \in \mathbb{R}$, we define the function h as:

$$h(u,R) := \frac{1}{2} \int_{\Sigma} e^{c^{\dagger} z} (u(y,z) - \bar{u}(y,z-R))^2 dx \ge 0.$$
(4.27)

In the following proposition, we show that the optimal approximation to u can be naturally introduced by minimizing h in (4.27) with respect to R.

PROPOSITION 4.4. For any $\delta > 0$ sufficiently small there exists $\varepsilon > 0$ such that, for any $u \in H^1_{c^{\dagger}}(\Sigma)$ satisfying $||u - \bar{u}||_{L^2_{c^{\dagger}}(\Sigma)} \leq \varepsilon$ the function $h(u, \cdot)$ attains its global minimum. Furthermore, this minimum is unique, is contained in $(-\delta, \delta)$ and there are no other critical points in $(-\delta, \delta)$.

14

Proof. First of all, observe that $\eta \mapsto \bar{u}(y, z - \eta)$ is a twice differentiable curve in $L^2_{c^{\dagger}}$, thanks to Theorem 2. By assumption

$$\inf_{R \in \mathbb{R}} h(u, R) \le h(u, 0) \le \varepsilon^2.$$
(4.28)

Furthermore, from the lower bound in Lemma 4.3 we get that

$$C\min\{1, |R|\} \le ||\mathbf{T}_R \bar{u} - \bar{u}||_{L^2_{c^{\dagger}}(\Sigma)}$$
$$\le ||u - \bar{u}||_{L^2_{c^{\dagger}}(\Sigma)} + \sqrt{2h(u, R)} \le \varepsilon + \sqrt{2h(u, R)},$$
(4.29)

for some C > 0. Therefore, by continuity of $h(u, \cdot)$ its minimum is attained and lies in $(-\delta, \delta)$ for any $\delta > 0$, provided that ε is sufficiently small.

We now calculate the first and the second derivative of $h(u, \cdot)$ with respect to R:

$$h'(u,R) = \int_{\Sigma} e^{c^{\dagger} z} (u(y,z) - \bar{u}(y,z-R)) \bar{u}_z(y,z-R) \, dx, \qquad (4.30)$$

$$h''(u,R) = c^{\dagger}h'(u,R) + \int_{\Sigma} e^{c^{\dagger}z} u_z(y,z)\bar{u}_z(y,z-R) \, dx, \qquad (4.31)$$

where from now on the prime denotes the derivative with respect to R. Recalling Lemma 4.3, for all $|R| \le \delta \le 1$ we have

$$|h'(u,R)| \leq C ||u - \mathcal{T}_R \bar{u}||_{L^2_{c^{\dagger}}(\Sigma)}$$

$$\leq C \left(||u - \bar{u}||_{L^2_{c^{\dagger}}(\Sigma)} + ||\bar{u} - \mathcal{T}_R \bar{u}||_{L^2_{c^{\dagger}}(\Sigma)} \right)$$

$$\leq C \left(\varepsilon + |R|\right), \qquad (4.32)$$

for some C > 0. Now, observe that upon integration by parts we have

$$\int_{\Sigma} e^{c^{\dagger} z} \bar{u}_{z}(y, z - R) (u_{z}(y, z) - \bar{u}_{z}(y, z - R)) dx$$

= $-\int_{\Sigma} e^{c^{\dagger} z} (\bar{u}_{zz}(y, z - R) + c^{\dagger} \bar{u}_{z}(y, z - R)) (u(y, z) - \bar{u}(y, z - R)) dx.$

Therefore, since $\bar{u}_z \in H^1_{c^{\dagger}}(\Sigma)$ by Theorem 2, applying Cauchy-Schwarz inequality we obtain

$$\left| \int_{\Sigma} e^{c^{\dagger} z} \bar{u}_{z}(y, z - R) (u_{z}(y, z) - \bar{u}_{z}(y, z - R)) \, dx \right| \\ \leq C e^{c^{\dagger} R/2} ||u - \mathcal{T}_{R} \bar{u}||_{L^{2}_{a^{\dagger}}(\Sigma)}, \tag{4.33}$$

for some C > 0. Applying this estimate to (4.31) and combining it with the estimates in (4.32), we obtain

$$h''(u,R) \ge e^{c^{\dagger}R} ||\bar{u}_z||^2_{L^2_{c^{\dagger}}(\Sigma)} - C(\varepsilon + |R|),$$
(4.34)

for some constant C > 0. This implies that $h''(u, R) \ge M$ for some M > 0 and all $|R| \le \delta$, provided that δ and ε are small enough. Hence $h(u, \cdot)$ is a strictly convex function on $[-\delta, \delta]$, and the minimum of $h(u, \cdot)$ is the unique critical point in $(-\delta, \delta)$. \Box

Recalling the comment following Lemma 4.1, we can now formulate a nonlinear analog of the result of that lemma.

REMARK 4.5. Suppose that u is sufficiently close to \bar{u} in $L^2_{c^{\dagger}}(\Sigma) \cap L^{\infty}(\Sigma)$. Then by Proposition 4.4 there exists $R_0 \in \mathbb{R}$, such that the function h(u, R) in (4.27) is minimized with respect to R at $R = R_0$. Therefore, we have that $u - T_{R_0}\bar{u}$ is orthogonal to $T_{R_0}\bar{u}$ in $L^2_{c^{\dagger}}(\Sigma)$, and so by Lemma 4.1, hypothesis (H2) and the minimizing property of \bar{u} we have $\Phi_{c^{\dagger}}[u] \geq \frac{1}{2}K||u - T_{R_0}\bar{u}||^2_{L^2_{c^{\dagger}}(\Sigma)}$, where K > 0 is as in Lemma 4.1. Hence \bar{u} is, in fact, a strict local minimizer of $\Phi_{c^{\dagger}}$ in the above sense.

We now conclude the proof of Theorem 3.

Proof of Theorem 3. Let $\delta > 0$ be sufficiently small, so that Proposition 4.4 applies with $u = u_0$ and all $0 < \varepsilon \leq \varepsilon_0$, for some $\varepsilon_0 > 0$. Then by Propositions 3.1 and 4.4 there exists $T_0 > 0$, such that there exists a minimizer R(t) of $h(u(\cdot, t), R)$ in R for each $t \in [0, T_0]$. Furthermore, R(t) is the unique critical point of $h(u(\cdot, t), R)$ in $(-\delta, \delta)$. In fact, R(t) is a continuously differentiable function of t on $[0, T_0]$. Indeed, since R(t) minimizes $h(u(\cdot, t), R)$ in R, we have

$$h'(u(\cdot,t),R) = \int_{\Sigma} e^{c^{\dagger}z} (u(y,z,t) - \bar{u}(y,z-R))\bar{u}_z(y,z-R) \, dx = 0, \quad (4.35)$$

whenever R = R(t). In view of the continuity of $u_t(\cdot, t)$ in $L^2_{c^{\dagger}}(\Sigma)$ guaranteed by Proposition 3.1, as well as Theorem 2 and Lemma 4.3, the function in (4.35) is continuously differentiable in R and t in some small neighborhood of the origin. Then, arguing as in Proposition 4.4 one can see that $h''(u(\cdot, t), R) > 0$ there, so we can apply the implicit function theorem to (4.35). Furthermore, after some algebra we obtain

$$\frac{dR(t)}{dt} = -\frac{\int_{\Sigma} e^{c^{\dagger} z} u_t(y, z, t) \bar{u}_z(y, z - R(t)) \, dx}{\int_{\Sigma} e^{c^{\dagger} z} u_z(y, z, t) \bar{u}_z(y, z - R(t)) \, dx}.$$
(4.36)

For $t \in (0, T_0]$ and u solving (4.2) we define

$$w(y, z, t) := u(y, z, t) - \bar{u}(y, z - R(t)).$$
(4.37)

The function w satisfies the equation

$$w_t = \Delta w + c^{\dagger} w_z + \frac{dR}{dt} \bar{u}_z + f_u(\tilde{u}, y)w, \qquad (4.38)$$

for some \tilde{u} , with $|\tilde{u} - T_{R(t)}\bar{u}| \leq |w|$. Also, by construction we have

$$\int_{\Sigma} e^{c^{\dagger} z} w(y, z, t) \bar{u}_z(y, z - R(t)) \, dx = 0.$$
(4.39)

We now introduce

$$m(t) := \int_{\Sigma} e^{c^{\dagger} z} w^2(x, t) \, dx \qquad t \ge 0.$$

so that $m(0) \leq \varepsilon^2$. Multiplying (4.38) with $e^{c^{\dagger}z}w$ and integrating over Σ , we obtain

$$\frac{dm(t)}{dt} = -2\int_{\Sigma} e^{c^{\dagger}z} \left(|\nabla w|^2 - f_u(\tilde{u}, y)w^2 \right) dx \tag{4.40}$$

where we used (4.39) to erase the term multiplying dR/dt. By Lemma 4.1 we have

$$\int_{\Sigma} e^{c^{\dagger} z} \left(|\nabla w|^2 - f_u(\tilde{u}, y) w^2 \right) dx = \int_{\Sigma} e^{c^{\dagger} z} \left(|\nabla w|^2 - f_u(\bar{u}, y) w^2 \right) dx + \int_{\Sigma} e^{c^{\dagger} z} \left(f_u(\bar{u}, y) - f_u(\tilde{u}, y) \right) w^2 dx \geq K m(t) + \int_{\Sigma} e^{c^{\dagger} z} \left(f_u(\bar{u}, y) - f_u(\tilde{u}, y) \right) w^2 dx.$$
(4.41)

Possibly reducing δ and recalling assumption (H2), we have

$$\int_{\{|w|<\delta\}} e^{c^{\dagger}z} |f_u(\bar{u}, y) - f_u(\tilde{u}, y)| \, w^2 dx \le \frac{K}{2} m(t),$$

which, combined with (4.41), gives

$$\begin{split} \int_{\Sigma} e^{c^{\dagger} z} \left(|\nabla w|^2 - f_u(\tilde{u}, y) w^2 \right) dx &\geq \frac{K}{2} m(t) - \int_{\{|w| \geq \delta\}} e^{c^{\dagger} z} \left| f_u(\bar{u}, y) - f_u(\tilde{u}, y) \right| w^2 dx \\ &\geq \frac{K}{2} m(t) - \int_{-\infty}^{z_{\delta}(t)} \int_{\Omega} e^{c^{\dagger} z} \left| f_u(\bar{u}, y) - f_u(\tilde{u}, y) \right| w^2 dy \, dz \\ &\geq \frac{K}{2} m(t) - C e^{c^{\dagger} z_{\delta}(t)}, \end{split}$$

for some C > 0, where $z_{\delta}(t)$ is defined in (4.7).

We can now apply Proposition 4.2 with $z_0 = \frac{4}{c^{\dagger}} \log \omega$, which yields some $\varepsilon > 0$ and $\eta > 0$, with η independent of ω , and $T \in (0, T_0]$ depending on η . We then get

$$\int_{\Sigma} e^{c^{\dagger} z} \left(|\nabla w|^2 - f_u(\tilde{u}, y) w^2 \right) dx \ge \frac{K}{2} m(t) - C e^{c^{\dagger}(z_0 - bt)} \ge \frac{K}{2} m(t) - C \omega^4 e^{-c^{\dagger} bt}$$
(4.42)

for some b > 0 and C > 0 and all $t \in [0, T]$. From (4.40) and (4.42) we, therefore, obtain

$$\frac{dm(t)}{dt} \le -Km(t) + 2C\omega^4 e^{-c^{\dagger}bt} \qquad \forall t \in [0,T],$$
(4.43)

which gives

$$||w(\cdot,t)||_{L^2_{\sigma^{\dagger}}(\Sigma)} \le M\omega^2 e^{-\sigma t} \qquad \forall t \in [0,T],$$

$$(4.44)$$

for some $\sigma > 0$ and M > 0, provided that ε is small enough.

To estimate the behavior of R(t), we substitute $u_t = w_t - \bar{u}_z dR/dt$ into (4.36) and take into account (4.38) and (1.5) differentiated in z, noting that $\bar{u}_z \in H_c^2(\Sigma)$ by Theorem 2. After a few integrations by parts we obtain

$$\frac{dR(t)}{dt} = \frac{\int_{\Sigma} e^{c^{\dagger} z} (f_u(\bar{u}(y, z - R(t)), y) - f_u(\tilde{u}, y)) w(y, z, t) \bar{u}_z(y, z - R(t)) dx}{\int_{\Sigma} e^{c^{\dagger} z} (\bar{u}_z(y, z - R(t)) + w_z(y, z, t)) \bar{u}_z(y, z - R(t)) dx}.$$
 (4.45)

By the same argument as the one leading to (4.33), we have

$$\left| \int_{\Sigma} e^{c^{\dagger} z} \bar{u}_{z}(y, z - R(t)) w_{z}(y, z, t) \, dx \right| \le C e^{c^{\dagger} R(t)/2} ||w(\cdot, t)||_{L^{2}_{c^{\dagger}}(\Sigma)}.$$
(4.46)

With hypothesis (H2), this leads to the following estimate for dR/dt:

$$\left|\frac{dR(t)}{dt}\right| \le \frac{C||w(\cdot,t)||_{L^2_{c^{\dagger}}(\Sigma)}}{\widetilde{C}e^{c^{\dagger}R(t)/2} - ||w(\cdot,t)||_{L^2_{c^{\dagger}}(\Sigma)}}$$
(4.47)

for some constants $C, \tilde{C} > 0$, provided that ω is so small that by (4.44) the denominator in (4.47) is positive for all $t \in [0, T]$. Then we have

$$\left. \frac{dR(t)}{dt} \right| \le C\omega^2 e^{-\sigma t} \qquad \forall t \in [0, T], \tag{4.48}$$

and hence

$$|R(t)| \le \widetilde{M}\omega^2 \le \omega \qquad \forall t \in [0, T], \tag{4.49}$$

for some $\widetilde{M} > 0$ and ω small enough. Moreover, since η and δ are independent of ω , (4.44) also implies that (4.9) holds uniformly in T for ω small enough, whence $T = T_0$. Indeed, if $T_1 < T_0$ is the maximum value of T for which Proposition 4.4 can be applied, then by (4.44) the left-hand side of (4.9) is bounded by $\eta/2$ at $t = T_1$, provided that ω is sufficiently small. Therefore, by continuity of $w(\cdot, t)$ in $L^2_{c^{\dagger}}(\Sigma)$ guaranteed by Proposition 3.1, the inequality in (4.9) also holds for some interval beyond T_1 , contradicting the maximality of T_1 .

Moreover, by (4.49) the function R(t) is in fact defined and continuously differentiable for all $t \ge 0$. Indeed, let us take T_0 to be the largest possible value for which $||u(\cdot,t)-\bar{u}||_{L^2_{c^{\dagger}}(\Sigma)} \le \varepsilon_0$ for all $t \in [0,T_0]$, so that Proposition 4.4 still applies. In view of Lemma 4.3, (4.44) and (4.49), we have

$$\begin{aligned} ||u(\cdot,t) - \bar{u}||_{L^{2}_{c^{\dagger}}(\Sigma)} &\leq ||u(\cdot,t) - \mathcal{T}_{R(t)}\bar{u}||_{L^{2}_{c^{\dagger}}(\Sigma)} \\ &+ ||\bar{u} - \mathcal{T}_{R(t)}\bar{u}||_{L^{2}_{c^{\dagger}}(\Sigma)} \\ &\leq M\omega^{2} \quad \forall t \in [0,T_{0}], \end{aligned}$$

$$(4.50)$$

for some M > 0. Therefore, choosing ω so small that the right-hand side of (4.50) is bounded by $\frac{1}{2}\varepsilon_0$ and, once again, taking into account continuity of $w(\cdot, t)$ in $L^2_{c^{\dagger}}(\Sigma)$, we can then make sure that the assumptions of Proposition 4.4 are satisfied on some interval beyond T_0 , contradicting maximality of T_0 . We thus proved that we can take an arbitrarily large $T_0 > 0$ in all the arguments above.

Finally, using (4.44) and (4.47) again and keeping in mind that by (4.49) the denominator in (4.47) is bounded away from zero, we finally obtain that the limit $R_{\infty} := \lim_{t \to +\infty} R(t)$ exists, and recalling Lemma 4.3 we have

$$||u(\cdot,t) - \mathcal{T}_{R_{\infty}}\bar{u}||_{L^{2}_{\star}(\Sigma)} \le \omega e^{-\sigma t} \qquad \forall t \ge 0,$$

$$(4.51)$$

for some $\sigma > 0$, provided that ω is small enough, yielding the thesis of the theorem.

5. Proof of the main result. We will prove Theorem 1 in the reference frame moving with speed c^{\dagger} , that is, we will prove that if u is the solution of (4.2) with the initial datum satisfying the assumptions of Theorem 1, then it converges in $H_{c^{\dagger}}^2(\Sigma)$ to $T_{R_{\infty}}\bar{u}$ for some R_{∞} as $t \to \infty$. The result then follows by noting that $T_{c^{\dagger}t}u$ solves (1.3) with the same initial condition, upon applying $T_{-R_{\infty}}$.

From now on, u always refers to the solution of (4.2). We divide the proof into five steps.

Step 1. We begin by constructing an appropriate pair of barrier solutions of (4.2) to ensure that the solution of the initial value problem for (4.2) does not move too far towards the ends of the cylinder. The barriers are obtained by considering the solutions \bar{u}^{\pm} of (4.2) with the initial data

$$\bar{u}_0^-(y,z) = \min\{u_0(y,z-R), \bar{u}(y,z)\},\tag{5.1}$$

$$\bar{u}_0^+(y,z) = \max\{u_0(y,z+R), \bar{u}(y,z)\},\tag{5.2}$$

where R > 0 is so big that both \bar{u}_0^{\pm} satisfy the assumptions of Theorem 3, provided that α in (2.3) is small enough. Indeed, by definition and (3.6) the assumption in (2.3) is satisfied for both \bar{u}_0^{\pm} . Moreover, for $\bar{u}_0^+ - \bar{u} = \max(T_{-R}u_0 - \bar{u}, 0)$ we have

$$0 \le \bar{u}_0^+ - \bar{u} \le \mathcal{T}_{-R} u_0 \to 0 \quad \text{in } L^2_{c^{\dagger}}(\Sigma) \quad \text{as } R \to +\infty,$$
(5.3)

so $\bar{u}_0^+ \to \bar{u}$ in $L^2_{c^{\dagger}}(\Sigma)$ as $R \to +\infty$. By a similar argument for $\bar{u} - \bar{u}_0^- = \max(\bar{u} - T_R u_0, 0)$ we have

$$0 \le \bar{u} - \bar{u}_0^- \le \bar{u} \to 0 \quad \text{in } L^2_{c^{\dagger}}(\Omega \times (M, +\infty)) \quad \text{as } M \to +\infty, \tag{5.4}$$

uniformly in R. At the same time, by boundedness of \bar{u}_0^- and \bar{u} we have $||T_R u_0 - \bar{u}||_{L^2_{c^{\dagger}}(\Omega \times (-\infty, -M))} \to 0$ as $M \to +\infty$, again, uniformly in R. Finally, in view of (2.3), (3.6) and the Hopf lemma, for every M > 0 and R > 0 large enough we have $|\{x \in \Omega \times (-M, M) : \bar{u}_0^-(x) < \bar{u}(x)\}| \leq C\alpha$, with some C = C(M) > 0, for small enough α . Therefore, it is possible to choose M large enough, then R large enough, and then α small enough, so that $||\bar{u} - \bar{u}_0^-||_{L^2_{c^{\dagger}}(\Sigma)}$ can be made as small as desired. Note that both functions \bar{u}_0^{\pm} obtained above satisfy (2.3) uniformly in R.

We now claim that $\bar{u}^{\pm}(y, z \mp R, t)$, i.e., the solutions of (4.2) with initial data $\bar{u}_0^{\pm}(y, z \mp R)$, are the appropriate barrier solutions. Indeed, by construction the initial data u_0 is sandwiched between $\bar{u}^{\pm}(y, z \mp R, t)$ at t = 0, hence by parabolic comparison principle [26] the solution of (4.2) will remain so for all times. By Theorem 3 we know that there exist R_{∞}^{\pm} such that

$$||\bar{u}^{\pm}(y, z \mp R, t) - \bar{u}(y, z \mp R_{\infty}^{\pm})||_{L^{2}_{c^{\dagger}}(\Sigma)} \le e^{-\sigma t},$$
 (5.5)

for some $\sigma > 0$ and any $z_0 \in \mathbb{R}$, provided that α is sufficiently small and R is sufficiently large.

Step 2. We now use the functional $\Phi_{c^{\dagger}}$ as a Lyapunov functional to establish existence of a sequence $t_n \to +\infty$ on which $u(\cdot, t_n)$ converges to a translate of \bar{u} . Indeed, multiplying (4.2) by a test function $\varphi \in C_0^{\infty}(\mathbb{R}^n)$ vanishing on $\partial \Sigma_{\pm}$ and integrating over Σ , we can write (4.2) in the weak form as

$$\int_{\Sigma} e^{c^{\dagger} z} \varphi \, u_t \, dx = -\int_{\Sigma} e^{c^{\dagger} z} \left(\nabla u \cdot \nabla \varphi - f(u, y) \varphi \right) dx, \tag{5.6}$$

where the integral in the right-hand side is the Gâteaux derivative of $\Phi_{c^{\dagger}}$ at $u(\cdot, t)$ in the direction of φ . Therefore, (4.2) is the gradient flow generated by $\Phi_{c^{\dagger}}$ in $L^2_{c^{\dagger}}(\Sigma)$, and in view of (3.3) for all $t_2 \geq t_1 > 0$, we have

$$\Phi_{c^{\dagger}}[u(\cdot,t_1)] - \Phi_{c^{\dagger}}[u(\cdot,t_2)] = \int_{t_1}^{t_2} \|u_t(\cdot,t)\|_{L^2_{c^{\dagger}}(\Sigma)}^2 dt.$$
(5.7)

Letting $t_2 \to +\infty$ and recalling that $\Phi_{c^{\dagger}}[u]$ is bounded below by Theorem 2, from (5.7) it follows that there exists a sequence $t_n \to +\infty$ such that

$$\lim_{n \to +\infty} \|u_t(\cdot, t_n)\|_{L^2_{c^{\dagger}}(\Sigma)} = 0.$$
(5.8)

Also note that, since $0 \leq u(y, z, t) \leq \overline{u}^+(y, z - R, t)$ for all $(y, z) \in \Sigma$ and $t \geq 0$, and $\overline{u}^+(\cdot, t)$ is uniformly bounded in $L^2_{c^{\dagger}}(\Sigma)$ by Theorem 3, in view of hypotheses (H1)–(H2) we have that

$$\|u\|_{H^{1}_{c^{\dagger}}(\Sigma)}^{2} \leq 2\Phi_{c^{\dagger}}[u] + C\|u\|_{L^{2}_{c^{\dagger}}(\Sigma)}^{2} \leq M(t_{0}) \qquad \forall t \geq t_{0} > 0$$
(5.9)

for some constants $C, M(t_0) > 0$.

From (5.8) and (5.9), up to a possible subsequence, we can pass to the limit in (5.6) and get that $u(\cdot, t_n)$ converges to a critical point of $\Phi_{c^{\dagger}}$ weakly in $H^1_{c^{\dagger}}(\Sigma)$. In fact, the limit must be a non-trivial critical point u_{∞} of $\Phi_{c^{\dagger}}$, in view of Step 1, hence a translate of \bar{u} by [20, Propositions 3.2 and 3.5] and Theorem 2.

Step 3. We now prove that $u(\cdot, t_n) \to u_{\infty}$ in $L^2_{c^{\dagger}}(\Sigma)$. Notice first that since both u and u_{∞} are uniformly bounded, for a given $\varepsilon > 0$ we can find M such that

$$\|u(\cdot, t_n) - u_{\infty}\|_{L^2_{c^{\dagger}}(\Omega \times (-\infty, -M])} \le \varepsilon.$$
(5.10)

Moreover, since $u(y, z, t) \leq \overline{u}^+(y, z - R, t)$, from (5.5) it also follows that

$$\begin{aligned} \|u(\cdot,t_{n})\|_{L^{2}_{c^{\dagger}}(\Omega\times(M,+\infty))} &\leq \|\bar{u}^{+}(\cdot,t_{n})\|_{L^{2}_{c^{\dagger}}(\Omega\times(M,+\infty))} \\ &\leq \|\bar{u}(y,z-R^{+}_{\infty})\|_{L^{2}_{c^{\dagger}}(\Omega\times(M,+\infty))} \\ &+ \|\bar{u}^{+}(\cdot,t_{n}) - \bar{u}(y,z-R^{+}_{\infty})\|_{L^{2}_{c^{\dagger}}(\Sigma)} \\ &\leq \varepsilon, \end{aligned}$$
(5.11)

for M big enough and all $n \geq N$, for some $N = N(M) \in \mathbb{N}$. Recalling that $H^1_{c^{\dagger}}(\Sigma)$ compactly embeds into $L^2_{c^{\dagger}}(\Omega \times (-M, M))$, from Proposition 3.1, (5.10) and (5.11) we obtain that

$$u(\cdot, t_n) \to u_{\infty}$$
 in $L^2_{c^{\dagger}}(\Sigma)$.

Step 4. Take n big enough so that

$$\|u(\cdot, t_n) - u_{\infty}\|_{L^2_{c^{\dagger}}(\Sigma)} \le \varepsilon \qquad \forall t \ge t_n \,, \tag{5.12}$$

where ε is the same as the one corresponding to $\omega = 1$ in Theorem 3. On the other hand, for every $\alpha' > 0$ it is possible to choose $\delta \leq \alpha'$ in Proposition 4.2, such that the subsolution u^- constructed there satisfies $u^-(\cdot, z, t) \geq v - \alpha'$ for all $z \leq \overline{z}_0$, with some $\overline{z}_0 \in \mathbb{R}$ independent of ε and R in the definition of \overline{u}^- , and all $t \geq 0$, if α is sufficiently small. Therefore, we have $\overline{u}^- \geq u^-$ in $\Omega \times (-\infty, \overline{z}_0] \times [0, +\infty)$, and since $\overline{u}^- \leq u$ for all $t \geq 0$, the same inequality holds for u. So we can apply Theorem 3 to $u(\cdot, t_n)$ in place of u_0 (also applying suitable translations in z and t), and obtain

$$\|u(\cdot,t) - u_{\infty}\|_{L^{2}_{*}(\Sigma)} \le e^{-\sigma(t-t_{n})}$$
(5.13)

for some $\sigma > 0$ independent of u_0 and all $t \ge t_n$.

Step 5. We now demonstrate that the exponential convergence of (5.13) also holds in spaces of higher regularity. We first show this for $H^1_{c^{\dagger}}(\Sigma)$, and then for $H^2_{c^{\dagger}}(\Sigma)$. In the following, we denote by $\mathcal{A} : \mathcal{D}(\mathcal{A}) \to L^2_{c^{\dagger}}(\Sigma)$ the sectorial operator $\mathcal{A} = \Delta + c^{\dagger}\partial_z$, with domain $\mathcal{D}(\mathcal{A}) = H^2_{c^{\dagger}}(\Sigma)$ dense in $L^2_{c^{\dagger}}(\Sigma)$ (see also [24, 21]).

Letting $w(\cdot, t) := u(\cdot, t) - u_{\infty}$, we have

$$w_t = \mathcal{A}w + g(x, t)w, \tag{5.14}$$

where $g(y, z, t) = f_u(\tilde{u}(y, z, t), y)$ for some \tilde{u} such that $|\tilde{u} - u_\infty| \leq |w|$, i.e., g is such that $|g| \leq C$, for some C > 0. As a consequence, by parabolic regularity theory [34, Chapter 15] (see also [21, Proposition 2.1.1 and Theorem 3.1.1]) and recalling (5.13), for all $t \geq 1$ we have

$$\|u(\cdot,t) - u_{\infty}\|_{H^{1}_{c^{\dagger}}(\Sigma)} \leq C \left(\|w(\cdot,t-1)\|_{L^{2}_{c^{\dagger}}(\Sigma)} + \int_{t-1}^{t} \frac{\|w(\cdot,s)\|_{L^{2}_{c^{\dagger}}(\Sigma)}}{\sqrt{t-s}} \, ds \right)$$

$$\leq C e^{-\sigma t},$$
(5.15)

for some C > 0. In particular, from [20, Proposition 3.2], (5.15), the minimizing property of u_{∞} and hypothesis (H2) we get

$$\begin{split} \Phi_{c^{\dagger}}[u(\cdot,t)] &= \Phi_{c^{\dagger}}[u(\cdot,t)] - \Phi_{c^{\dagger}}[u_{\infty}] \\ &= \int_{\Sigma} e^{c^{\dagger}z} \left(\frac{1}{2} |\nabla w|^2 + V(u_{\infty} + w, y) - V(u_{\infty}, y) - V'(u_{\infty}, y)w\right) dx \\ &\leq C \|w\|_{H^{1}_{c^{\dagger}}(\Sigma)}^2 \leq C e^{-2\sigma t}, \end{split}$$
(5.16)

for all $t \ge t_0$, with any $t_0 > 0$ and some $C = C(t_0) > 0$.

Let us now rewrite (5.14) in the form

 $||h(\cdot,$

$$w_t = \mathcal{A}w + h(\cdot, t), \qquad h(\cdot, t) := f(u(\cdot, t)) - f(u_\infty).$$
(5.17)

Recalling hypothesis (H2), (5.7) and (5.16), for all $t_2 \ge t_1 \ge t_0 > 0$ we have

$$\begin{split} t_{2}) - h(\cdot, t_{1}) \|_{L^{2}_{c^{\dagger}}(\Sigma)} &= \|f(u(\cdot, t_{2})) - f(u(\cdot, t_{1}))\|_{L^{2}_{c^{\dagger}}(\Sigma)} \\ &\leq C \|u(\cdot, t_{2}) - u(\cdot, t_{1})\|_{L^{2}_{c^{\dagger}}(\Sigma)} \\ &\leq C \int_{t_{1}}^{t_{2}} \|u_{t}(\cdot, s)\|_{L^{2}_{c^{\dagger}}(\Sigma)} ds \\ &\leq C \sqrt{(t_{2} - t_{1})\Phi_{c^{\dagger}}[u(\cdot, t_{1})]} \leq C \sqrt{t_{2} - t_{1}} e^{-\sigma t_{1}}, \end{split}$$

for some $C = C(t_0) > 0$. Then, reasoning as in [21, Theorem 4.3.1] with $t_1 = t - 1$ and $t_2 = t$ and using (5.15), we have

$$\begin{split} \|w(\cdot,t)\|_{H^{2}_{c^{\dagger}}(\Sigma)} &\leq C(\|\mathcal{A}w(\cdot,t)\|_{L^{2}_{c^{\dagger}}(\Sigma)} + \|w(\cdot,t)\|_{H^{1}_{c^{\dagger}}(\Sigma)}) \\ &\leq C\bigg(\|w(\cdot,t-1)\|_{L^{2}_{c^{\dagger}}(\Sigma)} + \|w(\cdot,t)\|_{H^{1}_{c^{\dagger}}(\Sigma)} \\ &\quad + \int_{t-1}^{t} \frac{\|h(\cdot,s) - h(\cdot,t)\|_{L^{2}_{c^{\dagger}}(\Sigma)}}{t-s} \, ds\bigg) \\ &\leq C \, e^{-\sigma t}, \end{split}$$
(5.18)

for all $t \ge t_0 + 1$, for any $t_0 > 0$ and some $C = C(t_0) > 0$. In writing (5.18), we used the same reasoning as in the standard estimate of the H^2 -norm of a function in terms of the L^2 -norm of the Laplacian to obtain the inequality in the first line. This gives (2.4) and concludes the proof of Theorem 1. \square Acknowledgements. The authors wish to thank the University of Tours and the Research Institute *Le Studium* for the kind hospitality and support (M.N.). We would also like to acknowledge valuable discussions with V. Moroz.

REFERENCES

- G. I. BARENBLATT AND Y. B. ZELDOVICH, On the stability of flame propagation, Prikladnaya Matematika i Mekhanika (Applied Mathematics and Mechanics (PMM)), 21 (1957), pp. 856–859. (in Russian).
- H. BERESTYCKI, B. LARROUTUROU, AND J.-M. ROQUEJOFFRE, Stability of travelling fronts in a model for flame propagation. I. Linear analysis, Arch. Rational Mech. Anal., 117 (1992), pp. 97–117.
- [3] H. BERESTYCKI AND L. NIRENBERG, Traveling fronts in cylinders, Ann. Inst. H. Poincaré Anal. Non Linéaire, 9 (1992), pp. 497–572.
- [4] M. BRAMSON, Convergence of solutions of the Kolmogorov equation to travelling waves, Amer. Math. Soc., Providence, RI, 1983.
- [5] L. C. EVANS, Partial Differential Equations, vol. 19 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 1998.
- [6] P. C. FIFE, Mathematical Aspects of Reacting and Diffusing Systems, Springer-Verlag, Berlin, 1979.
- [7] P. C. FIFE AND J. B. MCLEOD, The approach of solutions of nonlinear diffusion equations to traveling front solutions, Arch. Rat. Mech. Anal., 65 (1977), pp. 335–361.
- [8] T. GALLAY AND R. JOLY, Global stability of travelling fronts for a damped wave equation with bistable nonlinearity, Ann. Sci. Éc. Norm. Supér. (4), 42 (2009), pp. 103–140.
- T. GALLAY AND E. RISLER, A variational proof of global stability for bistable travelling waves, Differential Integral Equations, 20 (2007), pp. 901–926.
- [10] D. GILBARG AND N. S. TRUDINGER, Elliptic Partial Differential Equations of Second Order, Springer-Verlag, Berlin, 1983.
- S. HEINZE, Travelling waves for semilinear parabolic partial differential equations in cylindrical domains, SFB 123 Preprint 506, University of Heidelberg, 1989.
- [12] —, A variational approach to traveling waves., Tech. Rep. 85, Max Planck Institute for Mathematical Sciences, Leipzig, 2001.
- [13] D. HENRY, Perturbation of the boundary in boundary-value problems of partial differential equations, vol. 318 of London Mathematical Society Lecture Note Series, Cambridge University Press, Cambridge, 2005.
- [14] Y. I. KANEL', The behavior of solutions of the Cauchy problem when the time tends to infinity, in the case of quasilinear equations arising in the theory of combustion, Soviet Math. Dokl., 1 (1960), pp. 533–536.
- [15] —, On the stabilization of solutions of the Cauchy problem for the equations arising in the theory of combusion, Mat. Sbornik, 59 (1962), pp. 245–288.
- [16] K. KIRCHGÄSSNER, On the nonlinear dynamics of travelling fronts, J. Differential Equations, 96 (1992), pp. 256–278.
- [17] A. N. KOLMOGOROV, I. G. PETROVSKY, AND N. S. PISKUNOV, Study of the diffusion equation with growth of the quantity of matter and its application to a biological problem, Bull. Univ. Moscow, Ser. Int., Sec. A, 1 (1937), pp. 1–25.
- [18] P.-L. LIONS, The concentration-compactness principle in the calculus of variations. The locally compact case. I, Ann. Inst. H. Poincaré Anal. Non Linéaire, 1 (1984), pp. 109–145.
- [19] M. LUCIA, C. B. MURATOV, AND M. NOVAGA, Linear vs. nonlinear selection for the propagation speed of the solutions of scalar reaction-diffusion equations invading an unstable equilibrium, Commun. Pure Appl. Math., 57 (2004), pp. 616–636.
- [20] _____, Existence of traveling waves of invasion for Ginzburg-Landau-type problems in infinite cylinders, Arch. Rational Mech. Anal., 188 (2008), pp. 475–508.
- [21] A. LUNARDI, Analytic semigroups and optimal regularity in parabolic problems, vol. 16 of Progress in Nonlinear Differential Equations and their Applications, Birkhäuser, Basel, 1995.
- [22] J.-F. MALLORDY AND J.-M. ROQUEJOFFRE, A parabolic equation of the KPP type in higher dimensions, SIAM J. Math. Anal., 26 (1995), pp. 1–20.
- [23] C. B. MURATOV, A global variational structure and propagation of disturbances in reactiondiffusion systems of gradient type, Discrete Cont. Dyn. S., Ser. B, 4 (2004), pp. 867–892.
- [24] C. B. MURATOV AND M. NOVAGA, Front propagation in infinite cylinders. I. A variational approach, Comm. Math. Sci., 6 (2008), pp. 799–826.

- [25] —, Front propagation in infinite cylinders. II. The sharp reaction zone limit, Calc. Var. PDE, 31 (2008), pp. 521–547.
- [26] M. H. PROTTER AND H. F. WEINBERGER, Maximum principles in differential equations, Springer-Verlag, New York, 1984.
- [27] E. RISLER, Global convergence toward traveling fronts in nonlinear parabolic systems with a gradient structure, Ann. Inst. H. Poincaré Anal. Non Linéaire, 25 (2008), pp. 381–424.
- [28] J.-M. ROQUEJOFFRE, Stability of travelling fronts in a model for flame propagation. II. Nonlinear stability, Arch. Rational Mech. Anal., 117 (1992), pp. 119–153.
- [29] , Convergence to travelling waves for solutions of a class of semilinear parabolic equations, J. Differ. Equations, 108 (1994), pp. 262–295.
- [30] —, Eventual monotonicity and convergence to traveling fronts for the solutions of parabolic equations in cylinders, Ann. Inst. H. Poincaré Anal. Non Linéaire, 14 (1997), pp. 499–552.
- [31] F. ROTHE, Convergence to pushed fronts, Rocky Mountain J. Math., 11 (1981), pp. 617-633.
- [32] D. H. SATTINGER, On the stability of waves of nonlinear parabolic systems, Advances in Math., 22 (1976), pp. 312–355.
- [33] M. STRUWE, Variational methods: applications to nonlinear partial differential equations and Hamiltonian systems, Springer, Berlin, 2000.
- [34] M. E. TAYLOR, Partial Differential Equations III: Nonlinear Equations, Springer-Verlag, Berlin, 1996.
- [35] K. UCHIYAMA, The behavior of solutions of some non-linear diffusion equations for large time, J. Math. Kyoto Univ., 18 (1977), pp. 453–508.
- [36] J. M. VEGA, The asymptotic behavior of the solutions of some semilinear elliptic equations in cylindrical domains, J. Differ. Equations, 102 (1993), pp. 119–152.
- [37] —, On the uniqueness of multidimensional travelling fronts of some semilinear equations, J. Math. Anal. Appl., 177 (1993), pp. 481–490.
- [38] A. I. VOLPERT, V. A. VOLPERT, AND V. A. VOLPERT, Traveling wave solutions of parabolic systems, AMS, Providence, 1994.
- [39] J. XIN, Front propagation in heterogeneous media, SIAM Review, 42 (2000), pp. 161–230.