

# ATOMISTIC-TO-CONTINUUM COUPLING APPROXIMATION OF A ONE-DIMENSIONAL TOY MODEL FOR DENSITY FUNCTIONAL THEORY

B. LANGWALLNER, C. ORTNER, AND E. SÜLI

**ABSTRACT.** We consider an atomistic model defined through an interaction field satisfying a variational principle, and can therefore be considered a toy model of (orbital free) density functional theory. We investigate atomistic-to-continuum coupling mechanisms for this atomistic model, paying special attention to the dependence of the atomistic subproblem on the atomistic region boundary and the boundary conditions. We rigorously prove first-order error estimates for two related coupling mechanisms.

## 1. INTRODUCTION

The quasicontinuum (QC) method and, more generally, atomistic/continuum coupling (a/c) methods, are numerical coarse-graining techniques for the efficient simulation of phenomena and processes in materials at the nano-scale, such as defects, fracture, grain boundaries, or nano-indentation [18, 19, 16, 11]. Incompatibilities between the treatment of forces in atomistic and continuum models lead to difficulties in defining coupling mechanisms that do not introduce additional errors. Substantial effort has been made to understand this problem and to construct efficient and accurate a/c methods; see [17, 4, 15, 9, 20] for examples of formulations of computational methods and [1, 2, 3, 12, 13, 14] and references therein for examples of analytical treatments. Formulations of a/c methods for atomistic models based on quantum mechanics were proposed in [8, 6], but, to the best of our knowledge, no rigorous analysis of these methods exists.

In the present article we formulate and analyze one-dimensional a/c methods for an atomistic model that is defined through an interaction field satisfying a linear variational principle. Our results are related to two classes of a/c methods: Firstly, our work can be viewed as an analysis of (a simplified version of) the a/c method proposed by Iyer and Gavini [9], who use field-based versions of classical potentials to formulate their method. Secondly, the atomistic model we formulate can be considered a toy model of (orbital free) density functional theory, and hence our work represents a preliminary step towards a rigorous analysis of the a/c methods described in [8, 6].

The article is structured as follows. In Section 1 we formally motivate the atomistic model, and introduce the necessary notation. In Section 2 we give a precise formulation of the model with periodic boundary conditions and derive a “weak formulation” for the resulting forces on the particles. Section 3 is devoted to the analysis of the model in a bounded domain when the fields are subjected to Dirichlet boundary conditions. The Cauchy–Born continuum model is derived and analyzed in Section 4. Finally, in Sections 5 and 6 we propose two possible

---

*Date:* January 12, 2021.

*2000 Mathematics Subject Classification.* 65N12, 65N15, 70C20.

*Key words and phrases.* atomistic models, quasicontinuum method, coarse graining.

This work was supported by the EPSRC Critical Mass Programme “New Frontiers in the Mathematics of Solids” (OxMoS) and by the EPSRC Grant “Analysis of Atomistic-to-Continuum Coupling Methods”.

constructions of a/c methods based on different exchange of boundary conditions between an atomistic and continuum region, and establish error estimates.

**1.1. Field-based formulation of pair interactions.** The following outline follows ideas presented in [9]. Let  $\mathbf{y} = (y_1, \dots, y_N) \in \mathbb{R}^N$  represent the coordinates of  $N$  particles in one dimension. We consider an atomistic energy based on a pair-potential  $V$ ,

$$\mathcal{E}(\mathbf{y}) = \frac{1}{2} \sum_{\substack{i,j=1 \\ i \neq j}}^N V(|y_i - y_j|).$$

The force on particle  $i$  is given by

$$-D_{y_i} \mathcal{E}(\mathbf{y}) = - \sum_{\substack{j=1 \\ j \neq i}}^N \text{sign}(y_i - y_j) V'(|y_i - y_j|).$$

We note that the forces are nonlocal expressions in the sense that their computation involves summation over the other  $N - 1$  particles.

Next, we make a few modifications to this model. First, we replace the pointwise particles with smooth, nonnegative, and compactly supported particle densities  $\delta_\varepsilon(\cdot - y_i)$  (such that  $\int_{\mathbb{R}} \delta_\varepsilon(x) dx = 1$ ). This leads to

$$\mathcal{E}(\mathbf{y}) \approx \frac{1}{2} \sum_{\substack{i,j=1 \\ i \neq j}}^N \int_{\mathbb{R}} \int_{\mathbb{R}} \delta_\varepsilon(z - y_i) V(|z - x|) \delta_\varepsilon(x - y_j) dz dx.$$

To simplify the presentation further, we include the self-energies of the individual particle densities and define

$$\mathcal{E}_\varepsilon(\mathbf{y}) = \frac{1}{2} \sum_{i,j=1}^N \int_{\mathbb{R}} \int_{\mathbb{R}} \delta_\varepsilon(z - y_i) V(|z - x|) \delta_\varepsilon(x - y_j) dz dx.$$

This additional self-energy contribution does not affect the forces. It can be computed explicitly and subtracted from the energy later on. Upon introducing the field  $\phi : \mathbb{R} \rightarrow \mathbb{R}$ ,

$$\phi(x) = \int_{\mathbb{R}} \rho_{\mathbf{y}}(z) V(|x - z|) dz, \quad \text{where} \quad \rho_{\mathbf{y}}(z) = \sum_{i=1}^N \delta_\varepsilon(z - y_i), \quad (1.1)$$

to rewrite the energy  $\mathcal{E}_\varepsilon(\mathbf{y})$  in the form

$$\mathcal{E}_\varepsilon(\mathbf{y}) = \frac{1}{2} \int_{\mathbb{R}} \rho_{\mathbf{y}}(x) \phi(x) dx.$$

It is now easy to see that the forces are given by the *local* expression

$$-D_{\mathbf{y}} \mathcal{E}_\varepsilon(\mathbf{y}) = - \int_{\mathbb{R}} D_{\mathbf{y}} \rho_{\mathbf{y}}(z) \phi(z) dz.$$

Hence, if the field  $\phi$  is known, then it becomes unnecessary to compute nonlocal sums over particles. The nonlocality of the interaction has been encoded in the field  $\phi$ . However, it is now necessary to compute the field  $\phi$ , which is defined via the convolution (1.1).

Suppose that the pair-potential  $V$  is the Green's function for a linear differential operator  $L_V(\nabla)$ ; then,  $\phi$  can alternatively be computed by solving the differential equation

$$L_V(\nabla) \phi = \rho_{\mathbf{y}}.$$

As an example we consider the Yukawa potential in one space dimension

$$V(x) = \frac{1}{2m} e^{-m|x|} = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{1}{k^2 + m^2} e^{ikx} dk.$$

In this case  $\phi$  can be obtained as the solution to

$$-\Delta\phi + m^2\phi = \rho_{\mathbf{y}}$$

or, equivalently, as a solution to the minimization problem

$$\phi = \arg \min_{\varphi} \left\{ \frac{1}{2} \int_{\mathbb{R}} |\nabla\varphi|^2 + m^2\varphi^2 dx - \int_{\mathbb{R}} \rho_{\mathbf{y}}\varphi dx \right\}.$$

The resulting interaction potential  $\mathcal{E}_{\varepsilon}$  can also be written in the form

$$\mathcal{E}_{\varepsilon}(\mathbf{y}) = - \min_{\varphi} \left\{ \frac{1}{2} \int_{\mathbb{R}} |\nabla\varphi|^2 + m^2\varphi^2 dx - \int_{\mathbb{R}} \rho_{\mathbf{y}}\varphi dx \right\}. \quad (1.2)$$

The present work is devoted to the analysis of a/c approximations of (1.2) in a periodic one-dimensional setting. What distinguishes this analysis from previous analyses of a/c methods is that the coupling is achieved through an exchange of boundary conditions for the interaction field  $\phi$ , rather than ghost-force removal ideas such as [4, 15].

**Remark 1.** The interaction defined by (1.2) is purely repulsive. A purely attractive interaction can be obtained by changing the outer minus sign in the definition of  $\mathcal{E}_{\varepsilon}$  to a plus sign. We could combine two energies of the form (1.2) with different parameters  $m$  to model an interaction similar to the Morse potential  $V(|x|) = e^{-2|x|} - 2e^{-|x|}$  [9].  $\square$

**1.2. Notation.** We consider an infinite chain of atoms on the one-dimensional lattice  $\widehat{\mathbf{X}} = \varepsilon\mathbb{Z}$ , where  $\varepsilon = 2/(2N + 1)$  is the reference lattice spacing. Moreover, to keep the analysis simple, we admit only  $(2N + 1)$ -periodic displacements from the reference lattice (cf. [14]). Hence, we define the spaces of admissible displacements and deformations, respectively, by

$$\mathcal{U} = \left\{ \mathbf{u} \in \mathbb{R}^{\mathbb{Z}} : u_{j+(2N+1)} = u_j \quad \forall j \in \mathbb{Z}, \quad \sum_{j=-N}^N u_j = 0 \right\}, \quad \text{and}$$

$$\mathcal{Y} = F\widehat{\mathbf{X}} + \mathcal{U},$$

where  $F > 0$  is a prescribed *macroscopic strain*. A deformation  $\mathbf{y} \in \mathcal{Y}$  defines the computational domain

$$\Omega = (y_{-N-1}, y_N)$$

for the field variable  $\phi$ . We note that the length of the interval is independent of  $\mathbf{y}$ .

We define the finite differences  $\mathbf{y}', \mathbf{y}'' \in \mathcal{U}$  for  $\mathbf{y} \in \mathcal{Y}$  or  $\mathcal{U}$  by their respective components

$$y'_j = \frac{y_j - y_{j-1}}{\varepsilon}, \quad y''_j = \frac{y_{j+1} - 2y_j + y_{j-1}}{\varepsilon^2}.$$

Let us also define the weighted  $\ell^2$  scalar product and norm by

$$(\mathbf{u}, \mathbf{v})_{\varepsilon} = \varepsilon \sum_{\nu=-N}^N u_{\nu} v_{\nu} \quad \forall \mathbf{u}, \mathbf{v} \in \mathcal{U}, \quad \|\mathbf{u}\|_{\ell^2_{\varepsilon}} := (\mathbf{u}, \mathbf{u})_{\varepsilon}^{1/2} \quad \forall \mathbf{u} \in \mathcal{U}. \quad (1.3)$$

The  $\ell^{\infty}$ -norm is defined in the obvious way

$$\|\mathbf{u}\|_{\ell^{\infty}} = \max_{\nu=-N, \dots, N} |u_{\nu}| \quad \forall \mathbf{u} \in \mathcal{U}.$$

The space  $\mathcal{U}$  equipped with the discrete Sobolev seminorm  $\|\mathbf{u}\|_{\mathcal{U}^{1,2}} = \|\mathbf{u}'\|_{\ell^2_{\varepsilon}}$  will be denoted by  $\mathcal{U}^{1,2}$  and its topological dual space by  $\mathcal{U}^{-1,2}$ . The norm on  $\mathcal{U}^{-1,2}$  is given by

$$\|T\|_{\mathcal{U}^{-1,2}} = \sup_{\mathbf{u} \in \mathcal{U}^{1,2}} \frac{T\mathbf{u}}{\|\mathbf{u}\|_{\mathcal{U}^{1,2}}}.$$




FIGURE 2.1. Sketch of the basic atomistic problem: the field  $\phi$  is periodic in  $\Omega = (y_{-N-1}, y_N)$  and  $\rho_{\mathbf{y}}$  is a smooth particle density representing the atoms with positions given by  $\mathbf{y} \in \mathcal{Y}$ .

For monotonically increasing  $\mathbf{y} \in \mathcal{Y}$  (which we will write as  $\mathbf{y}' > 0$ ) we denote by  $S(\mathbf{y}) \subset H^1(\Omega)$  the space of continuous functions that are linear on every interval  $Q_i = (y_{i-1}, y_i)$ ,  $i \in \{-N, \dots, N\}$ . Furthermore, we define  $S_{\#}(\mathbf{y}) = S(\mathbf{y}) \cap H_{\#}^1(\Omega)$  to be the subset of all periodic functions in  $S(\mathbf{y})$ .

## 2. PERIODIC BOUNDARY CONDITIONS

We now put the field-based interaction potential that was outlined above in a precise mathematical framework. Let the functional  $I : H_{\#}^1(\Omega) \times \mathcal{Y} \rightarrow \mathbb{R}$  be defined by

$$I(\varphi, \mathbf{y}) = \int_{\Omega} \left( \frac{1}{2} \varepsilon^2 |\nabla \varphi|^2 + \frac{1}{2} m^2 \varphi^2 \right) dx - \int_{\Omega} \rho_{\mathbf{y}} \varphi dx, \quad \text{where}$$

$$\rho_{\mathbf{y}}(x) = \varepsilon \sum_{j \in \mathbb{Z}} \delta_{\varepsilon}(x - y_j), \quad \text{and} \quad \delta_{\varepsilon}(x) = \varepsilon^{-1} \delta_1(x/\varepsilon).$$

Here,  $\delta_1$  is a symmetric, nonnegative, regularized delta distribution with compact support  $[-\frac{\varsigma_0}{2}, \frac{\varsigma_0}{2}]$ , where  $\varsigma_0 > 0$  and  $\int_{\mathbb{R}} \delta_1 dx = 1$ ; see Figure 2.1. We will frequently refer to the parameter  $\varsigma_0$ , which is fixed throughout the paper.

We then define the interaction potential  $\mathcal{E} : \mathcal{Y} \rightarrow \mathbb{R}$  by

$$\mathcal{E}(\mathbf{y}) = - \min_{\varphi \in H_{\#}^1(\Omega)} I(\varphi, \mathbf{y}). \quad (2.1)$$

The respective minimizer (see Figure 2.1)

$$\phi = \arg \min_{\varphi \in H_{\#}^1(\Omega)} I(\varphi, \mathbf{y})$$

is the periodic solution to the Euler–Lagrange equation

$$-\varepsilon^2 \Delta \phi + m^2 \phi = \rho_{\mathbf{y}} \quad \text{in } \Omega. \quad (2.2)$$

Although  $\phi$  depends on  $\mathbf{y}$ , we will usually suppress this in our notation. It will always be clear from the context, which configuration  $\phi$  belongs to. It follows from (2.2) and integration by parts that

$$\mathcal{E}(\mathbf{y}) = \frac{1}{2} \int_{\Omega} \phi \rho_{\mathbf{y}} dx.$$

To determine equilibrium configurations subject to a given external force  $\mathbf{f} \in \mathcal{U}^{-1,2}$  we need to minimize the total potential energy  $E_{\mathbf{f}} : \mathcal{Y} \rightarrow \mathbb{R}$  defined by

$$E_{\mathbf{f}}(\mathbf{y}) = \mathcal{E}(\mathbf{y}) + (\mathbf{f}, \mathbf{y})_{\varepsilon}. \quad (2.3)$$

A minimizer  $\bar{\mathbf{y}} \in \mathcal{Y}$  of (2.3) satisfies the following Euler–Lagrange equation in  $\mathcal{U}^{-1,2}$ :

$$DE_{\mathbf{f}}(\bar{\mathbf{y}}) = D\mathcal{E}(\bar{\mathbf{y}}) + \mathbf{f} = \mathbf{0}.$$

In the following we analyze the derivatives of  $\mathcal{E}$ . In particular, we obtain a “weak” formulation for the first derivative  $D\mathcal{E}$  that acts as a natural connection point for the coupling with a continuum model.

**Proposition 1.1.** *The potential  $\mathcal{E} : \mathcal{Y} \rightarrow \mathbb{R}$  defined by (2.1) is twice continuously Fréchet differentiable. The components of the first derivative are given by*

$$D_{y_j}\mathcal{E}(\mathbf{y}) = -\varepsilon \int_{\Omega} \nabla \delta_{\varepsilon}(x - y_j) \phi(x) \, dx \quad (2.4)$$

for  $j \in \{-N, \dots, N-1\}$  and by

$$D_{y_N}\mathcal{E}(\mathbf{y}) = -\varepsilon \int_{\Omega} (\nabla \delta_{\varepsilon}(x - y_{-N-1}) + \nabla \delta_{\varepsilon}(x - y_N)) \phi(x) \, dx. \quad (2.5)$$

*Proof.* The proof of this result is standard and can be found in [9], for example.  $\square$

We stress the fact that the forces  $-D_{\mathbf{y}}\mathcal{E}(\mathbf{y})$  are local expressions. To calculate the force on atom  $j$  it is necessary to know  $\phi$  in  $\text{supp} \delta_{\varepsilon}(\cdot - y_j)$  but there is no need to sum over all remaining atoms. This nonlocality is encoded in the field  $\phi$ .

Next we establish the *weak formulation* for the forces on particles. This very much resembles the structure of the continuum equations and will be the basis for the a/c coupling in Section 5. A version of this calculation was already shown in [7], which used an interpolant for the displacement that is constant on the support of every  $\delta_{\varepsilon}(\cdot - y_j)$ . To avoid this restriction, we modify and extend the argument in [7].

For simplicity we assume that the supports of the densities of different particles do not intersect:

$$\text{supp} \delta_{\varepsilon}(\cdot - y_i) \cap \text{supp} \delta_{\varepsilon}(\cdot - y_j) = \emptyset \quad \forall i, j \in \mathbb{Z}, \quad i \neq j.$$

Since,  $|\text{supp} \delta_{\varepsilon}(\cdot - y_i)| = \varepsilon \varsigma_0$ , this is equivalent to  $|y_j - y_i| > \varepsilon \varsigma_0$  for  $i \neq j$  or, if  $\mathbf{y}$  is an increasing sequence,  $y'_j > \varsigma_0$  for all  $j \in \mathbb{Z}$ .

**Lemma 1.2.** *Let  $\mathbf{y} \in \mathcal{Y}$  satisfy  $\mathbf{y}' > \varsigma_0$  and let  $\phi \in H_{\#}^1(\Omega)$  be the associated field, defined by (2.2). Let  $\mathbf{u} = (u_j)_{j \in \mathbb{Z}} \in \mathcal{U}$  be a test vector and  $u \in S_{\#}(\mathbf{y})$  the periodic piecewise linear interpolant of  $\mathbf{u}$ , that is,  $u(y_j) = u_j$  for  $j \in \mathbb{Z}$ . Then,*

$$D\mathcal{E}(\mathbf{y}) \cdot \mathbf{u} = \sum_{j=-N}^N D_{y_j}\mathcal{E}(\mathbf{y}) \cdot u_j = \int_{\Omega} \sigma_{\mathbf{y}}(x) \nabla u(x) \, dx, \quad (2.6)$$

where  $\sigma_{\mathbf{y}} = \sigma_{\mathbf{y},1} + \sigma_{\mathbf{y},2}$  and

$$\begin{aligned} \sigma_{\mathbf{y},1}(x) &= \frac{1}{2} \varepsilon^2 |\nabla \phi|^2 - \frac{1}{2} m^2 \phi^2 + \rho_{\mathbf{y}} \phi, \\ \sigma_{\mathbf{y},2}(x) &= \varepsilon \sum_{j=-N-1}^N \phi(x) \nabla \delta_{\varepsilon}(x - y_j) (x - y_j). \end{aligned} \quad (2.7)$$

*Proof.* We begin by multiplying the derivative (2.4) for  $j \in \{-N, \dots, N-1\}$  by the component  $u_j$ :

$$\begin{aligned}
D_{y_j} \mathcal{E}(\mathbf{y}) u_j &= -\varepsilon u_j \int_{\Omega} \nabla \delta_{\varepsilon}(x - y_j) \phi(x) \, dx \\
&= -\varepsilon \int_{\Omega} u(x) \nabla \delta_{\varepsilon}(x - y_j) \phi(x) \, dx + \varepsilon \int_{\Omega} (u(x) - u_j) \nabla \delta_{\varepsilon}(x - y_j) \phi(x) \, dx \\
&= \varepsilon \int_{\Omega} \delta_{\varepsilon}(x - y_j) u(x) \nabla \phi(x) \, dx + \varepsilon \int_{\Omega} \delta_{\varepsilon}(x - y_j) \phi(x) \nabla u(x) \, dx \\
&\quad + \varepsilon \int_{\Omega} (u(x) - u_j) \nabla \delta_{\varepsilon}(x - y_j) \phi(x) \, dx =: T_1^{(j)} + T_2^{(j)} + T_3^{(j)}.
\end{aligned}$$

Here we have used integration by parts but there are no boundary terms since  $u$ ,  $\phi$  and  $\rho_{\mathbf{y}}$  are periodic on  $\Omega$ . Using (2.5) we obtain a similar expression for  $D_{y_N} \mathcal{E}(\mathbf{y}) u_N$ . Summing over  $j = -N, \dots, N$  we obtain

$$D\mathcal{E}(\mathbf{y}) \cdot \mathbf{u} = \sum_{j=-N}^N D_{y_j} \mathcal{E}(\mathbf{y}) \cdot u_j = T_1 + T_2 + T_3, \quad (2.8)$$

where  $T_i = \sum_{j=-N}^N T_i^{(j)}$ ,  $i \in \{1, 2, 3\}$ . From  $\rho_{\mathbf{y}} = \varepsilon \sum_{j \in \mathbb{Z}} \delta_{\varepsilon}(\cdot - y_j)$  it immediately follows that

$$T_2 = \int_{\Omega} \rho_{\mathbf{y}}(x) \phi(x) \nabla u(x) \, dx.$$

For  $T_1$  we can carry out the following rearrangements

$$\begin{aligned}
T_1 &= \int_{\Omega} \rho_{\mathbf{y}} u \nabla \phi \, dx = \int_{\Omega} (-\varepsilon^2 \Delta \phi + m^2 \phi) u \nabla \phi \, dx \\
&= \int_{\Omega} (-\varepsilon^2 \nabla \phi \Delta \phi + m^2 \phi \nabla \phi) u \, dx = \frac{1}{2} \int_{\Omega} \nabla (-\varepsilon^2 |\nabla \phi|^2 + m^2 \phi^2) u \, dx \\
&= \frac{1}{2} \int_{\Omega} (\varepsilon^2 |\nabla \phi|^2 - m^2 \phi^2) \nabla u \, dx.
\end{aligned}$$

Here, we have again used integration by parts and the periodicity of all functions involved. We deduce that

$$T_1 + T_2 = \int_{\Omega} \sigma_{\mathbf{y},1}(x) \nabla u(x) \, dx$$

with  $\sigma_{\mathbf{y},1}$  as defined in (2.7).

Before turning to  $T_3$  we first note that, since  $u$  is piecewise linear,

$$\begin{aligned}
u(x) &= u_j + \frac{x - y_j}{y_j - y_{j-1}} (u_j - u_{j-1}) = u_j + (x - y_j) \nabla u(x) \quad \text{for } x \in Q_j = (y_{j-1}, y_j), \\
u(x) &= u_j + \frac{x - y_j}{y_{j+1} - y_j} (u_{j+1} - u_j) = u_j + (x - y_j) \nabla u(x) \quad \text{for } x \in Q_{j+1} = (y_j, y_{j+1}).
\end{aligned}$$

Hence,  $T_3$  in the above equation (2.8) can be written as

$$\begin{aligned}
T_3 &= \varepsilon \sum_{j=-N-1}^N \int_{\Omega} \phi(x) \nabla \delta_{\varepsilon}(x - y_j) (u(x) - u_j) \, dx \\
&= \varepsilon \sum_{j=-N-1}^N \int_{\Omega} \phi(x) \nabla \delta_{\varepsilon}(x - y_j) (x - y_j) \nabla u(x) \, dx = \varepsilon \int_{\Omega} \sigma_{\mathbf{y},2}(x) \nabla u \, dx,
\end{aligned}$$

with  $\sigma_{\mathbf{y},2}$  as defined in (2.7), which concludes the proof.  $\square$

**Remark 2.** 1. In more than one space dimension the above calculations can be generalized if a triangular, respectively, tetrahedral mesh with the atomic positions as nodes is constructed. For example, this leads to

$$\sigma_{\mathbf{y},1}(x) = \left(-\frac{1}{2}\varepsilon^2|\nabla\phi|^2 - \frac{1}{2}m^2\phi^2 + \rho_{\mathbf{y}}\phi\right) \text{id} + \varepsilon^2\nabla\phi \otimes \nabla\phi.$$

2. A closer look at the calculations in the proof of Lemma 1.2 shows that the weak form can be obtained for semilinear models  $-\varepsilon^2\Delta\phi + F'(\phi) = \rho_{\mathbf{y}}$  with any convex function  $F$ . Even a fourth-order model of the form  $\varepsilon^4\Delta^2\phi - \varepsilon^2\Delta\phi + F'(\phi) = \rho_{\mathbf{y}}$  admits a similar weak formulation.  $\square$

As already suggested in the introduction the Green's function for the differential operator  $-\varepsilon^2\Delta + m^2\text{id}$  acting on functions defined on  $\mathbb{R}$  is given by

$$G_\varepsilon(x) = \frac{1}{2\varepsilon m} e^{-\frac{m}{\varepsilon}|x|}. \quad (2.9)$$

We therefore get the following explicit formulas for the function values  $\phi(x)$  and  $\nabla\phi(x)$  for  $x \in \Omega$ .

**Proposition 2.1.** *Let  $\mathbf{y} \in \mathcal{Y}$  and let  $\phi = \arg \min_{\varphi \in H_{\#}^1(\Omega)} I(\varphi, \mathbf{y})$  be the corresponding interaction field. Then, for every  $x \in \Omega$ ,*

$$\phi(x) = \int_{\mathbb{R}} G_\varepsilon(x-z)\rho_{\mathbf{y}}(z) dz = \frac{1}{2m} \sum_{k \in \mathbb{Z}} \int_{\mathbb{R}} \delta_\varepsilon(z-y_k) e^{-\frac{m}{\varepsilon}|x-z|} dz, \quad (2.10)$$

$$\nabla\phi(x) = \int_{\mathbb{R}} G_\varepsilon(x-z)\nabla\rho_{\mathbf{y}}(z) dz = \frac{1}{2m} \sum_{k \in \mathbb{Z}} \int_{\mathbb{R}} \nabla\delta_\varepsilon(z-y_k) e^{-\frac{m}{\varepsilon}|x-z|} dz. \quad (2.11)$$

*Proof.* The proof of this proposition is similar to the one of [5, Thm. 2.1]; see also [10, Prop. 2.4].  $\square$

The following is a consequence of the simple exponential form of the Yukawa potential and some elementary properties of the exponential function in one dimension. Let  $y_i, y_j \in \mathbb{R}$  satisfy  $y_j > y_i + \varepsilon\varsigma_0$ , so that the supports of particle densities representing the atoms  $i$  and  $j$  do not intersect. Then,

$$\begin{aligned} \int_{\mathbb{R}} \int_{\mathbb{R}} \delta_\varepsilon(z-y_j) e^{-\frac{m}{\varepsilon}|z-x|} \delta_\varepsilon(x-y_i) dx dz &= \int_{\mathbb{R}} \int_{\mathbb{R}} \delta_\varepsilon(z-y_j) e^{-\frac{m}{\varepsilon}(z-x)} \delta_\varepsilon(x-y_i) dx dz \\ &= e^{-\frac{m}{\varepsilon}(y_j-y_i)} \int_{\mathbb{R}} e^{-\frac{m}{\varepsilon}(z-y_j)} \delta_\varepsilon(z-y_j) dz \cdot \int_{\mathbb{R}} e^{-\frac{m}{\varepsilon}(y_i-x)} \delta_\varepsilon(y_i-x) dx \\ &= \mu^2 e^{-\frac{m}{\varepsilon}(y_j-y_i)}, \end{aligned} \quad (2.12)$$

where we have defined

$$\mu = \int_{\mathbb{R}} \delta_\varepsilon(x) e^{-\frac{m}{\varepsilon}x} dx = \int_{\mathbb{R}} \delta_\varepsilon(x) e^{\frac{m}{\varepsilon}x} dx = \int_{\mathbb{R}} \delta_1(x) e^{mx} dx.$$

Although we will frequently use this property, it is not essential for our reasoning. It merely makes some calculations more convenient.




FIGURE 3.1. The atomistic model in the domain  $\Omega_a$  with Dirichlet boundary conditions  $g = [g_L \ g_R]^T$ .

### 3. DIRICHLET BOUNDARY CONDITIONS

In this section we consider a version of the model (2.1) in the domain  $\Omega_a = (a_L, a_R) \subset \mathbb{R}$  subject to Dirichlet instead of periodic boundary conditions. This concept will be used later on, for the formulation of a/c methods, as the atomistic subproblem. We set  $a = [a_L \ a_R]^T \in \mathbb{R}^2$  and  $\Delta a = a_R - a_L$ . Throughout Section 3 we think of  $\mathbf{y} = (y_{-K}, \dots, y_K)$  as an ordered element of  $\Omega_a^{2K+1}$  such that  $a_L < y_{-K} < \dots < y_K < a_R$ . The particle density  $\rho_{\mathbf{y}}$  is canonically defined by

$$\rho_{\mathbf{y}} = \varepsilon \sum_{j=-K}^K \delta_\varepsilon(\cdot - y_j).$$

For simplicity we assume that the  $y_j$  are separated and lie well inside  $\Omega_a$  in the sense that  $\text{supp } \rho_{\mathbf{y}} \cap \partial\Omega_a = \emptyset$  or, equivalently,

$$\begin{aligned} y'_i &\geq \varepsilon_0, \text{ for } i = -K+1, \dots, K, \\ a_R - y_K &> \varepsilon_0/2 \quad \text{and} \quad y_{-K} - a_L > \varepsilon_0/2. \end{aligned} \quad (3.1)$$

We impose the following boundary conditions on the resulting field  $\phi : \Omega_a \rightarrow \mathbb{R}$ :

$$\phi(a_L) = g_L, \quad \phi(a_R) = g_R;$$

i.e.,  $\phi|_{\partial\Omega_a} = g$  with  $g = [g_L \ g_R]^T \in \mathbb{R}^2$ . The interaction potential  $\mathcal{E}_{a,g} : \Omega_a^{2K+1} \rightarrow \mathbb{R}$  is defined by

$$\mathcal{E}_{a,g}(\mathbf{y}) = - \min_{\substack{\varphi \in H^1(\Omega_a) \\ \varphi|_{\partial\Omega_a} = g}} I_a(\varphi, \mathbf{y}), \quad (3.2)$$

where  $I_a : H^1(\Omega_a) \times \Omega_a^{2K+1} \rightarrow \mathbb{R}$  is given by

$$I_a(\varphi, \mathbf{y}) = \int_{a_L}^{a_R} \left( \frac{1}{2} \varepsilon^2 |\nabla \varphi|^2 + \frac{1}{2} m^2 \varphi^2 \right) dx - \int_{a_L}^{a_R} \rho_{\mathbf{y}} \varphi dx. \quad (3.3)$$

For given  $\mathbf{y}$  the minimizer  $\phi$  is the weak solution to

$$\begin{aligned} -\varepsilon^2 \Delta \phi + m^2 \phi &= \rho_{\mathbf{y}} \quad \text{in } \Omega_a, \\ \phi|_{\partial\Omega_a} &= g. \end{aligned} \quad (3.4)$$

We will frequently use the decomposition

$$\phi = \phi_0 + \xi_{a,g}, \quad (3.5)$$

where  $\phi_0 \in H_0^1(\Omega_a)$  and  $\xi_{a,g} \in H^1(\Omega_a)$ , respectively, solve the boundary-value problems

$$\begin{aligned} -\varepsilon^2 \Delta \phi_0 + m^2 \phi_0 &= \rho_{\mathbf{y}} \quad \text{in } \Omega_a, \\ \phi_0|_{\partial\Omega_a} &= 0 \end{aligned}$$

and

$$\begin{aligned} -\varepsilon^2 \Delta \xi_{a,g} + m^2 \xi_{a,g} &= 0 \quad \text{in } \Omega_a, \\ \xi_{a,g}|_{\partial\Omega_a} &= g. \end{aligned} \tag{3.6}$$

This last boundary-value problem can be solved explicitly, which yields the following lemma.

**Lemma 2.1.** *The solution  $\xi_{a,g}$  of (3.6) is given by*

$$\xi_{a,g}(x) = c_L(a, g) e^{-\frac{m}{\varepsilon}(x-a_L)} + c_R(a, g) e^{-\frac{m}{\varepsilon}(a_R-x)}, \tag{3.7}$$

where the coefficients  $c_L(a, g)$  and  $c_R(a, g)$  are given by

$$c(a, g) = \begin{bmatrix} c_L(a, g) \\ c_R(a, g) \end{bmatrix} = \begin{bmatrix} 1 & \tau \\ \tau & 1 \end{bmatrix}^{-1} \begin{bmatrix} g_L \\ g_R \end{bmatrix} =: T_a^{-1} \cdot g \tag{3.8}$$

and we have defined  $\tau = \exp(-\frac{m}{\varepsilon} \Delta a)$ .

Note that, for  $\Delta a \gg \varepsilon$ ,  $\tau$  is exponentially small; hence we will often neglect terms of that order of magnitude. We will write  $\mathcal{O}(\tau)$  for a quantity or function that is (uniformly) bounded above by  $C\tau$  in modulus, where  $C$  is independent of  $\varepsilon$  and  $\Delta a$ . For example, we have  $c(a, g) = g + \mathcal{O}(\tau)$ .

Next, we compute the derivative of  $\mathcal{E}_{a,g}$  with respect to the atomic coordinates. For these derivatives, we obtain a “weak” formulation of the same shape as in the periodic case (see Proposition 1.1).

If  $\mathbf{y}' > 0$ , then we denote by  $\mathcal{S}(\mathbf{y} \cup a)$  the set of continuous, piecewise affine functions over the mesh given by the nodes  $a_L, y_{-K}, \dots, y_K, a_R$ . Moreover,  $\mathcal{S}_0(\mathbf{y} \cup a) = \mathcal{S}(\mathbf{y} \cup a) \cap H_0^1(\Omega_a)$ .

**Proposition 2.2.** *Let  $a, g \in \mathbb{R}^2$ ,  $a_L < a_R$ ; then  $\mathcal{E}_{a,g} : \mathcal{Y} \rightarrow \mathbb{R}$  defined by (3.2) is continuously Fréchet differentiable at  $\mathbf{y}$ .*

(i) *The components of the first derivative are given by*

$$D_{y_j} \mathcal{E}_{a,g}(\mathbf{y}) = -\varepsilon \int_{\Omega_a} \nabla \delta_\varepsilon(x - y_j) \phi(x) dx \quad \text{for } i = -K, \dots, K. \tag{3.9}$$

(ii) *Let  $\mathbf{u} \in \mathcal{U}$  be a test vector,  $u \in \mathcal{S}_0(\mathbf{y} \cup a)$  its interpolant, and let  $\min \mathbf{y}' \geq \varsigma_0$ ; then*

$$D_{\mathbf{y}} \mathcal{E}_{a,g}(\mathbf{y}) \cdot \mathbf{u} = \int_{\Omega_a} \sigma_{\mathbf{y}}(x) \nabla u(x) dx, \tag{3.10}$$

where  $\sigma_{\mathbf{y}}$  is given by (2.7).

*Proof.* The derivatives with respect to the coordinates  $\mathbf{y}$  are easy to calculate along the same lines as in the proof of Proposition 1.1. The weak formulation can be obtained as in the periodic case (Lemma 1.2) using the fact that the interpolant  $u$  vanishes on  $\partial\Omega_a$ .  $\square$

**Remark 3.** We point out that, in general,

$$\mathcal{E}_{a,g}(\mathbf{y}) \neq \frac{1}{2} \int_{\Omega_a} \rho_{\mathbf{y}} \phi dx.$$

However, we will see below that  $\mathcal{E}_{a,g}(\mathbf{y})$  can be written as the sum of a boundary data contribution and a term that is independent of  $g$ .  $\square$

With a view to the subsequent derivation of a/c methods we will from now on interpret  $a$  and  $g$  as arguments to  $\mathcal{E}_{a,g}$  rather than fixed parameters entering its definition. We consider the map  $\Omega_a^{2K+1} \times \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$ ,  $(\mathbf{y}, a, g) \mapsto \mathcal{E}_{a,g}(\mathbf{y})$ , and derive the derivatives of this map with respect to the boundary  $a$  and the boundary data  $g$ .

**3.1. Dependence on the boundary positions.** When formulating a/c methods in Section 5 we will let the boundary  $a$  of the atomistic subdomain depend on the configuration  $\mathbf{y}$ . It is therefore necessary to understand the dependence of the energy  $\mathcal{E}_{a,g}(\mathbf{y})$  on  $a$ . Our main result is that the derivative  $D_a \mathcal{E}_{a,g}(\mathbf{y})$  can be combined with  $D_{\mathbf{y}} \mathcal{E}_{a,g}(\mathbf{y})$  into a weak formulation reminiscent of (2.6). This will be a central building block for a/c methods.

**Proposition 3.1.** *Suppose that  $\mathbf{y} \in \mathcal{Y}$ ,  $\min \mathbf{y}' \geq \varsigma_0$ . Let  $h = [h_L \ h_R]^\top \in \mathbb{R}^2$  and  $\mathbf{u} = (u_{-K}, \dots, u_K) \in \mathbb{R}^{2K+1}$  be test vectors, and let  $u \in \mathcal{S}(\mathbf{y} \cup a)$  denote the interpolant of  $\mathbf{u}$  and  $h$  in the sense that*

$$u(a_L) = h_L, \quad u(a_R) = h_R, \quad \text{and} \quad u(y_j) = u_j \quad \forall j \in \{-K, \dots, K\}.$$

Then,

$$D_a \mathcal{E}_{a,g}(\mathbf{y}) \cdot h + D_{\mathbf{y}} \mathcal{E}_{a,g}(\mathbf{y}) \cdot \mathbf{u} = \int_{\Omega_a} \sigma_{\mathbf{y}}(x) \nabla u(x) \, dx.$$

*Proof.* This is a direct consequence of the following two lemmas.  $\square$

In the first auxiliary lemma we compute the derivative of  $\mathcal{E}_{a,g}(\mathbf{y})$  with respect to  $a = [a_L \ a_R]^\top$  while keeping the relative distances between the atoms constant. In other words we consider the change in  $\mathcal{E}_{a,g}(\mathbf{y})$  when the whole domain  $\Omega_a$  is stretched with the atom positions following this stretching. For  $\mathbf{y} \in \Omega_a^{2K+1}$  let  $\mathbf{X} = (X_{-K}, \dots, X_K) \in (0, 1)^{2K+1}$  be given by  $y_j = a_L + \Delta a X_j$  for all  $j \in \{-K, \dots, K\}$ . For fixed  $g, \mathbf{X}$  we define

$$\begin{aligned} \tilde{\mathcal{E}}(a) &:= \mathcal{E}_{a,g}(a_L + (a_R - a_L)\mathbf{X}), \quad \text{and} \\ \tilde{D}_{a_R} \mathcal{E}_{a,g}(\mathbf{y}) &:= D_{a_R} \tilde{\mathcal{E}}(a). \end{aligned}$$

(We understand  $a_L + (a_R - a_L)\mathbf{X}$  in a componentwise manner:  $(a_L + \Delta a \mathbf{X})_j = a_L + \Delta a X_j$  for all  $j \in \{-K, \dots, K\}$ .) The derivative  $\tilde{D}_{a_L} \mathcal{E}_{a,g}(\mathbf{y})$  is defined analogously.

**Lemma 3.2.** *Let  $\mathbf{y} \in \Omega_a^{2K+1}$  satisfy (3.1); then*

$$-\tilde{D}_{a_L} \mathcal{E}_{a,g}(\mathbf{y}) = \tilde{D}_{a_R} \mathcal{E}_{a,g}(\mathbf{y}) = \frac{1}{\Delta a} \int_{\Omega_a} \sigma_{\mathbf{y}}(x) \, dx. \quad (3.11)$$

*Proof.* We fix  $\mathbf{X}$  and let  $\boldsymbol{\eta}(a) := a_L + \Delta a \mathbf{X}$ . We begin by transforming the problem to the unit interval  $(0, 1)$  using the transformation  $x \mapsto X(x) = (x - a_L)/(a_R - a_L)$ :

$$\begin{aligned} \tilde{\mathcal{E}}(a) &= \mathcal{E}_{a,g}(\boldsymbol{\eta}(a)) = \int_{\Omega_a} \left( -\frac{1}{2} \varepsilon^2 |\nabla \phi|^2 - \frac{1}{2} m^2 \phi^2 + \rho_{\boldsymbol{\eta}(a)} \phi \right) dx \\ &= \Delta a \int_0^1 \left( -\frac{\varepsilon^2}{2 \Delta a^2} |\nabla \hat{\phi}|^2 - \frac{m^2}{2} \hat{\phi}^2 + \hat{\rho}_{\boldsymbol{\eta}(a)} \hat{\phi} \right) dX. \end{aligned} \quad (3.12)$$

Here,  $\hat{\phi}(X) = \phi(x(X))$  and  $\hat{\rho}_{\boldsymbol{\eta}(a)}(X) = \rho_{\boldsymbol{\eta}(a)}(x(X))$ . It follows as in Proposition 1.1 that, to compute  $D_a \tilde{\mathcal{E}}(a)$ , it is sufficient to calculate the partial derivatives of the right-hand side with respect to  $a_R$  (the derivative of  $\phi$  or  $\hat{\phi}$  with respect to  $a_R$  does not appear since  $\phi$  is a minimizer of  $I_a(\cdot, \mathbf{y})$ ). This leads to

$$\begin{aligned} D_{a_R} \tilde{\mathcal{E}}(a) &= \int_0^1 \left( -\frac{\varepsilon^2}{2 \Delta a^2} |\nabla \hat{\phi}|^2 - \frac{m^2}{2} \hat{\phi}^2 + \hat{\rho}_{\boldsymbol{\eta}(a)} \hat{\phi} \right) dX + \Delta a \int_0^1 \frac{\varepsilon^2}{\Delta a^3} |\nabla \hat{\phi}|^2 dX \\ &\quad + \Delta a \int_0^1 \hat{\phi} D_{a_R} \hat{\rho}_{\boldsymbol{\eta}(a)} dX. \end{aligned}$$

Transforming the first two integrals on the right-hand side back to the interval  $\Omega_a$  we arrive at

$$\frac{1}{\Delta a} \mathcal{E}_{a,g}(\mathbf{y}) + \frac{\varepsilon^2}{\Delta a} \int_{\Omega_a} |\nabla \phi|^2 dx = \frac{1}{\Delta a} \int_{\Omega_a} \sigma_{\mathbf{y},1}(x) dx,$$

where  $\sigma_{\mathbf{y},1}$  was given in (2.7).

It remains to differentiate  $\widehat{\rho}_{\eta(a)}$  with respect to  $a_R$ . By the definition of the transformation  $x \mapsto X(x)$  we have

$$\begin{aligned} D_{a_R} \widehat{\rho}_{\eta(a)}(X) &= \varepsilon D_{a_R} \sum_{j=-K}^K \delta_\varepsilon((a_R - a_L)(X - X_j)) \\ &= \varepsilon \sum_{j=-K}^K (X - X_j) \nabla \delta_\varepsilon((a_R - a_L)(X - X_j)). \end{aligned}$$

Using  $\Delta a(X - X_j) = (x - y_j)$  we therefore get

$$\begin{aligned} \Delta a \int_0^1 \widehat{\phi} D_{a_R} \widehat{\rho}_{\eta(a)} dX &= \frac{\varepsilon}{\Delta a} \sum_{j=-K}^K \int_{\Omega_a} (x - y_j) \nabla \delta_\varepsilon(x - y_j) \phi(x) dx \\ &= \frac{1}{\Delta a} \int_{\Omega_a} \sigma_{\mathbf{y},2}(x) dx \end{aligned}$$

with  $\sigma_{\mathbf{y},2}(x)$  as given in (2.7).

To see that  $D_{a_L} \widetilde{\mathcal{E}} = -D_{a_R} \widetilde{\mathcal{E}}$  we simply note that  $\mathcal{E}(a)$  depends only on  $\Delta a$ , which can be seen from (3.12) and the definition of  $\widehat{\rho}_{\eta(a)}(X)$ .  $\square$

We define  $\theta_R \in \mathcal{S}(\mathbf{y} \cup a)$  to be the piecewise linear function with

$$\theta_R(a_R) = 1, \quad \theta_R(a_L) = 0, \quad \theta_R(y_j) = 0 \quad \text{for all } j \in \{-K, \dots, K\}.$$

The function  $\theta_L \in \mathcal{S}(\mathbf{y} \cup a)$  is defined analogously.

**Lemma 3.3.** *Let  $\mathbf{y} \in \Omega_a^{2K+1}$  satisfy (3.1); then, the derivatives of  $\mathcal{E}_{a,g}(\mathbf{y})$  with respect to  $a_L$ ,  $a_R$  (for fixed  $\mathbf{y}$  and  $g$ ) satisfy*

$$\begin{aligned} D_{a_L} \mathcal{E}_{a,g}(\mathbf{y}) &= \int_{\Omega_a} \sigma_{\mathbf{y}}(x) \nabla \theta_L(x) dx, \\ D_{a_R} \mathcal{E}_{a,g}(\mathbf{y}) &= \int_{\Omega_a} \sigma_{\mathbf{y}}(x) \nabla \theta_R(x) dx. \end{aligned}$$

*Proof.* Let  $\Theta_R$  be the affine function defined on  $\Omega_a$  with  $\Theta_R(a_L) = 0$ ,  $\Theta_R(a_R) = 1$ . Since  $\nabla \Theta_R(x) = \frac{1}{\Delta a}$ , Lemma 3.2 yields

$$\widetilde{D}_{a_R} \mathcal{E}_{a,\mathbf{y}}(\mathbf{y}) = \int_{\Omega_a} \sigma_{\mathbf{y}} \nabla \Theta_R dx = \int_{\Omega_a} \sigma_{\mathbf{y}} \nabla (\Theta_R - \theta_R) dx + \int_{\Omega_a} \sigma_{\mathbf{y}} \nabla \theta_R dx. \quad (3.13)$$

Now, we have  $\Theta_R - \theta_R \in \mathcal{S}_0(\mathbf{y} \cup a)$  and hence, by Proposition 2.2,

$$\int_{\Omega_a} \sigma_{\mathbf{y}}(x) \nabla (\Theta_R - \theta_R) dx = \sum_{j=-K}^K D_{y_j} \mathcal{E}_{a,g}(\mathbf{y}) \Theta_R(y_j). \quad (3.14)$$

However,  $\tilde{D}_{a_R}\mathcal{E}_{a,g}(\mathbf{y})$  was defined as the derivative with respect to  $a_R$ , while the relative distances of the atoms are kept constant. This can be formulated as

$$\tilde{D}_{a_R}\mathcal{E}_{a,g}(\mathbf{y}) = D_{a_R}\mathcal{E}_{a,g}(\mathbf{y}) + \sum_{j=-K}^K D_{y_j}\mathcal{E}_{a,g}(\mathbf{y})\Theta_R(y_j).$$

Inserting this into (3.13) and using (3.14) then gives

$$\int_{\Omega_a} \sigma_{\mathbf{y}}(x)\nabla\theta_R dx = D_{a_R}\mathcal{E}_{a,g}(\mathbf{y}).$$

Similarly, we can show the expression stated for  $D_{a_L}\mathcal{E}_{a,g}(\mathbf{y})$ .  $\square$

**3.2. Dependence on the boundary conditions.** Next, we compute the derivative of  $\mathcal{E}_{a,g}(\mathbf{y})$  with respect to the boundary conditions  $g$  when the configuration  $\mathbf{y}$  and the boundary  $a$  are kept fixed. We define

$$\gamma_L(\mathbf{y}, a) = 2 \int_{\Omega_a} \rho_{\mathbf{y}}(x)G_\varepsilon(x - a_L) dx, \quad \text{and} \quad \gamma_R(\mathbf{y}, a) = 2 \int_{\Omega_a} \rho_{\mathbf{y}}(x)G_\varepsilon(a_R - x) dx. \quad (3.15)$$

**Lemma 3.4.** *The partial derivative of  $\mathcal{E}_{a,g}(\mathbf{y})$  with respect to  $g$  is given by:*

$$D_g\mathcal{E}_{a,g}(\mathbf{y}) = -m\varepsilon \left( (1 - \tau^2) \begin{bmatrix} c_L(a, g) \\ c_R(a, g) \end{bmatrix} - \begin{bmatrix} \gamma_L(\mathbf{y}, a) \\ \gamma_R(\mathbf{y}, a) \end{bmatrix} \right)^T \cdot T_a^{-1},$$

where  $T_a$ ,  $c(a, g) = [c_L(a, g) \ c_R(a, g)]^T$  and  $\tau = e^{-\frac{m}{\varepsilon}\Delta a}$  are defined in Lemma 2.1.

*Proof.* Throughout the proof we suppress the arguments of  $\gamma_L$ ,  $\gamma_R$ , and  $c$  for ease of readability. We recall the additive decomposition  $\phi = \phi_0 + \xi_{a,g}$  from (3.5). From  $\phi_0 \in H_0^1(\Omega)$ , and from the equation  $-\varepsilon^2\Delta\xi_{a,g} + m^2\xi_{a,g} = 0$  it follows that  $\varepsilon^2(\nabla\xi_{a,g}, \nabla\phi_0) + m^2(\xi_{a,g}, \phi_0) = 0$ . Hence, a short calculation shows that the energy  $\mathcal{E}_{a,g}(\mathbf{y})$  can be rewritten as

$$\mathcal{E}_{a,g}(\mathbf{y}) = -I_a(\phi, \mathbf{y}) = -I_a(\phi_0, \mathbf{y}) - I_a(\xi_{a,g}, \mathbf{y}). \quad (3.16)$$

The first term on the right-hand side does not depend on the boundary conditions  $g$  and the second term is known explicitly: using  $-\varepsilon^2\Delta\xi_{a,g} + m^2\xi_{a,g} = 0$ , integration by parts, and the explicit formula (3.7) for  $\xi_{a,g}$ , we obtain

$$\begin{aligned} I_a(\xi_{a,g}, \mathbf{y}) &= \int_{\Omega_a} \frac{1}{2}(\varepsilon^2|\nabla\xi_{a,g}|^2 + m^2\xi_{a,g}^2) dx - \int_{\Omega_a} \rho_{\mathbf{y}}\xi_{a,g} dx \\ &= \frac{\varepsilon^2}{2}(-\xi_{a,g}(a_L)\nabla\xi_{a,g}(a_L) + \xi_{a,g}(a_R)\nabla\xi_{a,g}(a_R)) - \int_{\Omega_a} \rho_{\mathbf{y}}\xi_{a,g} dx \\ &= \frac{\varepsilon m}{2}(c_L^2 + c_R^2)(1 - e^{-2\frac{m}{\varepsilon}\Delta a}) - \int_{\Omega_a} \rho_{\mathbf{y}}(c_L e^{-\frac{m}{\varepsilon}(x-a_L)} + c_R e^{-\frac{m}{\varepsilon}(a_R-x)}) dx \\ &= m\varepsilon \left( \frac{c_L^2 + c_R^2}{2}(1 - \tau^2) - \frac{2}{2m\varepsilon} \int_{\Omega_a} \rho_{\mathbf{y}}(c_L e^{-\frac{m}{\varepsilon}(x-a_L)} + c_R e^{-\frac{m}{\varepsilon}(a_R-x)}) dx \right) \\ &= m\varepsilon \left( \frac{c_L^2 + c_R^2}{2}(1 - \tau^2) - (c_L\gamma_L + c_R\gamma_R) \right). \end{aligned}$$

Here we have used the Green's function  $G_\varepsilon$  from (2.9). Differentiating this expression with respect to  $c_L$  and  $c_R$  and applying the chain rule with  $D_g c = T_a^{-1}$  yield the result.  $\square$

**Remark 4.** 1. We remark that  $D_g \mathcal{E}_{a,g}(\mathbf{y}) = 0$  if and only if  $c_L(a, g) = \gamma_L(\mathbf{y}, a)/(1 - \tau^2)$  and  $c_R(a, g) = \gamma_R(\mathbf{y}, a)/(1 - \tau^2)$ . According to (3.8) this corresponds to the “optimal” boundary conditions

$$g_L^* = \frac{1}{1 - \tau} \frac{\gamma_L + \tau \gamma_R}{1 + \tau}, \quad \text{and} \quad g_R^* = \frac{1}{1 - \tau} \frac{\tau \gamma_L + \gamma_R}{1 + \tau}. \quad (3.17)$$

That is, the boundary conditions are weighted averages of the values  $\frac{1}{1 - \tau} \gamma_L$  and  $\frac{1}{1 - \tau} \gamma_R$ .

2. As can be seen from Lemma 3.4 the boundary data contribution  $I_a(\xi_{a,g}, \mathbf{y})$  to the energy  $\mathcal{E}_{a,g}(\mathbf{y})$  is quadratic in  $g$ . For fixed configuration  $\mathbf{y}$  and domain  $\Omega_a$  the boundary conditions  $g = g^*(\mathbf{y}, a)$  minimize the boundary data contribution  $I_a(\xi_{a,g}, \mathbf{y})$  to the energy  $\mathcal{E}_{a,g}(\mathbf{y})$ . This is equivalent to minimizing  $I_a(\cdot, \mathbf{y})$  over  $H^1(\Omega_a)$  and therefore leads to homogeneous Neumann boundary conditions for  $\phi$  on  $\partial\Omega_a$ .

3. If  $\Delta a \gg \varepsilon$ , i.e.,  $\tau \ll 1$ , then we have  $\gamma_{L/R} = g_{L/R}^* + \mathcal{O}(\tau)$ , and hence we can simplify

$$\begin{aligned} I_a(\xi_{a,g}, \mathbf{y}) &= m\varepsilon \left( \frac{1}{2} (g_L^2 + g_R^2) - (g_L g_L^* + g_R g_R^*) \right) + \mathcal{O}(\varepsilon\tau), \quad \text{and} \\ D_g \mathcal{E}_{a,g}(\mathbf{y}) &= m\varepsilon (g^* - g) + \mathcal{O}(\varepsilon\tau). \end{aligned} \quad (3.18) \quad \square$$

A useful auxiliary result for the analysis of a/c methods is the global Lipschitz continuity of the field  $\phi$  with respect to variations in the boundary conditions  $g$ .

**Lemma 4.1.** *Let  $\phi_1, \phi_2 \in H^1(\Omega_a)$  be minimizers of  $I_a(\cdot, \mathbf{y})$  subject to the boundary conditions  $g_1 \in \mathbb{R}^2$ , respectively,  $g_2 \in \mathbb{R}^2$ . Then,*

$$\begin{aligned} |\phi_1(x) - \phi_2(x)| &\leq \sqrt{2} |T_a^{-1}(g_1 - g_2)| e^{-\frac{m}{\varepsilon} d_a(x)}, \quad \text{and} \\ \varepsilon |\nabla \phi_1(x) - \nabla \phi_2(x)| &\leq \sqrt{2} m |T_a^{-1}(g_1 - g_2)| e^{-\frac{m}{\varepsilon} d_a(x)}, \end{aligned}$$

where  $d_a(x) := \min(x - a_L, a_R - x)$  denotes the distance to the boundary of  $\Omega_a$ , for  $x \in \Omega_a$ .

*Proof.* We write both functions in the form  $\phi_i = \phi_0 + \xi_{a,g_i}$ ,  $i \in \{1, 2\}$ . For  $i = 1, 2$ , let  $c_i = T_a^{-1} g_i$  be the respective coefficients entering  $\xi_{a,g_i}$ ; then

$$|\phi_1(x) - \phi_2(x)| = |\xi_{a,g_1}(x) - \xi_{a,g_2}(x)| \leq |c_{1,L} - c_{2,L}| e^{-\frac{m}{\varepsilon}(x - a_L)} + |c_{1,R} - c_{2,R}| e^{-\frac{m}{\varepsilon}(a_R - x)}.$$

This immediately yields the first bound. The bound for the derivatives is obtained similarly.  $\square$

**3.3. A special case.** We now take a closer look at the interaction potential  $\mathcal{E}_{a,g}$  from (3.2) with the  $\mathbf{y}$ -dependent boundary conditions  $g = g^*(\mathbf{y}, a)$  defined in Remark 4.

**Proposition 4.2.** *Let  $\mathbf{y} \in \Omega_a^{2K+1}$ . Then,*

$$\begin{aligned} \mathcal{E}_{a,g^*(\mathbf{y},a)}(\mathbf{y}) &= \frac{1}{4m\varepsilon} \int_{\Omega_a} \int_{\Omega_a} \rho_{\mathbf{y}}(x) e^{-\frac{m}{\varepsilon}|x-z|} \rho_{\mathbf{y}}(z) dz dx + \tau M_\tau(\gamma_L, \gamma_R) \\ &\quad + \frac{1}{4m\varepsilon} \int_{\Omega_a} \int_{\Omega_a} \rho_{\mathbf{y}}(x) \left( e^{-\frac{m}{\varepsilon}(2a_R - x - z)} + e^{-\frac{m}{\varepsilon}(x + z - 2a_L)} \right) \rho_{\mathbf{y}}(z) dz dx, \end{aligned} \quad (3.19)$$

where  $M_\tau(\gamma_L, \gamma_R)$  depends quadratically on  $\gamma_L$  and  $\gamma_R$ .

Expression (3.19) can be interpreted as the energy of the atoms represented by  $\mathbf{y}$  interacting with each other plus the interaction with mirror atoms outside  $\Omega_a$ . This mirror interaction was introduced by means of the boundary conditions  $g = g^*$ .

For the proof of the proposition it is convenient to use an explicit formula for the function values of  $\phi_0 \in H_0^1(\Omega_a)$  from the decomposition (3.5). By Proposition 2.1, the Green’s function for the equation  $-\varepsilon^2 \Delta \phi + m^2 \phi = \rho_{\mathbf{y}}$  in  $\mathbb{R}$  is given by  $G_\varepsilon(x, y) = \frac{1}{2m\varepsilon} e^{-\frac{m}{\varepsilon}|x-y|}$ . We will now

construct the Green's function  $G_{\varepsilon,a}$  for the operator  $-\varepsilon^2\Delta + m^2\text{id}$  subject to homogeneous Dirichlet conditions on  $\partial\Omega_a$ .

**Lemma 4.3.** *Let  $\phi_0 \in H_0^1(\Omega_a)$  satisfy  $-\varepsilon^2\Delta\phi_0 + m^2\phi_0 = \rho_{\mathbf{y}}$  in  $\Omega_a$ . Then,*

$$\phi_0(x) = \int_{\Omega_a} G_{\varepsilon,a}(x,z)\rho_{\mathbf{y}}(z) dz \quad \forall x \in \Omega_a, \quad (3.20)$$

where  $G_{\varepsilon,a} = G_{\varepsilon,a}^{(1)} + \tau G_{\varepsilon,a}^{(2)}$ , with  $G_{\varepsilon,a}^{(i)}$ ,  $i = 1, 2$ , given by

$$\begin{aligned} G_{\varepsilon,a}^{(1)}(x,z) &= \frac{1}{2m\varepsilon} \left( e^{-\frac{m}{\varepsilon}|x-z|} - e^{-\frac{m}{\varepsilon}(x+z-2a_L)} - e^{-\frac{m}{\varepsilon}(2a_R-x-z)} \right), \\ G_{\varepsilon,a}^{(2)}(x,z) &= -\frac{1}{2m\varepsilon} \frac{1}{1-\tau^2} \left( \tau e^{-\frac{m}{\varepsilon}(x+z-2a_L)} + \tau e^{-\frac{m}{\varepsilon}(2a_R-x-z)} \right. \\ &\quad \left. - e^{-\frac{m}{\varepsilon}(x-z+a_R-a_L)} - e^{-\frac{m}{\varepsilon}(z-x+a_R-a_L)} \right). \end{aligned}$$

*Proof.* The proof of this result is standard [5, Chapter 2.2.4]; see also [10, Lemma 3.10].  $\square$

We remark that  $G_{\varepsilon,a} = G_{\varepsilon,a}^{(1)} + \mathcal{O}(\tau)$ .

*Proof of Proposition 4.2.* We have already seen in (3.16) that for any choice of boundary data  $g \in \mathbb{R}^2$  the energy  $\mathcal{E}_{a,g}(\mathbf{y})$  can be written as the sum of two terms

$$\mathcal{E}_{a,g}(\mathbf{y}) = -I_a(\phi, \mathbf{y}) = -I_a(\phi_0, \mathbf{y}) - I_a(\xi_{a,g}, \mathbf{y}),$$

where  $I_a(\phi_0, \mathbf{y})$  is independent of the boundary conditions.

*Calculation of  $I_a(\phi_0, \mathbf{y})$ .* Since the function  $\phi_0$  is a minimizer of  $I_a(\cdot, \mathbf{y})$  over  $H_0^1(\Omega)$ , we have with the expression (3.20) for  $\phi_0(x)$  that

$$I_a(\phi_0, \mathbf{y}) = -\frac{1}{2} \int_{\Omega_a} \int_{\Omega_a} \rho_{\mathbf{y}}\phi_0 dx = -\frac{1}{2} \int_{\Omega_a} \int_{\Omega_a} \rho_{\mathbf{y}}(x)G_{\varepsilon,a}(x,z)\rho_{\mathbf{y}}(z) dz dx. \quad (3.21)$$

By the definition (3.15) of  $\gamma_L$  and  $\gamma_R$  we have

$$\begin{aligned} \frac{1}{4m\varepsilon} \int_{\Omega_a} \int_{\Omega_a} \rho_{\mathbf{y}}(x)e^{-\frac{m}{\varepsilon}(2a_R-x-z)}\rho_{\mathbf{y}}(z) dx dz &= \frac{m\varepsilon}{4} \gamma_R^2, \\ \frac{1}{4m\varepsilon} \int_{\Omega_a} \int_{\Omega_a} \rho_{\mathbf{y}}(x)e^{-\frac{m}{\varepsilon}(x+z-2a_L)}\rho_{\mathbf{y}}(z) dx dz &= \frac{m\varepsilon}{4} \gamma_L^2, \\ \frac{1}{4m\varepsilon} \int_{\Omega_a} \int_{\Omega_a} \rho_{\mathbf{y}}(x)e^{-\frac{m}{\varepsilon}(z-x+a_R-a_L)}\rho_{\mathbf{y}}(z) dx dz &= \frac{m\varepsilon}{4} \gamma_L\gamma_R. \end{aligned} \quad (3.22)$$

Inserting the expression  $G_{\varepsilon,a} = G_{\varepsilon,a}^{(1)} + \tau G_{\varepsilon,a}^{(2)}$  into (3.21) and using these equalities yields

$$\begin{aligned} I_a(\phi_0, \mathbf{y}) &= -\frac{1}{2} \int_{\Omega_a} \int_{\Omega_a} \rho_{\mathbf{y}}(x)G_{\varepsilon}(x,z)\rho_{\mathbf{y}}(z) dz dx + \frac{m\varepsilon}{4} (\gamma_L^2 + \gamma_R^2) \\ &\quad + \frac{m\varepsilon}{4} \frac{\tau}{1-\tau^2} (\tau\gamma_L^2 + \tau\gamma_R^2 - 2\gamma_L\gamma_R). \end{aligned}$$

*Calculation of  $I_a(\xi_{a,g^*(\mathbf{y},a)}, \mathbf{y})$ .* From Lemma 3.4 we know that for general  $g \in \mathbb{R}^2$

$$I_a(\xi_{a,g}, \mathbf{y}) = m\varepsilon \left( \frac{c_L^2 + c_R^2}{2} (1-\tau^2) - (c_L\gamma_L + c_R\gamma_R) \right).$$

If  $g = g^*(\mathbf{y}, a)$ , then  $c_L = \gamma_L/(1-\tau^2)$  and  $c_R = \gamma_R/(1-\tau^2)$  as seen in Remark 4. Hence,

$$I_a(\xi_{a,g^*(\mathbf{y},a)}, \mathbf{y}) = -\frac{m\varepsilon}{2} \frac{1}{1-\tau^2} (\gamma_L^2 + \gamma_R^2).$$




FIGURE 4.1. The Cauchy–Born approximation: independent periodic problems are solved on the cells  $Q_j = (y_{j-1}, y_j)$  leading to locally defined fields  $\psi^{(j)}$ .

Isolating the dependence on  $\tau$  gives

$$I_a(\xi_{a,g^*}(\mathbf{y}, a), \mathbf{y}) = -\frac{m\varepsilon}{2}(\gamma_L^2 + \gamma_R^2) - \frac{m\varepsilon}{2} \frac{\tau^2}{1 - \tau^2}(\gamma_L^2 + \gamma_R^2). \quad (3.23)$$

*Conclusion.* Adding  $-I_a(\xi_{a,g^*}(\mathbf{y}, a), \mathbf{y})$  as just obtained and  $-I_a(\phi_0, \mathbf{y})$  from above we arrive at

$$\begin{aligned} \mathcal{E}_{a,g^*}(\mathbf{y}, a)(\mathbf{y}) &= \frac{1}{4m\varepsilon} \int_{\Omega_a} \int_{\Omega_a} \rho_{\mathbf{y}}(x) e^{-\frac{m}{\varepsilon}|x-z|} \rho_{\mathbf{y}}(z) dz dx + \frac{m\varepsilon}{4} (\gamma_L^2 + \gamma_R^2) \\ &\quad - \frac{m\varepsilon}{4} \frac{\tau}{1 - \tau^2} (\tau\gamma_L^2 + 2\gamma_L\gamma_R + \tau\gamma_R^2). \end{aligned} \quad (3.24)$$

Defining  $\tau M_\tau(\gamma_L, \gamma_R)$  to be the third term on the right-hand side and applying (3.22) yields (3.19).  $\square$

#### 4. THE CAUCHY–BORN APPROXIMATION

The next building block for the design of a/c methods based on the model (2.1) is the respective continuum model. Let  $\mathbf{y} \in \mathcal{Y}$  satisfy  $\min \mathbf{y}' > \varsigma_0$ . The Cauchy–Born approximation is obtained by computing the energy of the cells  $Q_j = (y_{j-1}, y_j)$  independently from one another, by treating each of them as part of a homogeneous chain (see Figure 4.1). We define the Cauchy–Born energy of the cell  $Q_j$  by

$$\mathcal{E}_j^{\text{cb}}(\mathbf{y}) = - \min_{\psi \in H_{\#}^1(Q_j)} \left( \int_{Q_j} \left( \frac{1}{2} \varepsilon^2 |\nabla \psi|^2 + \frac{1}{2} m^2 \psi^2 \right) dx - \int_{Q_j} \rho_{\mathbf{y}} \psi dx \right). \quad (4.1)$$

Note that this energy only depends on the distance  $(y_j - y_{j-1})$ . The minimizer  $\psi^{(j)}$  of (4.1) satisfies the equation  $-\varepsilon^2 \Delta \psi^{(j)} + m^2 \psi^{(j)} = \rho_{\mathbf{y}}$  in  $Q_j$  and its  $|Q_j|$ -periodic extension to  $\mathbb{R}$ :

$$-\varepsilon^2 \Delta \psi^{(j)} + m^2 \psi^{(j)} = \rho_{\mathbf{y}^{(j)}} \quad \text{in } \mathbb{R}. \quad (4.2)$$

Here we have defined the positions  $\mathbf{y}^{(j)} = (y_k^{(j)})_{k \in \mathbb{Z}}$  of an infinite chain of equidistant atoms by

$$y_k^{(j)} = y_j + (k - j)(y_j - y_{j-1}) \quad \forall k \in \mathbb{Z}. \quad (4.3)$$

The Cauchy–Born approximation  $\mathcal{E}^{\text{cb}}(\mathbf{y})$  of the atomistic energy  $\mathcal{E}(\mathbf{y})$  is then given by the sum over all cells

$$\mathcal{E}^{\text{cb}}(\mathbf{y}) = \sum_{j=-N}^N \mathcal{E}_j^{\text{cb}}(\mathbf{y}) = \frac{1}{2} \sum_{j=-N}^N \int_{Q_j} \rho_{\mathbf{y}} \psi^{(j)} dx. \quad (4.4)$$

In the Cauchy–Born model we seek to minimize the total potential energy  $E_{\mathbf{f}}^{\text{cb}} : \mathcal{Y} \rightarrow \mathbb{R}$  defined by

$$E_{\mathbf{f}}^{\text{cb}}(\mathbf{y}) = \mathcal{E}^{\text{cb}}(\mathbf{y}) + (\mathbf{f}, \mathbf{y})_\varepsilon. \quad (4.5)$$

Whether the Cauchy–Born model is a good approximation to the exact atomistic model strongly depends on the regularity properties of minimizers of (4.5).

Let  $\mathbf{u} \in \mathcal{U}$  be a test vector and  $u \in \mathcal{S}_{\#}(\mathbf{y})$  an interpolant of  $\mathbf{u}$ , i.e.,  $u(y_j) = u_j$  for  $j \in \mathbb{Z}$ . It follows as in Lemma 3.2 that the derivative of  $\mathcal{E}_j^{\text{cb}}(\mathbf{y})$  can be written in the form

$$D_{\mathbf{y}}\mathcal{E}_j^{\text{cb}}(\mathbf{y}) \cdot \mathbf{u} = \frac{u_j - u_{j-1}}{y_j - y_{j-1}} \int_{Q_j} \sigma_{j,\mathbf{y}}^{\text{cb}}(x) \, dx = \int_{Q_j} \sigma_{j,\mathbf{y}}^{\text{cb}}(x) \nabla u(x) \, dx, \quad (4.6)$$

where the local continuum stress function  $\sigma_{j,\mathbf{y}}^{\text{cb}}$ , in direct correspondence with (2.7), is

$$\begin{aligned} \sigma_{j,\mathbf{y}}^{\text{cb}}(x) &= \frac{1}{2} \varepsilon^2 |\nabla \psi^{(j)}(x)|^2 - \frac{1}{2} m^2 \psi^{(j)}(x)^2 + \rho_{\mathbf{y}}(x) \psi^{(j)}(x) \\ &\quad + \varepsilon \sum_{j=-N-1}^N \psi^{(j)}(x) \nabla \delta_{\varepsilon}(x - y_j)(x - y_j). \end{aligned} \quad (4.7)$$

Furthermore, we define the Cauchy–Born stress function  $\sigma_{\mathbf{y}}^{\text{cb}} : \Omega \rightarrow \mathbb{R}$  by

$$\sigma_{\mathbf{y}}^{\text{cb}}(x) = \sigma_{j,\mathbf{y}}^{\text{cb}}(x) \quad \text{if } x \in \Omega_j$$

for all  $x \in \Omega$ .

**4.1. Consistency.** Next, we turn to the consistency analysis of the Cauchy–Born approximation, for which we thoroughly analyze the modelling error incurred. From (2.6) and (4.6) we deduce that

$$\begin{aligned} |(D\mathcal{E}(\mathbf{y}) - D\mathcal{E}^{\text{cb}}(\mathbf{y})) \cdot \mathbf{u}| &\leq \int_{\Omega} |\sigma_{\mathbf{y}}(x) - \sigma_{\mathbf{y}}^{\text{cb}}(x)| |\nabla u(x)| \, dx \\ &= \sum_{j=-N}^N \int_{Q_j} |\sigma_{\mathbf{y}}(x) - \sigma_{j,\mathbf{y}}^{\text{cb}}(x)| |\nabla u(x)| \, dx, \end{aligned}$$

where the stress functions  $\sigma_{\mathbf{y}}$  and  $\sigma_{j,\mathbf{y}}^{\text{cb}}$  are given by (2.7) and (4.7), respectively. To investigate the modelling error  $|\sigma_{\mathbf{y}}(x) - \sigma_{j,\mathbf{y}}^{\text{cb}}(x)|$  incurred by going from the atomistic description to the Cauchy–Born approximation it is therefore sufficient to analyze  $|\phi - \psi^{(j)}|$  and  $|\nabla \phi - \nabla \psi^{(j)}|$  in  $Q_j$  for every  $j \in \{-N, \dots, N\}$ .

**Lemma 4.1.** *Let  $\mathbf{y} \in \ell^{\infty}(\mathbb{Z})$  and define  $\mathbf{y}^{(j)} = (y_k^{(j)})_{k \in \mathbb{Z}}$  by  $y_k^{(j)} = y_j + \varepsilon y'_j(k - j)$  for all  $k \in \mathbb{Z}$ ; then*

$$\begin{aligned} |y_n - y_n^{(j)}| &\leq (n - j) \varepsilon^2 \|\mathbf{y}''\|_{\ell^1([j, n-1])} \quad \text{for } n > j, \\ |y_n - y_n^{(j)}| &\leq (j - 1 - n) \varepsilon^2 \|\mathbf{y}''\|_{\ell^1([n+1, j-1])}, \quad \text{for } n < j - 1. \end{aligned}$$

*Proof.* Assume, without loss of generality that  $n > j$ . Since  $y_{j-1} = y_{j-1}^{(j)}$  and  $y_j = y_j^{(j)}$ ,

$$y_n - y_n^{(j)} = \varepsilon \sum_{k=j+1}^n (y'_k - (y_k^{(j)})') = \varepsilon^2 \sum_{k=j+1}^n \sum_{l=j}^{k-1} (y''_l - (y_l^{(j)})'') = \varepsilon^2 \sum_{k=j+1}^n \sum_{l=j}^{k-1} y''_l,$$

where we have used that  $(\mathbf{y}^{(j)})'$  is constant. Changing the order of summation we get

$$|y_n - y_n^{(j)}| \leq \varepsilon^2 \sum_{l=j}^{n-1} \sum_{k=l+1}^n |y''_l| = \varepsilon^2 \sum_{l=j}^{n-1} (n - l) y''_l \leq (n - j) \varepsilon^2 \|\mathbf{y}''\|_{\ell^1([j, n-1])}. \quad \square$$

In the next result we estimate the errors  $|\phi(x) - \psi^{(j)}(x)|$ ,  $|\nabla\phi(x) - \nabla\psi^{(j)}(x)|$  for  $x$  in the cell  $Q_j$ . As anticipated by Lemma 4.1 they depend on the second difference  $\mathbf{y}''$ .

**Lemma 4.2.** *Let  $\mathbf{y} \in \mathcal{Y}$  satisfy  $\min \mathbf{y}' > \varsigma_0$ . Let  $\phi \in H_{\#}^1(\Omega)$  satisfy (2.2) and  $\psi^{(j)} \in H_{\#}^1(Q_j)$  satisfy (4.2), respectively. Then,*

$$\begin{aligned} \|\phi - \psi^{(j)}\|_{L^\infty(Q_j)} &\leq \mu\varepsilon \sum_{n=1}^{\infty} \|\mathbf{y}''\|_{\ell^1([j-n, j+n-1])} n e^{-mn \min \mathbf{y}'}, \quad \text{and} \\ \|\varepsilon\nabla\phi - \varepsilon\nabla\psi^{(j)}\|_{L^\infty(Q_j)} &\leq m\mu\varepsilon \sum_{n=1}^{\infty} \|\mathbf{y}''\|_{\ell^1([j-n, j+n-1])} n e^{-mn \min \mathbf{y}'}. \end{aligned}$$

*Proof.* From Proposition 2.1 we immediately deduce that, for all  $x \in Q_j$ ,

$$\begin{aligned} \phi(x) &= \frac{1}{2m} \int_{\mathbb{R}} \sum_{k \in \mathbb{Z}} \delta_\varepsilon(z - y_k) e^{-\frac{m}{\varepsilon}|x-z|} dz, \\ \psi^{(j)}(x) &= \frac{1}{2m} \int_{\mathbb{R}} \sum_{k \in \mathbb{Z}} \delta_\varepsilon(z - y_k^{(j)}) e^{-\frac{m}{\varepsilon}|x-z|} dz. \end{aligned} \tag{4.8}$$

Since  $y_j^{(j)} = y_j$  and  $y_{j-1}^{(j)} = y_{j-1}$ , the respective terms in the sums cancel. Hence, we get for  $x \in Q_j$ :

$$\phi(x) - \psi^{(j)}(x) = \frac{1}{2m} \sum_{\substack{k \in \mathbb{Z} \\ k \neq j-1, j}} \int_{\mathbb{R}} (\delta_\varepsilon(z - y_k) - \delta_\varepsilon(z - y_k^{(j)})) e^{-\frac{m}{\varepsilon}|x-z|} dz.$$

We now derive bounds on the individual terms in the sum. Note that (2.12) simplifies the following calculations but due to the smoothness of the Green's function similar bounds can be obtained without it.

Let  $k > j$ . Then we have  $|x - z| = z - x$  for all  $z \in \text{supp} \delta_\varepsilon(\cdot - y_k)$  and all  $z \in \text{supp} \delta_\varepsilon(\cdot - y_k^{(j)})$ . Thus, with (2.12),

$$\frac{1}{2m} \int_{\mathbb{R}} (\delta_\varepsilon(z - y_k) - \delta_\varepsilon(z - y_k^{(j)})) e^{-\frac{m}{\varepsilon}|x-z|} dz = \frac{\mu}{2m} (e^{-\frac{m}{\varepsilon}(y_k - x)} - e^{-\frac{m}{\varepsilon}(y_k^{(j)} - x)}). \tag{4.9}$$

If  $y_k^{(j)} \geq y_k$ , then

$$\begin{aligned} \left| \frac{1}{2m} \int_{\mathbb{R}} (\delta_\varepsilon(z - y_k) - \delta_\varepsilon(z - y_k^{(j)})) e^{-\frac{m}{\varepsilon}|x-z|} dz \right| &\leq \frac{\mu}{2m} e^{-\frac{m}{\varepsilon}(y_k - x)} (1 - e^{-\frac{m}{\varepsilon}(y_k^{(j)} - y_k)}) \\ &\leq \frac{\mu}{2m} e^{-\frac{m}{\varepsilon}(y_k - x)} \frac{m}{\varepsilon} (y_k^{(j)} - y_k). \end{aligned}$$

Using  $(y_k - x) \geq (k - j)\varepsilon \min \mathbf{y}'$  for all  $x \in Q_j$  and applying Lemma 4.1 leads to

$$\frac{\mu}{2\varepsilon} e^{-\frac{m}{\varepsilon}(y_k - x)} |y_k - y_k^{(j)}| \leq \frac{\mu\varepsilon}{2} \|\mathbf{y}''\|_{\ell^1([j, k-1])} (k - j) e^{-(k-j)m \min \mathbf{y}'}$$

The same bound on (4.9) can be obtained if  $y_k^{(j)} \leq y_k$ .

For any  $k < j - 1$  we can use the same techniques to obtain that

$$\begin{aligned} \left| \frac{1}{2m} \int_{\mathbb{R}} (\delta_\varepsilon(z - y_k) - \delta_\varepsilon(z - y_k^{(j)})) e^{-\frac{m}{\varepsilon}|x-z|} dz \right| &\leq \frac{\mu\varepsilon}{2} \|\mathbf{y}''\|_{\ell^1([k+1, j-1])} (j - k - 1) e^{-(j-k-1)m \min \mathbf{y}'}. \end{aligned}$$

Summing over all  $k \in \mathbb{Z} \setminus \{j-1, j\}$  we deduce that

$$|\phi(x) - \psi^{(j)}(x)| \leq \mu\varepsilon \sum_{n=1}^{\infty} \|\mathbf{y}''\|_{\ell^1([j-n, j+n-1])} n e^{-mn \min \mathbf{y}'}$$

The proof for the derivatives  $\nabla\phi, \nabla\psi^{(j)}$  is analogous.  $\square$

We wish to prove modelling error estimates on  $\|\sigma_{\mathbf{y}} - \sigma_{j,\mathbf{y}}^{\text{cb}}\|_{L^\infty(Q_j)}$  in terms of  $\|\phi - \psi^{(j)}\|_{L^\infty(Q_j)}$  and  $\|\nabla\phi - \nabla\psi^{(j)}\|_{L^\infty(Q_j)}$ . Since the stress functions  $\sigma_{\mathbf{y}}$  and  $\sigma_{j,\mathbf{y}}^{\text{cb}}$  are quadratic in the fields  $\phi$  and  $\psi^{(j)}$  we need  $L^\infty$ -bounds, which we establish in the next lemma.

**Lemma 4.3.** *Let  $\mathbf{y} \in \mathcal{Y}$ ,  $\mathbf{y}' > \varsigma_0$ , and let  $\phi = \arg \min_{\varphi \in H_{\#}^1(\Omega)} I(\varphi, \mathbf{y})$  be the corresponding field. Then, there are continuous functions  $K_0, K_1$ , that depend implicitly on  $m$  (but are independent of  $\varepsilon$  and  $\mathbf{y}$ ), such that*

$$\|\phi\|_{L^\infty(\Omega)} \leq K_0(m \min \mathbf{y}'), \quad \text{and} \quad \varepsilon \|\nabla\phi\|_{L^\infty(\Omega)} \leq K_1(m \min \mathbf{y}').$$

*Proof.* The stated estimates follow in a straightforward manner from the integral representation of the solution  $\phi$ ; see [10, Lemma 4.4] for the details.  $\square$

We can now prove the following modelling error estimates.

**Lemma 4.4.** *Let  $\sigma_{\mathbf{y}}$  and  $\sigma_{j,\mathbf{y}}^{\text{cb}}$  be given by (2.7), respectively, (4.7); then*

$$\|\sigma_{\mathbf{y}} - \sigma_{j,\mathbf{y}}^{\text{cb}}\|_{L^\infty(Q_j)} \leq C(\varepsilon \|\nabla\phi - \nabla\psi^{(j)}\|_{L^\infty(Q_j)} + \|\phi - \psi^{(j)}\|_{L^\infty(Q_j)}), \quad j = 1, \dots, N,$$

where the constant  $C$  only depends on  $\delta_1, K_i = K_i(m\mathbf{y}')$ , and on  $m$ .

*Proof.* From the definitions of the atomistic and continuum stress function we deduce that

$$\begin{aligned} \sigma_{\mathbf{y}}(x) - \sigma_{j,\mathbf{y}}^{\text{cb}}(x) &= -\frac{1}{2}(\varepsilon\nabla\phi(x) - \varepsilon\nabla\psi^{(j)}(x))(\varepsilon\nabla\phi(x) + \varepsilon\nabla\psi^{(j)}(x)) \\ &\quad + \frac{1}{2}m^2(\phi(x) - \psi^{(j)}(x))(\phi(x) + \psi^{(j)}(x)) - \rho_{\mathbf{y}}(x)(\phi(x) - \psi^{(j)}(x)) \\ &\quad - (\phi(x) - \psi^{(j)}(x)) \sum_{i=j-1}^j \varepsilon\nabla\delta_\varepsilon(x - y_i)(x - y_i) \end{aligned}$$

for all  $x \in Q_j$ . With  $\delta_\varepsilon(x) = \varepsilon^{-1}\delta_1(x/\varepsilon)$ , the  $L^\infty$ -bound on  $\phi$  from Lemma 4.3, and the analogous bound for  $\psi^{(j)}$  we get

$$\begin{aligned} \frac{1}{2}|\varepsilon\nabla\phi(x) + \varepsilon\nabla\psi^{(j)}(x)| &\leq K_1(m \min \mathbf{y}'), \\ \frac{m^2}{2}|\phi(x) + \psi^{(j)}(x)| &\leq m^2 K_0(m \min \mathbf{y}'), \\ \|\rho_{\mathbf{y}}\|_{L^\infty} &\leq \|\delta_1\|_{L^\infty}, \\ |\varepsilon\nabla\delta_\varepsilon(x - y_i)(x - y_i)| &\leq \|\nabla\delta_1 \text{id}\|_{L^\infty}, \end{aligned}$$

which implies the stated result.  $\square$

**4.2. Stability.** Besides consistency, the second crucial property of an approximation to a given model is its stability. The following auxiliary result will play a role in the stability analysis of a/c methods.

**Lemma 4.5.** *Let  $\mathbf{y} \in \mathcal{Y}$  satisfy  $\min \mathbf{y}' > \varsigma_0$ . Then, for all  $j \in \{-N, \dots, N\}$ ,*

$$D^2 \mathcal{E}_j^{\text{cb}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}] \geq \frac{m^2 \mu^2}{2} e^{-m \max \mathbf{y}'} \varepsilon |u'_j|^2 \quad \forall \mathbf{u} \in \mathcal{U}.$$

*Proof.* We first recall that  $\mathcal{E}_j^{\text{cb}}(\mathbf{y}) = \frac{1}{2} \int_{Q_j} \rho_{\mathbf{y}} \psi^{(j)} dx$  because  $\psi^{(j)}$  is a minimizer of (4.1). Extending  $\psi^{(j)}$   $|Q_j|$ -periodically to  $\mathbb{R}$  and using the symmetry of the cell problem, we can rewrite this as

$$\mathcal{E}_j^{\text{cb}}(\mathbf{y}) = \frac{\varepsilon}{2} \int_{\mathbb{R}} \delta_\varepsilon(x - y_j) \psi^{(j)}(x) dx.$$

We now insert the explicit formula (4.8) for  $\psi^{(j)}(x)$  and apply (2.12) to get

$$\begin{aligned} \mathcal{E}_j^{\text{cb}}(\mathbf{y}) &= \frac{\varepsilon}{4m} \sum_{k \in \mathbb{Z}} \int_{\mathbb{R}} \int_{\mathbb{R}} \delta_\varepsilon(x - y_j) \delta_\varepsilon(z - y_k^{(j)}) e^{-\frac{m}{\varepsilon} |x-z|} dz dx \\ &= \frac{\mu^2 \varepsilon}{4m} \sum_{\substack{k \in \mathbb{Z} \\ k \neq j}} e^{-\frac{m}{\varepsilon} |y_j - y_k^{(j)}|} + \mathcal{E}_{\text{self}} = \frac{\mu^2 \varepsilon}{2m} \sum_{\nu=1}^{\infty} e^{-m \nu y'_j} + \mathcal{E}_{\text{self}}, \end{aligned}$$

where the constant  $\mathcal{E}_{\text{self}}$  coming from  $k = j$  in the sum represents the self-energies of the atoms in the cell  $Q_j$ . Here we have also used that  $|y_k^{(j)} - y_j| = |k - j| y'_j$  for all  $k \in \mathbb{Z}$ . Differentiating twice leads to

$$\begin{aligned} D^2 \mathcal{E}_j^{\text{cb}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}] &= \frac{m \mu^2}{2} \varepsilon \sum_{\nu=1}^{\infty} \nu^2 e^{-\nu m y'_j} |u'_j|^2 \\ &\geq \frac{m \mu^2}{2} \varepsilon |u'_j|^2 \sum_{\nu=1}^{\infty} \nu^2 e^{-\nu m \max \mathbf{y}'} \geq \frac{m \mu^2}{2} e^{-m \max \mathbf{y}'} \varepsilon |u'_j|^2. \end{aligned}$$

In the last step we have only kept the term for  $\nu = 1$ , which represents the nearest neighbour interactions.  $\square$

## 5. ATOMISTIC-TO-CONTINUUM COUPLING

The computation of the original atomistic energy  $\mathcal{E}(\mathbf{y})$  involves the solution of the optimization problem (2.1) posed in the whole of  $\Omega = (y_{-N-1}, y_N)$ . Our goal is the construction of computationally cheaper, approximate energies  $\mathcal{E}^{\text{ac}}(\mathbf{y})$  such that  $\mathcal{E}(\mathbf{y}) \approx \mathcal{E}^{\text{ac}}(\mathbf{y})$  for all relevant  $\mathbf{y}$  and minimizers  $\bar{\mathbf{y}}^{\text{ac}} \in \mathcal{Y}$  of

$$E_{\mathbf{f}}^{\text{ac}}(\mathbf{y}) = \mathcal{E}^{\text{ac}}(\mathbf{y}) + (\mathbf{f}, \mathbf{y})_\varepsilon,$$

are good approximations of minimizers  $\bar{\mathbf{y}}$  of the energy  $E_{\mathbf{f}}$  from (2.3).

Following the philosophy of a/c methods we approximate  $\mathcal{E}(\mathbf{y})$  by the continuum model where  $\mathbf{y}$  is smooth and a version of the atomistic model where  $\mathbf{y}$  is nonsmooth. In the following we will implicitly assume that the configurations  $\mathbf{y} \in \mathcal{Y}$  under consideration are smooth except in the segment  $y_{-K}, \dots, y_K$  for some  $K < N$ . We divide  $\Omega$  into an atomistic subdomain  $\Omega^{\text{at}}$  such that  $y_j \in \Omega^{\text{at}}$  for all  $j \in \{-K, \dots, K\}$  and the continuum domain  $\Omega^{\text{cb}} = \Omega \setminus \Omega^{\text{at}}$ . In  $\Omega^{\text{cb}}$  we will use the Cauchy–Born approximation on a cell-by-cell basis. In  $\Omega^{\text{at}}$  we will use the atomistic model with Dirichlet boundary conditions as discussed in Section 3.

This basic setting gives rise to a variety of possibilities including the precise choice of  $\partial \Omega^{\text{at}}$  and the boundary conditions imposed on the atomistic subproblem. Both will in general depend




FIGURE 5.1. An illustration of the first a/c method. In  $\Omega^{\text{at}} = (a_L(\mathbf{y}), a_R(\mathbf{y}))$  the atomistic problem is solved with the Dirichlet boundary conditions  $g^*(\mathbf{y})$ . Outside  $\Omega^{\text{at}}$  the Cauchy–Born approximation is used in all cells  $Q_j$ .

on the configuration  $\mathbf{y}$ . Our main objective for  $\mathcal{E}^{\text{ac}}$  is the existence of a weak formulation in the sense that

$$D\mathcal{E}^{\text{ac}}(\mathbf{y}) \cdot \mathbf{u} = \int_{\Omega} \sigma_{\mathbf{y}}^{\text{ac}}(x) \nabla u(x) \, dx,$$

where  $u \in \mathcal{S}_{\#}(\mathbf{y})$  is a piecewise linear interpolant of  $\mathbf{u} \in \mathcal{U}$  and  $\sigma_{\mathbf{y}}^{\text{ac}}$  is a stress function to be determined. If this weak formulation can be obtained, the consistency analysis reduces to error estimates on fields, as already seen in Lemma 4.4.

Throughout this section,  $\phi \in H_{\#}^1(\Omega)$  denotes the solution of the original minimization problem (2.1) for a given configuration  $\mathbf{y} \in \mathcal{Y}$ .

**5.1. An a/c method with optimal boundary conditions.** We place the boundary  $a$  of the atomistic subproblem halfway between the interface atoms, that is  $a = a(\mathbf{y}) = [a_L(\mathbf{y}) \ a_R(\mathbf{y})]^{\text{T}}$ , where

$$a_L(\mathbf{y}) = \frac{y_{-K-1} + y_{-K}}{2}, \quad a_R(\mathbf{y}) = \frac{y_K + y_{K+1}}{2}.$$

Let  $\Omega^{\text{at}} = (a_L(\mathbf{y}), a_R(\mathbf{y}))$  and  $\Omega^{\text{cb}} = \Omega \setminus \Omega^{\text{at}}$ . We write the a/c energy  $\mathcal{E}^{\text{ac}}(\mathbf{y})$  as the sum of a continuum and an atomistic part

$$\mathcal{E}^{\text{ac}}(\mathbf{y}) = \mathcal{E}_*^{\text{cb}}(\mathbf{y}) + \mathcal{E}_*^{\text{at}}(\mathbf{y}), \quad (5.1)$$

which are introduced below.

Due to the choice of  $a(\mathbf{y})$  there are two half cells,  $(y_{-K-1}, a_L(\mathbf{y}))$  and  $(a_R(\mathbf{y}), y_{K+1})$ , in the continuum region  $\Omega^{\text{cb}}$  (see Figure 5.1). Since the cell problems are symmetric, the Cauchy–Born energies of these half cells are given by  $\frac{1}{2}\mathcal{E}_{-K}^{\text{cb}}(\mathbf{y})$  and  $\frac{1}{2}\mathcal{E}_{K+1}^{\text{cb}}(\mathbf{y})$ , respectively. Hence, the continuum contribution to the energy  $\mathcal{E}^{\text{ac}}$  is defined by

$$\mathcal{E}_*^{\text{cb}}(\mathbf{y}) = \sum_{j=-N+1}^{-K-1} \mathcal{E}_j^{\text{cb}}(\mathbf{y}) + \frac{1}{2}\mathcal{E}_{-K}^{\text{cb}}(\mathbf{y}) + \frac{1}{2}\mathcal{E}_{K+1}^{\text{cb}}(\mathbf{y}) + \sum_{j=K+2}^N \mathcal{E}_j^{\text{cb}}(\mathbf{y}). \quad (5.2)$$

The coordinates of the atoms in the atomistic region  $\Omega^{\text{at}}$  are represented by

$$\mathbf{y}_{\text{at}} = (y_{-K}, \dots, y_K)^{\text{T}}.$$

For the definition of  $\mathcal{E}_*^{\text{at}}(\mathbf{y})$  we consider the minimization problem (3.2) on the atomistic domain  $\Omega^{\text{at}}$  subject to the Dirichlet boundary conditions  $g^*(\mathbf{y}) = [g_L^*(\mathbf{y}) \ g_R^*(\mathbf{y})]^{\text{T}}$ . In correspondence with Remark 4 and Section 3.3 they are given by

$$g_L^*(\mathbf{y}) = \frac{1}{1-\tau} \frac{\gamma_L(\mathbf{y}) + \tau\gamma_R(\mathbf{y})}{1+\tau}, \quad g_R^*(\mathbf{y}) = \frac{1}{1-\tau} \frac{\tau\gamma_L(\mathbf{y}) + \gamma_R(\mathbf{y})}{1+\tau},$$

where  $\tau = e^{-\frac{m}{\varepsilon}\Delta a(\mathbf{y})}$ , and  $\gamma_L, \gamma_R$  are defined in (3.15). The energy contribution from the atomistic subproblem is thus given by

$$\mathcal{E}_*^{\text{at}}(\mathbf{y}) = \mathcal{E}_{a(\mathbf{y}), g^*(\mathbf{y})}(\mathbf{y}_{\text{at}}) = -\inf \left\{ I_{a(\mathbf{y})}(\varphi, \mathbf{y}_{\text{at}}) : \varphi \in H^1(\Omega^{\text{at}}), \varphi|_{\partial\Omega^{\text{at}}} = g^*(\mathbf{y}) \right\},$$

where  $I_{a(\mathbf{y})}$  is defined as in (3.3). We denote the solution of this optimization problem by  $\phi_{\text{at}}^* \in H^1(\Omega^{\text{at}})$ . It satisfies the boundary-value problem

$$\begin{aligned} -\varepsilon^2 \Delta \phi_{\text{at}}^* + m^2 \phi_{\text{at}}^* &= \rho_{\mathbf{y}} \text{ in } \Omega^{\text{at}}, \\ \phi_{\text{at}}^*|_{\partial\Omega^{\text{at}}} &= g^*(\mathbf{y}). \end{aligned}$$

From a computational point of view  $g^*(\mathbf{y})$  is also a convenient choice since this is equivalent to homogeneous Neumann boundary conditions. In Section 3.3 we deduced a clear interpretation of the effect of this choice of boundary data: besides the interaction among themselves, the atoms in  $\Omega^{\text{at}}$  interact with mirror atoms outside  $\Omega^{\text{at}}$ . This is closely related to the geometric reconstruction idea for classical potentials [17, 4].

In analogy to (2.3) we search for minimizers of the total potential energy

$$E_{\mathbf{f}}^{\text{ac}}(\mathbf{y}) = \mathcal{E}^{\text{ac}}(\mathbf{y}) + (\mathbf{f}, \mathbf{y})_{\varepsilon} \quad (5.3)$$

in  $\mathcal{Y}$ , where  $\mathbf{f} \in \mathcal{U}^{-1,2}$  represents an external force. Formally, a minimizer  $\bar{\mathbf{y}}^{\text{ac}}$  satisfies the following Euler–Lagrange equation in  $\mathcal{U}^{-1,2}$ :

$$DE_{\mathbf{f}}^{\text{ac}}(\mathbf{y}) = D\mathcal{E}^{\text{ac}}(\mathbf{y}) + \mathbf{f} = \mathbf{0}.$$

Throughout the remainder of this article we assume that the atomistic domain  $\Omega^{\text{at}}$  is large compared with  $\varepsilon$ , that is  $\Delta a \gg \varepsilon$  and hence terms of order  $\mathcal{O}(\tau)$  are exponentially small. To keep the notation more compact we will not give precise estimates of  $\tau$ -dependent terms arising from the atomistic domain explicitly but include an  $\mathcal{O}(\tau)$  where necessary.

**5.2. Consistency.** In order to study the consistency properties of the a/c energy  $\mathcal{E}^{\text{ac}}(\mathbf{y})$  from (5.1) we first need to calculate its derivative. Having established weak formulations for the derivatives of  $\mathcal{E}$ ,  $\mathcal{E}^{\text{cb}}$ , as well as  $\mathcal{E}_{a,g}$ , we will prove that the a/c energy  $\mathcal{E}^{\text{ac}}$  admits a similar reformulation of  $DE^{\text{ac}}(\mathbf{y}) \cdot \mathbf{u}$ . For this we have to take into account that both the boundary of the atomistic domain  $\Omega^{\text{at}}$  and the boundary conditions depend on  $\mathbf{y}$ . The necessary preparations were carried out in Section 3.

**Lemma 4.1.** *Let  $\mathbf{y} \in \mathcal{Y}$  satisfy  $\min \mathbf{y}' > \varsigma_0$ . Furthermore, let  $\mathbf{u} \in \mathcal{U}$  be a test vector and  $u \in S_{\#}(\mathbf{y})$  an interpolant of  $\mathbf{u}$ ; then,*

$$D\mathcal{E}^{\text{ac}}(\mathbf{y}) \cdot \mathbf{u} = \int_{\Omega} \sigma_{\mathbf{y}}^{\text{ac}}(x) \nabla u(x) \, dx, \quad \text{where } \sigma_{\mathbf{y}}^{\text{ac}}(x) = \begin{cases} \sigma_{\mathbf{y}}^{\text{cb}}(x) & \text{if } x \in \Omega^{\text{cb}}, \\ \sigma_{\mathbf{y},*}^{\text{at}}(x) & \text{if } x \in \Omega^{\text{at}}, \end{cases} \quad (5.4)$$

and  $\sigma_{\mathbf{y},*}^{\text{at}}(x)$  is given by (2.7) with  $\phi = \phi_{\text{at}}^*$ .

*Proof. 1. Continuum Contribution.* From Section 4 we already have the equality

$$D\mathcal{E}_j^{\text{cb}}(\mathbf{y}) \cdot \mathbf{u} = \int_{Q_j} \sigma_{\mathbf{y},j}^{\text{cb}}(x) \nabla u(x) \, dx,$$

$j \in \{-N, \dots, -K-1\} \cup \{K+2, \dots, N\}$ . For the contribution  $\frac{1}{2}\mathcal{E}_{-K}^{\text{cb}}(\mathbf{y})$  from the half cell  $(y_{-K-1}, a_L(\mathbf{y}))$  we make use of the symmetry of the cell problems. Since  $\nabla u|_{Q_{-K}}$  is constant,

$a_L(\mathbf{y})$  is the midpoint of  $Q_{-K} = (y_{-K-1}, y_{-K})$ , and  $\sigma_{\mathbf{y}, -K}^{\text{cb}}$  is symmetric in  $Q_{-K}$ , we deduce that

$$\frac{1}{2} D\mathcal{E}_{-K}^{\text{cb}}(\mathbf{y}) \cdot \mathbf{u} = \frac{1}{2} \int_{Q_{-K}} \sigma_{\mathbf{y}, -K}^{\text{cb}}(x) \nabla u(x) \, dx = \int_{y_{-K-1}}^{a_L(\mathbf{y})} \sigma_{\mathbf{y}, -K}^{\text{cb}}(x) \nabla u(x) \, dx.$$

We treat  $\frac{1}{2} \mathcal{E}_{K+1}^{\text{cb}}(\mathbf{y})$  analogously. Hence,

$$D\mathcal{E}_*^{\text{cb}}(\mathbf{y}) \cdot \mathbf{u} = \int_{\Omega^{\text{cb}}} \sigma_{\mathbf{y}}^{\text{cb}}(x) \nabla u(x) \, dx,$$

where  $\sigma_{\mathbf{y}}^{\text{cb}}(x) = \sigma_{\mathbf{y}, j}^{\text{cb}}(x)$  if  $x \in Q_j$ .

2. *Atomistic Contribution.* To calculate the derivative  $D\mathcal{E}_*^{\text{at}}(\mathbf{y})$  we use the chain rule and the derivatives that were provided in Section 3. Applying Proposition 3.1 (with  $h_L = (u_{-K-1} + u_{-K})/2$ ,  $h_R = (u_K + u_{K+1})/2$  because of  $D_{\mathbf{y}} a(\mathbf{y}) \cdot \mathbf{u} = a(\mathbf{u})$ ), we get

$$\begin{aligned} D\mathcal{E}_*^{\text{at}}(\mathbf{y}) \cdot \mathbf{u} &= D_{\mathbf{y}} \mathcal{E}_{a(\mathbf{y}), g^*(\mathbf{y})}(\mathbf{y}_{\text{at}}) \cdot \mathbf{u}_{\text{at}} + D_a \mathcal{E}_{a(\mathbf{y}), g^*(\mathbf{y})}(\mathbf{y}_{\text{at}}) \cdot D_{\mathbf{y}} a(\mathbf{y}) \cdot \mathbf{u} \\ &= \int_{\Omega^{\text{at}}} \sigma_{\mathbf{y}, * }^{\text{at}}(x) \nabla u(x) \, dx, \end{aligned} \quad (5.5)$$

where the stress  $\sigma_{\mathbf{y}, * }^{\text{at}}$  is given by (2.7) with  $\phi = \phi_{\text{at}}^*$  and  $\mathbf{u}_{\text{at}} = (u_{-K}, \dots, u_K) \in \mathbb{R}^{2K+1}$  is the section of  $\mathbf{u}$  corresponding to the atoms in the atomistic region. Note that the choice of boundary conditions implies  $D_g \mathcal{E}_{a(\mathbf{y}), g^*(\mathbf{y})}(\mathbf{y}_{\text{at}}) = 0$ ; cf. Remark 4.  $\square$

**Remark 5.** The weak form (5.4) of the derivative  $D\mathcal{E}^{\text{ac}}$  already implies that there are no ghost forces for homogeneous deformations  $\mathbf{y}$ . If the atoms are equidistant, then  $g_L^*(\mathbf{y}) = \phi(a_L)$  and  $g_R^*(\mathbf{y}) = \phi(a_R)$  and thus also  $\phi_{\text{at}}^* = \phi$  in  $\Omega^{\text{at}}$ . Moreover, it is clear that  $\psi^{(j)} = \phi$  for all  $j$ . Hence, we obtain that  $\sigma_{\mathbf{y}}^{\text{ac}}(x) = \sigma_{\mathbf{y}}(x)$  for all  $x \in \Omega$ , which implies that  $D\mathcal{E}^{\text{ac}}(\mathbf{y}) = D\mathcal{E}(\mathbf{y}) = 0$  for all  $\mathbf{y} = F\widehat{\mathbf{X}} \in \mathcal{Y}$  representing homogeneous deformations (i.e., that the method exhibits no ghost forces).  $\square$

Absence of ghost forces does not immediately imply *consistency* of the a/c method, but has to be shown separately. This we do next. Because of the structure of the weak formulation (5.4), the analysis boils down to estimating the errors between the field  $\phi$  coming from the original atomistic model and the fields  $\psi^{(j)}$ , respectively,  $\phi_{\text{at}}^*$ .

**Theorem 5.1.** *Let  $\mathbf{y} \in \mathcal{Y}$  be such that  $\min \mathbf{y}' \geq s_0 > s_0$ ; then, for all  $\mathbf{u} \in \mathcal{U}$  with interpolants  $u \in \mathcal{S}_{\#}(\mathbf{y})$ ,*

$$|(D\mathcal{E}(\mathbf{y}) - D\mathcal{E}^{\text{ac}}(\mathbf{y})) \cdot \mathbf{u}| \leq C(\varepsilon \|\mathbf{y}''\|_{\ell_{w, s_0}^2} + \tau) \|\nabla u\|_{L^2},$$

where  $C = C(s_0)$  and the weighted  $\ell_{w, s_0}^2$ -norm is defined by

$$\|\mathbf{y}''\|_{\ell_{w, s_0}^2}^2 := \varepsilon \sum_{j=-N}^N w_j |y_j''|^2, \quad (5.6)$$

with weights  $w_j := \max(1, e^{-ms_0 \text{dist}(j, \{-K, K\})})$ .

*Proof.* Using the weak formulation (5.4) of  $D\mathcal{E}^{\text{ac}}(\mathbf{y})$  we obtain

$$\begin{aligned} |(D_{\mathbf{y}} \mathcal{E}(\mathbf{y}) - D_{\mathbf{y}} \mathcal{E}^{\text{ac}}(\mathbf{y})) \cdot \mathbf{u}| &= \left| \int_{\Omega} (\sigma_{\mathbf{y}}(x) - \sigma_{\mathbf{y}}^{\text{qc}}(x)) \nabla u(x) \, dx \right| \\ &\leq \|\sigma_{\mathbf{y}} - \sigma_{\mathbf{y}}^{\text{qc}}\|_{L^2(\Omega)} \|\nabla u\|_{L^2} \\ &\leq \left( \sum_{j=-N}^N \varepsilon \|\sigma_{\mathbf{y}} - \sigma_{\mathbf{y}}^{\text{qc}}\|_{L^\infty(Q_j)}^2 \right)^{1/2} \cdot \|\nabla u\|_{L^2}. \end{aligned} \quad (5.7)$$

For  $Q_j$  belonging to the continuum region Lemma 4.4 and Lemma 4.2 imply

$$\begin{aligned}
\|\sigma_{\mathbf{y}} - \sigma_{\mathbf{y}}^{\text{qc}}\|_{L^\infty(Q_j)} &\leq C\varepsilon \sum_{n=1}^{\infty} \|\mathbf{y}''\|_{\ell^1([j-n, j+n-1])} n e^{-mns_0} \\
&\leq C\varepsilon \sum_{n=1}^{\infty} \|\mathbf{y}''\|_{\ell^2([j-n, j+n-1])} n^{3/2} e^{-mns_0} \\
&\leq C\varepsilon \left( \sum_{n=1}^{\infty} \|\mathbf{y}''\|_{\ell^2([j-n, j+n-1])}^2 e^{-mns_0} \right)^{1/2}, \tag{5.8}
\end{aligned}$$

where we have employed the Cauchy–Schwarz inequality twice and used the fact that the series  $\sum_{n=1}^{\infty} n^3 e^{-mns_0}$  is convergent.

Summing over all cells belonging to the continuum region and interchanging the order of summation we obtain

$$\begin{aligned}
\sum_{\substack{j \in \{-N, \dots, N\} \\ \setminus \{-K+1, \dots, K\}}} \varepsilon \|\sigma_{\mathbf{y}} - \sigma_{\mathbf{y}}^{\text{qc}}\|_{L^\infty(Q_j)}^2 &\leq C\varepsilon^3 \sum_{\substack{j \in \{-N, \dots, N\} \\ \setminus \{-K+1, \dots, K\}}} \sum_{n=1}^{\infty} \|\mathbf{y}''\|_{\ell^2([j-n, j+n-1])}^2 e^{-mns_0} \\
&\leq C\varepsilon^3 \sum_{k=-N}^N w'_k |y''_k|^2,
\end{aligned}$$

where

$$w'_k = \sum_{\substack{j \in \{-N, \dots, N\} \\ \setminus \{-K+1, \dots, K\}}} \sum_{\substack{n=1, \dots, \infty \\ k \in [j-n, j+n-1]}} e^{-mns_0}.$$

This is a geometric series from which we can factor out  $e^{-ms_0 \text{dist}(k, \{-K, K\})}$ , and hence we obtain  $w'_k \leq Cw_k$ , which gives

$$\sum_{j \notin \{-K+1, \dots, K\}} \varepsilon \|\sigma_{\mathbf{y}} - \sigma_{\mathbf{y}}^{\text{qc}}\|_{L^\infty(Q_j)}^2 \leq C\varepsilon^3 \sum_{k=-N}^N w_k |y''_k|^2. \tag{5.9}$$

To compute the consistency error of the weak form in the atomistic region, we need to bound the difference  $\|\sigma_{\mathbf{y}} - \sigma_{\mathbf{y}}^{\text{qc}}\|_{L^\infty(Q_j)} = \|\sigma_{\mathbf{y}} - \sigma_{\mathbf{y},*}^{\text{at}}\|_{L^\infty(Q_j)}$  for  $Q_j \subset \Omega^{\text{at}}$ . Using the same arguments as in the proof of Lemma 4.4 we obtain

$$\|\sigma_{\mathbf{y}} - \sigma_{\mathbf{y},*}^{\text{at}}\|_{L^\infty(Q_j)} \leq C(\|\phi - \phi_{\text{at}}^*\|_{L^\infty(Q_j)} + \varepsilon \|\nabla \phi - \nabla \phi_{\text{at}}^*\|_{L^\infty(Q_j)}).$$

Lemma 4.1 implies

$$\|\sigma_{\mathbf{y}} - \sigma_{\mathbf{y},*}^{\text{at}}\|_{L^\infty(Q_j)} \leq C(|\phi(a_L) - g_L^*(\mathbf{y})| + |\phi(a_R) - g_R^*(\mathbf{y})|) e^{-\frac{m}{\varepsilon} \min_{x \in Q_j} d_a(x)}.$$

Next, we recall from Remark 4 that  $g_R^* = \gamma_R^* + \mathcal{O}(\tau)$ , which is given by (cf. Remark 4)

$$\gamma_R(\mathbf{y}) = \int_{\mathbb{R}} \rho_{\mathbf{y}}^{\text{refl}}(x) G_\varepsilon(a_R - x) dx + \mathcal{O}(\tau),$$

where  $\rho_{\mathbf{y}}^{\text{refl}}(z) = \sum_{j \in \mathbb{Z}} \delta_\varepsilon(z - y_j^{\text{refl}})$  and  $\mathbf{y}^{\text{refl}}$  is a reflected and periodized extension of  $(y_j)_{j=-K}^K$ . Hence, we obtain

$$\begin{aligned}
|\phi(a_R) - g_R^*(\mathbf{y})| &= |\phi(a_R) - \gamma_L(\mathbf{y})| + \mathcal{O}(\tau) \\
&\leq \frac{1}{2m\varepsilon} \left| \int_{\mathbb{R}} (\rho_{\mathbf{y}}(z) - \rho_{\mathbf{y}}^{\text{refl}}(z)) e^{-\frac{m}{\varepsilon} |a_R - z|} dz \right| + \mathcal{O}(\tau).
\end{aligned}$$

Minor modifications of the proofs of Lemma 4.2 and Lemma 4.1 yield

$$\begin{aligned} |\phi(a_R) - g_R^*(\mathbf{y})| &\leq \frac{C}{\varepsilon} \sum_{j=K+1}^{\infty} |y_j^{\text{refl}} - y_j| e^{-\frac{m}{\varepsilon}(\min(y_j^{\text{refl}}, y_j) - a_R)} + \mathcal{O}(\tau) \\ &\leq C\varepsilon \sum_{n=1}^{\infty} \|\mathbf{y}''\|_{\ell^1([K-n+1, K+n])} n e^{-mns_0} + \mathcal{O}(\tau). \end{aligned} \quad (5.10)$$

An analogous result holds for  $|\phi(a_L) - g_L^*(\mathbf{y})|$ . It is now straightforward to see that the consistency error committed in the atomistic region can be bounded above in the same way as the consistency error committed in the continuum region (in fact it is dominated by (5.9)). This completes the proof.  $\square$

**5.3. Stability.** The special choice  $g^*(\mathbf{y})$  of boundary conditions for the atomistic subproblem allows for an elementary stability analysis of  $\mathcal{E}^{\text{ac}}(\mathbf{y})$  that draws from the ideas we used in Section 3.3. We recall that

$$\begin{aligned} \mathcal{E}_*^{\text{at}}(\mathbf{y}) &= \frac{1}{4m\varepsilon} \int_{\Omega^{\text{at}}} \int_{\Omega^{\text{at}}} \rho_{\mathbf{y}}(x) \left( e^{-\frac{m}{\varepsilon}|x-z|} + e^{-\frac{m}{\varepsilon}(2a_R(\mathbf{y})-x-z)} \right. \\ &\quad \left. + e^{-\frac{m}{\varepsilon}(x+z-2a_L(\mathbf{y}))} \right) \rho_{\mathbf{y}}(z) \, dz \, dx + \mathcal{O}(\tau). \end{aligned}$$

The next result addresses the differentiability of  $\gamma_L$  and  $\gamma_R$ . We show that the derivatives satisfy certain bounds.

**Lemma 5.2.** *Let  $\mathbf{y} \in \Omega_a^{2K+1}$  satisfy  $y_{i+1} - y_i > \varepsilon s_0$  for all  $i \in \{-K+1, \dots, K\}$ ,  $a_R - y_K > \varepsilon s_0/2$ , and  $y_{-K} - a_L > \varepsilon s_0/2$ . Then,  $\gamma_L(\mathbf{y})$  is twice continuously differentiable with respect to  $\mathbf{y}$  and  $a$  and there exists  $C(m \min \mathbf{y}')$  (independent of  $\varepsilon$ ) such that*

$$\begin{aligned} |D\gamma_L(\mathbf{y}, a) \cdot (\mathbf{u}, h)| &\leq C(m \min \mathbf{y}') \left( \left( \frac{u_{-K} - h_L}{\varepsilon} \right)^2 + \sum_{k=-K+1}^K (u'_k)^2 \right)^{1/2}, \\ |D^2\gamma_L(\mathbf{y}, a) \cdot [(\mathbf{u}, h), (\mathbf{u}, h)]| &\leq C(m \min \mathbf{y}') \left( \left( \frac{u_{-K} - h_L}{\varepsilon} \right)^2 + \sum_{k=-K+1}^K (u'_k)^2 \right) \end{aligned}$$

for all  $\mathbf{u} \in \mathcal{U}$  and  $h \in \mathbb{R}^2$ . Analogous bounds hold for  $\gamma_R(\mathbf{y}, a)$ .

*Proof.* The proof is based on the observation that

$$\gamma_L(\mathbf{y}, a) = \frac{1}{m} \sum_{j=-K}^K \int_{\Omega_a} e^{-\frac{m}{\varepsilon}(x-a_L)} \delta_{\varepsilon}(x - y_j) \, dx = \frac{\mu}{m} e^{-\frac{m}{\varepsilon}(y_{-K}-a_L)} \sum_{j=-K}^K e^{-\frac{m}{\varepsilon}(y_j-y_{-K})}.$$

The rest of the proof is a straightforward computation; see [10, Lemma 5.3] for the details.  $\square$

The  $\tau$ -dependent terms in  $\mathcal{E}_*^{\text{at}}(\mathbf{y}) = \mathcal{E}_{a(\mathbf{y}), g^*(\mathbf{y})}(\mathbf{y}_{\text{at}})$  from (3.24) only contain  $\gamma_L(\mathbf{y})$  and  $\gamma_R(\mathbf{y})$ , whose derivatives are bounded by Lemma 5.2. The derivatives of these  $\tau$ -dependent terms are therefore still of order  $\mathcal{O}(\tau)$  and will be neglected in the proof of the following result.

**Lemma 5.3.** *Let  $\mathbf{y} \in \mathcal{Y}$  satisfy  $\min \mathbf{y}' > s_0$ . Then,*

$$D^2\mathcal{E}^{\text{ac}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}] \geq \left( \frac{m\mu^2}{2} e^{-m \max \mathbf{y}'} - \mathcal{O}(\tau) \right) \|\mathbf{u}'\|_{\ell^2_{\varepsilon}}^2 \quad \forall \mathbf{u} \in \mathcal{U}.$$

*Proof.* We treat continuum and atomistic contributions independently and start with the former. Lemma 4.5 states that

$$D^2 \mathcal{E}_j^{\text{cb}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}] \geq \frac{m^2 \mu^2}{2} e^{-m \max \mathbf{y}' \varepsilon |u'_j|^2}$$

for all  $j = -N, \dots, N$ . Hence, the definition (5.2) of  $\mathcal{E}_*^{\text{cb}}$  directly implies that

$$D^2 \mathcal{E}_*^{\text{cb}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}] \geq e^{-m \max \mathbf{y}' \frac{m^2 \mu^2}{2} \varepsilon \left( \sum_{j=-N}^{-K-1} |u'_j|^2 + \frac{1}{2} (|u'_{-K}|^2 + |u'_{K+1}|^2) + \sum_{j=K+2}^N |u'_j|^2 \right)}.$$

Let us now turn to the atomistic part  $\mathcal{E}_*^{\text{at}}(\mathbf{y})$ . From Section 3.3 we know that for the given choice of boundary conditions and  $a(\mathbf{y})$  we can write the energy of the atomistic part as

$$\begin{aligned} \mathcal{E}_*^{\text{at}}(\mathbf{y}) &= \frac{\varepsilon}{4m} \sum_{i,j=-K}^K \int_{\Omega^{\text{at}}} \int_{\Omega^{\text{at}}} \delta_\varepsilon(x - y_i) \left( e^{-\frac{m}{\varepsilon} |x-z|} + e^{-\frac{m}{\varepsilon} (x+z-y_{K-1}-y_{-K})} \right. \\ &\quad \left. + e^{-\frac{m}{\varepsilon} (y_{K+1}+y_{K-x-z})} \right) \delta_\varepsilon(z - y_j) dz dx \\ &= \frac{\varepsilon \mu^2}{4m} \sum_{i,j=-K}^K \left( e^{-\frac{m}{\varepsilon} |y_i - y_j|} + e^{-\frac{m}{\varepsilon} (y_i + y_j - y_{-K} - y_{K-1})} \right. \\ &\quad \left. + e^{-\frac{m}{\varepsilon} (y_{K+1} + y_{K-x-z} - y_i - y_j)} \right) + \mathcal{E}_{\text{self}} + \mathcal{O}(\tau), \end{aligned} \quad (5.11)$$

where the constant  $\mathcal{E}_{\text{self}}$  accounts for the self-energies of the atoms  $\{-K, \dots, K\}$ . Differentiating twice and keeping only contributions from nearest neighbour interactions leads directly to

$$D^2 \mathcal{E}_*^{\text{at}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}] \geq e^{-m \max \mathbf{y}' \frac{m \mu^2}{2} \varepsilon \left( \frac{1}{2} |u'_{-K}|^2 + \sum_{i=-K+1}^K |u'_i|^2 + \frac{1}{2} |u'_{K+1}|^2 \right) - \mathcal{O}(\tau)}.$$

Adding the lower bounds for  $D^2 \mathcal{E}_*^{\text{cb}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}]$  and  $D^2 \mathcal{E}_*^{\text{at}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}]$  we arrive at

$$D^2 \mathcal{E}^{\text{ac}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}] = (D^2 \mathcal{E}_*^{\text{cb}}(\mathbf{y}) + D^2 \mathcal{E}_*^{\text{at}}(\mathbf{y})) \cdot [\mathbf{u}, \mathbf{u}] \geq \left( e^{-m \max \mathbf{y}' \frac{m \mu^2}{2} \varepsilon} - \mathcal{O}(\tau) \right) \|\mathbf{u}'\|_{\ell_\varepsilon^2}^2,$$

for all  $\mathbf{u} \in \mathcal{U}$ , as desired.  $\square$

**5.4. Error Estimates.** Combining the consistency and stability results we obtain the following error estimates. We note that the upper bound on the error depends on the smoothness of  $\bar{\mathbf{y}}$  in the continuum region, but that the dependence on  $\bar{\mathbf{y}}$  in the atomistic region decays exponentially with distance to the a/c interface. In realistic higher-dimensional models such an estimate would make it possible to allow defects in the atomistic region without affecting the error estimate.

**Theorem 5.4.** *Suppose that  $\bar{\mathbf{y}} \in \arg \min E_f$  and  $\bar{\mathbf{y}}_{\text{ac}} \in \arg \min E_f^{\text{ac}}$  satisfy*

$$\min \bar{\mathbf{y}}', \min \bar{\mathbf{y}}'_{\text{ac}} \geq s_0 \geq s_0, \quad \text{and} \quad \max \bar{\mathbf{y}}', \max \bar{\mathbf{y}}'_{\text{ac}} \leq S_0 < +\infty. \quad (5.12)$$

*There exist constants  $c$  and  $C = C(s_0, S_0)$  such that, if  $\Delta a \geq c \log(S_0)$ , then*

$$\|\bar{\mathbf{y}}' - \bar{\mathbf{y}}'_{\text{ac}}\|_{\ell_\varepsilon^2} \leq C \left( \varepsilon \|\bar{\mathbf{y}}''\|_{\ell_{w,s_0}^2} + \tau \right). \quad (5.13)$$

*Proof.* From Lemma 5.3 it is clear that there exists a constant  $c$  such that, for  $\Delta a \geq c$ , we have

$$D^2 \mathcal{E}^{\text{ac}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}] \geq \frac{m \mu^2}{4} e^{-m S_0} \|\mathbf{u}'\|_{\ell_\varepsilon^2}^2 \quad \forall \mathbf{u} \in \mathcal{U}, \quad \forall \mathbf{y} \in \mathcal{Y}, \mathbf{y}' \leq S_0.$$

In particular, this holds for all  $\mathbf{y} \in \text{conv}\{\bar{\mathbf{y}}, \bar{\mathbf{y}}_{\text{ac}}\}$ . Let  $c_0 = \frac{m \mu^2}{4} e^{-m S_0}$ .




FIGURE 6.1. Illustration of the problem in the interval  $Q_J^R = (y_{K-J}, 2a_R(\mathbf{y}) - y_{K-J})$  used to compute  $g_R(\mathbf{y})$ .

Let  $\mathbf{u} = \bar{\mathbf{y}} - \bar{\mathbf{y}}_{\text{ac}}$ ; then we can choose  $\mathbf{y} \in \text{conv}\{\bar{\mathbf{y}}, \bar{\mathbf{y}}_{\text{ac}}\}$  such that

$$c_0 \|\mathbf{u}'\|_{\ell_x^2}^2 \leq D^2 \mathcal{E}^{\text{ac}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}] = (D\mathcal{E}^{\text{ac}}(\bar{\mathbf{y}}) - D\mathcal{E}^{\text{ac}}(\bar{\mathbf{y}}_{\text{ac}}))[\mathbf{u}].$$

Employing the consistency estimate of Theorem 5.1 we obtain the stated result.  $\square$

**Remark 6.** With some additional work it is possible to avoid assuming the existence of  $\bar{\mathbf{y}}_{\text{ac}}$ , but deduce it from an inverse function theorem type argument [14, 10].  $\square$

## 6. BOUNDARY CONDITIONS FROM CELL PROBLEMS

The boundary conditions  $g^*(\mathbf{y})$  we imposed on the atomistic subproblem in Section 5.1 gave rise to a method without ghost forces, and whose analysis was relatively straightforward. The reasons for this is the clean weak formulation (5.4) of  $D\mathcal{E}^{\text{ac}}$  and the convenient stability properties established in Lemma 5.3. We now investigate how this situation changes if computationally cheaper boundary conditions are chosen. The following construction may also provide a starting point for generalisations to higher dimensions.

For example, a canonical choice, which requires no additional computational effort, is

$$g_L(\mathbf{y}) = \psi^{(-K)}(a_L) \quad \text{and} \quad g_R(\mathbf{y}) = \psi^{(K+1)}(a_R), \quad (6.1)$$

where we still assume  $a_L(\mathbf{y}) = \frac{1}{2}(y_{-K-1} + y_{-K})$  and  $a_R(\mathbf{y}) = \frac{1}{2}(y_K + y_{K+1})$ . In this case, we have the following result, which suggests that the additional error committed can be controlled.

**Lemma 6.1.** *Let  $\min \mathbf{y} \geq s_0 \geq s_0$  and let  $g_{L/R}$  be given by (6.1); then,*

$$|g(\mathbf{y}) - g^*(\mathbf{y})| \leq C(\varepsilon^{1/2} \|\mathbf{y}''\|_{\ell_{w,s_0}^2} + \tau). \quad (6.2)$$

*Proof.* Without loss of generality we focus on  $g_R$  only. Upon first estimating

$$|g_R(\mathbf{y}) - g_R^*(\mathbf{y})| \leq |\psi^{(K+1)}(a_R) - \phi(a_R)| + |\phi(a_R) - g_R^*(\mathbf{y})|,$$

and then employing (5.10) and Lemma 4.2, we obtain

$$|g_R(\mathbf{y}) - g_R^*(\mathbf{y})| \leq C\varepsilon \sum_{n=1}^{\infty} \|\mathbf{y}''\|_{\ell^1([K-n+1, K+n])} n e^{-mn s_0} + \mathcal{O}(\tau).$$

Using the same argument as in (5.8) to (5.9), we obtain the upper bound (6.2).  $\square$

Motivated by Lemma 6.1 we define a second a/c energy  $\mathcal{E}^{\text{ac}}(\mathbf{y})$  by

$$\mathcal{E}^{\text{ac}}(\mathbf{y}) = \mathcal{E}_*^{\text{cb}}(\mathbf{y}) + \mathcal{E}^{\text{at}}(\mathbf{y}), \quad (6.3)$$

where  $\mathcal{E}_*^{\text{cb}}(\mathbf{y})$  is the same as in the method discussed in Section 5.1 (see (5.2)) and

$$\begin{aligned} \mathcal{E}^{\text{at}}(\mathbf{y}) &= \mathcal{E}_{a(\mathbf{y}), g(\mathbf{y})}(\mathbf{y}_{\text{at}}) \\ &= - \inf \left\{ J_{a(\mathbf{y})}(\varphi, \mathbf{y}_{\text{at}}) : \varphi \in H^1(\Omega^{\text{at}}), \varphi|_{\partial\Omega^{\text{at}}} = g(\mathbf{y}) \right\}. \end{aligned}$$

We denote the minimizer for given  $\mathbf{y}$  by  $\phi_{\text{at}} \in H^1(\Omega^{\text{at}})$ .

Before we embark on the analysis of this new method, we establish a useful auxiliary result.

**Lemma 6.2.** *Let  $\min \mathbf{y}' \geq s_0 \geq s_0$ . Let  $g_R(\mathbf{y})$  be defined by (6.1); then, it can be equivalently written as*

$$g_R(\mathbf{y}) = \frac{1}{2m} \sum_{k \in \mathbb{Z}} \int_{\mathbb{R}} \delta_\varepsilon(z - (k + \frac{1}{2})\varepsilon y'_{K+1}) e^{-\frac{m}{\varepsilon}|z|} dz. \quad (6.4)$$

In particular,  $g_R$  is twice Fréchet differentiable with respect to  $\mathbf{y}$ , and there exists a constant  $C = C(s_0)$  such that, for all  $\mathbf{y} \in \mathcal{Y}$  with  $\min \mathbf{y}' \geq s_0 \geq s_0$ ,

$$|D_{\mathbf{y}} g_R(\mathbf{y}) \cdot \mathbf{u}| \leq C |u'_{K+1}| \quad \text{and} \quad |D_{\mathbf{y}}^2 g_R(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}]| \leq C |u'_{K+1}|^2 \quad \forall \mathbf{u} \in \mathcal{U}.$$

Analogous results hold for  $g_L(\mathbf{y})$ .

*Proof.* Recall from (4.8) that

$$g_R(\mathbf{y}) = \psi^{(K+1)}(a_R) = \frac{1}{2m} \sum_{k \in \mathbb{Z}} \int_{\mathbb{R}} \delta_\varepsilon(z - y_{K+k}^{(K+1)}) e^{-\frac{m}{\varepsilon}|a_R - z|} dz,$$

where  $y_j^{(K+1)}$  denotes the periodic extension defined in (4.3). We use the identities

$$y_{K+k}^{(K+1)} = y_K + k\varepsilon y'_{K+1} \quad \text{and} \quad a_R = y_K + \frac{1}{2}\varepsilon y'_{K+1}$$

to obtain

$$g_R(\mathbf{y}) = \frac{1}{2m} \sum_{k \in \mathbb{Z}} \int_{\mathbb{R}} \delta_\varepsilon(z - y_K - k\varepsilon y'_{K+1}) e^{-\frac{m}{\varepsilon}|y_K + \frac{1}{2}\varepsilon y'_{K+1} - z|} dz.$$

Shifting the integration by  $(y_K + \frac{1}{2}\varepsilon y'_{K+1})$ , we obtain (6.4).

The bound on the first and second derivatives follows as in Lemma 5.2; the key observation being that  $\delta_\varepsilon = \mathcal{O}(\varepsilon^{-1})$  is balanced against the  $\varepsilon$  preceding  $y'_{K+1}$  in its argument.  $\square$

**6.1. Consistency.** A crucial difference between the a/c energy (6.3) and the energy from Section 5.1 is that now the derivative of the atomistic energy with respect to the boundary conditions does not vanish.

Since the continuum contribution to  $D\mathcal{E}^{\text{ac}}(\mathbf{y}) \cdot \mathbf{u}$  is the same as in Section 5.1 we only need to analyze  $D\mathcal{E}^{\text{at}}(\mathbf{y})$ . Using the chain rule we obtain

$$\begin{aligned} D\mathcal{E}^{\text{at}}(\mathbf{y}) \cdot \mathbf{u} &= D_{\mathbf{y}_{\text{at}}} \mathcal{E}_{a(\mathbf{y}), g(\mathbf{y})}(\mathbf{y}_{\text{at}}) \cdot \mathbf{u}_{\text{at}} + D_a \mathcal{E}_{a(\mathbf{y}), g(\mathbf{y})}(\mathbf{y}_{\text{at}}) \cdot a(\mathbf{u}) \\ &\quad + D_g \mathcal{E}_{a(\mathbf{y}), g(\mathbf{y})}(\mathbf{y}_{\text{at}}) \cdot (D_{\mathbf{y}} g(\mathbf{y}) \cdot \mathbf{u}). \end{aligned}$$

The same reasoning as in Section 5.1 gives for the first two terms on the right-hand side

$$D_{\mathbf{y}_{\text{at}}} \mathcal{E}_{a(\mathbf{y}), g(\mathbf{y})}(\mathbf{y}_{\text{at}}) \cdot \mathbf{u}_{\text{at}} + D_a \mathcal{E}_{a(\mathbf{y}), g(\mathbf{y})}(\mathbf{y}_{\text{at}}) \cdot a(\mathbf{u}) = \int_{\Omega^{\text{at}}} \sigma_{\mathbf{y}}^{\text{at}}(x) \nabla u(x) dx, \quad (6.5)$$

where  $\sigma_{\mathbf{y}}^{\text{at}}(x)$  is given by (2.7) with  $\phi = \phi_{\text{at}}$ .

Next, we turn our attention to the term  $D_g \mathcal{E}_{a(\mathbf{y}), g(\mathbf{y})}(\mathbf{y}_{\text{at}}) \cdot (D_{\mathbf{y}} g(\mathbf{y}) \cdot \mathbf{u})$ . We recall from Lemma 3.4 that (for  $\Delta a \gg \varepsilon$ )

$$D_g \mathcal{E}_{a(\mathbf{y}), g(\mathbf{y})}(\mathbf{y}_{\text{at}}) = -m\varepsilon [g_L(\mathbf{y}) - g_L^*(\mathbf{y}), \quad g_R(\mathbf{y}) - g_R^*(\mathbf{y})] + \mathcal{O}(\varepsilon\tau).$$

Combining this result with Lemma 6.2 and Lemma 6.1, we obtain

$$\begin{aligned} |D_g \mathcal{E}_{a(\mathbf{y}), g(\mathbf{y})}(\mathbf{y}_{\text{at}}) \cdot (D_{\mathbf{y}} g(\mathbf{y}) \cdot \mathbf{u})| &\leq C\varepsilon \left( |g_L - g_L^*| + |g_R - g_R^*| + \tau \right) (|u'_{-K}|^2 + |u'_{K+1}|^2)^{1/2} \\ &\leq C \left( \varepsilon \| \mathbf{y}'' \|_{\ell_{u, s_0}^2} + \tau \right) \cdot \| \mathbf{u}' \|_{\ell_\varepsilon^2}, \end{aligned} \quad (6.6)$$

where  $C = C(\min \mathbf{y}')$ , and we have estimated  $\varepsilon^{1/2}\tau \leq \tau$ . Equipped with these estimates, we obtain the following consistency result.

**Lemma 6.3.** *Let  $\mathbf{y} \in \mathcal{Y}$  with  $\min \mathbf{y}' \geq s_0 \geq s_0$ ; then, there exists a constant  $C = C(s_0)$  such that*

$$\left| D\mathcal{E}(\mathbf{y}) \cdot \mathbf{u} - D\mathcal{E}^{\text{at}}(\mathbf{y}) \cdot \mathbf{u} \right| \leq C \left( \varepsilon \|\mathbf{y}''\|_{\ell_{w,s_0}^2} + \tau \right) \|\nabla u\|_{L^2} \quad \forall \mathbf{u} \in \mathcal{U},$$

where we have used the same notation as in Theorem 5.1.

*Proof.* From (6.5) and (6.6) we obtain that

$$\left| D\mathcal{E}(\mathbf{y}) \cdot \mathbf{u} - D\mathcal{E}^{\text{at}}(\mathbf{y}) \cdot \mathbf{u} \right| \leq \left[ \left( \varepsilon \sum_{\substack{j=-N,\dots,N \\ j \notin \{-K+1,\dots,K\}}} \|\sigma_{\mathbf{y}} - \sigma_{j;\mathbf{y}}^{\text{cb}}\|_{L^\infty(Q_j)}^2 + \|\sigma_{\mathbf{y}} - \sigma_{\mathbf{y}}^{\text{at}}\|_{L^2(a_L, a_R)}^2 \right)^{1/2} + C(\varepsilon \|\mathbf{y}''\|_{\ell_{w,s_0}^2} + \tau) \right] \cdot \|\nabla u\|_{L^2}.$$

The first group in the upper bound was already estimated in the proof of Theorem 5.1, and the second group,  $\|\sigma_{\mathbf{y}} - \sigma_{\mathbf{y}}^{\text{at}}\|_{L^2(a_L, a_R)}$  can be treated analogously to the term  $\|\sigma_{\mathbf{y}} - \sigma_{\mathbf{y},*}^{\text{at}}\|_{L^2(a_L, a_R)}$  in the proof of Theorem 5.1.  $\square$

**6.2. Stability.** We wish to compute a convenient lower bound on  $D^2\mathcal{E}^{\text{ac}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}]$  for some given  $\mathbf{y} \in \mathcal{Y}$  with  $s_0 \leq s_0 \leq \mathbf{y}' \leq S_0$ . Since the continuum part of the energy is the same as in the first method, we only address the stability of the atomistic subproblem with the given choice of boundary data. We write the second derivative of the energy  $\mathcal{E}^{\text{at}}$  in the form

$$D^2\mathcal{E}^{\text{at}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}] = D^2\mathcal{E}_*^{\text{at}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}] + (D^2\mathcal{E}^{\text{at}}(\mathbf{y}) - D^2\mathcal{E}_*^{\text{at}}(\mathbf{y})) \cdot [\mathbf{u}, \mathbf{u}]$$

and use the coercivity of  $D^2\mathcal{E}_*^{\text{at}}(\mathbf{y})$ : we know from Lemma 5.3 that

$$D^2\mathcal{E}_*^{\text{at}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}] \geq e^{-m \max \mathbf{y}'} \frac{m\mu^2}{2} \varepsilon \left( \frac{1}{2} |u'_{-K}|^2 + \sum_{i=-K+1}^K |u'_i|^2 + \frac{1}{2} |u'_{K+1}|^2 \right) - \mathcal{O}(\tau)$$

for all  $\mathbf{u} \in \mathcal{U}$ ; hence we are left to analyze the difference  $D^2\mathcal{E}^{\text{at}}(\mathbf{y}) - D^2\mathcal{E}_*^{\text{at}}(\mathbf{y})$ . We will *not* show that this difference is small, but will only be able to bound it below by a controllable quantity. This is reminiscent of similar observations made in [14].

**Lemma 6.4.** *Let  $\mathbf{y} \in \mathcal{Y}$  such that  $\min \mathbf{y}' \geq s_0 \geq s_0$ ; then there exists a constant  $C = C(s_0)$  such that*

$$(D^2\mathcal{E}^{\text{at}}(\mathbf{y}) - D^2\mathcal{E}_*^{\text{at}}(\mathbf{y})) \cdot [\mathbf{u}, \mathbf{u}] \geq -C(\varepsilon^{1/2} \|\mathbf{y}''\|_{\ell_{w,s_0}^2} + \tau).$$

*Proof.* The difference between the energies  $\mathcal{E}^{\text{at}}(\mathbf{y})$  and  $\mathcal{E}_*^{\text{at}}(\mathbf{y})$  only consists of effects from the boundary conditions. We have, by (3.18),

$$\mathcal{E}^{\text{at}}(\mathbf{y}) - \mathcal{E}_*^{\text{at}}(\mathbf{y}) = -I_{a(\mathbf{y})}(\xi_{a(\mathbf{y}),g(\mathbf{y})}, \mathbf{y}) + I_{a(\mathbf{y})}(\xi_{a(\mathbf{y}),g^*(\mathbf{y})}, \mathbf{y}) = \frac{m\varepsilon}{2} |g(\mathbf{y}) - g^*(\mathbf{y})|^2 + \mathcal{O}(\varepsilon\tau).$$

As in Section 5.3, one can verify that the  $\mathcal{O}(\tau)$  term remains of that same order in the first and second derivatives. This implies that

$$\begin{aligned} (D^2\mathcal{E}^{\text{at}}(\mathbf{y}) - D^2\mathcal{E}_*^{\text{at}}(\mathbf{y})) \cdot [\mathbf{u}, \mathbf{u}] &= m\varepsilon (g(\mathbf{y}) - g^*(\mathbf{y}))^\top [(D^2g(\mathbf{y}) - D^2g^*(\mathbf{y})) \cdot [\mathbf{u}, \mathbf{u}]] \\ &\quad + 2m\varepsilon |(Dg(\mathbf{y}) - Dg^*(\mathbf{y})) \cdot \mathbf{u}|^2 + \mathcal{O}(\tau \|\mathbf{u}'\|_{\ell_{\frac{\varepsilon}{2}}^2}^2) \\ &\geq m\varepsilon (g(\mathbf{y}) - g^*(\mathbf{y}))^\top [(D^2g(\mathbf{y}) - D^2g^*(\mathbf{y})) \cdot [\mathbf{u}, \mathbf{u}]] + \mathcal{O}(\tau \|\mathbf{u}'\|_{\ell_{\frac{\varepsilon}{2}}^2}^2). \end{aligned} \quad (6.7)$$

We now employ Lemma 6.2 to bound  $D^2g(\mathbf{y})$ , Lemma 5.2 to bound  $D^2g^*$  (up to another  $\mathcal{O}(\tau)$  error), and Lemma 6.1 to bound  $g - g^*$ , which yields

$$(D^2\mathcal{E}^{\text{at}}(\mathbf{y}) - D^2\mathcal{E}_*^{\text{at}}(\mathbf{y})) \cdot [\mathbf{u}, \mathbf{u}] \geq -C(\varepsilon^{1/2}\|\mathbf{y}''\|_{\ell_{w,s_0}^2} + \tau)\|\mathbf{u}'\|_{\ell_\varepsilon^2},$$

where  $C = C(\min \mathbf{y}')$ .  $\square$

From Lemma 6.4 and Lemma 5.3 we immediately obtain the following corollary, which states that, if  $S_0$  is moderate,  $\mathbf{y}$  “smooth” in a neighbourhood of the interfaces  $a_{L/R}$  and in the continuum region, and if the atomistic region is sufficiently large, then  $D^2\mathcal{E}^{\text{ac}}(\mathbf{y})$  is stable.

**Corollary 6.5.** *Let  $\mathbf{y} \in \mathcal{Y}$  satisfy  $\min \mathbf{y}' \geq s_0 \geq s_0$  and  $\max \mathbf{y}' \leq S_0$ ; then there exists a constant  $C = C(s_0)$  such that*

$$D^2\mathcal{E}^{\text{ac}}(\mathbf{y}) \cdot [\mathbf{u}, \mathbf{u}] \geq \left( \frac{m\mu^2}{2} e^{-mS_0} - C(\varepsilon^{1/2}\|\mathbf{y}''\|_{\ell_{w,s_0}^2} + \tau) \right) \|\mathbf{u}'\|_{\ell_\varepsilon^2}^2 \quad \forall \mathbf{u} \in \mathcal{U}.$$

**Remark 7.** The scaling  $\varepsilon^{1/2}$  is due to the fact that the additional error committed is concentrated in a region of length  $\varepsilon$ .  $\square$

**6.3. Error Estimates.** Repeating the proof of Theorem 5.4 verbatim, but replacing the consistency and stability estimates from Section 5 with those derived in Lemma 6.3 and Corollary 6.5, we obtain the following error estimates for the modified a/c method.

**Theorem 7.1.** *Recall the notation introduced in Theorem 5.1. Suppose that  $\bar{\mathbf{y}} \in \arg \min E_{\mathbf{f}}$  and  $\bar{\mathbf{y}}_{\text{ac}} \in \arg \min E_{\mathbf{f}}^{\text{ac}}$ , where  $\mathcal{E}^{\text{ac}}$  is defined in (6.3), satisfy*

$$\min \bar{\mathbf{y}}', \min \bar{\mathbf{y}}'_{\text{ac}} \geq s_0 \geq s_0, \quad \text{and} \quad \max \bar{\mathbf{y}}', \max \bar{\mathbf{y}}'_{\text{ac}} \leq S_0 < +\infty. \quad (6.8)$$

*There exist constants  $c$  and  $C = C(s_0, S_0)$  such that, if  $\tau + \varepsilon^{1/2}\|\mathbf{y}''\|_{\ell_{w,s_0}^2} \leq ce^{-mS_0}$  (in particular,  $K$  must be sufficiently large), then*

$$\|\bar{\mathbf{y}}' - \bar{\mathbf{y}}'_{\text{ac}}\|_{\ell_\varepsilon^2} \leq C(\varepsilon\|\bar{\mathbf{y}}''\|_{\ell_{w,s_0}^2} + \tau). \quad (6.9)$$

## 7. CONCLUSIONS AND OUTLOOK

We have presented a rigorous error analysis of an atomistic-to-continuum coupling method for a field-based interaction potential in one space dimension. The starting point for the design of coupling methods was a weak formulation of the forces arising from the atomistic model. This provided a natural connection point to the corresponding continuum model. We believe that the present work in a comparably simple setting addresses several important questions relevant for a/c coupling in the presence of fields, most prominently the dependence of the a/c methods on choice of the boundary and the boundary data for the interaction fields.

For the two a/c methods we discussed we chose  $\mathbf{y}$ -dependent boundaries  $a(\mathbf{y})$  of the atomistic subdomain  $\Omega^{\text{at}}$ . In other words we fixed the position of the boundary in the Lagrangian domain. This leads to convenient weak formulations of  $D\mathcal{E}^{\text{ac}}(\mathbf{y})$ . An obvious alternative (particularly relevant for higher dimensions) is the choice of  $\mathbf{y}$ -independent  $a$ . We have not investigated this further, however, see [10] for some preliminary remarks.

We also remark that we heavily utilized the one-dimensional setting in several places in the analysis. A generalisation both of the numerical methods and their analysis is therefore non-trivial. In particular, we can see no straightforward generalisation of the reflection boundary conditions  $g^*(\mathbf{y})$ . A possible way forward would be to give an alternative analysis of the second method described in Section 6 that does not utilize these reflection techniques.

## REFERENCES

- [1] X. Blanc, C. Le Bris, and F. Legoll. Analysis of a prototypical multiscale method coupling atomistic and continuum mechanics. *M2AN Math. Model. Numer. Anal.*, 39(4):797–826, 2005.
- [2] M. Dobson and M. Luskin. An analysis of the effect of ghost force oscillation on quasicontinuum error. *M2AN Math. Model. Numer. Anal.*, 43(3):591–604, 2009.
- [3] M. Dobson, M. Luskin, and C. Ortner. Accuracy of quasicontinuum approximations near instabilities. *J. Mech. Phys. Solids*, 58(10):1741–1757, 2010.
- [4] W. E, J. Lu, and J. Z. Yang. Uniform accuracy of the quasicontinuum method. *Phys. Rev. B*, 74(21):214115, Dec 2006.
- [5] L. C. Evans. *Partial Differential Equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1998.
- [6] C. J. García-Cervera, J. Lu, and W. E. A sub-linear scaling algorithm for computing the electronic structure of materials. *Commun. Math. Sci.*, 5(4):999–1026, 2007.
- [7] V. Gavini. Configurational forces in field formulation of quasicontinuum. unpublished manuscript.
- [8] V. Gavini, K. Bhattacharya, and M. Ortiz. Quasi-continuum orbital-free density-functional theory: a route to multi-million atom non-periodic DFT calculation. *J. Mech. Phys. Solids*, 55(4):697–718, 2007.
- [9] M. Iyer and V. Gavini. A field theoretic approach to the quasi-continuum method. *J. Mech. Phys. Solids*, 59:1506–1535, 2011.
- [10] B. Langwallner, C. Ortner, and E. Süli. Quasicontinuum coupling for a field-based interaction potential. OxMOS Report 34/2011, <http://www2.maths.ox.ac.uk/oxmos/reports/>.
- [11] R. Miller, E. B. Tadmor, R. Phillips, and M. Ortiz. Quasicontinuum simulation of fracture at the atomic scale. *Modelling and Simulation in Materials Science and Engineering*, 6:607, 1998.
- [12] P. Ming and J. Z. Yang. Analysis of a one-dimensional nonlocal quasi-continuum method. *Multiscale Model. Simul.*, 7(4):1838–1875, 2009.
- [13] C. Ortner. The role of the patch test in 2d atomistic-to-continuum coupling methods. arXiv:1101.5256, to appear in *M2AN Math. Model. Numer. Anal.*
- [14] C. Ortner. A priori and a posteriori analysis of the quasinonlocal quasicontinuum method in 1D. *Math. Comp.*, 80(275):1265–1285, 2011.
- [15] A. V. Shapeev. Consistent energy-based atomistic/continuum coupling for two-body potentials in one and two dimensions. *Multiscale Model. Simul.*, 9(3):905–932, 2011.
- [16] V. B. Shenoy, R. Miller, E. B. Tadmor, R. Phillips, and M. Ortiz. Quasicontinuum models of interfacial structure and deformation. *Physical Review Letters*, 80(4):742–745, 1998.
- [17] T. Shimokawa, J. J. Mortensen, J. Schiøtz, and K. W. Jacobsen. Matching conditions in the quasicontinuum method: Removal of the error introduced at the interface between the coarse-grained and fully atomistic region. *Phys. Rev. B*, 69(21):214104, 2004.
- [18] E. B. Tadmor, M. Ortiz, and R. Phillips. Quasicontinuum analysis of defects in solids. *Philosophical Magazine A*, 73(6):1529–1563, 1996.
- [19] E. B. Tadmor, R. Phillips, and M. Ortiz. Mixed atomistic and continuum models of deformation in solids. *Langmuir*, 12(19):4529–4534, 1996.
- [20] S. P. Xiao and T. Belytschko. A bridging domain method for coupling continua with molecular dynamics. *Computer methods in applied mechanics and engineering*, 193(17-20):1645–1669, 2004.

B. LANGWALLNER, MATHEMATICAL INSTITUTE, 24-29 ST GILES', OXFORD OX1 3LB, UK  
*E-mail address:* langwallner@maths.ox.ac.uk

C. ORTNER, MATHEMATICS INSTITUTE, ZEEMAN BUILDING, UNIVERSITY OF WARWICK, COVENTRY CV4 7AL, UK  
*E-mail address:* christoph.ortner@warwick.ac.uk

E. SÜLI, MATHEMATICAL INSTITUTE, 24-29 ST GILES', OXFORD OX1 3LB, UK  
*E-mail address:* suli@maths.ox.ac.uk