Pre-asymptotic error Analysis of CIP-FEM and FEM for Helmholtz Equation with high Wave Number. Part II: *hp* version

Lingxue Zhu^{*} Haijun Wu[†]

Abstract

In this paper, which is part II in a series of two, the pre-asymptotic error analysis of the continuous interior penalty finite element method (CIP-FEM) and the FEM for the Helmholtz equation in two and three dimensions is continued. While part I contained results on the linear CIP-FEM and FEM, the present part deals with approximation spaces of order $p \ge 1$. By using a modified duality argument, pre-asymptotic error estimates are derived for both methods under the condition of $\frac{kh}{p} \le C_0(\frac{p}{k})^{\frac{1}{p+1}}$, where k is the wave number, h is the mesh size, and C_0 is a constant independent of k, h, p, and the penalty parameters. It is shown that the pollution errors of both methods in H^1 -norm are $O(k^{2p+1}h^{2p})$ if p = O(1) and are $O\left(\frac{k}{p^2}\left(\frac{kh}{\sigma p}\right)^{2p}\right)$ if the exact solution $u \in H^2(\Omega)$ which coincide with existent dispersion analyses for the FEM on Cartesian grids. Here σ is a constant independent of k, h, p, and the penalty parameters. Moreover, it is proved that the CIP-FEM is stable for any k, h, p > 0 and penalty parameters with positive imaginary parts. Besides the advantage of the absolute stability of the CIP-FEM compared to the FEM, the penalty parameters may be tuned to reduce the pollution effects.

Key words. Helmholtz equation, large wave number, pre-asymptotic error estimates, continuous interior penalty finite element methods, finite element methods

AMS subject classifications. 65N12, 65N15, 65N30, 78A40

1 Introduction

This is the second installment in a series (cf. [47]) which is devoted to pre-asymptotic stability and error estimates of hp versions of some continuous interior penalty finite element method (CIP-FEM) and the finite element method (FEM) for the following Helmholtz problem:

(1.1)
$$-\Delta u - k^2 u = f \qquad \text{in } \Omega$$

(1.2)
$$\frac{\partial u}{\partial n} + \mathbf{i}ku = g \qquad \text{on I}$$

where $\Omega \subset \mathbb{R}^d$, d = 2, 3 is a domain with smooth boundary, $\Gamma := \partial \Omega$, $\mathbf{i} = \sqrt{-1}$ denotes the imaginary unit, and *n* denotes the unit outward normal to $\partial \Omega$. The above Helmholtz problem is an approximation of the following acoustic scattering problem (with time dependence $e^{\mathbf{i}\omega t}$):

(1.3)
$$-\Delta u - k^2 u = f \qquad \text{in } \mathbb{R}^d,$$

(1.4)
$$\sqrt{r} \Big(\frac{\partial (u - u^{\text{inc}})}{\partial r} + \mathbf{i} k (u - u^{\text{inc}}) \Big) \to 0 \quad \text{as } r = |x| \to \infty,$$

where u^{inc} is the incident wave and k is known as the wave number. The Robin boundary condition (1.2) is known as the first order approximation of the radiation condition (1.4) (cf. [24]). We remark

^{*}Department of Mathematics, Nanjing University, Jiangsu, 210093, P.R. China. (zhulingxue@163.com).

[†]Department of Mathematics, Nanjing University, Jiangsu, 210093, P.R. China. (hjw@nju.edu.cn). The work of the second author was partially supported by the National Magnetic Confinement Fusion Science Program under grant 2011GB105003 and by the NSF of China grants 10971096, 11071116, 91130004.

that the Helmholtz problem (1.1)–(1.2) also arises in applications as a consequence of frequency domain treatment of attenuated scalar waves (cf. [23]).

It is well-known that the error bound of the finite element solution to the Helmholtz problem (1.1)–(1.2) usually consists of two parts, one is the same order as the error of the best approximation of u from the finite element space, another one is worse than the convergence order of the best approximation and dominates the error bound of the finite element solution for large wave number k (relative to h and p) [1, 5, 6, 21, 25, 26, 27, 31, 34, 35, 45, 47]. The second part is the so-called pollution error in the literature. We recall that, the term "asymptotic error estimate" refers to the error estimate without pollution error and the term "pre-asymptotic error estimate" refers to the estimate with non-negligible pollution effect. The latest asymptotic error analysis was given by Melenk and Sauter [39, 40]. It is shown that, the hp-FEM is pollution-free in the H^1 -norm if

(1.5)
$$\frac{kh}{p} + k \left(\frac{kh}{\sigma p}\right)^p \text{ is small enough,}$$

where h is the mesh size, p is the polynomial degree of finite element space, and σ is some positive constant independent of k, h, and p. In particular, the hp-FEM is pollution-free under either of the following two conditions:

(1.6)
$$p = O(1)$$
 fixed independent of k and $k^{1+\frac{1}{p}}h$ is small enough,

(1.7)
$$p \ge c \ln k \text{ and } \frac{k h}{p} \text{ is small enough},$$

where c is some positive constant independent of k, h, and p. Although the above results improve greatly the previous results which require $\frac{k^2h}{p}$ small enough (cf. [34, 35]), the pre-asymptotic error estimates are still worth to be studied based on but not limited to the following considerations:

- For fixed *p*, the pollution effect can be reduced substantially but cannot be avoided in principle (cf. [6, 5]).
- Implementations of high order FEMs, e.g. $p \ge \ln k$ for large k, are not easy for problems with complicated geometries or different materials.
- Given a tolerance ε , one wish to use as small as possible number of degrees of freedom, or in other words, as large as possible mesh size h, to achieve this tolerance. We denote the maximum mesh size by $h(k,\varepsilon)$. Then $h(k,\varepsilon)$ always locates in the pre-asymptotic range for large k because the pollution error dominates the error bound. For example, when p = 1, for large k, the pollution term is $O(k^3h^2)$ (cf. [34, 47]) and hence $h(k,\varepsilon) = O((\varepsilon/k^3)^{1/2})$ which is not in the asymptotic range $(0, C/k^2)$ implied by the condition (1.6).
- The pre-asymptotic error analysis of FEM is hard for higher dimensional problems. To the best of the authors' knowledge, besides the first part of this series [47] for the linear FEM, no work has been done ever since the pioneer works of Ihlenburg and Babuška [34, 35] for one dimensional problems.

The purpose of this paper and the companion paper [47] is twofold: One is to derive pre-asymptotic error estimates for the FEM for the Helmholtz equation in two and three dimensions, under the condition of $\frac{kh}{p} \leq C_0 \left(\frac{p}{k}\right)^{\frac{1}{p+1}}$ for some positive constant C_0 independent of k, h, and p. Clearly, this condition extends the asymptotic range given by (1.5). For example, when p = O(1), the above condition is satisfied if $k^{1+\frac{1}{p+1}}h$ is small enough. This would be the condition obtained for the (p + 1)-th order FEM by "standard" arguments (cf. (1.6)). Another purpose is to analyze some CIP-FEM which is absolutely stable (that is, stable for any k, h, and p) and capable of achieving much less pollution effect than the FEM for fixed p. The CIP-FEM, which was first proposed by Douglas and Dupont [22] for elliptic and parabolic problems in 1970's and then successfully applied to convection-dominated problems as a stabilization technique [13, 14, 15, 16, 17], modifies the sesquilinear form of the FEM by adding the following least squares term penalizing the jump of the gradient of the discrete solution at mesh interfaces:

(1.8)
$$J(u,v) := \sum_{e \in \mathcal{E}_h^I} \mathbf{i} \gamma \frac{h_e}{p^2} \left\langle \left\lfloor \frac{\partial u}{\partial n_e} \right\rfloor, \left\lfloor \frac{\partial v}{\partial n_e} \right\rfloor \right\rangle_e.$$

Note that the penalty parameter $i\gamma$ is chosen as a complex number in this paper and [47] instead of the real numbers as usual. The paper [47] has considered the linear case p = 1 and this paper will be devoted to the hp-version. To be precise, we obtain the following results:

(i) There exists a constant $C_0 > 0$ independent of k, h, and γ , such that if $k \ge 1, 0 \le \gamma \le 1$, and

(1.9)
$$\frac{kh}{p} \le C_0 \left(\frac{p}{k}\right)^{\frac{1}{p+1}},$$

then the following pre-asymptotic error estimates hold for both the CIP-FEM and the FEM:

$$\|u - u_h\|_{H^1(\Omega)} \le \begin{cases} C_1 \left(\frac{h}{p} + \frac{1}{p} \left(\frac{kh}{\sigma p}\right)^p\right) + C_2 \frac{k}{p^2} \left(\frac{kh}{\sigma p}\right)^{2p}, & \text{if } u \in H^2(\Omega), \\ C_1(kh)^p + C_2 k(kh)^{2p}, & \text{if } p = O(1) \text{ and } \|u\|_{H^{p+1}(\Omega)} \lesssim k^p. \end{cases}$$

Here σ is a constant independent of k, h, p, and the penalty parameters. Note that the pollution term is $O(k^{2p+1}h^{2p})$ if p = O(1) and $||u||_{H^{p+1}(\Omega)} \leq k^p$ and is $O\left(\frac{k}{p^2}\left(\frac{kh}{\sigma p}\right)^{2p}\right)$ if $u \in H^2(\Omega)$.

- (ii) The CIP-FEM attains a unique solution for any k > 0, h > 0, p > 0, and $\gamma > 0$.
- (iii) Estimates in the L^2 -norm are also obtained.

We remark that the numerical tests in [47] for the linear CIP-FEM show that the penalty parameter may be tuned to greatly reduce the pollution errors.

Error analysis and dispersion analysis are two main tools to understand numerical behaviors in short wave computations. The later one, which is usually performed on *structured* meshes, estimates the error between the wave number k of the continuous problem and some discrete wave number (denoted by ω) of the numerical scheme [1, 2, 21, 31, 34, 35, 45, 46]. In particular, it is shown for the hp-FEM (cf. [1, 35]) that

$$k - \omega = \begin{cases} O(k^{2p+1}h^{2p}) & \text{if } kh \ll 1, \\ O\left(\frac{k}{p}\left(\frac{ekh}{4p}\right)^{2p}\right) & \text{if } kh \gg 1 \text{ and } 2p+1 > \frac{ekh}{2}. \end{cases}$$

By contrast, our pre-asymptotic error analysis, which works also for *unstructured* meshes, gives error estimates between the exact solution u and the discrete solution u_h . Clearly, our pollution error bounds in H^1 -norm coincide with the phase difference $|k - \omega|$ as above.

Our analysis relies on two novel tricks and the profound stability estimates for the continuous problem given in [39, 40]. The first one is a modified duality argument (or Aubin-Nitsche trick). The traditional duality argument is a crucial step in asymptotic error analyses of FEM for scattering problems, which is usually used to estimate the L^2 -error of the finite element solution by its H^1 -error (cf. [4, 23, 35, 39, 40, 42]). Our key idea is to use some special designed elliptic projections in the duality-argument step so that we can bound the L^2 -error of the discrete solution by using the errors of the elliptic projections of the exact solution u and thus obtain the L^2 -error estimate directly by the modified duality argument. This helps us to derive pre-asymptotic error estimates for the CIP-FEM and the FEM under the condition (1.9). The second trick is used to prove the absolute stability of the CIP-FEM. One of the authors and Feng [26, 27] have developed an approach to analyze the stability of the interior penalty discontinuous Galerkin methods which makes use of the special test function $v_h := \nabla u_h \cdot (x - x_\Omega)$ (defined element-wise) and a local version of the Rellich identity (for the Laplacian) and mimics the stability analysis for the PDE solutions given in [38, 20, 32]. Here x_Ω is a point such that the domain Ω is strictly star-shaped with respect to it. But the function v_h is not in the approximation space of CIP-FEM which is the same as that of the FEM, i.e., the space of *continuous* piecewise (mapped) polynomials, and thus, it can not server as a test function for the CIP-FEM. We circumvent this difficulty by using the L^2 projection of v_h onto the approximation space of CIP-FEM instead. By using the so-called "Oswald interpolation" [15, 33, 36, 41] we may show that the difference between v_h and its L^2 projection is estimated by the jump of v_h at mesh interfaces, and hence, controlled by using the jump term (1.8) in the CIP-FEM. We remark that the technique for deriving the stability of the linear CIP-FEM developed in the first part of the series [47] does not work for higher order methods because it relies on the fact of $\Delta u_h = 0$ on each element.

The remainder of this paper is organized as follows. The CIP-FEM is introduced in Section 2. Some preliminary results, including the stability of the continuous solution, the approximation properties of the hp-finite element space, and estimates of some elliptic projections, are cited or proved in Section 3. In Section 4, by using an improved duality argument, some pre-asymptotic error estimates in H^1 - and L^2 -norms are given for both CIP-FEM and FEM under the condition that $\left(\frac{k}{p}\right)^{\frac{1}{p+1}}\frac{kh}{p}$ is small enough. In Section 5, the CIP-FEM is shown to be stable for any k > 0, h > 0, p > 0, and $\gamma > 0$. In Section 6, pre-asymptotic error estimates of the CIP-FEM are proved for k > 0, h > 0, p > 0, and $\gamma > 0$ by utilizing the error estimates for the elliptic projection, the stability results for the CIP-FEM, and the triangle inequality. The proofs of two preliminary lemmas on hp-approximation properties are given in Appendix A.

Throughout the paper, C is used to denote a generic positive constant which is independent of h, p, k, f, g, and the penalty parameters. We also use the shorthand notation $A \leq B$ and $B \geq A$ for the inequality $A \leq CB$ and $B \geq CA$. $A \simeq B$ is a shorthand notation for the statement $A \leq B$ and $B \leq A$. We assume that $k \gtrsim 1$ since we are considering high-frequency problems. For the ease of presentation, we assume that k is constant on the domain Ω . We also assume that Ω is a strictly star-shaped domain with an analytic boundary. Here "strictly star-shaped" means that there exist a point $x_{\Omega} \in \Omega$ and a positive constant c_{Ω} depending only on Ω such that

(1.10)
$$(x - x_{\Omega}) \cdot n \ge c_{\Omega}, \quad \forall x \in \partial \Omega.$$

Sure the theory of this paper can be extended to the case of polyhedral domains. The extensions to other types of boundary conditions, such as PML absorbing boundary condition (cf. [11, 10, 18]) and DtN boundary condition (cf. [39]), will be addressed in future works.

2 Formulation of CIP-FEM

To formulate our CIP-FEM, we first introduce some notation. The standard space, norm and inner product notation are adopted. Their definitions can be found in [12, 19]. In particular, $(\cdot, \cdot)_Q$ and $\langle \cdot, \cdot \rangle_{\Sigma}$ for $\Sigma \subset \partial Q$ denote the L^2 -inner product on complex-valued $L^2(Q)$ and $L^2(\Sigma)$ spaces, respectively. Denote by $(\cdot, \cdot) := (\cdot, \cdot)_\Omega$ and $\langle \cdot, \cdot \rangle := \langle \cdot, \cdot \rangle_{\partial\Omega}$.

For the triangulation of Ω we adopt the setting of [39, 40]. The triangulation \mathcal{T}_h consists of elements which are the image of the reference triangle (in two dimensions) or the reference tetrahedron (in three dimensions). We do not allow hanging nodes and assume—as is standard—that the element maps of elements sharing an edge or a face induce the same parametrization on that edge or face. Additionally, the element maps F_K from the reference element \hat{K} to $K \in \mathcal{T}_h$ satisfy the assumption 5.1 in [40]. For any triangle/tetrahedron $K \in \mathcal{T}_h$, we define $h_K := \operatorname{diam}(K)$. Similarly, for each edge/face e of $K \in \mathcal{T}_h$, define $h_e := \operatorname{diam}(e)$. Let $h = \max_{K \in \mathcal{T}_h} h_K$. We define

$$\mathcal{E}_h^I := \text{set of all interior edges/faces of } \mathcal{T}_h,$$

 $\mathcal{E}_h^B := \text{set of all boundary edges/faces of } \mathcal{T}_h \text{ on } \Gamma.$

We also define the jump [v] of v on an interior edge/face $e = \partial K \cap \partial K'$ as

$$[v]|_e := \begin{cases} v|_K - v|_{K'}, & \text{if the global label of } K \text{ is bigger}, \\ v|_{K'} - v|_K, & \text{if the global label of } K' \text{ is bigger}. \end{cases}$$

For every $e = \partial K \cap \partial K' \in \mathcal{E}_h^I$, let n_e be the unit outward normal to edge/face e of the element K if the global label of K is bigger and of the element K' if the other way around. For every $e \in \mathcal{E}_h^B$, let $n_e = n$ the unit outward normal to $\partial \Omega$.

Now we define the "energy" space V and the sesquilinear form $a_h(\cdot, \cdot)$ on $V \times V$ as follows:

(2.1)

$$V := H^{1}(\Omega) \cap \prod_{K \in \mathcal{T}_{h}} H^{2}(K),$$

$$a_{h}(u, v) := (\nabla u, \nabla v) + J(u, v) \quad \forall u, v \in V,$$

where

(2.2)
$$J(u,v) := \sum_{e \in \mathcal{E}_h^I} \mathbf{i} \gamma_e \frac{h_e}{p^2} \left\langle \left[\frac{\partial u}{\partial n_e}\right], \left[\frac{\partial v}{\partial n_e}\right] \right\rangle_e,$$

and $\gamma_e, e \in \mathcal{E}_h^I$ are nonnegative numbers to be specified later.

Remark 2.1. (a) The terms in J(u, v) are so-called penalty terms. The penalty parameter in J(u, v) is $i\gamma_e$. So it is a pure imaginary number with positive imaginary part. It turns out that if it is replaced by a complex number with positive imaginary part, the ideas of the paper still apply. Here we set their real parts to be zero partly because the terms from real parts do not help much (and do not cause any problem either) in our theoretical analysis and partly for the ease of presentation.

(b) Penalizing the jumps of normal derivatives was used early by Douglas and Dupont [22] for second order PDEs and by Babuška and Zlámal [8] for fourth order PDEs in the context of C^0 finite element methods, by Baker [9] for fourth order PDEs and by Arnold [3] for second order parabolic PDEs in the context of IPDG methods.

(c) In this paper we consider the scattering problem with time dependence $e^{i\omega t}$, that is, the signs before **i**'s in the Sommerfeld radiation condition (1.4) and its first order approximation (1.2) are positive. If we consider the scattering problem with time dependence $e^{-i\omega t}$, that is, the signs before **i**'s in (1.4) and (1.2) are negative, then the penalty parameters should be complex numbers with negative imaginary parts.

It is clear that J(u, v) = 0 if $u \in H^2(\Omega)$ and $v \in V$. Therefore, if $u \in H^2(\Omega)$ is the solution of (1.1)-(1.2), then

(2.3)
$$a_h(u,v) - k^2(u,v) + \mathbf{i}k \langle u,v \rangle = (f,v) + \langle g,v \rangle, \quad \forall v \in V.$$

Let V_h be the *hp*-CIP approximation space, that is,

$$V_h := \left\{ v_h \in H^1(\Omega) : v_h \mid_K \circ F_K \in \mathcal{P}_p(\hat{K}), \forall K \in \mathcal{T}_h \right\},\$$

where $\mathcal{P}_p(\hat{K})$ denote the set of all polynomials whose degrees do not exceed p on \hat{K} . Then our CIP-FEMs are defined as follows: Find $u_h \in V_h$ such that

(2.4)
$$a_h(u_h, v_h) - k^2(u_h, v_h) + \mathbf{i}k \langle u_h, v_h \rangle = (f, v_h) + \langle g, v_h \rangle, \quad \forall v_h \in V_h.$$

We remark that if the parameters $\gamma_e \equiv 0$, then the above CIP-FEM becomes the standard FEM.

The following (semi-)norms on the space V are useful for the subsequent analysis:

(2.5)
$$\|v\|_{1,h} := \left(\|\nabla v\|_{L^{2}(\Omega)}^{2} + \sum_{e \in \mathcal{E}_{h}^{I}} \gamma_{e} \frac{h_{e}}{p^{2}} \left\| \left[\frac{\partial v}{\partial n_{e}} \right] \right\|_{L^{2}(e)}^{2} \right)^{1/2}$$

(2.6)
$$||v||| := \left(||v||_{1,h}^2 + k ||v||_{L^2(\Gamma)}^2 \right)^{\frac{1}{2}}$$

In the next sections, we shall consider the pre-asymptotic stability and error analysis for the above CIP-FEM and the FEM. For the ease of presentation, we assume that $\gamma_e \simeq \gamma$ for some positive constant γ and that $h_K \simeq h$.

3 Preliminary lemmas

In this section, we first recall the stability estimates of the continuous problem from Melenk and Sauter [39, 40]. Then we consider the *hp*-approximation estimates of the discrete space V_h . These estimates improve slightly the estimates in [40] when $u \in H^2(\Omega)$. Finally, we analyze some elliptic projections which are crucial for our pre-asymptotic analysis.

3.1 Stability estimates of the continuous problem

Let ∇^n stand for derivatives of order n; more precisely, for a function $u: \Omega \to \mathbb{R}^d$, $\Omega \subset \mathbb{R}^d$, $|\nabla^n u(x)|^2 = \sum_{\alpha \in \mathbb{N}_0^d: |\alpha|=n} \frac{n!}{\alpha!} |D^{\alpha}u(x)|^2$. The following lemma (cf. [40, Theorem 4.10]) says that the solution u to the continuous problem (1.1)–(1.2) can be decomposed into the sum of an elliptic part and an analytic part $u = u_{\mathcal{E}} + u_{\mathcal{A}}$ where $u_{\mathcal{E}}$ is usually non-smooth but the H^2 -bound of $u_{\mathcal{E}}$ is independent of k and $u_{\mathcal{A}}$ is oscillatory but the H^j -bound of $u_{\mathcal{A}}$ is available for any integer $j \geq 0$.

Lemma 3.1. The solution u to the problem (1.1)–(1.2) can be written as $u = u_{\mathcal{E}} + u_{\mathcal{A}}$, and satisfies

- (3.1) $\|u_{\mathcal{E}}\|_{H^{2}(\Omega)} + k|u_{\mathcal{E}}|_{H^{1}(\Omega)} + k^{2}\|u_{\mathcal{E}}\|_{L^{2}(\Omega)} \leq CC_{f,g},$
- $(3.2) |u_{\mathcal{A}}|_{H^1(\Omega)} + k ||u_{\mathcal{A}}||_{L^2(\Omega)} \le CC_{f,g},$
- (3.3) $\forall p \in \mathbb{N}_0, \quad \|\nabla^{p+2}u_{\mathcal{A}}\|_{L^2(\Omega)} \le C\lambda^p k^{-1} \max\{p,k\}^{p+2} C_{f,q}.$

Here, $\lambda > 1$ independent of k and $C_{f,g} := \|f\|_{L^2(\Omega)} + \|g\|_{H^{1/2}(\Gamma)}$.

Remark 3.1. An direct consequence of the above lemma is that

 $\|u\|_{H^2(\Omega)} \lesssim kC_{f,g}.$

This estimate was proved in [20, 32, 38].

3.2 Approximation properties

In this subsection we consider to approximate the solution u to the problem (1.1)–(1.2) by finite element functions in V_h . We give two types of approximation estimates in both L^2 -norm and the norm $||| \cdot |||$ defined in (2.6). The first type of estimates can be applied to smooth solution and gives higher order convergences both in h and p. The second type of estimates assumes only H^2 regularity and gives first order convergences (even for higher order elements), but it uses the decomposition given in Lemma 3.1 and is more subtle than the first one. The proofs are a little long and will be given in the appendix.

Lemma 3.2. Let $s \ge 2$ and $\mu = \min\{p+1, s\}$. Suppose $u \in H^s(\Omega)$. Then there exists $\hat{u}_h \in V_h$ such that

(3.4)
$$\|u - \hat{u}_h\|_{L^2(\Omega)} \lesssim \frac{h^{\mu}}{p^s} \|u\|_{H^s(\Omega)},$$

(3.5)
$$|||u - \hat{u}_h||| \lesssim C_{\text{err}} \frac{h^{\mu-1}}{p^{s-1}} ||u||_{H^s(\Omega)},$$

where $C_{\text{err}} := \left(1 + \gamma + \frac{kh}{p}\right)^{\frac{1}{2}}$.

The following lemma gives approximation estimates of the solution to the problem (1.1)-(1.2).

Lemma 3.3. Let u be the solution to the problem (1.1)-(1.2). Then there exist $\hat{u}_h \in V_h$ and a constant $\sigma > 0$ independent of k, p, h, and the penalty parameters, such that

(3.6)
$$||u - \hat{u}_h||_{L^2(\Omega)} \lesssim \left(\frac{h^2}{p^2} + \frac{h}{p^2} \left(\frac{kh}{\sigma p}\right)^p\right) C_{f,g},$$

(3.7)
$$|||u - \hat{u}_h||| \lesssim C_{\operatorname{err}} \left(\frac{h}{p} + \frac{1}{p} \left(\frac{kh}{\sigma p}\right)^p\right) C_{f,g},$$

where $C_{f,g} := \|f\|_{L^2(\Omega)} + \|g\|_{H^{1/2}(\Gamma)}$ and $C_{\text{err}} := \left(1 + \gamma + \frac{kh}{p}\right)^{\frac{1}{2}}$.

Remark 3.2. The above estimates are based on the results from [39, 40] and a duality argument. Similar estimates have been given in [39, 40] as follows

$$\inf_{v_h \in V_h} \left(\left\| \nabla (u - v_h) \right\|_{L^2(\Omega)} + k \left\| u - v_h \right\|_{L^2(\Omega)} \right) \lesssim \left(1 + \frac{kh}{p} \right) \left(\frac{h}{p} + \left(\frac{kh}{\sigma p} \right)^p \right) C_{f,g}.$$

Our estimates in this lemma improve a little the above estimate in terms of p. See also Remark A.1.

3.3 Elliptic projections

In this subsection, we introduce two kind of elliptic projections and estimate the approximation errors of them. For any $u \in V$, we define its elliptic projections $u_h^+ \in V_h$ and $u_h^- \in V_h$ by the following two formulations, respectively.

(3.8)
$$a_h(u_h^+, v_h) + \mathbf{i}k\langle u_h^+, v_h \rangle = a_h(u, v_h) + \mathbf{i}k\langle u, v_h \rangle \quad \forall v_h \in V_h,$$

(3.9)
$$a_h(v_h, u_h^-) + \mathbf{i}k\langle v_h, u_h^-\rangle = a_h(v_h, u) + \mathbf{i}k\langle v_h, u\rangle \quad \forall v_h \in V_h.$$

Let us take a look at the projections from other points of view. Define

$$a_h^{\pm}(u,v) := (\nabla u, \nabla v) \pm \sum_{e \in \mathcal{E}_h^I} \mathbf{i} \gamma_e \frac{h_e}{p^2} \left\langle \left[\frac{\partial u}{\partial n_e} \right], \left[\frac{\partial v}{\partial n_e} \right] \right\rangle_e.$$

Then $a_h^+ = a_h$, $a_h^-(u, v) = \overline{a_h(v, u)}$ (cf. (2.1)), and clearly, $u_h^\pm \in V_h$ satisfies

(3.10)
$$a_h^{\pm}(u_h^{\pm}, v_h) \pm \mathbf{i}k\langle u_h^{\pm}, v_h \rangle = a_h^{\pm}(u, v_h) \pm \mathbf{i}k\langle u, v_h \rangle \quad \forall v_h \in V_h.$$

In other words, u_h^{\pm} is an CIP finite element approximation to the solution u of the following (complexvalued) Poisson problem:

$$-\triangle u = F \quad \text{in } \Omega,$$
$$\frac{\partial u}{\partial n} \pm \mathbf{i}ku = \psi \quad \text{on } \Gamma,$$

for some given function F and ψ which are determined by u.

Next, we estimate the errors of u_h^{\pm} . From (3.10) we have the following Galerkin orthogonality:

(3.11)
$$a_h^{\pm}(u - u_h^{\pm}, v_h) \pm \mathbf{i}k\langle u - u_h^{\pm}, v_h \rangle = 0 \quad \forall v_h \in V_h.$$

We state the following continuity and coercivity properties for the sesquilinear form $a_h(\cdot, \cdot)$. Since they follow easily from (2.1)-(2.5), so we omit their proofs to save space.

Lemma 3.4. For any $v, w \in V$,

$$(3.12) |a_h^{\pm}(v,w)|, |a_h^{\pm}(w,v)| \le \|v\|_{1,h} \|w\|_{1,h},$$

(3.13)
$$\operatorname{Re} a_{h}^{\pm}(v,v) \pm \operatorname{Im} a_{h}^{\pm}(v,v) = \|v\|_{1,h}^{2}$$

Lemma 3.5. Suppose u is any function in $H^2(\Omega)$. Then there hold the following estimates:

(3.14)
$$\left\| \left\| u - u_h^{\pm} \right\| \right\| \lesssim \inf_{z_h \in V_h} \left\| \left\| u - z_h \right\| \right\|,$$

(3.15)
$$\|u - u_h^{\pm}\|_{L^2(\Omega)} \lesssim C_{\text{err}} \frac{h}{p} \inf_{z_h \in V_h} \|u - z_h\|,$$

where $C_{\text{err}} := \left(1 + \gamma + \frac{kh}{p}\right)^{\frac{1}{2}}$.

8

Proof. We only prove the estimates for u_h^+ since the proof for u_h^- follows almost the same procedure. For any $z_h \in V_h$, let $\eta = u - u_h^+$, $\eta_h = u_h^+ - z_h$. From $\eta_h + \eta = u - z_h$ and (3.11), we have

(3.16)
$$a_h(\eta,\eta) + \mathbf{i}k\langle\eta,\eta\rangle = a_h(\eta,u-z_h) + \mathbf{i}k\langle\eta,u-z_h\rangle$$

Applying Lemma 3.4 and (3.16) we can obtain that

$$\begin{split} \|\eta\|_{1,h}^2 &= \operatorname{Re} a_h(\eta,\eta) + \operatorname{Im} a_h(\eta,\eta) \\ &= \operatorname{Re}(a_h(\eta,\eta) + \mathbf{i}k\langle\eta,\eta\rangle) + \operatorname{Im}(a_h(\eta,\eta) + \mathbf{i}k\langle\eta,\eta\rangle) - k\langle\eta,\eta\rangle \\ &= \operatorname{Re}(a_h(\eta,u-z_h) + \mathbf{i}k\langle\eta,u-z_h\rangle) \\ &+ \operatorname{Im}(a_h(\eta,u-z_h) + \mathbf{i}k\langle\eta,u-z_h\rangle) - k\|\eta\|_{L^2(\Gamma)} \\ &\leq C\Big(\|\eta\|_{1,h}\|u-z_h\|_{1,h} + k\|\eta\|_{L^2(\Gamma)}\|u-z_h\|_{L^2(\Gamma)}\Big) - k\|\eta\|_{L^2(\Gamma)}. \end{split}$$

Therefore,

(3.17)
$$\|\eta\|_{1,h}^2 + k\|\eta\|_{L^2(\Gamma)}^2 \lesssim \|u - z_h\|_{1,h}^2 + k\|u - z_h\|_{L^2(\Gamma)}^2$$

That is, (3.14) holds.

To show (3.15), we use the Nitsche's duality argument (cf. [12, 19]). Consider the following auxiliary problem:

$$(3.18) \qquad \qquad -\triangle w = \eta \quad \text{ in } \Omega$$

(3.19)
$$\frac{\partial w}{\partial n} - \mathbf{i}kw = 0 \quad \text{on } \Gamma$$

It can be shown that w satisfies

(3.20)
$$||w||_{H^2(\Omega)} \lesssim ||\eta||_{L^2(\Omega)}.$$

As a matter of fact, the PDE theory shows that $|w|_{H^2(\Omega)} \leq C(\Omega) \left(\|\Delta w\|_{L^2(\Omega)} + \|w\|_{H^1(\Omega)} \right)$ (cf. [37]), and testing (3.18) by the conjugate of w and taking the real part and imaginary part imply that $\|w\|_{H^1(\Omega)} \leq \|\eta\|_{L^2(\Omega)}$.

Let $\hat{w}_h \in V_h$ be defined in Lemma 3.2 (with *u* replaced by *w*). From (3.11),

$$a_h(\eta, \hat{w}_h) + \mathbf{i}k\langle \eta, \hat{w}_h \rangle = 0$$

Testing the conjugated (3.18) by η and using the above orthogonality we get

$$\begin{split} \|\eta\|_{L^{2}(\Omega)}^{2} &= -(u - u_{h}^{+}, \Delta w) = a_{h}(u - u_{h}^{+}, w) + \mathbf{i}k\langle u - u_{h}^{+}, w\rangle \\ &= a_{h}(\eta, w - \hat{w}_{h}) + \mathbf{i}k\langle \eta, w - \hat{w}_{h}\rangle \\ &\lesssim \|\eta\|_{1,h} \|w - \hat{w}_{h}\|_{1,h} + k\|\eta\|_{L^{2}(\Gamma)} \|w - \hat{w}_{h}\|_{L^{2}(\Gamma)} \\ &\lesssim \|\|\eta\|\| \|w - \hat{w}_{h}\|\| \\ &\lesssim \|\|\eta\|\| C_{\mathrm{err}} \frac{h}{p} \|w\|_{H^{2}(\Omega)}, \end{split}$$

which together with (3.14) and (3.20) gives (3.15). The proof is completed.

By combining Lemma 3.5 and Lemma 3.3 we have the following lemma which gives the error estimates between the solution to the problem (1.1)-(1.2) and its elliptic projections.

Lemma 3.6. Let u be the solution to the problem (1.1)-(1.2). Then there hold the following estimates:

(3.21)
$$|||u - u_h^{\pm}||| \lesssim C_{\mathrm{err}} \Big(\frac{h}{p} + \frac{1}{p} \Big(\frac{kh}{\sigma p}\Big)^p\Big) C_{f,g},$$

(3.22)
$$||u - u_h^{\pm}||_{L^2(\Omega)} \lesssim C_{\text{err}}^2 \Big(\frac{h^2}{p^2} + \frac{h}{p^2} \Big(\frac{kh}{\sigma p}\Big)^p\Big) C_{f,g}$$

where $C_{\text{err}} := \left(1 + \gamma + \frac{kh}{p}\right)^{\frac{1}{2}}$.

4 Pre-asymptotic error estimates by duality argument

One crucial step in asymptotic error analyses of FEM for scattering problems is performing the duality argument (or Aubin-Nitsche trick) (cf. [4, 23, 35, 39, 40, 42]). This argument is usually used to estimate the L^2 -error of the finite element solution by its H^1 -error. In this section we use a modified duality argument to derive pre-asymptotic error estimates under the condition that $\left(\frac{k}{p}\right)^{\frac{1}{p+1}}\frac{kh}{p}$ is sufficiently small. The key idea is to used the elliptic projections from the previous section in the duality-argument step so that we can bound the L^2 -error of the discrete solution by using the errors of the elliptic projections of the exact solution u and thus obtain the L^2 -error estimate directly by the modified duality argument.

Theorem 4.1. Let u and u_h denote the solutions of (1.1)-(1.2) and (2.4), respectively. Suppose $0 \le \gamma \le 1$. Then there exist constants C_0 and $\sigma > 0$ independent of k, h, p, and the penalty parameters, such that if

(4.1)
$$\frac{kh}{p} \le C_0 \left(\frac{p}{k}\right)^{\frac{1}{p+1}}$$

then the following error estimates hold:

(4.2)
$$|||u - u_h||| \lesssim \left(1 + \frac{k}{p} \left(\frac{kh}{\sigma p}\right)^p\right) \inf_{z_h \in V_h} |||u - z_h|||$$

(4.3)
$$\|u - u_h\|_{L^2(\Omega)} \lesssim \left(\frac{h}{p} + \frac{1}{p} \left(\frac{kh}{\sigma p}\right)^p\right) \inf_{z_h \in V_h} \|u - z_h\| .$$

Proof. First, we estimate the L^2 -error by introducing the dual problem and using the elliptic projections of the solution to the original continuous problem and of the solution to the dual problem. Let $e_h := u - u_h$. Consider the following dual problem:

(4.4)
$$-\Delta w - k^2 w = e_h \quad \text{in } \Omega,$$

(4.5)
$$\frac{\partial w}{\partial n} - \mathbf{i}kw = 0 \quad \text{on } \Gamma$$

Let u_h^+ be the elliptic projection of u defined by (3.8) and let $w_h^- \in V_h$ be the elliptic projection defined as (3.9). From (2.3) and (2.4) we have the following orthogonality,

(4.6)
$$a_h(e_h, v_h) - k^2(e_h, v_h) + \mathbf{i}k\langle e_h, v_h \rangle = 0, \quad \forall v_h \in V_h.$$

Testing the conjugated (4.4) by e_h and using (4.6) with $v_h = w_h^-$, (3.8)–(3.9) we get

(4.7)
$$\|e_{h}\|_{L^{2}(\Omega)}^{2} = (\nabla e_{h}, \nabla w) - k^{2}(e_{h}, w) + \mathbf{i}k\langle e_{h}, w \rangle$$
$$= a_{h}(e_{h}, w) - k^{2}(e_{h}, w) + \mathbf{i}k\langle e_{h}, w \rangle$$
$$= a_{h}(e_{h}, w - w_{h}^{-}) + \mathbf{i}k\langle e_{h}, w - w_{h}^{-} \rangle - k^{2}(e_{h}, w - w_{h}^{-})$$
$$= a_{h}(u - u_{h}^{+}, w - w_{h}^{-}) + \mathbf{i}k\langle u - u_{h}^{+}, w - w_{h}^{-} \rangle - k^{2}(e_{h}, w - w_{h}^{-}).$$

Similar to Lemma 3.6, we may show that

(4.8)
$$\left\| \left\| w - w_h^- \right\| \right\| \lesssim C_{\operatorname{err}} \left(\frac{h}{p} + \frac{1}{p} \left(\frac{kh}{\sigma p} \right)^p \right) \|e_h\|_{L^2(\Omega)}$$

(4.9)
$$\|w - w_h^-\|_{L^2(\Omega)} \lesssim C_{\text{err}}^2 \left(\frac{h^2}{p^2} + \frac{h}{p^2} \left(\frac{kh}{\sigma p}\right)^p\right) \|e_h\|_{L^2(\Omega)}$$

From (4.7), Lemma 3.4, Lemma 3.5, and (4.8)-(4.9), we have

$$\begin{split} \|e_{h}\|_{L^{2}(\Omega)}^{2} &\leq \|u - u_{h}^{+}\|_{1,h} \|w - w_{h}^{-}\|_{1,h} + k \left\|u - u_{h}^{+}\right\|_{L^{2}(\Gamma)} \left\|w - w_{h}^{-}\right\|_{L^{2}(\Gamma)} \\ &+ k^{2} \left\|e_{h}\right\|_{L^{2}(\Omega)} \left\|w - w_{h}^{-}\right\|_{L^{2}(\Omega)} \\ &\leq CC_{\mathrm{err}} \left(\frac{h}{p} + \frac{1}{p} \left(\frac{kh}{\sigma p}\right)^{p}\right) \|e_{h}\|_{L^{2}(\Omega)} \inf_{z_{h} \in V_{h}} \||u - z_{h}\|| \\ &+ CC_{\mathrm{err}}^{2} k^{2} \left(\frac{h^{2}}{p^{2}} + \frac{h}{p^{2}} \left(\frac{kh}{\sigma p}\right)^{p}\right) \|e_{h}\|_{L^{2}(\Omega)}^{2} . \end{split}$$

Noting that $C_{\text{err}} = 1 + \gamma + \frac{kh}{p} \lesssim 1 + \frac{kh}{p}$, there exists a constant C_0 independent of k, h, p, and the penalty parameters such that

(4.10) if
$$\frac{kh}{p} \le C_0 \left(\frac{p}{k}\right)^{\frac{1}{p+1}}$$
 then $CC_{\text{err}}^2 k^2 \left(\frac{h^2}{p^2} + \frac{h}{p^2} \left(\frac{kh}{\sigma p}\right)^p\right) \le CC_{\text{err}}^2 \left(C_0^2 \left(\frac{p}{k}\right)^{\frac{2}{p+1}} + \left(\frac{C_0}{\sigma}\right)^{p+1}\right) \le \frac{1}{2},$

and as a consequence,

(4.11)
$$\|e_h\|_{L^2(\Omega)} \lesssim \left(\frac{h}{p} + \frac{1}{p}\left(\frac{kh}{\sigma p}\right)^p\right) \inf_{z_h \in V_h} \|\|u - z_h\|\|.$$

That is, (4.3) holds.

Next we turn to prove (4.2). Let $u_h^{\pm} \in V_h$ be the elliptic projections of u defined by (3.8)–(3.9) and denote by $\zeta^{\pm} = u - u_h^{\pm}$. It follows from Lemma 3.4 and (4.6) that

$$\begin{split} \|e_{h}\|_{1,h}^{2} &= \operatorname{Re} a_{h}(e_{h}, e_{h}) + \operatorname{Im} a_{h}(e_{h}, e_{h}) \\ &= \operatorname{Re}(a_{h}(e_{h}, e_{h}) - k^{2}(e_{h}, e_{h}) + \mathbf{i}k\langle e_{h}, e_{h}\rangle) + k^{2}(e_{h}, e_{h}) \\ &+ \operatorname{Im}(a_{h}(e_{h}, e_{h}) - k^{2}(e_{h}, e_{h}) + \mathbf{i}k\langle e_{h}, e_{h}\rangle) - k\|e_{h}\|_{L^{2}(\Gamma)}^{2} \\ &= \operatorname{Re}(a_{h}(e_{h}, \zeta^{-}) - k^{2}(e_{h}, \zeta^{-}) + \mathbf{i}k\langle e_{h}, \zeta^{-}\rangle) + k^{2}\|e_{h}\|_{L^{2}(\Omega)}^{2} \\ &+ \operatorname{Im}(a_{h}(e_{h}, \zeta^{-}) - k^{2}(e_{h}, \zeta^{-}) + \mathbf{i}k\langle e_{h}, \zeta^{-}\rangle) - k\|e_{h}\|_{L^{2}(\Gamma)}^{2} \\ &= \operatorname{Re}(a_{h}(\zeta^{+}, \zeta^{-}) - k^{2}(e_{h}, \zeta^{-}) + \mathbf{i}k\langle \zeta^{+}, \zeta^{-}\rangle) - k\|e_{h}\|_{L^{2}(\Omega)}^{2} \\ &+ \operatorname{Im}(a_{h}(\zeta^{+}, \zeta^{-}) - k^{2}(e_{h}, \zeta^{-}) + \mathbf{i}k\langle \zeta^{+}, \zeta^{-}\rangle) - k\|e_{h}\|_{L^{2}(\Omega)}^{2} \\ &+ \operatorname{Im}(a_{h}(\zeta^{+}, \zeta^{-}) - k^{2}(e_{h}, \zeta^{-}) + \mathbf{i}k\langle \zeta^{+}, \zeta^{-}\rangle) - k\|e_{h}\|_{L^{2}(\Gamma)}^{2} \\ &\leq 2\|\zeta^{+}\|_{1,h}\|\zeta^{-}\|_{1,h} + k^{2}\|\zeta^{-}\|_{L^{2}(\Omega)}^{2} + 2k\|\zeta^{+}\|_{L^{2}(\Gamma)}\|\zeta^{-}\|_{L^{2}(\Gamma)} + 2k^{2}\|e_{h}\|_{L^{2}(\Omega)}^{2} - k\|e_{h}\|_{L^{2}(\Gamma)}^{2}. \end{split}$$

Therefore, from Lemma 3.5 and noting that $\frac{kh}{p}$, $\gamma \leq 1$,

$$|||e_h||| \lesssim \inf_{z_h \in V_h} |||u - z_h||| + k ||e_h||_{L^2(\Omega)}.$$

which together with (4.3) implies (4.2). This completes the proof of the theorem.

Remark 4.1. (i) Noting that if $\gamma = 0$ then the CIP-FEM becomes the FEM, the above theorem and the three corollaries below hold for the standard FEM.

(ii) Noting that the wave length is $\frac{2\pi}{k}$, the condition (4.1) roughly says that about $\frac{2\pi}{C_0} \left(\frac{k}{p}\right)^{\frac{1}{p+1}}$ degrees of freedom are needed in one wave length.

(iii) From (4.10), clearly, the theorem holds under the following condition which is a little more general (and a little more complicated as well) than (4.1):

(4.12)
$$\frac{k^2h^2}{p^2} + \frac{k}{p}\left(\frac{kh}{\sigma p}\right)^{p+1} \text{ is sufficiently small.}$$

(iv) If

(4.13)
$$\frac{kh}{p} + \frac{k}{p} \left(\frac{kh}{\sigma p}\right)^p \text{ is sufficiently small,}$$

then CIP-FEM, as well as the FEM, is pollution free in the norm $\||\cdot|\|$, i.e., $\||u - u_h\|| \lesssim \inf_{z_h \in V_h} \||u - z_h\||$. The condition (4.13) improves a little the condition of $\frac{kh}{p} + k \left(\frac{kh}{\sigma p}\right)^p$ small enough given in [39] for the FEM, because, for example, if we choose $k \left(\frac{kh}{\sigma p}\right)^p = c$ for some fixed constant c so that the latter condition does not hold, then $\frac{kh}{p} + \frac{k}{p} \left(\frac{kh}{\sigma p}\right)^p = \left(\frac{c}{k}\right)^{\frac{1}{p}} + \frac{c}{p}$ may be small enough for an appropriate choice of p and large k.

(v) The pre-asymptotic error estimates for the linear CIP-FEM and FEM (p = 1) in two and three dimensions are given in the first part of this series [47]. For the pre-asymptotic error estimates for the linear and hp version of the FEM in one dimension, we refer to [34] and [35], respectively. To the best of the authors' knowledge, there have been no pre-asymptotic error estimates for the higher order FEM and CIP-FEM $(p \ge 2)$ in two and three dimensions given in the literature so far.

Pre-asymptotic Analysis of CIP-FEM and FEM

From Remark 4.1(iv) we have the following corollary which gives practical sufficient conditions for asymptotic estimates of the CIP-FEM and the FEM.

Corollary 4.1. Let u and u_h denote the solutions of (1.1)-(1.2) and (2.4), respectively. Suppose $0 \le \gamma \lesssim$ 1. Then the following asymptotic error estimate

$$|||u - u_h||| \lesssim \inf_{z_h \in V_h} |||u - z_h|||$$

holds under any of the following three conditions:

(4.14)
$$\left(\frac{k}{p}\right)^{\frac{1}{p}}\frac{kh}{p}$$
 is sufficiently small,

(4.15)
$$p \gtrsim (\ln k)^{\frac{1}{2}} \text{ and } \left(\frac{k}{p}\right)^{\frac{1}{p+1}} \frac{kh}{p} \text{ is sufficiently small,}$$

(4.16)
$$p \gtrsim \ln k \text{ and } \frac{k h}{p} \text{ is sufficiently small.}$$

Proof. (4.14) implies (4.13) and the conditions (4.15)–(4.16) are all sufficient conditions of (4.14). \Box

Remark 4.2. (i) Our asymptotic estimates hold for both the CIP-FEM and the FEM. Melenk and Sauter [39, 40] proved recently the asymptotic error estimates for the FEM under either of the condition (4.16) and the following one:

(4.17)
$$p = O(1)$$
 fixed independent of k and $k^{1+\frac{1}{p}}h$ is sufficiently small.

The above condition is similar to the condition (4.14) but we prefer the later one because it does not require p = O(1).

(ii) The condition (4.15) says that the theorem 4.1 gives actually asymptotic estimates in the norm $\|\|\cdot\|\|$ under the further condition of $p \gtrsim (\ln k)^{\frac{1}{2}}$. This condition is not strict from the practical point of view. For example if we take $p \simeq (\ln k)^{\frac{1}{2}}$ then p grows slowly as k increases, and so is $(\frac{k}{p})^{\frac{1}{p+1}} \simeq k^{\frac{1}{\sqrt{\ln k}}}$ which is $o(k^{\varepsilon})$ for any positive constant ε .

(iii) The condition (4.14) is convenient for fixed p. The condition (4.16) requires $p \gtrsim \ln k$ and is perfect when implementations of sufficiently high order CIP-FEM or FEM are available. And the condition (4.15) which requires merely $p \gtrsim (\ln k)^{\frac{1}{2}}$ is suitable when only "medium" order methods can be used.

From Theorem 4.1 and Lemma 3.3, we have the following corollary which gives estimates for H^2 regular solutions.

Corollary 4.2. Suppose the solution $u \in H^2(\Omega)$. Under the conditions of Theorem 4.1, there hold the following estimates:

(4.18)
$$||u - u_h||_{1,h} \lesssim \left(\frac{h}{p} + \frac{1}{p}\left(\frac{kh}{\sigma p}\right)^p + \frac{k}{p^2}\left(\frac{kh}{\sigma p}\right)^{2p}\right) C_{f,g},$$

(4.19)
$$\|u - u_h\|_{L^2(\Omega)} \lesssim \left(\frac{h^2}{p^2} + \frac{1}{p^2} \left(\frac{kh}{\sigma p}\right)^{2p}\right) C_{f,g}.$$

Remark 4.3. (i) If $p = o((\ln k)^{\frac{1}{2}})$ then the condition (4.1) is weaker than the condition (4.14) for asymptotic error estimates, and the estimate (4.18) is a pre-asymptotic one with pollution term $O\left(\frac{k}{p^2}\left(\frac{kh}{\sigma p}\right)^{2p}\right)$.

(ii) Pre-asymptotic error analysis and dispersion analysis are two main tools to understand numerical behaviors in short wave computations. The later one which is usually performed on structured meshes estimates the error between the wave number k of the continuous problem and some discrete wave number ω [1, 2, 21, 31, 34, 35, 46, 45]. In particular, it is shown for the hp-FEM (cf. [1, 35]) that

$$k - \omega = \begin{cases} O(k^{2p+1}h^{2p}) & \text{if } kh \ll 1, \\ O\left(\frac{k}{p}\left(\frac{ekh}{4p}\right)^{2p}\right) & \text{if } kh \gg 1 \text{ and } 2p+1 > \frac{ekh}{2}. \end{cases}$$

By contrast, our pre-asymptotic error analysis gives the error between the exact solution u and the discrete solution u_h and works for unstructured meshes. Clearly, our pollution error bounds in H^1 -norm coincide with the phase difference $|k - \omega|$ as above.

(iii) Our pre-asymptotic estimates in Theorem 4.1 and Corollary 4.2–4.3 hold in particular under the following condition

(4.20)
$$p = O(1)$$
 fixed independent of k and $k^{1+\frac{1}{p+1}}h$ is small enough,

which extends the asymptotic range given by (4.17).

By combining Theorem 4.1 and Lemma 3.2, we have the following corollary which gives estimates for H^s regular solutions (s > 2).

Corollary 4.3. Suppose the solution $u \in H^s(\Omega)$, s > 2. Let $\mu = \min\{p+1, s\}$. Under the conditions of Theorem 4.1, there hold the following estimates:

(4.21)
$$\|u - u_h\|_{1,h} \lesssim \left(1 + \frac{k}{p} \left(\frac{kh}{\sigma p}\right)^p\right) \frac{h^{\mu-1}}{p^{s-1}} \|u\|_{H^s(\Omega)},$$

(4.22)
$$\|u - u_h\|_{L^2(\Omega)} \lesssim \left(\frac{h}{p} + \frac{1}{p} \left(\frac{kh}{\sigma p}\right)^p\right) \frac{h^{\mu - 1}}{p^{s - 1}} \|u\|_{H^s(\Omega)},$$

where the invisible constants are independent on p, h, k, and the penalty parameters, but may depend on s. In particular, if $u \in H^{p+1}(\Omega)$ is an oscillating solution in the sense of [35, Definition 3.2], i.e.

$$\|u\|_{H^{p+1}(\Omega)} \lesssim k^p$$

then the following estimates hold:

(4.23)
$$\|u - u_h\|_{1,h} \lesssim C(p) \left(\left(\frac{kh}{\sigma p}\right)^p + \frac{k}{p} \left(\frac{kh}{\sigma p}\right)^{2p} \right),$$

(4.24)
$$\|u - u_h\|_{L^2(\Omega)} \lesssim \frac{C(p)}{k} \left(\left(\frac{kh}{\sigma p}\right)^{p+1} + \frac{k}{p} \left(\frac{kh}{\sigma p}\right)^{2p} \right).$$

Remark 4.4. (i) Corollary 4.3 certainly holds for s = 2. We exclude the the case s = 2 in this corollary because the estimates in (4.21)–(4.22) with s = 2 are worse than those in (4.18)–(4.19). For s > 2, Corollary 4.3 gives optimal convergence order in h and p while Corollary 4.2 gives only first order convergence in h and p.

(ii) If p = O(1) and u is an oscillating solution then (4.23) shows clearly that the pollution error in H^1 -norm is again bounded by $O(k^{2p+1}h^{2p})$.

By combining Lemma 3.1 and Corollary 4.2 we have the following stability estimates for the CIP-FEM (FEM).

Corollary 4.4. Suppose the solution $u \in H^2(\Omega)$. Under the conditions of Theorem 4.1, there hold the following estimates:

$$||u_h||_{1,h} + k ||u_h||_{L^2(\Omega)} \lesssim C_{f,g},$$

and hence the CIP-FEM is well-posed.

Remark 4.5. If $\gamma = 0$, then this corollary gives the following stability estimate for the standard FEM under the same condition.

$$\|\nabla u_h\|_{L^2(\Omega)} + k \,\|u_h\|_{L^2(\Omega)} \lesssim C_{f,g}.$$

5 Stability estimates for the CIP-FEM

In the previous section we have shown that the CIP-FEM and the FEM are stable under the condition that $\frac{kh}{p} \leq C_0 \left(\frac{p}{k}\right)^{\frac{1}{p+1}}$ and $0 \leq \gamma \lesssim 1$. The goal of this section is to derive stability estimates (or a priori estimates) for the CIP-FEM (2.4) for any k, h, p, and $\gamma > 0$.

Some stability estimates have been proved for the discontinuous Galerkin methods (cf. [26, 27, 28]) and the spectral-Galerkin methods (cf. [44]). Their analyses mimic the stability analysis for the PDE solutions given in [20, 32, 38]. The key idea is to use the test functions $v = u_h$ and $v = (x - x_\Omega) \cdot \nabla u_h$, respectively, and use the Rellich identity(for the Laplacian), where x_Ω is a point such that the domain Ω is strictly star-shaped with respect to it (see (1.10)). As for our CIP-FEM (2.4), although the test function $v_h = u_h$ can be still be used, the test function $v_h = (x - x_\Omega) \cdot \nabla u_h$ does not apply since it is discontinuous and hence not in the test space V_h . We surround this difficulty by taking the L^2 projection of $(x - x_\Omega) \cdot \nabla u_h$ to the finite element space V_h . To analyze the error of the L^2 projection, we recall the so-called Oswald interpolation operator \mathcal{I}_{Os} , which has been analyzed in [15, 33, 36, 41]. Let W_h be the space of (discontinuous) piecewise mapped polynomials:

(5.1)
$$W_h := \prod_{K \in \mathcal{T}_h} \mathcal{P}_p(F_K^{-1}(K)).$$

Lemma 5.1. There exist an operator $\mathcal{I}_{Os} : W_h \mapsto V_h$ and a constant C_p depending only on p, such that, for all $K \in \mathcal{T}_h$, the following estimate holds:

(5.2)
$$\forall v_h \in W_h, \quad \|v_h - \mathcal{I}_{\text{Os}} v_h\|_{L^2(K)} \lesssim C_p h_K^{\frac{1}{2}} \sum_{e \in \mathcal{E}_h^I \cap K} \|[v_h]\|_{L^2(e)},$$

where $C_p = p^{-1}$ for Cartesian meshes and $C_p = p^{\frac{d-3}{2}}$ for triangulations (d = 2, 3).

Lemma 5.2. Let Q_h be the L^2 projection onto V_h and let C_p be the constant in Lemma 5.1. Then the following estimate holds:

(5.3)
$$\forall v_h \in W_h, \quad \|v_h - Q_h v_h\|_{L^2(\mathcal{T}_h)} \lesssim C_p \bigg(\sum_{e \in \mathcal{E}_h^I} h_e \| [v_h] \|_{L^2(e)}^2 \bigg)^{\frac{1}{2}},$$

(5.4)
$$\forall v_h \in W_h, \quad \|\nabla(v_h - Q_h v_h)\|_{L^2(\mathcal{T}_h)} \lesssim \frac{p^2 C_p}{h} \left(\sum_{e \in \mathcal{E}_h^I} h_e \|[v_h]\|_{L^2(e)}^2\right)^{\frac{1}{2}}.$$

Proof. (5.3) follows from $||v_h - Q_h v_h||_{L^2(\mathcal{T}_h)} \leq ||v_h - \mathcal{I}_{OS} v_h||_{L^2(\mathcal{T}_h)}$ and Lemma 5.1. (5.4) follows from the inverse inequality and (5.3).

We cite the following lemma [26, Lemma 4.1], which establishes two integral identities and play a crucial role in our analysis.

Lemma 5.3. Let
$$\alpha(x) := x - x_{\Omega}, v \in \prod_{K \in \mathcal{T}_h} H^2(K), K \in \mathcal{T}_h \text{ and } e \in \mathcal{E}_h^I$$
. Then there hold

(5.5)
$$d\|v\|_{L^2(K)}^2 + 2\operatorname{Re}(v, \alpha \cdot \nabla v)_K = \int_{\partial K} \alpha \cdot n_K |v|^2,$$

(5.6)
$$(d-2) \|\nabla v\|_{L^2(K)}^2 + 2\operatorname{Re}(\nabla v, \nabla(\alpha \cdot \nabla v))_K = \int_{\partial K} \alpha \cdot n_K |\nabla v|^2.$$

Here x_{Ω} is a point such that the domain Ω is strictly star-shaped with respect to it (see (1.10)).

Remark 5.1. The identity (5.6) can be viewed as a local version of the Rellich identity for the Laplacian \triangle (cf. [20]).

Lemma 5.4. Let $u_h \in V_h$ solve (2.4). Then

(5.7)
$$\|\nabla u_h\|_{L^2(\Omega)}^2 - k^2 \|u_h\|_{L^2(\Omega)}^2 \le 2 \|f\|_{L^2(\Omega)} \|u_h\|_{L^2(\Omega)} + \frac{1}{k} \|g\|_{L^2(\Gamma)}^2,$$

(5.8)
$$\sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \left[\frac{\partial u_h}{\partial n_e} \right] \right\|_{L^2(e)}^2 + k \|u_h\|_{L^2(\Gamma)}^2 \le 2 \|f\|_{L^2(\Omega)} \|u_h\|_{L^2(\Omega)} + \frac{1}{k} \|g\|_{L^2(\Gamma)}^2.$$

Proof. Taking $v_h = u_h$ in (2.4) yields

(5.9)
$$a_h(u_h, u_h) - k^2 \|u_h\|_{L^2(\Omega)}^2 + \mathbf{i}k\|u_h\|_{L^2(\Gamma)}^2 = (f, u_h) + \langle g, u_h \rangle$$

Therefore, taking real part and imaginary part of the above equation, we get

(5.10)
$$\|\nabla u_h\|_{L^2(\Omega)}^2 - k^2 \|u_h\|_{L^2(\Omega)}^2 \le |(f, u_h) + \langle g, u_h \rangle|,$$

(5.11)
$$\sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \left[\frac{\partial u_h}{\partial n_e} \right] \right\|_{L^2(e)}^2 + k \|u_h\|_{L^2(\Gamma)}^2 \le |(f, u_h) + \langle g, u_h \rangle|.$$

From (5.11),

$$\begin{split} \sum_{e \in \mathcal{E}_{h}^{I}} \gamma_{e} \frac{h_{e}}{p^{2}} \left\| \left[\frac{\partial u_{h}}{\partial n_{e}} \right] \right\|_{L^{2}(e)}^{2} + k \|u_{h}\|_{L^{2}(\Gamma)}^{2} \leq |(f, u_{h}) + \langle g, u_{h} \rangle| \\ \leq \|f\|_{L^{2}(\Omega)} \|u_{h}\|_{L^{2}(\Omega)} + \frac{1}{2k} \|g\|_{L^{2}(\Gamma)}^{2} + \frac{k}{2} \|u_{h}\|_{L^{2}(\Gamma)}^{2} \end{split}$$

which implies (5.8). From (5.10) and (5.8),

$$\begin{aligned} \|\nabla u_h\|_{L^2(\Omega)}^2 - k^2 \|u_h\|_{L^2(\Omega)}^2 &\leq \|f\|_{L^2(\Omega)} \|u_h\|_{L^2(\Omega)} + \frac{1}{2k} \|g\|_{L^2(\Gamma)}^2 + \frac{k}{2} \|u_h\|_{L^2(\Gamma)}^2 \\ &\leq 2 \|f\|_{L^2(\Omega)} \|u_h\|_{L^2(\Omega)} + \frac{1}{k} \|g\|_{L^2(\Gamma)}^2 . \end{aligned}$$

That is, (5.7) holds. This completes the proof of the lemma.

From (5.7) and (5.8) we can bound $\|\nabla u_h\|_{L^2(\Omega)}$ and the jumps of $\frac{\partial u_h}{\partial n_e}$ across $e \in \mathcal{E}_h^I$. In order to get the desired a priori estimates, we need to derive a reverse inequality whose coefficients can be controlled. Such a reverse inequality, which is often difficult to get under practical mesh constraints, and stability estimates for u_h will be derived next.

Theorem 5.1. Let $u_h \in V_h$ solve (2.4) and suppose $\gamma_e \simeq \gamma$, h_K , $h_e \simeq h$. Then

(5.12)
$$k \|u_h\|_{L^2(\Omega)} + \|u_h\|_{1,h} \lesssim C_{\text{sta}} M(f,g),$$

where

(5.13)
$$M(f,g) := \|f\|_{L^2(\Omega)} + \|g\|_{L^2(\Gamma)},$$

(5.14)
$$C_{\text{sta}} := 1 + \frac{1}{k} \left(\frac{\gamma p^4}{h^2} + \frac{p^6 C_p^2}{\gamma h^2} \right)$$

Proof. We divide the proof into three steps.

Step 1: Derivation of a representation identity for $||u_h||_{L^2(\Omega)}$. Define v_h by $v_h|_K = \alpha \cdot \nabla u_h|_K$ for every $K \in \mathcal{T}_h$, denote by $w_h = Q_h v_h$, hence $w_h \in V_h$. Using this w_h as a test function in (2.4) and taking the real part of resulted equation we get

(5.15)
$$-k^2 \operatorname{Re}(u_h, w_h) = \operatorname{Re}((f, w_h) + \langle g, w_h \rangle - a_h(u_h, w_h) - \mathbf{i}k \langle u_h, w_h \rangle).$$

It follows from (5.5), (5.9), and (5.15) that

$$(5.16) \qquad 2k^{2} \|u_{h}\|_{L^{2}(\Omega)}^{2} = k^{2} \sum_{K \in \mathcal{T}_{h}} \int_{\partial K} \alpha \cdot n_{K} |u_{h}|^{2} - (d-2)k^{2} \|u_{h}\|_{L^{2}(\Omega)}^{2} - 2k^{2} \operatorname{Re}(u_{h}, w_{h}) - 2k^{2} \operatorname{Re}(u_{h}, v_{h} - w_{h}) = k^{2} \sum_{K \in \mathcal{T}_{h}} \int_{\partial K} \alpha \cdot n_{K} |u_{h}|^{2} + (d-2) \operatorname{Re}((f, u_{h}) + \langle g, u_{h} \rangle - a_{h}(u_{h}, u_{h})) + 2 \operatorname{Re}((f, w_{h}) + \langle g, w_{h} \rangle - a_{h}(u_{h}, w_{h}) - \mathbf{i}k \langle u_{h}, w_{h} \rangle) = k^{2} \sum_{K \in \mathcal{T}_{h}} \int_{\partial K} \alpha \cdot n_{K} |u_{h}|^{2} + (d-2) \operatorname{Re}((f, u_{h}) + \langle g, u_{h} \rangle) + 2 \operatorname{Re}((f, w_{h}) + \langle g, w_{h} \rangle) - \sum_{K \in \mathcal{T}_{h}} ((d-2) \|\nabla u_{h}\|_{L^{2}(K)}^{2} + 2 \operatorname{Re}(\nabla u_{h}, \nabla w_{h})_{K}) - 2 \operatorname{Re} J(u_{h}, w_{h}) + 2k \operatorname{Im}\langle u_{h}, w_{h} \rangle.$$

Since u_h is continuous, we have

(5.17)
$$\sum_{K \in \mathcal{T}_h} \int_{\partial K} \alpha \cdot n_K |u_h|^2 = 2 \sum_{e \in \mathcal{E}_h^I} \operatorname{Re}\langle \alpha \cdot n_e\{u_h\}, [u_h]\rangle_e + \langle \alpha \cdot n_\Omega, |u_h|^2 \rangle$$
$$= \langle \alpha \cdot n_\Omega, |u_h|^2 \rangle.$$

Using the identity $|a|^2 - |b|^2 = \operatorname{Re}(a+b)(\bar{a}-\bar{b})$ followed by the Rellich identity (5.6) we get

$$(5.18) \qquad \sum_{K \in \mathcal{T}_{h}} ((d-2) \| \nabla u_{h} \|_{L^{2}(K)}^{2} + 2 \operatorname{Re}(\nabla u_{h}, \nabla w_{h})_{K}) \\ = \sum_{K \in \mathcal{T}_{h}} ((d-2) \| \nabla u_{h} \|_{L^{2}(K)}^{2} + 2 \operatorname{Re}(\nabla u_{h}, \nabla v_{h})_{K}) \\ + 2 \sum_{K \in \mathcal{T}_{h}} \operatorname{Re}(\nabla u_{h}, \nabla (w_{h} - v_{h}))_{K} \\ = \sum_{K \in \mathcal{T}_{h}} \int_{\partial K} \alpha \cdot n_{K} | \nabla u_{h} |^{2} + 2 \sum_{K \in \mathcal{T}_{h}} \operatorname{Re}(\nabla u_{h}, \nabla (w_{h} - v_{h}))_{K} \\ = 2 \sum_{e \in \mathcal{E}_{h}^{I}} \operatorname{Re}\langle \alpha \cdot n_{e} \{ \nabla u_{h} \}, [\nabla u_{h}] \rangle_{e} + \langle \alpha \cdot n_{\Omega}, |\nabla u_{h}|^{2} \rangle \\ + 2 \sum_{K \in \mathcal{T}_{h}} \operatorname{Re}(\nabla u_{h}, \nabla (w_{h} - v_{h}))_{K}.$$

Plugging (5.17) and (5.18) into (5.16) gives

$$(5.19) \qquad 2k^2 \|u_h\|_{L^2(\Omega)}^2 = (d-2)\operatorname{Re}((f,u_h) + \langle g, u_h \rangle) + 2\operatorname{Re}((f,v_h) + \langle g, v_h \rangle) + k^2 \langle \alpha \cdot n_\Omega, |u_h|^2 \rangle + 2k\operatorname{Im}\langle u_h, v_h \rangle - \langle \alpha \cdot n_\Omega, |\nabla u_h|^2 \rangle - 2\sum_{e \in \mathcal{E}_h^I} \operatorname{Re}\langle \alpha \cdot n_e \{\nabla u_h\}, [\nabla u_h] \rangle_e - 2\operatorname{Re} J(u_h, v_h) - 2\operatorname{Re} J(u_h, w_h - v_h) - 2\sum_{K \in \mathcal{T}_h} \operatorname{Re}(\nabla u_h, \nabla(w_h - v_h))_K + 2k\operatorname{Im}\langle u_h, w_h - v_h \rangle + 2\operatorname{Re}((f, w_h - v_h) + \langle g, w_h - v_h \rangle).$$

Step 2: Derivation of a reverse inequality. We bound each term on the right-hand side of (5.19). We have

(5.20)
$$2\operatorname{Re}((f, v_h) + \langle g, v_h \rangle) \leq C \|f\|_{L^2(\Omega)}^2 + \frac{1}{8} \|\nabla u_h\|_{L^2(\Omega)}^2 + C \|g\|_{L^2(\Gamma)}^2 + \frac{c_\Omega}{4} \sum_{e \in \mathcal{E}_h^B} \|\nabla u_h\|_{L^2(e)}^2.$$

It is clear that

(5.21)
$$k^2 \langle \alpha \cdot n_{\Omega}, |u_h|^2 \rangle \le Ck^2 ||u_h||_{L^2(\Gamma)}^2.$$

It follows from the star-shaped assumption on Ω that

(5.22)
$$2k \operatorname{Im}\langle u_h, v_h \rangle - \langle \alpha \cdot n_{\Omega}, |\nabla u_h|^2 \rangle$$
$$\leq Ck \sum_{e \in \mathcal{E}_h^B} \|u_h\|_{L^2(e)} \|\nabla u_h\|_{L^2(e)} - c_{\Omega} \sum_{e \in \mathcal{E}_h^B} \|\nabla u_h\|_{L^2(e)}^2$$
$$\leq Ck^2 \|u_h\|_{L^2(\Gamma)}^2 - \frac{c_{\Omega}}{2} \sum_{e \in \mathcal{E}_h^B} \|\nabla u_h\|_{L^2(e)}^2.$$

For an edge/face $e \in \mathcal{E}_h^I$, let K_e and K'_e denote the two elements in \mathcal{T}_h that share e. We have

$$(5.23) \qquad -2\sum_{e\in\mathcal{E}_{h}^{I}}\operatorname{Re}\langle\alpha\cdot n_{e}\{\nabla u_{h}\}, [\nabla u_{h}]\rangle_{e} \leq C\sum_{e\in\mathcal{E}_{h}^{I}}\left(\frac{p^{2}}{h}\right)^{\frac{1}{2}}\|\nabla u_{h}\|_{L^{2}(K_{e}\bigcup K_{e}')}\left\|\left[\frac{\partial u_{h}}{\partial n_{e}}\right]\right\|_{L^{2}(e)}$$
$$\leq \frac{1}{8}\|\nabla u_{h}\|_{L^{2}(\Omega)}^{2} + C\frac{p^{4}}{\gamma h^{2}}\sum_{e\in\mathcal{E}_{h}^{I}}\gamma_{e}\frac{h_{e}}{p^{2}}\left\|\left[\frac{\partial u_{h}}{\partial n_{e}}\right]\right\|_{L^{2}(e)}^{2}.$$

From (2.2), the trace and inverse inequalities (cf. Lemma A.1), we have

$$(5.24) -2\operatorname{Re} J(u_h, v_h) = -2\operatorname{Re} \sum_{e \in \mathcal{E}_h^I} \mathbf{i} \gamma_e \frac{h}{p^2} \left\langle \left[\frac{\partial u_h}{\partial n_e} \right], \left[\frac{\partial v_h}{\partial n_e} \right] \right\rangle_e \\ \lesssim \sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \left[\frac{\partial u_h}{\partial n_e} \right] \right\|_{L^2(e)} \left\| \left[\frac{\partial v_h}{\partial n_e} \right] \right\|_{L^2(e)} \\ \lesssim \sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \left[\frac{\partial u_h}{\partial n_e} \right] \right\|_{L^2(e)} \left(\frac{p^2}{h} \right)^{\frac{1}{2}} \sum_{K=K_e, K'_e} |v_h|_{H^1(K)} \\ \lesssim \sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \left[\frac{\partial u_h}{\partial n_e} \right] \right\|_{L^2(e)} \left(\frac{p^2}{h} \right)^{\frac{1}{2}} \sum_{K=K_e, K'_e} \frac{p^2}{h} \| \nabla u_h \|_{L^2(K)} \\ \le \frac{1}{8} \left\| \nabla u_h \right\|_{L^2(\Omega)}^2 + \frac{\gamma p^4}{h^2} \sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \left[\frac{\partial u_h}{\partial n_e} \right] \right\|_{L^2(e)}^2.$$

Next we estimate the terms containing $v_h - w_h = v_h - Q_h v_h$. From Lemma 5.2 and the definition of v_h ,

(5.25)
$$\|v_h - w_h\|_{L^2(\mathcal{T}_h)} \lesssim \frac{pC_p}{\gamma^{\frac{1}{2}}} \bigg(\sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \bigg[\frac{\partial u_h}{\partial n_e} \bigg] \right\|_{L^2(e)}^2 \bigg)^{\frac{1}{2}},$$

(5.26)
$$|v_h - w_h|_{H^1(\mathcal{T}_h)} \lesssim \frac{p^3 C_p}{\gamma^{\frac{1}{2}} h} \left(\sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \left[\frac{\partial u_h}{\partial n_e} \right] \right\|_{L^2(e)}^2 \right)^{\frac{1}{2}}.$$

From (5.26) and the trace inequality in Lemma A.1,

$$(5.27) -2\operatorname{Re} J(u_h, w_h - v_h) \leq \sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \left[\frac{\partial u_h}{\partial n_e} \right] \right\|_{L^2(e)} \left\| \left[\nabla(w_h - v_h) \right] \right\|_{L^2(e)} \\ \leq C \sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \left[\frac{\partial u_h}{\partial n_e} \right] \right\|_{L^2(e)} \sum_{K = K_e, K'_e} \left(\frac{p^2}{h} \right)^{\frac{1}{2}} \left\| \nabla(w_h - v_h) \right\|_{L^2(K)} \\ \leq C \gamma^{\frac{1}{2}} \left(\sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \left[\frac{\partial u_h}{\partial n_e} \right] \right\|_{L^2(e)}^2 \right)^{\frac{1}{2}} \left\| v_h - w_h \right\|_{H^1(\mathcal{T}_h)} \\ \leq C \frac{p^3 C_p}{h} \sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \left[\frac{\partial u_h}{\partial n_e} \right] \right\|_{L^2(e)}^2.$$

And we have,

(5.28)
$$-2\sum_{K\in\mathcal{T}_{h}}\operatorname{Re}(\nabla u_{h},\nabla(w_{h}-v_{h}))_{K} \leq 2\sum_{K\in\mathcal{T}_{h}}\|\nabla u_{h}\|_{L^{2}(K)}\|\nabla(w_{h}-v_{h})\|_{L^{2}(K)}$$
$$\leq \frac{1}{8}\|\nabla u_{h}\|_{L^{2}(\Omega)}^{2}+C|v_{h}-w_{h}|_{H^{1}(\mathcal{T}_{h})}^{2}$$
$$\leq \frac{1}{8}\|\nabla u_{h}\|_{L^{2}(\Omega)}^{2}+C\frac{p^{6}C_{p}^{2}}{\gamma h^{2}}\sum_{e\in\mathcal{E}_{h}^{I}}\gamma_{e}\frac{h_{e}}{p^{2}}\left\|\left[\frac{\partial u_{h}}{\partial n_{e}}\right]\right\|_{L^{2}(e)}^{2}.$$

From (5.25) and Lemma A.1,

(5.29)
$$2k \operatorname{Im}\langle u_{h}, w_{h} - v_{h} \rangle \leq Ck \sum_{e \in \mathcal{E}_{h}^{B}} \|u_{h}\|_{L^{2}(e)} \|w_{h} - v_{h}\|_{L^{2}(e)}$$
$$\leq Ck \|u_{h}\|_{L^{2}(\Gamma)} \left(\frac{p^{2}}{h}\right)^{\frac{1}{2}} \|w_{h} - v_{h}\|_{L^{2}(\mathcal{T}_{h})}$$
$$\leq Ck \|u_{h}\|_{L^{2}(\Gamma)} \left(\frac{p^{2}}{h}\right)^{\frac{1}{2}} \frac{pC_{p}}{\gamma^{\frac{1}{2}}} \left(\sum_{e \in \mathcal{E}_{h}^{I}} \gamma_{e} \frac{h_{e}}{p^{2}} \left\|\left[\frac{\partial u_{h}}{\partial n_{e}}\right]\right\|_{L^{2}(e)}^{2}\right)^{\frac{1}{2}}$$
$$\leq Cp^{2}C_{p} \left(\frac{k}{\gamma h}\right)^{\frac{1}{2}} \left(k \|u_{h}\|_{L^{2}(\Gamma)}^{2} + \sum_{e \in \mathcal{E}_{h}^{I}} \gamma_{e} \frac{h_{e}}{p^{2}} \left\|\left[\frac{\partial u_{h}}{\partial n_{e}}\right]\right\|_{L^{2}(e)}^{2}\right).$$

From (5.25), (5.26), and Lemma A.1,

(5.30)
$$2\operatorname{Re}((f, w_{h} - v_{h}) + \langle g, w_{h} - v_{h} \rangle) \\ \lesssim \|f\|_{L^{2}(\Omega)} \|w_{h} - v_{h}\|_{L^{2}(\mathcal{T}_{h})} + \|g\|_{L^{2}(\Gamma)} \left(\frac{p^{2}}{h}\right)^{\frac{1}{2}} \|w_{h} - v_{h}\|_{L^{2}(\mathcal{T}_{h})} \\ \lesssim \left(\|f\|_{L^{2}(\Omega)} + \|g\|_{L^{2}(\Gamma)}\right) \frac{p^{2}C_{p}}{(\gamma h)^{\frac{1}{2}}} \left(\sum_{e \in \mathcal{E}_{h}^{I}} \gamma_{e} \frac{h_{e}}{p^{2}} \left\|\left[\frac{\partial u_{h}}{\partial n_{e}}\right]\right\|_{L^{2}(e)}^{2}\right)^{\frac{1}{2}} \\ \lesssim \|f\|_{L^{2}(\Omega)}^{2} + \|g\|_{L^{2}(\Gamma)}^{2} + \frac{p^{4}C_{p}^{2}}{\gamma h} \sum_{e \in \mathcal{E}_{h}^{I}} \gamma_{e} \frac{h_{e}}{p^{2}} \left\|\left[\frac{\partial u_{h}}{\partial n_{e}}\right]\right\|_{L^{2}(e)}^{2}.$$

Plugging (5.20)–(5.24) and (5.27)–(5.30) into (5.19) we obtain

$$\begin{aligned} 2k^2 \|u_h\|_{L^2(\Omega)}^2 &\leq |(f, u_h) + \langle g, u_h \rangle| + C(\|f\|_{L^2(\Omega)}^2 + \|g\|_{L^2(\Gamma)}^2) \\ &+ \frac{1}{2} \|\nabla u_h\|_{L^2(\Omega)}^2 - \frac{c_\Omega}{4} \sum_{e \in \mathcal{E}_h^B} \|\nabla u_h\|_{L^2(e)}^2 + C\left(k + p^2 C_p\left(\frac{k}{\gamma h}\right)^{\frac{1}{2}}\right) k \|u_h\|_{L^2(\Gamma)}^2 \\ &+ C\left(\frac{p^4}{\gamma h^2} + \frac{\gamma p^4}{h^2} + \frac{p^3 C_p}{h} + \frac{p^6 C_p^2}{\gamma h^2} + p^2 C_p\left(\frac{k}{\gamma h}\right)^{\frac{1}{2}}\right) \sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \left[\frac{\partial u_h}{\partial n_e}\right] \right\|_{L^2(e)}^2. \end{aligned}$$

From (5.8) and the facts

$$C_p \ge p^{-1}, \quad \frac{p^3 C_p}{h} \lesssim \gamma + \frac{p^6 C_p^2}{\gamma h^2}, \quad p^2 C_p \left(\frac{k}{\gamma h}\right)^{\frac{1}{2}} \lesssim \frac{p^6 C_p^2}{\gamma h^2} + \frac{kh}{p^2},$$

we have

$$2k^{2} \|u_{h}\|_{L^{2}(\Omega)}^{2} + \frac{c_{\Omega}}{4} \sum_{e \in \mathcal{E}_{h}^{B}} \|\nabla u_{h}\|_{L^{2}(e)}^{2}$$

$$\leq -\frac{1}{2} \|\nabla u_{h}\|_{L^{2}(\Omega)}^{2} + \|\nabla u_{h}\|_{L^{2}(\Omega)}^{2} + C(\|f\|_{L^{2}(\Omega)}^{2} + \|g\|_{L^{2}(\Gamma)}^{2})$$

$$+ C\left(k + 1 + \frac{\gamma p^{4}}{h^{2}} + \frac{p^{6}C_{p}^{2}}{\gamma h^{2}} + \frac{kh}{p^{2}}\right) \left(\|f\|_{L^{2}(\Omega)} \|u_{h}\|_{L^{2}(\Omega)} + \frac{1}{k} \|g\|_{L^{2}(\Gamma)}^{2}\right).$$

Step 3: Finishing up. It follows from (5.7), (5.13), (5.14), and the above inequality that

$$\begin{aligned} &2k^2 \|u_h\|_{L^2(\Omega)}^2 + \frac{c_\Omega}{4} \sum_{e \in \mathcal{E}_h^B} \|\nabla u_h\|_{L^2(e)}^2 \\ &\leq k^2 \|u_h\|_{L^2(\Omega)}^2 - \frac{1}{2} \|\nabla u_h\|_{L^2(\Omega)}^2 + C(\|f\|_{L^2(\Omega)}^2 + \|g\|_{L^2(\Gamma)}^2) \\ &+ CkC_{\text{sta}} \left(\|f\|_{L^2(\Omega)} \|u_h\|_{L^2(\Omega)} + \frac{1}{k} \|g\|_{L^2(\Gamma)}^2\right) \\ &\leq \frac{3k^2}{2} \|u_h\|_{L^2(\Omega)}^2 - \frac{1}{2} \|\nabla u_h\|_{L^2(\Omega)}^2 + CC_{\text{sta}}^2 M(f,g)^2, \end{aligned}$$

which together with (5.8) implies (5.12). The proof is completed.

Since scheme (2.4) is a linear complex-valued system, an immediate consequence of the stability estimates is the following well-posedness theorem for (2.4).

Theorem 5.2. The CIP-FEM (2.4) has a unique solution for k > 0, h > 0, $p \ge 1$ and $\gamma > 0$.

Remark 5.2. (i) For the general case when the meshes may be nonuniform, Theorem 5.1 and Theorem 5.2 still hold with h replaced by $\underline{h} = \min_{K \in \mathcal{T}_h} h_K$. The proof is similar and is omitted.

(ii) When p = 1, a better stability estimate is derived in [47] by another approach taking advantage of $\Delta u_h = 0$ on each element.

(iii) The stability may be enhanced by an over-penalized technique, just like what is done for the hp-IPDG method (cf. [27]). More precisely, if we replace $a_h(u_h, v_h)$ in the CIP-FEM (2.4) by the following sesquilinear

$$a_{h}^{q}(u_{h},v_{h}) = (\nabla u_{h},\nabla v_{h}) + \sum_{j=1}^{q} \sum_{e \in \mathcal{E}_{h}^{j}} \mathbf{i} \gamma_{j,e} \left(\frac{h_{e}}{p^{2}}\right)^{2j-1} \left\langle \left[\frac{\partial^{j} u_{h}}{\partial n_{e}^{j}}\right], \left[\frac{\partial^{j} v_{h}}{\partial n_{e}^{j}}\right] \right\rangle_{e},$$

with $\gamma_{j,e} \simeq \gamma_j > 0$, $j = 1, 2, \cdots, q$, $2 \le q \le p$, then, after some tedious but similar derivations, we may show that (5.12) holds with C_{sta} replaced by the following stability constant:

$$C_{\text{sta}}^{q} = 1 + \frac{1}{k} \Big(\frac{p^{4} C_{p}^{2}}{\gamma_{1} h} + \frac{p^{4} C_{p}^{2}}{(\gamma_{1} \gamma_{2})^{\frac{1}{2}} h} + \frac{p^{3} C_{p}}{h} \Big(\sum_{j=1}^{q} \frac{\gamma_{j}}{\gamma_{1}} \Big)^{\frac{1}{2}} + \frac{p^{2}}{h} \max_{1 \le j \le q-1} \Big(\frac{\gamma_{j}}{\gamma_{j+1}} \Big)^{\frac{1}{2}} + \gamma_{q} \frac{p^{4}}{h^{2}} \Big).$$

We omit the details here. Clearly, if we choose $\gamma_j \simeq (C_p^2 h)^{\frac{2(j-1)}{2q-1}}, 1 \leq j \leq q$, then $C_{\text{sta}}^q \lesssim \frac{p^4 C_p^{\frac{4q}{2q-1}}}{kh^{\frac{2q}{2q-1}}}$, where C_p is defined in Lemma 5.1. Recall that the best stability constant obtained so far for the hp-IPDG method is $O\left(\frac{p^{\frac{8}{3}}}{kh^{\frac{4}{3}}}\right)$ (cf. [27, Theorem 3.3]).

6 Pre-asymptotic error estimates for the CIP-FEM by using the stability

In this subsection we shall derive error estimates for scheme (2.4). This will be done by exploiting the linearity of the Helmholtz equation and making use of the stability estimates derived in Theorem 5.1 and the projection error estimates established in Lemma 3.6.

Let u and u_h denote the solutions of (1.1)-(1.2) and (2.4), respectively. Recall that $e_h = u - u_h$. Let u_h^+ be the elliptic projection of u as defined in (3.8). Write $e_h = \eta - \xi$ with $\eta := u - u_h^+, \xi := u_h - u_h^+$. From the Galerkin orthogonality (4.6) and (3.8) we get

(6.1)
$$a_h(\xi, v_h) - k^2(\xi, v_h) + \mathbf{i}k\langle\xi, v_h\rangle_{\Gamma} = a_h(\eta, v_h) - k^2(\eta, v_h) + \mathbf{i}k\langle\eta, v_h\rangle_{\Gamma}$$
$$= -k^2(\eta, v_h) \quad \forall v_h \in V_h.$$

The above equation implies that $\xi \in V_h$ is the solution of scheme (2.4) with sources terms $f = -k^2 \eta$ and $g \equiv 0$. Then an application of Theorem 5.1 and Lemma 3.5 immediately gives the following lemma.

Lemma 6.1. $\xi = u_h - u_h^+$ satisfies the following estimate:

(6.2)
$$k\|\xi\|_{L^{2}(\Omega)} + \|\xi\|_{1,h} \lesssim C_{\mathrm{sta}}C_{\mathrm{err}}\frac{k^{2}h}{p}\inf_{z_{h}\in V_{h}}\|u-z_{h}\|.$$

We are ready to state our error estimate results for scheme (2.4), which follows from Lemma 3.5, Lemma 6.1 and an application of the triangle inequality.

Theorem 6.1. Let u and u_h denote the solution of (1.1)-(1.2) and (2.4), respectively. Suppose $u \in H^s(\Omega), s \geq 2$. Then

(6.3)
$$\|u - u_h\|_{1,h} \lesssim \left(1 + C_{\operatorname{err}} C_{\operatorname{sta}} \frac{k^2 h}{p}\right) \inf_{z_h \in V_h} \|u - z_h\|,$$

(6.4)
$$||u - u_h||_{L^2(\Omega)} \lesssim C_{\text{err}} (1 + kC_{\text{sta}}) \frac{h}{p} \inf_{z_h \in V_h} ||u - z_h||,$$

where $C_{\text{err}} := \left(1 + \gamma + \frac{kh}{p}\right)^{\frac{1}{2}}$ and C_{sta} is defined in Theorem 5.1.

Remark 6.1. (i) For the linear CIP-FEM, we have shown that [47], if $\gamma \simeq 1$, then the following preasymptotic error estimate hold for any k, h > 0:

$$||u - u_h||_{1,h} \le (C_1 kh + C_2 \min\{k^3 h^2, 1\})C_{f,g}.$$

(ii) The theorem can also be extended to the over-penalized CIP-FEM. In particular (cf. Remark 5.2 (iii)), if $p \ge q \ge 2$, and $\gamma_j \simeq (C_p^2 h)^{\frac{2(j-1)}{2q-1}}, 1 \le j \le q$, then the following pre-asymptotic error estimate holds for any k, h, p > 0:

$$\|u - u_h\|_{1,h} \le C_1 \inf_{z_h \in V_h} \||u - z_h|| + C_2 \left(1 + \frac{kh}{p}\right)^{\frac{1}{2}} kp^3 C_p^{\frac{4q-4}{2q-1}} h^{\frac{-1}{2q-1}} \inf_{z_h \in V_h} \||u - z_h|\|.$$

The details will be reported in a separate work.

A Approximation by *hp*-finite element

In this appendix, we prove Lemma 3.2 and Lemma 3.3.

A.1 Proof of Lemma 3.2

We need the following trace and inverse inequalities (cf. [43, 15]).

Pre-asymptotic Analysis of CIP-FEM and FEM

Lemma A.1. For any $K \in \mathcal{T}_h$ and $z \in \mathcal{P}_p(F_K^{-1}(K))$,

$$||z||_{L^{2}(\partial K)} \lesssim ph^{-\frac{1}{2}} ||z||_{L^{2}(K)},$$

$$||\nabla z||_{L^{2}(K)} \lesssim p^{2}h^{-1} ||z||_{L^{2}(K)}.$$

We recall the following well-known hp approximation properties (cf. [7, 29, 30, 39]):

Lemma A.2. Let $\mu = \min\{p+1, s\}$. Suppose $u \in H^s(\Omega)$. • There exists $\check{u}_h \in W_h$ such that,

(A.1)
$$\|u - \check{u}_h\|_{H^j(\mathcal{T}_h)} := \left(\sum_{K \in \mathcal{T}_h} \|u - \check{u}_h\|_{H^2(K)}^2\right)^{\frac{1}{2}} \lesssim \frac{h^{\mu - j}}{p^{s - j}} \|u\|_{H^s(\Omega)}, \quad j = 0, 1, 2.$$

• There exists $\hat{u}_h \in V_h$ such that

(A.2)
$$\|u - \hat{u}_h\|_{H^j(\Omega)} \lesssim \frac{h^{\mu-j}}{p^{s-j}} \|u\|_{H^s(\Omega)}, \quad j = 0, 1.$$

Here the invisible constants in the two inequalities above depend on s but are independent of k, h, and p.

Clearly, (3.4) holds (cf. (A.2)). It remains to prove (3.5). It follows from the (A.2) and the trace inequality that

(A.3)
$$\|u - \hat{u}_h\|_{L^2(\Gamma)} \lesssim \|u - \hat{u}_h\|_{L^2(\Omega)}^{\frac{1}{2}} \|u - \hat{u}_h\|_{H^1(\Omega)}^{\frac{1}{2}} \lesssim \frac{h^{\mu - \frac{1}{2}}}{p^{s - \frac{1}{2}}} \|u\|_{H^s(\Omega)}.$$

From the local trace inequality

(A.4)
$$\|v\|_{L^{2}(\partial K)}^{2} \lesssim h_{K}^{-1} \|v\|_{L^{2}(K)}^{2} + \|v\|_{L^{2}(K)} \|\nabla v\|_{L^{2}(K)},$$

we have

$$\sum_{K \in \mathcal{T}_h} \sum_{e \in \partial K} \left\| \frac{\partial (u - \hat{u}_h)}{\partial n_e} \right\|_{L^2(e)}^2 \lesssim h^{-1} \| u - \hat{u}_h \|_{H^1(\mathcal{T}_h)}^2 + \| u - \hat{u}_h \|_{H^1(\mathcal{T}_h)} \| u - \hat{u}_h \|_{H^2(\mathcal{T}_h)}.$$

On the other hand, from Lemma A.1,

$$\begin{aligned} \|u - \hat{u}_h\|_{H^2(\mathcal{T}_h)} &\leq \|u - \check{u}_h\|_{H^2(\mathcal{T}_h)} + \|\check{u}_h - \hat{u}_h\|_{H^2(\mathcal{T}_h)} \\ &\lesssim \|u - \check{u}_h\|_{H^2(\mathcal{T}_h)} + \frac{p^2}{h} \|\check{u}_h - \hat{u}_h\|_{H^1(\mathcal{T}_h)} \end{aligned}$$

Therefore, from (A.1) and (A.2),

$$\sum_{K\in\mathcal{T}_h}\sum_{e\in\partial K}\left\|\frac{\partial(u-\hat{u}_h)}{\partial n_e}\right\|_{L^2(e)}^2 \lesssim \frac{p^2}{h}\frac{h^{2\mu-2}}{p^{2s-2}}\|u\|_{H^s(\Omega)}^2.$$

Then it follows from (A.2) and (2.5) that

$$\begin{aligned} \|u - \hat{u}_h\|_{1,h}^2 &\lesssim \|\nabla(u - \hat{u}_h)\|_{L^2(\Omega)}^2 + \sum_{e \in \mathcal{E}_h^I} \gamma_e \frac{h_e}{p^2} \left\| \left[\frac{\partial(u - \hat{u}_h)}{\partial n_e} \right] \right\|_{L^2(e)}^2 \\ &\lesssim (1 + \gamma) \frac{h^{2\mu - 2}}{p^{2s - 2}} \|u\|_{H^s(\Omega)}^2. \end{aligned}$$

By combining the above estimate, (A.3), and (2.6) we conclude that (3.5) holds. This completes the proof of Lemma 3.2.

A.2 Proof of Lemma 3.3

Lemma A.2 can also be used to approximate the elliptic part of the solution u. The following lemma gives an approximation to the analytic part of the solution u.

Lemma A.3. Suppose $u_{\mathcal{A}}$ is analytic and satisfies (3.2)–(3.3).

• There exists $\check{u}_{\mathcal{A},h} \in W_h$ such that

(A.5)
$$\|u_{\mathcal{A}} - \check{u}_{\mathcal{A},h}\|_{H^{j}(\mathcal{T}_{h})} := \left(\sum_{K \in \mathcal{T}_{h}} \|u_{\mathcal{A}} - \check{u}_{\mathcal{A},h}\|_{H^{2}(K)}^{2}\right)^{\frac{1}{2}} \\ \lesssim \left\{\frac{h^{2-j}}{kp^{2-j}} + \frac{h^{1-j}}{p} \left(\frac{kh}{\sigma p}\right)^{p}\right\} C_{f,g}, \quad j = 0, 1, 2$$

• There exists $\hat{u}_{\mathcal{A},h} \in V_h$ such that

(A.6)
$$\|u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}\|_{H^{j}(\Omega)} \lesssim \left\{\frac{h^{2-j}}{kp^{2-j}} + \frac{h^{1-j}}{p^{2-j}} \left(\frac{kh}{\sigma p}\right)^{p}\right\} C_{f,g}, \quad j = 0, 1.$$

Here $\sigma > 0$ is some constant independent of k, h, and p.

Remark A.1. The following estimate is proved in [39, Theorem 5.5] (see also [40, Proposition 5.3]):

$$\inf_{v_h \in V_h} \left(\left\| \nabla (u_{\mathcal{A}} - v_h) \right\|_{L^2(\Omega)} + k \left\| u_{\mathcal{A}} - v_h \right\|_{L^2(\Omega)} \right) \lesssim \left(1 + \frac{kh}{p} \right) \left(\frac{h}{p} + \left(\frac{kh}{\sigma p} \right)^p \right).$$

Clearly, a combination of our H^1 -estimate and L^2 -estimate in (A.6) gives better upper bound than the above estimate. This is because our L^2 -estimate improves that implied in the proof of [39, Theorem 5.5].

Proof. Following the proof of [39, Theorem 5.5], we start by defining for each element $K \in \mathcal{T}_h$ the constant C_K by

(A.7)
$$C_K^2 := \sum_{p \in \mathbb{N}_0} \frac{\|\nabla^p u_{\mathcal{A}}\|_{L^2(K)}^2}{(2\lambda \max\{p, k\})^{2p}}$$

and we note

(A.8)
$$\|\nabla^p u_{\mathcal{A}}\|_{L^2(K)} \le (2\lambda \max\{p,k\})^p C_K \quad \forall p \in \mathbb{N}_0,$$

(A.9)
$$\sum_{K \in \mathcal{T}_h} C_K^2 \lesssim \frac{1}{k^2} C_{f,g}^2.$$

Then, [39, Lemma C.2] and a scaling argument provides an approximant $\check{u}_{\mathcal{A},h} \in W_h$ which satisfies for j = 0, 1, 2,

$$\|u_{\mathcal{A}} - \check{u}_{\mathcal{A},h}\|_{H^{j}(K)} \lesssim h^{-j} C_{K} \Big\{ \Big(\frac{h}{h+\sigma}\Big)^{p+1} + \Big(\frac{kh}{\sigma p}\Big)^{p+1} \Big\}.$$

Summation over all elements $K \in \mathcal{T}_h$ gives

$$\begin{aligned} \|u_{\mathcal{A}} - \check{u}_{\mathcal{A},h}\|_{H^{j}(\mathcal{T}_{h})}^{2} &:= \Big(\sum_{K \in \mathcal{T}_{h}} \|u_{\mathcal{A}} - \check{u}_{\mathcal{A},h}\|_{H^{j}(K)}^{2}\Big) \\ &\lesssim h^{-2j} \Big[\Big(\frac{h}{h+\sigma}\Big)^{2p+2} + \Big(\frac{kh}{\sigma p}\Big)^{2p+2} \Big] \sum_{K \in \mathcal{T}_{h}} C_{K}^{2} \end{aligned}$$

The combination of the above inequality and (A.9) yields

(A.10)
$$\|u_{\mathcal{A}} - \check{u}_{\mathcal{A},h}\|_{H^{j}(\mathcal{T}_{h})} \lesssim \left[\frac{1}{k}h^{-j}\left(\frac{h}{h+\sigma}\right)^{p+1} + \frac{h^{1-j}}{p}\left(\frac{kh}{\sigma p}\right)^{p}\right]C_{f,g}$$

Pre-asymptotic Analysis of CIP-FEM and FEM

Furthermore, we estimate using $h \leq \text{diam } \Omega$ and $\sigma > 0$ (independent of h)

(A.11)
$$h^{-j} \left(\frac{h}{h+\sigma}\right)^{p+1} = \frac{h^{2-j}}{p^{2-j}} \frac{h^{p-1}}{(h+\sigma)^{p+1}} p^{2-j} \le \frac{h^{2-j}}{p^{2-j}} \left(\frac{h}{h+\sigma}\right)^{p-1} \frac{p^2}{\sigma^2} \le \frac{h^{2-j}}{p^{2-j}} \left(\frac{\operatorname{diam}\,\Omega}{\sigma+\operatorname{diam}\,\Omega}\right)^{p-1} \frac{p^2}{\sigma^2} \le \frac{h^{2-j}}{p^{2-j}},$$

we therefore arrive at

$$\|u_{\mathcal{A}} - \check{u}_{\mathcal{A},h}\|_{H^{j}(\mathcal{T}_{h})} \lesssim \Big[\frac{1}{k}\frac{h^{2-j}}{p^{2-j}} + \frac{h^{1-j}}{p}\Big(\frac{kh}{\sigma p}\Big)^{p}\Big]C_{f,g}$$

That is, (A.5) holds.

Similarly, [39, Lemma C.3] and a scaling argument provides an approximant $u_{\mathcal{A},h} \in V_h$ which satisfies for j = 1,

$$\|u_{\mathcal{A}} - u_{\mathcal{A},h}\|_{H^1(K)} \lesssim h^{-1} C_K \left\{ \left(\frac{h}{h+\sigma}\right)^{p+1} + \left(\frac{kh}{\sigma p}\right)^{p+1} \right\}$$

Summation over all elements $K \in \mathcal{T}_h$ gives

$$\|u_{\mathcal{A}} - u_{\mathcal{A},h}\|_{H^1(\Omega)}^2 \lesssim \left[h^{-2} \left(\frac{h}{h+\sigma}\right)^{2p+2} + \frac{k^2}{p^2} \left(\frac{kh}{\sigma p}\right)^{2p}\right] \sum_{K \in \mathcal{T}_h} C_K^2.$$

The combination of the above inequality and (A.9), (A.11) yields

$$\|u_{\mathcal{A}} - u_{\mathcal{A},h}\|_{H^1(\Omega)} \lesssim \Big[\frac{h}{kp} + \frac{1}{p}\Big(\frac{kh}{\sigma p}\Big)^p\Big]C_{f,g}.$$

We introduce the sesquilinear form $a(\cdot, \cdot) := (\nabla u, \nabla v) + (u, v)$ and define the elliptic projection $\hat{u}_{\mathcal{A},h} \in V_h$ of $u_{\mathcal{A}}$ by

(A.12)
$$a(\hat{u}_{\mathcal{A},h},v_h) = a(u_{\mathcal{A}},v_h) \quad \forall v_h \in V_h.$$

Then,

$$\begin{aligned} \|u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}\|_{H^{1}(\Omega)}^{2} &= a(u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}, u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}) = a(u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}, u_{\mathcal{A}} - u_{\mathcal{A},h}) \\ &\lesssim \|u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}\|_{H^{1}(\Omega)} \|u_{\mathcal{A}} - u_{\mathcal{A},h}\|_{H^{1}(\Omega)}. \end{aligned}$$

Therefore,

(A.13)
$$\|u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}\|_{H^1(\Omega)} \lesssim \|u_{\mathcal{A}} - u_{\mathcal{A},h}\|_{H^1(\Omega)} \lesssim \left[\frac{h}{kp} + \frac{1}{p}\left(\frac{kh}{\sigma p}\right)^p\right] C_{f,g}$$

To show (A.6), we use the Nitsche's duality argument and consider the following auxiliary problem:

(A.14)
$$-\triangle w + w = u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h} \quad \text{in } \Omega,$$

(A.15)
$$\frac{\partial w}{\partial n} = 0$$
 on Γ .

It can be shown that w satisfies

(A.16)
$$\|w\|_{H^2(\Omega)} \lesssim \|u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}\|_{L^2(\Omega)}$$

Let $\hat{w}_h \in V_h$ be defined in Lemma A.2. Testing the conjugated (A.14) by $u_A - \hat{u}_{A,h}$ we get

$$\begin{aligned} \|u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}\|_{L^{2}(\Omega)} &= (u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}, -\bigtriangleup w + w) \\ &= (\nabla(u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}), \nabla w) + (u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}, w) \\ &= a(u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}), w) = a(u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}, w - \hat{w}_{h}) \\ &\lesssim \|u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}\|_{H^{1}(\Omega)} \|w - \hat{w}_{h}\|_{H^{1}(\Omega)} \\ &\lesssim \|u_{\mathcal{A}} - \hat{u}_{\mathcal{A},h}\|_{H^{1}(\Omega)} \frac{h}{n} \|w\|_{H^{2}(\Omega)}, \end{aligned}$$

which together with (A.16) and (A.13) gives (A.6). The proof is completed.

Next we prove Lemma 3.3. We consider to approximate $u_{\mathcal{E}}$ and $u_{\mathcal{A}}$, respectively. First, from Lemma A.2, there are two functions $\hat{u}_{\mathcal{E},h} \in V_h$ and $\check{u}_{\mathcal{E},h} \in W_h$ such that

(A.17)
$$\|u_{\mathcal{E}} - \hat{u}_{\mathcal{E},h}\|_{H^{j}(\Omega)} \lesssim \frac{h^{2-j}}{p^{2-j}} C_{f,g}, \quad j = 0, 1,$$

(A.18)
$$\|u_{\mathcal{E}} - \check{u}_{\mathcal{E},h}\|_{H^{j}(\mathcal{T}_{h})} \lesssim \frac{h^{2-j}}{p^{2-j}} C_{f,g}, \quad j = 0, 1, 2.$$

On the other hand, from Lemma A.3, there exist $\hat{u}_{\mathcal{A},h} \in V_h$ and $\check{u}_{\mathcal{A},h} \in W_h$ satisfying (A.6) and (A.5) respectively. Let $\hat{u}_h = \hat{u}_{\mathcal{E},h} + \hat{u}_{\mathcal{A},h}$, $\check{u}_h = \check{u}_{\mathcal{E},h} + \check{u}_{\mathcal{A},h}$. It follows from the triangle inequality that

(A.19)
$$||u - \hat{u}_h||_{H^j(\Omega)} \lesssim \left(\frac{h^{2-j}}{p^{2-j}} + \frac{h^{1-j}}{p^{2-j}} \left(\frac{kh}{\sigma p}\right)^p\right) C_{f,g}, \quad j = 0, 1,$$

(A.20)
$$\|u - \check{u}_h\|_{H^j(\mathcal{T}_h)} \lesssim \left(\frac{h^{2-j}}{p^{2-j}} + \frac{h^{1-j}}{p}\left(\frac{kh}{\sigma p}\right)^p\right) C_{f,g}, \quad j = 0, 1, 2.$$

Therefore, (3.6) holds.

(3.7) can be proved by following the same procedure as that for deriving (3.5). We omit the details. This completes the proof of Lemma 3.3.

Acknowledgments. The author would like to thank Professor J.M. Melenk for helpful communications.

References

- M AINSWORTH, Discrete dispersion relation for hp-version finite element approximation at high wave number, SIAM J. Numer. Anal., 42 (2004), pp. 553–575.
- [2] M. AINSWORTH AND H. A. WAJID, Dispersive and dissipative behavior of the spectral element method, SIAM J. Numer. Anal., 47 (2009), pp. 3910–3937.
- [3] D. ARNOLD, An interior penalty finite element method with discontinuous elements, SIAM J. Numer. Anal., 19 (1982), pp. 742–760.
- [4] A.K. AZIZ AND R.B. KELLOGG, A scattering problem for the Helmholtz equation, in Advances in Computer Methods for Partial Differential Equations-III, vol. 1, 1979, pp. 93–95.
- [5] I.M. BABUSKA AND S.A. SAUTER, Is the pollution effect of the FEM avoidable for the Helmholtz equation considering high wave numbers?, SIAM review, 42 (2000), pp. 451–484.
- [6] I. BABUŠKA, F. IHLENBURG, E.T. PAIK, AND S.A. SAUTER, A generalized finite element method for solving the Helmholtz equation in two dimensions with minimal pollution, Computer methods in applied mechanics and engineering, 128 (1995), pp. 325–359.
- [7] I. BABUŠKA AND M. SURI., The h-p version of the finite element method with quasiuniform meshes., Math. Modeling Numer. Anal., 21 (1987), pp. 199–238.
- [8] I. BABUŠKA AND M. ZLÁMAL, Nonconforming elements in the finite element method with penalty, SIAM Journal on Numerical Analysis, 10 (1973), pp. pp. 863–875.
- G.A. BAKER, Finite element methods for elliptic equations using nonconforming elements, Math. Comp., 31 (1977), pp. 44–59.
- [10] G. BAO AND H. WU, Convergence analysis of the pml problems for time-harmonic maxwell's equations, SIAM J. Numer. Anal., 43 (2005), pp. 2121–2143.
- J.-P. BERENGER, A perfectly matched layer for the absorption of electromagnetic waves, J. Comput. Phys., 114 (1994), pp. 185–200.

- [12] S.C. BRENNER AND L.R. SCOTT, The mathematical theory of finite element methods, Springer-Verlag, third ed., 2008.
- [13] E. BURMAN, A unified analysis for conforming and nonconforming stabilized finite element methods using interior penalty, SIAM journal on numerical analysis, 43 (2005), pp. 2012–2033.
- [14] E. BURMAN AND A. ERN, Stabilized Galerkin approximation of convection-diffusion-reaction equations: discrete maximum principle and convergence, Mathematics of computation, 74 (2005), p. 1637.
- [15] —, Continuous interior penalty hp-finite element methods for advection and advection-diffusion equations, Math. Comp., 259 (2007), pp. 1119–1140.
- [16] E. BURMAN, M.A. FERNÁNDEZ, AND P. HANSBO, Continuous interior penalty finite element method for Oseen's equations, SIAM journal on numerical analysis, 44 (2006), pp. 1248–1274.
- [17] E. BURMAN AND P. HANSBO, Edge stabilization for Galerkin approximations of convectiondiffusion-reaction problems, Computer methods in applied mechanics and engineering, 193 (2004), pp. 1437–1453.
- [18] Z. CHEN AND H. WU, An adaptive finite element method with perfectly matched absorbing layers for the wave scattering by periodic structures, SIAM J. Numer. Anal., 41 (2003), pp. 799–826.
- [19] P. G. CIARLET, The Finite Element Method for Elliptic Problems, North-Holland, Amsterdam, 1978.
- [20] P. CUMMINGS AND X. FENG, Sharp regularity coefficient estimates for complex-valued acoustic and elastic Helmholtz equations, Mathematical Models and Methods in Applied Sciences, 16 (2006), pp. 139–160.
- [21] A. DERAEMAEKER, I. BABUŠKA, AND P. BOUILLARD, Dispersion and pollution of the FEM solution for the Helmholtz equation in one, two and three dimensions, Int. J. Numer. Meth. Engng, 46 (1999), pp. 471–499.
- [22] J. DOUGLAS JR AND T. DUPONT, Interior Penalty Procedures for Elliptic and Parabolic Galerkin methods, Lecture Notes in Phys. 58, Springer-Verlag, Berlin, 1976.
- [23] J. DOUGLAS JR, JE SANTOS, AND D. SHEEN, Approximation of scalar waves in the space-frequency domain, Math. Mod. Meth. Appl. Sci, 4 (1994), pp. 509–531.
- [24] B. ENGQUIST AND A. MAJDA, Radiation boundary conditions for acoustic and elastic wave calculations, Communications on Pure and Applied Mathematics, 32 (1979), pp. 313–357.
- [25] B. ENGQUIST AND L. YING, Sweeping preconditioner for the helmholtz equation: moving perfectly matched layers, Multiscale Modeling and Simulation, 9 (2011), p. 686.
- [26] X. FENG AND H. WU, Discontinuous Galerkin methods for the Helmholtz equation with large wave numbers., SIAM J. Numer. Anal., 47 (2009), pp. 2872-2896, also downloadable at http://arXiv.org/abs/0810.1475.
- [27] —, hp-discontinuous Galerkin methods for the Helmholtz equation with large wave number, Math. Comp., (2011, posted online).
- [28] X. FENG AND Y. XING, Absolutely stable local discontinuous Galerkin methods for the Helmholtz equation with large wave number, Arxiv preprint arXiv:1010.4563, (2010).
- [29] B. GUO, Approximation theory for the p-version of the finite element method in three dimensions, part 1: approximabilities of singular functions in the framework of the jacobi-weighted Besov and Sobolev spaces., SIAM J. Numer. Anal., 44 (2006), pp. 246–269.
- [30] B. GUO AND W. SUN, The optimal convergence of the h-p version of the finite element method with quasi-uniform meshes., SIAM J. Numer. Anal., 45 (2007), pp. 698–730.

- [31] I. HARARI, Reducing spurious dispersion, anisotropy and reflection in finite element analysis of timeharmonic acoustics, Computer methods in applied mechanics and engineering, 140 (1997), pp. 39–58.
- [32] U. HETMANIUK, Stability estimates for a class of Helmholtz problems, Commun. Math. Sci, 5 (2007), pp. 665–678.
- [33] R.H.W HOPPE AND B. WOHLMUTH, Element-oriented and edge-oriented local error estimators for nonconforming finite element methods, Math. Model. Numer. Anal, 30 (1996), pp. 237–263.
- [34] F. IHLENBURG AND I. BABUŠKA, Finite element solution of the Helmholtz equation with high wave number. I. The h-version of the FEM, Comput. Math. Appl., 30 (1995), pp. 9–37.
- [35] —, Finite element solution of the Helmholtz equation with high wave number. II. The h-p version of the FEM, SIAM J. Numer. Anal., 34 (1997), pp. 315–358.
- [36] O.A. KARAKASHIAN AND F. PASCAL, A posteriori error estimates for a discontinuous galerkin approximation of second-order elliptic problems, SIAM Journal on Numerical Analysis, (2004), pp. 2374–2399.
- [37] J. KIM AND D. SHEEN, A priori estimates for elliptic boundary value problems with nonlinear boundary conditions, to appear, (1995).
- [38] J.M. MELENK, On Generalized Finite Element Methods, PhD thesis, University of Maryland at College Park, 1995.
- [39] J. M. MELENK AND S. SAUTER, Convergence analysis for finite element discretizations of the Helmholtz equation with Dirichlet-to-Neumann boundary conditions, Math. Comp., 79 (2010), pp. 1871–1914.
- [40] —, Wavenumber explicit convergence analysis for Galerkin discretizations of the Helmholtz equation, SIAM J. Numer. Anal., 49 (2011), pp. 1210–1243.
- [41] P. OSWALD, On a BPX-preconditioner for p1 elements, Computing, 51 (1993), pp. 125–133.
- [42] A.H. SCHATZ, An observation concerning Ritz-Galerkin methods with indefinite bilinear forms, Math. Comp., 28 (1974), pp. 959–962.
- [43] C. SCHWAB, P- and hp-Finite Element Methods, Oxford University Press, 1998.
- [44] J. SHEN AND L.L. WANG, Analysis of a spectral-Galerkin approximation to the Helmholtz equation in exterior domains, SIAM J. Numer. Anal, 45 (2007), pp. 1954–1978.
- [45] L.L. THOMPSON, A review of finite-element methods for time-harmonic acoustics, J. Acoust. Soc. Am., 119 (2006), pp. 1315–1330.
- [46] L.L. THOMPSON AND P.M. PINSKY, Complex wavenumber fourier analysis of the p-version finite element method, Computational Mechanics, 13 (1994), pp. 255–275.
- [47] H. WU, Pre-asymptotic error analysis of CIP-FEM and FEM for Helmholtz equation with high wave number. Part I: Linear version, to appear.