# SUM OF SQUARES CERTIFICATES FOR CONTAINMENT OF $\mathcal{H}$ -POLYTOPES IN $\mathcal{V}$ -POLYTOPES

KAI KELLNER AND THORSTEN THEOBALD

ABSTRACT. Given an  $\mathcal{H}$ -polytope P and a  $\mathcal{V}$ -polytope Q, the decision problem whether P is contained in Q is co-NP-complete. This hardness remains if P is restricted to be a standard cube and Q is restricted to be the affine image of a cross polytope. While this hardness classification by Freund and Orlin dates back to 1985, for general dimension there seems to be only limited progress on that problem so far.

Based on a formulation of the problem in terms of a bilinear feasibility problem, we study sum of squares certificates to decide the containment problem. These certificates can be computed by a semidefinite hierarchy. As a main result, we show that under mild and explicitly known preconditions the semidefinite hierarchy converges in finitely many steps. In particular, if P is contained in a large  $\mathcal{V}$ -polytope Q (in a well-defined sense), then containment is certified by the first step of the hierarchy.

#### 1. INTRODUCTION

Convex polytopes (polytopes, for short) can be represented as the convex hull of finitely many points (" $\mathcal{V}$ -polytopes") or as the intersection of finitely many halfspaces (" $\mathcal{H}$ -polytopes"). For  $a \in \mathbb{R}^k$ ,  $A \in \mathbb{R}^{k \times d}$ , and  $B = [b^{(1)}, \ldots, b^{(l)}] \in \mathbb{R}^{d \times l}$  let

$$P = P_A = \{x \in \mathbb{R}^d \mid a - Ax \ge 0\}$$
 and  $Q = Q_B = \text{conv}(B) = \text{conv}(b^{(1)}, \dots, b^{(l)})$ 

be an  $\mathcal{H}$ -polytope and a  $\mathcal{V}$ -polytope, respectively. The subscript in the notion of P and Q indicates the dependency on the specific representation of the polytopes involved. However, if there is no risk of confusion, we often state P and Q without subscript. The following two problems are prominent problems in algorithmic polytope theory (see Kaibel and Pfetsch [15]). We always assume that the polytope data is given in terms of rational numbers.

POLYTOPE VERIFICATION: **Input:**  $d \in \mathbb{N}$ , an  $\mathcal{H}$ -polytope  $P \subseteq \mathbb{R}^d$  and a  $\mathcal{V}$ -polytope  $Q \subseteq \mathbb{R}^d$ . **Task:** Decide whether P = Q.

POLYTOPE CONTAINMENT (or  $\mathcal{H}$ -IN- $\mathcal{V}$  CONTAINMENT): **Input:**  $d \in \mathbb{N}$ , an  $\mathcal{H}$ -polytope  $P \subseteq \mathbb{R}^d$  and a  $\mathcal{V}$ -polytope  $Q \subseteq \mathbb{R}^d$ . **Task:** Decide whether  $P \subseteq Q$ .

While the complexity status of the first problem is open, the second problem is co-NPcomplete (see Freund and Orlin [7]); note that it is trivial to decide the converse question  $Q \subseteq P$ . It is well known that the problem of enumerating all facets of a polytope given by

Date: August 23, 2021.

a finite set of points (or, equivalently, enumerating all vertices of a polytope given by a finite number of halfspaces) can be polynomially reduced to the POLYTOPE VERIFICATION problem (see Avis et. al. [2], Kaibel and Pfetsch [15]). Note that enumerating the vertices of an (unbounded) polyhedron is hard [18]. While in fixed dimension, enumeration of the vertices of P can be done in polynomial time and gives a polynomial time algorithm for POLYTOPE CONTAINMENT (cf. Theorem 2.2), progress on approximation results for the latter problem in general dimension seems to be limited so far.

In this paper we study the POLYTOPE CONTAINMENT problem. Our main focus is to consider the problem from the viewpoint of the transition from linear/polyhedral problems to low-degree semialgebraic problems. To that end, we formulate the problem as a disjointly constrained bilinear feasibility problem and consider semialgebraic certificates. In particular, the reformulation as a bilinear program allows to effectively apply Putinar's Positivstellensatz [30], which is the primary theorem underlying Lasserre's hierarchy [21] and gives sum of squares certificates; see also Laurent's extensive survey [23]. The sum of squares can be computed by a hierarchy of semidefinite programs.

The basic idea of the approach – which is meanwhile common in polynomial optimization, but whose understanding of the particular potential for concrete problems is often challenging – can be explained as follows. One point of view towards linear programming is as an application of Farkas' Lemma which characterizes the (non-)solvability of a system of linear inequalities. The affine form of Farkas' Lemma [32, Corollary 7.1h] characterizes linear polynomials which are nonnegative on a given polyhedron. By omitting the linearity condition, one gets a polynomial nonnegativity question, leading to so called Positivstellensätze (or, more precisely, Nichtnegativstellensätze). These Positivstellensätze provide a *certificate* for the positivity of a polynomial function in terms of a polynomial identity. As in the linear case, the Positivstellensätze are the foundation of polynomial optimization and relaxation methods (see [21, 22, 23]).

The general machinery from polynomial optimization automatically implies convergence results, but often these results come with restrictions or technical assumptions.

#### Our contributions:

1. Based on a formulation of the POLYTOPE CONTAINMENT problem in terms of a bilinear problem (Proposition 3.1), we characterize geometric properties of a natural bilinear programming reformulation; see Corollary 3.3.

2. We study the application of sums of squares techniques on the bilinear programming formulation. An important point is whether the hierarchy always converges in finitely many steps. While in the case of strong containment (as defined in Section 3) this property is implied by Putinar's Positivstellensatz (Theorem 4.2), in the case of non-strong containment this is a critical issue. As a main result of this paper, we show that under mild and explicitly known conditions, the semidefinite hierarchy converges in finitely many steps (Theorem 4.3) based on results by Marshall ([24, 25]; see also Nie [28]).

3. We exhibit structural differences between conventional methods for the POLYTOPE CONTAINMENT (such as vertex tracking methods) and our approach. Theorem 4.10 shows that the containment of polytopes in "large" polytopes (as quantified in the theorem) can already be certified in the initial step of the semidefinite hierarchy and thus by computing a semidefinite program of polynomial size in the input.

While it is a fundamental geometric problem by itself, we mention some exemplary application scenarios in which the POLYTOPE CONTAINMENT problem occurs. Generally, many applications in data analysis or shape analysis of point clouds involve the convex hull of point sets (see, e.g., [4]), and a POLYTOPE CONTAINMENT problem can be used to answer questions about certain (polyhedral) properties on the set.

A specific example is theorem proving in linear real arithmetic: A sub-branch in theorem proving is based on formulas in linear real arithmetic; see, e.g., [5, 27]. Given a set of (quantifier-free) linear inequalities of the form  $L_i(x_1, \ldots, x_d) \ge 0$  in the real variables  $x_1, \ldots, x_d$  specifying the assumptions of a certain theorem, one may ask whether all these solutions satisfy a certain property Q. If Q is described as the convex hull of a finite number of points, then the theorem proving problem corresponds to a POLYTOPE CONTAINMENT problem.

Let us briefly mention some related problems. Finding the largest simplex in a  $\mathcal{V}$ -polytope is an NP-hard problem [12]. However for that problem Packer has given a polynomial-time approximation [29]. Recently, Gouveia et. al. have studied the question which nonnegative matrices are slack matrices [9], and they establish equivalence of the decision problem to the polyhedral verification problem. For containment of polytopes and spectrahedra see [16, 17]. Joswig and Ziegler [14] showed that the POLYTOPE VERIFICA-TION problem is polynomially equivalent to a geometric polytope completeness problem.

The paper is structured as follows. After introducing the relevant notation in Section 2, we study geometric properties of a natural bilinear programming formulation in Section 3. Section 4 deals with sum of squares certificates for the POLYTOPE CONTAINMENT problem. Finally, Section 5 lists several open questions.

#### 2. Preliminaries

Recall that a polyhedron P is the intersection of finitely many affine halfspaces in  $\mathbb{R}^d$ and a bounded polyhedron is called a polytope [33]. Denote by V(P) the set of vertices of a polytope P, and by F(P) the set of facets. By McMullen's Upper bound Theorem [26], any *d*-polytope with k vertices (resp. facets) has at most

$$\binom{k - \lfloor \frac{1}{2}(d+1) \rfloor}{k - d} + \binom{k - \lfloor \frac{1}{2}(d+2) \rfloor}{k - d}$$

facets (resp. vertices). This bound is sharp for neighborly polytopes such as cyclic polytopes.

Our model of computation is the binary Turing machine: polytopes are given in terms of rational numbers, and the size of the input is defined as the length of the binary encoding of the input data (see, e.g., [10]). It is well-known that the complexity of deciding containment of one polytope in another one strongly depends on the type of input representations. In particular, the following hardness statement is known.

**Proposition 2.1** ([7, 10]). The POLYTOPE CONTAINMENT problem is co-NP-complete.

This hardness remains if P is restricted to be a standard cube and Q is restricted to be the affine image of a cross polytope.

If the dimension is fixed, then the problem of deciding whether an  $\mathcal{H}$ -polytope is contained in a  $\mathcal{V}$ -polytope can be decided in polynomial time.

Note that the result for fixed dimension can be slightly strengthened.

**Corollary 2.2.** If the dimension of P or the dimension of Q is fixed, then containment of an  $\mathcal{H}$ -polytope P in a  $\mathcal{V}$ -polytope Q can be decided in polynomial time.

*Proof.* If the dimension of P is fixed, then first compute the affine hull of P. This can be done in polynomial time. Taking that affine hull as ambient space in fixed dimension, P can be transformed into a  $\mathcal{V}$ -representation in polynomial time. It remains to decide containment of a  $\mathcal{V}$ -polytope in a  $\mathcal{V}$ -polytope, which can be done in polynomial time.

Similarly, if the dimension of Q is fixed, an  $\mathcal{H}$ -representation of Q can be computed in polynomial time, and the resulting problem of deciding whether an  $\mathcal{H}$ -polytope is contained in an  $\mathcal{H}$ -polytope can be decided in polynomial time.

Throughout the paper, we assume boundedness and nonemptyness of P as well as  $0 \in \operatorname{int} Q$ , where  $\operatorname{int} Q$  denotes the interior of Q. All these properties can be tested in polynomial time [19]. If Q is full-dimensional and  $0 \notin \operatorname{int} Q$ , one can translate Q and P by the centroid of the vertices of Q. Recall that the polar polyhedron of Q is

$$Q^{\circ} = \left\{ z \in \mathbb{R}^d \mid \mathbb{1}_l - B^T z \ge 0 \right\},\$$

where  $\mathbb{1}_l$  denotes the all-1-vector in  $\mathbb{R}^l$ . Since we assume  $0 \in \operatorname{int} Q$ ,  $Q^\circ$  is a polytope (i.e., bounded) and  $Q^{\circ\circ} = Q$ .

# 3. A BILINEAR APPROACH TO THE POLYTOPE CONTAINMENT PROBLEM

We first collect some geometric properties of the POLYTOPE CONTAINMENT problem. Our starting point is the following reformulation of the POLYTOPE CONTAINMENT problem as a bilinear problem.

**Proposition 3.1.** Let the  $\mathcal{H}$ -polytope  $P = \{x \in \mathbb{R}^d \mid a - Ax \ge 0\}$  be nonempty and the  $\mathcal{V}$ -polytope  $Q = \operatorname{conv}(B) = \operatorname{conv}(b^{(1)}, \ldots, b^{(l)})$  containing the origin in its interior.

(1) P is contained in Q if and only if  $x^T z \leq 1$  for all  $(x, z) \in P \times Q^\circ$ . That is,  $P \subseteq Q$  if and only if the maximum

(3.1) 
$$\mu^* := \max\{x^T z \mid (x, z) \in P \times Q^\circ\}$$

is at most 1.

(2) We have  $\mu^* = 1$  if and only if  $P \subseteq Q$  and  $\partial P \cap \partial Q \neq \emptyset$ .

Motivated by the second statement, we say that P is strongly contained in Q if  $P \subseteq Q$ and  $\partial P \cap \partial Q = \emptyset$ . Since Q is full-dimensional, this is equivalent to  $P \subseteq$  int Q. Note that strong containment differs from (set-theoretic) strict inclusion as the latter allows common boundary points while strong containment does not. *Proof.* To (1): If  $P \subseteq Q$  then for any  $x \in P$  we have  $x^T z \leq 1$  for all  $z \in Q^\circ$ . Conversely, if  $x^T z \leq 1$  holds for all  $z \in Q^\circ$ , then for any  $x \in P$  we have  $x \in Q^{\circ\circ} = Q$ .

To (2): Let  $P \subseteq Q$  and  $\partial P \cap \partial Q$  be nonempty. Then there exists a vertex  $v \in V(P)$  and a facet  $F \in F(Q)$  such that  $v \in F$ . Since  $0 \in \text{int } Q$ , F defines a vertex f of the polar  $Q^{\circ}$ . Further  $f^T v = 1$  implies that the maximum is at least one. By part (1) of the statement, the maximum must be exactly one.

Conversely, if the maximum is one, then  $x^T z \leq 1$  for all  $(x, z) \in P \times Q^\circ$ . Therefore, since the set  $P \times Q^\circ$  is compact, there exists a point  $(\bar{x}, \bar{z}) \in P \times Q^\circ$  such that  $\bar{x}^T \bar{z} = 1$ . Hence  $\bar{x}^T z \leq 1$  for all  $z \in Q^\circ$  and  $\bar{x}^T \bar{z} = 1$ , i.e.,  $\bar{x}$  defines a supporting hyperplane of  $Q^\circ$ . Thus  $\bar{x}$  is a boundary point of Q. Similarly,  $x^T \bar{z} \leq 1$  for all  $x \in P$  and  $\bar{x}^T \bar{z} = 1$ , implying  $\bar{x} \in \partial P$ . Consequently,  $\bar{x} \in \partial Q \cap \partial P$ .

The following characterization of the optimal solutions to (3.1) is a slight extension of a result by Konno [20] on bilinear programming.

**Proposition 3.2.** Let int  $P \neq \emptyset$  and  $0 \in \text{int } Q$ . Then the set of optimal solutions to (3.1) is a set of proper faces  $F \times G$  of  $P \times Q^{\circ}$ , and the maximum is attained at a pair of vertices of P and  $Q^{\circ}$ .

For the convenience of the reader, we recall the short proof.

Proof. Let  $(\bar{x}, \bar{z}) \in P \times Q^{\circ}$  be an optimal solution. Then the set  $G = \{z \in Q^{\circ} | \bar{x}^T z = \bar{x}^T \bar{z}\}$  is a non-empty face of  $Q^{\circ}$ . For all  $\hat{z} \in G$ , let F be the set of maximizers of  $\max\{x^T \hat{z} | x \in P\}$ . Consequently, for  $(\hat{x}, \hat{z}) \in F \times G$  we have  $\hat{x}^T \hat{z} = \bar{x}^T \hat{z}$  and, by the optimality of  $(\bar{x}, \bar{z}), F \times G$  is contained in the set of optimal solutions. To complete the proof of the first part of the statement, note that every boundary point of a polytope has a unique minimal face containing it.

Since the set of optimal solutions is a set of proper faces of  $P \times Q^{\circ}$ , there exists an optimal pair of vertices of P and  $Q^{\circ}$ .

There is a nice geometric interpretation of the latter proposition. Since, in the case  $0 \in \operatorname{int} Q$ , each vertex of  $Q^{\circ}$  corresponds to a facet of Q and vice versa, an optimal solution  $(x, z) \in V(P) \times V(Q^{\circ})$  of (3.1) yields a pair of a vertex of P and a facet defining normal vector of Q. However, since computing the set of vertices  $V(Q^{\circ})$  is an NP-hard problem, it is not reasonable to reduce the problem to the set of vertices in general.

The optimal value of problem (3.1) might be attained by other boundary points than vertices and, moreover, there might be infinitely many optimal solutions. From a geometric point of view, this only occurs in somewhat degenerate cases.

**Corollary 3.3.** Let int  $P \neq \emptyset$  and  $0 \in \text{int } Q$ . Problem (3.1) has finitely many optimal solutions if and only if every optimal solution of Problem (3.1) is a pair of vertices of P and  $Q^{\circ}$ .

As there always exists a pair of vertices that is an optimal solution to (3.1), it is natural to ask for vertex tracking algorithms. This is the approach in, e.g., [8, 20]. So far, no converging algorithm is known based on this approach. Note that the formulation as a bilinear programming problem from Proposition 3.1 also allows to apply existing nonlinear programming techniques for non-convex quadratic optimization (see, e.g., [6]). In the next section, we study the bilinear reformulation of the POLYTOPE CONTAINMENT from the viewpoint of algebraic certificates and semidefinite relaxations yielding a generically convergent algorithm.

# 4. Sum of squares certificates

In this section, we study sum of squares techniques for the POLYTOPE CONTAINMENT problem. Our main goal is to show that in the situation of Corollary 3.3, the corresponding semidefinite hierarchy yields a certificate for containment after finitely many steps; see Theorems 4.2 and 4.3.

4.1. Putinar's Positivstellensatz. Consider a set of polynomials  $G = \{g_1, \ldots, g_k\} \subseteq \mathbb{R}[x]$  in the variables  $x = (x_1, \ldots, x_d)$ . The quadratic module generated by G is defined as

$$QM(G) = \left\{ \sigma_0 + \sum_{i=1}^k \sigma_i g_i \mid \sigma_i \in \Sigma[x] \right\},\$$

where  $\Sigma[x] \subseteq \mathbb{R}[x]$  is the set of sum of squares polynomials. Here, a polynomial  $p \in \mathbb{R}[x]$  is called *sum of squares* (sos) if it can be written in the form  $p = \sum_i h_i(x)^2$  for some  $h_i \in \mathbb{R}[x]$ . Equivalently, p has the form  $[x]^T Q[x]$ , where [x] is the vector of all monomials in x up to half the degree of p and Q is a positive semidefinite matrix of appropriate size. Checking whether a polynomial is sos is a semidefinite feasibility problem.

Obviously, every element in QM(G) is nonnegative on the semialgebraic set  $S = \{x \in \mathbb{R}^d \mid g(x) \ge 0 \ \forall g \in G\}$ . In [30] Putinar showed that for positive polynomials the converse is true under some regularity assumption.

A quadratic module QM(G) is called Archimedean if there is a polynomial  $p \in QM(G)$ such that the level set  $\{x \in \mathbb{R}^d \mid p(x) \ge 0\}$  is compact, or, equivalently, the polynomial  $N - (x_1^2 + \dots + x_d^2) \in QM(G)$  for some positive integer N; see Marshall's book [24] for more equivalent characterizations.

**Proposition 4.1** (Putinar's Positivstellensatz [30]. See also [24, Theorem 5.6.1]). Let  $S = \{x \in \mathbb{R}^d \mid g(x) \ge 0 \ \forall g \in G\}$  for some finite subset  $G \subseteq \mathbb{R}[x]$ . If the quadratic module QM(G) is Archimedean, then QM(G) contains every polynomial  $f \in \mathbb{R}[x]$  positive on S.

The Archimedean condition in the proposition is not very restrictive. Especially, in our case of interest where all polynomials  $g_i$  are linear and S is compact, the condition is always fulfilled; see [24, Theorem 7.1.3].

In order to apply Putinar's Positivstellensatz to polynomial optimization, consider an optimization problem

(4.1) 
$$\sup \{f(x) \mid g_i(x) \ge 0, \ i = 1, \dots, k\}$$

with  $f, g_1, \ldots, g_k \in \mathbb{R}[x]$ . Clearly, this is the same as to find the infimum of  $\mu$  such that  $\mu - f(x) \geq 0$  on the set S. A common way to tackle the latter problem is to replace the nonnegativity condition by an sos condition. This is a semi-infinite program since deciding membership can be rephrased as a semi-infinite feasibility problem. In order to

get a (finite-dimensional) semidefinite program, we truncate the quadratic module QM(G) by considering only monomials up to a certain degree 2t,

$$QM_t(G) = \left\{ \sigma_0 + \sum_{i=1}^k \sigma_i g_i \mid \sigma_i \in \Sigma[x] \text{ with } \deg(\sigma_0) \le 2t \text{ and } \deg(\sigma_i g_i) \le 2t \right\}.$$

The *t*-th sos program has the form

(4.2) 
$$\mu(t) = \inf \{ \mu \mid \mu - f(x) \in QM_t(G) \}.$$

Clearly, the sequence of truncated quadratic modules is increasing with respect to inclusion as t grows. Thus the sequence of optimal values  $\mu(t)$  is monotone decreasing and bounded from below by the optimal value of (4.1). Generally, the infimum is not attained in (4.2).

The dual problem to (4.2) can be formulated in terms of moment matrices, again leading to an SDP relaxation of the polynomial optimization problem (4.1). From a computational point of view it is often easier (i.e., faster) to compute the dual side. This is because of the time consuming process of extracting coefficients in a formal sos representation. It is known that there is no duality gap between the primal and dual problem, whenever the quadratic module is Archimedean and S contains an interior point [22, Theorem 5.21]. Since we do not use the dual side here, we refer interested readers to Lasserre's fundamental work [21].

4.2. Sum of squares certificates for POLYTOPE CONTAINMENT. To keep notation simple, we denote the (truncated) quadratic module generated by the linear constraints a - Ax and  $1 - B^T z$  by  $QM_t(A, B)$ . The sos formulation of problem (3.1) reads as

(4.3) 
$$\mu(t) = \inf \left\{ \mu \mid \mu - x^T z \in \mathrm{QM}_t(A, B) \right\}.$$

Denote the *i*-th constraint defining  $P \times Q^{\circ}$  by  $g_i$ . Let  $\mu - x^T z = \sigma_0 + \sum_{i=1}^{k+l} \sigma_i g_i$  be an sos representation. Assume t = 1. Then monomials of degree at most 2 appear, i.e.,  $\deg(\sigma_0) \in \{0, 2\}$  and  $\deg(\sigma_i g_i) \leq 2$ . Since  $\deg(g_i) = 1$  and  $\sigma_i$  is sos,  $\sigma_i$  must be constant (otherwise monomials of degree greater than 2 appear). Thus  $\deg(\sum_i \sigma_i g_i) \leq 1$ . Moreover, if  $\deg(\sigma_0) = 2$ , then purely quadratic terms like  $x_j^2$  or  $z_j^2$  appear for some j on the right-hand side while the coefficients of these terms are zero on the left-hand side. As a consequence, the first order of the hierarchy making sense is t = 2. We call t = 2 the *initial step of the hierarchy*.

Asymptotic convergence of the hierarchy in the general case and finite convergence in the strong containment case follow easily from the general theory.

**Theorem 4.2.** Let P be a nonempty  $\mathcal{H}$ -polytope and Q be a  $\mathcal{V}$ -polytope with  $0 \in \operatorname{int} Q$ .

- (1) If  $\mu(t) \leq 1$  for some integer  $t \geq 2$ , then  $P \subseteq Q$ .
- (2) The hierarchy (4.3) converges asymptotically from above to the optimal value  $\mu^*$  of problem (3.1).
- (3) If P is strongly contained in Q, then the hierarchy (4.3) decides the POLYTOPE CONTAINMENT problem in finitely many steps.

*Proof.* The first statement is clear by construction of the hierarchy.

Consider the second statement. Since all constraints are linear in x, z and the feasible region is bounded, the quadratic module generated by the constraints of problem (3.1) is Archimedean [24, Theorem 7.1.3] and thus contains all polynomials  $f(x, z) \in \mathbb{R}[x, z]$ positive on  $P \times Q^{\circ}$  by Putinar's Positivstellensatz 4.1. Let  $\mu^*$  be the optimal value of problem (3.1). Then  $\mu^* - x^T z \ge 0$  on  $P \times Q^{\circ}$  and hence  $\mu^* + \epsilon - x^T z \in \text{QM}(A, B)$  for all  $\epsilon > 0$ .

If P is strongly contained in Q, then  $\mu^* < 1$  and thus  $1 - x^T z \in QM(A, B)$ .

A priori it is not clear whether in the non-strong case finite convergence holds. In fact, for general polynomials, there are examples where finite convergence is not possible. As our main result, we provide a partial extension of Theorem 4.2 to the case where the bilinear optimization problem (3.1) has only finitely many optimal solutions (as characterized in Corollary 3.3).

**Theorem 4.3.** Let  $P = \{x \in \mathbb{R}^d \mid a - Ax \ge 0\}$  be a full-dimensional  $\mathcal{H}$ -polytope and let  $Q = \operatorname{conv}(B)$  be a  $\mathcal{V}$ -polytope containing the origin in its interior. Assume that one of the equivalent statements in Corollary 3.3 holds. Then  $\mu^* - x^T z \in \operatorname{QM}(A, B)$ , and thus the hierarchy (4.3) decides the POLYTOPE CONTAINMENT problem in finitely many steps.

To prepare for the proof, we show that in the semidefinite hierarchy for the POLYTOPE CONTAINMENT problem, the sos formulation is invariant under redundant constraints, i.e., redundant inequalities in the  $\mathcal{H}$ -representation of P or redundant points in the  $\mathcal{V}$ representation of Q. Note that for a general semialgebraic constraint set this is not always true, even in the case of optimizing a linear function over it; see [13, Section 5.2] for a well-known example (cf. also [1]). Recall that every  $\mathcal{H}$ -representation of a certain polytope contains the facet defining halfspaces. Similarly, the vertices are part of each  $\mathcal{V}$ -representation.

**Lemma 4.4** (Redundant constraints). Let  $P_A = \{x \in \mathbb{R}^d \mid a - Ax \ge 0\}$  and  $Q_B = \operatorname{conv}(B)$  be nonempty polytopes with  $a \in \mathbb{R}^{k+1}$ ,  $A \in \mathbb{R}^{(k+1) \times d}$ , and  $B \in \mathbb{R}^{d \times (l+1)}$ .

- (1) If  $(a Ax)_{k+1} \ge 0$  is a redundant inequality in the  $\mathcal{H}$ -representation of  $P_A$ , then it is also redundant in the sos representation (4.3), i.e., the inclusion  $P_A \subseteq Q_B$  is certified by a certain step of the hierarchy if and only if  $P_{A\setminus A_{k+1}} \subseteq Q_B$  is certified by the same step.
- (2) If  $b^{(l+1)}$  is a redundant point in the  $\mathcal{V}$ -representation of  $Q_B$ , then it is also redundant in the sos representation (4.3), i.e.,  $P_A \subseteq Q_B$  is certified by a certain step of the hierarchy if and only if  $P_A \subseteq Q_{B\setminus b^{(l+1)}}$  is certified by the same step.

*Proof.* We only prove statement (1), the proof of part (2) is analog. Consider an sos representation of  $\mu(t) - x^T z$  for some  $t \ge 2$ ,

$$\mu(t) - x^T z = \sigma_0 + \sum_{i=1}^{k+1} \sigma_i \left(a - Ax\right)_i + \sum_{i=1}^l \sigma_{k+1+i} \left(1 - B^T z\right)_i \in \text{QM}(A, B)$$

where  $\sigma_0, \ldots, \sigma_{k+l+1} \in \Sigma[x, z]$  are sos polynomials with deg  $\sigma_0 \leq 2t$  and deg  $\sigma_i \leq 2t - 2$ for  $i \in \{1, \ldots, k+l+1\}$ . Since  $(a - Ax)_{k+1}$  is redundant in the description of  $P_A$ , we can

8

write it as a conic combination of the remaining linear polynomials,

$$(a - Ax)_{k+1} = \lambda_0 + \lambda^T (a - Ax), \ \lambda \in \mathbb{R}^k_+, \ \lambda_0 \in \mathbb{R}_+.$$

Replacing  $\sigma_{k+1}(a - Ax)_{k+1}$  in the sos representation yields

$$\mu(t) - x^T z = \sigma'_0 + \sum_{i=1}^k \sigma'_i (a - Ax)_i + \sum_{i=1}^l \sigma_{k+1+i} \left( 1 - B^T z \right)_i \in \text{QM}(A \setminus A_{k+1}, B),$$

where  $\sigma'_i = \lambda_i \sigma_{k+1} + \sigma_i \in \Sigma[x, z]$  with degree  $\deg(\sigma'_i) = \max\{\deg(\lambda_i \sigma_{k+1}), \deg(\sigma_i)\} \le 2t - 2$ for  $i \in \{0, \dots, k\}$ .

To prove Theorem 4.3, we introduce a sufficient convergence condition by Marshall (see [24, 25]) which is based on a boundary Hessian condition.

Given  $g_1, \ldots, g_k \in \mathbb{R}[x]$  and a boundary point  $\bar{x}$  of  $S = \{x \in \mathbb{R}^d \mid g_i(x) \ge 0, i = 1, \ldots, k\}$ . We assume that (say, by an application of the inverse function theorem) there exists a local parameterization for  $\bar{x}$  in the following sense: There exist open sets  $U, V \subseteq \mathbb{R}^d$  such that  $\bar{x} \in U, \phi : U \to V, x \mapsto t := (t_1, \ldots, t_d)$  is bijective, the inverse  $\phi^{-1} : V \to U$  is a continuously differentiable function on V, and for some  $r \in \{1, \ldots, d\}$  let  $t_1 = g_1, \ldots, t_r = g_r$  on U.

**Condition 4.5** (Boundary Hessian condition, BHC). Given a polynomial  $f \in \mathbb{R}[x]$ , denote by  $f_1$  and  $f_2$  the linear and quadratic part of f in the localizing parameters  $t_1, \ldots, t_d$ , respectively. Let  $R = \{(t_1, \ldots, t_d) \in \mathbb{R}^d | t_1 \ge 0, \ldots, t_r \ge 0\}$ . If the linear form  $f_1 = c_1t_1 + \cdots + c_rt_r$  has only negative coefficients and the quadratic form  $f_2(0, \ldots, 0, t_{r+1}, \ldots, t_d)$  is negative definite, then the restriction  $f_{|R}$  has a local maximum at  $\bar{x}$ .

Using this condition, the following generalization of Putinar's Theorem can be stated.

**Proposition 4.6** ([24, Theorem 9.5.3], see also [31, Theorem 3.1.7]). Let  $f, g_1, \ldots, g_k \in \mathbb{R}[x]$ , and suppose that the quadratic module QM(G) generated by  $G = \{g_1, \ldots, g_k\}$  is Archimedean. Further assume that for each global maximizer  $\bar{x}$  of f over  $S = \{x \in \mathbb{R}^d \mid g(x) \ge 0 \ \forall g \in G\}$  there exists an index set  $I \subseteq \{1, \ldots, d\}$  such that (after renaming the variables w.r.t. the indices in I and w.r.t. the indices not in I) f satisfies BHC at  $\bar{x}$ . Denote by  $f_{\max}$  the global maximum of f on S. In this situation,  $f_{\max} - f \in QM(G)$ .

Our goal is to show that under the assumptions of Theorem 4.3 the boundary Hessian condition holds. We will use the following version of the Karush-Kuhn-Tucker conditions adapted to the bilinear situation.

**Lemma 4.7** ([3, Section 5.1]). Let  $f(x, z) \in \mathbb{R}[x, z]$  be a continuously differentiable function and let  $\mathbb{P} := P_A \times P_B = \{(x, z) \in \mathbb{R}^{2d} \mid a - Ax \ge 0, b - Bz \ge 0\}$  be the product of two nonempty polytopes. If f attains a local maximum at  $(\bar{x}, \bar{z})$  on  $\mathbb{P}$ , then there exists  $(\alpha, \beta)$  such that

(4.4) 
$$\nabla f(\bar{x}, \bar{z}) = \begin{bmatrix} A^T & 0\\ 0 & B^T \end{bmatrix} \begin{pmatrix} \alpha\\ \beta \end{pmatrix}$$
$$0 = \alpha_i (a - A\bar{x})_i = \beta_j (b - B\bar{z})_j, \quad i = 1, \dots, k, \ j = 1, \dots, l$$
$$\alpha \ge 0, \ \beta \ge 0.$$

In the lemma, only multipliers corresponding to active constraints can be positive, since otherwise one of the equations (4.4) is violated.

We are now able to prove Theorem 4.3. In a more general setting, Nie used the Karush-Kuhn-Tucker optimality conditions to certify the BHC; see [28]. Because of the special structure of problem (3.1), we do not need the whole machinery used by Nie. In particular, the local parameterization needed for the BHC (see the paragraph before Condition 4.5) comes from an affine variable transformation. As a consequence, for POLYTOPE CON-TAINMENT, our direct approach allows to prove a stronger result than we would obtain just by applying Nie's Theorem. Specifically, we obtain a geometric characterization of the degenerate situations as given in Theorem 4.3.

Proof (of Theorem 4.3). Let  $(\bar{x}, \bar{z}) \in P \times Q^{\circ}$  be an arbitrary but fixed optimal solution. By Lemma 4.7 there exists  $(\alpha, \beta) \in \mathbb{R}^{k+l}$  such that

(4.5) 
$$\begin{aligned} (\bar{z},\bar{x}) &= (A^T \alpha, B\beta) \\ 0 &= \alpha_i (a - A\bar{x})_i = \beta_j (\mathbb{1} - B^T \bar{z})_j, \quad i = 1, \dots, k, \ j = 1, \dots, l \\ \alpha \geq 0, \ \beta \geq 0. \end{aligned}$$

As mentioned before, only multipliers corresponding to active constraints can be positive. Denote by I the collection of index sets of active constraints in  $\bar{x}$  with positive multipliers in (4.5). Assume |I| < d holds for any such index set  $I \in I$ . Then  $\bar{z}$  is a positive combination of at most d-1 active constraints in  $\bar{x}$ . That is,  $\bar{z}$  does not lie in the interior of the outer normal cone of the vertex  $\bar{x}$ . Equivalently,  $\bar{z}$  lies in the outer normal cone of an at least one-dimensional face F of P containing  $\bar{x}$ . Then  $x^T \bar{z} = \bar{x}^T \bar{z}$  for all  $x \in F$ , in contradiction to the assumption of the theorem and Corollary 3.3. Thus there must be an index set  $\bar{I} \in I$  of cardinality of at least d. By a symmetric argument, there exists an index set  $\bar{J}$  of active constraints in  $\bar{z}$  with positive multipliers in (4.5) that has cardinality  $|\bar{J}| \geq d$ .

If existent, we pick such index sets I and J with |I| = |J| = d. Otherwise, we proceed as follows, where  $A_i$  denotes the *i*-th row of A. As  $\bar{x}$  is a vertex of P, the cone pos $\{A_i^T | i \in I\}$  is full-dimensional and contains  $\bar{z}$  in its interior. There exist linearly independent  $v^{(1)}, \ldots, v^{(d)} \in \text{pos}\{A_i^T | i \in I\}$  generating a simplicial subcone with  $\bar{z} \in \text{int pos}\{v^{(1)}, \ldots, v^{(d)}\} = \{\sum_{i=1}^d \mu_i v^{(i)} | \mu_i > 0\}.$ 

Indeed, introducing the vectors  $v^{(i)}$  corresponds to adding redundant inequalities to the  $\mathcal{H}$ -polytope P which are active in  $\bar{x}$ . By Lemma 4.4, if we show the statement for this redundant representation of P, it is also applicable to the original set. Thus, after possibly introducing these redundancies, there exists an index set I, |I| = d, of linearly independent active constraints with positive coefficients. And analogously for the subset J and the representation of  $Q^{\circ}$ . We apply the affine variable transformation  $\phi$ :  $\mathbb{R}^{2d} \to \mathbb{R}^{2d}$  defined by

$$\phi(x,z) = \begin{bmatrix} (a - Ax)_I \\ (\mathbb{1}_l - B^T z)_J \end{bmatrix}$$

and denote the new variables by  $(s,t) := (s_1, \ldots, s_d, t_1, \ldots, t_d) = (\phi_1(x, z), \ldots, \phi_{2d}(x, z))$ . Clearly,  $\phi$  is a local parameterization at  $(\bar{x}, \bar{z})$  in the sense of Condition 4.5. The inverse of  $\phi$  is given by

$$(s,t) \mapsto \begin{bmatrix} A_I^{-1}(a_I - s) \\ (B_J^T)^{-1}(\mathbb{1}_J - t) \end{bmatrix}.$$

Setting  $M := B_J^{-1} A_I^{-1}$ , the objective  $x^T z$  has the form

$$f(s,t) := \left(A_I^{-1}(a_I - s)\right)^T \left((B_J^T)^{-1}(\mathbb{1}_J - t)\right) = s^T M^T t - s^T M^T \mathbb{1}_J - a_I^T M^T t + a_I^T M^T \mathbb{1}_J$$

in the local parameterization space. Denote by  $f_1$  the homogeneous part of degree 1. Then  $(\bar{x}, \bar{z}) = \phi^{-1}(0) = (A_I^{-1}a_I, (B_J^T)^{-1}\mathbb{1}_J)$  implies

$$\nabla_{s,t} f_1(0) = (-\mathbb{1}_J^T B_J^{-1} A_I^{-1}, -a_I^T (A_I^T)^{-1} (B_J^T)^{-1}) = (-\bar{z}^T A_I^{-1}, -\bar{x}^T (B_J^T)^{-1}) = (-\alpha_I^T, -\beta_J^T),$$

where the last equation follows from the first identity in (4.5). Thus the first part of Condition 4.5 is satisfied. Since |I| + |J| = r = 2d (where r is from Condition 4.5), the second assumption in Condition 4.5 is obsolete. Therefore, by Proposition 4.6,  $\mu^* - x^T z \in QM(A, B)$ .

Geometrically, the proof uses that in a global maximizer of the bilinear problem (which by Corollary 3.3 is a vertex of the polytope  $P \times Q^{\circ}$ ) traversing along one of the outgoing edges strictly decreases the objective function, making the second assumption in Condition 4.5 obsolete.

4.3. A sufficient criterion and examples. To illustrate the behavior of the approach, we discuss some properties and a sufficient criterion. It is helpful to start from the following two structured examples.

**Example 4.8.** Let P be the cube  $P = \{-1 \leq x_i \leq 1, i = 1, ..., d\} \subseteq \mathbb{R}^d$ , and let  $Q^\circ = \{-1 \leq ez_i \leq 1, i = 1, ..., d\} \subseteq \mathbb{R}^d$ , i.e., Q is a d-dimensional cross polytope scaled by a positive integer e. Clearly,  $P \subseteq Q$  if and only if  $e \geq d$ .

Consider the sos representation of order t = 2

$$\frac{d}{e} - x^T z = \frac{1}{8e} \sum_{i=1}^a \left[ (1 - x_i) [(1 + x_i)^2 + (1 + ez_i)^2] + (1 + x_i) [(1 - x_i)^2 + (1 - ez_i)^2] \right] \\ + \frac{1}{8e} \sum_{i=1}^d \left[ (1 - ez_i) [(1 + x_i)^2 + (1 + ez_i)^2] + (1 + ez_i) [(1 - x_i)^2 + (1 - ez_i)^2] \right].$$

If  $e \ge d$ , then  $1 - x^T z \ge \frac{d}{e} - x^T z \ge 0$ , certifying the containment  $P \subseteq Q$  (with strongness if e > d). If e < d, then  $1 - x^T z < \frac{d}{e} - x^T z$ . This is not a certificate for non-containment, since there might be a different sos representation. However, in this case this is not possible since  $e \ge d$  is a necessary condition for containment. Note that the necessary order is low and the number of terms is linear in the dimension.

$d \setminus t$	2	3	4	5
2	0.7071	$\begin{array}{c} 0.9937 \\ 0.8819 \\ 0.7906 \\ 0.7211 \end{array}$	0.9994	0.9999
3	0.5774	0.8819	0.9949	0.9994
4	0.5000	0.7906	0.9461	
5	0.4472	0.7211		

TABLE 1. Computational test of containment of an r-scaled  $\mathcal{H}$ -cube in a  $\mathcal{V}$ -cube as described in Example 4.9. The entries denote the maximal r (rounded to four decimal places) such that containment in dimension d is certified by the order t.

**Example 4.9.** Let P be the d-dimensional cube in  $\mathcal{H}$ -representation as in Example 4.8 and  $Q = \operatorname{conv}(\{-1,1\}^d)$  be the d-dimensional cube in  $\mathcal{V}$ -representation. Denote by  $rP := \{x \in \mathbb{R}^d \mid -r \leq x_i \leq r, i = 1, \ldots, d\}$  the r-scaled cube with edge length 2r. Clearly,  $rP \subseteq Q$  if and only if  $0 \leq r \leq 1$ . This containment problem is combinatorially hard since the number of inequalities is equal to  $2d + 2^d$  and thus exponential in the dimension.

We are interested in the maximal r such that the containment  $rP \subseteq Q$  is certified by a certain step t. We also ask for the minimal t such that  $P = 1P \subseteq Q$  is certified. Note that for r = 1 a priori the existence of such a t is not clear since neither Theorem 4.2 nor Theorem 4.3 applies.

For  $r = \sqrt{d}/d$ ,  $rP \subseteq Q$  is certified by the sos representation

$$1 - x^{T}z = \frac{1}{2} \sum_{i=1}^{d} (x_{i} - z_{i})^{2} + \frac{1}{2^{d+1}} \sum_{v \in \{-1,1\}^{d}} (1 + v^{T}z)^{2} (1 - v^{T}z)$$
  
+  $\frac{1}{4r} \sum_{i=1}^{d} ((r + x_{i})^{2} (r - x_{i}) + (r - x_{i})^{2} (r + x_{i})).$ 

We are not aware of a more compact sos representation. Numerically, for t = 2 and  $d \le 5$ , we get  $r(d) = \sqrt{d/d}$ ; see Table 1.

Note that the variable transformations  $x_i \mapsto \frac{1}{\lambda_i} x'_i$  and  $z_i \mapsto \lambda_i z'_i$  give a certificate for the containment of the box  $[-\lambda_1, \lambda_1] \times \cdots \times [-\lambda_d, \lambda_d]$  in the box  $\frac{d}{\sqrt{d}}Q$ , where  $Q = \text{conv}(\{-\lambda_1, \lambda_1\} \times \cdots \times \{-\lambda_d, \lambda_d\})$ .

The consideration of the box leads to the following sufficient criterion for the existence of a certificate in the initial relaxation step. The criterion implies that the containment of any polytope within any other "sufficiently large" polytope is certified already in the initial relaxation step.

**Theorem 4.10.** Let P be an  $\mathcal{H}$ -polytope and Q be a  $\mathcal{V}$ -polytope in  $\mathbb{R}^d$ , and assume that there exists a box  $S = \prod_{i=1}^d [-\lambda_i, \lambda_i]$  with  $\lambda_i > 0$  and  $P \subseteq S \subseteq \frac{\sqrt{d}}{d}Q$ . Then the inclusion  $P \subseteq Q$  is certified in the initial relaxation step.

To prepare for the proof, we first provide a transitivity property.

### Lemma 4.11 (Transitivity).

- Given a V-polytope Q and H-polytopes P and P' such that P' ⊆ P ⊆ Q. If for a certain t ≥ 2 the t-th step of hierarchy (4.3) certifies containment of P in Q, then it also certifies containment of P' in Q.
- (2) Given  $\mathcal{V}$ -polytopes Q and Q', and an  $\mathcal{H}$ -polytope P such that  $P \subseteq Q \subseteq Q'$ . If for a certain  $t \geq 2$  the t-th step of hierarchy (4.3) certifies containment of P in Q, then it also certifies containment of P in Q'.

*Proof.* Assume first that  $P' = P \cap \{x \in \mathbb{R}^d \mid f(x) \ge 0\}$  for an affine function  $f : \mathbb{R}^d \to \mathbb{R}$ . Given an sos representation of  $\mu(t) - x^T z$  w.r.t. P, by setting the additional sos polynomial  $\sigma_{k+1}$  to the zero-polynomial, i.e.  $\sigma_{k+1} \equiv 0$ , this yields an sos representation w.r.t. P'.

In the general case, starting with P, incorporate the defining inequalities of P' into the representation of P step-by-step. In every step the lower bound of the optimal value in (4.3) can not increase. At the end of this process the defining inequalities of P are all redundant (since  $P' \subseteq P$ ) and thus can be dropped, by Lemma 4.4. This proves part (1) of the statement. The proof of (2) is analog.

Proof of Theorem 4.10. Consider S as an  $\mathcal{H}$ -polytope and let S' be the box  $S' = \frac{d}{\sqrt{d}}S = \frac{d}{\sqrt{d}}[-\lambda,\lambda]^n$  in  $\mathcal{V}$ -representation. By example 4.9, the inclusion  $S \subseteq S'$  is certified in the initial relaxation step. Since  $P \subseteq S$  and  $S' \subseteq Q$  the transitivity statement in Lemma 4.11 implies that the inclusion  $P \subseteq Q$  is certified in the initial relaxation step.  $\Box$ 

From an optimization viewpoint, such as considering smallest enclosing balls of a polytope with regard to a polyhedral norm, it is natural to consider scaled containment problems (cf. [11]). Theorem 4.10 implies the following version of a scaled containment.

**Corollary 4.12** (Scaled containment). Let P be an  $\mathcal{H}$ -polytope and Q be a  $\mathcal{V}$ -polytope in  $\mathbb{R}^d$ , both containing 0 in the interior. Then there exists a  $\lambda > 0$  such that the containment  $\lambda P \subseteq Q$  is certified in the initial relaxation step.


We conclude with the numerical behavior of relaxation (4.3) for two non-symmetric examples.

**Example 4.13.** Consider the  $\mathcal{H}$ -polytope  $P = \{x \in \mathbb{R}^2 \mid \mathbb{1}_4 - Ax \ge 0\}$  and the  $\mathcal{V}$ -polytopes  $Q_1 = \operatorname{conv} B_1$  and  $Q_2 = \operatorname{conv} B_2$  defined by

$$A = \begin{bmatrix} -1 & -1 \\ 0 & -1 \\ 1 & 0 \\ -1 & 1 \end{bmatrix}, \quad B_1 = \begin{bmatrix} -1 & 0 & 2 & 2 & -1 \\ 1 & 3 & 1 & -1 & -1 \end{bmatrix}, \quad B_2 = \begin{bmatrix} -1 & -2 & 1 & 2 & 1 \\ 2 & 0 & -2 & 1 & 2 \end{bmatrix}.$$

P is contained in both  $Q_1$  and  $Q_2$  but not strongly contained.  $Q_1$  and P share infinitely many boundary points, in fact, the boundary of  $Q_1$  contains a facet of P.  $Q_2$  and Pintersect in a single vertex. See Figure 1. Thus in both examples we are not in the situation of Theorem 4.3.

The first problem  $P \subseteq Q_1$  is not certified in the initial relaxation step t = 2. If we scale P, the maximum scaling factor for which containment is certified is r = 0.9271. For the



(A) P (grey area) in  $Q_1$  (dotted). (B) P (grey area) in  $Q_2$  (dashed).

FIGURE 1. Two non-symmetric examples as defined in Example 4.13.

second problem  $P \subseteq Q_2$ , numerically the initial relaxation does only for a scaling up to r = 0.9996.

# 5. Open questions

In this paper, we studied algebraic certificates for POLYTOPE CONTAINMENT coming from a sum of squares approach. We close with a short discussion of open questions. We believe that these questions will be very relevant in improving the understanding of sum of squares methods for low-degree geometric problems, such as the one studied here.

For the POLYTOPE CONTAINMENT problem, can the structure of the certificates be better characterized? Such as, what are suitable degree bounds with regard to POLYTOPE CONTAINMENT or, somewhat more general, with regard to general bilinear programming problems?

The finite convergence result 4.3 is an essential prerequisite for potential combinatorial accesses to sum of squares certificates of low-degree problems. How are  $\mathcal{H}$ -to- $\mathcal{V}$  conversion algorithms (such as the Fourier-Motzkin-elimination) related to the algebraic certificates of POLYTOPE CONTAINMENT? Since Theorem 4.10 provides a very efficient certification of POLYTOPE CONTAINMENT in large polytopes, the question also arises in how far the sum of squares techniques can be effectively combined with existing combinatorial techniques such as Fourier-Motzkin.

Acknowledgments. A preliminary version of this paper appeared as a regular contributed talk for the conference MEGA 2015 (thanks to one of the conference reviewers for providing an improved sos representation in Example 4.9). We would like to thank all the anonymous referees for careful reading and detailed comments.

#### References

- M.F. Anjos, B. Ghaddar, and J.C. Vera. An iterative scheme for valid polynomial inequality generation in binary polynomial programming. In O. Günlük and G.J. Woeginger, editors, *Integer Programming and Combinatorial Optimization*, volume 6655 of *Lecture Notes in Comput. Sci.*, pages 207–222. Springer, Heidelberg, 2011.
- [2] D. Avis, D. Bremner, and R. Seidel. How good are convex hull algorithms? Comput. Geom., 7(5-6):265-301, 1997.
- [3] M.S. Bazaraa, H.D. Sherali, and C.M. Shetty. *Nonlinear Programming*. Wiley-Interscience, Hoboken, NJ, third edition, 2006.
- [4] C. Böhm and H.-P. Kriegel. Determining the convex hull in large multidimensional databases. In Y. Kambayashi, W. Winiwarter, and M. Arikawa, editors, *Data Warehousing and Knowledge Dis*covery, volume 2114, pages 294–306. Springer, 2001.
- [5] V. Chandru and J.-L. Lassez. Qualitative theorem proving in linear constraints. In N. Dershowitz and Z. Manna, editors, *Verification: Theory and Practice*, volume 2772, pages 395–406. Springer, 2003.
- [6] J. Chen and S. Burer. Globally solving nonconvex quadratic programming problems via completely positive programming. *Math. Program. Comput.*, 4(1):33–52, 2012.
- [7] R.M. Freund and J.B. Orlin. On the complexity of four polyhedral set containment problems. *Math. Program.*, 33(2):139–145, 1985.
- [8] G. Gallo and A. Ülkücü. Bilinear programming: An exact algorithm. Math. Program., 12(1):173–194, 1977.
- [9] J. Gouveia, R. Grappe, V. Kaibel, K. Pashkovich, R. Robinson, and R. Thomas. Which nonnegative matrices are slack matrices? *Linear Algebra Appl.*, 439(10):2921–2933, 2013.
- [10] P. Gritzmann and V. Klee. Computational complexity of inner and outer *j*-radii of polytopes in finite-dimensional normed spaces. *Math. Program.*, 59(2, Ser. A):163–213, 1993.
- P. Gritzmann and V. Klee. On the complexity of some basic problems in computational convexity.
  I. Containment problems. *Discrete Math.*, 136(1-3):129–174, 1994.
- [12] P. Gritzmann, V. Klee, and D. Larman. Largest j-simplices in n-polytopes. Discrete Comput. Geom., 13(3-4):477–515, 1995.
- [13] D. Henrion. On semidefinite representations of plane quartics. Technical Report 08444, LAAS-CNRS, 2008.
- [14] M. Joswig and G.M. Ziegler. Convex hulls, oracles, and homology. J. Symb. Comp., 38(4):1247 1259, 2004.
- [15] V. Kaibel and M.E. Pfetsch. Some algorithmic problems in polytope theory. In M. Joswig and N. Takayama, editors, *Algebra, Geometry, and Software Systems*, pages 23–47. Springer, Berlin, 2003.
- [16] K. Kellner, T. Theobald, and C. Trabandt. Containment problems for polytopes and spectrahedra. SIAM J. Optim., 23(2):1000–1020, 2013.
- [17] K. Kellner, T. Theobald, and C. Trabandt. A semidefinite hierarchy for containment of spectrahedra. SIAM J. Optim., 25(2):1013–1033, 2015.
- [18] L. Khachiyan, E. Boros, K. Borys, K. Elbassioni, and V. Gurvich. Generating all vertices of a polyhedron is hard. *Discrete Comput. Geom.*, 39(1-3):174–190, 2008.
- [19] L.G. Khachiyan. Polynomial algorithms in linear programming. USSR Computational Mathematics and Mathematical Physics, 20(1):53-72, 1980.
- [20] H. Konno. A cutting plane algorithm for solving bilinear programs. Math. Program., 11(1):14–27, 1976.
- [21] J.B. Lasserre. Global optimization with polynomials and the problem of moments. SIAM J. Optim., 11(3):796, 2001.
- [22] J.B. Lasserre. Moments, Positive Polynomials and their Applications. Imperial College Press, London, 2010.

- [23] M. Laurent. Sums of squares, moment matrices and optimization over polynomials. In M. Putinar and S. Sullivant, editors, *Emerging Applications of Algebraic Geometry*, volume 149 of *IMA Vol. Math. Appl.*, pages 157–270. Springer, New York, 2009.
- [24] M. Marshall. Positive Polynomials and Sums of Squares. American Mathematical Society, Providence, Rhode Island, 2008.
- [25] M. Marshall. Representation of non-negative polynomials, degree bounds and applications to optimization. Canad. J. Math., 61(205-221), 2009.
- [26] P. McMullen. The maximum numbers of faces of a convex polytope. Mathematika, 17:179–184, 1970.
- [27] D. Monniaux. A quantifier elimination algorithm for linear real arithmetic. In I. Cervesato, H. Veith, and A. Voronkov, editors, *Logic for Programming, Artificial Intelligence, and Reasoning*, volume 5330, pages 243–257. Springer, 2008.
- [28] J. Nie. Optimality conditions and finite convergence of Lasserre's hierarchy. Math. Program., 146(1, Ser. A):97–121, 2014.
- [29] A. Packer. Polynomial-time approximation of largest simplices in V-polytopes. Discrete Appl. Math., 134(1-3):213–237, 2004.
- [30] M. Putinar. Positive polynomials on compact semi-algebraic sets. Indiana Univ. Math. J., 42(3):969– 984, 1993.
- [31] C. Scheiderer. Positivity and sums of squares: a guide to recent results. In M. Putinar and S. Sullivant, editors, *Emerging Applications of Algebraic Geometry*, volume 149 of *IMA Vol. Math. Appl.*, pages 271–324. Springer, New York, 2009.
- [32] A. Schrijver. Theory of Linear and Integer Programming. Wiley-Interscience Series in Discrete Mathematics. John Wiley & Sons Ltd., Chichester, 1986.
- [33] G.M. Ziegler. Lectures on Polytopes. Graduate Texts in Mathematics. Springer, New York, 1995.

Goethe-Universität, FB 12 – Institut für Mathematik, Postfach 11 19 32, D–60054 Frankfurt am Main, Germany

*E-mail address*: {kellner,theobald}@math.uni-frankfurt.de