ASYMPTOTICS FOR INFINITE SYSTEMS OF DIFFERENTIAL EQUATIONS

LASSI PAUNONEN AND DAVID SEIFERT

ABSTRACT. This paper investigates the asymptotic behaviour of solutions to certain infinite systems of ordinary differential equations. In particular, we use results from ergodic theory and the asymptotic theory of C_0 -semigroups to obtain a characterisation, in terms of convergence of certain Cesàro averages, of those initial values which lead to convergent solutions. Moreover, we obtain estimates on the *rate* of convergence for solutions whose initial values satisfy a stronger ergodic condition. These results rely on a detailed spectral analysis of the operator describing the system, which is made possible by certain structural assumptions on the operator. The resulting class of systems is sufficiently broad to cover a number of important applications, including in particular both the so-called *robot rendezvous problem* and an important class of *platoon systems* arising in control theory. Our method leads to new results in both cases.

1. INTRODUCTION

The purpose of this paper is to study the asymptotic behaviour of solutions to infinite systems of coupled ordinary differential equations. In particular, given $m \in \mathbb{N}$, we consider time-dependent vectors $x_k(t)$ satisfying

(1.1)
$$\dot{x}_k(t) = A_0 x_k(t) + A_1 x_{k-1}(t), \quad k \in \mathbb{Z}, \ t \ge 0,$$

for $m \times m$ matrices A_0 and A_1 , and we assume that the initial values $x_k(0) \in \mathbb{C}^m$, $k \in \mathbb{Z}$, are known. The characteristic feature of this class of systems is that the dynamics of each subsystem depend not only on the state of the subsystem itself but also the state of the previous subsystem. Systems of this type arise naturally in applications, and indeed our investigation of such models is motivated by two important examples.

The first is the so-called robot rendezvous problem [9, 10], where m = 1, $A_0 = -1$ and $A_1 = 1$. In this case the equations in (1.1) can be thought of as describing the motion in the complex plane of countably many vehicles, or robots, indexed by the integers $k \in \mathbb{Z}$, following the rule that robot k moves in the direction of robot k-1 with speed equal to their separation. A second

²⁰¹⁰ Mathematics Subject Classification. 34A30, 34D05 (34H15, 47D06, 47A10, 47A35). Key words and phrases. System, ordinary differential equations, asymptotic behaviour, rates of convergence, C_0 -semigroup, spectrum, ergodic theory.

Part of this work was carried out while the first author visited Oxford in March and April 2015. The visit was supported by the EPSRC grant EP/J010723/1 held by Professor C.J.K. Batty (Oxford) and Professor Y. Tomilov (Warsaw). The first author is funded by the Academy of Finland grant number 298182. Both authors would like to thank Professor Batty for his useful comments on an earlier version of this paper.

important example in which the general model (1.1) arises is the study of *platoon systems* in control theory; see for instance [17, 19, 21]. Here we begin with a more realistic dynamical model of our vehicles by associating with each a position in the complex plane as well as a velocity and an acceleration, and the control objective is to steer the vehicles towards a state in which, for each $k \in \mathbb{Z}$, vehicle k is a certain target separation $c_k \in \mathbb{C}$ away from vehicle k - 1 and all vehicles are moving at a target velocity $v \in \mathbb{C}$. This model too can be written in the form (1.1) for m = 3 and suitable 3×3 matrices A_0 and A_1 which involve certain control parameters that need to be fixed. In both cases the key question is whether solutions converge to a limit as $t \to \infty$. Thus in the robot rendezvous problem we would like to know whether the positions of the robots converge to a mutual meeting, or *rendezvous*, point, and in the platoon system we ask whether we can choose the control parameters in such a way that the vehicles asymptotically approach their target state.

We present a unified approach to the study of these problems by first reformulating the system (1.1) as the abstract Cauchy problem

(1.2)
$$\begin{cases} \dot{x}(t) = Ax(t), & t \ge 0, \\ x(0) = x_0 \in X, \end{cases}$$

on the space $X = \ell^p(\mathbb{C}^m)$ with $m \in \mathbb{N}$ and $1 \leq p \leq \infty$. Note that (1.2) indeed becomes (1.1) if we let $x(t) = (x_k(t))_{k \in \mathbb{Z}}$ for $t \geq 0$, $x_0 = (x_k(0))_{k \in \mathbb{Z}}$ and take the bounded linear operator A to act by sending a sequence $(x_k)_{k \in \mathbb{Z}} \in X$ to

$$Ax = (A_0 x_k + A_1 x_{k-1})_{k \in \mathbb{Z}}.$$

Systems of this form are examples of what are sometimes called "spatially invariant systems", where in general it is possible for the dynamics of each subsystem to depend on more than just one other subsystem; see for instance [4]. Our main objective is to investigate whether or not the solution x(t), $t \geq 0$, of (1.2) converges to a limit as $t \to \infty$ and, if so, what can be said about the rate of convergence. Most of the existing research into such systems is confined to the Hilbert space case p = 2. For instance, it is shown in [8] using Fourier transform techniques that solutions $x(t), t \ge 0$, of some spatially invariant systems of the form (1.1) on the space $X = \ell^2(\mathbb{C}^2)$ satisfy $x(t) \to 0$ as $t \to \infty$ for all initial values $x_0 \in X$, but that there exists no uniform rate of decay. Since the Fourier transform approach is specific to the Hilbert space setting, we develop a new approach to studying the asymptotic behaviour of solutions of (1.2) in the case where the matrices A_0 and A_1 satisfy certain additional assumptions. Specifically, we assume throughout that $A_1 \neq 0$ to avoid the trivial uncoupled case, but more importantly we suppose that there exists a rational function ϕ such that

(1.3)
$$A_1(\lambda - A_0)^{-1}A_1 = \phi(\lambda)A_1, \qquad \lambda \in \mathbb{C} \setminus \sigma(A_0)$$

When such a function ϕ exists we call it the *characteristic function* of our system. Both the robot rendezvous problem and the platoon system fall into this special class, as indeed do many other systems. For systems having this property we develop techniques allowing us to handle the full range $1 \leq p \leq \infty$ rather than just the case p = 2, and in particular we include the cases p = 1 and $p = \infty$, where it turns out no longer to be the case that

all solutions converge to a limit. In fact our approach, which is based on a detailed analysis of the operator A and the C_0 -semigroup it generates, leads to a complete understanding of which initial values do and which do not lead to convergent solutions in these cases, and moreover gives an estimate on the *rate* of convergence for a certain subset of initial values.

The paper is organised as follows. Our main theoretical results are presented in Sections 2, 3, and 4. In Section 2 we examine the spectral properties of A, and the main results are Theorem 2.3, which among other things provides a very simple characterisation of the set $\sigma(A) \setminus \sigma(A_0)$ in terms of the characteristic function ϕ , namely

$$\sigma(A) \setminus \sigma(A_0) = \{ \lambda \in \mathbb{C} \setminus \sigma(A_0) : |\phi(\lambda)| = 1 \},\$$

and Proposition 2.5, which describes the behaviour of the resolvent operator of A in the neighbourhood of spectral points. In Section 3 we turn to the delicate issue of whether the semigroup generated by A is uniformly bounded. The main result here is Theorem 3.1, which gives a sufficient condition for uniform boundedness involving the derivatives of ϕ . In Section 4, we then combine the results of Sections 2 and 3 with known results in ergodic theory and recent results in the theory of C_0 -semigroups [6, 7, 16] in order to obtain our main result, Theorem 4.3, which describes the asymptotic behaviour of solutions to general systems in our class. For instance, it is a consequence of Theorem 4.3 that there exists an even integer $n \ge 2$ determined solely by the characteristic function ϕ such that for all $x_0 \in X$ the derivative of the solution $x(t), t \ge 0$, of (1.2) satisfies the quantified decay estimate

(1.4)
$$\|\dot{x}(t)\| = O\left(\left(\frac{\log t}{t}\right)^{1/n}\right), \quad t \to \infty,$$

and the logarithm can be omitted if p = 2. Moreover, for $1 not only the derivative of each solution but also the solution itself decays to zero as <math>t \to \infty$, but this is no longer true when p = 1 or $p = \infty$. In these cases, Theorem 4.3 gives a characterisation, in terms of convergence of certain Cesàro means, of those initial values $x_0 \in X$ which do lead to convergent solutions, and the result also shows that under a supplementary condition the convergence of solutions to their limit can be quantified in a form analogous to (1.4).

In Sections 5 and 6 we return to the motivating examples. First, in Section 5, we apply the general result in the setting of the platoon system, which leads to extensions of results obtained previously in [8, 21] for the Hilbert space case p = 2. In particular, the main result in this section, Theorem 5.1, shows that the platoon system approaches its target for all $x_0 \in X$ not just for p = 2, as was shown in [21], but more generally when 1 . We also show that for <math>p = 1 and $p = \infty$ this statement is no longer true but our Theorem 5.1 provides a simple ergodic condition on the initial *displacements* of the vehicles which is necessary and sufficient for the solution to converge to a limit. Then, in Section 6 we return to the robot rendezvous problem and use our general result, Theorem 4.3, to settle several questions left open in [9, 10]. We conclude in Section 7 by mentioning several topics which remain subjects for future research.

The notation we use is more or less standard throughout. In particular, given a complex Banach space X, the norm on X will typically be denoted by $\|\cdot\|$ and occasionally, in order to avoid ambiguity, by $\|\cdot\|_X$. In particular, for $m \in \mathbb{N}$ and $1 \leq p \leq \infty$, we let $\ell^p(\mathbb{C}^m)$ denote the space of doubly infinite sequences $(x_k)_{k\in\mathbb{Z}}$ such that $x_k\in\mathbb{C}^m$ for all $k\in\mathbb{Z}$ and $\sum_{k\in\mathbb{Z}}\|x_k\|^p<\infty$ if $1 \leq p < \infty$ and $\sup_{k \in \mathbb{Z}} ||x_k|| < \infty$ if $p = \infty$. Here and in all that follows we endow the finite-dimensional space \mathbb{C}^m with the standard Euclidean norm and we consider $\ell^p(\mathbb{C}^m)$ with the norm given for $x = (x_k)_{k \in \mathbb{Z}}$ by ||x|| = $(\sum_{k\in\mathbb{Z}} \|x_k\|^p)^{1/p}$ if $1 \leq p < \infty$ and $\|x\| = \sup_{k\in\mathbb{Z}} \|x_k\|$ if $p = \infty$. With respect to this norm $\ell^p(\mathbb{C}^m)$ is a Banach space for $1 \leq p \leq \infty$ and a Hilbert space when p = 2. We write X^* for the dual space of X, and given $\phi \in X^*$ the action of ϕ on $x \in X$ is written as $\langle x, \phi \rangle$. Moreover we write $\mathcal{B}(X)$ for the space of bounded linear operators on X, and given $A \in \mathcal{B}(X)$ we write $\operatorname{Ker}(A)$ for the kernel and $\operatorname{Ran}(A)$ for the range of A. Moreover, we let $\sigma(A)$ denote the spectrum of A and, for $\lambda \in \mathbb{C} \setminus \sigma(A)$ we write $R(\lambda, A)$ for the resolvent operator $(\lambda - A)^{-1}$. We write $\sigma_p(A)$ for the point spectrum and $\sigma_{ap}(A)$ for the approximate point spectrum of A. Given $A \in \mathcal{B}(X)$ we denote the dual operator of A by A'. If A is a matrix, we write A^T for the transpose of A. Given two functions f and g taking values in $(0,\infty)$, we write $f(t) = O(q(t)), t \to \infty$, if there exists a constant C > 0 such that $f(t) \leq Cg(t)$ for all sufficiently large values of t. If f(t) = O(g(t)) and g(t) = O(f(t)) as $t \to \infty$, or more generally as the argument t tends to some point in the extended complex plane, we write $f(t) \simeq g(t)$ in the limit. Given two real-valued quantities a and b, we write $a \leq b$ if there exists a constant C > 0 such that $a \leq Cb$ for all values of the parameters that are free to vary in a given situation. Finally, we denote the open right/left half plane by $\mathbb{C}_{\pm} = \{\lambda \in \mathbb{C} : \operatorname{Re} \lambda \geq 0\}$, and we use a horizontal bar over a set to denote its closure.

2. Spectral theory

We begin by stating two standing assumptions on the matrices A_0 , A_1 appearing in (1.1).

Assumptions 2.1. We assume that

(A1)
$$A_1 \neq 0.$$

Moreover we assume that there exists a function ϕ such that

(A2)
$$A_1 R(\lambda, A_0) A_1 = \phi(\lambda) A_1, \qquad \lambda \in \mathbb{C} \setminus \sigma(A_0).$$

If this assumption is satisfied we call ϕ the *characteristic function*.

Remark 2.2. It is clear that if (A2) is satisfied then the characteristic function ϕ is a rational function whose poles belong to the set $\sigma(A_0)$. Note also that for $|\lambda| > ||A_0||$ we have

$$\|\phi(\lambda)\|\|A_1\| \le \frac{\|A_1\|^2}{|\lambda| - \|A_0\|}.$$

In particular, when (A1) and (A2) both hold it follows that $|\phi(\lambda)| \to 0$ as $|\lambda| \to \infty$. It is straightforward to show that both (A1) and (A2) are satisfied whenever rank $(A_1) = 1$.

In this section we characterise the spectrum of the operator A under our standing assumptions (A1) and (A2). The following is the main result. It essentially characterises the spectrum of A in terms of the characteristic function ϕ . Here and in what follows we use the notation

$$\Omega_{\phi} = \{\lambda \in \mathbb{C} \setminus \sigma(A_0) : |\phi(\lambda)| = 1\}.$$

Theorem 2.3. Let $1 \le p \le \infty$ and $m \in \mathbb{N}$, and suppose that (A1), (A2) hold. Then the spectrum of A satisfies

(2.1)
$$\sigma(A) \setminus \sigma(A_0) = \Omega_{\phi}.$$

Moreover, the following hold:

- (a) If $1 \le p < \infty$, then $\sigma(A) \setminus \sigma(A_0) \subset \sigma_{ap}(A) \setminus \sigma_p(A)$.
- (b) If $p = \infty$, then $\sigma(A) \setminus \sigma(A_0) \subset \sigma_p(A)$ and, given $\lambda \in \sigma(A) \setminus \sigma(A_0)$,

(2.2)
$$\operatorname{Ker}(\lambda - A) = \left\{ (\phi(\lambda)^k x_0)_{k \in \mathbb{Z}} : x_0 \in \operatorname{Ran}(R(\lambda, A_0)A_1) \right\}$$

In particular, dim Ker $(\lambda - A) = \operatorname{rank}(A_1)$ for all $\lambda \in \sigma(A) \setminus \sigma(A_0)$.

Furthermore, for $\lambda \in \sigma(A) \setminus \sigma(A_0)$ the range of $\lambda - A$ is dense in X if and only if 1 .

Remark 2.4. The points $\sigma(A_0)$ may be in either $\sigma(A)$ or $\rho(A)$ depending on the matrices A_0 and A_1 . Note for instance that, given $\lambda \in \mathbb{C}$, any vector $x = (x_k)_{k \in \mathbb{Z}}$ with $x_0 \in \operatorname{Ker}(\lambda - A_0) \cap \operatorname{Ker}(A_1)$ and $x_k = 0$ for $k \neq 0$ satisfies $x \in \operatorname{Ker}(\lambda - A)$. In particular, $\lambda \in \sigma(A_0) \cap \sigma(A)$ whenever $\operatorname{Ker}(\lambda - A_0) \cap \operatorname{Ker}(A_1) \neq \{0\}$. Moreover, if $\lambda \in \mathbb{C}$ is such that $\operatorname{Ran}(\lambda - A_0) +$ $\operatorname{Ran}(A_1) \neq \mathbb{C}^m$ then it is easy to see that any sequence $(x_k)_{k \in \mathbb{Z}} \in X$ such that $x_k \notin \operatorname{Ran}(\lambda - A_0) + \operatorname{Ran}(A_1)$ for some $k \in \mathbb{Z}$ has an open neighbourhood which is disjoint from $\operatorname{Ran}(\lambda - A)$, so $\operatorname{Ran}(\lambda - A)$ cannot be dense in X and once again $\lambda \in \sigma(A_0) \cap \sigma(A)$. In Sections 5 and 6 we will see examples in which, by contrast, we have $\sigma(A_0) \cap \sigma(A) = \emptyset$.

Proof of Theorem 2.3. We begin by showing that every $\lambda \in \mathbb{C} \setminus \sigma(A_0)$ such that $|\phi(\lambda)| \neq 1$ belongs to $\rho(A)$. Indeed, given $\lambda \in \mathbb{C} \setminus \sigma(A_0)$, let $R_{\lambda} = R(\lambda, A_0)$. Supposing first that $|\phi(\lambda)| < 1$, we consider the operator $R(\lambda) \in \mathcal{B}(X)$ given by

(2.3)
$$R(\lambda)x = \left(R_{\lambda}x_{k} + R_{\lambda}A_{1}R_{\lambda}\sum_{\ell=0}^{\infty}\phi(\lambda)^{\ell}x_{k-\ell-1}\right)_{k\in\mathbb{Z}}$$

for all $x = (x_k)_{k \in \mathbb{Z}} \in X$, noting that this gives a well-defined element of Xby Young's inequality. Using the fact that $(A_1 R_\lambda)^{\ell} = \phi(\lambda)^{\ell-1} A_1 R_\lambda$ for all $\ell \in \mathbb{N}$ as a consequence of assumption (A1), it is straightforward to verify that $(\lambda - A)R(\lambda)x = R(\lambda)(\lambda - A)x = x$ for all $x \in X$, and hence $\lambda \in \rho(A)$ and $R(\lambda, A) = R(\lambda)$. A completely analogous argument goes through for $\lambda \in \mathbb{C} \setminus \sigma(A_0)$ such that $|\phi(\lambda)| > 1$, with the only difference that the operator $R(\lambda) \in \mathcal{B}(X)$ is now defined by

$$R(\lambda)x = \left(R_{\lambda}x_{k} - R_{\lambda}A_{1}R_{\lambda}\sum_{\ell=0}^{\infty}\phi(\lambda)^{-\ell-1}x_{k+\ell}\right)_{k\in\mathbb{Z}}$$

for all $x \in X$. This shows that $\sigma(A) \setminus \sigma(A_0) \subset \Omega_{\phi}$.

Suppose now that $1 \leq p < \infty$ and let $\lambda \in \Omega_{\phi}$. We will first show that $\lambda \notin \sigma_p(A)$. To this end, let $x \in X$ be such that $(\lambda - A)x = 0$. Then a simple calculation shows that $x_k = \phi(\lambda)^{k-\ell-1}R_{\lambda}A_1x_{\ell}$ for all $k, \ell \in \mathbb{Z}$ with $k > \ell$, and in particular $||x_k|| = ||R_{\lambda}A_1x_{\ell}||$ for all $k > \ell$. Hence the assumption that $x \in X$ implies that x = 0 and therefore $\lambda \notin \sigma_p(A)$, as required. In order to show that $\lambda \in \sigma_{ap}(A)$, choose $y_0 \in \mathbb{C}^m$ such that $A_1y_0 \neq 0$ and, for $n \in \mathbb{N}$, define the sequence $x^n = (x_k^n)_{k \in \mathbb{Z}} \in X$ by

$$x_k^n = \frac{\phi(\lambda)^k R_\lambda A_1 y_0}{(2n+1)^{1/p} \|R_\lambda A_1 y_0\|}, \quad |k| \le n,$$

and $x_k^n = 0$ otherwise. Then $||x^n||_p = 1$ for all $n \in \mathbb{N}$, and a direct computation shows that

$$\|(\lambda - A)x^n\|^p = \frac{\|y_0\|^p + \|A_1R_\lambda A_1y_0\|^p}{(2n+1)\|R_\lambda A_1y_0\|^p} \to 0, \quad n \to \infty.$$

Thus $\lambda \in \sigma_{ap}(A)$, which establishes (a).

Now suppose that $p = \infty$ and let $\lambda \in \Omega_{\phi}$. We will prove that (2.2) holds, from which (b) follows. Note first that if $x_0 \in \operatorname{Ran}(R_{\lambda}A_1)$ then a simple calculation shows that $(\phi(\lambda)^k x_0)_{k \in \mathbb{Z}} \in \operatorname{Ker}(\lambda - A)$. On the other hand, if $x = (x_k)_{k \in \mathbb{Z}} \in \operatorname{Ker}(\lambda - A)$, then

$$(\lambda - A_0)x_k - A_1x_{k-1} = 0$$

and hence $x_k = R_{\lambda}A_1x_{k-1} \in \operatorname{Ran}(R_{\lambda}A_1)$ for all $k \in \mathbb{Z}$. Since

$$x_k = (R_\lambda A_1)^2 x_{k-2} = \phi(\lambda) R_\lambda A_1 x_{k-2} = \phi(\lambda) x_{k-1}, \quad k \in \mathbb{Z},$$

by assumption (A1) we obtain that $x_k = \phi(\lambda)^k x_0$ for all $k \in \mathbb{Z}$. Thus (b) follows, and by combining (a) and (b) with the fact that $\sigma(A) \setminus \sigma(A_0) \subset \Omega_{\phi}$ we obtain (2.1). It remains to prove the final statement.

Suppose first that $1 and let <math>q = p(p-1)^{-1}$ be the Hölder conjugate of p. Moreover let $\lambda \in \Omega_{\phi}$ and that $y = (y_k)_{k \in \mathbb{Z}} \in X^* = \ell^q(\mathbb{C}^m)$ is such that $\langle (\lambda - A)x, y \rangle = 0$ for all $x \in X$. Then $y \in \operatorname{Ker}(\lambda - A')$, where the dual operator A' of A is given by $A'y = (A_0^T y_k + A_1^T y_{k+1})_{k \in \mathbb{Z}}$ for all $y = (y_k)_{k \in \mathbb{Z}} \in X^*$. Since by assumption (A1) we have $(A_1 R_\lambda A_1)^T = \phi(\lambda) A_1^T$, a direct computation shows that

$$y_k = \phi(\lambda)^{\ell - k - 1} R_\lambda^T A_1^T y_\ell$$

for all $k, \ell \in \mathbb{Z}$ with $k < \ell$. As in the above argument showing that $\lambda \notin \sigma_p(A)$ we obtain that y = 0, and hence $\operatorname{Ran}(\lambda - A)$ is dense in X by a standard corollary of the Hahn-Banach theorem. On the other hand, if p = 1 and $\lambda \in \Omega_{\phi}$ we can consider the element $y = (y_k)_{k \in \mathbb{Z}} \in X^* = \ell^{\infty}(\mathbb{C}^m)$ with entries

$$y_k = \phi(\lambda)^{-k} R_\lambda^T A_1^T y_0, \quad k \in \mathbb{Z},$$

where $y_0 \in \mathbb{C}^m$ is chosen in such a way that $A_1^T y_0 \neq 0$. A simple verification shows $y \in \operatorname{Ker}(\lambda - A')$ and hence that $\langle (\lambda - A)x, y \rangle = 0$ for all $x \in X$, so $\operatorname{Ran}(\lambda - A)$ cannot be dense in X. Finally, suppose that $p = \infty$ and that $\lambda \in \Omega_{\phi}$. Let $y = (\phi(\lambda)^k y_0)_{k \in \mathbb{Z}} \in X$, where $y_0 \in \mathbb{C}^m$ is such that $R_{\lambda}A_1R_{\lambda}y_0 \neq 0$. We show that y lies outside the closure of $\operatorname{Ran}(\lambda - A)$. Indeed, let $0 < \varepsilon < ||R_{\lambda}A_1R_{\lambda}y_0||/||R_{\lambda}A_1R_{\lambda}||$ and suppose for the sake of contradiction that there exists $x \in X$ such that

$$\|(\lambda - A)x - y\| = \sup_{k \in \mathbb{Z}} \|(\lambda - A_0)x_k - A_1x_{k-1} - y_k\| < \varepsilon.$$

Let $z_k = (\lambda - A_0)x_k - A_1x_{k-1} - y_k$, so that $||z_k|| < \varepsilon$ for all $k \in \mathbb{Z}$. A simple inductive argument shows that for all $n \in \mathbb{N}$ we have

$$x_0 = \phi(\lambda)^{n-1} R_{\lambda} A_1 x_{-n} + R_{\lambda} (y_0 + z_0) + R_{\lambda} A_1 R_{\lambda} \sum_{\ell=1}^{n-1} \phi(\lambda)^{\ell-1} (y_{-\ell} + z_{-\ell}).$$

Since

$$\left\| R_{\lambda}A_{1}R_{\lambda}\sum_{\ell=1}^{n}\phi(\lambda)^{\ell-1}y_{-\ell} \right\| = n\|R_{\lambda}A_{1}R_{\lambda}y_{0}\|, \quad n \in \mathbb{N},$$

we obtain that

$$\|x_0\| \ge n \big(\|R_{\lambda}A_1R_{\lambda}y_0\| - \varepsilon \|R_{\lambda}A_1R_{\lambda}\| \big) - \|R_{\lambda}\| \big(\|y_0\| + \varepsilon \big) - \|R_{\lambda}A_1\| \|x\|$$

for all $n \in \mathbb{N}$. However, by the choice of ε this is absurd. Hence no such $x \in X$ exists and in particular the range of $\lambda - A$ is not dense in X. This completes the proof.

The next result establishes a useful estimate for the norm of the resolvent operator in the neighbourhood of singular points.

Proposition 2.5. Fix $1 \le p \le \infty$ and $m \in \mathbb{N}$, and suppose that (A1), (A2) hold. If $\lambda \in \mathbb{C} \setminus \sigma(A_0)$ is such that $|\phi(\lambda)| \ne 1$, then

$$\left| \|R(\lambda, A)\| - \frac{\|R(\lambda, A_0)A_1R(\lambda, A_0)\|}{|1 - |\phi(\lambda)||} \right| \le \|R(\lambda, A_0)\|$$

In particular, for $\lambda_0 \in \mathbb{C} \setminus \sigma(A_0)$ such that $|\phi(\lambda_0)| = 1$ we have

$$\|R(\lambda, A)\| \asymp \frac{1}{|1 - |\phi(\lambda)||}$$

as $\lambda \to \lambda_0$ in the region $\{\lambda \in \mathbb{C} \setminus \sigma(A_0) : |\phi(\lambda)| \neq 1\}.$

Proof. As in the proof of Theorem 2.3, we let $R_{\lambda} = R(\lambda, A_0)$ for $\lambda \in \mathbb{C} \setminus \sigma(A_0)$. We consider the case where $0 < |\phi(\lambda)| < 1$; the case $|\phi(\lambda)| > 1$ follows similarly, as in the proof of Theorem 2.3. From (2.3) we see that for $\lambda \in \mathbb{C} \setminus \sigma(A_0)$ such that $|\phi(\lambda)| < 1$ we have $R(\lambda, A) = D(\lambda) + Q(\lambda)$, where $D(\lambda)x = (R_{\lambda}x_k)_{k\in\mathbb{Z}}$ and

$$Q(\lambda)x = \left(R_{\lambda}A_{1}R_{\lambda}\sum_{\ell=0}^{\infty}\phi(\lambda)^{\ell}x_{k-\ell-1}\right)_{k\in\mathbb{Z}}$$

for all $x = (x_k)_{k \in \mathbb{Z}} \in X$. Note that $||D(\lambda)|| = ||R_{\lambda}||$, so the result will follow from the triangle inequality once we have established that

(2.4)
$$\|Q(\lambda)\| = \frac{\|R_{\lambda}A_1R_{\lambda}\|}{1 - |\phi(\lambda)|}.$$

In fact, since

$$\|Q(\lambda)\| \leq \frac{\|R_{\lambda}A_{1}R_{\lambda}\|}{1 - |\phi(\lambda)|}$$

for $1 \le p \le \infty$ by a straightforward estimate, it suffices to prove the converse inequality.

Suppose first that $p = \infty$ and consider the sequence $x = (e^{ik\theta}y_0)_{k\in\mathbb{Z}} \in X$, where $\theta = \arg \phi(\lambda)$ and $y_0 \in \mathbb{C}^m$ is such that $||x_0|| = 1$ and $||R_{\lambda}A_1R_{\lambda}y_0|| =$ $||R_{\lambda}A_1R_{\lambda}||$. Then ||x|| = 1 and

$$\|Q(\lambda)x\| = \sup_{k \in \mathbb{Z}} \left\| R_{\lambda} A_1 R_{\lambda} \sum_{\ell=0}^{\infty} \phi(\lambda)^{\ell} x_{k-\ell-1} \right\| = \frac{\|R_{\lambda} A_1 R_{\lambda}\|}{1 - |\phi(\lambda)|},$$

thus establishing (2.4). Now suppose that $1 \leq p < \infty$. Once again let $\theta = \arg \phi(\lambda)$ and let $y_0 \in \mathbb{C}^m$ be such that $||y_0|| = 1$ and $||R_{\lambda}A_1R_{\lambda}y_0|| = ||R_{\lambda}A_1R_{\lambda}||$. Furthermore, let $\varepsilon \in (0, 1)$ and let $M, N \in \mathbb{N}$ be such that

$$\sum_{\ell=M+1}^{\infty} |\phi(\lambda)|^{\ell} < \varepsilon \quad \text{and} \quad \frac{N-M}{N} > (1-\varepsilon)^{p}.$$

Consider the sequence $x = (x_k)_{k \in \mathbb{Z}} \in X$ with entries $x_k = e^{ik\theta} \alpha_k y_0$, where $\alpha_k = N^{-1/p}$ for $-N \leq k \leq -1$ and $\alpha_k = 0$ otherwise. Then ||x|| = 1 and

$$\|Q(\lambda)x\|^p = \|R_{\lambda}A_1R_{\lambda}\|^p \sum_{k \in \mathbb{Z}} \left(\sum_{\ell=0}^{\infty} |\phi(\lambda)|^\ell \alpha_{k-\ell-1}\right)^p,$$

and hence by our choices of M and N we obtain that

$$\begin{aligned} \|Q(\lambda)x\| &\geq \frac{\|R_{\lambda}A_{1}R_{\lambda}\|}{N^{1/p}} \left(\sum_{M-N+1\leq k\leq 0} \left(\sum_{\ell=0}^{M} |\phi(\lambda)|^{\ell}\right)^{p}\right)^{1/p} \\ &> (1-\varepsilon) \frac{\|R_{\lambda}A_{1}R_{\lambda}\|}{1-|\phi(\lambda)|} - \varepsilon(1-\varepsilon) \|R_{\lambda}A_{1}R_{\lambda}\|. \end{aligned}$$

1 /

Since $\varepsilon \in (0, 1)$ was arbitrary, (2.4) follows and the proof is complete. \Box

We conclude this section with a refinement of Proposition 2.5 in an important special case.

Lemma 2.6. Fix $1 \le p \le \infty$ and $m \in \mathbb{N}$, and suppose that (A1), (A2) hold, and that $0 \in \Omega_{\phi} \subset \mathbb{C}_{-} \cup \{0\}$. Then there exists an even integer $n \ge 2$ such that $1 - |\phi(is)| \asymp |s|^n$ as $|s| \to 0$.

Proof. The rational function ϕ is of the form $\phi(\lambda) = p(\lambda)/q(\lambda)$, where p and q are coprime polynomials and the roots of q are contained in the set $\sigma(A_0) \subset \mathbb{C}_-$. Since $|\phi(0)| = 1$ and $|\phi(\lambda)| \to 0$ as $|\lambda| \to \infty$, we have that $|\phi(is)| < 1$ for $s \neq 0$ and hence

$$1 - |\phi(is)| = \frac{|q(is)|^2 - |p(is)|^2}{|q(is)|(|p(is)| + |q(is)|)}, \quad s \neq 0.$$

The denominator of the right-hand side is bounded from above and from below near s = 0. Thus the rate at which $1 - |\phi(is)| \to 0$ is equal to that at which $r(s) = |q(is)|^2 - |p(is)|^2 \to 0$ as $|s| \to 0$. Since r is a real polynomial satisfying r(0) = 0 and r(s) > 0 for $s \neq 0$, we have that $r(s) = s^n r_0(s)$, $s \in \mathbb{R}$, where $n \in \mathbb{N}$ is even and r_0 is a polynomial satisfying $r_0(0) > 0$. The claim now follows. **Remark 2.7.** Note that $n = n_{\phi}$ is determined by the characteristic function ϕ . We call n_{ϕ} the resolvent growth parameter.

3. Uniform boundedness of the semigroup

Consider our general model and assume that assumptions (A1) and (A2) are satisfied. In this section we present conditions on the characteristic function ϕ under which the semigroup generated by A is uniformly bounded or even contractive. Since uniform boundedness necessarily requires that $\sigma(A) \subset \overline{\mathbb{C}}_{-}$, Theorem 2.3 shows that it is necessary to assume that $\Omega_{\phi} \subset \overline{\mathbb{C}}_{-}$, where $\Omega_{\phi} = \{\lambda \in \mathbb{C} \setminus \sigma(A_0) : |\phi(\lambda)| = 1\}$. Note also that, since $|\phi(\lambda)| \to 0$ as $|\lambda| \to \infty$ by Remark 2.2, we must have $|\phi(\lambda)| < 1$ for all $\lambda \in \mathbb{C}_{+}$ in this case. The following theorem is the main result of this section.

Theorem 3.1. Let $1 \le p \le \infty$ and $m \in \mathbb{N}$. Suppose that assumptions (A1), (A2) hold, that $\sigma(A_0) \subset \mathbb{C}_-$ and that $\Omega_{\phi} \subset \overline{\mathbb{C}_-}$. If furthermore

$$(3.1) \quad \sup_{0<\lambda\leq 1} \frac{\lambda}{1-|\phi(\lambda)|} < \infty \quad and \quad \sup_{n\in\mathbb{N}} \sup_{\lambda>0} \frac{\lambda^{n+1}}{n!} \sum_{\ell=1}^{\infty} \left| \frac{d^n}{d\lambda^n} \phi(\lambda)^\ell \right| < \infty,$$

then the semigroup generated by A is uniformly bounded. If

(3.2)
$$\sup_{\lambda>0} \left(\lambda \|R(\lambda, A_0)\| + \lambda \frac{\|R(\lambda, A_0)A_1R(\lambda, A_0)\|}{1 - |\phi(\lambda)|}\right) \le 1,$$

then the semigroup generated by A is contractive.

Proof. Both parts of the result are consequences of the Hille-Yosida theorem. We thus aim to establish a uniform upper bound for $\|\lambda^n R(\lambda, A)^n\|$ as $\lambda > 0$ and $n \in \mathbb{N}$ are allowed to vary. For $\lambda > 0$ we let $R_{\lambda} = R(\lambda, A_0)$. Then by (2.3) in the proof of Theorem 2.3 and by standard properties of resolvent operators we have that

$$R(\lambda, A)^n x = (R^n_\lambda x_k)_{k \in \mathbb{Z}} + \left(\frac{(-1)^{n-1}}{(n-1)!} \sum_{\ell=0}^{\infty} \frac{d^{n-1}}{d\lambda^{n-1}} (\phi(\lambda)^\ell R_\lambda A_1 R_\lambda) x_{k-\ell-1} \right)_{k \in \mathbb{Z}}$$

for all $x = (x_k)_{k \in \mathbb{Z}} \in X$, and hence

$$(3.3) \quad \|\lambda^n R(\lambda, A)^n\| \le \|\lambda^n R_\lambda^n\| + \frac{\lambda^n}{(n-1)!} \sum_{\ell=0}^{\infty} \left\| \frac{d^{n-1}}{d\lambda^{n-1}} \left(\phi(\lambda)^\ell R_\lambda A_1 R_\lambda \right) \right\|$$

for all $\lambda > 0$ and all $n \in \mathbb{N}$. Now since $\sigma(A_0) \subset \mathbb{C}_-$, there exists $\varepsilon > 0$ such that $A_0 + \varepsilon$ generates a uniformly bounded semigroup, and in particular

(3.4)
$$\sup_{n \in \mathbb{N}} \sup_{\lambda > 0} \| (\lambda + \varepsilon)^n R^n_\lambda \| < \infty.$$

Thus the first term on the right-hand side of (3.3) is uniformly bounded as $\lambda > 0$ and $n \in \mathbb{N}$ are allowed to vary. It remains to consider the second term. Let $\phi_{\ell}(\lambda) = \phi(\lambda)^{\ell}$ and observe that, for $\lambda > 0$ and $\ell, n \in \mathbb{Z}_+$,

$$\frac{1}{n!}\frac{d^n}{d\lambda^n}\left(\phi(\lambda)^\ell R_\lambda A_1 R_\lambda\right) = \sum_{k=0}^n \frac{\phi_\ell^{(k)}(\lambda)}{k!} \frac{1}{(n-k)!} \frac{d^{n-k}}{d\lambda^{n-k}} (R_\lambda A_1 R_\lambda)$$

and by (3.4)

$$\left\|\frac{1}{n!}\frac{d^n}{d\lambda^n}(R_{\lambda}A_1R_{\lambda})\right\| = \left\|\sum_{j=0}^n R_{\lambda}^{j+1}A_1R_{\lambda}^{n-j+1}\right\| \lesssim \frac{n+1}{(\lambda+\varepsilon)^{n+2}}$$

for all $\lambda > 0$ and $n \in \mathbb{Z}_+$. It follows that

$$\frac{\lambda^n}{(n-1)!} \sum_{\ell=0}^{\infty} \left\| \frac{d^{n-1}}{d\lambda^{n-1}} \left(\phi(\lambda)^\ell R_\lambda A_1 R_\lambda \right) \right\| \lesssim \lambda^n \sum_{k=0}^{n-1} \sum_{\ell=0}^{\infty} \frac{|\phi_\ell^{(k)}(\lambda)|}{k!} \frac{n-k}{(\lambda+\varepsilon)^{n-k+1}}$$

for all $\lambda > 0$ and $n \in \mathbb{N}$. Using the first part of assumption (3.1) for the interval $0 < \lambda \leq 1$ and the fact that $\sup_{\lambda > 1} |\phi(\lambda)| < 1$ for the interval $1 < \lambda < \infty$, it is straightforward to see that the first term on the right-hand side, corresponding to k = 0, is uniformly bounded above by

$$\sup_{n\in\mathbb{N}}\sup_{\lambda>0}\ \frac{1}{1-|\phi(\lambda)|}\frac{n\lambda^n}{(\lambda+\varepsilon)^{n+1}}<\infty.$$

Using the second part of assumption (3.1) the remaining terms on the righthand side can be estimated, for all $\lambda > 0$ and $n \in \mathbb{N}$, by

$$\lambda^n \sum_{k=1}^{n-1} \sum_{\ell=0}^{\infty} \frac{|\phi_{\ell}^{(k)}(\lambda)|}{k!} \frac{n-k}{(\lambda+\varepsilon)^{n-k+1}} \lesssim \sum_{k=1}^{n-1} \frac{k\lambda^{k-1}}{(\lambda+\varepsilon)^{k+1}} \le \varepsilon^{-2}.$$

Combining the last two estimates with (3.4) in (3.3) shows that

$$\sup_{n \in \mathbb{N}} \sup_{\lambda > 0} \|\lambda^n R(\lambda, A)^n\| < \infty,$$

and hence the semigroup generated by A is uniformly bounded by the Hille-Yosida theorem.

For the second statement we note that if (3.2) holds, then Proposition 2.5 shows that $\lambda ||R(\lambda, A)|| \leq 1$ for all $\lambda > 0$, and thus the semigroup generated by A is contractive by the Hille-Yosida theorem.

The next lemma shows that the assumptions in (3.1) are satisfied in a simple but important special case.

Lemma 3.2. Let $\zeta > 0$ and $k \in \mathbb{N}$ be given, and suppose that

$$\phi(\lambda) = \frac{\zeta^k}{(\lambda+\zeta)^k}, \quad \lambda \in \mathbb{C} \setminus \{-\zeta\}.$$

Then both conditions in (3.1) are satisfied.

Proof. Note first that

$$\sup_{0<\lambda\leq 1}\frac{\lambda}{1-|\phi(\lambda)|}=\sup_{0<\lambda\leq 1}\frac{\lambda(\lambda+\zeta)^k}{(\lambda+\zeta)^k-\zeta^k}\leq \sup_{0<\lambda\leq 1}\frac{(\lambda+\zeta)^k}{k\zeta^{k-1}}<\infty,$$

so the first part of (3.1) certainly holds. For $n, \ell \in \mathbb{N}$ and $\lambda > 0$ we have that

$$\left|\frac{d^n}{d\lambda^n}\phi(\lambda)^\ell\right| = \zeta^{k\ell}\frac{(k\ell+n-1)!}{(k\ell-1)!|\lambda+\zeta|^{k\ell+n}}.$$

Given $\lambda > 0$ let $z = (\lambda + \zeta)/\zeta$. Then z > 1 and

$$\sum_{\ell=1}^{\infty} \left| \frac{d^n}{d\lambda^n} \phi(\lambda)^{\ell} \right| \le \frac{1}{\zeta^n} \sum_{\ell=1}^{\infty} \frac{(k\ell+n-1)!}{(k\ell-1)! z^{k\ell+n}} \le \frac{1}{\zeta^n} \sum_{\ell=1}^{\infty} \frac{(\ell+n-1)!}{(\ell-1)! z^{\ell+n}}$$

Since

$$\sum_{\ell=1}^{\infty} \frac{(\ell+n-1)!}{(\ell-1)! z^{\ell+n}} = \sum_{\ell=1}^{\infty} (-1)^n \frac{d^n}{dz^n} \frac{1}{z^\ell} = \frac{d^n}{dz^n} \left(\frac{1}{z-1}\right) = \frac{n!}{(z-1)^{n+1}}$$

and $z - 1 = \lambda/\zeta$, we obtain that

$$\sup_{n\in\mathbb{N}}\sup_{\lambda>0}\frac{\lambda^{n+1}}{n!}\sum_{\ell=1}^{\infty}\left|\frac{d^n}{d\lambda^n}\phi(\lambda)^{\ell}\right|\leq\zeta,$$

and hence ϕ also satisfies the second part of (3.1), as required.

4. Asymptotic behaviour

We now turn to the asymptotic behaviour of solutions to our system (1.2). For this we require, in addition to our earlier assumptions (A1) and (A2), three further assumptions. Recall that $\Omega_{\phi} = \{\lambda \in \mathbb{C} \setminus \sigma(A_0) : |\phi(\lambda)| = 1\}$, where ϕ is the characteristic function of our system.

Assumptions 4.1. We introduce the further assumptions that

(A3)
$$\sigma(A_0) \subset \mathbb{C}_-$$

(A4) $0 \in \Omega_{\phi} \subset \mathbb{C}_{-} \cup \{0\} \text{ and } \phi'(0) \neq 0,$

(A5)
$$\sup_{t\geq 0} \|T(t)\| < \infty,$$

where T is the semigroup generated by A.

Remark 4.2. Differentiating the identity in assumption (A2) gives

$$-A_1 R(\lambda, A_0)^2 A_1 = \phi'(\lambda) A_1, \quad \lambda \in \mathbb{C} \setminus \sigma(A_0).$$

In particular, if (A3) holds then $-A_1A_0^{-2}A_1 = \phi'(0)A_1$, and now assumption (A4) implies that $A_1A_0^{-1}$ restricts to an isomorphism from $\operatorname{Ran}(A_0^{-1}A_1)$ onto $\operatorname{Ran}(A_1)$.

In what follows we write L for the inverse of this isomorphism appearing in Remark 4.2, so that L maps $\operatorname{Ran}(A_1)$ isomorphically onto $\operatorname{Ran}(A_0^{-1}A_1)$. Moreover, having fixed $1 \leq p \leq \infty$ and $m \in \mathbb{N}$, we let

(4.1)
$$Y = \left\{ x_0 \in X : \lim_{t \to \infty} x(t) \text{ exists} \right\},$$

where x(t), $t \ge 0$, is the solution of (1.2) with initial condition $x(0) = x_0$. Furthermore, we denote the right-shift operator on X by S, so that $Sx = (x_{k-1})_{k\in\mathbb{Z}}$ for all $x = (x_k)_{k\in\mathbb{Z}} \in X$. Recall finally that n_{ϕ} denotes the resolvent growth parameter of our system; see Remark 2.7. The aim in this section is to prove the following theorem.

Theorem 4.3. Let $1 \le p \le \infty$, $m \in \mathbb{N}$ and assume that (A1)–(A5) hold. Define the operator $M \in \mathcal{B}(X)$ by $M(x_k) = (A_1 A_0^{-1} x_k)$, and let the operator L and the space Y be defined as above.

- (a) We have Y = X if and only if 1
 (i) If 1 0</sub> ∈ X.
 - (ii) If p = 1 and $x_0 \in X$ then $x_0 \in Y$ if and only if

(4.2)
$$\left\|\frac{1}{n}\sum_{k=1}^{n}\phi(0)^{k}S^{k}Mx_{0}\right\| \to 0, \quad n \to \infty,$$

and if this holds then $x(t) \to 0$ as $t \to \infty$.

(iii) If $p = \infty$ and $x_0 \in X$ then $x_0 \in Y$ if and only if there exists $y_0 \in \operatorname{Ran}(A_1)$ such that for $y = (\phi(0)^k y_0)$ we have

(4.3)
$$\left\|\frac{1}{n}\sum_{k=1}^{n}\phi(0)^{k}S^{k}Mx_{0}-y\right\| \to 0, \quad n \to \infty,$$

and if this holds then $x(t) \to z$ as $t \to \infty$, where $z = (\phi(0)^k L y_0)$. (b) Let n_{ϕ} be the resolvent growth parameter of the system.

(i) If $1 \le p < \infty$ and the decay in (4.2) is like $O(n^{-1})$ as $n \to \infty$ then

(4.4)
$$||x(t)|| = O\left(\left(\frac{(\log t)^{|1-2/p|}}{t}\right)^{1/n_{\phi}}\right), \quad t \to \infty.$$

(ii) If $p = \infty$ and the decay in (4.3) is like $O(n^{-1})$ as $n \to \infty$ then

$$\|x(t) - z\| = O\left(\left(\frac{\log t}{t}\right)^{1/n_{\phi}}\right), \quad t \to \infty.$$

(c) For $1 \le p \le \infty$ and all $x_0 \in X$ we have

$$\|\dot{x}(t)\| = O\left(\left(\frac{(\log t)^{|1-2/p|}}{t}\right)^{1/n_{\phi}}\right), \quad t \to \infty.$$

The proof of Theorem 4.3 is based on a number of general results. Given a C_0 -semigroup T on a complex Banach space X, let

(4.5)
$$Y = \left\{ x \in X : \lim_{t \to \infty} T(t)x \text{ exists} \right\},$$

noting that this notation is consistent with (4.1).

Proposition 4.4. Let T be a uniformly bounded C_0 -semigroup on a complex Banach space X and suppose that the generator A of T satisfies $\sigma(A) \cap i\mathbb{R} = \{0\}$. Then the set Y defined in (4.5) satisfies $Y = X_0 \oplus X_1$, where $X_0 = \text{Ker}(A)$ and X_1 denotes the closure of Ran(A). Moreover, if $x \in Y$ and $T(t)x \to y$ as $t \to \infty$, then y = Px, where $P \in \mathcal{B}(Y)$ is the projection onto X_0 along X_1 .

Proof. If $x \in X_0$, then T(t)x = x for all $t \ge 0$ and hence $x \in Y$. Thus $X_0 \subset Y$. Now define the function $f \in L^1(\mathbb{R}_+)$ by $f(t) = (t-1)e^{-t}$, then the Laplace transform F of f is given by

$$F(\lambda) = -\frac{\lambda}{(1+\lambda)^2}, \quad \operatorname{Re} \lambda \ge 0,$$

and we can define the operator $Q \in \mathcal{B}(X)$ by

$$Qx = \int_0^\infty f(t)T(t)x \,\mathrm{d}t, \quad x \in X,$$

noting that $Q = AR(1, A)^2$. Since F vanishes on the set $\sigma(A) \cap i\mathbb{R} = \{0\}$ and since singleton sets are of spectral synthesis, it follows from the Katznelson-Tzafriri theorem [20, Theorem 3.2] that $||T(t)Q|| \to 0$ as $t \to \infty$, and hence $\operatorname{Ran}(Q) \subset Y$. A simple argument shows that $\operatorname{Ran}(Q) = \operatorname{Ran}(A) \cap D(A)$, where D(A) denotes the domain of A. In particular, $\operatorname{Ran}(Q)$ is dense in X_1 , so by uniform boundedness of T we obtain that $X_1 \subset Y$. Thus $X_0 + X_1 \subset Y$. Next we show that the sum is direct. Suppose that $x \in X_0 \cap X_1$. Since $x \in X_0$, ||x|| = ||T(t)x|| for all $t \ge 0$. On the other hand, since $x \in X_1$, $||T(t)x|| \to 0$ as $t \to \infty$. It follows that x = 0, and hence $X_0 \cap X_1 = \{0\}$, as required.

Now suppose that $x \in Y$. Then there exists $y \in X$ such that $y = \lim_{s\to\infty} T(s)x$. For $t \ge 0$ we have $T(t)y = \lim_{s\to\infty} T(t)T(s)x = y$, which implies that $y \in X_0$. Let z = x - y. Then

(4.6)
$$||T(t)z|| = ||T(t)x - y|| \to 0, \quad t \to \infty.$$

Suppose that $z \in X \setminus X_1$. It follows from a standard application of the Hahn-Banach theorem that there exists $\phi \in X^*$ such that $\langle z, \phi \rangle = 1$ and $\phi|_{X_1} = 0$. In particular, $\phi|_{\text{Ran}(A)} = 0$ and hence $\phi \in \text{Ker}(A')$. It follows that $T(t)'\phi = \phi$ for all $t \geq 0$, and therefore

$$\langle T(t)z,\phi\rangle = \langle z,T(t)'\phi\rangle = \langle z,\phi\rangle = 1, \quad t \ge 0.$$

This contradicts (4.6), so $z \in X_1$. Thus $x = y + z \in X_0 + X_1$ and consequently $Y = X_0 \oplus X_1$. Furthermore,

$$\lim_{t \to \infty} T(t)x = y = Px,$$

where $P: Y \to Y$ is the projection onto X_0 along X_1 . Since both X_0 and X_1 are closed, P is bounded and the proof is complete.

Remark 4.5. Note that if $x \in Y$ and $T(t)x \to y$ as $t \to \infty$, then

(4.7)
$$\lim_{t \to \infty} \frac{1}{t} \int_0^t T(s) x \, \mathrm{d}s = y.$$

It is well known that the set of $x \in X$ for which (4.7) holds is given by $X_0 \oplus X_1$, where X_0 and X_1 are as in Proposition 4.4. The result is therefore Tauberian in flavour, showing as it does that (4.7) implies $\lim_{t\to\infty} T(t)x = y$. Note that this ergodic approach also leads to the further equivalent characterisation of the set Y as

$$Y = \left\{ x \in X : \lim_{\lambda \to 0+} R(\lambda, A) x \text{ exists} \right\}.$$

For details of the above results see for instance [2, Section 4.3], and for a more general result related to Proposition 4.4 see [2, Theorem 5.5.4].

As observed in Remark 4.5, the characterisation of the set Y obtained in Proposition 4.6 can be interpreted as the set of mean ergodic vectors of the semigroup T. We now collect some important facts about the set of mean ergodic vectors of certain bounded linear operators, which will then be used to obtain descriptions of the set Y in the case where the semigroup T has a suitable bounded generator.

Proposition 4.6. Let X be the dual space of a complex Banach space X_* and consider the operator A = B - C, where $B, C \in \mathcal{B}(X)$. Suppose that C is invertible and that the operator $Q = C^{-1}B$ is power-bounded and satisfies Q = U' for some $U \in \mathcal{B}(X_*)$. Let $Z = X_0 \oplus X_1$, where $X_0 = \text{Ker}(A)$ and X_1 denotes the closure of Ran(A), and let $Z_0 = X_0 \oplus \text{Ran}(A)$. Then, given $x \in X$, we have $x \in Z$ if and only if there exists $y \in X_0$ such that

(4.8)
$$\left\|\frac{1}{n}\sum_{k=1}^{n}Q^{k}C^{-1}(x-y)\right\| \to 0, \quad n \to \infty.$$

Furthermore, Z_0 consists of all those $x \in Z$ for which the convergence in (4.8) is like $O(n^{-1})$ as $n \to \infty$.

Proof. Note first that $X_0 = \text{Fix}(Q)$ and that $\text{Ran}(A) = \{Cx : x \in \text{Ran}(I - Q)\}$. Hence $X_1 = \{Cx : x \in X_2\}$, where X_2 denotes the closure of Ran(I - Q). By [13, Theorem 1.3 of Section 2.1] and power-boundedness of Q,

$$X_{2} = \left\{ x \in X : \lim_{n \to \infty} \frac{1}{n} \sum_{k=1}^{n} Q^{k} x = 0 \right\}.$$

Hence, given $x \in X$, we have $x \in Z$ if and only if there exists $y \in X_0$ such that $x - y \in X_1$, which is equivalent to $C^{-1}(x - y) \in X_2$. This shows that (4.8) holds. The characterisation of Z_0 follows similarly using [14, Theorem 5], and it is here that the duality assumptions are needed.

Remark 4.7. The characterisation of the space Z in fact holds on arbitrary complex Banach spaces and when the condition of power-boundedness is replaced by the weaker assumptions that Q is *Cesàro bounded*, which is to say

$$\sup_{n\geq 1} \left\| \frac{1}{n} \sum_{k=1}^{n} Q^k \right\| < \infty,$$

and that $||Q^n x|| = o(n)$ as $n \to \infty$ for each $x \in X$. A characterisation of Z_0 in this more general setting can be deduced from the results in [14].

We now seek to combine Propositions 4.4 and 4.6 in to obtain a characterisation of the set Y defined in (4.5) when the semigroup T is generated by a suitable bounded operator A. In particular, we hope to deduce from Proposition 4.6 a statement about the *rate* at which certain semigroup orbits converge to a limit. This requires two abstract results.

Theorem 4.8. Let X be a complex Banach space and suppose T is a uniformly bounded C_0 -semigroup on X whose generator $A \in \mathcal{B}(X)$ satisfies $\sigma(A) \cap i\mathbb{R} = \{0\}$. Suppose that

$$||R(is, A)|| \le m(|s|), \quad 0 < |s| \le 1,$$

for some continuous non-increasing function $m: (0,1] \rightarrow [1,\infty)$. Then for any $c \in (0,1)$

(4.9)
$$||AT(t)|| = O\left(m_{\log}^{-1}(ct)\right), \quad t \to \infty,$$

where m_{\log}^{-1} is the inverse function of the map $m_{\log}: (0,1] \to (0,\infty)$ given by

(4.10)
$$m_{\log}(r) = m(r) \log\left(1 + \frac{m(r)}{r}\right), \quad 0 < r \le 1.$$

Proof. The result is a consequence of [7, Corollary 2.12]. Indeed, since T is norm-continuous and hence differentiable, it follows from [5, Theorem 5.6] that the non-analytic growth bound $\zeta(T)$ of T satisfies $\zeta(T) = -\infty$, and in particular $\zeta(T) < 0$. Thus [7, Corollary 2.12] shows that

$$||T(t)AR(1,A)|| = O\left(m_{\log}^{-1}(ct)\right), \quad t \to \infty,$$

and (4.9) follows by applying the bounded linear operator I - A.

- **Remark 4.9.** (a) The unquantified version of the above result, namely that $||AT(t)|| \to 0$ as $t \to \infty$ when T is bounded and $\sigma(A) \cap i\mathbb{R} = \{0\}$ is shown in the more general setting of eventually differentiable semigroups in [3, Theorem 3.10]. The result can also be deduced from the Katznelson-Tzafriri theorem. Indeed, it was shown that in the proof of Proposition 4.4 that $||T(t)AR(1,A)^2|| \to 0$ as $t \to \infty$, from which the claim follows easily; see also [6, Remark 6.3]
- (b) As is shown in [7, Corollary 2.12], the result in fact holds more generally for bounded semigroups whose generator is not necessarily bounded. When X is a Hilbert space, it follows from [5, Theorem 5.4] that the condition $\zeta(T) < 0$ can be replaced by the condition $\sup_{|s|\geq 1} ||R(is, A)|| < \infty$. A more direct way of showing that $\zeta(T) = -\infty$ when the semigroup T has bounded generator is to observe that in this case

$$T(t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} A^n, \quad t \ge 0,$$

with the sum converging in operator norm. In particular, T itself extends to an analytic and exponentially bounded operator-valued family on a sector containing $(0, \infty)$, and the claim follows from the definition of $\zeta(T)$. See [5] for details on the non-analytic growth bound $\zeta(T)$.

(c) Theorem 4.8 can also be deduced from [16, Proposition 3.1] with the function $M : [1, \infty) \to (0, \infty)$ taken to be constant, since in this case both the M_{\log}^{-1} -term and the t^{-1} -term are dominated by the m_{\log}^{-1} -term.

The next result is a special case of Theorem 4.8 dealing with the case of polynomial resolvent growth, and it contains a sharper estimate in the Hilbert space setting.

Theorem 4.10. Let X be a complex Banach space and suppose T is a uniformly bounded C_0 -semigroup on X whose generator $A \in \mathcal{B}(X)$ satisfies $\sigma(A) \cap i\mathbb{R} = \{0\}$ and $||R(is, A)|| = O(|s|^{-\alpha})$ as $|s| \to 0$ for some $\alpha \ge 1$. Then

$$||AT(t)|| = O\left(\left(\frac{\log t}{t}\right)^{1/\alpha}\right), \quad t \to \infty.$$

Moreover, if X is a Hilbert space then the logarithm can be omitted.

Proof. The first statement is a consequence of Theorem 4.8 with the choice $m(r) = Cr^{-\alpha}$, $0 < r \leq 1$, for a suitable constant $C \geq 1$, since in this case

$$m_{\log}^{-1}(ct) = O\left(\left(\frac{\log t}{t}\right)^{1/\alpha}\right), \quad t \to \infty,$$

for all $c \in (0, 1)$. The second statement is a direct consequence of [6, Theorem 7.6] and boundedness of the generator A.

Remark 4.11. It follows from [6, Corollary 6.11] that if in Theorem 4.10 we in fact have $||R(is, A)|| \approx |s|^{-\alpha}$ as $|s| \to 0$, then there exists a constant c > 0 such that

$$||AT(t)|| \ge \frac{c}{t^{1/\alpha}}, \quad t \ge 1,$$

provided that $||sR(is, A)|| \to \infty$ as $|s| \to 0$. Since the resolvent growth parameter n_{ϕ} is always strictly greater than 1 by Lemma 2.6, it follows from Proposition 2.5 that the latter condition is always satisfied in Theorem 4.3.

We now combine the previous results in this section in order to obtain a general result about the asymptotics of semigroups whose generators are suitable bounded operators. Recall from (4.5) that

$$Y = \left\{ x \in X : \lim_{t \to \infty} T(t)x \text{ exists} \right\}.$$

Theorem 4.12. Let T be a uniformly bounded C_0 -semigroup on a space Xwhich is the dual of a complex Banach space X_* . Suppose that the generator A of T satisfies A = B - C, where $B, C \in \mathcal{B}(X)$, C is invertible, the operator $Q = C^{-1}B$ is power-bounded and satisfies Q = U' for some $U \in \mathcal{B}(X_*)$. Suppose furthermore that $\sigma(A) \cap i\mathbb{R} = \{0\}$ and that

$$|R(is, A)|| \le m(|s|), \quad 0 < |s| \le 1,$$

for some continuous non-increasing function $m: (0,1] \to [1,\infty)$.

Then, given $x \in X$, we have $x \in Y$ if and only if there exists $y \in Fix(Q)$ such that

(4.11)
$$\left\|\frac{1}{n}\sum_{k=1}^{n}Q^{k}C^{-1}(x-y)\right\| \to 0, \quad n \to \infty,$$

and if (4.11) holds then $T(t)x \to y$ as $t \to \infty$. Moreover, if the convergence in (4.11) is like $O(n^{-1})$ as $n \to \infty$, then for each $c \in (0, 1)$

(4.12)
$$||T(t)x - y|| = O(m_{\log}^{-1}(cn)), \quad t \to \infty$$

where m_{\log} is as defined in (4.10). In particular, if $||R(is, A)|| = O(|s|^{-\alpha})$ for some $\alpha \ge 1$ as $|s| \to 0$, then

(4.13)
$$||T(t)x - y|| = O\left(\left(\frac{\log t}{t}\right)^{1/\alpha}\right), \quad t \to \infty,$$

and the logarithm can be omitted if X is a Hilbert space.

Proof. The description of the set Y follows immediately by combining Propositions 4.4 and 4.6. If the convergence in (4.11) is like $O(n^{-1})$ as $n \to \infty$, then by Proposition 4.6 we have that x - y = Az for some $z \in X$, and hence

$$T(t)x - y = T(t)(x - y) = T(t)Az, \quad t \ge 0.$$

Thus (4.12) follows from Theorem 4.8, and (4.13) and the statement after it follow from Theorem 4.10. $\hfill \Box$

Remark 4.13. By Remark 4.7 it is possible to obtain the unquantified statements of Theorem 4.12 under weaker assumptions.

We now come to the proof of Theorem 4.3.

Proof of Theorem 4.3. Note first that X has a predual X_* for each choice of p. Indeed, if $1 then X is the dual of <math>X_* = \ell^q(\mathbb{Z}; \mathbb{C}^m)$, where q is the Hölder conjugate of p, and if p = 1 then X is the dual of $X_* = c_0(\mathbb{Z}; \mathbb{C}^m)$, the space of \mathbb{C}^m -valued sequences $(x_k)_{k\in\mathbb{Z}}$ such that $|x_k| \to 0$ as $k \to \pm\infty$. Since $\sigma(A_0)$ is contained in the open left half-plane by assumption (A3), A_0 is invertible and hence so is M_0 . Moreover, $\sigma(A) \cap i\mathbb{R} = \{0\}$ by assumption (A4) and Theorem 2.3, and by Proposition 2.5 we have that $||R(is, A)|| \approx$ $|s|^{-n_{\phi}}$ as $|s| \to 0$. For j = 0, 1 let $M_j \in \mathcal{B}(X)$ denote the operator given by $M_i(x_k) = (A_i x_k)$, noting that both M_0 and M_1 commute with the right-shift operator S on X. Moreover, let $M, N \in \mathcal{B}(X)$ be given by $M = M_1(-M_0)^{-1}$ and $N = (-M_0)^{-1} M_1$ so that $Mx = (A_1 R_0 x_k)_{k \in \mathbb{Z}}$ and $Nx = (R_0 A_1 x_k)_{k \in \mathbb{Z}}$ for all $x = (x_k)_{k \in \mathbb{Z}} \in X$. Then A = B - C with $B = SM_1$ and $C = -M_0$. Let $Q = C^{-1}B$. Then Q = SN and in particular Q = U', where $U \in \mathcal{B}(X_*)$ is given by $Ux = (A_1^T R_0^T x_{k+1})_{k \in \mathbb{Z}}$. Moreover, for $n \ge 1$, it follows from our assumption on the matrices A_0 , A_1 that $N^n = \phi(0)^{n-1}N$, and hence $Q^n = \phi(0)^{n-1} S^n N$. Note also that $|\phi(0)| = 1$ since $0 \in \Omega_{\phi}$. In particular,

$$||Q^n|| = ||S^n N|| \le ||N||, \quad n \ge 1,$$

so Q is power-bounded. Moreover,

$$Q^n C^{-1} = (-M_0)^{-1} \phi(0)^{n-1} S^n M, \quad n \ge 1.$$

Suppose that $1 \le p < \infty$ and let $x_0 \in X$. Then $\operatorname{Ker}(A) = \operatorname{Fix}(Q) = \{0\}$, and since M_0 is an isomorphism and $|\phi(0)| = 1$, it follows from Theorem 4.12 that $x_0 \in Y$ if and only if (4.2) holds, and that $x(t) \to 0$ as $t \to \infty$ whenever this is the case. If $p = \infty$ then by Theorem 2.3 any $z \in \text{Ker}(A)$ has the form $z = (\phi(0)^k z_0)$ for some $z_0 \in \operatorname{Ran}(A_0^{-1}A_1)$. For such a $z \in \operatorname{Ker}(A)$ let y = Mz. Then $y = (\phi(0)^k y_0)$, where $y_0 = A_1 A_0^{-1} z_0$. In particular, $y_0 \in \operatorname{Ran}(A_1)$ and $z_0 = Ly_0$. Moreover, $\phi(0)^n S^n M z = y$ for all $n \ge 1$ and hence, given $x_0 \in X$, Theorem 4.12 implies that $x_0 \in Y$ if and only if (4.3) holds for some $y_0 \in \operatorname{Ran}(A_1)$, and that $x(t) \to z$ as $t \to \infty$ whenever this is the case. When p = 1 and when $p = \infty$, it is straightforward to see that (4.2) and, respectively, (4.3) are not satisfied for all $x_0 \in X$, whereas (4.2) does hold for all $x_0 \in X$ when 1 , as can be seen by considering thedense subspace of finitely supported sequences. Thus Y = X if and only if 1 and part (a) is established. For part (b) note that (ii) followsimmediately from Theorem 4.12, while if $1 \leq p < \infty$ and convergence in (4.2) is like $O(n^{-1})$ as $n \to \infty$, Theorem 4.12 shows that

$$||x(t)|| = O\left(\left(\frac{\log t}{t}\right)^{1/n_{\phi}}\right), \quad t \to \infty,$$

and that the logarithm can be omitted when p = 2. The estimate in (4.4) now follows by appealing to the Riesz-Thorin theorem [11, Theorem 9.3.3]

to interpolate these bounds for $1 and <math>2 . Part (c) follows similarly using the fact that <math>\dot{x}(t) = AT(t)x_0$ for all $x_0 \in X$ and $t \ge 0$. \Box

- **Remark 4.14.** (a) The statement in part (a)(i) can also be deduced from the well-known Arendt-Batty-Lyubich-Vũ theorem; see [1, 15]. Indeed, the semigroup T is uniformly bounded by assumption (A5), and by Theorem 2.3 the other assumptions ensure that the generator A of Thas no residual spectrum on the imaginary axis. This argument can be extended to obtain strong stability of T also in the case where Ω_{ϕ} meets the imaginary axis in several (but necessarily at most finitely many) points.
- (b) It follows from Remark 4.11 together with Lemma 2.6 and an application of the uniform boundedness principle that the rates in Theorem 4.3 are optimal when p = 2 and worse than optimal by at most a logarithmic term when $p \neq 2$. We expect that the quantified statements in Theorem 4.3 remain true without the logarithms even when $p \neq 2$, but we leave it as an open problem whether this is indeed the case; see also Remark 5.2(a) and Theorem 6.1 below.

5. The platoon model

In this section we study a linearised model of an infinitely long platoon of vehicles. The objective is to drive the solution of the system to a configuration in which all of the vehicles are moving at a given constant velocity $v \in \mathbb{C}$ and the separation between the vehicles k and k-1 is equal to $c_k \in \mathbb{C}$, $k \in \mathbb{Z}$. For $k \in \mathbb{Z}$ and $t \geq 0$, we write $d_k(t)$ for the separation between vehicles k and k-1 at time t, $v_k(t)$ for the velocity of vehicle k at time t and $a_k(t)$ for the acceleration of vehicle k at time t. Furthermore, we let $y_k(t) = c_k - d_k(t)$ denote the deviation of the actual separation from the target separation of vehicles k and k-1 at time t, and we similarly let $w_k(t) = v_k(t) - v$ stand for the excess velocity of vehicle k at time t. Note in particular that, as the variables are allowed to be complex, they can be used to describe the dynamics of the vehicles in the complex plane and not just along a straight line. On the other hand, if all the variables are constrained to be real, the same model can be used to study the behaviour of an infinitely long chain of vehicles.

As the basis of our study we consider a linear model which has been used to study infinitely long chains of cars on a highway in [17, 18, 21], namely

(5.1)
$$\begin{pmatrix} \dot{y}_k(t) \\ \dot{w}_k(t) \\ \dot{a}_k(t) \end{pmatrix} = \begin{pmatrix} w_k(t) - w_{k-1}(t) \\ a_k(t) \\ -\tau^{-1}a_k(t) + \tau^{-1}u_k(t) \end{pmatrix}, \quad k \in \mathbb{Z}, \ t \ge 0,$$

where $\tau > 0$ is a parameter and $u_k(t)$ is the control input of vehicle k. In the above references the model (5.1) was studied on the space $X = \ell^2(\mathbb{C}^3)$, and it has in particular been shown that the system is not exponentially stabilisable [12, 21] but that strong stability can be achieved [8, 12]. In this paper we study model (5.1) on the spaces $X = \ell^p(\mathbb{C}^3)$ for $1 \le p \le \infty$, and in particular we include the case $p = \infty$ argued in [12] to be the most realistic. We begin by rewriting the problem in the form of (1.1) for the state vectors

$$x_k(t) = \begin{pmatrix} y_k(t) \\ w_k(t) \\ a_k(t) \end{pmatrix}, \quad k \in \mathbb{Z}, \ t \ge 0,$$

by applying an identical state feedback

$$u_k(t) = \beta_1 y_k(t) + \beta_2 w_k(t) + \beta_3 a_k(t), \quad k \in \mathbb{Z}, \ t \ge 0,$$

to each of the vehicles, where $\beta_1, \beta_2, \beta_3 \in \mathbb{C}$ are constants. This control law requires that the state vectors $x_k(t)$ are known and available for feedback. Equations (5.1) can then be written in the form (1.1) with matrices

$$A_0 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -\alpha_0 & -\alpha_1 & -\alpha_2 \end{pmatrix} \quad \text{and} \quad A_1 = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

where $\alpha_0 = -\beta_1/\tau$, $\alpha_1 = -\beta_2/\tau$, and $\alpha_2 = (1-\beta_3)/\tau$ can be freely assigned by choosing appropriate feedback parameters $\beta_1, \beta_2, \beta_3 \in \mathbb{C}$. This in turn allows us to choose the eigenvalues of the matrix A_1 . Since rank $A_1 = 1$, we know from Remark 2.2 that conditions (A1) and (A2) of Assumption 2.1 are satisfied, and the characteristic function ϕ is given by the formula

$$\phi(\lambda) = \frac{\alpha_0}{p(\lambda)}, \qquad \lambda \in \mathbb{C} \setminus \sigma(A_0),$$

where $p(\lambda) = \lambda^3 + \alpha_2 \lambda^2 + \alpha_1 \lambda + \alpha_0$ is the characteristic polynomial of A_0 . Note that $\phi(0) = 1$ and hence $0 \in \sigma(A)$ by Theorem 2.3. It follows that the platoon system cannot be stabilised exponentially. Our main goal is to choose the parameters $\alpha_0, \alpha_1, \alpha_2 \in \mathbb{C}$ in such a way that the platoon system achieves good stability properties. The simplest possible characteristic polynomial is $p(\lambda) = (\lambda - \lambda_0)^3$ corresponding to the choices $\alpha_0 = -\lambda_0^3, \alpha_1 = 3\lambda_0^2$, and $\alpha_2 = -3\lambda_0$ for a fixed $\lambda_0 \in \mathbb{C}$. In this case

$$\Omega_{\phi} = \big\{ \lambda \in \mathbb{C} : |\lambda - \lambda_0| = |\lambda_0| \big\},\$$

so in order for conditions (A3) and (A4) of Assumptions 4.1 to be satisfied, so that $\sigma(A) \subset \mathbb{C}_- \cup \{0\}$, it is necessary to choose $\lambda_0 = -\zeta$ for some $\zeta > 0$. It is possible in principle to derive more general necessary geometric conditions on the roots of p which ensure that (A3) and (A4) are satisfied. We restrict ourselves here to exhibiting, in Figure 1, several examples of level sets Ω_{ϕ} and spectra $\sigma(A)$ for different choices of the parameters $\alpha_0, \alpha_1, \alpha_2 \in \mathbb{C}_-$.

We now consider the Cauchy problem (1.2) for the platoon problem with a characteristic function ϕ having just one pole. Our main asymptotic result is a consequence of the general results proved in the preceding sections. Recall that for $1 \leq p \leq \infty$ we denote the set of initial states $x(0) = x_0$ leading to convergent solutions x(t) of the platoon system by

$$Y = \left\{ x_0 \in X : \lim_{t \to \infty} x(t) \text{ exists} \right\}.$$

Theorem 5.1. Let $1 \le p \le \infty$ and consider the platoon model with the choices $\alpha_0 = \zeta^3$, $\alpha_1 = 3\zeta^2$ and $\alpha_2 = 3\zeta$, where $\zeta > 0$ is a fixed real number.

- (a) We have Y = X if and only if 1 . More specifically:
 - (i) If 1 then <math>Y = X and $x(t) \to 0$ for all $x_0 \in X$.




FIGURE 1. The level set Ω_{ϕ} and $\sigma(A_0)$ for various choices of A_0 .

(ii) If p = 1 and $x_0 \in X$ then $x_0 \in Y$ if and only if the vector $y_0 = (y_k(0))_{k \in \mathbb{Z}} \in \ell^1(\mathbb{Z})$ of initial deviations is such that

(5.2)
$$\left\|\frac{1}{n}\sum_{k=1}^{n}S^{k}y_{0}\right\|_{\ell^{1}(\mathbb{Z})}\to 0, \quad n\to\infty,$$

and if this holds then $x(t) \to 0$ as $t \to \infty$.

(iii) If $p = \infty$ and $x_0 \in X$ then $x_0 \in Y$ if and only if there exists $c \in \mathbb{C}$ such that for $y = (\dots, c, c, c, \dots)$ we have

(5.3)
$$\left\| \frac{1}{n} \sum_{k=1}^{n} S^{k} y_{0} - y \right\|_{\ell^{\infty}(\mathbb{Z})} \to 0, \quad n \to \infty,$$

and if this holds then $x(t) \to z$ as $t \to \infty$, where

$$z = \left(\dots, \begin{pmatrix} c \\ -\zeta c/3 \\ 0 \end{pmatrix}, \begin{pmatrix} c \\ -\zeta c/3 \\ 0 \end{pmatrix}, \begin{pmatrix} c \\ -\zeta c/3 \\ 0 \end{pmatrix}, \dots \right).$$

(b) (i) If $1 \le p < \infty$ and the decay in (5.2) is like $O(n^{-1})$ as $n \to \infty$ then

$$||x(t)|| = O\left(\left(\frac{(\log t)^{|1-2/p|}}{t}\right)^{1/2}\right), \quad t \to \infty.$$

(i) If $p = \infty$ and the decay in (5.3) is like $O(n^{-1})$ as $n \to \infty$ then

$$||x(t) - z|| = O\left(\left(\frac{\log t}{t}\right)^{1/2}\right), \quad t \to \infty.$$

(c) For $1 \le p \le \infty$ and all $x_0 \in X$ we have

$$\|\dot{x}(t)\| = O\left(\left(\frac{(\log t)^{|1-2/p|}}{t}\right)^{1/2}\right), \quad t \to \infty.$$

Proof. Note that (A1) holds and that (A2) is satisfied for the function

$$\phi(\lambda) = \frac{\zeta^3}{(\lambda+\zeta)^3}, \quad \lambda \neq -\zeta.$$

As above, we have $\sigma(A_0) = \{-\zeta\}$, so that (A3) holds, and since

$$\Omega_{\phi} = \{\lambda \in \mathbb{C} : |\lambda + \zeta| = \zeta\},\$$

we see that (A4) holds as well. Furthermore, (A5) holds by Lemma 3.2 and the first part of Theorem 3.1. A simple calculation based on the ideas used in Lemma 2.6 shows that $n_{\phi} = 2$. Noting that $\phi(0) = 1$ and that

$$A_1(-A_0)^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \qquad (-A_0)^{-1}A_1 = \begin{pmatrix} 0 & -3/\zeta & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

the result follows from Theorem 4.3.

- **Remark 5.2.** (a) We do not know whether the logarithms in the decay estimates of Theorem 5.1 are needed when $p \neq 2$. We suspect not; see also Remark 4.14(b) above and Theorem 6.1 below.
- (b) It follows from straightforward estimates that the semigroup T generated by the operator A in the platoon model is in general not contractive, even when p = 2.
- (c) Note that the above analysis can also be used in the setting considered in [18], where the objective of attaining given target separations as $t \to \infty$ is replaced by the objective that the separations should approach $c_k + hv_k(t)$, where h > 0 and $c_k \in \mathbb{C}$ are constants and $v_k(t)$ is the velocity of vehicle $k \in \mathbb{Z}$ at time $t \ge 0$.

6. The robot rendezvous problem

We now return to the robot rendezvous problem, which corresponds in the general setting of (1.1) to the choices m = 1, $A_0 = -1$ and $A_1 = 1$. In particular, the Banach space we are working in is $X = \ell^p(\mathbb{Z})$, where $1 \leq p \leq \infty$. The following result, which can be viewed as an extension of the results in [9], is in large part a consequence of Theorem 4.3 but with slightly sharper estimates on the rates of decay. For $1 \leq p \leq \infty$, we once again use the notation

$$Y = \left\{ x_0 \in X : \lim_{t \to \infty} x(t) \text{ exists} \right\},\$$

where $x(t), t \ge 0$, now denotes the solution of the robot rendezvous problem with initial condition $x(0) = x_0$.

Theorem 6.1. Let $1 \le p \le \infty$ and consider the robot rendezvous problem.

- (a) We have Y = X if and only if 1 . More specifically:
 - (i) If 1 then <math>Y = X and $x(t) \to 0$ for all $x_0 \in X$.
 - (ii) If p = 1 and $x_0 \in X$ then $x_0 \in Y$ if and only if

(6.1)
$$\left\|\frac{1}{n}\sum_{k=1}^{n}S^{k}x_{0}\right\| \to 0, \quad n \to \infty,$$

and if this holds then $x(t) \to 0$ as $t \to \infty$.

(iii) If $p = \infty$ and $x_0 \in X$ then $x_0 \in Y$ if and only if there exists a constant sequence $z \in X$ such that

(6.2)
$$\left\|\frac{1}{n}\sum_{k=1}^{n}S^{k}x_{0}-z\right\| \to 0, \quad n \to \infty,$$

and if this holds then $x(t) \to z$ as $t \to \infty$.

(b) (i) If $1 \le p < \infty$ and the decay in (6.1) is like $O(n^{-1})$ as $n \to \infty$ then

(6.3)
$$||x(t)|| = O(t^{-1/2}), \quad t \to \infty.$$

(i) If $p = \infty$ and the decay in (6.2) is like $O(n^{-1})$ as $n \to \infty$ then

(6.4)
$$||x(t) - z|| = O(t^{-1/2}), \quad t \to \infty$$

(c) For $1 \le p \le \infty$ and all $x_0 \in X$ we have

(6.5)
$$\|\dot{x}(t)\| = O(t^{-1/2}), \quad t \to \infty.$$

Finally, the rate $t^{-1/2}$ in (6.3), (6.4) and (6.5) is optimal.

Proof. Note that (A1) holds and that (A2) is satisfied for the function

$$\phi(\lambda) = \frac{1}{\lambda+1}, \quad \lambda \neq -1.$$

We also have $\sigma(A_0) = \{-1\}$, so that (A3) holds, and since

$$\Omega_{\phi} = \{\lambda \in \mathbb{C} : |\lambda + 1| = 1\},\$$

we see that (A4) holds as well. Assumption (A5) again holds by Lemma 3.2 and Theorem 3.1, and indeed the second part of the latter result even shows that the semigroup is contractive. As in the proof of Theorem 5.1, a simple calculation shows that $n_{\phi} = 2$, so all of the statements follow from Theorem 4.3 except for the rates in equations (6.3), (6.4) and (6.5) and the final statement concerning optimality. The latter follows as in Remark 4.14(b). In order to obtain the sharper rates we require a better estimate on the asymptotic behaviour of ||AT(t)|| as $t \to \infty$ than is given in Theorem 4.10 for the general case.

For $t \ge 0$, let $y(t) \in \ell^1(\mathbb{Z})$ be the scalar-valued sequence whose k-th term is given by

$$y_k(t) = \frac{t^k}{k!} - \frac{t^{k+1}}{(k+1)!}, \quad k \ge 0,$$

and $y_k(t) = 0$ for k < 0. It is shown in the proof of [9, Theorem 3] that, given $x_0 \in X$,

$$AT(t)x_0 = e^{-t}(y(t) * z_0 - x_0), \quad t \ge 0,$$

where $z_0 = Sx_0$ with S being the right-shift. Then $||z_0|| = ||x_0||$ and it follows from Young's inequality that

$$||AT(t)x_0|| \le e^{-t} (1 + ||y(t)||_{\ell^1(\mathbb{Z})}) ||x_0||, \quad t \ge 0.$$

In particular,

$$||AT(t)|| \le e^{-t} (1 + ||y(t)||_{\ell^1(\mathbb{Z})}) \quad t \ge 0.$$

As explained in the proof of [9, Theorem 3], for each $t \ge 0$ there exists an integer $n(t) \ge 0$ such that $n(t) \le t \le n(t) + 1$ and

$$||y(t)||_{\ell^1(\mathbb{Z})} \le 2\frac{t^{n(t)}}{n(t)!}.$$

By Stirling's approximation,

$$n(t)! \ge \sqrt{2\pi n(t)} \left(\frac{n(t)}{e}\right)^{n(t)}, \quad t \ge 1.$$

Now straightforward estimates show that

(6.6)
$$||AT(t)|| \le \frac{C}{t^{1/2}}, \quad t \ge 2,$$

for some C > 0, and the result follows as in the proof of Theorem 4.3. \Box

Remark 6.2. (a) Note that in the above setting, the semigroup has an explicit representation. Indeed, if $T(t)(x_k) = (y_k(t))$ for $(x_k), (y_k) \in X$ and $t \ge 0$, then

$$y_k(t) = e^{-t} \sum_{n=0}^{\infty} \frac{t^n}{n!} x_{k-n}, \quad k \in \mathbb{Z}, \ t \ge 0.$$

In particular, an application of Young's inequality gives an alternative, more direct proof of the fact that the semigroup T is contractive in this case for $1 \le p \le \infty$; cf. Remark 5.2(b).

(b) The above proof can be refined to give an explicit constant C in (6.6). For instance, it is straightforward to show that the value

$$C = \frac{t_0^{1/2}}{e^{t_0}} + \left(\frac{2}{\pi}\right)^{1/2} \left(1 - \frac{1}{t_0}\right)^{-t_0 - 1/2}$$

gives the inequality for the range $t \ge t_0 > 1$. In particular, C = 4.705 works for $t \ge 2$, C = 2.191 works for $t \ge 100$, and as $t_0 \to \infty$ the value of the constant approaches $e(2/\pi)^{1/2} \approx 2.169$.

(c) For further discussion of the robot rendezvous problem and in particular its connection with the theory of Borel summability, see [9, 10].

We conclude by briefly considering an interesting generalisation of the robot rendezvous problem in which the original differential equations are replaced by

$$\dot{x}_k(t) = x_{k-1}(t) + \alpha_k x_k(t), \quad k \in \mathbb{Z}, \ t \ge 0,$$

where for each $k \in \mathbb{Z}$ either $\alpha_k = -1$ or $\operatorname{Re} \alpha_k < -1$. The original robot rendezvous problem corresponds to the choice $\alpha_k = -1$ for all $k \in \mathbb{Z}$. If we again let $X = \ell^p(\mathbb{C})$ for $1 \leq p \leq \infty$, we are led to consider the semigroup Tgenerated by the operator $A \in \mathcal{B}(X)$ given by $Ax = (x_{k-1} + \alpha_k x_k)_{k \in \mathbb{Z}}$ for all $(x_k)_{k \in \mathbb{Z}} \in X$. We restrict ourselves to stating a result about the decay of ||AT(t)|| as $t \to \infty$, which could be used to obtain statements about orbits and their derivatives as in Sections 4 and 5.

Theorem 6.3. In the modified robot rendezvous problem considered above, let $\Omega = \{\alpha_k : k \in \mathbb{Z}\}$ and suppose that $-1 \in \Omega$. Then, for all $c \in (0, 1)$,

$$||AT(t)|| = O\left(m_{\log}^{-1}(ct)\right), \ t \to \infty,$$

where $m: (0,1] \to [1,\infty)$ is defined by

(6.7)
$$m(r) = \sup_{r \le |s| \le 1} \frac{1}{\operatorname{dist}(is, \Omega) - 1}, \quad 0 < r \le 1,$$

and m_{\log} is as in Theorem 4.8.

Proof. Let $\lambda \in \mathbb{C}$ be such that $\operatorname{dist}(\lambda, \Omega) > 1$ and define $R(\lambda) \in \mathcal{B}(X)$ by

$$R(\lambda)x = \left(\sum_{\ell=0}^{\infty} \prod_{j=0}^{\ell} \frac{x_{k-\ell}}{\lambda - \alpha_{k-j}}\right)_{k \in \mathbb{Z}}$$

for all $x = (x_k)_{k \in \mathbb{Z}} \in X$. Straightforward computations show that $R(\lambda)(\lambda - A)x = (\lambda - A)R(\lambda)x = x$ for all $x \in X$, and hence $\lambda \notin \sigma(A)$ and $R(\lambda) = R(\lambda, A)$ for all $\lambda \in \mathbb{C}$ such that $\operatorname{dist}(\lambda, \Omega) > 1$. Furthermore, it is straightforward to verify that if $\operatorname{dist}(\lambda, \Omega) > 1$ and $\|x\| = 1$, then

$$||R(\lambda, A)x|| \le \sum_{\ell=0}^{\infty} \frac{1}{\operatorname{dist}(\lambda, \Omega)^{\ell+1}} = \frac{1}{\operatorname{dist}(\lambda, \Omega) - 1},$$

and hence $||R(\lambda, A)|| \leq (\operatorname{dist}(\lambda, \Omega) - 1)^{-1}$ for all $\lambda \in \mathbb{C}$ such that $\operatorname{dist}(\lambda, \Omega) > 1$. In particular, $||R(is, A)|| \leq m(|s|)$ for $0 < |s| \leq 1$, where $m : (0, 1] \rightarrow [1, \infty)$ is as in (6.7), so the result follows from Theorem 4.8. \Box

Remark 6.4. Note that the conclusion of Theorem 6.3 remains true whenever Ω is replaced by any set Ω' such that

$$\{\alpha_k : k \in \mathbb{Z}\} \subset \Omega' \subset \{\lambda \in \mathbb{C} : \operatorname{Re} \lambda < -1\} \cup \{-1\}.$$

In particular, we can take $\Omega' = \Omega_{\psi}$, where $\Omega_{\psi} = \{\lambda \in \mathbb{C} : \operatorname{Re} \lambda \leq -\psi(|\operatorname{Im} \lambda|)\}$ for some non-decreasing and continuously differentiable function $\psi: [0,1] \to [1,\infty)$ satisfying $\psi(0) = 1$ and $\psi(s) > 1$ for $s \in (0,1]$; see Figure 2. Then (6.7) becomes

(6.8)
$$m(r) = \frac{1}{\operatorname{dist}(ir, \Omega_{\psi}) - 1}, \quad 0 < r \le 1.$$

If $\psi'(0) > 0$, it is easy to see that $m(r) \approx r^{-2}$ as $r \to 0+$ and therefore

(6.9)
$$||AT(t)|| = O\left(\left(\frac{\log t}{t}\right)^{1/2}\right), \quad t \to \infty$$

On the other hand if $\psi'(0) = 0$ it follows from geometric considerations that

$$m(r) = \left(\psi(q(r))\left(1 + \psi'(q(r))^2\right)^{1/2} - 1\right)^{-1}, \quad 0 < r \le 1,$$

where $q: (0,1] \to (0,1]$ is the inverse function of the map $p(r) = r + \psi(r)\psi'(r), 0 < r \leq 1$.

- **Example 6.5.** (a) In the original robot rendezvous problem, where $\Omega = \{-1\}$, it follows from Theorem 6.3 that (6.9) holds. The proof of Theorem 6.1 shows that the logarithm can be omitted.
- (b) In the context of Remark 6.4, if $\psi(s) = 1 + s^{\alpha}$ for some $\alpha \ge 1$, then crude estimates show that $m(r) \approx r^{-2}$ as $r \to 0+$ if $1 \le \alpha < 2$ and




FIGURE 2. The region Ω_{ψ} .

 $m(r) \approx r^{-\alpha}$ as $r \to 0+$ if $\alpha \geq 2$. Hence in the first case (6.9) holds, while for $\alpha \geq 2$

$$||AT(t)|| = O\left(\left(\frac{\log t}{t}\right)^{1/\alpha}\right), \quad t \to \infty.$$

If p = 2, so that X is a Hilbert space, it follows from Theorem 4.10 that the logarithm can be omitted in both cases.

7. CONCLUSION

The main result of this paper, Theorem 4.3, is a powerful tool for studying the asymptotic behaviour of solutions to a rather general class of infinite systems of coupled differential equations. The versatility of the general theory is illustrated by the applications presented in Sections 5 and 6 to two important special cases: the platoon model and the robot rendezvous problem. Underlying Theorem 4.3 are a number of abstract results from operator theory and in particular the asymptotic theory of operator semigroups. It is striking how effective the results obtained by these abstract techniques are even in particular examples, shedding new light both on the platoon model and the robot rendezvous problem. Nevertheless, a number of important questions remain open. The first question, namely whether the logarithmic factors are needed in Theorem 4.3 when $p \neq 2$, was already raised in Remark 4.14(b). Here it would already be of interest to have an affirmative answer in certain special cases, for instance the platoon model dealt with in Theorem 5.1; see Remark 5.2(a). Another aspect of the theory which would benefit from further development is the condition for uniform boundedness of the semigroup presented in Theorem 3.1, since in its present state this condition is rather difficult to verify except for relatively simple characteristic functions. Furthermore, it remains to be determined to what extent the results obtained here can be extended to situations involving more complicated coupling, such as systems in which the evolution of each subsystem

depends on the states of several other subsystems rather than just one. Finally, it would seem worth investigating the corresponding questions in the discrete-time setting, both from an applications perspective and in view of the fact that the corresponding abstract theory is equally well developed as in the continuous-time setting. We hope to address some of these issues in future publications.

References

- W. Arendt and C. J. K. Batty. Tauberian theorems and stability of one-parameter semigroups. *Trans. Amer. Math. Soc.*, 306:837–841, 1988.
- [2] W. Arendt, C.J.K. Batty, M. Hieber, and F. Neubrander. Vector-valued Laplace transforms and Cauchy problems. Birkhäuser, Basel, second edition, 2011.
- [3] W. Arendt and J. Prüss. Vector-valued Tauberian theorems and asymptotic behavior of linear Volterra equations. SIAM J. Math. Anal., 23(2):412–448, 1992.
- [4] B. Bamieh, F. Paganini, and M.A. Dahleh. Distributed control of spatially invariant systems. *IEEE Trans. Automat. Control*, 47(7):1091–1107, 2002.
- [5] C.J.K. Batty, M.D. Blake, and S. Srivastava. A non-analytic growth bound for Laplace transforms and semigroups of operators. *Integral Equations Operator Theory*, 45:125– 154, 2003.
- [6] C.J.K. Batty, R. Chill, and Y. Tomilov. Fine scales of decay of operator semigroups. J. Europ. Math. Soc., 18(4):853–929, 2016.
- [7] R. Chill and D. Seifert. Quantified versions of Ingham's theorem. Submitted (http://arxiv.org/abs/1504.00385), April 2015.
- [8] R. Curtain, O. Iftime, and H. Zwart. System theoretic properties of a class of spatially invariant systems. Automatica, 45(7):1619 – 1627, 2009.
- [9] A. Feintuch and B. Francis. Infinite chains of kinematic points. Automatica, 48(5):901–908, 2012.
- [10] A. Feintuch and B. Francis. An infinite string of ants and Borel's method of summability. *Math. Intelligencer*, 34(2):15–18, 2012.
- [11] D.J.H. Garling. Inequalities: a journey into linear analysis. Cambridge University Press, Cambridge, 2007.
- [12] M.R. Jovanovic and B. Bamieh. On the ill-posedness of certain vehicular platoon control problems. *IEEE Trans. Automat. Control*, 50(9):1307–1321, Sept 2005.
- [13] U. Krengel. Ergodic Theorems. Walter de Gruyter, Berlin, 1985.
- [14] M. Lin and R. Sine. Ergodic theory and the functional equation (I T)x = y. J. Operator Theory, 10(1):153–166, 1983.
- [15] Yu. I. Lyubich and Q.P. Vũ. Asymptotic stability of linear differential equations in Banach spaces. *Studia Math.*, 88:37–42, 1988.
- [16] M.M. Martínez. Decay estimates of functions through singular extensions of vectorvalued Laplace transforms. J. Math. Anal. Appl., 375(1):196–206, 2011.
- [17] J. Ploeg, B.T.M. Scheepers, E. van Nunen, N. van de Wouw, and H. Nijmeijer. Design and experimental evaluation of cooperative adaptive cruise control. In *Proceedings* of the 14th International IEEE Conference on Intelligent Transportation Systems (ITSC), pages 260–265, 2011.
- [18] J. Ploeg, N. van de Wouw, and H. Nijmeijer. \mathcal{L}_p string stability of cascaded systems: Application to vehicle platooning. *IEEE Trans. Control Syst. Technol.*, 22(2):786–793, 2014.
- [19] D. Swaroop and J.K. Hedrick. String stability of interconnected systems. *IEEE Trans. Automat. Control*, 41(3):349–357, 1996.
- [20] Q.P. Vũ. Theorems of Katznelson-Tzafriri type for semigroups of operators. J. Funct. Anal., 103:74–84, 1992.
- [21] H. Zwart, A. Firooznia, J. Ploeg, and N. van de Wouw. Optimal control for nonexponentially stabilizable spatially invariant systems with an application to vehicular platooning. In *Proceedings of the 52nd IEEE Conference on Decision and Control*, Florence, Italy, December 10–13, 2013.

DEPARTMENT OF MATHEMATICS, TAMPERE UNIVERSITY OF TECHNOLOGY, PO. Box 553, 33101 TAMPERE, FINLAND *E-mail address*: lassi.paunonen@tut.fi

ST JOHN'S COLLEGE, ST GILES, OXFORD OX1 3JP, UNITED KINGDOM *E-mail address:* david.seifert@sjc.ox.ac.uk