

CURVATURE AND HIGHER ORDER BUSER INEQUALITIES FOR THE GRAPH CONNECTION LAPLACIAN*

SHIPING LIU[†], FLORENTIN MÜNCH[‡], AND NORBERT PEYERIMHOFF[§]

Abstract. We study the eigenvalues of the connection Laplacian on a graph with an orthogonal group or unitary group signature. We establish higher order Buser type inequalities, i.e., we provide upper bounds for eigenvalues in terms of Cheeger constants in the case of nonnegative Ricci curvature. In this process, we discuss the concepts of Cheeger type constants and a discrete Ricci curvature for connection Laplacians and study their properties systematically. The Cheeger constants are defined as mixtures of the expansion rate of the underlying graph and the frustration index of the signature. The discrete curvature, which can be computed efficiently via solving semidefinite programming problems, has a characterization by the heat semigroup for functions combined with a heat semigroup for vector fields on the graph.

Key words. connection Laplacian, Cheeger constants, discrete curvature, Buser inequality, semidefinite programming, Carstesian product

AMS subject classifications. 05C50, 53C23, 58J35, 05C76, 90C34

DOI. 10.1137/16M1056353

1. Introduction. A graph structure with its Laplacian matrix provides a mathematical tool to analyze the similarities between data points: those points with large enough similarities are connected by an edge. One can also assign edge weights to quantify such similarities. In many applications, it is noticed that the representation of the data set can be vastly improved by endowing the edges of the graph additionally with linear transformations [7, 23, 46]. For example, when the graph is representing a social network, we hope to attach to each edge an element from the one-dimensional orthogonal group $O(1) = \{\pm 1\}$ to indicate two kinds of opposite relationships between members of the network (vertices). When the graph is representing a higher dimensional data set, e.g., two-dimensional photos of a three-dimensional object from different views, one would like to assign to each edge an element of the orthogonal group $O(2)$ which optimally rotationally aligns photos when comparing their similarity (see, e.g., [7, 46]). In theoretical research, assigning linear transformations to the edges of a graph also provides mathematical structures that have been found very useful in various topics, e.g., the study of the Heawood map-coloring problem [21, 22], the construction of Ramanujan graphs [11, 39], and the study of a discrete analogue of magnetic operators [45, 47]. Agarwal, Kolla, and Madan [1, section 1.1] pointed out relations of these additional structures with the famous unique game conjecture of Khot and their relevance for the synchronization problem is explained in [19].

*Received by the editors January 11, 2016; accepted for publication (in revised form) December 17, 2018; published electronically February 5, 2019.

<http://www.siam.org/journals/sidma/33-1/M105635.html>

Funding: This work was supported by EPSRC grant EP/K016687/1, “Topology, Geometry and Laplacians of Simplicial Complexes.”

[†]School of Mathematical Sciences, University of Science and Technology of China, Hefei 230026, China (spliu@ustc.edu.cn).

[‡]Max Planck Institute for Mathematics in the Sciences, Inselstrasse 22, 04103 Leipzig, Germany (florentin.muench@mis.mpg.de).

[§]Department of Mathematical Sciences, Durham University, DH1 3LE Durham, United Kingdom (norbert.peyerimhoff@durham.ac.uk).

We consider such additional structures on the oriented edges of an undirected graph assuming only values in the orthogonal or unitary groups $O(d)$ or $U(d)$ and refer to them as d -dimensional signatures and denote them by σ throughout this paper. The corresponding Laplacian of a graph G with such an additional structure σ is called the *connection Laplacian* Δ^σ , defined by Singer and Wu [46]. The restriction to the groups $O(d)$ and $U(d)$ guarantee that Δ^σ is self-adjoint and has real eigenvalues.

In fact, the connection Laplacian of a graph yields a very elegant and general mathematical framework for the analysis of massive data sets, which includes several extensively studied graph operators as particular cases, e.g., the classical Laplacian, the signless Laplacian [15], the Laplacian for Harary's signed graphs [4, 51], and the discrete magnetic Laplacian [32, 45, 47].

In this paper, we study the spectra of the graph connection Laplacian, which are closely related to the geometric structure of the underlying graph G with a signature σ comprising the linear transformations attached to its edges. We describe this geometric structure by introducing two types of quantities, Cheeger type constants (denoted by $h_k^\sigma(G)$), and a discrete Ricci curvature. The global Ricci curvature condition is expressed by the notion $CD^\sigma(K, n)$, where $K \in \mathbb{R}$ is a lower curvature bound and $n > 0$ is an upper dimension bound (meaning that $CD^\sigma(K, n)$ implies $CD^\sigma(K', n')$ for any $K' \leq K$ and $n' \geq n$). Our main theorem is concerned with higher order Buser type inequalities, showing the close relations between eigenvalues λ_k^σ of the connection Laplacian Δ^σ and the Cheeger constants h_k^σ assuming nonnegativity of the discrete Ricci curvature. In the special case of an unweighted D -regular graph G , our result takes the following explicit form (the objects in this theorem require lengthier introductions which will be provided in the following subsections).

THEOREM 1.1 (higher order Buser inequalities). *Assume that a D -regular unweighted graph G with N vertices and a d -dimensional signature σ satisfies $CD^\sigma(0, \infty)$. Then for each natural number $1 \leq k \leq N$, we have the upper estimates*

$$(1.1) \quad \lambda_{kd}^\sigma \leq 16D(kd)^2 \log(2kd)(h_k^\sigma)^2$$

for the eigenvalues of the associated connection Laplacian Δ^σ .

We also obtain a lower bound estimate of the first nonzero eigenvalue of the connection Laplacian by the lower Ricci curvature bound, i.e., we show a Lichnerowicz type eigenvalue estimate (see Theorem 1.9 later in the introduction). In this process, the properties of the Cheeger constants and discrete Ricci curvature are explored systematically. In particular, our eigenvalues estimates help us to deepen the understanding of these geometric quantities.

For the interested reader we start with some background information in the setting of Riemannian manifolds. The following subsection can be skipped but we think that it is helpful to understand the involved notions in their wider context.

1.1. Background about Ricci curvature and eigenvalue estimates in Riemannian geometry. We provide a brief survey on the background in Riemannian geometry of the curvature notion and types of eigenvalue estimates that will be developed in this paper. This explains the motivation of the curvature notion and terminology like *higher order Cheeger and Buser inequalities* and *Lichnerowicz inequality*.

Let $(M, \langle \cdot, \cdot \rangle)$ be a Riemannian manifold of dimension n and $\Delta := \operatorname{div} \circ \operatorname{grad} \leq 0$ be the Laplace–Beltrami operator of $(M, \langle \cdot, \cdot \rangle)$. Bochner's formula states that for any smooth functions f on M and any point $x \in M$,

$$\frac{1}{2}\Delta|\operatorname{grad}f|^2(x) = |\operatorname{Hess}f|^2(x) + \langle \operatorname{grad}\Delta f(x), \operatorname{grad}f(x) \rangle + \operatorname{Ric}(\operatorname{grad}f(x)),$$

where Hess stands for the Hessian and Ric stands for the Ricci tensor. If the Ricci curvature of M is lower bounded by K , i.e., $\operatorname{Ric}(v) \geq K|v|^2$ for all $v \in T_x M$ and all $x \in M$, and, using the estimates $\operatorname{Hess}f^2(x) \geq \frac{1}{n}(\Delta f(x))^2$, we obtain for all $x \in M$

$$(1.2) \quad \frac{1}{2}\Delta|\operatorname{grad}f|^2(x) - \langle \operatorname{grad}\Delta f(x), \operatorname{grad}f(x) \rangle \geq \frac{1}{n}(\Delta f(x))^2 + K|\operatorname{grad}f(x)|^2.$$

This inequality relates the Ricci curvature and the Laplace–Beltrami operator.

Bakry and Émery [5, 6] clarify this relation by the following deep observation. They introduce the following symmetric bilinear forms Γ and Γ_2 for two smooth functions f, g , on M :

$$(1.3) \quad 2\Gamma(f, g) := \Delta(fg) - f\Delta g - (\Delta f)g,$$

$$(1.4) \quad 2\Gamma_2(f, g) := \Delta(\Gamma(f, g)) - \Gamma(f, \Delta g) - \Gamma(\Delta f, g).$$

Notice that $\Gamma(f, g) = \langle \operatorname{grad}f, \operatorname{grad}g \rangle$ and $\Gamma_2(f, g) = \frac{1}{2}\Delta|\operatorname{grad}f|^2 - \langle \operatorname{grad}\Delta f, \operatorname{grad}g \rangle$. Therefore, the inequality (1.2) can be rewritten as

$$(1.5) \quad \Gamma_2(f, f)(x) \geq \frac{1}{n}(\Delta f(x))^2 + K\Gamma(f, f)(x) \quad \text{for all smooth } f : M \rightarrow \mathbb{R}.$$

In conclusion, an n -dimensional Riemannian manifold $(M, \langle \cdot, \cdot \rangle)$ with Ricci curvature bounded from below by K satisfies the inequality (1.5). Notice from the definition of Γ and Γ_2 that (1.5) is expressed solely in terms of the Laplace–Beltrami operator Δ . This suggests a way to define, indirectly, a Ricci curvature notion for a metric space via the help of a Laplace–Beltrami operator. This will be the Ricci curvature approach for our discrete setting of graphs.

Suppose M be closed (i.e., compact without boundary). The spectrum of Δ is discrete and can be listed with multiplicity as

$$0 = \lambda_1(M) < \lambda_2(M) \leq \lambda_3(M) \leq \cdots \leq \lambda_k(M) \leq \cdots \nearrow \infty.$$

One of the main topics in spectral geometry is to explore lower or upper bounds of those eigenvalues in terms of geometric quantities. The following estimate is due to Lichnerowicz. (See, e.g., [20, Chapter 4.G.4].)

THEOREM 1.2 (Lichnerowicz estimate). *Let $(M, \langle \cdot, \cdot \rangle)$ be a closed Riemannian manifold of dimension n , and suppose that Ricci curvature is lower bounded by K . Then the first nonzero eigenvalue $\lambda_2(M)$ satisfies*

$$\lambda_2(M) \geq \frac{n}{n-1}K.$$

The so-called Cheeger constant is defined as

$$h(M) := \inf_S \frac{\operatorname{vol}(S)}{\min\{\operatorname{vol}(M_1), \operatorname{vol}(M_2)\}},$$

where S runs over hypersurfaces of M dividing M into two submanifolds M_1 and M_2 with boundary S . The following estimate is due to Cheeger. (See, e.g., [20, Chapter 4.G.4].)

THEOREM 1.3 (Cheeger inequality). *Let $(M, \langle \cdot, \cdot \rangle)$ be a closed Riemannian manifold. Then we have*

$$\lambda_2(M) \geq \frac{1}{4}h(M)^2.$$

In 1982, Peter Buser [12] showed that $\lambda_2(M)$ can also be bounded from *above* by $h(M)^2$, up to a constant involving Ricci curvature. This result was further improved by Ledoux [33] and can be stated as follows.

THEOREM 1.4 (Buser inequality). *Let $(M, \langle \cdot, \cdot \rangle)$ be a closed Riemannian manifold with nonnegative Ricci curvature. Then there exists a universal constant C such that*

$$\lambda_2(M) \leq Ch(M)^2.$$

It is natural to ask whether there are similar lower and upper bounds for the higher eigenvalues $\lambda_k(M)$, $k \geq 3$, involving some kind of multiway Cheeger constants. In fact, we can define such multiway Cheeger constants in the following way. For any Borel subset $A \subset M$ with positive measure, define

$$(1.6) \quad \phi(A) := \frac{\text{vol}(\partial A)}{\text{vol}(A)}.$$

Then we set for any natural number k ,

$$(1.7) \quad h_k(M) := \inf_{\{A_i\}_{i=1}^k} \max_{1 \leq i \leq k} \phi(A_i),$$

where the infimum is taken over all nonempty, disjoint subsets $\{A_i\}_{i=1}^k$ of M such that each A_i has positive measure. One can check that

$$h_2(M) = h(M).$$

Lower bound estimates of $\lambda_k(M)$ in terms of $h_k(M)^2$ are called *higher order Cheeger inequalities*, and upper bound estimates of $\lambda_k(M)$ in terms of $h_k(M)^2$ under assuming nonnegative Ricci curvature are called *higher order Buser inequalities*. Higher order Buser inequalities were first proved by Funano [18] and later improved in [36].

Theorem 1.1 presented in the previous subsection is such a higher order Buser inequality for the connection Laplacian in the discrete setting of a graph with a signature. Note that the appearance of a signature requires an adaption of the curvature condition and of the involved multiway Cheeger constants. These notions are introduced for graphs in the next two subsections.

1.2. Signatures, graph connection Laplacians, and Cheeger constants.

The aim of this subsection is to introduce central notions appearing in our main result, Theorem 1.1 above, in more detail: signatures, associated connection Laplacians, and multiway Cheeger constants.

We first fix some relevant notation. Let $G = (V, E)$ be an undirected simple finite graph with vertex set V and edge set E . For simplicity, we restrict ourselves to *unweighted D -regular graphs* in this introduction even though our results are proved later without this restriction. Let H be a group. For each edge $\{x, y\} \in E$, we assign an element $\sigma_{xy} \in H$ to it, such that

$$(1.8) \quad \sigma_{yx} = \sigma_{xy}^{-1}.$$

Actually, we are defining a map $\sigma : E^{or} \rightarrow H$, where $E^{or} := \{(x, y), (y, x) \mid \{x, y\} \in E\}$ is the set of all *oriented edges*. In this paper, we restrict the group H to be the

d -dimensional orthogonal group $O(d)$ or unitary group $U(d)$ and we call σ a d -dimensional signature of the graph G .

The (normalized) connection Laplacian Δ^σ , as a matrix, is given by

$$(1.9) \quad \Delta^\sigma := \frac{1}{D} A^\sigma - I_{Nd},$$

where D is the (constant) vertex degree and I_{Nd} is a $(Nd) \times (Nd)$ -identity matrix, N is the size of vertex set V , and A^σ is the $(Nd) \times (Nd)$ -matrix, blockwisely defined as

$$(1.10) \quad (A^\sigma)_{xy} = \begin{cases} 0 & \text{if } \{x, y\} \notin E; \\ \sigma_{xy} & \text{if } (x, y) \in E^{or}. \end{cases}$$

Due to the special choice of H and (1.8), Δ^σ is Hermitian. Hence all eigenvalues of the matrix Δ^σ are real. Note that the connection Laplacian Δ^σ in (1.9) is defined as a negative semidefinite matrix for our later purpose of defining the discrete curvature, due to a convention originating from Riemannian geometry. However, we still want to deal with nonnegative eigenvalues. Hence, when we speak of eigenvalues of the connection Laplacian Δ^σ , we mean the eigenvalue of the matrix $-\Delta^\sigma$. They can be listed (counting multiplicity) as

$$(1.11) \quad 0 \leq \lambda_1^\sigma \leq \lambda_2^\sigma \leq \cdots \leq \lambda_d^\sigma \leq \cdots \leq \lambda_{(N-1)d+1}^\sigma \leq \lambda_{(N-1)d+2}^\sigma \leq \cdots \leq \lambda_{Nd}^\sigma \leq 2.$$

Observe that two different signatures do not necessarily lead to different spectra. Given a function $\tau : V \rightarrow H$ and a signature $\sigma : E^{or} \rightarrow H$, we consider the new signature σ^τ defined by

$$(1.12) \quad \sigma_{xy}^\tau := \tau(x)^{-1} \sigma_{xy} \tau(y) \quad \forall (x, y) \in E^{or}.$$

Then the corresponding connection Laplacians Δ^σ and Δ^{σ^τ} are unitarily equivalent and hence share the same spectra. Indeed, it is easy to check that

$$(1.13) \quad \Delta^{\sigma^\tau} = (M_\tau)^{-1} \Delta^\sigma M_\tau,$$

where M_τ stands for a diagonal matrix given blockwisely by $(M_\tau)_{xx} := \tau(x)$. We call the function τ a *switching function*. Two signatures σ and σ' are said to be *switching equivalent* if there exists a switching function τ such that $\sigma' = \sigma^\tau$. It follows from (1.13) that the eigenvalues of the connection Laplacian Δ^σ are switching invariant.

The Cheeger type constants $\{h_k^\sigma \mid k = 1, 2, \dots, N\}$ and the discrete Ricci curvature $K_\infty(\sigma)$ that we are going to introduce are also switching invariant. A signature σ is said to be *balanced* if it is switching equivalent to the trivial signature $\sigma_{\text{triv}} : E^{or} \rightarrow \{id\} \in H$. In fact, the constants $\{h_k^\sigma \mid k = 1, 2, \dots, N\}$ are quantifying the connectivity of the graph and the unbalancedness of the signature σ . The latter is described by the *frustration index* $\iota^\sigma(S)$ of the signature σ restricted to the induced subgraph of $S \subseteq V$, with the property that

$$\iota^\sigma(S) = 0 \Leftrightarrow \sigma \text{ restricted on } S \text{ is balanced.}$$

By abuse of notation, we will also use S to denote its induced subgraph and denote by the E_S the edges of this induced subgraph. In particular, for an $U(1)$ signature $\sigma : E^{or} \rightarrow U(1)$, $\iota^\sigma(S)$ is defined as

$$(1.14) \quad \iota^\sigma(S) := \min_{\tau: S \rightarrow U(1)} \sum_{\{x, y\} \in E_S} |\sigma_{xy} \tau(y) - \tau(x)|,$$

where the minimum is taken over all switching functions on S . For higher dimensional signatures, we need to choose a matrix norm to define $\iota^\sigma(S)$; see Definition 4.3 in section 4.1. For $U(1)$ signatures, we show that there is an easier way to calculate its frustration index using *spanning trees* (see subsection 4.2).

Denote by $|E(S, V \setminus S)|$ the number of edges connecting S and its complement $V \setminus S$. We then define

$$(1.15) \quad \phi^\sigma(S) := \frac{\iota^\sigma(S) + |E(S, V \setminus S)|}{D \cdot |S|},$$

where $|S|$ is the cardinality of the set S . Note that the term $|E(S, V \setminus S)|$ corresponds to $\text{vol}(\partial A)$ and $D \cdot |S|$ corresponds to $\text{vol}(A)$ in (1.6). The quotient $\frac{|E(S, V \setminus S)|}{D \cdot |S|}$ can be understood as an *expansion rate* of the set S , that is, the ratio between the number of edges connecting S with the “outside world” and the size of the set S . The contribution of the signature is concentrated in the frustration index $\iota^\sigma(S)$ which does not appear in the setting without a signature.

Analogously as in (1.7), the Cheeger constants h_k^σ are then defined as

$$(1.16) \quad h_k^\sigma := \min_{\{S_i\}_{i=1}^k} \max_{1 \leq i \leq k} \phi^\sigma(S_i),$$

where the minimum is taken over all nonempty, pairwise disjoint subsets $\{S_i\}_{i=1}^k$ of the vertex set V . Choosing $k = 2$ and $\sigma_{\text{triv}} : E^{\text{or}} \rightarrow \{1\} \in O(1)$, we recover the classical Cheeger constant $h = h_2^{\sigma_{\text{triv}}}$, a measure for the connectedness of the graph G and crucial in the definition of *expander graphs* (see, e.g., [30]). Multiway Cheeger constants with trivial signature $h_k^{\sigma_{\text{triv}}}$ are used in the definition of *k-way expanders* (see, e.g., [40, 48]). Note that even though these are higher order concepts, the underlying spaces are still one-dimensional graphs in contrast to the recently very intensively studied topic of *higher dimensional expanders* (see, e.g., [28] and references therein), where expansion properties are generalized to higher dimensional simplicial complexes.

Concerning the involvement of a nontrivial signature σ , the definitions of the Cheeger constants h_k^σ are natural extensions of the constants in [4] and [32], where $H = O(1)$ and $U(1)$, respectively, and it is closely related to the $O(d)$ frustration ℓ^1 constant in [7] (see Remark 4.7 for a detailed explanation).

1.3. Curvature dimension inequalities with signatures. The nonnegativity of the discrete Ricci curvature $K_\infty(\sigma)$, or the *curvature dimension inequality with a signature*, $CD^\sigma(0, \infty)$, is an extension of the classical curvature dimension inequality à la Bakry and Émery [5, 6] on graphs, which has been studied extensively in recent years; see, e.g., [13, 17, 26, 27, 29, 34, 37, 44]. For related notions of curvature dimension inequalities and their strong implications in establishing various Li-Yau inequalities for heat semigroups on graphs, we refer to [8, 25, 41, 42]. The definition of $CD^\sigma(0, \infty)$ uses both the connection Laplacian Δ^σ and the graph Laplacian Δ , capturing the structure of the graph (especially its cycles) and the signature (especially the signature of cycles) locally around each vertex (see Proposition 3.12). The curvature condition $CD^\sigma(0, \infty)$ can be characterized by properties of the classical heat semigroup $P_t := e^{t\Delta}$ for functions and the heat semigroup $P_t^\sigma := e^{t\Delta^\sigma}$ for vector fields (vector valued functions) of the underlying graph (see Theorem 3.20). Another appealing feature of this curvature notion is that it can be calculated very efficiently. Indeed, calculating this curvature is equivalent to solving semidefinite programming problems.

It actually looks more natural to define the curvature dimension inequality with a signature using both matrices Δ and Δ^σ when we come back to the general setting: For a d -dimensional signature σ , the connection Laplacian Δ^σ , as an operator, acts on vector fields, i.e., functions $f : V \rightarrow \mathbb{K}^d$, where $\mathbb{K} = \mathbb{R}$ or \mathbb{C} .

DEFINITION 1.5. For any two functions $f, g : V \rightarrow \mathbb{K}^d$, we define

$$(1.17) \quad 2\Gamma^\sigma(f, g) := \Delta(f^T \bar{g}) - f^T (\overline{\Delta^\sigma g}) - (\Delta^\sigma f)^T \bar{g}$$

and

$$(1.18) \quad 2\Gamma_2^\sigma(f, g) := \Delta\Gamma^\sigma(f, g) - \Gamma^\sigma(f, \Delta^\sigma g) - \Gamma^\sigma(\Delta^\sigma f, g).$$

Note that $\Gamma^\sigma(f, g)$ and $\Gamma_2^\sigma(f, g)$ are \mathbb{K} -valued functions on V . We also write $\Gamma^\sigma(f) := \Gamma^\sigma(f, f)$ and $\Gamma_2^\sigma(f) := \Gamma_2^\sigma(f, f)$, for short. In (1.17) and (1.18), we use the graph Laplacian whenever we deal with a \mathbb{K} -valued function, and we use the graph connection Laplacian whenever we deal with a \mathbb{K} -vector valued function.

DEFINITION 1.6 ($CD^\sigma(K, \infty)$ inequality). Let $K \in \mathbb{R}$. We say the graph G with a signature σ satisfies the curvature dimension inequality $CD^\sigma(K, \infty)$ if we have for any vector field $f : V \rightarrow \mathbb{K}^d$ and any vertex $x \in V$,

$$(1.19) \quad \Gamma_2^\sigma(f)(x) \geq K\Gamma^\sigma(f)(x).$$

The precise ∞ -dimensional Ricci curvature lower bound $K_\infty(\sigma)$ is defined as the largest constant K such that (1.19) holds for all f .

In section 3.6, we show that the above curvature condition $CD^\sigma(0, \infty)$ can be characterized in terms of the corresponding heat semigroups $P_t := e^{t\Delta}$ and $P_t^\sigma = e^{t\Delta^\sigma}$ as follows:

$$CD^\sigma(K, \infty) \Leftrightarrow \Gamma^\sigma(P_t^\sigma f) \leq e^{-2Kt} P_t(\Gamma^\sigma(f)) \quad \forall f : V \rightarrow \mathbb{K}^d, \quad \forall t \geq 0.$$

This is very useful for the proof of Theorem 1.1.

It turns out that every graph G with a signature σ satisfies $CD^\sigma(\frac{2}{D} - 1, \infty)$ (see Corollary 3.8). This is shown by considering the switching invariance of $CD^\sigma(K, \infty)$ and CD^σ inequalities of covering graphs (see sections 3.2 and 3.3). In particular, every (unweighted) cycle graph with a signature $\sigma : E^{or} \rightarrow O(d)$ or $U(d)$ satisfies $CD^\sigma(0, \infty)$.

Given a graph G and a signature σ , the curvature $K_\infty(\sigma)$ can be computed very efficiently by reformulating the $CD^\sigma(K, \infty)$ inequality as linear matrix inequalities at local neighborhoods of all vertices (see section 3.4). Computing the precise Ricci curvature lower bound $K_\infty(\sigma)$ is then equivalent to solving semidefinite programming problems. In particular, we derive the precise formula of $K_\infty(\sigma)$ for a triangle (3-cycle) graph with $\sigma : E^{or} \rightarrow U(1)$ in section 3.5.

Moreover, the class of graphs with signatures satisfying $CD^\sigma(0, \infty)$ inequalities is rich since this curvature property is preserved by taking Cartesian products: Given two graphs $G_i = (V_i, E_i)$, $i = 1, 2$, with signatures $\sigma_i : E_i^{or} \rightarrow H_i = O(d_i)$ or $U(d_i)$, $i = 1, 2$, denote their Cartesian product graph by $G_1 \times G_2 = (V_1 \times V_2, E_{12})$. Let us assign a signature $\hat{\sigma}_{12} : E_{12}^{or} \rightarrow H_1 \otimes H_2$ to $G_1 \times G_2$ as follows:

$$\begin{aligned} \hat{\sigma}_{12, (x_1, y)(x_2, y)} &:= \sigma_{1, x_1 x_2} \otimes I_{d_2} \quad \text{for any } (x_1, x_2) \in E_1^{or}, y \in V_2; \\ \hat{\sigma}_{12, (x, y_1)(x, y_2)} &:= I_{d_1} \otimes \sigma_{2, y_1 y_2} \quad \text{for any } (y_1, y_2) \in E_2^{or}, x \in V_1. \end{aligned}$$

Then we have the following theorem (see Theorem A.3 and Remark A.4).

THEOREM 1.7. *Let $G_i, i = 1, 2$ with signatures $\sigma_i, i = 1, 2$, satisfy $CD^{\sigma_1}(K_1, \infty)$ and $CD^{\sigma_2}(K_2, \infty)$, respectively. Then the Cartesian product graph $G_1 \times G_2$ with the signature $\hat{\sigma}_{12}$ satisfies $CD^{\hat{\sigma}_{12}}(\frac{1}{2} \min\{K_1, K_2\}, \infty)$.*

In Appendix A, we discuss a similar behavior of the curvature dimension inequality on the Cartesian product $G_1 \times G_2$ when we assign to it various choices of edge weights, vertex measures, and signatures. The behavior of frustration indices and the Cheeger constant under taking Cartesian products is also discussed in Appendix A.

1.4. Motivation via a dual Buser inequality. Having introduced all relevant notions, let us have a second look at our main result presented earlier.

THEOREM 1.1 (higher order Buser inequalities). *Assume that a D -regular unweighted graph G with N vertices and a d -dimensional signature σ satisfies $CD^\sigma(0, \infty)$. Then for each natural number $1 \leq k \leq N$, we have the upper estimates*

$$(1.20) \quad \lambda_{kd}^\sigma \leq 16D(kd)^2 \log(2kd)(h_k^\sigma)^2$$

for the eigenvalues of the associated connection Laplacian Δ^σ .

Note that λ_{kd}^σ should be considered as the maximal value of the group of eigenvalues $\{\lambda_{(k-1)d+1}^\sigma, \dots, \lambda_{kd}^\sigma\}$. There are N different groups of eigenvalues and N Cheeger constants, correspondingly.

A Buser inequality on graphs satisfying the classical curvature dimension inequality $CD(0, \infty)$ has been established in [29] by extending an argument of Ledoux [33]. In fact, Theorem 1.1 reduces to their result (see (1.21) below) up to a constant, when $k = 2$, $d = 1$, and the signature σ is balanced.

Higher order Buser inequalities for graph Laplacians were proved in [37], via showing an eigenvalue ratio estimate. However, the method in [36, 37] does not extend to the connection Laplacian for a general signature $\sigma : E^{or} \rightarrow H = O(d)$ or $U(d)$, except for the very special case $O(1)$ (see Example 7.5). We discuss extensions of the methods in [36, 37] for $H = O(1)$ signatures in section 7. For general signatures, our proof neatly extends Ledoux's [33] argument for Buser's inequality and provides new ideas for establishing higher order Buser inequalities.

In the remainder of this section, we like to discuss a special consequence of Theorem 1.1, namely, a *dual Buser inequality*, which can also be viewed as a motivation for our general result.

Let us start by presenting some known results about Cheeger and dual Cheeger constants of a graph G and the eigenvalues of the classical graph Laplacian Δ . We have $\Delta := \frac{1}{D}A - I_N$, where A is the adjacency matrix of G and its real eigenvalues can be ordered in size with multiplicities by

$$0 = \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_N \leq 2.$$

We can also view Δ as a connection Laplacian with the trivial signature $\sigma_{\text{triv}} : E^{or} \rightarrow \{1\} \in O(1)$.

We already mentioned that if we assign to G the trivial $O(1)$ signature $\sigma_{\text{triv}} : E^{or} \rightarrow \{1\} \in O(1)$, then the constant $h_2^{\sigma_{\text{triv}}}$ coincides with the classical Cheeger constant of G . If, instead, we assign to G the signature $-\sigma_{\text{triv}} : E^{or} \rightarrow \{-1\} \in O(1)$, then the constant $h_1^{-\sigma_{\text{triv}}}$ reduces to the bipartiteness ratio of Trevisan [49], or to one minus the dual Cheeger constant of Bauer and Jost [9]. For details, we refer to [4]. In fact, we have the following relations between eigenvalues, Cheeger constants, and structural properties of the underlying graph:

$$\begin{aligned}\lambda_2 = 0 &\Leftrightarrow h_2^{\sigma_{\text{triv}}} = 0 \Leftrightarrow G \text{ has at least two connected components;} \\ 2 - \lambda_N = 0 &\Leftrightarrow h_1^{-\sigma_{\text{triv}}} = 0 \Leftrightarrow G \text{ has a bipartite connected component.}\end{aligned}$$

The Cheeger [2, 3, 16] and dual Cheeger [9, 49] inequalities assert that

$$\frac{(h_2^{\sigma_{\text{triv}}})^2}{2} \leq \lambda_2 \leq 2h_2^{\sigma_{\text{triv}}} \quad \text{and} \quad \frac{(h_1^{-\sigma_{\text{triv}}})^2}{2} \leq 2 - \lambda_N \leq 2h_1^{-\sigma_{\text{triv}}}.$$

For many purposes, it is very useful to have further relations between λ_2 ($2 - \lambda_N$, resp.) and $h_2^{\sigma_{\text{triv}}}$ ($h_1^{-\sigma_{\text{triv}}}$, resp.). The authors of [29] prove the following *Buser inequality*: If G satisfies the curvature dimension inequality $CD(0, \infty)$, then

$$(1.21) \quad \lambda_2 \leq 16D(h_2^{\sigma_{\text{triv}}})^2.$$

Note that the condition $CD(0, \infty)$ can be understood as a special case of $CD^\sigma(0, \infty)$ by choosing $\sigma = \sigma_{\text{triv}}$.

In particular, every cycle graph \mathcal{C}_N with N vertices satisfies $CD(0, \infty)$. Moreover, we have for the graph \mathcal{C}_N (see, e.g., [35, Proposition 7.4]),

$$(1.22) \quad (h_2^{\sigma_{\text{triv}}})^2 \leq \lambda_2(\mathcal{C}_N) \leq 5(h_2^{\sigma_{\text{triv}}})^2,$$

which is in line with the Cheeger inequality and Buser inequality, and also

$$(1.23) \quad 0.3(h_1^{-\sigma_{\text{triv}}})^2 \leq 2 - \lambda_N(\mathcal{C}_N) \leq 5(h_1^{-\sigma_{\text{triv}}})^2.$$

A natural question then arises: Is there a similar generalization of the right-hand side of (1.23)? That is, we are asking for a possible *dual Buser inequality* for the graph Laplacian Δ .

Observe that the first eigenvalue of the connection Laplacian $\Delta^{-\sigma_{\text{triv}}}$, also known as the signless Laplacian [15], is equal to $2 - \lambda_N$. Indeed, one can check that

$$-\Delta^{-\sigma_{\text{triv}}} = 2I_N - (-\Delta) = I_N + \frac{1}{D}A.$$

Therefore, Theorem 1.1 implies the following result.

COROLLARY 1.8 (dual Buser inequality). *Assume that a D -regular graph G with N vertices satisfies $CD^{-\sigma_{\text{triv}}}(0, \infty)$. Then we have for the largest eigenvalues of the graph Laplacian $\Delta = \frac{1}{D}A - I_N$*

$$(1.24) \quad 2 - \lambda_N \leq 16(\log 2)D(h_1^{-\sigma_{\text{triv}}})^2.$$

This provides a “dual” version of the Buser inequality in (1.21). We would like to mention that every cycle graph \mathcal{C}_N also fulfills the inequality $CD^{-\sigma_{\text{triv}}}(0, \infty)$.

Note that the inequality $CD^{-\sigma_{\text{triv}}}(0, \infty)$ is not solely defined by replacing the Laplacian Δ in (1.3) and (1.4) by $\Delta^{-\sigma_{\text{triv}}}$. In fact, our definition of $CD^{-\sigma_{\text{triv}}}(0, \infty)$ involves both matrices Δ and $\Delta^{-\sigma_{\text{triv}}}$, as was explained in subsection 1.3. A reason why both operators are needed in the curvature condition is that the corresponding heat semigroup $P_t^{-\sigma_{\text{triv}}} := e^{t\Delta^{-\sigma_{\text{triv}}}}$ does not possess a probability kernel (the operator $P_t^{-\sigma_{\text{triv}}}$ is not even nonnegative), a property which is essential for the proofs in [29, 33].

1.5. Lichnerowicz inequality and the jump of the curvature. Let λ^σ be the first nonzero eigenvalue of the connection Laplacian Δ^σ . Suppose the graph G is connected. We observed that when σ is unbalanced, $\lambda_1^\sigma \neq 0$, and hence $\lambda^\sigma = \lambda_1^\sigma$. Moreover, when $\iota^\sigma(V)$ becomes very small, i.e., when σ is very close to being balanced, $\lambda^\sigma = \lambda_1^\sigma$ becomes very close to 0. Once σ becomes balanced, $\lambda_1^\sigma = 0$, and $\lambda^\sigma = \lambda_2^\sigma > 0$. We say that the quantity λ^σ *jumps* when σ becomes balanced.

We show the following Lichnerowicz type eigenvalue estimate in section 6.

THEOREM 1.9 (Lichnerowicz inequality). *Assume that the graph G with a signature σ satisfies $CD^\sigma(K, \infty)$. Then the first nonzero eigenvalue λ^σ satisfies*

$$(1.25) \quad \lambda^\sigma \geq K.$$

For another Lichnerowicz type eigenvalue estimate for the eigenvalues λ_2 and $2 - \lambda_N$ of the graph Laplacian Δ in terms of the coarse Ricci curvature bound due to Ollivier [43], we refer to [10]. An interesting application of Theorem 1.9 is the following: The jump phenomenon of the quantity λ^σ imposes a similar jump phenomenon on the curvature.

Figure 1 illustrates the jumps of the first nonzero eigenvalue λ^σ and the curvature $K_\infty(\sigma)$ of the particular example of a triangle graph \mathcal{C}_3 with $\sigma : E^{or} \rightarrow U(1)$, when σ becomes balanced. In Figure 1, the complex variable $s = \text{Sgn}(\mathcal{C}_3) \in U(1)$ is the signature of the triangle (see (2.2) for the definition). The signature σ is balanced if and only if $\text{Re}(s) = 1$. See section 3.5 for details.

Moreover, Theorem 1.9 also establishes direct relations between Cheeger constants and the discrete Ricci curvature; see section 6.

1.6. Organization of the paper. In section 2, we set up our general setting of a graph with edge weights, a general vertex measure, and a signature and discuss the associated connection Laplacian. In section 3, we discuss various basic properties of the curvature dimension inequalities with signatures and also their equivalent definitions. In section 4, we introduce multiway Cheeger constants with signatures and discuss some of the fundamental properties. Section 5 is devoted to the proof of our main result, that is, higher order Buser inequalities. In section 6, we prove a




FIG. 1. Curvature and eigenvalues of a signed triangle.

Lichnerowicz type eigenvalue estimate and discuss its applications. The special case of graphs with $O(1)$ signatures is treated in section 7, where an eigenvalue ratio estimate is obtained. In Appendix A, we provide a detailed discussion about the behavior on Cartesian product graphs of the two concepts, curvature dimension inequalities and Cheeger constants with signatures. Appendix B contains a counterexample showing that a combinatorial expression of the frustration index via spanning trees, which we established in section 4 for graphs with $U(1)$ signatures, no longer holds for $U(d)$ signatures with $d > 1$.

2. The connection Laplacian. In this section, we introduce the basic setting of a graph with edge weights, a vertex measure and a signature, and the corresponding connection Laplacian.

2.1. Basic setting. Throughout the paper, $G = (V, E, w)$ denotes an undirected weighted simple finite graph with vertex set V and edge set E . If two vertices $x, y \in V$ are connected by an edge, we write $x \sim y$ and denote this edge by $\{x, y\}$. To each edge $\{x, y\} \in E$, we associate a positive symmetric weight $w_{xy} = w_{yx}$. Let

$$d_x := \sum_{y, y \sim x} w_{xy}$$

be the (weighted) vertex degree of $x \in V$.

For the vertex set V , we assign a finite positive measure $\mu : V \rightarrow \mathbb{R}_{>0}$. The following two quantities D_G^{non} and D_G^{nor} will appear naturally in our arguments:

$$(2.1) \quad D_G^{non} := \max_{x \in V} \frac{d_x}{\mu(x)}, \quad \text{and} \quad D_G^{nor} := \max_{x \in V} \max_{y, y \sim x} \frac{\mu(x)}{w_{xy}}.$$

Typically, one chooses $\mu(x) = 1$ for all $x \in V$ ($\mu = \mathbf{1}_V$ for short), or $\mu(x) = d_x$ for all $x \in V$ ($\mu = \mathbf{d}_V$ for short). The superscripts in (2.1) are abbreviations for “nonnormalized” and “normalized,” respectively. Observe that, $D_G^{non} = \max_{x \in V} d_x$ for the measure $\mu = \mathbf{1}_V$, while $D_G^{nor} = \max_{x \in V} d_x$ for the measure $\mu = \mathbf{d}_V$ and $w_{xy} = 1$ for all $\{x, y\} \in E$.

We write (G, μ, σ) to denote a graph $G = (V, E, w)$ with the vertex measure μ and the signature $\sigma : E^{or} \rightarrow H$, where H is a group (recall (1.8)).

Recall from the introduction that σ is balanced if it is switching equivalent to the trivial signature $\sigma_{\text{triv}} : E^{or} \rightarrow id \in H$. Actually, the original definition of balancedness of a signature by Harary [23] is defined via the signature of cycles of the underlying graph. Let \mathcal{C} be a cycle of G , i.e., a subgraph composed of a sequence $(x_1, x_2), (x_2, x_3), \dots, (x_{\ell-1}, x_\ell), (x_\ell, x_1)$ of distinct edges. Then the signature $Sgn(\mathcal{C})$ of \mathcal{C} is defined as the conjugacy class of the element

$$(2.2) \quad \sigma_{x_1 x_2} \sigma_{x_2 x_3} \cdots \sigma_{x_{\ell-1} x_\ell} \sigma_{x_\ell x_1} \in H.$$

Note that the signature of any cycle is switching invariant. Harary [23] (see also [50]) defines a signature $\sigma : E^{or} \rightarrow H$ to be balanced if the signature of every cycle of G is (the conjugacy class of the) identity element $id \in H$. In fact, the above two definitions of balancedness of a signature are equivalent; see [50, Corollary 3.3 and section 9].

For more historical background about signatures of graphs, we refer the reader to [38, section 3].

2.2. Connection Laplacian. Let $\mathbb{K} = \mathbb{R}$ or \mathbb{C} . Throughout the paper we restrict H to be the orthogonal group $O(d)$ or the unitary group $U(d)$, of dimension

d , $d \in \mathbb{Z}_{>0}$, when $\mathbb{K} = \mathbb{R}$ or \mathbb{C} , respectively. For every edge $(x, y) \in E^{or}$, σ_{xy} is a $(d \times d)$ -orthogonal or unitary matrix and we have $\sigma_{yx} = \sigma_{xy}^{-1} = \overline{\sigma_{xy}^T}$.

For any vector-valued functions $f : V \rightarrow \mathbb{K}^d$ and any vertex $x \in V$, the graph connection Laplacian Δ^σ is defined via

$$\Delta^\sigma f(x) := \frac{1}{\mu(x)} \sum_{y, y \sim x} w_{xy} (\sigma_{xy} f(y) - f(x)) \in \mathbb{K}^d.$$

Note that a function $f : V \rightarrow \mathbb{K}^d$ can also be considered as an (Nd) -dimensional column vector, which we denote by $\vec{f} \in \mathbb{K}^{Nd}$. This vector is well defined once we enumerate the vertices in V . The Laplacian can then be written as

$$(2.3) \quad \Delta^\sigma = (\text{diag}_\mu)^{-1} (A^\sigma - \text{diag}_D),$$

where diag_μ and diag_D are $(Nd) \times (Nd)$ -diagonal matrices with the diagonal blocks $(\text{diag}_\mu)_{xx} = \mu(x)I_d$ and $(\text{diag}_D)_{xx} = d_x I_d$ for $x \in V$, respectively. Here we use I_d for a $(d \times d)$ -identity matrix. The matrix A^σ is defined blockwise as follows. For $x, y \in V$, the $(d \times d)$ -block of it is given by

$$(2.4) \quad (A^\sigma)_{xy} = \begin{cases} 0 & \text{if } \{x, y\} \notin E; \\ w_{xy} \sigma_{xy}, & (x, y) \in E. \end{cases}$$

Then we have $\overline{\Delta^\sigma \vec{f}} = (\text{diag}_\mu)^{-1} (A^\sigma - \text{diag}_D) \vec{f}$.

If every edge has the trivial signature $1 \in O(1)$, Δ^σ reduces to the graph Laplacian Δ . When $H = U(1)$, Δ^σ coincides with the discrete magnetic Laplacian [32, 45, 47].

Given two functions $f, g : V \rightarrow \mathbb{K}^d$, locally at a vertex x the Hermitian inner product of $f(x)$ and $g(x)$ is given by $f(x)^T \overline{g(x)}$. The corresponding norm of $f(x)$ is denoted by $|f(x)| := \sqrt{f(x)^T \overline{f(x)}}$. Globally, we have the following inner product between f and g :

$$(2.5) \quad \langle f, g \rangle_\mu := \sum_{x \in V} \mu(x) f(x)^T \overline{g(x)}.$$

We denote by $\ell^2(V, \mathbb{K}^d; \mu)$ the corresponding Hilbert space of functions. The ℓ^2 norm corresponding to (2.5) is denoted by $\|\cdot\|_{2, \mu}$. Note that Δ^σ is a self-adjoint operator on $\ell^2(V, \mathbb{K}^d; \mu)$, i.e.,

$$(2.6) \quad \langle \Delta^\sigma f, g \rangle_\mu = \langle f, \Delta^\sigma g \rangle_\mu.$$

We call $\lambda^\sigma \in \mathbb{R}$ an eigenvalue of Δ^σ if there exists a nonzero function $f : V \rightarrow \mathbb{K}^d$ such that $\Delta^\sigma f = -\lambda^\sigma f$. In fact, all Nd eigenvalues of Δ^σ lie in the interval $[0, 2D_G^{non}]$.

Let Σ be the group generated by the elements of $\{\sigma_{xy} \mid (x, y) \in E^{or}\}$. We call Σ the *signature group* of the graph (G, σ) . If the action of Σ on \mathbb{K}^d is reducible, we have an orthogonal decomposition of \mathbb{K}^d , i.e.,

$$\mathbb{K}^d = U_1 \oplus U_2 \oplus \cdots \oplus U_r \text{ for some } r,$$

where the U_i 's are pairwise orthogonal w.r.t. the Hermitian inner product of \mathbb{K}^d and each U_i is an Σ -invariant subspace of \mathbb{K}^d of dimension d_i such that $\sum_{i=1}^r d_i = d$. Then there exist signatures $\sigma_i : E^{or} \rightarrow O(d_i)$ or $U(d_i)$, $i = 1, 2, \dots, r$, such that we can write

$$\Delta^\sigma = \Delta^{\sigma_1} \oplus \Delta^{\sigma_2} \oplus \cdots \oplus \Delta^{\sigma_r}$$

by identifying each U_i with the vector space \mathbb{K}^{d_i} .

3. Curvature dimension inequalities with signatures. In this section, we introduce the $CD^\sigma(K, n)$ inequality for $K \in \mathbb{R}$ and $n \in \mathbb{R}_+$ and discuss its basic properties. We will characterize the CD^σ inequality in terms of linear matrix inequalities and also in terms of heat semigroups for functions and vector fields.

3.1. Definitions. We start by discussing several basic properties of the operators Γ^σ and Γ_2^σ defined in Definition 1.5 (of course, we are using the Laplacians in our current general setting). First, observe that they have the following Hermitian properties:

$$(3.1) \quad \Gamma^\sigma(f, g) = \overline{\Gamma^\sigma(g, f)}, \quad \Gamma_2^\sigma(f, g) = \overline{\Gamma_2^\sigma(g, f)} \quad \forall f, g : V \rightarrow \mathbb{K}^d.$$

Since the graph Laplacian Δ satisfies

$$(3.2) \quad \sum_{x \in V} \mu(x) \Delta(f^T \bar{g})(x) = 0,$$

the definition (1.17) of Γ^σ and the self-adjointness (2.6) of Δ^σ lead to the following summation by part formula:

$$(3.3) \quad \sum_{x \in V} \mu(x) \Gamma^\sigma(f, g)(x) = -\langle f, \Delta^\sigma g \rangle_\mu = -\langle \Delta^\sigma f, g \rangle_\mu.$$

Moreover, we have the following properties.

PROPOSITION 3.1. *For any two functions $f, g : V \rightarrow \mathbb{K}^d$ and any $x \in V$, we have*

(i)

$$\Gamma^\sigma(f, g)(x) = \frac{1}{2\mu(x)} \sum_{y, y \sim x} w_{xy} (\sigma_{xy} f(y) - f(x))^T (\overline{\sigma_{xy} g(y) - g(x)});$$

(ii)

$$|\Gamma^\sigma(f, g)(x)| \leq \sqrt{\Gamma^\sigma(f)(x)} \sqrt{\Gamma^\sigma(g)(x)}.$$

Proof. The formula (i) follows from a direct calculation; (ii) is a consequence of (i) by applying the Cauchy–Schwarz inequality. \square

DEFINITION 3.2 (CD^σ inequality). *Let $K \in \mathbb{R}$ and $n \in \mathbb{R}_+$. We say that (G, μ, σ) satisfies the CD^σ inequality $CD^\sigma(K, n)$ if we have for any vector field $f : V \rightarrow \mathbb{K}^d$ and any vertex $x \in V$*

$$(3.4) \quad \Gamma_2^\sigma(f)(x) \geq \frac{1}{n} |\Delta^\sigma f(x)|^2 + K \Gamma^\sigma(f)(x).$$

We call K a lower curvature bound of (G, μ, σ) and n a dimension parameter. We define the n -dimensional Ricci curvature $K_n(G, \mu, \sigma; x)$ of (G, μ, σ) at the vertex $x \in V$ to be the largest K that the inequality (3.4) holds for a given dimension parameter n . We further define the precise n -dimensional Ricci curvature lower bound $K_n(G, \mu, \sigma)$ of (G, μ, σ) as

$$(3.5) \quad K_n(G, \mu, \sigma) := \min_{x \in V} K_n(G, \mu, \sigma; x).$$

We also simply write $K_n(\sigma; x)$ and $K_n(\sigma)$ when the setting (G, μ) is clear.

Note that for given $K \in \mathbb{R}$, and $n_1, n_2 \in \mathbb{R}_+$ with $n_1 \leq n_2$, the inequality $CD^\sigma(K, n_1)$ implies $CD^\sigma(K, n_2)$. In other words, $CD^\sigma(K, n)$ provides a lower curvature bound K and an upper dimension bound n of the graph.

We also remark that rescaling the measure μ by a constant $c > 0$ leads to

$$(3.6) \quad K_n(G, c\mu, \sigma; x) = \frac{1}{c} K_n(G, \mu, \sigma; x).$$

We will be particularly interested in graphs satisfying $CD^\sigma(K, \infty)$ in this paper.

The classical curvature-dimension inequality $CD(K, n)$ à la Bakry and Émery [6] on graphs is defined as follows: For any real-valued function $f : V \rightarrow \mathbb{R}$ and any vertex x , we have

$$(3.7) \quad \Gamma_2(f)(x) \geq \frac{1}{n} |\Delta f(x)|^2 + K\Gamma(f)(x).$$

Recall the definitions of Γ and Γ_2 from (1.3) and (1.4).

When $\sigma = \sigma_{\text{triv}} : E^{\text{or}} \rightarrow id \in U(d)$ is the trivial signature, the graph (G, μ, σ) satisfies the inequality $CD^\sigma(K, n)$ if and only if (G, μ) satisfies the inequality $CD(K, n)$. In fact, this follows immediately from the following general result.

PROPOSITION 3.3. *Assume that the action of the signature group Σ of the graph (G, μ, σ) is decomposable, i.e., we have*

$$\Delta^\sigma = \Delta^{\sigma_1} \oplus \Delta^{\sigma_2} \oplus \cdots \oplus \Delta^{\sigma_r},$$

where $\sigma_i : E^{\text{or}} \rightarrow U(d_i)$ or $O(d_i)$, $i = 1, 2, \dots, r$. Then the graph (G, μ, σ) satisfies the inequality $CD^\sigma(K, n)$ if and only if (G, μ, σ_i) satisfies $CD^{\sigma_i}(K, n)$ for each $i = 1, 2, \dots, r$.

Proof. By assumption, for any function $f : V \rightarrow \mathbb{K}^d$, there exist functions $f_i : V \rightarrow U_i \cong \mathbb{K}^{d_i}$, $i = 1, 2, \dots, r$, such that

$$f^T \bar{f} = f_1^T \bar{f}_1 + f_2^T \bar{f}_2 + \cdots + f_r^T \bar{f}_r$$

and

$$\Delta^\sigma f = \Delta^{\sigma_1} f_1 \oplus \Delta^{\sigma_2} f_2 \oplus \cdots \oplus \Delta^{\sigma_r} f_r.$$

Hence, for any $x \in V$, we obtain by Definition 1.5,

$$\Gamma^\sigma(f)(x) = \sum_{i=1}^r \Gamma^{\sigma_i}(f_i)(x), \quad \text{and} \quad \Gamma_2^\sigma(f)(x) = \sum_{i=1}^r \Gamma_2^{\sigma_i}(f_i)(x).$$

We also have

$$|\Delta^\sigma f(x)|^2 = \sum_{i=1}^r |\Delta^{\sigma_i} f_i(x)|^2.$$

Therefore, the inequality

$$\Gamma_2^\sigma(f)(x) \geq \frac{1}{n} |\Delta^\sigma f(x)|^2 + K\Gamma^\sigma(f)(x) \quad \forall x \in V$$

is equivalent to the inequality

$$\sum_{i=1}^r \Gamma_2^{\sigma_i}(f_i)(x) \geq \sum_{i=1}^r \left(\frac{1}{n} |\Delta^{\sigma_i} f_i(x)|^2 + \Gamma^{\sigma_i}(f_i)(x) \right),$$

and the proposition follows immediately. \square

Given a graph (G, μ, σ) , where $\sigma : E^{or} \rightarrow O(d_1)$ or $U(d_1)$, we have a natural new signature

$$\begin{aligned}\sigma \otimes I_{d_2} : E^{or} &\rightarrow O(d_1 d_2) \text{ or } U(d_1 d_2), \\ (x, y) &\mapsto \sigma_{xy} \otimes I_{d_2},\end{aligned}$$

where I_{d_2} stands for the identity matrix of size $d_2 \times d_2$. The following observation will be useful in our later discussion about the CD^σ inequalities on Cartesian products of graphs in Appendix A.

COROLLARY 3.4. *A graph (G, μ, σ) satisfies $CD^\sigma(K, n)$ if and only if $(G, \mu, \sigma \otimes I_{d_2})$ satisfies $CD^{\sigma \otimes I_{d_2}}(K, n)$.*

Proof. We observe that the action of the signature group of $(G, \sigma \otimes I_{d_2})$ on $\mathbb{K}^{d_1 d_2}$ admits an orthogonal decomposition and, therefore, we have

$$\Delta^{\sigma \otimes I_{d_2}} = \underbrace{\Delta^\sigma \oplus \cdots \oplus \Delta^\sigma}_{d_2 \text{ times}}.$$

Corollary 3.4 is then a direct consequence of Proposition 3.3. \square

3.2. Switching invariance. The CD^σ inequality is switching invariant.

PROPOSITION 3.5. *If (G, μ, σ) satisfies $CD^\sigma(K, n)$, then (G, μ, σ^τ) satisfies $CD^{\sigma^\tau}(K, n)$ for any switching function $\tau : V \rightarrow H$.*

Proof. Recalling (1.13), we check that we have for any $\tau : V \rightarrow H$ and $f, g : V \rightarrow \mathbb{K}^d$,

$$(3.8) \quad \Gamma^{\sigma^\tau}(f, g) = \Gamma^\sigma(\tau^{-1}f, \tau^{-1}g) \text{ and } \Gamma_2^{\sigma^\tau}(f, g) = \Gamma_2^\sigma(\tau^{-1}f, \tau^{-1}g),$$

using $\overline{\tau(x)^T} = \tau^{-1}(x)$. The proposition then follows immediately from (1.13) and (3.8). \square

The arguments in the above proof show also that $K_n(G, \mu, \sigma; x)$, introduced in Definition 3.2, is the switching invariant for any given n .

We denote by dist the canonical graph distance and define the ball of radius r centered at $x \in V$ by

$$B_r(x) := \{y \in V \mid \text{dist}(x, y) \leq r\}.$$

PROPOSITION 3.6. *Let (G, μ, σ) be given. If the signature of every cycle of length 3 or 4 is equal to (the conjugate class of) $\text{id} \in H$, then (G, μ, σ) satisfies $CD^\sigma(K, n)$ if and only if (G, μ) satisfies $CD(K, n)$.*

Proof. Let $x \in V$ be a vertex. Since all cycles of 3 or 4 have trivial signature, we can switch all the signatures of edges in the subgraph induced by the ball $B_2(x)$ to be trivial. Note that the inequality (3.4) only involves the vertices in the ball $B_2(x)$. Then the proposition follows from Propositions 3.5 and 3.3. \square

3.3. Coverings and a general lower curvature bound. Let $(\tilde{G}, \tilde{\mu}, \tilde{\sigma})$ and (G, μ, σ) be two graphs. Let $\pi : (\tilde{G}, \tilde{\mu}, \tilde{\sigma}) \rightarrow (G, \mu, \sigma)$ be a graph homomorphism, namely, $\pi : \tilde{V} \rightarrow V$ is surjective, and if $\{\tilde{x}, \tilde{y}\} \in \tilde{E}$, then $\{\pi(\tilde{x}), \pi(\tilde{y})\} \in E$. Moreover, we require

$$(3.9) \quad \tilde{\sigma}_{\tilde{x}\tilde{y}} = \sigma_{\pi(\tilde{x})\pi(\tilde{y})}, \quad \tilde{w}_{\tilde{x}\tilde{y}} = w_{\pi(\tilde{x})\pi(\tilde{y})}, \quad \text{and} \quad \tilde{\mu}(\tilde{x}) = \mu(\pi(\tilde{x})).$$

Such a map π is called a *covering map* if, furthermore, the subgraph of \tilde{G} induced by the ball $B_1(\tilde{x})$ centered at each vertex $\tilde{x} \in \tilde{V}$ is mapped bijectively to the subgraph of G induced by the ball $B_1(x)$. If a covering map $\pi : (\tilde{G}, \tilde{\mu}, \tilde{\sigma}) \rightarrow (G, \mu, \sigma)$ exists, we call the graph $(\tilde{G}, \tilde{\mu}, \tilde{\sigma})$ a *covering graph* of (G, μ, σ) .

THEOREM 3.7. *Let $(\tilde{G}, \tilde{\mu}, \tilde{\sigma})$ be a covering graph of (G, μ, σ) . If $(\tilde{G}, \tilde{\mu}, \tilde{\sigma})$ satisfies $CD^{\tilde{\sigma}}(K, n)$, then (G, μ, σ) satisfies $CD^{\sigma}(K, n)$.*

Proof. For any function $f : V \rightarrow \mathbb{K}^d$, we can find a corresponding function $\tilde{f} : \tilde{V} \rightarrow \mathbb{K}^d$ such that

$$(3.10) \quad \tilde{f}(\tilde{x}) := f(\pi(\tilde{x})) \quad \forall \tilde{x} \in \tilde{V},$$

where π is the covering map from $(\tilde{G}, \tilde{\mu}, \tilde{\sigma})$ to (G, μ, σ) .

For any $x \in V$, and any $\tilde{x} \in \pi^{-1}(x)$, we can check by definition of a covering map that

$$(3.11) \quad |\Delta^{\tilde{\sigma}} \tilde{f}(\tilde{x})|^2 = |\Delta^{\sigma} f(x)|^2, \quad \Gamma^{\tilde{\sigma}}(\tilde{f})(\tilde{x}) = \Gamma^{\sigma}(f)(x), \quad \Gamma_2^{\tilde{\sigma}}(\tilde{f})(\tilde{x}) = \Gamma_2^{\sigma}(f)(x).$$

Since $(\tilde{G}, \tilde{\mu}, \tilde{\sigma})$ satisfies $CD^{\tilde{\sigma}}(K, n)$, we obtain that for any $f : V \rightarrow \mathbb{K}^d$, and any vertex $x \in V$,

$$(3.12) \quad \Gamma_2^{\tilde{\sigma}}(\tilde{f})(\tilde{x}) \geq \frac{1}{n} |\Delta^{\tilde{\sigma}} \tilde{f}(\tilde{x})|^2 + K \Gamma^{\tilde{\sigma}}(\tilde{f})(\tilde{x}).$$

Combining this with (3.11) completes the proof. \square

COROLLARY 3.8. *Any graph (G, μ, σ) satisfies the inequality*

$$CD^{\sigma} \left(\frac{2}{D_G^{nor}} - D_G^{non}, 2 \right).$$

In particular, any unweighted cycle graph with constant vertex measure $\mu \equiv \nu_0 \cdot \mathbf{1}_V$ and any signature $\sigma : E^{or} \rightarrow H$ satisfies

$$CD^{\sigma}(0, 2).$$

Proof. Let $(T_G, \tilde{\mu}, \tilde{\sigma})$ be the universal covering of (G, μ, σ) , i.e., T_G is a tree. It is shown in [34, Theorem 1.2] (see also [27, Theorem 8]) that $(T_G, \tilde{\mu})$ satisfies the CD inequality

$$CD \left(\frac{2}{D_{T_G}^{nor}} - D_{T_G}^{non}, 2 \right).$$

Due to Proposition 3.6, we know that $(T_G, \tilde{\mu}, \tilde{\sigma})$ satisfies

$$CD^{\tilde{\sigma}} \left(\frac{2}{D_{T_G}^{nor}} - D_{T_G}^{non}, 2 \right),$$

since a tree has no cycles. By the definition of a covering graph, we have $D_G^{nor} = D_{T_G}^{nor}$ and $D_G^{non} = D_{T_G}^{non}$. Then the corollary follows directly from Theorem 3.7. \square

3.4. CD^{σ} inequality as linear matrix inequalities. In this subsection, we present an equivalent formulation of the CD^{σ} inequality via linear matrix inequalities. As a consequence, the problem of calculating the Ricci curvature of a graph is reduced to solving semidefinite programming problems. In this process, we explore the geometrical information captured by the CD^{σ} inequality of a graph.

By Definition 1.5, the operators Γ^σ and Γ_2^σ can be considered as two symmetric sesquilinear forms. Hence they can be represented by Hermitian matrices. For our purpose, we are interested in considering the two symmetric sesquilinear forms locally at every vertex $x \in V$. There exist two $(Nd) \times (Nd)$ -Hermitian matrices $\Gamma^\sigma(x)$ and $\Gamma_2^\sigma(x)$ such that for any two functions $f, g : V \rightarrow \mathbb{K}^d$,

$$(3.13) \quad \Gamma^\sigma(f, g)(x) = \vec{f}^T \Gamma^\sigma(x) \vec{g} \quad \text{and} \quad \Gamma_2^\sigma(f, g)(x) = \vec{f}^T \Gamma_2^\sigma(x) \vec{g}.$$

Denote by $|B_r(x)|$ the cardinality of the set $B_r(x)$. Observe that the matrix $\Gamma^\sigma(x)$ only has a nontrivial block of size $|B_1(x)| \times |B_1(x)|$, while the matrix Γ_2^σ only has a nontrivial block of size $|B_2(x)| \times |B_2(x)|$.

We denote by $\Delta^\sigma(x)$ the $(d \times Nd)$ -matrix such that $\Delta^\sigma f(x) = \Delta^\sigma(x) \vec{f}$ for all functions $f : V \rightarrow \mathbb{K}^d$. Given two Hermitian matrices M_1 and M_2 , the inequality $M_1 \geq M_2$ means that the matrix $M_1 - M_2$ is positive semidefinite. Then we have the following equivalent definition of CD^σ inequality.

DEFINITION 3.9 (CD^σ inequality as linear matrix inequalities). *Let $K \in \mathbb{R}$ and $n \in \mathbb{R}_+$. A graph (G, μ, σ) satisfies the CD^σ inequality $CD^\sigma(K, n)$ if and only if, for any vertex $x \in V$, the following linear matrix inequality holds:*

$$(3.14) \quad \Gamma_2(x) \geq \frac{1}{n} \Delta^\sigma(x)^T \overline{\Delta^\sigma(x)} + K \Gamma^\sigma(x).$$

A direct consequence is the following proposition.

PROPOSITION 3.10 (semidefinite programming). *The n -dimensional Ricci curvature $K_n(G, \mu, \sigma; x)$ of the graph (G, μ, σ) at the vertex $x \in V$ is the solution of the following semidefinite programming:*

$$\begin{aligned} & \text{maximize } K \\ & \text{subject to } \Gamma_2^\sigma(x) - \frac{1}{n} \Delta^\sigma(x)^T \overline{\Delta^\sigma(x)} \geq K \Gamma^\sigma(x). \end{aligned}$$

In the following, we describe the explicit structure of the matrices $\Delta^\sigma(x)$, $\Gamma^\sigma(x)$, and $\Gamma_2^\sigma(x)$. For simplicity, we restrict to the setting

$$(3.15) \quad \mu = \mathbf{1}_V, \text{ i.e., } \mu(x) = 1 \quad \forall x \in V,$$

and

$$(3.16) \quad w_{xy} = 1 \quad \forall \{x, y\} \in E.$$

Given a vertex $x \in V$, let us denote its neighbors by y_1, y_2, \dots, y_{d_x} . By abuse of notation, we still write $\Delta^\sigma(x)$ and $\Gamma^\sigma(x)$ for their nontrivial blocks corresponding to the vertices x, y_1, \dots, y_{d_x} . Then it is easy to see that

$$(3.17) \quad \Delta^\sigma(x) = \begin{pmatrix} -d_x I_d & \sigma_{xy_1} & \cdots & \sigma_{xy_{d_x}} \end{pmatrix}$$

and

$$(3.18) \quad 2\Gamma^\sigma(x) = \begin{pmatrix} d_x I_d & -\overline{\sigma_{xy_1}} & \cdots & -\overline{\sigma_{xy_{d_x}}} \\ -\sigma_{xy_1}^T & I_d & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ -\sigma_{xy_{d_x}}^T & 0 & \cdots & I_d \end{pmatrix}.$$

For the matrix $\Gamma_2^\sigma(x)$, the structure of the subgraph induced by $B_2(x)$ is of relevance. We denote the sphere of radius r centered at a vertex $x \in V$ by

$$S_r(x) := \{y \in V \mid \text{dist}(x, y) = r\}.$$

Then, the ball $B_2(x)$ has the decomposition $B_2(x) = \{x\} \sqcup S_1(x) \sqcup S_2(x)$.

We first introduce some natural geometric quantities before we present the entries of the matrix $\Gamma_2^\sigma(x)$. For any vertex $y \in S_1(x)$, we have

$$(3.19) \quad |S_1(y) \cap S_2(x)| := \sum_{z, z \sim y, z \not\sim x, z \neq x} 1$$

and

$$(3.20) \quad \sharp_\Delta(x, y) := |S_1(y) \cap S_1(x)| := \sum_{z, z \sim y, z \sim x} 1.$$

Note that (3.20) is the number of triangles (i.e., 3-cycles) which contain the two neighbors x and y . This justifies the notation $\sharp_\Delta(x, y)$.

For any vertex $z \in S_2(x)$, we have

$$(3.21) \quad |S_1(z) \cap S_1(x)| := \sum_{y, y \sim x, y \sim z} 1.$$

Note that (3.21) is related to the number of 4-cycles which contain the two vertices x and z .

Remark 3.11. The above three geometric quantities are all closely related to the growth rate of the cardinality of $B_r(x)$ (in other words, the volume of $B_r(x)$ w.r.t. the measure $\mu = \mathbf{1}_V$) when the radius r increases.

The quantity $\sharp_\Delta(x, y)$ counts the number of 3-cycles regardless of their signatures. A signed version of this quantity is also important, and we define the following quantity describing the unbalancedness of the triangles containing the two neighbors x and y :

$$(3.22) \quad \sharp_\Delta^\sigma(x, y) := \sum_{z, z \sim y, z \sim x} (\mathbf{I}_d - \overline{\sigma_{xz}\sigma_{zy}\sigma_{yx}}).$$

Note that the balanced triangles containing x and y do not contribute to the expression in (3.22).

PROPOSITION 3.12. *Under the setting of (3.15) and (3.16), the nontrivial block of $\Gamma_2^\sigma(x)$, which is Hermitian and of size $|B_2(x)| \times |B_2(x)|$, is given by the following blocks:*

$$(3.23) \quad (4\Gamma_2^\sigma(x))_{xx} = (3d_x + d_x^2)\mathbf{I}_d;$$

$$(3.24) \quad (4\Gamma_2^\sigma(x))_{xy} = -(3 + d_x + |S_1(y) \cap S_2(x)| + \sharp_\Delta^\sigma(x, y)) \overline{\sigma_{xy}} \\ \text{for any } y \in S_1(x);$$

$$(3.25) \quad (4\Gamma_2^\sigma(x))_{xz} = \sum_{y, y \sim x, y \sim z} \overline{\sigma_{xy}\sigma_{yz}}, \text{ for any } z \in S_2(x);$$

$$(3.26) \quad (4\Gamma_2^\sigma(x))_{yy} = (5 - d_x + 3|S_1(y) \cap S_2(x)| + 4\sharp_\Delta(x, y))\mathbf{I}_d \\ \text{for any } y \in S_1(x);$$

- (3.27) $(4\Gamma_2^\sigma(x))_{y_1 y_2} = 2\overline{\sigma_{y_1 x} \sigma_{x y_2}} - 4\overline{\sigma_{y_1 y_2}}$
 for any $y_1, y_2 \in S_1(x)$, $y_1 \neq y_2$, where we use $\sigma_{y_1 y_2} = 0$ if $\{y_1, y_2\} \notin E$;
- (3.28) $(4\Gamma_2^\sigma(x))_{yz} = -2\overline{\sigma_{yz}}$
 for any $y \in S_1(x)$ and $z \in S_2(x)$, where we use $\sigma_{yz} = 0$ if $\{y, z\} \notin E$;
- (3.29) $(4\Gamma_2^\sigma(x))_{zz} = |S_1(z) \cap S_1(x)| I_d$ for any $z \in S_2(x)$;
- (3.30) $(4\Gamma_2^\sigma(x))_{z_1 z_2} = 0$ for any $z_1, z_2 \in S_2(x)$, $z_1 \neq z_2$.

Proof. This follows from a direct expansion of the identity

$$\Gamma_2^\sigma(f, g)(x) = \vec{f}^T \Gamma_2^\sigma(x) \vec{g} \quad \text{for any } f, g : V \rightarrow \mathbb{K}^d.$$

We omit the details here. \square

Remark 3.13. (i) The block $(4\Gamma_2^\sigma(x))_{xz}$ above is a signed version of the quantity $|S_1(z) \cap S_1(x)|$ in (3.21).

- (ii) When $y_1, y_2 \in S_1(x)$ are neighbors, i.e., $\{y_1, y_2\} \in E$, we have a triangle containing x, y_1 , and y_2 . Then the block $(4\Gamma_2^\sigma(x))_{y_1 y_2}$ can be rewritten as

$$-2(I_d + (I_d - \overline{\sigma_{y_1 x} \sigma_{x y_2} \sigma_{y_2 y_1}})) \overline{\sigma_{y_1 y_2}},$$

which describes the unbalancedness of this triangle.

3.5. Example of a signed triangle. We consider a particular example of a triangle graph \mathcal{C}_3 , which consists of three vertices x, y , and z , as shown in Figure 2. We set

$$(3.31) \quad \mu(x) = \mu(y) = \mu(z) = 2 \quad \text{and} \quad w_{xy} = w_{xz} = w_{yz} = 1.$$

Let $\sigma : E^{or} \rightarrow U(1) := \{z \in \mathbb{C}, |z| = 1\}$ be a signature on \mathcal{C}_3 . Assume that the signature of the cycle \mathcal{C}_3 is equal to (the conjugacy class of) $s \in U(1)$. Then σ is switching equivalent to the signature given in Figure 2, i.e.,

$$\sigma_{xy} = \sigma_{xz} = 1 \quad \text{and} \quad \sigma_{yz} = s.$$

PROPOSITION 3.14. *Let $(\mathcal{C}_3, \mu, \sigma)$ be as above and $s = \text{Sgn}(\mathcal{C}_3)$. Then it has constant ∞ -dimensional Ricci curvature at every vertex. As a function of s , $K_\infty(s) := K_\infty(\mathcal{C}_3, \mu, \sigma)$ is given by*

$$(3.32) \quad K_\infty(s) = \begin{cases} \frac{5}{4} & \text{if } s = 1; \\ \frac{5 - \sqrt{17 + 8\text{Re}(s)}}{8} & \text{otherwise.} \end{cases}$$





FIG. 2. A signed triangle.

FIG. 3. ∞ -dimensional Ricci curvature of a signed triangle.

Remark 3.15. The curvature (3.32) is illustrated in Figure 3 as a function of the variable $\operatorname{Re}(s)$. The function $K_\infty(s)$ “jumps” at $s = 1$. That is,

$$(3.33) \quad \lim_{s \rightarrow 1} K_\infty(s) = 0 \quad \text{but} \quad K_\infty(1) = \frac{5}{4} > 0.$$

We will show that such a “jump” appears in a more general setting in section 6.

Proof of Proposition 3.14. Since the curvature is switching invariant, we can switch the signature σ to be as shown in Figure 2 before calculating the curvature $K_\infty(\sigma; x)$ at x . In fact, one can do similar operations for calculating $K_\infty(y)$ and $K_\infty(z)$. So $(\mathcal{C}_3, \mu, \sigma)$ has constant curvature and we only need to calculate the curvature at x .

By the fact (3.18) and Proposition 3.12, we can obtain the corresponding matrices $\Gamma^\sigma(x)$ and $\Gamma_2^\sigma(x)$. Note that in this example, we choose a different measure (3.31) from that in (3.15). Hence these matrices differ by a scaling of $1/2$ and $1/4$, respectively. Therefore, under the current setting (3.31), we have

$$2\Gamma^\sigma(x) = \frac{1}{2} \begin{pmatrix} 2 & -1 & -1 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{pmatrix} \quad \text{and} \quad 4\Gamma_2^\sigma(x) = \frac{1}{4} \begin{pmatrix} 10 & -6+s & -6+\bar{s} \\ -6+\bar{s} & 7 & 2-4\bar{s} \\ -6+s & 2-4s & 7 \end{pmatrix}.$$

By Proposition 3.10, we need to solve the following semidefinite programming:

$$(3.34) \quad \begin{aligned} &\text{maximize } K \\ &\text{subject to } \Gamma_2^\sigma(x) \geq K\Gamma^\sigma(x). \end{aligned}$$

Inequality (3.34) is equivalent to positive semidefiniteness of the following matrix:

$$(3.35) \quad 16\Gamma_2^\sigma(x) - 16K\Gamma^\sigma(x) = \begin{pmatrix} 10-8K & -6+s+4K & -6+\bar{s}+4K \\ -6+\bar{s}+4K & 7-4K & 2-4\bar{s} \\ -6+s+4K & 2-4s & 7-4K \end{pmatrix}.$$

By Sylvester's criterion, this is equivalent to nonnegativity of all principle minors of the above matrix. Calculating these principle minors, we translate the semidefinite programming to the following problem:

$$\begin{aligned} & \text{maximize } K \\ & \text{subject to } 10 - 8K \geq 0, \quad 7 - 4K \geq 0, \\ & \quad 16K^2 - 8(6 + \operatorname{Re}(s))K + (33 + 12\operatorname{Re}(x)) \geq 0, \\ & \quad 16K^2 - 56K + (16\operatorname{Re}(s) + 29) \geq 0, \\ & \quad 8(1 - \operatorname{Re}(s))K^2 - 10(1 - \operatorname{Re}(s))K + (1 - \operatorname{Re}(s))^2 \geq 0. \end{aligned}$$

Rewriting the above inequalities, we obtain

$$\begin{aligned} & \text{maximize } K \\ & \text{subject to } K \leq 5/4, \quad K \leq 7/4, \\ & \quad K \geq (5 + \sqrt{17 + 8\operatorname{Re}(s)})/8 \text{ or } K \leq (5 - \sqrt{17 + 8\operatorname{Re}(s)})/8, \\ & \quad K \geq (7 + 2\sqrt{5 - 4\operatorname{Re}(s)})/4 \text{ or } K \leq (7 - 2\sqrt{5 - 4\operatorname{Re}(s)})/4, \\ & \quad K \geq (6 + \operatorname{Re}(s) + \sqrt{\operatorname{Re}(s)^2 + 3})/4 \text{ or } K \leq (6 + \operatorname{Re}(s) - \sqrt{\operatorname{Re}(s)^2 + 3})/4. \end{aligned}$$

One can check directly that (3.32) is the solution of this optimization problem. \square

Similarly, one can calculate the ∞ -dimensional Ricci curvature of longer cycles \mathcal{C}_N for $N \geq 4$.

PROPOSITION 3.16. *Let $(\mathcal{C}_N, \mu, \sigma)$ be a cycle of length N with the edge weights and measure μ given in (3.31) and $s = \operatorname{Sgn}(\mathcal{C}_N)$. Then $(\mathcal{C}_N, \mu, \sigma)$ has constant ∞ -dimensional Ricci curvature at every vertex. Moreover, we have*

$$(3.36) \quad K_\infty(\mathcal{C}_4, \mu, \sigma) = \begin{cases} 1 & \text{if } s = 1; \\ 0 & \text{otherwise,} \end{cases}$$

and, for $N \geq 5$,

$$(3.37) \quad K_\infty(\mathcal{C}_N, \mu, \sigma) = 0.$$

We remark that new examples of graphs (G, σ) satisfying the $CD^\sigma(0, \infty)$ inequality can be constructed by taking Cartesian products of known examples for various choices of the signature, edge weights, and vertex measure on the product graph. We refer to Appendix A for full details about the behavior of CD^σ inequalities on Cartesian product graphs.

3.6. Heat semigroup characterizations of CD^σ inequalities. In this subsection, we derive characterizations of the CD^σ inequality via the solution of the following associated continuous time heat equation,

$$(3.38) \quad \begin{cases} \frac{\partial u(x, t)}{\partial t} = \Delta^\sigma u(x, t), \\ u(x, 0) = f(x), \end{cases}$$

where $f : V \rightarrow \mathbb{K}^d$ is an initial function. The solution $u : V \times [0, \infty) \rightarrow \mathbb{K}^d$ is given by $P_t^\sigma f := e^{t\Delta^\sigma} f$, where P_t^σ is a linear operator on the space $\ell^2(V, \mathbb{K}^d; \mu)$. Clearly, we have $P_0^\sigma f = f$. It is straightforward to check the following properties of P_t^σ .

PROPOSITION 3.17. *The operator $P_t^\sigma, t \geq 0$, satisfies the following properties:*

- (i) P_t^σ is a self-adjoint operator on the space $\ell^2(V, \mathbb{K}^d; \mu)$;
- (ii) P_t^σ commutes with Δ^σ , i.e., $P_t^\sigma \Delta^\sigma = \Delta^\sigma P_t^\sigma$;
- (iii) $P_t^\sigma P_s^\sigma = P_{t+s}^\sigma$ for any $t, s \geq 0$.

The solution of the heat equation corresponding to the graph Laplacian Δ is simply denoted by $P_t := e^{t\Delta}$. The matrix P_t has the following additional properties besides the ones listed in Proposition 3.17.

- PROPOSITION 3.18. (i) *All matrix entries of P_t are real and nonnegative.*
(ii) *For any constant function c on V , we have $P_t c = c$.*

In particular, the above properties imply that for a function $f : V \rightarrow \mathbb{R}$ with $0 \leq f(x) \leq c$ for all $x \in V$, we have $0 \leq P_t f(x) \leq c$ for all $x \in V$.

Proof. Recall that Δ can be written as the matrix $(\text{diag}_\mu)^{-1}(A - \text{diag}_D)$, where diag_D and diag_μ are the diagonal matrices with $(\text{diag}_D)_{xx} = d_x$ and $(\text{diag}_\mu)_{xx} = \mu(x)$ for all $x \in V$, and A is the weighted adjacency matrix. Now we exploit the fact that

$$(3.39) \quad \text{all off-diagonal entries of } \Delta \text{ are nonnegative,}$$

and, therefore, we can choose $C > 0$ such that $\Delta + C \cdot \mathbf{I}_N$ is entrywise nonnegative. Then $e^{\Delta + C \cdot \mathbf{I}_N}$ is also entrywise nonnegative, which implies that the same holds for $P_t = e^{\Delta + C \cdot \mathbf{I}_N} \cdot e^{-C \cdot \mathbf{I}_N}$.

For the constant function c , we have

$$(3.40) \quad \Delta c = 0.$$

Therefore, we have $\frac{\partial}{\partial t} P_t c = 0$, which implies $P_t c = c$. \square

Remark 3.19. Note that the two facts (3.39) and (3.40) do not extend to general P_t^σ , even when σ only takes values from $O(1) = \{\pm 1\}$. Therefore Proposition 3.18 does not hold for the more general operators P_t^σ .

If $n = \infty$, the CD^σ inequality is equivalent to the following local functional inequalities of $P_t^\sigma f$.

THEOREM 3.20. *Let (G, μ, σ) be given. Then the following are equivalent:*

- (i) *The inequality $CD^\sigma(K, \infty)$ holds, i.e., for any function $f : V \rightarrow \mathbb{K}^d$, we have*

$$\Gamma_2^\sigma(f) \geq K \Gamma^\sigma(f).$$

- (ii) *For any function $f : V \rightarrow \mathbb{K}^d$ and $t \geq 0$, we have*

$$\Gamma^\sigma(P_t^\sigma f) \leq e^{-2Kt} P_t(\Gamma^\sigma(f)).$$

- (iii) *For any function $f : V \rightarrow \mathbb{K}^d$ and $t \geq 0$, we have*

$$P_t(|f|^2) - |P_t^\sigma f|^2 \geq \frac{1}{K}(e^{2Kt} - 1)\Gamma^\sigma(P_t^\sigma f),$$

where we replace $(e^{2Kt} - 1)/K$ by $2t$ in the case $K = 0$.

Remark 3.21. Theorem 3.20 is similar in spirit to [5, Proposition 3.3]. Note that Proposition 3.18(i), which is crucial for the proof of [5, Proposition 3.3], is not true for P_t^σ in general. However, with our definitions of the operators Γ^σ and Γ_2^σ , we avoid this difficulty.

Proof. (i) \Rightarrow (ii): For any $0 \leq s \leq t$, we consider

$$(3.41) \quad F(s) := e^{-2Ks} P_s(\Gamma^\sigma(P_{t-s}^\sigma f)).$$

Since $F(0) = \Gamma^\sigma(P_t^\sigma f)$ and $F(t) = e^{-2Kt} P_t(\Gamma^\sigma(f))$, it is enough to prove $\frac{d}{ds} F(s) \geq 0$. We calculate

$$e^{2Ks} \frac{d}{ds} F(s) = -2K P_s(\Gamma^\sigma(P_{t-s}^\sigma f)) + \Delta P_s(\Gamma^\sigma(P_{t-s}^\sigma f)) + P_s \left(\frac{d}{ds} \Gamma^\sigma(P_{t-s}^\sigma f) \right)$$

and

$$\frac{d}{ds} \Gamma^\sigma(P_{t-s}^\sigma f) = -\Gamma^\sigma(\Delta^\sigma P_{t-s}^\sigma f, P_{t-s}^\sigma f) - \Gamma^\sigma(P_{t-s}^\sigma f, \Delta^\sigma P_{t-s}^\sigma f).$$

Therefore, $\Delta P_s = P_s \Delta$ and the inequality $CD^\sigma(K, \infty)$ imply

$$\frac{d}{ds} F(s) = 2e^{-2Ks} P_s [\Gamma_2^\sigma(P_{t-s}^\sigma f) - K \Gamma^\sigma(P_{t-s}^\sigma f)] \geq 0,$$

where we used Proposition 3.18(i). This proves (ii).

(ii) \Rightarrow (iii): For $0 \leq s \leq t$, we consider

$$(3.42) \quad G(s) := P_s(|P_{t-s}^\sigma f|^2).$$

Note that $G(0) = |P_t^\sigma f|^2$ and $G(t) = P_t(|f|^2)$. Using the estimate (ii) and Proposition 3.17, we have

$$\begin{aligned} \frac{d}{ds} G(s) &= \Delta P_s(|P_{t-s}^\sigma f|^2) + P_s [-(P_{t-s}^\sigma f)^T (\overline{\Delta^\sigma P_{t-s}^\sigma f}) - (\Delta^\sigma P_{t-s}^\sigma f)^T \overline{P_{t-s}^\sigma f}] \\ &= 2P_s(\Gamma^\sigma(P_{t-s}^\sigma f)) \geq 2e^{2Ks} \Gamma^\sigma(P_t^\sigma f). \end{aligned}$$

Therefore, we obtain

$$G(t) - G(0) = \int_0^t \frac{d}{ds} G(s) ds \geq 2\Gamma^\sigma(P_t^\sigma f) \int_0^t e^{2Ks} ds = \frac{e^{2Kt} - 1}{K} \Gamma^\sigma(P_t^\sigma f).$$

This proves (iii).

(iii) \Rightarrow (i): Here, we consider the inequality (iii) at $t = 0$ and use the expansion

$$P_t^\sigma = \text{Id} + t\Delta^\sigma + \frac{t^2}{2}(\Delta^\sigma)^2 + o(t^2).$$

Dividing (iii) by $2t^2$ and letting t tend to zero, we obtain

$$\begin{aligned} &\frac{1}{4} \Delta^2(|f|^2) - \frac{1}{4} f^T \left(\overline{(\Delta^\sigma)^2 f} \right) - \frac{1}{4} ((\Delta^\sigma)^2 f)^T \bar{f} - \frac{1}{2} (\Delta^\sigma f)^T (\overline{\Delta^\sigma f}) \\ &\geq K \Gamma^\sigma(f) + \Gamma^\sigma(f, \Delta^\sigma f) + \Gamma^\sigma(\Delta^\sigma f, f). \end{aligned}$$

Using Definition 1.5, the above inequality simplifies to

$$\Gamma_2^\sigma(f) \geq K \Gamma^\sigma(f),$$

which shows (i). \square

4. Multiway cheeger constants with signatures. In this section, we introduce multiway Cheeger constants with signatures for graphs (G, μ, σ) .

4.1. Cheeger constants with signatures. Following the ideas of [32], we introduce a Cheeger type constant of (G, μ, σ) as a mixture of a frustration index and the expansion rate. For any nonempty subset $S \subseteq V$, the frustration index $\iota^\sigma(S)$ is a measure of the unbalancedness of the signature σ on the induced subgraph of S . For that purpose, we need to choose a norm on H , to measure the distance between different elements in H .

DEFINITION 4.1. *Given a $(d \times d)$ -matrix $A = (a_{ij})$, we define the average $(2, 1)$ -norm $|A|_{2,1}$ of A as*

$$(4.1) \quad |A|_{2,1} := \frac{1}{d} \sum_{i=1}^d \left(\sum_{j=1}^d |a_{ij}|^2 \right)^{\frac{1}{2}}.$$

If we denote the vector of the i th column of A by A^i , this norm can be rewritten as $|A|_{2,1} := \frac{1}{d} \sum_{i=1}^d |A^i|$. Recall that $|A^i|^2 := (A^i)^T A^i$.

Remark 4.2.

- (i) The average $(2, 1)$ -norm is smaller or equal to the Frobenius norm (alternatively called the Hilbert–Schmidt norm), i.e., we have for any $(d \times d)$ -matrix $A = (a_{ij})$,

$$(4.2) \quad |A|_{2,1} \leq \frac{1}{\sqrt{d}} |A|_F,$$

where $|A|_F := (\sum_{i,j=1}^d |a_{ij}|^2)^{\frac{1}{2}}$ denotes the Frobenius norm of A . This is a straightforward consequence of the Cauchy–Schwarz inequality directly.

- (ii) The average $(2, 1)$ -norm is not submultiplicative in general, i.e., $|AB|_{2,1} \leq |A|_{2,1} |B|_{2,1}$ is not necessarily true for any $(d \times d)$ -matrices A and B . However, if $B \in O(d)$ or $U(d)$, we have

$$(4.3) \quad |BA|_{2,1} = |A|_{2,1}.$$

Note that in this case, $|B|_{2,1} = 1$.

DEFINITION 4.3 (frustration index). *Let (G, μ, σ) be given. We define the frustration index $\iota^\sigma(S)$ for $\emptyset \neq S \subseteq V$ as*

$$\begin{aligned} \iota^\sigma(S) &:= \min_{\tau: S \rightarrow H} \sum_{\{x,y\} \in E_S} w_{xy} |\sigma_{xy} \tau(y) - \tau(x)|_{2,1} \\ &= \min_{\tau: S \rightarrow H} \sum_{\{x,y\} \in E_S} w_{xy} |\sigma_{xy}^\tau - id|_{2,1}, \end{aligned}$$

where E_S is the edge set of the induced subgraph of S in G .

Remark 4.4.

- (i) By (4.3), the quantity

$$|\sigma_{xy} \tau(y) - \tau(x)|_{2,1} = |\sigma_{yx} \tau(x) - \tau(y)|_{2,1}$$

is independent of the orientation of the edge $\{x, y\} \in E$. Hence, the summation $\sum_{\{x,y\} \in E_S} w_{xy} |\sigma_{xy} \tau(y) - \tau(x)|_{2,1}$ is well defined.

- (ii) In the definition of the frustration index, we are taking the infimum over all possible switching functions. Hence, the frustration index is a switching invariant quantity.

- (iii) The average $(2, 1)$ -norm is only one possible choice which can be used in the definition of the frustration index. A more canonical norm to be used is the Frobenius norm. However, having the aim to present the strongest Buser type inequalities (5.1) in section 5, we choose the average $(2, 1)$ -norm here (recall (4.2)).

We denote by $|E(S, V \setminus S)|$ the boundary measure of $S \subseteq V$, which is given by

$$|E(S, V \setminus S)| := \sum_{x \in S} \sum_{y \notin S} w_{xy}.$$

In the above, we use the convention that $w_{xy} = 0$ if $x \not\sim y$. The μ -volume of S is given by

$$\mu(S) := \sum_{x \in S} \mu(x).$$

DEFINITION 4.5. We call k subsets $\{S_i\}_{i=1}^k$ of V a nontrivial k -subpartition of V if all S_i are nonempty and pairwise disjoint.

Now we are prepared to define the Cheeger constant.

DEFINITION 4.6 (Cheeger constant). Let (G, μ, σ) be given. The k -way Cheeger constant h_k^σ is defined as

$$h_k^\sigma := \min_{\{S_i\}_{i=1}^k} \max_{1 \leq i \leq k} \phi^\sigma(S_i),$$

where the minimum is taken over all possible nontrivial k -subpartitions $\{S_i\}_{i=1}^k$ of V and

$$\phi^\sigma(S) := \frac{\iota^\sigma(S) + |E(S, V \setminus S)|}{\mu(S)}.$$

Note that the multiway Cheeger constants defined above are switching invariant. Definition 4.6 is a natural extension of the Cheeger constants developed in [4, 32] and is related to the constants discussed in [7].

Remark 4.7 (relations with Bandeira, Singer, and Spielman's constants). In [7], Bandeira, Singer, and Spielman introduced the so-called $O(d)$ frustration ℓ_1 constant $\nu_{G,1}$ as follows:

$$\nu_{G,1} := \min_{\tau: V \rightarrow O(d)} \frac{1}{\sqrt{d}\mu(V)} \sum_{x,y \in V} w_{xy} |\sigma_{xy}\tau(y) - \tau(x)|_F,$$

where $|\cdot|_F$ denotes the Frobenius norm of a matrix. Modifying $\nu_{G,1}$ by also allowing zero matrices in the image of τ , we obtain

$$\nu_{G,1}^* := \min_{\tau: V \rightarrow H \cup \{0\}} \frac{\sum_{x,y \in V} w_{xy} |\sigma_{xy}\tau(y) - \tau(x)|_F}{\sum_{x \in V} \mu(x) |\tau(x)|_F},$$

where we denote the $(d \times d)$ -zero matrix by 0. Note that $|\tau(x)|_F = \sqrt{d}$, for $\tau(x) \in H$.

We observe the following relation between our Cheeger constant h_1^σ and the constant $\nu_{G,1}^*$:

$$h_1^\sigma \leq \frac{1}{2} \nu_{G,1}^*,$$

which is verified by the calculation

$$\begin{aligned} h_1^\sigma &= \min_{\emptyset \neq S \subseteq V} \frac{\iota^\sigma(S) + |E(S, V \setminus S)|}{\mu(S)} \\ &= \min_{\tau: V \rightarrow H \cup \{0\}} \frac{\sum_{\{x,y\} \in E} w_{xy} |\sigma_{xy} \tau(y) - \tau(x)|_{2,1}}{\sum_{x \in V} \mu(x) |\tau(x)|_{2,1}} \\ &\leq \min_{\tau: V \rightarrow H \cup \{0\}} \frac{\sum_{\{x,y\} \in E} w_{xy} |\sigma_{xy} \tau(y) - \tau(x)|_F}{\sqrt{d} \sum_{x \in V} \mu(x) |\tau(x)|_{2,1}} \\ &= \frac{1}{2} \nu_{G,1}^*. \end{aligned}$$

In the inequality above, we used (4.2).

For convenience, we call $\{S_i\}_{i=1}^k$ a *connected*, nontrivial k -subpartition of V if all sets $S_i \subseteq V$ are nonempty and pairwise disjoint and if every subgraph induced by S_i is connected. Then the Cheeger constants introduced in Definition 4.6 do not change if we restrict our considerations to connected, nontrivial k -subpartitions:

LEMMA 4.8. *Let (G, μ, σ) be given. Then we have*

$$h_k^\sigma = \min_{\{S_i\}_{i=1}^k} \max_{1 \leq i \leq k} \phi^\sigma(S_i),$$

where the minimum is taken over all possible connected, nontrivial k -subpartitions $\{S_i\}_{i=1}^k$ of V .

Proof. Let $\{S_i\}_{i=1}^k$ be a possibly nonconnected, nontrivial k -subpartition achieving h_k^σ . Suppose S_i has the connected components $W_i^1, \dots, W_i^{n(i)}$. Then,

$$\phi^\sigma(S_i) \mu(S_i) = \sum_{j=1}^{n(i)} \phi^\sigma(W_i^j) \mu(W_i^j)$$

and $\mu(S_i) = \sum_{j=1}^n \mu(W_i^j)$. Hence, there exists $j(i) \in \{1, 2, \dots, n(i)\}$ such that $\phi^\sigma(W_i^{j(i)}) \leq \phi^\sigma(S_i)$. Consequently,

$$\max_{1 \leq i \leq k} \phi^\sigma(W_i^{j(i)}) \leq \max_{1 \leq i \leq k} \phi^\sigma(S_i) = h_k^\sigma,$$

and thus $\{W_i^{j(i)}\}_{i=1}^k$ is a connected, nontrivial k -subpartition of V with

$$h_k^\sigma = \max_{1 \leq i \leq k} \phi^\sigma(W_i^{j(i)}).$$

This implies the lemma. \square

4.2. Frustration index via spanning trees. This subsection is motivated by the following question: Is there any easier way to calculate the frustration index $\iota^\sigma(S)$ of a subset $S \subseteq V$? We will provide an affirmative answer in the case $H = U(1)$.

Note that the average (2,1)-norm reduces to the absolute value of a complex number, and the frustration index $\iota^\sigma(S)$ for $S \subseteq V$ simplifies to

$$\iota^\sigma(S) := \min_{\tau: S \rightarrow U(1)} \sum_{\{x,y\} \in E_S} w_{xy} |\sigma_{xy} \tau(y) - \tau(x)|,$$

where E_S is the edge set of the induced subgraph of S . Here our aim is to make it easier to calculate $\iota^\sigma(S)$ by considering all spanning trees of the induced subgraph of S and taking the minimum over so-called *constant functions on these trees with*

respect to the signature. This reduces the original minimization problem to a finite combinatorial problem. We will show in Appendix B via a counterexample that this reduction is no longer possible in the case of higher dimensional signature groups.

DEFINITION 4.9. Let (G, σ) be a finite, connected graph the signature $\sigma : E^{or} \rightarrow H$. A function $\tau : V \rightarrow H$ is constant on G with respect to σ if, for all $(x, y) \in E^{or}$, we have

$$\sigma_{xy}\tau(y) = \tau(x).$$

In other words, τ is a switching function such that σ^τ is trivial, i.e., $\sigma_{xy}^\tau = id \in H$ for all $(x, y) \in E^{or}$.

Let $T = (S, E_T)$, $E_T \subseteq E_S$, be a spanning tree of the induced subgraph of S . We write $C_T(S) := \{\tau : S \rightarrow U(1) : \tau \text{ is constant on } T \text{ with respect to } \sigma\}$. Moreover, we define \mathbb{T}_S as the set of all spanning trees of the induced subgraph of S .

Since T is a tree, the set $C_T(S)$ is not empty. Since T is a spanning tree, we have $C_T(S) = \tau U(1) := \{\tau z : S \rightarrow U(1) \mid z \in U(1)\}$ for any $\tau \in C_T(S)$.

THEOREM 4.10. Let $S \subseteq V$ be a nonempty subset of V which induces a connected subgraph. Then,

$$(4.4) \quad \iota^\sigma(S) = \min_{T \in \mathbb{T}_S} \sum_{\{x, y\} \in E_S} w_{xy} |\sigma_{xy} \tau_T(y) - \tau_T(x)|,$$

where τ_T denotes an arbitrary representative of $C_T(S)$.

Moreover, if a function $\tau : S \rightarrow U(1)$ satisfies $\sum_{\{x, y\} \in E_S} w_{xy} |\sigma_{xy} \tau(y) - \tau(x)| = \iota^\sigma(S)$, then there is a spanning tree $T = (S, E_T)$ such that τ is constant on T with respect to σ .

We remark that in (4.4) we are taking the minimum over a finite set. Moreover, given a spanning tree $T \in \mathbb{T}_S$, only terms associated to edges of E_S not belonging to the spanning tree contribute to the sum.

Theorem 4.10 can be considered an extension of [24, Theorem 2], where Harary and Kabell derived this result on unweighted graphs for the case $H = O(1) = \{\pm 1\}$. Their proof depends in an essential way on the fact that the frustration index in their setting (which they called *line index of balance*) only assumes integer values. Therefore, their proof cannot be extended to the current general setting.

We first prove a basic lemma.

LEMMA 4.11. Let $Z := \{z_1, \dots, z_n\} \subset U(1)$ and $w_1, \dots, w_n > 0$. Then we have

$$(4.5) \quad \min_{z \in U(1)} \sum_{k=1}^n w_k |z - z_k| = \min_{z \in Z} \sum_{k=1}^n w_k |z - z_k|.$$

Moreover, if $z \in U(1) \setminus Z$, then

$$\sum_{k=1}^n w_k |z - z_k| > \min_{z \in Z} \sum_{k=1}^n w_k |z - z_k|.$$

Proof. The minimum over on the left-hand side of (4.5) exists, since $U(1)$ is compact and $\sum_{k=1}^n w_k |z - z_k|$ is continuous in z . Suppose that the minimum is assumed in $z_0 = e^{it_0}$ with $z_0 \notin Z$. That is, the function $\phi : \mathbb{R} \rightarrow \mathbb{R}, t \mapsto \sum_{k=1}^n w_k |e^{it} - z_k|$ assumes a local minimum in t_0 . Since $z_0 \notin Z$, the second derivative ϕ'' exists at t_0 and is not negative due to the minimum property. But for all $k \in \{1, \dots, n\}$, we can set $z_k = e^{it_k}$ and compute

$$\frac{d^2}{dt^2} |e^{it} - e^{it_k}|(t_0) = 2 \frac{d^2}{dt^2} \left| \sin \frac{t - t_k}{2} \right| (t_0) < 0.$$

This is a contradiction and, hence, $z_0 \in Z$. This finishes the proof of the lemma. \square

Now, we prove the theorem with the help of the lemma.

Proof of Theorem 4.10. First, we notice that the expression $w_{xy}|\sigma_{xy}\tau_T(y) - \tau_T(x)|$ does not depend on the choice of $\tau_T \in C_T(S)$ since $C_T(S) = \tau_T U(1)$. Hence, the restriction to the representative of $C_T(S)$ makes sense.

Let $\tau_0 : S \rightarrow U(1)$ be a minimizer of $\sum_{\{x,y\} \in E_S} w_{xy}|\sigma_{xy}\tau(y) - \tau(x)|$. Denote by

$$E_0 := \{\{x, y\} \in E_S : \sigma_{xy}\tau_0(y) = \tau_0(x)\}$$

the set of edges where τ_0 is constant with respect to σ . It is sufficient to show that $G_0 = (S, E_0)$ is connected since then there is a spanning tree T_0 of G_0 such that τ_0 is constant on T_0 with respect to σ .

Suppose G_0 is not connected. Then there is a connected component $W \subsetneq S$. Denote $\partial_S W := \{(x, y) \in E_S^{or} : x \in W, y \in S \setminus W\}$. We have $\partial_S W \neq \emptyset$ since S is connected. Moreover, we have $\sigma_{xy}\tau_0(y) \neq \tau_0(x)$ for all $(x, y) \in \partial_S W$, since W is a connected component and, otherwise, y would also belong to W , contradicting $(x, y) \in \partial_S W$.

The previous lemma states that

$$\min_{z \in U(1)} \sum_{(x,y) \in \partial_S W} w_{xy}|\sigma_{xy}\tau_0(y) - z\tau_0(x)| = \min_{z \in U(1)} \sum_{(x,y) \in \partial_S W} w_{xy}|\sigma_{xy}\tau_0(y)\overline{\tau_0(x)} - z|$$

achieves the minimum only in elements of the set $\{\sigma_{xy}\tau_0(y)\overline{\tau_0(x)} : (x, y) \in \partial_S W\}$. But $1 \notin \{\sigma_{xy}\tau_0(y)\overline{\tau_0(x)} : (x, y) \in \partial_S W\}$, since $\sigma_{xy}\tau_0(y) \neq \tau_0(x)$ for all $(x, y) \in \partial_S W$. Hence, there exists $z_0 \in U(1)$ such that

$$(4.6) \quad \sum_{(x,y) \in \partial_S W} w_{xy}|\sigma_{xy}\tau_0(y) - z_0\tau_0(x)| < \sum_{(x,y) \in \partial_S W} w_{xy}|\sigma_{xy}\tau_0(y) - \tau_0(x)|.$$

We define $\tau_1 : S \rightarrow U(1)$,

$$\tau_1(x) := \begin{cases} z_0\tau_0(x) & \text{if } x \in W; \\ \tau_0(x) & \text{if } x \in S \setminus W. \end{cases}$$

Consequently,

$$\begin{aligned} & \sum_{\{x,y\} \in E_S} w_{xy}|\sigma_{xy}\tau_1(y) - \tau_1(x)| \\ &= \sum_{\{x,y\} \in E_W \cup E_{S \setminus W}} w_{xy}|\sigma_{xy}\tau_1(y) - \tau_1(x)| + \sum_{(x,y) \in \partial_S W} w_{xy}|\sigma_{xy}\tau_1(y) - \tau_1(x)| \\ &= \sum_{\{x,y\} \in E_W \cup E_{S \setminus W}} w_{xy}|\sigma_{xy}\tau_0(y) - \tau_0(x)| + \sum_{(x,y) \in \partial_S W} w_{xy}|\sigma_{xy}\tau_0(y) - z_0\tau_0(x)| \\ &\stackrel{(4.6)}{<} \sum_{\{x,y\} \in E_W \cup E_{S \setminus W}} w_{xy}|\sigma_{xy}\tau_0(y) - \tau_0(x)| + \sum_{(x,y) \in \partial_S W} w_{xy}|\sigma_{xy}\tau_0(y) - \tau_0(x)| \\ &= \sum_{\{x,y\} \in E_S} w_{xy}|\sigma_{xy}\tau_0(y) - \tau_0(x)|. \end{aligned}$$

This is a contradiction to the fact that τ_0 is a minimizer of $\sum_{\{x,y\} \in E_S} w_{xy} |\sigma_{xy} \tau(y) - \tau(x)|$. Thus, G_0 has to be connected. This finishes the proof. \square

Recall from Lemma 4.8 that the Cheeger constant h_k^σ is the minimum of $\max_{1 \leq i \leq k} \phi^\sigma(S_i)$ over all possible *connected*, nontrivial k -subpartitions $\{S_i\}_{i=1}^k$. Therefore, Theorem 4.10 implies that the calculation of h_k^σ reduces to a finite combinatorial minimization problem if $H = U(1)$.

5. Buser inequalities. In this section, we prove our main theorem, namely, higher order Buser type inequalities for nonnegatively curved graphs (cf. Theorem 1.1 in the introduction).

THEOREM 5.1 (main theorem). *Let (G, μ, σ) satisfy $CD^\sigma(0, \infty)$. Then for all $1 \leq k \leq N$, we have*

$$(5.1) \quad \sqrt{\lambda_{kd}^\sigma} \leq 4\sqrt{D_G^{nor}} \left(kd\sqrt{\log(2kd)} \right) h_k^\sigma.$$

Before we present the proof, we first discuss the following two lemmata. We will use the following notation for the $\ell^p(V, \mathbb{K}^d; \mu)$ norm of functions, $1 \leq p \leq \infty$,

$$\|f\|_{p,\mu} := \left(\sum_{x \in V} \mu(x) |f(x)|^p \right)^{\frac{1}{p}}.$$

For simplicity, we omit the subscript μ in the following arguments.

LEMMA 5.2. *Assume that (G, μ, σ) satisfies $CD^\sigma(0, \infty)$. Then for any function $f : V \rightarrow \mathbb{K}^d$ and $t \geq 0$, we have*

$$(5.2) \quad \|f - P_t^\sigma f\|_1 \leq \sqrt{2t} \|\sqrt{\Gamma^\sigma(f)}\|_1.$$

Proof. First, the equivalent formulation of the $CD^\sigma(0, \infty)$ inequality in Theorem 3.20(iii) implies that

$$(5.3) \quad \|\sqrt{P_t}(|f|^2)\|_\infty \geq \sqrt{2t} \|\sqrt{\Gamma^\sigma(P_t^\sigma f)}\|_\infty.$$

The inequality (5.2) is actually a dual version of the above one. We set

$$g(x) := \begin{cases} 0 & \text{if } f(x) - P_t^\sigma f(x) = 0; \\ (f(x) - P_t^\sigma f(x))/|f(x) - P_t^\sigma f(x)| & \text{otherwise,} \end{cases}$$

and calculate

$$\begin{aligned} \|f - P_t^\sigma f\|_1 &= \langle f - P_t^\sigma f, g \rangle_\mu = \left\langle - \int_0^t \frac{\partial}{\partial s} P_s^\sigma f ds, g \right\rangle_\mu \\ &= - \int_0^t \langle \Delta^\sigma f, P_s^\sigma g \rangle_\mu ds \\ &= \int_0^t \sum_{x \in V} \mu(x) \Gamma^\sigma(f, P_s^\sigma g)(x) ds, \end{aligned}$$

where we used the self-adjointness of P_t^σ and the summation by part formula (3.3). We further apply Proposition 3.1 and the estimate (5.3) to derive

$$\begin{aligned}
\|f - P_t^\sigma f\|_1 &\leq \int_0^t \sum_{x \in V} \mu(x) \sqrt{\Gamma^\sigma(f)(x)} \sqrt{\Gamma^\sigma(P_s^\sigma g)(x)} ds \\
&\leq \int_0^t \|\sqrt{\Gamma^\sigma(f)}\|_1 \|\sqrt{\Gamma^\sigma(P_s^\sigma g)}\|_\infty ds \\
&\leq \|\sqrt{\Gamma^\sigma(f)}\|_1 \int_0^t \frac{1}{\sqrt{2s}} \|\sqrt{P_s(|g|^2)}\|_\infty ds \\
&\leq \sqrt{2t} \|\sqrt{\Gamma^\sigma(f)}\|_1.
\end{aligned}$$

In the last inequality we used the fact $P_s(|g|^2) \leq \|g\|^2_\infty = 1$, which follows from Proposition 3.18. \square

We still need the following technical lemma.

LEMMA 5.3. *For any function $f : V \rightarrow \mathbb{K}^d$, we have*

$$(5.4) \quad \|\sqrt{\Gamma^\sigma(f)}\|_1 \leq \sqrt{2D_G^{nor}} \sum_{\{x,y\} \in E} w_{xy} |\sigma_{xy} f(y) - f(x)|.$$

Proof. It is straightforward to calculate

$$\begin{aligned}
\|\sqrt{\Gamma^\sigma(f)}\|_1 &= \sum_{x \in V} \mu(x) \sqrt{\frac{1}{2\mu(x)} \sum_{y, y \sim x} w_{xy} |\sigma_{xy} f(y) - f(x)|^2} \\
&\leq \sum_{x \in V} \sqrt{\frac{\mu(x)}{2}} \sum_{y, y \sim x} \sqrt{w_{xy}} |\sigma_{xy} f(y) - f(x)| \\
&\leq \sqrt{\frac{D_G^{nor}}{2}} \sum_{x \in V} \sum_{y, y \sim x} w_{xy} |\sigma_{xy} f(y) - f(x)|.
\end{aligned}$$

This simplifies to (5.4), since the summands above are symmetric w.r.t. x and y . \square

Now, we have all ingredients for the proof of the Buser type inequality (5.1).

Proof of Theorem 5.1. Let $\{S_i\}_{i=1}^k$ be an arbitrary nontrivial k -subpartition of V . For each S_i , let $\tau_i : S_i \rightarrow H$ be the function achieving the values $\iota^\sigma(S_i)$ introduced in Definition 4.3. We extend each τ_i trivially to a function on V by assigning zero matrices to the vertices in $V \setminus S_i$. By abuse of notation, we denote this extension, again, by $\tau_i : V \rightarrow H$. Each τ_i gives rise to d pairwise orthogonal functions in $\ell^2(V, \mathbb{K}^d; \mu)$:

$$(5.5) \quad \tau_i^l : V \rightarrow \mathbb{K}^d, \quad x \mapsto (\tau_i(x))^l, \quad l = 1, 2, \dots, d,$$

where $(\tau_i(x))^l$ denotes the l th column vector of the matrix $\tau_i(x)$. Note that for $x \in S_i$, we have $|\tau_i^l(x)| = 1$.

For every $1 \leq i \leq k$, we apply Lemma 5.3 to obtain

$$\begin{aligned}
&\frac{1}{d} \sum_{l=1}^d \|\sqrt{\Gamma^\sigma(\tau_i^l)}\|_1 \\
&\leq \frac{1}{d} \sum_{l=1}^d \sqrt{2D_G^{nor}} \left(\sum_{\{x,y\} \in E_{S_i}} w_{xy} |\sigma_{xy} \tau_i^l(y) - \tau_i^l(x)| + |E(S_i, V \setminus S_i)| \right) \\
(5.6) \quad &\leq \sqrt{2D_G^{nor}} (\iota^\sigma(S_i) + |E(S_i, V \setminus S_i)|).
\end{aligned}$$

On the other hand, we have by Lemma 5.2,

$$\begin{aligned}\sqrt{2t}\|\sqrt{\Gamma^\sigma(\tau_i^l)}\|_1 &\geq \sum_{x \in V} \mu(x) |\tau_i^l(x) - P_t^\sigma \tau_i^l(x)| \\ &\geq \sum_{x \in V} \mu(x) |\tau_i^l(x) - P_t^\sigma \tau_i^l(x)| \cdot |\tau_i^l(x)| \\ &\geq \sum_{x \in V} \mu(x) \operatorname{Re} \left((\tau_i^l(x))^T \overline{(\tau_i^l(x) - P_t^\sigma \tau_i^l(x))} \right),\end{aligned}$$

where $\operatorname{Re}(\cdot)$ denotes the real part of a complex number, and we used the Cauchy–Schwarz inequality in the last inequality. By Proposition 3.17, we continue to calculate

$$(5.7) \quad \sqrt{2t}\|\sqrt{\Gamma^\sigma(\tau_i^l)}\|_1 \geq \operatorname{Re} \left(\langle \tau_i^l, \tau_i^l - P_t^\sigma \tau_i^l \rangle_\mu \right) = \|\tau_i^l\|_2^2 - \|P_{t/2}^\sigma \tau_i^l\|_2^2.$$

Let $\{\psi_n\}_{n=1}^{Nd}$ be an orthonormal basis of $\ell^2(V, \mathbb{K}^d; \mu)$ consisting of the eigenfunctions corresponding to $\{\lambda_n^\sigma\}_{n=1}^{Nd}$, respectively. Setting

$$\alpha_{i,n}^l := \langle \tau_i^l, \psi_n \rangle_\mu,$$

we have

$$(5.8) \quad \sum_{n=1}^{Nd} |\alpha_{i,n}^l|^2 = \|\tau_i^l\|_2^2 = \mu(S_i)$$

and

$$(5.9) \quad \|P_{t/2}^\sigma \tau_i^l\|_2^2 = \sum_{n=1}^{Nd} e^{-t\lambda_n^\sigma} |\alpha_{i,n}^l|^2.$$

Now (5.6), (5.7), (5.8), and (5.9) together imply, for each $1 \leq i \leq k$,

$$\begin{aligned}(5.10) \quad 2\sqrt{D_G^{nor}} t \phi^\sigma(S_i) &\geq \frac{1}{d} \sum_{l=1}^d \left(1 - \sum_{n=1}^{Nd} e^{-t\lambda_n^\sigma} \frac{|\alpha_{i,n}^l|^2}{\mu(S_i)} \right) \\ &\geq \frac{1}{d} \sum_{l=1}^d \left(1 - \sum_{n=1}^{kd-1} \frac{|\alpha_{i,n}^l|^2}{\mu(S_i)} - e^{-t\lambda_{kd}^\sigma} \sum_{n=kd}^{Nd} \frac{|\alpha_{i,n}^l|^2}{\mu(S_i)} \right) \\ &\geq 1 - \frac{1}{d} \sum_{l=1}^d \sum_{n=1}^{kd-1} \frac{|\alpha_{i,n}^l|^2}{\mu(S_i)} - e^{-t\lambda_{kd}^\sigma}.\end{aligned}$$

By (5.8), we know

$$1 - \frac{1}{d} \sum_{l=1}^d \sum_{n=1}^{kd-1} \frac{|\alpha_{i,n}^l|^2}{\mu(S_i)} \geq 0,$$

but our aim is to show that for some $i_0 \in \{1, 2, \dots, k\}$ this expression is strictly positive. We rewrite the summands as follows:

$$(5.11) \quad \frac{|\alpha_{i,n}^l|^2}{\mu(S_i)} = \left| \left\langle \frac{\tau_i^l}{\sqrt{\mu(S_i)}}, \psi_n \right\rangle \right|^2.$$

Since the functions $\tau_i^l / \sqrt{\mu(S_i)}$, $i = 1, 2, \dots, k$, $l = 1, 2, \dots, d$, are orthonormal in the space $\ell^2(V, \mathbb{K}^d; \mu)$, we obtain

$$(5.12) \quad \sum_{i=1}^k \sum_{l=1}^d \frac{|\alpha_{i,n}^l|^2}{\mu(S_i)} \leq \|\psi_n\|_2^2 = 1.$$

Summation over n yields

$$(5.13) \quad \sum_{i=1}^k \sum_{l=1}^d \sum_{n=1}^{kd-1} \frac{|\alpha_{i,n}^l|^2}{\mu(S_i)} \leq kd - 1.$$

Consequently, there exists an $i_0 \in \{1, 2, \dots, k\}$ such that

$$(5.14) \quad \frac{1}{d} \sum_{l=1}^d \sum_{n=1}^{kd-1} \frac{|\alpha_{i_0,n}^l|^2}{\mu(S_{i_0})} \leq 1 - \frac{1}{kd}.$$

We insert this estimate into inequality (5.10) to obtain

$$(5.15) \quad 2\sqrt{D_G^{nor}t} \max_{1 \leq i \leq k} \phi^\sigma(S_i) \geq \frac{1}{kd} - e^{-t\lambda_{kd}^\sigma}.$$

Since the k -subpartition was chosen arbitrarily, we have

$$(5.16) \quad 2\sqrt{D_G^{nor}t} \cdot h_k^\sigma \geq \frac{1}{kd} - e^{-t\lambda_{kd}^\sigma}.$$

For $\lambda_{kd}^\sigma \neq 0$, we choose $t = \log(2dk)/\lambda_{kd}^\sigma$ to obtain

$$(5.17) \quad 4\sqrt{D_G^{nor}kd} \sqrt{\log(2dk)} h_k^\sigma \geq \sqrt{\lambda_{kd}^\sigma}.$$

This completes the proof. \square

Recall from Corollary 3.8 that any graph (G, μ, σ) has a specific finite lower curvature bound. In case of a negative lower curvature bound, we have the following result. For a subset $S \subseteq V$, we define the following constant, which is no greater than $\iota^\sigma(S)$:

$$(5.18) \quad \tilde{\iota}^\sigma(S) := \min_{\substack{f: S \rightarrow \mathbb{K}^d \\ |f(x)|=1, \forall x \in S}} \sum_{\{x,y\} \in E_S} w_{xy} |\sigma_{xy} f(y) - f(x)|.$$

Using this constant, we have the following isoperimetric type inequality.

THEOREM 5.4. *Let (G, μ, σ) satisfy $CD^\sigma(-K, \infty)$, for $K \geq 0$. Then for any subset $\emptyset \neq S \subseteq V$,*

$$(5.19) \quad \tilde{\iota}^\sigma(S) + |E(S, V \setminus S)| \geq \frac{1}{2\sqrt{2D_G^{nor}}} \min \left\{ (1 - e^{-1})\sqrt{\lambda_1^\sigma}, \frac{\lambda_1^\sigma}{2\sqrt{2K}} \right\} \mu(S).$$

Proof. Modifying the proof of Lemma 5.2 for $K \geq 0$, we derive from the inequality $CD^\sigma(-K, \infty)$ that for any function $f: V \rightarrow \mathbb{K}^d$,

$$(5.20) \quad \|f - P_t^\sigma f\|_1 \leq \int_0^t \sqrt{\frac{K}{1 - e^{-2Ks}}} ds \|\sqrt{\Gamma^\sigma(f)}\|_1.$$

Using $1 - e^{-u} \geq u/2$ for $0 \leq u \leq 1$, we have for $0 \leq t \leq 1/(2K)$,

$$(5.21) \quad \|f - P_t^\sigma f\|_1 \leq 2\sqrt{t} \|\sqrt{\Gamma^\sigma(f)}\|_1.$$

Let S be an arbitrary nonempty subset of V . Let $f^0 : S \rightarrow \mathbb{K}^d$ with $|f^0(x)| = 1$ for all $x \in S$ be the function achieving the value of $\tilde{\iota}^\sigma(S)$. By similar reasoning as in the proof of Theorem 5.1, we obtain for $0 \leq t \leq 1/(2K)$,

$$(5.22) \quad 2\sqrt{2D_G^{nor}t} (\tilde{\iota}^\sigma(S) + |E(S, V \setminus S)|) \geq \mu(S)(1 - e^{-t\lambda_1^\sigma}).$$

If $\lambda_1^\sigma \geq 2K$, we set $t = 1/\lambda_1^\sigma$ and obtain

$$(5.23) \quad 2\sqrt{2D_G^{nor}} (\tilde{\iota}^\sigma(S) + |E(S, V \setminus S)|) \geq \mu(S)(1 - e^{-1})\sqrt{\lambda_1^\sigma}.$$

If $\lambda_1^\sigma < 2K$, we set $t = 1/(2K)$ and obtain

$$(5.24) \quad 2\sqrt{\frac{D_G^{nor}}{K}} (\tilde{\iota}^\sigma(S) + |E(S, V \setminus S)|) \geq \mu(S)(1 - e^{-\frac{\lambda_1^\sigma}{2K}}) \geq \mu(S) \frac{\lambda_1^\sigma}{4K}.$$

Combining both cases completes the proof. \square

We now define the following Cheeger type constant \widetilde{h}_1^σ corresponding to $\tilde{\iota}^\sigma(S)$.

DEFINITION 5.5. Let (G, μ, σ) be given. The constant \widetilde{h}_1^σ is defined as

$$\widetilde{h}_1^\sigma = \min_{\emptyset \neq S \subseteq V} \frac{\tilde{\iota}^\sigma(S) + |E(S, V \setminus S)|}{\mu(S)}.$$

By definition, we observe that $\widetilde{h}_1^\sigma \leq h_1^\sigma$. Theorem 5.4 implies the following estimate immediately.

COROLLARY 5.6. Let (G, μ, σ) satisfy $CD^\sigma(-K, \infty)$, $K \geq 0$. Then we have

$$(5.25) \quad \lambda_1^\sigma \leq 8 \max\{(e/(e-1))^2 D_G^{nor} (\widetilde{h}_1^\sigma)^2, \sqrt{D_G^{nor} K} \widetilde{h}_1^\sigma\}.$$

Note that, for the constant \widetilde{h}_1^σ , the following Cheeger type inequality is proved in [7, Theorem 4.1] (see also [32, Theorem 4.6 and Remark 4.9]).

THEOREM 5.7 (see [7]). Let (G, μ, σ) be given. Then we have

$$(5.26) \quad \frac{2}{5D_G^{nor}} \widetilde{h}_1^{\sigma^2} \leq \lambda_1^\sigma \leq 2\widetilde{h}_1^\sigma.$$

Example 5.8 (signed triangle). We revisit the example of a signed triangle discussed in section 3.5 (see Figure 2). In this case we have $H = U(1)$ and, therefore, $h_1^\sigma = \widetilde{h}_1^\sigma$. Using Theorem 4.10, we can check

$$h_1^\sigma = \frac{|s-1|}{6} = \frac{\sqrt{2(1-\operatorname{Re}(s))}}{6}.$$

The Buser type inequality Theorem 5.1 tells us

$$(5.27) \quad \lambda_1^\sigma \leq 32 \log 2 (h_1^\sigma)^2,$$

while the Cheeger type inequality Theorem 5.7 gives

$$(5.28) \quad \frac{2}{5} (h_1^\sigma)^2 \leq \lambda_1^\sigma \leq 2h_1^\sigma.$$

The comparison of the estimates (5.27) and (5.28) is shown in Figure 4, where we treat the quantities λ_1^σ and h_1^σ as functions of the variable $\operatorname{Re}(s)$.




FIG. 4. Comparison of Cheeger and Buser estimates for a signed triangle.

6. Lichnerowicz estimate and applications. We have the following Lichnerowicz type eigenvalue estimate (cf. Theorem 1.9 in the introduction).

THEOREM 6.1 (Lichnerowicz inequality). *Assume that (G, μ, σ) satisfies $CD^\sigma(K, n)$ for $K \in \mathbb{R}$ and $n \in \mathbb{R}_+$. Then we have for any nonzero eigenvalue λ^σ of Δ^σ ,*

$$(6.1) \quad \frac{n-1}{n} \lambda^\sigma \geq K,$$

where we use the convention $(n-1)/n = 1$ in the case $n = \infty$.

Proof. Let $\psi : V \rightarrow \mathbb{K}^d$ be the corresponding eigenfunction of λ^σ with unit $\ell^2(V, \mathbb{K}^d; \mu)$ norm. Integrating the inequality $CD^\sigma(K, n)$ over the measure μ , we obtain

$$(6.2) \quad \sum_{x \in V} \mu(x) \Gamma_2^\sigma(\psi)(x) \geq \frac{1}{n} \sum_{x \in V} \mu(x) |\Delta^\sigma \psi(x)|^2 + K \sum_{x \in V} \mu(x) \Gamma^\sigma(\psi)(x).$$

By (3.2), we have

$$\sum_{x \in V} \mu(x) \Gamma_2^\sigma(\psi)(x) = - \sum_{x \in V} \mu(x) \operatorname{Re}(\Gamma^\sigma(\psi, \Delta^\sigma \psi)(x)) = \lambda^\sigma \sum_{x \in V} \mu(x) \Gamma^\sigma(\psi)(x).$$

Recalling the summation by part formula (3.3), we have

$$\sum_{x \in V} \mu(x) \Gamma^\sigma(\psi)(x) = - \langle \psi, \Delta^\sigma \psi \rangle_\mu = \lambda^\sigma.$$

Therefore, (6.2) tells us that

$$(6.3) \quad (\lambda^\sigma)^2 \geq \frac{1}{n} (\lambda^\sigma)^2 + \lambda^\sigma K.$$

This implies (6.1) in the case $\lambda^\sigma \neq 0$. \square

Consequently, we have the following estimates about the lower curvature bound of a graph.

COROLLARY 6.2. Let (G, μ, σ) satisfy $CD^\sigma(K, n)$, for some $K \in \mathbb{R}$ and $n \in \mathbb{R}_+$. Then we have the following facts:

(i) If $n = 1$, we have

$$K \leq 0.$$

(ii) If $0 < n < 1$, we have $K < 0$. If, furthermore, σ is not balanced, we have

$$K \leq -\frac{2(1-n)}{5nD_G^{non}}(\widetilde{h}_1^\sigma)^2.$$

(iii) If $1 < n \leq \infty$ and σ is not balanced, we have

$$K \leq \frac{2(n-1)}{n}\widetilde{h}_1^\sigma \leq \frac{2(n-1)}{n}h_1^\sigma.$$

Proof. The estimate (i) follows directly from Theorem 6.1. Note that λ_1^σ is positive when σ is not balanced. Hence we can combine Theorems 6.1 and 5.7 to conclude estimates (ii) and (iii). \square

If the graph has a nonnegative lower curvature bound, we can improve the estimate Corollary 6.2(iii) by applying Corollary 5.6.

COROLLARY 6.3. Let (G, μ, σ) satisfy $CD^\sigma(K, n)$ for some $K \geq 0$ and $1 < n \leq \infty$. If the signature σ is not balanced, then we have

$$(6.4) \quad K \leq \frac{n-1}{n} \min \left\{ 2\widetilde{h}_1^\sigma, \frac{8e^2}{(e-1)^2} D_G^{nor}(\widetilde{h}_1^\sigma)^2 \right\}.$$

Proof. Recall that $CD^\sigma(K, n)$ implies $CD^\sigma(K, \infty)$. Hence, Corollary 5.6 is applicable here. \square

Remark 6.4 (jump of the curvature around a balanced signature). Suppose that a graph (G, μ) with a balanced signature has positive n -dimensional Ricci curvature, i.e., $K_n(\sigma_{\text{triv}}) > 0$. (Recall that every balanced signature is switching equivalent to σ_{triv} . Note that by Corollary 6.2, $K_n(\sigma_{\text{triv}}) > 0$ is possible only when $1 < n \leq \infty$.) Then by Corollary 6.2, we observe that the curvature $K_n(\sigma)$ of (G, μ, σ) , as a function of the signature σ , has the following “jump” phenomenon: For unbalanced signatures σ , when they are close to the balanced signature σ_{triv} ,

$$(6.5) \quad \limsup_{\iota^\sigma(V) \rightarrow 0} K_n(\sigma) \leq 0, \text{ but } K_n(\sigma_{\text{triv}}) > 0.$$

In the above expression, we use $\iota^\sigma(V)$ as a measure for the difference between σ and σ_{triv} .

The jump of the curvature is closely related to the jump phenomenon of the first non-zero eigenvalue of Δ^σ . When the signature σ of a connected graph becomes balanced, the first nonzero eigenvalue jumps from λ_1^σ to λ_2^σ .

Example 6.5 (signed triangle). We consider the example of a signed triangle again. Recall that we have observed the jump phenomenon of the curvature of a signed triangle in Remark 3.15 (see Figure 3). In Figure 1 of the introduction, the jumps of the ∞ -dimensional Ricci curvature and the first nonzero eigenvalue of a signed triangle are illustrated in the same diagram.

We conclude this section by an interesting application of the jump phenomenon of the curvature.

THEOREM 6.6. *Suppose that a graph G has at least one cycle, but no cycles of length 3 or 4. Then, for any signature σ , any edge weights, and any vertex measure μ , we have*

$$K_n(G, \mu, \sigma) \leq 0 \quad \text{for any } n \in \mathbb{R}_+.$$

Proof. Since G contains at least one cycle, there exist unbalanced signatures on G . On the other hand, we have

$$(6.6) \quad K_n(G, \mu, \sigma) = K_n(G, \mu, \sigma_{\text{triv}})$$

by Proposition 3.6, as G has no cycles of length 3 or 4. Therefore, if $K_n(G, \mu, \sigma_{\text{triv}}) > 0$, the equality (6.6) leads to a contradiction to the jump of the curvature observed in Remark 6.4. Hence we must have $K_n(G, \mu, \sigma) \leq 0$. \square

Note that the conditions on the graph in Theorem 6.6 are purely combinatorial, whereas the curvature estimate holds for any edge weights and vertex measures.

Combining Theorem 6.6 and Corollary 3.8, we obtain an indirect verification of (3.37). Actually, we obtain the following more general result.

COROLLARY 6.7. *Let $N \geq 5$ and $(\mathcal{C}_N, \mu, \sigma)$ be an unweighted cycle with constant vertex measure $\mu = \nu_0 \cdot \mathbf{1}_V$. Then we have*

$$K_n(\mathcal{C}_N, \mu, \sigma) = 0 \quad \text{for any } n \geq 2.$$

7. Eigenvalue ratios of graphs with $O(1)$ signatures. In this section, we restrict our considerations to the setting of a graph (G, μ) with a signature

$$\sigma : E^{\text{or}} \rightarrow O(1) = \{\pm 1\}.$$

We show that Theorem 5.1 can be applied to derive an upper bound for the ratio of the k th eigenvalue λ_k^σ to the first eigenvalue λ_1^σ when (G, μ, σ) satisfies $CD^\sigma(0, \infty)$.

Note that the connection Laplacian reduces to an operator on $\ell^2(V, \mathbb{R}; \mu)$. That is, for any real function $f : V \rightarrow \mathbb{R}$ and any vertex $x \in V$, we have

$$(7.1) \quad \Delta^\sigma f(x) := \frac{1}{\mu(x)} \sum_{y, y \sim x} w_{xy} (\sigma_{xy} f(y) - f(x)) \in \mathbb{R}.$$

The eigenvalues of Δ^σ can be listed as

$$0 \leq \lambda_1^\sigma \leq \dots \leq \lambda_k^\sigma \leq \dots \leq \lambda_N^\sigma \leq 2D_G^{\text{non}}.$$

In [4, Theorem 3], Atay and Liu prove the following estimate.

THEOREM 7.1 (see [4]). *For any graph (G, μ, σ) with $\sigma : E^{\text{or}} \rightarrow O(1)$ and any natural number $1 \leq k \leq N$, we have*

$$(7.2) \quad h_1^\sigma \leq 16 \sqrt{2D_G^{\text{non}}} k \frac{\lambda_1^\sigma}{\sqrt{\lambda_k^\sigma}}.$$

This result is an extension of the so-called improved Cheeger inequality due to Kwok et al. [31] for the graph Laplacian Δ . We also mention that in the current case of $O(1)$ signatures, the multiway Cheeger constants, given in Definition 4.6, have more explicit combinatorial expressions. We refer to [4] for more details.

As an application of Theorem 5.1, we prove the following eigenvalue ratio estimates.

THEOREM 7.2. For any graph (G, μ, σ) with $\sigma : E^{or} \rightarrow O(1)$ satisfying $CD^\sigma(0, \infty)$ and any natural number $1 \leq k \leq N$, there exists an absolute constant C such that

$$(7.3) \quad \lambda_k^\sigma \leq CD_G^{nor} D_G^{non} k^2 \lambda_1^\sigma.$$

Proof. Since the inequality $CD^\sigma(0, \infty)$ is satisfied, we have by Theorem 5.1,

$$\sqrt{\lambda_1^\sigma} \leq 4\sqrt{(\log 2) D_G^{nor} h_1^\sigma}.$$

Combining this with Theorem 7.1, we obtain

$$(7.4) \quad \sqrt{\lambda_1^\sigma} \leq 64\sqrt{2 \log 2} \sqrt{D_G^{nor} D_G^{non}} k \frac{\lambda_1^\sigma}{\sqrt{\lambda_k^\sigma}}.$$

This implies (7.3) immediately. \square

A direct corollary is the following Buser type inequality.

COROLLARY 7.3. Let (G, μ, σ) with $\sigma : E^{or} \rightarrow O(1)$ satisfy $CD^\sigma(0, \infty)$. Then for all $1 \leq k \leq N$, there exists an absolute constant C such that

$$(7.5) \quad \sqrt{\lambda_k^\sigma} \leq C \sqrt{D_G^{non} D_G^{nor}} k h_1^\sigma.$$

Proof. Combining (7.4) with Theorem 7.2 leads to this result immediately. \square

Remark 7.4. Comparing this result with Theorem 5.1, the advantage of the estimate (7.5) lies in the fact that $h_1^\sigma \leq h_k^\sigma$ and that the order of k in (7.5) is lower. However, in the estimate (7.5), the orders of the degrees D_G^{nor} and D_G^{non} are higher than in Theorem 5.1.

Finally, we observe that Theorem 7.1 and hence the estimate in Corollary 7.3 cannot be true for general signatures $\sigma : E^{or} \rightarrow H$, even in the one-dimensional case $H = U(1)$. To explain the reason, let us revisit the example of a signed triangle.

Example 7.5 (signed triangle). The example of a signed triangle, discussed in section 3.5, carries a $U(1)$ signature σ is assigned (see Figure 2). If $\operatorname{Re}(s)$ tends to 1, i.e., if the signature on the triangle tends to be balanced, we observe that λ_2^σ has a positive lower bound (see Figure 1), while both λ_1^σ and h_1^σ tend to zero, but at a different rate (see Figure 4). In fact, by Theorem 5.1, we have

$$(7.6) \quad \lambda_1^\sigma \leq 32 \log 2 (h_1^\sigma)^2.$$

Assume that Theorem 7.1 holds in this case for $k = 2$. Combining this with (7.6), we obtain $1 \leq C h_1^\sigma$ for some absolute constant $C > 0$. This is a contradiction. Hence, Theorem 7.1 cannot hold for more general signatures.

Appendix A. Curvature and Cheeger constants on Cartesian products.

In this section, we discuss the CD^σ inequality and the Cheeger constants on the Cartesian product of two graphs. For two graphs, $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$, their Cartesian product $G_1 \times G_2 = (V_1 \times V_2, E_{12})$ is defined as follows. Any two vertices $(x_1, y_1), (x_2, y_2) \in V_1 \times V_2$ are connected by an edge in E_{12} if and only if

$$\text{either } \{x_1, x_2\} \in E_1, y_1 = y_2 \text{ or } x_1 = x_2, \{y_1, y_2\} \in E_2.$$

A.1. Curvature on Cartesian products. We first discuss the simpler case of graphs with constant vertex measures.

A.1.1. Graphs with constant vertex measures. Given two graphs $G_1 = (V_1, E_1, w_1)$ and $G_2 = (V_2, E_2, w_2)$, we assign the following edge weights to the Cartesian product $G_1 \times G_2 = (V_1 \times V_2, E_{12})$:

$$(A.1) \quad \begin{aligned} w_{12, (x_1, y)(x_2, y)} &:= w_{1, x_1 x_2} \quad \text{for any } \{x_1, x_2\} \in E_1, y \in V_2; \\ w_{12, (x, y_1)(x, y_2)} &:= w_{2, y_1 y_2} \quad \text{for any } \{y_1, y_2\} \in E_2, x \in V_1. \end{aligned}$$

Let $\sigma_i : E_i^{or} \rightarrow H_i, i = 1, 2$, be the signatures of $G_i, i = 1, 2$, respectively. We need to find a proper construction of the signature on the Cartesian product graph $G_1 \times G_2$.

We first consider the case that

$$H_1 = H_2 := H = O(d) \quad \text{or} \quad U(d) \quad \text{for some } d \in \mathbb{Z}_{>0}.$$

In this case, we define the signature $\sigma_{12} : E_{12}^{or} \rightarrow H$ as follows:

$$(A.2) \quad \begin{aligned} \sigma_{12, (x_1, y)(x_2, y)} &:= \sigma_{1, x_1 x_2} \quad \text{for any } (x_1, x_2) \in E_1^{or}, y \in V_2; \\ \sigma_{12, (x, y_1)(x, y_2)} &:= \sigma_{2, y_1 y_2} \quad \text{for any } (y_1, y_2) \in E_2^{or}, x \in V_1. \end{aligned}$$

Let Σ_i be the *signature group* of the graph G_i with σ_i (recall the definition of the signature group at the end of section 2.2). We say the two subgroups Σ_1 and Σ_2 of H commute if for any $s_1 \in \Sigma_1, s_2 \in \Sigma_2$, we have $s_1 s_2 = s_2 s_1$.

THEOREM A.1. *Let $(G_1, 1_{V_1}, \sigma_1)$ and $(G_2, 1_{V_2}, \sigma_2)$ be two graphs with $\sigma_i : E_i^{or} \rightarrow H, i = 1, 2$. Assume that they satisfy*

$$CD^{\sigma_1}(K_1, n_1) \quad \text{and} \quad CD^{\sigma_2}(K_2, n_2),$$

respectively. If their signature groups Σ_1 and Σ_2 commute, then their Cartesian product graph $(G_1 \times G_2, 1_{V_1 \times V_2}, \sigma_{12})$, with the edge weight w_{12} given in (A.1), satisfies

$$CD^{\sigma_{12}}(K_1 \wedge K_2, n_1 + n_2),$$

where $K_1 \wedge K_2 := \min\{K_1, K_2\}$.

Note that the commutativity restriction of Σ_1 and Σ_2 is a very natural condition. The intuition behind the above result is that the “mixed structure” in the Cartesian product is “flat.” To be precise, we want for two balanced signatures σ_1, σ_2 on G_1, G_2 that σ_{12} on $G_1 \times G_2$ is also balanced. In Figure 5, we show a typical new cycle created in the process of taking the Cartesian product, where $\{x_i, x\} \in E_1$ and $\{y_k, y\} \in E_2$. Since Σ_1 and Σ_2 commute, the signature of this cycle, given by $\sigma_{1, x x_i} \sigma_{2, y y_k} \sigma_{1, x x_i}^{-1} \sigma_{2, y y_k}^{-1}$, is trivial.

Proof of Theorem A.1. Let $f : V_1 \times V_2 \rightarrow \mathbb{K}^d$ be any \mathbb{K}^d valued function on the product graph. For fixed $y \in V_2$, we define $f_y(\cdot) := f(\cdot, y)$ to be a \mathbb{K}^d valued function on G_1 . Similarly, we define the function $f^x(\cdot) := f(x, \cdot) : V_2 \rightarrow \mathbb{K}^d$.

By the construction of σ_{12} , it is straightforward to check that

$$(A.3) \quad \Delta^{\sigma_{12}} f(x, y) = \Delta^{\sigma_1} f_y(x) + \Delta^{\sigma_2} f^x(y),$$

$$(A.4) \quad \Gamma^{\sigma_{12}}(f)(x, y) = \Gamma^{\sigma_1}(f_y)(x) + \Gamma^{\sigma_2}(f^x)(y).$$

For the operator $\Gamma_2^{\sigma_{12}}$, we claim that

$$(A.5) \quad \Gamma_2^{\sigma_{12}}(f)(x, y) \geq \Gamma_2^{\sigma_1}(f_y)(x) + \Gamma_2^{\sigma_2}(f^x)(y).$$




FIG. 5. A typical cycle in the Cartesian product graph.

Once inequality (A.5) is verified, we apply the CD^σ inequalities on G_1 and G_2 to estimate

$$\begin{aligned}
 \Gamma_2^{\sigma_{12}}(f)(x, y) &\geq \frac{1}{n_1} |\Delta^{\sigma_1} f_y(x)|^2 + K_1 \Gamma^{\sigma_1}(f_y)(x) + \frac{1}{n_2} |\Delta^{\sigma_2} f^x(y)|^2 + K_2 \Gamma^{\sigma_2}(f^x)(y) \\
 &\geq \frac{1}{n_1 + n_2} (|\Delta^{\sigma_1} f_y(x)| + |\Delta^{\sigma_2} f^x(y)|)^2 + (K_1 \wedge K_2) (\Gamma^{\sigma_1}(f_y)(x) + \Gamma^{\sigma_2}(f^x)(y)) \\
 &\geq \frac{1}{n_1 + n_2} |\Delta^{\sigma_{12}} f(x, y)|^2 + (K_1 \wedge K_2) \Gamma^{\sigma_{12}}(f)(x, y).
 \end{aligned}
 \tag{A.6}$$

In the third inequality above, we used the triangle inequality and the equalities (A.3) and (A.4).

Hence, it only remains to prove the claim (A.5). Recall that

$$(A.7) \quad 2\Gamma_2^{\sigma_{12}}(f)(x, y) = \Delta \Gamma^{\sigma_{12}}(f)(x, y) - \Gamma^\sigma(f, \Delta^{\sigma_{12}} f)(x, y) - \Gamma^\sigma(\Delta^{\sigma_{12}} f, f)(x, y).$$

For simplicity, we denote the neighbors of x in V_1 by x_i and the neighbors of y in V_2 by y_k . We will then write, for short,

$$w_{1,i} := w_{1,xx_i}, w_{2,k} := w_{2,yy_k}, \text{ and } \sigma_{1,i} := \sigma_{1,xx_i}, \sigma_{2,k} := \sigma_{2,yy_k},$$

and \sum_{x_i} (\sum_{y_k} , resp.) the summation over all neighbors of $x \in V_1$ ($y \in V_2$, resp.). We first calculate

$$\begin{aligned}
 \Delta \Gamma^{\sigma_{12}}(f)(x, y) &= \underbrace{\sum_{x_i} w_{1,i} (\Gamma^{\sigma_{12}}(f)(x_i, y) - \Gamma^{\sigma_{12}}(f)(x, y))}_{=: L_1} + \underbrace{\sum_{y_k} w_{2,k} (\Gamma^{\sigma_{12}}(f)(x, y_k) - \Gamma^{\sigma_{12}}(f)(x, y))}_{=: L_2}.
 \end{aligned}$$

Applying (A.4), we obtain

$$\begin{aligned}
 L_1 &= \sum_{x_i} w_{1,i} (\Gamma^{\sigma_1}(f_y)(x_i) + \Gamma^{\sigma_2}(f^{x_i})(y) - \Gamma^{\sigma_1}(f_y)(x) - \Gamma^{\sigma_2}(f^x)(y)) \\
 &= \Delta \Gamma^{\sigma_1}(f_y)(x) + \frac{1}{2} \sum_{x_i, y_k} w_{1,i} w_{2,k} (|\sigma_{2,k} f(x_i, y_k) - f(x_i, y)|^2 \\
 &\quad - |\sigma_{2,k} f(x, y_k) - f(x, y)|^2).
 \end{aligned}
 \tag{A.8}$$

Similarly, we have

$$(A.9) \quad L_2 = \Delta \Gamma^{\sigma_2}(f^x)(y) + \frac{1}{2} \sum_{x_i, y_k} w_{1,i} w_{2,k} (|\sigma_{1,i} f(x_i, y_k) - f(x, y_k)|^2 - |\sigma_{i,i} f(x_i, y) - f(x, y)|^2).$$

Now we calculate the remaining terms in (A.7).

$$(A.10) \quad \begin{aligned} & \Gamma^{\sigma_{12}}(f, \Delta^{\sigma_{12}} f)(x, y) \\ &= \frac{1}{2} \sum_{x_i} w_{1,i} (\sigma_{1,i} f(x_i, y) - f(x, y))^T (\overline{\sigma_{1,i} \Delta^{\sigma_{12}} f(x_i, y) - \Delta^{\sigma_{12}} f(x, y)}) \\ & \quad + \frac{1}{2} \sum_{y_k} w_{2,k} (\sigma_{2,k} f(x, y_k) - f(x, y))^T (\overline{\sigma_{2,k} \Delta^{\sigma_{12}} f(x, y_k) - \Delta^{\sigma_{12}} f(x, y)}) \\ &=: M_1 + M_2. \end{aligned}$$

Applying the equality (A.3), we obtain

$$(A.11) \quad \begin{aligned} M_1 &= \Gamma^{\sigma_1}(f_y, \Delta^{\sigma_1} f_y)(x) \\ &+ \frac{1}{2} \sum_{x_i} w_{1,i} (\sigma_{1,i} f(x_i, y) - f(x, y))^T (\overline{\sigma_{1,i} \Delta^{\sigma_2} f^{x_i}(y) - \Delta^{\sigma_2} f^x(y)}). \end{aligned}$$

Hence, we get

$$(A.12) \quad \begin{aligned} & M_1 - \Gamma^{\sigma_1}(f_y, \Delta^{\sigma_1} f_y)(x) \\ &= \frac{1}{2} \sum_{x_i, y_k} w_{1,i} w_{2,k} [(\sigma_{1,i} f(x_i, y) - f(x, y))^T (\overline{\sigma_{1,i} \sigma_{2,k} f(x_i, y_k) - \sigma_{2,k} f(x, y_k)}) \\ & \quad - |\sigma_{1,i} f(x_i, y) - f(x, y)|^2]. \end{aligned}$$

Similarly, we have

$$(A.13) \quad \begin{aligned} & M_2 - \Gamma^{\sigma_2}(f^x, \Delta^{\sigma_2} f^x)(y) \\ &= \frac{1}{2} \sum_{x_i, y_k} w_{1,i} w_{2,k} [(\sigma_{2,k} f(x, y_k) - f(x, y))^T (\overline{\sigma_{2,k} \sigma_{1,i} f(x_i, y_k) - \sigma_{1,i} f(x_i, y)}) \\ & \quad - |\sigma_{2,k} f(x, y_k) - f(x, y)|^2]. \end{aligned}$$

Combining (A.8) and (A.13), we arrive at

$$(A.14) \quad \begin{aligned} & (L_1 - \Delta \Gamma^{\sigma_1}(f_y)(x)) - (M_2 - \Gamma^{\sigma_2}(f^x, \Delta^{\sigma_2} f^x)(y)) - (\overline{M_2 - \Gamma^{\sigma_2}(f^x, \Delta^{\sigma_2} f^x)(y)}) \\ &= \frac{1}{2} \sum_{x_i, y_k} w_{1,i} w_{2,k} [|\sigma_{2,k} f(x_i, y_k) - f(x_i, y)|^2 + |\sigma_{2,k} f(x, y_k) - f(x, y)|^2 \\ & \quad + (\sigma_{2,k} f(x, y_k) - f(x, y))^T (\overline{\sigma_{2,k} \sigma_{1,i} f(x_i, y_k) - \sigma_{1,i} f(x_i, y)}) \\ & \quad + (\overline{\sigma_{2,k} f(x, y_k) - f(x, y)})^T (\sigma_{2,k} \sigma_{1,i} f(x_i, y_k) - \sigma_{1,i} f(x_i, y))]. \end{aligned}$$

Since Σ_1 and Σ_2 commute, we have

$$\sigma_{2,k} \sigma_{1,i} = \sigma_{1,i} \sigma_{2,k}.$$

Therefore, we obtain

$$\begin{aligned} & (L_1 - \Delta \Gamma^{\sigma_1}(f_y)(x)) - (M_2 - \Gamma^{\sigma_2}(f^x, \Delta^{\sigma_2} f^x)(y)) - \overline{(M_2 - \Gamma^{\sigma_2}(f^x, \Delta^{\sigma_2} f^x)(y))} \\ &= \frac{1}{2} \sum_{x_i, y_k} w_{1,i} w_{2,k} |\sigma_{1,i} \sigma_{2,k} f(x_i, y_k) - \sigma_{1,i} f(x_i, y) - \sigma_{2,k} f(x, y_k) + f(x, y)|^2. \end{aligned} \quad (\text{A.15})$$

Similarly, by combining (A.9) and (A.12), we obtain

$$\begin{aligned} & (L_2 - \Delta \Gamma^{\sigma_2}(f^x)(y)) - (M_1 - \Gamma^{\sigma_1}(f_y, \Delta^{\sigma_1} f_y)(x)) - \overline{(M_1 - \Gamma^{\sigma_1}(f_y, \Delta^{\sigma_1} f_y)(x))} \\ &= \frac{1}{2} \sum_{x_i, y_k} w_{1,i} w_{2,k} |\sigma_{2,k} \sigma_{1,i} f(x_i, y_k) - \sigma_{2,k} f(x, y_k) - \sigma_{1,i} f(x_i, y) + f(x, y)|^2. \end{aligned} \quad (\text{A.16})$$

Adding (A.15) and (A.16), and using (3.1), we get

$$\begin{aligned} & 2\Gamma_2^{\sigma_{12}}(f)(x, y) - 2\Gamma_2^{\sigma_1}(f_y)(x) - 2\Gamma_2^{\sigma_2}(f^x)(y) \\ &= \sum_{x_i, y_k} w_{1,i} w_{2,k} |\sigma_{2,k} \sigma_{1,i} f(x_i, y_k) - \sigma_{2,k} f(x, y_k) - \sigma_{1,i} f(x_i, y) + f(x, y)|^2 \geq 0. \end{aligned} \quad (\text{A.17})$$

This proves (A.5). \square

Remark A.2 (tightness of Theorem A.1). The estimate in Theorem A.1 is tight at least in the case of taking the Cartesian product of $(G, \mathbf{1}_V, \sigma)$ with itself, assuming that its signature group Σ is abelian. That is, for any given $n \in \mathbb{R}_+$, the precise lower curvature bounds satisfy

$$K_{2n}(G \times G, \mathbf{1}_{V \times V}, \sigma_{12}) = K_n(G, \mathbf{1}_V, \sigma).$$

Note that the tightness of Theorem A.1 lies in the tightness of (A.6) and (A.17). By assumption, there exists a function $f : V \rightarrow \mathbb{K}^d$ and a vertex $x \in V$ such that

$$\Gamma_2^\sigma(f)(x) = \frac{1}{n} |\Delta^\sigma f(x)|^2 + K \Gamma^\sigma(f)(x) \quad \text{and} \quad \Gamma^\sigma f(x) \neq 0.$$

Then we can choose a function $F : V \times V \rightarrow \mathbb{K}^d$ satisfying, locally, around the vertex $(x, x) \in V \times V$,

- (i) $F(x, x) := f(x)$;
 - (ii) $F(x_i, x) := f(x_i)$ for all $x_i \sim x$;
 - (iii) $F(x, x_k) := f(x_k)$ for all $x_k \sim x$;
 - (iv) $F(x_i, x_k) := \sigma_{xx_i}^{-1} \sigma_{xx_k}^{-1} (\sigma_{xx_k} f(x_k) + \sigma_{xx_i}^{-1} f(x_i) - f(x))$ for all $x_i \sim x$ and $x_k \sim x$.
- Note that (i)–(iii) implies $\Delta^\sigma F_x(x) = \Delta^\sigma F^x(x)$ and, hence, (A.6) holds with equalities. Property (iv) ensures that (A.17) holds also with equality. This shows the tightness of the result.

Next, we discuss the situation when the two groups H_1 and H_2 are different. We assume that

$$H_1 = O(d_1), H_2 = O(d_2), \quad \text{or} \quad H_1 = U(d_1), H_2 = U(d_2), \quad \text{for some } d_1, d_2 \in \mathbb{Z}_{>0},$$

where d_1, d_2 can be different integers. In such a general situation, we construct the signature $\widehat{\sigma}_{12} : E_{12}^{or} \rightarrow H_1 \otimes H_2$ on the Cartesian product graph in the following way:

$$(A.18) \quad \begin{aligned} \widehat{\sigma}_{12, (x_1, y)(x_2, y)} &:= \sigma_{1, x_1 x_2} \otimes I_{d_2} \quad \text{for any } (x_1, x_2) \in E_1^{or}, y \in V_2; \\ \widehat{\sigma}_{12, (x, y_1)(x, y_2)} &:= I_{d_1} \otimes \sigma_{2, y_1 y_2} \quad \text{for any } (y_1, y_2) \in E_2^{or}, x \in V_1, \end{aligned}$$

where I_{d_i} is the identity matrix of size $d_i \times d_i$, $i = 1, 2$.

THEOREM A.3. *Let $(G_1, \mathbf{1}_{V_1}, \sigma_1)$ and $(G_2, \mathbf{1}_{V_2}, \sigma_2)$ be two graphs with $\sigma_i : E_i^{or} \rightarrow H_i$, $i = 1, 2$. Assume that they satisfy*

$$CD^{\sigma_1}(K_1, n_1) \quad \text{and} \quad CD^{\sigma_2}(K_2, n_2),$$

respectively. Then their Cartesian product graph $(G_1 \times G_2, \mathbf{1}_{V_1 \times V_2}, \widehat{\sigma}_{12})$ with the edge weight w_{12} given in (A.1) satisfies

$$CD^{\widehat{\sigma}_{12}}(K_1 \wedge K_2, n_1 + n_2).$$

This is an immediate consequence of Theorem A.1 and Corollary 3.4.

Proof of Theorem A.3. By Corollary 3.4, we know that $(G_1, \mathbf{1}_{V_1}, \sigma_1 \otimes I_{d_2})$ and $(G_2, \mathbf{1}_{V_2}, I_{d_1} \otimes \sigma_2)$ satisfy $CD^{\sigma_1 \otimes I_{d_2}}(K_1, n_1)$ and $CD^{I_{d_1} \otimes \sigma_2}(K_2, n_2)$, respectively. Note that for any $(x, x_i) \in E_1^{or}$ and $(y, y_k) \in E_2^{or}$, we have

$$(A.19) \quad (\sigma_{1, x x_i} \otimes I_{d_2})(I_{d_1} \otimes \sigma_{2, y y_k}) = (I_{d_1} \otimes \sigma_{2, y y_k})(\sigma_{1, x x_i} \otimes I_{d_2}).$$

That is, the corresponding signature groups of $(G_1, \mathbf{1}_{V_1}, \sigma_1 \otimes I_{d_2})$ and $(G_2, \mathbf{1}_{V_2}, I_{d_1} \otimes \sigma_2)$ commute. Hence, we can apply Theorem A.1 and finish the proof. \square

Remark A.4 (vertex measure). In Theorems A.1 and A.3, we use the particular vertex measure $\mu(x) = 1$ for all vertices x . In fact, we have more flexibility about those measures. Assume that the vertex measures of $G_1, G_2, G_1 \times G_2$ take constant values $\nu_1, \nu_2, \nu_{12} \in \mathbb{R}$, respectively. Then under the assumptions of Theorem A.1 (replacing $\mathbf{1}_{V_i}$ by $\nu_i \cdot \mathbf{1}_{V_i}$), we have that the graph $(G_1 \times G_2, \nu_{12} \cdot \mathbf{1}_{V_1 \times V_2}, \sigma_{12})$ satisfies

$$CD^{\sigma_{12}} \left(\frac{1}{\nu_{12}} (\nu_1 K_1 \wedge \nu_2 K_2), n_1 + n_2 \right).$$

The result in Theorem A.3 can be generalized similarly.

A.1.2. Graphs with nonconstant vertex measures. For two graphs (G_1, μ_1) and (G_2, μ_2) whose vertex measures are not necessarily constant, we modify the definition of the edge weights of their Cartesian product. In [14], Chung and Tetali introduced the edge weight for the Cartesian product $G_1 \times G_2 = (V_1 \times V_2, E_{12})$

$$(A.20) \quad \begin{aligned} w_{12, (x_1, y)(x_2, y)}^{\square} &:= w_{1, x_1 x_2} \mu_2(y) \quad \text{for any } \{x_1, x_2\} \in E_1, y \in V_2; \\ w_{12, (x, y_1)(x, y_2)}^{\square} &:= w_{2, y_1 y_2} \mu_1(x) \quad \text{for any } \{y_1, y_2\} \in E_2, x \in V_1, \end{aligned}$$

and the specific vertex measure

$$(A.21) \quad 2\mu_1 \mu_2 : V_1 \times V_1 \ni (x, y) \mapsto 2\mu_1(x) \mu_2(y) \in \mathbb{R}.$$

Observe that, in the case $\mu_i = \mathbf{d}_{V_i}$, $i = 1, 2$, we have

$$2\mu_1(x) \mu_2(y) = \sum_{x_i} w_{12, (x_i, y)(x, y)}^{\square} + \sum_{y_k} w_{12, (x, y_k)(x, y)}^{\square}.$$

The definitions (A.20) and (A.21) lead to a Laplacian associated to a natural random walk on the product graph $G_1 \times G_2$.

THEOREM A.5. Let (G_1, μ_1, σ_1) and (G_2, μ_2, σ_2) be two graphs with $\sigma_i : E_i^{or} \rightarrow H$, $i = 1, 2$. Assume that they satisfy

$$CD^{\sigma_1}(K_1, n_1) \quad \text{and} \quad CD^{\sigma_2}(K_2, n_2),$$

respectively. If their signature groups Σ_1 and Σ_2 commute, then their Cartesian product graph $(G_1 \times G_2, 2\mu_1\mu_2, \sigma_{12})$, with the edge weight w_{12}^\square given in (A.20), satisfies

$$CD^{\sigma_{12}}\left(\frac{1}{2}(K_1 \wedge K_2), n_1 + n_2\right),$$

where $K_1 \wedge K_2 := \min\{K_1, K_2\}$.

Proof. The proof is analogous to the proof of Theorem A.1. We only mention here that, this time, we have

$$\begin{aligned}\Delta^{\sigma_{12}}f(x, y) &= \frac{1}{2}\Delta^{\sigma_1}f_y(x) + \frac{1}{2}\Delta^{\sigma_2}f^x(y), \\ \Gamma^{\sigma_{12}}(f)(x, y) &= \frac{1}{2}\Gamma^{\sigma_1}(f_y)(x) + \frac{1}{2}\Gamma^{\sigma_2}(f^x)(y),\end{aligned}$$

and

$$\Gamma_2^{\sigma_{12}}(f)(x, y) \geq \frac{1}{4}\Gamma_2^{\sigma_1}(f_y)(x) + \frac{1}{4}\Gamma_2^{\sigma_2}(f^x)(y).$$

The last inequality above is derived from

$$\begin{aligned}2\mu(x)\mu(y) &(4\Gamma_2^{\sigma_{12}}(f)(x, y) - \Gamma_2^{\sigma_1}(f_y)(x) - \Gamma_2^{\sigma_2}(f^x)(y)) \\ &= \sum_{x_i, y_k} w_{1,i}w_{2,k}|\sigma_{2,k}\sigma_{1,i}f(x_i, y_k) - \sigma_{2,k}f(x, y_k) - \sigma_{1,i}f(x_i, y) + f(x, y)|^2 \geq 0. \quad \square\end{aligned}$$

Remark A.6. Let us assign a general vertex measure μ_{12} to the Cartesian product graph. Then, under the assumption of Theorem A.5, a proof analogous to the proof of Theorem A.1 yields that $(G_1 \times G_2, \mu_{12}, \sigma_{12})$, with the edge weight w_{12}^\square given in (A.20), satisfies

$$CD^{\sigma_{12}}\left(\min_{(x,y) \in V_1 \times V_2} \frac{\mu_1(x)\mu_2(y)}{\mu_{12}(x,y)}(K_1 \wedge K_2), n_1 + n_2\right).$$

Note that this general result also includes Theorem A.1 as a particular case.

A result similar to Theorem A.3 follows immediately from Theorem A.5.

THEOREM A.7. Let (G_1, μ_1, σ_1) and (G_2, μ_2, σ_2) be two graphs with $\sigma_i : E_i^{or} \rightarrow H_i$, $i = 1, 2$. Assume that they satisfy

$$CD^{\sigma_1}(K_1, n_1) \quad \text{and} \quad CD^{\sigma_2}(K_2, n_2),$$

respectively. Then their Cartesian product graph $(G_1 \times G_2, \mu_{12}, \widehat{\sigma}_{12})$, with the edge weight w_{12}^\square given in (A.20), satisfies

$$CD^{\widehat{\sigma}_{12}}\left(\min_{(x,y) \in V_1 \times V_2} \frac{\mu_1(x)\mu_2(y)}{\mu_{12}(x,y)}(K_1 \wedge K_2), n_1 + n_2\right).$$

A.2. Cheeger constants on Cartesian products. In this subsection, we discuss relations between the Cheeger constants on two graphs and on their Cartesian products.

Recall (4.3), i.e., for any $(d \times d)$ -matrix A and any $B \in O(d)$ or $U(d)$, their average $(2, 1)$ -norm satisfies

$$(A.22) \quad |BA|_{2,1} = |A|_{2,1}.$$

This ensures the following relation between the Cheeger constants.

THEOREM A.8. *Let $(G_1, 1_{\mathbf{v}_1}, \sigma_1)$ and $(G_2, 1_{\mathbf{v}_2}, \sigma_2)$ be two graphs with $\sigma_i : E_i^{or} \rightarrow H_i$, $i = 1, 2$. Suppose that H_1 and H_2 are embedded in a group H_{12} such that H_1 and H_2 commute. Define σ_{12} as in (A.2). Then the kl -way Cheeger constant $h_{kl}^{\sigma_{12}}$ of their Cartesian product graph $(G_1 \times G_2, 1_{\mathbf{v}_1 \times \mathbf{v}_2}, \sigma_{12})$, with the edge weight w_{12} given in (A.1), satisfies*

$$h_{kl}^{\sigma_{12}} \leq h_k^{\sigma_1} + h_l^{\sigma_2}.$$

We first show the following lemma.

LEMMA A.9. *For any subsets $S_i \subseteq V_i$, $i = 1, 2$, we have*

$$(A.23) \quad \iota^{\sigma_{12}}(S_1 \times S_2) \leq |S_1| \iota^{\sigma_2}(S_2) + |S_2| \iota^{\sigma_1}(S_1).$$

Proof. Let $\tau_i : S_i \rightarrow H_i$ be the function that achieves the frustration index $\iota^{\sigma_i}(S_i)$. Set $\tau := \tau_1 \tau_2 : S_1 \times S_2 \rightarrow H_{12}$. Then, by definition, we calculate

$$\begin{aligned} \iota^{\sigma_{12}}(S_1 \times S_2) &\leq \sum_{y \in S_2} \sum_{\{x, x'\} \in E_{S_1}} w_{1,xx'} |\sigma_{1,xx'} \tau_1(x') \tau_2(y) - \tau_1(x) \tau_2(y)|_{2,1} \\ &\quad + \sum_{x \in S_1} \sum_{\{y, y'\} \in E_{S_2}} w_{2,yy'} |\sigma_{2,yy'} \tau_1(x) \tau_2(y') - \tau_1(x) \tau_2(y)|_{2,1} \\ &= |S_2| \sum_{\{x, x'\} \in E_{S_1}} w_{1,xx'} |\sigma_{1,xx'} \tau_1(x') - \tau_1(x)|_{2,1} \\ &\quad + |S_1| \sum_{\{y, y'\} \in E_{S_2}} w_{2,yy'} |\sigma_{2,yy'} \tau_2(y') - \tau_2(y)|_{2,1}. \end{aligned}$$

In the last equality, we used that H_1 and H_2 commute and (A.22). This implies the lemma immediately. \square

Proof of Theorem A.8. For any two subsets $S_i \subseteq V_i$, $i = 1, 2$, it is straightforward to check that

$$|E(S_1 \times S_2, V_1 \times V_2 \setminus S_1 \times S_2)| \leq |S_2| |E(S_1, V_1 \setminus S_1)| + |S_1| |E(S_2, V_2 \setminus S_2)|.$$

Combining this with Lemma A.9, and using the fact $|S_1 \times S_2| = |S_1| |S_2|$, we obtain

$$\phi^{\sigma_{12}}(S_1 \times S_2) \leq \phi^{\sigma_1}(S_1) + \phi^{\sigma_2}(S_2).$$

Then the theorem follows immediately, by definition, since every nontrivial k -subpartition $\{S_1^{(i)}\}_{i=1}^k$ of V_1 and every nontrivial l -subpartition $\{S_2^{(j)}\}_{j=1}^l$ of V_2 induce a nontrivial kl -subpartition $\{S_1^{(i)} \times S_2^{(j)}\}_{i=1, j=1}^{k,l}$ of $V_1 \times V_2$. \square

Note that Theorem A.8 can be applied in the following particular case: Given two signatures $\sigma_i : E_i^{or} \rightarrow H_i$, $i = 1, 2$, we can embed them into $H_1 \otimes H_2$ by identifying H_1 with $H_1 \otimes I_{d_2}$ and H_2 with $I_{d_1} \otimes H_2$, respectively. Then the signature σ_{12} on $G_1 \times G_2$ coincides with the signature $\widehat{\sigma}_{12}$ given in (A.18).

For graphs with nonconstant vertex measures, we can extend the above proof to obtain the following result.

THEOREM A.10. *Let (G_1, μ_1, σ_1) and (G_2, μ_2, σ_2) be two graphs with $\sigma_i : E_i^{or} \rightarrow H_i$, $i = 1, 2$. Then the kl -way Cheeger constant $h_{kl}^{\widehat{\sigma}_{12}}$ of their Cartesian product graph $(G_1 \times G_2, 2\mu_1\mu_2, \widehat{\sigma}_{12})$, with the edge weight w_{12}^\square given in (A.20), satisfies*

$$h_{kl}^{\widehat{\sigma}_{12}} \leq \frac{1}{2}(h_k^{\sigma_1} + h_l^{\sigma_2}).$$

Appendix B. Frustration index and spanning trees. In section 4.2, we showed that, in the case $H = U(1)$, there is an easier way to calculate the frustration index of a subset $S \subseteq V$. Recall that the frustration index $\iota^\sigma(S)$ is defined as the minimum of $\sum_{\{x,y\} \in S} w_{xy} |\sigma_{xy}\tau(y) - \tau(x)|_{2,1}$ over all possible switching functions τ on S . Theorem 4.10 tells us that it is enough to take the minimum of

$$(B.1) \quad \sum_{\{x,y\} \in S} w_{xy} |\sigma_{xy}\tau_T(y) - \tau_T(x)|_{2,1}$$

over all spanning trees T of S , where τ_T is an arbitrary representative of the set

$$C_T(S) = \{\tau : S \rightarrow U(1) : \tau \text{ is constant on } T \text{ w.r.t. } \sigma\}.$$

Recall that (B.1) is well defined because for any two $\tau_1, \tau_2 \in C_T(S)$, there exists $z \in U(1)$ such that $\tau_1 = \tau_2 z$, and hence

$$(B.2) \quad |\sigma_{xy}\tau_1(y) - \tau_1(x)|_{2,1} = |(\sigma_{xy}\tau_2(y) - \tau_2(x))z|_{2,1} = |\sigma_{xy}\tau_2(y) - \tau_2(x)|_{2,1}.$$

That is, the quantity $|\sigma_{xy}\tau_T(x) - \tau_T(y)|_{2,1}$ does not depend on the choice of $\tau_T \in C_T(S)$.

It is natural to ask whether Theorem 4.10 can be generalized to higher dimensional signatures, i.e., $H = U(d)$, for $d \geq 2$. We first observe that, for the signature $\sigma : E^{or} \rightarrow U(d)$, $d \geq 2$, the quantity (B.1) is not well defined since it depends on the representatives! Note that for any $(d \times d)$ -matrices A and $B \in U(d)$, we do not always have $|AB|_{2,1} = |A|_{2,1}$ (recall that $|BA|_{2,1} = |A|_{2,1}$). But this is needed in the verification of (B.2).

However, if we use the Frobenius norm $|\cdot|_F$ instead, (B.1) is still well defined. In this section, we present a counterexample to show that Theorem 4.10 does not hold for higher dimensional signatures, even if we use the Frobenius norm in the definition of the frustration index.

Recall that Lemma 4.11, which is a statement about the metric space $S^1 = U(1)$, plays a crucial role in the proof of Theorem 4.10. This lemma does not generalize to higher dimensional spheres. Already in S^2 , we have the following counterexample: For three equidistributed points P_1, P_2, P_3 on a meridian close to the north pole N , we have

$$d(P_1, P_2) + d(P_1, P_3) > d(N, P_1) + d(N, P_2) + d(N, P_3),$$

where d denotes the intrinsic distance in S^2 . Lifting this example into $U(2)$ by using the Hopf fibration $S^1 \rightarrow SU(2) \cong S^3 \rightarrow S^2$, we obtain the following matrices in $U(2)$:

$$(B.3) \quad A_0 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad A_j = \begin{pmatrix} re^{i\alpha_k} & -\sqrt{1-r^2} \\ \sqrt{1-r^2} & re^{-i\alpha_k} \end{pmatrix}, \quad k = 1, 2, 3,$$

where $r \in [0, 1]$, $\alpha_k = 2(k-1)\pi/3$. We check that

$$|A_k - A_l|_F = \sqrt{6}r \quad \forall 1 \leq k \neq l \leq 3$$

and

$$|A_0 - A_k|_F = 2\sqrt{1 - \sqrt{1-r^2}} \quad \forall k = 1, 2, 3.$$

Therefore, for small r (e.g., when $r \leq 0.85$),

$$(B.4) \quad \sum_{k=1}^3 |A_0 - A_k|_F < \min_{A \in \{A_1, A_2, A_3\}} \sum_{k=1}^3 |A - A_k|_F.$$

Equation (B.4) implies that the generalization of Lemma 4.11 does not hold in $U(2)$.

For later purposes, we transform one of the matrices $\{A_1, A_2, A_3\}$ to be the identity matrix I_2 . Set

$$B_0 = A_3^{-1}A_0, B_k = A_3^{-1}A_k, \quad k = 1, 2, 3.$$

Then we have $B_3 = I_2$. Using the definition of the Frobenius norm, we obtain

$$(B.5) \quad |B_1 - B_2|_F = |B_2 - B_3|_F = |B_3 - B_1|_F = \sqrt{6}r,$$

and, for small r ,

$$(B.6) \quad \sum_{k=1}^3 |B_0 - B_k|_F < \min_{B \in \{B_1, B_2, B_3\}} \sum_{k=1}^3 |B - B_k|_F \leq |B_1 - I_2|_F + |B_2 - I_2|_F.$$

Let us consider the graph shown in Figure 6. This is a graph with vertex set $V = \{x, y, z, w\}$, edge set $E = \{\{x, y\}, \{y, z\}, \{z, w\}, \{w, x\}, \{y, w\}\}$, and a signature $\sigma : E^{or} \rightarrow U(2)$ as shown in the figure.

PROPOSITION B.1. *For the graph as shown in Figure 6, we have*

$$\iota_F^\sigma(V) := \min_{\tau: V \rightarrow U(2)} \sum_{\{x, y\} \in E_S} w_{xy} |\sigma_{xy} \tau(y) - \tau(x)|_F < \min_{T \in \mathbb{T}_S} \sum_{\{x, y\} \in E} |\sigma_{xy} \tau_T(y) - \tau_T(x)|_F,$$

where τ_T is a representative of the set

$$C_T(V) = \{\tau : V \rightarrow U(2) : \tau \text{ is constant on } T \text{ w.r.t. } \sigma\}.$$




FIG. 6. A counterexample.

Proof. Observe that $T_1 = (V, \{\{x, y\}, \{y, w\}, \{y, z\}\})$ is a spanning tree and that the function $\tau_{T_1} \equiv I_2$ is a constant function on T_1 with respect to σ . We calculate

$$\begin{aligned} \sum_{\{x,y\} \in E} |\sigma_{xy} \tau_{T_1}(y) - \tau_{T_1}(x)|_F &= |B_1 - I_2|_{2,1} + |B_2 - I_2|_F \\ &\stackrel{(B.6)}{>} \sum_{k=1}^3 |B_0 - B_k|_F \\ &= \sum_{\{x,y\} \in E} |\sigma_{xy} \tau_0(y) - \tau_0(x)|_F, \end{aligned}$$

where the switching function τ_0 is defined via $\tau_0(x) = \tau_0(y) = \tau_0(z) = B_0$, $\tau_0(w) = I_2$. Therefore, by definition, we have

$$(B.7) \quad \iota_F^\sigma(V) < \sum_{\{x,y\} \in E} |\sigma_{xy} \tau_{T_1}(y) - \tau_{T_1}(x)|_F.$$

The graph in Figure 6 has eight spanning trees, which we denote by T_i , $i = 1, 2, \dots, 8$. We claim that

$$(B.8) \quad \sum_{\{x,y\} \in E} |\sigma_{xy} \tau_{T_i}(y) - \tau_{T_i}(x)|_F = |B_1 - I_2|_F + |B_2 - I_2|_F \quad \forall i = 1, 2, \dots, 8.$$

The proposition then follows immediately from (B.7) and (B.8).

Claim (B.8) can be checked directly with the help of (B.5) for all choices of spanning trees. \square

Acknowledgments. The second author acknowledges the hospitality of the Department of Mathematical Sciences of Durham University during his visit in February 2015. Parts of this work were completed while the first and second authors were attending the *Spring School: Discrete Ricci Curvature* at Institut Henri Poincaré, Paris, in May 2015, and while the first and third authors were visiting the Max-Planck-Institut für Mathematik, Bonn, in July 2015. We thank these institutes for their hospitality. Finally, we thank the referees for their helpful comments.

REFERENCES

- [1] N. AGARWAL, A. KOLLA, AND V. MADAN, *Small Lifts of Expander Graphs Are Expanding*, arXiv:1311.3268v1, 2013.
- [2] N. ALON, *Eigenvalues and expanders*, Combinatorica, 6 (1986), pp. 83–96.
- [3] N. ALON AND V. MILMAN, λ_1 , *isoperimetric inequalities for graphs, and superconcentrators*, J. Combin. Theory Ser. B, 38 (1985), pp. 73–88.
- [4] F. M. ATAY AND S. LIU, *Cheeger Constants, Structural Balance, and Spectral Clustering Analysis for Signed Graphs*, arXiv:1411.3530, 2017.
- [5] D. BAKRY, *Functional inequalities for Markov semigroups*, in Probability Measures on Groups: Recent Directions and Trends, S. G. Dani and P. Graczyk, eds., Tata Institute of Fundamental Research Mumbai, 2006, pp. 91–147.
- [6] D. BAKRY AND M. ÉMERY, *Diffusions hypercontractives*, Séminaire de probabilités, XIX, 1983/84, Lecture Notes in Math. 1123, J. Azéma and M. Yor, eds., Springer, Berlin, 1985, pp. 177–206.
- [7] A. S. BANDEIRA, A. SINGER, AND D. A. SPIELMAN, *A Cheeger inequality for the graph connection Laplacian*, SIAM J. Matrix Anal. Appl., 34 (2013), pp. 1611–1630.
- [8] F. BAUER, P. HORN, Y. LIN, G. LIPPNER, D. MANGOUBI, AND S.-T. YAU, *Li-Yau inequality on graphs*, J. Differential Geom., 99 (2015), pp. 359–405.

- [9] F. BAUER AND J. JOST, *Bipartite and neighborhood graphs and the spectrum of the normalized graph Laplacian*, Comm. Anal. Geom., 21 (2013), pp. 787–845.
- [10] F. BAUER, J. JOST, AND S. LIU, *Ollivier-Ricci curvature and the spectrum of the normalized graph Laplace operator*, Math. Res. Lett., 19 (2012), pp. 1185–1205.
- [11] Y. BILU AND N. LINIAL, *Lifts, discrepancy and nearly optimal spectral gap*, Combinatorica, 26 (2006), pp. 495–519.
- [12] P. BUSER, *A note on the isoperimetric constant*, Ann. Sci. École Norm. Sup. (4), 15 (1982), pp. 213–230.
- [13] F. R. K. CHUNG, Y. LIN, AND S.-T. YAU, *Harnack inequalities for graphs with non-negative Ricci curvature*, J. Math. Anal. Appl., 415 (2014), pp. 25–32.
- [14] F. R. K. CHUNG AND P. TETALI, *Isoperimetric inequalities for Cartesian products of graphs*, Combin. Probab. Comput., 7 (1998), pp. 141–148.
- [15] D. CVETKOVIĆ AND S. K. SIMIĆ, *Towards a spectral theory of graphs based on the signless Laplacian*, I, Publ. Inst. Math. (Beograd) (N.S.), 85 (2009), pp. 19–33.
- [16] J. DODZIUK, *Difference equations, isoperimetric inequality and transience of certain random walks*, Trans. Amer. Math. Soc., 284 (1984), pp. 787–794.
- [17] K. D. ELWORTHY, *Manifolds and graphs with mostly positive curvatures*, in: Stochastic Analysis and Applications (Lisbon, 1989), Progr. Probab. 26, Birkhäuser Boston, Boston, MA, 1991, pp. 96–110.
- [18] K. FUNANO, *Eigenvalues of Laplacian and Multi-way Isoperimetric Constants on Weighted Riemannian Manifolds*, arXiv:1307.3919v1, 2013.
- [19] T. GAO, J. BRODZKI, AND S. MUKHERJEE, *The Geometry of Synchronization Problems and Learning Group Actions*, arXiv:1610.09051, 2016.
- [20] S. GALLOT, D. HULIN, AND J. LAFONTAINE, *Riemannian Geometry*, 3rd ed., Universitext, Springer-Verlag, Berlin, 2004.
- [21] J. L. GROSS, *Voltage graphs*, Discrete Math., 9 (1974), pp. 239–246.
- [22] J. L. GROSS AND T. W. TUCKER, *Quotients of complete graphs: Revisiting the Heawood map-coloring problem*, Pacific J. Math., 55 (1974), pp. 391–402.
- [23] F. HARARY, *On the notion of balance of a signed graph*, Michigan Math. J., 2 (1953), pp. 143–146.
- [24] F. HARARY AND J. A. KABELL, *A simple algorithm to detect balance in signed graphs*, Math. Social Sci., 1 (1980/81), pp. 131–136.
- [25] P. HORN, Y. LIN, S. LIU, AND S.-T. YAU, *Volume doubling, Poincaré inequality and Gaussian heat kernel estimate for nonnegative curvature graphs*, J. Reine Angew. Math., to appear, doi:10.1515/crelle-2017-0038.
- [26] B. HUA AND Y. LIN, *Stochastic completeness for graphs with curvature dimension conditions*, Adv. Math., 306 (2017), pp. 279–302.
- [27] J. JOST AND S. LIU, *Ollivier’s Ricci curvature, local clustering and curvature-dimension inequalities on graphs*, Discrete Comput. Geom., 51 (2014), pp. 300–322.
- [28] T. KAUFMAN, D. KAZHDAN, AND A. LUBOTZKY, *Isoperimetric inequalities for Ramanujan complexes and topological expanders*, Geom. Funct. Anal., 26 (2016), pp. 250–287.
- [29] B. KLARTAG, G. KOZMA, P. RALLI, AND P. TETALI, *Discrete curvature and abelian groups*, Canad. J. Math., 68 (2016), pp. 655–674.
- [30] M. KREBS AND A. SHAHEEN, *Expander Families and Cayley Graphs, A Beginner’s Guide*, Oxford University Press, Oxford, UK, 2011.
- [31] T.-C. KWOK, L.-C. LAU, Y.-T. LEE, S. OVEIS GHARAN, AND L. TREVISAN, *Improved Cheeger’s inequality: Analysis of spectral partitioning algorithms through higher order spectral gap*, in Proceedings of the 2013 ACM Symposium on Theory of Computing, ACM, New York, 2013, pp. 11–20.
- [32] C. LANGE, S. LIU, N. PEYERIMHOFF, AND O. POST, *Frustration index and Cheeger inequalities for discrete and continuous magnetic Laplacians*, Calc. Var. Partial Differential Equations, 54 (2015), pp. 4165–4196.
- [33] M. LEDOUX, *Spectral gap, logarithmic Sobolev constant, and geometric bounds*, in Eigenvalues of Laplacians and other Geometric Operators, Surv. Differ. Geom., 9, A. Grigor’yan and S.-T. Yau, ed., International Press, Somerville, MA, 2004, pp. 219–240.
- [34] Y. LIN AND S.-T. YAU, *Ricci curvature and eigenvalue estimate on locally finite graphs*, Math. Res. Lett., 17 (2010), pp. 343–356.
- [35] S. LIU, *Multi-way dual Cheeger constants and spectral bounds of graphs*, Adv. Math., 268 (2015), pp. 306–338.
- [36] S. LIU, *An Optimal Dimension-Free Upper Bound for Eigenvalue Ratios*, arXiv:1405.2213, 2014.
- [37] S. LIU AND N. PEYERIMHOFF, *Eigenvalue ratios of nonnegatively curved graphs*, Combin. Probab. Comput., 27 (2018), pp. 829–850, doi:10.1017/S0963548318000214.

- [38] S. LIU, N. PEYERIMHOFF, AND A. VDOVINA, *Signatures, Lifts and Eigenvalues of Graphs*, arXiv:1412.6841, 2014.
- [39] A. W. MARCUS, D. A. SPIELMAN, AND N. SRIVASTAVA, *Interlacing families I: Bipartite Ramanujan graphs of all degrees*, Ann. of Math., 182 (2015), pp. 307–325.
- [40] M. MIMURA, *Multi-way expanders and imprimitive group actions on graphs*, Int. Math. Res. Note IMRN, 8 (2016), pp. 2522–2543.
- [41] F. MÜNCH, *Li-Yau inequality on finite graphs via non-linear curvature dimension conditions*, J. Math. Pures Appl. (9), 120 (2018), pp. 130–164.
- [42] F. MÜNCH, *Remarks on curvature dimension conditions on graphs*, Calc. Var. Partial Differential Equations, 56 (2017), 11.
- [43] Y. OLLIVIER, *Ricci curvature of Markov chains on metric spaces*, J. Funct. Anal., 256 (2009), pp. 810–864.
- [44] M. SCHMUCKENSCHLÄGER, *Curvature of nonlocal Markov generators*, in Convex Geometric Analysis (Berkeley, CA, 1996), Math. Sci. Res. Inst. Publ. 34, Cambridge University Press, Cambridge, UK, 1999, pp. 189–197.
- [45] M. A. SHUBIN, *Discrete magnetic Laplacian*, Comm. Math. Phys., 164 (1994), pp. 259–275.
- [46] A. SINGER AND H.-T. WU, *Vector diffusion maps and the connection Laplacian*, Comm. Pure Appl. Math., 65 (2012), pp. 1067–1144.
- [47] T. SUNADA, *A discrete analogue of periodic magnetic Schrödinger operators*, in Geometry of the Spectrum (Seattle, WA, 1993), Contemp. Math. 173, AMS, Providence, RI, 1994, pp. 283–299.
- [48] M. TANAKA, *Multi-way Expansion Constants and Partitions of a Graph*, arXiv:1112.3434, 2013.
- [49] L. TREVISAN, *Max cut and the smallest eigenvalue*, SIAM J. Comput., 41 (2012), pp. 1769–1786.
- [50] T. ZASLAVSKY, *Signed graphs*, Discrete Appl. Math., 4 (1982), pp. 47–74.
- [51] T. ZASLAVSKY, *Matrices in the theory of signed simple graphs*, in Advances in Discrete Mathematics and Applications (Mysore, 2008), Ramanujan Math. Soc. Lect. Notes Ser. 13, B. D. Acharya, G. O. H. Katona, and J. Nešetřil, ed., Ramanujan Mathematical Society, Mysore, 2010, pp. 207–229.