TEMPLATES FOR BINARY MATROIDS

KEVIN GRACE AND STEFAN H. M. VAN ZWAM

ABSTRACT. A binary frame template is a device for creating binary matroids from graphic or cographic matroids. Such matroids are said to conform or coconform to the template. We introduce a preorder on these templates and determine the nontrivial templates that are minimal with respect to this order. As an application of our main result, we determine the eventual growth rates of certain minor-closed classes of binary matroids, including the class of binary matroids with no minor isomorphic to PG(3, 2). Our main result applies to all highly-connected matroids in a class, not just those of maximum size. As a second application, we characterize the highly-connected 1-flowing matroids.

1. INTRODUCTION

Geelen, Gerards, and Whittle [1] recently announced a structure theorem describing the highly connected members of any proper minorclosed class of matroids representable over a given finite field. In this paper we study some consequences of their result. To state a first, rough version of their result, we need the following definitions.

A matroid M is vertically k-connected if, for each partition (X, Y)of the ground set of M with r(X) + r(Y) - r(M) < k - 1, either X or Y is spanning. We denote the unique prime subfield of \mathbb{F} by $\mathbb{F}_{\text{prime}}$. We say that a matroid M_2 is a rank- $(\leq t)$ perturbation of a matroid M_1 if there exist matrices A_1 and A_2 over \mathbb{F} such that $r(M(A_1 - A_2)) \leq t$ and such that $M_1 \cong M(A_1)$ and $M_2 \cong M(A_2)$.

We now restate [1, Theorem 3.3]. Its proof is forthcoming in a future paper by Geelen, Gerards, and Whittle.

Theorem 1.1. Let \mathbb{F} be a finite field and let m_0 be a positive integer. Then there exist $k, n, t \in \mathbb{Z}_+$ such that, if M is a matroid representable over \mathbb{F} such that M or M^* is vertically k-connected and such that M

¹⁹⁹¹ Mathematics Subject Classification. 05B35.

The first author was supported by a Huel D. Perkins Fellowship from the Louisiana State University Graduate School. The second author was supported by the National Science Foundation, grant 1500343.

has an $M(K_n)$ -minor but no $PG(m_0 - 1, \mathbb{F}_{prime})$ -minor, then M is a rank- $(\leq t)$ perturbation of a frame matroid representable over \mathbb{F} .

Let us consider a very simple example of a rank-1 perturbation. Let A_1 be the binary matrix

1	0	0	0	1	1	1	0	0	0
0	1	0	0	1	0	0	1	1	0
0	0	1	0	0	1	0	1	0	1
0	0	0	1	0	0	1	0	1	1

and let A_2 be the binary matrix

Note that A_2 is the result of adding the rank-1 matrix

1	1	1	1	0	0	0	0	0	0
1	1	1	1	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0

to A_1 . Therefore, the vector matroid $M(A_2)$ is a rank-1 perturbation of $M(A_1)$.

Theorem 1.1 is essentially a simplified version of a much more complex structure theorem [1, Theorem 4.2]. Geelen, Gerards, and Whittle introduced the concept of a *template* as a tool to capture much of this complexity.

Our focus in this paper is on the binary case. Roughly speaking, a binary frame template can be thought of as a recipe for constructing a representable matroid from a graphic or cographic matroid. A matroid constructed in this way is said to *conform* or *coconform* to the template.

In the example above, we may think of $M(A_2)$ as the matroid obtained from the vector matroid of the following matrix by contracting the element indexing the final column. Note that the large submatrix on the bottom left is A_1 :

[1	1	1	1	0	0	0	0	0	0	1
1	0	0	0	1	1	1	0	0	0	1
0	1	0	0	1	0	0	1	1	0	1
0	0	1	0	0	1	0	1	0	1	0
0	0	0	1	0	0	1	0	1	1	0

In fact, for any matrix A of the following form, where v and w are arbitrary binary vectors, the matroid M(A)/c conforms to the template Φ_{CX} , which we will define in Section 3:



Let $\mathcal{M}(\Phi)$ denote the set of matroids representable over a field \mathbb{F} that conform to a frame template Φ . Theorem 1.2 below is a slight modification of [1, Theorem 4.2]. The modification is explained in Section 2.

Theorem 1.2. Let \mathbb{F} be a finite field, let m be a positive integer, and let \mathcal{M} be a minor-closed class of matroids representable over \mathbb{F} . Then there exist $k, l \in \mathbb{Z}_+$ and frame templates $\Phi_1, \ldots, \Phi_s, \Psi_1, \ldots, \Psi_t$ such that

- \mathcal{M} contains each of the classes $\mathcal{M}(\Phi_1), \ldots, \mathcal{M}(\Phi_s)$,
- \mathcal{M} contains the duals of the matroids in each of the classes $\mathcal{M}(\Psi_1), \ldots, \mathcal{M}(\Psi_t)$, and
- if M is a simple vertically k-connected member of M with at least l elements and with no PG(m - 1, F_{prime}) minor, then either M is a member of at least one of the classes M(Φ₁),..., M(Φ_s), or M^{*} is a member of at least one of the classes M(Ψ₁),..., M(Ψ_t).

Our contribution is to shed some light on how these templates are related to each other. We define a preorder on the set of frame templates. Our main result, Theorem 3.19, is a list of nontrivial binary frame templates that are minimal with respect to this preorder.

One application of this result involves growth rates of minor-closed classes of binary matroids. The growth rate function of a minor-closed class \mathcal{M} is the function whose value at an integer $r \geq 0$ is given by the maximum number of elements in a simple matroid in \mathcal{M} of rank at most r. We prove that a minor-closed class of binary matroids has a growth rate that is eventually equal to the growth rate of the class of graphic matroids if and only if it contains all graphic matroids but does not contain the class of matroids conforming to a certain template. The class of matroids conforming to this template is exactly the class of matroids having an even-cycle representation with a blocking pair. Geelen and Nelson also proved this result in [5]. We also prove the following theorem. Here, $\mathcal{EX}(F)$ denotes the class of binary matroids with no *F*-minor. If *f* and *g* are functions, we write $f(r) \approx g(r)$ if f(r) = g(r) for all but finitely many *r*.

Theorem 1.3. The growth rate function for $\mathcal{EX}(PG(3,2))$ is

 $h_{\mathcal{EX}(PG(3,2))} \approx r^2 - r + 1.$

Note that $r^2 - r + 1$ is the growth rate of the class of even-cycle matroids.

Our main result goes beyond growth rates because it gives information about all highly-connected matroids in a minor-closed class, not just the maximum-sized matroids. This is illustrated by our second application, involving 1-flowing matroids. The 1-flowing property is a generalization of the max-flow min-cut property of graphs. We prove the following.

Theorem 1.4. There exist $k, l \in \mathbb{Z}_+$ such that every simple, vertically k-connected, 1-flowing matroid with at least l elements is either graphic or cographic.

We use templates to study a minor-closed class \mathcal{M} by describing the highly-connected matroids in the class. This analysis follows a certain pattern:

- (1) Find a matroid N not in \mathcal{M} .
- (2) Find all templates such that N is not a minor of any matroid conforming to that template.
- (3) If all matroids conforming to these templates are in \mathcal{M} , then the analysis is complete.
- (4) Otherwise, repeat Step (1).

From the definition of conforming to a template, which we will give in Section 2, it will not be difficult to see that for each binary frame template Φ , there are integers t_1 and t_2 such that every matroid conforming to Φ is a rank-($\leq t_1$) perturbation of a graphic matroid and every matroid coconforming to Φ is a rank-($\leq t_2$) perturbation of a cographic matroid. Thus, by Theorem 1.2, the highly connected matroids in a minor-closed class of binary matroids are "close" to being graphic or cographic. In this regard, the work regarding templates resembles work done by Robertson and Seymour concerning minor-closed classes of graphs. In Theorem 1.3 of [9], Robertson and Seymour showed that highly-connected graphs in a minor-closed class are in some sense "close" to being embeddable in some surface.

Section 2 of this paper repeats the necessary definitions found in [1]. In Section 3, we prove our main result, as well as giving some machinery leading up to it. Section 4 applies our result to growth rates of minor-closed classes of binary matroids, and in Section 5, we prove Theorem 1.4.

2. Preliminaries

We repeat here several definitions concerning highly connected matroids which can be found in Geelen, Gerards, and Whittle [1]. Although the results found in [1] are technically about matrices rather than matroids, it suffices for our purposes to state the results in terms of their immediate matroid consequences.

Let A be a matrix over a field \mathbb{F} . Then A is a *frame matrix* if each column of A has at most two nonzero entries. We let \mathbb{F}^{\times} denote the multiplicative group of \mathbb{F} . Let Γ be a subgroup of \mathbb{F}^{\times} . A Γ -frame matrix is a frame matrix A such that:

- Each column of A with a nonzero entry contains a 1.
- If a column of A has a second nonzero entry, then that entry is $-\gamma$ for some $\gamma \in \Gamma$.

In the case where Γ is the multiplicative group of one element, a matrix is a Γ -frame matrix if and only if it is the signed incidence matrix of a graph, with possibly a row removed. In particular, a binary matroid is graphic if and only if it can be represented by a matrix over GF(2) in which no column has more than two nonzero entries.

To facilitate the description of their structure theorem, Geelen, Gerards, and Whittle capture capture much of the complexity with the concept of a "template." Let \mathbb{F} be a finite field. A *frame template* over \mathbb{F} is a tuple $\Phi = (\Gamma, C, X, Y_0, Y_1, A_1, \Delta, \Lambda)$ such that the following hold¹:

- (i) Γ is a subgroup of \mathbb{F}^{\times} .
- (ii) C, X, Y_0 and Y_1 are disjoint finite sets.
- (iii) $A_1 \in \mathbb{F}^{X \times (C \cup Y_0 \cup Y_1)}$.
- (iv) Λ is a subgroup of the additive group of \mathbb{F}^X and is closed under scaling by elements of Γ .
- (v) Δ is a subgroup of the additive group of $\mathbb{F}^{C \cup Y_0 \cup Y_1}$ and is closed under scaling by elements of Γ .

Let $\Phi = (\Gamma, C, X, Y_0, Y_1, A_1, \Delta, \Lambda)$ be a frame template. Let *B* and *E* be finite sets, and let $A' \in \mathbb{F}^{B \times E}$. We say that *A'* respects Φ if the following hold:

- (i) $X \subseteq B$ and $C, Y_0, Y_1 \subseteq E$.
- (ii) $A'[X, C \cup Y_0 \cup Y_1] = A_1.$

¹The authors of [1] divided our set X into two separate sets which they called X and D. Their set X can be absorbed into Y_0 , therefore we omit it.

		Z	$Y_0 Y_1 C$
X	columns from Λ	0	A_1
	Γ-frame matrix	unit columns	rows from Δ

FIGURE 1.

- (iii) There exists a set $Z \subseteq E (C \cup Y_0 \cup Y_1)$ such that A'[X, Z] = 0, each column of A'[B - X, Z] is a unit vector, and $A'[B - X, E - (C \cup Y_0 \cup Y_1 \cup Z)]$ is a Γ -frame matrix.
- (iv) Each column of $A'[X, E (C \cup Y_0 \cup Y_1 \cup Z)]$ is contained in Λ .
- (v) Each row of $A'[B X, C \cup Y_0 \cup Y_1]$ is contained in Δ .

Figure 1 shows the structure of A'.

Suppose that A' respects Φ and that Z satisfies (iii) above. Now suppose that $A \in \mathbb{F}^{B \times E}$ satisfies the following conditions:

- (i) A[B, E Z] = A'[B, E Z]
- (ii) For each $i \in Z$ there exists $j \in Y_1$ such that the *i*-th column of A is the sum of the *i*-th and the *j*-th columns of A'.

We say that any such matrix *conforms* to Φ .

Let M be a matroid representable over \mathbb{F} . We say that M conforms to Φ if there is a matrix A that conforms to Φ such that M is isomorphic to $M(A)/C \setminus Y_1$.

Let $\mathcal{M}(\Phi)$ denote the set of matroids representable over \mathbb{F} that conform to Φ . Recall that a matroid M is vertically k-connected if, for each partition (X, Y) of the ground set of M with r(X)+r(Y)-r(M) < k-1, either X or Y is spanning. We denote the unique prime subfield of \mathbb{F} by $\mathbb{F}_{\text{prime}}$. Geelen, Gerards, and Whittle will prove Theorem 1.2 in a future paper. This theorem is actually a slight modification of the theorem found in [1]. In that paper, there is no mention of the requirement that a matroid have size at least l. However, Geelen (personal communication) has stated that this is necessary to ensure that adding a finite number of matroids to the class \mathcal{M} does not add any templates to the list $\Phi_1, \ldots, \Phi_s, \Psi_1, \ldots, \Psi_t$.

Although the term coconform does not appear in [1], we define it in the following obvious way.

Definition 2.1. A matroid M coconforms to a template Φ if its dual M^* conforms to Φ .

To simplify the proofs in this paper, it will be helpful to expand the concept of conforming slightly.

Definition 2.2. Let A' be a matrix that respects Φ , as defined above, except that we allow columns of A'[B-X, Z] to be either unit columns or zero columns. Let A be a matrix that is constructed from A' as described above. Thus, A[B, E-Z] = A'[B, E-Z], and for each $i \in Z$ there exists $j \in Y_1$ such that the *i*-th column of A is the sum of the *i*-th and the *j*-th columns of A'. Let M be isomorphic to $M(A)/C \setminus Y_1$. We say that A and M virtually conform to Φ and that A' virtually respects Φ . If M^* virtually conforms to Φ , we say that M virtually coconforms to Φ .

We will denote the set of matroids representable over \mathbb{F} that virtually conform to Φ by $\mathcal{M}_v(\Phi)$ and the set of matroids representable over \mathbb{F} that virtually coconform to Φ by $\mathcal{M}_v^*(\Phi)$.

The following notation will be used throughout this paper. We denote an empty matrix by $[\emptyset]$. We denote a group of one element by $\{0\}$ or $\{1\}$, if it is an additive or multiplicative group, respectively. If S' is a subset of a set S and G is a subgroup of the additive group \mathbb{F}^S , we denote by G|S' the projection of G into $\mathbb{F}^{S'}$. Similarly, if $\bar{x} \in G$, we denote the projection of \bar{x} into $\mathbb{F}^{S'}$ by $\bar{x}|S'$.

Unexplained notation and terminology will generally follow Oxley [8]. One exception is that we denote the vector matroid of a matrix A by M(A), rather than M[A].

3. Reducing a Template

In this section, we will introduce reductions and show that every template reduces to one of several basic templates.

Since templates are used to study minor-closed classes of matroids, a natural question to ask is whether the set of matroids conforming to a particular template is minor-closed. The answer is no, in general. For example, if $|Y_0| = 1$, then a matroid conforms to the following binary frame template if and only if it is a graphic matroid with a loop:

$$\{\{1\}, \emptyset, \emptyset, Y_0, \emptyset, [\emptyset], \{0\}, \{0\}\}$$

Clearly, this is not a minor-closed class.

Another question to ask is whether there might be some sort of minor relationship between a pair of templates, where every matroid conforming to one template is a minor of a matroid conforming to the other. These questions motivate the following discussion. **Definition 3.1.** A reduction is an operation on a frame template Φ that produces a frame template Φ' such that $\mathcal{M}(\Phi') \subseteq \mathcal{M}(\Phi)$.

Proposition 3.2. The following operations are reductions on a frame template Φ :

- (1) Replace Γ with a proper subgroup.
- (2) Replace Λ with a proper subgroup closed under multiplication by elements from Γ .
- (3) Replace Δ with a proper subgroup closed under multiplication by elements from Γ .
- (4) Remove an element y from Y_1 . (More precisely, replace A_1 with $A_1[X, Y_0 \cup (Y_1 y) \cup C]$ and replace Δ with $\Delta|(Y_0 \cup (Y_1 y) \cup C)$.
- (5) For all matrices A' respecting Φ , perform an elementary row operation on A'[X, E]. (Note that this alters the matrix A_1 and performs a change of basis on Λ .)
- (6) If there is some element x ∈ X such that, for every matrix A' respecting Φ, we have that A'[{x}, E] is a zero row vector, remove x from X. (This simply has the effect of removing a zero row from every matrix conforming to Φ.)
- (7) Let $c \in C$ be such that $A_1[X, \{c\}]$ is a unit column whose nonzero entry is in the row indexed by $x \in X$, and let either $\lambda_x = 0$ for each $\lambda \in \Lambda$ or $\delta_c = 0$ for each $\delta \in \Delta$. Let Δ' be the result of adding $-\delta_c A_1[\{x\}, Y_0 \cup Y_1 \cup C]$ to each element $\delta \in \Delta$. Replace Δ with Δ' , and then remove c from C and d from D. (More precisely, replace A_1 with $A_1[X - x, Y_0 \cup Y_1 \cup (C - c)]$, replace Λ with $\Lambda|(X-x)$, and replace Δ with $\Delta'|(Y_0 \cup Y_1 \cup (C-c))$.)
- (8) Let $c \in C$ be such that $A_1[X, \{c\}]$ is a zero column and $\delta_c = 0$ for all $\delta \in \Delta$. Then remove c from C. (More precisely, replace A_1 with $A_1[X, Y_0 \cup Y_1 \cup (C c)]$, and replace Δ with $\Delta|(Y_0 \cup Y_1 \cup (C c)).)$

Proof. Let Φ' be the template that results from performing one of operations (1)-(8) on Φ .

For (1)-(3), every matrix A' respecting Φ' also respects Φ .

For (4), let A' be a matrix respecting Φ' , and let M be the matroid $M(A)/C \setminus Y_1$, where A is a matrix conforming to Φ' that has been constructed from A' respecting Φ' as described in Section 2. Since Y_1 is deleted to produce M, the only effect of Y_1 on M is that for each $i \in Z$ there exists $j \in Y_1$ such that the *i*-th column of A is the sum of the *i*-th and the *j*-th columns of A'. But each $j \in Y_1$ in the template Φ' is also contained in Y_1 in the template Φ . Therefore, A conforms to Φ , as does M.

For (5) and (6), elementary row operations and removing zero rows produce isomorphic matroids.

Operations (7) and (8) have the effect of contracting c from $M(A) \setminus Y_1$ for every matrix A conforming to Φ . Since all of C is contracted to produce a matroid M conforming to Φ , the matroids we produce by performing either of these operations still conform to Φ .

For $i \in \{1, \ldots, 8\}$, we call operation (i) above a reduction of type i.

The operations listed in the definition below are not reductions as defined above, but we continue the numbering from Proposition 3.2 for ease of reference.

Definition 3.3. A template Φ' is a *template minor* of Φ if Φ' is obtained from Φ by repeatedly performing the following operations:

- (9) Performing a reduction of type 1-8 on Φ .
- (10) Removing an element y from Y_0 , replacing A_1 with $A_1[X, (Y_0 y) \cup Y_1 \cup C]$, and replacing Δ with $\Delta|((Y_0 y) \cup Y_1 \cup C))$. (This has the effect of deleting y from every matroid conforming to Φ .)
- (11) Let $x \in X$ with $\lambda_x = 0$ for every $\lambda \in \Lambda$, and let $y \in Y_0$ be such that $(A_1)_{x,y} \neq 0$. Then contract y from every matroid conforming to Φ . (More precisely, perform row operations on A_1 so that $A_1[X, \{y\}]$ is a unit column with $(A_1)_{x,y} =$ 1. Then replace every element $\delta \in \Delta$ with the row vector $-\delta_y A_1[\{x\}, Y_0 \cup Y_1 \cup C] + \delta$. This induces a group homomorphism $\Delta \to \Delta'$, where Δ' is also a subgroup of the additive group of $\mathbb{F}^{C \cup Y_0 \cup Y_1}$ and is closed under scaling by elements of Γ . Finally, replace A_1 with $A_1[X - x, (Y_0 - y) \cup Y_1 \cup C]$, project Λ into \mathbb{F}^{X-x} , and project Δ' into $\mathbb{F}^{(Y_0-y)\cup Y_1\cup C}$. The resulting groups play the roles of Λ and Δ , respectively in Φ' .)
- (12) Let $y \in Y_0$ with $\delta_y = 0$ for every $\delta \in \Delta$. Then contract y from every matroid conforming to Φ . (More precisely, if $A_1[X, \{y\}]$ is a zero vector, this is the same as simply removing y from Y_0 . Otherwise, choose some $x \in X$ such that $(A_1)_{x,y} \neq 0$. Then for every matrix A' that respects Φ , perform row operations so that $A_1[X, \{y\}]$ is a unit column with $(A_1)_{x,y} = 1$. This induces a group isomorphism $\Lambda \to \Lambda'$ where Λ' is also a subgroup of the additive group of \mathbb{F}^X and is closed under scaling by elements of Γ . Finally, replace A_1 with $A_1[X - x, (Y_0 - y) \cup Y_1 \cup C]$, project Λ' into \mathbb{F}^{X-x} , and project Δ into $\mathbb{F}^{(Y_0-y)\cup Y_1\cup C}$. The resulting groups play the roles of Λ and Δ , respectively in Φ' .)

Let Φ' be a template minor of Φ , and let A' be a matrix that virtually respects Φ' . Let A be a matrix that virtually conforms to Φ' , and let M be a matroid that virtually conforms to Φ' . We say that A' weakly respects Φ and that A and M weakly conform to Φ . Let $\mathcal{M}_w(\Phi)$ denote the set of matroids representable over \mathbb{F} that weakly conform to Φ , and let $\mathcal{M}_w^*(\Phi)$ denote the set of matroids representable over \mathbb{F} whose duals weakly conform to Φ . If $M \in \mathcal{M}_w^*(\Phi)$, we say that M weakly coconforms to Φ .

Lemma 3.4. If a matroid M weakly conforms to a template Φ , then M is a minor of a matroid that conforms to Φ .

Proof. Let Φ' be a template minor of Φ . As we can see from Definition 3.3, every matroid M weakly conforming to Φ' is a minor of a matroid virtually conforming to Φ . It remains to analyze the case where M virtually conforms to Φ ; so M is isomorphic to $M(K)/C \setminus Y_1$, where K is built from a matrix K' that virtually respects Φ . Consider the following matrix A' obtained from K' by adding a row r and a column c.

	c			Z	$Y_0 Y_1 C$
X	0	columns from Λ		0	A_1
	0	Γ-frame matrix	0	unit columns	rows from Δ
r	1	0	$1 \cdots 1$	0	0

From A', we can obtain a matrix A conforming to Φ such that M is isomorphic to $M(A)/C \setminus Y_1/c$. So M is a minor of a matroid conforming to Φ .

An easy consequence of Lemma 3.4 is that Theorem 1.2, which deals with minor-closed classes, can be restated in terms of weak conforming.

Corollary 3.5. Let \mathbb{F} be a finite field, let m be a positive integer, and let \mathcal{M} be a minor-closed class of matroids representable over \mathbb{F} . Then there exist $k, l \in \mathbb{Z}_+$ and frame templates $\Phi_1, \ldots, \Phi_s, \Psi_1, \ldots, \Psi_t$ such that

- \mathcal{M} contains each of the classes $\mathcal{M}_w(\Phi_1), \ldots, \mathcal{M}_w(\Phi_s)$,
- \mathcal{M} contains the duals of the matroids in each of the classes $\mathcal{M}_w(\Psi_1), \ldots, \mathcal{M}_w(\Psi_t)$, and

10

 if M is a simple vertically k-connected member of M with at least l elements and with no PG(m − 1, F_{prime}) minor, then either M is a member of at least one of the classes M_v(Φ₁),..., M_v(Φ_s) or M^{*} is a member of at least one of the classes M_v(Ψ₁),..., M_v(Ψ_t).

Proof. Let $\Phi_1, \ldots, \Phi_s, \Psi_1, \ldots, \Psi_t$ be the templates whose existence is implied by Theorem 1.2. For $\Phi \in {\Phi_1, \ldots, \Phi_s}$, Lemma 3.4 implies that any matroid $M \in \mathcal{M}_w(\Phi)$ is a minor of a matroid $N \in \mathcal{M}(\Phi)$. Since \mathcal{M} contains $\mathcal{M}(\Phi)$ and is minor-closed, \mathcal{M} contains $\mathcal{M}_w(\Phi)$ as well. Similarly, \mathcal{M} contains the duals of the matroids in each of the classes $\mathcal{M}_w(\Psi_1), \ldots, \mathcal{M}_w(\Psi_t)$. The third condition above holds since every matroid conforming to a template also virtually conforms to it. \Box

If $\mathcal{M}_w(\Phi) = \mathcal{M}_w(\Phi')$, we say that Φ is *equivalent* to Φ' and write $\Phi \sim \Phi'$. It is clear that \sim is indeed an equivalence relation.

Definition 3.6. Let $T_{\mathbb{F}}$ be the set of all frame templates over \mathbb{F} . We define a preorder \leq on $T_{\mathbb{F}}$ as follows. We say $\Phi \leq \Phi'$ if $\mathcal{M}_w(\Phi) \subseteq \mathcal{M}_w(\Phi')$. This is indeed a preorder since reflexivity and transitivity follow from the subset relation. We may obtain a partial order by considering equivalence classes of templates, with equivalence as defined above. However, the templates themselves, rather than equivalence classes, are the objects we work with in this paper.

Let Φ_0 be the frame template with all groups trivial and all sets empty. We call this template the *trivial template*. In general, we say that a template Φ is *trivial* if $\Phi \leq \Phi_0$. It is easy to see that for any template Φ , we have $\Phi_0 \leq \Phi$. Therefore, if $\Phi \leq \Phi_0$, then actually $\Phi \sim \Phi_0$.

Our desire is to find a collection of minimal nontrivial templates. For the remainder of this paper, we restrict our attention to binary frame templates: those frame templates where $\mathbb{F} = \mathrm{GF}(2)$ and Γ is the group of one element.

Definition 3.7.

- Let Φ_C be the template with all groups trivial and all sets empty except that |C| = 1 and $\Delta \cong \mathbb{Z}/2\mathbb{Z}$.
- Let Φ_X be the template with all groups trivial and all sets empty except that |X| = 1 and $\Lambda \cong \mathbb{Z}/2\mathbb{Z}$.
- Let Φ_{Y_0} be the template with all groups trivial and all sets empty except that $|Y_0| = 1$ and $\Delta \cong \mathbb{Z}/2\mathbb{Z}$.
- Let Φ_{CX} be the template with $Y_0 = Y_1 = \emptyset$, with |C| = |X| = 1, with $\Delta \cong \Lambda \cong \mathbb{Z}/2\mathbb{Z}$, with Γ trivial, and with $A_1 = [1]$.

• Let Φ_{Y_1} be the template with all groups trivial, with $C = Y_0 = \emptyset$, with $|Y_1| = 3$ and |X| = 2, and with $A_1 = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}$.

It is not too difficult to see that the Fano matroid F_7 virtually conforms to each of Φ_C , Φ_X , Φ_{CX} , Φ_{Y_0} , and Φ_{Y_1} . Therefore, these templates are nontrivial. In fact, one can see that $\mathcal{M}(\Phi_{Y_0})$ is the set of graft matroids, that $\mathcal{M}(\Phi_C)$ is the class of matroids obtained by closing the set of graft matroids under minors, and that $\mathcal{M}(\Phi_X)$ is the class of even-cycle matroids. In Lemma 4.5, we will show that $\mathcal{M}_v(\Phi_{Y_1})$ is the class of matroids having an even-cycle representation with a blocking pair.

Our goal in defining reductions and weak conforming was essentially to perform operations on matrices while leaving the Γ -frame submatrix intact. The following lemma does not contribute to that goal; so we will only make occasional use of it.

Lemma 3.8. The following relations hold:

(1) $\Phi_{Y_1} \preceq \Phi_X$ (2) $\Phi_{Y_1} \preceq \Phi_C$ (3) $\Phi_{Y_0} \preceq \Phi_C$ (4) $\Phi_C \preceq \Phi_{CX}$ (5) $\Phi_X \preceq \Phi_{CX}$

Proof. For (1), note that any simple matroid M of rank r virtually conforming to Φ_{Y_1} is a restriction of the vector matroid of a matrix A of the following form:

0	1	0	1	$1 \cdots 1$	$0 \cdots 0$	$1 \cdots 1$
0	0	1	1	$0 \cdots 0$	$1 \cdots 1$	$1 \cdots 1$
Γ-frame matrix		0		Ι	Ι	Ι

If we label the sets of rows and columns of A as B and E respectively, and the first row as x, then we see that A[B-x, E] is a Γ -frame matrix. If we let $X = \{x\}$, then we see that M conforms to Φ_X .

For (2), consider the matrix A above. Note that it is obtained by contracting c in the following matrix:

c

							C
	0	0	1	$0 \cdots 0$	$0 \cdots 0$	$1 \cdots 1$	1
0	1	0	0	$1 \cdots 1$	$0 \cdots 0$	$0 \cdots 0$	1
	0	1	0	$0 \cdots 0$	$1 \cdots 1$	$0 \cdots 0$	1
Γ-frame matrix		0		Ι	Ι	Ι	0

Removing c from this matrix, we obtain a Γ -frame matrix. Therefore, M conforms to Φ_C .

12

For (3), any matroid M conforming to Φ_{Y_0} is the vector matroid of a matrix of the following form, where v is an arbitrary column vector:



Let A be the matrix below. Label its sets of rows and columns as B and E respectively, and let c be the last column, with $C = \{c\}$.



Note that M is isomorphic to M(A)/C. Since A[B, E - C] is a Γ -frame matrix, we see that M conforms to Φ_C .

For (4), let A be a matrix conforming to Φ_C and let M = M(A)/Cbe the corresponding matroid conforming to Φ_C . If the column of A indexed by C is a zero column, then construct the matrix \overline{A} by adding a unit row, indexed by X, whose nonzero entry is in the column indexed by C. One readily sees that \overline{A} conforms to Φ_{CX} and that the corresponding matroid $M(\overline{A})/C$ is equal to M. Otherwise, if the column of A indexed by C has a nonzero entry, then one readily sees that A conforms to Φ_{CX} by considering the row containing the nonzero entry to be indexed by X.

For (5), any matroid M conforming to Φ_D is the vector matroid of a matrix of the following form, where v is an arbitrary row vector:



Consider the following matrix A, whose last column is indexed by $\{c\} = C$:

v	1
0	1
Γ-frame matrix	0

The matroid M is isomorphic to M(A)/c, which conforms to Φ_{CX} . \Box

		Z	$Y_0 Y_1$	C_0	C_1
X_0	columns from ΛX_0	0	۰	Ι	*
X_1	columns from ΛX_1	0	*	C)
	Γ-frame matrix	unit or zero columns	rows from	iΔ	

FIGURE 2. Standard form

Lemma 3.9. Let Φ be a template with $y \in Y_1$. Let Φ' be the template obtained from Φ by removing y from Y_1 and placing it in Y_0 . Then $\Phi' \preceq \Phi$.

Proof. Any matrix respecting Φ' virtually respects Φ by adding column y only to the zero Z column. Thus, any matroid conforming to Φ' weakly conforms to Φ .

We call the operation described in Lemma 3.9 a y-shift.

Definition 3.10. Let $\Phi = (\Gamma, C, X, Y_0, Y_1, A_1, \Delta, \Lambda)$ be a frame template over a finite field \mathbb{F} . We say that Φ is in *standard form* if there are disjoint sets C_0, C_1, X_0 , and X_1 such that $C = C_0 \cup C_1$, such that $X = X_0 \cup X_1$, such that $A_1[X_0, C_0]$ is an identity matrix, and such that $A_1[X_1, C]$ is a zero matrix.

Figure 2, with the stars representing arbitrary matrices, shows a matrix that virtually respects a template in standard form. Note that if Φ is in standard form, $|C_0| = |X_0|$. Also note that any of C_0, C_1, X_0 , or X_1 may be empty. Finally, note that we have defined standard form for frame templates over any finite field, not just binary frame templates.

Lemma 3.11. Every frame template $\Phi = (\Gamma, C, X, Y_0, Y_1, A_1, \Delta, \Lambda)$ is equivalent to a frame template in standard form.

Proof. Choose a basis C_0 for $M(A_1[X, C])$, and let $C_1 = C - C_0$. Repeatedly perform operation (5) to obtain a template Φ' where $A_1[X, C_0]$ consists of an identity matrix on top of a zero matrix. Each use of operation (5) results in an equivalent template; therefore, $\Phi \sim \Phi'$. Let $X_0 \subseteq X$ index the rows of the identity matrix, and let $X_1 \subseteq X$ index the rows of the zero matrix. Since C_0 is a basis for $M(A_1[X, C])$, the matrix $A_1[X, C_1]$ must be a zero matrix as well. Thus, Φ' is in standard form.

Throughout the rest of this paper, we will implicitly use Lemma 3.11 to assume that all templates are in standard form. Also, the operations (1)-(12) to which we will refer throughout the rest of this paper are the operations (1)-(8) from Proposition 3.2 and (9)-(12) from Definition 3.3.

Lemma 3.12. If $\Phi = (\{1\}, C, X, Y_0, Y_1, A_1, \Delta, \Lambda)$ is a binary frame template with $\Lambda | X_1$ nontrivial, then $\Phi_X \preceq \Phi$.

Proof. Perform operations (2) and (3) on Φ to obtain the following template, where λ is an element of Λ with $\lambda_x \neq 0$ for some $x \in X_1$:

$$(\{1\}, C, X, Y_0, Y_1, A_1, \{0\}, \{\mathbf{0}, \lambda\}).$$

On this template, repeatedly perform operation (7), then (8), then (4), and then (10) until the following template is obtained:

$$(\{1\}, \emptyset, X_1, \emptyset, \emptyset, [\emptyset], \{0\}, \{0, \lambda\}).$$

On this template, repeatedly perform operation (5) to obtain a template that is identical to the previous one except that the support of λ contains only one element of X_1 . On this template, repeatedly perform operation (6) to obtain the following template, where $x \in X_1$:

$$(\{1\}, \emptyset, \{x\}, \emptyset, \emptyset, [\emptyset], \{0\}, \mathbb{Z}/2\mathbb{Z}).$$

This template is Φ_X .

Lemma 3.13. If $\Phi = (\{1\}, C, X, Y_0, Y_1, A_1, \Delta, \Lambda)$ is a binary frame template, then either $\Phi_C \preceq \Phi$ or Φ is equivalent to a template with $C_1 = \emptyset$.

Proof. Suppose there is an element $\delta \in \Delta | C$ that is not in the row space of $A_1[X, C]$. Repeatedly perform operations (4) and (10) on Φ until the following template is obtained:

$$(\{1\}, C, X, \emptyset, \emptyset, A_1[X, C], \Delta | C, \Lambda).$$

On this template, perform operations (2) and (3) to obtain the following template:

 $(\{1\}, C, X, \emptyset, \emptyset, A_1[X, C], \{\mathbf{0}, \delta\}, \{0\}).$

Every matrix virtually respecting this template is row equivalent to a matrix virtually respecting a template that is identical to the previous template except that there is the additional condition that $\delta | C_0$ is a zero vector. Note that $\delta | C_1$ is nonzero since, in the previous template, δ was not in the row space of $A_1[X, C]$. Now, on the current template, repeatedly perform operation (7) and then operation (6) to obtain the following template:

$$\Phi' = (\{1\}, C_1, \emptyset, \emptyset, \emptyset, [\emptyset], \{\mathbf{0}, \delta | C_1\}, \{0\}).$$

Now, any matroid M conforming to Φ' is obtained by contracting C_1 from M(A), where A is a matrix conforming to Φ' . By contracting any single element $c \in C_1$, where $\delta_c = 1$, we turn the rest of the elements of C_1 into loops. So $C_1 - c$ is deleted to obtain M. Thus, M conforms to the template

$$(\{1\}, \{c\}, \emptyset, \emptyset, \emptyset, [\emptyset], \mathbb{Z}/2\mathbb{Z}, \{0\}),$$

which is Φ_C . Similarly, the converse is true that any matroid conforming to Φ_C conforms to Φ' . Thus, $\Phi_C \sim \Phi' \preceq \Phi$.

Now suppose that every element of $\Delta | C$ is in the row space of $A_1[X, C]$. Thus, contraction of C_0 turns the elements of C_1 into loops, and contraction of C_1 is the same as deletion of C_1 . By deleting C_1 from every matrix virtually conforming to Φ , we see that Φ is equivalent to a template with $C_1 = \emptyset$.

Lemma 3.14. If $\Phi = (\{1\}, C, X, Y_0, Y_1, A_1, \Delta, \Lambda)$ is a binary frame template, then one of the following is true:

- $\Phi_C \preceq \Phi$
- Φ is equivalent to a template with $\Lambda|X_1$ nontrivial and $\Phi_X \preceq \Phi$
- Φ is equivalent to a template with $\Lambda | X_0$ nontrivial and $\Phi_{CX} \preceq \Phi$
- Φ is equivalent to a template with Λ trivial and $C = \emptyset$.

Proof. By Lemmas 3.12 and 3.13, we may assume that $\Lambda|X_1$ is trivial and that $C_1 = \emptyset$.

First, suppose there exist elements $\delta \in \Delta | C_0$ and $\lambda \in \Lambda | X_0$ such that there are an odd number of natural numbers *i* with $\delta_i = \lambda_i = 1$. Thus, $\Lambda | X_0$ is nontrivial. Repeatedly perform operations (4) and (10) on Φ until the following template is obtained:

$$(\{1\}, C_0, X, \emptyset, \emptyset, A_1[X, C_0], \Delta | C_0, \Lambda).$$

On this template, repeatedly perform operation (6) to obtain the following template:

$$\Phi' = (\{1\}, C_0, X_0, \emptyset, \emptyset, A_1[X_0, C_0], \Delta | C_0, \Lambda | X_0).$$

Perform operations (2) and (3) on Φ' to obtain the following template:

 $(\{1\}, C_0, X_0, \emptyset, \emptyset, A_1[X_0, C_0], \{\mathbf{0}, \delta\}, \{\mathbf{0}, \lambda\}).$

Any matroid conforming to this template is obtained by contracting C_0 . If δ is in the row labeled by r and λ is in the column labeled by c, then when C_0 is contracted, 1 is added to the entry of the Γ -frame matrix in row r and column c. Otherwise, the entry remains unchanged when C is contracted. We see then that this template is equivalent to Φ_{CX} , where 1s are used to replace δ and λ .

Thus, we may assume that for every element $\delta \in \Delta | C_0$ and $\lambda \in \Lambda | X_0$, there are an even number of natural numbers *i* such that $\delta_i = \lambda_i = 1$. This implies that contraction of *C* has no effect on the Γ -frame matrix. So Φ is equivalent to a template with $\Lambda | X_0$ trivial. Therefore, since $\Lambda | X_1$ is trivial, we see that Λ is trivial. Note that operation (7) is a reduction that produces an equivalent template, since *C* must be contracted to produce a matroid that conforms to a template. By repeatedly performing operation (7), we obtain a template equivalent to Φ with $C = \emptyset$.

Lemma 3.15. If $\Phi = (\{1\}, C, X, Y_0, Y_1, A_1, \Delta, \Lambda)$ is a binary frame template with Λ trivial and with $C = \emptyset$, then either $\Phi_{Y_0} \preceq \Phi$ or Φ is equivalent to a template with Δ trivial.

Proof. First, suppose there is an element $\delta \in \Delta$ that is not in the row space of $A_1 = A_1[X_1, (Y_0 \cup Y_1)]$. Recall that a *y*-shift is the operation described in Lemma 3.9. Repeatedly perform *y*-shifts to obtain the following template, where $Y'_0 = Y_0 \cup Y_1$:

$$(\{1\}, \emptyset, X, Y'_0, \emptyset, A_1, \Delta, \{0\}).$$

On this template, perform operation (3) to obtain the following template:

 $(\{1\}, \emptyset, X, Y'_0, \emptyset, A_1, \{\mathbf{0}, \delta\}, \{0\}).$

Choose a basis B' for $M(A_1)$. By performing elementary row operations on every matrix virtually respecting Φ , we may assume that $A_1[X, B']$ consists of an identity matrix with zero rows below it and that $\delta|B'$ is the zero vector. By assumption, there is some element $y \in (Y'_0 - B')$ such that δ_y is nonzero. Thus, we can repeatedly perform operation (10) to obtain the following template:

$$(\{1\}, \emptyset, X, B' \cup y, \emptyset, A_1[X, B' \cup y], \{\mathbf{0}, \delta | (B' \cup y)\}, \{0\}).$$

Now, we can repeatedly perform operation (6) and then operation (12) to obtain the following template:

$$(\{1\}, \emptyset, \emptyset, \{y\}, \emptyset, [\emptyset], \mathbb{Z}/2\mathbb{Z}, \{0\}),$$

which is Φ_{Y_0} .

Now suppose that every element $\delta \in \Delta$ is in the row space of $A_1 = A_1[X, (Y_0 \cup Y_1)]$. Since Λ is trivial, by performing elementary row operations on every matrix virtually respecting Φ , we obtain a template equivalent to Φ with Δ trivial.

Lemma 3.16. Let $\Phi = (\{1\}, C, X, Y_0, Y_1, A_1, \Delta, \Lambda)$ be a binary frame template with Λ and Δ trivial. If $M(A_1[X_1, (Y_0 \cup Y_1)])$ has a circuit Y' with $|Y' \cap Y_1| \geq 3$, then $\Phi_{Y_1} \preceq \Phi$.

Proof. Any matroid conforming to Φ is obtained by contracting C. Since Λ and Δ are trivial, we may assume that $C = X_0 = \emptyset$ and therefore that $X = X_1$. Repeatedly perform operation (4) and then operation (10) on Φ to obtain the following template:

$$(\{1\}, \emptyset, X, Y_0 \cap Y', Y_1 \cap Y', A_1[X, Y'], \{0\}, \{0\}).$$

Choose any 3-element subset of $Y' \cap Y_1$ and call it Y''. Repeatedly perform y-shifts to obtain the following template:

$$(\{1\}, \emptyset, X, Y' - Y'', Y'', A_1[X, Y'], \{0\}, \{0\}).$$

On this template, repeatedly perform operation (11) to obtain the following template:

$$(\{1\}, \emptyset, X', \emptyset, Y'', A_1[X', Y''], \{0\}, \{0\}),$$

where X' is the subset of X that remains after Y' - Y'' is contracted. On this template, repeatedly perform operations (5) and (6) to obtain the following template, where X'' is a 2-element subset of X':

$$(\{1\}, \emptyset, X'', \emptyset, Y'', \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}, \{0\}, \{0\}).$$

This template is Φ_{Y_1} .

Lemma 3.17. If Φ is a frame template with Δ trivial, then Φ is equivalent to a template Φ' where $A_1[X, Y_1]$ is a matrix with every column nonzero and where no column is a copy of another. Moreover, if Φ is a binary frame template, then $M(A_1[X, Y_1])$ is simple.

Proof. Let A be a matrix that virtually conforms to Φ . Since Δ is trivial, the columns of A indexed by elements of Z are formed by placing a column of $A_1[X, Y_1]$ on top of a unit column or a zero column. These columns can be made using any copy of the same column of $A_1[X, Y_1]$; so only one copy is needed. If any column of $A_1[X, Y_1]$ is a zero column, then any column indexed by an element of Z that is made with this zero column can also be made as a column indexed by an element of $E - (Z \cup Y_0 \cup Y_1 \cup C)$ and choosing for the element of Λ the zero vector. Thus, no zero columns of $A_1[X, Y_1]$ are needed.

In the binary case, $M(A_1[X, Y_1])$ has no parallel elements because any such elements index copies of the same column. Also, $M(A_1[X, Y_1])$ has no loops because every column of $A_1[X, Y_1]$ is nonzero. Therefore, $M(A_1[X, Y_1])$ is simple.

Lemma 3.18. Let Φ be a binary frame template. Then at least one of the following is true:

(i) $\Phi_0 \sim \Phi$

18

- (ii) $\Phi' \preceq \Phi$ for some $\Phi' \in \{\Phi_X, \Phi_C, \Phi_{CX}, \Phi_{Y_0}, \Phi_{Y_1}\}$
- (iii) Φ is equivalent to a template where C = Ø, where Λ and Δ are trivial, and where A₁ is of the following form, with Y₀ = V₀∪V₁, with L an arbitrary binary matrix, and with each column of H containing at most two nonzero entries:

Y_1	V_0	V_1	
Ι	0	Η	
0	Ι	L	

Proof. Suppose neither (i) nor (ii) holds. By Lemma 3.14, we may assume that Λ is trivial and $C = \emptyset$. By Lemma 3.15, we may assume that Δ is trivial. By Lemma 3.16, every dependent set of M(A) = $M(A_1[X_1, (Y_0 \cup Y_1)])$ has an intersection with Y_1 with size at most 2. So by elementary row operations, we may assume that A_1 is of the following form, where $Y_0 = V_0 \cup V_1$, where L is an arbitrary binary matrix, where K consists of unit and zero columns, and where each column of H contains at most two nonzero entries:

J	Y_1	V_0	V_1	
Ι	K	0	Η	
0	0	Ι	L	

However, by Lemma 3.17, we may assume that K is an empty matrix. Thus, (iii) holds.

Theorem 3.19. Let Φ be a binary frame template. Then at least one of the following is true:

- (i) $\Phi_0 \sim \Phi$
- (ii) $\Phi' \preceq \Phi$ for some $\Phi' \in \{\Phi_X, \Phi_C, \Phi_{CX}, \Phi_{Y_0}, \Phi_{Y_1}\}$
- (iii) There exist $k, l \in \mathbb{Z}_+$ such that no simple, vertically k-connected matroid with at least l elements either virtually conforms or virtually coconforms to Φ .

Proof. Suppose for contradiction that none of outcomes (i)-(iii) hold for Φ . By Lemma 3.18, outcome (iii) of that lemma holds. Note that any simple matroid N virtually conforming to Φ is a restriction of a matroid M represented by a matrix of the following form, where $Z = Z_0 \cup Z_1$, where $Y_0 = V_0 \cup V_1$, and where the Γ -frame matrix has n rows and has a vector matroid isomorphic to the cycle matroid of the graph K_{n+1} :

			Z_0			Z_1	V_0	V_1
X	0	$1 \cdots 1$	$1 \cdots 1$		$1\cdots 1$	Ι	0	Η
			0			0	Ι	L
	Γ -frame matrix	Ι	Ι	•••	Ι	0	0	0

Also recall from the definition of conforming to a template that $Y_0 \subseteq E(N)$.

We see that

$$\lambda_N(Y_0 \cup (Z_1 \cap E(N))) \le \lambda_M(Y_0 \cup Z_1)$$

= $r_M(Y_0 \cup Z_1) + r_M(E - (Y_0 \cup Z_1)) - r(M)$
= $|V_0| + |Y_1| + |Y_1| + n - (|Y_1| + |V_0| + n)$
= $|Y_1|.$

Note that each column of the above matrix, except possibly those columns indexed by V_1 , has at most two nonzero entries. Thus, Mis graphic and Φ is trivial if $V_0 = \emptyset$. Since (i) does not hold, Φ is nontrivial. Therefore, $V_0 \neq \emptyset$, and $E(N) - (Y_0 \cup Z_1)$ is not spanning. Thus, if $k > |Y_1| + 1$, then N is not vertically k-connected unless $Y_0 \cup (Z_1 \cap E(N))$ is spanning in N. This implies that n = 0; in that case, N is only simple if the Γ -frame matrix is a 0×0 matrix. This implies that $|E(N)| \leq |Y_0 \cup Y_1|$. So if $l > |Y_0 \cup Y_1|$, then no simple, vertically k-connected matroid with at least l elements virtually conforms to Φ .

Now, consider a simple matroid N^* which virtually coconforms to Φ . Then N is a restriction of M with $Y_0 \subseteq E(N)$. Since a matroid and its dual have the same connectivity function, we have $\lambda_{N^*}(Y_0 \cup (Z_1 \cap E(N)) \leq |Y_1|$. So if $k > |Y_1| + 1$, then N^* is not vertically k-connected unless either $Y_0 \cup (Z_1 \cap E(N))$ or $E(N) - (Z_1 \cup Y_0)$ is spanning in N^* , implying that either $E(N) - (Z_1 \cup Y_0)$ or $Y_0 \cup (Z_1 \cap E(N))$ is independent in N. If $E(N) - (Z_1 \cup Y_0)$ is independent in N, then

$$|E(N) - (Z_1 \cup Y_0)| = r_N(E(N) - (Z_1 \cup Y_0))$$

$$\leq r_M(E(M) - (Z_1 \cup Y_0))$$

$$= |Y_1| + n.$$

By the formula for corank, we have

$$r_{N^*}(E(N) - (Z_1 \cup Y_0)) \le r_{M^*}(E(N) - (Z_1 \cup Y_0))$$

= $|E(N) - (Z_1 \cup Y_0)| + r_M(Z_1 \cup Y_0) - r(M)$
 $\le |Y_1| + n + |Y_1| + |V_0| - (|Y_1| + |V_0| + n)$
= $|Y_1|.$

Since N^* is simple and binary, we have $|E(N) - (Z_1 \cup Y_0)| \le 2^{|Y_1|} - 1$. This implies that $|E(N)| \le 2^{|Y_1|} - 1 + |Y_1| + |Y_0|$. Thus, if we set l greater than this value, then no simple, vertically k-connected matroid with at least l elements virtually coconforms to Φ unless $Y_0 \cup (Z_1 \cap E(N))$ is independent in N. Since (iii) does not hold, this must be true for some matroid N. In particular, $Y_0 = V_0 \cup V_1$ is independent in N, implying that H is a linearly independent matrix.

Let P denote the matrix

$$P = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 1 \\ \hline 1 & 1 \end{bmatrix}.$$

Suppose $A_1[X, V_1]$ has P as a submatrix, with the first three rows of P contained in H and the last row of P contained in L. Then A_1 contains the following submatrix, with the first three columns contained in $A_1[X, Y_1]$ and the last two contained in $A_1[X, V_1]$:

1	0	0	1	0 -	
0	1	0	0	1	
0	0	1	0	1	•
0	0	0	1	1	

After contracting all other elements of Y_1 by repeatedly performing *y*-shifts and operation (12), the columns of this submatrix form a circuit in $M(A_1)$ whose intersection with Y_1 has size 3. However, we have already deduced by Lemma 3.16 that this is impossible. Therefore, A_1 does not contain P as a submatrix, with the first three rows of P contained in H and the last row of P contained in L. We will refer to this fact by saying that A_1 has no P-configuration.

Let $\{1, 2, \ldots, m\}$ be the rows of L. (So $|V_0| = m$.) Let S_i be the submatrix of H obtained by restricting H to the columns j such that $L_{i,j} = 1$. Recall that H, and therefore S_i , contain at most two nonzero entries per column. Also, since H is linearly independent, each column has at least one nonzero entry, and no column is a copy of another. Suppose a column e of S_i contains exactly two nonzero entries. Since

 A_1 has no *P*-configuration, all other columns of S_i must contain a nonzero entry in exactly one of the same rows as *e*. Suppose that there are columns *f* and *g* in S_i such that *f* contains a nonzero entry in one of the same rows as *e*, but *g* contains a nonzero entry in the other row. Then S_i contains the following submatrix:

$$\begin{array}{ccc} e & f & g \\ \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} .$$

Since H is a linearly independent matrix, f or g (say f) must have an additional nonzero entry in H. To avoid f and g forming a Pconfiguration, g must have an additional nonzero entry in the same row as f. Therefore, S_i contains the following submatrix:

$$\begin{array}{cccc} e & f & g \\ \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}$$

Since each column of H contains at most two nonzero entries, $\{e, f, g\}$ is a dependent set of columns, contradicting the assumption that H is linearly independent.

Therefore, we deduce that each S_i either consists entirely of unit columns or contains a row s_i consisting entirely of 1s. Note that each S_i is the incidence matrix of a star, with possibly one row removed. We will call s_i the *star center* of row *i*. If S_i consists entirely of unit columns, then we define its star center to be $s_i = \emptyset$.

If the sets of columns of all the S_i are pairwise disjoint, then by adding each row *i* to its star center s_i , we see that every matroid virtually conforming to Φ can be represented by a matrix with at most two nonzero entries per column. Thus, Φ is trivial, contradicting the assumption that (i) does not hold. Also, if *i* and *j* are distinct rows of *L* with distinct star centers s_i and s_j , then S_i and S_j can have at most one column in common because otherwise, the columns they have in common form a linearly dependent set in *H*.

Now suppose there are S_i and S_j with $s_i = s_j$. Also, suppose that neither S_i nor S_j is a submatrix of the other. Then A_1 contains the following submatrix. In fact, after repeatedly performing y-shifts, operation (11), and operation (10), we may assume that A_1 is the following matrix, with the first three columns indexed by Y_1 , the next two indexed by V_0 , and the last three by V_1 :

Γ	1	0	0	0	0	1	1	1	
	0	1	0	0	0	0	1	0	
	0	0	1	0	0	0	0	1	.
ľ	0	0	0	1	0	1	1	0	
	0	0	0	0	1	0	1	1	

Add the fourth row to the first, and swap the fourth and sixth columns to obtain the following matrix:

1	0	0	0	0	1	0	1
0	1	0	0	0	0	1	0
0	0	1	0	0	0	0	1
0	0	0	1	0	1	1	0
0	0	0	0	1	0	1	1

The last two columns of this matrix contain a *P*-configuration.

Now suppose there are matrices S_i and S_j so that S_j is a submatrix of S_i . Then A_1 contains a submatrix obtained by deleting columns from a matrix of the following form, where the left portion comes from the set V_0 , the upper-right portion comes from the matrix H, the lower-left portion comes from the matrix L, and x is 1 or 0 depending on whether or not the last column is contained in S_i :



Choose any column contained in S_j and perform row operations so that this column becomes a unit column with nonzero entry in L. Then we obtain the following matrix:

0	1	1	•••	1	0	0	• • •	0	x+1
0	0	1							0
:	÷		۰.						÷
0	0			1					0
0	1				0	1	•••	1	x
0	0					1			0
:	÷						·		÷
0	0							1	0
1	1	1	•••	1	0	•••	•••	0	x+1
0	1	0	•••	0	1	• • •	•••	1	x

Now, by swapping the appropriate columns, we obtain the following:

Γ	0	0	1	•••	1	1	0	•••	0	x+1
(0	0	1							0
	:	÷		۰.						÷
(0	0			1					0
(0	0				1	1	•••	1	x
(0	0					1			0
	:	÷						·		÷
(0	0							1	0
	1	0	1	•••	1	1	0	•••	0	x+1
	0	1	0	• • •	0	1	• • •	•••	1	x

We see that in this new matrix, S_i and S_j have only one column in common and $s_i \neq s_j$. The last column is in S_i if x = 0 and S_j if x = 1. Thus, this case reduces to the final case that remains to be checked: for all *i* and *j*, we have $s_i \neq s_j$ and S_i and S_j have at most one column in common. Since each column of *H* contains at most two nonzero entries, and since all S_i have distinct star centers, we see that a column of *H* can be contained in at most two S_i . By adding each row *i* to its star center s_i , one can see that every matrix virtually conforming to Φ can be rewritten so that every column contains at most two nonzero entries. Therefore, Φ is trivial, and (i) holds.

This completes the contradiction and proves the result.

Outcome (iii) of Theorem 3.19 only occurs in very specific situations. In fact, due to connectivity considerations, it is not needed in order to use Corollary 3.5. **Definition 3.20.** Let \mathcal{M} be a minor-closed class of binary matroids, and suppose there exist $k, l, m \in \mathbb{Z}_+$ and a set $\mathcal{T}_{\mathcal{M}} = \{\Phi_1, \ldots, \Phi_s, \Psi_1, \ldots, \Psi_t\}$ of binary frame templates such that

- \mathcal{M} contains each of the classes $\mathcal{M}_w(\Phi_1), \ldots, \mathcal{M}_w(\Phi_s),$
- \mathcal{M} contains the duals of the matroids in each of the classes $\mathcal{M}_w(\Psi_1), \ldots, \mathcal{M}_w(\Psi_t),$
- if M is a simple vertically k-connected member of \mathcal{M} with at least l elements and with no PG(m-1,2) minor, then either Mis a member of at least one of the classes $\mathcal{M}_v(\Phi_1), \ldots, \mathcal{M}_v(\Phi_s)$ or M^* is a member of at least one of the classes $\mathcal{M}_v(\Psi_1), \ldots, \mathcal{M}_v(\Psi_t)$, and
- for each template $\Phi \in \mathcal{T}_{\mathcal{M}}$, either Φ is trivial or $\Phi' \preceq \Phi$ for some $\Phi' \in \{\Phi_X, \Phi_C, \Phi_{CX}, \Phi_{Y_0}, \Phi_{Y_1}\}.$

We say that $\mathcal{T}_{\mathcal{M}}$ describes \mathcal{M} .

By combining Corollary 3.5 with Theorem 3.19, one can observe that every proper minor-closed class \mathcal{M} of binary matroids can be described by a set of templates. Moreover, that set is nonempty if and only if \mathcal{M} contains all graphic matroids or all cographic matroids.

Corollary 3.21. Let \mathcal{M} be a minor-closed class of binary matroids, and let $\{\Phi_1, \ldots, \Phi_s, \Psi_1, \ldots, \Psi_t\}$ be a set of templates describing \mathcal{M} . If any of these templates is nontrivial, then \mathcal{M} contains $\mathcal{M}(\Phi_{Y_0})$, $\mathcal{M}(\Phi_{Y_1}), \mathcal{M}^*(\Phi_{Y_0}), \text{ or } \mathcal{M}^*(\Phi_{Y_1}).$

Proof. Let Φ be a nontrivial template in the set $\{\Phi_1, \ldots, \Phi_s\}$. By Definition 3.20 and Lemma 3.8, either $\Phi_{Y_0} \preceq \Phi$ or $\Phi_{Y_1} \preceq \Phi$. If $\Phi_{Y_0} \preceq \Phi$, then

$$\mathcal{M}(\Phi_{Y_0}) \subseteq \mathcal{M}_v(\Phi_{Y_0}) \subseteq \mathcal{M}_v(\Phi) \subseteq \mathcal{M},$$

where the first containment holds because every matroid conforming to a template also virtually conforms to it, the second containment holds by definition of \leq , and the third containment holds by Definition 3.20. In the case where $\Phi_{Y_1} \leq \Phi$, a similar argument shows that $\mathcal{M}(\Phi_{Y_1}) \subseteq \mathcal{M}$.

If Ψ is a nontrivial template in the set $\{\Psi_1, \ldots, \Psi_s\}$, a similar argument shows that either $\mathcal{M}^*(\Phi_{Y_0}) \subseteq \mathcal{M}$, or $\mathcal{M}^*(\Phi_{Y_1}) \subseteq \mathcal{M}$. \Box

4. Growth Rates

Let \mathcal{M} be a minor-closed class of matroids. Let $h_{\mathcal{M}}(r)$ denote the growth rate function of \mathcal{M} : the function whose value at an integer $r \geq 0$ is given by the maximum number of elements in a simple matroid in \mathcal{M} of rank at most r. For a matroid M, we denote by $\varepsilon(M)$ the size

of the simplification of M, that is the number of rank-1 flats of M. By combining the main result in [4] with earlier results of Geelen and Whittle [3] and Geelen and Kabell [2], Geelen, Kung, and Whittle proved the following:

Theorem 4.1 (Growth Rate Theorem). If \mathcal{M} is a nonempty minorclosed class of matroids, then there exists $c \in \mathbb{R}$ such that either:

- (1) $h_{\mathcal{M}}(r) \leq cr \text{ for all } r,$ (2) $\binom{r+1}{2} \leq h_{\mathcal{M}}(r) \leq cr^2 \text{ for all } r \text{ and } \mathcal{M} \text{ contains all graphic}$ matroids,
- (3) there is a prime-power q such that $\frac{q^r-1}{q-1} \leq h_{\mathcal{M}}(r) \leq cq^r$ for all r and \mathcal{M} contains all GF(q)-representable matroids, or
- (4) $h_{\mathcal{M}}$ is infinite and \mathcal{M} contains all simple rank-2 matroids.

If outcome (2) of the Growth Rate Theorem holds for a minor-closed class \mathcal{M} , then \mathcal{M} is said to be *quadratically dense*. In this section, we will consider growth rates of some quadratically dense classes of binary matroids. Let $\mathcal{EX}(F)$ denote the class of binary matroids with no Fminor. If f and g are functions, we write $f(r) \approx g(r)$ if f(r) = g(r) for all but finitely many r.

Since the growth rate function for the class of graphic matroids is $\binom{r+1}{2}$, the Growth Rate Theorem implies that, if F is a nongraphic binary matroid,

$$h_{\mathcal{EX}(F)}(r) \ge \binom{r+1}{2}.$$

Kung et. al. [6] pose the following question: For which nongraphic binary matroids F of rank 4 does equality hold above for all but finitely many r? Geelen and Nelson answer this question in [5]. Let N_{12} be the matroid formed by deleting a three-element independent set from PG(3,2). The nongraphic binary matroids F of rank 4 for which $h_{\mathcal{EX}(F)}(r) \approx {r+1 \choose 2}$ are exactly the nongraphic restrictions of N_{12} . We present here an alternate proof. Both proofs allow us to answer the question when F is a matroid of any rank, not just rank 4. We will prove the following theorem after proving several lemmas.

Theorem 4.2. Let \mathcal{M} be a minor-closed class of binary matroids. Then $h_{\mathcal{M}}(r) \approx \binom{r+1}{2}$ if and only if \mathcal{M} contains all graphic matroids but does not contain $\mathcal{M}_v(\Phi_{Y_1})$.

Our proof of Theorem 4.2 will depend on the following theorem, proved by Geelen and Nelson in [5]:

26

Theorem 4.3. Let \mathcal{M} be a quadratically dense minor-closed class of matroids and let p(x) be a real quadratic polynomial with positive leading coefficient. If $h_{\mathcal{M}}(n) > p(n)$ for infinitely many $n \in \mathbb{Z}^+$, then for all integers $r, s \geq 1$ there exists a vertically s-connected matroid $M \in \mathcal{M}$ satisfying $\varepsilon(M) > p(r(M))$ and $r(M) \geq r$.

An even-cycle matroid is a binary matroid of the form $M = M {w \choose D}$, where $D \in GF(2)^{V \times E}$ is the vertex-edge incidence matrix of a graph G = (V, E) and $w \in GF(2)^E$ is the characteristic vector of a set $W \subseteq E$. The pair (G, W) is an *even-cycle representation* of M. The edges in W are called *odd* edges, and the other edges are *even* edges. An *odd cycle* of (G, W) is a cycle of G with an odd number of odd edges. A blocking *pair* of (G, W) is a pair of vertices u, v of G so that every odd cycle passes through at least one of these vertices. Resigning at a vertex uof G occurs when all the edges incident with u are changed from even to odd and vice-versa. It is easy to see that this corresponds to adding the row of the matrix corresponding to u to the characteristic vector of W. Therefore, resigning at a vertex does not change an even-cycle matroid. It is also easy to see that if an even-cycle representation has a blocking pair, then we can resign so that every odd edge is incident with at least one vertex in the blocking pair. For our purposes, it will be convenient to think of a blocking pair in this way.

For $r \geq 2$, let A_r be the following binary matrix, where we choose for the Γ -frame matrix the matrix representation of $M(K_{r-1})$, so that the identity matrices are $(r-2) \times (r-2)$ matrices.

0	$\begin{array}{c} 1 \\ 0 \end{array}$	0 1	1 1	$\begin{array}{c} 1 \cdots 1 \\ 0 \cdots 0 \end{array}$	$\begin{array}{c} 0 \cdots 0 \\ 1 \cdots 1 \end{array}$	$\begin{array}{c} 1 \cdots 1 \\ 1 \cdots 1 \end{array}$
Γ -frame matrix		0		Ι	Ι	Ι

Note that $M(A_r)$ is the largest simple matroid of rank r that virtually conforms to Φ_{Y_1} .

Definition 4.4. Let X_r be the largest simple matroid of rank r that virtually conforms to Φ_{Y_1} . Equivalently, $X_1 = U_{1,1}$, and for $r \ge 2$, we have $X_r = M(A_r)$.

Lemma 4.5. The class $\mathcal{M}_v(\Phi_{Y_1})$ is the class of matroids having an even-cycle representation with a blocking pair. This class is minor-closed.

Proof. Any simple matroid M virtually conforming to Φ_{Y_1} is a restriction of X_r for some r.

Label the rows of A_r as $1, \ldots, r$. Add to the matrix row r+1, which is the sum of rows $2, \ldots, r$. This does not change the matroid X_r . We

see that X_r is an even-cycle matroid (G, W), where row 1 is the characteristic vector of W and rows $2, \ldots, r+1$ form the incidence matrix of G. Moreover, every edge in W is incident with the vertex corresponding to either row 2 or row r + 1. Thus, every matroid virtually conforming to Φ_{Y_1} has an even-cycle representation with a blocking pair. Conversely, every matroid that has an even-cycle representation with a blocking pair $\{u, v\}$ virtually conforms to Φ_{Y_1} , by making ucorrespond to the second row and making v correspond to row r + 1, which can be removed without changing the matroid.

By resigning whenever we wish to contract an element represented by an odd edge, it is not difficult to see that the class of matroids having an even-cycle representation with a blocking pair is minor-closed. \Box

Lemma 4.6. Any simple, rank-r matroid M that is a minor of a matroid virtually conforming to Φ_{Y_1} is a restriction of X_r .

Proof. From the preceding lemma, M is a restriction of some $X_{r'}$. So M has an even-cycle representation (G, W) with a blocking pair $\{u, v\}$. Let w be the characteristic vector of W. There are r' - r rows in the matrix $A_{r'}[(V \cup w) - v, E(M)]$ whose deletion does not alter the matroid M. After these rows are deleted, the resulting matrix is a submatrix of A_r .

Lemma 4.7. Every matroid virtually conforming to Φ_{Y_1} is a minor of a matroid conforming to Φ_{Y_0} .

Proof. By Lemma 3.8, we have $\Phi_{Y_1} \preceq \Phi_C$. Every matroid conforming to Φ_C is obtained by contracting an element from a matroid conforming to Φ_{Y_0} .

Lemma 4.8. Let k be a positive integer. Then there are at most finitely many integers r such that the complete graphic matroid $M(K_{r+1})$ is a rank- $(\leq k)$ perturbation of a cographic matroid.

Proof. Let N be a cographic matroid. Observe that adding a rank-1 matrix to a matrix representation of a binary matroid N changes $\varepsilon(N)$ by a factor of at most 2. This occurs when, in every rank-1 flat of N, there is at least one nonloop element indexing a column that is changed by adding the rank-1 matrix and at least one nonloop element indexing a column that remains unchanged when the rank-1 matrix is added. Thus, if M is a rank-($\leq t$) perturbation of N, we have $\varepsilon(M) \leq 2^t \varepsilon(N)$.

Let r = r(M). Recall that a cographic matroid N has $\varepsilon(N) \leq 3r(N) - 3$. Therefore, $\varepsilon(M) \leq 2^t(3r(N) - 3) \leq 2^t(3(r+t) - 3)$. For fixed t and sufficiently large r, this expression is less than $\binom{r+1}{2} = \varepsilon(M(K_{r+1}))$.

Lemma 4.9. Let \mathcal{M} be a quadratically dense minor-closed class of matroids representable over a given field \mathbb{F} . Let $\{\Phi_1, \ldots, \Phi_s, \Psi_1, \ldots, \Psi_t\}$ be a set of templates describing \mathcal{M} . For sufficiently large r, the growth rate $h_{\mathcal{M}}(r)$ is equal to the size of the largest simple matroid of rank r that virtually conforms to any template in $\{\Phi_1, \ldots, \Phi_s\}$.

Proof. Let $h'_{\mathcal{M}}(r)$ denote the size of the largest simple matroid of rank r that virtually conforms to any template in $\{\Phi_1, \ldots, \Phi_s\}$. So $h_{\mathcal{M}}(r) \geq h'_{\mathcal{M}}(r)$. The size of the largest simple matroid of rank r that virtually conforms to any particular template is a quadratic polynomial in r. Thus, for sufficiently large r, the function $h'_{\mathcal{M}}(r)$ is a quadratic polynomial as well.

By Definition 3.20, there exist $k, l \in \mathbb{Z}_+$ so that every simple vertically k-connected member of \mathcal{M} with at least l elements either weakly conforms to a template in $\{\Phi_1, \ldots, \Phi_s\}$ or weakly coconforms to some template in $\{\Psi_1, \ldots, \Psi_t\}$. Suppose, for contradiction, that $h_{\mathcal{M}}(r) >$ $h'_{\mathcal{M}}(r)$ for infinitely many r. Theorem 4.3, with $h'_{\mathcal{M}}(r)$ playing the role of p(r), implies that there is a sequence M_1, M_2, \ldots of vertically k-connected matroids in \mathcal{M} such that $\varepsilon(M_i) > h'_{\mathcal{M}}(i)$ and $r(M_i) \ge i$. Thus, in this sequence, there are infinitely many matroids that are vertically k-connected and have size at least l. Since these matroids are too large to virtually conform to any template in $\{\Phi_1, \ldots, \Phi_t\}$, there is at least one nontrivial template $\Psi \in \{\Psi_1, \ldots, \Psi_t\}$ such that infinitely many vertically k-connected matroids in \mathcal{M} coconform to Ψ . However, since \mathcal{M} contains all graphic matroids and since every complete graphic matroid has infinite vertical connectivity (hence vertical k-connectivity), we have that infinitely many complete graphic matroids coconform to Ψ . For some t depending on Ψ , every matroid coconforming to Ψ is a rank-($\leq t$) perturbation of a cographic matroid. This contradicts Lemma 4.8. By contradiction, the result holds.

Proof of Theorem 4.2. First, suppose $h_{\mathcal{M}}(r) \approx \binom{r+1}{2}$. By the Growth Rate Theorem, \mathcal{M} contains all graphic matroids. For $r \geq 1$, we have $|X_r| = \binom{r-1}{2} + 3r - 3$, which for r > 2 is greater than $\binom{r+1}{2}$. Thus, \mathcal{M} does not contain $\mathcal{M}_v(\Phi_{Y_1})$.

Now, suppose \mathcal{M} contains all graphic matroids but does not contain $\mathcal{M}_v(\Phi_{Y_1})$. Since \mathcal{M} contains all graphic matroids, there is a nonempty set $\{\Phi_1, \ldots, \Phi_s, \Psi_1, \ldots, \Psi_t\}$ of binary frame templates describing \mathcal{M} . By Lemma 4.9, $h_{\mathcal{M}}(r)$ is equal to the size of the largest simple matroid of rank r that conforms to any template in $\{\Phi_1, \ldots, \Phi_s\}$. Suppose Φ is a nontrivial template in $\{\Phi_1, \ldots, \Phi_s\}$. By Corollary 3.21, either $\Phi_{Y_0} \preceq \Phi$ or $\Phi_{Y_1} \preceq \Phi$. Since \mathcal{M} does not contain $\mathcal{M}_v(\Phi_{Y_1})$, we must

have $\Phi_{Y_0} \leq \Phi$. However, by Lemma 4.7, this implies $\mathcal{M}_v(\Phi_{Y_1}) \subseteq \mathcal{M}$. Therefore, we conclude that $h_{\mathcal{M}}(r) \approx \binom{r+1}{2}$, completing the proof. \Box

Corollary 4.10. Let F be a simple, binary matroid of rank r. Then $h_{\mathcal{EX}(F)} \approx \binom{r+1}{2}$ if and only if F is a nongraphic restriction of X_r .

Proof. By Theorem 4.2, $h_{\mathcal{EX}(F)} \approx {\binom{r+1}{2}}$ if and only if $\mathcal{EX}(F)$ contains all graphic matroids but does not contain $\mathcal{M}_v(\Phi_{Y_1})$. The condition that $\mathcal{EX}(F)$ contains all graphic matroids is equivalent to the condition that F is nongraphic. By Lemma 4.6, the condition that $\mathcal{EX}(F)$ does not contain $\mathcal{M}_v(\Phi_{Y_1})$ is equivalent to the condition that F is a restriction of X_r .

Note that $X_4 = N_{12}$; so this answers the question posed in [6].

We now consider the growth rate of $\mathcal{EX}(PG(3,2))$. We will prove Theorem 1.3, which we restate below.

Theorem 4.11. The growth rate function for $\mathcal{EX}(PG(3,2))$ is

$$h_{\mathcal{EX}(PG(3,2))} \approx r^2 - r + 1.$$

We will use the following.

Lemma 4.12. Let $\mathcal{T}_{\mathcal{EX}(PG(3,2))} = \{\Phi_1, \dots, \Phi_s, \Psi_1, \dots, \Psi_t\}$. If $\Phi \in \{\Phi_1, \dots, \Phi_s\}$, then either $\Phi = \Phi_X$ or Φ is a template with $C = \emptyset$ and with Λ and Δ trivial.

Proof. The class of matroids conforming to Φ_X is exactly the class of even-cycle matroids. This class is minor-closed. The largest simple, even-cycle matroid of rank r has an even-cycle representation obtained from the graph K_r by adding to each even edge an odd edge in parallel as well as adding one odd loop to the graph. Therefore, the class of even-cycle matroids has growth rate $2\binom{r}{2} + 1 = r^2 - r + 1$. So the largest simple, even-cycle matroid of rank 4 has size 13. Since PG(3, 2) has size 15, we have $\mathcal{M}(\Phi_X) \subseteq \mathcal{EX}(PG(3, 2))$. Therefore, we may assume that $\Phi_X \in \mathcal{T}_{\mathcal{EX}(PG(3,2))}$.

Since $\Phi_0 \leq \Phi_X$, we may assume that $\Phi_0 \notin \{\Phi_1, \dots, \Phi_s\}$. Let

$$\Phi = (\{1\}, C, X, Y_0, Y_1, A_1, \Delta, \Lambda)$$

be a nontrivial template such that $\Phi \neq \Phi_X$ and $\Phi \in {\Phi_1, \ldots, \Phi_s}$. Consider the graft matroid $M(K_6, V(K_6))$. A straightforward computation shows that, by contracting the nongraphic element, we obtain PG(3,2). Therefore, $\Phi_{Y_0} \not\preceq \Phi$. By Lemma 3.8, we also have $\Phi_C \not\preceq \Phi$ and $\Phi_{CX} \not\preceq \Phi$.

Now, we may assume that Φ is in standard form. Since $\Phi_C \not\preceq \Phi$, by Lemma 3.13 we may assume that $C_1 = \emptyset$. Also, by Lemma 3.14, since

 $\Phi_{CX} \not\preceq \Phi$ and $\Phi_C \not\preceq \Phi$, either $\Lambda | X_1$ is nontrivial and $\Phi_X \preceq \Phi$ or Λ is trivial and $C = \emptyset$.

First, suppose that Λ is trivial and $C = \emptyset$. Since $\Phi_{Y_0} \not\preceq \Phi$, Lemma 3.15 implies that Φ is equivalent to a template with Δ trivial. So we may assume

$$\Phi = (\{1\}, \emptyset, X, Y_0, Y_1, A_1, \{0\}, \{0\}),$$

which is one of the possible conclusions of the lemma.

Thus, we may assume that $\Lambda | X_1$ is nontrivial and $\Phi_X \preceq \Phi$. Suppose $|\Lambda | X_1 | > 2$. On the template

$$\Phi = (\{1\}, C_0, Y_0, Y_1, A_1, \Delta, \Lambda),$$

perform operation (3) and then repeatedly perform operations (4) and (10) to obtain the template

$$(\{1\}, C_0, X, \emptyset, \emptyset, A_1[X, C_0], \{0\}, \Lambda).$$

Then repeatedly perform operation (7) to obtain

$$(\{1\}, \emptyset, X_1, \emptyset, \emptyset, [\emptyset], \{0\}, \Lambda | X_1).$$

Since $\Lambda | X_1$ has characteristic 2 and size greater than 2, it contains a subgroup Λ' isomorphic to $(\mathbb{Z}/2\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z})$. Perform operation (2) to obtain the template

$$(\{1\}, \emptyset, X_1, \emptyset, \emptyset, [\emptyset], \{0\}, \Lambda');$$

then repeatedly perform operations (5) and (6) to obtain

$$(\{1\}, \emptyset, X', \emptyset, \emptyset, [\emptyset], \{0\}, \Lambda''),$$

where |X'| = 2 and Λ'' is the additive group generated by $\begin{bmatrix} 1\\ 0 \end{bmatrix}$ and $\begin{bmatrix} 0\\ 1 \end{bmatrix}$. One readily sees that PG(3,2) conforms to this template. Therefore, $|\Lambda| = 2$. We may perform row operations so that Λ is generated by $[1, 0..., 0]^T$. Let Σ be the element of X such that $\Lambda | \{\Sigma\}$ is nonzero.

Now, suppose there is an element $\bar{x} \in \Delta$ that is not in the row space of A_1 . Perform operations (2) and (3) on Φ to obtain

$$(\{1\}, C_0, X, Y_0, Y_1, A_1, \{0, \bar{x}\}, \{0\}).$$

Now, by a similar argument to the one used in the proof of Lemma 3.15, we have $\Phi_{Y_0} \preceq \Phi$. Since we already know this is not the case, we deduce that every element of Δ is in the row space of A_1 .

Let $\bar{x} \in \Delta | C_0$ and $\bar{y} \in \Lambda$ be such that there are an odd number of natural numbers *i* such that $\bar{x}_i = \bar{y}_i = 1$. Then we call the ordered pair (\bar{x}, \bar{y}) a pair of odd type. Otherwise, (\bar{x}, \bar{y}) is a pair of even type.

Suppose (\bar{x}, \bar{y}) is a pair of odd type with $\bar{y}|X_1$ a zero vector. By performing operations (2) and (3) and repeatedly performing operations (4) and (10), we obtain

$$(\{1\}, C_0, X, \emptyset, \emptyset, A_1[X, C], \{0, \bar{x}\}, \{0, \bar{y}\}),$$

which is equivalent to Φ_{CX} . We already know this is not the case. Therefore, for every pair (\bar{x}, \bar{y}) of odd type, $\bar{y}|X_1 = [1, 0, \dots, 0]^T$.

Suppose $\bar{x} \in \Delta | C$ and $\bar{y}_1, \bar{y}_2 \in \Lambda$ are such that $\bar{y}_1 | X_1 = \bar{y}_2 | X_1 = [1, 0, \dots, 0]^T$, such that (\bar{x}, \bar{y}_1) is a pair of odd type, and such that (\bar{x}, \bar{y}_2) is a pair of even type. Then $(\bar{y}_1 + \bar{y}_2) | X_1$ is a zero vector, and $(\bar{x}, \bar{y}_1 + \bar{y}_2)$ is a pair of odd type. Therefore, either all pairs $(\bar{x}, \bar{y}) \in \Delta | C \times \Lambda$ are of even type, in which case Φ is equivalent to a template with $\Lambda | X_0$ trivial and $C = \emptyset$, or if (\bar{x}, \bar{y}) is a pair of odd type, then (\bar{x}, \bar{z}) is of odd type for every $\bar{z} \in \Lambda$ with $\bar{z} | X_1$ nonzero. In this case, consider any matrix virtually conforming to Φ . After contracting C, we can restore the Γ -frame matrix by adding Σ to each row where the Γ -frame matrix has been altered. Therefore, Φ is equivalent to a template with $\Lambda | X_0$ trivial and $C = \emptyset$.

So we now have that

 $\Phi = (\{1\}, \emptyset, X, Y_0, Y_1, A_1, \Delta, \Lambda),$

with Λ generated by $[1, 0, ..., 0]^T$ and with every element of Δ in the row space of A_1 . We will now show that, in fact, Φ is equivalent to a template with Δ trivial. On Φ , perform *y*-shifts to obtain the following template, where $Y'_0 = Y_0 \cup Y_1$:

$$\Phi' = (\{1\}, \emptyset, X, Y'_0, \emptyset, A_1, \Delta, \Lambda).$$

By repeatedly performing operation (5) and then operation (6) on this template, we may assume that A_1 has the following form, with the star representing an arbitrary binary matrix and \bar{v} representing an arbitrary row vector:

$$\begin{bmatrix} 0\cdots 0 & \bar{v} \\ \hline I_{|X|-1} & * \end{bmatrix}.$$

Also, since $\Lambda|(D - \{\Sigma\})$ is trivial, we may perform row operations on every matrix conforming to Φ' to obtain a template

$$\Phi'' = (\{1\}, \emptyset, X, Y'_0, \emptyset, A_1, \Delta'', \Lambda),$$

so that every element of Δ'' has 0 for its first |X|-1 entries. Since every element of Δ was in the row space of A_1 , the only possible nonzero element of Δ'' is the row vector with 0 for its first |X| - 1 entries and whose last $|Y'_0| - |X| + 1$ entries form the row vector \bar{v} . Note that operations (5) and (6) and the row operations we performed on every matrix conforming to Φ' each changes a template to an equivalent template. Thus, we may assume that \bar{v} is nonzero and that $\Delta'' = \{\mathbf{0}, \bar{v}\}$ because otherwise, Φ is equivalent to a template with Δ trivial. So, for some $y \in Y'_0$, we have $\bar{v}_y = 1$. On the template Φ'' , repeatedly perform operation (11) and then operation (10) to obtain the following template:

$$\Phi''' = (\{1\}, \emptyset, \{\Sigma\}, \{y\}, \emptyset, [1], \mathbb{Z}/2\mathbb{Z}, \mathbb{Z}/2\mathbb{Z}).$$

The following matrix conforms to Φ''' :

0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1]
1	0	0	0	1	1	1	0	0	0	0	0	0	1	0	1	
0	1	0	0	1	0	0	1	1	0	0	0	1	0	0	1	
0	0	1	0	0	1	0	1	0	1	0	1	0	0	0	1	
0	0	0	1	0	0	1	0	1	1	1	0	0	0	0	1	

By contracting y, we obtain PG(3,2). Thus, we have shown that Φ must be equivalent to a template with Δ trivial. So we may assume

$$\Phi = (\{1\}, \emptyset, X, Y_0, Y_1, A_1, \{0\}, \Lambda),$$

with Λ generated by $[1, 0, \ldots, 0]^T$.

Now, let us consider the structure of the matrix A_1 . By repeated use of operation (5), we may assume that A_1 is of the following form, with the top row indexed by Σ , with * representing an arbitrary row vector, with $Y_0 = V_0 \cup V_1$, and with each L_i representing an arbitrary binary matrix:

	Y_1		V_0	V_1
$0 \cdots 0$	$0 \cdots 0$	$1 \cdots 1$	$0 \cdots 0$	*
Ι	L_0	L_1	0	L_2
0	0	0	Ι	L_3

Suppose either L_0 or L_1 has a column with two or more nonzero entries. Let y be the element of Y_1 that indexes that column, and let Y' be the union of $\{y\}$ with the subset of Y_1 that indexes the columns of the identity submatrix of $A_1[X, Y_1]$. Repeatedly perform operations (4) and (10) on Φ to obtain

$$(\{1\}, \emptyset, X, \emptyset, Y', A_1, \{0\}, \Lambda).$$

On this template, repeatedly perform y-shifts, operation (11), and operation (6) to obtain

$$(\{1\}, \emptyset, X', \emptyset, Y'', \begin{bmatrix} 0 & 0 & x \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}, \{0\}, \Lambda),$$

where x = i if y indexes a column of L_i and where X' and Y'' index the set of rows and columns, respectively, of the matrix $\begin{bmatrix} 0 & 0 & x \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}$.

The following matrix conforms to this template. By contracting the columns printed in bold, we obtain PG(3, 2).

Γ	0	0	0	0	0	0	0	0	0	0	1	1	1	1	0	0	x	$x \rceil$
	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1
	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1
	1	0	0	0	1	1	1	0	0	0	0	0	0	1	0	0	1	0
	0	1	0	0	1	0	0	1	1	0	0	0	0	1	0	0	0	1
l	0	0	1	0	0	1	0	1	0	1	1	0	1	0	1	0	0	0
L	0	0	0	1	0	0	1	0	1	1	0	1	1	0	0	1	0	0]

This shows that L_0 and L_1 consist entirely of unit and zero columns. Thus, by Lemma 3.17, L_0 is an empty matrix and L_1 consists entirely of distinct unit columns. Therefore, A_1 is of the following form:

	Y_1		V_0	V_1
$0 \cdots 0$	$0 \cdots 0$	$1 \cdots 1$	$0 \cdots 0$	*
Ι	0	Ι	0	Q_1
0	Ι	0	0	Q_2
0	0	0	Ι	Q_3

with each Q_i representing an arbitrary binary matrix.

Let M be any matroid conforming to Φ with rank and connectivity functions r and λ , respectively. Let r' be the rank of the submatrix of A_1 consisting of Q_1 , Q_2 , and the row vector we have denoted with a star. Then $r(Y_0) = |V_0| + r'$ and $r(E(M) - Y_0) = r(M) - |V_0|$. Thus, $\lambda(Y_0) = r'$. So if k > r' + 1, then M is not vertically k-connected unless Y_0 or $E(M) - Y_0$ is spanning. If Y_0 is spanning in M, then the Γ -frame matrix used to construct M has 0 rows. Thus, M is not simple unless $|E(M)| \leq |Y_0| + |Y_1| + 1$, with the 1 coming from the element $[1, 0 \cdots, 0]^T$ of Λ . Thus, if we set $l > |Y_0| + |Y_1| + 1$, then no simple, vertically k-connected matroid with at least l elements conforms to Φ unless $E(M) - V_0$ is spanning in M. Therefore, we have $V_0 = \emptyset$.

Let Q be the submatrix of A_1 consisting of Q_1 and Q_2 . If every column of Q has at most two nonzero entries, then $\Phi \leq \Phi_X$, and as we deduced above, we may assume $\Phi = \Phi_D$. Therefore, we assume that Q has a column c, indexed by the element $y \in Y_0$ with three or more nonzero entries.

Repeatedly perform operation (10) on Φ to obtain the template

$$\Phi' = (\{1\}, \emptyset, X, \{y\}, Y_1, A_1[D, Y_1 \cup \{y\}], \{0\}, \Lambda).$$

Let $c = \left\lfloor \frac{c_1}{c_2} \right\rfloor$, with c_1 a column of Q_1 and c_2 a column of Q_2 . Consider the following cases:

Case 1. The vector c_1 has three nonzero entries.

- Case 2. The vector c_1 has two nonzero entries, and c_2 has one nonzero entry.
- Case 3. The vector c_1 has one nonzero entry, and c_2 has two nonzero entries.

Case 4. The vector c_2 has three nonzero entries.

In Case i, repeatedly perform y-shifts and operation (11) to obtain the template

$$\Phi_i'' = (\{1\}, \emptyset, X', \{y\}, Y_1', A_{1,i}, \{0\}, \Lambda),$$

where $A_{1,i}$ is the matrix defined below with rows indexed by X' and columns indexed by $Y'_1 \cup \{y\}$. In each case, the last column is indexed by y, and it turns out that the value of x does not matter.

$$A_{1,1} = \begin{bmatrix} 0 & 0 & 0 & 1 & 1 & 1 & x \\ \hline 1 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 1 & 1 \end{bmatrix} A_{1,2} = \begin{bmatrix} 0 & 0 & 0 & 1 & 1 & x \\ \hline 1 & 0 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{bmatrix}$$
$$A_{1,3} = \begin{bmatrix} 0 & 0 & 0 & 1 & x \\ \hline 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 \end{bmatrix} A_{1,4} = \begin{bmatrix} 0 & 0 & 0 & x \\ \hline 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{bmatrix}$$

In Case *i*, the matrix below virtually conforms to Φ_i'' . By contracting the columns printed in bold, we obtain PG(3, 2).

Case 1:

1	1	0	0	0	0	1	1	1	0	0	0	1	1	1	x
0	0	0	1	0	0	1	0	0	1	0	0	1	0	0	1
0	0	0	0	1	0	0	1	0	0	1	0	0	1	0	1
0	0	0	0	0	1	0	0	1	0	0	1	0	0	1	1
1	0	1	1	1	1	1	1	1	0	0	0	0	0	0	0

Case 2:

ſ	1	0	0	0	1	1	1	0	0	0	1	1	0	0	1	1	\boldsymbol{x}
	0	0	0	0	0	0	0	1	0	0	1	0	1	0	1	0	1
	0	0	0	0	0	0	0	0	1	0	0	1	0	1	0	1	1
	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1
	0	1	0	1	1	0	1	1	1	1	1	1	0	0	0	0	0
	0	0	1	1	0	1	1	0	0	0	0	0	1	1	1	1	0

Case 3:

	- 0 0 0	0 0 0	0 0 0	1 0 0	1 0 0	0 1 0	$\begin{array}{c} 0 \\ 0 \\ 1 \end{array}$	0 0 0	$\begin{array}{c} 0 \\ 1 \\ 0 \end{array}$	$\begin{array}{c} 0 \\ 0 \\ 1 \end{array}$	0 0 0	1 1 0	$\begin{array}{c} 0 \\ 0 \\ 1 \end{array}$	0 0 0	$\begin{array}{c} 0 \\ 1 \\ 0 \end{array}$	$\begin{array}{c} 0 \\ 0 \\ 1 \end{array}$	0 0 0	$egin{array}{c c} x & - \ 1 & 1 & 1 \ 1 & 1 & 1 \end{array}$
	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	1	1
	1	0	1	1	0	1	1	1	0	0	0	0	0	0	0	0	0	0
	0	1	1	0	1	0	0	0	1	1	1	1	0	0	0	0	0	0
	Ο	Ο	Ω	0	1	0	0	0	0	0	0	0	1	1	0	0	Ο	0
L	_ U	0	0	U	-	U.	0	0	0	0	0	0	-	-	0	0	0	U _

Case 4:

Γ	1	0	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	$x \urcorner$
I	0	0	0	0	0	1	0	0	1	0	0	1	0	0	1	0	0	1
	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	1	0	1
	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	1	$1 \mid$
	0	0	1	0	0	1	1	1	0	0	0	0	0	0	0	0	0	0
l	0	0	0	0	1	0	0	0	1	1	1	0	0	0	0	0	0	0
	0	1	0	1	1	0	0	0	0	0	0	1	1	1	0	0	0	0]

By contradiction, this completes the proof.

Proof of Theorem 1.3. Let $\mathcal{M} = \mathcal{EX}(PG(3,2))$, and let $\mathcal{T}_{\mathcal{M}} = \{\Phi_1, \ldots, \Phi_s, \Psi_1, \ldots, \Psi_t\}$. By Lemma 4.9, for sufficiently large r, we have $h_{\mathcal{M}}(r)$ equal to the size of the largest simple matroid of rank r that virtually conforms to any template in $\Phi \in \{\Phi_1, \ldots, \Phi_s\}$. If $\Phi \in \{\Phi_1, \ldots, \Phi_s\}$, then by Lemma 4.12 either $\Phi = \Phi_X$ or Φ is of the form $(\{1\}, \emptyset, X, Y_0, Y_1, A_1, \{0\}, \{0\})$, for some matrix A_1 and some sets X, Y_0 , and Y_1 . Moreover, by operation (5), we may assume that A_1 is of the following form, with $Y_0 = V_0 \cup V_1$ and with the stars representing arbitrary binary matrices:

Y	1	V_0	V_1	
Ι	*	0	*	
0	0	Ι	*	

The largest simple matroid of rank r that virtually conforms to Φ is obtained by taking for the Γ -frame matrix a matrix representation of $M(K_{n+1})$, where $n = r - r(M(A_1[X, Y_1])) - |V_0|$. Thus, the largest simple matroid of rank r that virtually conforms to Φ has size $\binom{n+1}{2} +$ $|Y_1|n + |Y_1| + |Y_0|$. Substituting $r - r(M(A_1[X, Y_1])) - |V_0|$ for n, one sees that for sufficiently large r, this expression is less than $r^2 - r + 1$. Since the class of matroids virtually conforming to Φ_X is the class of even-cycle matroids, which has growth rate $r^2 - r + 1$, the result holds.

36

5. 1-FLOWING MATROIDS

In this section, we prove Theorem 1.4. The 1-flowing property is a generalization of the max-flow min-cut property of graphs. See Seymour [10] or Mayhew [7] for more of the background and motivation concerning 1-flowing matroids. We follow the notation and exposition of [7].

Definition 5.1. Let e be an element of a matroid M. Let c_x be a non-negative integral capacity assigned to each element $x \in E(M) - e$. A flow is a function f that assigns to each circuit C containing e a non-negative real number f_C with the constraint that for each $x \in E - e$, the sum of f_C over all circuits containing both e and x is at most c_x . We say that M is *e*-flowing if, for every assignment of capacities, there is a flow whose sum over all circuits containing e is equal to

$$\min\{\sum_{x\in C^*-e} c_x | C^* \text{ is a cocircuit containing } e\}.$$

If M is e-flowing for each $e \in E(M)$, then M is 1-flowing.

The matroid T_{11} is the even-cycle matroid obtained from K_5 by adding a loop and making every edge odd, including the loop. In [10], Seymour showed the following.

Proposition 5.2. The The class of 1-flowing matroids is minor-closed. Moreover, AG(3,2), $U_{2,4}$, T_{11} , and T_{11}^* are excluded minors for the class of 1-flowing matroids.

Seymour [10] conjectured that these are the only excluded minors.

Conjecture 5.3 (Seymour's 1-flowing Conjecture). The set of excluded minors for the class of 1-flowing matroids consists of AG(3,2), $U_{2,4}$, T_{11} , and T_{11}^* .

Since $U_{2,4}$ is an excluded minor for the class of 1-flowing matroids, all such matroids are binary. Therefore, the results in this paper apply to 1-flowing matroids. We will now prove Theorem 1.4, which we restate below.

Theorem 5.4. There exist $k, l \in \mathbb{Z}_+$ such that every simple, vertically k-connected, 1-flowing matroid with at least l elements is either graphic or cographic.

Proof. The matroid AG(3,2) conforms to Φ_{Y_1} since it is a restriction of N_{12} . Indeed, consider the matrix representing N_{12} that virtually conforms to Φ_{Y_1} . Add the rows labeled by X in this matrix

to one of the other rows. Then we can see the matrix representation $[I_4|J_4 - I_4]$ of AG(3,2) as a restriction of N_{12} . Also, it is not difficult to see that AG(3,2) can be obtained from a matroid conforming to Φ_{Y_0} by contracting Y_0 . Thus, $\mathcal{EX}(AG(3,2))$ contains neither $\mathcal{M}(\Phi_{Y_0})$ nor $\mathcal{M}(\Phi_{Y_1})$. Since AG(3,2) is self-dual, $\mathcal{EX}(AG(3,2))$ does not contain $\mathcal{M}^*(\Phi_{Y_0})$, or $\mathcal{M}^*(\Phi_{Y_1})$ either. Therefore, by Corollary 3.21, $\mathcal{EX}(AG(3,2))$ is described by the trivial template. Thus, since AG(3,2)is an excluded minor for the class of 1-flowing matroids, there exist $k, l \in \mathbb{Z}_+$ such that every simple, vertically k-connected, 1-flowing matroid with at least l elements either conforms or coconforms to the trivial template. The result follows. \Box

Acknowledgements

We thank the two anonymous referees for carefully reading the manuscript. In particular, we thank the first referee for giving many suggestions that improved the manuscript, including one that greatly simplified the proof of Lemma 4.8.

References

- [1] Jim Geelen, Bert Gerards, and Geoff Whittle, The highly connected matroids in minor-closed classes, *Annals of Combinatorics* **19** (2015), 107–123.
- [2] Jim Geelen, and Kasper Kabell, Projective geometries in dense matroids, Journal of Combinatorial Theory, Series B 99 (2009), 1–8.
- [3] Jim Geelen, and Geoff Whittle, Cliques in dense GF(q)-representable matroids, Journal of Combinatorial Theory, Series B 87 (2003), 264–269.
- [4] Jim Geelen, Joseph P.S. Kung, and Geoff Whittle, Growth rates of minorclosed classes of matroids, *Journal of Combinatorial Theory*, Series B 99 (2009), 420–427.
- [5] Jim Geelen and Peter Nelson, Matroids denser than a clique, Journal of Combinatorial Theory, Series B 114 (2015), 51–69.
- [6] Joseph P. S. Kung, Dillon Mayhew, Irene Pivotto, and Gordon F. Royle, Maximum size binary matroids with no AG(3, 2)-minor Are graphic, SIAM Journal on Discrete Mathematics 28 (2014), 1559–1577.
- [7] Dillon Mayhew, Seymour's 1-flowing Conjecture III, The Matroid Union, WordPress, 19 Mar. 2015, Web, 19 May 2016.
- [8] James Oxley, *Matroid Theory*, Second Edition, Oxford University Press, New York, 2011.
- [9] Neil Robertson and P.D. Seymour, Graph Minors. XVI. Excluding a non-planar graph, Journal of Combinatorial Theory, Series B 89 (2003), 43–75.
- [10] P.D. Seymour, Matroids and Multicommodity Flows. European Journal of Combinatorics 2 (1981), 257-290.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA

E-mail address: kgrace3@lsu.edu

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA

E-mail address: svanzwam@math.lsu.edu