LYAPUNOV-LIKE FUNCTIONS INVOLVING LIE BRACKETS

MONICA MOTTA AND FRANCO RAMPAZZO

ABSTRACT. For a given closed target we embed the dissipative relation that defines a control Lyapunov function in a more general differential inequality involving Hamiltonians built from iterated Lie brackets. The solutions of the resulting extended relation, here called *degree-k control Lyapunov functions* $(k \ge 1)$, turn out to be still sufficient for the system to be globally asymptotically controllable to the target. Furthermore, we work out some examples where no standard (i.e., degree-1) smooth control Lyapunov functions exist while a C^{∞} degree-k control Lyapunov function does exist, for some k > 1. The extension is performed under very weak regularity assumptions on the system, to the point that, for instance, (set valued) Lie brackets of locally Lipschitz vector fields are considered as well.

1. INTRODUCTION

A control Lyapunov function (shortly, CLF) for a control system

(1)
$$\begin{cases} \dot{y} = f(y, a) \\ y(0) = x \in I\!\!R^n \backslash \mathcal{T} \end{cases}$$

-where the control parameter *a* ranges over a compact set of controls, and the (closed) subset $\mathcal{T} \subset \mathbb{R}^n$ is regarded as a *target*- is a positive definite function $U: \overline{\mathbb{R}^n \setminus \mathcal{T}} \to \mathbb{R}$

such that, at each point $x \in \mathbb{R}^n \setminus \mathcal{T}$, the dynamics f(x, a) points in a direction along which U is strictly decreasing, for a suitable choice of $a \in A$. A wide literature investigates the links between the existence of a CLF and some properties of the system-target pair. Standard regularity assumptions include local semiconcavity of U in the interior of the domain of U, which, in particular, allows defining the set of limiting gradients ${}^1 D^* U(x)$ at each

²⁰¹⁰ Mathematics Subject Classification. 34A26, 93D30, 93C15.

Key words and phrases. Lyapunov functions, Lie brackets, global asymptotic controllability, partial differential inequalities.

This research is partially supported by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM), Italy; and by Padova University grant PRAT 2015 "Control of dynamics with reactive constraints".

¹See Definition 2.3. Under this hypothesis, $D^*U(x)$ coincides with the *limiting subdif*ferential $\partial_L U(x)$, largely used in the literature on Lyapunov functions.

 $x \in \mathbb{R}^n \setminus \mathcal{T}$. Therefore, the monotonicity of U along suitable directions of f can be expressed by means of the *dissipative* differential inequality

(2)
$$H(x, D^*U(x)) < 0 \qquad \forall x \in \mathbb{R}^n \setminus \mathcal{T},$$

where

$$H(x,p) := \inf_{a \in A} \left\langle p, f(x,a) \right\rangle.$$

Relation (2) has to be interpreted as the occurrence, at each x, of the inequality H(x, p) < 0 for every $p \in D^*U(x)$. Since U is assumed to be (proper and) positive definite, by choosing controls verifying (2) one is ideally looking for trajectories that run closer and closer to the target. More precisely, one has:

Theorem 1.1. If there exists a CLF, system (1) is GAC to \mathcal{T} .

As customary, GAC to \mathcal{T} is acronym of globally asymptotically controllable to \mathcal{T} (see Definition 1.2), which means that for any initial point x there exists a system trajectory $y(\cdot)$, y(0) = x, approaching the target \mathcal{T} (in possibly infinite time), uniformly with respect to the distance $\mathbf{d}(x, \mathcal{T})$.

Results like Theorem 1.1 –of which some "inverse" versions exist as well – lie at the basis of various constructions dealing, in particular, with stabilizability (see e.g. [S2], [Ri] and the references therein). Nonsmoothness is crucial for control Lyapunov functions: though relation (2) is a partial differential inequality –so admitting many more solutions than the corresponding Hamilton-Jacobi equation– in general no smooth control Lyapunov functions exist. A great deal of effective ideas has been flourishing during the last four decades to deal with this unavoidable lack of regularity (see e.g. [CLSS], [MaRS], the books [CLSW], [BR] and the references therein). Nevertheless, the regularity issue is of obvious interest from a numerical point of view. In addition, any feedback stabilizing strategy would likely benefit from smoothness properties of a CLF (or of some suitable CLF's replacement)– in particular, in reference with sensitivity to data errors.

As an attempt to reduce the unavoidability of nonsmoothness, in the present paper we replace relation (2) with a less demanding inequality which involves Lie brackets ². Let us assume that the dynamics is driftless control affine, namely:

(3)
$$\begin{cases} \dot{y} = \sum_{i=1,\dots,m} a_i f_i(y) \\ y(0) = x \in \mathbb{R}^n \setminus \mathcal{T}, \end{cases}$$

²We remind that the *Lie bracket* of two C^1 vector fields X, Y is defined (on any coordinate chart) as $[X, Y] := DY \cdot X - DX \cdot Y$.

and let $A := \{\pm e_1, \ldots, \pm e_m\}^3$. Assume the vector fields f_1, \ldots, f_m are of class C^{k-1} for some integer $k \ge 1$. We will define ⁴ the *degree-k Hamiltonian* $H^{(k)}(x,p)$ by setting

(4)
$$H^{(k)}(x,p) := \inf_{v \in \mathcal{F}^{(k)}(x)} \langle p, v \rangle \qquad \forall (x,p) \in (I\!\!R^n \backslash \mathcal{T}) \times I\!\!R^n,$$

where $\mathcal{F}^{(k)}$ denotes the family of iterated Lie brackets of degree $\leq k$ of the vector fields f_1, \ldots, f_m . (Notice, in particular, that $H^{(1)} = H$).

A function $U : \overline{\mathbb{R}^n \setminus \mathcal{T}} \to \mathbb{R}$ will be called a *degree-k control Lyapunov* function –shortly, degree-k CLF– if (it is positive definite, proper, semiconcave on domain's interior, and) it verifies inequality

(5)
$$H^{(k)}(x, D^*U(x)) < 0 \qquad \forall x \in \mathbb{R}^n \setminus \mathcal{T}.$$

Observe that, because of

(6)
$$H^{(k)} \le H^{(k-1)} \dots \le H^{(1)}$$

relation (5) is weaker than (2).

Still, in view of Theorem 1.2 below, the inequality (5) is sufficient for the system to be GAC to \mathcal{T} , as stated in the following result:

Theorem 1.2. Let a degree-k CLF exist, for some positive integer k. Then system (3) is GAC to \mathcal{T} .

The use of Lie brackets as *higher order directions* is widespread in Control Theory, both within necessary conditions for optimality and within sufficient conditions for various kinds of controllability (see e.g. [AgSa], [BP], [Co], [K], [S1], [Su], [FHT]). Furthermore, they are involved in boundary conditions ensuring uniqueness for Hamilton-Jacobi equations, e.g. in relation with continuity properties of the corresponding value function (see e.g. [BCD], [So]). However, here Lie brackets are directly involved in the proposed differential inequalities.

As for the regularity issue, we wish to remark that a degree-k control Lyapunov function, k > 1, may happen to be more regular than a standard (i.e., degree-1) control Lyapunov function. It may even occur the case where $H^{(k)}(x, D^*U(x)) < 0$ for some C^{∞} function U, while no smooth U satisfies the standard inequality $H(x, D^*U(x)) < 0$ (see Examples 2.1-2.3 below). Let us observe that the two reasons why a (degree-1) control Lyapunov function may result discontinuous are : i) the *shape* of the target's boundary $\partial \mathcal{T}$; ii) the *shortage* of dynamics' directions. While there is nothing one can do to remedy i), the introduction of Hamiltonians $H^{(k)}$ (k > 1), which are minima over larger sets of directions, is a way to reduce the effects of ii).

The regularity hypotheses in the case of degree-2 control Lyapunov functions are relaxed in Section 4 in order to include Lipschitz continuous vector

³More general control systems can be considered: see Remark 2.2 and subsection 5.3.

⁴See Section 4 for an extension of the notion of $H^{(2)}$ when the vector fields are locally Lipschitz but not *smooth*.

fields. Since the classical brackets $[f_i, f_j]$ may happen to be not even defined at possibly infinitely many points, we make use of the generalized, set-valued brackets defined in [RS1]. Accordingly, the Hamiltonian $H^{(2)}$ is computed as a min-max value. Let us remark that the degree-2 control Lyapunov function of Example 4.1 is C^{∞} despite the fact the vector fields are not even C^1 .

1.1. Preliminaries and notation. For the reader convenience, some classical concepts, like global asymptotic controllability to a set \mathcal{T} , in short GAC to \mathcal{T} , and a few technical definitions are here recalled.

Given an integer $k \geq 1$ and an open subset $\Omega \subseteq \mathbb{R}^n$, we write $C^k(\Omega)$ to denote the set of vector fields of class C^k on Ω , namely, $C^k(\Omega) := C^k(\Omega, \mathbb{R}^n)$. The subset $C_b^k(\Omega) \subset C^k(\Omega)$ of functions with bounded derivatives (up to the order k) will be endowed with the norm

$$||f||_k := \sum_{i=0,\dots,k} \sup_{x \in \Omega} |f^{(i)}(x)| \qquad (f^{(0)} := f)$$

(which makes it a Banach space). Similarly, $C^{k-1,1}(\Omega) \subset C^{k-1}(\Omega)$ denotes the subset of vector fields whose k - 1-th derivative is locally Lipschitz continuous and $C_b^{k-1,1}(\Omega)$ is the subset of $C_b^{k-1}(\Omega)$ with (globally) Lipschitz continuous k - 1-th derivative.

Definition 1.1. Let $k \geq 1$ be an integer, and let f_1, \ldots, f_m be vector fields belonging to $C^{k-1}(\mathbb{R}^n \setminus \mathcal{T})$. For any initial condition $x \in \mathbb{R}^n \setminus \mathcal{T}$ and any measurable control $\alpha : [0, +\infty) \to A$, a trajectory-control pair $(y, \alpha)(\cdot)$ will be called admissible if there exists $T \leq +\infty$ such that $y(\cdot)$ is a solution of (3) defined on [0, T) and

$$\lim_{t \to T} \mathbf{d}(y(t)) = 0,$$

where $\mathbf{d}(\cdot) := \mathbf{d}(\cdot, \mathcal{T})$. When k > 1, we will use $y_x(\cdot, \alpha)$ to denote the unique (possibly local) forward solution to the Cauchy problem (3).

Remark 1.1. The main object of the paper consists in establishing relations involving Lie brackets, so that a certain regularity is necessary when k > 1 (see also Section 4). However, observe that as soon as k = 1 the vector fields f_1, \ldots, f_m are just continuous, so that solutions of the Cauchy problem (3) for a given control may be not unique.

To give the notion of global asymptotic controllability, we recall that $\mathcal{K}L$ is used to denote the set of continuous functions $\beta : [0, +\infty) \times [0, +\infty) \rightarrow [0, +\infty)$ such that: (1) $\beta(0, s) = 0$ and $\beta(\cdot, s)$ is strictly increasing and unbounded for each $s \geq 0$; (2) $\beta(\delta, \cdot)$ is decreasing for each $\delta \geq 0$; (3) $\beta(\delta, s) \rightarrow 0$ as $s \rightarrow +\infty$ for each $\delta \geq 0$.

Definition 1.2. The control system in (3) is globally asymptotically controllable to \mathcal{T} -shortly, (3) is GAC to \mathcal{T} - provided there is a function $\beta \in \mathcal{K}L$

such that, for each initial state $x \in \mathbb{R}^n \setminus \mathcal{T}$, there exists an admissible trajectory-control pair $(y, \alpha)(\cdot)$ such that

(7)
$$\mathbf{d}(y(t)) \le \beta \big(\mathbf{d}(x), t \big) \qquad \forall t \in [0, +\infty).$$
⁵

Let us recall that if g_1, g_2 are C^1 vector fields on a differential manifold (of class C^2), their *Lie bracket* $[g_1, g_2]$ is the (continuous) vector field which is defined (on coordinate charts) by

$$[g_1, g_2] = Dg_2 \cdot g_1 - Dg_2 \cdot g_1.$$

Since $[g_1, g_2]$ turns out to be a vector field, provided sufficient regularity is assumed, one can iterate the bracketing process so obtaining *iterated Lie* brackets. We call degree of a given iterated bracket B the number of objects appearing in B (regarded as a formal object) when commas and left and right brackets are deleted. For instance, the degrees of $[[g_2, g_3], g_2],$ $[[g_2, g_3], [g_2, g_4]],$ and $[g_4, [g_4, [g_4, [g_4, g_6]]]]$ are 3, 4, and 5, respectively.

Let us summarize some basic notions in nonsmooth analysis (see e.g. [CS], [CLSW] for a thorough treatment).

Definition 1.3 (Positive definite and proper functions). A continuous function $F : \overline{\mathbb{R}^n \setminus \mathcal{T}} \to \mathbb{R}$ is said positive definite on $\mathbb{R}^n \setminus \mathcal{T}$ if F(x) > 0 $\forall x \in \mathbb{R}^n \setminus \mathcal{T}$ and $F(x) = 0 \ \forall x \in \partial \mathcal{T}$. The function F is called proper on $\mathbb{R}^n \setminus \mathcal{T}$ if the pre-image $F^{-1}(K)$ of any compact set $K \subset [0, +\infty]$ is compact.

Definition 1.4. (Semiconcavity). Let $\Omega \subset \mathbb{R}^n$. A continuous function $F: \Omega \to \mathbb{R}$ is said to be semiconcave on Ω if there exist $\rho > 0$ such that

$$F(z_1) + F(z_2) - 2F\left(\frac{z_1 + z_2}{2}\right) \le \rho |z_1 - z_2|^2,$$

for all $z_1, z_2 \in \Omega$ such that $[z_1, z_2] \subset \Omega$. The constant ρ above is called a semiconcavity constant for F in Ω . F is said to be locally semiconcave on Ω if it semiconcave on every compact subset of Ω .

Let us remind that locally semiconcave functions are locally Lipschitz. Actually, they are twice differentiable almost everywhere.

Definition 1.5. (Limiting gradient). Let $\Omega \subset \mathbb{R}^n$ be an open set, and let $F: \Omega \to \mathbb{R}$ be a locally Lipschitz function. For every $x \in \Omega$ let us set

$$D^*F(x) \doteq \left\{ w \in \mathbb{R}^n : w = \lim_k \nabla F(x_k), x_k \in DIFF(F) \setminus \{x\}, \lim_k x_k = x \right\}$$

where ∇ denotes the classical gradient operator and DIFF(F) is the set of differentiability points of F. $D^*F(x)$ is called the set of limiting gradients of F at x.

⁵ By convention, we fix an arbitrary $\bar{x} \in \partial \mathcal{T}$ and formally establish that, if $T < +\infty$, the trajectory $y(\cdot)$ is prolonged to $[0, +\infty)$, by setting $y(t) = \bar{x}$ for all $t \ge T$.

The set-valued map $x \mapsto D^*F(x)$ is upper semicontinuous on Θ , with nonempty, compact values. Notice that $D^*F(x)$ is not convex. When F is a locally semiconcave function, D^*F coincides with the limiting subdifferential $\partial_L F$, namely,

$$D^*F(x) = \partial_L F(x) := \{\lim p_i : p_i \in \partial_P F(x_i), \lim x_i = x\} \quad \forall x \in \Theta,$$

where $\partial_P F$ denotes the proximal subdifferential, largely used in the literature on Lyapunov functions.

2. Degree-k control Lyapunov functions

2.1. The main result. Let $k \geq 1$ be an integer. Throughout the whole paper we assume that the target $\mathcal{T} \subset \mathbb{R}^n$ is a closed set with compact boundary and that f_1, \ldots, f_m are vector fields belonging to $C_b^{k-1}(\Omega \setminus \mathcal{T})$ for any open, bounded subset $\Omega \subset \mathbb{R}^n$ (see Subsection 1.1).

Definition 2.1. Let us consider the family of vector fields

$$\mathcal{F}^{(1)} := \left\{ f = \sum_{i=1}^{m} a_i f_i, \quad a \in A \right\} = \left\{ \pm f_i, \quad i = 1, \dots, m \right\}.$$

Moreover, if k > 1, for every positive integer h such that $2 \le h \le k$, set

 $\mathcal{F}^{(h)} := \{B, B \text{ is a iterated Lie bracket of degree} \le h \text{ of } f_1, \dots, f_m \}$

Clearly, every element of $\mathcal{F}^{(h)}$ is a vector field belonging to $C_b^{k-h}(\Omega \setminus \mathcal{T})$ for any open, bounded subset $\Omega \subset \mathbb{R}^n$. Notice that

(8)
$$\mathcal{F}^{(1)} \subseteq \mathcal{F}^{(2)} \subseteq \cdots \subseteq \mathcal{F}^{(k)}$$

For every h = 1, ..., k, let us introduce the set-valued map

$$\mathcal{F}^{(h)}(x) := \left\{ X(x), \quad X \in \mathcal{F}^{(h)} \right\} \qquad \forall x \in \mathbb{R}^n \setminus \mathcal{T}.$$

Definition 2.2. For any integer $1 \leq h \leq k$, let us define the degree-h Hamiltonian $H^{(h)}$ corresponding to the control system (3), by setting

$$H^{(h)}(x,p) := \inf_{v \in \mathcal{F}^{(h)}(x)} \left\langle p, v \right\rangle \qquad \forall (x,p) \in (\mathbb{R}^n \setminus \mathcal{T}) \times \mathbb{R}^n.$$

Under the above hypotheses the Hamiltonians $H^{(h)}$ are well defined and continuous. As already mentioned in the Introduction, the degree-1 Hamiltonian $H^{(1)}$ coincides with the standard Hamiltonian:

$$H^{(1)}(x,p) = H(x,p) := \inf_{a \in A} \left\langle p, \sum_{i=1}^{m} a_i f_i(x) \right\rangle.$$

Morever, by (8) one gets

(9)
$$H^{(k)} \le H^{(k-1)} \le \dots \le H^{(1)}$$

Definition 2.3. We call degree-k control Lyapunov function -in short, degree-k CLF- any continuous function $U : \mathbb{R}^n \setminus \mathcal{T} \to \mathbb{R}$ such that the restriction to $\mathbb{R}^n \setminus \mathcal{T}$ is locally semiconcave, positive definite, proper, and verifies

(10)
$$H^{(k)}(x, D^*U(x)) < 0 \quad \forall x \in I\!\!R^n \setminus \mathcal{T},$$

the latter inequality meaning $H^{(k)}(x,p) < 0$ for each $p \in D^*U(x)$.

In Theorem 2.1 below we prove that the existence of a degree-k control Lyapunov function, k > 1, is sufficient for the system to be globally asymptotically controllable to \mathcal{T} (GAC to \mathcal{T} , see Definition 1.2), as in the classical case k = 1.

Theorem 2.1. Let us assume that, for some integer $k \ge 1$, a degree-k control Lyapunov function exists. Then system (3) is GAC to \mathcal{T} .

We postpone the proof of Theorem 2.1 to the next section and make some general remarks. Furthermore, we give some examples where, in particular, the distance function is a (possibly smooth) degree-k CLF for some k > 1 and *is not* a degree-1 CLF.

2.2. Remarks and examples.

Remark 2.1. The regularity assumptions can be slightly weakened in some cases by observing that, in order that certain degree-k brackets (k > 3) to be defined, it is not necessary that the vector fields are k-1 times differentiable. For instance, the bracket $[[f_1, f_2], [f_3, f_4]]$ is well defined as soon as the vector fields f_1, f_2, f_3, f_4 are two times differentiable.

Remark 2.2. By suitably rescaling time, one can easily generalize Theorem 2.1 to the case when the control set A contains a ball of \mathbb{R}^m with positive radius. By means of linear algebraic and relaxation arguments one can also try to extend the result up to the point of admitting sets A such that 0 is contained in the interior of the convex hull co(A).

Remark 2.3. It is easy to adapt Theorem 2.1 to the case when the state space is an open set $\Omega \subset \mathbb{R}^n$, $\Omega \supset \mathcal{T}$. In fact, the thesis keeps unchanged as soon as one requires the degree-k CLF $U : \Omega \setminus \overset{\circ}{\mathcal{T}} \to \mathbb{R}$ to verify all the assumptions in Definition 2.3 in Ω , plus the following one:

$$\exists U_0 \in (0, +\infty] : \lim_{x \to x_0, \ x \in \Omega} U(x) = U_0 \ \forall x_0 \in \partial\Omega; \quad U(x) < U_0 \quad \forall x \in \Omega \setminus \overset{\circ}{\mathcal{T}}.$$

Remark 2.4. While the fact that U is a degree-k control Lyapunov function implies that U is also a degree- \bar{k} control Lyapunov function for every $\bar{k} > k$, the converse is in general false (see Example 2.1). On the other hand, coupling Theorem 2.1 with an *inverse Lyapunov* result like in [S2], [Ri], it is easy to verify that the existence of a degree-k control Lyapunov function, k > 1, implies the existence of a standard (i.e., degree-1) control Lyapunov function. **Remark 2.5.** As in the case of standard (i.e. degree-1) CLF, the notion of degree-k CLF is intrinsic, for vector fields, their Lie brackets, and the set of limiting gradients D^*U are chart independent. In particular the results in Theorems 2.1 and 4.1 are fit to be extended to Riemannian manifolds (where, of course, the notion of distance should coincide with the considered Riemannian metric). Incidentally, let us notice that we can define the Hamiltonians $H^{(k)}$ in terms of Poisson brackets⁶. Indeed, setting, for every vector field X

$$H_X(x,p) := \langle p, X(x) \rangle$$

one has

$$H^{(1)}(x,p) = \inf \left\{ -|H_{f_i}(x,p)|, \ i = 1, \dots, m \right\},$$
$$H^{(2)}(x,p) = \inf \left\{ H^{(1)}(x,p), -|\{H_{f_i}, H_{f_j}\}(x,p)|, \ i,j = 1, \dots, m \right\},$$
$$H^{(3)}(x,p) = \inf \left\{ H^{(2)}(x,p), -|\{H_{f_i}, \{H_{f_j}, H_{f_\ell}\}\}(x,p)|, \ i,j,\ell = 1, \dots, m \right\}$$
and similarly for higher degrees

and similarly for higher degrees.

Example 2.1. Consider the so-called *nonholonomic integrator*

$$\dot{y} = a_1 f_1(y) + a_2 f_2(y) ,$$

where $f_1 := \frac{\partial}{\partial x_1} - x_2 \frac{\partial}{\partial x_3}, f_2 := \frac{\partial}{\partial x_2} + x_1 \frac{\partial}{\partial x_3}.$
By $[f_1, f_2] = 2 \frac{\partial}{\partial x_3}$ we get
 $H^{(1)}(x, p) = -\max\{|p_1 - p_3 x_2|, |p_2 + p_3 x_1|\}$

and

$$H^{(2)}(x,p) = -\max\left\{ \left| p_1 - p_3 x_2 \right|, \left| p_2 + p_3 x_1 \right|, \left| 2 \right| p_3 \right| \right\}.$$

Let \mathcal{T} be a compact target and let $U(\cdot)$ coincide with the distance $\mathbf{d}(\cdot)$ from \mathcal{T} . If there exists a point $\bar{x} \in (\{0\} \times \{0\} \times \mathbb{R}) \cap (\mathbb{R}^3 \setminus \mathcal{T})$ such that $D^*(U)(x) \cap (\{0\} \times \{0\} \times \mathbb{R}) \neq \emptyset$, then $H^{(1)}(\bar{x}, D^*U(\bar{x})) = 0$. Therefore, in this case the distance function U fails to be a degree-1 CLF. For instance, this is the case when $\mathcal{T} = \{x, ||x| \leq \rho\}$ for some $\rho \geq 0$. Indeed,

$$H^{(1)}((0,0,x_3), D^*U((0,0,x_3))) = 0,$$

for all $|x_3| > \rho$. In fact, when $\rho = 0$ no degree-1 CLF of class C^1 exist (see [BR] and [Ri]).

$$\{H, K\}(x, p) := \sum_{i=1}^{k} \left(\frac{\partial H}{\partial x_i} \frac{\partial K}{\partial p_i} - \frac{\partial H}{\partial p_i} \frac{\partial K}{\partial x_i} \right) (x, p).$$

⁶If H(x,p) and K(x,p) are differentiable functions, the Poisson bracket $\{H,K\}$ is defined by

3:

Yet, U is a degree-2 CLF, for whichever compact target \mathcal{T} . Indeed, for every $x \in \mathbb{R}^3 \setminus \mathcal{T}$ one has |p| = 1 for all $p \in D^*U(x)$, which implies

$$H^{(2)}(x, D^*U(x)) \le \max_{|p|=1} \left\{ -\max\left\{ |p_1 - p_3 x_2|, |p_2 + p_3 x_1|, 2|p_3| \right\} \right\} < 0,$$

for all $x \in \mathbb{R}^3 \setminus \mathcal{T}$. In the case when $\mathcal{T} = \{x, ||x| \leq \rho\}$ for some $\rho \geq 0, U$ is a C^1 -actually, C^{∞} -degree-2 CLF (hence $D^*U(x) = \{\nabla U(x)\}$). Furthermore, one has

(11)
$$H^{(2)}(x, D^*U(x)) \le -\frac{2}{3} \qquad \forall x \in \mathbb{R}^3 \setminus \mathcal{T}.$$

Example 2.2. Now let us consider the system

$$\dot{y} = a_1 f_1(y) + a_2 f_2(y)$$
where $f_1 := \frac{\partial}{\partial x_2} + x_2^2 \frac{\partial}{\partial x_3}, f_2 := \frac{\partial}{\partial x_2} + x_1^2 \frac{\partial}{\partial x_3}.$
Let us compute the brackets of degree less than or equal to

$$[f_1, f_2](x) = 2(x_1 - x_2)\frac{\partial}{\partial x_3}, \quad [f_1, [f_1, f_2]](x) = -[f_2, [f_1, f_2]](x) = 2\frac{\partial}{\partial x_3}.$$

Therefore,

$$H^{(1)}(x,p) = -\max\left\{ |p_1 + p_3 x_2^2|, |p_2 + p_3 x_1^2| \right\},\$$

$$H^{(2)}(x,p) = -\max\left\{ |p_1 + p_3 x_2^2|, |p_2 + p_3 x_1^2|, 2|p_3(x_1 - x_2)| \right\},\$$

and

$$H^{(3)}(x,p) = -\max\left\{ |p_1 + p_3 x_2^2|, |p_2 + p_3 x_1^2|, 2|p_3(x_1 - x_2)|, 2|p_3| \right\}.$$

For simplicity let us consider only the target $\mathcal{T} = \{0\}$. Once again, the distance function U(x) := |x| is not a degree-1 CLF, since $H^{(1)}(x, D^*U(x)) = 0$ for all $x \in \{0\} \times \{0\} \times (\mathbb{R} \setminus \{0\})$. U is not even a degree-2 CLF, for $H^{(2)}(x, D^*U(x)) = 0$ for all $x \in \{0\} \times \{0\} \times (\mathbb{R} \setminus \{0\})$. However, $|\nabla U(x)| = 1$ (and $D^*U(x) = \{\nabla U(x)\}$) so that

$$H^{(3)}(x, D^*U(x)) \le \max_{|p|=1} H^{(3)}(x, p) < 0,$$

and the distance U is a (C^{∞}) degree-3 CLF.

Remark 2.6. The control systems in Examples 2.1 and 2.2 verify a Lie algebra rank condition at each point ⁷. Hence, by Chow-Rashevsky's Theorem, they are small time locally controllable at every point x, that is, the interior of the reachable set from x at any time contains x. Actually, with akin arguments it is not difficult to prove the following general fact:

Let $\mathcal{T} \subset \mathbb{R}^n$ be any target with compact boundary. If a system verifies the Lie algebra rank condition at every point by means of brackets of degree $\leq k$ the distance function $\mathbf{d}(\cdot)$ from \mathcal{T} is a degree-k CLF.

⁷A system verifies the Lie algebra rank condition at x if the iterated Lie brackets linearly span \mathbb{R}^n .

Indeed, since |p| = 1 for every $p \in D^* \mathbf{d}(x)$ and every $x \in \mathbb{R}^n \setminus \mathcal{T}$, the Lie algebra rank condition implies that for every such x and p there must exist $w \in \mathcal{F}^k(x)$ such that $\langle p, w \rangle < 0$.

While in the previous examples the minimum time function is finite at each point, this is not the case of the following example, where no trajectories issuing from points (x_1, x_2, x_3) such that $x_3 \neq 0$ can reach the target (in finite time). Notice incidentally that the Lie algebra rank condition is violated at each point belonging to the plane $x_3 = 0$.

Example 2.3. Consider the system

$$\dot{y} = a_1 f_1(y) + a_2 f_2(y) \,,$$

where

$$f_1 := \frac{\partial}{\partial x_1} - x_2 \phi(x_3) \frac{\partial}{\partial x_3} \quad f_2 := \frac{\partial}{\partial x_2} + x_1 \phi(x_3) \frac{\partial}{\partial x_3},$$

 $\phi : \mathbb{R} \to [0, +\infty[$ being a C^1 function such that $\phi(x_3) = 0$ if and only if $x_3 = 0$. By $[f_1, f_2](x) = 2\phi(x_3)\frac{\partial}{\partial x_3}$ one gets

$$H^{(1)}(x,p) = -\max\left\{ \left| p_1 - p_3 x_2 \phi(x_3) \right|, \left| p_2 + p_3 x_1 \phi(x_3) \right| \right\}$$

and

$$H^{(2)}(x,p) = -\max\left\{ \left| p_1 - p_3 x_2 \phi(x_3) \right|, \left| p_2 + p_3 x_1 \phi(x_3) \right|, \left| 2\phi(x_3) \right| p_3 \right| \right\}.$$

Let us consider again the target $\mathcal{T} = \{0\}$. Also in this case the distance function U(x) := |x| is not a degree-1 CLF, since

$$H^{(1)}(x, D^*U(x)) = 0 \quad \forall x \in \{0\} \times \{0\} \times (I\!\!R \setminus \{0\}),$$

and U is a still degree-2 CLF, since

$$H^{(2)}(x, D^*U(x)) < 0$$

for all $x \in \{0\} \times \{0\} \times \{R \setminus \{0\}\}$.

Of course the fact that the Lie algebra rank condition is verified almost everywhere –as in the previous examples– is far from being necessary for a CLF of whatever degree to exist. In fact, a system that fails to be small time locally controllable on large areas of its domain might not have any C^1 degree-1 CLF while admitting a smooth degree-k CLF, for some k > 1, as illustrated in the following example:

Example 2.4. Let $\varphi, \psi : [0, +\infty) \to [0, 1]$ be C^{∞} maps such that, for any $q \in \mathbb{N}$,

$$\begin{aligned} \varphi(r) &= 1 \quad \text{if } r \in [2q, 2q+1], \\ \varphi(r) &= 0 \quad \text{if } r \in [2q+(5/4), 2q+(7/4)]; \\ \psi(r) &= 1 \quad \text{if } r \in [2q+(7/8), 2q+(17/8)], \\ \psi(r) &= 0 \quad \text{if } r \in [2q+(1/4), 2q+(3/4)] \cup [0, (1/4)] \end{aligned}$$

Let us consider the control system

$$\dot{y} = a_1 f_1(y) + a_2 f_2(y) + a_3 f_3(y),$$

11

where

$$f_1 = \varphi(|x|) \left(\frac{\partial}{\partial x_1} - x_2 \frac{\partial}{\partial x_3} \right), \quad f_2 = \varphi(|x|) \left(\frac{\partial}{\partial x_2} + x_1 \frac{\partial}{\partial x_3} \right),$$
$$f_3 = \psi(|x|) \left(x_1 \frac{\partial}{\partial x_1} + x_2 \frac{\partial}{\partial x_2} + x_3 \frac{\partial}{\partial x_3} \right).$$

Clearly the system is not small time locally controllable at every point x such that $2q + (5/4) \leq |x| \leq 2q + (7/4)$. Let the target \mathcal{T} coincide with the origin $\{0\}$ and, again, let us set $U(x) := \mathbf{d}(x) = |x|$. For every $q \in \mathbb{N}$ one has $H^{(1)}(x, D^*U(x)) = -|x|$ for all x such that $2q + (5/4) \leq |x| \leq 2q + (7/4)$. Furthermore, $H^{(1)}(x, D^*U(x)) = 0$ for every x such that $x_1 = x_2 = 0$ and $|x_3| \leq 1$ or $2q + (1/4) \leq |x_3| \leq 2q + (3/4), q \geq 1^8$. However, one easily checks that

$$H^{(2)}(x, D^*U(x)) = -\frac{1}{|x|} \max\left\{ |x_1 - x_3 x_2|, |x_2 + x_3 x_1|, 2|x_3| \right\} \le -\frac{2}{3},$$

for all $x \in \mathbb{R}^3 \setminus \mathcal{T}$, so that U is a (C^{∞}) degree-2 Lyapunov function.

3. Proof of Theorem 2.1

The case when k = 1 has already been proved in [MR], where the hypotheses are even weaker than the ones assumed here (for instance, vector fields are allowed to be unbounded near the target). So we will always assume k > 1: in particular, there will be a unique trajectory $y_x(\cdot, \alpha)$ corresponding to an initial condition x and a control $\alpha(\cdot)$.

3.1. Preliminary facts. To begin with, let us point out that the 0 in the dissipative relation can be replaced by a nonnegative function of U:

Proposition 3.1. Let $U : \overline{\mathbb{R}^n \setminus \mathcal{T}} \to \mathbb{R}$ be a continuous function, such that U is locally semiconcave, positive definite and proper on $\mathbb{R}^n \setminus \mathcal{T}$. Then the conditions (i) and (ii) below are equivalent:

(i) U verifies

(12)
$$H^{(2)}(x, D^*U(x)) < 0$$

for all $x \in \mathbb{R}^n \setminus \mathcal{T}$;

(ii) for every $\sigma > 0$ there exists a continuous, strictly increasing, function $\gamma : [0, +\infty) \rightarrow : [0, +\infty)$ such that

(13)
$$H^{(2)}(x, D^*U(x)) \le -\gamma(U(x))$$

for all $x \in U^{-1}((0, 2\sigma])$.

⁸Actually, there are no C^1 degree-1 CLF, as it can be proved by noticing that the system coincides with the nonholonomic integrator in a whole neighborhood of the target.

Notice that the only non trivial implication, namely (i) \implies (ii), is a simple consequence of the upper semicontinuity of the set-valued map $x \mapsto D^*U(x)$ on the compact sets $U^{-1}([u, 2\sigma])$ ($u \in (0, 2\sigma)$) and of the upper semicontinuity of $H^{(k)}$. For a detailed proof, we refer to [MR, Proposition 3.1].

Remark 3.1. In a good deal of literature on control Lyapunov functions one utilizes the proximal subdifferential $\partial_P U(x)$ as a nonsmooth substitute for the derivative of U (see [CLSW]). However, in view of Proposition 3.1, our using the set of limiting gradients $D^*U(x)$ (which coincides with the limiting subdifferential $\partial_L U(x)$, see Subsection 1.1) is equivalent to using the proximal subdifferential. Indeed, for any N > 0 condition $H^{(k)}(x, D^*U(x)) \leq -\gamma(Ux)$) is equivalent to $H^{(k)}(x, \partial_L U(x)) \leq -\gamma(Ux)$) for any $x \in U^{-1}((0, N])$. On the other hand, by the construction of $\partial_L U(x)$ and by the continuity of $H^{(k)}(\cdot)$, this holds true as soon as $H^{(k)}(x, \partial_P U(x)) \leq -\gamma(Ux)$).

Secondly, basic properties of the semiconcave functions imply the following fact (see e.g. [CS]):

Lemma 3.1. Let $U : \overline{\mathbb{R}^n \setminus \mathcal{T}} \to \mathbb{R}$ be a continuous function, such that U is locally semiconcave, positive definite, and proper on $\mathbb{R}^n \setminus \mathcal{T}$. Then for any compact set $\mathcal{K} \subset \mathbb{R}^n \setminus \mathcal{T}$ there exist some positive constants L and ρ such that, for any $x \in \mathcal{K}^{9}$,

 $|p| \le L \quad \forall p \in D^* U(x),$

$$U(\hat{x}) - U(x) \le \langle p, \hat{x} - x \rangle + \rho |\hat{x} - x|^2$$

for any point $\hat{x} \in \mathcal{K}$ such that $[x, \hat{x}] \subset \mathcal{K}$.

Moreover, if $\mathcal{K}_1, \mathcal{K}_2 \subset \mathbb{R}^n \setminus \mathcal{T}$ are compact subsets and $\mathcal{K}_1 \subseteq \mathcal{K}_2$, we can choose the corresponding constants L_1 , ρ_1 and L_2 , ρ_2 such that $L_1 \leq L_2$ and $\rho_1 \leq \rho_2$.

3.2. A degree-k "feedback". Let us introduce a notion of degree-k feedback. For a given $\sigma > 0$, let γ be a function as in Proposition 3.1, and let $x \mapsto p(x)$ be a selection of $x \mapsto D^*U(x)$ on $U^{-1}((0, 2\sigma])$, so that

$$H^{(k)}(x, p(x)) \le -\gamma(U(x)) \qquad \forall x \in U^{-1}((0, 2\sigma]).$$

Definition 3.1. For a given $\sigma > 0$, let $\gamma(\cdot)$, and $p(\cdot)$ be chosen as above. A selection

$$\mathbf{v}: U^{-1}((0, 2\sigma]) \to \mathbb{R}^n, \qquad x \mapsto \mathbf{v}(x) \in \mathcal{F}^{(k)}(x)$$

(14)

⁹The inequality (14) is usually formulated with the proximal superdifferential $\partial^P F$. However, this does not make a difference here since $\partial^P F = \partial_C F = coD^*F$ as soon as F is locally semiconcave. Hence (14) is true in particular for D^*U .

is called a degree-k feedback (corresponding to U, σ , $\gamma(\cdot)$, and $p(\cdot)$) if for every $x \in U^{-1}((0, 2\sigma])$ there exists a positive integer $h \leq k$ such that

(15)
$$\begin{cases} \mathbf{v}(x) \in \mathcal{F}^{(h)}(x), \\ \left\langle p(x), \mathbf{v}(x) \right\rangle \leq -\gamma(U(x)), \\ and, \text{ if } h > 1: \quad H^{(h-1)}(x, p(x)) > -\gamma(U(x)). \end{cases}$$

The number h will be called the degree of the feedback \mathbf{v} at x.

Let us momentarily assume that there exists $M \ge 0$ such that

(16)
$$||f_i||_{k-1} \le \widehat{M} \qquad \forall i = 1, \dots, m$$

in the whole set $\mathbb{R}^n \setminus \mathcal{T}$, which, in view of the compactness of the control set A, implies that there is $M \geq 0$ verifying

$$(17) ||X||_0 \le M$$

for any iterated bracket X in $\mathcal{F}^{(k)}$. Under this assumption one can regard each vector $\mathbf{v}(x)$ as a *tangent* vector to a curve that is a suitable composition of flows, as stated in the following result (see e.g. [FR1], [FR2]):

Lemma 3.2. Under assumption (16) there exists a real constant c > 0such that for any $x \in \mathbb{R}^n \setminus \mathcal{T}$, any feedback $\mathbf{v}(\cdot)$ of degree h at x, and any t > 0, one can find a control $\alpha_t : [0, t] \to A$ such that

(i)
$$\alpha_t(\cdot)$$
 is constant on intervals $\left\lfloor \frac{jt}{r}, \frac{(j+1)t}{r} \right\rfloor$, $j = 0, \dots, r-1$;

(ii) the estimate

(18)
$$\left| y_x(t,\alpha_t) - x - \frac{\mathbf{v}(x)}{r^h} t^h \right| \le \frac{c}{r^h} t^{h+1}$$

holds true, where r is an integer depending on the formal Lie bracket corresponding to $\mathbf{v}(x)$ and is increasing with the degree.

For instance, r = 1, 4, 10 if h = 1, 2, 3, respectively. In particular, if $\mathbf{v}(x) = [[f_1, f_2], f_3](x)$ one sets

$$\alpha_t(s) := \begin{cases} e_1 & \text{if } s \in [0, t/10) \cup [6t/10, 7t/10) \\ e_2 & \text{if } s \in [t/10, 2t/10) \cup [5t/10, 6t/10) \\ e_3 & \text{if } s \in [4t/10, 5t/10) \\ - & e_1 & \text{if } s \in [2t/10, 3t/10) \cup [8t/10, 9t/10) \\ - & e_2 & \text{if } s \in [3t/10, 4t/10) \cup [7t/10, 8t/10) \\ - & e_3 & \text{if } s \in [9t/10, t). \end{cases}$$

Let us point out that one can have different r's for feedbacks having the same degree 10.

¹⁰Precisely: For each formal bracket B, the corresponding r = r(B) is defined recursively: one sets r(B) = 1 if B has degree 1, while, if $B = [B_1, B_2]$ and $r_1 = r(B_1)$,

3.3. A step of degree $h \leq k$. Now let us choose $z \in U^{-1}((0, \sigma])$ and a feedback **v** of degree k (surely existing by (10)). Let the feedback **v** have degree h at z. We shall rely on the following result.

Claim 3.1. Let $U(\cdot)$, σ , $\gamma(\cdot)$ and $p(\cdot)$ as above. Furthermore let $\mathbf{v}(\cdot)$ be a degree-k feedback corresponding to these data. Then there exists a time-valued function

$$\tau: (0,\sigma] \times \{1,\ldots,k\} \to (0,1],$$

such that

- i) $j \mapsto \tau(u,j)^j$ and $j \mapsto \tau(u,j)^{j-1}$ are decreasing for every $u \in]0,\sigma]$,
- ii) $u \mapsto \tau(u, j)$ is increasing for every $j \in \{1, \ldots, k\}$, and

iii) for all $z \in U^{-1}((0,\sigma])$ with a feedback $\mathbf{v}(\cdot)$ of degree h at z, one has

$$U(y_z(t,\alpha_t)) - U(z) \le -\frac{\gamma(U(z))}{2} \left(\frac{t}{r}\right)^h \quad \forall t \in [0, \, \tau(U(z), h)] ,$$

where r and $y_z(\cdot, \alpha_t)$ are an integer and a trajectory associated to $\mathbf{v}(z)$ as in Lemma 3.2.

Proof. Let $\nu > 0$ be such that $U^{-1}((0, 2\sigma]) \subset B\left(\mathcal{T}, \frac{\nu}{2}\right)$ and fix $z \in U^{-1}((0, \sigma])$ with a feedback $\mathbf{v}(\cdot)$ of degree h at z. To begin with we wish to choose a time $\overline{\tau}$ such that, for any $t \in [0, \overline{\tau}]$;

- i) $y(t) \in B(\mathcal{T}, \nu)$ for any system's trajectory $y(\cdot)$ issuing from a point of $U^{-1}((0, \sigma])$;
- ii) $y_z^t(t) \in B\left(z, \frac{\mathbf{d}(z)}{2}\right)$ for any trajectory $y_z^t(\cdot) := y_z(\cdot, \alpha_t)$ associated to $\mathbf{v}(z)$ as in Lemma 3.2.

For this purpose, it is clearly sufficient to set

$$\bar{\tau}(u,j) := \min\left\{\frac{\nu}{2M}, \sqrt[j]{\frac{\mathbf{d}(U^{-1}(u))}{2M}}\right\} \quad \forall (u,j) \in (0,\sigma] \times \{1,\dots,k\},$$

and to choose

$$\bar{\tau} := \bar{\tau}(U(z), h).$$

Because of $\mathbf{d}(U^{-1}(U(z))) \leq \mathbf{d}(z)$ and (18), the distance $\mathbf{d}([z, y_z^t(t)])$ between the segment $[z, y_z^t(t)]$ and the target \mathcal{T} verifies

$$\mathbf{d}([z, y_z^t(t)]) \ge \frac{\mathbf{d}(z)}{2} \ge \frac{\mathbf{d}(U^{-1}(U(z)))}{2},$$

for every $t \in [0, \bar{\tau}]$. For every $u \in (0, \sigma]$, in relation with the compact set

$$\mathcal{K}(u) := \left\{ x : \frac{\mathbf{d}(U^{-1}(u))}{2} \le \mathbf{d}(x) \le \nu \right\},\$$

 $r_2 = r(B_2)$ one sets $r(B) := 2(r_1 + r_2)$. For instance, $r([g_1, g_2]) = 4$, $r([g_1, [g_2, g_3]]) = 10$, $r([g_1, [g_2, [g_3, g_4]]]) = 22$ and $r([[g_1, g_2], [g_3, g_4]] = 16$.

let L(u) and $\rho(u)$ be a Lipschitz continuity and a semiconcavity constant, whose existence is stated in Lemma 3.1. Let us set L := L(U(z)) and $\rho := \rho(U(z))$. By (18), for any $t \in [0, \overline{\tau}]$ we get

$$U(y_{z}^{t}(t)) - U(z) \leq \left\langle p(z), y_{z}^{t}(t) - z \right\rangle + \rho \left| y_{z}^{t}(t) - z \right|^{2} \leq \left\langle p(z), \mathbf{v}(z) \right\rangle \left(\frac{t}{r} \right)^{h} + |p(z)|ct \left(\frac{t}{r} \right)^{h} + \rho \left(\frac{t}{r} \right)^{2h} (M + ct)^{2} \leq -\gamma(U(z)) \left(\frac{t}{r} \right)^{h} + |p(z)|ct \left(\frac{t}{r} \right)^{h} + \rho \left(\frac{t}{r} \right)^{2h} (M + ct)^{2} \leq \left(-\gamma(U(z)) + Lct + \rho \left(\frac{t}{r} \right)^{h} (M + ct)^{2} \right) \left(\frac{t}{r} \right)^{h}$$

Let us observe that $\left(\frac{t}{r}\right)^h \leq t \leq 1$ as soon as $t \leq 1$. Therefore, if we define, for every u,

(20)
$$\check{\tau}(u) := \frac{\gamma(u)}{2[L(u)c + \rho(u)(M+c)^2]},$$

and we set

(21)
$$\tau(u,j) := \min\{1, \bar{\tau}(u,j), \check{\tau}(u)\} \quad \forall (u,j) \in (0,\sigma] \times \{1, \dots, k\},$$

we get

$$Lct + \rho\left(\frac{t}{r}\right)^h (M + ct)^2 \le t[Lc + \rho(M + c)^2] \le \frac{\gamma(U(z))}{2}.$$

as soon as $t \in [0, \tau], \tau := \tau(U(z), h).$

Therefore, with this choice of τ we obtain

(22)
$$U(y_z^t(t)) - U(z) \le -\frac{\gamma(U(z))}{2} \left(\frac{t}{r}\right)^h \quad \forall t \in [0,\tau].$$

Moreover, $u \mapsto \tau(u, j)$ is increasing for every j: indeed, by Lemma 3.1, the constants L(u) and $\rho(u)$ turn out to be decreasing in u. Finally, the fact that $j \mapsto \tau(u, j)^j$ and $j \mapsto \tau(u, j)^{j-1}$ are decreasing is an easy consequence of the definition of $\tau(u, j)$ in (21). The claim is now proved.

3.4. Piecewise C^1 trajectories approaching the target. Now, let us define recursively a sequence of times $(t_j)_{j\geq 0}$, of trajectory-control pairs $(y_j(\cdot), \alpha_j(\cdot)) : [s_{j-1}, s_j] \to \mathbb{R}^n \times A, j \geq 1, s_0 := 0, s_j := s_{j-1} + t_i$, and points x_j as follows:

- $t_0 := s_0 = 0, \ x_1 := x;$
- if $j \ge 1$, $t_j := \tau(U(x_j), h_j)$, where h_j is the degree of the feedback **v** at x_j and $\tau(\cdot, \cdot)$ is as in Claim 3.1;
- $(y_1, \alpha_1) : [s_0, s_1] \to I\!\!R^n \times A$ is the trajectory-control pair defined as $(y_1, \alpha_1) := (y_{x_1}^{t_1}, \alpha_{t_1});$

• for every j > 1, $y_j(s_{j-1}) := y_{j-1}(s_{j-1}) := x_j$, and the pair $(y_j(\cdot), \alpha_j(\cdot)) : [s_{j-1}, s_j] \to \mathbb{R}^n \times A$ is given by $(y_j, \alpha_j) := (y_{x_j}^{t_j}, \alpha_{t_j})(s - s_{j-1})$ for every $s \in [s_{j-1}, s_j]$.

Let us consider the real sequence

$$u_j := U(x_j) \qquad j \in \mathbb{N}$$

and let us show that

$$\lim_{j \to \infty} u_j = 0.$$

Indeed, the degree h_j of the feedback **v** at every x_j is bounded by k. Moreover, if we use r_j to denote the positive integer appearing in formula (18) in relation with the feedback **v** at x_j , we get

$$r_j^{h_j} \le (r(k))^k,$$

if we set

$$r(k) := \max\{r_j, \ j \in \mathbb{N}\}^{11}$$

Therefore, by Claim 3.1 we obtain

(23)
$$u_{j+1} - u_j = U(x_{j+1}) - U(x_j) \le -\frac{\gamma(u_j)}{2} \left(\frac{\tau(u_j, h_j)}{r_j}\right)^{n_j} \le -\frac{\gamma(u_j)}{2} \left(\frac{\tau(u_j, k)}{r(k)}\right)^k < 0$$

for all $j \ge 1$. Hence the sequence (u_j) is positive and decreasing, so there exists the limit

$$\lim_{j \to \infty} u_j = \eta \ge 0.$$

Let us show that $\eta = 0$. If, on the contrary, η were strictly positive, by Claim 3.1 one would have $\lim_{j\to\infty} \tau(u_j, k) \ge \tau(\eta, k) > 0$. Hence taking the limit in (23) one would obtain

$$0 = \eta - \eta \le -\lim_{j \to \infty} \frac{\gamma(u_j)\tau^k(u_j, k)}{2(r(k))^k} \le -\frac{\gamma(\eta)\tau^k(\eta, k)}{2(r(k))^k} < 0,$$

a contradiction. Therefore

$$\lim_{j \to \infty} U(x_j) = \lim_{j \to \infty} u_j = 0.$$

Hence, setting

$$S := \lim_{j \to \infty} s_j = \sum_{i=1}^{\infty} t_i$$

and

$$(y,\alpha)(s) := (y_j,\alpha_j)(s) \quad \forall j \ge 1, \ \forall s \in [s_{j-1},s_j],$$

one finds that

$$\lim_{j \to \infty} \mathbf{d} \left(y(s_j) \right) = 0.$$

16

¹¹This maximum clearly exists and depends (monotonically) on k. For instance. $r(2) = 4, r(3) = 10, r(4) = \max\{22, 16\} = 22.$

Actually the stronger limit relation

$$\lim_{s \to S^-} \mathbf{d} \left(y(s) \right) = 0$$

holds, as it follows from the construction of the function β below.

3.5. Construction of a bounding $\mathcal{K}L$ function. In order to conclude the proof that the system is GAC to \mathcal{T} , we have to establish the existence of a $\mathcal{K}L$ function β such that $\mathbf{d}(y(s)) \leq \beta(\mathbf{d}(y(0)), s)$ for every $s \geq 0$, as in Definition 1.2.

By Claim 3.1, for any $t_j = \tau(u_j, h_j)$ one has $t_j^{h_j-1} \ge \tau^{k-1}(u_j, k)$. Moreover, as already remarked, $(r_j)^{h_j} \le (r(k))^k$ (recall that we are using r_j to denote the positive integer appearing in formula (18) in relation with the feedback **v** at x_j). Hence, for any $j \ge 1$, we have:

$$U(y_j(s_j)) - U(y_j(s_{j-1})) = u_{j+1} - u_j$$

$$\leq -\frac{\gamma(u_j)}{2} \left(\frac{t_j}{r_j}\right)^{h_j} \leq -\frac{\gamma(u_j)\tau^{k-1}(u_j,k)}{2(r(k))^k} t_j$$

Let us define the function $\tilde{\gamma}: (0, \sigma] \to \mathbb{R}$ by setting

(24)
$$\tilde{\gamma}(u) := \frac{\gamma(u)\,\tau^{k-1}(u,k)}{2(r(k))^k}$$

Clearly, by the monotonicity of $u \mapsto \tau(u,k)$, $\tilde{\gamma}$ is (positive and) strictly increasing. Therefore, since $U(y(s_j)) \leq U(y(s_i))$ for every $i = 1, \ldots, j$, we get

$$U(y(s_j)) - U(z) = [U(y(s_j)) - U(y(s_{j-1}))] + [U(y(s_{j-1})) - U(y(s_{j-2}))] + \dots$$

 $+[U(y(s_1)) - U(y(0))] \le -\sum_{i=1}^{j} \tilde{\gamma}(U(y(s_i))) [s_i - s_{i-1}] \le -\tilde{\gamma}(U(y(s_j))) s_j.$

In particular, we have

(25)
$$U(y(s_j)) + \tilde{\gamma}(U(y(s_j))) s_j \le U(z).$$

We now replace the function $\tilde{\gamma}$ with the slightly different function $\hat{\gamma}$: $[0, +\infty) \rightarrow [0, +\infty)$ defined by $\hat{\gamma}(u) \doteq \min\{u, \tilde{\gamma}(u)\}$ for all $u \in [0, +\infty)$. Notice that $\hat{\gamma}$ is continuous, strictly increasing and $\hat{\gamma}(u) > 0 \quad \forall u > 0$, $\hat{\gamma}(0) = 0$. Then, for any $j \ge 1$,

$$\hat{\gamma}(U(y(s_j)))(1+s_j) \le U(z),$$

so that

(26)
$$U(y(s_j)) \le \hat{\gamma}^{-1} \left(\frac{U(z)}{1+s_j} \right)$$

Let $\delta_{-}, \delta_{+}: [0, +\infty) \to [0, +\infty)$ be the continuous, strictly increasing, unbounded functions defined by

(27)
$$\delta_{-}(u) \doteq \min\{\mathbf{d}(x): U(x) \ge u\}, \quad \delta_{+}(u) \doteq \max\{\mathbf{d}(x): U(x) \le u\}$$

and let us set $\hat{\delta}_{-}(u) := \min\{\delta_{-}(u), u\}$. Notice that $\hat{\delta}_{-}(0) = \delta^{+}(0) = 0$, and $\hat{\delta}_{-}(U(x)) \leq \mathbf{d}(x) \leq \delta^{+}(U(x))$,

 $\forall x \in U^{-1}((0,\sigma])$. Therefore, setting

(28)
$$\hat{\beta}(\delta,s) := \delta_+ \circ \hat{\gamma}^{-1} \left(\frac{\hat{\delta}_-^{-1}(\delta)}{1+s} \right) \quad \forall (\delta,s) \in [0,+\infty) \times [0,+\infty),$$

by (26) we get

(29)
$$\mathbf{d}(y(s_j)) \le \delta_+ \left(U(y(s_j)) \le \delta_+ \left(\hat{\gamma}^{-1} \left(\frac{U(z)}{1+s_j} \right) \right) \le \hat{\beta}(\mathbf{d}(z), s_j),$$

for every $j \geq 1$. The estimate (29) says that the function $\hat{\beta}$ bounds the distance of the trajectory $y(\cdot)$ from the target \mathcal{T} at the discrete times s_j . Hence, in order to get a bound at all times, we need to slightly modify $\hat{\beta}$. For this purpose, given any $x \in \mathbb{R}^n \setminus \mathcal{T}$, let us select a point $\pi(x) \in \mathcal{T}$ such that $\mathbf{d}(x) = |x - \pi(x)|$. Notice that for any $s \in [s_j, s_{j+1}]$, one has

$$\mathbf{d}(y(s)) \le |y(s) - \pi(y(s_j))| \le |y(s) - y(s_j)| + |y(s_j) - \pi(y(s_j))|$$

$$\leq M[s_{j+1} - s_j] + \mathbf{d}(y(s_j)).$$

Furthermore, by the definition of τ (see Claim 3.1) it follows that

(30)
$$s_{j+1} - s_j = t_{j+1} \le \tau(u_{j+1}, k) \le \tau(\delta_-^{-1}(\mathbf{d}(y(s_j))), k).$$

Therefore,

$$\begin{aligned} \mathbf{d}(y(s)) &\leq M\tau(\delta_{-}^{-1}(\mathbf{d}(y(s_j))), k) + \mathbf{d}(y(s_j)) \\ &\leq M\tau(\delta_{-}^{-1}(\hat{\beta}(\mathbf{d}(z), s_j)), k) + \hat{\beta}(\mathbf{d}(z), s_j) \,. \end{aligned}$$

Since for all δ the function $s \mapsto \hat{\beta}(\delta, s)$ is decreasing, one obtains

$$\mathbf{d}(y(s)) \le \beta(\mathbf{d}(z), s) \qquad \forall s \in [0, +\infty[,$$

where we have set, for all $s \in [0, \tau(\delta_{-}^{-1}(\delta), k)],$

$$\beta(\delta,s) := M\tau(\delta_{-}^{-1}(\hat{\beta}(\delta,0)),k) + \hat{\beta}(\delta,0)$$

and, if $s > \tau(\delta_{-}^{-1}(\delta), k)$,

$$\beta(\delta,s) := M\tau(\delta_-^{-1}(\hat{\beta}(\delta,s-\tau(\delta_-^{-1}(\delta),k))),k) + \hat{\beta}(\delta,s-\tau(\delta_-^{-1}(\delta),k)).$$

3.6. Removal of the fictitious C^{k-1} -bound. Let us see now that, by means of a cut-off argument, we can remove the auxiliary boundedness hypothesis (17). Let $\psi : \mathbb{R}^n \to [0,1]$ be a C^{∞} map such that

(31)
$$\psi = 1$$
 on $\overline{B(\mathcal{T}, \nu) \setminus \mathcal{T}}$, $\psi = 0$ on $\mathbb{R}^n \setminus B(\mathcal{T}, 2\nu)$

and consider the control system

(32)
$$\xi' = \sum_{i=1}^{m} a_i \; (\psi f_i(\xi))$$

Notice that the functions (ψf_i) belong to $C_b^{k-1}(\mathbb{R}^n \setminus \mathcal{T})$ because of the cut-off factor ψ . Moreover, any trajectory $\xi(\cdot)$ of (32) with the initial condition $z \in U^{-1}((0,\sigma])$, exists globally and cannot exit the compact set $\overline{B(\mathcal{T}, 2\nu) \setminus \mathcal{T}}$. Owing to the previous step, there exists a trajectory ξ which approaches asymptotically the target and verifies $\mathbf{d}(\xi(s)) \leq \beta(\mathbf{d}(z), s) \quad \forall s \in [0, +\infty[$. Moreover, $\xi(s)$ belongs to $B(\mathcal{T}, \nu)$ for every $s \geq 0$. Therefore ξ is a solution of the original system, proving that (3) has the GAC property in $U^{-1}((0, \sigma])$.

By the arbitrariness of $\sigma > 0$, it is easy to extend these constructions from $U^{-1}((0,\sigma])$ to the whole set $\mathbb{R}^n \setminus \mathcal{T}$. This concludes the proof of (2.1). \Box

4. The case of nonsmooth dynamics

Let us begin with an example, where the vector fields are not C^1 .

Example 4.1. Let us consider the system

(33)
$$\dot{y} = a_1 f_1(y) + a_2 f_2(y),$$

where

$$f_1 := \frac{\partial}{\partial x_1} + (|x_2| - 2x_2)\frac{\partial}{\partial x_3}, \quad f_2 := \frac{\partial}{\partial x_2} + (|x_1| + 2x_1)\frac{\partial}{\partial x_3},$$

and let the target \mathcal{T} coincide with the origin. For the same reason as in the nonholonomic integrator, a smooth degree-1 CLF does not exist (see Example 2.1). On the other hand, the given notion of degree-2 control Lyapunov function is not even meaningful here, in that the classical bracket $[f_1, f_2]$ is not defined at points x such that $x_1 = 0$ or $x_2 = 0$. Yet, in the open, dense set $\{x : x_1 \neq 0, x_2 \neq 0\}$ the bracket is well defined and, furthermore, the Lie algebra rank condition is verified. So it is reasonable to look for a Lyapunov-like condition involving *somehow* Lie brackets.

One one hand, in the definition of degree-k CLF one requires the vector fields f_1, \ldots, f_m to be of class C^{k-1} (see also Remark 2.1), for this guarantees that the Lie brackets up to the degree k are well defined and continuous. On the other hand, as remarked in the Introduction, the non smoothness of a degree-1 CLF is more related to a shortage of dynamics' directions than to the regularity of the involved vector fields.

Let us introduce the notion of set-valued bracket for locally Lipschitz vector fields.

Definition 4.1 ([RS1]). Let $\Omega \subset \mathbb{R}^n$ be an open set and let f, g be vector fields belonging to $C^{0,1}(\Omega)$. For every $x \in \Omega$, let us set

$$[f,g]_{set}(x) :=$$

$$co\Big\{v \in \mathbb{R}^n, v = \lim_{x_n \to x} [f, g](x_n) \quad (x_n)_{n \in \mathbb{N}} \subset DIFF(f) \cap DIFF(g)\Big\},\$$

where co means "convex hull", and DIFF(f), DIFF(g) denote the subsets of differentiability points of f and g, respectively. Let us observe that DIFF(f) and DIFF(g) have full measure, hence they are dense in Ω . Notice that, as in the regular case, one has $[f, f]_{set}(x) = \{0\}$ and $[f, g]_{set}(x) = -[g, f]_{set}(x)$ for every $x \in \Omega$ (for any $E \subset \mathbb{R}^n$ we use the notation $-E = \{-v : v \in E\}$).

Let us consider the control system

(34)
$$\begin{cases} \dot{y} = \sum_{i=1,\dots,m} a_i f_i(y) \\ y(0) = x \in I\!\!R^n \setminus \mathcal{T} \end{cases}$$

and let us assume that f_1, \ldots, f_m belong to $C_b^{0,1}(\Omega \setminus \mathcal{T})$ for any bounded, open set $\Omega \subset \mathbb{R}^n$ (see Subsection 1.1).

The families $\mathcal{F}^{(1)}$ and $\mathcal{F}^{(2)}$ are formally defined as in the regular (i.e., C^1) case, except that their elements are set-valued vector fields ¹²:

$$\mathcal{F}^{(1)} := \left\{ \{ f_{\ell}(\cdot) \}, \{ -f_{\ell}(\cdot) \} : \quad \ell = 1, \dots, m \right\},\$$

and

$$\mathcal{F}^{(2)} := \mathcal{F}^{(1)} \cup \Big\{ [f_i, f_j]_{set}(\cdot) : \quad i, j = 1, \dots, m \Big\}.$$

As in the regular case, for every $x \in \mathbb{R}^n \setminus \mathcal{T}$, we set

$$\mathcal{F}^{(1)}(x) := \left\{ \left\{ f_{\ell}(x) \right\}, \left\{ -f_{\ell}(x) \right\} : \quad \ell = 1, \dots, m \right\},\$$

and

$$\mathcal{F}^{(2)}(x) := \mathcal{F}^{(1)}(x) \cup \Big\{ [f_i, f_j]_{set}(x) : \quad i, j = 1, \dots, m \Big\}.$$

Accordingly, for h = 1, 2, we define the *degree-h* Hamiltonian $H^{(h)}$ by setting

$$H^{(h)}(x,p) := \inf_{v \in \mathcal{F}^{(h)}(x)} \sup_{w \in v} \left\langle p, w \right\rangle,$$

for all $(x,p) \in (\mathbb{I} \mathbb{R}^n \setminus \mathcal{T}) \times \mathbb{I} \mathbb{R}^n$. More explicitly, one has

$$H^{(1)}(x,p) = \inf \{ - |\langle p, f_{\ell}(x) \rangle | : \ell = 1, \dots, m \},\$$

and

$$H^{(2)}(x,p) = \inf_{\ell,i,j} \left\{ -\left| \left\langle p, f_{\ell}(x) \right\rangle \right|, \sup_{w \in [f_i,f_j]_{set}(x)} \left\langle p, w \right\rangle : \quad \ell, i, j = 1, \dots, m \right\}.$$

We can now state, for the case k = 2, a generalization of Theorem 2.1 to the nonsmooth case:

Theorem 4.1. Let us assume that a degree-2 control Lyapunov function exists. Then system (34) is GAC to \mathcal{T} .

20

 $^{{}^{12}\}mathcal{F}^{(1)}$ practically coincides with the one in the regular case, for its elements are setvalued maps which are singletons, indeed.

Example 4.1 (continued). Let us come back to the system in (33) and compute the bracket $[f_1, f_2]_{set}$. It turns out that

$$[f_1, f_2]_{set}(x) = I(x)\frac{\partial}{\partial x_3},$$

where

$$I(x) := \begin{cases} \{4\} & \text{if } x_1 x_2 > 0, \\ \{2\} & \text{if } x_1 < 0 \text{ and } x_2 > 0, \\ \{6\} & \text{if } x_1 > 0 \text{ and } x_2 < 0, \\ [2,4] & \text{if either } x_1 = 0 \text{ and } x_2 > 0, \text{ or } x_1 < 0 \text{ and } x_2 = 0, \\ [4,6] & \text{if either } x_1 = 0 \text{ and } x_2 < 0, \text{ or } x_1 > 0 \text{ and } x_2 = 0, \\ [2,6] & \text{if } x_1 = x_2 = 0. \end{cases}$$

The distance function U(x) = |x| is not a degree-1 CLF. Indeed, by

$$H^{(1)}(x,p) = \inf \left\{ -|p_1 + p_3(|x_2| - 2x_2)|, -|p_2 + p_3(|x_1| + 2x_1)| \right\}$$

one obtains

$$H^{(1)}((0,0,x_3), DU(0,0,x_3)) = 0$$
 for every $x_3 \neq 0$.

Yet, the distance function U happens to be a (C^{∞}) degree-2 CLF. Indeed,

$$H^{(2)}(x,p) = \inf \left\{ H^{(1)}(x,p), \sup_{w \in I(x)} w p_3, \sup_{w \in -I(x)} w p_3 \right\} = -\sup \left\{ |p_1 + p_3(|x_2| - 2x_2)|, |p_2 + p_3(|x_1| + 2x_1)|, 2|p_3| \right\}$$

and, since |DU(x)| = 1 for every $x \neq 0$, one gets

(35)
$$H^{(2)}(x, DU(x)) < 0.$$

Notice that, for the validity of the strict inequality in (35), it is crucial that $0 \notin [f_1, f_2]_{set}(x)$ for every $x \neq 0$. Furthermore, arguing as in Remark 2.4, we know that a (possibly nonsmooth) degree-1 CLF does exist. Actually, the function

$$U(x) = \max\left\{\sqrt{x_1^2 + x_2^2}, |x_3| - \sqrt{x_1^2 + x_2^2}\right\}$$

introduced in [Ri] as a control Lyapunov function for the nonholonomic integrator is a degree-1 CLF also for this system.

4.1. **Proof of Theorem 4.1.** The proof is akin to the proof of Theorem 2.1. Yet, because of the new kind of brackets and Hamiltonians here involved, some changes are needed.

As in the regular case, the 0 in the dissipative relation can be replaced by a nonnegative function of U:

Proposition 4.1. Let $U : \overline{\mathbb{R}^n \setminus \mathcal{T}} \to \mathbb{R}$ be a continuous function, such that U is locally semiconcave, positive definite and proper on $\mathbb{R}^n \setminus \mathcal{T}$. Then the conditions (i) and (ii) below are equivalent:

(i) U verifies

$$H^{(2)}(x, D^*U(x)) < 0 \quad for \ all \ x \in \mathbb{R}^n \setminus \mathcal{T};$$

(ii) for every σ > 0 there exists a continuous, strictly increasing, function γ: [0, +∞) →: [0, +∞) such that

$$H^{(2)}(x, D^*U(x)) \le -\gamma(U(x))$$
 for all $x \in U^{-1}((0, 2\sigma])$.

Proof. By [RS1], for every i, j, the set-valued map $x \mapsto [f_i, g_j]_{set}(x)$ is upper semicontinuous, with compact, convex values. Moreover, basic results on marginal functions imply that $(x, p) \mapsto \sup_{w \in [f_i, g_j]_{set}(x)} \langle p, w \rangle$ is upper semicontinuous (see e.g. [AC]). As an easy consequence, the Hamiltonian

$$(x,p) \mapsto H^{(2)}(x,p)$$

turns out to be upper semicontinuous. At this point one can conclude, arguing exactly as in Proposition 3.1. $\hfill \Box$

We now need to adapt the notion of degree-2 feedback to the case when Lie brackets are set-valued. For a given $\sigma > 0$, let γ be a function as in Proposition 4.1, and let $x \mapsto p(x)$ be a selection of $x \mapsto D^*U(x)$ on $U^{-1}((0, 2\sigma])$, so that

$$H^{(2)}(x, p(x)) \le -\gamma(U(x)) \qquad \forall x \in U^{-1}((0, 2\sigma]).$$

Definition 4.2. For a given $\sigma > 0$, let $\gamma(\cdot)$, and $p(\cdot)$ be chosen as above. A selection

$$\mathbf{v}: U^{-1}((0, 2\sigma]) \to 2^{\mathbb{R}^n}, \qquad x \mapsto \mathbf{v}(x) \in \mathcal{F}^{(2)}(x)$$

is called a degree-2 feedback (corresponding to U, σ , γ , and $p(\cdot)$) if for every $x \in U^{-1}((0, 2\sigma])$ there exists $h \in \{1, 2\}$ such that

(36)
$$\begin{cases} \mathbf{v}(x) \in \mathcal{F}^{(h)}(x), \\ \sup_{w \in \mathbf{v}(x)} \left\langle p(x), w \right\rangle \leq -\gamma(U(x)), \\ and, if h = 2, \quad H^{(1)}(x, p(x)) > -\gamma(U(x)). \end{cases}$$

The number h will be called the degree of the feedback \mathbf{v} at x.

More explicitly, when h = 1, for some $\ell = 1, \ldots, m$ one has

$$\begin{cases} \mathbf{v}(x) = \{f_{\ell}(x)\} \text{ or } \mathbf{v}(x) = -\{f_{\ell}(x)\} \\ -\left|\left\langle p(x), f_{\ell}(x)\right\rangle\right| \leq -\gamma(U(x)); \end{cases}$$

22

if h = 2, for some i, j = 1, ..., m $(i \neq j)$ one has

$$\begin{cases} \mathbf{v}(x) = [f_i, f_j]_{set}(x), \\ \left\langle p(x), w \right\rangle \le -\gamma(U(x)) \quad \forall w \in [f_i, f_j]_{set}(x), \\ \text{and} \quad H^{(1)}(x, p(x)) > -\gamma(U(x)). \end{cases}$$

The following claim is a version of Claim 3.1 adapted to the set-valued notion of feedback:

Claim 4.1. Let $U(\cdot)$, σ , $\gamma(\cdot)$ and $p(\cdot)$ as above. Furthermore let $\mathbf{v}(\cdot)$ be a (set-valued) degree-2 feedback corresponding to these data. Then there exists a time-valued function

$$\tau: (0,\sigma] \times \{1,2\} \to (0,1],$$

such that

- i) $j \mapsto \tau(u,j)^j$ and $j \mapsto \tau(u,j)^{j-1}$ are decreasing for every $u \in (0,\sigma]$,
- ii) $u \mapsto \tau(u, j)$ is increasing for every $j \in \{1, 2\}$, and
- iii) for all $z \in U^{-1}((0,\sigma])$ with a feedback $\mathbf{v}(\cdot)$ of degree h at z, one has

$$U(y_z(t,\alpha_t)) - U(z) \le -\frac{\gamma(U(z))}{2} \left(\frac{t}{r}\right)^h \quad \forall t \in [0, \, \tau(U(z), h)] ,$$

where r and $y_z(\cdot, \alpha_t)$ are an integer and a trajectory associated to $\mathbf{v}(z)$ according to Lemma 4.1.

The proof of Claim 3.1 was based the on the asymptotic formulas stated in Lemma 3.2. Similarly, Claim 4.1 results proved as soon as one applies the asymptotic formula stated in Lemma 4.1 below. Precisely, once arrived to formula (19), one simply replaces estimate (18) with (42), and observes that, by the definition of $H^{(2)}$, the inequality $\langle p(z), w \rangle \leq -\gamma(U(z))$ is verified for all $w \in [f_i, f_j]_{set}(z)$ (see Definition 4.2).

Lemma 4.1 ([RS2], [FR2]). If $f_1, \ldots, f_m \in C_b^{0,1}(\mathbb{R}^n \setminus \mathcal{T})$, there exists a constant c > 0 such that for any $x \in \mathbb{R}^n \setminus \mathcal{T}$, any feedback $\mathbf{v}(x) = [f_i, f_j]_{set}(x)$, and any t > 0, setting

$$\alpha_t(s) = e_i \chi_{[0,t/4[}(s) + e_j \chi_{[t/4,t/2[}(s) - e_i \chi_{[t/2,3t/4[}(s) - e_j \chi_{[3t/4,t]}(s), \quad s \in [0,t]],$$

the estimate

(37)
$$d\left(y_x(t,\alpha_t) - x, [f_i, f_j]_{set}(x) \frac{t^2}{16}\right) \le \frac{c}{16} t^3$$

holds true. In particular, for any t > 0 there exists some $w(t) \in \mathbf{v}(x)$ such that

(38)
$$\left| y_x(t,\alpha_t) - x - w(t) \frac{t^2}{16} \right| \le \frac{c}{16} t^3.$$

M. MOTTA AND F. RAMPAZZO

Let us point out that Claim 3.1 is the starting point for the construction of an admissible trajectory-control pair by means of the recursive procedure described in the proof of Theorem 2.1. Through exactly the same arguments and in view of Claim 4.1, the proof of Theorem 4.1 can now be completed.

5. Concluding Remarks

5.1. Feedback constructions. Degree-1 control Lyapunov functions are used as primary ingredient for the construction of feedback stabilizing strategies, a classical question that is mainly concerned with the definition of an appropriate notion of solution for discontinuous ODE's (see e.g. [CLSS], [CLRS], [MaRS] and [AB]).

It might be interesting to associate some concept of feedback strategy also in relation with a degree-k control Lyapunov function, k > 1, all the more so as it may happen to be smoother than the degree-1 CLF. As a matter of fact, the proofs of Theorems 2.1 and 4.1 seem to suggest a notion of "feedback" such that, in dependence of the bracket that minimizes the Hamiltonian $H^{(2)}$, singles out a suitable finite sequence of constant controls to be implemented along small time intervals.

5.2. **Degree-**k **CLF as viscosity supersolutions.** One is obviously tempted to refer to a degree-k control Lyapunov function as a *strict supersolution* of the Hamilton-Jacobi equation

$$-H^{(k)}(x, DU(x)) = 0.$$

Actually this holds true as soon as we consider, for instance, the notion of viscosity solution. More precisely, one has:

Let $U: \mathbb{R}^n \setminus \mathcal{T} \to \mathbb{R}$ be a continuous function which, furthermore, is locally semiconcave, positive, and proper on $\mathbb{R}^n \setminus \mathcal{T}$. Then U is a degree-k control Lyapunov function if and only for any N > 0 there is some continuous, strictly increasing function $\gamma: [0, +\infty) \to [0, +\infty)$ such that U is a viscosity supersolution ¹³ of

(39)
$$-H^{(k)}(x, DU(x)) = \gamma(U(x)) \quad in \ U^{-1}((0, N)).$$

Indeed, in the case of a locally semiconcave function U, at every $x \in DIFF(U)$ the subdifferential $D^-U(x) = \{\nabla U(x)\}$ coincides with $D^*U(x)$, while $D^-U(x)$ is empty if $x \notin D^-U(x)$. Therefore, thanks to Proposition 3.1, (39) follows from the inequality

(40)
$$H^{(k)}(x, D^*U(x)) < 0,$$

which defines the notion of degree-k CLF. To obtain the converse implication, for any $x \in U^{-1}((0, N))$ and $p \in D^*U(x)$, let $(x_n)_n \subset U^{-1}((0, N)) \cap$

¹³Namely, $-H^{(k)}(x,p) \ge \gamma(U(x))$ for all $x \in \mathbb{R}^n \setminus \mathcal{T}$ and p in the subgradient $DU^-(x)$ of U at x (see [BCD]).

DIFF(U) be such that $\lim_{n \to \infty} (x_n, \nabla U(x_n)) = (x, p)$. Then, by hypothesis (39), one has

$$-H^{(k)}(x_n, \nabla U(x_n)) \ge \gamma(U(x_n)) \qquad \forall n \in \mathbb{N},$$

so, passing to the limit, one gets (40).

5.3. Generalizations to larger classes of systems. As it is well known, the lack of symmetry poses non trivial problems for controllability. The same kind of difficulty is therefore encountered in the attempt to define a reasonable notion of degree-k CLF for systems

(41)
$$\begin{cases} \dot{y} = f_0(y) + \sum_{i=1,\dots,m} a_i f_i(y) \\ y(0) = x \in \mathbb{R}^n \setminus \mathcal{T} \end{cases} \quad a \in \{0, \pm e_1, \dots, e_m\}$$

having a non zero drift f_0 . Let us assume that f_0, \ldots, f_m belong to $C_b^1(\Omega \setminus \mathcal{T})$ for any open, bounded set $\Omega \subset \mathbb{R}^n$.

Let us examine the case k = 2. Intuition coming from controllability literature suggests that a notion of degree-2 control Lyapunov function should be shaped in such a way that it would be allowed to violate the standard dissipative inequality only at the points where the drift f_0 vanishes. Accordingly, let us redefine the classes of vector fields

$$\mathcal{F}^{(1)} := \left\{ f_0, f_0 - f_i, f_0 + f_i \quad i = 1 \dots, m \right\},$$
$$\mathcal{F}^{(2)} := \mathcal{F}^{(1)} \cup \left\{ \pm [f_0, f_i] \cdot \chi_{\{f_0 = 0\}}, [f_j, f_\ell] \cdot \chi_{\{f_0 = 0\}} \quad i, j, \ell = 1 \dots, m \right\},$$

and the Hamiltonians

$$H^{(h)}(x,p) := \inf_{g \in \mathcal{F}^{(h)}(x)} \left\langle p, g \right\rangle \quad h = 1, 2.$$

Accordingly, one might call degree-2 control Lyapunov function any continuous function $U: \overline{\mathbb{R}^n \setminus \mathcal{T}} \to \mathbb{R}$ such that its restriction to $\mathbb{R}^n \setminus \mathcal{T}$ is locally semiconcave, positive definite, proper, and verifies

$$H^{(2)}(x, D^*U(x)) < 0 \quad \forall x \in \mathbb{R}^n \setminus \mathcal{T}.$$

In particular, U is a degree-2 CLF if, at each point $x \in \mathbb{R}^n \setminus \mathcal{T}$, either

$$\min\left\{\langle D^*U(x), f_0(x)\rangle, \langle D^*U(x), (f_0 - f_i)(x)\rangle, \langle D^*U(x), (f_0 + f_j)(x)\rangle, i, j = 1, \dots, m\right\} < 0,$$

or
$$\left(\int_{0}^{\infty} f_j(x) - 0\right)$$

$$\begin{cases} f_0(x) = 0,\\ \min\left\{ -|\langle D^*U(x), [f_0, f_i](x)\rangle|, -|\langle D^*U(x), [f_j, f_\ell(x)\rangle| \quad i, j, \ell = 1, \dots, m \right\} < 0. \end{cases}$$
With these settings, we get the following result:

With these settings, we get the following result:

Theorem 5.1. Let a degree-2 control Lyapunov function exist. Then system (41) is GAC to \mathcal{T} .

A proof of this result can be deduced by first observing that Lemma 3.2 has a counterpart in an asymptotic formula for brackets of the form $[f_0, f_i]$, $[f_j, f_\ell]$ valid at all points x where $f_0(x) = 0$. Precisely, through standard Taylor expansions one can prove the following result:

Lemma 5.1. If $f_0, f_1, \ldots, f_m \in C_b^1(\mathbb{R}^n \setminus \mathcal{T})$, there exists a constant c > 0 such that for any $x \in \mathbb{R}^n \setminus \mathcal{T}$ where $f_0(x) = 0$, any $j = 1, \ldots, m$, and any t > 0, the following estimates hold true:

(42)
$$d\left(y_x(t,\alpha_t) - x, [f_0, f_i](x) \frac{t^2}{4}\right) \le \frac{c}{4}t^3 \quad \forall s \in [0, t]$$

with

$$\alpha_t(s) := e_i \chi_{[0,t/2[}(s) - e_i \chi_{[t/2,t[}(s));$$

(43)
$$d\left(y_x(t,\hat{\alpha}_t) - x, [f_i, f_0](x)\frac{t^2}{4}\right) \le \frac{c}{4}t^3, \forall s \in [0, t]$$

with

$$\hat{\alpha}_t(s) := -e_i \chi_{[0,t/2[}(s) + e_i \chi_{[t/2,t[}(s);$$

and

(44)
$$d\left(y_x(t,\tilde{\alpha}_t) - x, [f_j, f_\ell](x) \frac{t^2}{16}\right) \le \frac{c}{16} t^3, \quad s \in [0, t]$$

with

$$\tilde{\alpha}_t(s) = e_j \chi_{[0,t/4[}(s) + e_\ell \chi_{[t/4,t/2[}(s) - e_j \chi_{[t/2,3t/4[}(s) - e_\ell \chi_{[3t/4,t]}(s).$$

Hence, provided one gives a (obvious) notion of degree-2 feedback at the points x where $f_0(x) = 0$, the proof of Theorem 5.1 can be easily achieved by means of the same arguments as in the proof of Theorem 2.1.

Example 5.1. Consider the so-called *soft landing* problem, $\dot{y}_1 = y_2$, $\dot{y}_2 = a$, $a \in \{0, -1, 1\}$, $\mathcal{T} = \{(0, 0)\}$. The distance function U(x) = |x| –as well as $|x|^{\alpha}$, $\alpha > 1$ – fails to be a degree-1 CLF (because the required inequality is not strict on the x_1 axis). However, U(x) is a degree-2 CLF.

A reasonable and useful notion of degree-k CLF can be likely given for the general case $k \ge 1$ provided only suitable subsets of brackets are allowed¹⁴.

¹⁴For instance, one can consider "good" brackets (see e.g. [Co]), of which $[f_0, f_i]$, $[f_j, f_\ell]$ are degree 2 instances. Among degree-3 brackets, $[f_0, [f_0, f_i]]$ is good for every $i = 1, \ldots, m$, while $[f_0, [f_j, f_\ell]]$ is not good, for all $j, \ell = 1, \ldots, m$

References

- [AB] F. Ancona & A. Bressan (1999) Patchy vector fields and asymptotic stabilization. ESAIM Control Optim. Calc. Var. 4, 445–471.
- [AgSa] A. A. Agrachev, Y. L. Sachkov, (2004) Control theory from the geometric viewpoint. Encyclopaedia of Mathematical Sciences, 87. Control Theory and Optimization, II. Springer-Verlag, Berlin.
- [AC] J.P. Aubin & A. Cellina, (1984) Differential inclusions. Set-valued maps and viability theory. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 264. Springer-Verlag, Berlin.
- [BR] A. Bacciotti & L. Rosier, (2005) Lyapunov functions and stability in control theory (2nd edition), Ed. Springer
- [BCD] M. Bardi & I. Capuzzo Dolcetta, (1997) Optimal control and viscosity solutions of Hamilton-Jacobi-Bellman equations, Ed. Birkhäuser, Boston.
- [BP] A. Bressan & B. Piccoli, (2007) Introduction to the mathematical theory of control. AIMS Series on Applied Mathematics, 2. American Institute of Mathematical Sciences (AIMS), Springfield, MO.
- [CS] P. Cannarsa & C. Sinestrari (2004) Semiconcave functions, Hamilton-Jacobi equations, and optimal control. Progress in Nonlinear Differential Equations and their Applications, 58. Birkhäuser Boston, Inc., Boston, MA.
- [CLSW] F. H. Clarke; Yu. S. Ledyaev; R. J. Stern; P. R. Wolenski, (1998) Nonsmooth analysis and control theory. Graduate Texts in Mathematics, 178. Springer-Verlag, New York.
- [CLSS] F. Clarke, Y. Ledyaev, E. Sontag, A. Subbotin, (1997) Asymptotic controllability implies feedback stabilization. IEEE Trans. Automat. Control 42, no. 10, 1394–1407.
- [CLRS] F. Clarke, Y. Ledyaev, L. Rifford, R.J. Stern, (2000), Feedback stabilization and Lyapunov functions. SIAM J. Control Optim. 39, no. 1, 25–48.
- [Co] J.-M. Coron, (2007), Control and nonlinearity. Mathematical Surveys and Monographs, 136. American Mathematical Society, Providence, RI.
- [FR1] E. Feleqi, F. Rampazzo, (2015), Integral representations for bracket-generating multi-flows. Discrete Contin. Dyn. Syst. 35, no. 9, 4345–4366.
- [FR2] E. Feleqi, F. Rampazzo (2016) Iterated Lie brackets for nonsmooth vector fields, Preprint.
- [FHT] H. Frankowska, D. Hoehener, D. Tonon, (2013), A second-order maximum principle in optimal control under state constraints. Serdica Math. J. 39, no. 3-4, 233–270.
- [K] H.-W. Knobloch, (1981), Higher order necessary conditions in optimal control theory. Lecture Notes in Computer Science, 34. Springer-Verlag, Berlin-New York.
- [Kr] M. Krastanov, (2009), High-order variations and small-time local attainability. Control Cybernet. 38, no. 4B, 1411–1427.
- [KrQ] M. Krastanov, M. Quincampoix, (2001), Local small time controllability and attainability of a set for nonlinear control system. ESAIM Control Optim. Calc. Var. 6, 499–516.
- [MaRS] M. Malisoff, L. Rifford, E. Sontag, (2004), Global asymptotic controllability implies input-to-state stabilization. SIAM J. Control Optim. 42, no. 6, 2221–2238.
- [Mar] A. Marigonda, S. Rigo, (2015), Controllability of some nonlinear systems with drift via generalized curvature properties. SIAM J. Control Optim. 53, no. 1, 434–474.
- [MR] M. Motta, F. Rampazzo, (2013), Asymptotic controllability and optimal control, J. Differential Equations 254, no. 7, 2744–2763.
- [RS1] F. Rampazzo, H.J. Sussmann, (2001), Set-valued differentials and a nonsmooth version of Chow's theorem, in: Proc. of the 40th IEEE Conf. on Decision and Control, Orlando, FL, December 2001, vol. 3, IEEE Publications, New York, pp. 2613–2618.
- [RS2] F. Rampazzo, H.J. Sussmann, (2007), Commutators of flow maps of nonsmooth vector fields. J. Differential Equations 232, no. 1, 134–175.

- [Ri] L. Rifford, (2000), Existence of Lipschitz and semiconcave control-Lyapunov functions. SIAM J. Control Optim. 39, no. 4, 1043–1064.
- [So] P. Soravia, (1992), Hölder continuity of the minimum-time function for C1-manifold targets. J. Optim. Theory Appl. 75, no. 2, 401–421.
- [Su] H.J. Sussmann, (2012) Geometry and optimal control, in "Mathematical control theory", John B. Baillieul, J.C. Willems (Eds), Springer Science & Business Media.
- [S1] E. Sontag, (1998), Mathematical control theory. Deterministic finite-dimensional systems. Second edition. Texts in Applied Mathematics, 6. Springer-Verlag, New York.
- [S2] E. Sontag, (1983), A Lyapunov-like characterization of asymptotic controllability. SIAM J. Control Optim. 21, no. 3, 462-471.

M. Motta, Dipartimento di Matematica, Università di Padova, Via Trieste, 63, Padova 35121, Italy

E-mail address: motta@math.unipd.it

F. RAMPAZZO, DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PADOVA, VIA TRIESTE, 63, PADOVA35121,ITALY

E-mail address: rampazzo@math.unipd.it