The crossing number of the cone of a graph

Carlos A. Alfaro¹*, Alan Arroyo²**, Marek Derňár³, and Bojan Mohar⁴***

- ¹ Banco de México, Ciudad de México, México
- ² Department of Combinatorics and Optimization, University of Waterloo, Canada
- ³ Faculty of Informatics, Masaryk University, Brno, Czech Republic
- ⁴ Department of Mathematics, Simon Fraser University, Burnaby, Canada

Abstract. Motivated by a problem asked by Richter and by the long standing Harary-Hill conjecture, we study the relation between the crossing number of a graph G and the crossing number of its cone CG, the graph obtained from G by adding a new vertex adjacent to all the vertices in G. Simple examples show that the difference cr(CG) - cr(G) can be arbitrarily large for any fixed k = cr(G). In this work, we are interested in finding the smallest possible difference, that is, for each non-negative integer k, find the smallest f(k) for which there exists a graph with crossing number at least k and cone with crossing number f(k). For small values of k, we give exact values of f(k) when the problem is restricted to simple graphs, and show that $f(k) = k + \Theta(\sqrt{k})$ when multiple edges are allowed.

1 Introduction

Little is known on the relation between the crossing number and the chromatic number. In this sense Albertson's conjecture (see [2]), that if $\chi(G) \geq r$, then $cr(G) \geq cr(K_r)$, has taken a great interest. Albertson's conjecture has been proved [2,4,15] for $r \leq 16$. It is related to Hajós' Conjecture that every r-chromatic graph contains a subdivision of K_r . If G contains a subdivision of K_r , then $cr(G) \geq cr(K_r)$. Thus Albertson's conjecture is weaker than Hajós' conjecture, however Hajós' conjecture is false for any $r \geq 7$ [7].

The cone of a graph G is the graph CG obtained from G by adding an apex, a new vertex that is adjacent to each vertex in G. Many properties of a graph are automatically transferred to its cone. For example, if G is r-coloring-critical, then CG is (r+1)-coloring-critical. During the Crossing Numbers Workshop in 2013, in an attempt to understand Alberston's conjecture, Richter proposed the following problem: Given an integer $n \geq 5$ and a graph G with crossing number at least $cr(K_n)$, does it follow that the crossing number of its cone CG is at least

^{*} Supported by SNI and CONACYT grant 166059.

 $^{^{\}star\star}$ Suported by CONACYT.

^{***} Supported in part by an NSERC Discovery Grant (Canada), by the Canada Research Chairs program, and by a Research Grant of ARRS (Slovenia). On leave from: IMFM, Department of Mathematics, Ljubljana, Slovenia.

 $cr(K_{n+1})$? There are examples where these two values can differ arbitrarily (for instance, if G is the disjoint union of K_4 's and K_5 's). What is less clear is how close these values can be.

The answer to Richter's question is positive for the first interesting case when n=5: Kuratowski's theorem implies that the cone of any graph with crossing number at least $cr(K_5)=1$ contains a subdivision of CK_5 or $CK_{3,3}$, and each of these graphs has crossing number at least $cr(K_6)=3$. Unfortunately, the answer is negative for the next case, as the graph in Figure 1 shows. This graph has crossing number 3, and a cone with crossing number at most 6, and this is less than $cr(K_7)=9$. This motivated us to investigate the following question.

Problem 1. For each $k \geq 0$, find the smallest integer f(k) for which there is a graph G with crossing number at least k and its cone has cr(CG) = f(k).



Fig. 1. A counterexample to Richter's question when n = 6.

Note that f(k) can also be defined as the largest integer such that every graph with $cr(G) \geq k$, has $cr(CG) \geq f(k)$. An upper bound to the function f(k) is obtained from the graph in Figure 1, by changing the multiplicity of each edge to r. Any drawing of the new graph has at least $3r^2$ crossings, and its cone has crossing number $3r^2 + 3r$. This shows that $f(k) \leq k + \sqrt{3k}$. Our main result shows that this is close to be best possible.

Theorem 2. Let G be a graph with $cr(G) \ge k$. Then $cr(CG) \ge k + \sqrt{k/2}$.

Thus we have the following:

Corollary 3. For multigraphs we have $f(k) = k + \Theta(\sqrt{k})$.

The paper is organized as follows. Page drawings, a concept intimately related to drawings of the cone of a graph, are defined in Section 2 and used throughout the subsequent sections. Although, there seems to be a connection between 1-page drawings and drawings of the cone, their exact relationship is much more subtle. Our proofs are instructive in this manner and provide further understanding of these concepts.

The proof of our main result, Theorem 2 is provided in Section 3. In Section 4, we restrict Problem 1 to the case of simple graphs. To distinguish between these two problems we use $f_s(k)$ instead of f(k). Along this paper, a graph is

allowed to have multiple edges but no loops; when our graphs have no multiple edges, then we refer them as simple graphs. We find the smallest values of f_s by showing that $f_s(1) = 3$, $f_s(2) = 5$, $f_s(3) = 6$, $f_s(4) = 8$ and $f_s(5) = 10$. These initial values may suggest that $f_s(k) \ge 2k$. However, in Section 5 we show that

$$f_s(k) = k + o(k),$$

and provide additional justification for a more specific conjecture that

$$f_s(k) = k + \sqrt{2} k^{3/4} (1 + o(1)).$$

2 Page drawings

In this section we describe a perspective provided from considering page drawings of graphs, a concept that has been studied in its own and has interesting applications. The relation between 1- and 2-page drawings has shown to be handy as it is used in the proofs of Theorems 2 and 7. A more detailed discussion on the relevant aspects of this section can be found in [3,6,14].

For an integer $k \geq 1$, a k-page book consists of k half planes sharing their boundary line ℓ (spine). A k-page-drawing is a drawing of a graph in which vertices are placed in the spine of a k-page book, and each edge arc is contained in one page. A convenient way to visualize a k-page drawing is by means of the circular model. In this model each page is represented by a unit 2-dimensional disk, so that the vertices are arranged identically on each disk boundary and each edge is drawn entirely in exactly one disk. In this work we are only interested in 1 and 2-page drawings, and, to be more precise, in the following problem.

Problem 4. Given a 1-page drawing of a graph G with k crossings, find an upper bound on the number of crossings of an optimal 2-page drawing of G while having the order of vertices of G on the spine unchanged.

In other words, if the drawing of G in the plane is such that all the vertices are incident to the outer-face (which is equivalent to having a 1-page drawing), what is the most efficient way to redraw some edges in the outer-face to reduce the number of crossings? For this purpose, we define the *circle graph* C_D of any 1-page drawing D of G as the graph whose vertices are the edges of G, and any two elements are adjacent if they cross in D. Note that C_D depends only on the cyclic order of the vertices of G in the spine.

A related problem was previously formulated by Kainen in [12], where he studied the *outerplanar crossing number* of a graph as the minimum number of crossings in any drawing of G so that all its vertices are incident to the same face. Clearly, the crossing number of CG is at most the outer-planar crossing number of G. Although, Kainen was interested in finding an n-vertex graph that has the largest difference between its crossing number and its outer-planar crossing number, for us it will be useful to consider drawings in which the vertices are incident to the same face.

Turning a 1-page drawing into a 2-page drawing is equivalent to finding a bipartition $(X, V(C_D) \setminus X)$ of the vertices of C_D , each part representing the set of edges of G drawn in one of the pages. Minimizing the number of crossings in the obtained 2-page drawing of G is equivalent to maximize the number edges in C_D between X and $V(C_D) \setminus X$. This last problem is known as the max-cut problem, and if the considered graph C_D has m edges, then, a well-known result of Erdős [8] states that its maximum edge-cut has size more than m/2. Improvements to this general bound are known (see [9], [10] and a more recent survey [5]). For our purpose the following bound of Edwards will be useful.

Lemma 5 (Edwards [9,10]). Suppose that G is a graph of order n with $m \ge 1$ edges. Then G contains a bipartite subgraph with at least $\frac{1}{2}m + \sqrt{\frac{1}{8}m + \frac{1}{64} - \frac{1}{8}} > \frac{1}{2}m$ edges.

In our context, this result translates to the following observation that we will use.

Corollary 6. Let D be a 1-page drawing of a graph G with $k \geq 1$ crossings. Then some edges of G can be redrawn in a new page, obtaining a 2-page drawing with at most $\frac{1}{2}k - \sqrt{\frac{1}{8}k + \frac{1}{64}} + \frac{1}{8}$ crossings. Such a drawing can be found in time O(|E(G)| + k).

The proof of Corollary 6 will be provided in the full version.

3 Lower bound on the crossing number of the cone

This section contains the proof of our main result.

Proof (of Theorem 2). Let \widehat{D} be an optimal drawing of the cone CG of G with apex a, and suppose \widehat{D} has less than $k + \sqrt{k/2}$ crossings. We consider $D = \widehat{D}|_{G}$, the drawing of G induced by \widehat{D} . If we let t to be the number of crossings in D, then we have

$$k \le t < k + \sqrt{k/2}.\tag{1}$$

For each vertex $v \in V(G) \cup \{a\}$, let s_v be the number of crossings in \widehat{D} involving edges incident with v. Using that $cr(\widehat{D}) < k + \sqrt{k/2}$ and the left-hand side inequality in (1), we obtain that $s_a < \sqrt{k/2}$.

Consider x_1, \ldots, x_{s_a} , the crossings involving edges incident with a. Since \widehat{D} is optimal, each of these crossings is between an edge incident to a and an edge in G. Let e_1, \ldots, e_{s_a} be the list of edges in G (we allow repetitions) so that x_i is the crossing between e_i and an edge incident with a. We subdivide each edge e_i in D using two points close to the crossing x_i , and we remove the edge segment σ_i joining these new two vertices, in order to obtain a drawing D_0 of a graph G_0 with t crossings (see Figure 2).

The obtained drawing D_0 has all its vertices incident to the face of D_0 containing the point corresponding to the apex vertex a of CG in \widehat{D} . For simplicity,




Fig. 2. A drawing where the crossed edges are cut.

we may assume that this is the unbounded face of D_0 . It follows that there exists a simple closed curve ℓ in the closure of this face, containing all the vertices of G_0 . Thus, D_0 gives rise to a 1-page drawing of G_0 with spine ℓ .

Now construct a new drawing of G as follows:

- 1. Start with the 1-page drawing D_0 . Partition the edges of G_0 according to Corollary 6, and draw the edges of one part in page 2 outside ℓ .
- 2. Reinsert edge segments $\sigma_1, \ldots, \sigma_{s_a}$ as they where drawn in D, to obtain a drawing D_1 (of a subdivision) of G. These segments do not cross each other, but they may cross some of the edges of G_0 that we placed in page 2 in step 1

Now we estimate the number of crossings in D_1 . According to Corollary 6, after step 1 we obtain a 2-page drawing D_0 with less than $t/2 - \sqrt{t/8} + 1/8$ crossings. After step 2 we gain some new crossings between the added segments $\sigma_1, \ldots, \sigma_{s_a}$ and the edges of G_0 drawn on page 2 in step 1.

Claim. The number of new crossings between $\sigma_1, \ldots, \sigma_{s_a}$ and the edges drawn on page 2 in step 1 is at most (k-1)/2.

Proof. We may assume that, for each $v \in V(G)$, $s_v < \sqrt{k/2}$. Otherwise, by removing v and all the edges incident to v, we obtain a drawing of CG - v containing a subdrawing of G, in which v is represented as the apex, and this drawing has less than k crossings, a contradiction.

Let $e \in E(G)$ be an edge having ends $u, v \in V(G)$. Suppose that $ay_1, ..., ay_{r_e}$ are the edges incident to a that cross e in \widehat{D} . We may assume that, for every i, j with $1 \le i < j \le r_e$, when we traverse e from u to v, the crossing $x_i = e \cap ay_i$ precedes the crossing $x_j = e \cap ay_j$. It is convenient to let $x_0 = u$ and $x_{r_e+1} = v$.

The edges of G_0 included in D[e] are the segments of $D[e] - \{\sigma_1, \ldots, \sigma_{s_a}\}$. We enumerate these edges as $\tau_0^e, \ldots, \tau_{r_e}^e$, so that τ_i^e is included in the $x_i x_{i+1}$ -arc of D[e]. Note that τ_1^e is incident to u, while τ_{r_e} is incident to v.

Let $T = \{\tau_i^e : e \in E(G) \text{ and } 0 \le i \le r_e\}$ be the set of edges of G_0 . In Step 1, when we apply Corollary 2.2 to the edges in D_0 , we obtain a partition

 $T_1 \cup T_2$ of T. Instead of counting how many crossings are between the segments in $\sigma_1, \ldots, \sigma_{s_a}$ and the edges in one of the T_i 's when we redraw T_i in page 2, we estimate the number m of crossings between $\sigma_1, \ldots, \sigma_{s_a}$ and the edges in T when we draw all the crossing edges in T in page 2. This will show that one of the two parts, either T_1 or T_2 , can be drawn in page 2 creating at most m/2 crossings with the segments $\sigma_1, \ldots, \sigma_{s_a}$. To show our claim, it suffices to prove that $m \leq k-1$, and this is what we do next.

For every point p distinct from a and contained in an edge f incident to a, the $depth\ h(p)$ of p is the number of crossings in \widehat{D} , contained in the open subarc of f connecting a to p. When we redraw an edge τ_i^e in page 2, we can draw it so that it crosses at most $h(x_i) + h(x_{i+1})$ segments in $\sigma_1, \ldots, \sigma_{s_a}$. Such new drawing of τ_i^e is obtained from letting the segment of τ_i^e near to x_i follow the same dual path in D that x_i follows to reach a via ay_i . Likewise the new end of τ_i^e near x_{i+1} is defined. The new τ_i^e is obtained from connecting the two end segments of τ_i^e inside the face of D containing a.

Let X(a) be the set of crossings involving edges incident to a. For every $x \in X(a)$, there are precisely two elements in T, so that when they are redrawn in page 2, one of its end segments mimics the arc between x and a inside the edge including x and a. Each $v \in V(G)$ is incident to at most s_v edges crossing in D_0 . Then, for every $v \in V(G)$, there are are most s_v edges in T, so that when we redraw them in page 2, one of their ends mimics the dual path followed by the edge $\widehat{D}[xa]$. These two observations together imply that

$$\begin{split} m &\leq \sum_{x \in X(a)} 2h(x) + \sum_{v \in V} h(v) s_v \\ &< 2 \sum_{v \in V} \left(1 + 2 + \ldots + (h(v) - 1) \right) + \sqrt{k/2} \sum_{v \in V} h(v) \\ &\leq \sum_{v \in V} h(v)^2 + (\sqrt{k/2}) s_a \leq \left(\sum_{v \in V} h(v) \right)^2 + k/2 \\ &= s_a^2 + k/2 < k. \end{split}$$

Because m is an integer less than $k, m \leq k-1$ as desired.

At the end, we obtained a drawing D_1 of (a subdivision of) G with less than $t/2 - \sqrt{t/8} + 1/8 + (k-1)/2$ crossings. Using (1) it follows that

$$cr(D_1) < \frac{1}{2}(k + \sqrt{k/2}) - \sqrt{t/8} + 1/8 + k/2 - 1/2 = k + \sqrt{k/8} - \sqrt{t/8} - 3/8 < k$$
, contradicting the fact that $cr(D_1) \ge cr(G) \ge k$.

4 Exact values of the crossing number of the cone for simple graphs

In this section, we investigate the minimum crossing number of a cone, with the restriction of only considering simple graphs. We are interested in finding the

smallest integer $f_s(k)$ for which there is a simple graph with crossing number at least k, whose cone has crossing number $f_s(k)$. On one hand, we describe below a family of simple graphs that shows that $f_s(k) \leq 2k$. Our best general lower bound is obtained from Theorem 2. The main result in this section, Theorem 7, help us to obtain exact values on $f_s(k)$ for cases when k is small.

Theorem 7. Let G be a simple graph with crossing number k. Then

- (1) if $k \geq 2$, then $cr(CG) \geq k + 3$;
- (2) if $k \ge 4$, then $cr(CG) \ge k + 4$; and
- (3) if $k \geq 5$, then $cr(CG) \geq k + 5$.

Before proving Theorem 7, we describe a family of examples that is used to find an upper bound for $f_s(k)$, that is exact for the values k=3,4,5. Given an integer $k\geq 3$, the graph F_k (Figure 3) is obtained from two disjoint cycles $C_1=x_0\dots x_{k-1}x_0$ and $C_2=y_0\dots y_{2k-1}y_0$ by adding, for each $i=0,\dots,k-1$, the edges $x_iy_{2i-2}, x_iy_{2i-1}, x_iy_{2i}, x_iy_{2i+1}$ (where the indices of the vertices y_j are taken modulo 2k). It is not hard to see that F_k has crossing number k: a drawing with k crossings is shown in Figure 3. To show that $cr(F_k)\geq k$, for $i\in\{0,\dots,k-1\}$, consider L^i , the K_4 induced by the vertices in $\{x_i,x_{i+1},y_{2i},y_{2i+1}\}$. Every L^i is a subgraph of a K_5 subdivision of F_k , thus, in an optimal drawing of F_k , at least one of the edges in L^i is crossed. This only guarantees that $cr(F_k)\geq k/2$, as two edges from distinct L^i 's might be crossed. However, if an edge from L^i crosses an edge e_j from some other L^j , then F_k-e_j has a K_5 subdivision including L^i , exhibiting a new crossing in some edge in L^i . Therefore, every L^i either has a crossing not involving an edge in another L^j , or there are least two crossings involving edges in L^i . This shows that $cr(F_k) \geq k$.




Fig. 3. The graph F_k .

The graph shown in Figure 4 has crossing number 2, and its cone has crossing number 5. This shows that $f_s(2) \leq 5$. On the other hand, F_3 , F_4 , and F_5 serve

as examples to show that $f_s(k) \leq 2k$ for k = 3, 4, 5. These bounds are tight for $2 \leq k \leq 5$ by Theorem 7.




Fig. 4. A graph with crossing number 2 whose cone has crossing number 5.

Proof (Proof of Theorem 7). Suppose G is a graph with cr(G) = k. Let \widehat{D} be an optimal drawing of the cone CG, D its restriction to G, and F_a be the face of D containing the apex a. The vertices of G incident to F_a are the planar neighbors of a.

Assume that $k \geq 2$, and suppose \widehat{D} has exactly k+t crossings. Theorem 2 guarantees that $t \geq 1$. Since each edge from a to a non-planar-neighbor introduce at least one crossing, the apex a has either 0, 1, 2, 3 or 4 non-planar neighbors (if a has more than 4 non-planar neighbours, then any of the items in Theorem 7 is satisfied).

We start by assuming that a has no non-planar neighbors. In this case, D is a 1-page drawing of G. Corollary 6 implies that we can obtain a new drawing of G with less than (k+t)/2 crossings. Thus (k+t)/2 > cr(G) = k, which implies that $t \ge k+1$. In any case of the theorem, this implies the conclusion, thus we may now assume that a has at least one and at most t non-planar neighbors.

(1) Let us now assume that k > 2 and t < 2.

Suppose a has exactly one non-planar neighbor u. Then cr(D) has at most k+1 crossings. At least one edge incident to u is crossed in D, otherwise, all the crossed edges have ends in F_a , and using Corollary 6, we obtain a drawing of G with less than (k+1)/2 crossings, contradicting that cr(G) = k. If at least two crossings in D involve edges incident to u, or if D has k crossings, then by redrawing u in F_a , and adding all the edges to its neighbors without creating any crossings, we obtain a drawing of G with less than k crossings. Therefore D has k+1 crossings, and exactly k of them involve edges not incident to u. Again, we apply Corollary 6 to obtain a drawing of G with at most $\frac{1}{2}(k-1) < k$ crossings (this time we are more careful by setting our two pages in such way that the edge not incident to u that crosses an edge incident to u is redrawn in the page contained in F_a).

Finally, suppose a has exactly two distinct non-planar neighbors u and v. Then, \widehat{D} has k+2 crossings; D has k crossings, and the edges au, av are crossed exactly once. Notice that any crossed edge in D is incident to either u or v; otherwise, we can redraw such edge inside F_a , obtaining a drawing of G with less than k crossings. Redraw v in $\widehat{D}[a]$ (where $\widehat{D}[a]$ denotes the point representing a in \widehat{D}); draw the edge uv (if it exists in G) as the arc $\widehat{D}[au]$, and draw the edges from v to its neighbors distinct from u, inside F_a without creating new crossings. Since every crossing in D involves an edge incident with v, we obtain a drawing of G with at most one crossing, a contradiction.

(2) Now, suppose that $k \geq 4$ and that t = 3.

The case when the apex a has only one non-planar neighbor u is similar to the above. If at least three crossings in D involve edges incident with u, then by redrawing u and the edges incident to u in F_a , we obtain a drawing with less than k crossings, a contradiction. Thus, at most two crossings involve edges incident to u. We redraw the remaining crossed edges according to Corollary 6 (if there is an edge e that crosses an edge incident to u, in order to remove an extra crossing, we may choose this new drawing so that e is redrawn in the page contained in F_a). If 2 crossings involve edges incident to u, then the obtained drawing has at most $\frac{k}{2} + 1$ crossings, where the +1 comes from the fact that e was drawn in the page contained in F_a . If at most one of the edges at u is crossed, then the new drawing has at most (k+1)/2 crossings. In any case, since $k \geq 4$, the new drawing has less than k crossings, a contradiction.

Let us now consider the case when the apex has two non-planar neighbors u and v. In this case, the drawing D has either k or k+1 crossings, and one of $\{au,av\}$, say au, is crossed only once. Let L be the set of crossed edges in D that are not incident to u or v. Suppose there are at least two crossings involving only edges in L. Then, either there are two edges in L that do not cross, or L has an edge e that crosses two other edges in L. In the former case, we redraw such pair of edges in F_a ; in the latter case, we redraw e in F_a . Any of these modifications yield a drawing with less than k crossings. Thus, we may assume that at most one crossing in D involves two edges not incident to u or v. Redraw v in $\widehat{D}[a]$; draw the edge vu (if such edge exists in G) as $\widehat{D}[au]$; and the remaining edges from v to its neighbors distinct from u without creating new crossings. The new drawing of G has at most two crossings: possibly one in $\widehat{D}[av]$ and another between edges in L, a contradiction.

Finally suppose that the apex a has three non-planar neighbors u, v, w. In this case D has precisely k crossings, and the edges au, av, aw are crossed exactly once. Observe that any crossed edge in D is incident to one of $\{u, v, w\}$, otherwise we can redraw such edge in F_a , obtaining a drawing of G with less than k crossings.

Let H be the graph induced by $\{u, v, w\}$. If, for $x \in \{u, v, w\}$, $d_H(x)$ denotes the degree of x in H, then at most $d_H(x)$ crossings involve edges at x. Otherwise, by redrawing x in $\widehat{D}[a]$; drawing the edges from x to its neighbors in H by using the respective edges from a; and, by drawing the remaining edges at x in F_a without creating new crossings, we obtain a drawing of G with less than K crossings. So for each vertex $K \in \{u, v, w\}$, there are at most two crossings involving edges at K. Hence K has at most three crossings, a contradiction.

(3) The proof can be found in Apendix A, and it will be included in the full version of the paper. \Box

5 Asymptotics for simple graphs

Lastly, we try to understand the behaviour of $f_s(k)$ when k is large. The important part is the increase of the crossing number after adding the apex, thus we define

$$\phi_s(k) = f_s(k) - k$$
.

We have proved that $\phi(k) = f(k) - k \ge \frac{1}{2}k^{1/2}$. The term $k^{1/2}$ is asymptotically tight in the case when we allow multiple edges. However, it is unclear how large $\phi_s(k)$ is. This question is treated next.

Theorem 8. $\phi_s(k) = O(k^{3/4})$.

Proof. Let us consider a positive integer k and let n be the smallest integer such that $cr(K_n) \ge k$. Then $G = K_n$ has a crossing number at least k and its cone is K_{n+1} .

To find an upper bound for $cr(K_{n+1})$ in terms of $cr(K_n)$, start with a drawing of K_n with $cr(K_n)$ crossings. Then clone a vertex, that is, place a new vertex very close to an original vertex, and draw the new edges along the original edges. Each edge incident to the new vertex cross $O(n^2)$ edges, thus the obtained drawing has $cr(K_n) + O(n^3)$ crossings. Therefore

$$\phi_s(k) \le cr(K_{n+1}) - cr(K_n) \le O(n^3).$$

It is known [13] that

$$\frac{3}{10} \binom{n}{4} \le cr(K_n) \le \frac{3}{8} \binom{n}{4}.$$

(The constant 3/10 in the lower bound has been recently improved to 0.32025, see [13] for more information.) Then $\phi_s(k) = O(n^3) = O(k^{3/4})$.

The Harary-Hill Conjecture [11] states that

$$cr(K_n) = \begin{cases} \frac{1}{64}n(n-2)^2(n-4), & n \text{ is even;} \\ \frac{1}{64}(n-1)^2(n-3)^2, & n \text{ is odd.} \end{cases}$$

Proposition 9. If the Harary-Hill conjecture holds, then

$$\phi_s(k) \le \sqrt{2} k^{3/4} (1 + o(1)).$$

Proof. As in the proof of Theorem 8, but with a slight twist for added precision, we take n such that $cr(K_{n-1}) < k \le cr(K_n)$. We also take n_1 such that for $k_1 = k - cr(K_{n-1})$ we have $cr(K_{n_1-1}) < k_1 \le cr(K_{n_1})$. Let $G = K_{n-1} \cup K_{n_1}$.

Then $cr(G) = cr(K_{n-1}) + cr(K_{n_1}) \ge k$ and $cr(CG) = cr(K_n) + cr(K_{n_1+1})$. Therefore,

$$\phi_s(k) \le cr(K_n) + cr(K_{n_1+1}) - cr(K_{n-1}) - cr(K_{n_1})$$

$$\le cr(K_n) - cr(K_{n-1}) + cr(K_{n_1+1}) - cr(K_{n_1-1}).$$

By inserting the values for the crossing number from the Harary-Hill Conjecture, we obtain (the calculation given is for odd n and odd n_1 , it is similar when n or n_1 is even):

$$cr(K_n) - cr(K_{n-1}) = \frac{1}{64}((n-1)^2(n-3)^2 - (n-1)(n-3)^2(n-5)) = \frac{1}{16}n^3(1 + o(1))$$

and

$$cr(K_{n_1+1}) - cr(K_{n_1-1}) = \frac{1}{64}((n_1+1)(n_1-1)^2(n_1-3) - (n_1-1)(n_1-3)^2(n_1-5))$$
$$= \frac{1}{8}n_1^3(1+o(1)).$$

Noticing that $k = \frac{1}{64}n^4(1+o(1))$ and $k_1 = \frac{1}{64}n_1^4(1+o(1)) = O(n^3)$ because $k_1 \le cr(K_n) - cr(K_{n-1})$, we conclude that $n_1^3 = O(n^{9/4}) = o(k^{3/4})$ and henceforth

$$\phi_s(k) \le \frac{1}{16}n^3(1+o(1)) + \frac{1}{8}n_1^3(1+o(1)) = \sqrt{2}k^{3/4}(1+o(1)).$$

The above proof works even under a weaker hypothesis that $cr(K_n) = \alpha n^4 + \beta n^3 (1 + o(1))$, where α and β are constants. This would imply that $\phi_s(k) = O(k^{3/4})$. Our conjecture is that (9) gives the precise asymptotics.

Conjecture 10.
$$\phi_s(k) = \sqrt{2} k^{3/4} (1 + o(1)).$$

A reviewer noted that this asymptotic is matched when the graph we are considering is dense.

Remark 11. Let G be a graph with n vertices, m edges, cr(G) = k and such that $m \ge 4n$. If $m = \Omega(n^2)$, then $cr(CG) \ge k + \Omega(k^{3/4})$.

The details will be provided in the full version.

Summary

To put the results of this paper into context, let us overview some of the motivation and some of directions for future work. The starting point of this paper was an attempt to understand Albertson's conjecture. The results of the paper (and their proofs) show that the crossing number behavior when adding an apex vertex is intimately related to 1-page drawings, but the exact relationship is quite subtle. There is some evidence that the minimal increase of the crossing number when an apex is added should be achieved with very dense graphs, close to the

complete graphs. Our Conjecture 10 entails this problem. Although very dense graphs have fewer vertices than sparser graphs with the same crossing number and thus need fewer connections to be made from the apex to their vertices, their near optimal drawings are far from 1-page drawings and therefore more crossings are needed. The full understanding of this antinomy would shed new light on the Harary-Hill conjecture.

Finally, it is worth pointing out that neither exact nor approximation algorithm is known for computing the crossing number of graphs of bounded treewidth. Adding an apex to a graph increases the tree-width of the graph by 1, thus understanding the crossing number of the cone is an important special case that would need to be understood before devising an algorithm for general graphs of bounded tree-width.

References

- M. Ajtai, V. Chvátal, M. M Newborn and E. Szemerédi, Crossing- free subgraphs, Theory and Practice of Combinatorics, North-Holland Math. Stud. 60, North-Holland, (1982), 9-12.
- M. O. Albertson, D. W. Cranston and J. Fox, Crossings, Colorings and Cliques, Electron. J. Combin. 16 (2009),# R45.
- 3. Bannister, M. J., Eppstein, D. (2014, September). Crossing minimization for 1-page and 2-page drawings of graphs with bounded treewidth. In International Symposium on Graph Drawing (pp. 210-221). Springer Berlin Heidelberg.
- J. Barát, G. Tóth, Towards the Albertson conjecture, Electron. J. Combin. 17 (2010) # R73.
- B. Bollobás, A.D. Scott, Better bounds for Max Cut, Contemporary Combinatorics, 185–246, Bolyai Soc. Math. Stud. 10, János Bolyai Math. Soc., Budapest, 2002.
- C. Buchheim and L. Zheng, Fixed Linear Crossing Minimization by Reduction to the Maximum Cut Problem, Computing and Combinatorics (2006) 507–516.
- 7. P.A. Catlin, Hajos' graph-coloring conjecture: variations and counterexamples. Journal of Combinatorial Theory Series B 26 (1979) 268–274.
- 8. P. Erdős, Gráfok páros körüljárású résgráfjairól (On bipartite subgraphs of graphs, in Hungarian), *Mat. Lapok* **18** (1967) 283–288.
- 9. C. S. Edwards. Some extremal properties of bipartite subgraphs. *Canad. J. Math.* **25** (1973) 475–485.
- 10. C. S. Edwards. An improved lower bound for the number of edges in a largest bipartite subgraph. In Recent Advances in Graph Theory (1975) 167–181.
- 11. F. Harary and A. Hill, On the number of crossings in a complete graph, *Proc. Edinburgh Math. Soc.* **13** (1963) 333–338.
- 12. P.C. Kainen, The book thickness of a graph II. Congr. Numer. 71 (1990) 127–132.
- E. De Klerk, Dmitrii V. Pasechnik, and Alexander Schrijver. Reduction of symmetric semidefinite programs using the regular *-representation, Math. Progr. 109 (2007) 613–624.
- E. de Klerk, D. Pasechnik, G. Salazar, Improved lower bounds on book crossing numbers of complete graphs, SIAM J. Discrete Math. 27 (2013) 619–633.
- B. Oporowski and D. Zhao, Coloring graphs with crossings. Discrete Math. 309 (2009) 2948–2951.

A Appendix: Proof of part (3) of Theorem 7

In this section we complete the proof of Theorem 7 by showing that if cr(G) = k and $k \ge 5$, then $cr(CG) \ge k + 5$.

Proof. Suppose G is a graph with cr(G) = k. Let \widehat{D} be an optimal drawing of the cone CG, D its restriction to G, and F_a be the face of D containing the apex a. The vertices of G incident to F_a are the *planar neighbors* of a. If u is a non-planar neighbor of a, then we let s_{au} to be the number crossings involving the edge au in \widehat{D} .

Assume that $k \geq 5$, and suppose \widehat{D} has less than k+5 crossings. By part (2) of Theorem 4.1 we know that $cr(\widehat{D}) = k+4$. Suppose that cr(D) = k+t. Let N denote the set of non-planar neighbors of a. As we did before, we split into cases depending on the size of N.

If the apex a has only planar neighbors, then, applying Corollary 6, we can obtain a drawing with less than $\frac{k+4}{2} < k$ crossings, a contradiction. Thus $|N| \ge 1$.

We need the following observation.

Claim. If $u \in N$, then the following holds:

- (i) At most $4 s_{au}$ crossings in D involve edges incident to u.
- (ii) The number of crossing in which both edges involved are incident to some vertex in N is at most $2|N| \lceil s/2 \rceil$.

Proof. (i) If there are more than $4 - s_{au}$ crossings involving edges at u, then we redraw u in the place of $\widehat{D}[a]$, join u to its neighbors using the corresponding edges from a to V(G). This is a drawing with less than

$$k+4-s-(4-s_{au})+s-s_{au}=k$$

crossings, a contradiction.

(ii) From (i), we know that for each $u \in N$, there are at most $4-s_{au}$ crossings in D involving edges at u. Let us count the number of pairs (u, x) where $u \in N$ and x is a crossing involving an edge incident with u. By (i), the number of such pairs is at most

$$\sum_{u \in N} (4 - s_{au}) = 4|N| - s. \tag{2}$$

This in particular implies (ii).

Case 1. The apex a has exactly one planar neighbor.

From Item (i) in the previous Claim, we know that there are at most $4-s \le 3$ crossings involving edges incident with $u \in N$. So at least k crossings involve crossing pairs that are not incident to u. We apply Corollary 6 to redraw some crossing edges not incident with u in F_a , and we are careful by choosing our

two pages so that we draw one edge crossing an edge at u (if such edge exists) in F_a to remove an extra crossing. Then we obtained a drawing with at most $k+4-s-(\frac{k}{2}+\frac{1}{2})-1< k$ crossings.

Case 2. The apex a has two non-planar neighbors.

Let us define t = 4 - s for brevity. Then $0 \le t \le 2$. Let X be the set of crossings involving an edge not incident to a vertex in N, and E_X be the set of crossing edges not incident to any vertex in N. We claim that $|X| \le t$.

This is easy to see when t = 0, as if $|X| \ge 1$, then there is an edge in E_X that we can redraw in F_a to obtain a drawing with less than k crossings. Suppose that t = 1. If $|X| \ge 2$, then, either there is an edge $e \in E_X$ including two crossings, or there is a pair of edges in E_X that are not crossed. In the former case we redraw e in F_a , in the latter we redraw the pair in F_a , to obtain a drawing with less than k crossings.

Finally, suppose that t=2. Any edge in E_X is involved in at most 2 crossings, otherwise we could redraw it in F_a to obtain a drawing with less than k crossings. If $|X| \geq 3$, then, either there is an edge $e \in E_X$ crossed twice and an edge $f \in E_X$ not crossing e, or every edge in E_X is crossed at most once. In the former case, we redraw e and f in F_a , in the latter, for each crossing in X we pick an edge E_X involved in the crossing and redraw it in F_a . In any case we obtain a drawing of G with less than k crossings. Therefore $|X| \leq 2$.

In any case we know that $|X| \leq t \leq 2$, and by Item (ii), the number of crossings in D is at most

$$2 + t/2 + |X| \le 2 + 3t/2 \le 5$$
.

Since $cr(G) \ge 5$, we have that cr(D) = 5, |X| = t = 2, and $E_X \ne \emptyset$. However, if we redraw any edge from E_X in F_a , we obtain a drawing of G with less than 5 crossings.

Case 3. The apex a has three non-planar neighbors.

In this case cr(D) is either k or k+1, so t=0 or t=1. The argument given in the previous case shows that there are at most t crossings involving an edge not incident to a vertex in N. Using Item (ii), we know that D has at most 4+t/2+|X| crossings, where X is defined as in the previous case. Since $cr(D) \geq 5$, this shows that D has exactly 5 crossings, and that |X| = 1. In particular cr(D) = k and thus t = 0, which contradicts that $|X| \leq t$.

Case 4. The apex has four non-planar neighbors.

In this case s=4, cr(D)=k and $s_{ay}=1$ for every $y\in N$. Let $N=\{u,v,w,x\}$. Note that each crossing edge is incident to a vertex in N. By Item (i), there are at most 3 crossings involving edges incident to a fixed vertex in N, and by (ii), $cr(D) \leq 6$. Moreover, the count (2) in the proof of (ii) shows that cr(D)=5 if there is an edge with both ends in N that is involved in a crossing.

Let H be the graph induced by N. We will split into two cases depending on whether D[H] is a crossing K_4 or not.

Subcase 1 D[H] is not a crossing K_4 .

If $H = K_4$, then D[H] is a planar K_4 . This implies that there is a 3-cycle composed of vertices in N, separating a fourth vertex in N from a, and this contradicts that $s_{ay} = 1$ for every $y \in N$. Therefore, there is a pair of vertices in H, say u and v, with $uv \notin E(G)$.

If, for $y \in N$, $d_H(y)$ denotes the degree of y in H, then at most $d_H(y)$ crossings involve edges at a. Otherwise, by redrawing y in $\widehat{D}[a]$; drawing the edges from y to its neighbors in H by using the respective edges from a; and, by drawing the remaining edges at y in F_a without creating new crossings, we obtain a drawing of G with less than k crossings. Since $d_H(u), d_H(v) \leq 2$ and $d_H(w), d_H(x) \leq 3$, cr(D) is at most $\left(\sum_{y \in N} d_H(y)\right)/2 = 5$. Because $cr(D) \geq 5$, this implies that cr(D) = 5, $d_H(u) = d_H(v) = 2$ and that $d_H(w) = d_H(x) = 3$. This also shows that for each $y \in N$, the number of crossings involving edges at y is exactly $d_H(y)$. Also note that none of the edges in H are crossed, as otherwise, at least three of the four vertices involved in some crossing belong to N, and a refined version of our previous counting would exhibit that D has at most 4 crossings.

Let H' be the drawing induced by $N \cup \{a\}$. Our previous observations imply that the drawing of D[H'] is isomorphic to the drawing of the cone of a planarly drawn K_4 minus one edge, where the apex is drawn in the face bounded by the 4-cycle of $K_4 - e$, and the edges incident to the apex connect directly to the boundary of the 4-cycle. Moreover, the only crossings of H' in \widehat{D} , are those between the edges at a and the boundary of F_a in D. This restricted drawing of H' implies that the ends of a crossing pair of edges have exactly one element in $\{u,v\}$, exactly one element in $\{w,x\}$, and none of the two edges has both ends in N. However, this is not possible, as there are 4 crossings involving edges incident to one of u or v, while there are 6 crossings involving edges incident to one of w or x.

Subcase 2 D[H] is a K_4 with a crossing.

Suppose that uv and wx is the crossing pair in D[H], and that \times is the crossing between uv and wx. Following the same argument given in the previous case, it is easy to see that for every $y \in N$, there are exactly 2 crossings distinct from \times involving edges at y; cr(D) = 5; if H' is graph is induced by $N \cup \{a\}$, then its drawing D[H'] is isomorphic to the drawing of the cone of a crossing K_4 , where the apex is drawn in the face bounded by the 4-cycle of the K_4 , and the edges incident to the apex connect directly to the boundary of the 4-cycle; and, the only crossings of \widehat{D} in H', distinct from \times , are those between the edges at a and the boundary of F_a in D.

The restrictions on H' show that the ends of a crossing pair of edges distinct from uv and wx, have exactly one element in $\{u, v\}$, exactly one element in $\{w, x\}$, and none of the two edges has both ends in N.

The boundary walk of F_a contains a cycle C that in \widehat{D} separates a from N. There are two internally disjoint subarcs α and β of D[C] connecting the

crossings between D[C] and each of aw and ax. We label α and β so that α includes the crossing between D[C] and au, and β includes the crossing between D[C] and av. The restrictions imposed by the crossings in H' imply that all the neighbors of u not in N are contained in α , and likewise the neighbors of v not in N are contained in β .

We obtain a drawing of G with 4 crossings as follows. Redraw u in the place of $\widehat{D}[a]$; join a to each of w and x using the corresponding edges from a to each of w and x. Draw the edges from u to its neighbors not in N without creating new crossings. Now redraw v near u in the face bounded by β and the two segments of the new uw, ux edges. Connect v to each of w and x by following arcs near the new uw, ux edges. Connect v to the rest of its neighbors without creating new crossings. Since $cr(G) \geq 5$ and this drawing has 4 crossings, this is a contradiction.

In any case we obtained a contradiction. Thus $cr(CG) \ge k + 5$ when $k \ge 5$.