An HDG Method for Dirichlet Boundary Control of Convection Dominated Diffusion PDE

Gang Chen*

John R. Singler[†]

Yangwen Zhang[‡]

November 27, 2018

Abstract

We first propose a hybridizable discontinuous Galerkin (HDG) method to approximate the solution of a *convection dominated* Dirichlet boundary control problem. Dirichlet boundary control problems and convection dominated problems are each very challenging numerically due to solutions with low regularity and sharp layers, respectively. Although there are some numerical analysis works in the literature on *diffusion dominated* convection diffusion Dirichlet boundary control problems, we are not aware of any existing numerical analysis works for convection dominated boundary control problems. Moreover, the existing numerical analysis techniques for convection dominated PDEs are not directly applicable for the Dirichlet boundary control problem because of the low regularity solutions. In this work, we obtain an optimal a priori error estimate for the control under some conditions on the domain and the desired state. We also present some numerical experiments to illustrate the performance of the HDG method for convection dominated Dirichlet boundary control problems.

1 Introduction

Let $\Omega \subset \mathbb{R}^d$ (d = 2, 3) be a Lipschitz polyhedral domain with boundary $\Gamma = \partial \Omega$. We consider the following Dirichlet boundary control problem:

$$\min J(y,u) = \frac{1}{2} \|y - y_d\|_{L^2(\Omega)}^2 + \frac{\gamma}{2} \|u\|_{L^2(\Gamma)}^2, \quad \gamma > 0,$$
(1.1)

subject to

$$-\varepsilon \Delta y + \nabla \cdot (\beta y) + \sigma y = f \quad \text{in } \Omega,$$

$$y = u \quad \text{on } \Gamma,$$
 (1.2)

where $f \in L^2(\Omega), \varepsilon \ll 1$, and we make other assumptions on β and σ for our analysis.

Researchers have performed numerical analysis of computational methods for Dirichlet boundary control problems for over a decade. Many researchers considered the standard finite element method and obtained an error estimate for the optimal control of order h^s for all $s < \min\{1, \pi/2\omega\}$, where ω is the largest angle of the boundary polygon; see, e.g., [8, 36, 37]. Apel et al. in [1] considered special meshes and obtained an optimal convergence rate with $s < \min\{3/2, \pi/\omega - 1/2\}$.

^{*}School of Mathematics Sciences, University of Electronic Science and Technology of China, Chengdu, China (cglwdm@uestc.edu.cn).

[†]Department of Mathematics and Statistics, Missouri University of Science and Technology, Rolla, MO, USA (singlerj@mst.edu).

[‡]Department of Mathematics Science, University of Delaware, Newark, DE, USA (ywzhangf@udel.edu).

Some mixed finite element methods have also been used for Dirichlet boundary control problems because the essential Dirichlet boundary condition becomes natural, i.e., the Dirichlet boundary data directly enters the variational setting. In [22], Gong et al. used a standard mixed method to obtain an error estimate for all $s < \min\{1, \pi/2\omega\}$. Recently, we used an HDG method to obtain an optimal convergence rate for all $s < \min\{3/2, \pi/\omega - 1/2\}$ without using higher order elements or a special mesh [30]. Moreover, the number of degrees of freedom are lower for HDG methods than standard mixed methods.

All of the above works focus on Dirichlet boundary control of the Poisson equation. However, Dirichlet boundary control problems play an important role in many applications governed by more complicated models, such as the Navier-Stokes equations; see, e.g., [21, 24–27]. In order to work towards numerical analysis results for more difficult PDEs, one essential and necessary step is to fully understand the convection diffusion Dirichlet boundary control problem. Benner and Yücel in [4] used a local discontinuous Galerkin (LDG) method and they obtained an error estimate for the control of order $\mathcal{O}(h^s)$ for all $s < \min\{1, \pi/2\omega\}$. Also, very recently, we proposed a new HDG method to study this problem and obtained an optimal convergence rate $\mathcal{O}(h^s)$ for all $s < \min\{3/2, \pi/\omega - 1/2\}$; see [23, 29] for more details.

However, the previous works only approximated solutions of convection diffusion Dirichlet boundary control problems in the *diffusion dominated* case. They did not consider the more difficult *convection dominated* case, i.e., $\varepsilon \ll |\beta|$. Even without the Dirichlet boundary control, solutions of convection dominated diffusion PDEs typically have layers; therefore, designing a robust numerical scheme for this problem is a major difficulty difficulty and has been considered in many works; see, e.g., [7, 19, 32, 38] and the references therein. Discontinuous Galerkin (DG) methods have proved very useful for solving convection dominated PDEs; see, e.g., [6, 9, 12, 13, 16, 33, 41] for standard DG methods and [20, 31] for HDG methods. For more information on HDG methods; see, e.g., [10, 11, 14, 15, 17, 18, 39, 40]. Moreover, there are some existing convection dominated diffusion *distributed* optimal control numerical analysis works; see, e.g., [3, 28, 34]. However, the techniques in the above works are *not* applicable for convection dominated Dirichlet boundary control problems since the solutions of (1.1)-(1.2) frequently have low regularity, i.e., $y \in H^{1+s}(\Omega)$ with $0 \leq s < 1/2$.

Formally, the optimal control $u \in L^2(\Gamma)$ and the optimal state $y \in L^2(\Omega)$ minimizing the cost functional satisfy a mixed weak formulation of the optimality system

$$-\varepsilon \Delta y + \nabla \cdot (\beta y) + \sigma y = f \qquad \text{in } \Omega, \qquad (1.3a)$$

$$y = u$$
 on Γ , (1.3b)

$$-\varepsilon \Delta z - \nabla \cdot (\boldsymbol{\beta} z) + (\nabla \cdot \boldsymbol{\beta} + \sigma) z = y - y_d \quad \text{in } \Omega,$$
(1.3c)

$$z = 0 \qquad \text{on } \Gamma, \tag{1.3d}$$

$$\gamma u - \varepsilon \nabla z \cdot \boldsymbol{n} = 0 \qquad \text{on } \Gamma. \tag{1.3e}$$

In this work, we use polynomials of degree k to approximate the state y, dual state z and their fluxes $\mathbf{q} = -\varepsilon \nabla y$ and $\mathbf{p} = -\varepsilon \nabla z$, respectively. Moreover, we also use polynomials of degree k to approximate the numerical trace of the state and dual state on the edges (or faces) of the spatial mesh, which are the only globally coupled unknowns. The HDG method considered here is different from the HDG method we considered for convection diffusion Dirichlet boundary control problems in [23,29]. A major difference is that the HDG method here has a lower computational cost.

In Section 4, we obtain an optimal convergence rate for the optimal control in 2D under certain basic assumptions on the desired state y_d and the domain Ω ; specifically, we prove

$$\|u - u_h\|_{\Gamma} \le Ch^s,\tag{1.4}$$

for all $s < \min\{3/2, \pi/\omega - 1/2\}$, and the constant *C* only depends on the exact solution, the domain and the polynomial degree. To prove the estimate (1.4), we cannot use the numerical analysis strategy from [4, 23, 29] because the constants in their error estimates may blow up as ε approaches zero. In order to obtain the estimate (1.4) with the constant *C* independent of ε , we follow a strategy from [20] and use weighted test functions in an energy argument. However, the techniques used in [20] are not directly applicable for solutions with low regularity. Moreover, unlike all the previous Dirichlet boundary control numerical analysis works, we only assume the mesh is shape regular, not quasi-uniform. We present numerical results in Section 5 to illustrate the performance of the HDG method.

2 Optimality system, regularity and HDG formulation

We begin with some notation. For any bounded domain $\Lambda \subset \mathbb{R}^d$ (d = 2, 3), let $H^m(\Lambda)$ and $H_0^m(\Lambda)$ denote the usual *m*th-order Sobolev spaces on Λ , and let $\|\cdot\|_{m,\Lambda}$, $|\cdot|_{m,\Lambda}$ denote the norm and seminorm on these spaces. We use $(\cdot, \cdot)_{m,\Lambda}$ to denote the inner product on $H^m(\Lambda)$, and set $(\cdot, \cdot)_{\Lambda} := (\cdot, \cdot)_{0,\Lambda}$. When $\Lambda = \Omega$, we denote $\|\cdot\|_m := \|\cdot\|_{m,\Omega}$, and $|\cdot|_m := |\cdot|_{m,\Omega}$. Also, when Λ is the boundary of a set in \mathbb{R}^d , we use $\langle \cdot, \cdot \rangle_{\Lambda}$ to replace $(\cdot, \cdot)_{\Lambda}$. Bold face fonts will be used for vector Sobolev spaces along with vector-valued functions. In addition, we introduce the following space:

$$\boldsymbol{H}(\operatorname{div},\Lambda) := \{ \boldsymbol{v} \in [L^2(\Lambda)]^d : \nabla \cdot \boldsymbol{v} \in L^2(\Lambda) \}.$$

We now present the optimality system for problem (1.1)-(1.2) and give a regularity result.

2.1 Optimality system and regularity

Throughout the paper, we suppose Ω is a convex polygonal domain, and let $\omega \in [\pi/3, \pi)$ denote its largest interior angle. The optimal control u is determined by the optimality system for the state y and the dual state z. For the HDG method, we use a mixed formulation of the optimality system; therefore we introduce the primary flux $\boldsymbol{q} = -\varepsilon \nabla y$ and the dual flux $\boldsymbol{p} = -\varepsilon \nabla z$. The well-posedness and regularity of the mixed formulation of the optimality system is contained in the result below. The proof of Theorem 1 is omitted here since it is very similar with a proof of a similar result in [29].

Theorem 1. If $y_d \in H^{t^*}(\Omega)$ for some $0 \leq t^* < 1$, $\sigma \in L^{\infty}(\Omega) \cap H^1(\Omega)$, f = 0 and the velocity vector field β satisfies

$$\boldsymbol{\beta} \in [L^{\infty}(\Omega)]^2, \quad \nabla \cdot \boldsymbol{\beta} \in L^{\infty}(\Omega), \quad \sigma + \frac{1}{2} \nabla \cdot \boldsymbol{\beta} \ge 0, \quad \nabla \nabla \cdot \boldsymbol{\beta} \in [L^2(\Omega)]^2,$$
(2.1)

then problem (1.1)-(1.2) has a unique solution $u \in L^2(\Gamma)$. Moreover, for any s > 0 satisfying $s \leq 1/2 + t^*$ and $s < \min\{3/2, \pi/\omega - 1/2\}$, we have

$$(u, \boldsymbol{q}, \boldsymbol{p}, y, z) \in H^{s}(\Gamma) \times \boldsymbol{H}^{s-\frac{1}{2}}(\Omega) \times \boldsymbol{H}^{s+\frac{1}{2}}(\Omega) \times H^{s+\frac{1}{2}}(\Omega) \times (H^{s+\frac{3}{2}}(\Omega) \cap H^{1}_{0}(\Omega)),$$

is the unique solution of

$$\varepsilon^{-1}(\boldsymbol{q},\boldsymbol{r})_{\Omega} - (\boldsymbol{y},\nabla\cdot\boldsymbol{r})_{\Omega} + \langle \boldsymbol{u},\boldsymbol{r}\cdot\boldsymbol{n}\rangle_{\Gamma} = 0, \qquad (2.2a)$$

$$(\nabla \cdot (\boldsymbol{q} + \boldsymbol{\beta} y), w)_{\Omega} + (\sigma y, w)_{\Omega} = 0, \qquad (2.2b)$$

$$\varepsilon^{-1}(\boldsymbol{p},\boldsymbol{r})_{\Omega} - (z,\nabla\cdot\boldsymbol{r})_{\Omega} = 0, \qquad (2.2c)$$

$$(\nabla \cdot (\boldsymbol{p} - \boldsymbol{\beta} z), w)_{\Omega} + ((\nabla \cdot \boldsymbol{\beta} + \sigma)y, w)_{\Omega} = (y - y_d, w)_{\Omega},$$
(2.2d)

$$\langle \gamma u + \boldsymbol{p} \cdot \boldsymbol{n}, v \rangle_{\Gamma} = 0,$$
 (2.2e)

for all $(\boldsymbol{r}, w, v) \in \boldsymbol{H}(\operatorname{div}, \Omega) \times L^2(\Omega) \times L^2(\Gamma)$. Furthermore, we have $\Delta y \in L^2(\Omega)$.

2.2 The HDG formulation

Let $\mathcal{T}_h = \bigcup\{T\}$ be a conforming simplex mesh that partitions the domain Ω . For any $T \in \mathcal{T}_h$, we let h_T be the diameter of T and denote the mesh size by $h := \max_{T \in \mathcal{T}_h} h_T$. Denote the edges of T by E, let \mathcal{E}_h be the set of all edges E, let \mathcal{E}_h^∂ be the set of edges E such that $E \subset \Gamma$, and set $\mathcal{E}_h^o = \mathcal{E}_h \setminus \mathcal{E}_h^\partial$. Let h_E denote the diameter of E. The mesh dependent inner products are denoted by

$$(w,v)_{\mathcal{T}_h} = \sum_{T \in \mathcal{T}_h} (w,v)_T, \qquad \qquad \langle \zeta, \rho \rangle_{\partial \mathcal{T}_h} = \sum_{T \in \mathcal{T}_h} \langle \zeta, \rho \rangle_{\partial T}.$$

We use ∇ and ∇ to denote the broken gradient and broken divergence with respect to \mathcal{T}_h . For an integer $k \geq 0$, $\mathcal{P}_k(\Lambda)$ denotes the set of all polynomials defined on Λ with degree not greater than k. We introduce the discontinuous finite element spaces.

$$\begin{aligned} \boldsymbol{V}_h &:= \{ \boldsymbol{v} \in [L^2(\Omega)]^2 : \boldsymbol{v}|_T \in [\mathcal{P}_k(T)]^d, \forall T \in \mathcal{T}_h \}, \\ W_h &:= \{ w \in L^2(\Omega) : w|_T \in \mathcal{P}_k(T), \forall T \in \mathcal{T}_h \}, \\ M_h^o &:= \{ \mu \in L^2(\mathcal{E}_h) : \mu|_E \in \mathcal{P}_k(E), \forall E \in \mathcal{E}_h, \text{ and } \mu|_{\Gamma} = 0 \}, \\ M_h^\partial &:= \{ \mu \in L^2(\mathcal{E}_h^\partial) : \mu|_E \in \mathcal{P}_k(E), \forall E \in \mathcal{E}_h^\partial \}. \end{aligned}$$

In our earlier works [23,29], we used a \mathcal{P}_{k+1} local space for the spaces W_h and M_h . In this work, we use polynomial degree k for all spaces. Since the globally coupled degrees of freedom depend on the space M_h , the computational cost of the HDG method in this paper is much lower than the HDG method in [23,29].

The HDG method for mixed weak form of the optimality system (2.2) is to find $(\boldsymbol{q}_h, y_h, \hat{y}_h^o, \boldsymbol{p}_h, z_h, \hat{z}_h^o, u_h) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2 \times M_h^\partial$ such that

$$\varepsilon^{-1}(\boldsymbol{q}_h, \boldsymbol{r}_h)_{\mathcal{T}_h} - (y_h, \nabla \cdot \boldsymbol{r}_h)_{\mathcal{T}_h} + \langle \hat{y}_h^o, \boldsymbol{r}_h \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_h} = -\langle u_h, \boldsymbol{r}_h \cdot \boldsymbol{n} \rangle_{\mathcal{E}_h^\partial},$$
(2.3a)

for all $\boldsymbol{r}_h \in \boldsymbol{V}_h$,

$$-(w_{h}, \nabla \cdot \boldsymbol{q}_{h})_{\mathcal{T}_{h}} + \langle \widehat{w}_{h}^{o}, \boldsymbol{q}_{h} \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_{h}} - \langle \tau_{1}(y_{h} - \widehat{y}_{h}^{o}), w_{h} - \widehat{w}_{h}^{o} \rangle_{\partial \mathcal{T}_{h}} + (y_{h}, \boldsymbol{\beta} \cdot \nabla w_{h})_{\mathcal{T}_{h}} - \langle \widehat{y}_{h}^{o}, \boldsymbol{\beta} \cdot \boldsymbol{n} w_{h} \rangle_{\partial \mathcal{T}_{h}} - (\sigma y_{h}, w_{h})_{\mathcal{T}_{h}}$$
(2.3b)
$$= -(f, w_{h})_{\mathcal{T}_{h}} - \langle (\tau_{1} - \boldsymbol{\beta} \cdot \boldsymbol{n}) u_{h}, v_{h} \rangle_{\mathcal{E}_{h}^{\partial}},$$

for all $(w_h, \widehat{w}_h^o) \in W_h \times M_h^o$,

$$\varepsilon^{-1}(\boldsymbol{p}_h, \boldsymbol{r}_h)_{\mathcal{T}_h} - (z_h, \nabla \cdot \boldsymbol{r}_h)_{\mathcal{T}_h} + \langle \hat{z}_h^o, \boldsymbol{r}_h \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_h} = 0, \qquad (2.3c)$$

for all $r_h \in V_h$,

$$-(w_{h}, \nabla \cdot \boldsymbol{p}_{h})_{\mathcal{T}_{h}} + \langle \widehat{w}_{h}^{o}, \boldsymbol{p}_{h} \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_{h}} - \langle \tau_{2}(z_{h} - \widehat{z}_{h}^{o})_{\mathcal{T}_{h}}, w_{h} - \widehat{w}_{h}^{o} \rangle_{\partial \mathcal{T}_{h}} -(z_{h}, \boldsymbol{\beta} \cdot \nabla w_{h})_{\mathcal{T}_{h}} + \langle \widehat{z}_{h}^{o}, \boldsymbol{\beta} \cdot \boldsymbol{n} w_{h} \rangle_{\partial \mathcal{T}_{h}} - ((\sigma + \nabla \cdot \boldsymbol{\beta}) z_{h}, w_{h})_{\mathcal{T}_{h}}$$
(2.3d)
$$= -(y_{h} - y_{d}, w_{h})_{\mathcal{T}_{h}},$$

for all $(w_h, \widehat{w}_h^o) \in W_h \times M_h^o$,

$$\langle \gamma u_h + \boldsymbol{p}_h \cdot \boldsymbol{n} + \tau_2 (z_h - \hat{z}_h^o), \, \widehat{w}_h^\partial \rangle_{\varepsilon_h^\partial} = 0,$$
 (2.3e)

for all $\widehat{w}_h^{\partial} \in M_h^{\partial}$. Here, the positive stabilization functions τ_1 and τ_2 are chosen as

$$\tau_1|_{\partial T} = \|\boldsymbol{\beta} \cdot \boldsymbol{n}\|_{0,\infty,\partial T} + \frac{1}{2}\boldsymbol{\beta} \cdot \boldsymbol{n} + \varepsilon h_T^{-1}, \qquad (2.3f)$$

$$\tau_2|_{\partial T} = \|\boldsymbol{\beta} \cdot \boldsymbol{n}\|_{0,\infty,\partial T} - \frac{1}{2}\boldsymbol{\beta} \cdot \boldsymbol{n} + \varepsilon h_T^{-1}.$$
(2.3g)

To simplify the presentation later, we define

$$\tau = \frac{\tau_1 + \tau_2}{2} = \|\boldsymbol{\beta} \cdot \boldsymbol{n}\|_{0,\infty,\partial T} + \varepsilon h_T^{-1}.$$
(2.3h)

2.3 A compact formulation

To simplify the notation, for $(\boldsymbol{q}_h, y_h, \widehat{y}_h^o, \boldsymbol{p}_h, z_h, \widehat{z}_h^o, \boldsymbol{r}_h, w_h, \widehat{w}_h^o) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^3$, we denote

$$\mathcal{B}_{1}(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}; \boldsymbol{r}_{h}, w_{h}, \widehat{w}_{h}^{o}) \\
= \varepsilon^{-1}(\boldsymbol{q}_{h}, \boldsymbol{r}_{h})_{\mathcal{T}_{h}} - (y_{h}, \nabla \cdot \boldsymbol{r}_{h})_{\mathcal{T}_{h}} + \langle \widehat{y}_{h}^{o}, \boldsymbol{r}_{h} \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_{h}} \\
- (w_{h}, \nabla \cdot \boldsymbol{q}_{h})_{\mathcal{T}_{h}} + \langle \widehat{w}_{h}^{o}, \boldsymbol{q}_{h} \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_{h}} - \langle \tau_{1}(y_{h} - \widehat{y}_{h}^{o}), w_{h} - \widehat{w}_{h}^{o} \rangle_{\partial \mathcal{T}_{h}} \\
+ (y_{h}, \boldsymbol{\beta} \cdot \nabla w_{h})_{\mathcal{T}_{h}} - \langle \widehat{y}_{h}^{o}, \boldsymbol{\beta} \cdot \boldsymbol{n} w_{h} \rangle_{\partial \mathcal{T}_{h}} - (\sigma y_{h}, w_{h})_{\mathcal{T}_{h}}, \qquad (2.4)$$

$$=\varepsilon^{-1}(\boldsymbol{p}_{h},\boldsymbol{r}_{h},\boldsymbol{z}_{h},\boldsymbol{z}_{h},\boldsymbol{w}_{h},\boldsymbol{w}_{h})$$

$$=\varepsilon^{-1}(\boldsymbol{p}_{h},\boldsymbol{r}_{h})_{\mathcal{T}_{h}}-(z_{h},\nabla\cdot\boldsymbol{r}_{h})_{\mathcal{T}_{h}}+\langle\hat{z}_{h}^{o},\boldsymbol{r}_{h}\cdot\boldsymbol{n}\rangle_{\partial\mathcal{T}_{h}}$$

$$-(w_{h},\nabla\cdot\boldsymbol{p}_{h})_{\mathcal{T}_{h}}+\langle\hat{w}_{h}^{o},\boldsymbol{p}_{h}\cdot\boldsymbol{n}\rangle_{\partial\mathcal{T}_{h}}-\langle\tau_{2}(z_{h}-\hat{z}_{h}^{o}),w_{h}-\hat{w}_{h}^{o}\rangle_{\partial\mathcal{T}_{h}}$$

$$-(z_{h},\boldsymbol{\beta}\cdot\nabla w_{h})_{\mathcal{T}_{h}}+\langle\hat{z}_{h}^{o},\boldsymbol{\beta}\cdot\boldsymbol{n}w_{h}\rangle_{\partial\mathcal{T}_{h}}-((\sigma+\nabla\cdot\boldsymbol{\beta})z_{h},w_{h})_{\mathcal{T}_{h}}.$$
(2.5)

Then we can rewrite (2.3) as follows: find $(\boldsymbol{q}_h, y_h, \hat{y}_h^o, \boldsymbol{p}_h, z_h, \hat{z}_h^o, u_h) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2 \times M_h^\partial$ such that

$$\mathcal{B}_1(\boldsymbol{q}_h, y_h, \widehat{y}_h^o; \boldsymbol{r}_1, w_1, \widehat{w}_1^o) = -(f, w_1)_{\mathcal{T}_h} - \langle u_h, \tau_2 w_1 + \boldsymbol{r}_1 \cdot \boldsymbol{n} \rangle_{\mathcal{E}_h^\partial},$$
(2.6a)

$$\mathcal{B}_{2}(\boldsymbol{p}_{h}, z_{h}, \widehat{z}_{h}^{o}; \boldsymbol{r}_{2}, w_{2}, \widehat{w}_{2}^{o}) = -(y_{h} - y_{d}, w_{2})_{\mathcal{T}_{h}}, \qquad (2.6b)$$

$$\langle \boldsymbol{p}_h \cdot \boldsymbol{n} + \tau_2(z_h - \hat{z}_h^o), \hat{w}_h^\partial \rangle_{\varepsilon_h^\partial} = \langle \gamma u_h, \hat{w}_h^\partial \rangle_{\varepsilon_h^\partial},$$
(2.6c)

for all $(\mathbf{r}_1, w_1, \widehat{w}_1^o, \mathbf{r}_2, w_2, \widehat{w}_2^o, \widehat{w}_h^\partial) \in [\mathbf{V}_h \times W_h \times M_h^o]^2 \times M_h^\partial.$

The following basic result, which is similar to results in [23, 29], is crucial to the proof of the well-posedness of the discrete optimality system (2.3a)-(2.3e), and is also a very important part of the final stage of numerical analysis (see the proof of Lemma 14).

Lemma 1. For all $(\boldsymbol{q}_h, y_h, \widehat{y}_h^o, \boldsymbol{r}_h, w_h, \widehat{w}_h^o) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2$, we have

$$\mathcal{B}_1(\boldsymbol{q}_h, y_h, \widehat{y}_h^o; \boldsymbol{r}_h, w_h, \widehat{w}_h^o) = \mathcal{B}_2(\boldsymbol{r}_h, w_h, \widehat{w}_h^o; \boldsymbol{q}_h, y_h, \widehat{y}_h^o).$$
(2.7)

Proof. Using the definitions in (2.4)-(2.5) and integration by parts give

$$\begin{aligned} \mathcal{B}_{1}(\boldsymbol{q}_{h},y_{h},\widehat{y}_{h}^{o};\boldsymbol{r}_{h},w_{h},\widehat{w}_{h}^{o}) &- \mathcal{B}_{2}(\boldsymbol{r}_{h},w_{h},\widehat{w}_{h}^{o};\boldsymbol{q}_{h},y_{h},\widehat{y}_{h}^{o}) \\ &= -\langle \tau_{1}(y_{h}-\widehat{y}_{h}^{o}),w_{h}-\widehat{w}_{h}^{o}\rangle_{\partial\mathcal{T}_{h}} \\ &+ \langle \tau_{2}(y_{h}-\widehat{y}_{h}^{o}),w_{h}-\widehat{w}_{h}^{o}\rangle_{\partial\mathcal{T}_{h}} \\ &+ (y_{h},\boldsymbol{\beta}\cdot\nabla w_{h})\tau_{h}-\langle\widehat{y}_{h}^{o},\boldsymbol{\beta}\cdot\boldsymbol{n}w_{h}\rangle_{\partial\mathcal{T}_{h}} - (\sigma y_{h},w_{h})\tau_{h} \\ &+ (w_{h},\boldsymbol{\beta}\cdot\nabla y_{h})\tau_{h}-\langle\widehat{w}_{h}^{o},\boldsymbol{\beta}\cdot\boldsymbol{n}y_{h}\rangle_{\partial\mathcal{T}_{h}} + ((\sigma+\nabla\cdot\boldsymbol{\beta})w_{h},y_{h})\tau_{h} \\ &= -\langle\boldsymbol{\beta}\cdot\boldsymbol{n}(y_{h}-\widehat{y}_{h}^{o}),w_{h}-\widehat{w}_{h}^{o}\rangle_{\partial\mathcal{T}_{h}} \\ &+ \langle y_{h},\boldsymbol{\beta}\cdot\boldsymbol{n}w_{h}\rangle_{\partial\mathcal{T}_{h}} - \langle\widehat{y}_{h}^{o},\boldsymbol{\beta}\cdot\boldsymbol{n}w_{h}\rangle_{\partial\mathcal{T}_{h}} - \langle\widehat{w}_{h}^{o},\boldsymbol{\beta}\cdot\boldsymbol{n}y_{h}\rangle_{\partial\mathcal{T}_{h}} \\ &= 0, \end{aligned}$$

where we used $\langle \boldsymbol{\beta} \cdot \boldsymbol{n}, \hat{y}_h^o \hat{w}_h^o \rangle_{\partial \mathcal{T}_h} = 0$. This proves our result.

3 Stability

To perform the stability and error analysis for the convection dominated boundary control problem, we need to assume some conditions on the velocity vector field $\boldsymbol{\beta}$ and the effective reaction function $\bar{\sigma} := \sigma + \frac{1}{2} \nabla \cdot \boldsymbol{\beta}$.

(A1) $\bar{\sigma}$ has a nonnegative lower bound, i.e,

$$\sigma_0 := \inf_{\boldsymbol{x} \in \Omega} \bar{\sigma} \ge 0. \tag{3.1}$$

(A2) β has no closed curves and

$$|\boldsymbol{\beta}(\boldsymbol{x})| \neq 0$$
 for all $\boldsymbol{x} \in \Omega$.

(A3) $\varepsilon < \min_{T \in \mathcal{T}_h} \{h_T\}.$

We note that we have already assumed (A1) in Equation (2.1) in Theorem 1. We repeat the assumption here to highlight it. Also, since we are interested in the convection dominated case, (A3) is a reasonable assumption. As shown in [2], assumption (A2) implies for any integer $k \ge 0$, there exists a function $\psi \in W^{k+1,\infty}(\Omega)$ such that for all $x \in \Omega$, we have

$$\boldsymbol{\beta} \cdot \nabla \psi \ge 2\beta_0 > 0, \tag{3.2}$$

where $\beta_0 := \|\beta\|_{0,\infty}/L$ and L is the diameter of Ω . We use assumption (A3) in the analysis to remove the assumption on the meshes. Specifically, in the proofs of Lemma 11 and Lemma 15, we use assumption (A3) and a local inverse inequality to replace a global inverse inequality that has been used in all previous Dirichlet boundary control works. Therefore, we only assume $\{\mathcal{T}_h\}$ is a conforming simplex partition of Ω . All previous works on Dirichlet boundary control problems required a conforming quasi-uniform mesh. In the future, we hope to performed an *a posteriori* error analysis for the convection dominated boundary control problem.

Remark 1. If $\sigma_0 \ge 0$, then assumption (A2) is the minimal known requirement that can be used to establish stability and error analysis results for numerical methods; see, e.g., [2, 20]. If instead $\sigma_0 > 0$, then we don't need to assume (A2) and the numerical analysis is less technical. Specifically, we don't need to prove Theorem 2 below if $\sigma_0 \ge 0$.

3.1 Preliminary material

For any nonnegative integer j, we define the L^2 -projections Π_j^o and Π_j^∂ as follows: for any $T \in \mathcal{T}_h$, $E \subset \partial T$, $v \in L^2(T)$, $q \in L^2(E)$, find $\Pi_j^o v \in \mathcal{P}_j(T)$ and $\Pi_j^\partial q \in \mathcal{P}_j(E)$ satisfying

$$(\Pi_j^o v, w_j)_T = (v, w_j)_T, \quad \forall w_j \in \mathcal{P}_j(T),$$
(3.3a)

$$\langle \Pi_{j}^{\partial} q, r_{j} \rangle_{E} = \langle q, r_{j} \rangle_{E}, \quad \forall r_{j} \in \mathcal{P}_{j}(E).$$
 (3.3b)

We also define $\widetilde{\Pi}_k^{\partial}$ as

$$\widetilde{\Pi}_{k}^{\partial}|_{E} = \begin{cases} \Pi_{k}^{\partial}|_{E}, & E \in \mathcal{E}_{h}^{o}, \\ 0, & E \in \mathcal{E}_{h}^{\partial}. \end{cases}$$

Then $\widetilde{\Pi}^\partial_k$ is an operator mapping $L^2(\mathcal{E}_h)$ to $M^o_h.$

We first give an approximation property from [5, Theorem 4.3.8, Proposition 4.1.9], and then we prove the basic stability and approximation properties for L^2 projections.

Lemma 2. Let $m \ge 1$ be an integer. For any $T \in \mathcal{T}_h$, $v \in H^m(T)$ and integer s satisfying $0 \le s \le m$, there exists $I_{m-1}v \in \mathcal{P}_{m-1}(T)$ such that

$$|v - I_{m-1}v|_{s,T} \le Ch_T^{m-s}|v|_{m,T}, \tag{3.4a}$$

$$\|v - I_{m-1}v\|_{0,\infty,T} \le Ch_T |v|_{1,\infty,T}.$$
(3.4b)

Lemma 3. Let s be a real number. For any nonnegative integer j, let m be a real number satisfying $\frac{1}{2} < m \leq j + 1$ and let $\ell \in \{0, 1\}$. For all $T \in \mathcal{T}_h$, $E \in \mathcal{E}_h$, it holds

$$|\Pi_{\ell}^{o}v|_{j,T} \le C|v|_{j,T}, \qquad \forall v \in H^{j}(T), \qquad (3.5a)$$

$$\|\Pi_j^{\partial} v\|_E \le \|v\|_E, \qquad \qquad \forall v \in L^2(E), \qquad (3.5b)$$

$$|v - \Pi_j^o v|_{s,T} \le Ch_T^{m-s} |v|_{m,T}, \qquad \forall v \in H^m(T), \ 0 \le s \le m,$$
(3.5c)

$$|v - \Pi_0^o v|_{\ell,\infty,T} \le Ch_T^{1-\ell} |v|_{1,\infty,T}, \qquad \forall v \in W^{1,\infty}(T), \qquad (3.5d)$$

$$|v - \Pi_{j}^{o}v|_{s,\partial T} \leq Ch_{T}^{m-s-1/2} |v|_{m,T}, \qquad \forall v \in H^{m}(T), 0 \leq s+1 \leq m,$$

$$||w||_{\partial T} \leq Ch_{T}^{-1/2} ||w||_{T}, \qquad \forall w \in W_{h}.$$

$$(3.5e)$$

Proof. Equation (3.5a) follows from Equation (3.5c); Equation (3.5b) follows from the definition of L^2 projection; Equation (3.5e) follows from Equation (3.5c) and the trace inequality; and Equation (3.5f) follows from the trace inequality and inverse inequality. The only thing left is to prove Equation (3.5c) and Equation (3.5d).

For Equation (3.5c), in view of Equation (3.4a), an inverse inequality, and the fact that $\|\Pi_i^o v\|_{0,T} \leq \|v\|_{0,T}$, for $1 \leq m \leq j+1$ we have

$$\begin{aligned} |v - \Pi_{j}^{o}v|_{s,T} &\leq |v - I_{m-1}v|_{s,T} + |I_{m-1}v - \Pi_{j}^{o}v|_{s,T} \\ &= |v - I_{m-1}v|_{s,T} + |\Pi_{j}^{o}(I_{m-1}v - v)|_{s,T} \\ &\leq |v - I_{m-1}v|_{s,T} + Ch_{T}^{-s} \|\Pi_{j}^{o}(I_{m-1}v - v)\|_{0,T} \\ &\leq |v - I_{m-1}v|_{s,T} + Ch_{T}^{-s} \|I_{m-1}v - v\|_{0,T} \\ &\leq Ch_{T}^{m-s} \|v\|_{m,T}. \end{aligned}$$

As for Equation (3.5d), $\ell = 1$ is obvious and therefore we set $\ell = 0$. By a standard scaling argument, the following stability result holds:

$$\|\Pi_0^o v\|_{0,\infty,T} \le C \|v\|_{0,\infty,T}.$$
(3.6)

By an inverse inequality, (3.6), and (3.4b) we get

$$\begin{aligned} \|v - \Pi_0^o v\|_{0,\infty,T} &\leq \|v - I_0 v\|_{0,\infty,T} + \|I_0 v - \Pi_0^o v\|_{0,\infty,T} \\ &= \|v - I_0 v\|_{0,\infty,T} + \|\Pi_0^o (I_0 v - v)\|_{0,\infty,T} \\ &\leq C \|v - I_0 v\|_{0,\infty,T} \\ &\leq C h_T |v|_{1,\infty,T}. \end{aligned}$$

In addition, we have the following super-approximation results; a similar result can be found in [2].

Lemma 4. Let $T \in \mathcal{T}_h, E \subset \partial T$. Then, for any $u_h \in \mathcal{P}_k(T)$ and $\eta \in W^{1,\infty}(T)$, there holds:

$$\|\eta u_h - \Pi_k^o(\eta u_h)\|_T \le Ch_T |\eta|_{1,\infty,T} \|u_h\|_T,$$
(3.7a)

$$|\eta u_h - \Pi_k^o(\eta u_h)|_{1,T} \le C |\eta|_{1,\infty,T} ||u_h||_T,$$
(3.7b)

$$\|\eta u_h - \Pi_k^o(\eta u_h)\|_{\partial T} \le C h_T^{1/2} |\eta|_{1,\infty,T} \|u_h\|_T,$$
(3.7c)

$$\|\eta u_h - \Pi_k^{\partial}(\eta u_h)\|_E \le Ch_T |\eta|_{1,\infty,T} \|u_h\|_E.$$
(3.7d)

Proof. We notice that (3.7c) follows from (3.7a), (3.7b), and the trace inequality. Next, for $j \in \{0, 1\}$, we have

This proves (3.7a) and (3.7b). Similarly, for $E \subset \partial T$, we have

$$\begin{aligned} \|\eta u_{h} - \Pi_{k}^{\partial}(\eta u_{h})\|_{E} &\leq \|\eta u_{h} - (\Pi_{0}^{o}\eta)u_{h}\|_{E} + \|(\Pi_{0}^{o}\eta)u_{h} - \Pi_{k}^{\partial}(\eta u_{h})\|_{E} \\ &= \|(\eta - \Pi_{0}^{o}\eta)u_{h}\|_{E} + \|\Pi_{k}^{\partial}((\Pi_{0}^{o}\eta)u_{h} - \eta u_{h})\|_{E} \\ &\leq C\|(\eta - \Pi_{0}^{o}\eta)u_{h}\|_{E} \\ &\leq C\|\eta - \Pi_{0}^{o}\eta\|_{0,\infty,T}\|u_{h}\|_{E} \\ &\leq Ch_{T}|\eta|_{1,\infty,T}\|u_{h}\|_{E} \end{aligned}$$
by (3.5d).

This proves (3.7d).

For the analysis of the low regularity case, we need the following result from [35]:

Lemma 5. If $\ell \geq 1$ is an integer that is large enough, then there exists an interpolation operator $\mathcal{I}_h^c: W_h \times M_h^o \to H_0^1(\Omega) \cap \mathcal{P}_{k+\ell}^{\mathcal{T}_h}$ such that for all $(w_h, \widehat{w}_h^o, v_h, \widehat{v}_h^o) \in [W_h \times M_h^o]^2$, for all $T \in \mathcal{T}_h$ and for all $E \in \mathcal{E}_h$, we have

$$(\mathcal{I}_h^c(w_h, \widehat{w}_h^o), v_h)_T = (w_h, v_h)_T, \tag{3.8a}$$

$$\langle \mathcal{I}_h^c(w_h, \widehat{w}_h^o), \widehat{v}_h^o \rangle_E = \langle \widehat{w}_h^o, \widehat{v}_h^o \rangle_E, \tag{3.8b}$$

$$\|\nabla \mathcal{I}_h^c(w_h, \widehat{w}_h^o)\|_{\mathcal{T}_h} \le C\left(\|\nabla w_h\|_{\mathcal{T}_h} + \|h_E^{-1/2}(w_h - \widehat{w}_h^o)\|_{\partial \mathcal{T}_h}\right),\tag{3.8c}$$

$$\|w_h - \mathcal{I}_h^c(w_h, \widehat{w}_h^o)\|_{\mathcal{T}_h} \le Ch\left(\|\nabla w_h\|_{\mathcal{T}_h} + \|h_E^{-1/2}(w_h - \widehat{w}_h^o)\|_{\partial \mathcal{T}_h}\right),\tag{3.8d}$$

where $\mathcal{P}_{k+\ell}^{\mathcal{T}_h} = \{ w_h \in L^2(\Omega) : w_h |_T \in \mathcal{P}_{k+\ell}(T), \ \forall T \in \mathcal{T}_h \}.$

3.2 Proof of the stability of (2.6)

Next, we present the stability of the above HDG method for the convection dominated Dirichlet boundary control problem. We follow a similar strategy to [2, 20]. We first collect some basic equalities and inequalities, which are used frequently in our paper.

Lemma 6. For all $(w_h, \widehat{w}_h^o) \in W_h \times M_h^o$, we have

$$(w_h, \boldsymbol{\beta} \cdot \nabla w_h)_{\mathcal{T}_h} = \frac{1}{2} \langle \boldsymbol{\beta} \cdot \boldsymbol{n} w_h, w_h \rangle_{\partial \mathcal{T}_h} - \frac{1}{2} (\nabla \cdot \boldsymbol{\beta} w_h, w_h)_{\mathcal{T}_h}, \qquad (3.9a)$$

$$\frac{1}{2}\langle \boldsymbol{\beta} \cdot \boldsymbol{n} w_h, w_h \rangle_{\partial \mathcal{T}_h} - \langle \widehat{w}_h^o, \boldsymbol{\beta} \cdot \boldsymbol{n} w_h \rangle_{\partial \mathcal{T}_h} = \frac{1}{2}\langle \boldsymbol{\beta} \cdot \boldsymbol{n} (w_h - \widehat{w}_h^o), w_h - \widehat{w}_h^o \rangle_{\partial \mathcal{T}_h},$$
(3.9b)

$$\|w_h\|_{\mathcal{T}_h}^2 \le C \|\nabla w_h\|_{\mathcal{T}_h}^2 + C \sum_{T \in \mathcal{T}_h} h_T^{-1} \|w_h - \widehat{w}_h^o\|_{\partial T}^2.$$
(3.9c)

The identity (3.9a) can be obtained by integration by parts and the proof of (3.9b) follows from the fact $\langle \boldsymbol{\beta} \cdot \boldsymbol{n} \hat{w}_h^o, \hat{w}_h^o \rangle_{\partial \mathcal{T}_h} = 0$. For the last inequality (3.9c), we refer to [39, page 354] for the proof. Next, we define some seminorms:

Definition 1. For all $(\boldsymbol{q}_h, y_h, \widehat{y}_h^o) \in \boldsymbol{V}_h \times W_h \times M_h^o$, define

$$\|(y_h, \hat{y}_h^o)\|_{W,w}^2 := \varepsilon \|\nabla y_h\|_{\mathcal{T}_h}^2 + \|\tau^{1/2}(y_h - \hat{y}_h^o)\|_{\partial \mathcal{T}_h}^2 + \|\bar{\sigma}^{1/2}y_h\|_{\mathcal{T}_h}^2,$$
(3.10a)

$$\|(y_h, \hat{y}_h^o)\|_W^2 := \varepsilon \|\nabla y_h\|_{\mathcal{T}_h}^2 + \|\tau^{1/2}(y_h - \hat{y}_h^o)\|_{\partial \mathcal{T}_h}^2 + \|(\beta_0 + \bar{\sigma})^{1/2}y_h\|_{\mathcal{T}_h}^2, \tag{3.10b}$$

$$\|(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o})\|_{w}^{2} \coloneqq \varepsilon^{-1} \|\boldsymbol{q}_{h}\|_{\mathcal{T}_{h}}^{2} + \|(y_{h}, \widehat{y}_{h}^{o})\|_{W, w}^{2}, \qquad (3.10c)$$

$$\|(\boldsymbol{q}_h, y_h, \hat{y}_h^o)\|^2 := \varepsilon^{-1} \|\boldsymbol{q}_h\|_{\mathcal{T}_h}^2 + \|(y_h, \hat{y}_h^o)\|_W^2.$$
(3.10d)

It is easy to see that the seminorm $\|(\cdot, \cdot)\|_W$ is a norm since $\beta_0 > 0$, hence $\|(\cdot, \cdot, \cdot)\|$ is also a norm. To prove the seminorms $\|(\cdot, \cdot)\|_{W,w}$ and $\|(\cdot, \cdot, \cdot)\|_w$ are norms, we just need to show $\|(\cdot, \cdot)\|_{W,w}$ is a norm.

Lemma 7. $\|(\cdot, \cdot)\|_{W,w}$ is a norm for the space $W_h \times M_h^o$.

 (\boldsymbol{r}_h, w_h)

Proof. It is obvious that we only need to show that $||(y_h, \hat{y}_h^o)||_{W,w} = 0$ implies $\hat{y}_h^o|_{\Gamma} = y_h = 0$. This is true because y_h is piecewise constant on \mathcal{T}_h and $y_h = \hat{y}_h^o$ on \mathcal{E}_h ; therefore, $y_h = \hat{y}_h^o$ are constants. Since $\hat{y}_h^o|_{\Gamma} = 0$, we have $y_h = \hat{y}_h^o = 0$.

Lemma 8. (Stability in weak norm) For all $(\boldsymbol{q}_h, y_h, \widehat{y}_h^o) \in \boldsymbol{V}_h \times W_h \times M_h^o$, the following stability results hold:

$$\sup_{\widehat{w}_{h}^{o})\in \boldsymbol{V}_{h}\times W_{h}\times M_{h}^{o}}\frac{\mathcal{B}_{1}(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}; \boldsymbol{r}_{h}, w_{h}, \widehat{w}_{h}^{o})}{\left\|(\boldsymbol{r}_{h}, w_{h}, \widehat{w}_{h}^{o})\right\|_{w}} \geq C\left\|(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o})\right\|_{w},$$
(3.11a)

$$\sup_{(\boldsymbol{r}_h, w_h, \widehat{w}_h^o) \in \boldsymbol{V}_h \times W_h \times M_h^o} \frac{\mathcal{B}_2(\boldsymbol{q}_h, y_h, \widehat{y}_h^o; \boldsymbol{r}_h, w_h, \widehat{w}_h^o)}{\left\| (\boldsymbol{r}_h, w_h, \widehat{w}_h^o) \right\|_w} \ge C \left\| (\boldsymbol{q}_h, y_h, \widehat{y}_h^o) \right\|_w.$$
(3.11b)

Proof. We only prove the first inequality; the second can be obtained by the same argument. First, let $(\mathbf{r}_h, w_h, \widehat{w}_h^o) = (q_h, -y_h, -\widehat{y}_h^o)$ in the definition of \mathcal{B}_1 in (2.4) to get

$$\begin{aligned} \mathcal{B}_{1}(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}; \boldsymbol{q}_{h}, -y_{h}, -\widehat{y}_{h}^{o}) \\ &= \varepsilon^{-1} \|\boldsymbol{q}_{h}\|_{\mathcal{T}_{h}}^{2} + \langle \tau_{1}(y_{h} - \widehat{y}_{h}^{o}), y_{h} - \widehat{y}_{h}^{o} \rangle_{\partial \mathcal{T}_{h}} \\ &- (y_{h}, \boldsymbol{\beta} \nabla y_{h})_{\mathcal{T}_{h}} + \langle \widehat{y}_{h}^{o}, \boldsymbol{\beta} \cdot \boldsymbol{n} y_{h} \rangle_{\partial \mathcal{T}_{h}} + (\sigma y_{h}, y_{h})_{\mathcal{T}_{h}} \\ &= \varepsilon^{-1} \|\boldsymbol{q}_{h}\|_{\mathcal{T}_{h}}^{2} + \langle \tau_{1}(y_{h} - \widehat{y}_{h}^{o}), y_{h} - \widehat{y}_{h}^{o} \rangle_{\partial \mathcal{T}_{h}} \\ &- \frac{1}{2} \langle \boldsymbol{\beta} \cdot \boldsymbol{n} y_{h}, y_{h} \rangle_{\partial \mathcal{T}_{h}} + \langle \widehat{y}_{h}^{o}, \boldsymbol{\beta} \cdot \boldsymbol{n} y_{h} \rangle_{\partial \mathcal{T}_{h}} + ((\sigma + \frac{1}{2} \nabla \cdot \boldsymbol{\beta}) y_{h}, y_{h})_{\mathcal{T}_{h}} \quad \text{by (3.9a),} \\ &= \varepsilon^{-1} \|\boldsymbol{q}_{h}\|_{\mathcal{T}_{h}}^{2} + \langle \tau_{1}(y_{h} - \widehat{y}_{h}^{o}), y_{h} - \widehat{y}_{h}^{o} \rangle_{\partial \mathcal{T}_{h}} \\ &- \frac{1}{2} \langle \boldsymbol{\beta} \cdot \boldsymbol{n}(y_{h} - \widehat{y}_{h}^{o}), y_{h} - \widehat{y}_{h}^{o} \rangle_{\partial \mathcal{T}_{h}} + ((\sigma + \frac{1}{2} \nabla \cdot \boldsymbol{\beta}) y_{h}, y_{h})_{\mathcal{T}_{h}} \quad \text{by (3.9b),} \\ &= \varepsilon^{-1} \|\boldsymbol{q}_{h}\|_{\mathcal{T}_{h}}^{2} + \|\sqrt{\tau}(y_{h} - \widehat{y}_{h}^{o})\|_{\partial \mathcal{T}_{h}}^{2} + \|\overline{\sigma}^{1/2} y_{h}\|_{\mathcal{T}_{h}}^{2} \quad \text{by (3.1).} \end{aligned}$$

Next, let $(\boldsymbol{r}_h, w_h, \widehat{w}_h^o) = (\varepsilon \nabla y_h, 0, 0)$ to get

$$\mathcal{B}_{1}(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}; \varepsilon \nabla y_{h}, 0, 0) = (\boldsymbol{q}_{h}, \nabla y_{h})_{\mathcal{T}_{h}} - \varepsilon (y_{h}, \nabla \cdot \nabla y_{h})_{\mathcal{T}_{h}} + \varepsilon \langle \widehat{y}_{h}^{o}, \nabla y_{h} \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_{h}} = \varepsilon \|\nabla y_{h}\|_{\mathcal{T}_{h}}^{2} + \varepsilon \langle \widehat{y}_{h}^{o} - y_{h}, \nabla y_{h} \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_{h}} + (\boldsymbol{q}_{h}, \nabla y_{h})_{\mathcal{T}_{h}} \\ \geq \varepsilon \|\nabla y_{h}\|_{\mathcal{T}_{h}}^{2} - \varepsilon^{1/2} h^{-1/2} \|\widehat{y}_{h}^{o} - y_{h}\|_{\partial \mathcal{T}_{h}} \varepsilon^{1/2} \|\nabla y_{h}\|_{\mathcal{T}_{h}} - \|\boldsymbol{q}_{h}\|_{\mathcal{T}_{h}} \|\nabla y_{h}\|_{\mathcal{T}_{h}} \\ \geq \frac{\varepsilon}{2} \|\nabla y_{h}\|_{\mathcal{T}_{h}}^{2} - C_{0} (\|\boldsymbol{\tau}^{1/2}(\widehat{y}_{h}^{o} - y_{h})\|_{\partial \mathcal{T}_{h}}^{2} + \varepsilon^{-1} \|\boldsymbol{q}_{h}\|_{\mathcal{T}_{h}}^{2}), \qquad (3.13)$$

where C_0 is a fixed positive constant. The definitions of $\|(\cdot, \cdot, \cdot)\|$ in (3.10d) and $\|(\cdot, \cdot, \cdot)\|_w$ in (3.10c) imply

$$\|(\varepsilon \nabla y_h, 0, 0)\| = \|(\varepsilon \nabla y_h, 0, 0)\|_w \le C \|(\boldsymbol{q}_h, y_h, \widehat{y}_h^o)\|_w.$$
(3.14)

Finally, we take $(\mathbf{r}_h, w_h, \widehat{w}_h^o) = (\frac{1}{2} + C_0)(\mathbf{q}_h, -y_h, -\widehat{y}_h^o) + (\varepsilon \nabla y_h, 0, 0)$ to obtain

$$\begin{aligned} &\mathcal{B}_{1}(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}; \boldsymbol{r}_{h}, w_{h}, \widehat{w}_{h}^{o}) \\ &\geq \frac{1}{2} \left\| (\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}) \right\|_{w}^{2} & \text{by (3.12) and (3.13),} \\ &\geq C \left\| (\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}) \right\|_{w} \left\| (\boldsymbol{r}_{h}, w_{h}, \widehat{w}_{h}^{o}) \right\|_{w} & \text{by (3.14).} \end{aligned}$$

This completes our proof.

For later use, by (3.14), for any $(\boldsymbol{q}_h, y_h, \hat{y}_h^o) \in \boldsymbol{V}_h \times W_h \times M_h^o$, we have

$$\|(\boldsymbol{r}_h, w_h, \widehat{w}_h^o)\| \le C \|(\boldsymbol{q}_h, y_h, \widehat{y}_h^o)\| + \|(\varepsilon \nabla y_h, 0, 0)\| \le C \|(\boldsymbol{q}_h, y_h, \widehat{y}_h^o)\|.$$
(3.16)

Remark 2. The existence of a unique solution to the HDG discretization (2.6) of the optimality system now follows similarly to [29]; we omit the details. Also, to obtain the L^2 error estimates for the state y_h , Lemma 8 is not sufficient since the effective reaction term $\bar{\sigma}^{1/2}$ can equal zero at some points; therefore, it is possible for the term $\|\bar{\sigma}^{1/2}y_h\|_{\mathcal{T}_h}$ in the definition of $\|(\boldsymbol{q}_h, y_h, \hat{y}_h^o)\|_w$ to equal zero for some y_h . Therefore, we need a refined analysis technique to derive a strong stability result that contains the norm $\|y_h\|_{\mathcal{T}_h}$.

Theorem 2. (Stability in strong norm) If assumptions (A1) and (A2) hold, then there exists h_0 , independent of ε , such that the following stability results hold: for all $(\mathbf{q}_h, y_h, \hat{y}_h^o) \in \mathbf{V}_h \times W_h \times M_h^o$ with $h \leq h_0$,

$$\sup_{(\boldsymbol{r}_h, w_h, \widehat{w}_h^o) \in \boldsymbol{V}_h \times W_h \times M_h^o} \frac{\mathcal{B}_1(\boldsymbol{q}_h, y_h, \widehat{y}_h^o; \boldsymbol{r}_h, w_h, \widehat{w}_h^o)}{\left\| (\boldsymbol{r}_h, w_h, \widehat{w}_h^o) \right\|} \ge C \left\| (\boldsymbol{q}_h, y_h, \widehat{y}_h^o) \right\|,$$
(3.17a)

$$\sup_{(\boldsymbol{r}_h, w_h, \widehat{w}_h^o) \in \boldsymbol{V}_h \times W_h \times M_h^o} \frac{\mathcal{B}_2(\boldsymbol{q}_h, y_h, \widehat{y}_h^o; \boldsymbol{r}_h, w_h, \widehat{w}_h^o)}{\left\| (\boldsymbol{r}_h, w_h, \widehat{w}_h^o) \right\|} \ge C \left\| (\boldsymbol{q}_h, y_h, \widehat{y}_h^o) \right\|.$$
(3.17b)

Proof. We only prove (3.17a), and we split the proofs into two steps.

Step 1: Let $\psi \in W^{1,\infty}(\Omega)$ satisfy (3.2). We take

$$(\mathbf{r}_h, w_h, \widehat{w}_h^o) = (\mathbf{r}_1, w_1, \widehat{w}_1) = (\mathbf{0}, -e^{-\psi} y_h, -e^{-\psi} \widehat{y}_h^o)$$
(3.18)

in the definition of \mathcal{B}_1 in (2.4) to obtain

$$\begin{aligned} \mathcal{B}_{1}(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}; \boldsymbol{r}_{1}, w_{1}, \widehat{w}_{1}) \\ &= \left[(e^{-\psi}y_{h}, \nabla \cdot \boldsymbol{q}_{h})_{\mathcal{T}_{h}} - \langle e^{-\psi}\widehat{y}_{h}^{o}, \boldsymbol{q}_{h} \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_{h}} \right] + \langle \tau_{1}(y_{h} - \widehat{y}_{h}^{o}), e^{-\psi}y_{h} - e^{-\psi}\widehat{y}_{h}^{o} \rangle_{\partial \mathcal{T}_{h}} \\ &+ \left[-(y_{h}, \boldsymbol{\beta} \cdot \nabla (e^{-\psi}y_{h}))_{\mathcal{T}_{h}} + \langle \widehat{y}_{h}^{o}, \boldsymbol{\beta} \cdot \boldsymbol{n} e^{-\psi}y_{h} \rangle_{\partial \mathcal{T}_{h}} \right] + (\sigma y_{h}, e^{-\psi}y_{h})_{\mathcal{T}_{h}} \\ &= S_{1} + S_{2} + S_{3} + S_{4}. \end{aligned}$$

Next, we estimate $\{S_i\}_{i=1}^4$ term by term. First,

$$S_{1} = -(\nabla(e^{-\psi}y_{h}), \mathbf{q}_{h})_{\mathcal{T}_{h}} - \langle e^{-\psi}(\hat{y}_{h}^{o} - y_{h}), \mathbf{q}_{h} \cdot \mathbf{n} \rangle_{\partial \mathcal{T}_{h}}$$

$$= -(y_{h}\nabla e^{-\psi} + e^{-\psi}\nabla y_{h}, \mathbf{q}_{h})_{\mathcal{T}_{h}} - \langle e^{-\psi}(\hat{y}_{h}^{o} - y_{h}), \mathbf{q}_{h} \cdot \mathbf{n} \rangle_{\partial \mathcal{T}_{h}}$$

$$\leq C(\varepsilon^{1/2} \|y_{h}\|_{\mathcal{T}_{h}} + \varepsilon^{1/2} \|\nabla y_{h}\|_{\mathcal{T}_{h}} + \|\tau^{1/2}(\hat{y}_{h}^{o} - y_{h})\|_{\partial \mathcal{T}_{h}})\varepsilon^{-1/2} \|\mathbf{q}_{h}\|_{\mathcal{T}_{h}}$$

$$\leq C(\varepsilon^{1/2} \|\nabla y_{h}\|_{\mathcal{T}_{h}} + \|\tau^{1/2}(\hat{y}_{h}^{o} - y_{h})\|_{\partial \mathcal{T}_{h}})\varepsilon^{-1/2} \|\mathbf{q}_{h}\|_{\mathcal{T}_{h}} \qquad \text{by (3.9c),}$$

$$\leq C \|(y_{h}, \hat{y}_{h}^{o})\|_{W,w} \varepsilon^{-1/2} \|\mathbf{q}_{h}\|_{\mathcal{T}_{h}} \qquad \text{by (3.10a),}$$

$$\leq C \|(\mathbf{q}_{h}, y_{h}, \hat{y}_{h}^{o})\|_{w}^{2} \qquad \text{by (3.10c).}$$

$$\leq O ||(\mathbf{q}h, gh, g_h)||_w$$

Second, to estimate the term S_3 , let $\varphi = e^{-\psi}$ in (3.9a) to obtain

$$-(y_h, \boldsymbol{eta} \cdot
abla (e^{-\psi}y_h))_{\mathcal{T}_h} = rac{1}{2} (
abla \cdot \boldsymbol{eta} e^{-\psi}y_h, y_h)_{\mathcal{T}_h} + rac{1}{2} (y_h, e^{-\psi}y_h \boldsymbol{eta} \cdot
abla \psi)_{\mathcal{T}_h}
onumber \ -rac{1}{2} \langle y_h, \boldsymbol{eta} \cdot \boldsymbol{n} e^{-\psi}y_h
angle_{\partial \mathcal{T}_h}.$$

Hence,

$$\begin{split} S_{3} &= -(y_{h}, \boldsymbol{\beta} \cdot \nabla(e^{-\psi}y_{h}))_{\mathcal{T}_{h}} + \langle \widehat{y}_{h}^{o}, \boldsymbol{\beta} \cdot \boldsymbol{n} e^{-\psi}y_{h} \rangle_{\partial \mathcal{T}_{h}} \\ &= \frac{1}{2} (\nabla \cdot \boldsymbol{\beta} e^{-\psi}y_{h}, y_{h})_{\mathcal{T}_{h}} + \frac{1}{2} (y_{h}, e^{-\psi}y_{h} \boldsymbol{\beta} \cdot \nabla \psi)_{\mathcal{T}_{h}} \\ &- \frac{1}{2} \langle y_{h}, \boldsymbol{\beta} \cdot \boldsymbol{n} e^{-\psi}y_{h} \rangle_{\partial \mathcal{T}_{h}} + \langle \widehat{y}_{h}^{o}, \boldsymbol{\beta} \cdot \boldsymbol{n} e^{-\psi}y_{h} \rangle_{\partial \mathcal{T}_{h}} \\ &= \frac{1}{2} (\nabla \cdot \boldsymbol{\beta} e^{-\psi}y_{h}, y_{h})_{\mathcal{T}_{h}} + \frac{1}{2} (e^{-\psi}\boldsymbol{\beta} \cdot \nabla \psi y_{h}, y_{h})_{\mathcal{T}_{h}} \\ &- \frac{1}{2} \langle \boldsymbol{\beta} \cdot \boldsymbol{n} e^{-\psi}(y_{h} - \widehat{y}_{h}^{o}), y_{h} - \widehat{y}_{h}^{o} \rangle_{\partial \mathcal{T}_{h}} \\ &\geq \frac{1}{2} (\nabla \cdot \boldsymbol{\beta} e^{-\psi}y_{h}, y_{h})_{\mathcal{T}_{h}} + (\beta_{0} e^{-\psi}y_{h}, y_{h})_{\mathcal{T}_{h}} \\ &- \frac{1}{2} \langle \boldsymbol{\beta} \cdot \boldsymbol{n} e^{-\psi}(y_{h} - \widehat{y}_{h}^{o}), y_{h} - \widehat{y}_{h}^{o} \rangle_{\partial \mathcal{T}_{h}}. \end{split}$$
by (3.2),

Therefore, by the definition of τ in (2.3h), there exist a positive constant C_0 such that

$$S_{2} + S_{3} + S_{4} \geq \|e^{-\psi/2}(\beta_{0} + \bar{\sigma})^{1/2}y_{h}\|_{\mathcal{T}_{h}}^{2} + \|e^{-\psi/2}\tau^{1/2}(y_{h} - \hat{y}_{h}^{o})\|_{\partial\mathcal{T}_{h}}^{2}$$

$$\geq C_{0}\|(\beta_{0} + \bar{\sigma})^{1/2}y_{h}\|_{\mathcal{T}_{h}}^{2}.$$

This implies that there exist positive constants C_1 and C_2 such that

$$\mathcal{B}_{1}(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}; \boldsymbol{r}_{1}, w_{1}, \widehat{w}_{1}) \geq C_{1} \| (\beta_{0} + \bar{\sigma})^{1/2} y_{h} \|_{\mathcal{T}_{h}}^{2} - C_{2} \| (\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}) \|_{w}^{2}.$$
(3.19)

Moreover, we have

$$\begin{aligned} \|(\boldsymbol{r}_{1}, w_{1}, \widehat{w}_{1})\|^{2} &= \left\| (\boldsymbol{0}, -e^{-\psi}y_{h}, -e^{-\psi}\widehat{y}_{h}^{o}) \right\| & \text{by (3.18),} \\ &= \varepsilon \|\nabla(e^{-\psi}y_{h})\|_{\mathcal{T}_{h}}^{2} + \|\tau^{1/2}e^{-\psi}(y_{h} - \widehat{y}_{h}^{o})\|_{\partial\mathcal{T}_{h}}^{2} \\ &+ \|(\beta_{0} + \overline{\sigma})^{1/2}e^{-\psi}y_{h}\|_{\mathcal{T}_{h}}^{2} & \text{by (3.10d),} \\ &\leq C \|(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o})\|^{2} & \text{by (3.9c),} \end{aligned}$$

where we used $\nabla(e^{-\psi}y_h) = y_h \nabla e^{-\psi} + e^{-\psi} \nabla y_h$ in (3.20). **Step 2:** Let $R_k^o = \mathbb{I} - \Pi_k^o$ and $\widetilde{R}_k^\partial = \mathbb{I} - \widetilde{\Pi}_k^\partial$, where \mathbb{I} is the identity operator. We take

$$(\mathbf{r}_{h}, w_{h}, \widehat{w}_{h}^{o}) = (\mathbf{r}_{2}, w_{2}, \widehat{w}_{2}) = (\mathbf{0}, R_{k}^{o}(e^{-\psi}y_{h}), \widetilde{R}_{k}^{\partial}(e^{-\psi}\widehat{y}_{h}^{o}))$$
(3.21)

in the definition of \mathcal{B}_1 and use the orthogonality properties of Π_k^o and $\widetilde{\Pi}_k^\partial$, integration by parts, and $\tau_1 = \tau_2 + \boldsymbol{\beta} \cdot \boldsymbol{n}$ to get

$$\mathcal{B}_{1}(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}; \boldsymbol{r}_{2}, w_{2}, \widehat{w}_{2}) = -(R_{k}^{o}(e^{-\psi}y_{h}), \nabla \cdot \boldsymbol{q}_{h})_{\mathcal{T}_{h}} + \langle \widetilde{R}_{k}^{\partial}(e^{-\psi}\widehat{y}_{h}^{o}), \boldsymbol{q}_{h} \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_{h}} \\ - \langle \tau_{1}(y_{h} - \widehat{y}_{h}^{o}), R_{k}^{o}(e^{-\psi}y_{h}) - \widetilde{R}_{k}^{\partial}(e^{-\psi}\widehat{y}_{h}^{o}) \rangle_{\partial \mathcal{T}_{h}} \\ + (y_{h}, \boldsymbol{\beta} \cdot \nabla R_{k}^{o}(e^{-\psi}y_{h}))_{\mathcal{T}_{h}} - \langle \widehat{y}_{h}^{o}, \boldsymbol{\beta} \cdot \boldsymbol{n} R_{k}^{o}(e^{-\psi}y_{h}) \rangle_{\partial \mathcal{T}_{h}} \\ - (\sigma y_{h}, R_{k}^{o}(e^{-\psi}y_{h}))_{\mathcal{T}_{h}}.$$
(3.22)

The definitions of Π^o_k and $\widetilde{\Pi}^\partial_k$ in (3.3) imply

$$(R_k^o(e^{-\psi}y_h), \nabla \cdot \boldsymbol{q}_h)_{\mathcal{T}_h} = 0, \text{ and } \langle \widetilde{R}_k^\partial(e^{-\psi}\widehat{y}_h^o), \boldsymbol{q}_h \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_h} = 0.$$

Next, integration by parts gives

$$\begin{split} (y_h, \boldsymbol{\beta} \cdot \nabla R_k^o(e^{-\psi}y_h))_{\mathcal{T}_h} \\ &= \langle \boldsymbol{\beta} \cdot \boldsymbol{n} y_h, R_k^o(e^{-\psi}y_h) \rangle_{\partial \mathcal{T}_h} - (\boldsymbol{\beta} \nabla y_h, R_k^o(e^{-\psi}y_h))_{\mathcal{T}_h} - (\nabla \cdot \boldsymbol{\beta} y_h, R_k^o(e^{-\psi}y_h))_{\mathcal{T}_h} \\ &= \langle \boldsymbol{\beta} \cdot \boldsymbol{n} (y_h - \hat{y}_h^o), R_k^o(e^{-\psi}y_h) \rangle_{\partial \mathcal{T}_h} + \langle \boldsymbol{\beta} \cdot \boldsymbol{n} \hat{y}_h^o, R_k^o(e^{-\psi}y_h) \rangle_{\partial \mathcal{T}_h} \\ &- (\boldsymbol{\beta} \cdot \nabla y_h, R_k^o(e^{-\psi}y_h))_{\mathcal{T}_h} - (\nabla \cdot \boldsymbol{\beta} y_h, R_k^o(e^{-\psi}y_h))_{\mathcal{T}_h} \\ &= \langle \boldsymbol{\beta} \cdot \boldsymbol{n} (y_h - \hat{y}_h^o), R_k^o(e^{-\psi}y_h) - \widetilde{R}_k^\partial(e^{-\psi} \hat{y}_h^o) \rangle_{\partial \mathcal{T}_h} + \langle \boldsymbol{\beta} \cdot \boldsymbol{n} (y_h - \hat{y}_h^o), \widetilde{R}_k^\partial(e^{-\psi} \hat{y}_h^o) \rangle_{\partial \mathcal{T}_h} \\ &+ \langle \boldsymbol{\beta} \cdot \boldsymbol{n} \hat{y}_h^o, R_k^o(e^{-\psi}y_h) \rangle_{\partial \mathcal{T}_h} - (\boldsymbol{\beta} \cdot \nabla y_h, R_k^o(e^{-\psi}y_h))_{\mathcal{T}_h} - (\nabla \cdot \boldsymbol{\beta} y_h, R_k^o(e^{-\psi}y_h))_{\mathcal{T}_h}. \end{split}$$

Using $\tau_1 = \tau_2 + \boldsymbol{\beta} \cdot \boldsymbol{n}$ along with (3.22) and the above equalities give

$$\mathcal{B}_{1}(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}; \boldsymbol{r}_{2}, w_{2}, \widehat{w}_{2})
= -\langle \tau_{2}(y_{h} - \widehat{y}_{h}^{o}), R_{k}^{o}(e^{-\psi}y_{h}) - \widetilde{R}_{k}^{\partial}(e^{-\psi}\widehat{y}_{h}^{o})\rangle_{\partial\tau_{h}}
+ \langle y_{h} - \widehat{y}_{h}^{o}, \boldsymbol{\beta} \cdot \boldsymbol{n}\widetilde{R}_{k}^{\partial}(e^{-\psi}\widehat{y}_{h}^{o})\rangle_{\partial\tau_{h}} - (\boldsymbol{\beta} \cdot \nabla y_{h}, R_{k}^{o}(e^{-\psi}y_{h}))_{\tau_{h}}
- ((\boldsymbol{\sigma} + \nabla \cdot \boldsymbol{\beta})y_{h}, R_{k}^{o}e^{-\psi}y_{h})_{\tau_{h}}
= T_{1} + T_{2} + T_{3} + T_{4}.$$
(3.23)

Before we estimate $\{T_i\}_{i=1}^4$, we first define $R_k^\partial = \mathbb{I} - \Pi_k^\partial$ and estimate the following term:

$$\begin{aligned} \| (R_{k}^{o}(e^{-\psi}y_{h}) - R_{k}^{\partial}(e^{-\psi}\widehat{y}_{h}^{o})) \|_{\partial \mathcal{T}_{h}}^{2} \\ &\leq C \sum_{T \in \mathcal{T}_{h}} \left(\| R_{k}^{o}(e^{-\psi}y_{h}) \|_{\partial T}^{2} + \| R_{k}^{\partial}(e^{-\psi}\widehat{y}_{h}^{o}) \|_{\partial T}^{2} \right) \\ &\leq C \sum_{T \in \mathcal{T}_{h}} \left(h_{T} \| y_{h} \|_{T}^{2} + h_{T}^{2} \| y_{h} - \widehat{y}_{h}^{o} \|_{\partial T}^{2} \right) \qquad \text{by } (3.7c) - (3.7d), \\ &\leq Ch^{2} \| y_{h} - \widehat{y}_{h}^{o} \|_{\partial \mathcal{T}_{h}}^{2} + Ch \| y_{h} \|_{\mathcal{T}_{h}}^{2}. \end{aligned}$$

Therefore,

$$|T_1| \le Ch^2 \|\tau^{1/2} (y_h - \hat{y}_h^o)\|_{\partial \mathcal{T}_h}^2 + Ch \|(\beta_0 + \bar{\sigma})^{1/2} y_h\|_{\mathcal{T}_h}^2.$$
(3.25)

Next, by the definition of ${\cal R}^o_k,$ we have

$$T_{2} + T_{3} + T_{4} = (\boldsymbol{\beta} \cdot \nabla y_{h}, R_{k}^{o}(e^{-\psi}y_{h}))_{\mathcal{T}_{h}} + \langle y_{h} - \widehat{y}_{h}^{o}, \boldsymbol{\beta} \cdot \boldsymbol{n}\widetilde{R}_{k}^{\partial}(e^{-\psi}\widehat{y}_{h}^{o})\rangle_{\partial\mathcal{T}_{h}}$$
$$- ((\boldsymbol{\sigma} + \nabla \cdot \boldsymbol{\beta})y_{h}, R_{k}^{o}(e^{-\psi}y_{h}))_{\mathcal{T}_{h}}$$
$$= (R_{0}^{o}(\boldsymbol{\beta}) \cdot \nabla y_{h}, R_{k}^{o}(e^{-\psi}y_{h}))_{\mathcal{T}_{h}} + \langle y_{h} - \widehat{y}_{h}^{o}, \boldsymbol{\beta} \cdot \boldsymbol{n}\widetilde{R}_{k}^{\partial}(e^{-\psi}\widehat{y}_{h}^{o})\rangle_{\partial\mathcal{T}_{h}}$$
$$+ ((\boldsymbol{\sigma} + \nabla \cdot \boldsymbol{\beta})y_{h}, R_{k}^{o}(e^{-\psi}y_{h})).$$

Hence,

$$T_{2} + T_{3} + T_{4} \leq C \left(\sum_{T \in \mathcal{T}_{h}} h_{T}^{2} \| \nabla y_{h} \|_{T}^{2} \right)^{1/2} h \| y_{h} \|_{\mathcal{T}_{h}} \qquad \text{by (3.7a),}$$

$$+ C \left(\sum_{T \in \mathcal{T}_{h}} \| y_{h} - \widehat{y}_{h}^{o} \|_{\partial T}^{2} \right)^{1/2} \left(\sum_{T \in \mathcal{T}_{h}} h_{T}^{2} \| y_{h} \|_{\partial T}^{2} \right)^{1/2} \qquad \text{by (3.7d),}$$

$$+ Ch(\| \boldsymbol{\beta} \|_{1,\infty} + \| \bar{\sigma} \|_{0,\infty}) \| y_{h} \|_{\mathcal{T}_{h}}^{2} \qquad \text{by (3.7a),}$$

$$\leq Ch \| (\boldsymbol{\beta} - \| \bar{\sigma} \|_{0,\infty}) \| y_{h} \|_{\mathcal{T}_{h}}^{2} + \| \bar{\sigma}^{1/2} (y_{h} - \widehat{\sigma}^{2}) \|^{2} \qquad (2.26)$$

$$\leq Ch \| (\beta_0 + \bar{\sigma})^{1/2} y_h \|_{\mathcal{T}_h}^2 + \| \tau^{1/2} (y_h - \hat{y}_h^o) \|_{\partial \mathcal{T}_h}^2.$$
(3.26)

From (3.23), (3.25) and (3.26), we get

$$\mathcal{B}_{1}(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}; \boldsymbol{r}_{2}, w_{2}, \widehat{w}_{2}^{o}) \geq -C_{3}h \|(\beta_{0} + \bar{\sigma})^{1/2}y_{h}\|_{\mathcal{T}_{h}}^{2} - C_{4} \|(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o})\|_{w}^{2}.$$
(3.27)

Using (3.24), we have

By (3.16), there exists a $(\boldsymbol{r}_0, w_0, \widehat{w}_0^o) \in \boldsymbol{V}_h \times W_h \times M_h^o$ such that

$$\mathcal{B}_{1}(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}; \boldsymbol{r}_{0}, w_{0}, \widehat{w}_{0}^{o}) \geq \|(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o})\|_{w}^{2}, \qquad (3.29a)$$

 $\|(\boldsymbol{r}_{0}, w_{0}, \widehat{w}_{0}^{o})\| \leq C \|(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o})\|.$ (3.29b)

Take h small enough so that we have $C_3h \leq C_1/2$. Set $C_* = C_1 + C_2 + C_4$ and

$$(\mathbf{r}_h, w_h, \widehat{w}_h^o) = C_*(\mathbf{r}_0, w_0, \widehat{w}_0^o) + (\mathbf{r}_1, w_1, \widehat{w}_1^o) + (\mathbf{r}_2, w_2, \widehat{w}_2^o).$$
(3.30)

By (3.19), (3.27) and (3.29), we get

$$\mathcal{B}_{1}(\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}; \boldsymbol{r}_{h}, w_{h}, \widehat{w}_{h}^{o}) \geq C \left\| (\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}) \right\|^{2} \geq C \left\| (\boldsymbol{q}_{h}, y_{h}, \widehat{y}_{h}^{o}) \right\| \cdot \left\| (\boldsymbol{r}_{h}, w_{h}, \widehat{w}_{h}^{o}) \right\|,$$

which implies (3.17a).

4 Error analysis

Next, we perform a convergence analysis for the convection dominated Dirichlet boundary control problem.

4.1 Assumptions and main result

Throughout, we assume Ω is a bounded convex polyhedral domain. Therefore, in the 2D case the largest interior angle ω satisfies $\pi/3 \leq \omega < \pi$. Moreover, we assume the velocity vector field β and σ satisfy

$$\boldsymbol{\beta} \in [C(\overline{\Omega})]^d, \nabla \cdot \boldsymbol{\beta} \in L^{\infty}(\Omega), \sigma + \frac{1}{2} \nabla \cdot \boldsymbol{\beta} \ge 0, \nabla \nabla \cdot \boldsymbol{\beta} \in [L^2(\Omega)]^d, \sigma \in L^{\infty}(\Omega) \cap H^1(\Omega).$$
(4.1)

We assume the solution has the following regularity properties:

$$y \in H^{r_y}(\Omega), \quad z \in H^{r_z}(\Omega), \quad \boldsymbol{q} \in [H^{r_q}(\Omega)]^d, \quad \boldsymbol{p} \in [H^{r_p}(\Omega)]^d,$$
(4.2a)

$$r_y \ge 1, \quad r_z \ge 2, \quad r_q \ge 0, \quad r_p \ge 1.$$
 (4.2b)

In the 2D case, Theorem 1 guarantees this condition is satisfied.

It is worthwhile to mention that if \boldsymbol{q} has a well-defined boundary trace in $L^2(\Gamma)$, i.e., $r_{\boldsymbol{q}} > 1/2$, then we refer to this as the high regularity case for the boundary control problem; otherwise, if $r_{\boldsymbol{q}} \in [0, 1/2]$, then we say this is the low regularity case. In 2D, by Theorem 1, if $y_d \in H^{t^*}(\Omega)$ for some $t^* \in (1/2, 1)$, and $\pi/3 \leq \omega < 2\pi/3$, then we are guaranteed to be in the high regularity case. However, if one of the above assumptions concerning y_d or ω is not satisfied, then \boldsymbol{q} is no longer guaranteed to have a well-defined boundary trace.

For the *diffusion dominated* boundary control problem, we gave a rigorous error analysis of a different HDG method for the high regularity case in [29,30] and for the low regularity case in [23]. In this work, we are interested in the *convection dominated* case. However, existing numerical analysis works for convection dominated diffusion PDEs only consider the high regularity case; see, e.g. [2,20]. To the best of our knowledge, there is no existing error analysis work on convection dominated PDEs with low regularity solutions.

We now state our main convergence result.

Theorem 3. Let $s_y = \min\{r_y, k+1\}$, $s_z = \min\{r_z, k+1\}$, (u, y, z) and (u_h, y_h, z_h) be the solutions of (1.3) and (2.3), respectively. If assumptions **(A1)-(A3)** hold, then there exists h_0 , independent of ε , such that for all $h \leq h_0$ we have

$$\begin{aligned} \|u - u_h\|_{\mathcal{E}_h^{\partial}} &\leq C\left(h^{s_y - 1/2} \|y\|_{s_y} + h^{s_z - 1/2} \|z\|_{s_z} + \delta(s_y)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}\right),\\ \|y - y_h\|_{\mathcal{T}_h} &\leq C\left(h^{s_y - 1/2} \|y\|_{s_y} + h^{s_z - 1/2} \|z\|_{s_z} + \delta(s_y)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}\right),\\ \|z - z_h\|_{\mathcal{T}_h} &\leq C\left(h^{s_y - 1/2} \|y\|_{s_y} + h^{s_z - 1/2} \|z\|_{s_z} + \delta(s_y)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}\right),\end{aligned}$$

where $\delta(t) = 1$ if $t \leq 3/2$, otherwise $\delta(t) = 0$.

Remark 3. If $s_y \leq 3/2$, then we have $\|\Delta y\|_{\mathcal{T}_h}$ in the error estimates. This term is finite by Theorem 1.

Specializing to the 2D case gives the following result:

Corollary 1. Suppose d = 2, f = 0, $y_d \in H^{t^*}(\Omega)$ for some $t^* \in [0, 1)$ and assumptions (A1)-(A3) hold. Let $\pi/3 \leq \omega < \pi$ be the largest interior angle of Γ , and let r > 0 satisfy

$$r \leq r_d := \frac{1}{2} + t^* \in [1/2, 3/2), \text{ and } r < r_{\Omega} := \min\left\{\frac{3}{2}, \frac{\pi}{\omega} - \frac{1}{2}\right\} \in (1/2, 3/2]$$

If k = 1, then there exists h_0 , independent of ε , such that for all $h \leq h_0$ we have

$$\begin{aligned} \|u - u_h\|_{\varepsilon_h^{\partial}} &\leq Ch^r(\|y\|_{H^{r+1/2}(\Omega)} + \|z\|_{H^{r+3/2}(\Omega)} + \delta(r)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}), \\ \|y - y_h\|_{\mathcal{T}_h} &\leq Ch^r(\|y\|_{H^{r+1/2}(\Omega)} + \|z\|_{H^{r+3/2}(\Omega)} + \delta(r)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}), \\ \|z - z_h\|_{\mathcal{T}_h} &\leq Ch^r(\|y\|_{H^{r+1/2}(\Omega)} + \|z\|_{H^{r+3/2}(\Omega)} + \delta(r)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}). \end{aligned}$$

Furthermore, if k = 0, then there exists h_1 , independent of ε , such that for all $h \leq h_1$ we have

$$\begin{aligned} \|u - u_h\|_{\varepsilon_h^{\partial}} &\leq Ch^{1/2}(\|y\|_{H^1(\Omega)} + \|z\|_{H^1(\Omega)} + \delta(r)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}), \\ \|y - y_h\|_{\mathcal{T}_h} &\leq Ch^{1/2}(\|y\|_{H^1(\Omega)} + \|z\|_{H^1(\Omega)} + \delta(r)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}), \\ \|z - z_h\|_{\mathcal{T}_h} &\leq Ch^{1/2}(\|y\|_{H^1(\Omega)} + \|z\|_{H^1(\Omega)} + \delta(r)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}). \end{aligned}$$

Similar to [29,30], the convergence rates are optimal for the control when k = 1 and suboptimal when k = 0. However, if $y_d \in L^2(\Omega)$, then $u \in H^{1/2}(\Gamma)$ only and the convergence rate for the control is optimal when k = 0.

4.2 Proof of Theorem 3

We introduce an auxiliary problem with the approximate control u_h in the HDG discretized optimality system (2.6a) replaced by a projection of the exact optimal control, and split the proof into seven steps.

We first bound the error between the solution of the optimality system (2.2a)-(2.2d) and $(\boldsymbol{q}_h(u), y_h(u), \hat{y}_h^o(u), \boldsymbol{p}_h(u), z_h(u), \hat{z}_h^o(u)) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2$ satisfying the auxiliary problem

$$\mathcal{B}_1(\boldsymbol{q}_h(\boldsymbol{u}), y_h(\boldsymbol{u}); \boldsymbol{\hat{y}}_h(\boldsymbol{u}); \boldsymbol{r}_1, w_1, \widehat{w}_1^o) = -(f, w_1)_{\mathcal{T}_h} - \langle \Pi_k^\partial \boldsymbol{u}, \tau_2 w_1 + \boldsymbol{r}_1 \cdot \boldsymbol{n} \rangle_{\mathcal{E}_k^\partial}, \tag{4.3a}$$

$$\mathcal{B}_{2}(\boldsymbol{p}_{h}(u), z_{h}(u), \widehat{z}_{h}(u); \boldsymbol{r}_{2}, w_{2}, \widehat{w}_{2}^{o}) = -(y_{h}(u) - y_{d}, w_{2})_{\mathcal{T}_{h}},$$
(4.3b)

for all $(\boldsymbol{r}_1, w_1, \widehat{w}_1^o, \boldsymbol{r}_2, w_2, \widehat{w}_2^o) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2$.

4.2.1 Step 1: errors between the auxiliary problem (4.3) and the continuous problem (2.2)

Lemma 9. Let $(\boldsymbol{q}, y, \boldsymbol{p}, z, u)$ be the solution of (2.2). Then for all $(\boldsymbol{r}_1, w_1, \widehat{w}_1^o, \boldsymbol{r}_2, w_2, \widehat{w}_2^o) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2$, we have

$$\mathcal{B}_{1}(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{q},\boldsymbol{\Pi}_{k}^{o}\boldsymbol{y},\boldsymbol{\Pi}_{k}^{\partial}\boldsymbol{y};\boldsymbol{r}_{1},\boldsymbol{w}_{1},\boldsymbol{\widehat{w}}_{1}^{o}) = -(f,\boldsymbol{w}_{1})_{\mathcal{T}_{h}} - \langle \boldsymbol{\Pi}_{k}^{\partial}\boldsymbol{u},\boldsymbol{\tau}_{2}\boldsymbol{w}_{1} + \boldsymbol{r}_{1}\cdot\boldsymbol{n}\rangle_{\mathcal{E}_{h}^{\partial}} + E_{1}(\boldsymbol{q},\boldsymbol{y};\boldsymbol{w}_{1},\boldsymbol{\widehat{w}}_{1}^{o}),$$

$$(4.4a)$$

$$\mathcal{B}_{2}(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{p},\boldsymbol{\Pi}_{k}^{o}z,\widetilde{\boldsymbol{\Pi}}_{k}^{\partial}z;\boldsymbol{r}_{2},w_{2},\widehat{w}_{2}^{o}) = -(y-y_{d},w_{2})_{\mathcal{T}_{h}} + E_{2}(\boldsymbol{p},z;w_{2},\widehat{w}_{2}^{o}),$$
(4.4b)

where

$$E_{1}(\boldsymbol{q}, \boldsymbol{y}; \boldsymbol{w}_{1}, \widehat{\boldsymbol{w}}_{1}^{o}) = (\Pi_{k}^{o}\boldsymbol{y} - \boldsymbol{y}, \boldsymbol{\beta} \cdot \nabla \boldsymbol{w}_{1})_{\mathcal{T}_{h}} - (\sigma(\Pi_{k}^{o} - \mathbb{I})\boldsymbol{y}, \boldsymbol{w}_{1})_{\mathcal{T}_{h}} + \langle \widehat{\boldsymbol{w}}_{1}^{o} - \boldsymbol{w}_{1}, (\boldsymbol{\Pi}_{k}^{o}\boldsymbol{q} - \boldsymbol{q}) \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_{h}} + \langle \boldsymbol{y} - \boldsymbol{\Pi}_{k}^{\partial}\boldsymbol{y}, \boldsymbol{\beta} \cdot \boldsymbol{n}(\boldsymbol{w}_{1} - \widehat{\boldsymbol{w}}_{1}^{o}) \rangle_{\partial \mathcal{T}_{h}} - \langle \tau_{1}(\Pi_{k}^{o}\boldsymbol{y} - \Pi_{k}^{\partial}\boldsymbol{y}), \boldsymbol{w}_{1} - \widehat{\boldsymbol{w}}_{1}^{o} \rangle_{\partial \mathcal{T}_{h}}, \qquad (4.5)$$
$$E_{2}(\boldsymbol{p}, \boldsymbol{z}; \boldsymbol{w}_{2}, \widehat{\boldsymbol{w}}_{2}^{o}) = -(\Pi_{k}^{o}\boldsymbol{z} - \boldsymbol{z}, \boldsymbol{\beta} \cdot \nabla \boldsymbol{w}_{2})_{\mathcal{T}_{h}} - (\sigma(\boldsymbol{\Pi}_{k}^{o} - \mathbb{I})(\boldsymbol{z} + \nabla \cdot \boldsymbol{\beta}), \boldsymbol{w}_{2})_{\mathcal{T}_{h}} + \langle \widehat{\boldsymbol{w}}_{2}^{o} - \boldsymbol{w}_{2}, (\boldsymbol{\Pi}_{k}^{o}\boldsymbol{p} - \boldsymbol{p}) \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_{h}} - \langle \boldsymbol{z} - \boldsymbol{\Pi}_{k}^{\partial}\boldsymbol{z}, \boldsymbol{\beta} \cdot \boldsymbol{n}(\boldsymbol{w}_{2} - \widehat{\boldsymbol{w}}_{2}^{o}) \rangle_{\partial \mathcal{T}_{h}} - \langle \tau_{2}(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{z} - \boldsymbol{\Pi}_{k}^{\partial}\boldsymbol{z}), \boldsymbol{w}_{2} - \widehat{\boldsymbol{w}}_{2}^{o} \rangle_{\partial \mathcal{T}_{h}}. \qquad (4.6)$$

Proof. We only give a proof of (4.4a). By the definition of \mathcal{B}_1 in (2.4), one gets

$$\begin{split} \mathcal{B}_{1}(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{q},\boldsymbol{\Pi}_{k}^{o}y,\widetilde{\boldsymbol{\Pi}}_{k}^{\partial}y;\boldsymbol{r}_{1},w_{1},\widehat{w}_{1}^{o}) \\ &=\varepsilon^{-1}(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{q},\boldsymbol{r}_{1})_{\mathcal{T}_{h}}-(\boldsymbol{\Pi}_{k}^{o}y,\nabla\cdot\boldsymbol{r}_{1})_{\mathcal{T}_{h}}+\langle\widetilde{\boldsymbol{\Pi}}_{k}^{\partial}y,\boldsymbol{r}_{1}\cdot\boldsymbol{n}\rangle_{\partial\mathcal{T}_{h}} \\ &-(w_{1},\nabla\cdot\boldsymbol{\Pi}_{k}^{o}\boldsymbol{q})_{\mathcal{T}_{h}}+\langle\widehat{w}_{1}^{o},\boldsymbol{\Pi}_{k}^{o}\boldsymbol{q}\cdot\boldsymbol{n}\rangle_{\partial\mathcal{T}_{h}}-\langle\tau_{1}(\boldsymbol{\Pi}_{k}^{o}y-\widetilde{\boldsymbol{\Pi}}_{k}^{\partial}y),w_{1}-\widehat{w}_{1}^{o}\rangle_{\partial\mathcal{T}_{h}} \\ &+(\boldsymbol{\Pi}_{k}^{o}y,\boldsymbol{\beta}\cdot\nabla w_{1})_{\mathcal{T}_{h}}-\langle\widetilde{\boldsymbol{\Pi}}_{k}^{\partial}y,\boldsymbol{\beta}\cdot\boldsymbol{n}w_{1}\rangle_{\partial\mathcal{T}_{h}}-(\sigma\boldsymbol{\Pi}_{k}^{o}y,w_{1})_{\mathcal{T}_{h}}. \end{split}$$

By the orthogonality properties of Π_k^o , Π_k^o , $\widetilde{\Pi}_k^\partial$, and the fact y = u on \mathcal{E}_h^∂ , we have

$$\begin{split} \mathcal{B}_{1}(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{q},\boldsymbol{\Pi}_{k}^{o}\boldsymbol{y},\boldsymbol{\Pi}_{k}^{\partial}\boldsymbol{y};\boldsymbol{r}_{1},w_{1},\widehat{w}_{1}^{o}) \\ &=\varepsilon^{-1}(\boldsymbol{q},\boldsymbol{r}_{1})_{\mathcal{T}_{h}}-(\boldsymbol{y},\nabla\cdot\boldsymbol{r}_{1})_{\mathcal{T}_{h}}+\langle\boldsymbol{y},\boldsymbol{r}_{1}\cdot\boldsymbol{n}\rangle_{\partial\mathcal{T}_{h}}-\langle\boldsymbol{\Pi}_{k}^{\partial}\boldsymbol{u},\boldsymbol{r}_{1}\cdot\boldsymbol{n}\rangle_{\mathcal{E}_{h}^{\partial}} \\ &+(\nabla w_{1},\boldsymbol{q})_{\mathcal{T}_{h}}+\langle\widehat{w}_{1}^{o}-w_{1},\boldsymbol{\Pi}_{k}^{o}\boldsymbol{q}\cdot\boldsymbol{n}\rangle_{\partial\mathcal{T}_{h}}-\langle\boldsymbol{\tau}_{1}(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{y}-\boldsymbol{\Pi}_{k}^{\partial}\boldsymbol{y}),w_{1}-\widehat{w}_{1}^{o}\rangle_{\partial\mathcal{T}_{h}} \\ &+(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{y}-\boldsymbol{y},\boldsymbol{\beta}\cdot\nabla w_{1})_{\mathcal{T}_{h}}-\langle\boldsymbol{\Pi}_{k}^{\partial}\boldsymbol{y}-\boldsymbol{y},\boldsymbol{\beta}\cdot\boldsymbol{n}w_{1}\rangle_{\partial\mathcal{T}_{h}}-(\sigma(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{y}-\boldsymbol{y}),w_{1})_{\mathcal{T}_{h}} \\ &+(\boldsymbol{y},\boldsymbol{\beta}\cdot\nabla w_{1})_{\mathcal{T}_{h}}-\langle\boldsymbol{y},\boldsymbol{\beta}\cdot\boldsymbol{n}w_{1}\rangle_{\partial\mathcal{T}_{h}}-(\sigma\boldsymbol{y},w_{1})_{\mathcal{T}_{h}}+\langle\boldsymbol{\Pi}_{k}^{\partial}\boldsymbol{u},\boldsymbol{\beta}\cdot\boldsymbol{n}w_{1}\rangle_{\mathcal{E}_{h}^{\partial}}. \end{split}$$

By integration by parts, and the fact $\langle \hat{w}_1^o, \boldsymbol{q} \cdot \boldsymbol{n} \rangle_{\partial \mathcal{T}_h} = 0$, we arrive at

$$\begin{split} \mathcal{B}_{1}(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{q},\boldsymbol{\Pi}_{k}^{o}y,\boldsymbol{\Pi}_{k}^{\partial}y;\boldsymbol{r}_{1},w_{1},\boldsymbol{\widehat{w}}_{1}^{o}) \\ &=\varepsilon^{-1}(\boldsymbol{q},\boldsymbol{r}_{1})_{\mathcal{T}_{h}}+(\nabla y,\boldsymbol{r}_{1})_{\mathcal{T}_{h}}-\langle\boldsymbol{\Pi}_{k}^{\partial}u,\boldsymbol{r}_{1}\cdot\boldsymbol{n}\rangle_{\mathcal{E}_{h}^{\partial}}-(w_{1},\nabla\cdot\boldsymbol{q})_{\mathcal{T}_{h}} \\ &+\langle\boldsymbol{\widehat{w}}_{1}^{o}-w_{1},(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{q}-\boldsymbol{q})\cdot\boldsymbol{n}\rangle_{\partial\mathcal{T}_{h}}-\langle\boldsymbol{\tau}_{1}(\boldsymbol{\Pi}_{k}^{o}y-\boldsymbol{\Pi}_{k}^{\partial}y),w_{1}-\boldsymbol{\widehat{w}}_{1}^{o}\rangle_{\partial\mathcal{T}_{h}} \\ &-\langle\boldsymbol{\tau}_{1}\boldsymbol{\Pi}_{k}^{\partial}u,w_{1}\rangle_{\mathcal{E}_{h}^{\partial}}+(\boldsymbol{\Pi}_{k}^{o}y-y,\boldsymbol{\beta}\cdot\nabla w_{1})_{\mathcal{T}_{h}}+\langle y-\boldsymbol{\Pi}_{k}^{\partial}y,\boldsymbol{\beta}\cdot\boldsymbol{n}w_{1}\rangle_{\partial\mathcal{T}_{h}} \\ &-(\nabla\cdot(\boldsymbol{\beta}y),w_{1})_{\mathcal{T}_{h}}+\langle\boldsymbol{\beta}\cdot\boldsymbol{n}\boldsymbol{\Pi}_{k}^{\partial}u,w_{1}\rangle_{\mathcal{E}_{h}^{\partial}}-(\sigma y,w_{1})_{\mathcal{T}_{h}}-(\sigma(\boldsymbol{\Pi}_{k}^{o}-\boldsymbol{\mathbb{I}})y,w_{1})_{\mathcal{T}_{h}}. \end{split}$$

Then by the facts $\varepsilon^{-1} q = -\nabla y$ and $\nabla \cdot q + \nabla \cdot (\beta y) + \sigma y = f$, we have

$$\mathcal{B}_{1}(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{q},\boldsymbol{\Pi}_{k}^{o}\boldsymbol{y},\boldsymbol{\Pi}_{k}^{o}\boldsymbol{y};\boldsymbol{r}_{1},w_{1},\hat{w}_{1}^{o}) \\
= -(f,w_{1})_{\mathcal{T}_{h}} - \langle \boldsymbol{\Pi}_{k}^{\partial}\boldsymbol{u},\tau_{2}w_{1} + \boldsymbol{r}_{1} \cdot \boldsymbol{n} \rangle_{\mathcal{E}_{h}^{\partial}} \\
+ (\boldsymbol{\Pi}_{k}^{o}\boldsymbol{y} - \boldsymbol{y},\boldsymbol{\beta} \cdot \nabla w_{1})_{\mathcal{T}_{h}} - (\sigma(\boldsymbol{\Pi}_{k}^{o} - \mathbb{I})\boldsymbol{y},w_{1})_{\mathcal{T}_{h}} \\
+ \langle \hat{w}_{1}^{o} - w_{1}, (\boldsymbol{\Pi}_{k}^{o}\boldsymbol{q} - \boldsymbol{q}) \cdot \boldsymbol{n} \rangle_{\partial\mathcal{T}_{h}} - \langle \tau_{1}(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{y} - \boldsymbol{\Pi}_{k}^{\partial}\boldsymbol{y}), w_{1} - \hat{w}_{1}^{o} \rangle_{\partial\mathcal{T}_{h}} \\
+ \langle \boldsymbol{y} - \boldsymbol{\Pi}_{k}^{\partial}\boldsymbol{y}, \boldsymbol{\beta} \cdot \boldsymbol{n}(w_{1} - \hat{w}_{1}^{o}) \rangle_{\partial\mathcal{T}_{h}},$$
(4.7)

where we used $\langle y - \Pi_k^{\partial} y, \boldsymbol{\beta} \cdot \boldsymbol{n} \widehat{w}_1^{o} \rangle_{\partial \mathcal{T}_h} = 0$ in (4.7).

By (4.4) and (4.3) we have the following error equations.

Lemma 10. Let $(\boldsymbol{q}, y, \boldsymbol{p}, z, u)$ and $(\boldsymbol{q}_h(u), y_h(u), \hat{y}_h^o(u), \boldsymbol{p}_h(u), z_h(u), \hat{z}_h^o(u)) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2$ be the solutions of (2.2) and (4.3), respectively. Then for all $(\boldsymbol{r}_1, w_1, \hat{w}_1^o, \boldsymbol{r}_2, w_2, \hat{w}_2^o) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2$, we have

$$\mathcal{B}_1(\boldsymbol{\Pi}_k^o \boldsymbol{q} - \boldsymbol{q}_h(u), \boldsymbol{\Pi}_k^o y - y_h(u), \widetilde{\boldsymbol{\Pi}}_k^\partial y - \widehat{y}_h^o(u); \boldsymbol{r}_1, w_1, \widehat{w}_1^o) = E_1(\boldsymbol{q}, y; w_1, \widehat{w}_1^o),$$
(4.8a)

$$\mathcal{B}_{2}(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{p} - \boldsymbol{p}_{h}(u), \boldsymbol{\Pi}_{k}^{o}z - z_{h}(u), \widetilde{\boldsymbol{\Pi}}_{k}^{\partial}z - \widehat{z}_{h}^{o}(u); \boldsymbol{r}_{2}, w_{2}, \widehat{w}_{2}^{o}) = -(y - y_{h}(u), w_{2})_{\mathcal{T}_{h}} + E_{2}(\boldsymbol{p}, z; w_{2}, \widehat{w}_{2}^{o}),$$
(4.8b)

where E_1 and E_2 are defined in Lemma 9.

Lemma 11. Let (q, y, p, z) be the solution of (2.2). Then for all $(w_1, w_2, \widehat{w}_1^o, \widehat{w}_2^o) \in [W_h \times M_h^o]^2$, we have

$$|E_1(\boldsymbol{q}, y; w_1, \widehat{w}_1^o)| \leq C \left(h^{s_y - 1/2} \|y\|_{s_y} + \delta(s_y) \varepsilon^{1/2} h \|\Delta y\|_{\mathcal{T}_h} \right) \|(w_1, \widehat{w}_1^o)\|_W,$$
(4.9)

$$|E_2(\boldsymbol{p}, z; w_2, \widehat{w}_2^o)| \leq C \quad h^{s_z - 1/2} ||z||_{s_z} ||(w_2, \widehat{w}_2^o)||_W.$$
(4.10)

Proof. Since the proof for (4.10) is similar to the proof of (4.9), we only prove (4.9). To simplify notation, we write E_1 from Lemma 9 as $E_1(\boldsymbol{q}, y; v_h, \hat{v}_h^o) = \sum_{i=1}^5 R_i$. For the term R_1 , since $((\mathbb{I} - \Pi_k^o)y, \mathbf{\Pi}_0^o \boldsymbol{\beta} \cdot \nabla w_1)_{\mathcal{T}_h} = 0$, we get

For the term R_2 , by a direct estimate, we get

$$\begin{aligned} |R_{2}| &\leq C \|\sigma\|_{0,\infty}^{1/2} \|(\mathbb{I} - \Pi_{k}^{o})y\|_{\mathcal{T}_{h}} \|\sigma^{1/2}w_{1}\|_{\mathcal{T}_{h}} \\ &\leq Ch^{s_{y}} \|\sigma\|_{0,\infty}^{1/2} \|y\|_{s_{y}} \|(w_{1},\widehat{w}_{1}^{o})\|_{W} \\ &\leq Ch^{s_{y}} (\beta_{0}^{1/2} + \sigma_{0}^{1/2}) \|y\|_{s_{y}} \|(w_{1},\widehat{w}_{1}^{o})\|_{W} \\ &\leq Ch^{s_{y}} \|y\|_{s_{y}} \|(w_{1},\widehat{w}_{1}^{o})\|_{W}. \end{aligned}$$
 by (3.10b),

For the term R_3 , we need a refined analysis since this term relates to the boundary trace of the gradient of y. Below, we use $\mathbf{q} = -\varepsilon \nabla y$ and $\varepsilon < \min_{T \in \mathcal{T}_h} \{h_T\}$.

If $s_y > 3/2$, we have

$$\begin{aligned} |R_3| &= \varepsilon |\langle \boldsymbol{n} \cdot (\nabla y - \boldsymbol{\Pi}_k^o \nabla y), w_1 - \widehat{w}_1^o \rangle_{\partial \mathcal{T}_h}| \\ &\leq C h^{s_y - 1} \varepsilon^{1/2} \|y\|_{s_y} \left(\sum_{T \in \mathcal{T}_h} \varepsilon h_T^{-1} \|w_1 - \widehat{w}_1^o\|_{\partial T}^2 \right)^{1/2} \\ &\leq C h^{s_y - 1/2} \|y\|_{s_y} \|(w_1, \widehat{w}_1^o)\|_W. \end{aligned}$$

If $s_y \leq 3/2$, use $\langle \boldsymbol{n} \cdot \nabla y, \hat{w}_1^o \rangle_{\partial \mathcal{T}_h} = 0$ and integration by parts to obtain

$$\begin{aligned} |R_3| &= \varepsilon |\langle \boldsymbol{n} \cdot \nabla \boldsymbol{y}, w_1 \rangle_{\partial \mathcal{T}_h} - \langle \boldsymbol{n} \cdot \boldsymbol{\Pi}_k^o \nabla \boldsymbol{y}, w_1 - \widehat{w}_1^o \rangle_{\partial \mathcal{T}_h}| \\ &= \varepsilon |(\Delta \boldsymbol{y}, w_1)_{\mathcal{T}_h} + (\boldsymbol{\Pi}_k^o \nabla \boldsymbol{y}, \nabla w_1)_{\mathcal{T}_h} - \langle \boldsymbol{n} \cdot \boldsymbol{\Pi}_k^o \nabla \boldsymbol{y}, w_1 - \widehat{w}_1^o \rangle_{\partial \mathcal{T}_h}|. \end{aligned}$$

We use integration by parts again, and also (3.8b) and (3.8a) to get

$$\begin{aligned} |R_3| &= \varepsilon |(\Delta y, w_1)_{\mathcal{T}_h} - (\nabla \cdot \mathbf{\Pi}_k^o \nabla y, w_1)_{\mathcal{T}_h} + \langle \boldsymbol{n} \cdot \mathbf{\Pi}_k^o \nabla y, \widehat{w}_1^o \rangle_{\partial \mathcal{T}_h} | \\ &= \varepsilon |(\Delta y, w_1)_{\mathcal{T}_h} - (\nabla \cdot \mathbf{\Pi}_k^o \nabla y, \mathcal{I}_h(w_1, \widehat{w}_1^o))_{\mathcal{T}_h} + \langle \boldsymbol{n} \cdot \mathbf{\Pi}_k^o \nabla y, \mathcal{I}_h(w_1, \widehat{w}_1^o) \rangle_{\partial \mathcal{T}_h} | \\ &= \varepsilon |(\Delta y, w_1)_{\mathcal{T}_h} + (\mathbf{\Pi}_k^o \nabla y, \nabla \mathcal{I}_h(w_1, \widehat{w}_1^o))_{\mathcal{T}_h} |. \end{aligned}$$

Therefore, by the triangle inequality, integration by parts, (3.8c) and (3.8d) we have

$$\begin{aligned} |R_3| &\leq \varepsilon |(\Delta y, \mathcal{I}_h(w_1, \widehat{w}_1^o))_{\mathcal{T}_h} + (\mathbf{\Pi}_k^o \nabla y, \mathcal{I}_h(w_1, \widehat{w}_1^o))_{\mathcal{T}_h}| + \varepsilon |(\Delta y, w_1 - \mathcal{I}_h(w_1, \widehat{w}_1^o))_{\mathcal{T}_h}| \\ &= \varepsilon |(\nabla y - \mathbf{\Pi}_k^o \nabla y, \nabla \mathcal{I}_h(w_1, \widehat{w}_1^o))_{\mathcal{T}_h}| + \varepsilon |(\Delta y, w_1 - \mathcal{I}_h(w_1, \widehat{w}_1^o))_{\mathcal{T}_h}| \\ &\leq C \varepsilon^{1/2} (h^{s_y - 1} ||y||_{s_y} + h ||\Delta y||_{\mathcal{T}_h}) ||(w_1, \widehat{w}_1^o)||_W \\ &\leq C (h^{s_y - 1/2} ||y||_{s_y} + \varepsilon^{1/2} h ||\Delta y||_{\mathcal{T}_h}) ||(w_1, \widehat{w}_1^o)||_W. \end{aligned}$$

For the terms R_4 and R_5 , use the Cauchy-Schwarz inequality and (3.2) to get

$$\begin{aligned} |R_4| &= |\langle \boldsymbol{\beta} \cdot \boldsymbol{n}(y - \Pi_k^{\partial} y), w_1 - \widehat{w}_1^{o} \rangle_{\partial \mathcal{T}_h}| \\ &\leq C h^{s_y - 1/2} \|\boldsymbol{\beta}\|_{0,\infty}^{1/2} \|y\|_{s_y} \left(\sum_{T \in \mathcal{T}_h} |\boldsymbol{\beta} \cdot \boldsymbol{n}|^{1/2} (w_1 - \widehat{w}_1^{o})\|_{\partial T}^2 \right)^{1/2} \\ &\leq C h^{s_y - 1/2} \|y\|_{s_y} \|(w_1, \widehat{w}_1^{o})\|_W, \\ |R_5| &= |\langle \tau_1(\Pi_k^o - \Pi_k^{\partial}) y, w_1 - \widehat{w}_1^o \rangle_{\partial \mathcal{T}_h}| \\ &\leq C h^{s_y - 1} (\beta_0^{1/2} h^{1/2} + \varepsilon^{1/2}) \|y\|_{s_y} \|(w_1, \widehat{w}_1^o)\|_W \\ &\leq C h^{s_y - 1/2} \|y\|_{s_y} \|(w_1, \widehat{w}_1^o)\|_W. \end{aligned}$$

From all the estimates above we get our final result.

Lemma 12. Let (q, y, p, z) and $(q_h(u), y_h(u), \hat{y}_h^o(u), p_h(u), z_h(u), \hat{z}_h^o(u)) \in [V_h \times W_h \times M_h^o]^2$ be the solutions of (2.2) and (4.3), respectively. If assumptions (A1) and (A2) hold, then there exists $h_0 > 0$, independent of ε , such that for all $h \leq h_0$ we have the error estimates

$$\begin{split} \|y - y_h(u)\|_{\mathcal{T}_h} &\leq C \left(h^{s_y - 1/2} \|y\|_{s_y} + \delta(s_y) \varepsilon^{1/2} h \|\Delta y\|_{\mathcal{T}_h} \right), \\ \|z - z_h(u)\|_{\mathcal{T}_h} &\leq C \left(h^{s_y - 1/2} \|y\|_{s_y} + h^{s_z - 1/2} \|z\|_{s_z} + \delta(s_y) \varepsilon^{1/2} h \|\Delta y\|_{\mathcal{T}_h} \right), \\ \|\boldsymbol{p} - \boldsymbol{p}_h(u)\|_{\mathcal{T}_h} &\leq C \varepsilon^{1/2} \left(h^{s_y - 1/2} \|y\|_{s_y} + h^{s_z - 1/2} \|z\|_{s_z} + \delta(s_y) \varepsilon^{1/2} h \|\Delta y\|_{\mathcal{T}_h} \right). \end{split}$$

Proof. By Theorem 2, (4.8a) and (4.9) we get

$$\begin{split} \| (\mathbf{\Pi}_{k}^{o} \boldsymbol{q} - \boldsymbol{q}_{h}(u), \Pi_{k}^{o} y - y_{h}(u), \Pi_{k}^{o} y - \widehat{y}_{h}^{o}(u)) \| \\ & \leq C \sup_{(\boldsymbol{r}_{1}, w_{1}, \widehat{w}_{1}^{o}) \in \boldsymbol{V}_{h} \times W_{h} \times M_{h}^{o}} \frac{\mathcal{B}_{1}(\mathbf{\Pi}_{k}^{o} \boldsymbol{q} - \boldsymbol{q}_{h}(u), \Pi_{k}^{o} y - y_{h}(u), \widetilde{\Pi}_{k}^{\partial} y - \widehat{y}_{h}^{o}(u); \boldsymbol{r}_{1}, w_{1}, \widehat{w}_{1}^{o}) \| \\ & \leq C \sup_{(\boldsymbol{r}_{1}, w_{1}, \widehat{w}_{1}^{o}) \in \boldsymbol{V}_{h} \times W_{h} \times M_{h}^{o}} \frac{E_{1}(\boldsymbol{q}, y; v_{h}, \widehat{v}_{h}^{o})}{\|(\boldsymbol{r}_{1}, w_{1}, \widehat{w}_{1}^{o})\|} \\ & \leq C \left(h^{s_{y} - 1/2} \|y\|_{s_{y}} + \delta(s_{y}) \varepsilon^{1/2} h \|\Delta y\|_{\mathcal{T}_{h}}\right). \end{split}$$

 ~ 0

Therefore,

$$\begin{aligned} \|y - y_{h}(u)\|_{\mathcal{T}_{h}} \\ &\leq \|y - \Pi_{k}^{o}y\|_{\mathcal{T}_{h}} + \|\Pi_{k}^{o}y - y_{h}(u)\|_{\mathcal{T}_{h}} \\ &\leq \|y - \Pi_{k}^{o}y\|_{\mathcal{T}_{h}} + C\|(\mathbf{\Pi}_{k}^{o}\boldsymbol{q} - \boldsymbol{q}_{h}(u), \Pi_{k}^{o}y - y_{h}(u), \widetilde{\Pi}_{k}^{\partial}y - \widehat{y}_{h}^{o}(u))\| \\ &\leq C\left(h^{s_{y}-1/2}\|y\|_{s_{y}} + \delta(s_{y})\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_{h}}\right). \end{aligned}$$
(4.11)

By Theorem 2, (4.8b), (4.10) and estimate (4.11) we get

$$\begin{aligned} \| (\mathbf{\Pi}_{k}^{o} \mathbf{p} - \mathbf{p}_{h}(u), \mathbf{\Pi}_{k}^{o} z - z_{h}(u), \widetilde{\mathbf{\Pi}}_{k}^{\partial} z - \widehat{z}_{h}^{o}(u)) \| \\ &\leq C \sup_{(\mathbf{r}_{2}, w_{2}, \widehat{w}_{2}^{o}) \in \mathbf{V}_{h} \times W_{h} \times M_{h}^{o}} \frac{\mathcal{B}_{2}(\mathbf{\Pi}_{k}^{o} \mathbf{p} - \mathbf{p}_{h}(u), \mathbf{\Pi}_{k}^{o} z - z_{h}(u), \widetilde{\mathbf{\Pi}}_{k}^{\partial} z - \widehat{z}_{h}^{o}(u); \mathbf{r}_{2}, w_{2}, \widehat{w}_{2}^{o})}{\| (\mathbf{r}_{2}, w_{2}, \widehat{w}_{2}^{o}) \|} \\ &\leq C \sup_{(\mathbf{r}_{2}, w_{2}, \widehat{w}_{2}^{o}) \in \mathbf{V}_{h} \times W_{h} \times M_{h}^{o}} \frac{E_{2}(\mathbf{p}, z; w_{2}, \widehat{w}_{2}^{o}) - (y - y_{h}(u), w_{2})}{\| (\mathbf{r}_{2}, w_{2}, \widehat{w}_{2}^{o}) \|} \\ &\leq C \left(h^{s_{z} - 1/2} \| z \|_{s_{z}} + h^{s_{y} - 1/2} \| y \|_{s_{y}} + \delta(s_{y}) \varepsilon^{1/2} h \| \Delta y \|_{\mathcal{T}_{h}} \right). \end{aligned}$$

$$(4.12)$$

Therefore, the triangle inequality gives

$$\begin{aligned} \|z - z_h(u)\|_{\mathcal{T}_h} \\ &\leq \|z - \Pi_k^o z\|_{\mathcal{T}_h} + \|\Pi_k^o z - z_h(u)\|_{\mathcal{T}_h} \\ &\leq \|z - \Pi_k^o z\|_{\mathcal{T}_h} + (\beta_0 + \sigma_0)^{-1} \|(\mathbf{\Pi}_k^o \mathbf{p} - \mathbf{p}_h(u), \Pi_k^o z - z_h(u), \widetilde{\Pi}_k^\partial z - \widehat{z}_h^o(u))\| \\ &\leq C \left(h^{s_z - 1/2} \|z\|_{s_z} + h^{s_y - 1/2} \|y\|_{s_y} + \delta(s_y) \varepsilon^{1/2} h \|\Delta y\|_{\mathcal{T}_h} \right). \end{aligned}$$

Next, we use the triangle inequality, $\boldsymbol{p} = -\varepsilon \nabla z$, and $\varepsilon < \min_{T \in \mathcal{T}_h} \{h_T\}$ to get

$$\begin{aligned} \|\boldsymbol{p} - \boldsymbol{p}_{h}(u)\|_{\mathcal{T}_{h}} &\leq \|\boldsymbol{p} - \boldsymbol{\Pi}_{k}^{o}\boldsymbol{p}\|_{\mathcal{T}_{h}} + \|\boldsymbol{\Pi}_{k}^{o}\boldsymbol{p} - \boldsymbol{p}_{h}(u)\|_{\mathcal{T}_{h}} \\ &\leq \|\boldsymbol{p} - \boldsymbol{\Pi}_{k}^{o}\boldsymbol{p}\|_{\mathcal{T}_{h}} + \varepsilon^{1/2} \|(\boldsymbol{\Pi}_{k}^{o}\boldsymbol{p} - \boldsymbol{p}_{h}(u), \boldsymbol{\Pi}_{k}^{o}\boldsymbol{z} - \boldsymbol{z}_{h}(u), \widetilde{\boldsymbol{\Pi}}_{k}^{\partial}\boldsymbol{z} - \widehat{\boldsymbol{z}}_{h}^{o}(u))\| \\ &\leq C\varepsilon^{1/2} \left(h^{s_{z}-1/2} \|\boldsymbol{z}\|_{s_{z}} + h^{s_{y}-1/2} \|\boldsymbol{y}\|_{s_{y}} + \delta(s_{y})\varepsilon^{1/2}h\|\Delta\boldsymbol{y}\|_{\mathcal{T}_{h}} \right). \end{aligned}$$

4.2.2 Step 2: errors between the auxiliary problem (4.3) and the discrete problem (2.6)

By (2.6) and (4.3) we have the following error equations.

Lemma 13. Let $(\boldsymbol{q}_h, y_h, \hat{y}_h^o, \boldsymbol{p}_h, z_h, \hat{z}_h^o, u_h) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2 \times M_h^\partial$ and $(\boldsymbol{q}_h(u), y_h(u), \hat{y}_h^o(u), \boldsymbol{p}_h(u), z_h(u), \hat{z}_h^o(u)) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2$ be the solutions of (2.6) and (4.3), respectively. Then for all $(\boldsymbol{r}_1, w_1, \hat{w}_1^o, \boldsymbol{r}_2, w_2, \hat{w}_2^o) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2$, we have

$$\mathcal{B}_{1}(\boldsymbol{q}_{h}-\boldsymbol{q}_{h}(u),y_{h}-y_{h}(u),\widehat{y}_{h}^{o}-\widehat{y}_{h}^{o}(u);\boldsymbol{r}_{1},w_{1},\widehat{w}_{1}^{o}) = -\langle u_{h}-\Pi_{k}^{\partial}u,\tau_{2}w_{1}+\boldsymbol{r}_{1}\cdot\boldsymbol{n}\rangle_{\mathcal{E}_{h}^{\partial}}, \qquad (4.13a)$$

$$\mathcal{B}_{2}(\boldsymbol{p}_{h} - \boldsymbol{p}_{h}(u), z_{h} - z_{h}(u), \widehat{z}_{h}^{o} - \widehat{z}_{h}^{o}(u); \boldsymbol{r}_{2}, w_{2}, \widehat{w}_{2}^{o}) = -(y_{h} - y_{h}(u), w_{2})_{\mathcal{T}_{h}}.$$
(4.13b)

Lemma 14. Let $(\boldsymbol{q}_h, y_h, \hat{y}_h^o, \boldsymbol{p}_h, z_h, \hat{z}_h^o) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2$ and $(\boldsymbol{q}_h(u), y_h(u), \hat{y}_h^o(u), z_h(u), \hat{z}_h^o(u)) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2$ be the solutions of (2.6) and (4.3), respectively. Then we have

$$\begin{aligned} \|y_h - y_h(u)\|_{\mathcal{T}_h}^2 + \gamma \|u_h - \Pi_k^{\partial} u\|_{\mathcal{E}_h^{\partial}}^2 \\ &= \langle u_h - \Pi_k^{\partial} u, \Pi_k^{\partial} (\boldsymbol{p} \cdot \boldsymbol{n}) - \boldsymbol{p}_h(u) \cdot \boldsymbol{n} - \tau_2 (z_h(u) - \widehat{z}_h^o(u)) \rangle_{\mathcal{E}_h^{\partial}} \end{aligned}$$

Proof. Take $(\mathbf{r}_1, w_1, \widehat{w}_1^o) = (\mathbf{p}_h - \mathbf{p}_h(u), z_h - z_h(u), \widehat{z}_h^o - \widehat{z}_h^o(u))$ and $(\mathbf{r}_2, w_2, \widehat{w}_2^o) = (\mathbf{q}_h - \mathbf{q}_h(u), y_h - y_h(u), \widehat{y}_h^o - \widehat{y}_h^o(u))$ in (4.13a) and (4.13b), respectively, and use Lemma 1 to get

$$\begin{aligned} &-\|y_{h} - y_{h}(u)\|_{\mathcal{T}_{h}}^{2} \\ &= \mathcal{B}_{2}(\boldsymbol{p}_{h} - \boldsymbol{p}_{h}(u), z_{h} - z_{h}(u), \widehat{z}_{h}^{o} - \widehat{z}_{h}^{o}(u); \boldsymbol{q}_{h} - \boldsymbol{q}_{h}(u), y_{h} - y_{h}(u), \widehat{y}_{h}^{o} - \widehat{y}_{h}^{o}(u)) \\ &= \mathcal{B}_{1}(\boldsymbol{q}_{h} - \boldsymbol{q}_{h}(u), y_{h} - y_{h}(u), \widehat{y}_{h}^{o} - \widehat{y}_{h}^{o}(u); \boldsymbol{p}_{h} - \boldsymbol{p}_{h}(u), z_{h} - z_{h}(u), \widehat{z}_{h}^{o} - \widehat{z}_{h}^{o}(u)) \\ &= -\langle u_{h} - \Pi_{k}^{\partial} u, (\boldsymbol{p}_{h} - \boldsymbol{p}_{h}(u)) \cdot \boldsymbol{n} + \tau_{2}(z_{h} - z_{h}(u)) \rangle_{\mathcal{E}_{h}^{\partial}}. \end{aligned}$$

Therefore, (2.3e), (2.2) and $\widehat{z}_h^o=\widehat{z}_h^o(u)=0$ on \mathcal{E}_h^∂ give

$$\begin{split} \|y_{h} - y_{h}(u)\|_{\mathcal{T}_{h}}^{2} + \gamma \|u_{h} - \Pi_{k}^{\partial}u\|_{\mathcal{E}_{h}^{\partial}}^{2} \\ &= \langle u_{h} - \Pi_{k}^{\partial}u, \boldsymbol{p}_{h} \cdot \boldsymbol{n} + \tau_{2}z_{h} + \gamma u_{h} \rangle_{\mathcal{E}_{h}^{\partial}} \\ &- \langle u_{h} - \Pi_{k}^{\partial}u, \boldsymbol{p}_{h}(u) \cdot \boldsymbol{n} + \tau_{2}z_{h}(u) + \gamma \Pi_{k}^{\partial}u \rangle_{\mathcal{E}_{h}^{\partial}} \\ &= \langle u_{h} - \Pi_{k}^{\partial}u, \Pi_{k}^{\partial}(\boldsymbol{p} \cdot \boldsymbol{n}) - \boldsymbol{p}_{h}(u) \cdot \boldsymbol{n} - \tau_{2}(z_{h}(u) - \widehat{z}_{h}^{o}(u)) \rangle_{\mathcal{E}_{h}^{\partial}}. \end{split}$$

		_ 1
		_ 1
_	_	_

Lemma 15. Let $(\boldsymbol{q}_h, y_h, \hat{y}_h^o, \boldsymbol{p}_h, z_h, \hat{z}_h^o) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2$ and $(\boldsymbol{q}_h(u), y_h(u), \hat{y}_h^o(u), \boldsymbol{p}_h(u), z_h(u), \hat{z}_h^o(u)) \in [\boldsymbol{V}_h \times W_h \times M_h^o]^2$ be the solutions of (2.6) and (4.3), respectively. If assumptions (A1)-(A3) hold, then there exists h_0 , independent of ε such that for all $h \leq h_0$, we have the estimates

$$\begin{aligned} \|u - u_h\|_{\mathcal{E}_h^{\partial}} &\leq C\left(h^{s_y - 1/2} \|y\|_{s_y} + h^{s_z - 1/2} \|z\|_{s_z} + \delta(s_y)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}\right),\\ \|y_h - y_h(u)\|_{\mathcal{T}_h} &\leq C\left(h^{s_y - 1/2} \|y\|_{s_y} + h^{s_z - 1/2} \|z\|_{s_z} + \delta(s_y)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}\right),\\ \|z_h - z_h(u)\|_{\mathcal{T}_h} &\leq C\left(h^{s_y - 1/2} \|y\|_{s_y} + h^{s_z - 1/2} \|z\|_{s_z} + \delta(s_y)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}\right).\end{aligned}$$

Proof. By Lemma 14, the Cauchy-Schwarz inequality, and Young's inequality one gets

$$\begin{aligned} \|y_h - y_h(u)\|_{\mathcal{T}_h} + \gamma^{1/2} \|u_h - \Pi_k^{\partial} u\|_{\mathcal{E}_h^{\partial}} \\ &\leq C \|\Pi_k^{\partial} (\boldsymbol{p} \cdot \boldsymbol{n}) - \boldsymbol{p}_h(u) \cdot \boldsymbol{n}\|_{\mathcal{E}_h^{\partial}} + C \|\tau_2(z_h(u) - \hat{z}_h^o(u)\|_{\mathcal{E}_h^{\partial}}. \end{aligned}$$

By the triangle inequality, $p = -\varepsilon \nabla z$, an inverse inequality and the estimates in Lemma 15 we

have

$$\begin{split} \|\mathbf{\Pi}_{k}^{\partial}(\boldsymbol{p}\cdot\boldsymbol{n}) - \boldsymbol{p}_{h}(\boldsymbol{u})\cdot\boldsymbol{n}\|_{\mathcal{E}_{h}^{\partial}} \\ &\leq \|\mathbf{\Pi}_{k}^{\partial}(\boldsymbol{p}\cdot\boldsymbol{n}) - \mathbf{\Pi}_{k}^{o}(\boldsymbol{p}\cdot\boldsymbol{n})\|_{\mathcal{E}_{h}^{\partial}} + \|\mathbf{\Pi}_{k}^{o}(\boldsymbol{p}\cdot\boldsymbol{n}) - \boldsymbol{p}_{h}(\boldsymbol{u})\cdot\boldsymbol{n}\|_{\mathcal{E}_{h}^{\partial}} \\ &\leq C\varepsilon h^{s_{z}-3/2}\|\boldsymbol{z}\|_{s_{z}} + \sum_{T\in\mathcal{T}_{h},\partial T\cap\mathcal{E}_{h}^{\partial}\neq\emptyset} h_{T}^{-1/2}\|\mathbf{\Pi}_{k}^{o}\boldsymbol{p} - \boldsymbol{p}_{h}(\boldsymbol{u})\|_{T} \\ &\leq C\varepsilon h^{s_{z}-3/2}\|\boldsymbol{z}\|_{s_{z}} + \sum_{T\in\mathcal{T}_{h},\partial T\cap\mathcal{E}_{h}^{\partial}\neq\emptyset} h_{T}^{-1/2}\left(\|\mathbf{\Pi}_{k}^{o}\boldsymbol{p} - \boldsymbol{p}\|_{T} + \|\boldsymbol{p} - \boldsymbol{p}_{h}(\boldsymbol{u})\|_{T}\right) \\ &\leq Ch^{s_{z}-1/2}\|\boldsymbol{z}\|_{s_{z}} + \varepsilon^{-1/2}\sum_{T\in\mathcal{T}_{h},\partial T\cap\mathcal{E}_{h}^{\partial}\neq\emptyset} \|\boldsymbol{p} - \boldsymbol{p}_{h}(\boldsymbol{u})\|_{T} \\ &\leq Ch^{s_{z}-1/2}\|\boldsymbol{z}\|_{s_{z}} + \varepsilon^{-1/2}\|\boldsymbol{p} - \boldsymbol{p}_{h}(\boldsymbol{u})\|_{\mathcal{T}_{h}} \\ &\leq C\left(h^{s_{y}-1/2}\|\boldsymbol{y}\|_{s_{y}} + h^{s_{z}-1/2}\|\boldsymbol{z}\|_{s_{z}} + \delta(s_{y})\varepsilon^{1/2}h\|\Delta\boldsymbol{y}\|_{\mathcal{T}_{h}}\right). \end{split}$$

By the triangle inequality, z = 0 on Γ , and the estimate (4.12) we have

$$\begin{aligned} \|\tau_2(z_h(u) - \widehat{z}_h^o(u)\|_{\mathcal{E}_h^\partial} \\ &\leq C \|(z_h(u) - \widehat{z}_h^o(u) - \Pi_k^o z + \widetilde{\Pi}_k^\partial z)\|_{\mathcal{E}_h^\partial} + C \|(\Pi_k^o z - \widetilde{\Pi}_k^\partial z)\|_{\mathcal{E}_h^\partial} \\ &\leq C \|(\mathbf{\Pi}_k^o \boldsymbol{p} - \boldsymbol{p}_h(u), \Pi_k^o z - z_h(u), \widetilde{\Pi}_k^\partial z - \widehat{z}_h^o(u))\| + Ch^{s_z - 1/2} \|z\|_{s_z} \\ &\leq C \left(h^{s_y - 1/2} \|y\|_{s_y} + h^{s_z - 1/2} \|z\|_{s_z} + \delta(s_y)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}\right). \end{aligned}$$

This implies

$$\begin{aligned} \|y_h - y_h(u)\|_{\mathcal{T}_h} + \|u_h - \Pi_k^{\partial} u\|_{\mathcal{E}_h^{\partial}} \\ &\leq C\left(h^{s_y - 1/2} \|y\|_{s_y} + h^{s_z - 1/2} \|z\|_{s_z} + \delta(s_y)\varepsilon^{1/2}h\|\Delta y\|_{\mathcal{T}_h}\right). \end{aligned}$$

By the triangle inequality and the fact y = u on \mathcal{E}_h^∂ , we get

$$\begin{aligned} \|u - u_h\|_{\mathcal{E}_h^{\partial}} &\leq \|y - \Pi_k^{\partial} y\|_{\mathcal{E}_h^{\partial}} + \|\Pi_k^{\partial} u - u_h\|_{\mathcal{E}_h^{\partial}} \\ &\leq C\left(h^{s_y - 1/2} \|y\|_{s_y} + h^{s_z - 1/2} \|z\|_{s_z} + \delta(s_y)\varepsilon^{1/2} h\|\Delta y\|_{\mathcal{T}_h}\right). \end{aligned}$$

By Theorem 2 and (4.13b), one has

$$\begin{split} \| (\boldsymbol{p}_{h}(u) - \boldsymbol{p}_{h}, z_{h}(u) - z_{h}, \widehat{z}_{h}^{o}(u) - \widehat{z}_{h}^{o}) \| \\ &\leq C \sup_{(\boldsymbol{r}_{2}, w_{2}, \widehat{w}_{2}^{o}) \in \boldsymbol{V}_{h} \times W_{h} \times M_{h}^{o}} \frac{\mathcal{B}_{2}(\boldsymbol{p}_{h}(u) - \boldsymbol{p}_{h}, z_{h}(u) - z_{h}, \widehat{z}_{h}^{o}(u) - \widehat{z}_{h}^{o}; \boldsymbol{r}_{2}, w_{2}, \widehat{w}_{2}^{o})}{\| (\boldsymbol{r}_{2}, w_{2}, \widehat{w}_{2}^{o}) \|} \\ &\leq C \sup_{(\boldsymbol{r}_{2}, w_{2}, \widehat{w}_{2}^{o}) \in \boldsymbol{V}_{h} \times W_{h} \times M_{h}^{o}} \frac{(y_{h} - y_{h}(u), w_{2})\tau_{h}}{\| (\boldsymbol{r}_{2}, w_{2}, \widehat{w}_{2}^{o}) \|} \\ &\leq C \| y_{h} - y_{h}(u) \|_{\mathcal{T}_{h}} \\ &\leq C \left(h^{s_{y} - 1/2} \| y \|_{s_{y}} + h^{s_{z} - 1/2} \| z \|_{s_{z}} + \delta(s_{y}) \varepsilon^{1/2} h \| \Delta y \|_{\mathcal{T}_{h}} \right). \end{split}$$

Therefore,

$$\|z_h(u) - z_h\|_{\mathcal{T}_h} \le C\varepsilon^{1/2} \left(h^{s_y - 1/2} \|y\|_{s_y} + h^{s_z - 1/2} \|z\|_{s_z} + \delta(s_y)\varepsilon^{1/2} h \|\Delta y\|_{\mathcal{T}_h} \right).$$

$h/\sqrt{2}$	2^{-1}	2^{-2}	2^{-3}	2^{-4}	2^{-5}
$\ y-y_h\ _{0,\Omega}$	6.0299E-02	1.3188E-02	2.1788 E-03	4.8975E-04	1.1863E-04
order	-	2.1929	2.5976	2.1534	2.0456
$\ z-z_h\ _{0,\Omega}$	1.0572 E-01	2.6724E-02	6.2451E-03	1.5091 E-03	3.7092E-04
order	-	1.9841	2.0973	2.0491	2.0245
$\ u-u_h\ _{0,\Gamma}$	2.5537E-01	5.6029 E-02	1.2108E-02	2.8176E-03	6.7424E-04
order	-	2.1883	2.2102	2.1034	2.0631

Table 1: Smooth test with k = 1 and $\varepsilon = 10^{-7}$: Errors for the control u, state y and the adjoint state z.

5 Numerical Experiments

In this section, we report numerical experiments to illustrate our theoretical results. For all experiments, we take $\Omega = [0, 1] \times [0, 1] \subset \mathbb{R}^2$, $\gamma = 1$, and the stabilization functions are chosen as in (2.3f)-(2.3g).

5.1 Smooth test

In our first test, the state, dual state, and convection coefficient are chosen as

$$y = -\varepsilon \pi (\sin(\pi x_1) + \sin(\pi x_2)), \ z = \sin(\pi x_1) \sin(\pi x_2),$$
$$\beta = -[x_1^2 \sin(x_2), \cos(x_1)e^{x_2}],$$

and the source term f and the desired state y_d are generated using the optimality system (1.3) with the above data. We show the numerical results for k = 1 and $\varepsilon = 10^{-7}$ in Table 1.

5.2 Non-smooth test

Next, we choose the data as

$$y_d = x(1-x)y(1-y), \quad f = 0, \text{ and } \quad \beta = -[x_1^2\sin(x_2), \cos(x_1)e^{x_2}].$$

We tested 5 cases with different values for ε and we do not have exact solutions for these problems; we solved the problems numerically for a triangulation with approximately 1.5 million elements and compared these reference solutions against other solutions computed on meshes with larger h. The numerical results are shown in Table 2; the computed convergence rates are erratic and do not follow a clear pattern. The same phenomenon has been observed in another work on a convection dominated Dirichlet boundary control problem [4]. Also, we plot the state, dual state and boundary control in Figures 1 to 5. Furthermore, many works on convection dominated PDEs observe well-behaved convergence if they remove a small portion of the domain containing the layer; see [28, Section 6] for a convection dominated distributed optimal control problem and [20, Table 4 in Section 5.4] for a convection dominated PDEs. We did not to compute the rates by removing the layer is always on the boundary; see Figures 1 to 5.

6 Conclusion

In [23, 29], we studied an HDG method for a *diffuison dominated* convection diffusion Dirichlet boundary control problem. We obtained optimal convergence rates for the control under a high

An HDG Method for Dirichlet Boundary Control of Convection Dominated Diffusion PD	ЭE
---	----

ε	$\frac{h}{\sqrt{2}}$	$\ y-y_h\ _{\mathcal{T}_h}$		$\ z-z_h\ _{\mathcal{T}_h}$		$\ u-u_h\ _{\varepsilon_h^\partial}$	
		Error	Rate	Error	Rate	Error	Rate
1/10	2^{-1}	4.8530E-04		1.0783E-03		2.0808E-03	
	2^{-2}	1.6439E-04	1.56	5.3340E-04	1.01	7.2793E-04	1.51
	2^{-3}	5.5062 E-05	1.58	2.0652 E-04	1.37	2.6827E-04	1.44
	2^{-4}	1.5885E-05	1.79	6.4955 E-05	1.67	9.1834 E-05	1.54
	2^{-5}	4.1123E-06	1.95	$1.7687 \text{E}{-}05$	1.88	2.7451E-05	1.74
	2^{-1}	1.3406E-03		2.2114E-03		4.1984E-03	
	2^{-2}	4.9404E-04	1.44	1.3080E-03	0.75	1.8165E-03	1.21
1/50	2^{-3}	2.4238E-04	1.02	7.3661E-04	0.83	8.9345 E-04	1.02
,	2^{-4}	1.1777E-04	1.04	3.5878E-04	1.04	5.2980E-04	0.75
	2^{-5}	4.1387E-05	1.50	1.4389E-04	1.32	2.6837 E-04	0.98
1/100	2^{-1}	1.6216E-03		2.5742E-03		4.7465E-03	
	2^{-2}	6.6643E-04	1.28	1.5857 E-03	0.70	2.1124 E-03	1.17
	2^{-3}	2.8129E-04	1.24	9.2685 E-04	0.77	1.0435 E-03	1.02
	2^{-4}	1.7254 E-04	0.71	5.2512E-04	0.82	7.3691E-04	0.50
	2^{-5}	8.5868E-05	1.00	2.5477 E-04	1.04	5.0212 E-04	0.53
1/1000	2^{-1}	1.8500E-03		2.9440E-03		5.3059E-03	
	2^{-2}	8.3498E-04	1.14	2.0237 E-03	0.54	2.4554 E-03	1.11
	2^{-3}	3.9689E-04	1.07	1.3617E-03	0.57	1.0865 E-03	1.17
	2^{-4}	2.7140E-04	0.55	9.1122E-04	0.58	5.2906E-04	1.03
	2^{-5}	1.5180E-04	0.84	5.8115 E-04	0.65	4.5760 E-04	0.21
1/10000	2^{-1}	1.9431E-03		3.0087E-03		5.3675E-03	
	2^{-2}	8.4727E-04	1.20	2.0778 E-03	0.53	2.5082 E-03	1.10
	2^{-3}	3.9498E-04	1.10	1.4261 E-03	0.54	1.1393E-03	1.14
	2^{-4}	2.3736E-04	0.73	9.9958E-04	0.51	5.6195 E-04	1.01
	2^{-5}	1.5931E-04	0.58	6.9841 E-04	0.52	3.2155 E-04	0.81

Table 2: Non-smooth test with different ε and k = 1: Errors for the control u, state y and the adjoint state z.



Figure 1: Left is the state y_h and right is the control u_h for $\varepsilon = 1/10$.



Figure 2: Left is the state y_h and right is the control u_h for $\varepsilon = 1/50$.



Figure 3: Left is the state y_h and right is the control u_h for $\varepsilon = 1/100$.



Figure 4: Left is the state y_h and right is the control u_h for $\varepsilon = 1/1000$.



Figure 5: Left is the state y_h and right is the control u_h for $\varepsilon = 1/10000$.

regularity assumption in [29] and a low regularity assumption in [23]. In this work, we considered a different HDG method with a lower computational cost for a convection dominated convection diffusion boundary control problem under high and low regularity conditions and again proved optimal convergence rates for the control. All existing numerical analysis work on Dirichlet boundary control problems have assumed the mesh is quasi-uniform; however, we do not need to have this assumption here.

To the best of our knowledge, this work is the only existing numerical analysis exploration of this convection dominated diffusion Dirichlet control problem.

References

- T. Apel, M. Mateos, J. Pfefferer, and A. Rösch. Error estimates for Dirichlet control problems in polygonal domains: Quasi-uniform meshes. *Math. Control Relat. Fields*, 8(1), 2018.
- [2] Blanca Ayuso and L. Donatella Marini. Discontinuous Galerkin methods for advectiondiffusion-reaction problems. SIAM J. Numer. Anal., 47(2):1391–1420, 2009.
- [3] Roland Becker and Boris Vexler. Optimal control of the convection-diffusion equation using stabilized finite element methods. *Numer. Math.*, 106(3):349–367, 2007.
- [4] Peter Benner and Hamdullah Yücel. A Local Discontinuous Galerkin Method For Dirichlet Boundary Control Problems.
- [5] Susanne C. Brenner and L. Ridgway Scott. *The mathematical theory of finite element methods*, volume 15 of *Texts in Applied Mathematics*. Springer, New York, third edition, 2008.
- [6] A. Buffa, T. J. R. Hughes, and G. Sangalli. Analysis of a multiscale discontinuous Galerkin method for convection-diffusion problems. SIAM J. Numer. Anal., 44(4):1420–1440, 2006.
- [7] Erik Burman and Peter Hansbo. Edge stabilization for Galerkin approximations of convectiondiffusion-reaction problems. *Comput. Methods Appl. Mech. Engrg.*, 193(15-16):1437–1453, 2004.

- [8] E. Casas and J. P. Raymond. Error estimates for the numerical approximation of Dirichlet boundary control for semilinear elliptic equations. SIAM J. Control Optim., 45(5):1586–1611, 2006.
- [9] P. Castillo, B. Cockburn, D. Schötzau, and C. Schwab. Optimal a priori error estimates for the hp-version of the local discontinuous Galerkin method for convection-diffusion problems. *Math. Comp.*, 71(238):455–478, 2002.
- [10] A. Cesmelioglu, B. Cockburn, and W. Qiu. Analysis of a hybridizable discontinuous Galerkin method for the steady-state incompressible Navier-Stokes equations. *Math. Comp.*, 86(306):1643–1670, 2017.
- [11] Yanlai Chen, Bernardo Cockburn, and Bo Dong. Superconvergent HDG methods for linear, stationary, third-order equations in one-space dimension. *Math. Comp.*, 85(302):2715–2742, 2016.
- [12] Y. Cheng, X. Meng, and Q. Zhang. Application of generalized Gauss-Radau projections for the local discontinuous Galerkin method for linear convection-diffusion equations. *Math. Comp.*, 86(305):1233–1267, 2017.
- [13] Y. Cheng and C. W. Shu. Superconvergence of discontinuous Galerkin and local discontinuous Galerkin schemes for linear hyperbolic and convection-diffusion equations in one space dimension. SIAM J. Numer. Anal., 47(6):4044–4072, 2010.
- [14] B. Cockburn, J. Gopalakrishnan, and R. Lazarov. Unified hybridization of discontinuous Galerkin, mixed, and continuous Galerkin methods for second order elliptic problems. *SIAM J. Numer. Anal.*, 47(2):1319–1365, 2009.
- [15] B. Cockburn, J. Gopalakrishnan, N.C. Nguyen, J. Peraire, and F. Sayas. Analysis of HDG methods for Stokes flow. *Math. Comp.*, 80(274):723–760, 2011.
- [16] B. Cockburn and C. W. Shu. The local discontinuous Galerkin method for time-dependent convection-diffusion systems. SIAM J. Numer. Anal., 35(6):2440–2463, 1998.
- [17] Bernardo Cockburn and Kassem Mustapha. A hybridizable discontinuous Galerkin method for fractional diffusion problems. *Numer. Math.*, 130(2):293–314, 2015.
- [18] Bernardo Cockburn and Jiguang Shen. A hybridizable discontinuous Galerkin method for the p-Laplacian. SIAM J. Sci. Comput., 38(1):A545–A566, 2016.
- [19] W. Eckhaus. Boundary layers in linear elliptic singular perturbation problems. SIAM Rev., 14:225–270, 1972.
- [20] G. Fu, W. Qiu, and W. Zhang. An analysis of HDG methods for convection-dominated diffusion problems. ESAIM Math. Model. Numer. Anal., 49(1):225–256, 2015.
- [21] A. V. Fursikov, M. D. Gunzburger, and L. Hou. Boundary value problems and optimal boundary control for the Navier-Stokes system: the two-dimensional case. SIAM J. Control Optim., 36(3):852–894, 1998.
- [22] W. Gong and N. Yan. Mixed finite element method for Dirichlet boundary control problem governed by elliptic PDEs. SIAM J. Control Optim., 49(3):984–1014, 2011.

- [23] Wei Gong, Weiwei Hu, Mariano Mateos, John Singler, Xiao Zhang, and Yangwen Zhang. A New HDG Method for Dirichlet Boundary Control of Convection Diffusion PDEs II: Low Regularity. SIAM J. Numer. Anal., 56(4):2262–2287, 2018.
- [24] M. D. Gunzburger, L. Hou, and T. P. Svobodny. Analysis and finite element approximation of optimal control problems for the stationary Navier-Stokes equations with Dirichlet controls. *RAIRO Modél. Math. Anal. Numér.*, 25(6):711–748, 1991.
- [25] M. D. Gunzburger, L. Hou, and T. P. Svobodny. Boundary velocity control of incompressible flow with an application to viscous drag reduction. SIAM J. Control Optim., 30(1):167–181, 1992.
- [26] M. D. Gunzburger and S. Manservisi. The velocity tracking problem for Navier-Stokes flows with bounded distributed controls. SIAM J. Control Optim., 37(6):1913–1945, 1999.
- [27] M. D. Gunzburger and S. Manservisi. Analysis and approximation of the velocity tracking problem for Navier-Stokes flows with distributed control. SIAM J. Numer. Anal., 37(5):1481– 1512, 2000.
- [28] Matthias Heinkenschloss and Dmitriy Leykekhman. Local error estimates for SUPG solutions of advection-dominated elliptic linear-quadratic optimal control problems. SIAM J. Numer. Anal., 47(6):4607–4638, 2010.
- [29] W. Hu, M. Mateos, J. R. Singler, and Y. Zhang. A New HDG Method for Dirichlet Boundary Control of Convection Diffusion PDEs I: High regularity.
- [30] W. Hu, J. Shen, J. R. Singler, Y. Zhang, and X. Zheng. A Superconvergent Hybridizable Discontinuous Galerkin Method for Dirichlet Boundary Control of Elliptic PDEs.
- [31] Y. Jin. Local analysis of HDG methods for convection-dominated diffusion problems. J. Comput. Appl. Math., 304:57–72, 2016.
- [32] C. Johnson, A. H. Schatz, and L. B. Wahlbin. Crosswind smear and pointwise errors in streamline diffusion finite element methods. *Math. Comp.*, 49(179):25–38, 1987.
- [33] Ji L, Y. Xu, and J.K. Ryan. Accuracy-enhancement of discontinuous Galerkin solutions for convection-diffusion equations in multiple-dimensions. *Math. Comp.*, 81(280):1929–1950, 2012.
- [34] Dmitriy Leykekhman and Matthias Heinkenschloss. Local error analysis of discontinuous Galerkin methods for advection-dominated elliptic linear-quadratic optimal control problems. SIAM J. Numer. Anal., 50(4):2012–2038, 2012.
- [35] Binjie Li and Xiaoping Xie. Analysis of a family of HDG methods for second order elliptic problems. J. Comput. Appl. Math., 307:37–51, 2016.
- [36] Mariano Mateos and Ira Neitzel. Dirichlet control of elliptic state constrained problems. Comput. Optim. Appl., 63(3):825–853, 2016.
- [37] S. May, R. Rannacher, and B. Vexler. Error analysis for a finite element approximation of elliptic Dirichlet boundary control problems. *SIAM J. Control Optim.*, 51(3):2585–2611, 2013.
- [38] Koichi Niijima. Pointwise error estimates for a streamline diffusion finite element scheme. Numer. Math., 56(7):707–719, 1990.

- [39] W. Qiu and K. Shi. An HDG method for convection diffusion equation. J. Sci. Comput., 66(1):346–357, 2016.
- [40] Weifeng Qiu and Ke Shi. A superconvergent HDG method for the incompressible Navier-Stokes equations on general polyhedral meshes. *IMA J. Numer. Anal.*, 36(4):1943–1967, 2016.
- [41] Jan Česenek and Miloslav Feistauer. Theory of the space-time discontinuous Galerkin method for nonstationary parabolic problems with nonlinear convection and diffusion. *SIAM J. Numer. Anal.*, 50(3):1181–1206, 2012.