Stability of strong solutions to the Navier–Stokes–Fourier system

Jan Březina Eduard Feireisl * Antonín Novotný May 23, 2022

> Tokyo Institute of Technology 2-12-1 Ookayama, Meguro-ku, Tokyo, 152-8550, Japan

Institute of Mathematics of the Academy of Sciences of the Czech Republic Žitná 25, CZ-115 67 Praha 1, Czech Republic

Université de Toulon, IMATH, EA 2134 BP 20132, 83957 La Garde, France

Abstract

We identify a large class of objects - dissipative measure-valued (DMV) solutions to the Navier-Stokes-Fourier system - in which the strong solutions are stable. More precisely, a DMV solution coincides with the strong solution emanating from the same initial data as long as the latter exists. The DMV solutions are represented by parameterized families of measures satisfying certain compatibility conditions. They can be seen as an analogue to the dissipative measure-valued solutions introduced earlier in the context of the (inviscid) Euler system.

Keywords: Navier–Stokes–Fourier system, measure–valued solution, weak–strong uniqueness, relative energy

^{*}The research of E.F. leading to these results has received funding from the Czech Sciences Foundation (GAČR), Grant Agreement 18–05974S. The Institute of Mathematics of the Academy of Sciences of the Czech Republic is supported by RVO:67985840.

1 Navier-Stokes-Fourier system

Motivated by our earlier work on the dissipative measure–valued solutions to the Euler system (see [2], [3], [4]), we consider the Navier–Stokes–Fourier system describing the time evolution of a general compressible, viscous, and heat conducting fluid. Our goal is to identify the largest class possible of objects that can be understood as "solutions" to this problem. These objects represent a generalization of classical solutions and will coincide with a classical solution emanating from the same initial data as long as the classical solution exists. They will be described by means of a parameterized family of probability measures $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$, where the vectors (t,x) range over the physical space - the space–time cylinder

$$Q_T = (0, T) \times \Omega$$
, $\Omega \subset \mathbb{R}^N$ a domain, $N = 1, 2, 3$.

We call these objects dissipative measure-valued (DMV) solutions although $\{V_{t,x}\}_{(t,x)\in Q_T}$ may not, in general, coincide with some Young measure generated by a weakly convergent family of state variables.

The concept of measure-valued solutions in the context of *inviscid* fluids has been used by several authors starting with the pioneering papers by DiPerna [9] and DiPerna and Majda [8], see also related results by Kroener and Zajaczkowski [20], Matušů-Nečasová and Novotný [24], Neustupa [26]. A nice summary of the "fluid mechanics" applications of this method is in the monograph by Málek et al. [23]. Recently, Tzavaras and coauthors adapted these ideas to problems in elastodynamics, thermoelasticity, and other related problems [5], [7], [21]. Another application to the Cucker-Smale equation is due to Mucha and Peszek [25]. Interesting results in numerical analysis have been obtained by Fjordholm et al. [16], [17], [18]. Here we apply the same technique to a model of viscous fluids. Accordingly, our main ambition is not to capture possible oscillations in the solution set which might be a highly unlikely phenomenon when viscosity is present, but rather to identify the largest class of objects in which the strong solutions represent a stable set. In particular, the dissipative measure-valued solutions introduced below will satisfy the weakstrong uniqueness principle. We expect these results will find applications whenever approximate solutions are constructed, in particular by means of a numerical scheme. It might be relatively easy to show that families of numerical approximations generate a measure-valued solution, the latter being quite general object. Such an observation together with the weak-strong uniqueness principle will finally provide a rigorous convergence proof as long as the strong solution is available. This philosophy has been successfully applied in [13] in the case of a simpler barotropic fluid flow. We expect that the results obtained in the present paper will give rise to similar applications in the context of physically relevant models of compressible, viscous, and heat conducting fluids.

1.1 Phase variables

The state of the fluid is determined by the standard phase variables: The mass density ϱ , the (absolute) temperature ϑ , and the (bulk) velocity $\mathbf{u} \in R^N$. In addition, in the context of viscous and heat conducting fluid, it is convenient to include $\nabla_x \mathbf{u}$ and $\nabla_x \vartheta$ in the family of state variables.

1.2 Thermodynamic functions

The constitutive properties of a given fluid will be expressed through thermodynamic functions: The pressure $p = p(\varrho, \vartheta)$, the internal energy $e = e(\varrho, \vartheta)$, and the entropy $s = s(\varrho, \vartheta)$, interrelated through Gibbs' equation:

$$\vartheta Ds = De + pD\left(\frac{1}{\varrho}\right). \tag{1.1}$$

1.3 Physical space

We use the Eulerian coordinate system, where the independent variables are the time $t \in [0, T]$, and the spatial position $x \in \Omega$, $\Omega \subset \mathbb{R}^N$ a domain.

1.4 Field equations

The time evolution of $\varrho = \varrho(t, x)$, $\vartheta = \vartheta(t, x)$, and $\mathbf{u} = \mathbf{u}(t, x)$ is determined by a system of partial differential equations expressing the basic physical principles:

• Mass conservation or equation of continuity

$$\partial_t \varrho + \operatorname{div}_x(\varrho \mathbf{u}) = 0;$$
 (1.2)

• Conservation of linear momentum or Newton's second law

$$\partial_t(\rho \mathbf{u}) + \operatorname{div}_x(\rho \mathbf{u} \otimes \mathbf{u}) + \nabla_x p(\rho, \vartheta) = \operatorname{div}_x \mathbb{S},$$
 (1.3)

where S is the viscous stress tensor;

Conservation of internal energy

$$\partial_t(\varrho e(\varrho,\vartheta)) + \operatorname{div}_x(\varrho e(\varrho,\vartheta)\mathbf{u}) + \operatorname{div}_x\mathbf{q} = \mathbb{S} : \nabla_x\mathbf{u} - p(\varrho,\vartheta)\operatorname{div}_x\mathbf{u}, \tag{1.4}$$

where \mathbf{q} is the internal energy flux.

1.5 Diffusion terms

To close the system (1.2–1.4), we suppose that $\mathbb{S} = \mathbb{S}(\varrho, \vartheta, \nabla_x \mathbf{u})$ is given by Newton's rheological law

$$\mathbb{S} = \mu(\varrho, \vartheta) \left(\nabla_x \mathbf{u} + \nabla_x^t \mathbf{u} - \frac{2}{N} \mathrm{div}_x \mathbf{u} \mathbb{I} \right) + \lambda(\varrho, \vartheta) \mathrm{div}_x \mathbf{u} \mathbb{I}; \tag{1.5}$$

while the internal energy flux obeys Fourier's law

$$\mathbf{q} = -\kappa(\varrho, \vartheta) \nabla_x \vartheta. \tag{1.6}$$

The system (1.2–1.6) is termed Navier–Stokes–Fourier system.

1.6 Boundary conditions

We suppose that $\Omega \subset \mathbb{R}^N$ is a bounded domain and consider the no-slip/no flux boundary conditions:

$$\mathbf{u}|_{\partial\Omega} = 0, \ \mathbf{q} \cdot \mathbf{n}|_{\partial\Omega} = \kappa \nabla_x \vartheta \cdot \mathbf{n}|_{\partial\Omega} = 0. \tag{1.7}$$

1.7 The basic laws of thermodynamics

The First law of thermodynamics requires the total energy of the fluid system to be conserved:

$$\partial_t \left(\frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e \right) + \operatorname{div}_x \left[\left(\frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e \right) \mathbf{u} \right] + \operatorname{div}_x (p(\varrho, \vartheta) \mathbf{u}) = \operatorname{div}_x (\mathbb{S} \cdot \mathbf{u}) - \operatorname{div}_x \mathbf{q}. \tag{1.8}$$

Note that (1.8) follows from (1.2–1.4) as long as all quantities are smooth.

The Second law of thermodynamics postulates the entropy production principle:

$$\partial_t(\varrho s) + \operatorname{div}_x(\varrho s \mathbf{u}) + \operatorname{div}_x\left(\frac{\mathbf{q}}{\vartheta}\right) = \sigma,$$
 (1.9)

where the entropy production rate σ is given as

$$\sigma = \frac{1}{\vartheta} \left(\mathbb{S} : \nabla_x \mathbf{u} - \frac{\mathbf{q} \cdot \nabla_x \vartheta}{\vartheta} \right). \tag{1.10}$$

Similarly to (1.8), the entropy balance equation (1.9), (1.10) can be derived directly from (1.2), (1.4), with help of Gibb's relation (1.1), as soon as all quantities are smooth.

The rest of the paper is organized as follows. In Section 2, we introduce the concept of dissipative measure—valued solution to the Navier—Stokes—Fourier system. Section 3 deals with the basic tool for studying stability – the relative energy and the associated relative energy inequality – expressed in terms of DMV solutions. In Section 4, we use the relative energy to compare the "distance" between a DMV solution and a smooth solution of the same problem. In Section 5, we establish the desired weak—strong uniqueness property under several restrictions imposed on both the DMV and the strong solution. In Section 6, we find sufficient conditions for a DMV solution to be a classical one by applying the regularity criterion of Sun, Wang, and Zhang [28]. The paper is concluded by a short discussion on possible extensions of the theory in Section 7.

2 Generalized solutions

The basic idea borrowed from [14] is to retain equations (1.2), (1.3) while (1.4) is replaced by

• total energy balance

$$\partial_t \int_{\Omega} \left(\frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e \right) dx = 0; \tag{2.1}$$

• the entropy inequality

$$\partial_{t}(\varrho s) + \operatorname{div}_{x}(\varrho s \mathbf{u}) + \operatorname{div}_{x}\left(\frac{\mathbf{q}}{\vartheta}\right) = \sigma,$$

$$\sigma \geq \frac{1}{\vartheta}\left(\mathbb{S}: \nabla_{x}\mathbf{u} - \frac{\mathbf{q}\cdot\nabla_{x}\vartheta}{\vartheta}\right).$$
(2.2)

As shown in [14, Chapter 2], any smooth weak (distributional) solution of (1.2), (1.3), (2.1), (2.2) satisfying the boundary conditions (1.7) is a classical solution of the original system (1.2–1.4). The basic theory in the framework of weak (distributional) solutions for the problem (1.2), (1.3), (2.1), (2.2) has been developed in [14]. The main advantage of a weak formulation based on (2.1), (2.2) rather than (1.8) is that the integrated energy balance (2.1) does not contain the convective terms present in (1.8), on which we have no control.

The measure-valued solutions are typically generated by a sequence of weak solutions, see e.g. the monograph by Málek et al. [23]. Here we adopt a slightly different approach advocated by Brennier, DeLellis, and Székelyhidi [1], Gwiazda et al. [12], [19] defining a measure-valued solution as an object independent of any generating procedure whatever the latter might be.

2.1 Phase space

A natural candidate for the phase space is of course the space based on the standard variables $[\varrho, \vartheta, \mathbf{u}]$. However, the entropy balance (2.2), or more specifically, the entropy production rate σ , contains also spatial gradients of (\mathbf{u}, ϑ) . A suitable phase space framework for the measure–valued solutions is therefore

$$\mathcal{F} = \left\{ [\varrho, \vartheta, \mathbf{u}, \mathbb{D}_u, \mathbf{D}_{\vartheta}] \mid \varrho \in [0, \infty), \ \vartheta \in [0, \infty), \ \mathbf{u} \in R^N, \ \mathbb{D}_u \in R^{N \times N}, \ \mathbf{D}_{\vartheta} \in R^N \ \right\},$$

where \mathbb{D}_u stands for the symmetric velocity gradient $\mathbb{D}[\mathbf{u}] \equiv \frac{1}{2} (\nabla_x \mathbf{u} + \nabla_x^t \mathbf{u})$, and \mathbf{D}_{ϑ} for $\nabla_x \vartheta$. Oscillations in sequences of (weak) solutions are usually described in terms of the associated Young measure. Analogously, we define a measure-valued solution to the Navier-Stokes-Fourier system on the space-time cylinder $Q_T = (0, T) \times \Omega$ as a family of probability measures $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$,

$$\mathcal{V}_{t,x} \in \mathcal{P}(\mathcal{F})$$
 for a.a. $(t,x) \in Q_T, \ (t,x) \in Q_T \mapsto \mathcal{M}(\mathcal{F})$ weakly-(*) measurable.

Indeed the measure $\mathcal{V}_{t,x}$ can be interpreted as probability that a solution $[\varrho, \vartheta, \mathbf{u}, \mathbb{D}_u, \mathbf{D}_{\vartheta}]$ at a point $(t,x) \in Q_T$ attains certain value in the phase space \mathcal{F} . Classical "single valued" solutions $[\varrho, \vartheta, \mathbf{u}]$ are represented by a Dirac mass,

$$[\varrho, \vartheta, \mathbf{u}, \mathbb{D}_u, \mathbf{D}_{\vartheta}] \approx \delta_{\varrho,\vartheta,\mathbf{u},\mathbb{D}[\mathbf{u}],\nabla_x\vartheta}.$$

In addition, we require a very natural compatibility condition,

$$\frac{1}{2} \left(\nabla_x \left\langle \mathcal{V}_{t,x}; \mathbf{u} \right\rangle + \nabla_x^t \left\langle \mathcal{V}_{t,x}; \mathbf{u} \right\rangle \right) = \left\langle \mathcal{V}_{t,x}; \mathbb{D}_u \right\rangle, \ \nabla_x \left\langle \mathcal{V}_{t,x}; \vartheta \right\rangle = \left\langle \mathcal{V}_{t,x}; \mathbf{D}_\vartheta \right\rangle \text{ in } \mathcal{D}'(Q_T),$$

more precisely

$$-\int_{0}^{T} \int_{\Omega} \langle \mathcal{V}_{t,x}; \mathbf{u} \rangle \cdot \operatorname{div}_{x} \mathbb{T} \, dx \, dt = \int_{0}^{T} \int_{\Omega} \langle \mathcal{V}_{t,x}; \mathbb{D}_{u} \rangle : \mathbb{T} \, dx \, dt \text{ for any } \mathbb{T} \in C^{1}(\overline{Q}_{T}; R_{\text{sym}}^{N \times N}),$$

$$-\int_{0}^{T} \int_{\Omega} \langle \mathcal{V}_{t,x}; \vartheta \rangle \, \operatorname{div}_{x} \boldsymbol{\varphi} \, dx \, dt = \int_{0}^{T} \int_{\Omega} \langle \mathcal{V}_{t,x}; \mathbf{D}_{\vartheta} \rangle \cdot \boldsymbol{\varphi} \, dx \, dt \text{ for any } \boldsymbol{\varphi} \in C^{1}(\overline{Q}_{T}; R^{N}),$$

$$\boldsymbol{\varphi} \cdot \mathbf{n}|_{\partial\Omega} = 0.$$

$$(2.3)$$

Note that (2.3) also reflects the no-slip condition $\mathbf{u}|_{\partial\Omega} = 0$.

2.2 Initial data

Similarly to the preceding part, the initial state of the system

$$[\varrho_0, \vartheta_0, \mathbf{u}_0] = [\varrho(0, \cdot), \vartheta(0, \cdot), \mathbf{u}(0, \cdot)]$$

can be expressed by means of a parameterized family of measures $\{\mathcal{V}_{0,x}\}_{x\in\Omega}$ acting on the phase space

$$\mathcal{F}_0 = \left\{ [\varrho_0, \vartheta_0, \mathbf{u}_0] \middle| \varrho_0 \in [0, \infty), \ \vartheta_0 \in [0, \infty), \mathbf{u}_0 \in \mathbb{R}^N \right\}.$$

To avoid ambiguity concerning the values of \mathbf{u}_0 on the vacuum, we always assume the initial density ϱ_0 to be bounded below away from zero.

2.3 Field equations

In accordance with the previous discussion, a generalized formulation of the equation of continuity (1.2) reads

$$\left[\int_{\Omega} \langle \mathcal{V}_{t,x}; \varrho \rangle \varphi(t,x) \, dx \right]_{t=0}^{t=\tau} = \int_{0}^{\tau} \int_{\Omega} \left[\langle \mathcal{V}_{t,x}; \varrho \rangle \, \partial_{t} \varphi(t,x) + \langle \mathcal{V}_{t,x}; \varrho \mathbf{u} \rangle \cdot \nabla_{x} \varphi(t,x) \right] \, dx \, dt \qquad (2.4)$$

for a.a. $\tau \in (0,T)$ and any $\varphi \in C_c^{\infty}([0,T) \times \overline{\Omega})$.

Similarly, the momentum equation (1.3) reads

$$\left[\int_{\Omega} \langle \mathcal{V}_{t,x}; \varrho \mathbf{u} \rangle \cdot \boldsymbol{\varphi}(t,x) \, dx\right]_{t=0}^{t=\tau}$$

$$= \int_{0}^{\tau} \int_{\Omega} \left[\langle \mathcal{V}_{t,x}; \varrho \mathbf{u} \rangle \cdot \partial_{t} \boldsymbol{\varphi}(t,x) + \langle \mathcal{V}_{t,x}; \varrho \mathbf{u} \otimes \mathbf{u} \rangle : \nabla_{x} \boldsymbol{\varphi}(t,x) + \langle \mathcal{V}_{t,x}; p(\varrho,\vartheta) \rangle \operatorname{div}_{x} \boldsymbol{\varphi}(t,x) \right] \, dx \, dt$$

$$+ \int_{0}^{\tau} \int_{\Omega} \langle \mathcal{V}_{t,x}; \mathbb{S}(\varrho,\vartheta,\mathbb{D}_{u}) \rangle : \nabla_{x} \boldsymbol{\varphi}(t,x) \, dx \, dt + \int_{0}^{\tau} \int_{\overline{\Omega}} \nabla_{x} \boldsymbol{\varphi} : d\nu_{C} \tag{2.5}$$

for a.a. $\tau \in (0,T)$ and any $\varphi \in C_c^{\infty}([0,T) \times \Omega; \mathbb{R}^N)$. The symbol ν_C in (2.5) stands for the concentration measure. More specifically, ν_C is a (tensor valued) signed Borel measure, $\nu_C \in \mathbb{R}^N$ $\mathcal{M}(\overline{Q}_T; R^{N \times N})$, that will be controlled by the concentration defect introduced below, cf. also [3].

The energy equation (1.8) is replaced by the total energy balance (2.1), more specifically

$$\int_{\Omega} \left\langle \mathcal{V}_{\tau,x}; \frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) \right\rangle dx \le \int_{\Omega} \left\langle \mathcal{V}_{0,x}; \frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) \right\rangle dx \tag{2.6}$$

for a.a. $\tau \in (0,T)$.

Finally, we reformulate the entropy inequality (2.2) as

$$\left[\int_{\Omega} \langle \mathcal{V}_{t,x}; \varrho s(\varrho, \vartheta) \rangle \varphi(t, x) \, dx \right]_{t=0}^{t=\tau}
= \int_{0}^{\tau} \int_{\Omega} \left[\langle \mathcal{V}_{t,x}; \varrho s(\varrho, \vartheta) \rangle \, \partial_{t} \varphi(t, x) + \left\langle \mathcal{V}_{t,x}; \varrho s(\varrho, \vartheta) \mathbf{u} - \frac{\kappa(\varrho, \vartheta)}{\vartheta} \mathbf{D}_{\vartheta} \right\rangle \cdot \nabla_{x} \varphi(t, x) \right] \, dx \, dt \qquad (2.7)
+ \int_{0}^{\tau} \int_{\overline{\Omega}} \varphi(t, x) \, d\sigma \text{ for a.a. } \tau \in (0, T),$$

and any $\varphi \in C_c^{\infty}([0,T) \times \overline{\Omega})$, where the entropy production rate $\sigma \in \mathcal{M}^+(\overline{Q}_T)$ is a non-negative Borel measure satisfying

$$\sigma \ge \left\langle \mathcal{V}_{t,x}; \frac{1}{\vartheta} \left(\mathbb{S}(\varrho, \vartheta, \mathbb{D}_u) : \mathbb{D}_u + \frac{\kappa(\varrho, \vartheta)}{\vartheta} |\mathbf{D}_{\vartheta}|^2 \right) \right\rangle. \tag{2.8}$$

Thus we may rewrite (2.7), (2.8) in a more concise form

$$\left[\int_{\Omega} \langle \mathcal{V}_{t,x}; \varrho s(\varrho, \vartheta) \rangle \varphi(t, x) \, dx \right]_{t=0}^{t=\tau} \\
\geq \int_{0}^{\tau} \int_{\Omega} \left[\langle \mathcal{V}_{t,x}; \varrho s(\varrho, \vartheta) \rangle \, \partial_{t} \varphi(t, x) + \left\langle \mathcal{V}_{t,x}; \varrho s(\varrho, \vartheta) \mathbf{u} - \frac{\kappa(\varrho, \vartheta)}{\vartheta} \mathbf{D}_{\vartheta} \right\rangle \cdot \nabla_{x} \varphi(t, x) \right] \, dx \, dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \frac{1}{\vartheta} \left(\mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) : \mathbb{D}_{u} + \frac{\kappa(\varrho, \vartheta)}{\vartheta} |\mathbf{D}_{\vartheta}|^{2} \right) \right\rangle \varphi(t, x) \, dx \, dt$$
(2.9)

for a.a. $\tau \in (0,T)$ and any $\varphi \in C_c^{\infty}([0,T) \times \overline{\Omega}), \varphi \geq 0$.

Remark 2.1. The functions $(\varrho, \vartheta, \mathbb{D}_u) \mapsto \frac{1}{\vartheta} \mathbb{S}(\varrho, \vartheta, \mathbb{D}_u) : \mathbb{D}_u$ and $(\varrho, \vartheta, \mathbf{D}_{\vartheta}) \mapsto \frac{\kappa(\varrho, \vartheta)}{\vartheta^2} |\mathbf{D}_{\vartheta}|^2$ are defined as

$$\frac{1}{\vartheta} \mathbb{S}(\varrho, \vartheta, \mathbb{D}_u) : \mathbb{D}_u = \lim_{\varepsilon \to 0+} \frac{1}{\vartheta + \varepsilon} \mathbb{S}(\varrho, \vartheta, \mathbb{D}_u) : \mathbb{D}_u \in [0, \infty],
\frac{\kappa(\varrho, \vartheta)}{\vartheta^2} |\mathbf{D}_{\vartheta}|^2 = \lim_{\varepsilon \to 0+} \frac{\kappa(\varrho, \vartheta)}{(\vartheta + \varepsilon)^2} |\mathbf{D}_{\vartheta}|^2 \in [0, \infty]$$

for any $\vartheta \in [0, \infty)$.

The term $\int_0^\tau \int_{\overline{\Omega}} \nabla_x \varphi : d\nu_C$ is understood as the value of the functional ν_C over the continuous function $\nabla_x \varphi$.

2.4 Compatibility conditions

Besides (2.3), further compatibility conditions will be imposed on the measure-valued solution $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$. Note that all compatibility conditions specified below (including (2.3)) are automatically satisfied as long as the parameterized measure $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$ is a Young measure generated by a family $\{\varrho_{\varepsilon}, \vartheta_{\varepsilon}, \mathbf{u}_{\varepsilon}, \mathbb{D}[\nabla_x \mathbf{u}_{\varepsilon}], \nabla_x \vartheta_{\varepsilon}\}_{\varepsilon>0}$ of functions satisfying the bounds

ess
$$\sup_{t \in (0,T)} \int_{\Omega} \left[\varrho_{\varepsilon} + \varrho_{\varepsilon} |\mathbf{u}_{\varepsilon}|^{2} + \varrho_{\varepsilon} e(\varrho_{\varepsilon}, \vartheta_{\varepsilon}) + \varrho_{\varepsilon} |s(\varrho_{\varepsilon}, \vartheta_{\varepsilon})| \right] dx$$

$$+ \int_{0}^{T} \int_{\Omega} \left[\frac{\mu(\varrho_{\varepsilon}, \vartheta_{\varepsilon})}{2\vartheta_{\varepsilon}} \left| \nabla_{x} \mathbf{u}_{\varepsilon} + \nabla_{x}^{t} \mathbf{u}_{\varepsilon} - \frac{2}{N} \operatorname{div}_{x} \mathbf{u}_{\varepsilon} \mathbb{I} \right|^{2} + \frac{\lambda(\varrho_{\varepsilon}, \vartheta_{\varepsilon})}{\vartheta_{\varepsilon}} |\operatorname{div}_{x} \mathbf{u}_{\varepsilon}|^{2} \right] dx dt \qquad (2.10)$$

$$+ \int_{0}^{T} \int_{\Omega} \kappa(\varrho_{\varepsilon}, \vartheta_{\varepsilon}) |\nabla_{x} \log(\vartheta_{\varepsilon})|^{2} dx dt \stackrel{\sim}{\sim} 1$$

uniformly for $\varepsilon \to 0$.

Here and hereafter, the symbol $a \stackrel{<}{\sim} b$ means $a \leq cb$ for a certain constant c > 0.

2.4.1 Concentration defect

Set

$$\mathcal{D}(\tau) = \int_{\Omega} \left\langle \mathcal{V}_{0,x}; \frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) \right\rangle dx - \int_{\Omega} \left\langle \mathcal{V}_{\tau,x}; \frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) \right\rangle dx.$$

The total energy balance (2.6) implies that $\mathcal{D} \geq 0$ a.a. in (0,T). Similarly to [3], we require

$$\int_0^T \psi(t) \int_{\overline{\Omega}} d|\nu_C| \stackrel{<}{\sim} \int_0^T \psi(t) \mathcal{D}(t) dt \text{ for any } \psi \in C[0, T], \ \psi \ge 0$$
 (2.11)

and $\mathcal{D} \in L^{\infty}(0,T)$.

2.4.2 Korn-Poincaré inequality

Let

$$\mathbb{T}[\mathbb{A}] \equiv \mathbb{A} + \mathbb{A}^t - \frac{2}{N} \operatorname{tr}[\mathbb{A}] \mathbb{I}.$$

Keeping in mind (2.3) we impose an analogue of the Korn–Poincaré inequality

$$\int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; |\mathbf{u} - \tilde{\mathbf{u}}(t,x)|^{2} \right\rangle dx dt \stackrel{\leq}{\sim} \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; |\mathbb{T}[\mathbb{D}_{u}] - \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}]|^{2} \right\rangle dx dt \qquad (2.12)$$

for any $\tilde{\mathbf{u}} \in L^2(0,T; W_0^{1,2}(\Omega; \mathbb{R}^N))$. Note that $\mathbb{T}[\mathbb{D}_u] = 2\left(\mathbb{D}_u - \frac{1}{N} \mathrm{tr}[\mathbb{D}_u]\mathbb{I}\right)$.

The following lemma shows that the Korn–Poincaré inequality is naturally "inherited" by weakly converging Sobolev velocity fields vanishing on the boundary $\partial\Omega$.

Lemma 2.2. Suppose that $\{\mathbf{v}_{\varepsilon}\}_{{\varepsilon}>0}$ is a sequence of functions bounded in $L^2(0,T;W_0^{1,2}(\Omega;R^N))$. Let $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$ be a Young measure on the phase space $[R^N;R^{N\times N}]$ generated by $\{\mathbf{v}_{\varepsilon},\nabla_x\mathbf{v}_{\varepsilon}\}$. Then the Korn-Poincaré inequality (2.12) holds.

Proof. As the fields \mathbf{v}_{ε} vanish on $\partial\Omega$, they can be extended to be zero outside Ω . Applying the standard Korn and Poincaré inequalities we get

$$\int_{\Omega} |\mathbf{v}_{\varepsilon}|^{\beta} dx \lesssim \int_{\Omega} |\nabla_{x} \mathbf{v}_{\varepsilon}|^{\beta} dx \lesssim \int_{\Omega} |\nabla_{x} \mathbf{v}_{\varepsilon} + \nabla_{x}^{t} \mathbf{v}_{\varepsilon} - \frac{2}{N} \operatorname{div}_{x} \mathbf{v}_{\varepsilon} \mathbb{I}|^{\beta} dx$$

uniformly in ε for any $1 < \beta < 2$.

Integrating in time and letting $\varepsilon \to 0$ we obtain

$$\int_0^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; |\mathbf{v}|^{\beta} \right\rangle \, \mathrm{d}x \, \mathrm{d}t \stackrel{\leq}{\sim} \int_0^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; |\mathbb{D}_v - \frac{1}{N} \mathrm{tr}[\mathbb{D}_v] \mathbb{I}|^{\beta} \right\rangle \, \mathrm{d}x \, \mathrm{d}t.$$

We conclude the proof by letting $\beta \nearrow 2$ and Monotone convergence theorem.

It is interesting to see that (2.12) follows from a weaker stipulation:

$$\int_{0}^{\tau} \int_{\Omega} \langle \mathcal{V}_{t,x}; |\mathbf{u} - \langle \mathcal{V}_{t,x}; \mathbf{u} \rangle|^{2} \rangle \, dx \, dt \stackrel{\sim}{\sim} \int_{0}^{\tau} \int_{\Omega} \langle \mathcal{V}_{t,x}; |\mathbb{T}[\mathbb{D}_{u}] - \langle \mathcal{V}_{t,x}; \mathbb{T}[\mathbb{D}_{u}] \rangle|^{2} \rangle \, dx \, dt, \qquad (2.13)$$

cf. a similar hypothesis in [12]. Indeed we have

$$\int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; |\mathbf{u} - \tilde{\mathbf{u}}(t,x)|^{2} \right\rangle dx dt$$

$$\stackrel{\sim}{\sim} \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; |\mathbf{u} - \left\langle \mathcal{V}_{t,x}; \mathbf{u} \right\rangle|^{2} \right\rangle dx dt + \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; |\left\langle \mathcal{V}_{t,x}; \mathbf{u} \right\rangle - \tilde{\mathbf{u}}(t,x)|^{2} \right\rangle dx dt$$

$$= \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; |\mathbf{u} - \left\langle \mathcal{V}_{t,x}; \mathbf{u} \right\rangle|^{2} \right\rangle dx dt + \int_{0}^{\tau} \int_{\Omega} |\left\langle \mathcal{V}_{t,x}; \mathbf{u} \right\rangle - \tilde{\mathbf{u}}(t,x)|^{2} dx dt$$

$$\stackrel{\sim}{\sim} \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; |\mathbb{T}[\mathbb{D}_{u}] - \left\langle \mathcal{V}_{t,x}; \mathbb{T}[\mathbb{D}_{u}] \right\rangle|^{2} \right\rangle dx dt + \int_{0}^{\tau} \int_{\Omega} |\left\langle \mathcal{V}_{t,x}; \mathbf{u} \right\rangle - \tilde{\mathbf{u}}(t,x)|^{2} dx dt.$$

Finally, by the standard Korn–Poincaré inequality and (2.3),

$$\int_0^{\tau} \int_{\Omega} |\langle \mathcal{V}_{t,x}; \mathbf{u} \rangle - \tilde{\mathbf{u}}(t,x)|^2 dx dt \stackrel{\leq}{\sim} \int_0^{\tau} \int_{\Omega} |\langle \mathcal{V}_{t,x}; \mathbb{T}[\mathbb{D}_u] \rangle - \mathbb{T}[\nabla_x \tilde{\mathbf{u}}](x)|^2 dx dt;$$

whence (2.12) follows as

$$\langle \mathcal{V}_{t,x}; |\mathbb{T}[\mathbb{D}_u] - \langle \mathcal{V}_{t,x}; \mathbb{T}[\mathbb{D}_u] \rangle|^2 \rangle + |\langle \mathcal{V}_{t,x}; \mathbb{T}[\mathbb{D}_u] \rangle - \mathbb{T}[\nabla_x \tilde{\mathbf{u}}](x)|^2 = \langle \mathcal{V}_{t,x}; |\mathbb{T}[\mathbb{D}_u] - \mathbb{T}[\nabla_x \tilde{\mathbf{u}}](x)|^2 \rangle.$$

2.5 Dissipative measure–valued solutions

We are ready to introduce the concept of dissipative measure-valued (DMV) solutions to the Navier-Stokes-Fourier system.

Definition 2.3. We say that a parameterized family of probability measures $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$ is a dissipative measure-valued (DMV) solution to the Navier-Stokes-Fourier system (1.2–1.7), with the initial conditions $\{\mathcal{V}_{0,x}\}_{x\in\Omega}$ if the following holds:

- $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$ is a weakly-(*) measurable family mapping the physical space Q_T in the set of probability measures on the phase space \mathcal{F} satisfying the compatibility condition (2.3).
- The integral identities (2.4–2.6), (2.9) hold true.
- $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$ complies with the compatibility conditions (2.11), (2.13).

Remark 2.4. Alternatively, a DMV solution could be defined as a Young measure $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$ associated to a family $\{\varrho_{\varepsilon}, \vartheta_{\varepsilon}, \mathbf{u}_{\varepsilon}, \mathbb{D}[\nabla_x \mathbf{u}_{\varepsilon}], \nabla_x \vartheta_{\varepsilon}\}_{\varepsilon>0}$ satisfying the uniform bound (2.10). Then it is sufficient to require $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$ to fulfill the integral identities (2.4–2.6), (2.9). As already pointed out such a definition might be more restrictive than Definition 2.3 but provide better information concerning the properties of $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$. We come back to this issue in the concluding Section 7.

3 Relative energy inequality

Let

$$H_{\tilde{\vartheta}} = \varrho \left[e(\varrho, \vartheta) - \tilde{\vartheta} s(\varrho, \vartheta) \right]$$

be the ballistic free energy, cf. Ericksen [10]. Motivated by [15], we introduce the relative energy

$$\mathcal{E}\left(\varrho,\vartheta,\mathbf{u}\Big|,\tilde{\varrho},\tilde{\vartheta},\tilde{\mathbf{u}}\right) \equiv \frac{1}{2}\varrho|\mathbf{u}-\tilde{\mathbf{u}}|^{2} + \frac{\partial H_{\tilde{\vartheta}}(\tilde{\varrho},\tilde{\vartheta})}{\partial\varrho}(\varrho-\tilde{\varrho}) - H_{\tilde{\vartheta}}(\tilde{\varrho},\tilde{\vartheta})$$

$$= \left[\frac{1}{2}\varrho|\mathbf{u}|^{2} + \varrho e(\varrho,\vartheta)\right] - \left[\frac{1}{2}\tilde{\varrho}|\tilde{\mathbf{u}}|^{2} + \tilde{\varrho}e(\tilde{\varrho},\tilde{\vartheta})\right] + \frac{1}{2}\tilde{\varrho}|\tilde{\mathbf{u}}|^{2} - \varrho\mathbf{u}\cdot\tilde{\mathbf{u}} + \frac{1}{2}\varrho|\tilde{\mathbf{u}}|^{2}$$

$$-\tilde{\vartheta}\left(\varrho s(\varrho,\vartheta) - \tilde{\varrho}s(\tilde{\varrho},\tilde{\vartheta})\right) - \left(e(\tilde{\varrho},\tilde{\vartheta}) - \tilde{\vartheta}s(\tilde{\varrho},\tilde{\vartheta}) + \frac{p(\tilde{\varrho},\tilde{\vartheta})}{\tilde{\varrho}}\right)(\varrho-\tilde{\varrho}),$$
(3.1)

where the last equality was shown in [2, Appendix]. It is worth noting that \mathcal{E} differs from the relative entropy introduced in the classical paper by Dafermos [6] by a multiplicative factor $\tilde{\vartheta}$, see [2]. The advantage of working with relative energy rather than entropy is that the present setting is compatible with the weak formulation (2.4–2.9).

Given a DMV solution $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$ of the Navier–Stokes–Fourier system, we consider the quantity

$$\int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mathcal{E}\left(\varrho, \vartheta, \mathbf{u} \middle|, \tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}\right) \right\rangle dx.$$

After a short inspection of all terms on the right-hand side of (3.1), it is easy to see that

$$\left[\int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mathcal{E} \left(\varrho, \vartheta, \mathbf{u} \middle|, \tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}} \right) \right\rangle \, \mathrm{d}x \right]_{t=0}^{t=\tau}$$

can be expressed by means of the integral identities (2.4–2.9) as soon as the functions $(\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}})$ are smooth and satisfy the "compatibility conditions"

$$\tilde{\varrho} > 0, \ \tilde{\vartheta} > 0, \ \tilde{\mathbf{u}}|_{\partial\Omega} = 0.$$

After a bit tedious but straightforward manipulation that has been performed in detail in [3] and [15], we deduce from (3.1) and (2.4–2.9) the relative energy inequality:

$$\left[\int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mathcal{E}\left(\varrho, \vartheta, \mathbf{u} \middle|, \tilde{\varrho}(t, x), \tilde{\vartheta}(t, x), \tilde{\mathbf{u}}(t, x)\right) \right\rangle \, dx\right]_{t=0}^{t=\tau} + \mathcal{D}(\tau) \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \frac{\tilde{\vartheta}(t, x)}{\vartheta} \mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) : \mathbb{D}_{u} \right\rangle + \tilde{\vartheta}(t, x) \left\langle \mathcal{V}_{t,x}; \frac{\kappa(\varrho, \vartheta)}{\vartheta^{2}} \middle| \mathbf{D}_{\vartheta} \middle|^{2} \right\rangle \right] \, dx \, dt \\
- \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) \right\rangle : \nabla_{x} \tilde{\mathbf{u}}(t, x) \, dx \, dt \\
\leq - \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \varrho s(\varrho, \vartheta) \right\rangle \partial_{t} \tilde{\vartheta}(t, x) + \left\langle \mathcal{V}_{t,x}; \left(\varrho s(\varrho, \vartheta) \mathbf{u} - \frac{\kappa(\varrho, \vartheta)}{\vartheta} \mathbf{D}_{\vartheta}\right) \right\rangle \cdot \nabla_{x} \tilde{\vartheta}(t, x) \right] \, dx \, dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \varrho(\tilde{\mathbf{u}}(t, x) - \mathbf{u}) \right\rangle \cdot \partial_{t} \tilde{\mathbf{u}}(t, x) + \left\langle \mathcal{V}_{t,x}; \varrho(\tilde{\mathbf{u}}(t, x) - \mathbf{u}) \otimes \mathbf{u} \right\rangle : \nabla_{x} \tilde{\mathbf{u}}(t, x) \right] \, dx \, dt \\
- \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \varrho(\varrho, \vartheta) \right\rangle \, div_{x} \tilde{\mathbf{u}}(t, x) \, dx \, dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \varrho \right\rangle \partial_{t} \tilde{\vartheta}(t, x) s(\tilde{\varrho}, \tilde{\vartheta})(t, x) + \left\langle \mathcal{V}_{t,x}; \varrho \mathbf{u} \right\rangle \cdot \nabla_{x} \tilde{\vartheta}(t, x) s(\tilde{\varrho}, \tilde{\vartheta})(t, x) \right] \, dx \, dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \varrho \right\rangle \partial_{t} \tilde{\vartheta}(t, x) s(\tilde{\varrho}, \tilde{\vartheta})(t, x) + \left\langle \mathcal{V}_{t,x}; \varrho \mathbf{u} \right\rangle \cdot \nabla_{x} \tilde{\vartheta}(t, x) s(\tilde{\varrho}, \tilde{\vartheta})(t, x) \right] \, dx \, dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \varrho(t, x) - \varrho \right\rangle \frac{1}{\tilde{\varrho}} \partial_{t} p(\tilde{\varrho}, \tilde{\vartheta})(t, x) - \left\langle \mathcal{V}_{t,x}; \varrho \mathbf{u} \right\rangle \cdot \frac{1}{\tilde{\varrho}} \nabla_{x} p(\tilde{\varrho}, \tilde{\vartheta})(t, x) \right] \, dx \, dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \varrho(t, x) - \varrho \right\rangle \frac{1}{\tilde{\varrho}} \partial_{t} p(\tilde{\varrho}, \tilde{\vartheta})(t, x) - \left\langle \mathcal{V}_{t,x}; \varrho \mathbf{u} \right\rangle \cdot \frac{1}{\tilde{\varrho}} \nabla_{x} p(\tilde{\varrho}, \tilde{\vartheta})(t, x) \right] \, dx \, dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \varrho(t, x) - \varrho \right\rangle \frac{1}{\tilde{\varrho}} \partial_{t} p(\tilde{\varrho}, \tilde{\vartheta})(t, x) - \left\langle \mathcal{V}_{t,x}; \varrho \mathbf{u} \right\rangle \cdot \frac{1}{\tilde{\varrho}} \nabla_{x} p(\tilde{\varrho}, \tilde{\vartheta})(t, x) \right] \, dx \, dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \varrho(t, x) - \varrho \right\rangle \frac{1}{\tilde{\varrho}} \partial_{t} p(\tilde{\varrho}, \tilde{\vartheta})(t, x) - \left\langle \mathcal{V}_{t,x}; \varrho \mathbf{u} \right\rangle \cdot \frac{1}{\tilde{\varrho}} \nabla_{x} p(\tilde{\varrho}, \tilde{\vartheta})(t, x) \right] \, dx \, dt$$

for any

$$\tilde{\varrho}, \tilde{\vartheta} \in C^1(\overline{\Omega}), \ \tilde{\varrho}, \tilde{\vartheta} > 0, \ \tilde{\mathbf{u}} \in C^1([0, T] \times \overline{\Omega}; R^N), \ \tilde{\mathbf{u}}|_{\partial\Omega} = 0.$$
 (3.3)

4 Weak vs. strong solutions

Following the approach of [15] we use the relative energy to compare a (DMV) solution with a strong solution starting from the same initial data. To this end, we introduce a cut-off function

$$\psi_{\delta} \in C_c^{\infty}((0,\infty)^2), \ 0 \le \psi_{\delta} \le 1, \ \psi_{\delta}(\varrho,\vartheta) = 1 \text{ whenever } \delta < \varrho < \frac{1}{\delta} \text{ and } \delta < \vartheta < \frac{1}{\delta} \text{ for some } \delta > 0.$$

For $h = h(\varrho, \vartheta, \mathbf{u}, \mathbb{D}_u, \mathbf{D}_{\vartheta})$ we consider a decomposition

$$h = h_{\text{ess}} + h_{\text{res}}, \ h_{\text{ess}} = \psi_{\delta}(\varrho, \vartheta) h(\varrho, \vartheta, \mathbf{u}, \mathbb{D}_u, \mathbf{D}_{\vartheta}), \ h_{\text{res}} = (1 - \psi_{\delta}(\varrho, \vartheta)) h(\varrho, \vartheta, \mathbf{u}, \mathbb{D}_u, \mathbf{D}_{\vartheta}).$$

Suppose that the thermodynamic functions are interrelated through (1.1), and, in addition, satisfy the *hypothesis of thermodynamic stability*:

$$\frac{\partial p(\varrho, \vartheta)}{\partial \varrho} > 0, \ \frac{\partial e(\varrho, \vartheta)}{\partial \vartheta} > 0 \text{ for any } \varrho, \vartheta > 0.$$
 (4.1)

Under these circumstances, it was shown in [14, Chapter 3, Proposition 3.2] that

$$\mathcal{E}\left(\varrho,\vartheta,\mathbf{u}\middle|\tilde{\varrho},\tilde{\vartheta},\tilde{\mathbf{u}}\right) \\
\geq c(\delta)\left\{\left[\left|\varrho-\tilde{\varrho}\right|^{2}+\left|\vartheta-\tilde{\vartheta}\right|^{2}+\left|\mathbf{u}-\tilde{\mathbf{u}}\right|^{2}\right]_{\text{ess}}+\left[1+\varrho+\varrho|s(\varrho,\vartheta)|+\varrho e(\varrho,\vartheta)+\varrho|\mathbf{u}|^{2}\right]_{\text{res}}\right\}$$
(4.2)

whenever $2\delta < \tilde{\varrho} < \frac{1}{\delta} - \delta$, $2\delta < \tilde{\vartheta} < \frac{1}{\delta} - \delta$ for some $\delta > 0$. Accordingly, the function $\mathcal{E}\left(\varrho,\vartheta,\mathbf{u}\middle|\tilde{\varrho},\tilde{\vartheta},\tilde{\mathbf{u}}\right)$ can be seen as a kind of distance between the triples $[\varrho,\vartheta,\mathbf{u}]$ and $[\tilde{\varrho},\tilde{\vartheta},\tilde{\mathbf{u}}]$.

4.1 Comparison between strong and (DMV) solutions

Let $[\tilde{\varrho}, \vartheta, \tilde{\mathbf{u}}]$ be a strong (classical) solution of the Navier–Stokes–Fourier system (1.1–1.7) on a time interval [0, T]. More specifically, the functions $[\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}]$ belong to the class

$$\tilde{\varrho},\ \tilde{\vartheta}\in C^1([0,T]\times\overline{\Omega}),\ \tilde{\mathbf{u}}\in C^1([0,T]\times\overline{\Omega};R^N),\ D_x^2\tilde{\vartheta},\ D_x^2\tilde{\mathbf{u}}\in C([0,T]\times\overline{\Omega})$$

and satisfy (1.1-1.7), and (3.3). Set

$$\varrho_0(x) = \tilde{\varrho}(0, x), \ \vartheta_0(x) = \tilde{\vartheta}(0, x), \ \mathbf{u}_0(x) = \tilde{\mathbf{u}}(0, x), \ x \in \Omega$$

and consider the initial data for (DMV) solutions in the form

$$\mathcal{V}_{0,x} = \delta_{\varrho_0(x),\vartheta_0(x),\mathbf{u}_0(x)}, \ x \in \Omega. \tag{4.3}$$

In what follows, we use the relative energy inequality (3.2) to compare a (DMV) solution $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$ emanating from the initial data (4.3) with the strong solution $[\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}]$. In view of (4.2), we fix $\delta > 0$ small enough so that

$$0 < 2\delta < \tilde{\varrho}(t,x), \tilde{\vartheta}(t,x) < \frac{1}{\delta} - \delta \text{ for all } (t,x) \in \overline{Q}_T.$$

Setting

$$\mathcal{H}(\tau) \equiv \int_{\Omega} \left\langle \mathcal{V}_{\tau,x}; \mathcal{E}\left(\varrho, \vartheta, \mathbf{u} \middle|, \tilde{\varrho}(\tau, x), \tilde{\vartheta}(\tau, x), \tilde{\mathbf{u}}(\tau, x)\right) \right\rangle dx + \mathcal{D}(\tau),$$

where \mathcal{D} is the concentration defect identified in Section 2.4.1, and using (4.3), together with the compatibility condition (2.11), we deduce from (3.2) that

$$\mathcal{H}(\tau) + \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \frac{\tilde{\vartheta}}{\vartheta} \mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) : \mathbb{D}_{u} \right\rangle + \tilde{\vartheta} \left\langle \mathcal{V}_{t,x}; \frac{\kappa(\varrho, \vartheta)}{\vartheta^{2}} | \mathbf{D}_{\vartheta} |^{2} \right\rangle \right] dx dt \\
- \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) \right\rangle : \nabla_{x}\tilde{\mathbf{u}} dx dt - \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \frac{\kappa(\varrho, \vartheta)}{\vartheta} \mathbf{D}_{\vartheta} \right\rangle \cdot \nabla_{x}\tilde{\vartheta} dx dt \\
\lesssim - \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \varrho s(\varrho, \vartheta) \right\rangle \partial_{t}\tilde{\vartheta} + \left\langle \mathcal{V}_{t,x}; \varrho s(\varrho, \vartheta) \mathbf{u} \right\rangle \cdot \nabla_{x}\tilde{\vartheta} \right] dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \varrho(\tilde{\mathbf{u}} - \mathbf{u}) \right\rangle \cdot \partial_{t}\tilde{\mathbf{u}} + \left\langle \mathcal{V}_{t,x}; \varrho(\tilde{\mathbf{u}} - \mathbf{u}) \otimes \mathbf{u} \right\rangle : \nabla_{x}\tilde{\mathbf{u}} \right] dx dt \\
- \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \varrho(\varrho, \vartheta) \right\rangle div_{x}\tilde{\mathbf{u}} dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \varrho \right\rangle \partial_{t}\tilde{\vartheta}s(\tilde{\varrho}, \tilde{\vartheta}) + \left\langle \mathcal{V}_{t,x}; \varrho \mathbf{u} \right\rangle \cdot \nabla_{x}\tilde{\vartheta}s(\tilde{\varrho}, \tilde{\vartheta}) \right] dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \varrho \right\rangle \partial_{t}\tilde{\vartheta}s(\tilde{\varrho}, \tilde{\vartheta}) + \left\langle \mathcal{V}_{t,x}; \varrho \mathbf{u} \right\rangle \cdot \nabla_{x}\tilde{\vartheta}s(\tilde{\varrho}, \tilde{\vartheta}) \right] dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \tilde{\varrho} - \varrho \right\rangle \frac{1}{\tilde{\varrho}} \partial_{t}p(\tilde{\varrho}, \tilde{\vartheta}) - \left\langle \mathcal{V}_{t,x}; \varrho \mathbf{u} \right\rangle \cdot \frac{1}{\tilde{\varrho}} \nabla_{x}p(\tilde{\varrho}, \tilde{\vartheta}) \right] dx dt \\
+ \int_{0}^{\tau} \mathcal{H}(t) dt. \tag{4.4}$$

The constant hidden in $\stackrel{<}{\sim}$ may depend on the norm of the strong solution $[\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}]$. Our goal in the remaining part of this section is to use the fact that $[\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}]$ represent a strong solution of the Navier–Stokes–Fourier system to rewrite (4.4) in a more concise form so that a Gronwall type argument may be applied to deduce $\mathcal{H} = 0$.

4.1.1 Step 1

Using the momentum equation

$$\partial_t \tilde{\mathbf{u}} + \tilde{\mathbf{u}} \cdot \nabla_x \tilde{\mathbf{u}} = -\frac{1}{\tilde{\varrho}} \nabla_x p(\tilde{\varrho}, \tilde{\vartheta}) + \frac{1}{\tilde{\varrho}} \operatorname{div}_x \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_x \tilde{\mathbf{u}})$$

we deduce

$$\begin{split} \left\langle \mathcal{V}_{t,x}; \varrho(\tilde{\mathbf{u}} - \mathbf{u}) \right\rangle \cdot \partial_{t} \tilde{\mathbf{u}} + \left\langle \mathcal{V}_{t,x}; \varrho(\tilde{\mathbf{u}} - \mathbf{u}) \otimes \mathbf{u} \right\rangle : \nabla_{x} \tilde{\mathbf{u}} \\ &= \left\langle \mathcal{V}_{t,x}; \varrho(\tilde{\mathbf{u}} - \mathbf{u}) \right\rangle \cdot \left(\partial_{t} \tilde{\mathbf{u}} + \tilde{\mathbf{u}} \cdot \nabla_{x} \tilde{\mathbf{u}} \right) + \left\langle \mathcal{V}_{t,x}; \varrho(\tilde{\mathbf{u}} - \mathbf{u}) \otimes \left(\mathbf{u} - \tilde{\mathbf{u}} \right) \right\rangle : \nabla_{x} \tilde{\mathbf{u}} \\ &\leq - \left\langle \mathcal{V}_{t,x}; \varrho(\tilde{\mathbf{u}} - \mathbf{u}) \right\rangle \cdot \left(\frac{1}{\tilde{\varrho}} \nabla_{x} p(\tilde{\varrho}, \tilde{\vartheta}) - \frac{1}{\tilde{\varrho}} \mathrm{div}_{x} \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x} \tilde{\mathbf{u}}) \right) + \left\langle \mathcal{V}_{t,x}; \mathcal{E}\left(\varrho, \vartheta, \mathbf{u} \middle| \tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}} \right) \right\rangle \\ &= - \left\langle \mathcal{V}_{t,x}; \varrho(\tilde{\mathbf{u}} - \mathbf{u}) \right\rangle \cdot \frac{1}{\tilde{\varrho}} \nabla_{x} p(\tilde{\varrho}, \tilde{\vartheta}) \\ &+ \left\langle \mathcal{V}_{t,x}; \left(\frac{\varrho}{\tilde{\varrho}} - 1\right) (\tilde{\mathbf{u}} - \mathbf{u}) \right\rangle \cdot \mathrm{div}_{x} \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x} \tilde{\mathbf{u}}) + \left\langle \mathcal{V}_{t,x}; (\tilde{\mathbf{u}} - \mathbf{u}) \right\rangle \cdot \mathrm{div}_{x} \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x} \tilde{\mathbf{u}}) \\ &+ \left\langle \mathcal{V}_{t,x}; \mathcal{E}\left(\varrho, \vartheta, \mathbf{u} \middle| \tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}} \right) \right\rangle. \end{split}$$

Next, using the compatibility condition (2.3) we may perform by parts integration

$$\int_{\Omega} \langle \mathcal{V}_{t,x}; (\tilde{\mathbf{u}} - \mathbf{u}) \rangle \cdot \operatorname{div}_{x} \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x} \tilde{\mathbf{u}}) \, dx = -\int_{\Omega} \langle \mathcal{V}_{t,x}; (\mathbb{D}[\tilde{\mathbf{u}}] - \mathbb{D}_{u}) \rangle : \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x} \tilde{\mathbf{u}}) \, dx.$$

In view of the previous discussion, the relation (4.4) reduces to

$$\mathcal{H}(\tau) + \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \frac{\tilde{\vartheta}}{\vartheta} \mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) : \mathbb{D}_{u} \right\rangle + \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x}\tilde{\mathbf{u}}) : \nabla_{x}\tilde{\mathbf{u}} \right] dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \tilde{\vartheta} \left\langle \mathcal{V}_{t,x}; \kappa(\varrho, \vartheta) \frac{\mathbf{D}_{\vartheta}}{\vartheta} \cdot \left(\frac{\mathbf{D}_{\vartheta}}{\vartheta} - \nabla_{x} \log(\tilde{\vartheta}) \right) \right\rangle dx dt \\
- \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) \right\rangle : \nabla_{x}\tilde{\mathbf{u}} dx dt - \int_{0}^{\tau} \int_{\Omega} \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x}\tilde{\mathbf{u}}) : \left\langle \mathcal{V}_{t,x}; \mathbb{D}_{u} \right\rangle dx dt \\
\sim - \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \varrho \left[s(\varrho, \vartheta) - s(\tilde{\varrho}, \tilde{\vartheta}) \right] \right\rangle \partial_{t}\tilde{\vartheta} + \left\langle \mathcal{V}_{t,x}; \varrho \left[s(\varrho, \vartheta) - s(\tilde{\varrho}, \tilde{\vartheta}) \right] \mathbf{u} \right\rangle \cdot \nabla_{x}\tilde{\vartheta} \right] dx dt \\
- \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \varrho(\varrho, \vartheta) \right\rangle div_{x}\tilde{\mathbf{u}} dx dt - \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \varrho\tilde{\mathbf{u}} \right\rangle \cdot \frac{1}{\tilde{\varrho}} \nabla_{x} p(\tilde{\varrho}, \tilde{\vartheta}) dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \tilde{\varrho} - \varrho \right\rangle \frac{1}{\tilde{\varrho}} \partial_{t} p(\tilde{\varrho}, \tilde{\vartheta}) dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; |\varrho - \tilde{\varrho}| |\mathbf{u} - \tilde{\mathbf{u}}| \right\rangle dx dt + \int_{0}^{\tau} \mathcal{H}(t) dt, \tag{4.5}$$

4.1.2 Step 2

We have

$$\left\langle \mathcal{V}_{t,x}; \varrho \left[s(\varrho, \vartheta) - s(\tilde{\varrho}, \tilde{\vartheta}) \right] \mathbf{u} \right\rangle \cdot \nabla_{x} \tilde{\vartheta} = \left\langle \mathcal{V}_{t,x}; \varrho \left[s(\varrho, \vartheta) - s(\tilde{\varrho}, \tilde{\vartheta}) \right] \right\rangle \tilde{\mathbf{u}} \cdot \nabla_{x} \tilde{\vartheta}$$

$$+ \left\langle \mathcal{V}_{t,x}; \varrho \left[s(\varrho, \vartheta) - s(\tilde{\varrho}, \tilde{\vartheta}) \right] (\mathbf{u} - \tilde{\mathbf{u}}) \right\rangle \cdot \nabla_{x} \tilde{\vartheta}.$$

Next, we get

$$\int_{\Omega} \langle \mathcal{V}_{t,x}; \varrho \tilde{\mathbf{u}} \rangle \cdot \frac{1}{\tilde{\varrho}} \nabla_{x} p(\tilde{\varrho}, \tilde{\vartheta}) \, dx$$

$$= \int_{\Omega} \langle \mathcal{V}_{t,x}; (\varrho - \tilde{\varrho}) \rangle \, \tilde{\mathbf{u}} \cdot \frac{1}{\tilde{\varrho}} \nabla_{x} p(\tilde{\varrho}, \tilde{\vartheta}) \, dx + \int_{\Omega} \tilde{\mathbf{u}} \cdot \nabla_{x} p(\tilde{\varrho}, \tilde{\vartheta}) \, dx$$

$$= \int_{\Omega} \langle \mathcal{V}_{t,x}; (\varrho - \tilde{\varrho}) \rangle \, \tilde{\mathbf{u}} \cdot \frac{1}{\tilde{\varrho}} \nabla_{x} p(\tilde{\varrho}, \tilde{\vartheta}) \, dx - \int_{\Omega} \operatorname{div}_{x} \tilde{\mathbf{u}} \cdot p(\tilde{\varrho}, \tilde{\vartheta}) \, dx.$$

Consequently, we rewrite (4.5) as

$$\mathcal{H}(\tau) + \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \frac{\tilde{\vartheta}}{\vartheta} \mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) : \mathbb{D}_{u} \right\rangle + \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x}\tilde{\mathbf{u}}) : \nabla_{x}\tilde{\mathbf{u}} \right] dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \tilde{\vartheta} \left\langle \mathcal{V}_{t,x}; \kappa(\varrho, \vartheta) \frac{\mathbf{D}_{\vartheta}}{\vartheta} \cdot \left(\frac{\mathbf{D}_{\vartheta}}{\vartheta} - \nabla_{x} \log(\tilde{\vartheta}) \right) \right\rangle dx dt \\
- \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) \right\rangle : \nabla_{x}\tilde{\mathbf{u}} dx dt - \int_{0}^{\tau} \int_{\Omega} \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x}\tilde{\mathbf{u}}) : \left\langle \mathcal{V}_{t,x}; \mathbb{D}_{u} \right\rangle dx dt \\
\lesssim - \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \tilde{\varrho} \left[s(\varrho, \vartheta) - s(\tilde{\varrho}, \tilde{\vartheta}) \right]_{\text{ess}} \right\rangle \left(\partial_{t}\tilde{\vartheta} + \tilde{\mathbf{u}} \cdot \nabla_{x}\tilde{\vartheta} \right) dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; p(\tilde{\varrho}, \tilde{\vartheta}) - p(\varrho, \vartheta) \right\rangle div_{x}\tilde{\mathbf{u}} dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \tilde{\varrho} - \varrho \right\rangle \frac{1}{\tilde{\varrho}} \left(\partial_{t}p(\tilde{\varrho}, \tilde{\vartheta}) + \tilde{\mathbf{u}} \cdot \nabla_{x}p(\tilde{\varrho}, \tilde{\vartheta}) \right) dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; |\varrho - \tilde{\varrho}| |\mathbf{u} - \tilde{\mathbf{u}}| \right\rangle + \left\langle \mathcal{V}_{t,x}; \varrho \left| s(\varrho, \vartheta) - s(\tilde{\varrho}, \tilde{\vartheta}) \right| |\mathbf{u} - \tilde{\mathbf{u}}| \right\rangle \right] dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \varrho \left| \left[s(\varrho, \vartheta) - s(\tilde{\varrho}, \tilde{\vartheta}) \right]_{\text{res}} \right| \right\rangle dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \varrho \left| \left[s(\varrho, \vartheta) - s(\tilde{\varrho}, \tilde{\vartheta}) \right]_{\text{res}} \right| \right\rangle dx dt \\
+ \int_{0}^{\tau} \mathcal{H}(t) dt,$$

4.1.3 Step 3

Using the equation of continuity (1.2), Gibbs' relation (1.1), and the entropy balance (1.9), (1.10), we deduce

$$\begin{aligned}
&\left(p(\tilde{\varrho},\tilde{\vartheta}) - p(\varrho,\vartheta)\right) \operatorname{div}_{x}\tilde{\mathbf{u}} + (\tilde{\varrho} - \varrho) \frac{1}{\tilde{\varrho}} \left(\partial_{t}p(\tilde{\varrho},\tilde{\vartheta}) + \tilde{\mathbf{u}} \cdot \nabla_{x}p(\tilde{\varrho},\tilde{\vartheta})\right) \\
&= \left(p(\tilde{\varrho},\tilde{\vartheta}) - \frac{\partial p(\tilde{\varrho},\tilde{\vartheta})}{\partial \varrho} (\tilde{\varrho} - \varrho) - \frac{\partial p(\tilde{\varrho},\tilde{\vartheta})}{\partial \vartheta} (\tilde{\vartheta} - \vartheta) - p(\varrho,\vartheta)\right) \operatorname{div}_{x}\tilde{\mathbf{u}} \\
&- \left[\tilde{\varrho} \frac{\partial s(\tilde{\varrho},\tilde{\vartheta})}{\partial \varrho} (\tilde{\varrho} - \varrho) + \tilde{\varrho} \frac{\partial s(\tilde{\varrho},\tilde{\vartheta})}{\partial \vartheta} (\tilde{\vartheta} - \vartheta)\right] \left(\partial_{t}\tilde{\vartheta} + \tilde{\mathbf{u}} \cdot \nabla_{x}\tilde{\vartheta}\right) \\
&+ (\tilde{\vartheta} - \vartheta) \operatorname{div}_{x} \left(\kappa(\tilde{\varrho},\tilde{\vartheta})\nabla_{x}\log\tilde{\vartheta}\right) + \left(1 - \frac{\vartheta}{\tilde{\vartheta}}\right) \left(\mathbb{S}(\tilde{\varrho},\tilde{\vartheta},\nabla_{x}\tilde{\mathbf{u}}) : \nabla_{x}\tilde{\mathbf{u}}\right) + (\tilde{\vartheta} - \vartheta)\kappa(\tilde{\varrho},\tilde{\vartheta})|\nabla_{x}\log\tilde{\vartheta}|^{2}.
\end{aligned}$$

Plugging this in (4.6) we conclude

$$\mathcal{H}(\tau) + \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \frac{\tilde{\vartheta}}{\vartheta} \mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) : \mathbb{D}_{u} \right\rangle + \left\langle \mathcal{V}_{t,x}; \frac{\vartheta}{\tilde{\vartheta}} \right\rangle \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x}\tilde{\mathbf{u}}) : \nabla_{x}\tilde{\mathbf{u}} \right] \, dx \, dt \\
+ \int_{0}^{\tau} \int_{\Omega} \tilde{\vartheta} \left\langle \mathcal{V}_{t,x}; \kappa(\varrho, \vartheta) \frac{\mathbf{D}_{\vartheta}}{\vartheta} \cdot \left(\frac{\mathbf{D}_{\vartheta}}{\vartheta} - \nabla_{x} \log(\tilde{\vartheta}) \right) \right\rangle \, dx \, dt \\
+ \int_{0}^{\tau} \int_{\Omega} \kappa(\tilde{\varrho}, \tilde{\vartheta}) \nabla_{x} \log(\tilde{\vartheta}) \cdot \left\langle \mathcal{V}_{t,x}; \vartheta \left(\nabla_{x} \log(\tilde{\vartheta}) - \frac{\mathbf{D}_{\vartheta}}{\vartheta} \right) \right\rangle \, dx \, dt \\
- \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) \right\rangle : \nabla_{x}\tilde{\mathbf{u}} \, dx \, dt - \int_{0}^{\tau} \int_{\Omega} \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x}\tilde{\mathbf{u}}) : \left\langle \mathcal{V}_{t,x}; \mathbb{D}_{u} \right\rangle \, dx \, dt \\
\tilde{\sim} \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \left| p(\tilde{\varrho}, \tilde{\vartheta}) - \frac{\partial p(\tilde{\varrho}, \tilde{\vartheta})}{\partial \varrho} (\tilde{\varrho} - \varrho) - \frac{\partial p(\tilde{\varrho}, \tilde{\vartheta})}{\partial \vartheta} (\tilde{\vartheta} - \vartheta) - p(\varrho, \vartheta) \right| \right\rangle \, dx \, dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \left| \varrho - \tilde{\varrho} \right| \, |\mathbf{u} - \tilde{\mathbf{u}}| \right\rangle + \left\langle \mathcal{V}_{t,x}; \varrho \left| s(\varrho, \vartheta) - s(\tilde{\varrho}, \tilde{\vartheta}) \right| \, |\mathbf{u} - \tilde{\mathbf{u}}| \right\rangle \right] \, dx \, dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \left| \varrho - \tilde{\varrho} \right| \, |\mathbf{u} - \tilde{\mathbf{u}}| \right\rangle + \left\langle \mathcal{V}_{t,x}; \varrho \left| s(\varrho, \vartheta) - s(\tilde{\varrho}, \tilde{\vartheta}) \right| \, |\mathbf{u} - \tilde{\mathbf{u}}| \right\rangle \right] \, dx \, dt \\
+ \int_{0}^{\tau} \mathcal{H}(t) \, dt.$$

Finally, as a consequence of (4.2), we have

$$\int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \left| p(\tilde{\varrho}, \tilde{\vartheta}) - \frac{\partial p(\tilde{\varrho}, \tilde{\vartheta})}{\partial \varrho} (\tilde{\varrho} - \varrho) - \frac{\partial p(\tilde{\varrho}, \tilde{\vartheta})}{\partial \vartheta} (\tilde{\vartheta} - \vartheta) - p(\varrho, \vartheta) \right| \right\rangle dx dt
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \left| \varrho - \tilde{\varrho} \right| \left| \mathbf{u} - \tilde{\mathbf{u}} \right| \right\rangle + \left\langle \mathcal{V}_{t,x}; \varrho \left| s(\varrho, \vartheta) - s(\tilde{\varrho}, \tilde{\vartheta}) \right| \left| \mathbf{u} - \tilde{\mathbf{u}} \right| \right\rangle \right] dx dt
+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \varrho \left| \left[s(\varrho, \vartheta) - s(\tilde{\varrho}, \tilde{\vartheta}) \right]_{\text{res}} \right| \right\rangle dx dt
\stackrel{\leq}{\sim} \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \left[\vartheta + \left| p(\varrho, \vartheta) \right| + \left| \mathbf{u} - \tilde{\mathbf{u}} \right| + \varrho \left| s(\varrho, \vartheta) \right| \left| \mathbf{u} \right| \right]_{\text{res}} \right\rangle dx dt + \int_{0}^{\tau} \mathcal{H}(t) dt.$$
(4.8)

Thus the relations (4.7), (4.8) give rise to:

$$\mathcal{H}(\tau) + \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \frac{\tilde{\vartheta}}{\vartheta} \mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) : \mathbb{D}_{u} \right\rangle + \left\langle \mathcal{V}_{t,x}; \frac{\vartheta}{\tilde{\vartheta}} \right\rangle \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x}\tilde{\mathbf{u}}) : \nabla_{x}\tilde{\mathbf{u}} \right] dx dt \\ - \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) \right\rangle : \nabla_{x}\tilde{\mathbf{u}} dx dt - \int_{0}^{\tau} \int_{\Omega} \mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x}\tilde{\mathbf{u}}) : \left\langle \mathcal{V}_{t,x}; \mathbb{D}_{u} \right\rangle dx dt \\ + \int_{0}^{\tau} \int_{\Omega} \tilde{\vartheta} \left\langle \mathcal{V}_{t,x}; \kappa(\varrho, \vartheta) \frac{\mathbf{D}_{\vartheta}}{\vartheta} \cdot \left(\frac{\mathbf{D}_{\vartheta}}{\vartheta} - \nabla_{x} \log(\tilde{\vartheta}) \right) \right\rangle dx dt \\ + \int_{0}^{\tau} \int_{\Omega} \kappa(\tilde{\varrho}, \tilde{\vartheta}) \nabla_{x} \log(\tilde{\vartheta}) \cdot \left\langle \mathcal{V}_{t,x}; \vartheta \left(\nabla_{x} \log(\tilde{\vartheta}) - \frac{\mathbf{D}_{\vartheta}}{\vartheta} \right) \right\rangle dx dt \\ \leq \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \left[\vartheta + |p(\varrho, \vartheta)| + |\mathbf{u} - \tilde{\mathbf{u}}| + \varrho |s(\varrho, \vartheta)| |\mathbf{u}| \right]_{\text{res}} \right\rangle dx dt + \int_{0}^{\tau} \mathcal{H}(t) dt.$$

Since S is given by Newton's rheological law (1.5), we have

$$\mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) : \mathbb{D}_{u} = \frac{\mu(\varrho, \vartheta)}{2} \mathbb{T}[\mathbb{D}_{u}] : \mathbb{T}[\mathbb{D}_{u}] + \lambda(\varrho, \vartheta) \operatorname{tr}^{2}[\mathbb{D}_{u}],
\mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x}\tilde{\mathbf{u}}) : \nabla_{x}\tilde{\mathbf{u}} = \frac{\mu(\tilde{\varrho}, \tilde{\vartheta})}{2} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] : \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] + \lambda(\tilde{\varrho}, \tilde{\vartheta}) (\operatorname{div}_{x}\tilde{\mathbf{u}})^{2},
\mathbb{S}(\tilde{\varrho}, \tilde{\vartheta}, \nabla_{x}\tilde{\mathbf{u}}) : \mathbb{D}_{u} = \frac{\mu(\tilde{\varrho}, \tilde{\vartheta})}{2} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] : \mathbb{T}[\mathbb{D}_{u}] + \lambda(\tilde{\varrho}, \tilde{\vartheta}) (\operatorname{div}_{x}\tilde{\mathbf{u}}) (\operatorname{tr}[\mathbb{D}_{u}]),
\mathbb{S}(\varrho, \vartheta, \mathbb{D}_{u}) : \nabla_{x}\tilde{\mathbf{u}} = \frac{\mu(\varrho, \vartheta)}{2} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] : \mathbb{T}[\mathbb{D}_{u}] + \lambda(\varrho, \vartheta) (\operatorname{div}_{x}\tilde{\mathbf{u}}) (\operatorname{tr}[\mathbb{D}_{u}]),$$

and hence we can rewrite (4.9) as

$$\mathcal{H}(\tau) + \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \frac{\tilde{\vartheta}\mu(\varrho,\vartheta)}{2\vartheta} \mathbb{T}[\mathbb{D}_{u}] : \mathbb{T}[\mathbb{D}_{u}] \right\rangle + \left\langle \mathcal{V}_{t,x}; \frac{\vartheta\mu(\tilde{\varrho},\tilde{\vartheta})}{2\tilde{\vartheta}} \right\rangle \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] : \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] \right] dx dt \\
- \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \frac{\mu(\varrho,\vartheta)}{2} \mathbb{T}[\mathbb{D}_{u}] \right\rangle : \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] dx dt - \int_{0}^{\tau} \int_{\Omega} \frac{\mu(\tilde{\varrho},\tilde{\vartheta})}{2} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] : \left\langle \mathcal{V}_{t,x}; \mathbb{T}[\mathbb{D}_{u}] \right\rangle dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left[\left\langle \mathcal{V}_{t,x}; \frac{\tilde{\vartheta}\lambda(\varrho,\vartheta)}{\vartheta} \operatorname{tr}^{2}[\mathbb{D}_{u}] \right\rangle + \left\langle \mathcal{V}_{t,x}; \frac{\vartheta\lambda(\tilde{\varrho},\tilde{\vartheta})}{\tilde{\vartheta}} \right\rangle (\operatorname{div}_{x}\tilde{\mathbf{u}})^{2} \right] dx dt \\
- \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \lambda(\varrho,\vartheta) \operatorname{tr}[\mathbb{D}_{u}] \right\rangle \operatorname{div}_{x}\tilde{\mathbf{u}} dx dt - \int_{0}^{\tau} \int_{\Omega} \lambda(\tilde{\varrho},\tilde{\vartheta}) \operatorname{div}_{x}\tilde{\mathbf{u}} \left\langle \mathcal{V}_{t,x}; \operatorname{tr}[\mathbb{D}_{u}] \right\rangle dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \tilde{\vartheta} \left\langle \mathcal{V}_{t,x}; \kappa(\varrho,\vartheta) \left| \frac{\mathbf{D}_{\vartheta}}{\vartheta} - \nabla_{x} \log(\tilde{\vartheta}) \right|^{2} \right\rangle dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \kappa(\tilde{\varrho},\tilde{\vartheta}) \nabla_{x} \log(\tilde{\vartheta}) \cdot \left\langle \mathcal{V}_{t,x}; (\vartheta - \tilde{\vartheta}) \left(\nabla_{x} \log(\tilde{\vartheta}) - \frac{\mathbf{D}_{\vartheta}}{\vartheta} \right) \right\rangle dx dt \\
- \int_{0}^{\tau} \int_{\Omega} \nabla_{x}\tilde{\vartheta} \cdot \left\langle \mathcal{V}_{t,x}; (\kappa(\varrho,\vartheta) - \kappa(\tilde{\varrho},\tilde{\vartheta})) \left(\nabla_{x} \log(\tilde{\vartheta}) - \frac{\mathbf{D}_{\vartheta}}{\vartheta} \right) \right\rangle dx dt \\
\lesssim \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \left[\vartheta + |p(\varrho,\vartheta)| + |\mathbf{u} - \tilde{\mathbf{u}}| + \varrho|s(\varrho,\vartheta)| |\mathbf{u}| \right]_{\text{res}} \right\rangle dx dt + \int_{0}^{\tau} \mathcal{H}(t) dt. \tag{4.10}$$

Finally, we rewrite (4.10) in the form

$$\mathcal{H}(\tau) + \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \frac{\mu(\varrho,\vartheta)}{2} \middle| \sqrt{\frac{\tilde{\vartheta}}{\vartheta}} \mathbb{T}[\mathbb{D}_{u}] - \sqrt{\frac{\vartheta}{\tilde{\vartheta}}} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] \middle|^{2} \right\rangle dx dt \\
- \frac{1}{2} \int_{0}^{\tau} \int_{\Omega} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] : \left\langle \mathcal{V}_{t,x}; \left(\mu(\varrho,\vartheta) - \mu(\tilde{\varrho},\tilde{\vartheta})\right) \left(\frac{\vartheta}{\tilde{\vartheta}} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] - \mathbb{T}[\mathbb{D}_{u}]\right) \right\rangle dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \lambda(\varrho,\vartheta) \middle| \sqrt{\frac{\tilde{\vartheta}}{\vartheta}} tr[\mathbb{D}_{u}] - \sqrt{\frac{\vartheta}{\tilde{\vartheta}}} div_{x}\tilde{\mathbf{u}} \middle|^{2} \right\rangle dx dt \\
- \int_{0}^{\tau} \int_{\Omega} div_{x}\tilde{\mathbf{u}} \left\langle \mathcal{V}_{t,x}; \left(\lambda(\varrho,\vartheta) - \lambda(\tilde{\varrho},\tilde{\vartheta})\right) \left(\frac{\vartheta}{\tilde{\vartheta}} div_{x}\tilde{\mathbf{u}} - tr[\mathbb{D}_{u}]\right) \right\rangle dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \tilde{\vartheta} \left\langle \mathcal{V}_{t,x}; \kappa(\varrho,\vartheta) \middle| \frac{\mathbf{D}_{\vartheta}}{\vartheta} - \nabla_{x}\log(\tilde{\vartheta}) \middle|^{2} \right\rangle dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \kappa(\tilde{\varrho},\tilde{\vartheta}) \nabla_{x}\log(\tilde{\vartheta}) \cdot \left\langle \mathcal{V}_{t,x}; (\vartheta - \tilde{\vartheta}) \left(\nabla_{x}\log(\tilde{\vartheta}) - \frac{\mathbf{D}_{\vartheta}}{\vartheta}\right) \right\rangle dx dt \\
- \int_{0}^{\tau} \int_{\Omega} \nabla_{x}\tilde{\vartheta} \cdot \left\langle \mathcal{V}_{t,x}; \left(\kappa(\varrho,\vartheta) - \kappa(\tilde{\varrho},\tilde{\vartheta})\right) \left(\nabla_{x}\log(\tilde{\vartheta}) - \frac{\mathbf{D}_{\vartheta}}{\vartheta}\right) \right\rangle dx dt \\
\tilde{\sim} \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \left[\vartheta + |p(\varrho,\vartheta)| + |\mathbf{u} - \tilde{\mathbf{u}}| + \varrho|s(\varrho,\vartheta)| |\mathbf{u}|\right]_{res} \right\rangle dx dt + \int_{0}^{\tau} \mathcal{H}(t) dt.$$

In the remaining part of this paper, we apply a Gronwall type argument to (4.11) to deduce $\mathcal{H} \equiv 0$ - the strong and the (DMV) solutions coincide in Q_T . This can be done under additional restrictions imposed either on the (DMV) solution and/or on the constitutive relations. The first rather easy observation is that the first integral on the right-hand side of (4.11) vanishes as long as the "residual" component is empty. More specifically, if there exists $\delta > 0$ such that

$$\mathcal{V}_{t,x}\left\{0<\delta<\varrho<\frac{1}{\delta},\ 0<\delta<\vartheta<\frac{1}{\delta}
ight\}=1 \text{ for a.a. } (t,x)\in Q_T.$$

Under these circumstances, the left-hand side of (4.11) can be bounded from below as

$$\int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \frac{\mu(\varrho,\vartheta)}{2} \left| \sqrt{\frac{\tilde{\vartheta}}{\vartheta}} \mathbb{T}[\mathbb{D}_{u}] - \sqrt{\frac{\vartheta}{\tilde{\vartheta}}} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] \right|^{2} \right\rangle dx dt \\
- \frac{1}{2} \int_{0}^{\tau} \int_{\Omega} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] : \left\langle \mathcal{V}_{t,x}; \left(\mu(\varrho,\vartheta) - \mu(\tilde{\varrho},\tilde{\vartheta})\right) \left(\frac{\vartheta}{\tilde{\vartheta}} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] - \mathbb{T}[\mathbb{D}_{u}]\right) \right\rangle dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \lambda(\varrho,\vartheta) \left| \sqrt{\frac{\tilde{\vartheta}}{\vartheta}} tr[\mathbb{D}_{u}] - \sqrt{\frac{\vartheta}{\tilde{\vartheta}}} div_{x}\tilde{\mathbf{u}} \right|^{2} \right\rangle dx dt \\
- \int_{0}^{\tau} \int_{\Omega} div_{x}\tilde{\mathbf{u}} \left\langle \mathcal{V}_{t,x}; \left(\lambda(\varrho,\vartheta) - \lambda(\tilde{\varrho},\tilde{\vartheta})\right) \left(\frac{\vartheta}{\tilde{\vartheta}} div_{x}\tilde{\mathbf{u}} - tr[\mathbb{D}_{u}]\right) \right\rangle dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \tilde{\vartheta} \left\langle \mathcal{V}_{t,x}; \kappa(\varrho,\vartheta) \left| \frac{\mathbf{D}_{\vartheta}}{\vartheta} - \nabla_{x} \log(\tilde{\vartheta}) \right|^{2} \right\rangle dx dt \\
+ \int_{0}^{\tau} \int_{\Omega} \kappa(\tilde{\varrho},\tilde{\vartheta}) \nabla_{x} \log(\tilde{\vartheta}) \cdot \left\langle \mathcal{V}_{t,x}; (\vartheta - \tilde{\vartheta}) \left(\nabla_{x} \log(\tilde{\vartheta}) - \frac{\mathbf{D}_{\vartheta}}{\vartheta}\right) \right\rangle dx dt \\
- \int_{0}^{\tau} \int_{\Omega} \nabla_{x}\tilde{\vartheta} \cdot \left\langle \mathcal{V}_{t,x}; \left(\kappa(\varrho,\vartheta) - \kappa(\tilde{\varrho},\tilde{\vartheta})\right) \left(\nabla_{x} \log(\tilde{\vartheta}) - \frac{\mathbf{D}_{\vartheta}}{\vartheta}\right) \right\rangle dx dt \\
\approx - \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; |\varrho - \tilde{\varrho}|^{2} + |\vartheta - \tilde{\vartheta}|^{2} \right\rangle dx dt.$$

Thus applying Gronwall's lemma to the resulting expression we obtain the following result:

Theorem 4.1. Let the transport coefficients $\kappa = \kappa(\varrho, \vartheta)$, $\mu(\varrho, \vartheta)$, and $\lambda(\varrho, \vartheta)$ be continuously differentiable and positive for $\varrho > 0$, $\vartheta > 0$. Let the thermodynamic functions p, e, and s be smooth for ϱ , $\vartheta > 0$, satisfying Gibbs' relation (1.1) and the hypothesis of thermodynamic stability (4.1).

Assume that $[\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}]$ is a classical solution to the Navier–Stokes–Fourier system (1.2–1.10) in $[0, T] \times \Omega$ emanating from the initial data $[\varrho_0, \vartheta_0, \mathbf{u}_0]$ with $\varrho_0, \vartheta_0 > 0$ in $\overline{\Omega}$. Assume further that $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$ is a (DMV) solution to the same problem in the sense of Definition 2.3 such that

$$\mathcal{V}_{t,x}\left\{0 < \delta < \varrho < \frac{1}{\delta}, \ 0 < \delta < \vartheta < \frac{1}{\delta}\right\} = 1 \text{ for a.a. } (t,x) \in Q_T$$

for some $\delta > 0$ and

$$\mathcal{V}_{0,x} = \delta_{[\varrho_0,\vartheta_0,\mathbf{u}_0]} \text{ for a.a. } x \in \Omega.$$

Then

$$\mathcal{V}_{t,x} = \delta_{\left[\tilde{\varrho}(t,x),\tilde{\vartheta}(t,x),\tilde{\mathbf{u}}(t,x),\mathbb{D}\left[\tilde{\mathbf{u}}\right](t,x),\nabla_{x}\tilde{\vartheta}(t,x)\right]} \text{ for a.a. } (t,x) \in Q_{T}.$$

5 Weak-strong uniqueness

Our goal is to extend validity of Theorem 4.1 to a larger class of (DMV) solutions. For the sake of simplicity, we restrict ourselves to the case of vanishing bulk viscosity

$$\lambda \equiv 0. \tag{5.1}$$

In the remaining part of this section, we impose a structural restriction

$$|p(\varrho,\vartheta)| \stackrel{<}{\sim} (1 + \varrho e(\varrho,\vartheta) + \varrho |s(\varrho,\vartheta)|).$$
 (5.2)

Note that this is not a restrictive hypotheses, at least for gases, for which $p \approx \varrho e$. In accordance with (4.2), (5.1), (5.2), relation (4.11) reduces to

$$\mathcal{H}(\tau) + \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \frac{\mu(\varrho,\vartheta)}{2} \middle| \sqrt{\frac{\tilde{\vartheta}}{\vartheta}} \mathbb{T}[\mathbb{D}_{u}] - \sqrt{\frac{\vartheta}{\tilde{\vartheta}}} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] \middle|^{2} \right\rangle dx dt$$

$$- \frac{1}{2} \int_{0}^{\tau} \int_{\Omega} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] : \left\langle \mathcal{V}_{t,x}; \left(\mu(\varrho,\vartheta) - \mu(\tilde{\varrho},\tilde{\vartheta})\right) \left(\frac{\vartheta}{\tilde{\vartheta}} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] - \mathbb{T}[\mathbb{D}_{u}]\right) \right\rangle dx dt$$

$$+ \int_{0}^{\tau} \int_{\Omega} \tilde{\vartheta} \left\langle \mathcal{V}_{t,x}; \kappa(\varrho,\vartheta) \middle| \frac{\mathbf{D}_{\vartheta}}{\vartheta} - \nabla_{x} \log(\tilde{\vartheta}) \middle|^{2} \right\rangle dx dt$$

$$+ \int_{0}^{\tau} \int_{\Omega} \kappa(\tilde{\varrho},\tilde{\vartheta}) \nabla_{x} \log(\tilde{\vartheta}) \cdot \left\langle \mathcal{V}_{t,x}; (\vartheta - \tilde{\vartheta}) \left(\nabla_{x} \log(\tilde{\vartheta}) - \frac{\mathbf{D}_{\vartheta}}{\vartheta}\right) \right\rangle dx dt$$

$$- \int_{0}^{\tau} \int_{\Omega} \nabla_{x}\tilde{\vartheta} \cdot \left\langle \mathcal{V}_{t,x}; \left(\kappa(\varrho,\vartheta) - \kappa(\tilde{\varrho},\tilde{\vartheta})\right) \left(\nabla_{x} \log(\tilde{\vartheta}) - \frac{\mathbf{D}_{\vartheta}}{\vartheta}\right) \right\rangle dx dt$$

$$\stackrel{\sim}{\sim} \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \left[\vartheta + |\mathbf{u} - \tilde{\mathbf{u}}| + \varrho|s(\varrho,\vartheta)| |\mathbf{u}|\right]_{\text{res}} \right\rangle dx dt + \int_{0}^{\tau} \mathcal{H}(t) dt.$$
(5.3)

5.1 Conditional results

We focus on the case with particular transport coefficients:

$$\kappa > 0 \text{ constant}, \ \mu(\vartheta) = \mu_0 + \mu_1 \vartheta, \ \mu_0 > 0, \ \mu_1 \ge 0.$$
(5.4)

In particular,

$$\frac{\mu_{1}}{2} \left\langle \mathcal{V}_{t,x}; \vartheta \left| \sqrt{\frac{\tilde{\vartheta}}{\vartheta}} \mathbb{T}[\mathbb{D}_{u}] - \sqrt{\frac{\vartheta}{\tilde{\vartheta}}} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] \right|^{2} \right\rangle - \frac{\mu_{1}}{2} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] : \left\langle \mathcal{V}_{t,x}; (\vartheta - \tilde{\vartheta}) \left(\frac{\vartheta}{\tilde{\vartheta}} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] - \mathbb{T}[\mathbb{D}_{u}] \right) \right\rangle \\
= \frac{\mu_{1}}{2} \left\langle \mathcal{V}_{t,x}; \tilde{\vartheta} \middle| \mathbb{T}[\mathbb{D}_{u}] - \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] \middle|^{2} \right\rangle - \frac{\mu_{1}}{2} \left\langle \mathcal{V}_{t,x}; (\vartheta - \tilde{\vartheta}) \left(\mathbb{T}[\mathbb{D}_{u}] - \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] \right) \right\rangle : \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}].$$

Consequently, relation (5.3) simplifies as

$$\mathcal{H}(\tau) + \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mu_{0} \left| \sqrt{\frac{\tilde{\vartheta}}{\vartheta}} \mathbb{T}[\mathbb{D}_{u}] - \sqrt{\frac{\vartheta}{\tilde{\vartheta}}} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] \right|^{2} \right\rangle dx dt$$

$$+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mu_{1} \middle| \mathbb{T}[\mathbb{D}_{u}] - \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] \middle|^{2} + \kappa \middle| \frac{\mathbf{D}_{\vartheta}}{\vartheta} - \nabla_{x} \log(\tilde{\vartheta}) \middle|^{2} \right\rangle dx dt$$

$$\lesssim \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \left[\vartheta^{2} + |\mathbf{u} - \tilde{\mathbf{u}}| + \varrho |s(\varrho, \vartheta)| |\mathbf{u}| \right]_{res} \right\rangle dx dt + \int_{0}^{\tau} \mathcal{H}(t) dt \text{ for a.a. } \tau \in (0, T).$$

$$(5.5)$$

5.1.1 Constant transport coefficients

We start with the simplest case when κ and μ satisfy (5.4) with $\mu_1 = 0$.

Theorem 5.1. Let κ and μ be positive constants. Let the thermodynamic functions p, e, and s be smooth functions of (ϱ, ϑ) satisfying Gibbs' relation (1.1), the hypothesis of thermodynamic stability (4.1), together with (5.2).

Assume that $[\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}]$ is a classical solution to the Navier–Stokes–Fourier system (1.2–1.10) in $[0, T] \times \Omega$ emanating from the initial data $[\varrho_0, \vartheta_0, \mathbf{u}_0]$ with $\varrho_0, \vartheta_0 > 0$ in $\overline{\Omega}$. Assume further that $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$ is a (DMV) solution to the same problem in the sense of Definition 2.3 such that

$$\mathcal{V}_{t,x}\left\{\vartheta \leq \overline{\vartheta}, |\mathbf{u}| \leq \overline{\mathbf{u}}\right\} = 1 \text{ for a.a. } (t,x) \in Q_T$$
 (5.6)

for some constants $\overline{\vartheta}$, $\overline{\mathbf{u}}$ and

$$\mathcal{V}_{0,x} = \delta_{[\varrho_0,\vartheta_0,\mathbf{u}_0]} \text{ for a.a. } x \in \Omega.$$

Then

$$\mathcal{V}_{t,x} = \delta_{[\tilde{\varrho}(t,x),\tilde{\vartheta}(t,x),\tilde{\mathbf{u}}(t,x),\mathbb{D}[\tilde{\mathbf{u}}](t,x),\nabla_x\tilde{\vartheta}(t,x)]} \text{ for a.a. } (t,x) \in Q_T.$$

Proof. By virtue of hypothesis (5.6), we have

$$\left\langle \mathcal{V}_{t,x}; \left[\vartheta^2 + |\mathbf{u} - \tilde{\mathbf{u}}| + \varrho |s(\varrho, \vartheta)| |\mathbf{u}| \right]_{\text{res}} \right\rangle \stackrel{<}{\sim} \left\langle \mathcal{V}_{t,x}; \left[1 + \varrho |s(\varrho, \vartheta)| \right]_{\text{res}} \right\rangle;$$

whence, in accordance with (4.2), the inequality (5.5) reduces to

$$\mathcal{H}(\tau) + \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mu_{0} \left| \sqrt{\frac{\tilde{\vartheta}}{\vartheta}} \mathbb{T}[\mathbb{D}_{u}] - \sqrt{\frac{\vartheta}{\tilde{\vartheta}}} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] \right|^{2} \right\rangle \, dx \, dt \\
+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \kappa \left| \frac{\mathbf{D}_{\vartheta}}{\vartheta} - \nabla_{x} \log(\tilde{\vartheta}) \right|^{2} \right\rangle \, dx \, dt \lesssim \int_{0}^{\tau} \mathcal{H}(t) \, dt \text{ for a.a. } \tau \in (0, T).$$

Thus the desired conclusion follows by applying the standard Gronwall lemma argument.

Remark 5.2. Note that the conclusions of Theorems 4.1 and 5.1 remain valid even if the hypothesis (2.13) (the analogue of Korn–Poincaré inequality) is omitted. If, in addition, $\lambda = 0$, we may replace \mathbb{D}_u by its traceless component $\mathbb{D}_u - \frac{1}{N} \text{tr}[\mathbb{D}_u] \mathbb{I}$. This might be of interest when applying these results to certain numerical schemes for which the Korn–Poincaré inequality does not hold.

5.1.2 Temperature dependent viscosity

In the case of temperature dependent viscosity, specifically if $\mu_1 > 0$, we may use the Korn–Poincaré inequality (2.12) in order to handle the velocity dependent terms on the right hand side of (5.5):

$$\int_{0}^{\tau} \int_{\Omega} \langle \mathcal{V}_{t,x}, | [\mathbf{u} - \tilde{\mathbf{u}}]_{\text{res}} | \rangle \, dx \, dt \lesssim \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}, \left[\frac{1}{2\delta} \right]_{\text{res}} + \delta |\mathbf{u} - \tilde{\mathbf{u}}|^{2} \right\rangle \, dx \, dt$$
$$\lesssim c(\delta) \int_{0}^{\tau} \mathcal{H}(t) \, dt + \delta \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; |\mathbf{u} - \tilde{\mathbf{u}}|^{2} \right\rangle \, dx \, dt.$$

In view of (2.12), the last integral may be absorbed by the left hand side of (5.5). We thereby obtain:

$$\mathcal{H}(\tau) + \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mu_{0} \left| \sqrt{\frac{\tilde{\vartheta}}{\vartheta}} \mathbb{T}[\mathbb{D}_{u}] - \sqrt{\frac{\vartheta}{\tilde{\vartheta}}} \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] \right|^{2} \right\rangle dx dt$$

$$+ \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \mu_{1} \left| \mathbb{T}[\mathbb{D}_{u}] - \mathbb{T}[\nabla_{x}\tilde{\mathbf{u}}] \right|^{2} + \kappa \left| \frac{\mathbf{D}_{\vartheta}}{\vartheta} - \nabla_{x} \log(\tilde{\vartheta}) \right|^{2} \right\rangle dx dt$$

$$\lesssim \int_{0}^{\tau} \int_{\Omega} \left\langle \mathcal{V}_{t,x}; \left[\vartheta^{2} + \varrho |s(\varrho, \vartheta)| |\mathbf{u}| \right]_{\text{res}} \right\rangle dx dt + \int_{0}^{\tau} \mathcal{H}(t) dt \text{ for a.a. } \tau \in (0, T).$$

$$(5.7)$$

Theorem 5.3. Let κ and μ satisfy (5.4) with $\mu_1 > 0$. Let the thermodynamic functions p, e, and s comply with the perfect gas constitutive relations

$$p(\varrho, \vartheta) = \varrho \vartheta, \ e(\varrho, \vartheta) = c_v \vartheta, \ s(\varrho, \vartheta) = \log \left(\frac{\vartheta^{c_v}}{\varrho}\right), \ c_v > 1.$$

Assume that $[\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}]$ is a classical solution to the Navier–Stokes–Fourier system (1.2–1.10) in $[0, T] \times \Omega$ emanating from the initial data $[\varrho_0, \vartheta_0, \mathbf{u}_0]$ with $\varrho_0, \vartheta_0 > 0$ in $\overline{\Omega}$. Assume further that $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$ is a (DMV) solution to the same problem in the sense of Definition 2.3 such that

$$\mathcal{V}_{t,x}\left\{|s(\varrho,\vartheta)| \le \overline{s}\right\} = 1 \text{ for } a.a. \ (t,x) \in Q_T$$

$$(5.8)$$

for a certain constant \overline{s} and

$$\mathcal{V}_{0,x} = \delta_{[\varrho_0,\vartheta_0,\mathbf{u}_0]} \text{ for a.a. } x \in \Omega.$$

Then

$$\mathcal{V}_{t,x} = \delta_{[\tilde{\varrho}(t,x),\tilde{\vartheta}(t,x),\tilde{\mathbf{u}}(t,x),\mathbb{D}[\tilde{\mathbf{u}}](t,x),\nabla_x\tilde{\vartheta}(t,x)]} \text{ for a.a. } (t,x) \in Q_T.$$

Proof. In view of (4.2), (5.7), and (5.8), we have to control only the integral

$$\int_0^\tau \int_{\Omega} \left\langle \mathcal{V}_{t,x}; [\vartheta^2]_{\rm res} \right\rangle \ \mathrm{d}x.$$

It follows from (5.8) that

$$\vartheta^{c_v} \stackrel{<}{\sim} \varrho$$
; whence $\vartheta^{c_v+1} \stackrel{<}{\sim} \varrho\vartheta = \frac{1}{c_v} \varrho e(\varrho, \vartheta)$.

Thus the desired conclusion follows from (4.2).

5.2 Unconditional results

Let the molecular pressure p_M of the gas and the associate internal energy e_M be interrelated through the monoatomic gas state equation,

$$p_M(\varrho, \vartheta) = \frac{2}{3} \varrho e_M(\varrho, \vartheta).$$

It follows from Gibbs' relation (1.1) that

$$p_M(\varrho,\vartheta) = \vartheta^{5/2} P\left(\frac{\varrho}{\vartheta^{3/2}}\right) \tag{5.9}$$

for some function P, see [14, Chapter 2]. Motivated by [14, Chapter 3], we further assume that $P \in C^1[0,\infty) \cap C^5(0,\infty)$,

$$P(0) = 0, P'(q) > 0 \text{ for all } q \ge 0,$$
 (5.10)

$$0 < \frac{\frac{5}{3}P(q) - P'(q)q}{q} < c \text{ for all } q > 0, \lim_{q \to \infty} \frac{P(q)}{q^{5/3}} = \overline{p} > 0.$$
 (5.11)

Note that these requirements basically follow from the thermodynamic stability hypothesis (4.1). Moreover, we set

$$e_M(\varrho, \vartheta) = \frac{3}{2}\vartheta\left(\frac{\vartheta^{3/2}}{\varrho}\right)P\left(\frac{\varrho}{\vartheta^{3/2}}\right),$$
 (5.12)

and, by virtue of Gibbs' relation (1.1),

$$s_M(\varrho, \vartheta) = S\left(\frac{\varrho}{\vartheta^{3/2}}\right),$$
 (5.13)

for some function S, see [14, Chapter 2], where

$$S'(q) = -\frac{3}{2} \frac{\frac{5}{3}P(q) - P'(q)q}{q^2} < 0.$$
 (5.14)

Finally, we impose the Third law of thermodynamics in the form

$$\lim_{q \to \infty} S(q) = 0. \tag{5.15}$$

Theorem 5.4. Let κ and μ satisfy (5.4) with $\mu_1 > 0$. Let the thermodynamic functions p, e, and s be given as

$$p(\rho,\vartheta) = p_M(\rho,\vartheta) + p_B(\rho,\vartheta), \ e(\rho,\vartheta) = e_M(\rho,\vartheta) + e_B(\rho,\vartheta), \ s(\rho,\vartheta) = s_M(\rho,\vartheta) + s_B(\rho,\vartheta),$$

where p_M , e_M , and s_M satisfy (5.9–5.15), and

$$p_R = a\vartheta^2, \ e_R = a\frac{\vartheta^2}{\rho}, \ s_R = 2a\frac{\vartheta}{\rho}, \ a > 0.$$
 (5.16)

Assume that $[\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}]$ is a classical solution to the Navier–Stokes–Fourier system (1.2–1.10) in $[0, T] \times \Omega$ emanating from the initial data $[\varrho_0, \vartheta_0, \mathbf{u}_0]$ with $\varrho_0, \vartheta_0 > 0$ in $\overline{\Omega}$. Assume further that $\{\mathcal{V}_{t,x}\}_{(t,x)\in Q_T}$ is a (DMV) solution to the same problem in the sense of Definition 2.3 such that

$$\mathcal{V}_{0,x} = \delta_{[\varrho_0,\vartheta_0,\mathbf{u}_0]} \text{ for a.a. } x \in \Omega.$$

Then

$$\mathcal{V}_{t,x} = \delta_{[\tilde{\varrho}(t,x),\tilde{\vartheta}(t,x),\tilde{\mathbf{u}}(t,x),\mathbb{D}[\tilde{\mathbf{u}}](t,x),\nabla_x\tilde{\vartheta}(t,x)]} \text{ for a.a. } (t,x) \in Q_T.$$

Remark 5.5. The quantity p_R corresponds to the "radiation" component of the pressure. Note that the standard theory of radiative fluids postulates $p_R = a\vartheta^4$ rather than (5.16), see Oxenius [27] or [14, Chapter 2]. The present setting is in the spirit of models of neutron stars studied by Lattimer et al. [22].

Proof. A short inspection of (5.7) reveals that the only term we have to handle is

$$\int_0^\tau \int_\Omega \langle \mathcal{V}_{t,x}; \left[\varrho |s_M(\varrho,\vartheta)|^2 + \varrho s_R |\mathbf{u}| \right]_{\text{res}} \rangle \, \mathrm{d}x.$$

First, it follows from hypothesis (5.16) that

$$\rho s_R |\mathbf{u}| \lesssim \rho s_R |\mathbf{u} - \tilde{\mathbf{u}}| + \rho s_R \lesssim \vartheta |\mathbf{u} - \tilde{\mathbf{u}}| + \rho s_R$$

and

$$\vartheta |\mathbf{u} - \tilde{\mathbf{u}}| \le c(\delta)\vartheta^2 + \delta |\mathbf{u} - \tilde{\mathbf{u}}|^2 \stackrel{<}{\sim} c(\delta)\varrho e_R + \delta |\mathbf{u} - \tilde{\mathbf{u}}|^2.$$

Second, it was shown in [4, Section 3.5.1] that the inequality

$$\varrho |s_M(\varrho,\vartheta)|^2 \stackrel{<}{\sim} 1 + \varrho + \varrho e_M(\varrho,\vartheta)$$

follows from the hypotheses (5.9–5.15).

We complete the proof by applying (2.12) and (4.2).

6 Conditional regularity

Combining the regularity criterion of Sun, Wang, and Zhang [28] with the (DMV)–strong uniqueness result stated in Theorem 5.1 we deduce the following result.

Theorem 6.1. Let $\kappa > 0$, $\mu > 0$, and $\lambda \geq 0$ be constant. Let the thermodynamic functions p, e, and s comply with the perfect gas constitutive relations

$$p(\varrho, \vartheta) = \varrho \vartheta, \ e(\varrho, \vartheta) = c_v \vartheta, \ s(\varrho, \vartheta) = \log \left(\frac{\vartheta^{c_v}}{\varrho}\right), \ c_v > 1.$$

Assume that $\{V_{t,x}\}_{(t,x)\in Q_T}$ is a (DMV) solution of the Navier-Stokes-Fourier system (1.2-1.10) in the sense of Definition 2.3 such that

$$\mathcal{V}_{t,x}\left\{0 < \underline{\varrho} \le \varrho \le \overline{\varrho}, \ \vartheta \le \overline{\vartheta}, \ |\mathbf{u}| \le \overline{u}\right\} = 1 \ \text{for a.a.} \ (t,x) \in Q_T$$
 (6.1)

for some constants $\underline{\varrho}$, $\overline{\varrho}$, $\overline{\vartheta}$, and $\overline{\mathbf{u}}$. Assume further that $\Omega \subset R^3$ is a bounded domain of class $C^{2+\nu}$, and

$$\mathcal{V}_{0,x} = \delta_{\rho_0(x),\vartheta_0(x),\mathbf{u}_0(x)}$$
 for a.a. $x \in \Omega$,

where $[\rho_0, \vartheta_0, \mathbf{u}_0]$ belong to the regularity class

$$\varrho_0, \vartheta_0 \in W^{3,2}(\Omega), \ \varrho, \ \vartheta > 0 \ in \ \overline{\Omega}, \ \mathbf{u}_0 \in W_0^{3,2}(\Omega; R^3)$$

 $\mathbf{u}_0|_{\partial\Omega} = 0, \ \nabla_x \vartheta_0 \cdot \mathbf{n}|_{\partial\Omega} = 0, \ \nabla_x p(\varrho_0, \vartheta_0) = \mathrm{div}_x \mathbb{S}(\nabla_x \mathbf{u}_0).$

Then

$$\mathcal{V}_{t,x} = \delta_{\left[\tilde{\varrho}(t,x),\tilde{\vartheta}(t,x),\tilde{\mathbf{u}}(t,x),\mathbb{D}\left[\tilde{\mathbf{u}}\right](t,x),\nabla_{x}\tilde{\vartheta}(t,x)\right]} \text{ for a.a. } (t,x) \in Q_{T},$$

where $[\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}]$ is a classical solution to the Navier–Stokes–Fourier system with the initial data $[\varrho_0, \vartheta_0, \mathbf{u}_0]$.

Proof. By virtue of the standard local existence results of Valli, and Valli and Zajaczkowski [29], [30], and [31] (among others), the Navier–Stokes–Fourier system (1.2–1.7) admits a local classical solution $[\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}]$ on a maximal existence interval $[0, T_{\text{max}}), T_{\text{max}} > 0$. It follows from hypothesis (6.1) and Theorem 5.1, that the (DMV) solution coincides with $[\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}]$ at least on any compact subinterval of $[0, T_{\text{max}})$.

On the other hand, it follows from hypothesis (6.1) and the fact that the strong and (DMV) solutions coincide that the classical solution $[\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}]$ complies with the regularity criterion established by Sun, Wang, and Zhang [28]. Consequently, the classical solution $[\tilde{\varrho}, \tilde{\vartheta}, \tilde{\mathbf{u}}]$ can be extended up to the time T, and, by Theorem 5.1 coincides with the (DMV) solution on Q_T .

7 Concluding remarks

We discuss briefly the case of more complex constitutive equations considered in the context of weak (distributional) solutions in [15]. The thermodynamic functions have the same structure as in Theorem 5.4:

$$p(\varrho, \vartheta) = p_M(\varrho, \vartheta) + p_R(\varrho, \vartheta), \ e(\varrho, \vartheta) = e_M(\varrho, \vartheta) + e_R(\varrho, \vartheta),$$

where the molecular components p_M , e_M , (and s_M) obey (5.9–5.15) while the radiation terms satisfy

 $p_R = a \frac{1}{3} \vartheta^4, \ e_R = a \frac{\vartheta^4}{\rho}, \ a > 0.$

The transport coefficients $\mu = \mu(\vartheta)$, $\eta = \eta(\vartheta)$ and $\kappa = \kappa(\vartheta)$ are continuously differentiable functions of the temperature ϑ ,

$$|\mu'(\vartheta)| \stackrel{<}{\sim} c, \ (1+\vartheta^{\alpha}) \stackrel{<}{\sim} \mu(\vartheta) \leq (1+\vartheta^{\alpha}), \ 0 \leq \eta(\vartheta) \stackrel{<}{\sim} (1+\vartheta^{\alpha}), \ \alpha \in \left(\frac{2}{5};1\right],$$

$$(1+\vartheta^3) \lesssim \kappa(\vartheta) \lesssim (1+\vartheta^3).$$

The above hypotheses guarantee the *existence* of a weak (distributional) solution to the Navier–Stokes–Fourier system, see [14, Chapter 3]. Moreover, as shown in [11], [15], the weak–strong uniqueness principle holds in this class of weak solutions. The proof requires validity of certain Sobolev embedding relations that can be verified for the weak solutions but that might be violated by the (DMV) solutions specified in Definition 2.3. To retain these properties and, consequently, to extend the results of [15] to the class of (DMV) solutions, the alternative definition proposed in Remark 2.4 must be adopted. We leave the details to the interested reader.

References

- [1] Y. Brenier, C. De Lellis, and L. Székelyhidi, Jr. Weak-strong uniqueness for measure-valued solutions. *Comm. Math. Phys.*, **305**(2):351–361, 2011.
- [2] J. Březina and E. Feireisl. Maximal dissipation principle for the complete Euler system. 2017. arxiv preprint 1712.04761.
- [3] J. Březina and E. Feireisl. Measure-valued solutions to the complete Euler system. 2017. arxiv preprint No. 1702.04878, to appear in J. Math. Soc. Japan.
- [4] J. Březina and E. Feireisl. Measure-valued solutions to the complete Euler system revisited. 2017. arxiv preprint No. 1710.10751.

- [5] C. Christoforou, M. Galanopoulou, and A.E. Tzavaras. A symmetrizable extension of polyconvex thermoelasticity and applications to zero-viscosity limits and weak-strong uniqueness. 2017. arxiv preprint No.1711.01582.
- [6] C.M. Dafermos. The second law of thermodynamics and stability. *Arch. Rational Mech. Anal.*, **70**:167–179, 1979.
- [7] S. Demoulini, D. M. A. Stuart, and A. E. Tzavaras. Weak-strong uniqueness of dissipative measure-valued solutions for polyconvex elastodynamics. *Arch. Ration. Mech. Anal.*, **205**(3):927–961, 2012.
- [8] R. J. DiPerna and A. J. Majda. Oscillations and concentrations in weak solutions of the incompressible fluid equations. *Comm. Math. Phys.*, **108**(4):667–689, 1987.
- [9] R.J. DiPerna. Measure-valued solutions to conservation laws. Arch. Rat. Mech. Anal., 88:223–270, 1985.
- [10] J.L. Ericksen. *Introduction to the thermodynamics of solids, revised ed.* Applied Mathematical Sciences, vol. 131, Springer-Verlag, New York, 1998.
- [11] E. Feireisl. Relative entropies in thermodynamics of complete fluid systems. *Discrete Contin. Dyn. Syst.*, **32**(9):3059–3080, 2012.
- [12] E. Feireisl, P. Gwiazda, A. Świerczewska-Gwiazda, and E. Wiedemann. Dissipative measure-valued solutions to the compressible Navier–Stokes system. *Calc. Var. Partial Differential Equations*, 55(6):55:141, 2016.
- [13] E. Feireisl and M. Lukáčová-Medvidová. Convergence of a mixed finite element finite volume scheme for the isentropic Navier–Stokes system via dissipative measure–valued solutions. 2017. Foundation Comput. Math.
- [14] E. Feireisl and A. Novotný. Singular limits in thermodynamics of viscous fluids. Birkhäuser-Verlag, Basel, 2009.
- [15] E. Feireisl and A. Novotný. Weak-strong uniqueness property for the full Navier-Stokes-Fourier system. *Arch. Rational Mech. Anal.*, **204**:683–706, 2012.
- [16] U. K. Fjordholm, R. Käppeli, S. Mishra, and E. Tadmor. Construction of approximate entropy measure valued solutions for hyperbolic systems of conservation laws. Foundations Comp. Math., pages 1–65, 2015.
- [17] U. S. Fjordholm, S. Mishra, and E. Tadmor. Arbitrarily high-order accurate entropy stable essentially nonoscillatory schemes for systems of conservation laws. *SIAM J. Numer. Anal.*, **50**(2):544–573, 2012.

- [18] U. S. Fjordholm, S. Mishra, and E. Tadmor. On the computation of measure-valued solutions. *Acta Numer.*, **25**:567–679, 2016.
- [19] P. Gwiazda, A. Świerczewska-Gwiazda, and E. Wiedemann. Weak-strong uniqueness for measure-valued solutions of some compressible fluid models. *Nonlinearity*, 28(11):3873–3890, 2015.
- [20] D. Kröner and W. M. Zajaczkowski. Measure-valued solutions of the Euler equations for ideal compressible polytropic fluids. Math. Methods Appl. Sci., 19(3):235–252, 1996.
- [21] C. Lattanzio and A. E. Tzavaras. Relative entropy in diffusive relaxation. SIAM J. Math. Anal., 45(3):1563–1584, 2013.
- [22] J.M. Lattimer, K.A. Van Riper, M. Prakash, and M. Prakash. Rapid cooling and the structure of neutron stars. Astrophysics J., 425:802–813, 1994.
- [23] J. Málek, J. Nečas, M. Rokyta, and M. Růžička. Weak and measure-valued solutions to evolutionary PDE's. Chapman and Hall, London, 1996.
- [24] Š. Matušů-Nečasová and A. Novotný. Measure-valued solution for nonnewtonian compressible isothermal monopolar fluid. *Acta Applicandae Mathematicae*, **37**:109–128, 1994.
- [25] P. B. Mucha and J. Peszek. The Cucker-Smale Equation: Singular Communication Weight, Measure-Valued Solutions and Weak-Atomic Uniqueness. Arch. Ration. Mech. Anal., 227(1):273–308, 2018.
- [26] J. Neustupa. Measure-valued solutions of the Euler and Navier-Stokes equations for compressible barotropic fluids. Math. Nachr., 163:217–227, 1993.
- [27] J. Oxenius. Kinetic theory of particles and photons. Springer-Verlag, Berlin, 1986.
- [28] Y. Sun, C. Wang, and Z. Zhang. A Beale-Kato-Majda criterion for three dimensional compressible viscous heat-conductive flows. *Arch. Ration. Mech. Anal.*, **201**(2):727–742, 2011.
- [29] A. Valli. A correction to the paper: "An existence theorem for compressible viscous fluids"
 [Ann. Mat. Pura Appl. (4) 130 (1982), 197–213; MR 83h:35112]. Ann. Mat. Pura Appl. (4), 132:399–400 (1983), 1982.
- [30] A. Valli. An existence theorem for compressible viscous fluids. Ann. Mat. Pura Appl. (4), 130:197–213, 1982.
- [31] A. Valli and M. Zajaczkowski. Navier-Stokes equations for compressible fluids: Global existence and qualitative properties of the solutions in the general case. Commun. Math. Phys., 103:259–296, 1986.