

# The convex hull of a quadratic constraint over a polytope

Asteriode Santana<sup>\*1</sup> and Santanu S. Dey<sup>†1</sup>

<sup>1</sup>School of Industrial and Systems Engineering, Georgia Institute of Technology

December 27, 2018

## Abstract

A quadratically constrained quadratic program (QCQP) is an optimization problem in which the objective function is a quadratic function and the feasible region is defined by quadratic constraints. Solving non-convex QCQP to global optimality is a well-known NP-hard problem and a traditional approach is to use convex relaxations and branch-and-bound algorithms. This paper makes a contribution in this direction by showing that the exact convex hull of a general quadratic equation intersected with any bounded polyhedron is second-order cone representable. We present a simple constructive proof of this result.

## 1 Introduction

A quadratically constrained quadratic program (QCQP) is an optimization problem in which the objective function is a quadratic function and the feasible region is defined by quadratic constraints. A variety of complex systems can be cast as an instance of a QCQP. Combinatorial problems like MAXCUT [24], engineering problems such as signal processing [23, 30], chemical process [28, 40, 4, 19, 26, 55] and power engineering problems such as the optimal power flow [11, 34, 15, 31] are just a few examples.

Solving non-convex QCQP to global optimality is a well-know NP-hard problem and a traditional approach is to use spacial branch-and-bound tree based algorithm. The computational success of any branch-and-bound tree based algorithm depends on the convexification scheme used at each node of the tree. Not surprisingly, there has been a lot of research on deriving strong convex relaxations for general-purpose QCQPs. The most common relaxations found in the literature are based on Linear programming (LP), second order cone programing (SOCP) or semi-definite programming (SDP). Reformulation-linearization technique (RLT) [48, 50] is a LP-based hierarchy, Lasserre hierarchy or the sum-of-square hierarchy [33] is a SDP-based hierarchy which exactly solves QCQPs under some minor technical conditions and, recently, new LP and SOCP-based alternatives to sum of squares optimization have also been proposed [2].

---

<sup>\*</sup>asteriode.santana@gatech.edu

<sup>†</sup>santanu.dey@isye.gatech.edu

While SDP relaxations are known to be strong, they don't always scale very well computationally. SOCP relaxations tend to be more computationally attractive, although they are often derived by further relaxing SDP relaxations [14].

Another direction of research focuses on convexification of functions, with the McCormick relaxation [37] being perhaps the most classic example. In this case, a constraint of the form  $f(x) = b$  is replaced with  $\check{f}(x) \leq b$  and  $\hat{f}(x) \geq b$ , where  $\check{f}$  is a convex lower approximation and  $\hat{f}$  is a concave upper approximation of  $f$ . While there have been a lot of works in function convexification (see for instance [3, 49, 5, 46, 35, 10, 38, 6, 8, 7, 41, 18, 47, 45, 39, 55, 56, 36, 12, 16, 1, 27, 51]) it is well-known that it does not necessarily yield the convex hull of the set  $\{x \mid f(x) = b\}$ . To the best of our knowledge, there have been much less work on explicit convexification of sets: [54, 42, 43, 53, 25, 32, 44, 17, 34, 13].

A related question when studying convex relaxations is that of representability of the exact convex hull of the feasible set: Is it LP, SOCP or SDP representable? In [20], we prove that the convex hull of the so-called bipartite bilinear constraint (which is a special case of a quadratic constraint) intersected with a box constraint is SOCP representable (SOCr). The proof yields a procedure to compute this convex hull exactly. Encouraging computational results are also reported in [20] in terms of obtaining dual bounds using this construction, which significantly outperform SDP and McCormick relaxations and also bounds produced by commercial solvers.

## 2 Our result

For an integer  $t \geq 1$ , we use  $[t]$  to describe the set  $\{1, \dots, t\}$ . For a set  $G \subseteq \mathbb{R}^n$ , we use  $\text{conv}(G)$ ,  $\text{extr}(G)$  to denote the convex hull of  $G$  and the set of extreme points of  $G$  respectively.

In this paper, we generalize one of the main results in [20]. Specifically, we show that the convex hull of a *general* quadratic equation intersected with *any* bounded polyhedron is SOCr. Moreover the proof is constructive, therefore adding to the literature on explicit convexification in the context of QCQPs. The formal result is as following:

**Theorem 1.** *Let*

$$S := \{x \in \mathbb{R}^n \mid x^\top Q x + \alpha^\top x = g, x \in P\}, \quad (1)$$


*where  $Q \in \mathbb{R}^{n \times n}$  is a symmetric matrix,  $\alpha \in \mathbb{R}^n$ ,  $g \in \mathbb{R}$  and  $P := \{x \mid Ax \leq b\}$  is a polytope. Then  $\text{conv}(S)$  is SOCr.*

Notice that we make no assumption regarding the structure or coefficients of the quadratic equation defining  $S$ . We require  $P$  to be a bounded polyhedron, which is not very restrictive given that in global optimization the variables are often assumed to be bounded to use branch-and-bound algorithms.


The result presented in Theorem 1 is somewhat unexpected since the sum-of-squares approach would build a sequence of SDP relaxations for (1) in order to optimize (exactly) a linear function over  $S$ , while even the SDP cone of three-by-three dimensional matrices is not SOCr [22]. Note that optimizing a linear function over  $S$  is NP-hard, therefore, while the convex hull is SOCr, the construction involves the introduction of an exponential number of variables.

Surprisingly, the proof of Theorem 1 is fairly straightforward and it introduces a technique (new, to the best of our knowledge) to compute convex hull of certain surfaces over a compact set. In the case of Theorem 1, the key observation is that the surface defined by the quadratic equation either:

1. is defined as the union of two convex surfaces (see Figure 1); or
2. it has the property that, through every point of the surface, there exists a *straight* line that is *entirely* contained in the surface (see Figure 2).



**Figure 1:** Two-sheets hyperboloid. The surface is the union of two convex peices.



**Figure 2:** One-sheet hyperboloid. Through every point of the surface, there exists a straight line that is entirely contained in the surface.

In Case 1, we can easily obtain that the convex hull of  $S$  is SOCr as we show in Section 3.3. In Case 2, no point in the interior of the polytope can be an extreme point of  $S$ . Observing that the convex hull of a compact set is also the convex hull of its extreme points, we intersect the surface with each facet of the polytope which will contain all the extreme points of  $S$ . Now, each such intersection leads to new sets with the same form as  $S$  but in one dimension lower. The argument then goes by recursion. The details of the proof are presented in Section 3.

After we had proved Theorem 1, we learned that the property described in Case 2 is known as “ruled surfaces” and it has been extensively studied from both algebraic and geometric perspectives [21]. To the best of our knowledge, however, no one from the global optimization community has ever exploited such results for convexification.

## 3 Proof of Theorem 1

### 3.1 Convex hulls via disjunctions

In this section, we describe a simple procedure to obtain the convex hull of a compact set  $S$  using a disjunctive argument. We use this procedure to prove Theorem 1 in Section 3.3. Let  $S$  be a compact set and let  $\text{extr}(S)$  be the set of extreme points of  $S$ . First, we partition the

extreme points of  $S$ . Specifically, suppose there exist  $B^1, \dots, B^k \subseteq S$  such that:

$$S \supseteq \bigcup_{i=1}^k B^i \supseteq \text{extr}(S). \quad (2)$$

We observe that (2) implies that

$$\text{conv}(S) \supseteq \text{conv}\left(\bigcup_{i=1}^k B^i\right) \supseteq \text{conv}(\text{extr}(S)) = \text{conv}(S), \quad (3)$$

where the last equality holds due to  $S$  being compact. Finally, we obtain that

$$\text{conv}(S) = \text{conv}\left(\bigcup_{i=1}^k B^i\right) = \text{conv}\left(\bigcup_{i=1}^k \text{conv}(B^i)\right). \quad (4)$$

**Observation 1.** *If  $\text{conv}(B^i)$  is SOCr for all  $i \in [k]$ , then the set*

$$\text{conv}\left(\bigcup_{i=1}^k \text{conv}(B^i)\right),$$

*is SOCr [9]. Thus, we obtain from (4) that  $\text{conv}(S)$  is SOCr. In addition, we obtain a constructive procedure to compute  $\text{conv}(S)$ .*

### 3.2 Reduction

In this section, we discuss how we can apply some transformations to the set  $S$  defined in (1) so as to re-write it in a “canonical” form where all the quadratic terms are squared terms. This will allow us to easily classify  $S$  into Case 1 and 2 as discussed in Section 2. We start with the following observation.

**Observation 2.** *Let  $S \subseteq \mathbb{R}^n$  and let  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be an affine map. Then*

$$\text{conv}(F(S)) = F(\text{conv}(S)),$$

*where  $F(S) := \{Fx \mid x \in S\}$ . Furthermore if  $\text{conv}(S)$  is SOCr, then  $\text{conv}(F(S))$  is also SOCr.*

Let  $S$  be the set defined in (1). Suppose, without loss of generality, that  $Q$  is a symmetric matrix. By the spectral theorem  $Q = V^\top \Sigma V$ , where  $\Sigma$  is a diagonal matrix and the columns of  $V$  are a set of orthogonal vectors. Letting  $w = Vx$ , we have that

$$S := V^{-1} \left( \{w \mid w^\top \Sigma w + \alpha^\top V^{-1} w = d, \ w \in \tilde{P}\} \right),$$

where  $\tilde{P} := \{w \mid AV^{-1}w \leq b\}$ .

Therefore, by Observation 2, it is sufficient to study the convex hull of a set of the form:

$$S := \left\{ (x, y, z) \in \mathbb{R}^n \mid \sum_{i=1}^{n_q} a_i x_i^2 + \sum_{i=1}^{n_q} \alpha_i x_i + \sum_{j=1}^{n_l} \beta_j y_j = g, (x, y, z) \in P \right\},$$

where  $z \in \mathbb{R}^{n_o}$  does not appear in the quadratic constraints,  $n_q + n_l + n_o = n$ ,  $a_i \neq 0$  for  $i \in [n_q]$  (i.e., the rank of  $Q$  is  $n_q$ ) and  $\beta_j \neq 0$  for  $j \in [n_l]$ . By completing squares, we may further write  $S$  as:

$$S := \{(x, y, z) \in \mathbb{R}^n \mid \sum_{i=1}^{n_q} \sigma(a_i) \left( \sqrt{|a_i|} x_i + \sigma(a_i) \frac{\alpha_i}{2\sqrt{|a_i|}} \right)^2 + \sum_{i=1}^{n_l} \beta_i y_i = g + \sum_{i=1}^{n_q} \frac{\alpha_i^2}{4a_i}, (x, y, z) \in P\},$$

where  $\sigma(a)$  denotes the sign of  $a$ . Now, since  $u_i = \left( \sqrt{|a_i|} x_i + \sigma(a_i) \frac{\alpha_i}{2\sqrt{|a_i|}} \right)$  for  $i \in [n_q]$  and  $v_i = \beta_i y_i$  for  $i \in [n_l]$  define linear bijections, it follows from Observation 2 that it is sufficient to study the convex hull of the following set:

$$S := \{(w, x, y, z) \in \mathbb{R}^{n_{q+}} \times \mathbb{R}^{n_{q-}} \times \mathbb{R}^{n_l} \times \mathbb{R}^{n_o} \mid \sum_{i=1}^{n_{q+}} w_i^2 - \sum_{j=1}^{n_{q-}} x_j^2 + \sum_{k=1}^{n_l} y_k = g, (w, x, y, z) \in P\}, \quad (5)$$

where we may further assume that  $g \geq 0$ , since otherwise we may multiply the equation by  $-1$  and apply suitable affine transformations to bring it back to the form of (5).

### 3.3 Recursive argument to prove Theorem 1

We begin by stating a variant of Observation 2 that we will use twice along the proof.

**Lemma 1.** *Let  $G = \{(x, w) \in \mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \mid x \in G_0, w = C^\top x + h\}$ , where  $G_0 \subseteq \mathbb{R}^{n_1}$  is bounded, and  $C^\top x + h$  is an affine function of  $x$ . Then,*

$$\text{conv}(G) = \{(x, w) \in \mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \mid x \in \text{conv}(G_0), w = C^\top x + h\}.$$

*Proof.* See Lemma 4 in [20]. □

#### 3.3.1 Dealing with low dimensional polytope

Let  $S$  and  $P$  be defined as in (1). Next, we show that we may assume without loss of generality that  $P$  is full dimension. In fact, if  $P$  is not full dimensional, then  $P$  is contained in a non-trivial affine subspace defined by a system of linear equations  $Mx = f$ . Without loss of generality, we may assume that  $M$  has full row-rank  $k$ ,  $1 \leq k < n$ . Let  $M = [M_B \ M_N]$  where  $M_B$  is invertible. Then, we may write this system as  $x_B = -M_B^{-1} M_N x_N + M_B^{-1} f$ , where  $x_B \in \mathbb{R}^k$ ,  $x_N \in \mathbb{R}^{n-k}$  and, for simplicity, we assume that  $x_B$  (resp.  $x_N$ ) correspond to the first  $k$  (resp. last  $n - k$ ) components of  $x$ . By defining  $C = -M_B^{-1} M_N$  and  $h = M_B^{-1} f$  to simplify notation, we obtain

$$x_B = Cx_N + h. \quad (6)$$

By partitioning  $Q$  in sub-matrices of appropriate sizes, we may explicitly write the quadratic equation defining  $S$  in terms of  $x_B$  and  $x_N$  as follows:

$$\begin{bmatrix} x_B^\top & x_N^\top \end{bmatrix} \begin{bmatrix} Q_{BB} & Q_{BN} \\ Q_{NB} & Q_{NN} \end{bmatrix} \begin{bmatrix} x_B \\ x_N \end{bmatrix} + \alpha^\top \begin{bmatrix} x_B \\ x_N \end{bmatrix} = g. \quad (7)$$

Using (6), we replace  $x_B$  in (7) to obtain

$$x_N^\top \tilde{Q} x_N + \tilde{\alpha}^\top x_N = \tilde{g},$$

where

$$\begin{aligned} \tilde{Q} &= C^\top Q_{BB} C + C^\top Q_{BN} + Q_{NB} C + Q_{NN}, \\ \tilde{\alpha} &= 2C^\top Q_{BB} h + Q_{BN}^\top h + Q_{NB} h + C^\top \alpha_B + \alpha_N, \\ \tilde{g} &= g - h^\top Q_{BB} h - \alpha_B^\top h. \end{aligned}$$

Therefore, we may write  $S$  as

$$S := \{(x_B, x_N) \in \mathbb{R}^n \mid x_N^\top \tilde{Q} x_N + \tilde{\alpha}^\top x_N = \tilde{g}, x_N \in \tilde{P}, x_B = C x_N + h\}, \quad (8)$$

where  $\tilde{P}$  is now a full dimensional polytope. Therefore, by Lemma 1, we may assume from now on that  $P$  is full dimensional.

### 3.3.2 Case 2: Sufficient conditions for points to not be extreme

Consider the set  $S$  as defined in (5).

**Lemma 2.** *Suppose  $n_o \geq 1$ . If  $(a, b, c, d) \in S \cap (\mathbb{R}^{n_{q+}} \times \mathbb{R}^{n_{q-}} \times \mathbb{R}^{n_l} \times \mathbb{R}^{n_o})$  where  $(a, b, c, d) \in \text{int}(P)$ , then  $(a, b, c, d)$  is not an extreme point of  $S$ .*

*Proof.* Since  $(a, b, c, d) \in \text{int}(P)$ , there exists a vector  $\delta \in \mathbb{R}^{n_o} \setminus \{0\}$  such that  $(a, b, c, d + \delta), (a, b, c, d - \delta) \in P$ . Clearly these points are in  $S$  as well and, therefore,  $(a, b, c, d)$  is not an extreme point of  $S$   $\square$

**Lemma 3.** *Suppose  $n_o = 0$  and  $n_l \geq 2$ . If  $(a, b, c) \in S \cap (\mathbb{R}^{n_{q+}} \times \mathbb{R}^{n_{q-}} \times \mathbb{R}^{n_l})$  where  $(a, b, c) \in \text{int}(P)$ , then  $(a, b, c)$  is not an extreme point of  $S$ .*

*Proof.* Since  $n_l \geq 2$ ,  $(a, b, c_1 \pm \lambda, c_2 \mp \lambda, \dots, c_{n_3})$  are feasible for sufficiently small positive values of  $\lambda$ . Therefore,  $(a, b, c)$  is not an extreme point.  $\square$

**Lemma 4.** *Suppose  $n_o = 0$ ,  $n_{q+}, n_{q-} \geq 1$  and  $n_l = 1$ . If  $(a, b, c) \in S \cap (\mathbb{R}^{n_{q+}} \times \mathbb{R}^{n_{q-}} \times \mathbb{R}^{n_l})$  where  $(a, b, c) \in \text{int}(P)$ , then  $(a, b, c)$  is not an extreme point of  $S$ .*

*Proof.* Since  $n_{q+}, n_{q-} \geq 1$ , and  $n_l = 1$ ,  $(a_1 + \lambda, a_2, \dots, a_{n_{q+}}, b_1 + \lambda, b_2, \dots, b_{n_{q-}}, c + 2\lambda(-a_1 + b_1))$  are feasible for sufficiently small positive and negative values of  $\lambda$ . Therefore,  $(a, b, c)$  is not an extreme point.  $\square$

**Lemma 5.** Suppose  $n_o = 0$ ,  $n_{q+} \geq 2$ ,  $n_{q-} \geq 1$  and  $n_l = 0$ . If  $(a, b) \in S \cap (\mathbb{R}^{n_{q+}} \times \mathbb{R}^{n_{q-}})$  where  $(a, b) \in \text{int}(P)$ , then  $(a, b)$  is not an extreme point of  $S$ .

*Proof.* We show that there exists a straight line through  $(a, b)$  that is entirely contained in the surface defined by the quadratic equation. More specifically, we prove that there exists a vector  $(u, v) \in (\mathbb{R}^{n_{q+}} \times \mathbb{R}^{n_{q-}}) \setminus \{0\}$  such that the line  $\{(a, b) + \lambda(u, v) \mid \lambda \in \mathbb{R}\}$  satisfies the quadratic equation and therefore,  $(a, b)$  being in the interior of  $P$  cannot be an extreme point of  $S$ . We consider two cases:

1.  $(a, b) \neq \mathbf{0}$ : Then observe that  $a \neq \mathbf{0}$ , since otherwise we would have  $a = \mathbf{0}$  and  $b = \mathbf{0}$ , because  $g \geq 0$ . Observe that

$$\sum_{i=1}^{n_{q+}} a_i^2 = g + \sum_{j=1}^{n_{q-}} b_j^2 \geq b_1^2 \Leftrightarrow \frac{|b_1|}{\|a\|_2} \leq 1. \quad (9)$$

Next, observe that:

$$\begin{aligned} g &= \sum_{i=1}^{n_{q+}} (a_i + \lambda u_i)^2 - \sum_{i=1}^{n_{q-}} (b_i + \lambda v_i)^2 \quad \forall \lambda \in \mathbb{R} \\ \Leftrightarrow g &= \left( \sum_{i=1}^{n_{q+}} a_i^2 - \sum_{i=1}^{n_{q-}} b_i^2 \right) + \lambda^2 \left( \sum_{i=1}^{n_{q+}} u_i^2 - \sum_{i=1}^{n_{q-}} v_i^2 \right) + 2\lambda \left( \sum_{i=1}^{n_{q+}} a_i u_i - \sum_{i=1}^{n_{q-}} b_i v_i \right) \quad \forall \lambda \in \mathbb{R} \\ \Leftrightarrow 0 &= \lambda \left( \sum_{i=1}^{n_{q+}} u_i^2 - \sum_{i=1}^{n_{q-}} v_i^2 \right) + 2 \left( \sum_{i=1}^{n_{q+}} a_i u_i - \sum_{i=1}^{n_{q-}} b_i v_i \right) \quad \forall \lambda \in \mathbb{R} \\ \Leftrightarrow &\sum_{i=1}^{n_{q+}} u_i^2 - \sum_{i=1}^{n_{q-}} v_i^2 = 0, \quad \sum_{i=1}^{n_{q+}} a_i u_i - \sum_{i=1}^{n_{q-}} b_i v_i = 0. \end{aligned} \quad (10)$$

Suppose we set  $v_1 = 1$  and  $v_j = 0$  for all  $j \in \{2, \dots, n_{q-}\}$ . Then satisfying (10) is equivalent to finding real values of  $u$  satisfying:

$$\sum_{i=1}^{n_{q+}} u_i^2 = 1, \quad \sum_{i=1}^{n_{q+}} a_i u_i = b_1.$$

This is the intersection of a circle of radius 1 in dimension two or higher (since  $n_{q+} \geq 2$  in this case) and a hyperplane whose distance from the origin is  $\frac{|b_1|}{\|a\|_2}$ . Since, by (9), we have that this distance is at most 1, the hyperplane intersects the circle and therefore we know that a real solution exists.

2.  $(a, b) = \mathbf{0}$ : In this case, observe that  $g = 0$  and then  $\mathbf{0}$  is a convex combination of

$$\left( \underbrace{\pm\lambda, 0, \dots, 0}_{\text{first } n_{q+} \text{ components}}, \underbrace{\pm\lambda, 0, \dots, 0}_{\text{second } n_{q-} \text{ components}} \right)$$

for sufficiently small  $\lambda > 0$ .

□

### 3.3.3 Case 1: Sufficient conditions for convex hull to be SOCr

In this section, we repeatedly use the following result from [52].

**Theorem 2.** *Let  $G \subseteq \mathbb{R}^n$  be a convex set and let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a continuous function. Then*

$$\text{conv}(\{G \cap \{x \mid f(x) = 0\}\}) = \text{conv}(\{G \cap \{x \mid f(x) \leq 0\}\}) \cap \text{conv}(\{G \cap \{x \mid f(x) \geq 0\}\}).$$

For the two lemmas that follows, consider the notation of  $S$  defined in (5).

**Lemma 6.** *Suppose  $n_o = 0$ ,  $n_l \leq 1$ . If  $n_{q+} = 0$  or  $n_{q-} = 0$ , then  $\text{conv}(S)$  is SOCr.*

*Proof.* We consider two cases.

1.  $n_{q-} = 0$ : Let  $(w, y) \in S \cap (\mathbb{R}^{n_{q+}} \times \mathbb{R}^{n_l})$ . Let  $y = y_1$  if  $n_l = 1$  and  $y = 0$  if  $n_l = 0$ . In this case,  $g - y$  is non-negative for all feasible values of  $y$  and we can use the identity  $t = \frac{(t+1)^2 - (t-1)^2}{4}$  to write  $S = S' \cap S''$ , where:

$$\begin{aligned} S' &:= \{(w, y) \in P \mid \|2w_1, \dots, 2w_{n_{q+}}, (g - y - 1)\| \leq (g - y + 1)\}, \\ S'' &:= \{(w, y) \in P \mid \|2w_1, \dots, 2w_{n_{q+}}, (g - y - 1)\| \geq (g - y + 1)\}. \end{aligned}$$

Notice that  $S'$  is a SOCr convex set. Also notice that  $S''$  is a reverse convex set intersected with a polytope and hence  $\text{conv}(S'' \cap P)$  is polyhedral and contained in  $P$  (see [29], Theorem 1). Therefore, by Theorem 2, we have that  $\text{conv}(S) = \text{conv}(S') \cap \text{conv}(S'')$  is SOCr.

2.  $n_{q+} = 0$ : Let  $(x, y) \in S \cap (\mathbb{R}^{n_{q+}} \times \mathbb{R}^{n_l})$ . Let  $y = y_1$  if  $n_l = 1$  and  $y = 0$  if  $n_l = 0$ . In this case,  $g - y$  is non-positive for all feasible values of  $y$  and may write  $S = S' \cap S''$ , where:

$$\begin{aligned} S' &:= \{(x, y) \in P \mid \|2x_1, \dots, 2x_{n_{q-}}, (y - g - 1)\| \leq (y - g + 1)\}, \\ S'' &:= \{(x, y) \in P \mid \|2x_1, \dots, 2x_{n_{q-}}, (y - g - 1)\| \geq (y - g + 1)\}. \end{aligned}$$

Therefore, as in the previous case,  $\text{conv}(S)$  is SOCr.

□

**Lemma 7.** *Suppose  $n_{q+} \leq 1$  and  $n_l = n_o = 0$ . Then  $\text{conv}(S)$  is SOCr.*

*Proof.* If  $n_{q+} = 0$ , then  $S$  is empty set or contains a single point, the origin.

Therefore, consider the case where  $n_{q+} = 1$ , thus  $w = w_1$ . Notice that  $S = S' \cap S''$ , where

$$\begin{aligned} S' &:= \{(w, x) \in \mathbb{R}^1 \times \mathbb{R}^{n_{q-}} \mid w^2 \geq g + \sum_{j=1}^{n_{q-}} x_j^2, (w, x) \in P\}, \\ S'' &:= \{(w, x) \in \mathbb{R}^1 \times \mathbb{R}^{n_{q-}} \mid w^2 \leq g + \sum_{j=1}^{n_{q-}} x_j^2, (w, x) \in P\}. \end{aligned}$$



By Theorem 2,  $\text{conv}(S) = \text{conv}(S') \cap \text{conv}(S'')$ . Next, we show that both  $\text{conv}(S')$  and  $\text{conv}(S'')$  are SOCr. Notice that  $S'$  is the union of the following two SOCr sets:

$$\begin{aligned}
S'_+ &:= \left\{ (w, x) \in \mathbb{R}^1 \times \mathbb{R}^{n_{q-}} \mid w \geq \left( g + \sum_{j=1}^{n_{q-}} x_j^2 \right)^{\frac{1}{2}}, w \geq 0, (w, x) \in P \right\}, \\
&= \text{Proj}_{w,x} \left( \left\{ (w, x, t) \in \mathbb{R}^1 \times \mathbb{R}^{n_{q-}} \times \mathbb{R} \mid w \geq \left( (\sqrt{g}t)^2 + \sum_{j=1}^{n_{q-}} x_j^2 \right)^{\frac{1}{2}}, x \geq 0, t = 1, (w, x) \in P \right\} \right), \\
S'_- &:= \left\{ (w, x) \in \mathbb{R}^1 \times \mathbb{R}^{n_{q-}} \mid -w \geq \left( g + \sum_{j=1}^{n_{q-}} x_j^2 \right)^{\frac{1}{2}}, w \leq 0, (w, x) \in P \right\} \\
&= \text{Proj}_{w,x} \left( \left\{ (w, x, t) \in \mathbb{R}^1 \times \mathbb{R}^{n_{q-}} \times \mathbb{R} \mid -w \geq \left( \sqrt{g}t^2 + \sum_{j=1}^{n_{q-}} x_j^2 \right)^{\frac{1}{2}}, w \leq 0, t = 1, (w, x) \in P \right\} \right).
\end{aligned}$$

Thus,  $\text{conv}(S') = \text{conv}(S'_+ \cup S'_-)$  is SOCr.

Notice that  $S'' = \{(w, x) \in \mathbb{R}^1 \times \mathbb{R}^{n_{q-}} \mid |w| \leq (g + \sum_{j=1}^{n_{q-}} x_j^2)^{\frac{1}{2}}, (w, x) \in P\}$  and is therefore the union of two sets:

$$\begin{aligned}
S''_+ &:= \left\{ (w, x) \in \mathbb{R}^1 \times \mathbb{R}^{n_{q-}} \mid w \leq \left( g + \sum_{j=1}^{n_{q-}} x_j^2 \right)^{\frac{1}{2}}, w \geq 0, (w, x) \in P \right\}, \\
S''_- &:= \left\{ (w, x) \in \mathbb{R}^1 \times \mathbb{R}^{n_{q-}} \mid -w \leq \left( g + \sum_{j=1}^{n_{q-}} x_j^2 \right)^{\frac{1}{2}}, w \leq 0, (w, x) \in P \right\},
\end{aligned}$$

each of them being a reverse convex set intersected with a polyhedron. Therefore,  $\text{conv}(S''_+)$  and  $\text{conv}(S''_-)$  are polyhedral and therefore  $\text{conv}(S'') = \text{conv}(\text{conv}(S''_+) \cup \text{conv}(S''_-))$  is a polyhedral set.  $\square$

### 3.3.4 Proof of Theorem 1

Finally, we bring the pieces together to prove Theorem 1.

*Proof.* (of Theorem 1) Let  $S(n)$  be defined as in (5), where  $n = n_{q+} + n_{q-} + n_l + n_o$  is the dimension of the space in which  $S$  is defined and without loss of generality  $P$  is full-dimensional (Section 3.3.1). The proof goes by induction on  $n$ . Notice that  $S(1)$  is a polytope and hence  $\text{conv}(S(1))$  is SOCr. Suppose  $S(n)$  is SOCr. We show that  $S(n+1)$  is SOCr as well. If  $n_o = 0$ ,  $n_l \leq 1$ , and  $n_{q+} = 0$  or  $n_{q-} = 0$ , then the result follows from Lemma 6. Similarly, if  $n_o = 0$ ,  $n_{q+} \leq 1$  and  $n_l = 0$ , then the result follows from Lemma 7. Otherwise, it follows from

Lemma 2, 3, 4 and 5 that no point in the interior of  $P$  can be an extreme point of  $S(n+1)$ . Let  $N$  be the number of facets of  $P$ , each of which given by one equation of the linear system  $Fx = f$ . Let  $B^i = S(n+1) \cap \{x \in \mathbb{R}^{n+1} \mid F_i x = f_i\}$  be the intersection of  $S(n+1)$  with the  $i$ th facet of  $P$ . By the discussion in Section 3.1, it is enough to show that the convex hull of each  $B^i$  is SOCr. Let  $i \in \{1, \dots, N\}$ . Choose  $j_0$  such that  $F_{ij_0} \neq 0$ . For simplicity, suppose  $j_0 = 1$ . Then, we may write  $B^i = \{x \in \mathbb{R}^{n+1} \mid (x_2, \dots, x_{n+1}) \in B_0^i, x_1 = b_i - \sum_{j=2}^{n+1} F_{ij} x_j\}$ , where  $B_0^i$  is obtained from  $B^i$  by replacing  $x_1 = f_i - \sum_{j=2}^{n+1} F_{ij} x_j$  in all the constraints defining  $S(n+1)$ . Now  $\text{conv}(B_0^i) \subseteq \mathbb{R}^n$  is SOCr by induction hypothesis. Therefore,  $\text{conv}(B^i)$  is SOCr by Lemma 1.  $\square$

## Acknowledgments

Funding: This work was supported by the NSF CMMI [grant number 1562578] and the CNPq-Brazil [grant number 248941/2013-5].

## References

- [1] Warren Adams, Akshay Gupte, and Yibo Xu. Error bounds for monomial convexification in polynomial optimization. *Mathematical Programming*, Mar 2018.
- [2] Amir Ali Ahmadi and Anirudha Majumdar. Dsos and sdsos optimization: Lp and socp-based alternatives to sum of squares optimization. In *Information Sciences and Systems (CISS), 2014 48th Annual Conference on*, pages 1–5. IEEE, 2014.
- [3] Faiz A. Al-Khayyal and James E. Falk. Jointly constrained biconvex programming. *Mathematics of Operations Research*, 8(2):273–286, 1983.
- [4] Mohammed Alfaki and Dag Haugland. Strong formulations for the pooling problem. *Journal of Global Optimization*, 56(3):897–916, 2013.
- [5] Ioannis P. Androulakis, Costas D. Maranas, and Christodoulos A Floudas.  $\alpha$ bb: A global optimization method for general constrained nonconvex problems. *Journal of Global Optimization*, 7(4):337–363, 1995.
- [6] Kurt M. Anstreicher and Samuel Burer. Computable representations for convex hulls of low-dimensional quadratic forms. *Mathematical programming*, 124(1-2):33–43, 2010.
- [7] Xiaowei Bao, Aida Khajavirad, Nikolaos V. Sahinidis, and Mohit Tawarmalani. Global optimization of nonconvex problems with multilinear intermediates. *Mathematical Programming Computation*, 7(1):1–37, 2015.
- [8] Pietro Belotti, Andrew J. Miller, and Mahdi Namazifar. Valid inequalities and convex hulls for multilinear functions. *Electronic Notes in Discrete Mathematics*, 36:805–812, 2010.

- [9] Aharon Ben-Tal and Arkadi Nemirovski. *Lectures on modern convex optimization: analysis, algorithms, and engineering applications*. SIAM, 2001.
- [10] Harold P. Benson. Concave envelopes of monomial functions over rectangles. *Naval Research Logistics (NRL)*, 51(4):467–476, 2004.
- [11] Subhonmesh Bose, Dennice F. Gayme, Kiananthra M. Chandy, and Steven H. Low. Quadratically constrained quadratic programs on acyclic graphs with application to power flow. *IEEE Transactions on Control of Network Systems*, 2(3):278–287, Sept 2015.
- [12] Christoph Buchheim and Claudia D’Ambrosio. Monomial-wise optimal separable under-estimators for mixed-integer polynomial optimization. *Journal of Global Optimization*, 67(4):759–786, 2017.
- [13] Samuel Burer and Fatma Kılınç-Karzan. How to convexify the intersection of a second order cone and a nonconvex quadratic. *Mathematical Programming*, 162(1):393–429, Mar 2017.
- [14] Samuel Burer, Sunyoung Kim, and Masakazu Kojima. Faster, but weaker, relaxations for quadratically constrained quadratic programs. *Computational Optimization and Applications*, 59(1):27–45, Oct 2014.
- [15] Chen Chen, Alper Atamtürk, and Shmuel S. Oren. Bound tightening for the alternating current optimal power flow problem. *IEEE Transactions on Power Systems*, 31(5):3729–3736, Sept 2016.
- [16] Yves Crama and Elisabeth Rodríguez-Heck. A class of valid inequalities for multilinear 0–1 optimization problems. *Discrete Optimization*, 25:28–47, 2017.
- [17] Danial Davarnia, Jean-Philippe P. Richard, and Mohit Tawarmalani. Simultaneous convexification of bilinear functions over polytopes with application to network interdiction. *SIAM Journal on Optimization*, 27(3):1801–1833, 2017.
- [18] Alberto Del Pia and Aida Khajavirad. A polyhedral study of binary polynomial programs. *Mathematics of Operations Research*, 42(2):389–410, 2016.
- [19] Santanu S. Dey and Akshay Gupte. Analysis of milp techniques for the pooling problem. *Operations Research*, 63(2):412–427, 2015.
- [20] Santanu S. Dey, Asteroide Santana, and Yang Wang. New socp relaxation and branching rule for bipartite bilinear programs. *Optimization and Engineering*, Sep 2018.
- [21] William L. Edge. *The Theory of Ruled Surfaces*. Cambridge University Press, 2011.
- [22] Hamza Fawzi. On representing the positive semidefinite cone using the second-order cone. *Mathematical Programming*, pages 1–10, 2018.

- [23] Ahmad Gharanjik, Bhavani Shankar, Mojtaba Soltanalian, and Björn Oftersten. An iterative approach to nonconvex qcqp with applications in signal processing. In *Sensor Array and Multichannel Signal Processing Workshop (SAM), 2016 IEEE*, pages 1–5. IEEE, 2016.
- [24] Michel X. Goemans and David P. Williamson. . 879-approximation algorithms for max cut and max 2sat. In *Proceedings of the twenty-sixth annual ACM symposium on Theory of computing*, pages 422–431. ACM, 1994.
- [25] Akshay Gupte. *Mixed integer bilinear programming with applications to the pooling problem*. PhD thesis, Georgia Institute of Technology, 2011.
- [26] Akshay Gupte, Shabbir Ahmed, Santanu S. Dey, and Myun Seok Cheon. Relaxations and discretizations for the pooling problem. *Journal of Global Optimization*, 67(3):631–669, 2017.
- [27] Akshay Gupte, Thomas Kalinowski, Fabian Rigterink, and Hamish Waterer. Extended formulations for convex hulls of graphs of bilinear functions. *arXiv preprint arXiv:1702.04813*, 2017.
- [28] Co A. Haverly. Studies of the behavior of recursion for the pooling problem. *Acm sigmap bulletin*, (25):19–28, 1978.
- [29] Richard J. Hillestad and Stephen E. Jacobsen. Linear programs with an additional reverse convex constraint. *Applied Mathematics and Optimization*, 6(1):257–269, Mar 1980.
- [30] Arash Khabbazi-basmenj and Sergiy A. Vorobyov. Generalized quadratically constrained quadratic programming for signal processing. In *2014 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, pages 7629–7633, May 2014.
- [31] Burak Kocuk, Santanu S. Dey, and Xu A. Sun. Strong socp relaxations for the optimal power flow problem. *Operations Research*, 64(6):1177–1196, 2016.
- [32] Burak Kocuk, Santanu S. Dey, and Xu A. Sun. Matrix minor reformulation and socp-based spatial branch-and-cut method for the ac optimal power flow problem. *arXiv preprint arXiv:1703.03050*, 2017.
- [33] Jean B. Lasserre. Global optimization with polynomials and the problem of moments. *SIAM Journal on Optimization*, 11(3):796–817, 2001.
- [34] Qifeng Li and Vijay Vittal. Convex hull of the quadratic branch ac power flow equations and its application in radial distribution networks. *IEEE Transactions on Power Systems*, 33(1):839–850, Jan 2018.
- [35] Leo Liberti and Constantinos C. Pantelides. Convex envelopes of monomials of odd degree. *Journal of Global Optimization*, 25(2):157–168, 2003.

- [36] Marco Locatelli. Polyhedral subdivisions and functional forms for the convex envelopes of bilinear, fractional and other bivariate functions over general polytopes. *Journal of Global Optimization*, 66(4):629–668, 2016.
- [37] Garth P. McCormick. Computability of global solutions to factorable nonconvex programs: Part i—convex underestimating problems. *Mathematical programming*, 10(1):147–175, 1976.
- [38] Clifford A. Meyer and Christodoulos A. Floudas. Trilinear monomials with mixed sign domains: Facets of the convex and concave envelopes. *Journal of Global Optimization*, 29(2):125–155, 2004.
- [39] Clifford A. Meyer and Christodoulos A. Floudas. Convex envelopes for edge-concave functions. *Mathematical programming*, 103(2):207–224, 2005.
- [40] Clifford A. Meyer and Christodoulos A. Floudas. Global optimization of a combinatorially complex generalized pooling problem. *AIChE journal*, 52(3):1027–1037, 2006.
- [41] Ruth Misener, James B. Smadbeck, and Christodoulos A. Floudas. Dynamically generated cutting planes for mixed-integer quadratically constrained quadratic programs and their incorporation into glomiqo 2. *Optimization Methods and Software*, 30(1):215–249, 2015.
- [42] Trang T. Nguyen, Jean-Philippe P. Richard, and Mohit Tawarmalani. Deriving the convex hull of a polynomial partitioning set through lifting and projection. Technical report, working paper, 2013.
- [43] Trang T. Nguyen, Mohit Tawarmalani, and Jean-Philippe P. Richard. Convexification techniques for linear complementarity constraints. In *IPCO*, volume 6655, pages 336–348. Springer, 2011.
- [44] Hamidur Rahman and Ashutosh Mahajan. Facets of a mixed-integer bilinear covering set with bounds on variables. *arXiv preprint arXiv:1707.06712*, 2017.
- [45] Anatoliy D. Rikun. A convex envelope formula for multilinear functions. *Journal of Global Optimization*, 10(4):425–437, Jun 1997.
- [46] Hong Seo Ryoo and Nikolaos V. Sahinidis. Analysis of bounds for multilinear functions. *Journal of Global Optimization*, 19(4):403–424, 2001.
- [47] Hanif D. Sherali. Convex envelopes of multilinear functions over a unit hypercube and over special discrete sets. *Acta mathematica vietnamica*, 22(1):245–270, 1997.
- [48] Hanif D. Sherali and Warren P. Adams. *A reformulation-linearization technique for solving discrete and continuous nonconvex problems*, volume 31. Springer Science & Business Media, 2013.

- [49] Hanif D. Sherali and Amine Alameddine. An explicit characterization of the convex envelope of a bivariate bilinear function over special polytopes. *Annals of Operations Research*, 25(1):197–209, 1990.
- [50] Hanif D. Sherali and Amine Alameddine. A new reformulation-linearization technique for bilinear programming problems. *Journal of Global optimization*, 2(4):379–410, 1992.
- [51] Emily Speakman and Jon Lee. Quantifying double mccormick. *Mathematics of Operations Research*, 42(4):1230–1253, 2017.
- [52] Mohit Tawarmalani and Jean-Philippe P. Richard. Decomposition techniques in convexification of inequalities. *Technical report*, 2013.
- [53] Mohit Tawarmalani, Jean-Philippe P. Richard, and Kwanghun Chung. Strong valid inequalities for orthogonal disjunctions and bilinear covering sets. *Mathematical Programming*, 124(1):481–512, 2010.
- [54] Mohit Tawarmalani, Jean-Philippe P. Richard, and Chuanhui Xiong. Explicit convex and concave envelopes through polyhedral subdivisions. *Mathematical Programming*, pages 1–47, 2013.
- [55] Mohit Tawarmalani and Nikolaos V Sahinidis. *Convexification and global optimization in continuous and mixed-integer nonlinear programming: theory, algorithms, software, and applications*, volume 65. Springer Science & Business Media, 2002.
- [56] Hoang Tuy. *Convex analysis and global optimization*, volume 110. Springer, 2016.