

# Hitting minors on bounded treewidth graphs.

## IV. An optimal algorithm<sup>1</sup>

Julien Baste<sup>2,3</sup>

Ignasi Sau<sup>3</sup>

Dimitrios M. Thilikos<sup>3</sup>

### Abstract

For a fixed finite collection of graphs  $\mathcal{F}$ , the  $\mathcal{F}$ -M-DELETION problem asks, given an  $n$ -vertex input graph  $G$ , for the minimum number of vertices that intersect all minor models in  $G$  of the graphs in  $\mathcal{F}$ . by Courcelle Theorem, this problem can be solved in time  $f_{\mathcal{F}}(\text{tw}) \cdot n^{\mathcal{O}(1)}$ , where  $\text{tw}$  is the treewidth of  $G$ , for some function  $f_{\mathcal{F}}$  depending on  $\mathcal{F}$ . In a recent series of articles, we have initiated the programme of optimizing asymptotically the function  $f_{\mathcal{F}}$ . Here we provide an algorithm showing that  $f_{\mathcal{F}}(\text{tw}) = 2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})}$  for every collection  $\mathcal{F}$ . Prior to this work, the best known function  $f_{\mathcal{F}}$  was double-exponential in  $\text{tw}$ . In particular, our algorithm vastly extends the results of Jansen et al. [SODA 2014] for the particular case  $\mathcal{F} = \{K_5, K_{3,3}\}$  and of Kociumaka and Pilipczuk [Algorithmica 2019] for graphs of bounded genus, and answers an open problem posed by Cygan et al. [Inf Comput 2017]. We combine several ingredients such as the machinery of boundaried graphs in dynamic programming via representatives, the Flat Wall Theorem, Bidimensionality, the irrelevant vertex technique, treewidth modulators, and protrusion replacement. Together with our previous results providing single-exponential algorithms for particular collections  $\mathcal{F}$  [Theor Comput Sci 2020] and general lower bounds [J Comput Syst Sci 2020], our algorithm yields the following complexity dichotomy when  $\mathcal{F} = \{H\}$  contains a single connected graph  $H$ , assuming the Exponential Time Hypothesis:  $f_H(\text{tw}) = 2^{\Theta(\text{tw})}$  if  $H$  is a contraction of the chair or the banner, and  $f_H(\text{tw}) = 2^{\Theta(\text{tw} \cdot \log \text{tw})}$  otherwise.

**Keywords:** parameterized complexity; graph minors; treewidth; hitting minors; Flat Wall Theorem; irrelevant vertex; dynamic programming; complexity dichotomy.

---

<sup>1</sup>An extended abstract of this article appeared in the *Proceedings of the 31st Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 951-970, Salt Lake City, Utah, U.S., January 2020. The first author was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - 388217545. The two last authors were supported by the ANR projects DEMOGRAPH (ANR-16-CE40-0028), ESIGMA (ANR-17-CE23-0010), ELIT (ANR-20-CE48-0008), the French-German Collaboration ANR/DFG Project UTMA (ANR-20-CE92-0027), and the French Ministry of Europe and Foreign Affairs, via the Franco-Norwegian project PHC AURORA.

<sup>2</sup>Univ. Lille, CNRS, Centrale Lille, UMR 9189 - CRISTAL - Centre de Recherche en Informatique Signal et Automatique de Lille, F-59000 Lille, France. Email: [julien.baste@univ-lille.fr](mailto:julien.baste@univ-lille.fr).

<sup>3</sup>LIRMM, Univ. Montpellier, CNRS, Montpellier, France. Emails: [ignasi.sau@lirmm.fr](mailto:ignasi.sau@lirmm.fr), [sedthilk@thilikos.info](mailto:sedthilk@thilikos.info).

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Overview of the algorithm</b>	<b>3</b>
<b>3</b>	<b>Preliminaries</b>	<b>6</b>
3.1	Basic definitions . . . . .	6
3.2	Formal definition of the problem . . . . .	7
3.3	Boundaried graphs, folios, and representatives . . . . .	8
<b>4</b>	<b>Flat walls</b>	<b>9</b>
4.1	Walls and subwalls . . . . .	10
4.2	Paintings and renditions . . . . .	11
4.3	Flat walls and flatness pairs . . . . .	13
4.4	Influence, regularity, and tilts of flatness pairs . . . . .	14
4.5	Homogeneous walls . . . . .	17
4.6	A parameter for affecting flat walls . . . . .	18
<b>5</b>	<b>Finding an irrelevant vertex</b>	<b>19</b>
5.1	A lemma for model taming . . . . .	19
5.2	Model rerouting in partially disk-embedded graphs . . . . .	22
5.3	Levelings and well-aligned flatness pairs . . . . .	27
5.4	Rerouting minors of small intrusion . . . . .	29
<b>6</b>	<b>Bounding the size of the representatives</b>	<b>36</b>
6.1	Finding a treewidth modulator of linear size . . . . .	36
6.2	Finding a linear protrusion decomposition and reducing protrusions . . . . .	38
<b>7</b>	<b>Further research</b>	<b>41</b>
<b>A</b>	<b>Illustration of the complexity dichotomy</b>	<b>46</b>
<b>B</b>	<b>An estimation of the constants depending on <math>\mathcal{F}</math> in our algorithm</b>	<b>46</b>
B.1	An improved version of <a href="#">Lemma 27</a> . . . . .	47
B.2	Upper bounds on the constants depending on the excluded minors . . . . .	48

# 1 Introduction

Let  $\mathcal{F}$  be a finite non-empty collection of non-empty graphs. In the  $\mathcal{F}$ -M-DELETION problem, we are given a graph  $G$  and an integer  $k$ , and the objective is to decide whether there exists a set  $S \subseteq V(G)$  with  $|S| \leq k$  such that  $G \setminus S$  does not contain any of the graphs in  $\mathcal{F}$  as a minor. This problem belongs to the family of *graph modification problems* and has a big expressive power, as instantiations of it correspond, for instance, to VERTEX COVER ( $\mathcal{F} = \{K_2\}$ ), FEEDBACK VERTEX SET ( $\mathcal{F} = \{K_3\}$ ), and VERTEX PLANARIZATION ( $\mathcal{F} = \{K_5, K_{3,3}\}$ ). Note that if  $\mathcal{F}$  contains a graph with at least one edge, then  $\mathcal{F}$ -M-DELETION is NP-hard [36].

We study the parameterized complexity of  $\mathcal{F}$ -M-DELETION in terms of the treewidth of the input graph (while the size of  $k$  may be unbounded). Since the property of containing a graph as a minor can be expressed in monadic second-order logic [32], by Courcelle Theorem [13],  $\mathcal{F}$ -M-DELETION can be solved in time<sup>1</sup>  $\mathcal{O}^*(f_{\mathcal{F}}(\text{tw}))$  on graphs with treewidth at most  $\text{tw}$ , where  $f_{\mathcal{F}}$  is some computable function depending on  $\mathcal{F}$ . As the function  $f_{\mathcal{F}}(\text{tw})$  given by Courcelle Theorem is typically enormous, our goal is to determine, for a fixed collection  $\mathcal{F}$ , which is the *best possible* such function  $f_{\mathcal{F}}$  that one can (asymptotically) hope for, subject to reasonable complexity assumptions. Besides being an interesting objective in its own, optimizing the running time of algorithms parameterized by treewidth has usually side effects. Indeed, black-box subroutines parameterized by treewidth are nowadays ubiquitous in parameterized [14], exact [19], and approximation [49] algorithms.

**Previous work.** This line of research has attracted considerable attention in the parameterized complexity community during the last years. For instance, VERTEX COVER is easily solvable in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw})})$ , called *single-exponential*, by standard dynamic programming techniques, and no algorithm with running time  $\mathcal{O}^*(2^{o(\text{tw})})$  exists, unless the Exponential Time Hypothesis (ETH) fails [26]. (The ETH implies that 3-SAT on  $n$  variables cannot be solved in time  $2^{o(n)}$ ; see [26] for more details.) For FEEDBACK VERTEX SET, standard dynamic programming techniques give a running time of  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$ , while the lower bound under the ETH [26] is again  $\mathcal{O}^*(2^{o(\text{tw})})$ . This gap remained open for a while, until Cygan et al. [16] presented an optimal (randomized) algorithm running in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw})})$ , introducing the celebrated *Cut & Count* technique. This article triggered several other (deterministic) techniques to obtain single-exponential algorithms for so-called *connectivity problems* on graphs of bounded treewidth, mostly based on algebraic tools [10, 20].

Concerning VERTEX PLANARIZATION, Jansen et al. [27] presented an algorithm running in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$  as a crucial subroutine in an algorithm running in time  $\mathcal{O}^*(2^{\mathcal{O}(k \cdot \log k)})$  where  $k$  is the solution size. Marcin Pilipczuk [39] proved afterwards that this running time is optimal under the ETH, by using the framework introduced by Lokshtanov et al. [37] for proving superexponential lower bounds.

Generalizing the above algorithm, the main technical contribution of the recent paper of Kociumaka and Pilipczuk [33] is an algorithm running in time  $\mathcal{O}^*(2^{\mathcal{O}((\text{tw}+g) \cdot \log(\text{tw}+g))})$  to solve the GENUS VERTEX DELETION problem, which consists in deleting the minimum number of vertices from an input graph in order to obtain a graph embeddable on a surface of Euler genus at most  $g$ .

Cygan et al. [15] studied the problem of hitting subgraphs (instead of minors), and proved that the problem of hitting all copies of a fixed *path* as a subgraph can be solved in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$ . As a path occurs as a subgraph if and only if it occurs as a minor, their result implies that  $\{P_h\}$ -M-DELETION can be solved in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$  for every fixed integer  $h \geq 2$ , where  $P_h$  is the path on  $h$  vertices. They left as an open problem whether the algorithm in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$  of Jansen et al. [27] for VERTEX PLANARIZATION could be generalized to more minor-closed graph classes, other than planar graphs.

In a recent series of three papers [6–8], we initiated a systematic study of the complexity of  $\mathcal{F}$ -M-

---

<sup>1</sup>The notation  $\mathcal{O}^*(\cdot)$  suppresses polynomial factors depending on the size of the input graph.

DELETION, parameterized by treewidth<sup>2</sup>. Before stating these results, we say that a collection  $\mathcal{F}$  is *connected* if it contains only connected graphs.

In [6] we showed that, for every fixed collection  $\mathcal{F}$ ,  $\mathcal{F}$ -M-DELETION can be solved in time  $\mathcal{O}^*(2^{2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})}})$  by a natural dynamic programming algorithm, and that if  $\mathcal{F}$  contains a planar graph, the running time can be improved<sup>3</sup> to  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$ . If the input graph  $G$  is planar or, more generally, embedded in a surface of bounded genus, then we showed that the running time can be further improved to  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw})})$ .

In [7] we provided single-exponential algorithms for  $\{H\}$ -M-DELETION when  $H$  is either  $P_4$ ,  $C_4$ , the claw, the paw, the chair, or the banner; see Figure 17 in Appendix A for an illustration of these graphs.

In [8] we focused on lower bounds under the ETH. We proved that for any connected  $\mathcal{F}$  containing graphs on at least two vertices,  $\mathcal{F}$ -M-DELETION cannot be solved in time  $\mathcal{O}^*(2^{o(\text{tw})})$ , even if the input graph  $G$  is planar. More notably, we proved that  $\mathcal{F}$ -M-DELETION cannot be solved in time  $\mathcal{O}^*(2^{o(\text{tw} \cdot \log \text{tw})})$  for collections  $\mathcal{F}$  satisfying some generic conditions. In particular, these conditions apply when  $\mathcal{F}$  contains a single connected graph  $H$  that is *not* a contraction of the chair or the banner. Note that the connected graphs  $H$  with  $|V(H)| \geq 2$  that are a contraction of the chair or the banner are those on the left in Figure 17, and for each of them  $\{H\}$ -M-DELETION can be solved in (optimal) single-exponential time [7, 8].

**Our results.** In this article we present an algorithm to solve  $\mathcal{F}$ -M-DELETION in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$  for every collection  $\mathcal{F}$ , thus making a significant step towards a complete classification of the complexity of the  $\mathcal{F}$ -M-DELETION problem parameterized by treewidth. That is, we drop the condition that  $\mathcal{F}$  contains a planar graph, which was critically needed in the algorithm presented in [6] in order to bound the treewidth of an  $\mathcal{F}$ -minor-free graph. Our algorithm can be interpreted as an exponential “collapse” of the natural dynamic programming algorithm running in time  $\mathcal{O}^*(2^{2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})}})$  given in [6]. Besides largely improving our previous results [6], this algorithm generalizes the ones for  $\mathcal{F} = \{K_5, K_{3,3}\}$  given by Jansen et al. [27] and for the GENUS VERTEX DELETION problem given by Kociumaka and Pilipczuk [33], which are based on embeddings, and answers positively the open problem of Cygan et al. [15] for every minor-closed graph class. Since the algorithm is quite involved, we provide an overview of it in Section 2.

This algorithm in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$  for every collection  $\mathcal{F}$ , together with the lower bounds under the ETH given in [8], the single-exponential algorithms given in [7], and the known cases  $\mathcal{F} = \{P_2\}$  [14, 26],  $\mathcal{F} = \{P_3\}$  [4, 48], and  $\mathcal{F} = \{C_3\}$  [10, 16], imply the following complexity dichotomy when  $\mathcal{F}$  consists of a single connected graph  $H$ , which we suppose to have at least one edge.

**Theorem 1.** *Let  $H$  be a connected graph. Under the ETH,  $\{H\}$ -M-DELETION is solvable in time<sup>4</sup>*

- $2^{\Theta(\text{tw})} \cdot n^{\mathcal{O}(1)}$ , if  $H$  is a contraction of the chair or the banner, and
- $2^{\Theta(\text{tw} \cdot \log \text{tw})} \cdot n^{\mathcal{O}(1)}$ , otherwise.

This dichotomy is depicted in Figure 17, containing all connected graphs  $H$  with  $2 \leq |V(H)| \leq 5$ ; note that if  $|V(H)| \geq 6$ , then  $H$  is not a contraction of the chair or the banner, and therefore the second item above applies. Note also that  $K_4$  and the diamond are the only graphs on at most four vertices for which the problem is solvable in time  $\mathcal{O}^*(2^{\Theta(\text{tw} \cdot \log \text{tw})})$  and that the chair and the banner are the only graphs on at least five vertices for which the problem is solvable in time  $\mathcal{O}^*(2^{\Theta(\text{tw})})$ .

The crucial role played by the chair and the banner in the complexity dichotomy may seem surprising at first sight. In fact, we realized a posteriori that the “easy” cases can be succinctly described in terms of the chair and the banner by taking a look at Figure 17. Note that the “easy” graphs can be equivalently

<sup>2</sup>In these papers [6–8], in some results we also considered the version of the problem where the graphs in  $\mathcal{F}$  are forbidden as *topological* minors; in the current paper we focus exclusively on the minor version.

<sup>3</sup>In the conference version of [6] we additionally required  $\mathcal{F}$  to be connected; in the journal version we proved this result without this assumption.

<sup>4</sup>We use  $n$  and  $\text{tw}$  for the number of vertices and the treewidth of the input graph, respectively.

characterized as those that are minors of the **banner**, with the exception of  $P_5$ . Nevertheless, there is some intuitive reason for which excluding the **chair** or the **banner** constitutes the horizon on the existence of single-exponential algorithms. Namely, focusing on the **banner**, every connected component (with at least five vertices) of a graph that excludes the **banner** as a minor is either a cycle (of any length) or a tree in which some vertices have been replaced by triangles; both such types of components can be maintained by a dynamic programming algorithm in single-exponential time [7]. A similar situation occurs when excluding the **chair**. It appears that if the characterization of the allowed connected components is enriched in some way, such as restricting the length of the allowed cycles or forbidding certain degrees, the problem becomes inherently more difficult, inducing a transition from time  $\mathcal{O}^*(2^{\Theta(\text{tw})})$  to  $\mathcal{O}^*(2^{\Theta(\text{tw} \cdot \log \text{tw})})$ .

**Organization of the paper.** In [Section 2](#) we provide a high-level overview of the algorithm running in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$ . In [Section 3](#) we give some preliminaries. In [Section 4](#) we deal with flat walls, in [Section 5](#) we apply the irrelevant vertex technique in the context of boundaried graphs, and in [Section 6](#) we use this in order to bound the size of the dynamic programming tables. We conclude the article in [Section 7](#) with some open questions for further research. In [Appendix B](#) we present an estimation of the constants depending on the (fixed) collection  $\mathcal{F}$  in our algorithm (cf. [Corollary 36](#)).

## 2 Overview of the algorithm

In order to obtain our algorithm of time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$  for every collection  $\mathcal{F}$ , our approach can be streamlined as follows. We use the machinery of boundaried graphs, equivalence relations, and representatives originating in the seminal work of Bodlaender et al. [11] and subsequently used, for instance, in [6, 21, 22, 32]. Let  $h$  be a constant depending only on the collection  $\mathcal{F}$  (to be defined in the formal description of the algorithm) and let  $t$  be a positive integer that is at most the treewidth of the input graph plus one. Skipping several technical details, a  $t$ -boundaried graph is a graph with a distinguished set of vertices –its *boundary*– labeled bijectively with integers from the set  $[t]$ . We say that two  $t$ -boundaried graphs are  $h$ -equivalent if for any other  $t$ -boundaried graph that we can “glue” to each of them, resulting in graphs  $G_1$  and  $G_2$ , and every graph  $H$  on at most  $h$  vertices,  $H$  is a minor of  $G_1$  if and only if it is a minor of  $G_2$  (cf. [Section 3](#) for the precise definitions). Let  $\mathcal{R}_h^{(t)}$  be a set of *minimum-sized* representatives of this equivalence relation. Since  $h$ -equivalent (boundaried) graphs have the same behavior in terms of eventual occurrences of minors of size up to  $h$ , there is a generic dynamic programming algorithm (already used in [6]) to solve  $\mathcal{F}$ -M-DELETION on a rooted tree decomposition of the input graph, via a typical bottom-up approach: at every bag  $B$  of the tree decomposition, naturally associated with a  $t$ -boundaried graph  $G_B$ , and for every representative  $R \in \mathcal{R}_h^{(t)}$ , store the minimum size of a set  $S \subseteq V(G_B)$  such that the graph  $G_B \setminus S$  is  $h$ -equivalent to  $R$  (cf. [Subsection 6.2](#) for some more details<sup>5</sup>). This yields an algorithm running in time  $\mathcal{O}^*(|\mathcal{R}_h^{(t)}|^2)$ , and therefore it suffices to prove that  $|\mathcal{R}_h^{(t)}| = 2^{\mathcal{O}_h(t \cdot \log t)}$ , where the notation ‘ $\mathcal{O}_h(\cdot)$ ’ means that the hidden constants depend only on  $h$ . Since we may assume that the graphs in  $\mathcal{R}_h^{(t)}$  exclude some graph on at most  $h$  vertices as a minor (as all those that do not are  $h$ -equivalent), hence they have a linear number of edges [38], it is enough to prove that, for every  $R \in \mathcal{R}_h^{(t)}$ , it holds that

$$|V(R)| = \mathcal{O}_h(t). \tag{1}$$

Note that this is indeed sufficient in order to obtain an algorithm within the claimed running time, as there are at most  $\binom{|V(R)|^2}{|E(R)|} = 2^{\mathcal{O}_h(|V(R)| \cdot \log |V(R)|)}$  representatives, and  $t! = 2^{\mathcal{O}(t \cdot \log t)}$  possible labelings for

<sup>5</sup>This step was the only reason for which in the conference version of this article we required the collection  $\mathcal{F}$  to be connected. As mentioned in [Section 1](#), in the full version of [6] we dropped the connectivity assumption, which implies that in the current article we can drop it as well.

the vertices in the boundary. In order to prove Equation 1, we combine a number of different techniques, which we proceed to discuss informally, and that are schematically summarized in Figure 1:




Figure 1: Diagram of the algorithm in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$  for any collection  $\mathcal{F}$ .

► We use the Flat Wall Theorem of Robertson and Seymour [42], in particular a version (Theorem 5) that has been recently proved in [45] (and is based on the framework of Kawarabayashi et al. [30]), which incorporates the so-called *regularity property*. In a nutshell, this theorem says that every  $K_h$ -minor-free graph  $G$  has a set of vertices  $A \subseteq V(G)$  –called *apices*– with  $|A| = \mathcal{O}_h(1)$  such that  $G \setminus A$  contains a *flat wall* of height  $\Omega_h(\text{tw}(G))$ . Here, the definition of “flat wall” is quite involved and is detailed in Section 4; it essentially means a subgraph that has a bidimensional grid-like structure, separated from the rest of the graph by its perimeter, and that is “close” to being planar, in the sense that it can be embedded in the plane in a way that its potentially non-planar pieces, called *flaps*, have a well-defined structure along larger pieces called *bricks*.

► A subwall of a flat wall is *h-homogeneous* if for every brick of the subwall, the flaps within that brick have the same variety of *h-folios*, that is, the same sets of “boundaried” minors of *detail* at most  $h$  (the detail of

a boundaried graph is the maximum between its number of edges and its number of non-boundary vertices). This notion is inspired (but is not the same) by the one defined by Robertson and Seymour in [42]. Using standard “zooming” arguments, we can prove that, given a flat wall, we can find a large  $h$ -homogeneous subwall inside it (Proposition 9). Homogeneous subwalls are very useful because, as we explain below, they permit the application of the irrelevant vertex technique adapted to our purposes.

► We say that a vertex set  $S$  *affects* a flat wall if some vertex within the wall has a neighbor in  $S$  that is *not* an apex. With these definitions at hand, we define a parameter, denoted by  $\mathbf{p}_{h,r}$  in this informal description, mapping every graph  $G$  to the smallest size of a vertex set that affects all  $h$ -homogeneous flat walls with at most  $h$  apices and height at least  $r$  in  $G$ . It is not hard to prove that the parameter  $\mathbf{p}_{h,r}$  has a “bidimensional” behavior [17, 18, 21], in the sense that its value on a flat wall depends quadratically on the height of the wall (Lemma 10) and it is *separable* [11, 18, 21] (Lemma 11).

► The most complicated step towards proving Equation 1 is to find an “irrelevant” vertex inside a sufficiently large (in terms of  $h$ ) flat wall of a boundaried graph that is *not* affected by its boundary (Theorem 23). Informally, here “irrelevant” means a non-boundary vertex of  $R$  that can be avoided by *any* minor model of a graph on at most  $h$  vertices and edges that traverses the boundary of  $R$ , *no matter* the graph that may be glued to it and *no matter how* this model traverses the boundary of  $R$ ; see Section 5 for the precise definition. The *irrelevant vertex technique* originated in the seminal work of Robertson and Seymour [42, 43] and has become a very useful tool used in various kinds of linkage and cut problems [1, 27, 33, 34, 40]. Nevertheless, given the nature of our setting, it is critical that the size of the flat wall where the irrelevant vertex appears does *not depend* on the boundary size. To the best of our knowledge, this property is not guaranteed by the existing results on the irrelevant vertex technique (such as [42, (10.2)] and its subsequent proof in [43]). To achieve it and, moreover, in order to make an estimation of the parametric dependencies, we develop a self-reliant theoretical framework that uses the following ingredients:

- With a flat wall  $W$  we associate a bipartite graph  $\tilde{W}$ , which we call its *leveling* as defined in [45]; cf. Subsection 5.3 for the precise definition. In particular, this graph has a vertex for every flap of the flat wall, and can be embedded in a disk in a planar way.
- It turns out to be more convenient to work with *topological* minor models instead of minor models; we can afford it since for every graph  $H$  there are at most  $f(H)$  different topological minor minimal graphs that contain  $H$  as a minor (Observation 3). The reason for this is that it is easier to deal with the branch vertices of a topological minor model in the analysis. Given a topological minor model, we say that a flap of a wall is *dirty* if it contains a branch vertex of the model, or there is an edge from the flap to an apex vertex of the wall. We also define the leveling of a topological minor model, and we equip its dirty flaps with colors that encode their  $h$ -folios. We now proceed to explain how to reroute the colored leveling of a topological minor model.
- In order to reroute (colored levelings of) topological minor models, it will be helpful to use *railed annuli*, a structure introduced in [28] that occurs as a subgraph inside a flat wall (Proposition 12) and that has the following nice property, recently proved in [25] (Proposition 13): if a railed annulus is large enough compared to  $h$ , every topological minor model of a graph on at most  $h$  vertices traversing it can be rerouted so that the branch vertices are preserved and such that, more importantly, the intersection of the new model with a large *prescribed* part of the railed annulus is *confined*, in the sense that it is only allowed to use a well-defined set of paths in that part, which does not depend on the original model.
- We also need a technical result with a graph drawing flavor (Lemma 16) guaranteeing that large enough railed annuli contain topological minor models of every graph of maximum degree three with the property, in particular, that certain vertices are pairwise far apart in the embedding. Using this result and the one proved in [25] mentioned above, we can finally prove (Theorem 17) that every

topological minor model of a graph  $H$  inside a graph with a large flat wall  $W$  can be “collapsed” inside the wall, in the following sense:  $G$  contains another topological minor model of a graph  $H'$ , such that  $H$  is a minor of  $H'$ , and such that the new model avoids the central part of the annulus; here is where the irrelevant vertex will be found.

- To conclude, it just remains to “lift” the constructed embedding of the colored leveling of the topological minor to an embedding of the “original” minor in the flat wall ([Theorem 23](#)). For that, we exploit the fact that we have rerouted the model inside an  $h$ -homogeneous subwall not affected by the boundary, which allows to mimic the behavior of the original minor inside the flaps of the wall, using that all bricks have the same variety of  $h$ -folios.

The above arguments, incorporated in the proof of [Theorem 23](#), imply that if  $R \in \mathcal{R}_h^{(t)}$  is a minimum-sized representative, then its boundary affects all large enough  $h$ -homogeneous flat walls, as otherwise we could remove an irrelevant vertex and find a smaller equivalent representative. In particular, it follows that, for every  $R \in \mathcal{R}_h^{(t)}$ , we have  $\mathbf{p}_{h,r}(R) \leq t$  ([Corollary 25](#)).

► Combining that the parameter  $\mathbf{p}_{h,r}$  is “bidimensional” and separable along with the fact that  $\mathbf{p}_{h,r}(R) \leq t$  for every  $R \in \mathcal{R}_h^{(t)}$ , we prove in [Lemma 27](#) that every representative  $R \in \mathcal{R}_h^{(t)}$  has a vertex subset  $S$  containing its boundary, with  $|S| = \mathcal{O}_h(t)$ , whose removal leaves a graph of treewidth bounded by a function of  $h$ ; such a set is called a *treewidth modulator*. (In [Appendix B](#) we provide an improved version of [Lemma 27](#), namely [Lemma 34](#), by adapting the proof of [[21](#), Lemma 3.6].)

► Once we have a treewidth modulator of size  $\mathcal{O}_h(t)$  of a representative  $R$ , all that remains is to pipeline it with known techniques to compute an appropriate *protrusion decomposition* [[32](#)] ([Lemma 29](#)) and to *reduce protrusions* to smaller equivalent ones of size bounded by a function of  $h$  –we use the version given in [[6](#)] adapted to the  $\mathcal{F}$ -M-DELETION problem– ([Lemma 30](#)), implying that  $|V(R)| = \mathcal{O}_h(t)$  for every every  $R \in \mathcal{R}_h^{(t)}$  and concluding the proof of [Equation 1](#).

It should be noted that all the items above do *not* need to be converted into an algorithm, they are just used in the analysis: the conclusion is that if  $R \in \mathcal{R}_h^{(t)}$  is a minimum-sized representative, then  $|V(R)| = \mathcal{O}_h(t)$ , as otherwise some reduction rule could be applied to it (either by removing an irrelevant vertex or by protrusion replacement), thus obtaining an equivalent representative of smaller size and contradicting its minimality. Our main result can be formally stated as follows.

**Theorem 2.** *Let  $\mathcal{F}$  be a finite non-empty collection of non-empty graphs. There exists a constant  $c_{\mathcal{F}}$  such that the  $\mathcal{F}$ -M-DELETION problem is solvable in time  $c_{\mathcal{F}}^{\text{tw} \cdot \log \text{tw}} \cdot n$  on  $n$ -vertex graphs of treewidth at most  $\text{tw}$ .*

In [Appendix B](#) we provide an estimation of the constant  $c_{\mathcal{F}}$  in the above theorem based on the parametric dependencies of the Unique Linkage Theorem [[31](#), [43](#)].

## 3 Preliminaries

### 3.1 Basic definitions

**Sets and integers.** We denote by  $\mathbb{N}$  the set of non-negative integers. Given two integers  $p, q$ , where  $p \leq q$ , we denote by  $[p, q]$  the set  $\{p, \dots, q\}$ . For an integer  $p \geq 1$ , we set  $[p] = [1, p]$  and  $\mathbb{N}_{\geq p} = \mathbb{N} \setminus [0, p-1]$ . Given a non-negative integer  $p$ , we denote by  $\text{odd}(p)$  the minimum odd number that is not smaller than  $p$ . For a set  $S$ , we denote by  $2^S$  the set of all subsets of  $S$  and by  $\binom{S}{2}$  the set of all subsets of  $S$  of size two. If  $S$  is a collection of objects where the operation  $\cup$  is defined, then we denote  $\mathbf{U}S = \bigcup_{X \in S} X$ .

**Basic concepts on graphs.** As a graph  $G$  we denote any pair  $(V, E)$  where  $V$  is a finite set and  $E \subseteq \binom{V}{2}$ , that is, all graphs of this paper are undirected, finite, and without loops or multiple edges. We also define  $V(G) = V$  and  $E(G) = E$ . Unless stated otherwise, we denote by  $n$  and  $m$  the number of vertices and edges, respectively, of the graph under consideration. We say that a pair  $(L, R) \in 2^{V(G)} \times 2^{V(G)}$  is a *separation* of  $G$  if  $L \cup R = V(G)$  and there is no edge in  $G$  between  $L \setminus R$  and  $R \setminus L$ . The *order* of a separation  $(L, R)$  is  $|L \cap R|$ . Given a vertex  $v \in V(G)$ , we denote by  $N_G(v)$  the set of vertices of  $G$  that are adjacent to  $v$  in  $G$ . Also, given a set  $S \subseteq V(G)$ , we set  $N_G(S) = (\bigcup_{v \in S} N_G(v)) \setminus S$ . A vertex  $v \in V(G)$  is *isolated* if  $N_G(v) = \emptyset$ . For  $S \subseteq V(G)$ , we set  $G[S] = (S, E \cap \binom{S}{2})$  and use  $G \setminus S$  to denote  $G[V(G) \setminus S]$ . Given an edge  $e = \{u, v\} \in E(G)$ , we define the *subdivision* of  $e$  to be the operation of deleting  $e$ , adding a new vertex  $w$ , and making it adjacent to  $u$  and  $v$ . Given two graphs  $H, G$ , we say that  $H$  is a *subdivision* of  $G$  if  $H$  can be obtained from  $G$  by subdividing edges. The *contraction* of an edge  $e = \{u, v\}$  of a simple graph  $G$  results in a simple graph  $G'$  obtained from  $G \setminus \{u, v\}$  by adding a new vertex  $w$  adjacent to all the vertices in the set  $N_G(u) \cup N_G(v) \setminus \{u, v\}$ . A graph  $G'$  is a *minor* of a graph  $G$  if  $G'$  can be obtained from a subgraph of  $G$  after a series of edge contractions. The *distance* between two vertices  $x$  and  $y$  of a graph  $G$  is the number of edges of a shortest path between  $x$  and  $y$  in  $G$ .

**Treewidth.** Let  $G = (V, E)$  be a graph. A *tree decomposition* of  $G$  is a pair  $(T, \mathcal{X} = \{X_t\}_{t \in V(T)})$  where  $T$  is a tree and  $\mathcal{X}$  is a collection of subsets of  $V$  such that

- $\bigcup_{t \in V(T)} X_t = V$ ,
- $\forall e = \{u, v\} \in E, \exists t \in V(T) : \{u, v\} \subseteq X_t$ , and
- $\forall v \in V, T[\{t \mid v \in X_t\}]$  is connected.

We call the vertices of  $T$  *nodes* and the sets in  $\mathcal{X}$  *bags* of the tree decomposition  $(T, \mathcal{X})$ . The *width* of  $(T, \mathcal{X})$  is equal to  $\max\{|X_t| - 1 \mid t \in V(T)\}$  and the *treewidth* of  $G$  is the minimum width over all tree decompositions of  $G$ . We denote the treewidth of a graph  $G$  by  $\text{tw}(G)$ .

For  $t \in \mathbb{N}$ , we say that a set  $S \subseteq V(G)$  is a  *$t$ -treewidth modulator* of  $G$  if  $\text{tw}(G \setminus S) \leq t$ .

### 3.2 Formal definition of the problem

Let  $\mathcal{F}$  be a finite non-empty collection of non-empty graphs; we call such a collection *proper*. We extend the minor relation to  $\mathcal{F}$  such that, given a graph  $G$ ,  $\mathcal{F} \preceq_m G$  if and only if there exists a graph  $H \in \mathcal{F}$  such that  $H \preceq_m G$ . We also denote  $\text{exc}_m(\mathcal{F}) = \{G \mid \mathcal{F} \not\preceq_m G\}$ , i.e.,  $\text{exc}_m(\mathcal{F})$  is the class of graphs that do *not* contain any graph in  $\mathcal{F}$  as a minor.

Let  $\mathcal{F}$  be a proper collection. We define the graph parameter  $\mathbf{m}_{\mathcal{F}}$  as the function that maps graphs to non-negative integers as follows:

$$\mathbf{m}_{\mathcal{F}}(G) = \min\{|S| \mid S \subseteq V(G) \wedge G \setminus S \in \text{exc}_m(\mathcal{F})\}.$$

The main objective of this paper is to study the problem of computing the parameter  $\mathbf{m}_{\mathcal{F}}$  for graphs of bounded treewidth. The corresponding decision problem is formally defined as follows.

**$\mathcal{F}$ -M-DELETION**  
**Input:** A graph  $G$  and an integer  $k \in \mathbb{N}$ .  
**Parameter:** The treewidth of  $G$ .  
**Output:** Is  $\mathbf{m}_{\mathcal{F}}(G) \leq k$ ?

### 3.3 Boundaried graphs, folios, and representatives

**Boundaried graphs.** Let  $t \in \mathbb{N}$ . A  $t$ -boundaried graph is a triple  $\mathbf{G} = (G, B, \rho)$  where  $G$  is a graph,  $B \subseteq V(G)$ ,  $|B| = t$ , and  $\rho : B \rightarrow [t]$  is a bijection. We say that two  $t$ -boundaried graphs  $\mathbf{G}_1 = (G_1, B_1, \rho_1)$  and  $\mathbf{G}_2 = (G_2, B_2, \rho_2)$  are *isomorphic* if there is an isomorphism from  $G_1$  to  $G_2$  that extends the bijection  $\rho_2^{-1} \circ \rho_1$ . The triple  $(G, B, \rho)$  is a *boundaried graph* if it is a  $t$ -boundaried graph for some  $t \in \mathbb{N}$ . As in [42], we define the *detail* of a boundaried graph  $\mathbf{G} = (G, B, \rho)$  as  $\text{detail}(\mathbf{G}) := \max\{|E(G)|, |V(G) \setminus B|\}$ . We denote by  $\mathcal{B}^{(t)}$  the set of all (pairwise non-isomorphic)  $t$ -boundaried graphs and by  $\mathcal{B}_h^{(t)}$  the set of all (pairwise non-isomorphic)  $t$ -boundaried graphs with detail at most  $h$ . We also set  $\mathcal{B} = \bigcup_{t \in \mathbb{N}} \mathcal{B}^{(t)}$ .

**Minors and topological minors of boundaried graphs.** We say that a  $t$ -boundaried graph  $\mathbf{G}_1 = (G_1, B_1, \rho_1)$  is a *minor* of a  $t$ -boundaried graph  $\mathbf{G}_2 = (G_2, B_2, \rho_2)$ , denoted by  $\mathbf{G}_1 \preceq_m \mathbf{G}_2$ , if there is a sequence of removals of non-boundary vertices, edge removals, and edge contractions in  $G_2$ , disallowing contractions of edges with both endpoints in  $B_2$ , that transforms  $\mathbf{G}_2$  to a boundaried graph that is isomorphic to  $\mathbf{G}_1$  (during edge contractions, boundary vertices prevail). Note that this extends the usual definition of minors in graphs without boundary.

We say that  $(M, T)$  is a *tm-pair* if  $M$  is a graph,  $T \subseteq V(M)$ , and all vertices in  $V(M) \setminus T$  have degree two. We denote by  $\text{diss}(M, T)$  the graph obtained from  $M$  by *dissolving* all vertices in  $V(M) \setminus T$ , that is, for every vertex  $v \in V(M) \setminus T$ , with neighbors  $u$  and  $w$ , we delete  $v$  and, if  $u$  and  $w$  are not adjacent, we add the edge  $\{u, w\}$ . A *tm-pair* of a graph  $G$  is a *tm-pair*  $(M, T)$  where  $M$  is a subgraph of  $G$ .

Given two graphs  $H$  and  $G$ , we say that a *tm-pair*  $(M, T)$  of  $G$  is a *topological minor model of  $H$  in  $G$*  if  $H$  is isomorphic to  $\text{diss}(M, T)$ . We denote this isomorphism by  $\sigma_{M, T} : V(H) \rightarrow T$ . We call the vertices in  $T$  *branch* vertices of  $(M, T)$ . We call each path in  $M$  between two distinct branch vertices and with no internal branch vertices a *subdivision path* of  $(M, T)$  and the internal vertices of such paths, i.e., the vertices of  $V(M) \setminus T$ , are the *subdivision* vertices of  $(M, T)$ . We also extend  $\sigma_{M, T}$  so to also map each  $e = \{x, y\} \in E(H)$  to the subdivision path of  $M$  with endpoints  $\sigma_{M, T}(x)$  and  $\sigma_{M, T}(y)$ . Furthermore, we extend  $\sigma_{M, T}$  so to also map each subgraph  $H'$  of  $H$  to the subgraph of  $M$  consisting of the vertices of  $\sigma_{M, T}(T)$  and the paths in  $\sigma_{M, T}(e)$ ,  $e \in E(H')$ .

If  $\mathbf{M} = (M, B, \rho) \in \mathcal{B}$  and  $T \subseteq V(M)$  with  $B \subseteq T$  and such that all vertices in  $V(M) \setminus T$  have degree two, we call  $(\mathbf{M}, T)$  a *btm-pair* and we define  $\text{diss}(\mathbf{M}, T) = (\text{diss}(M, T), B, \rho)$ . Note that we do not permit dissolution of boundary vertices, as we consider all of them to be branch vertices. If  $\mathbf{G} = (G, B, \rho)$  is a boundaried graph and  $(M, T)$  is a *tm-pair* of  $G$  where  $B \subseteq T$ , then we say that  $(\mathbf{M}, T)$ , where  $\mathbf{M} = (M, B, \rho)$ , is a *btm-pair of  $\mathbf{G} = (G, B, \rho)$* . Let  $\mathbf{G}_i = (G_i, B_i, \rho_i)$ ,  $i \in [2]$ . We say that  $\mathbf{G}_1$  is a *topological minor* of  $\mathbf{G}_2$ , denoted by  $\mathbf{G}_1 \preceq_{\text{tm}} \mathbf{G}_2$ , if  $\mathbf{G}_2$  has a *btm-pair*  $(\mathbf{M}, T)$  such that  $\text{diss}(\mathbf{M}, T)$  is isomorphic to  $\mathbf{G}_1$ .

Given a  $\mathbf{G} = (G, B, \rho) \in \mathcal{B}$ , we define  $\text{ext}(\mathbf{G})$  as the set containing every topological-minor-minimal boundaried graph  $\mathbf{G}' = (G', B, \rho)$  among those that contain  $\mathbf{G}$  as a minor. Notice that we insist that  $B$  and  $\rho$  are the same for all graphs in  $\text{ext}(\mathbf{G})$ . Moreover, we do not consider isomorphic boundaried graphs in  $\text{ext}(\mathbf{G})$  as different boundaried graphs. The set  $\text{ext}(\mathbf{G})$  helps us to express the minor relation in terms of the topological minor relation because of the following simple observation. Note that this definition extends naturally to graphs, seen as boundaried graphs with empty boundary.

*Observation 3.* If  $\mathbf{G}_1, \mathbf{G}_2 \in \mathcal{B}$ , then  $\mathbf{G}_1 \preceq_m \mathbf{G}_2 \iff \exists \mathbf{G} \in \text{ext}(\mathbf{G}_2) : \mathbf{G}_1 \preceq_{\text{tm}} \mathbf{G}$ . Moreover, if  $\mathbf{G}$  is a boundaried graph with detail  $h$ , then every graph in  $\text{ext}(\mathbf{G})$  has detail at most  $3h$ .

**Folios.** We define the *h-folio* of  $\mathbf{G} = (G, B, \rho) \in \mathcal{B}$  as

$$h\text{-folio}(\mathbf{G}) = \{\mathbf{G}' \in \mathcal{B} \mid \mathbf{G}' \preceq_{\text{tm}} \mathbf{G} \text{ and } \mathbf{G}' \text{ has detail at most } h\}.$$

Using the fact that an  $h$ -folio is a collection of  $K_{h+1}$ -minor-free boundaried graphs, it follows that the

$h$ -folio of a  $t$ -boundaried graph has at most  $2^{\mathcal{O}((h+t)\cdot\log(h+t))}$  elements. Therefore, the number of distinct  $h$ -folios of  $t$ -boundaried graphs is given by the following lemma (also observed in [6]).

**Lemma 4.** *There exists a function  $f_1 : \mathbb{N}^2 \rightarrow \mathbb{N}$  such that for every  $t, h \in \mathbb{N}$ ,  $|\{h\text{-folio}(\mathbf{G}) \mid \mathbf{G} \in \mathcal{B}_h^{(t)}\}| \leq f_1(t, h)$ . Moreover,  $f_1(t, h) = 2^{2^{\mathcal{O}((h+t)\cdot\log(h+t))}}$ .*

**Equivalent boundaried graphs and representatives.** We say that two boundaried graphs  $\mathbf{G}_1 = (G_1, B_1, \rho_1)$  and  $\mathbf{G}_2 = (G_2, B_2, \rho_2)$  are *compatible* if  $\rho_2^{-1} \circ \rho_1$  is an isomorphism from  $G_1[B_1]$  to  $G_2[B_2]$ . Given two compatible boundaried graphs  $\mathbf{G}_1 = (G_1, B_1, \rho_1)$  and  $\mathbf{G}_2 = (G_2, B_2, \rho_2)$ , we define  $\mathbf{G}_1 \oplus \mathbf{G}_2$  as the graph obtained if we take the disjoint union of  $G_1$  and  $G_2$  and, for every  $i \in [|B_1|]$ , we identify vertices  $\rho_1^{-1}(i)$  and  $\rho_2^{-1}(i)$ .

Given  $h \in \mathbb{N}$ , we say that two boundaried graphs  $\mathbf{G}_1$  and  $\mathbf{G}_2$  are  *$h$ -equivalent*, denoted by  $\mathbf{G}_1 \equiv_h \mathbf{G}_2$ , if they are compatible and, for every graph  $H$  on at most  $h$  vertices and  $h$  edges and every boundaried graph  $\mathbf{F}$  that is compatible with  $\mathbf{G}_1$  (hence, with  $\mathbf{G}_2$  as well), it holds that

$$H \preceq_m \mathbf{F} \oplus \mathbf{G}_1 \iff H \preceq_m \mathbf{F} \oplus \mathbf{G}_2. \quad (2)$$

Note that  $\equiv_h$  is an equivalence relation on  $\mathcal{B}$ . A minimum-sized (in terms of number of vertices) element of an equivalence class of  $\equiv_h$  is called *representative* of  $\equiv_h$ . For  $t \in \mathbb{N}$ , a *set of  $t$ -representatives* for  $\equiv_h$  is a collection containing a minimum-sized representative for each equivalence class of  $\equiv_h$  restricted to  $\mathcal{B}^{(t)}$ . Given  $t, h \in \mathbb{N}$ , we denote by  $\mathcal{R}_h^{(t)}$  a set of  $t$ -representatives for  $\equiv_h$ .

At this point, we wish to stress that the folio-equivalence defined in Equation 2 is related but is *not* the same as the one defined by “having the same  $h$ -folio”. Indeed, observe first that if  $\mathbf{G}_1$  and  $\mathbf{G}_2$  are compatible  $t$ -boundaried graphs and  $h\text{-folio}(\mathbf{G}_1) = h\text{-folio}(\mathbf{G}_2)$  then  $\mathbf{G}_1 \equiv_h \mathbf{G}_2$ , therefore the folio-equivalence is a refinement of  $\equiv_h$ . In fact, a dynamic programming procedure for solving  $\mathcal{F}$ -M-DELETION can also be based on the folio-equivalence, and this has already been done in the general algorithm in [6], which has a double-exponential parametric dependence due to the bound of Lemma 4. In this paper we build our dynamic programming on the equivalence  $\equiv_h$  and we essentially prove that  $\equiv_h$  is “coarse enough” so to reduce the double-exponential parametric dependence of the dynamic programming to a single-exponential one. In fact, this has already been done in [6] for the case where  $\mathcal{F}$  contains some planar graph, as this structural restriction directly implies an upper bound on the treewidth of the representatives. To deal with the general case, the only structural restriction for the (non-trivial) representatives is the exclusion of  $H$  as a minor. All the combinatorial machinery that we introduce in the next two sections is intended to deal with the structure of this general and (more entangled) setting.

## 4 Flat walls

In this section we deal with flat walls. More precisely, in Subsections 4.1, 4.2, and 4.3 we give the definition of a *flat wall* in the form of a *flatness pair*. In Subsection 4.4 we define the notion of *regular flatness pair* and we give a version of the Flat Wall Theorem of Robertson and Seymour [42] that has been recently proved in [45]. This version (Theorem 5) incorporates the regularity property and is based on the recent results and the terminology of Kawarabayashi et al. [30]. In Subsection 4.5 we define a notion of homogeneity of flat walls, also introduced in [45], that along with Theorem 5 will be the combinatorial framework for the proofs of Section 5. We stress that the notion of homogeneity that we use is different from that defined by Robertson and Seymour in [42] and can serve as an alternative for further applications based on the technology of flat walls (see e.g. [44, 46, 47]). In Subsection 4.6 we define a graph parameter related to flat walls and show that it enjoys a series of properties related to Bidimensionality (as introduced in [17] and further developed in [21]).

## 4.1 Walls and subwalls

We first introduce some basic concepts such as partially disk-embedded graphs, walls, subwalls, tilts, and layers (for an example of all the concepts defined in this subsection, see [Figure 2](#)).

**Partially disk-embedded graphs.** A *closed* (resp. *open*) *disk* is a set homeomorphic to the set  $\{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1\}$  (resp.  $\{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 1\}$ ). Let  $\Delta$  be an open or closed disk. We use  $\text{bd}(\Delta)$  to denote the boundary of  $\Delta$  and, if  $\Delta$  is closed, we use  $\text{int}(\Delta)$  to denote the open disk  $\Delta \setminus \text{bd}(\Delta)$ . Also, if  $\Delta$  is an open disk, we use  $\bar{\Delta} = \Delta \cup \text{bd}(\Delta)$  for the closure of  $\Delta$ . When we embed a graph  $G$  in the plane or in a disk, we treat  $G$  (both its vertex and edge sets) as a set of points. This permits us to make set operations between graphs and sets of points.

If  $\Delta$  is a closed disk, we say that a graph  $G$  is  $\Delta$ -*embedded* if  $G$  is embedded in  $\Delta$  without crossings such that the intersection of  $\text{bd}(\Delta)$  and  $G$  (seen as a set of points of  $\Delta$ ) is a subset of  $V(G)$ . We say that a graph  $G$  is *partially disk-embedded in some closed disk*  $\Delta$ , if there is some  $\Delta$ -embedded subgraph, say  $K$ , of  $G$  such that  $G \cap \Delta = K$  and  $(V(G) \cap \Delta, V(G) \setminus \text{int}(\Delta))$  is a separation of  $G$ . From now on, we use the term *partially  $\Delta$ -embedded graph*  $G$  to denote that a graph  $G$  is partially disk-embedded in some closed disk  $\Delta$ .

A *circle* of  $\Delta$  is any set homeomorphic to  $\{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$ . Given two distinct points  $x, y \in \Delta$ , an  $(x, y)$ -*arc* of  $\Delta$  is any subset of  $\Delta$  that is homeomorphic to the closed interval  $[0, 1]$ .




Figure 2: An 13-wall  $W$  along with a choice of pegs and corners. The six layers of  $W$  are colored alternatively in red and green and the central 5-subwall of  $W$  appears in grey. The pegs are the squared vertices while, among them, those that are black are the corners. The original vertices that are not pegs are turquoise circles while the subdivision vertices are the small yellow circles. The central vertices are the two 3-branch vertices that are surrounded by white squares. Notice that  $W$  has 144 bricks and, among them, 100 are internal.

**Walls.** Let  $k, r \in \mathbb{N}$ . The  $(k \times r)$ -*grid* is the graph whose vertex set is  $[k] \times [r]$  and two vertices  $(i, j)$  and  $(i', j')$  are adjacent if  $|i - i'| + |j - j'| = 1$ . An *elementary  $r$ -wall*, for some odd integer  $r \geq 3$ , is the graph obtained from a  $(2r \times r)$ -grid with vertices  $(x, y) \in [2r] \times [r]$ , after the removal of the “vertical” edges  $\{(x, y), (x, y + 1)\}$  for odd  $x + y$ , and then the removal of all vertices of degree one. Notice that, as  $r \geq 3$ , an elementary  $r$ -wall is a planar graph that has a unique (up to topological isomorphism) embedding in the plane  $\mathbb{R}^2$  such that all its finite faces are incident to exactly six edges. The *perimeter* of an elementary  $r$ -wall

is the cycle bounding its infinite face, while the cycles bounding its finite faces are called *bricks*. Also, the vertices in the perimeter of an elementary  $r$ -wall that have degree two are called *pegs*, while the vertices  $(1, 1)$ ,  $(2, r)$ ,  $(2r - 1, 1)$ , and  $(2r, r)$  are called *corners* (notice that the corners are also pegs).

An  $r$ -wall is any graph  $W$  obtained from an elementary  $r$ -wall  $\bar{W}$  after subdividing edges. A graph  $W$  is a wall if it is an  $r$ -wall for some odd  $r \geq 3$  and we refer to  $r$  as the *height* of  $W$ . Given a graph  $G$ , a wall of  $G$  is a subgraph of  $G$  that is a wall. We insist that, for every  $r$ -wall, the number  $r$  is always odd.

We call the vertices of degree three of a wall  $W$  *3-branch vertices*. The vertices that are created by subdivisions are called *subdivision vertices* while the rest are called *original vertices* of  $W$ . A cycle of  $W$  is a *brick* (resp. the *perimeter*) of  $W$  if its 3-branch vertices are the vertices of a brick (resp. the perimeter) of  $\bar{W}$ . We use  $D(W)$  in order to denote the perimeter of the wall  $W$ . A brick of  $W$  is *internal* if it is disjoint from  $D(W)$ . Note that every wall  $W$  has a unique (up to homeomorphism) embedding in the plane whose infinite face is bounded by the perimeter  $D(W)$  of the wall. Each time we consider a plane-embedded wall, we consider this embedding.

Given two vertices  $x$  and  $y$  of a plane graph  $G$ , we define their *face-distance* in  $G$  as the smallest integer  $i$  such that there exists an arc of the plane (i.e., a subset homeomorphic to the interval  $[0, 1]$ ) between  $x$  and  $y$  that does not cross the infinite face of the embedding, crosses no vertices of  $G$ , and intersects at most  $i$  faces of  $G$ . Note that two distinct vertices of a plane wall  $W$  are within face-distance one if and only if they belong to the same brick. Given two vertex sets  $X, Y$  of a plane graph  $G$ , we define the face-distance between  $X$  and  $Y$  as the minimum face-distance between a vertex in  $X$  and a vertex in  $Y$ .

**Subwalls.** Given an elementary  $r$ -wall  $\bar{W}$ , some  $i \in \{1, 3, \dots, 2r - 1\}$ , and  $i' = (i + 1)/2$ , the  $i'$ -th *vertical path* of  $\bar{W}$  is the one whose vertices, in order of appearance, are  $(i, 1), (i, 2), (i + 1, 2), (i + 1, 3), (i, 3), (i, 4), (i + 1, 4), (i + 1, 5), (i, 5), \dots, (i, r - 2), (i, r - 1), (i + 1, r - 1), (i + 1, r)$ . Also, given some  $j \in [2, r - 1]$  the  $j$ -th *horizontal path* of  $\bar{W}$  is the one whose vertices, in order of appearance, are  $(1, j), (2, j), \dots, (2r, j)$ .

A *vertical* (resp. *horizontal*) path of  $W$  is one that is a subdivision of a vertical (resp. horizontal) path of  $\bar{W}$ . Notice that the perimeter of an  $r$ -wall  $W$  is uniquely defined regardless of the choice of the elementary  $r$ -wall  $\bar{W}$ . A *subwall* of  $W$  is any subgraph  $W'$  of  $W$  that is an  $r'$ -wall, with  $r' \leq r$  and such the vertical (resp. horizontal) paths of  $W'$  are subpaths of the vertical (resp. horizontal) paths of  $W$ .

**Tilts.** The *interior* of a wall  $W$  is the graph obtained from  $W$  if we remove from it all edges of  $D(W)$  and all vertices of  $D(W)$  that have degree two in  $W$ . Given two walls  $W$  and  $\tilde{W}$  of a graph  $G$ , we say that  $\tilde{W}$  is a *tilt* of  $W$  if  $\tilde{W}$  and  $W$  have identical interiors.

**Layers.** The *layers* of an  $r$ -wall  $W$  are recursively defined as follows. The first layer of  $W$  is its perimeter. For  $i = 2, \dots, (r - 1)/2$ , the  $i$ -th layer of  $W$  is the  $(i - 1)$ -th layer of the subwall  $W'$  obtained from  $W$  after removing from  $W$  its perimeter and removing recursively all occurring vertices of degree one. The *central vertices* of an  $r$ -wall are its two 3-branch vertices that do not belong to any of its layers. See [Figure 2](#) for an illustration of the notions defined above. Given an  $r$ -wall  $W$  and an odd integer  $q$ , where  $3 \leq q \leq r$ , the *central  $q$ -subwall* of  $W$  is the subwall of  $W$  of height  $q$  whose central vertices are the central vertices of  $W$ .

## 4.2 Paintings and renditions

Before defining flat walls, we need to introduce paintings and renditions. Here we closely follow the terminology of [\[30\]](#).

**Paintings.** Let  $\Delta$  be a closed disk. A  $\Delta$ -*painting* is a pair  $\Gamma = (U, N)$  where

- $N$  is a finite set of points of  $\Delta$ ,
- $N \subseteq U \subseteq \Delta$ , and
- $U \setminus N$  has finitely many arcwise-connected components, called *cells*, where, for every cell  $c$ ,
  - the closure  $\bar{c}$  of  $c$  is a closed disk and
  - $|\tilde{c}| \leq 3$ , where  $\tilde{c} := \text{bd}(c) \cap N$ .

We use the notation  $U(\Gamma) := U$ ,  $N(\Gamma) := N$  and denote the set of cells of  $\Gamma$  by  $C(\Gamma)$ . For convenience, we may assume that each cell of  $\Gamma$  is an open disk of  $\Delta$ . See [Figure 3](#) for an example of a  $\Delta$ -painting.




Figure 3: A  $\Delta$ -painting  $\Gamma = (U, N)$ . The red circles are the points of  $N$  that are points of the boundary of  $\Delta$  (whose complement is drawn in grey) and the blue circles are those that lie in the interior of  $\Delta$ . The set  $U \setminus N$  is depicted in green.

Notice that, given a  $\Delta$ -painting  $\Gamma$ , the pair  $(N(\Gamma), \{\tilde{c} \mid c \in C(\Gamma)\})$  is a hypergraph whose hyperedges have cardinality at most three and  $\Gamma$  can be seen as a plane embedding of this hypergraph in  $\Delta$ .

**Renditions.** Let  $G$  be a graph, and let  $\Omega$  be a cyclic permutation of a subset of  $V(G)$  that we denote by  $V(\Omega)$ . By an  $\Omega$ -*rendition* of  $G$  we mean a triple  $(\Gamma, \sigma, \pi)$ , where

- (a)  $\Gamma$  is a  $\Delta$ -painting for some closed disk  $\Delta$ ,
- (b)  $\pi : N(\Gamma) \rightarrow V(G)$  is an injection, and
- (c)  $\sigma$  assigns to each cell  $c \in C(\Gamma)$  a subgraph  $\sigma(c)$  of  $G$ , such that
  - (1)  $G = \bigcup_{c \in C(\Gamma)} \sigma(c)$ ,
  - (2) for distinct  $c, c' \in C(\Gamma)$ ,  $\sigma(c)$  and  $\sigma(c')$  are edge-disjoint,
  - (3) for every cell  $c \in C(\Gamma)$ ,  $\pi(\tilde{c}) \subseteq V(\sigma(c))$ ,

- (4) for every cell  $c \in C(\Gamma)$ ,  $V(\sigma(c)) \cap \bigcup_{c' \in C(\Gamma) \setminus \{c\}} V(\sigma(c')) \subseteq \pi(\tilde{c})$ , and
- (5)  $\pi(N(\Gamma) \cap \text{bd}(\Delta)) = V(\Omega)$ , such that the points in  $N(\Gamma) \cap \text{bd}(\Delta)$  appear in  $\text{bd}(\Delta)$  in the same ordering as their images, via  $\pi$ , in  $\Omega$ .

Given an  $\Omega$ -rendition  $(\Gamma, \sigma, \pi)$  of a graph  $G$ , we call a cell  $c$  of  $\Gamma$  *trivial* if  $\pi(\tilde{c}) = V(\sigma(c))$ .

**Tight renditions.** We say that an  $\Omega$ -rendition  $(\Gamma, \sigma, \pi)$  of a graph  $G$  is *tight* if the following conditions are satisfied:

- (i) If there are two points  $x, y$  of  $N(\Gamma)$  such that  $e = \{\pi(x), \pi(y)\} \in E(G)$ , then there is a cell  $c \in C(\Gamma)$  such that  $\sigma(c)$  is the two-vertex connected graph  $(e, \{e\})$ ,
- (ii) for every  $c \in C(\Gamma)$ , every two vertices in  $\pi(\tilde{c})$  belong to some path of  $\sigma(c)$ ,
- (iii) for every  $c \in C(\Gamma)$  and every connected component  $C$  of the graph  $\sigma(c) \setminus \pi(\tilde{c})$ , if  $N_{\sigma(c)}(V(C)) \neq \emptyset$ , then  $N_{\sigma(c)}(V(C)) = \pi(\tilde{c})$ ,
- (iv) there are no two distinct non-trivial cells  $c_1$  and  $c_2$  such that  $\pi(\tilde{c}_1) = \pi(\tilde{c}_2)$ , and
- (v) for every  $c \in C(\Gamma)$  there are  $|\tilde{c}|$  vertex-disjoint paths in  $G$  from  $\pi(\tilde{c})$  to the set  $V(\Omega)$ .

As proved in [45], it is possible to transform any  $\Omega$ -rendition to a tight one. For this reason, in this paper, we always assume that  $\Omega$ -renditions are tight.

### 4.3 Flat walls and flatness pairs

We are now in position to define the notion of a flat wall. We further encode it into the concept of a *flatness pair* of a graph.

**Flat walls.** Let  $G$  be a graph and let  $W$  be an  $r$ -wall of  $G$ , for some odd integer  $r \geq 3$ . We say that a pair  $(P, C) \subseteq D(W) \times D(W)$  is a *choice of pegs and corners for  $W$*  if  $W$  is the subdivision of an elementary  $r$ -wall  $\bar{W}$  where  $P$  and  $C$  are the pegs and the corners of  $\bar{W}$ , respectively (clearly,  $C \subseteq P$ ). To get more intuition, notice that a wall  $W$  can occur in several ways from the elementary wall  $\bar{W}$ , depending on the way the vertices in the perimeter of  $\bar{W}$  are subdivided. Each of them gives a different selection  $(P, C)$  of pegs and corners of  $W$  (see Figure 2 for an example of a choice of pegs and corners  $(P, C)$  in a 13-wall  $W$ ).

We say that  $W$  is a *flat  $r$ -wall* of  $G$  if there is a separation  $(X, Y)$  of  $G$  and a choice  $(P, C)$  of pegs and corners for  $W$  such that:

- $V(W) \subseteq Y$ ,
- $P \subseteq X \cap Y \subseteq V(D(W))$ , and
- if  $\Omega$  is the cyclic ordering of the vertices  $X \cap Y$  as they appear in  $D(W)$ , then there exists an  $\Omega$ -rendition  $(\Gamma, \sigma, \pi)$  of  $G[Y]$ .




Figure 4: A graph  $G$  and a flatness pair  $(W, \mathfrak{R})$  of  $G$  where  $W$  is a 5-wall and  $\mathfrak{R} = (X, Y, P, C, \Gamma, \sigma, \pi)$  is a 7-tuple certifying the flatness of  $W$  in  $G$ . The edges of  $W$  are drawn in orange. In the corresponding separation  $(X, Y)$ , the vertices of  $X$  are green and yellow while the vertices in  $Y$  are all the non-green vertices. Consequently, the yellow vertices are the vertices in  $X \cap Y$ . The pegs and the corners are the squared vertices where the pegs that are not corners are purple and the the corners are black. The  $\Delta$ -painting of the  $\Omega$ -rendition  $(\Gamma, \sigma, \pi)$  of  $G' = G[Y]$  is the one depicted in Figure 3.

**Flatness pairs.** Given the above, we say that the choice of the 7-tuple  $\mathfrak{R} = (X, Y, P, C, \Gamma, \sigma, \pi)$  certifies that  $W$  is a flat wall of  $G$ . We call the pair  $(W, \mathfrak{R})$  a *flatness pair* of  $G$  and define the *height* of the pair  $(W, \mathfrak{R})$  to be the height of  $W$ . We use the term *cell* of  $\mathfrak{R}$  in order to refer to the cells of  $\Gamma$  (see Figure 4 for an example of a flatness pair  $(W, \mathfrak{R})$  of a graph).

We call the graph  $G[Y]$  the  $\mathfrak{R}$ -compass of  $W$  in  $G$ , denoted by  $\text{compass}_{\mathfrak{R}}(W)$  (see Figure 5 for the  $\mathfrak{R}$ -compass of  $W$ , corresponding to the flatness pair  $(W, \mathfrak{R})$  of Figure 4). We define the *flaps* of the wall  $W$  in  $\mathfrak{R}$  as  $\text{flaps}_{\mathfrak{R}}(W) := \{\sigma(c) \mid c \in C(\Gamma)\}$ . Given a flap  $F \in \text{flaps}_{\mathfrak{R}}(W)$ , we define its *base* as  $\partial F := V(F) \cap \pi(N(\Gamma))$ . A flap  $F \in \text{flaps}_{\mathfrak{R}}(W)$  is *trivial* if  $|\partial F| = 2$  and  $F$  consists of one edge between the two vertices in  $\partial F$ . We call the edges of the trivial flaps *short edges* of  $\text{compass}_{\mathfrak{R}}(W)$ . A cell  $c$  of  $\mathfrak{R}$  is *untidy* if  $\pi(\tilde{c})$  contains a vertex  $x$  of  $W$  such that two of the edges of  $W$  that are incident to  $x$  are edges of  $\sigma(c)$ . Notice that if  $c$  is untidy then  $|\tilde{c}| = 3$ . A cell is *tidy* if it is not untidy (in Figure 4 untidy cells are marked by green stars).

#### 4.4 Influence, regularity, and tilts of flatness pairs

We now introduce a classification of the cells of a flatness pair  $(W, \mathfrak{R})$ . This classification will be used in order to define the concepts of regularity and  $W'$ -tilts of flatness pairs that will be important for our proofs.

**Cell classification.** Given a cycle  $C$  of  $\text{compass}_{\mathfrak{R}}(W)$ , we say that  $C$  is  $\mathfrak{R}$ -normal if it is not a subgraph of a flap  $F \in \text{flaps}_{\mathfrak{R}}(W)$ . Given an  $\mathfrak{R}$ -normal cycle  $C$  of  $\text{compass}_{\mathfrak{R}}(W)$ , we call a cell  $c$  of  $\mathfrak{R}$   $C$ -perimetric if  $\sigma(c)$  contains some edge of  $C$ . Notice that if  $c$  is  $C$ -perimetric, then  $\pi(\tilde{c})$  contains two points  $p, q \in N(\Gamma)$  such that  $\pi(p)$  and  $\pi(q)$  are vertices of  $C$  where one, say  $P_c^{\text{in}}$ , of the two  $(\pi(p), \pi(q))$ -subpaths of  $C$  is a subgraph of  $\sigma(c)$  and the other, denoted by  $P_c^{\text{out}}$ ,  $(\pi(p), \pi(q))$ -subpath contains at most one internal vertex of  $\sigma(c)$ ,




Figure 5: The  $\mathfrak{R}$ -compass of the 5-wall  $W$  for the flatness pair  $(W, \mathfrak{R})$  depicted in Figure 4, and a subwall  $W'$  of  $W$  whose edges are depicted in bold. The red curve is the curve  $K_{W'}$ . The  $W'$ -internal cells are depicted in grey while the  $W'$ -perimetric cells are depicted in green.  $W'$ -marginal cells are marked with orange stars. The set  $\text{influence}_{\mathfrak{R}}(W')$  contains all the flaps that are drawn inside the grey or the green cells (the  $W'$ -external cells are not depicted).

which should be the (unique) vertex  $z$  in  $\partial\sigma(c) \setminus \{\pi(p), \pi(q)\}$ . We pick a  $(p, q)$ -arc  $A_c$  in  $\hat{c} := c \cup \tilde{c}$  such that  $\pi^{-1}(z) \in A_c$  if and only if  $P_c^{\text{in}}$  contains the vertex  $z$  as an internal vertex.

We consider the circle  $K_C = \bigcup\{A_c \mid c \text{ is a } C\text{-perimetric cell of } \mathfrak{R}\}$  and we denote by  $\Delta_C$  the closed disk bounded by  $K_C$  that is contained in  $\Delta$ . A cell  $c$  of  $\mathfrak{R}$  is called  $C$ -internal if  $c \subseteq \Delta_C$  and is called  $C$ -external if  $\Delta_C \cap c = \emptyset$ . Notice that the cells of  $\mathfrak{R}$  are partitioned into  $C$ -internal,  $C$ -perimetric, and  $C$ -external cells.

Let  $c$  be a tidy  $C$ -perimetric cell of  $\mathfrak{R}$  where  $|\tilde{c}| = 3$ . Notice that  $c \setminus A_c$  has two arcwise-connected components and one of them is an open disk  $D_c$  that is a subset of  $\Delta_C$ . If the closure  $\overline{D}_c$  of  $D_c$  contains only two points of  $\tilde{c}$  then we call the cell  $c$   $C$ -marginal.

**Influence.** For every  $\mathfrak{R}$ -normal cycle  $C$  of  $\text{compass}_{\mathfrak{R}}(W)$  we define the set

$$\text{influence}_{\mathfrak{R}}(C) = \{\sigma(c) \mid c \text{ is a cell of } \mathfrak{R} \text{ that is not } C\text{-external}\}.$$

A wall  $W'$  of  $\text{compass}_{\mathfrak{R}}(W)$  is  $\mathfrak{R}$ -normal if  $D(W')$  is  $\mathfrak{R}$ -normal. Notice that every wall of  $W$  (and hence every subwall of  $W$ ) is an  $\mathfrak{R}$ -normal wall of  $\text{compass}_{\mathfrak{R}}(W)$ . We denote by  $\mathcal{S}_{\mathfrak{R}}(W)$  the set of all  $\mathfrak{R}$ -normal walls of  $\text{compass}_{\mathfrak{R}}(W)$ . Given a  $W' \in \mathcal{S}_{\mathfrak{R}}(W)$  and a cell  $c$  of  $\mathfrak{R}$  we say that  $c$  is  $W'$ -perimetric/internal/external/marginal if  $c$  is  $D(W')$ -perimetric/internal/external/marginal (see Figure 5 for an example). We also use  $K_{W'}$ ,  $\Delta_{W'}$ ,  $\text{influence}_{\mathfrak{R}}(W')$  as shortcuts for  $K_{D(W')}$ ,  $\Delta_{D(W')}$ ,  $\text{influence}_{\mathfrak{R}}(D(W'))$ .

**Regularity.** Let  $(W, \mathfrak{R})$  be a flatness pair of a graph  $G$ . We call a flatness pair  $(W, \mathfrak{R})$  of a graph  $G$  regular if none of its cells is  $W$ -external,  $W$ -marginal, or untidy. Notice that the flatness pair of Figure 4 is not regular (for an example of a regular flatness pair of a graph that is a modification of the one in Figure 4, see Figure 9). The notion of regularity has been defined in [45] and will be useful in Subsection 5.3. In fact,

regularity permits the definition of a “well-allinged”  $\Delta$ -embedded representation of the  $\mathfrak{R}$ -compass that will be valuable in the proofs of [Subsection 5.4](#). The precise definition of the notion of well-allinged flatness pairs is given in [Subsection 5.3](#).

The next result has been proved in [45]. It can be seen as a version of the Flat Wall Theorem incorporating the concept of regularity, which is necessary for our proofs.

**Theorem 5.** *There exist two functions  $f_2, f_3 : \mathbb{N} \rightarrow \mathbb{N}$  such that for every graph  $G$ , every odd integer  $r \geq 3$ , and every  $q \in \mathbb{N}_{\geq 1}$ , one of the following is true:*

- $K_q$  is a minor of  $G$ ,
- $\text{tw}(G) \leq f_2(q) \cdot r$ , or
- there exist a set  $A \subseteq V(G)$ , where  $|A| \leq f_3(q)$ , and a regular flatness pair  $(W, \mathfrak{R})$  of  $G \setminus A$  of height  $r$ .

Moreover,  $f_2(q) = 2^{\mathcal{O}(q^2 \log q)}$  and  $f_3(q) = \mathcal{O}(q^{24})$ .

**Tilts of flatness pairs.** Let  $(W, \mathfrak{R})$  and  $(\tilde{W}', \tilde{\mathfrak{R}}')$  be two flatness pairs of a graph  $G$  and let  $W' \in \mathcal{S}_{\mathfrak{R}}(W)$ . We also assume that  $\mathfrak{R} = (X, Y, P, C, \Gamma, \sigma, \pi)$  and  $\tilde{\mathfrak{R}}' = (X', Y', P', C', \Gamma', \sigma', \pi')$ . We say that  $(\tilde{W}', \tilde{\mathfrak{R}}')$  is a  $W'$ -tilt of  $(W, \mathfrak{R})$  if

- $\tilde{\mathfrak{R}}'$  does not have  $\tilde{W}'$ -external cells,
- $\tilde{W}'$  is a tilt of  $W'$ ,
- the set of  $\tilde{W}'$ -internal cells of  $\tilde{\mathfrak{R}}'$  is the same as the set of  $W'$ -internal cells of  $\mathfrak{R}$  and their images via  $\sigma'$  and  $\sigma$  are also the same,
- $\text{compass}_{\tilde{\mathfrak{R}}'}(\tilde{W}')$  is a subgraph of  $\mathbf{U}\text{influence}_{\mathfrak{R}}(W')$ , and
- if  $c$  is a cell in  $C(\Gamma') \setminus C(\Gamma)$ , then  $|\tilde{c}| \leq 2$ .

The next observation follows from the definitions of regular flatness pairs and tilts.

*Observation 6.* If  $(W, \mathfrak{R})$  is a regular flatness pair, then for every  $W' \in \mathcal{S}_{\mathfrak{R}}(W)$ , every  $W'$ -tilt of  $(W, \mathfrak{R})$  is also regular.

We need one more observation, which follows from the third item above and the fact that the cells corresponding to flaps containing a central vertex of  $W'$  are all internal (recall that the height of a wall is always at least three).

*Observation 7.* The central vertices of  $W'$  belong to every  $W'$ -tilt of  $(W, \mathfrak{R})$ .

The need to define  $W'$ -tilts of flatness pairs emerges from the fact that not every subwall  $W'$  of a flat wall  $W$  is necessarily flat, recently observed in [45]. The next proposition, proved in [45], suggests that there is always a *slight modification* of  $W'$  in the  $\mathfrak{R}$ -compass of  $W$  that is indeed a flat wall. This “tilt” preserves the internal cells, and therefore the “essential” part of the influence of  $W'$ . That way, it permits us to define a notion of compass relative to a subwall of a flat wall.

**Proposition 8.** *For every flatness pair  $(W, \mathfrak{R})$  of a graph  $G$  and every  $W' \in \mathcal{S}_{\mathfrak{R}}(W)$ , there exists a flatness pair  $(\tilde{W}', \tilde{\mathfrak{R}}')$  of  $G$  that is a  $W'$ -tilt of  $(W, \mathfrak{R})$ .*

## 4.5 Homogeneous walls

Homogeneous walls were a basic ingredient of the seminal algorithm of Robertson and Seymour for the DISJOINT PATHS problem in [42]. This algorithm introduced the *Irrelevant Vertex Technique* that consisted in the identification of a vertex in an instance of the DISJOINT PATHS problem that is *irrelevant* in the sense that its removal does not change the YES/NO-status of the instance. The notion of wall homogeneity was given in [42] and was based on the concept of the *vision* of an “internal” flap of a flat wall. It was proved in [43] that the central vertices of a sufficiently big homogenous flat wall are indeed *irrelevant* with respect to the DISJOINT PATHS problem and therefore they could safely discarded. Our results are following the same technique. However, we need an alternative notion of homogeneity that we introduce in this subsection.

Let  $G$  be a graph, let  $A \subseteq V(G)$ , and let  $(W, \mathfrak{R})$  be a flatness pair of  $G \setminus A$ , where  $\mathfrak{R} = (X, Y, P, C, \Gamma, \sigma, \pi)$  and  $(\Gamma, \sigma, \pi)$  is an  $\Omega$ -rendition of  $G[Y]$ . Recall that  $\Gamma = (U, N)$  is a  $\Delta$ -painting of the closed disk  $\Delta$ .

**Augmented flaps.** For each flap  $F \in \text{flaps}_{\mathfrak{R}}(W)$  we consider a labeling  $\ell_F : \partial F \rightarrow \{1, 2, 3\}$  such that the set of labels assigned by  $\ell_F$  to  $\partial F$  is one of  $\{1\}$ ,  $\{1, 2\}$ ,  $\{1, 2, 3\}$ . We also consider a bijection  $\rho_A : A \rightarrow [a]$ , where  $a = |A|$ . The labelings in  $\mathcal{L} = \{\ell_F \mid F \in \text{flaps}_{\mathfrak{R}}(W)\}$  and the labeling  $\rho_A$  will be useful for defining a set of boundaried graphs that we will call augmented flaps. We first need some more definitions.

Given a flap  $F \in \text{flaps}_{\mathfrak{R}}(W)$ , we define an ordering  $\Omega(F) = (x_1, \dots, x_q)$ , with  $q \leq 3$ , of the vertices of  $\partial F$  so that

- $(x_1, \dots, x_q)$  is a counter-clockwise cyclic ordering of the vertices of  $\partial F$  as they appear in the corresponding cell of  $C(\Gamma)$ . Notice that this cyclic ordering is significant only when  $|\partial F| = 3$ , in the sense that  $(x_1, x_2, x_3)$  remains invariant under shifting, i.e.,  $(x_1, x_2, x_3)$  is the same as  $(x_2, x_3, x_1)$  but not under inversion, i.e.,  $(x_1, x_2, x_3)$  is not the same as  $(x_3, x_2, x_1)$ , and
- for  $i \in [q]$ ,  $\ell_F(x_i) = i$ .

Notice that the second condition is necessary for completing the definition of the ordering  $\Omega(F)$ , and this is the reason why we set up the labelings in  $\mathcal{L}$ .

For each  $F \in \text{flaps}_{\mathfrak{R}}(W)$  with  $t_F = |\partial F|$ , we fix  $\rho_F : \partial F \rightarrow [a+1, a+t_F]$  such that  $(\rho_F^{-1}(a+1), \dots, \rho_F^{-1}(a+t_F)) = \Omega(F)$ . Also, we define the boundaried graph

$$\mathbf{F}^A = (G[A \cup F], A \cup \partial F, \rho_A \cup \rho_F) \quad (3)$$

and we denote by  $F^A$  the underlying graph of  $\mathbf{F}^A$ . We call  $\mathbf{F}^A$  an *augmented flap* of the flatness pair  $(W, \mathfrak{R})$  of  $G \setminus A$  in  $G$ .

**Palettes and homogeneity.** For each cycle  $C$  of  $W$ , we define  $(A, \ell)$ -palette( $C$ ) =  $\{\ell\text{-folio}(\mathbf{F}^A) \mid F \in \text{influence}_{\mathfrak{R}}(C)\}$ . We say that the flatness pair  $(W, \mathfrak{R})$  of  $G \setminus A$  is  $\ell$ -homogeneous with respect to the pair  $(G, A)$  if every *internal* brick  $B$  of  $W$  (seen as a cycle of  $W$ ) has the *same*  $(A, \ell)$ -palette.

**Apex-wall triples.** Let  $G$  be a graph, let  $A \subseteq V(G)$  with  $|A| \leq a$ , and let  $(W, \mathfrak{R})$  be a regular flatness pair of  $G \setminus A$  such that  $W$  has height  $r$  and is  $\ell$ -homogenous with respect to  $(G, A)$  for some  $\ell \in \mathbb{N}$ . We call such a triple  $(A, W, \mathfrak{R})$  an  $(a, r, \ell)$ -apex-wall triple of  $G$ .

The next proposition, proved in [45], implies that it is possible to find an  $\ell$ -homogeneous flat wall inside the compass of a sufficiently big flat wall.

**Proposition 9.** *There exists a function  $f_4 : \mathbb{N}^2 \rightarrow \mathbb{N}$  such that if  $r \in \mathbb{N}_{\geq 3}$ ,  $G$  is a graph,  $A \subseteq V(G)$ , and  $(W, \mathfrak{R})$  is a flatness pair of  $G \setminus A$  of height  $f_4(r, w)$ , where  $w = f_1(|A| + 3, \ell)$ , then  $W$  contains some subwall  $W'$  of height  $r$  such that every  $W'$ -tilt of  $(W, \mathfrak{R})$  is  $\ell$ -homogeneous with respect to  $(G, A)$ . Moreover,  $f_4(r, w) = \mathcal{O}(r^w)$ .*

## 4.6 A parameter for affecting flat walls

We proceed to define a graph parameter that will be useful for our proofs. We prove that it satisfies some properties related to Bidimensionality theory [17, 18, 21] that will be used later in Subsection 6.2.

Let  $G$  be a graph and let  $(A, W, \mathfrak{R})$  be an  $(a, r, \ell)$ -apex-wall triple of  $G$ . We say that  $S$  affects  $(A, W, \mathfrak{R})$  if  $N_G[V(\text{compass}_{\mathfrak{R}}(W))] \cap (S \setminus A) \neq \emptyset$ . For  $a, r, \ell \in \mathbb{N}$ , we define

$$\mathbf{p}_{a,r,\ell}(G) = \min\{k \mid \exists S \subseteq V(G) : |S| \leq k \wedge S \text{ affects every } (a, r, \ell)\text{-apex-wall triple of } G\}.$$

Using Theorem 5, Proposition 8, and Proposition 9, we prove that the above parameter grows quadratically with its treewidth.

**Lemma 10.** *There is a function  $f_5 : \mathbb{N}^3 \rightarrow \mathbb{N}$  such that if  $q, r, \ell \in \mathbb{N}_{>0}$ , and  $G$  is a  $K_q$ -minor-free graph, then  $\text{tw}(G) \leq f_5(q, r, \ell) \cdot \max\{1, \sqrt{\mathbf{p}_{f_3(q),r,\ell}(G)}\}$ . In particular, one may choose  $f_5(q, r, \ell) = \mathcal{O}(f_2(q) \cdot r^w)$ , where  $w = f_1(f_3(q) + 3, \ell) = 2^{2^{\mathcal{O}((q^{24} + \ell) \cdot \log(q^{24} + \ell))}}$ .*

*Proof.* The lemma follows easily if we prove that, for every positive integer  $p$ ,  $\text{tw}(G) > f_2(q) \cdot ((r^w + 2) \cdot \lceil \sqrt{p+1} \rceil + 2)$  implies that  $\mathbf{p}_{f_3(q),r,\ell}(G) > p$ .

By Theorem 5, it follows that, for any  $q, r, \ell \in \mathbb{N}_{>0}$ , if  $G$  is  $K_q$ -minor-free and  $\text{tw}(G) > f_2(q) \cdot ((r^w + 2) \cdot \lceil \sqrt{p+1} \rceil + 2)$ , then  $G$  contains some vertex set  $A$  such that  $|A| \leq f_3(q)$  and  $G \setminus A$  has a regular flatness pair  $(W, \mathfrak{R})$  of height  $(r^w + 2) \cdot \lceil \sqrt{p+1} \rceil + 2$ . Let  $\hat{W}_1, \dots, \hat{W}'_{p+1} \in \mathcal{S}_{\mathfrak{R}}(W)$  be a collection of  $p+1$  pairwise disjoint subwalls of  $W$ , each of height  $r^w$ , such that there are no two vertices  $w_1, w_2$  in  $W$  of face-distance at most one such that  $w_1$  belongs to some  $\hat{W}_i$  and  $w_2$  belongs either in  $D(W)$  or in some other  $\hat{W}'_{i'}$ ,  $i' \neq i$ .

By Proposition 8, for every  $i \in [p+1]$ , there is a  $\hat{W}_i$ -tilt of  $(W, \mathfrak{R})$  that we denote by  $(W_i, \mathfrak{R}_i)$ . Since  $(W, \mathfrak{R})$  is regular, Observation 6 implies that  $(W_i, \mathfrak{R}_i)$  is also regular for every  $i \in [p+1]$ . Moreover, by the definition of a  $\hat{W}_i$ -tilt, it follows that, for every two distinct  $i, i' \in [p+1]$ ,  $N_G[V(\text{compass}_{\mathfrak{R}_i}(W_i))] \cap N_G[V(\text{compass}_{\mathfrak{R}_{i'}}(W_{i'}))] \subseteq A$ . Note that this is correct because the face-distance demand leaves a “buffer” among the flat walls and the perimeter of  $W$  to guarantee that the neighborhoods of their compasses do not intersect, except possibly at apex vertices. Let  $w = f_1(|A| + 3, \ell)$ . From Proposition 9, for each regular flatness pair  $(W_i, \mathfrak{R}_i)$  of  $G \setminus A$  there is a subwall  $\hat{W}'_i$  of height  $r$  such that every regular  $\hat{W}'_i$ -tilt of  $(W, \mathfrak{R})$  is  $\ell$ -homogeneous with respect to  $(G, A)$ . We denote this  $\hat{W}'_i$ -tilt by  $(W'_i, \mathfrak{R}'_i)$  and we conclude that  $(A, W'_i, \mathfrak{R}'_i)$  is an  $(a, r, \ell)$ -apex-wall triple of  $G$ , for every  $i \in [p+1]$ . As before,  $N_G[V(\text{compass}_{\mathfrak{R}'_i}(W'_i))] \cap N_G[V(\text{compass}_{\mathfrak{R}'_{i'}}(W'_{i'}))] \subseteq A$  for every  $i, i' \in [p+1]$ ,  $i \neq i'$ . Therefore, every set  $S \subseteq V(G)$  affecting every  $(f_3(q), r, \ell)$ -apex-wall triple of  $G$  needs to contain at least one vertex from each of the sets  $\{N_G[V(\text{compass}_{\mathfrak{R}_i}(W_i))] \mid i \in [p]\}$ , implying that  $\mathbf{p}_{f_3(q),r,\ell}(G) > p$ .  $\square$

We now prove that the parameter  $\mathbf{p}_{a,r,\ell}$  is *separable*, that is, that when considering a separation of a graph, the value of the parameter is “evenly” split along both sides of the separation, possibly with an offset bounded by the order of the separation.

**Lemma 11.** *Let  $a, r, \ell \in \mathbb{N}$ , let  $G$  be a graph, and let  $S \subseteq V(G)$  such that  $S$  affects every  $(a, r, \ell)$ -apex-wall triple of  $G$ . Then, for every separation  $(L, R)$  of  $S$  in  $G$ , the set  $L \cap (R \cup S)$  affects every  $(a, r, \ell)$ -apex-wall triple of  $G[L]$ .*

*Proof.* Suppose for contradiction that  $(A, W, \mathfrak{A})$  is an  $(a, r, \ell)$ -apex-wall triple of  $G[L]$  that is not affected by  $L \cap (R \cup S)$ . In particular, it holds that  $V(\text{compass}_{\mathfrak{A}}(W)) \subseteq L \setminus R$ . Since by assumption  $(A, W, \mathfrak{A})$  is affected by  $S$  but not by  $L \cap (R \cup S)$ , there should exist a vertex  $v \in S \cap (R \setminus L)$  with a neighbor in  $V(\text{compass}_{\mathfrak{A}}(W)) \subseteq L \setminus R$ , contradicting the hypothesis that  $(L, R)$  is a separation of  $G$ .  $\square$

## 5 Finding an irrelevant vertex

In this section we show how to find inside a sufficiently large flat wall of a boundaried graph  $\mathbf{G} = (G, B, \rho)$  a flat subwall whose compass is “irrelevant” with respect to the presence of a graph  $H$  in  $\mathcal{F}$  as a minor. Here the term “irrelevant” is not only related to  $G$  but to every graph  $\mathbf{K} \oplus \mathbf{G}$  that can be obtained by gluing  $\mathbf{G}$  with another boundaried graph  $\mathbf{K}$ . For this we need a stronger notion of irrelevancy, defined in [Subsection 5.4](#), that takes into account only the “essential part” of a topological minor model  $(M, T)$  of  $H$  that is “invading”  $\mathbf{G}$ .

We start in [Subsection 5.1](#), by detecting in every wall a *railed annulus*. This structure, introduced in [\[28\]](#) and reused later in [\[24, 25\]](#), turns out to be quite handy in order to guarantee a “taming property” of topological minor models (cf. [Proposition 13](#)). In [Subsection 5.2](#) we first use graph drawing tools to prove that we can assume that our model is embedded “nicely” inside a railed annulus, in the sense that certain vertices are sufficiently pairwise far apart (cf. [Lemma 16](#)); this will be helpful in order to reroute the model of every “invading” topological minor model  $(M, T)$  of  $H$ . With the help of [Proposition 13](#), we prove (cf. [Theorem 17](#)) that, given a partially disk-embedded graph that contains a railed annulus, the topological minor model  $(M, T)$  of a graph  $H$  can be rerouted so to obtain another topological minor model that can be contracted back to  $H$  and such that a “large enough” central region of the railed annulus is avoided. The rerouting of  $(M, T)$  will be done so that a prescribed subset of degree-3 vertices of the original model will not be affected by contractions.

Once we have all the above ingredients, we consider in [Subsection 5.4](#) a boundaried graph  $\mathbf{G} = (G, B, \rho)$  and an apex wall triple  $(A, W, \mathfrak{A})$  that is not affected by  $B$  and we show, in [Theorem 23](#), how every topological minor model  $(M, T)$  of a graph  $H$  in  $G$  can be rerouted away from the compass of the central subwall  $W'$  of  $W$ . This will permit us later to declare the whole compass of  $W'$  *irrelevant* and rule out the possibility that  $W$  has size exceeding some function depending on the “intrusion” of  $H$  in  $\mathbf{G}$ . The proof of [Theorem 23](#) is the most technical part of this paper. For this, we define an appropriate “flat” representation of the  $\mathfrak{A}$ -compass of  $W$ , called its *leveling*, and a representation of the wall  $W$  in the leveling that is “well-aligned”. This well-alignment property, defined in [Subsection 5.3](#), emerges from the regularity of the flatness pair  $(W, \mathfrak{A})$  and permits the representation of  $(M, T)$  by a topological minor model  $(\tilde{M}, \tilde{T})$  of the leveling, accompanied with a suitable encoding of the parts of  $(M, T)$  that have been suppressed by the leveling. This will permit us to obtain, using [Theorem 17](#), a rerouting  $(\hat{M}, \hat{T})$  of  $(\tilde{M}, \tilde{T})$  inside the leveling. Finally, using the homogeneity property, we will translate back  $(\hat{M}, \hat{T})$  to a rerouting of  $(M, T)$  that will avoid the compass of the central subwall  $W'$ .

### 5.1 A lemma for model taming

We introduce the concept of a railed annulus and present the main combinatorial result of [\[25\]](#).

**Railed annuli.** Let  $G$  be a partially  $\Delta$ -embedded graph and let  $\mathcal{C} = [C_1, \dots, C_r]$ ,  $r \geq 2$ , be a collection of vertex-disjoint cycles of the compass of  $G$ . We say that the sequence  $\mathcal{C}$  is a  *$\Delta$ -nested sequence of cycles* of  $G$  if every  $C_i$  is the boundary of an open disk  $D_i$  of  $\Delta$  such that  $\Delta \supseteq D_1 \supseteq \dots \supseteq D_r$ . From now on, each  $\Delta$ -nested sequence  $\mathcal{C} = [C_1, \dots, C_r]$  will be accompanied with the sequence  $[D_1, \dots, D_r]$  of the corresponding open disks as well as the sequence  $[\overline{D}_1, \dots, \overline{D}_r]$  of their closures. Given  $x, y \in [r]$  with  $x \leq y$ , we call the set




Figure 6: An example of a  $(5, 8)$ -railed annulus and its inner disk  $D_5$ .

$\overline{D}_x \setminus D_y$   $(x, y)$ -annulus of  $\mathcal{C}$  and we denote it by  $\text{ann}(\mathcal{C}, x, y)$ . Finally we say that  $\text{ann}(\mathcal{C}, 1, r)$  is the *annulus* of  $\mathcal{C}$  and we denote it by  $\text{ann}(\mathcal{C})$ .

Let  $r \in \mathbb{N}_{\geq 3}$  and  $q \in \mathbb{N}_{\geq 3}$  with  $r$  odd. An  $(r, q)$ -railed annulus of a  $\Delta$ -partially-embedded graph  $G$  is a pair  $\mathcal{A} = (\mathcal{C}, \mathcal{P})$  where  $\mathcal{C} = [C_1, \dots, C_r]$  is a  $\Delta$ -nested collection of cycles of  $G$  and  $\mathcal{P} = [P_1, \dots, P_q]$  is a collection of pairwise vertex-disjoint paths in  $G$ , called *rails*, such that

- for every  $j \in [q]$ ,  $P_j \subseteq \text{ann}(\mathcal{C})$ , and
- for every  $(i, j) \in [r] \times [q]$ ,  $C_i \cap P_j$  is a non-empty path that we denote by  $P_{i,j}$ .

See Figure 6 for an example of a  $(5, 8)$ -railed annulus. The following proposition states that large railed annuli can be found inside a modestly larger wall and will be used in the next section. A similar (but less precise) statement can be found in [28].

**Proposition 12.** *If  $x, z \geq 3$  are odd integers,  $y \geq 1$ , and  $W$  is an  $\text{odd}(2x + \max\{z, \frac{y}{4} - 1\})$ -wall, then*

- *there is a collection  $\mathcal{P}$  of  $y$  paths in  $W$  such that if  $\mathcal{C}$  is the collection of the first  $x$  layers of  $W$ , then  $(\mathcal{C}, \mathcal{P})$  is an  $(x, y)$ -railed annulus of  $W$  where the first cycle of  $\mathcal{C}$  is the perimeter of  $W$ , and*
- *the open disk defined by the  $x$ -th cycle of  $\mathcal{C}$  contains the vertices of the compass of the central  $z$ -subwall of  $W$ .*

*Proof.* Let  $\mathcal{C} = \{C_1, \dots, C_x\}$  be the collection of the first  $x$  layers of  $W$ . Notice that the open disk defined by  $C_x$  contains the vertices of the compass of the central  $w'$ -subwall, where  $w' := w - 2x = \text{odd}(\max\{z, \frac{y}{4} - 1\}) \geq z$ . On the other hand,  $W$  contains a collection  $\mathcal{P}$  of at least  $2w' + 2(w' + 2)$  pairwise vertex-disjoint paths from  $C_1$  to  $C_x$ . Since  $2w' + 2(w' + 2) = 4w' + 4 \geq y$ , the pair  $(\mathcal{C}, \mathcal{P})$  is an  $(x, y)$ -railed annulus of  $W$  where the first cycle of  $\mathcal{C}$  is the perimeter of  $W$  (see Figure 7).  $\square$

We define the *annulus* of  $\mathcal{A} = (\mathcal{C}, \mathcal{P})$  as the annulus of  $\mathcal{C}$ . We call  $C_1$  and  $C_r$  the *outer* and the *inner* cycle of  $\mathcal{A}$ , respectively. Also, if  $(i, i') \in [r]^2$  with  $i < i'$  then we define  $\mathcal{A}_{i,i'} = ([C_i, \dots, C_{i'}], \mathcal{P} \cap \text{ann}(\mathcal{C}, i, i'))$ .




Figure 7: A visualization of the proof of [Proposition 12](#) in the 33-wall where  $x = 9$ ,  $y = 64$ , and  $z = 15$ . For simplicity, the paths in  $\{P_{i,j} \mid (i,j) \in [33]^2\}$  are depicted as vertices.

The *union-graph* of an  $(r, q)$ -railed annulus  $\mathcal{A} = (\mathcal{C}, \mathcal{P})$  is defined as  $G(\mathcal{A}) := (\bigcup_{i \in [r]} C_i) \cup (\bigcup_{i \in [q]} P_i)$ . Clearly,  $G(\mathcal{A})$  is a planar graph and we always assume that its infinite face is the one whose boundary is the first cycle of  $\mathcal{C}$ .

Let  $\mathcal{A}$  be a  $(r, q)$ -railed annulus of a partially  $\Delta$ -embedded graph  $G$ . Let  $r = 2t + 1$ , for some  $t \geq 0$ . Let also  $s \in [r]$  where  $s = 2t' + 1$ , for some  $0 \leq t' \leq t$ . Given some  $I \subseteq [q]$ , we say that a subgraph  $M$  of  $G$  is  $(s, I)$ -confined in  $\mathcal{A}$  if

$$M \cap \text{ann}(\mathcal{C}, t - t', t + t') \subseteq \bigcup_{i \in I} P_i.$$

The following proposition has been recently proved by Golovach et al. [[25](#), Theorem 2.1], where it has been dubbed as the “Model Taming Lemma”.

**Proposition 13.** *There exist two functions  $f_6, f_7 : \mathbb{N}_{\geq 0} \rightarrow \mathbb{N}_{\geq 0}$  such that the images of  $f_7$  are even and such that if*

- $s$  is a positive odd integer,
- $H$  is a graph on at most  $\ell$  edges,
- $G$  is a  $\Delta$ -partially-embedded graph,
- $\mathcal{A} = (\mathcal{C}, \mathcal{P})$  is an  $(r, q)$ -railed annulus of  $G$ , where  $r = f_7(\ell) + 2 + s$  and  $q \geq 5/2 \cdot f_6(\ell)$ ,
- $(M, T)$  is a topological minor model of  $H$  in  $G$  such that  $T \cap \text{ann}(\mathcal{A}) = \emptyset$ , and
- $I \subseteq [q]$  where  $|I| > f_6(\ell)$ ,

then  $G$  contains a topological minor model  $(\tilde{M}, \tilde{T})$  of  $H$  in  $G$  such that

1.  $\tilde{T} = T$ ,
2.  $\tilde{M}$  is  $(s, I)$ -confined in  $\mathcal{A}$ , and
3.  $\tilde{M} \setminus \text{ann}(\mathcal{A}) \subseteq M \setminus \text{ann}(\mathcal{A})$ .

Moreover  $f_7(\ell) = \mathcal{O}((f_6(\ell))^2)$ .

*Remark 14.* It is worth mentioning here that the function  $f_6(\ell)$  of [Proposition 13](#) depends on the constants involved in the Unique Linkage Theorem [\[31, 43\]](#); see [Subsection B.2](#) for a more detailed discussion. At this point we just remark that, according to the results of Adler and Krause [\[2\]](#), we have that  $f_6(\ell) = 2^{\Omega(\ell)}$ . This permits us to henceforth make the (generous) assumption that  $\ell = \mathcal{O}(f_6(\ell))$ .

## 5.2 Model rerouting in partially disk-embedded graphs

Using classic results on how to optimally draw planar graphs of maximum degree three into grids (see e.g., [\[29\]](#)) one may easily derive the following.

**Proposition 15.** *There exists a function  $f_8 : \mathbb{N} \rightarrow \mathbb{N}$  such that for every  $\ell$ -vertex planar graph  $H$  with maximum degree three there is a tm-pair  $(M, T)$  of the  $(f_8(\ell) \times f_8(\ell))$ -grid, denoted by  $\Gamma$ , that is a topological minor model of  $H$  in  $\Gamma$ . Moreover, it holds that  $f_8(\ell) = \mathcal{O}(\ell)$ .*

Let  $\Gamma$  be an  $(r \times r)$ -grid for some  $r \geq 3$ . We see a  $\Gamma$ -grid as the union of  $r$  horizontal paths and  $r$  vertical paths. Given an  $i \in \lfloor \frac{r}{2} \rfloor$ , we define the  $i$ -th layer of  $\Gamma$  recursively as follows: the first layer of  $\Gamma$  is its perimeter, while, if  $i \geq 2$ , the  $i$ -th layer of  $\Gamma$  is the perimeter of the  $(r - 2(i - 1) \times r - 2(i - 1))$ -grid created if we remove from  $\Gamma$  its  $i - 1$  first layers. When we deal with a  $(r \times r)$ -grid  $\Gamma$ , we always consider its embedding where the infinite face is bounded by the first layer of  $\Gamma$ .

**Safely arranged models.** Let  $G$  be a plane graph. Given two subgraphs of  $G$ , we define their *face-distance* as the minimum face-distance between two of their vertices. We denote by  $\mathbf{F}_G^{(i)}(x)$  the set of all vertices of  $G$  that are within face-distance at most  $i$  from vertex  $x$ .

Given a  $c \geq 0$  and a tm-pair  $(M, T)$  of  $G$ , we say that  $(M, T)$  is *safely  $c$ -dispersed* in  $G$  if

- every two distinct vertices  $t, t' \in T$  are within face-distance at least  $2c + 1$  in  $G$ , and
- for every  $t \in T$  of degree  $d$  in  $M$ , the graph  $M[\mathbf{F}_G^{(c)}(t) \cap V(M)]$  consists of  $d$  paths with  $t$  as a unique common endpoint.

With [Proposition 15](#) at hand, we can prove the following useful lemma.

**Lemma 16.** *There exists a function  $f_9 : \mathbb{N}^3 \rightarrow \mathbb{N}$  such that the following holds. Let  $c, r, r', \ell \in \mathbb{N}$ ,  $r' \leq r$ ,  $H$  be a  $D$ -embedded  $(\ell + r')$ -vertex graph, and  $Z := \{z_1, \dots, z_{r'}\} \subseteq V(H)$  such that*

- *the vertices of  $H$  have degree at most three,*
- *$Z$  is an independent set of  $H$ ,*
- *all vertices of  $Z$  have degree one in  $H$ ,*
- *$\text{bd}(D) \cap H = Z$ , and*
- *$(z_1, \dots, z_{r'})$  is the cyclic ordering of the vertices of  $Z$  as they appear in the boundary of  $D$ .*

*Let also  $G$  be a  $\Delta$ -embedded graph,  $\mathcal{A} = (\mathcal{C}, \mathcal{P})$  be an  $(x, y)$ -railed annulus of  $G$ , where  $x, y$  are integers such that  $\min\{x, y\} \geq f_9(c, r, \ell) \geq r$ , and where  $\mathcal{C} = [C_1, \dots, C_x]$  and  $\mathcal{P} = [P_1, \dots, P_y]$ ,  $w_i$  be the endpoint of  $P_i$  that is contained in  $C_1$ , for  $i \in [r]$ , and  $I := \{i_1, \dots, i_{r'}\} \subseteq [r]$ . Then the union-graph  $G(\mathcal{A})$  of  $\mathcal{A}$  contains a tm-pair  $(M, T)$  that is a topological minor model of  $H$  in  $G(\mathcal{A})$  such that*

- *for each  $j \in [r']$ ,  $\sigma_{M, T}(z_j) = w_{i_j}$ ,*
- *the tm-pair  $(M, T)$  is safely  $c$ -dispersed in the union graph  $G(\mathcal{A})$ , and*

- none of the vertices of  $T \setminus \{w_{i_1}, \dots, w_{i_{r'}}\}$  is within face-distance less than  $c$  from some vertex in  $C_1$  or in  $C_r$ .

Moreover, it holds that  $f_9(c, r, \ell) = \mathcal{O}(cr(\ell + r))$ .

*Proof.* Using  $H$ , we construct a new graph  $H'$  as follows: consider a copy  $\tilde{H}$  of  $H$  where the copy of  $z_i$  in  $\tilde{H}$  is denoted by  $\tilde{z}_i$ , for each  $i \in [r']$ . We take the disjoint union of  $H$  and  $\tilde{H}$ , add the edges  $\{z_1, z_2\}, \dots, \{z_{r'}, z_1\}$ , forming a cycle  $C$ , subdivide the edges in  $C$ , add the edges  $\{\tilde{z}_1, \tilde{z}_2\}, \dots, \{\tilde{z}_r, \tilde{z}_1\}$ , forming a cycle  $\tilde{C}$ , subdivide the edges in  $\tilde{C}$ , and, given that, for  $i \in [r']$ ,  $x_i$  (resp.  $\tilde{x}_i$ ) is the vertex created after the subdivision of  $\{z_i, z_{i+1}\}$  (resp.  $\{\tilde{z}_i, \tilde{z}_{i+1}\}$ ) (here  $r' + 1$  is interpreted as 1), add the edges  $\{x_1, \tilde{x}_1\}, \dots, \{x_r, \tilde{x}_r\}$ . The resulting graph  $H'$  has  $2(\ell + r') \leq 2(\ell + r)$  vertices, is planar, and has maximum degree three. Let  $s = f_8(2(\ell + r))$ . By Proposition 15, there is a tm-pair  $(M', T')$  of the  $(s \times s)$ -grid  $\Gamma$  that is a topological minor model of  $H'$  in  $\Gamma$ . We now subdivide  $\bar{r} = 2(c + 1)r$  times each of the edges of  $\Gamma$  and see the resulting graph  $\Gamma'$  as a subgraph of a  $((\bar{r} + 1)(s + 2) \times (\bar{r} + 1)(s + 2))$ -grid  $\Gamma''$  in a way that none of the  $\bar{r}$  first layers of  $\Gamma''$  intersects  $\Gamma'$ . By also subdividing  $\bar{r}$  times each of the edges of  $M'$  we construct a tm-pair  $(M'', T')$  of  $\Gamma''$  that is a topological minor model of  $H'$  in  $\Gamma''$ .

Let  $w_1, \dots, w_r$  be the first  $r$  vertices of the lower path of  $\Gamma''$ . Recall that  $H$  is a subgraph of  $H'$ , therefore we can define  $M = \sigma_{M', T'}(H)$ . Let also  $T = \sigma_{M', T'}(V(H))$ . Notice that  $(M, T)$  is a tm-pair of  $\Gamma''$  that is a topological minor model of  $H$ . Let  $\hat{z}_i = \sigma_{M, T}(z_i)$ ,  $i \in [r']$ . We make two observations about the position of these vertices in  $\Gamma''$ . The first is that, because of the construction of  $H'$ ,  $\hat{z}_1, \dots, \hat{z}_{r'}$  appear, in this ordering, on a cycle of  $\Gamma'$  bounding a closed disk, say  $\Delta$ , that contains the whole  $M$ . The second is that, as  $M$  is a subgraph of  $\Gamma'$ , each pair  $\hat{z}_i, \hat{z}_j$ ,  $i \neq j$ , is at distance at least  $\bar{r} + 1$  in the graph  $\Gamma''' := \Gamma'' \setminus \text{int}(\Delta)$ . It is now easy to observe that the two previous observations permit to find in  $\Gamma'''$  pairwise disjoint paths joining  $\hat{z}_j$  with  $w_{i_j}$ , for  $j \in [r']$ . By adding these paths in  $M$  and including in  $T$  the set  $\{w_{i_1}, \dots, w_{i_{r'}}\}$ , we construct a tm-pair  $(M, T)$  that is a topological minor model of  $H$  in  $\Gamma'$  such that for each  $j \in [r']$ , the function  $\sigma_{M, T}$  maps the vertex  $z_j$  to  $w'_{i_j}$  and the intersection of  $V(M)$  and the upper path of  $\Gamma''$  is empty. Moreover, as we applied at least  $\bar{r} = (c + 2)r$  subdivisions, it also holds that the set  $T$  is safely  $2c$ -dispersed in  $\Gamma''$ . Moreover, it is easy to observe that none of the vertices of  $T \setminus \{w_{i_1}, \dots, w_{i_{r'}}\}$  is within face-distance less than  $c$  from some vertex in the perimeter of  $\Gamma''$ .

Consider now a  $(x, y)$ -railed annulus  $(\mathcal{C}, \mathcal{P})$  of some  $\Delta$ -embedded graph  $G$ , with  $\min\{x, y\} \geq q$ , where  $q = (\bar{r} + 1)(s + 2)$ . Let  $\mathcal{C} = [C_1, \dots, C_x]$  and  $\mathcal{P} = [P_1, \dots, P_y]$  as in the statement of the lemma. Let also  $\tilde{\Gamma} = G(\mathcal{A})$ . For every  $i \in [q]$ , we define  $F_{\mathcal{A}}^{(i)}$  as the edge set of the unique path in  $C_i$  with one endpoint in  $P_{i, q}$  and the other in  $P_{i, 1}$ , that does not contain internal vertices of the paths  $P_{i, q}$  or  $P_{i, 1}$ , and does not contain any vertex from  $P_2$ . We denote by  $F_{\mathcal{A}}^e$  (resp.  $F_{\mathcal{A}}^v$ ) the set of all edges (resp. internal vertices) of the paths  $F_{\mathcal{A}}^{(i)}$ ,  $i \in [q]$ . Notice that the grid  $\Gamma''$  occurs from  $(\tilde{\Gamma} \setminus F_{\mathcal{A}}^e) \setminus F_{\mathcal{A}}^v$  if, for every  $(i, j) \in [q]^2$ , we contract the path  $P_{i, j}$  defined by the intersection of the  $i$ -th horizontal path and the  $j$ -th vertical path of  $W$ . It is easy to see that if in  $\tilde{\Gamma}$  we uncontract each vertex, say  $(i, j)$  of  $M$  to the path  $P_{i, j}$ , one can transform  $(M, T)$  to a tm-pair of  $W$  that is a topological minor model of  $H$  in  $W$  and additionally, for each  $i \in [r']$ , the function  $\sigma_{M, T}$  maps the vertex  $z_j$  to  $w_{i_j}$ . This implies the first condition of the lemma. The second condition follows directly from the fact that the pair  $(M, T)$  was already safely  $2c$ -dispersed before applying the uncontractions and such uncontractions cannot reduce the distance to more than half of it. The third condition is also an obvious consequence of the uncontraction procedure. Therefore, the lemma holds if we set  $f_9(c, r, \ell) := q = \mathcal{O}(cr(\ell + r))$ .  $\square$

Let  $G$  be a partially  $\Delta$ -embedded graph and let  $\mathcal{C} = [C_1, \dots, C_r]$  be a  $\Delta$ -nested sequence of cycles of  $G$  and let  $[D_1, \dots, D_r]$  (resp.  $[\bar{D}_1, \dots, \bar{D}_r]$ ) be the sequences of the corresponding open (resp. closed) disks.

Let also  $(M, T)$  be a tm-pair of  $G$  and  $p \in [r]$ . We define the  $p$ -crop of  $(M, T)$  in  $\mathcal{C}$ , denoted by  $(M, T) \cap \bar{D}_p$ , as the tm-pair  $(M', T')$  where  $M' = M \cap \bar{D}_p$  and  $T' = (T \cap \bar{D}_p) \cup (V(C_p \cap M))$ .

Given a graph  $H$  a set  $Q \subseteq V(H)$  and a graph  $G$ , we say that  $\phi : V(H) \rightarrow 2^{V(G)}$  is a  $Q$ -respecting contraction-mapping of  $H$  to  $G$  if

- $\bigcup_{x \in V(H)} \phi(x) = V(G)$ ,
- $\forall x, y \in V(H)$ , if  $x \neq y$  then  $\phi(x) \cap \phi(y) = \emptyset$ ,
- $\forall x \in V(H)$ ,  $G[\phi(x)]$  is connected,
- $\forall \{x, y\} \in E(H)$ ,  $G[\phi(x) \cup \phi(y)]$  is connected, and
- $\forall x \in Q$ ,  $|\phi(x)| = 1$ .

The critical point in the above definitions is that vertices in  $Q$  are not “uncontracted” when transforming  $H$  to  $G$ .

**Intrusion of a topological minor model.** Let  $G$  be a graph, let  $S \subseteq V(G)$ , and let  $(M, T)$  be a **tm**-pair of  $G$ . We define the  $S$ -intrusion of  $(M, T)$  in  $G$  as the maximum value between  $|S \cap T|$  and the number of subdivision paths of  $(M, T)$  that contain vertices of  $S$ . It is important to notice that  $S$  can intersect many times a subdivision path of  $(M, T)$ , however the value of the  $S$ -intrusion counts each such a path only once.

Using [Proposition 12](#), [Proposition 13](#), and [Lemma 16](#) we prove the following.

**Theorem 17.** *There exist three functions  $f_{10} : \mathbb{N}^2 \rightarrow \mathbb{N}$ ,  $f_{11} : \mathbb{N}^2 \rightarrow \mathbb{N}$ , and  $f_{12} : \mathbb{N}^3 \rightarrow \mathbb{N}$  such that the following holds. Let  $c, \ell \in \mathbb{N}$ ,  $z \geq 3$  be an odd integer, and  $G$  be a partially  $\Delta$ -embedded graph, whose compass contains an  $f_{12}(c, z, \ell)$ -wall  $W$  with  $\text{bd}(\Delta)$  as perimeter. Let also  $\mathcal{C} = [C_1, \dots, C_{f_{10}(c, \ell)}]$  be the first  $f_{10}(c, \ell)$ -layers of  $W$  and  $D_1, \dots, D_{f_{10}(c, \ell)}$  be the open disks of  $\Delta$  that they define. If  $(M, T)$  is a **tm**-pair of  $G$  whose  $\Delta \cap V(G)$ -intrusion in  $G$  is at most  $\ell$  and  $Q$  is a subset of  $T$  containing vertices of degree at most three in  $M$ , then there is a **tm**-pair  $(\hat{M}, \hat{T})$  of  $G$  and an integer  $b \in [f_{10}(c, \ell)]$  such that*

1.  $\hat{M} \setminus D_b$  is a subgraph of  $M \setminus D_b$ ,
2.  $\text{ann}(\mathcal{C}, b, b + f_{11}(c, \ell) - 1) \cap (T \cup \hat{T}) = \emptyset$ ,
3.  $(\hat{M}, \hat{T}) \cap \overline{D}_{b+f_{11}(c, \ell)}$  is a **tm**-pair of  $W$  that is safely  $c$ -dispersed in  $W$  and none of the vertices of  $\hat{T} \cap \overline{D}_{b+f_{11}(c, \ell)}$  is within face-distance less than  $c$  in  $W$  from some vertex of  $C_{b+f_{11}(c, \ell)} \cup C_{f_{10}(c, \ell)}$ ,
4.  $\hat{M} \cap D_{f_{10}(c, \ell)} = \emptyset$ ,
5. the compass of the central  $z$ -subwall of  $W$  is a subset of  $D_{f_{10}(c, \ell)}$ , and
6. there is a  $Q$ -respecting contraction-mapping of  $\text{diss}(M, T)$  to  $\text{diss}(\hat{M}, \hat{T})$ .

Moreover, it holds that  $f_{10}(c, \ell) = \mathcal{O}(c \cdot (f_6(\ell))^3)$ ,  $f_{11}(c, \ell) = \mathcal{O}(c \cdot (f_6(\ell))^2)$ , and  $f_{12}(c, z, \ell) = \mathcal{O}(c \cdot (f_6(\ell))^3 + z)$ .

See [Figure 8](#) for an illustration of the conditions guaranteed by [Theorem 17](#).

*Proof.* Let  $r = f_6(\ell) + 1$ ,  $s = \text{odd}(f_9(c, r, 3\ell + r))$ ,  $x' = \text{odd}(f_7(\ell) + 2 + s)$ ,  $y = \max\{s, \lceil 5/2 \cdot f_6(\ell) \rceil\}$ ,  $x = \text{odd}((\ell + 1) \cdot x')$ , and  $w = \text{odd}(2x + \max\{z, \frac{y}{4} - 1\})$ . We will prove the theorem for  $f_{10}(c, \ell) = x$ ,  $f_{11}(\ell) = \frac{x' - s}{2}$ , and  $f_{12}(c, z, \ell) = w$ . Let  $G$  be a partially  $\Delta$ -embedded graph, whose compass contains a  $w$ -wall  $W$  with  $\text{bd}(\Delta)$  as perimeter. Let also  $\mathcal{C} = [C_1, \dots, C_x]$  be the first  $x$  layers of  $W$  and let  $[D_1, \dots, D_x]$  (resp.  $[\overline{D}_1, \dots, \overline{D}_x]$ ) be the sequences of the corresponding open (resp. closed) disks of  $\Delta$  bounded by the cycles in  $\mathcal{C}$ . From [Proposition 12](#) there is a collection  $\mathcal{P} = \{P_1, \dots, P_y\}$  of paths in  $W$  such that  $\mathcal{A} = (\mathcal{C}, \mathcal{P})$




Figure 8: A visualization of how a tm-pair  $(M, T)$  is rearranged to a new tm-pair  $(\hat{M}, \hat{T})$  as in [Theorem 17](#). The figure depicts in red the part of the tm-pair  $(\hat{M}, \hat{T})$  that intersects the disk  $\Delta$ . The cycles correspond to the first  $f_{10}(c, \ell)$  layers of  $W$ . The black vertices are the vertices in  $Q$ , while the circled vertices inside the turquoise area are the “new” branch vertices of  $\tilde{T}$  that are vertices of  $W$ . The “green clouds” are the non-singleton images of the  $Q$ -respecting contraction-mapping of  $\text{diss}(M, T)$  to  $\text{diss}(\hat{M}, \hat{T})$ . We stress that in this picture, the way the model enters the turquoise area does not reflect the fact  $\tilde{M}$  that is  $(s, I)$ -confined in  $\mathcal{A}'$ , as it is argued in the proof. We opted not to reflect this fact in the figure as we prioritized the visualization of other, more important, aspects of the proof.

is an  $(x, y)$ -railed annulus of  $W$  where the outer cycle of  $C$  is the perimeter of  $W$  and such that the vertices of the compass of the central  $z$ -subwall of  $W$  belong to  $D_x$ , and Property 5 follows.

Let  $\check{M}$  be the union of all subdivision paths of  $(M, T)$  that intersect  $\Delta \cap V(G)$  and let  $\check{T}$  be the endpoints of these paths. Moreover, we denote  $\check{H} = \text{diss}(\check{M}, \check{T})$  and observe that  $\check{H}$  is a subgraph of  $H$ . Intuitively,  $\check{H}$  is the subgraph of  $H$  whose topological minor model  $(\check{M}, \check{T})$  is the part of  $(M, T)$  that intersects the closed disk  $\Delta$ . As the  $\Delta \cap V(G)$ -intrusion of  $(M, T)$  in  $G$  is at most  $\ell$ , the same bound applies to the  $\Delta \cap V(G)$ -intrusion of  $(\check{M}, \check{T})$  in  $G$ . This in turn implies that  $|\check{T} \cap \Delta| \leq \ell$  and that  $|E(\check{H})| \leq \ell$ .

Since,  $x = (\ell + 1) \cdot x'$ , there is a  $b \leq \ell \cdot x' + 1 \leq x$  such that  $\mathbb{A} := \text{ann}(C, b, b + x' - 1)$  does not contain any vertex of  $T$ . We define  $T^{\text{out}} = \check{T} \setminus \overline{D}_b$  and  $T^{\text{in}} = \check{T} \cap D_{b+x'-1}$ . Clearly,  $\{T^{\text{out}}, T^{\text{in}}\}$  is a partition of  $\check{T}$ .

We set  $\mathcal{A}' = ([C_b, \dots, C_{b+x'-1}], \mathcal{P} \cap \mathbb{A})$ . By applying [Proposition 13](#) on  $s, \check{H}, g := \ell$ , the  $\Delta$ -bordered graph  $G$ , the  $(x', y)$ -railed annulus  $\mathcal{A}'$ , the tm-pair  $(\check{M}, \check{T})$ , and the set  $I = [r]$ , we have that  $G$  contains a

topological minor model  $(\tilde{M}, \tilde{T})$  of  $\tilde{H}$  in  $G$  such that  $\tilde{M}$  is  $(s, I)$ -confined in  $\mathcal{A}'$  and  $\tilde{M} \setminus \text{ann}(\mathcal{A}') \subseteq \tilde{M} \setminus \text{ann}(\mathcal{A}')$ . We enhance  $\tilde{M}$  by adding to it all subdivision paths of  $(M, T)$  that are not intersecting  $\Delta$ . That way, we have that  $(\tilde{M}, T)$  is a topological minor model of  $H$  in  $G$  such that  $\tilde{M}$  is  $(s, I)$ -confined in  $\mathcal{A}'$  and  $\tilde{M} \setminus \text{ann}(\mathcal{A}') \subseteq M \setminus \text{ann}(\mathcal{A}')$ .

Let  $p = b + \frac{x'-s}{2}$  and  $q = b + \frac{x'+s}{2} - 1$  and notice that  $q \leq x$ . We set  $\mathbb{A}' := \text{ann}(\mathcal{C}, p, q)$  and we define  $\mathcal{A}'' := ([C_p, \dots, C_q], \mathcal{P}')$  where  $\mathcal{P}' = \mathcal{P} \cap \mathbb{A}'$ . Let  $\mathcal{P}' = \{P'_1, \dots, P'_y\}$ . Observe that, from the second property of [Proposition 13](#), the connected components of  $\tilde{M} \cap \mathbb{A}'$  are some of the first  $r$  paths in  $\mathcal{P}'$ . This means that there is a subset of indices  $\{i_1, \dots, i_{r'}\} \subseteq I$  such that  $\tilde{M} \cap \mathbb{A}' = P'_{i_1} \cup \dots \cup P'_{i_{r'}}$ . Let  $Z = \{z_{i_1}, \dots, z_{i_{r'}}\}$  be the set of endpoints of the paths  $P'_{i_1}, \dots, P'_{i_{r'}}$  that are contained in  $C_p$ .

Let  $\tilde{M}^{\text{in}} = \tilde{M} \cap \overline{D}_p$ ,  $\tilde{M}^{\text{out}} = (\tilde{M} \setminus D_p) \setminus E(C_p)$ , and observe that  $\tilde{M} = \tilde{M}^{\text{in}} \cup \tilde{M}^{\text{out}}$  and that  $Z = V(\tilde{M}^{\text{in}}) \cap V(\tilde{M}^{\text{out}})$ . Moreover, all vertices of  $Z$  have degree one in both  $\tilde{M}^{\text{in}}$  and  $\tilde{M}^{\text{out}}$ . Let  $\tilde{H}^{\text{in}}$  (resp.  $\tilde{H}^{\text{out}}$ ) be the graph obtained from  $\tilde{M}^{\text{in}}$  (resp.  $\tilde{M}^{\text{out}}$ ) by dissolving all vertices except from those in  $T^{\text{in}} \cup Z$  (resp.  $T^{\text{out}} \cup Z$ ). Note that  $(\tilde{M}^{\text{in}}, T^{\text{in}} \cup Z)$  (resp.  $(\tilde{M}^{\text{out}}, T^{\text{out}} \cup Z)$ ) is a topological minor model of  $\tilde{H}^{\text{in}}$  (resp.  $\tilde{H}^{\text{out}}$ ).

Notice that  $\tilde{H}^{\text{in}}$  has vertex set  $T^{\text{in}} \cup Z$  and can be seen as a  $D$ -embedded graph, for some closed disk  $D$ , on at most  $\ell + r$  edges where  $\text{bd}(D) \cap H = Z$  and  $(z_{i_1}, \dots, z_{i_{r'}})$  is the ordering of the vertices of  $Z$  as they appear in  $C_p$ . Observe now that  $\tilde{H}^{\text{in}}$  can be seen as the contraction of another  $D$ -embedded graph  $\hat{H}^{\text{in}}$  with detail at most  $3\ell + r$  that has maximum degree at most three. Moreover, we can assume that the vertices of  $\tilde{H}^{\text{in}}$  that have degree at most three are also vertices of  $\hat{H}^{\text{in}}$  that are not affected by the contractions while transforming  $\hat{H}^{\text{out}}$  to  $\tilde{H}^{\text{out}}$ . This implies that there is a  $Q$ -respecting contraction-mapping of  $\tilde{H}^{\text{out}}$  to  $\hat{H}^{\text{out}}$ . Again, in the embedding of  $\hat{H}^{\text{in}}$  in  $D$ ,  $(z_{i_1}, \dots, z_{i_{r'}})$  is the ordering of the vertices of  $Z$  as they appear in  $\text{bd}(D)$ .

Keep in mind that  $\tilde{H}^+ := \tilde{H}^{\text{out}} \cup \tilde{H}^{\text{in}}$  is a minor of  $\hat{H}^+ := \hat{H}^{\text{out}} \cup \hat{H}^{\text{in}}$  and that, if we dissolve in  $\tilde{H}^+$  all the vertices in  $Z$ , we obtain  $H$ . Also let  $\hat{H}$  be the graph obtained if we dissolve in  $\hat{H}^+$  all the vertices in  $Z$ . Clearly,  $\hat{H}$  is a minor of  $H$ .

We now apply [Lemma 16](#) for  $c, r, r', 3\ell$ , the  $D$ -embedded graph  $\hat{H}^{\text{in}}$ , the set  $Z$ , and the  $(s, y)$ -railed annulus  $\mathcal{A}''$  of the  $\overline{D}_p$ -disk embedded graph  $G \cap \overline{D}_p$  and obtain a tm-pair  $(\hat{M}^{\text{in}}, \hat{T}^{\text{in}})$  of  $G(\mathcal{A}'')$  that is a topological minor model of  $\hat{H}^{\text{in}}$  and such that for each  $j \in [r']$ , the function  $\sigma_{\hat{M}^{\text{in}}, \hat{T}^{\text{in}}}$  maps vertex  $z_{i_j}$  to itself. Notice that  $G(\mathcal{A}'')$  is a subgraph of  $W \cap \text{ann}(\mathcal{C}, p, q)$ . From the second property of [Lemma 16](#),  $(\hat{M}^{\text{in}}, \hat{T}^{\text{in}})$  is safely  $c$ -dispersed in  $W \cap \text{ann}(\mathcal{C}, p, q)$ . From the third property of [Lemma 16](#), it follows that none of the vertices of  $\hat{T}^{\text{in}} \setminus \{w_{i_1}, \dots, w_{i_{r'}}\}$  is within face-distance less than  $c$  from some vertex of  $C_p \cup C_q$  in  $W \cap \text{ann}(\mathcal{C}, p, q)$ .

We now consider the graph  $\hat{M} = \hat{M}^{\text{in}} \cup \tilde{M}^{\text{out}}$ . Property 3 follows by the conclusions of the previous paragraph. Moreover,  $\hat{M}$  does not intersect  $D_q$  and, as  $q \leq x$ , it neither intersects  $D_x$ , hence Property 4 holds. Notice also that  $\tilde{M} \setminus \text{ann}(\mathcal{A}') \subseteq M \setminus \text{ann}(\mathcal{A}')$  implies  $\tilde{M} \setminus \overline{D}_b \subseteq M \setminus \overline{D}_b$ . This along with the fact that  $\hat{M} \setminus \overline{D}_b = \tilde{M} \setminus \overline{D}_b$ , yield Property 1.

Observe that  $(\hat{M}, \hat{T}^{\text{in}} \cup T^{\text{out}} \cup Z)$  is a topological minor model of  $\hat{H}^+$ , which in turn implies that  $(\hat{M}, \hat{T}^{\text{in}} \cup T^{\text{out}})$  is a topological minor model of  $\hat{H}$ . We now set  $\hat{T} = \hat{T}^{\text{in}} \cup T^{\text{out}}$ . As there is a  $Q$ -respecting contraction-mapping of  $\tilde{H}^{\text{out}}$  to  $\hat{H}^{\text{out}}$ , we also have that there is a  $Q$ -respecting contraction-mapping of  $H = \text{diss}(M, T)$  to  $\hat{H} = \text{diss}(\hat{M}, \hat{T})$  and Property 6 holds. As  $\hat{T}^{\text{in}} \subseteq \text{int}(\text{ann}(\mathcal{A}'')) \subseteq D_p = D_{b+f_{11}(\ell)}$  and  $T^{\text{out}} \subseteq G \setminus \overline{D}_b$ , we deduce that  $\hat{T} \in G \setminus \text{ann}(\mathcal{C}, b, b + f_{11}(\ell) - 1)$ , which together with the fact that  $\text{ann}(\mathcal{C}, b, b + x' - 1)$  does not contain any vertex of  $T$ , yield Property 2.




Figure 9: A graph  $G$  and a regular flatness pair  $(W, \mathfrak{R})$  of  $G$ .

To conclude the proof, let us provide upper bounds on the claimed functions. By definition, it holds that

$$\begin{aligned} f_{10}(c, \ell) &= \mathcal{O}(\ell \cdot (f_7(\ell) + f_9(c, f_6(\ell) + 1, 3\ell + f_6(\ell) + 1))), \\ f_{11}(c, \ell) &= \mathcal{O}(f_7(\ell) + f_9(c, f_6(\ell) + 1, 3\ell + f_6(\ell) + 1)), \text{ and} \\ f_{12}(c, z, \ell) &= \mathcal{O}(\ell \cdot (f_7(\ell) + f_9(c, f_6(\ell) + 1, 3\ell + f_6(\ell) + 1)) + \\ &\quad z + f_6(\ell) + f_9(c, f_6(\ell) + 1, 3\ell + f_6(\ell) + 1)). \end{aligned}$$

Since by [Proposition 13](#) we have that  $f_7(\ell) = \mathcal{O}((f_6(\ell))^2)$ , by [Lemma 16](#) we have that  $f_9(c, r, \ell) = \mathcal{O}(cr(\ell+r))$ , and by [Remark 14](#) we may assume that  $\ell = \mathcal{O}(f_6(\ell))$ , the above can be simplified to

$$\begin{aligned} f_{10}(c, \ell) &= \mathcal{O}(\ell \cdot ((f_6(\ell))^2 + c \cdot (f_6(\ell))^2)) \\ &= \mathcal{O}(c \cdot (f_6(\ell))^3), \\ f_{11}(c, \ell) &= \mathcal{O}(c \cdot (f_6(\ell))^2), \text{ and} \\ f_{12}(c, z, \ell) &= \mathcal{O}(\ell \cdot ((f_6(\ell))^2 + c \cdot (f_6(\ell))^2) + z) \\ &= \mathcal{O}(c \cdot (f_6(\ell))^3 + z), \end{aligned}$$

and the theorem follows.  $\square$

### 5.3 Levelings and well-aligned flatness pairs

Let  $G$  be a graph and let  $(W, \mathfrak{R})$  be a flatness pair of  $G$ . Let also  $\mathfrak{R} = (X, Y, P, C, \Gamma, \sigma, \pi)$ , where  $(\Gamma, \sigma, \pi)$  is an  $\Omega$ -rendition of  $G[Y]$  and  $\Gamma = (U, N)$  is a  $\Delta$ -painting. The *ground set* of  $W$  in  $\mathfrak{R}$  is  $\text{ground}_{\mathfrak{R}}(W) := \pi(N(\Gamma))$  and we refer to the vertices of this set as the *ground vertices* of the  $\mathfrak{R}$ -compass of  $W$  in  $G$ . Notice that  $\text{ground}_{\mathfrak{R}}(W)$  may contain vertices of  $\text{compass}_{\mathfrak{R}}(W)$  that are not necessarily vertices of  $W$ .

In the flatness pairs of [Figure 9](#) and [Figure 4](#) the ground vertices are the vertices on the boundaries of the green cells. (Notice also that the flatness pair in [Figure 9](#) is regular, while the one in [Figure 4](#) is not.)




Figure 10: The leveling  $W_{\mathfrak{R}}$  corresponding to the regular flatness pair  $(W, \mathfrak{R})$  in Figure 9. The ground-vertices of  $W_{\mathfrak{R}}$  are the circled vertices while the flap-vertices are the rhombic vertices. The representation  $R_W$  of  $W$  in  $W_{\mathfrak{R}}$  is obtained from  $W_{\mathfrak{R}}$  after removing the black squared vertices. The ground vertices in  $\text{bd}(\Delta) \cap W_{\mathfrak{R}} = \text{bd}(\Delta) \cap W_{\mathfrak{R}}$  are depicted in red.

**Levelings.** We define the  $\mathfrak{R}$ -leveling of  $W$  in  $G$ , denoted by  $W_{\mathfrak{R}}$ , as the bipartite graph where one part is the ground set of  $W$  in  $\mathfrak{R}$ , the other part is a set  $\text{vflaps}_{\mathfrak{R}}(W) = \{v_F \mid F \in \text{flaps}_{\mathfrak{R}}(W)\}$  containing one new vertex  $v_F$  for each flap  $F$  of  $W$  in  $\mathfrak{R}$ , and, given a pair  $(x, F) \in \text{ground}_{\mathfrak{R}}(W) \times \text{flaps}_{\mathfrak{R}}(W)$ , the set  $\{x, v_F\}$  is an edge of  $W_{\mathfrak{R}}$  if and only if  $x \in \partial F$ . We call the vertices of  $\text{ground}_{\mathfrak{R}}(W)$  (resp.  $\text{vflaps}_{\mathfrak{R}}(W)$ ) *ground-vertices* (resp. *flap-vertices*) of  $W_{\mathfrak{R}}$ . Notice that the incidence graph of the plane hypergraph  $(N(\Gamma), \{\tilde{c} \mid c \in C(\Gamma)\})$  is isomorphic to  $W_{\mathfrak{R}}$  via an isomorphism that extends  $\pi$  and, moreover, bijectively corresponds cells to flap-vertices. This permits us to treat  $W_{\mathfrak{R}}$  as a  $\Delta$ -embedded graph where  $\text{bd}(\Delta) \cap W_{\mathfrak{R}}$  is the set  $X \cap Y$ . As an example, see Figure 10 for the  $\mathfrak{R}$ -leveling corresponding to the flatness pair  $(W, \mathfrak{R})$  in Figure 9.

The following observation is a consequence of the definition of leveling and condition (v) of the tightness property of a rendition.

*Observation 18.* Let  $G$  be a graph, let  $(W, \mathfrak{R})$  be a flatness pair of  $G$ , and let  $W_{\mathfrak{R}}$  be the leveling of  $W$  in  $G$ . For every  $v_F \in \text{vflaps}_{\mathfrak{R}}(W)$  of degree  $r$  in  $W_{\mathfrak{R}}$ , there exist  $r$  internally vertex-disjoint paths in  $W_{\mathfrak{R}}$  from  $v_F$  to  $r$  distinct ground-vertices of  $W_{\mathfrak{R}}$  that belong to the perimeter of  $W$ .

**Well-aligned flatness pairs.** We denote by  $W^\bullet$  the graph obtained from  $W$  if we subdivide *once* every edge of  $W$  that is short in  $\text{compass}_{\mathfrak{R}}(W)$ . The graph  $W^\bullet$  is a “slightly richer variant” of  $W$  that is necessary for our definitions and proofs, namely to be able to associate every flap-vertex of an appropriate subgraph of  $W_{\mathfrak{R}}$  (that we will denote by  $R_W$ ) with a non-empty path of  $W^\bullet$ , as we proceed to formalize. We say that  $(W, \mathfrak{R})$  is *well-aligned* if the following holds:

$W_{\mathfrak{R}}$  contains as a subgraph an  $r$ -wall  $R_W$  where  $D(R_W) = D(W_{\mathfrak{R}})$  and  $W^\bullet$  is isomorphic to some subdivision of  $R_W$  via an isomorphism that maps each ground vertex to itself.

Suppose now that the flatness pair  $(W, \mathfrak{R})$  is well-aligned. We call the wall  $R_W$  in the above condition a *representation* of  $W$  in  $W_{\mathfrak{R}}$ . Note that, as  $R_W$  is a subgraph of  $W_{\mathfrak{R}}$ , it is bipartite as well. The above property

gives us a way to represent a flat wall by a wall of its leveling in a way that ground vertices are not altered. The following proposition, proved in [45], indicates that such a representation is yielded by regularity.

**Proposition 19.** *Every regular flatness pair  $(W, \mathfrak{R})$  of a graph  $G$  is well-aligned.*

Notice that both  $W_{\mathfrak{R}}$  and its subgraph  $R_W$  can be seen as  $\Delta$ -embedded graphs where  $\text{bd}(\Delta) \cap W_{\mathfrak{R}} = \text{bd}(\Delta) \cap R_W \subseteq V(D(W_{\mathfrak{R}})) = V(D(R_W))$ . This establishes a bijection from the set of cycles of  $W$  to the set of cycles of  $R_W$ , which allows to reinterpret the homogeneity property of a regular flatness pair in terms of its representation, as stated in the following observation. This translation will be used in the proof of [Theorem 23](#). Given the  $\Delta$ -embedded graph  $R_W$ , we define, for every brick  $B$  of  $R_W$ ,  $\text{vflaps}_{R_W}(B)$  as the flap-vertices of the leveling  $W_{\mathfrak{R}}$  that belong to the closed disk of the plane bounded by  $B$  disjoint from the infinite face. (Recall [Equation 3](#) for the definition of the augmented flap  $\mathbf{F}^A$  corresponding to a flap-vertex  $v_F$  of the leveling  $W_{\mathfrak{R}}$ .)

*Observation 20.* If  $(A, W, \mathfrak{R})$  is an  $(a, r, \ell)$ -apex-wall triple of a graph  $G$  and  $R_W$  is the representation of  $W$  in  $W_{\mathfrak{R}}$ , then for any two internal bricks  $B, B'$  of  $R_W$ , it holds that

$$\{\ell\text{-folio}(\mathbf{F}^A) \mid v_F \in \text{vflaps}_{R_W}(B)\} = \{\ell\text{-folio}(\mathbf{F}^A) \mid v_F \in \text{vflaps}_{R_W}(B')\}.$$

Note that, in the above equation, a flap  $F$  is notationally associated with both  $\mathbf{F}^A$  and  $v_F$ .

## 5.4 Rerouting minors of small intrusion

Let  $W$  be a plane-embedded  $r$ -wall and  $c \geq 1$ . We call a cycle  $C$  of  $W$   $c$ -internal if  $V(C)$  and  $V(D(W))$  are within face-distance at least  $c$ . Given a 1-internal cycle  $C$  of  $W$ , we define its *internal pegs* (resp. *external pegs*) as its vertices that are incident to edges of  $W$  that belong to the interior (resp. exterior) of  $C$  with respect to the embedding of the wall (we see edges as open sets). Notice that each vertex of  $C$  is either an internal or an external peg.

*Observation 21.* Let  $W$  be an  $r$ -wall and let  $C_1$  and  $C_2$  be four cycles of  $W$  within face-distance at least **four** and such that  $C_2$  is a subset of the closed disk bounded by  $C_1$ . Let  $y \in [3]$ , let  $p_1, \dots, p_y$  be internal pegs of  $C_1$  and  $\bar{p}_1, \dots, \bar{p}_y$  be external pegs of  $C_2$ , assuming that both these sets of vertices are ordered as they appear in their corresponding cycles in counter-clockwise order. Then there are  $y$  pairwise vertex-disjoint paths  $\hat{P}_1, \dots, \hat{P}_y$  such that, for  $i \in [y]$ ,  $\hat{P}_i$  joins  $p_i$  with  $\bar{p}_i$ ,  $V(\hat{P}_i) \cap V(C_1) = \{p_i\}$ , and  $V(\hat{P}_i) \cap V(C_2) = \{\bar{p}_i\}$ .

Given a 1-internal brick  $B$  of  $W$ , one can see the union of all bricks of  $W$  that have a common vertex with  $B$ , as a subdivision of the graph in the left part of [Figure 11](#). We call this subgraph  $X$  of  $W$  the *brick-neighborhood* of  $B$  in  $W$ . The *perimeter* of a brick-neighborhood is defined in the obvious way.

The next lemma is based on [Observation 18](#).

**Lemma 22.** *Let  $(W, \mathfrak{R})$  be a well-aligned flatness pair of a graph  $G$  and let  $R_W$  be its representation in the leveling  $W_{\mathfrak{R}}$  of  $W$ . For every 2-internal brick  $B$  of  $R_W$  and every flap vertex  $v_F \in \text{vflaps}_{\mathfrak{R}}(B)$ ,  $R_W$  contains  $|\partial F|$  internally vertex-disjoint paths from  $v_F$  to the external pegs of the perimeter of the brick-neighborhood of  $B$  in  $R_W$ . Moreover, these paths belong to the closed disk bounded by the perimeter of the brick-neighborhood of  $B$  in  $R_W$ .*

*Proof.* Let  $r = |\partial F|$ ,  $X$  be the brick-neighborhood of  $B$  in  $R_W$ , and  $P$  be the perimeter of  $X$  in  $R_W$ . We call *frontier-path* of  $X$  a subpath of  $P$  that joins two external pegs and does not contain any other external peg. Notice that  $P$  is the union of the frontier-paths of  $X$  and that there are exactly 12 such paths. Notice that for every frontier-path  $Q$  of  $X$  there is a path  $\hat{Q}$  of  $X$  such that

- its endpoints are in  $P$  but not in  $Q$ ,




Figure 11: On the left: the base graph for the definition of a brick-neighborhood – the external pegs of the perimeter of  $X$  are the black round vertices. On the right we depict a 2-internal brick  $B$  of an  $r$ -wall  $W$ ,  $r \geq 6$ , contained as a subgraph in a plane graph  $G$  along with three internally vertex-disjoint paths from a vertex  $F$  of  $G$  to the external pegs of the perimeter of the brick-neighborhood of  $B$ .

- it does not contain any internal vertex in  $P$ , and
- every path in  $W_{\mathfrak{R}}$  from  $v_F$  to a vertex of  $Q$  intersects some vertex of  $\hat{Q}$ .

See Figure 12 for two indicative examples of the above correspondence.




Figure 12: Two indicative examples for the choice of  $\hat{Q}$  (in blue), depending on the choice of  $Q$  (in green).

From [Observation 18](#), there are  $r$  internally vertex-disjoint paths  $P_1, \dots, P_r$  in  $W_{\mathfrak{R}}$  starting from  $v_F$  and finishing in vertices  $v_1, \dots, v_r$  of  $P$ . Moreover, we can assume that  $(P_1 \cup \dots \cup P_r) \cap P = \{v_1, \dots, v_r\}$ . The lemma is trivial in case  $r < 3$  as the paths  $P_1, \dots, P_r$  can easily be extended so to finish in external pegs of  $P$ . Moreover, the lemma also follows easily if  $r = 3$  and  $v_1, v_2, v_3$  do not all belong to some of the frontier-paths of  $X$ . It remains to examine the case where  $v_1, v_2, v_3$  are vertices of some frontier-path  $Q$  of  $X$ . Let  $\hat{Q}$  be as defined above and let  $\hat{q}$  be one of its endpoints. Let  $z_Q$  be the first vertex of  $P_1 \cup P_2 \cup P_3$  that is met while following  $\hat{Q}$  starting from  $\hat{q}$  and moving towards its other endpoint. The vertex  $z_Q$  exists because of the definition of  $\hat{Q}$ . W.l.o.g., we assume that  $z_Q$  is an internal vertex of  $P_1$ . We now define  $P'_1$  by first removing from  $P_1$  all vertices of its subpath from  $z_Q$  to  $v_i$ , except  $z_Q$  and then taking the union of the resulting path with the subpath of  $\hat{Q}$  between  $\hat{q}$  and  $z_Q$ . It is now easy to see that  $P'_1$  can be extended to some external peg that is different from the endpoints of  $P_2$  and  $P_3$ , while  $P_2$  and  $P_3$  can be extended towards the external pegs that are endpoints of  $\hat{Q}$ . This completes the proof of the lemma.  $\square$

**Irrelevant vertices in boundaried graphs.** Let  $G$  be a graph,  $H$  be a minor of  $G$ , and  $S \subseteq V(G)$ . We define the  $S$ -minor-intrusion of  $H$  in  $G$  as the minimum  $S$ -intrusion in  $G$  over all tm-pairs  $(M, T)$  of  $G$  such that  $(M, T)$  is a topological minor model in  $G$  where  $\text{diss}(M, T) \in \text{ext}(H)$ .

Let  $\mathbf{Z} = (Z, B, \rho)$  be a  $t$ -boundaried graph and let  $\ell \in \mathbb{N}$ . We say that a vertex set  $S \subseteq V(Z) \setminus B$  is  $\ell$ -irrelevant if for every boundaried graph  $\mathbf{K} = (K, B, \rho)$  that is compatible with  $\mathbf{Z}$ , every minor of  $\mathbf{K} \oplus \mathbf{Z}$  with  $(V(Z) \setminus B)$ -minor-intrusion at most  $\ell$ , is also a minor of  $\mathbf{K} \oplus (Z \setminus S, B, \rho)$ . Informally, an  $\ell$ -irrelevant

set of vertices can be removed without affecting the occurrences of any minor of minor-intrusion at most  $\ell$ , where the intrusion is defined without taking into account the terminal vertices in the boundary.

Using [Theorem 17](#), we can finally prove the main result of this section.

**Theorem 23.** *There exist two functions  $f_{13} : \mathbb{N}^3 \rightarrow \mathbb{N}$  and  $f_{14} : \mathbb{N}^2 \rightarrow \mathbb{N}$  such that, for every  $a, z, \ell \in \mathbb{N}$  and every boundaried graph  $\mathbf{Z} = (Z, B, \rho)$ , if  $(A, W, \mathfrak{R})$  is an  $(a, f_{13}(a, z, \ell), f_{14}(a, \ell))$ -apex-wall triple of  $Z$  that is not affected by  $B$ , then the vertex set of the compass of every  $W'$ -tilt of  $(W, \mathfrak{R})$ , where  $W'$  is the central  $z$ -subwall of  $W$ , is  $\ell$ -irrelevant. Moreover, it holds that  $f_{13}(a, z, \ell) = \mathcal{O}((f_6(16a + 12\ell))^3 + z)$  and  $f_{14}(a, \ell) = a + 3 + \ell$ .*

*Proof.* Let  $\tilde{\ell} = 16a + 12\ell$ ,  $\hat{\ell} = a + 3 + \ell$ ,  $r = f_{12}(6, z, \tilde{\ell})$ , and  $\tilde{r} = f_{10}(6, \tilde{\ell})$ . We prove the theorem for  $f_{13}(a, z, \ell) = r$  and  $f_{14}(a, \ell) = \hat{\ell}$ . Note that, by [Theorem 17](#),  $f_{13}(a, z, \ell) = \mathcal{O}((f_6(\tilde{\ell}))^3 + z)$ . Let  $\mathbf{Z} = (Z, B, \rho)$  and let  $(A, W, \mathfrak{R})$  be an  $(a, r, \hat{\ell})$ -apex-wall triple of the graph  $Z$  that is not affected by  $B$ . Let  $W'$  be the central  $z$ -subwall of  $W$ , and let  $(\tilde{W}', \tilde{\mathfrak{R}}')$  be a flatness pair of  $Z \setminus A$  that is a  $W'$ -tilt of  $(W, \mathfrak{R})$ . Our objective is to prove that  $V(\text{compass}_{\tilde{\mathfrak{R}}'}(\tilde{W}'))$  is  $\ell$ -irrelevant.

Let  $\mathbf{K} = (K, B, \rho)$  be a boundaried graph compatible with  $\mathbf{Z}$ . As  $(A, W, \mathfrak{R})$  is not affected by  $B$ , we have that  $(A, W, \mathfrak{R})$  is an  $(a, r, \hat{\ell})$ -apex-wall triple of the graph  $G := \mathbf{K} \oplus \mathbf{Z}$  as well. Let now  $H$  be a minor of  $G$  whose  $(V(Z) \setminus B)$ -minor-intrusion in  $G$  is at most  $\ell$ . Let also  $(M, T)$  be a tm-pair in  $G$  that is a topological minor model of a graph  $H' \in \text{ext}(H)$ , such that  $|T \setminus V(K)| = |T \cap (V(Z) \setminus B)| \leq \ell$  and with at most  $\ell$  subdivision paths intersecting  $V(Z) \setminus B$ . Our purpose is to find a tm-pair  $(\overline{M}, \overline{T})$  of  $G$  where  $V(\text{compass}_{\tilde{\mathfrak{R}}'}(\tilde{W}')) \cap V(\overline{M}) = \emptyset$  and such that  $H' = \text{diss}(M, T)$  is a minor of  $\text{diss}(\overline{M}, \overline{T})$ , which in turn implies that  $H$  is a minor of  $G \setminus V(\text{compass}_{\tilde{\mathfrak{R}}'}(\tilde{W}')) = \mathbf{K} \oplus (Z \setminus V(\text{compass}_{\tilde{\mathfrak{R}}'}(\tilde{W}'))), B, \rho)$ , as required.




Figure 13: An illustration of the graph  $G = \mathbf{K} \oplus \mathbf{Z}$  and a tm-model  $(M, T)$  in it. The apex set  $A$  is cyan. We also draw a single flap  $F$  and its magnification.

We proceed with the definition of a series of auxiliary graphs that will permit to work on a partially planarized version of  $G$ . Suppose that  $\mathfrak{R} = (X, Y, P, C, \Gamma, \sigma, \pi)$ . We first define the graph  $G_{\text{out}}$  as the graph obtained from  $G$  if we remove all the vertices of  $\text{compass}_{\mathfrak{R}}(W)$ , except from those in  $X \cap Y$ . We then define  $\tilde{G} := G_{\text{out}} \cup W_{\mathfrak{R}}$  where  $W_{\mathfrak{R}}$  is the leveling of  $W$ . Since  $(W, \mathfrak{R})$  is regular, it is also well-aligned by [Proposition 19](#), hence  $W_{\mathfrak{R}}$  contains a representation  $R_W$  of  $W$ , which is a subgraph of  $W_{\mathfrak{R}}$  that is isomorphic to some subdivision of  $W^\bullet$  via an isomorphism that maps each ground vertex to itself (recall that  $W^\bullet$  is

the graph obtained from  $W$  if we subdivide once every short edge in  $W$ ). Notice that  $\tilde{G}$  is a partially  $\Delta$ -embedded graph whose compass is  $W_{\mathfrak{R}}$  that, in turn, is the compass in  $\tilde{G}$  of the  $d$ -wall  $R_W$ . Recall that each flap  $F$  of  $W$  corresponds to a flap-vertex  $v_F$  of  $W_{\mathfrak{R}}$ .

We enhance  $(M, T)$  by defining another tm-pair  $(M_A, T_A)$ , where  $M_A = (V(M) \cup A, E(M))$  and  $T_A = T \cup A$ , i.e.,  $(M_A, T_A)$  is obtained from  $(M, T)$  by including all the apices in  $A$ . We set  $H_A = \text{diss}(M_A, T_A)$  and observe that  $H$  is a (topological) minor of  $H_A$ . We set  $T_{\text{in}} = A \cup (T_A \cap \text{compass}_{\mathfrak{R}}(W))$  and  $T_{\text{out}} = A \cup (T_A \setminus (\text{compass}_{\mathfrak{R}}(W) \setminus (X \cap Y)))$ . As  $(A, W, \mathfrak{R})$  is not affected by  $B$ , it follows that  $T_{\text{in}} \cap \text{compass}_{\mathfrak{R}}(W) \subseteq T \setminus (V(K) \cup B)$ , hence  $|T_{\text{in}} \cap \text{compass}_{\mathfrak{R}}(W)| \leq \ell$ .

We call a vertex  $v \in \text{compass}_{\mathfrak{R}}(W)$  an *apex-jump* vertex if there exists an edge  $\{v, w\}$  of  $M_A$  with  $w \in A$ . Notice that there are at most  $a + \ell$  apex-jump vertices. We define a set of  $(M_A, T_A)$ -dirty flaps of the flatness pair  $(W, \mathfrak{R})$  by applying the following definition: first we declare as  $(M_A, T_A)$ -dirty every flap  $F$  such that  $V(F) \setminus \partial F$  contains an apex-jump vertex or a vertex in  $T_{\text{in}}$ . Second, for every apex-jump vertex  $v \in \text{ground}(W)$  that is not in the boundary of some  $(M_A, T_A)$ -dirty flap, we arbitrarily pick a flap  $F$  with  $v \in \partial F$  and we declare it  $(M_A, T_A)$ -dirty. Observe that  $(W, \mathfrak{R})$  has at most  $a + \ell + |T_{\text{in}} \cap \text{compass}_{\mathfrak{R}}(W)| \leq a + 2\ell$   $(M_A, T_A)$ -dirty flaps.




Figure 14: An example of how part of the model  $(M_A, T_A)$  (in red edges) traverses the flaps of  $W$  along with the model  $(\tilde{M}_{\text{in}}, \tilde{T}_{\text{in}})$  of  $\tilde{W}$  (i.e., the leveling of  $(M_A, T_A)$ ). The dashed red edges are edges pointing to apices. The orange flaps are the  $(M_A, T_A)$ -dirty flaps. The black vertices are the ground vertices that do not belong to  $M$ .

We define  $\tilde{M}_{\text{in}}$  as the subgraph of the leveling  $W_{\mathfrak{R}}$  induced by the vertices in  $\text{ground}(W) \cap V(M_A)$  and all the flap-vertices  $v_F$  of  $W_{\mathfrak{R}}$  such that  $F$  is either  $(M_A, T_A)$ -dirty or contains an edge of  $M$  (in Figure 14, these latter flaps are turquoise). Note that the induced edges may increase the degree of some, say  $w$ , of the degree-2 ground vertices in  $V(M_A) \setminus T_A$  because of some edge  $e = \{v_F, w\}$  between a flap-vertex  $v_F$  of  $W_{\mathfrak{R}}$  and  $w$ ; we then remove from  $\tilde{M}_{\text{in}}$  all such edges (in Figure 14, these edges are the black dotted edges). Actually this last modification could be avoided, however it facilitates the presentation of the last part of the proof.

Notice that if a flap  $F$  is not  $(M_A, T_A)$ -dirty and contains an edge  $e$  of  $M$ , then this edge should belong to a subpath  $P$  of a subdivision path of  $M$  such that the endpoints of  $P$  are two of the vertices of  $\partial F$ . We call such flap-vertices of  $\tilde{M}_{\text{in}}$  *subdivision flap-vertices* of  $\tilde{M}_{\text{in}}$ . If a flap-vertex of  $\tilde{M}_{\text{in}}$  is not a subdivision flap-vertex, then we call it *branch flap-vertex* of  $\tilde{M}_{\text{in}}$ . We denote by  $Q$  the set of all the branch flap-vertices of  $\tilde{M}_{\text{in}}$ , and as each branch flap-vertex corresponds to a  $(M_A, T_A)$ -dirty flap, we have that  $|Q| \leq a + 2\ell$ . Notice that all vertices of  $Q$  have degree at most three in  $W_{\mathfrak{R}}$  (and therefore in  $\tilde{M}_{\text{in}}$  as well).

The ground-vertices of  $\tilde{M}_{\text{in}}$  that are apex-jump vertices or belong to  $T_{\text{in}} \setminus A$  are called *branch ground-vertices* of  $\tilde{M}_{\text{in}}$ , while the rest of the ground vertices of  $\tilde{M}_{\text{in}}$  are called *subdivision ground-vertices* of  $\tilde{M}_{\text{in}}$ . Notice that there are at most  $a + 2\ell$  branch ground-vertices in  $\tilde{M}_{\text{in}}$ . Let  $M_{\text{out}} = M_A \cap G_{\text{out}}$  and  $\tilde{M} = M_{\text{out}} \cup \tilde{M}_{\text{in}}$ . We define  $\tilde{T}_{\text{in}}$  as the union of the set of branch ground-vertices of  $\tilde{M}_{\text{in}}$  and the set  $Q$  of the branch flap-vertices of  $\tilde{M}_{\text{in}}$ . Observe that  $|\tilde{T}_{\text{in}}| \leq 2a + 4\ell$ . Moreover, we set  $T_{\text{out}} = T_A \cap V(M_{\text{out}})$  and  $\tilde{T} = T_{\text{out}} \cup \tilde{T}_{\text{in}}$ .

Notice that  $(\tilde{M}, \tilde{T})$  is a tm-pair of  $\tilde{G}$ ; we refer to it as the *leveling of the tm-pair*  $(M_A, T_A)$  of  $G$  with respect to  $(A, W, \mathfrak{R})$ . Clearly,  $\tilde{M}$  is a partially  $\Delta$ -embedded graph whose compass is  $\tilde{M}_{\text{in}}$ , i.e.,  $\tilde{M}_{\text{in}} = \Delta \cap \tilde{M}$ . Also, as  $|\tilde{T}_{\text{in}}| \leq 2a + 4\ell$ , the  $\Delta \cap V(G)$ -intrusion of  $(\tilde{M}, \tilde{T})$  in  $\tilde{G}$  can be bounded, using the fact that  $\tilde{M}_{\text{in}}$  is a subgraph of the planar graph  $\tilde{G}_{\text{in}}$ , by  $3(2a + 4\ell) = \tilde{\ell}$ .

Let us now give some intuition on the definition of  $(\tilde{M}, \tilde{T})$ . We see  $\tilde{G}$  and its tm-pair  $(\tilde{M}, \tilde{T})$  as a “projection” of the graph  $G$  and its tm-pair  $(M, T)$ , respectively, in what concerns the leveling  $W_{\mathfrak{R}}$  of  $W$ . This projection is losing some of the information of  $(\tilde{M}, \tilde{T})$ , however it will be valuable as now both  $\tilde{G}$  and  $\tilde{M}$  are partially  $\Delta$ -embedded graphs. The lost information is encoded, for every branch flap-vertex  $v_F$  in  $Q$ , by  $\ell$ -folio( $\mathbf{F}^A$ ). Recall that  $\mathbf{F}^A := (G[A \cup F], A \cup \partial F, \rho_A \cup \rho_F)$ . In what follows, we will use [Theorem 17](#) in order to draw in  $\tilde{G}$  a modification  $(\hat{M}, \hat{T})$  of  $(\tilde{M}, \tilde{T})$  so that  $(\hat{M}, \hat{T})$  does not go “too deeply” in the representation  $R_W$  of  $W$  in  $W_{\mathfrak{R}}$ , and the parts of  $(\hat{M}, \hat{T})$  that are different from  $(\tilde{M}, \tilde{T})$  are routed through  $R_W$  in a “dispersed enough” way. Moreover, again from [Theorem 17](#),  $(\tilde{M}, \tilde{T})$  can be seen as a contraction of  $(\hat{M}, \hat{T})$  that does not identify any of the vertices of  $Q$ . Maintaining the vertices in  $Q$  intact, while performing contractions in  $(\hat{M}, \hat{T})$ , is important as we need to keep the information given by  $\ell$ -folio( $\mathbf{F}^A$ ) for each branch flap-vertex  $v_F \in Q$ .

For each branch flap-vertex  $v_F$  of  $Q$  we define  $\mathbf{M}_F = (G[A \cup V(F)] \cap M_A, A \cup \partial F, \rho_A \cup \rho_F)$ , where the functions  $\rho_A$  and  $\rho_F$  are defined as explained in the beginning of [Subsection 4.5](#), and  $T_F = A \cup \partial F \cup (T_A \cap F)$ . Notice that  $(\mathbf{M}_F, T_F)$  is a btm-pair of  $\mathbf{F}^A$  such that  $\text{diss}(\mathbf{M}_F, T_F) \in \hat{\ell}$ -folio( $\mathbf{F}^A$ ), given that  $\hat{\ell} = a + 3 + \ell$ . This means that if  $v_{\bar{F}}$  is another flap-vertex of  $W_{\mathfrak{R}}$  with  $\hat{\ell}$ -folio( $\mathbf{F}^A$ ) =  $\hat{\ell}$ -folio( $\bar{\mathbf{F}}^A$ ), then there is a btm-pair  $(\mathbf{M}_{\bar{F}}, T_{\bar{F}})$  of  $\bar{\mathbf{F}}^A$  such that  $\text{diss}(\mathbf{M}_F, T_F) = \text{diss}(\mathbf{M}_{\bar{F}}, T_{\bar{F}})$ .

Recall that  $\tilde{G}$  is a partially  $\Delta$ -embedded graph whose compass is  $\tilde{G} \cap \Delta = W_{\mathfrak{R}}$  and that  $\tilde{M}$  can be seen as a partially  $\Delta$ -embedded graph whose compass is  $\tilde{M} \cap \Delta$ . Recall also that the  $\Delta \cap V(G)$ -intrusion of  $(\tilde{M}, \tilde{T})$  in  $\tilde{G}$  is at most  $\tilde{\ell}$ .

Let  $C_1, \dots, C_{\tilde{r}}$  be the first  $\tilde{r}$  layers of  $R_W$  and let  $[D_1, \dots, D_{\tilde{r}}]$  (resp.  $[\bar{D}_1, \dots, \bar{D}_{\tilde{r}}]$ ) be the sequences of the corresponding open (resp. closed) disks. We can now apply [Theorem 17](#) with input 6,  $\tilde{\ell}$ ,  $z$ ,  $\tilde{G}$ , the  $r$ -wall  $R_W$ , the tm-pair  $(\tilde{M}, \tilde{T})$  of  $\tilde{G}$ , and the set  $Q$  of branch flap-vertices defined above. Let  $d = f_{11}(6, \tilde{\ell})$ . [Theorem 17](#) guarantees the existence of a tm-pair  $(\hat{M}, \hat{T})$  of  $\tilde{G}$  and a  $b \in [\tilde{r}]$  such that

1.  $\hat{M} \setminus D_b$  is a subgraph of  $\tilde{M} \setminus D_b$ ,
2.  $\text{ann}(\mathcal{C}_{b, b+d-1}) \cap (\tilde{T} \cup \hat{T}) = \emptyset$ ,
3.  $(\hat{M}, \hat{T}) \cap \bar{D}_{b+d}$  is a tm-pair of  $R_W$  that is safely 6-dispersed in  $R_W$  and none of the vertices of  $\hat{T} \cap \bar{D}_{b+d}$  is within face-distance less than six in  $R_W$  from some vertex of  $C_{b+d} \cup C_{\tilde{r}}$ ,
4.  $\hat{M} \cap D_{\tilde{r}} = \emptyset$ ,
5. the compass of the central  $z$ -subwall of  $W$  is a subset of  $D_{\tilde{r}}$ , and
6. there is a  $Q$ -respecting contraction-mapping  $\phi$  of  $\text{diss}(\tilde{M}, \tilde{T})$  to  $\text{diss}(\hat{M}, \hat{T})$ .

Our next step is to modify the tm-pair  $(\hat{M}, \hat{T})$  to obtain another tm-pair  $(\hat{M}^+, \hat{T}^+)$  so that, in addition to Property 1 and Property 4 above (where  $\hat{M}$  is replaced by  $\hat{M}^+$ ),  $(\hat{M}^+, \hat{T}^+)$  satisfies the following stronger version of Property 6:

- 6<sup>+</sup>. there is a  $Q$ -respecting contraction-mapping  $\phi^+$  of  $\text{diss}(\tilde{M}, \tilde{T})$  to  $\text{diss}(\hat{M}^+, \hat{T}^+)$  such that for every branch flap-vertex  $v_F \in Q$ , if  $\phi^+(v_F) = \{v_{\hat{F}}\}$ , then  $\hat{\ell}\text{-folio}(\mathbf{F}^A) = \hat{\ell}\text{-folio}(\hat{\mathbf{F}}^A)$ .

To force Property 6<sup>+</sup>, we will make strong use of the fact that the flatness pair  $(W, \mathfrak{R})$  is  $\hat{\ell}$ -homogeneous with respect to  $(G, A)$ . This will permit us to modify the subdivision paths of  $(\hat{M}, \hat{T})$  by pointing to images of flaps in  $Q$  that have been displaced after the application of [Theorem 17](#).

We proceed as follows: let  $v_F \in Q$  be some flap-vertex, and let  $\phi(v_F) = \{w\}$  be its image given by the function  $\phi$  guaranteed by Property 6. From Property 2,  $w$  cannot belong to  $\text{ann}(\mathcal{C}_{b,b+d-1})$ . Also, from Property 4,  $w$  cannot belong to  $D_{\tilde{r}}$ . If  $w \in \text{ann}(\mathcal{C}_{1,b})$ , it follows from Property 1 that  $w = v_F$  and in this case Property 6<sup>+</sup> already holds trivially. The only remaining case is when  $w \in \text{ann}(\mathcal{C}_{b+d,\tilde{r}})$ . In this case, from Property 3,  $w$  is a vertex of  $R_W \cap \text{ann}(\mathcal{C}_{b+d,\tilde{r}})$  that is within face-distance at least six from both  $C_{b+d}$  and  $C_{\tilde{r}}$ .




Figure 15: Some part of  $R_W \cap \text{ann}(\mathcal{C}_{b+d,\tilde{r}})$  around a vertex  $w$  of  $\hat{T}$ . The interior of the brick  $\hat{R}$  is depicted in yellow. The cycle  $C_w$  is depicted in blue.

Let  $R$  and  $\hat{R}$  be bricks of  $R_W$  such that  $v_F \in \text{vflaps}_{R_W}(R)$  and  $w \in \text{vflaps}_{R_W}(\hat{R})$ , respectively. Certainly, both  $R$  and  $\hat{R}$  are internal bricks of  $R_W$ . Moreover,  $w$  is a vertex of the cycle  $\hat{R}$  (bounding the disk depicted in bright yellow in [Figure 15](#)). By [Observation 20](#), there exists a vertex  $v_{\hat{F}}$  in  $\hat{R}$  such that  $\hat{\ell}\text{-folio}(\mathbf{F}^A) = \hat{\ell}\text{-folio}(\hat{\mathbf{F}}^A)$ . Let  $y$  be the degree of  $w$  in  $\hat{M}$  (which equals the degree of  $v_F$  in  $\tilde{M}$ ) and let  $P_1, \dots, P_y$  be the subdivided paths of the model  $(\hat{M}, \hat{T})$  that have  $w$  as an endpoint, depicted as fat green lines in [Figure 15](#). Note that  $y \leq 3$ .

Let  $C_w$  be the cycle of  $R_W$  induced by the vertices that are within face-distance exactly six from  $w$  (in [Figure 15](#), this cycle is depicted with thick blue edges). From Property 3,  $C_w$  is within face-distance at least  $(2 \cdot 6 + 1) - 6 = 7$  from any other vertex in  $\hat{T} \cap \overline{D}_{b+d}$  in  $R_W$ . For  $i \in [y]$ , we define  $P'_i$  as the unique path, among the paths obtained from  $P_i$  by the removal the open disk bounded by  $C_w$ , that has only one endpoint in the cycle  $C_w$ . For  $i \in [y]$ , we denote by  $p_i$  the endpoint of  $P'_i$  that is not an endpoint of  $P_i$  (depicted in purple in [Figure 15](#)). Notice that  $p_1, \dots, p_y$  are internal pegs of  $C$  ordered as they appear in the perimeter in counter-clockwise order.

Consider now the brick-neighborhood, say  $X$ , of  $\hat{R}$ . Notice that the face-distance between the perimeter of  $X$  and  $C_w$  is at least four. From [Lemma 22](#), there are  $y$  internally vertex-disjoint paths  $\bar{P}_1, \dots, \bar{P}_y$  from  $v_{\hat{F}}$  to the external pegs of the perimeter  $C_X$  of  $X$  (depicted in red lines in [Figure 15](#)). Let these pegs be  $\bar{p}_1, \dots, \bar{p}_y$  ordered as they appear in the perimeter in counter-clockwise order. We are now in position to apply [Observation 21](#) to  $C_w$ ,  $C_X$ ,  $p_1, \dots, p_y$ , and  $\bar{p}_1, \dots, \bar{p}_y$  and find  $y$  pairwise vertex-disjoint paths  $\hat{P}_1, \dots, \hat{P}_y$  each joining  $p_i$  with  $\bar{p}_i$  (depicted in orange lines in [Figure 15](#)). We now remove from  $\hat{M}$  the open disk bounded by  $C_w$  and we add the graph  $\bar{P}_1 \cup \dots \cup \bar{P}_y \cup \hat{P}_1 \cup \dots \cup \hat{P}_y$ . We also modify  $\hat{T}$  by substituting  $w$  by  $v_{\hat{F}}$ . By repeating this procedure for every flap-vertex of  $Q$ , and defining  $(\hat{M}^+, \hat{T}^+)$  as the updated tm-pair obtained when this algorithm terminates, we enforce Property 6<sup>+</sup>. Namely the function  $\phi^+$  is defined from modifying  $\phi$  so that, whenever  $\phi(v_F) \neq \{v_F\}$ , we set  $\phi^+(v_F) = \{v_{\hat{F}}\}$  as above. Note that  $(\hat{M}^+, \hat{T}^+)$  indeed satisfies, besides Property 6<sup>+</sup>, Properties 1 and 4.

Our next (and last) step is to further modify the tm-pair  $(\hat{M}^+, \hat{T}^+)$  of  $\tilde{G}$  so to obtain a tm-pair  $(\bar{M}, \bar{T})$  of the original graph  $G$  so that  $\text{diss}(M, T)$  is a minor of  $\text{diss}(\bar{M}, \bar{T})$ . Notice that each edge of  $\hat{M}^+$  that is an edge of  $R_W$  (resp.  $W_{\mathfrak{R}}$ ) has one endpoint that is a ground-vertex of  $R_W$  (resp.  $W_{\mathfrak{R}}$ ) and another one that is a flap-vertex of  $R_W$  (resp.  $W_{\mathfrak{R}}$ ).

Let  $v_{\hat{F}}$  be a flap-vertex of  $\hat{M}^+$ . We modify  $(\hat{T}^+, \hat{M}^+)$  by distinguishing the following cases:

- $v_{\hat{F}} \notin \hat{T}^+$ . Then the neighbors of  $v_{\hat{F}}$  in  $R_W$  are two ground-vertices  $g$  and  $g'$ . From tightness property (ii) of a rendition, there is a path  $P_{\hat{F}}$  in  $\hat{F}$  with  $g$  and  $g'$  as endpoints. We substitute the edges  $\{g, \hat{F}\}$  and  $\{\hat{F}, g'\}$  of  $R_W$  with  $P_{\hat{F}}$ . Notice that  $P_{\hat{F}}$  is a path of  $\text{compass}_{\mathfrak{R}}(W)$ .
- $v_{\hat{F}} \in \hat{T}^+$  and  $v_{\hat{F}} \in \phi^+(x)$  for some  $x \notin Q$ . In this case, from tightness property (iii) of a rendition,  $\hat{F}$  has  $y \in [3]$  neighbors  $v_1, \dots, v_y$ , and a vertex  $z_{\hat{F}}$  that is connected to  $v_1, \dots, v_y$  via  $y$  internally vertex-disjoint paths. We substitute the edges  $\{z_{\hat{F}}, v_1\}, \dots, \{z_{\hat{F}}, v_y\}$  and the vertex  $z_{\hat{F}}$  of  $R_W$  with the union of these  $y$  internally vertex-disjoint paths. Notice that these paths are also paths of  $\text{compass}_{\mathfrak{R}}(W)$ . We also update  $\hat{T}^+ := \hat{T}^+ \setminus \{v_{\hat{F}}\} \cup \{z_{\hat{F}}\}$  and we call the vertex  $z_{\hat{F}}$  *replacement* of  $v_{\hat{F}}$ .
- $v_{\hat{F}} \in \hat{T}^+$  and  $\{v_{\hat{F}}\} = \phi^+(v_F)$  for some  $v_F \in Q$ . Let  $\mathbf{M}_F = (F \cap M, \partial F, \rho_A \cup \rho_F)$  and let  $T_F = T \cap F$ . From Property 6<sup>+</sup>,  $\hat{\ell}\text{-folio}(\hat{\mathbf{F}}^A) = \hat{\ell}\text{-folio}(\mathbf{F}^A)$ , which implies that there is a *btm-pair*  $(\mathbf{M}_{\hat{F}}, T_{\hat{F}})$  of  $\hat{\mathbf{F}}^A$  such that  $\text{diss}(\mathbf{M}_{\hat{F}}, T_{\hat{F}}) = \text{diss}(\mathbf{M}_F, T_F)$ . We denote by  $M_{\hat{F}}$  the underlying graph of  $\mathbf{M}_{\hat{F}}$  and we substitute the vertex  $v_{\hat{F}}$  and its incident edges in  $W_{\mathfrak{R}}$  with the graph  $M_{\hat{F}}$  (that is a subgraph of  $\text{compass}_{\mathfrak{R}}(W)$ ), by identifying the boundary vertices according to the functions  $\rho_A \cup \rho_F$ . We also update  $\hat{T}^+ := \hat{T}^+ \setminus \{v_{\hat{F}}\} \cup T_{\hat{F}}$ .

The above operations create a tm-pair of  $G$  that we denote henceforth by  $(\bar{M}, \bar{T})$ . Since  $(\bar{M}, \bar{T})$  satisfies Property 4 and  $\tilde{r} \geq z$ ,  $\bar{M} \cap \mathbf{U}\text{influence}_{\mathfrak{R}}(W') = \emptyset$ , and therefore  $(\bar{M}, \bar{T})$  is a tm-pair of  $G \setminus V(\text{compass}_{\mathfrak{R}}(\tilde{W}'))$ . It is worth mentioning that Property 1 implies the strong property that, throughout the rerouting procedure, the part of the topological minor model outside of the  $\mathfrak{R}$ -compass of  $W$  in  $G \setminus A$  can only be reduced; formally,  $\bar{M} \cap X \subseteq M \cap X$  (see [Remark 24](#) after the end of this proof).

As  $H$  is a minor of  $H' = \text{diss}(M, T)$ , hence a minor of  $\text{diss}(M_A, T_A)$  as well, and  $G \setminus V(\text{compass}_{\mathfrak{R}}(\tilde{W}')) = \mathbf{K} \oplus (Z \setminus V(\text{compass}_{\mathfrak{R}}(\tilde{W}')), B, \rho)$ , it remains to verify that  $\text{diss}(\bar{M}, \bar{T})$  contains  $\text{diss}(M_A, T_A)$  as a minor. For this, consider every  $x \in \tilde{T} \setminus Q$  and construct the set  $T_x$  by taking  $T_x := \phi^+(x)$  and substituting each branch flap-vertex  $v_{\hat{F}} \in T_x$  with its replacement  $z_{\hat{F}}$  defined as in the second case of the above case analysis. Notice that each set  $T_x$  is a subset of  $\bar{T}$ . We now construct  $(M^*, T^*)$  as follows:  $\tilde{M}^*$  is obtained by contracting, for each  $x \in \tilde{T} \setminus Q$ , all subdivision paths in  $\bar{M}$  that have as endpoints two vertices in  $T_x$  to a single vertex, which we again call  $x$ . This operation identifies, for each  $x \in \tilde{T} \setminus Q$ , all vertices of  $T_x$  into  $x$ , thus we set  $T^* = (\bar{T} \setminus (\bigcup_{x \in \tilde{T} \setminus Q} T_x)) \cup (\tilde{T} \setminus Q)$ . That way  $(M^*, T^*)$  can be seen as the tm-pair  $(\tilde{M}, \tilde{T})$  where each of its branch flap-vertices has been substituted as in the third case of the above case analysis. This permits us to




Figure 16: The tm-pairs considered in the proof of [Theorem 23](#). The cyan-shadowed pairs are tm-pairs of  $G$  and the pink-shadowed pairs are tm-pairs of  $\tilde{G}$ .

verify that  $\text{diss}(M_A, T_A)$  and  $\text{diss}(M^*, T^*)$  are isomorphic (see [Figure 16](#) for the relation between the models that we have defined). As  $\text{diss}(M^*, T^*)$  is a minor of  $\text{diss}(\overline{M}, \overline{T})$ , the theorem follows.  $\square$

*Remark 24.* In the above proof, the rerouted topological minor model  $(\overline{M}, \overline{T})$  obtained from the original model  $(M, T)$  satisfies the following property: the part of the topological minor model outside of the  $\mathfrak{A}$ -compass of  $W$  in  $G \setminus A$  can only be reduced with respect to the original one; formally,  $\overline{M} \cap X \subseteq M \cap X$ , where  $X$  is the “external” set corresponding to the separation of the considered apex-wall triple  $(A, W, \mathfrak{A})$ .

By definition of the set  $\mathcal{R}_h^{(t)}$ , its elements are of minimum size, and therefore a boundaried graph  $\mathbf{G} = (G, B, \rho) \in \mathcal{R}_h^{(t)}$  does not contain any  $3h$ -irrelevant vertex. To see this, recall that in [Equation 2](#) the equivalence is defined in terms of graphs  $H$  with detail at most  $h$  (i.e., with at most  $h$  vertices and at most  $h$  edges), and that by [Observation 3](#) every graph in  $\text{ext}(H)$  has detail at most  $3h$ . On the other hand, [Observation 7](#) implies that, in the setting of [Theorem 23](#), there is a vertex belonging to the compass of every  $W'$ -tilt of  $(W, \mathfrak{A})$ , where  $W'$  is the central  $z$ -subwall of  $W$ . Thus, from [Theorem 23](#) for the particular case  $z = 3$  and  $\ell = 3h$ ,  $B$  should affect every  $(a, f_{13}(a, 3, 3h), f_{14}(a, 3h))$ -apex-wall triple of  $G$ , for every value of  $a$ . As  $|B| = t$ , we conclude the following.

**Corollary 25.** *If  $t, h, a \in \mathbb{N}$  and  $\mathbf{G} = (G, B, \rho)$  is a boundaried graph in  $\mathcal{R}_h^{(t)}$ , then  $B$  affects every  $(a, f_{13}(a, 3, 3h), f_{14}(a, 3h))$ -apex-wall triple of  $G$ , in particular,  $\mathbf{p}_{a, f_{13}(a, 3, 3h), f_{14}(a, 3h)}(\mathbf{G}) \leq t$ .*

## 6 Bounding the size of the representatives

In this section we use the results obtained in the previous sections to prove that every representative in  $\mathcal{R}_h^{(t)}$  has size linear in  $t$ . For this, we first prove in [Subsection 6.1](#) that every representative in  $\mathcal{R}_h^{(t)}$  has a set of  $\mathcal{O}_h(t)$  vertices containing its boundary whose removal leaves a graph with treewidth bounded by a constant depending only on the collection  $\mathcal{F}$ ; such a set is called a *treewidth modulator*.

Once we have the treewidth modulator, we can use known results from the protrusion machinery to achieve our goal. Namely, in [Subsection 6.2](#) we show how to obtain a linear protrusion decomposition of a representative, and we reduce each of the linearly many protrusions in the decomposition to an equivalent protrusion of constant size. Once we have this, a dynamic programming algorithm similar to that of [\[6\]](#) yields [Theorem 33](#).

### 6.1 Finding a treewidth modulator of linear size

Given a graph  $G$  and a set  $S \subseteq V(G)$ , we say that a separation  $(L, R)$  of  $G$  is a *2/3-balanced separation* of  $S$  in  $G$  if  $|(L \setminus R) \cap S|, |(R \setminus L) \cap S| \leq \frac{2}{3}|S|$ . We need the following well-known property of graphs of bounded treewidth (see e.g. [\[9, 14\]](#)).

**Lemma 26.** *Let  $G$  be a graph and let  $S \subseteq V(G)$ . There is a  $2/3$ -balanced separation  $(L, R)$  of  $S$  in  $G$  of order at most  $\text{tw}(G) + 1$ .*

Using Lemma 10, Lemma 11, Corollary 25, and Lemma 26 we prove the following result, whose proof uses Akra-Bazzi Theorem [3], in particular its extended provided by Leighton [35]. We stress that  $\mathbf{p}$  is not a bidimensional parameter in the precise way that is defined in [17, 18, 21], therefore Lemma 27 cannot be derived by directly applying the results of [21].

**Lemma 27.** *There exist two functions  $f_{15}, f_{16} : \mathbb{N}^2 \rightarrow \mathbb{N}$  such that if  $t, q, h \in \mathbb{N}$  and  $\mathbf{G} = (G, B, \rho)$  is a  $K_q$ -minor-free boundaried graph in  $\mathcal{R}_h^{(t)}$ , then  $G$  contains an  $f_{15}(q, h)$ -treewidth modulator that contains  $B$  and has at most  $f_{16}(q, h) \cdot t$  vertices. Moreover, it holds that  $f_{15}(q, h) = \mathcal{O}((f_5(q, f_{13}(f_3(q), 3, 3h)), f_{14}(f_3(q), 3h)))^2$ .*

*Proof.* Let  $q, h \in \mathbb{N}$ . We use  $\mathbf{q}$  as a shortcut for the triple  $(f_3(q), f_{13}(f_3(q), 3, 3h), f_{14}(f_3(q), 3h))$  and we set  $s = f_5(q, f_{13}(f_3(q), 3, 3h), f_{14}(f_3(q), 3h))$ . Let  $t_0 = \max\{\min\{t' \mid s \cdot \sqrt{t'} + s + 1 \leq t' / \log^2 t'\}, 42534179953\}$  and let  $x = s \cdot \sqrt{t_0} + s$ . We define the function  $\mathbf{z} : \mathbb{N} \rightarrow \mathbb{N}$  where

$$\mathbf{z}(t) = \min\{z \mid \forall G \forall B \subseteq V(G) \text{ if } G \text{ is } K_q\text{-minor-free, } |B| \leq t, \text{ and } B \text{ affects every } \mathbf{q}\text{-apex-wall triple of } G, \\ \text{then } \exists Z \subseteq V(G) : |Z| \leq z, B \subseteq Z, \text{tw}(G \setminus Z) \leq x\}.$$

Let  $G$  be a  $K_q$ -minor-free graph and let  $B \subseteq V(G)$  such  $|B| \leq t$  and  $B$  affects every  $\mathbf{q}$ -apex-wall triple of  $G$ . From Lemma 10,  $\text{tw}(G) \leq s \cdot \max\{1, \sqrt{|B|}\} \leq s \cdot \max\{1, \sqrt{t}\} \leq s \cdot \sqrt{t} + s$ . From Lemma 26,  $G$  has a  $2/3$ -balanced separator  $(L, R)$  of  $B$  where  $|L \cap R| \leq s \cdot \sqrt{t} + s + 1$ . This means that  $|(L \setminus R) \cap B|, |(R \setminus L) \cap B| \leq \frac{2}{3}|B|$ . We set  $G_L = G[L]$ ,  $G_R = G[R]$ ,  $B_L = L \cap (R \cup B) \subseteq V(G_L)$ , and  $B_R = R \cap (L \cup B) \subseteq V(G_R)$  and observe that  $B \subseteq B_L \cup B_R$  and that both  $G_L$  and  $G_R$  are  $K_q$ -minor-free. Notice that  $B_L = ((L \setminus R) \cap B) \cup (L \cap R)$  and  $B_R = ((R \setminus L) \cap B) \cup (R \cap L)$ , therefore there is some  $\alpha \in [\frac{1}{2}, \frac{2}{3}]$ , such that  $|B_L| \leq \alpha \cdot t + s \cdot \sqrt{t} + s + 1$  and  $|B_R| \leq (1 - \alpha) \cdot t + s \cdot \sqrt{t} + s + 1$ . From Lemma 11,  $B_L$  affects every  $\mathbf{q}$ -apex-wall triple of  $G_L$  and  $B_R$  affects every  $\mathbf{q}$ -apex-wall triple of  $G_R$ . This means that there exists some  $Z_L \subseteq V(G_L)$  such that  $|Z_L| \leq \mathbf{z}(\alpha \cdot t + s \cdot \sqrt{t} + s + 1)$ ,  $B_L \subseteq Z_L$ , and  $\text{tw}(G_L \setminus Z_L) \leq x$ . Also, there exists some  $Z_R \subseteq V(G_R)$  such that  $|Z_R| \leq \mathbf{z}((1 - \alpha) \cdot t + s \cdot \sqrt{t} + s + 1)$ ,  $B_R \subseteq Z_R$ , and  $\text{tw}(G_R \setminus Z_R) \leq x$ . We set  $Z = Z_L \cup Z_R$  and observe that  $|Z| \leq \mathbf{z}(\alpha \cdot t + s \cdot \sqrt{t} + s + 1) + \mathbf{z}((1 - \alpha) \cdot t + s \cdot \sqrt{t} + s + 1)$ . Moreover,  $B \subseteq B_L \cup B_R \subseteq Z_L \cup Z_R = Z$  and, since  $L \cap R \subseteq Z$ , it holds that  $\text{tw}(G \setminus Z) \leq x$ . We obtain that

$$\mathbf{z}(t) \leq \mathbf{z}(\alpha \cdot t + s \cdot \sqrt{t} + s + 1) + \mathbf{z}((1 - \alpha) \cdot t + s \cdot \sqrt{t} + s + 1). \quad (4)$$

Let now  $f : \mathbb{N}_{\geq 1} \rightarrow \mathbb{N}$  be the solution of the following recurrence:

$$f(t) = \begin{cases} f(\frac{t}{3} + s \cdot \sqrt{t} + s + 1) + f(\frac{2t}{3} + s \cdot \sqrt{t} + s + 1) & \text{if } t > t_0 \\ t_0 & \text{if } 1 \leq t \leq t_0 \end{cases}$$

By the choice of  $t_0$ , it holds that  $t_0 = \Theta_s(1)$ . Also, the choice of  $t_0$  is made so that the conditions for applying the extended version of Akra-Bazzi Theorem [3] provided by Leighton [35] are satisfied<sup>6</sup>. Consequently, the solution of the above recurrence is  $f(t) = \Theta_s(t^\alpha)$  where  $\alpha$  is the unique solution of the equation  $(1/3)^\alpha + (2/3)^\alpha = 1$ . Therefore  $f(t) = \Theta_s(t)$ .

Note that, if  $1 \leq t \leq t_0$ , then from Lemma 10 we have that  $\mathbf{z}(t) \leq t \leq t_0 = f(t)$ . On the other hand, by convexity, the right part of (4) is upper-bounded by  $f(t)$ , so for all  $t > t_0$  we have that  $\mathbf{z}(t) \leq f(t)$ . Summarizing, we have that  $\mathbf{z}(t) \leq f(t)$  for all  $t \geq 1$ .

Let now  $\mathbf{G} = (G, B, \rho)$  be a  $K_q$ -minor-free boundaried graph in  $\mathcal{R}_h^{(t)}$ . Applying Corollary 25 with  $a = f_3(q)$ , we obtain that  $B$  affects every  $\mathbf{q}$ -apex-wall triple of  $G$ . Therefore  $G$  contains a  $x$ -treewidth modulator

<sup>6</sup>We verified these conditions using an elementary MATLAB program, from which the number 42534179953 was generated.

that contains  $B$  and has  $\mathbf{z}(t) \leq f(t) = \mathcal{O}_s(t)$  vertices, as required. Therefore,  $\mathbf{z}(t) \leq f_{16}(q, h) \cdot t$  for some function  $f_{16} : \mathbb{N}^2 \rightarrow \mathbb{N}$ . Observe that  $t_0 = \mathcal{O}(s^2)$ , therefore  $x = \mathcal{O}(s^2)$  as well. The lemma follows with  $f_{15}(q, h) := x = \mathcal{O}(s^2)$ .  $\square$

Note that the above proof does not give any estimation on the function  $f_{16}(q, h)$ . In the Appendix (Subsection B.1) we provide an improved version of Lemma 27, namely Lemma 34, with  $f_{16}(q, h) = 2$ . This will permit us to make a better estimation of the contribution of  $h$  in the running time of our algorithm (cf. Subsection B.2). The proof of Lemma 34 is an adaptation to our setting of the one of [21, Lemma 3.6].

## 6.2 Finding a linear protrusion decomposition and reducing protrusions

Equipped with Lemma 34, the next step is to construct an appropriate protrusion decomposition of a representative. We first need to define protrusions and protrusion decompositions of graphs and boundaried graphs.

**Protrusion decompositions of unboundaried graphs.** Given a graph  $G$ , a set  $X \subseteq V(G)$  is a  $\beta$ -*protrusion* of  $G$  if  $|\partial(X)| \leq \beta$  and  $\text{tw}(G[X]) \leq \beta - 1$ . Given  $\alpha, t \in \mathbb{N}$ , an  $(\alpha, \beta)$ -*protrusion decomposition* of  $G$  is a sequence  $\mathcal{P} = \langle R_0, R_1, \dots, R_\ell \rangle$  of pairwise disjoint subsets of  $V(G)$  such that

- $\bigcup_{i \in [\ell]} R_i = V(G)$ ,
- $\max\{\ell, |R_0|\} \leq \alpha$ ,
- for  $i \in [\ell]$ ,  $N[R_i]$  is a  $\beta$ -protrusion of  $G$ , and
- for  $i \in [\ell]$ ,  $N(R_i) \subseteq R_0$ .

We call the sets  $N[R_i]$   $i \in [\ell]$ , the *protrusions* of  $\mathcal{P}$  and the set  $R_0$  the *core* of  $\mathcal{P}$ .

The above notions can be naturally generalized to boundaried graphs, just by requiring that both boundaries –of the host graph and of the protrusion– behave as one should expect, namely that the intersection of the protrusion with the boundary of the considered graph is a subset of the boundary of the protrusion.

**Protrusions and protrusion decompositions of boundaried graphs.** Given a boundaried graph  $\mathbf{G} = (G, B, \rho)$ , a *tree decomposition* of  $\mathbf{G}$  is any tree decomposition of  $G$  with a bag containing  $B$ . The *treewidth* of a boundaried graph  $\mathbf{G}$ , denoted by  $\text{tw}(\mathbf{G})$ , is the minimum width of a tree decomposition of  $\mathbf{G}$ . A boundaried graph  $\mathbf{G}' = (G', B', \rho')$  is a  $\beta$ -*protrusion* of  $\mathbf{G}$  if

- $V(G')$  is a  $\beta$ -protrusion of  $G$ ,
- $\text{tw}(\mathbf{G}') \leq \beta - 1$ ,
- $\partial(V(G')) \subseteq B'$ , and
- $B \cap V(G') \subseteq B'$ .

Given a boundaried graph  $\mathbf{G} = (G, B, \rho)$  and  $\alpha, t \in \mathbb{N}$ , an  $(\alpha, \beta)$ -*protrusion decomposition* of  $\mathbf{G}$  is a sequence  $\mathcal{P} = \langle R_0, R_1, \dots, R_\ell \rangle$  of pairwise disjoint subsets of  $V(G)$  such that

- $\bigcup_{i \in [\ell]} R_i = V(G)$ ,
- $\max\{\ell, |R_0|\} \leq \alpha$ ,
- $B \subseteq R_0$ ,

- for  $i \in [\ell]$ ,  $(G(N[R_i]), \partial(N[R_i]), \rho|_{\partial(N[R_i])})$  is a  $\beta$ -protrusion of  $\mathbf{G}$ , and
- for  $i \in [\ell]$ ,  $N(R_i) \subseteq R_0$ .

As in the unboundaried case, we call the sets  $N[R_i]$   $i \in [\ell]$ , the *protrusions* of  $\mathcal{P}$  and the set  $R_0$  the *core* of  $\mathcal{P}$ .

The following theorem is a reformulation using our notation of one of the main results of Kim et al. [32], which is stronger than what we need, in the sense that also applies to *topological-minor-graphs*. It is worth mentioning that, for  $H$ -minor-free-graphs, an appropriate protrusion decomposition can also be found using the results in [21, Lemma 3.10].

**Theorem 28.** *Let  $c, \beta, t$  be positive integers, let  $H$  be a  $q$ -vertex graph, and let  $G$  be an  $n$ -vertex  $H$ -topological-minor-free graph. If we are given a set  $M \subseteq V(G)$  with  $|M| \leq c \cdot t$  such that  $\text{tw}(G - M) \leq \beta$ , then we can compute in time  $\mathcal{O}(n)$  an  $((\alpha_H \cdot \beta \cdot c) \cdot t, 2\beta + q)$ -protrusion decomposition  $\mathcal{P}$  of  $G$  with  $M$  contained in the core of  $\mathcal{P}$ , where  $\alpha_H$  is a constant depending only on  $H$  such that  $\alpha_H \leq 40q^2 2^{5q \log q}$ .*

Having stated the above definitions, the following lemma is an easy consequence of Lemma 34 and Theorem 28.

**Lemma 29.** *There exists a function  $f_{17} : \mathbb{N}^2 \rightarrow \mathbb{N}$  such that if  $t, q, h \in \mathbb{N}$  and  $\mathbf{G} = (G, B, \rho)$  is a  $K_q$ -minor-free boundaried graph in  $\mathcal{R}_h^{(t)}$ , then  $\mathbf{G}$  admits a  $(f_{17}(q, h) \cdot t, f_{17}(q, h))$ -protrusion decomposition. Moreover, it holds that  $f_{17}(q, h) = f_{15}(q, h) \cdot f_{16}(q, h) \cdot 2^{\mathcal{O}(q \log q)}$ .*

*Proof.* By Lemma 34,  $G$  contains an  $f_{21}(q, h)$ -treewidth modulator  $M$  that contains  $B$  and has at most  $2t$  vertices. We can now apply Theorem 28 to  $G$  and  $M$  with  $H = K_q$ ,  $c = f_{16}(q, h)$ , and  $\beta = f_{15}(q, h)$ , obtaining a  $(f_{17}(q, h) \cdot t, f_{17}(q, h))$ -protrusion decomposition  $\mathcal{P}$  of  $G$  with  $M$  contained in the core of  $\mathcal{P}$  and  $f_{17}(q, h) := f_{15}(q, h) \cdot f_{16}(q, h) \cdot 40q^2 2^{5q \log q}$ . Since  $B \subseteq M$  and  $M$  contained in the core of  $\mathcal{P}$ , it can be easily checked that  $\mathcal{P}$  is also a  $(f_{17}(q, h) \cdot t, f_{17}(q, h))$ -protrusion decomposition of  $\mathbf{G}$ .  $\square$

Once we have the protrusion decomposition given by Lemma 29, all that remains is to replace the protrusions by equivalent ones of size depending only on the collection  $\mathcal{F}$ . The protrusion replacement technique, which is nowadays part of the basic toolbox of parameterized complexity, originated in the meta-theorem of Bodlaender et al. [11], whose objective was to produce linear kernels for a wide family of problems on graphs of bounded genus. This technique was later extended to graphs excluding a fixed minor by Fomin et al. [21] and then to graphs excluding a fixed topological minor by Kim et al. [32]. We could directly apply the results of Fomin et al. [21] to the protrusion decomposition of a representative given by Lemma 29, hence reducing each protrusion to an equivalent one of size  $\mathcal{O}_{\mathcal{F}}(1)$ , yielding an equivalent representative of size  $\mathcal{O}_{\mathcal{F}}(t)$ . However, the drawback of the results in [21] (and also in [11, 32]) is that they do not provide *explicit* bounds on the hidden constants. In order to be able to do so (cf. Subsection B.2), we apply the protrusion replacement used by Baste et al. [6], which is suited for the  $\mathcal{F}$ -M-DELETION problem. This yields explicit constants because it uses ideas similar to the ones presented by Garnero et al. [22] (later generalized in [23]) for obtaining kernels with explicit constants.

Given a function  $\xi : \mathbb{N}^2 \rightarrow \mathbb{N}$  and a  $t$ -boundaried graph  $\mathbf{G}$ , we say that  $\mathbf{G}$  is  $\xi$ -protrusion-bounded if, for every  $t' \in \mathbb{N}$ , all  $\beta$ -protrusions of  $\mathbf{G}$  have at most  $\xi(\beta)$  vertices. The following lemma is again a reformulation using our notation of one of the results of Baste et al. [6]. Namely, it is a consequence of the proof<sup>7</sup> of [6, Lemma 7.2].

<sup>7</sup>In the statement of [6, Lemma 7.2] it is required that the family  $\mathcal{F}$  contains a planar graph, an assumption that is not true anymore in our case. However, in the proof this fact is only used to guarantee that the considered protrusion has treewidth bounded by a function depending only on  $\mathcal{F}$ . Thanks to Lemma 29, we can assume that this also holds in our setting.

**Lemma 30.** *There exists a function  $f_{18} : \mathbb{N}^2 \rightarrow \mathbb{N}$  such that if  $t, q, h \in \mathbb{N}$  and  $\mathbf{G} = (G, B, \rho)$  is a  $K_q$ -minor-free boundaried graph in  $\mathcal{R}_h^{(t)}$ , then  $\mathbf{G}$  is  $f_{18}(q, h)$ -protrusion-bounded. Moreover,  $f_{18}(q, h) = 2^{2^{\mathcal{O}(f_{17}(q, h) \cdot \log f_{17}(q, h))}}$ .*

Using Lemma 29 and Lemma 30, we can easily prove Theorem 31, that is the main result on which the algorithm of Theorem 2 is based (cf. Section 2). In particular, it implies Equation 1.

**Theorem 31.** *There exists a function  $f_{19} : \mathbb{N}^2 \rightarrow \mathbb{N}$  such that, for every  $t \in \mathbb{N}$  and  $q, h \in \mathbb{N}_{\geq 1}$ , if  $\mathbf{G} = (G, B, \rho)$  is a  $K_q$ -minor-free boundaried graph in  $\mathcal{R}_h^{(t)}$ , then  $|V(G)| \leq f_{19}(q, h) \cdot t$ . Moreover, it holds that  $f_{19}(q, h) \leq f_{17}(q, h) \cdot (f_{18}(q, h) + 1)$ .*

*Proof.* By Lemma 29,  $\mathbf{G}$  admits an  $(f_{17}(q, h) \cdot t, f_{17}(q, h))$ -protrusion decomposition  $\mathcal{P}$ . By Lemma 30, each of the protrusions of  $\mathcal{P}$  has at most  $f_{18}(q, h)$  vertices. Therefore,

$$|V(G)| \leq f_{17}(q, h) \cdot t + f_{17}(q, h) \cdot f_{18}(q, h) \cdot t,$$

and the theorem follows with  $f_{19}(q, h) := f_{17}(q, h) \cdot (f_{18}(q, h) + 1)$ .  $\square$

Let  $h := \max_{F \in \mathcal{F}} \{\max_{H \in \text{ext}(F)} \text{detail}(H)\}$ . The following corollary is an immediate consequence of Theorem 31, by using the fact that all  $t$ -representatives in  $\mathcal{R}_h^{(t)}$ , except one, are  $K_h$ -minor-free, hence they have  $\mathcal{O}(f_{19}(h, h) \cdot h \sqrt{\log h}) \cdot t$  edges; see for instance [38]. Note that there are at most  $\binom{n^2}{m} = 2^{\mathcal{O}(n \log m)}$  different graphs on  $n$  vertices and  $m$  edges and that, if  $(G, B, \rho) \in \mathcal{R}_h^{(t)}$ , then Theorem 31 implies that  $|V(G)| \leq f_{19}(h, h) \cdot t$ . Note also that there are  $\binom{|V(G)|}{t} = 2^{\mathcal{O}(t \log |V(G)|)}$  choices for  $B$ , and  $t! = 2^{\mathcal{O}(t \log t)}$  choices for  $\rho$ . Therefore,  $|\mathcal{R}_h^{(t)}| = 2^{\mathcal{O}(f_{19}(h, h) \cdot t \cdot \log(f_{19}(h, h) \cdot h \sqrt{\log h \cdot t}) + f_{19}(h, h) \cdot t \log(f_{19}(h, h) \cdot t) + t \log t)}$  and we can conclude the following.

**Corollary 32.** *There exists a function  $f_{20} : \mathbb{N} \rightarrow \mathbb{N}$  such that for every  $t \in \mathbb{N}_{\geq 1}$ ,  $|\mathcal{R}_h^{(t)}| \leq 2^{f_{20}(h) \cdot t \cdot \log t}$ . In particular, the relation  $\equiv_h$  partitions  $\mathcal{B}^{(t)}$  into at most  $2^{f_{20}(h) \cdot t \cdot \log t}$  equivalence classes. Moreover, it holds that  $f_{20}(h) = \mathcal{O}(f_{19}(h, h) \cdot \log(f_{19}(h, h) \cdot h \sqrt{\log h}))$ .*

**The dynamic programming algorithm.** Having proved Corollary 32, we can apply [6, Theorem 8.1] to compute the parameter  $\mathbf{m}_{\mathcal{F}}(G)$  within the claimed running time.

For the sake of completeness, let us comment some details of this dynamic programming algorithm, whose details can be found in [6, Section 8]. First of all, to run the algorithm we need to have the set  $\mathcal{R}_h^{(t)}$  of representatives at hand. This can be done easily relying on Theorem 31, by generating all  $t$ -boundaried graphs on at most  $f_{19}(h, h) \cdot t$  vertices and  $\mathcal{O}(f_{19}(h, h) \cdot h \sqrt{\log h}) \cdot t$  edges, partitioning them into equivalence classes according to  $\equiv_h$ , and picking an element of minimum size in each of them; see [6, proof of Lemma 7.1] for more details. To simplify the description of the dynamic programming update operations, the main algorithm in [6] is written in terms of *branchwidth* instead of *treewidth*. Without defining branchwidth here, it is enough to say that it is linearly equivalent to treewidth, in the sense that both parameters differ by a constant factor and whose corresponding decompositions can be easily transformed from one to the other [41]. Also, the main algorithm in [6] is written in terms of *topological* minors, that is, given a finite graph class  $\mathcal{F}'$  and a graph  $G$ , it computes  $\mathbf{tm}_{\mathcal{F}'}(G)$ , that is the minimum-size set of vertices  $S \subseteq V(G)$  whose removal leaves a graph without any of the graphs in a fixed collection  $\mathcal{F}'$  as a *topological* minor. This works for our purposes because of the translation of the question on minors to one on topological minors, provided by Observation 3. The dynamic algorithm computes, in a typical bottom-up manner, at every bag separator  $B$  of the branch decomposition associated with a  $t$ -boundaried graph  $\mathbf{G}_B$  and for every representative  $\mathbf{R} \in \mathcal{R}_h^{(t)}$ , the minimum size of a set  $S \subseteq V(\mathbf{G}_B)$  such that  $\mathbf{G}_B \setminus S \equiv_h \mathbf{R}$ . These values can be computed in a standard way by combining the values associated with the children of a given node; cf. [6, Theorem 8.1]. The overall running time is bounded by  $\mathcal{O}(|\mathcal{R}_h^{(t)}|^2 \cdot |E(G)|)$ , and taking into account that  $|E(G)| \leq \text{tw}(G) \cdot |V(G)|$ , from Corollary 32 we obtain the following theorem, which is a more precise reformulation of Theorem 2.

**Theorem 33.** *Let  $t, h \in \mathbb{N}$ ,  $\mathcal{F}$  be a proper collection of size at most  $h$ , and  $G$  be an  $n$ -vertex graph of treewidth at most  $t$ . Then  $\mathbf{m}_{\mathcal{F}}(G)$  can be computed by an algorithm that runs in  $2^{\mathcal{O}(f_{20}(h) \cdot t \log t)} \cdot n$  steps.*

In [Appendix B](#) we give upper bounds on the constants depending on the collection  $\mathcal{F}$  involved in our algorithm. These upper bounds depend explicitly on the parametric dependencies of the Unique Linkage Theorem [\[31, 43\]](#).

## 7 Further research

We presented an algorithm for solving the  $\mathcal{F}$ -M-DELETION problem in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$  for every collection  $\mathcal{F}$ . This algorithm together with the single-exponential algorithms and lower bounds presented in previous papers of this series [\[7, 8\]](#) yield a complete classification of the asymptotic complexity of  $\mathcal{F}$ -M-DELETION parameterized by treewidth assuming the ETH, when  $\mathcal{F} = \{H\}$  and  $H$  is connected ([Theorem 1](#)). However, we do not have a complete classification when  $|\mathcal{F}| \geq 2$ , even for connected  $\mathcal{F}$ . To ease the presentation, let us call a connected graph  $H$  *easy* (resp. *hard*) if  $\{H\}$ -M-DELETION is solvable in time  $\mathcal{O}^*(2^{\Theta(\text{tw})})$  (resp.  $\mathcal{O}^*(2^{\Theta(\text{tw} \cdot \log \text{tw})})$ ). Suppose that  $\mathcal{F} = \{H_1, H_2\}$  with both  $H_1$  and  $H_2$  being connected. Using the recent results of Baste [\[5\]](#), it is possible to prove that if both  $H_1$  and  $H_2$  are easy, then  $\mathcal{F}$  is easy as well (easiness of graph collections is defined in the obvious way). However, if both  $H_1$  and  $H_2$  are hard, then strange things may happen. For instance, Bodlaender et al. [\[12\]](#) presented an algorithm running in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw})})$  for PSEUDOFORREST DELETION, which consists in, given a graph  $G$  and an integer  $k$ , deciding whether one can delete at most  $k$  vertices from  $G$  to obtain a *pseudoforest*, i.e., a graph where each connected component contains at most one cycle. Note that PSEUDOFORREST DELETION is equivalent to  $\{\text{diamond}, \text{butterfly}\}$ -M-DELETION. While both the *diamond* and the *butterfly* are hard graphs (cf. [Figure 17](#)),  $\{\text{diamond}, \text{butterfly}\}$  is an easy collection. The cases where  $H_1$  is easy and  $H_2$  is hard seem even trickier. Obtaining (tight) lower bounds when  $\mathcal{F}$  may contain disconnected graphs is another challenging avenue for further research.

It is also interesting to consider the version of the problem where the graphs in  $\mathcal{F}$  are forbidden as *topological* minors; we call this problem  $\mathcal{F}$ -TM-DELETION. While most of the lower bounds that we presented in [\[8\]](#) also hold for  $\mathcal{F}$ -TM-DELETION, the algorithm in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$  of this paper does *not* work for topological minors. In this direction, the algorithm in time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$  for  $\mathcal{F}$ -M-DELETION when  $\mathcal{F}$  contains a *planar* graph given in [\[6\]](#) also works for  $\mathcal{F}$ -TM-DELETION, if we additionally require  $\mathcal{F}$  to contain a *subcubic* planar graph (in order to bound the treewidth of the representatives). The main obstacle for applying our approach in order to achieve a time  $\mathcal{O}^*(2^{\mathcal{O}(\text{tw} \cdot \log \text{tw})})$  for every collection  $\mathcal{F}$ , is that topological-minor-free graphs do not enjoy the flat wall structure that is omnipresent in our proofs. Another reason is that in our rerouting procedure, in order to find an irrelevant vertex ([Theorem 17](#)), we may find a *different* topological minor model that corresponds to the same minor. Nevertheless, we think that this latter difficulty can be overcome for planar graphs –or even minor-free graphs– by making use of the rerouting potential of [Proposition 13](#), as this is done in [\[25\]](#) for planar graphs.

Finally, it is worth mentioning that the algorithm presented in this paper, as well as the main combinatorial result ([Theorem 23](#)), have been used in [\[44\]](#) (see [\[47\]](#) for the full version) to obtain a fixed-parameter algorithm for the  $\mathcal{F}$ -M-DELETION problem parameterized by  $k$ . [Theorem 23](#) has also been used in [\[46\]](#) in order to provide explicit upper bounds on the size of the minor-obstructions of the set of yes-instances of the  $\mathcal{F}$ -M-DELETION problem, as a function of  $\mathcal{F}$  and  $k$ .

## References

- [1] Isolde Adler, Stavros G. Kolliopoulos, Philipp Klaus Krause, Daniel Lokshantov, Saket Saurabh, and Dimitrios M. Thilikos. Irrelevant vertices for the planar Disjoint Paths Problem. *Journal of Combina-*

- torial Theory, Series B*, 122:815–843, 2017. doi:10.1016/j.jctb.2016.10.001.
- [2] Isolde Adler and Philipp Klaus Krause. A lower bound on the tree-width of graphs with irrelevant vertices. *Journal of Combinatorial Theory, Series B*, 137:126–136, 2019. doi:10.1016/j.jctb.2018.12.008.
- [3] Mohamad A. Akra and Louay Bazzi. On the solution of linear recurrence equations. *Computational Optimization and Applications*, 10(2):195–210, 1998. doi:10.1023/A:1018373005182.
- [4] Zongwen Bai, Jianhua Tu, and Yongtang Shi. An improved algorithm for the vertex cover  $P_3$  problem on graphs of bounded treewidth. *Discrete Mathematics & Theoretical Computer Science*, 21(4), 2019. doi:10.23638/DMTCS-21-4-17.
- [5] Julien Baste. Composing dynamic programming tree-decomposition-based algorithms. Manuscript, 2019. arXiv:1904.12500.
- [6] Julien Baste, Ignasi Sau, and Dimitrios M. Thilikos. Hitting minors on bounded treewidth graphs. I. General upper bounds. *SIAM Journal on Discrete Mathematics*, 34(3):1623–1648, 2020. doi:10.1137/19M1287146.
- [7] Julien Baste, Ignasi Sau, and Dimitrios M. Thilikos. Hitting minors on bounded treewidth graphs. II. Single-exponential algorithms. *Theoretical Computer Science*, 814:135–152, 2020. doi:10.1016/j.tcs.2020.01.026.
- [8] Julien Baste, Ignasi Sau, and Dimitrios M. Thilikos. Hitting minors on bounded treewidth graphs. III. Lower bounds. *Journal of Computer and System Sciences*, 109:56–77, 2020. doi:10.1016/j.jcss.2019.11.002.
- [9] Hans L. Bodlaender. A Partial  $k$ -Arboretum of Graphs with Bounded Treewidth. *Theoretical Computer Science*, 209(1-2):1–45, 1998. doi:10.1016/S0304-3975(97)00228-4.
- [10] Hans L. Bodlaender, Marek Cygan, Stefan Kratsch, and Jesper Nederlof. Deterministic single exponential time algorithms for connectivity problems parameterized by treewidth. *Information and Computation*, 243:86–111, 2015. doi:10.1016/j.ic.2014.12.008.
- [11] Hans L. Bodlaender, Fedor V. Fomin, Daniel Lokshtanov, Eelko Penninkx, Saket Saurabh, and Dimitrios M. Thilikos. (Meta) Kernelization. *Journal of the ACM*, 63(5):44:1–44:69, 2016. doi:10.1145/2973749.
- [12] Hans L. Bodlaender, Hirotaka Ono, and Yota Otachi. A faster parameterized algorithm for pseudoforest deletion. *Discrete Applied Mathematics*, 236:42–56, 2018. doi:10.1016/j.dam.2017.10.018.
- [13] Bruno Courcelle. The Monadic Second-Order Logic of Graphs. I. Recognizable Sets of Finite Graphs. *Information and Computation*, 85(1):12–75, 1990. doi:10.1016/0890-5401(90)90043-H.
- [14] Marek Cygan, Fedor V. Fomin, Lukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michal Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015. doi:10.1007/978-3-319-21275-3.
- [15] Marek Cygan, Dániel Marx, Marcin Pilipczuk, and Michal Pilipczuk. Hitting forbidden subgraphs in graphs of bounded treewidth. *Information and Computation*, 256:62–82, 2017. doi:10.1016/j.ic.2017.04.009.

- [16] Marek Cygan, Jesper Nederlof, Marcin Pilipczuk, Michal Pilipczuk, Johan M. M. van Rooij, and Jakub Onufry Wojtaszczyk. Solving Connectivity Problems Parameterized by Treewidth in Single Exponential Time. In *Proc. of the 52nd IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 150–159, 2011. doi:10.1109/FOCS.2011.23.
- [17] Erik D. Demaine, Fedor V. Fomin, Mohammadtaghi Hajiaghayi, and Dimitrios M. Thilikos. Subexponential parameterized algorithms on bounded-genus graphs and  $H$ -minor-free graphs. *Journal of the ACM*, 52(6):866–893, 2005. doi:10.1145/1101821.1101823.
- [18] Fedor V. Fomin, Erik D. Demaine, Mohammad Taghi Hajiaghayi, and Dimitrios M. Thilikos. Bidimensionality. In *Encyclopedia of Algorithms*, pages 203–207. Springer, 2016. doi:10.1007/978-1-4939-2864-4\_47.
- [19] Fedor V. Fomin and Dieter Kratsch. *Exact Exponential Algorithms*. Texts in Theoretical Computer Science. An EATCS Series. Springer, 2010. doi:10.1007/978-3-642-16533-7.
- [20] Fedor V. Fomin, Daniel Lokshtanov, Fahad Panolan, and Saket Saurabh. Efficient computation of representative families with applications in parameterized and exact algorithms. *Journal of the ACM*, 63(4):29:1–29:60, 2016. doi:10.1145/2886094.
- [21] Fedor V. Fomin, Daniel Lokshtanov, Saket Saurabh, and Dimitrios M. Thilikos. Bidimensionality and kernels. *SIAM Journal on Computing*, 49(6):1397–1422, 2020. doi:10.1137/16M1080264.
- [22] Valentin Garnero, Christophe Paul, Ignasi Sau, and Dimitrios M. Thilikos. Explicit linear kernels via dynamic programming. *SIAM Journal on Discrete Mathematics*, 29(4):1864–1894, 2015. doi:10.1137/140968975.
- [23] Valentin Garnero, Christophe Paul, Ignasi Sau, and Dimitrios M. Thilikos. Explicit linear kernels for packing problems. *Algorithmica*, 81(4):1615–1656, 2019. doi:10.1007/s00453-018-0495-5.
- [24] Petr A. Golovach, Marcin Kaminski, Spyridon Maniatis, and Dimitrios M. Thilikos. The parameterized complexity of graph cyclability. *SIAM Journal on Discrete Mathematics*, 31(1):511–541, 2017. doi:10.1137/141000014.
- [25] Petr A. Golovach, Giannos Stamoulis, and Dimitrios M. Thilikos. Hitting topological minor models in planar graphs is fixed parameter tractable. In *Proc. of the 31st ACM-SIAM Symposium on Discrete Algorithms, (SODA)*, pages 931–950, 2020. doi:10.1137/1.9781611975994.56.
- [26] Russell Impagliazzo, Ramamohan Paturi, and Francis Zane. Which problems have strongly exponential complexity? *Journal of Computer and System Sciences*, 63(4):512–530, 2001. doi:10.1006/jcss.2001.1774.
- [27] Bart M. P. Jansen, Daniel Lokshtanov, and Saket Saurabh. A near-optimal planarization algorithm. In *Proc. of the 25th ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1802–1811, 2014. doi:10.1137/1.9781611973402.130.
- [28] Marcin Kaminski and Dimitrios M. Thilikos. Contraction checking in graphs on surfaces. In *Proc. of the 29th International Symposium on Theoretical Aspects of Computer Science, (STACS)*, volume 14 of *LIPICs*, pages 182–193, 2012. doi:10.4230/LIPICs.STACS.2012.182.
- [29] Goossen Kant. *Algorithm for Drawing Planar Graphs*. PhD thesis, Department of Information and Computing Sciences - Utrecht University, 1993. URL: <https://dspace.library.uu.nl/bitstream/handle/1874/842/full.pdf>.

- [30] Ken-ichi Kawarabayashi, Robin Thomas, and Paul Wollan. A new proof of the flat wall theorem. *Journal of Combinatorial Theory, Series B*, 129:204–238, 2018. doi:10.1016/j.jctb.2017.09.006.
- [31] Ken-ichi Kawarabayashi and Paul Wollan. A shorter proof of the graph minor algorithm: the unique linkage theorem. In *Proc. of the 42nd ACM Symposium on Theory of Computing (STOC)*, pages 687–694, 2010. doi:10.1145/1806689.1806784.
- [32] Eun Jung Kim, Alexander Langer, Christophe Paul, Felix Reidl, Peter Rossmanith, Ignasi Sau, and Somnath Sikdar. Linear kernels and single-exponential algorithms via protrusion decompositions. *ACM Transactions on Algorithms*, 12(2):21:1–21:41, 2016. doi:10.1145/2797140.
- [33] Tomasz Kociumaka and Marcin Pilipczuk. Deleting vertices to graphs of bounded genus. *Algorithmica*, 81(9):3655–3691, 2019. doi:10.1007/s00453-019-00592-7.
- [34] Stefan Kratsch and Magnus Wahlström. Representative sets and irrelevant vertices: New tools for kernelization. *Journal of the ACM*, 67(3):16:1–16:50, 2020. doi:10.1145/3390887.
- [35] Tom Leighton. Notes on Better Master Theorems for Divide-and-Conquer Recurrences, 1996. URL: <https://courses.csail.mit.edu/6.046/spring04/handouts/akrabazzi.pdf>.
- [36] John M. Lewis and Mihalis Yannakakis. The node-deletion problem for hereditary properties is NP-complete. *Journal of Computer and System Sciences*, 20(2):219–230, 1980. doi:10.1016/0022-0000(80)90060-4.
- [37] Daniel Lokshtanov, Dániel Marx, and Saket Saurabh. Slightly superexponential parameterized problems. *SIAM Journal on Computing*, 47(3):675–702, 2018. doi:10.1137/16M1104834.
- [38] Joseph Samuel Myers and Andrew Thomason. The extremal function for noncomplete minors. *Combinatorica*, 25(6):725–753, 2005. doi:10.1007/s00493-005-0044-0.
- [39] Marcin Pilipczuk. A tight lower bound for Vertex Planarization on graphs of bounded treewidth. *Discrete Applied Mathematics*, 231:211–216, 2017. doi:10.1016/j.dam.2016.05.019.
- [40] Bruce A. Reed. Rooted routing in the plane. *Discrete Applied Mathematics*, 57(2-3):213–227, 1995. doi:10.1016/0166-218X(94)00104-L.
- [41] Neil Robertson and Paul D. Seymour. Graph Minors. X. Obstructions to Tree decomposition. *Journal of Combinatorial Theory, Series B*, 52(2):153–190, 1991. doi:10.1016/0095-8956(91)90061-N.
- [42] Neil Robertson and Paul D. Seymour. Graph Minors. XIII. The Disjoint Paths Problem. *Journal of Combinatorial Theory, Series B*, 63(1):65–110, 1995. doi:10.1006/jctb.1995.1006.
- [43] Neil Robertson and Paul D. Seymour. Graph Minors. XXII. Irrelevant Vertices in Linkage Problems. *Journal of Combinatorial Theory, Series B*, 102(2):530–563, 2012. doi:10.1016/j.jctb.2007.12.007.
- [44] Ignasi Sau, Giannos Stamoulis, and Dimitrios M. Thilikos. An FPT-algorithm for recognizing  $k$ -apices of minor-closed graph classes. In *Proc. of the 47th International Colloquium on Automata, Languages, and Programming (ICALP)*, volume 168 of *LIPICs*, pages 95:1–95:20, 2020. doi:10.4230/LIPICs.ICALP.2020.95.
- [45] Ignasi Sau, Giannos Stamoulis, and Dimitrios M. Thilikos. A more accurate view of the Flat Wall Theorem. Manuscript, 2021. arXiv:2102.06463.

- [46] Ignasi Sau, Giannos Stamoulis, and Dimitrios M. Thilikos. *k*-apices of minor-closed graph classes. I. Bounding the obstructions. Manuscript, 2021. [arXiv:2103.00882](https://arxiv.org/abs/2103.00882).
- [47] Ignasi Sau, Giannos Stamoulis, and Dimitrios M. Thilikos. *k*-apices of minor-closed graph classes. II. Parameterized algorithms. Manuscript, 2021. [arXiv:2004.12692](https://arxiv.org/abs/2004.12692).
- [48] Jianhua Tu, Lidong Wu, Jing Yuan, and Lei Cui. On the vertex cover  $P_3$  problem parameterized by treewidth. *Journal of Combinatorial Optimization*, 34(2):414–425, 2017. doi:10.1007/s10878-016-9999-6.
- [49] David P. Williamson and David B. Shmoys. *The Design of Approximation Algorithms*. Cambridge University Press, 2011. URL: <https://www.designofapproxalgs.com>.

## A Illustration of the complexity dichotomy




Figure 17: Classification of the complexity of  $\{H\}$ -M-DELETION for all connected simple graphs  $H$  with  $2 \leq |V(H)| \leq 5$ , according to our results: for the nine graphs on the left, the problem is solvable in time  $2^{\Theta(\text{tw})} \cdot n^{\mathcal{O}(1)}$  under the ETH. For the 21 graphs on the right and for all the connected graphs on at least six vertices, the problem is solvable in time  $2^{\Theta(\text{tw} \cdot \log \text{tw})} \cdot n^{\mathcal{O}(1)}$  under the ETH.

## B An estimation of the constants depending on $\mathcal{F}$ in our algorithm

The main result of this paper is that the  $\mathcal{F}$ -M-DELETION problem can be solved in time  $2^{\mathcal{O}(f(h) \cdot \text{tw} \cdot \log \text{tw})} \cdot n$  on  $n$ -vertex graphs of treewidth at most  $\text{tw}$ , for some computable function  $f : \mathbb{N} \rightarrow \mathbb{N}$ , where here  $h$  is an upper bound on the size of the graphs in  $\mathcal{F}$ . This appendix is dedicated to an estimation of an upper bound on the function  $f$ . Notice that almost all the statements of the results in this paper are accompanied with specific bounds on the involved functions, usually in terms of functions defined in previous statements. However, there are two exceptions. The first one is the function  $f_6$  of [Proposition 35](#), which we discuss in [Subsection B.2](#). The second one is [Lemma 27](#), where no explicit bound for  $f_{16}(q, h)$  is given. This is because the existence of  $f_{16}(q, h)$  follows by applying Akra-Bazzi Theorem [\[3\]](#) as a black box and this does not provide any estimation of  $f_{16}$ . To circumvent this issue, in [Subsection B.1](#) we provide an improved version of [Lemma 27](#), namely [Lemma 34](#), whose proof uses a direct induction, without invoking the Akra-Bazzi Theorem [\[3, 35\]](#). This alternative proof is strongly based on the proof of [\[21, Lemma 3.6\]](#). Finally, in

[Subsection B.2](#) we provide an upper bound on the constants involving  $\mathcal{F}$  in our algorithm. For this, we will use the stronger version of [Lemma 27](#) given in [Subsection B.1](#).

## B.1 An improved version of [Lemma 27](#)

In this section we provide an improved version of [Lemma 27](#), whose proof is an adaptation of the proof of [\[21, Lemma 3.6\]](#).

**Lemma 34.** *There exists a function  $f_{21} : \mathbb{N}^2 \rightarrow \mathbb{N}$  such that if  $t, q, h \in \mathbb{N}$  and  $\mathbf{G} = (G, B, \rho)$  is a  $K_q$ -minor-free bounded graph in  $\mathcal{R}_h^{(t)}$ , then  $G$  contains an  $f_{21}(q, h)$ -treewidth modulator that contains  $B$  and has at most  $2t$  vertices. Moreover, it holds that  $f_{21}(q, h) = \mathcal{O}((f_5(q, f_{13}(f_3(q), 3, 3h), f_{14}(f_3(q), 3h)))^2)$ .*

*Proof.* For simplicity, we use  $q$  as a shortcut for the triple  $(f_3(q), f_{13}(f_3(q), 3, 3h), f_{14}(f_3(q), 3h))$ . We define the constants  $s = f_5(q, f_{13}(f_3(q), 3, 3h), f_{14}(f_3(q), 3h))$ ,  $t_0 = 256s^2$ , and  $c = s \cdot \sqrt{t_0}$ . We define the relation  $\leq_\circ$  so that  $a \leq_\circ b$  means that  $a \leq \max\{0, b\}$ . We first prove, by induction on  $t$ , the following statement.

*Claim:* For every non-negative integer  $t$  and every  $K_q$ -minor-free graph  $G$ , if  $\mathbf{p}_q(G) \leq t$  then  $G$  has a  $c$ -treewidth modulator  $Z$  with  $|Z| \leq_\circ t - 16s \cdot \sqrt{t}$ .

*Proof of claim:* In the base case we consider any  $t$  with  $0 \leq t \leq t_0$ . As  $\mathbf{p}_q(G) \leq t$ , [Lemma 10](#) implies that  $\text{tw}(G) \leq s \cdot \max\{\sqrt{t}, 1\} \leq s \cdot \sqrt{t_0}$ . Thus  $G$  has a  $c$ -treewidth modulator of size  $0 \leq_\circ t - 16s \cdot \sqrt{t}$ , and the claim follows.

For the inductive step, let  $t > t_0$  and suppose that the claim is true for every  $t'$  with  $0 \leq t' \leq t - 1$ . We prove that the claim holds also for  $t$ . Consider a graph  $G$  with  $\mathbf{p}_q(G) \leq t$  and let  $S$  be a set of at most  $t$  vertices affecting every  $q$ -apex-wall triple of  $G$ . Because of [Lemma 10](#),  $\mathbf{p}_q(G) \leq t$  implies that  $\text{tw}(G) \leq s \cdot \max\{\sqrt{t}, 1\} = s \cdot \sqrt{t}$ .

By applying [Lemma 26](#) to  $G$  and  $S$ , there is a  $2/3$ -balanced separation  $(L, R)$  of  $S$  in  $G$  such that  $|L \cap R| \leq \text{tw}(G) + 1 \leq s \cdot \sqrt{t} + 1$  and there exists some  $\alpha \in [\frac{1}{3}, \frac{2}{3}]$  such that  $|(L \setminus R) \cap S| \leq \alpha \cdot |S| \leq \alpha \cdot t$  and  $|(R \setminus L) \cap S| \leq (1 - \alpha) \cdot |S| \leq (1 - \alpha) \cdot t$ .

Since  $S$  affects every  $q$ -apex-wall triple of  $G$ , [Lemma 11](#) gives that the set  $L \cap (R \cup S)$  affects every  $q$ -apex-wall triple of  $G[L]$ . This implies

$$\mathbf{p}_q(G[L]) \leq |L \cap (R \cup S)| = |(L \setminus R) \cap S| + |L \cap R| \leq \alpha \cdot t + (s \cdot \sqrt{t} + 1) \leq \alpha \cdot t + 2s \cdot \sqrt{t}.$$

Here the last inequality follows from the assumption that  $t \geq t_0 \geq 1$ .

In order to apply the inductive hypothesis, note that  $t' := \alpha \cdot t + 2s \cdot \sqrt{t} \leq t - 1$  for  $t \geq t_0$ . This can be verified by using the fact that  $s \geq 1$ ,  $\alpha \leq \frac{2}{3}$ , and checking that  $\frac{2}{3}t + 2s \cdot \sqrt{t} \leq t - 1$  for  $t \geq t_0$ . Indeed, the inequality holds whenever  $\sqrt{t} \geq 3s + \frac{3}{2} \sqrt{(2s)^2 + \frac{4}{3}}$  and this is the case as  $\sqrt{t} \geq \sqrt{t_0} = 16s \geq 3 \cdot s + \frac{3}{2} \sqrt{(2s)^2 + \frac{4}{3}}$ .

Therefore we can apply the induction hypothesis to  $G[L]$  and  $t'$  and obtain a  $c$ -treewidth modulator  $Z_L$  of  $G[L]$ , such that

$$|Z_L| \leq_\circ t' - 16s \cdot \sqrt{t'} \leq_\circ (\alpha \cdot t + 2s \cdot \sqrt{t}) - 16s \cdot \sqrt{\alpha \cdot t + 2s \cdot \sqrt{t}} \leq_\circ (\alpha \cdot t + 2s \cdot \sqrt{t}) - 16s \cdot \sqrt{\alpha \cdot t}.$$

A symmetric argument applied to  $G[R]$  yields a treewidth modulator  $Z_R$  of  $G[R]$ , such that

$$|Z_R| \leq_\circ \left( (1 - \alpha) \cdot t + 2s \cdot \sqrt{t} \right) - 16s \cdot \sqrt{(1 - \alpha) \cdot t}.$$

We now construct a  $c$ -treewidth modulator  $Z$  of  $G$  as follows by setting  $Z := Z_L \cup (L \cap R) \cup Z_R$ . The set  $Z$  is a  $c$ -treewidth-modulator of  $G$  because every connected component of  $G - Z$  is a subset of either

$(L \setminus (L \cap R)) \setminus Z_L$  or  $(R \setminus (L \cap R)) \setminus Z_R$ , and  $Z_L$  and  $Z_R$  are  $c$ -treewidth modulators for  $G[L]$  and  $G[R]$  respectively. Finally we bound the size of  $Z$ .

$$\begin{aligned}
|Z| &\leq |Z_L| + |Z_R| + |S| \\
&\leq (\alpha \cdot t + 2s \cdot \sqrt{t}) - 16s \cdot \sqrt{t \cdot \alpha} + \left( (1 - \alpha) \cdot t + 2s \cdot \sqrt{t} \right) - 16s \cdot \sqrt{t \cdot (1 - \alpha)} + s \cdot \sqrt{t} + 1 \\
&\leq (\alpha \cdot t + 2s \cdot \sqrt{t}) - 16s \cdot \sqrt{t \cdot \alpha} + \left( (1 - \alpha) \cdot t + 2s \cdot \sqrt{t} \right) - 16s \cdot \sqrt{t \cdot (1 - \alpha)} + 2s \cdot \sqrt{t} \\
&\leq t - (16(\sqrt{\alpha} + \sqrt{1 - \alpha}) - 6) \cdot s \cdot \sqrt{t} \\
&\leq t - 16s \cdot \sqrt{t}.
\end{aligned}$$

The last inequality uses the fact that  $16(\sqrt{\alpha} + \sqrt{1 - \alpha}) - 6 \geq 16$ , for every  $\alpha \in [\frac{1}{3}, \frac{2}{3}]$ . The claim follows.

Suppose now that  $\mathbf{G} = (G, B, \rho)$  is a  $K_q$ -minor-free boundaried graph in  $\mathcal{R}_h^{(t)}$ . From [Corollary 25](#),  $\mathbf{p}_q(G) \leq t$  and, because of the above claim,  $G$  contains a  $c$ -treewidth modulator  $Z$  where  $|Z| \leq t - 16s \cdot \sqrt{t} \leq t$ . This, in turn, implies that  $B \cup Z$  is a  $c$ -treewidth modulator of  $G$  that contains  $B$  and has size  $2t$ . Also observe that  $c = 16s^2$ . Therefore, the lemma holds for  $f_{21}(q, h) = \mathcal{O}(s^2)$ .  $\square$

We stress that, as it is done in the proof of [\[21, Lemma 3.6\]](#), it is possible to find a modulator of size at most  $(1 + \varepsilon) \cdot t$ , for every positive real  $\varepsilon > 0$ . Nevertheless, we have provided the proof of [Lemma 34](#) for the particular case  $\varepsilon = 2$ , which is enough for our purposes.

## B.2 Upper bounds on the constants depending on the excluded minors

In this section we provide an estimation on the function  $f_{20}$  in [Theorem 33](#), or equivalently on the constant  $c_{\mathcal{F}}$  in [Theorem 2](#). We first provide some definitions in order to introduce the Unique Linkage Theorem [\[31, 43\]](#).

A *linkage* in a graph  $G$  is a subgraph  $L$  of  $G$  whose connected components are all non-trivial paths. The *paths* of a linkage  $L$  are its connected components and we denote them by  $\mathcal{P}(L)$ . The *size* of  $L$  is the number of its paths and is denoted by  $|L|$ . The *terminals* of a linkage  $L$ , denoted by  $T(L)$ , are the endpoints of the paths in  $\mathcal{P}(L)$ , and the *pattern* of  $L$  is the set

$$\{\{s, t\} \mid \mathcal{P}(L) \text{ contains some } (s, t)\text{-path}\}.$$

Two linkages  $L_1, L_2$  of  $G$  are *equivalent* if they have the same pattern and we denote this fact by  $L_1 \equiv L_2$ . We say that a linkage  $L$  in a graph  $G$  is *unique* if for every linkage  $L'$  that is equivalent to  $L$  it holds that  $V(L') = V(L)$ .

According to the proof of [Proposition 13](#) in [\[25\]](#), the function  $f_6$  emerges from the following result, known as the Unique Linkage Theorem.

**Proposition 35** ([\[31, 43\]](#)). *There exists a function  $f_6 : \mathbb{N}_{\geq 0} \rightarrow \mathbb{N}_{\geq 0}$  such that if  $G$  is a graph and  $L$  is a unique linkage of  $G$ , then  $\text{tw}(G) \leq f_6(|L|)$ .*

It is worth mentioning that [\[31, 43\]](#) do not provide the precise number of exponentiations involved in the function  $f_6$ , and therefore we will express our upper bounds in terms of this function. Namely, in order to provide an upper bound on  $f_{20}(h)$ , we backtrack the functions involved in the intermediate results of this paper as follows:

- According to [Corollary 32](#),  $f_{20}(h) = \mathcal{O}(f_{19}(h, h) \cdot \log(f_{19}(h, h) \cdot h\sqrt{\log h}))$ .
- By [Theorem 31](#),  $f_{19}(h, h) = \mathcal{O}(f_{17}(h, h) \cdot f_{18}(h, h))$ .
- By [Lemma 30](#),  $f_{18}(h, h) = 2^{2^{\mathcal{O}(f_{17}(h, h) \cdot \log f_{17}(h, h))}}$ .

- By Lemma 29,  $f_{17}(h, h) = f_{15}(h, h) \cdot f_{16}(h, h) \cdot 2^{\mathcal{O}(h \log h)}$ .
- By Lemma 34, we can take  $f_{16}(h, h) = 2$ .
- By Lemma 27 (and Lemma 34 as well),  $f_{15}(h, h) = \mathcal{O}((f_5(h, f_{13}(f_3(h), 3, 3h), f_{14}(f_3(h), 3h)))^2)$ .
- By Theorem 5,  $f_3(h) = \mathcal{O}(h^{24}) = h^{\mathcal{O}(1)}$ .
- By Theorem 23,  $f_{14}(f_3(h), 3h) = \mathcal{O}(h^{24}) = h^{\mathcal{O}(1)}$ .
- By Theorem 23,  $f_{13}(f_3(h), 3, 3h) = \mathcal{O}((f_6(h^{\mathcal{O}(1)}))^3)$ .
- By Lemma 10,  $f_5(h, r, \hat{\ell}) = f_2(h) \cdot r^{2^{\mathcal{O}((h^{24} + \hat{\ell}) \cdot \log(h^{24} + \hat{\ell}))}}$ , which is  $r^{2^{2(h + \hat{\ell})\mathcal{O}(1)}}$  by Theorem 5. If we set  $r = f_{13}(f_3(h), 3, 3h) = \mathcal{O}((f_6(h^{\mathcal{O}(1)}))^3)$  and  $\hat{\ell} = f_{14}(f_3(h), 3h) = h^{\mathcal{O}(1)}$ , we have that

$$f_{15}(h, h) = \mathcal{O}((f_5(h, r, \hat{\ell}))^2) = ((f_6(h^{\mathcal{O}(1)}))^3)^{2^{2(h + h^{\mathcal{O}(1)})\mathcal{O}(1)}} = (f_6(h^{\mathcal{O}(1)}))^{2^{2h\mathcal{O}(1)}}.$$

- We set  $\lambda = f_6(h^{\mathcal{O}(1)})$ . Given that  $f_{17}(h, h) = f_{15}(h, h) \cdot f_{16}(h, h) \cdot 2^{\mathcal{O}(h \log h)}$  and that  $f_{16}(h, h) = 2$ , we obtain that  $f_{17}(h, h) = \lambda^{2^{2h\mathcal{O}(1)}}$ . We now have that  $f_{18}(h, h) = 2^{2^{2\lambda 2^{2h\mathcal{O}(1)}}$ , which implies that  $f_{19}(h, h) = 2^{2^{2\lambda 2^{2h\mathcal{O}(1)}}$  and thus  $f_{20}(h) = 2^{2^{2\lambda 2^{2h\mathcal{O}(1)}}$ .

From Theorem 33 and the above discussion, we conclude the following corollary, which gives an explicit upper bound on the contribution of the maximum size of the graphs in  $\mathcal{F}$  in the complexity of our algorithm, depending on the function  $f_6$  given by Proposition 35.

**Corollary 36.** *Let  $\mathcal{F}$  be a collection of graphs each of size at most  $h$ , and let  $G$  be a graph. Then the parameter  $\mathbf{m}_{\mathcal{F}}(G)$  can be computed in time*

$$2^{\left(2^{2^{2\lambda 2^{2h\mathcal{O}(1)}}}\right) \cdot \text{tw}(G) \cdot \log(\text{tw}(G))} \cdot |V(G)|, \text{ where } \lambda = f_6(h^{\mathcal{O}(1)}).$$