

GAMMA-CONVERGENT PROJECTION-FREE FINITE ELEMENT METHODS FOR NEMATIC LIQUID CRYSTALS: THE ERICKSEN MODEL*

RICARDO H. NOCHETTO[†], MICHELE RUGGERI[‡], AND SHUO YANG[§]

Abstract. The Ericksen model for nematic liquid crystals couples a director field with a scalar degree of orientation variable and allows the formation of various defects with finite energy. We propose a simple but novel finite element approximation of the problem that can be implemented easily within standard finite element packages. Our scheme is projection-free and thus circumvents the use of weakly acute meshes, which are quite restrictive in three dimensions but are required by recent algorithms for convergence. We prove stability and Γ -convergence properties of the new method in the presence of defects. We also design an effective nested gradient flow algorithm for computing minimizers that controls the violation of the unit-length constraint of the director. We present several simulations in two and three dimensions that document the performance of the proposed scheme and its ability to capture quite intriguing defects.

Key words. liquid crystals, finite element method, Γ -convergence, gradient flow, defect, energy minimization

AMS subject classifications. 65N30, 35J70, 65Z05

DOI. 10.1137/21M1407495

1. Introduction.

1.1. Liquid crystals with variable degree of orientation. Liquid crystals (LCs) are a mesophase between crystalline solid and isotropic liquid. There are numerous potential applications in engineering and science, in particular in materials science [1, 5, 10]. Nematic LCs are made of rod-like molecules with no positional order that tend to point in a preferred direction. LC materials are thus anisotropic.

We consider the one-constant Ericksen model for nematic LCs with variable degree of orientation [17], which lies between the Oseen–Frank director model and the Landau–de Gennes Q -tensor model [16, 29]. The state of the LC is described in terms of a vector field \mathbf{n} and a scalar function s , which satisfy the constraints $|\mathbf{n}| = 1$ and $-1/(d-1) < s < 1$ for the space dimension $d = 2, 3$. The director \mathbf{n} indicates the

*Received by the editors March 30, 2021; accepted for publication (in revised form) November 12, 2021; published electronically April 19, 2022.

<https://doi.org/10.1137/21M1407495>

Funding: The work of the first and second authors started while they were in residence at the Institute for Computational and Experimental Research in Mathematics (ICERM) during the workshop Numerical Methods and New Perspectives for Extended Liquid Crystalline Systems in 2019, supported by National Science Foundation grant DMS-1439786. The work of the second author was partially supported by the Austrian Science Fund (FWF) through the special research program Taming Complexity in Partial Differential Systems, grant F65, and the Erwin Schrödinger International Institute for Mathematics and Physics (ESI), given during the workshop New Trends in the Variational Modeling and Simulation of Liquid Crystals. The work of the first and third authors was partially supported by National Science Foundation grant DMS-1908267.

[†]Department of Mathematics and Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742 USA (rhn@umd.edu).

[‡]Department of Mathematics and Statistics, University of Strathclyde, Glasgow G1 1XH, United Kingdom, and Institute of Analysis and Scientific Computing, TU Wien, 1040 Vienna, Austria (michele.ruggeri@strath.ac.uk).

[§]Yanqi Lake Beijing Institute of Mathematical Sciences and Applications, 101408 Beijing, China, and Yau Mathematical Sciences Center, Tsinghua University, 100084 Beijing, China (shuoyang@bimsa.cn).




FIG. 1. Schematic illustration of $\mathbf{n}(x)$ and $s(x)$, in microscopic scale near a fixed $x \in \Omega \subset \mathbb{R}^d$. Note that $s = 1$ represents the state of perfect alignment in which all molecules in the local ensemble are parallel to \mathbf{n} . Likewise, $s = -1/(d-1)$ represents the state of perpendicular alignment. The case $s = 0$ corresponds to a defect in the LC material, an isotropic distribution of molecules in the local ensemble that do not lie along any preferred direction.

preferred orientations of the LC molecules, while s represents the degree of alignment that the molecules have with respect to \mathbf{n} , both in the sense of local probabilistic average. A schematic illustration of their meaning is given in Figure 1. The equilibrium state is given by an admissible pair (s, \mathbf{n}) that minimizes the Ericksen energy

$$(1.1) \quad E[s, \mathbf{n}] = \frac{1}{2} \int_{\Omega} (\kappa |\nabla s|^2 + s^2 |\nabla \mathbf{n}|^2) + \int_{\Omega} \psi(s),$$

where $\kappa > 0$ is constant; the constraint on s is enforced by the double well potential ψ . We refer to [4, 21] for early analysis of the Ericksen model.

If s can be approximated by a nonvanishing constant, then the energy (1.1) reduces to the Oseen–Frank energy $E[\mathbf{n}] \propto \int_{\Omega} |\nabla \mathbf{n}|^2$, whose minimizers are harmonic maps and have been extensively studied, e.g., in [27, 14]. However, the simpler Oseen–Frank model has severe limitations in capturing defects: It only admits point defects with finite energy for $d = 3$. In contrast, the Ericksen model (1.1) allows for $\mathbf{n} \notin H^1(\Omega)$ and compensates blow-up of $|\nabla \mathbf{n}|$ by letting s vanish, which is the mechanism for the formation of a variety of line and plane defects; see, e.g., [21] for a proof of the fact that the singular set of a minimizer of (1.1) can have positive Hausdorff dimension. This physical process leads to a *degenerate* Euler–Lagrange equation for \mathbf{n} that poses serious difficulties in formulating mathematically sound algorithms to approximate (1.1) and study their convergence.

1.2. Numerical analysis of the Ericksen model. Several numerical methods for the Oseen–Frank model have been proposed [22, 3, 7, 9]. Finite element methods (FEMs) for the Ericksen model are designed in [6, 23, 24, 30, 15]; see also the recent review [12]. In contrast to [6], a fundamental structure of (1.1) is exploited in [23, 24] to design and analyze FEMs that handle the inherent degeneracy of (1.1) without regularization and enforce the constraint $|\mathbf{n}| = 1$ robustly. Stability and convergence properties via Γ -convergence are proved in [23, 24], pioneering results in this setting. They hinge on a clever discrete energy that mimics the structure of (1.1) discretely but, unfortunately, is cumbersome to implement in standard software packages and requires weakly acute meshes. The latter ensures that the projection of discrete director fields onto the unit sphere is energy decreasing, and thus compatible with the quasi-gradient flow, but is quite restrictive and difficult to implement for $d = 3$ and domains with nontrivial topology.

1.3. Contributions. In this work, we propose a projection-free FEM that avoids dealing with weakly acute meshes. Without the projection step, the unit-length constraint $|\mathbf{n}| = 1$ is no longer satisfied exactly but instead is relaxed at each step of our

iterative solver, a nested gradient flow. The latter guarantees control of the violation of $|\mathbf{n}| = 1$ and asymptotic enforcement of it. We summarize the chief novelties and advantages of our approach as follows.

- *Shape-regular meshes.* Partitions of Ω are assumed to be only shape-regular, which allows for the use of software with general mesh generators such as Netgen [26]. Avoiding weakly acute meshes is important in three dimensions (3D) to deal with interesting but nontrivial geometries as documented in section 5. An earlier work achieving this goal is [30], which presents a mass-lumped FEM with a consistent stabilization term involving $s^2 \nabla \mathbf{n}^\top \mathbf{n}$ for the generalized Ericksen energy.

- *Standard algorithm.* Our novel discretization of (1.1) is straightforward, requires no stabilization, and is easy to implement in standard software packages such as NGSolve [26]. In contrast to [23, 24], our FEM no longer exploits the structure of (1.1), but its analysis does.

- *Γ -convergence.* The analysis of our FEM hinges heavily on the underlying structure of (1.1), which is fully discussed in section 2 and relies on the notion of L^2 -gradient on \mathbf{n} [18, Theorem 6.2]; see Proposition 2.1 below. Such a notion was already used in [11] in the context of the uniaxial Q-tensor LC model. We prove stability and Γ -convergence. Our results are similar to those in [23, 24, 30], but the use of the discrete structure is new.

- *Linear solver.* We propose a nested gradient flow that, despite the nonlinear nature of the problem, is *fully linear* to compute minimizers. The inner loop to advance the director \mathbf{n} for fixed degree of orientation s is allowed to subiterate. This turns out to induce an acceleration mechanism for the computation and motion of defects. For a recent acceleration technique based on a domain decomposition approach, we refer to [15]. Our nested gradient flow iterations fall within the Γ -convergence framework provided a CFL-type condition is imposed on the discretization parameters (see Proposition 4.3). However, well-posedness and stability of the algorithm are guaranteed without any such CFL restriction.

- *Numerical experiments.* We present several simulations in section 5. Some aim to compare the new algorithm with the existing literature in terms of performance and ability to capture defects. Other experiments explore three-dimensional intriguing configurations such as the propeller defect and challenging variations of the Saturn ring defect.

- *Boundary conditions.* Since we do not impose the unit-length constraint $|\mathbf{n}| = 1$, the treatment of boundary data could be simplified and their properties weakened. This potentially affects the regularization procedure for the lim-sup property and the possible presence of defects at the boundary of Ω . We do not explore these issues in this paper but rather in future extension to the Q-tensor model.

1.4. Outline. The remainder of this work is organized as follows. In the next subsection, we collect some general notation used throughout the paper. In section 2, we describe the Ericksen model for LCs with variable degree of orientation and discuss its key structure. In section 3, we introduce our discretization of the model and state our Γ -convergence result. In section 4, we present our iterative scheme for the computation of discrete local minimizers. In section 5, we show numerical experiments illustrating effectiveness and efficiency of our method, as well as its flexibility to deal with complex defects in 3D. We postpone the proofs of most results to section 6.

1.5. General notation. We denote by $\mathbb{N} = \{1, 2, \dots\}$ the set of natural numbers and set $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$. For $d = 2, 3$, we denote the unit sphere in \mathbb{R}^d by $\mathbb{S}^{d-1} = \{x \in \mathbb{R}^d : |x| = 1\}$. We denote by $B_r(x)$ the ball of radius $r > 0$ centered at

$x \in \mathbb{R}^d$. For (spaces of) vector- or matrix-valued functions, we use bold letters, e.g., for a generic domain $\Omega \subset \mathbb{R}^d$, we denote both $L^2(\Omega; \mathbb{R}^d)$ and $L^2(\Omega; \mathbb{R}^{d \times d})$ by $\mathbf{L}^2(\Omega)$. We denote by $\langle \cdot, \cdot \rangle$ both the scalar product of $\mathbf{L}^2(\Omega)$ and the duality pairing between $\mathbf{H}^1(\Omega)$ and its dual, with the ambiguity being resolved by the arguments. We use the notation \lesssim to denote *smaller than or equal to up to a multiplicative constant*, i.e., we write $A \lesssim B$ if there exists a constant $c > 0$, which is clear from the context and always independent of the discretization parameters, such that $A \leq cB$.

2. Problem formulation. Let $\Omega \subset \mathbb{R}^d$ ($d = 2, 3$) be a bounded Lipschitz domain. In the Ericksen model, the state of the LC is described in terms of a unit-length vector field $\mathbf{n} : \Omega \rightarrow \mathbb{S}^{d-1}$ and a scalar function $s : \Omega \rightarrow (-1/(d-1), 1)$. Equilibrium configurations are minimizers of the energy $E[s, \mathbf{n}] = E_1[s, \mathbf{n}] + E_2[s]$ in (1.1), where

$$(2.1) \quad E_1[s, \mathbf{n}] := \frac{1}{2} \int_{\Omega} (\kappa |\nabla s|^2 + s^2 |\nabla \mathbf{n}|^2), \quad E_2[s] := \int_{\Omega} \psi(s).$$

The double well potential $\psi : (-1/(d-1), 1) \rightarrow \mathbb{R}_{\geq 0}$ satisfies the following properties [17]:

- $\psi \in C^2(-1/(d-1), 1)$,
- $\lim_{s \rightarrow 1^-} \psi(s) = +\infty = \lim_{s \rightarrow -1/(d-1)^+} \psi(s)$,
- $\psi(0) > \psi(s^*) = \min_{s \in (-1/(d-1), 1)} \psi(s) = 0$ for some $s^* \in (0, 1)$,
- $\psi'(0) = 0$.

In (2.1), $E_1[s, \mathbf{n}]$ is the one-constant approximation of the elastic energy proposed in [17], while $E_2[s]$ is a potential energy which confines the variable s within the physically admissible interval $(-1/(d-1), 1)$. The presence of the weight s^2 in the second term of $E_1[s, \mathbf{n}]$ allows for blow-up of $\nabla \mathbf{n}$, namely $\mathbf{n} \notin \mathbf{H}^1(\Omega)$, in the *singular set* Σ , where defects may occur:

$$(2.2) \quad \Sigma := \{x \in \Omega : s(x) = 0\}.$$

To complete the setting, we define the set of admissible functions where we seek minimizers of (2.1). Note that, allowing for a director $\mathbf{n} \notin \mathbf{H}^1(\Omega)$, one encounters at least two difficulties: On the one hand, it is not clear how to interpret the gradient of \mathbf{n} appearing in $E_1[s, \mathbf{n}]$. On the other hand, the trace of \mathbf{n} on the boundary of Ω is not well defined, so that one cannot impose Dirichlet conditions on \mathbf{n} in the standard way. To cope with these problems, following [4, 21], we introduce the auxiliary variable $\mathbf{u} = s\mathbf{n}$. Then, the product rule formally yields that

$$(2.3) \quad \nabla \mathbf{u} = \mathbf{n} \otimes \nabla s + s \nabla \mathbf{n}.$$

Since $|\mathbf{n}| = 1$, the identities $\nabla \mathbf{n}^\top \mathbf{n} = \mathbf{0}$ and $|\mathbf{n} \otimes \nabla s| = |\nabla s|$ are valid. It follows that the above decomposition of $\nabla \mathbf{u}$ is orthogonal, i.e.,

$$(2.4) \quad |\nabla \mathbf{u}|^2 = |\mathbf{n} \otimes \nabla s|^2 + s^2 |\nabla \mathbf{n}|^2 = |\nabla s|^2 + s^2 |\nabla \mathbf{n}|^2.$$

In particular, $E_1[s, \mathbf{n}]$ can be rewritten in terms of s and $\mathbf{u} = s\mathbf{n}$ as

$$(2.5) \quad E_1[s, \mathbf{n}] = \tilde{E}_1[s, \mathbf{u}] = \frac{1}{2} \int_{\Omega} ((\kappa - 1) |\nabla s|^2 + |\nabla \mathbf{u}|^2).$$

In the latter, the degree of orientation and the auxiliary field are decoupled. In particular, this reveals that, for (s, \mathbf{n}) such that $E_1[s, \mathbf{n}] < \infty$, $\mathbf{u} = s\mathbf{n} \in \mathbf{H}^1(\Omega)$ even though $\mathbf{n} \notin \mathbf{H}^1(\Omega)$.

We say that a triple $(s, \mathbf{n}, \mathbf{u})$ satisfies the *structural condition* if

$$(2.6) \quad -\frac{1}{d-1} < s < 1, \quad |\mathbf{n}| = 1, \quad \text{and} \quad \mathbf{u} = s\mathbf{n} \quad \text{a.e. in } \Omega.$$

In view of the above discussion, we are led to consider the following admissible class:

$$(2.7) \quad \mathcal{A} := \{(s, \mathbf{n}, \mathbf{u}) \in H^1(\Omega) \times \mathbf{L}^\infty(\Omega) \times \mathbf{H}^1(\Omega) : (s, \mathbf{n}, \mathbf{u}) \text{ satisfies (2.6)}\}.$$

For triples $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}$, it is possible to characterize the gradient of \mathbf{n} occurring in $E_1[s, \mathbf{n}]$ using a weaker notion of differentiability. To this end, we recall the following definition [18, Theorem 6.2]: We say that \mathbf{n} is L^2 -differentiable at $x \in \Omega$, and we denote its L^2 -gradient at x by $\nabla \mathbf{n}(x)$, if

$$\oint_{B_r(x)} |\mathbf{n}(y) - \mathbf{n}(x) - \nabla \mathbf{n}(x)(y - x)|^2 dy = o(r^2) \quad \text{as } r \rightarrow 0.$$

It is well known that the notion of L^2 -differentiability is weaker than the existence of an L^2 -integrable weak gradient, in the sense that every H^1 -function is L^2 -differentiable almost everywhere and its L^2 -gradient coincides with the weak gradient; see, e.g., [18, Theorem 6.2].

In the following proposition, we establish that if $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}$, then \mathbf{n} is L^2 -differentiable and the decomposition (2.4) holds almost everywhere outside of the singular set Σ in (2.2). Its proof will be presented in subsection 6.1.

PROPOSITION 2.1 (orthogonal decomposition). *Let $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}$. Then, \mathbf{n} is L^2 -differentiable a.e. in $\Omega \setminus \Sigma$. In particular, its L^2 -gradient is given by*

$$(2.8) \quad \nabla \mathbf{n} = s^{-1}(\nabla \mathbf{u} - \mathbf{n} \otimes \nabla s) \quad \text{a.e. in } \Omega \setminus \Sigma.$$

Moreover, the following identity holds:

$$(2.9) \quad |\nabla \mathbf{u}|^2 = |\nabla s|^2 + s^2 |\nabla \mathbf{n}|^2 \quad \text{a.e. in } \Omega \setminus \Sigma.$$

This allows us to give a precise meaning to $E_1[s, \mathbf{n}]$ in (2.1). Depending on the context, we interpret $\nabla \mathbf{n}$ in the sense of L^2 -gradient in $\Omega \setminus \Sigma$ and $\int_{\Sigma} s^2 |\nabla \mathbf{n}|^2 = 0$, or we alternatively replace Ω by $\Omega \setminus \Sigma$ as domain of integration or even use the representation $\tilde{E}_1[s, \mathbf{u}]$ of (2.5).

Turning to boundary conditions, let $\Gamma_D \subseteq \partial\Omega$ be a relatively open subset of the boundary such that $|\Gamma_D| > 0$, where we aim to impose Dirichlet boundary conditions. These, in the context of LCs, are usually referred to as *strong anchoring* conditions. To this end, given a triple $(g, \mathbf{q}, \mathbf{r}) \in W^{1,\infty}(\mathbb{R}^d) \times \mathbf{L}^\infty(\mathbb{R}^d) \times \mathbf{W}^{1,\infty}(\mathbb{R}^d)$ satisfying the structural condition (2.6), we consider the following restricted admissible class that incorporates boundary conditions:

$$(2.10) \quad \mathcal{A}(g, \mathbf{r}) := \{(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A} : s|_{\Gamma_D} = g|_{\Gamma_D} \text{ and } \mathbf{u}|_{\Gamma_D} = \mathbf{r}|_{\Gamma_D}\}.$$

Overall, we are interested in the following constrained minimization problem: Find $(s^*, \mathbf{n}^*, \mathbf{u}^*) \in \mathcal{A}(g, \mathbf{r})$ such that

$$(2.11) \quad (s^*, \mathbf{n}^*, \mathbf{u}^*) = \arg \min_{(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}(g, \mathbf{r})} E[s, \mathbf{n}].$$

To conclude this section, let $\delta_0 > 0$ be sufficiently small. Some of our results below will require the following technical assumptions on the Dirichlet data, namely

$$(2.12) \quad -\frac{1}{d-1} + \delta_0 \leq g(x) \leq 1 - \delta_0 \quad \text{for all } x \in \mathbb{R}^d,$$

$$(2.13) \quad g \geq \delta_0 \quad \text{on } \Gamma_D,$$

and on the double well potential, namely

$$(2.14) \quad \begin{aligned} \psi(s) &\geq \psi(1 - \delta_0) && \text{for all } s \geq 1 - \delta_0, \\ \psi(s) &\geq \psi\left(-\frac{1}{d-1} + \delta_0\right) && \text{for all } s \leq -\frac{1}{d-1} + \delta_0, \end{aligned}$$

and ψ is monotone in $(-1/(d-1), -1/(d-1) + \delta_0)$ and in $(1 - \delta_0, 1)$. Note that (2.13) implies that $\mathbf{q} = g^{-1}\mathbf{r}$ is $\mathbf{W}^{1,\infty}$ on Γ_D so that imposing the Dirichlet condition $\mathbf{n}|_{\Gamma_D} = \mathbf{q}|_{\Gamma_D}$ on the physical variable \mathbf{n} is equivalent to imposing $\mathbf{u}|_{\Gamma_D} = \mathbf{r}|_{\Gamma_D}$ on the auxiliary variable \mathbf{u} . Finally, the property (2.14) is consistent with the fact that $\psi(s) \rightarrow +\infty$ as $s \rightarrow -1/(d-1)$ and $s \rightarrow 1$.

3. Γ -convergent finite element discretization. We assume Ω to be a polytopal domain and consider a shape-regular family $\{\mathcal{T}_h\}$ of simplicial meshes of Ω (in the sense of, e.g., [8, Definitions 3.3–3.4]) parametrized by the mesh size $h = \max_{K \in \mathcal{T}_h} h_K$, where $h_K = \text{diam}(K)$. We stress that we do not require any mesh to be weakly acute (we refer to [8, Remark 3.12] for a discussion of this assumption for $d = 2, 3$). We denote by \mathcal{N}_h the set of vertices of \mathcal{T}_h . For any $K \in \mathcal{T}_h$, we denote by $\mathcal{P}^1(K)$ the space of first-order polynomials on K . We consider the space of \mathcal{T}_h -piecewise affine and globally continuous functions

$$V_h := \{v_h \in C^0(\bar{\Omega}) : v_h|_K \in \mathcal{P}^1(K) \text{ for all } K \in \mathcal{T}_h\}.$$

Let $\mathbf{V}_h := (V_h)^d$ be the corresponding space of vector-valued polynomials. We denote by I_h both the nodal interpolant $I_h : C^0(\bar{\Omega}) \rightarrow V_h$ and its vector-valued counterpart $I_h : \mathbf{C}^0(\bar{\Omega}) \rightarrow \mathbf{V}_h$.

For $s_h \in V_h$ and $\mathbf{n}_h \in \mathbf{V}_h$, let the discrete energy be $E^h[s_h, \mathbf{n}_h] = E_1^h[s_h, \mathbf{n}_h] + E_2^h[s_h]$ with

$$(3.1) \quad E_1^h[s_h, \mathbf{n}_h] := \frac{1}{2} \int_{\Omega} (\kappa |\mathbf{n}_h \otimes \nabla s_h|^2 + s_h^2 |\nabla \mathbf{n}_h|^2), \quad E_2^h[s_h] := \int_{\Omega} \psi(s_h).$$

Note that E^h is consistent, in the sense that $E^h[s, \mathbf{n}] = E[s, \mathbf{n}]$ if $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}(g, \mathbf{r})$.

We say that a triple $(s_h, \mathbf{n}_h, \mathbf{u}_h) \in V_h \times \mathbf{V}_h \times \mathbf{V}_h$ satisfies the *discrete structural condition* if

$$(3.2) \quad -\frac{1}{d-1} < s_h(z) < 1, \quad |\mathbf{n}_h(z)| \geq 1, \quad \text{and} \quad \mathbf{u}_h(z) = s_h(z) \mathbf{n}_h(z) \quad \text{for all } z \in \mathcal{N}_h.$$

In (3.2), the requirements prescribed by the continuous structural condition (2.6) are imposed only at the vertices of the mesh, which is practical. Moreover, the unit-length constraint for the director is relaxed, since \mathbf{n}_h may attain also values outside of the unit sphere.

Let $\varepsilon > 0$, $g_h = I_h[g]$, and $\mathbf{r}_h = I_h[\mathbf{r}]$. We consider the following discrete minimization problem: Find $(s_h^*, \mathbf{n}_h^*, \mathbf{u}_h^*) \in \mathcal{A}_{h,\varepsilon}(g_h, \mathbf{r}_h)$ such that

$$(3.3) \quad (s_h^*, \mathbf{n}_h^*, \mathbf{u}_h^*) = \arg \min_{(s_h, \mathbf{n}_h, \mathbf{u}_h) \in \mathcal{A}_{h,\varepsilon}(g_h, \mathbf{r}_h)} E_h[s_h, \mathbf{n}_h],$$

where the discrete restricted admissible class is defined as

$$(3.4) \quad \mathcal{A}_{h,\varepsilon}(g_h, \mathbf{r}_h) := \{(s_h, \mathbf{n}_h, \mathbf{u}_h) \in V_h \times \mathbf{V}_h \times \mathbf{V}_h : \\ (s_h, \mathbf{n}_h, \mathbf{u}_h) \text{ satisfies (3.2), } \|I_h[|\mathbf{n}_h|^2] - 1\|_{L^1(\Omega)} \leq \varepsilon, \\ s_h(z) = g_h(z), \text{ and } u_h(z) = r_h(z) \text{ for all } z \in \mathcal{N}_h \cap \Gamma_D\}.$$

In the following theorem, we show that the discrete energy (3.1) converges toward the continuous one (2.1) in the sense of Γ -convergence.

THEOREM 3.1 (Γ -convergence). *Suppose that $\varepsilon \rightarrow 0$ as $h \rightarrow 0$. Then, the following two properties are satisfied:*

(i) *Lim-sup inequality (consistency): Let $\Gamma_D = \partial\Omega$. Let the assumptions (2.12)–(2.14) hold. If $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}(g, \mathbf{r})$, then there exists a sequence $\{(s_h, \mathbf{n}_h, \mathbf{u}_h)\} \subset \mathcal{A}_{h,\varepsilon}(g_h, \mathbf{r}_h)$ such that $\|\mathbf{n}_h\|_{L^\infty(\Omega)} = 1$, $s_h \rightarrow s$ in $H^1(\Omega)$, $\mathbf{n}_h \rightarrow \mathbf{n}$ in $L^2(\Omega \setminus \Sigma)$, $\mathbf{u}_h \rightarrow \mathbf{u}$ in $H^1(\Omega)$, as $h \rightarrow 0$, and*

$$(3.5) \quad E[s, \mathbf{n}] \geq \limsup_{h \rightarrow 0} E^h[s_h, \mathbf{n}_h].$$

(ii) *Lim-inf inequality (stability): Let $\{(s_h, \mathbf{n}_h, \mathbf{u}_h)\} \subset \mathcal{A}_{h,\varepsilon}(g_h, \mathbf{r}_h)$ be a sequence such that $E^h[s_h, \mathbf{n}_h] \leq C$ and $\|\mathbf{n}_h\|_{L^\infty(\Omega)} \leq C$, where $C \geq 1$ is a constant independent of h . Then, there exist $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}(g, \mathbf{r})$ and a subsequence of $\{(s_h, \mathbf{n}_h, \mathbf{u}_h)\}$ (not relabeled) such that $s_h \rightharpoonup s$ in $H^1(\Omega)$, $\mathbf{n}_h \rightarrow \mathbf{n}$ in $L^2(\Omega \setminus \Sigma)$, $\mathbf{u}_h \rightharpoonup \mathbf{u}$ in $H^1(\Omega)$ as $h \rightarrow 0$, and*

$$(3.6) \quad E[s, \mathbf{n}] \leq \liminf_{h \rightarrow 0} E^h[s_h, \mathbf{n}_h].$$

The proof of Theorem 3.1 is deferred to subsections 6.2 and 6.3. The assumption $\Gamma_D = \partial\Omega$ in part (i) is needed to apply a regularization result from [23] (see Lemma 6.2 below). The properties established in Theorem 3.1 are slight variations of the properties required by the standard definition of Γ -convergence; see, e.g., [13, Definition 1.5]. However, they still allow us to prove the convergence of discrete global minimizers.

COROLLARY 3.2 (convergence of discrete global minimizers). *Let $\Gamma_D = \partial\Omega$ and suppose that the assumptions (2.12)–(2.14) hold. Let $\{(s_h, \mathbf{n}_h, \mathbf{u}_h)\} \subset \mathcal{A}_{h,\varepsilon}(g_h, \mathbf{r}_h)$ be a sequence of global minimizers of the discrete energy (3.1) such that $\|\mathbf{n}_h\|_{L^\infty(\Omega)} \leq C$, where $C \geq 1$ is a constant independent of h . Then, every cluster point $(s, \mathbf{n}, \mathbf{u})$ belongs to $\mathcal{A}(g, \mathbf{r})$ and is a global minimizer of the continuous energy (2.1).*

4. Computation of discrete local minimizers. In this section, we propose an effective algorithm to compute discrete local minimizers of (3.1). The method is based on a discretization of the (nonphysical) energy-decreasing dynamics driven by the system of gradient flows

$$\begin{aligned} \partial_t \mathbf{n} + \delta_{\mathbf{n}} E^h[s, \mathbf{n}] &= 0, \\ \partial_t s + \delta_s E^h[s, \mathbf{n}] &= 0, \end{aligned}$$

where $\delta_{\mathbf{n}} E^h[s, \mathbf{n}]$ and $\delta_s E^h[s, \mathbf{n}]$ denote the Gâteaux derivatives of the energy with respect to the order parameters, i.e.,

$$\begin{aligned} \langle \delta_{\mathbf{n}} E^h[s, \mathbf{n}], \phi \rangle &= \langle \delta_{\mathbf{n}} E_1^h[s, \mathbf{n}], \phi \rangle = \kappa \langle \mathbf{n} \otimes \nabla s, \phi \otimes \nabla s \rangle + \langle s \nabla \mathbf{n}, s \nabla \phi \rangle, \\ \langle \delta_s E^h[s, \mathbf{n}], w \rangle &= \langle \delta_s E_1^h[s, \mathbf{n}], w \rangle + \langle \delta_s E_2^h[s, \mathbf{n}], w \rangle \\ &= \kappa \langle \mathbf{n} \otimes \nabla s, \mathbf{n} \otimes \nabla w \rangle + \langle s \nabla \mathbf{n}, w \nabla \mathbf{n} \rangle + \langle \psi'(s), w \rangle. \end{aligned}$$

Let us introduce the ingredients of the scheme. First, let

$$V_{h,D} := \{v_h \in V_h : v_h(z) = 0 \text{ for all } z \in \mathcal{N}_h \cap \Gamma_D\} \quad \text{and} \quad \mathbf{V}_{h,D} := (V_{h,D})^d$$

be the spaces of discrete functions satisfying homogeneous Dirichlet conditions on Γ_D . Given $\mathbf{n}_h \in \mathbf{V}_h$, we consider the subspace of $\mathbf{V}_{h,D}$ consisting of all discrete functions with nodal values orthogonal to those of \mathbf{n}_h at all vertices:

$$\mathcal{K}_h[\mathbf{n}_h] := \{\phi_h \in \mathbf{V}_{h,D} : \mathbf{n}_h(z) \cdot \phi_h(z) = 0 \text{ for all } z \in \mathcal{N}_h\}.$$

The space $\mathcal{K}_h[\mathbf{n}_h]$ can be interpreted as a discretization of the space of tangential variations $\mathcal{K}[\mathbf{n}] := \{\phi \in \mathbf{H}^1(\Omega) : \mathbf{n} \cdot \phi = 0 \text{ a.e. in } \Omega\}$, which naturally occurs in the variational formulation of problems with a unit-length constraint; see, e.g., [8, Lemma 7.1] for the harmonic map equation.

For the treatment of the double well potential, we follow a convex splitting approach (see, e.g., [31]): We assume the splitting $\psi = \psi_c - \psi_e$, where ψ_c and ψ_e are both convex and ψ_c is quadratic.

The *time* discretization of the gradient flow for the director and the degree of orientation are based on the constant time-step sizes $\tau_n > 0$ and $\tau_s > 0$, respectively. Moreover, we consider the difference quotient $d_t s_h^{i+1} := (s_h^{i+1} - s_h^i)/\tau_s$.

In the following algorithm, we state the proposed numerical scheme for the computation of discrete local minimizers of (3.1). We assume that (2.13) is satisfied so that imposing Dirichlet boundary conditions directly for the director is allowed. Let $\text{tol} > 0$ denote a tolerance.

Algorithm 4.1 Alternating direction discrete gradient flow

Input: $s_h^0 \in V_h$, $\mathbf{n}_h^0 \in \mathbf{V}_h$ such that $|\mathbf{n}_h^0(z)| = 1$ for all $z \in \mathcal{N}_h$, $\mathbf{n}_h^0(z) = \mathbf{r}_h(z)/g_h(z)$ and $s_h^0(z) = g_h(z)$ for all $z \in \mathcal{N}_h \cap \Gamma_D$.

Outer loop: For all $i \in \mathbb{N}_0$, iterate (i)–(ii):

(i) Inner loop: Given (\mathbf{n}_h^i, s_h^i) , let $\mathbf{n}_h^{i,0} = \mathbf{n}_h^i$. For all $\ell \in \mathbb{N}_0$, iterate (i-a)–(i-b):

(i-a) Compute $\mathbf{t}_h^{i,\ell} \in \mathcal{K}_h[\mathbf{n}_h^{i,\ell}]$ such that

$$(4.1) \quad \begin{aligned} \langle \mathbf{t}_h^{i,\ell}, \phi_h \rangle_* + \tau_n \kappa \langle \mathbf{t}_h^{i,\ell} \otimes \nabla s_h^i, \phi_h \otimes \nabla s_h^i \rangle + \tau_n \langle s_h^i \nabla \mathbf{t}_h^{i,\ell}, s_h^i \nabla \phi_h \rangle \\ = -\kappa \langle \mathbf{n}_h^{i,\ell} \otimes \nabla s_h^i, \phi_h \otimes \nabla s_h^i \rangle - \langle s_h^i \nabla \mathbf{n}_h^{i,\ell}, s_h^i \nabla \phi_h \rangle \end{aligned}$$

for all $\phi_h \in \mathcal{K}_h[\mathbf{n}_h^{i,\ell}]$;

(i-b) Update $\mathbf{n}_h^{i,\ell+1} := \mathbf{n}_h^{i,\ell} + \tau_n \mathbf{t}_h^{i,\ell}$;

until

$$(4.2) \quad |E_1^h[s_h^i, \mathbf{n}_h^{i,\ell+1}] - E_1^h[s_h^i, \mathbf{n}_h^{i,\ell}]| < \text{tol}.$$

If $\ell_i \in \mathbb{N}_0$ denotes the smallest integer for which the stopping criterion (4.2) is satisfied, define $\mathbf{n}_h^{i+1} := \mathbf{n}_h^{i,\ell_i+1}$.

(ii) Compute $s_h^{i+1} \in V_h$ such that $s_h^{i+1}(z) = g_h(z)$ for all $z \in \mathcal{N}_h \cap \Gamma_D$ and

$$(4.3) \quad \begin{aligned} \langle d_t s_h^{i+1}, w_h \rangle + \kappa \langle \mathbf{n}_h^{i+1} \otimes \nabla s_h^{i+1}, \mathbf{n}_h^{i+1} \otimes \nabla w_h \rangle \\ + \langle s_h^{i+1} \nabla \mathbf{n}_h^{i+1}, w_h \nabla \mathbf{n}_h^{i+1} \rangle + \langle \psi'_c(s_h^{i+1}), w_h \rangle = \langle \psi'_e(s_h^i), w_h \rangle \end{aligned}$$

for all $w_h \in V_{h,D}$.

Output: Sequence of approximations $\{(s_h^i, \mathbf{n}_h^i)\}_{i \in \mathbb{N}_0}$.

In Algorithm 4.1, $\langle \cdot, \cdot \rangle_*$ denotes the scalar product of the metric used in the discrete gradient flow (4.1) for the director. In this work, we consider the following two choices for $\langle \cdot, \cdot \rangle_*$, dictated by numerical convenience:

$$(4.4) \quad \langle \phi, \psi \rangle_* = \langle \phi, \psi \rangle \quad (L^2\text{-metric}),$$

$$(4.5) \quad \langle \phi, \psi \rangle_* = \langle h^\alpha \nabla \phi, \nabla \psi \rangle \quad \text{with } 0 < \alpha \leq 2 \quad (\text{weighted } H^1\text{-metric}).$$

In (4.5), the H^1 -metric is weakened by a positive power of the mesh size h . Note that the choice $\alpha = 0$ corresponds to a full H^1 -gradient flow, which is not appropriate since the director does not belong to $\mathbf{H}^1(\Omega)$ in general (e.g., in the presence of defects). On the other hand, if $\alpha = 2$, a standard scaling argument shows that the resulting metric is equivalent to the L^2 -metric in (4.4).

In the convex splitting of ψ , we adopt a semi-implicit approach: The convex and quadratic part ψ_c is treated implicitly, while the concave one ψ_e is treated explicitly. This leads to a linear and positive definite contribution to the left-hand side of (4.3). Moreover, the resulting discretization is unconditionally stable (see, e.g., [23, Lemma 4.1]). Altogether, both (4.1) and (4.3) are linear symmetric positive definite systems in the unknowns $\mathbf{t}_h^{i,\ell}$ and s_h^{i+1} . The orthogonality constraint in (4.1) can be imposed at the linear algebraic level by introducing a Lagrange multiplier associated with it (see, e.g., the discussion in [8, section 7.2.5]) or via a null-space method as done, e.g., in [25, 23, 20]. In this work, we implement it using a Lagrange multiplier.

Although in most of our numerical experiments we will set $\tau_{\mathbf{n}} = \tau_s$, we observed that in some situations the flexibility of choosing different time-step sizes in (4.1) and (4.3) is decisive in order to move defects in numerical simulations (see, e.g., the experiment in subsection 5.3 below).

In the following proposition, we prove well-posedness and an energy-decreasing property of Algorithm 4.1.

PROPOSITION 4.1 (properties of Algorithm 4.1). *Algorithm 4.1 is well-posed and energy decreasing. Specifically, for all $i \in \mathbb{N}_0$, the following assertions hold:*

- (i) *For all $\ell \in \mathbb{N}_0$, (4.1) admits a unique solution $\mathbf{t}_h^{i,\ell} \in \mathcal{K}_h[\mathbf{n}_h^{i,\ell}]$.*
- (ii) *The inner loop terminates in a finite number of iterations, i.e., there exists $\ell \in \mathbb{N}_0$ such that the stopping criterion (4.2) is met.*
- (iii) *(4.3) admits a unique solution $s_h^{i+1} \in V_h$ such that $s_h^{i+1}(z) = g_h(z)$ for all $z \in \mathcal{N}_h \cap \Gamma_D$.*
- (iv) *There holds*

$$(4.6) \quad \begin{aligned} E^h[s_h^{i+1}, \mathbf{n}_h^{i+1}] - E^h[s_h^i, \mathbf{n}_h^i] &\leq - \left(\tau_s \|d_t s_h^{i+1}\|_{L^2(\Omega)}^2 + \tau_{\mathbf{n}} \sum_{\ell=0}^{\ell_i} \|\mathbf{t}_h^{i,\ell}\|_*^2 \right) \\ &\quad - \left(\tau_s^2 E_1^h[d_t s_h^{i+1}, \mathbf{n}_h^{i+1}] + \tau_{\mathbf{n}}^2 \sum_{\ell=0}^{\ell_i} E_1^h[s_h^i, \mathbf{t}_h^{i,\ell}] \right). \end{aligned}$$

In particular, there holds $E^h[s_h^{i+1}, \mathbf{n}_h^{i+1}] \leq E^h[s_h^i, \mathbf{n}_h^i]$ and equality holds if and only if $(s_h^{i+1}, \mathbf{n}_h^{i+1}) = (s_h^i, \mathbf{n}_h^i)$ (equilibrium state).

Remark 4.2 (energy decrease). The right-hand side of (4.6) characterizes the energy decrease guaranteed by each step of Algorithm 4.1 and comprises two contributions: The term

$$- \left(\tau_s \|d_t s_h^{i+1}\|_{L^2(\Omega)}^2 + \tau_n \sum_{\ell=0}^{\ell_i} \|\mathbf{t}_h^{i,\ell}\|_*^2 \right)$$

is the energy decrease due to the gradient-flow nature of Algorithm 4.1. The term

$$- \left(\tau_s^2 E_1^h[d_t s_h^{i+1}, \mathbf{n}_h^{i+1}] + \tau_n^2 \sum_{\ell=0}^{\ell_i} E_1^h[s_h^i, \mathbf{t}_h^{i,\ell}] \right)$$

is the numerical dissipation due to the backward Euler methods used for the time discretization.

In practical implementations of Algorithm 4.1, the outer loop is terminated when

$$(4.7) \quad |E^h[s_h^{i+1}, \mathbf{n}_h^{i+1}] - E^h[s_h^i, \mathbf{n}_h^i]| < \text{tol}.$$

Since the algorithm fulfills a monotone energy-decreasing property, the stopping criterion is met in a finite number of iterations.

The approximations \mathbf{n}_h^{i+1} of the director generated by Algorithm 4.1 do not satisfy the unit-length constraint at the vertices of the mesh, as in [23, 24]. However, the following proposition, proved in subsection 6.4, shows that violation of this constraint can be controlled by the time-step size τ_n , independently of the number of iterations. Moreover, the uniform boundedness in $\mathbf{L}^\infty(\Omega)$ of the sequence required by the Γ -convergence result (cf. Theorem 3.1(ii)) can be guaranteed if the discretization parameters satisfy a suitable CFL-type condition. However, we stress that such a condition is not necessary for the well-posedness and the stability of the algorithm.

PROPOSITION 4.3 (properties of discrete director field). *Let $j \geq 1$. The following holds.*

(i) *Suppose that the norm induced by the metric $\langle \cdot, \cdot \rangle_*$ used in (4.1) is an upper bound for the L^2 -norm, i.e., there exists $C_* > 0$ such that*

$$(4.8) \quad \|\phi_h\|_{L^2(\Omega)} \leq C_* \|\phi_h\|_* \quad \text{for all } \phi_h \in \mathbf{V}_{h,D}.$$

Then, the approximations generated by Algorithm 4.1 satisfy

$$(4.9) \quad \|I_h[|\mathbf{n}_h^j|^2 - 1]\|_{L^1(\Omega)} \leq C_1 \tau_n E^h[s_h^0, \mathbf{n}_h^0],$$

where $C_1 > 0$ depends only on C_ and the shape-regularity of $\{\mathcal{T}_h\}$.*

(ii) *Suppose τ_n fulfills the following CFL-type condition:*

$$(4.10) \quad \begin{aligned} \tau_n h_{\min}^{-d} &\leq C^* && \text{if } \langle \cdot, \cdot \rangle_* \text{ is chosen as (4.4),} \\ \tau_n h_{\min}^{2-d-\alpha} |\log h_{\min}|^2 &\leq C^* && \text{if } \langle \cdot, \cdot \rangle_* \text{ is chosen as (4.5),} \end{aligned}$$

where $h_{\min} := \min_{K \in \mathcal{T}_h} h_K$ and $C^ > 0$ is arbitrary. Then, the approximations generated by Algorithm 4.1 satisfy*

$$(4.11) \quad \|\mathbf{n}_h^j\|_{L^\infty(\Omega)} \leq 1 + C_2 E^h[s_h^0, \mathbf{n}_h^0],$$

where $C_2 > 0$ is proportional to $C^ > 0$ in (4.10) with proportionality constant depending on the shape-regularity of $\{\mathcal{T}_h\}$.*

To conclude this section, we discuss the design of Algorithm 4.1 with special emphasis on its nested structure and distinct roles of τ_n and τ_s . Obviously, τ_n controls

the violation of the unit-length constraint according to (4.9), but the roles of subiterations in (4.1) and τ_s in (4.3) are more subtle and deserve further elaboration. The presence of defects is associated with values $s_h^i(x_j)$ close to zero at nodes x_j , which in turn act as weights in (4.1) for the tangential updates $\mathbf{t}_h^{i,\ell}$ of the director field $\mathbf{n}_h^{i,\ell}$. The fast decrease to zero of $s_h^i(x_j)$, relative to the growth of $\nabla \mathbf{n}_h^i$ in its vicinity, impedes further changes of $\mathbf{n}_h^i(x_j)$ because they are not energetically favorable: The defect is thus pinned at the same location x_j for many iterations. Experiments with Algorithm 4.1 reveal defect pinning if $\tau_n = \tau_s$ and one step of (4.1) per step of (4.3) is utilized. The subiterations within the inner loop (4.1) allow $\mathbf{n}_h^{i,\ell}$ to adjust to the current value of s_h^i . This mimics an approximate optimization step but with unit length and max norm control dictated by Proposition 4.3. In contrast, full optimization has been proposed in [23, 24, 30] instead of (4.1), followed by nodal projection onto the unit sphere, whereas one step of a weighted gradient flow (4.1) has been advocated in [11] for the Q -tensor model. On the other hand, since τ_s penalizes changes of s_h^i , smaller values of τ_s relative to τ_n delay changes of s_h^i in favor of changes of \mathbf{n}_h^i . This does not fix the stiff character of (4.1), studied in [15], but does remove defect pinning. Several numerical experiments in section 5 document this finding.

5. Numerical experiments. In this section, we present a series of numerical experiments that explore the accuracy of Algorithm 4.1 and its ability to approximate rather complex defects of nematic LCs in 2D and 3D. In both cases, these results complement the theory of sections 3 and 4 and extend it.

We have implemented Algorithm 4.1 within the high performance multiphysics finite element software Netgen/NGSolve [26]. To solve the constrained variational problem (4.1), we adopt a saddle point approach. The ensuing linear systems are solved using the built-in conjugate gradient solver of Netgen/NGSolve, while the visualization relies on ParaView [2].

All pictures below obey the following rules. The vector field depicts the director \mathbf{n} , whereas the color scale refers to the degree of orientation s . Blue regions indicate areas with values of s close to zero, which signify the occurrence of defects, while the red ones indicate regions with largest values of s ($s \approx 0.75$ in our simulations), where the director encodes the local orientation of the LC molecules. We generate unstructured, generally non-weakly acute, meshes within Netgen with desirable mesh size h_0 but the effective maximum size h of tetrahedra in 3D may only satisfy $h \approx h_0$. For the sake of reproducibility, we will specify h_0 when dealing with unstructured three-dimensional meshes.

We stress that, unlike FEMs proposed in previous works [24, 23], the energy-decreasing property of Algorithm 4.1 (cf. Proposition 4.1) does not rely on meshes being weakly acute. Except for simple three-dimensional geometries, such meshes are hard, to impossible, to construct. This is the case of the cylinder domain in subsection 5.3 and the Saturn ring configurations in subsection 5.5, for which mesh flexibility is of fundamental importance to capture topologically complicated defects.

Throughout this section, we consider the double well potential $\psi(s) = c_{\text{dw}}(\psi_c(s) - \psi_e(s))$ with

$$(5.1) \quad \psi_c(s) := 63s^2, \quad \psi_e(s) := -16s^4 + \frac{64}{3}s^3 + 57s^2 - 0.5625,$$

where $c_{\text{dw}} \geq 0$. Note that, for $c_{\text{dw}} > 0$, ψ has a local minimum at $s = 0$ and a global minimum at $s = \hat{s} := 0.750025$ such that $\psi(\hat{s}) = 0$. Moreover, in view of

Proposition 4.3, we measure the violation of the unit-length constraint in terms of the quantity

$$(5.2) \quad \text{err}_{\mathbf{n}} := \|I_h[|\mathbf{n}_h^N|^2 - 1]\|_{L^1(\Omega)},$$

where \mathbf{n}_h^N denotes the final approximation of the director generated by Algorithm 4.1. Furthermore, unless otherwise specified, we choose the L^2 -metric (4.4) in (4.1), and we set the tolerance $\text{tol} = 10^{-6}$ in both (4.2) and (4.7).

5.1. Point defect in 2D. In striking contrast with the Oseen–Frank model, the Ericksen model allows point defects to have finite energy in 2D: The blow-up of $|\nabla \mathbf{n}|$ near a defect is compensated by infinitesimal values of s for the energy $E[s, \mathbf{n}]$ in (1.1) to stay bounded. We examine this basic mechanism with simulations of a point defect in 2D and study the influence of the discretization parameters on the performance of Algorithm 4.1.

We consider the unit square $\Omega = (0, 1)^2$ and set $\kappa = 2$ in (1.1) as well as $c_{\text{dw}} = 0.1(0.3)^{-2}$ in (5.1). We impose Dirichlet boundary conditions for s and \mathbf{n} on $\partial\Omega$, namely

$$(5.3) \quad g = \hat{s} \quad \text{and} \quad \mathbf{q} = \mathbf{r}/g = \frac{(x - 0.5, y - 0.5)}{|(x - 0.5, y - 0.5)|} \quad \text{on } \partial\Omega.$$

To initialize Algorithm 4.1, we consider a constant degree of orientation $s_h^0 = \hat{s}$ in Ω and a director \mathbf{n}_h^0 exhibiting an off-center point defect located at $(0.24, 0.24)$. Due to the imposed boundary conditions and for symmetry reasons, we expect that an energy-decreasing dynamics moves the defect to the center of the square; see Figure 2.

In our first experiment, we consider a uniform mesh \mathcal{T}_h of the unit square consisting of 2048 right triangles. The resulting mesh size is $h = \sqrt{2}2^{-5}$. Moreover, we set $\tau_{\mathbf{n}} = \tau_s = 0.1$ and compare the results obtained for different choices of the metric $\langle \cdot, \cdot \rangle_*$ in (4.1); cf. (4.4)–(4.5). Table 5.1 displays the outputs for each run. On the one hand, we observe that using the L^2 -metric leads to the fastest dynamics in terms of both number of iterations and CPU time. On the other hand, the violation of the unit-length constraint is smaller for the weighted H^1 -metrics. For smaller values of α in the weighted H^1 -metric, Algorithm 4.1 terminates with a configuration exhibiting defect pinning at an off-center location. The expected equilibrium state, depicted in Figure 2 (right), can be restored when reducing the time-step size τ_s .




FIG. 2. Point defect experiment of subsection 5.1: Plot of the approximation (s_h^1, \mathbf{n}_h^1) after the first iteration (left) and of the final approximation (s_h^N, \mathbf{n}_h^N) (right). The gradient flow algorithm moves the defect to the center of the domain.

TABLE 5.1

Point defect experiment of subsection 5.1: Final outputs of Algorithm 4.1 for different choices of metric $\langle \cdot, \cdot \rangle_*$, namely total number of iterations N , value of the energy $E^h[s_h^N, \mathbf{n}_h^N]$ for the equilibrium state, smallest value of the final s_h^N , error in the unit-length constraint in (5.2), and the CPU time.

Metric	N	$E^h[s_h^N, \mathbf{n}_h^N]$	$\min(s_h^N)$	$\text{err}_{\mathbf{n}}$	CPU time
L^2	60	2.984	0.0757	0.0404	64.83
weighted H^1 , $\alpha = 2.0$	67	2.944	0.0750	0.0370	98.65
weighted H^1 , $\alpha = 1.9$	65	2.938	0.0754	0.0362	111.69
weighted H^1 , $\alpha = 1.8$	67	2.932	0.0755	0.0353	130.17
weighted H^1 , $\alpha = 1.7$	80	2.926	0.0760	0.0342	154.92

TABLE 5.2

Point defect experiment of subsection 5.1: final outputs of Algorithm 4.1 for different uniform meshes with mesh size h and time steps $\tau_{\mathbf{n}} = Ch^2$ (top) and different time-step sizes $\tau_{\mathbf{n}}$ with fixed mesh size $h = \sqrt{2}2^{-5}$ (bottom).

h	N	$E^h[s_h^N, \mathbf{n}_h^N]$	$\min(s_h^N)$	$\text{err}_{\mathbf{n}}$	CPU time
$\sqrt{2}2^{-5}$	60	2.984	0.0757	0.0404	64.83
$\sqrt{2}2^{-6}$	61	2.940	0.0422	0.0232	592.23
$\sqrt{2}2^{-7}$	133	2.939	0.0289	0.0100	7919.25

$\tau_{\mathbf{n}}$	$\text{err}_{\mathbf{n}}$
$(0.1)2^{-5}$	0.00610
$(0.1)2^{-6}$	0.00346
$(0.1)2^{-7}$	0.001927

In our second experiment, we investigate the effect of mesh refinement and changes of the time-step size on the results. To this end, we first repeat the simulation using three uniform meshes with $h = \sqrt{2}2^{-5-\ell}$ ($\ell = 0, 1, 2$); we set $\tau_{\mathbf{n}} = 0.12^{-2\ell}$, in agreement with the CFL condition in (4.10) for the L^2 -metric and $d = 2$. We collect the results of computations in Table 5.2 (top) and observe that both $\min(s_h^N)$ and $\text{err}_{\mathbf{n}}$ decrease about linearly with h , whereas the energy $E^h[s_h^N, \mathbf{n}_h^N]$ slightly decreases. The linear decay of $\text{err}_{\mathbf{n}}$ with respect to $\tau_{\mathbf{n}}$ established in (4.9) is not observed. This can be explained by the increase of $E^h[s_h^0, \mathbf{n}_h^0]$ upon refinement, attributable to the fact that \mathbf{n}_h^0 has a point defect while s_h^0 is constant and does not compensate the blow-up of $\nabla \mathbf{n}_0$.

In our third experiment, we aim to empirically confirm the first-order convergence of the error $\text{err}_{\mathbf{n}}$ in (5.2) with respect to $\tau_{\mathbf{n}}$ established in Proposition 4.3; see (4.9). To this end, we consider a fixed mesh with $h = \sqrt{2}2^{-5}$, and we set $\tau_{\mathbf{n}} = (0.1)2^{-5-\ell}$ ($\ell = 0, 1, 2$) as well as $\text{tol} = 10^{-5}\tau_{\mathbf{n}}$ in both (4.2) and (4.7). The computational results, collected in Table 5.2 (bottom), confirm the expected linear decay of the error.

5.2. Plane defect in 3D. We simulate a plane defect in the unit cube $\Omega = (0, 1)^3$ located at $\{z = 0.5\}$, according to [29, section 6.4]. We set $\kappa = 0.2$ in (1.1) and $c_{\text{dw}} = 0$ in (5.1). We impose Dirichlet boundary conditions on the top and bottom faces Γ_D of the cube

$$\begin{aligned} g &= \hat{s}, \quad \mathbf{q} = \mathbf{r}/g = (1, 0, 0) \text{ on } \partial\Omega \cap \{z = 0\}, \\ g &= \hat{s}, \quad \mathbf{q} = \mathbf{r}/g = (0, 1, 0) \text{ on } \partial\Omega \cap \{z = 1\}. \end{aligned}$$




FIG. 3. Plane defect of subsection 5.2: plots of the three components of \mathbf{n}_h^k (first row) and plots of s_h^k (second row) for iterations $k = 1, 31, 79$. In the final configuration ($k = N = 79$), the energy is $E_h[s_h^N, \mathbf{n}_h^N] = 0.247$, $\min(s_h^N) = 0.0101$, and $\text{err}_{\mathbf{n}} = 0.0556$. Moreover, there is a transition layer between about $z = 0.4$ and $z = 0.6$, and s_h is almost linear in $(0, 0.4)$ and $(0.6, 1)$.

The exact solution is $\mathbf{n}(z) = (1, 0, 0)$ for $z < 0.5$ and $\mathbf{n}(z) = (0, 1, 0)$ for $z > 0.5$, while $s(z) = 0$ on $z = 0.5$ and linear on $(0, 0.5) \cup (0.5, 1)$ [29, section 6.4]. Our numerical results are consistent with those in [23, section 5.3]. To initialize Algorithm 4.1, we set $s_h^0 = \hat{s}$ and \mathbf{n}_h^0 to be a regularized point defect away from the center of the cube. Figure 3 displays the three components of \mathbf{n}_h^k and s_h^k evaluated along the vertical line $(0.5, 0.5, z)$ for iterations $k = 1, 31, 79$ computed on a uniform mesh with $h = \sqrt{3}0.05$ and $\tau_{\mathbf{n}} = \tau_s = 0.01$.

5.3. Effect of κ on equilibria. The value of the constant $\kappa > 0$ in (1.1) plays a crucial role in the formation of defects. For large values of κ , the dominant term in $E_1[s, \mathbf{n}]$ is $\int_{\Omega} \kappa |\nabla s|^2$ that prevents variations of s . Typically s tends to be close to a (usually positive) constant and the model behaves much like the simpler Oseen–Frank model, where defects are less likely to occur (and no defects with finite energy beyond point defects are allowed in 3D). On the other hand, for small values of κ , the energy is dominated by $\int_{\Omega} s^2 |\nabla \mathbf{n}|^2$, which allows s to become zero to compensate large gradients of \mathbf{n} , and defects are then more likely to occur. In this section, we investigate this dichotomy numerically.

We consider a cylindrical domain Ω in 3D with lateral boundary Γ_D

$$\begin{aligned} \Omega &= \{(x, y, z) \in \mathbb{R}^3 : (x - 0.5)^2 + (y - 0.5)^2 < 0.5^2, 0 < z < 1\}, \\ \Gamma_D &= \{(x, y, z) \in \mathbb{R}^3 : (x - 0.5)^2 + (y - 0.5)^2 = 0.5^2, 0 < z < 1\}, \end{aligned}$$

and impose the Dirichlet conditions on Γ_D

$$(5.4) \quad g = \hat{s} \quad \text{and} \quad \mathbf{q} = \mathbf{r}/g = \frac{(x - 0.5, y - 0.5, 0)}{|(x - 0.5, y - 0.5, 0)|},$$

The top and bottom faces of Ω are treated as free boundaries and the double well potential ψ is neglected, i.e., $c_{\text{dw}} = 0$ in (5.1). The analysis in [29, section 6.5] predicts




FIG. 4. Effect of κ in subsection 5.3: equilibria for $\kappa = 0.2$ (left) and $\kappa = 2$ (right). Both pictures show s_h^N and \mathbf{n}_h^N on the slices $z = 0.2, 0.5, 0.8$. If $\kappa = 0.2$, the final configuration exhibits a line defect along the central axis of the cylinder; the final energy is $E^h[s_h^N, \mathbf{n}_h^N] = 0.806$, $\min(s_h^N) = -7.33 \times 10^{-4}$, $\text{err}_{\mathbf{n}} = 0.0778$, and $N = 226$. If $\kappa = 2$, the z -component of the director is not zero. This behavior is usually referred to as the fluting effect or escape to the third dimension [29, section 6.5.1]. Moreover, the degree of orientation is bounded well away from zero; the final energy is $E^h[s_h^N, \mathbf{n}_h^N] = 2.635$, $\min(s_h^N) = 0.224$, $\text{err}_{\mathbf{n}} = 0.044$, and $N = 17$.

that minimizers of the energy exhibit a line defect along the central axis of the cylinder if κ is sufficiently small, whereas they are smooth (no defects) if κ is sufficiently large.

Figure 4 displays the final configurations obtained for $\kappa = 0.2$ and $\kappa = 2$. To discretize Ω , we consider an unstructured mesh generated by Netgen with $h_0 = 0.05$. For both values of κ , we set \hat{s} as initial condition for the degree of orientation. For $\kappa = 0.2$, we set $\tau_{\mathbf{n}} = 0.1$ and $\tau_s = 10^{-3}$ and take as initial condition for the director field an off-center point defect located at the slice $z = 0.5$. For $\kappa = 2$, we set $\tau_{\mathbf{n}} = \tau_s = 0.01$ and initialize \mathbf{n}_h^0 as an off-center point defect located at the slice $z = 0.25$. These computational results are consistent with those in [23] and confirm the predicted effect of κ [29, section 6.5].

5.4. Propeller defect. In this section, we investigate a new defect discovered in [23, section 5.4]. We consider a setup similar to the one discussed in subsection 5.3, except that the domain is the unit cube $\Omega = (0, 1)^3$, and we again set $c_{\text{dw}} = 0$ in (5.1). The top and bottom faces of the cube are treated as free boundary, while the same strong anchoring conditions as in (5.4) are imposed on the vertical faces Γ_D of the cube (lateral boundary). The initial conditions are $s_h^0 = \hat{s}$ for the degree of orientation and an off-center point defect located on the slice $z = 0.5$ for the director. The domain is discretized using an unstructured mesh generated by Netgen with $h_0 = 0.025$, and we set $\tau_{\mathbf{n}} = 0.02$. We consider the values $\kappa = 2$ and $\kappa = 0.1$. For $\kappa = 2$ and $\tau_s = 0.2$, the computational results agree with those of subsection 5.3: The equilibrium state is smooth and is characterized by a nonzero z -component (fluting effect).

For $\kappa = 0.1$, the final configuration reported in [23, section 5.4, Figure 5] consists of two plane defects intersecting at the vertical symmetry axis of the cube, the so-called propeller defect. Whether this was a numerical artifact due to the inherent symmetries of the structured uniform weakly acute meshes used in [23] for simulation was an intriguing open question that we now answer. Owing to the flexibility of our approach regarding meshes, we repeated the experiment using an unstructured non-symmetric mesh with $\tau_s = 10^{-4}$. Our computational results confirm the emergence of the propeller defect in Figure 5, which in turn displays the director field \mathbf{n}_h^k at iterations $k = 0, 1, 2766$ with colors indicating the size of s_h^k .




FIG. 5. Propeller defect of subsection 5.4: evolution of the order parameters on the top face of the cube ($z = 1$). Plots of the initial state (s_h^0, \mathbf{n}_h^0) (left), of the intermediate approximation (s_h^1, \mathbf{n}_h^1) obtained after the first iteration (middle), and of the equilibrium state (s_h^N, \mathbf{n}_h^N) after 2766 iterations (right). In the initial state, due to the off-center point defect at $z = 0.5$, there is a corresponding region on the slice for $z = 1$ where \mathbf{n} is aligned with the z -direction. After the first iteration, in which \mathbf{n} is minimized for fixed $s = \hat{s}$, by symmetry the defect has moved to the center on $z = 0.5$. Correspondingly, on the top surface of the cube, the region where \mathbf{n} is aligned with the z -axis has moved to the center. The final state is a propeller defect consisting of a planar X -like configuration extruded in the z -direction. The final energy is $E^h[s_h^N, \mathbf{n}_h^N] = 0.592$, $\min(s_h^N) = -1.575 \times 10^{-4}$, $\text{err}_{\mathbf{n}} = 0.0265$, and $N = 2766$.

5.5. Colloidal effects in nematic LCs. Colloidal particles suspended in a nematic LC can induce interesting topological defects and distortions [19, 28]. One prominent example is the so-called Saturn ring defect, a director configuration characterized by a circular ring singularity surrounding a spherical particle and located around its equator. Such defects are typically nonorientable and captured within the Landau-de Gennes Q -tensor model [11, 12], but the Ericksen model yields similar orientable defects under suitable boundary conditions [24]. We confirm the ability of Algorithm 4.1 to produce similar configurations.

In this section, we exploit the flexibility of Algorithm 4.1 regarding meshes, together with the built-in constructive solid geometry approach of Netgen/NGSolve, to explore numerically the formation of Saturn-ring-like defects induced by nonspherical or multiple particles.

5.5.1. One ellipsoidal particle. Let $\Omega_c = (0, 1)^3$ be the unit cube and let $\Omega_s \subset \Omega_c$ be an ellipsoid centered at $(0.5, 0.5, 0.5)$ with axes parallel to the coordinate axes and semiaxis lengths equal to 0.3 (x -direction), 0.075 (y -direction), and 0.075 (z -direction); Ω_s has an aspect ratio 1 : 4. The computational domain is then $\Omega := \Omega_c \setminus \overline{\Omega_s}$. We set $\kappa = 1$ in (1.1) as well as $c_{\text{dw}} = 0.2$ in (5.1). On $\partial\Omega = \partial\Omega_c \cup \partial\Omega_s$, we impose strong anchoring conditions

$$(5.5) \quad g = \hat{s} \text{ on } \partial\Omega, \quad \mathbf{q} = \mathbf{r}/g = \boldsymbol{\nu} \text{ on } \partial\Omega_s, \quad \text{and} \quad \mathbf{q} = \mathbf{r}/g = \mathbf{n}_{sr} \text{ on } \partial\Omega_c,$$

where $\boldsymbol{\nu} : \partial\Omega_s \rightarrow \mathbb{S}^2$ denotes the outward-pointing unit normal vector of Ω_s and $\mathbf{n}_{sr} : \partial\Omega_s \rightarrow \mathbb{S}^2$ smoothly interpolates between the constant values $(0, 0, -1)$ on the bottom face and $(0, 0, 1)$ on the top face of the cube (see [24, Figure 11]). These boundary conditions are essential in order to induce the defect. The initial conditions for Algorithm 4.1 are given by

$$(5.6) \quad s_h^0 = \hat{s} \text{ in } \Omega \quad \text{and} \quad \mathbf{n}_h^0(z) = \begin{cases} (0, 0, 1), & z \in \Omega \text{ and } z_3 \geq 0.5, \\ (0, 0, -1), & z \in \Omega \text{ and } z_3 < 0.5, \\ \mathbf{q}(z), & z \in \partial\Omega, \end{cases}$$




FIG. 6. *Saturn ring experiment of subsection 5.5.1. Three different perspectives of the Saturn ring defect around an ellipsoidal particle: slice $z = 0.5$ (left), a three-dimensional view clipped at $y = 0.5$ (middle), and a three-dimensional view clipped at $x = 0.5$ (right). The blue ring surrounding the particle, the iso-surface for $s = 0.15$, provides a good approximation of the defect. We stress that neither the distance between the defect and the particle nor the defect diameter is constant, which is a consequence of the anisotropic shape of the particle. The final energy is $E^h[s_h^N, \mathbf{n}_h^N] = 7.263$, $\min(s_h^N) = 0.0128$, $\text{err}_{\mathbf{n}} = 0.145$, and $N = 33$.*




FIG. 7. *Two-particle experiment of subsection 5.5.2. Fat figure “8” defect around two spherical colloids viewed from different perspectives: slice $y = 0.5$ (left), slice $z = 0.5$ (middle), and a three-dimensional view clipped at $y = 0.5$ (right). The blue ring surrounding the particle is the iso-surface for $s = 0.12$, which provides a good approximation of the defect. The final energy is $E^h[s_h^N, \mathbf{n}_h^N] = 7.656$, $\min(s_h^N) = 0.0146$, $\text{err}_{\mathbf{n}} = 0.0972$, and $N = 57$.*

for $z = (z_1, z_2, z_3) \in \mathcal{N}_h$. Figure 6 displays cuts of the final configuration obtained using Algorithm 4.1 with an unstructured mesh with $h_0 = 0.05$ and time-step sizes $\tau_{\mathbf{n}} = \tau_s = 0.01$.

5.5.2. Multiple spherical particles. We conclude this section with two novel and challenging simulations involving multiple spherical colloidal particles. In both cases, the domain has the form $\Omega := \Omega_c \setminus \overline{\Omega_s}$, where $\Omega_c \subset \mathbb{R}^3$ denotes a simply connected domain (representing the LC container), whereas $\Omega_s \subset \Omega_c$ denotes the region occupied by spherical colloidal particles. We set $\kappa = 1$ in (1.1) and $c_{\text{dw}} = 0.2$ in (5.1). Moreover, boundary and initial conditions are suitable extensions to the multiple particle case of (5.5) and (5.6) considered in subsection 5.5.1.

Figure 7 shows the equilibrium state corresponding to $\Omega_c = (0, 1)^3$ and a pair of disjoint spherical colloids Ω_s with radii 0.1 and centered at $(0.3, 0.5, 0.5)$ and $(0.7, 0.5, 0.5)$. Algorithm 4.1 employs an unstructured mesh with $h_0 = 0.025$ and time-step sizes $\tau_{\mathbf{n}} = \tau_s = 0.0025$. A novel *fat figure “8”* defect forms.

Figure 8 depicts the equilibrium state corresponding to $\Omega_c = (-0.1, 1.1)^3$ and a colloidal region consisting of six spheres. The latter have radii 0.1 and centers located at $(0.2, 0.5, 0.5)$, $(0.8, 0.5, 0.5)$, $(0.5, 0.2, 0.5)$, $(0.5, 0.8, 0.5)$, $(0.5, 0.5, 0.2)$, and $(0.5, 0.5, 0.8)$ distributed symmetrically with respect to the cube center. Algorithm 4.1 utilizes an unstructured mesh with $h_0 = 0.05$ and time-step sizes $\tau_{\mathbf{n}} = \tau_s = 0.005$.




FIG. 8. Six-particle experiment of subsection 5.5.2. Defect around six spherical colloids viewed from different perspectives: slice $y = 0.5$ (left), slice $z = 0.5$ (middle), and a three-dimensional view clipped at $y = 0.5$ (right); the slice $x = 0.5$ is similar to $y = 0.5$. The blue ring surrounding the particles is the iso-surface for $s = 0.22$, which provides a good approximation of the defect. Therefore the defect appears to be a combination of a large Saturn ring defect around particles with center in the plane $z = 0.5$ and a planar X-like configuration with axis $x = 0.5, y = 0.5, -0.1 < z < 1$. The final energy is $E^h[s_h^N, \mathbf{n}_h^N] = 12.562$, $\min(s_h^N) = -0.0079$, $\text{err}_{\mathbf{n}} = 0.163$, and $N = 61$.

6. Proofs. In this section, we present the proofs of the results discussed in sections 2 to 4.

6.1. L^2 -differentiability of admissible directors. We now prove that any admissible director field, despite not being in $\mathbf{H}^1(\Omega)$, is L^2 -differentiable in $\Omega \setminus \Sigma$. We refer to [11] for a similar argument for a line field.

Proof of Proposition 2.1. Since $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}$, we have that $s \in H^1(\Omega)$ and $\mathbf{u} = s\mathbf{n} \in \mathbf{H}^1(\Omega)$. Then, for almost all $x \in \Omega$ (specifically, for all Lebesgue points of $(s, \mathbf{u}, \nabla s, \nabla \mathbf{u})$), s and \mathbf{u} are L^2 -differentiable and their L^2 -gradients coincide with their respective weak gradients for a.e. $x \in \Omega$, i.e., as $r \rightarrow 0$, it holds that

$$\begin{aligned} \int_{B_r(x)} |s(y) - s(x) - \nabla s(x) \cdot (y - x)|^2 dy &= o(r^2), \\ \int_{B_r(x)} |\mathbf{u}(y) - \mathbf{u}(x) - \nabla \mathbf{u}(x)(y - x)|^2 dy &= o(r^2); \end{aligned}$$

see [18, Theorem 6.2]. For almost all $x \in \Omega \setminus \Sigma$ (specifically, for all Lebesgue points of $(s, \mathbf{n}, \mathbf{u}, \nabla s, \nabla \mathbf{u})$ in $x \in \Omega \setminus \Sigma$), in view of the identity (2.3), we define the quantity

$$(6.1) \quad \nabla \mathbf{n}(x) := \frac{\nabla \mathbf{u}(x) - \mathbf{n}(x) \otimes \nabla s(x)}{s(x)}.$$

Let $r > 0$. It holds that

$$\begin{aligned} & \int_{B_r(x)} |\mathbf{n}(y) - \mathbf{n}(x) - \nabla \mathbf{n}(x)(y - x)|^2 dy \\ & \leq \frac{1}{s(x)^2} \int_{B_r(x)} |\mathbf{u}(y) - \mathbf{u}(x) - \nabla \mathbf{u}(x)(y - x)|^2 dy \\ & \quad + \frac{1}{s(x)^2} \int_{B_r(x)} |s(y) - s(x) - \nabla s(x) \cdot (y - x)|^2 |\mathbf{n}(y)|^2 dy \\ & \quad + \frac{|\nabla s(x)|^2}{s(x)^2} \int_{B_r(x)} |\mathbf{n}(y) - \mathbf{n}(x)|^2 |y - x|^2 dy = o(r^2) \end{aligned}$$

as $r \rightarrow 0$. This shows that $\nabla \mathbf{n}(x)$ is the L^2 -gradient of \mathbf{n} at x . Moreover, (2.9) follows from a direct computation. In fact, in view of (6.1), there holds that

$$\begin{aligned} s(x)^2 |\nabla \mathbf{n}(x)|^2 &= |\nabla \mathbf{u}(x) - \mathbf{n}(x) \otimes \nabla s(x)|^2 \\ &= |\nabla \mathbf{u}(x)|^2 + |\mathbf{n}(x) \otimes \nabla s(x)|^2 - 2 \nabla \mathbf{u}(x) : [\mathbf{n}(x) \otimes \nabla s(x)] \\ &= |\nabla \mathbf{u}(x)|^2 - |\nabla s(x)|^2, \end{aligned}$$

where the last equality follows from the identities

$$|\mathbf{n}(x) \otimes \nabla s(x)|^2 = \sum_{i,j=1}^d n_i(x)^2 (\partial_j s(x))^2 = \sum_{j=1}^d (\partial_j s(x))^2 = |\nabla s(x)|^2$$

and for a.e. $x \in \Omega \setminus \Sigma$

$$\begin{aligned} \nabla \mathbf{u}(x) : [\mathbf{n}(x) \otimes \nabla s(x)] &= \sum_{i,j=1}^d \partial_j u_i(x) n_i(x) \partial_j s(x) = \frac{1}{s(x)} \sum_{i,j=1}^d \partial_j u_i(x) u_i(x) \partial_j s(x) \\ &= \frac{1}{2s(x)} \sum_{i,j=1}^d \partial_j |u_i(x)|^2 \partial_j s(x) = \frac{1}{2s(x)} \sum_{j=1}^d \partial_j |\mathbf{u}(x)|^2 \partial_j s(x) \\ &= \frac{1}{2s(x)} \sum_{j=1}^d \partial_j (s(x)^2) \partial_j s(x) = \sum_{j=1}^d (\partial_j s(x))^2 = |\nabla s(x)|^2. \end{aligned}$$

This concludes the proof. \square

6.2. Lim-sup inequality (consistency). We start with two results from [23] that we state without proofs. The first one shows that the degree of orientation s can be truncated near the end points of the domain of definition $(-1/(d-1), 1)$ of ψ without increasing the energy $E[s, \mathbf{n}]$. We refer to [23, Lemma 3.1] for a proof.

LEMMA 6.1 (truncation of s). *Let the assumptions (2.12) and (2.14) hold. Let $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}(g, \mathbf{r})$. For all $0 < \rho \leq \delta_0$, define*

$$s_\rho(x) := \min \left\{ 1 - \rho, \max \left\{ -\frac{1}{d-1} + \rho, s(x) \right\} \right\} \quad \text{and} \quad \mathbf{u}_\rho(x) := s_\rho(x) \mathbf{n}(x)$$

for a.e. $x \in \Omega$. Then, $(s_\rho, \mathbf{n}, \mathbf{u}_\rho) \in \mathcal{A}(g, \mathbf{r})$ and $E_1[s_\rho, \mathbf{n}] \leq E_1[s, \mathbf{n}]$, $E_2[s_\rho] \leq E_2[s]$.

A simple consequence of Lemma 6.1, based on the convergence of the characteristic function $\chi_{\{s_\rho=s\}} \rightarrow \chi_\Omega$ as $\rho \rightarrow 0$, is that $\|(s, \mathbf{u}) - (s_\rho, \mathbf{u}_\rho)\|_{H^1(\Omega)^{1+d}} \rightarrow 0$ as $\rho \rightarrow 0$. The second result is about regularization of admissible functions but preserving the structural condition (2.6) and boundary values. This is a rather tricky two-scale process fully discussed in [23, Proposition 3.2].

LEMMA 6.2 (regularization of functions in $\mathcal{A}(g, \mathbf{r})$). *Let the assumptions (2.12) and (2.13) hold, and suppose that $\Gamma_D = \partial\Omega$. Let $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}(g, \mathbf{r})$ and $\rho \leq \delta_0$ such that $-1/(d-1) + \rho \leq s(x) \leq 1 - \rho$ for a.e. $x \in \Omega$. Then, for all $\sigma > 0$, there exists a triple $(s_\sigma, \mathbf{n}_\sigma, \mathbf{u}_\sigma) \in \mathcal{A}(g, \mathbf{r})$ such that $s_\sigma \in W^{1,\infty}(\Omega)$ and $\mathbf{u}_\sigma \in \mathbf{W}^{1,\infty}(\Omega)$. Moreover, there holds $\|(s, \mathbf{u}) - (s_\sigma, \mathbf{u}_\sigma)\|_{H^1(\Omega)^{1+d}} \leq \sigma$, $\|\mathbf{n} - \mathbf{n}_\sigma\|_{L^2(\Omega \setminus \Sigma)} \leq \sigma$, and $-1/(d-1) + \rho \leq s_\sigma(x) \leq 1 - \rho$ for all $x \in \Omega$.*

We recall the following classical local inverse estimates (see, e.g., [8, Lemma 3.5]), which will be used in several points of the upcoming analysis: For all $v_h \in V_h$, $1 \leq p, r \leq \infty$, and $K \in \mathcal{T}_h$, there holds

$$\|\nabla v_h\|_{\mathbf{L}^p(K)} \leq Ch_K^{-1} \|v_h\|_{L^p(K)} \quad \text{and} \quad \|v_h\|_{L^p(K)} \leq Ch_K^{d(r-p)/(pr)} \|v_h\|_{L^r(K)},$$

where $C > 0$ is a constant depending only on the shape-regularity of $\{\mathcal{T}_h\}$.

It is well known that the Lagrange interpolation operator $I_h : C(\bar{\Omega}) \rightarrow V_h$ is not stable in $H^1(\Omega)$ unless $d = 1$. We exploit stability in $L^\infty(\Omega)$ to derive stability in $W^{1,p}(\Omega)$ for $p > d$.

LEMMA 6.3 ($W^{1,p}$ -stability of Lagrange interpolant). *Let $v \in W^{1,p}(\Omega)$ for $d < p \leq \infty$. Then*

$$(6.2) \quad \|\nabla I_h[v]\|_{\mathbf{L}^p(K)} \leq C \|\nabla v\|_{\mathbf{L}^p(K)} \quad \text{for all } K \in \mathcal{T}_h,$$

where $C > 0$ depends only on the shape-regularity of $\{\mathcal{T}_h\}$.

Proof. Let $K \in \mathcal{T}_h$ be an arbitrary element and let $\bar{v}_K = f_K v$. An inverse estimate gives

$$\|\nabla I_h[v]\|_{\mathbf{L}^p(K)}^p \leq |K| \|\nabla I_h[v - \bar{v}_K]\|_{\mathbf{L}^\infty(K)}^p \lesssim h_K^{d-p} \|v - \bar{v}_K\|_{L^\infty(K)}^p.$$

The Bramble–Hilbert estimate yields $\|v - \bar{v}_K\|_{L^\infty(K)} \lesssim h_K^{1-d/p} \|\nabla v\|_{\mathbf{L}^p(K)}$ and ends the proof. \square

Applying a standard density argument in $W^{1,p}(\Omega)$, for $d < p < \infty$, we deduce

$$(6.3) \quad \lim_{h \rightarrow 0} \|v - I_h[v]\|_{W^{1,p}(\Omega)} = 0 \quad \text{for all } v \in W^{1,p}(\Omega).$$

We have collected all the ingredients to show the existence of a recovery sequence (lim-sup inequality).

Proof of Theorem 3.1(i). For the sake of clarity, we decompose the proof into seven steps.

Step 1: Regularization. Let $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}(g, \mathbf{r})$. For all $k \in \mathbb{N}$ such that $1/k \leq \delta_0$, let $0 < \sigma_k \leq 1/k$ be sufficiently small. Applying successively Lemma 6.1 (with $\rho = 1/k$) and Lemma 6.2 (with $\sigma = \sigma_k$), we obtain $(s_k, \mathbf{n}_k, \mathbf{u}_k) \in \mathcal{A}(g, \mathbf{r})$ satisfying $(s_k, \mathbf{u}_k) \in [W^{1,\infty}(\Omega)]^{1+d}$ and $-1/(d-1) + 1/k \leq s_k \leq 1 - 1/k$ in Ω for all k . Moreover, we have that

$$\|(s, \mathbf{u}) - (s_k, \mathbf{u}_k)\|_{H^1(\Omega)^{1+d}} \leq \sigma_k \quad \text{and} \quad \|\mathbf{n} - \mathbf{n}_k\|_{\mathbf{L}^2(\Omega \setminus \Sigma)} \leq \sigma_k.$$

Since $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}(g, \mathbf{r})$, Proposition 2.1 guarantees that \mathbf{n} is L^2 -differentiable a.e. in $\Omega \setminus \Sigma$, with its L^2 -gradient given by (2.8) and that the identity (2.9) holds. The same result is valid for \mathbf{n}_k a.e. in $\Omega \setminus \Sigma_k$, where $\Sigma_k := \{x \in \Omega : s_k(x) = 0\}$.

We have convergence of the energy: $E[s_k, \mathbf{n}_k] \rightarrow E[s, \mathbf{n}]$ as $k \rightarrow \infty$. To see this, we first observe that, thanks to (2.5), we have that

$$\begin{aligned} E_1[s_k, \mathbf{n}_k] &= \tilde{E}_1[s_k, \mathbf{u}_k] = \frac{1}{2} \int_{\Omega} (\kappa - 1) |\nabla s_k|^2 + |\nabla \mathbf{u}_k|^2 \\ &\rightarrow \frac{1}{2} \int_{\Omega} (\kappa - 1) |\nabla s|^2 + |\nabla \mathbf{u}|^2 = \tilde{E}_1[s, \mathbf{u}] = E_1[s, \mathbf{n}] \quad \text{as } k \rightarrow \infty. \end{aligned}$$




FIG. 9. A schematic illustration of the mutual relations of the sets defined in Steps 1–2 of the proof of Theorem 3.1(i) (lim-sup inequality). Note that the set $\Sigma_k \subset \Omega$ is closed, as it is the preimage of a closed set with respect to the continuous function s_k , but it might be more topologically complicated than in the picture.

Moreover, the monotonicity of ψ in $(-1/(d-1), -1/(d-1) + \delta_0)$ and in $(1 - \delta_0, 1)$ translates into $\psi(s_k) \geq 0$ increasing and converging pointwise to $\psi(s)$, whence the monotone convergence theorem gives

$$E_2[s_k] = \int_{\Omega} \psi(s_k) \rightarrow \int_{\Omega} \psi(s) = E_2[s] \quad \text{as } k \rightarrow \infty.$$

Let $\epsilon > 0$ be arbitrary. The above convergences guarantee the existence of $k \in \mathbb{N}$ such that $\sigma_k < \epsilon$, $|E[s_k, \mathbf{n}_k] - E[s, \mathbf{n}]| < \epsilon$, and $|(\Sigma_k \setminus \Sigma) \cup (\Sigma \setminus \Sigma_k)| < \epsilon$. Let such a $k \in \mathbb{N}$ be fixed for the rest of the proof.

Step 2: Discretization. Let $s_{k,h} := I_h[s_k]$ and $\mathbf{u}_{k,h} := I_h[\mathbf{u}_k]$. Let $\mathbf{n}_{k,h} \in \mathbf{V}_h$ be defined as

$$\mathbf{n}_{k,h}(z) := \begin{cases} \mathbf{u}_{k,h}(z)/s_{k,h}(z) = \mathbf{u}_k(z)/s_k(z) & \text{if } z \in \mathcal{N}_h \cap (\Omega \setminus \Sigma_k), \\ \text{an arbitrary unit vector} & \text{if } z \in \mathcal{N}_h \cap \Sigma_k. \end{cases}$$

Note that, by construction, $(s_{k,h}, \mathbf{n}_{k,h}, \mathbf{u}_{k,h})$ satisfies the discrete structural condition (3.2) and $\|\mathbf{n}_{k,h}\|_{L^\infty(\Omega)} = 1$. Moreover, since $0 = \|I_h[|\mathbf{n}_{k,h}|^2] - 1\|_{L^1(\Omega)} \leq \varepsilon$ as well as $s_{k,h}(z) = g_h(z)$ and $\mathbf{u}_{k,h}(z) = \mathbf{r}_h(z)$ for all $z \in \mathcal{N}_h \cap \Gamma_D$, we deduce that $(s_{k,h}, \mathbf{n}_{k,h}, \mathbf{u}_{k,h}) \in \mathcal{A}_{h,\varepsilon}(g_h, \mathbf{r}_h)$.

Given $\delta > 0$, we consider the sets

$$\Sigma_{k,\delta} := \{x \in \Omega : |s_k(x)| \leq \delta\} \quad \text{and} \quad \Omega_{k,\delta}^h := \bigcup \{K \in \mathcal{T}_h : K \cap \Sigma_{k,\delta} = \emptyset\}.$$

Note that, by construction, there holds $\Omega_{k,\delta}^h \subset \Omega \setminus \Sigma_{k,\delta}$; see Figure 9.

Let $K \in \mathcal{T}_h$ such that $K \cap \Sigma_{k,\delta} \neq \emptyset$. In particular, there exists $x_0 \in K \cap \Sigma_{k,\delta}$. For $x_1 \in K$ arbitrary, the Lipschitz continuity of s_k yields

$$|s_k(x_1)| \leq |s_k(x_0)| + |s_k(x_1) - s_k(x_0)| \leq \delta + \|\nabla s_k\|_{L^\infty(\Omega)} h.$$

Hence, $\Omega \setminus \Omega_{k,\delta}^h \subset \Sigma_{k,2\delta}$ provided h and δ are such that $h\|\nabla s_k\|_{L^\infty(\Omega)} \leq \delta$; see Figure 9.

Now, for any $x \in \Omega_{k,\delta}^h$, we infer that

$$|s_k(x) - s_{k,h}(x)| \leq \|s_k - s_{k,h}\|_{L^\infty(\Omega_{k,\delta}^h)} = \|s_k - I_h[s_k]\|_{L^\infty(\Omega_{k,\delta}^h)} \lesssim h \|\nabla s_k\|_{L^\infty(\Omega_{k,\delta}^h)},$$

whence

$$|s_{k,h}(x)| \geq |s_k(x)| - |s_k(x) - s_{k,h}(x)| > \delta - Ch \|\nabla s_k\|_{L^\infty(\Omega_{k,\delta}^h)} > \delta/2$$

provided h and δ are such that $Ch \|\nabla s_k\|_{L^\infty(\Omega)} < \delta/2$. Hence, for this range of parameters, we can define $\tilde{\mathbf{n}}_k := \mathbf{u}_{k,h}/s_{k,h}$ in $\Omega_{k,\delta}^h$. Note that, by definition, the relation $\mathbf{n}_{k,h} = I_h[\tilde{\mathbf{n}}_k]$ in $\Omega_{k,\delta}^h$ holds.

To conclude this step, we observe that the L^2 -gradient $\nabla \mathbf{n}_k$ of \mathbf{n}_k exists a.e. in $\Omega \setminus \Sigma_k$ and

$$(6.4) \quad \int_{\Omega_{k,\delta}^h} |\nabla \mathbf{n}_k - \nabla \mathbf{n}_{k,h}|^2 \lesssim \int_{\Omega_{k,\delta}^h} |\nabla \mathbf{n}_k - \nabla \tilde{\mathbf{n}}_k|^2 + \int_{\Omega_{k,\delta}^h} |\nabla \tilde{\mathbf{n}}_k - \nabla \mathbf{n}_{k,h}|^2,$$

where $\nabla \tilde{\mathbf{n}}_k$ and $\nabla \mathbf{n}_{k,h}$ denote the weak gradients of $\tilde{\mathbf{n}}_k$ and $\mathbf{n}_{k,h}$, respectively, which coincide elementwise with their classical gradients in $\Omega_{k,\delta}^h$. In the following two steps, we will show that, for fixed $k \in \mathbb{N}$ (cf. Step 1), both terms on the right-hand side of (6.4) converge to 0 if $h, \delta \rightarrow 0$ in an appropriate way (note that we are completely free to choose the speed of convergence of the parameters).

Step 3: Proof of $\lim_{h,\delta \rightarrow 0} \int_{\Omega_{k,\delta}^h} |\tilde{\mathbf{n}}_k - \mathbf{n}_{k,h}|^2 + |\nabla \tilde{\mathbf{n}}_k - \nabla \mathbf{n}_{k,h}|^2 = 0$. Since $\mathbf{n}_{k,h} = I_h[\tilde{\mathbf{n}}_k]$ in $\Omega_{k,\delta}^h$, a classical local interpolation estimate yields that

$$\int_{\Omega_{k,\delta}^h} |\nabla \tilde{\mathbf{n}}_k - \nabla \mathbf{n}_{k,h}|^2 = \sum_{\substack{K \in \mathcal{T}_h \\ K \cap \Sigma_k, \delta = \emptyset}} \int_K |\nabla(\tilde{\mathbf{n}}_k - I_h[\tilde{\mathbf{n}}_k])|^2 \lesssim \sum_{\substack{K \in \mathcal{T}_h \\ K \cap \Sigma_k, \delta = \emptyset}} h_K^2 \|D^2 \tilde{\mathbf{n}}_k\|_{L^2(K)}^2.$$

Similarly, there holds

$$\int_{\Omega_{k,\delta}^h} |\tilde{\mathbf{n}}_k - \mathbf{n}_{k,h}|^2 \lesssim \sum_{\substack{K \in \mathcal{T}_h \\ K \cap \Sigma_k, \delta = \emptyset}} h_K^4 \|D^2 \tilde{\mathbf{n}}_k\|_{L^2(K)}^2.$$

Moreover, in view of $\tilde{\mathbf{n}}_k = \mathbf{u}_{k,h}/s_{k,h}$ in $\Omega_{k,\delta}^h$, explicit computations reveal that

$$\begin{aligned} \partial_i \tilde{\mathbf{n}}_k &= s_{k,h}^{-1} \partial_i \mathbf{u}_{k,h} - s_{k,h}^{-2} \partial_i s_{k,h} \mathbf{u}_{k,h} = s_{k,h}^{-1} (\partial_i \mathbf{u}_{k,h} - \partial_i s_{k,h} \tilde{\mathbf{n}}_k), \\ \partial_j \partial_i \tilde{\mathbf{n}}_k &= s_{k,h}^{-1} (s_{k,h}^{-1} \partial_j s_{k,h} \partial_i s_{k,h} \tilde{\mathbf{n}}_k - \partial_i s_{k,h} \partial_j \tilde{\mathbf{n}}_k - s_{k,h}^{-1} \partial_j s_{k,h} \partial_i \mathbf{u}_{k,h}), \end{aligned}$$

for all $1 \leq i, j \leq d$ (note that the second derivatives of the piecewise affine functions $s_{k,h}$ and $\mathbf{u}_{k,h}$ vanish). Several applications of the generalized Hölder inequality, stability (6.2) in $W^{1,p}$ of the Lagrange interpolation operator I_h when $d < p \leq \infty$, in conjunction with the lower bound $|s_{k,h}| > \delta/2$ in $\Omega_{k,\delta}^h$ and the uniform boundedness of $\mathbf{u}_{k,h}$ in $L^\infty(\Omega)$, thus yield

$$\begin{aligned} \|D^2 \tilde{\mathbf{n}}_k\|_{L^2(K)} &\lesssim \delta^{-3} \|\nabla s_{k,h}\|_{L^s(K)}^2 \|\mathbf{u}_{k,h}\|_{L^4(K)} \\ &\quad + \delta^{-1} \|\nabla s_{k,h}\|_{L^4(K)} \left(\delta^{-1} \|\nabla \mathbf{u}_{k,h}\|_{L^4(K)} + \delta^{-2} \|\mathbf{u}_{k,h}\|_{L^s(K)} \|\nabla s_{k,h}\|_{L^s(K)} \right) \\ &\quad + \delta^{-2} \|\nabla s_{k,h}\|_{L^4(K)} \|\nabla \mathbf{u}_{k,h}\|_{L^4(K)} \\ &\lesssim |K|^{1/2} (\delta^{-3} \|\nabla s_k\|_{L^\infty(K)}^2 + \delta^{-2} \|\nabla s_k\|_{L^\infty(K)} \|\nabla \mathbf{u}_k\|_{L^\infty(K)}). \end{aligned}$$

Altogether, we thus obtain the estimate

$$\begin{aligned} \int_{\Omega_{k,\delta}^h} |\nabla \tilde{\mathbf{n}}_k - \nabla \mathbf{n}_{k,h}|^2 + \sum_{\substack{K \in \mathcal{T}_h \\ K \cap \Sigma_{k,\delta} = \emptyset}} h_K^{-2} \int_K |\tilde{\mathbf{n}}_k - \mathbf{n}_{k,h}|^2 &\lesssim \sum_{\substack{K \in \mathcal{T}_h \\ K \cap \Sigma_{k,\delta} = \emptyset}} h_K^2 \|D^2 \tilde{\mathbf{n}}_k\|_{L^2(K)}^2 \\ &\lesssim h^2 (\delta^{-3} \|\nabla s_k\|_{L^\infty(\Omega)}^2 + \delta^{-2} \|\nabla s_k\|_{L^\infty(\Omega)} \|\nabla \mathbf{u}_k\|_{L^\infty(\Omega)})^2, \end{aligned}$$

which yields the desired convergence, if $h\delta^{-3}$ goes to 0 as $h, \delta \rightarrow 0$.

Step 4: Proof of $\lim_{h,\delta \rightarrow 0} \int_{\Omega_{k,\delta}^h} |\mathbf{n}_k - \tilde{\mathbf{n}}_k|^2 + |\nabla \mathbf{n}_k - \nabla \tilde{\mathbf{n}}_k|^2 = 0$. We first observe that

$$\begin{aligned} \|\tilde{\mathbf{n}}_k - \mathbf{n}_k\|_{L^p(\Omega_{k,\delta}^h)} &= \|s_{k,h}^{-1} \mathbf{u}_{k,h} - s_k^{-1} \mathbf{u}_k\|_{L^p(\Omega_{k,\delta}^h)} \\ &\lesssim \delta^{-2} \|s_k - s_{k,h}\|_{L^p(\Omega_{k,\delta}^h)} \|\mathbf{u}_{k,h}\|_{L^\infty(\Omega_{k,\delta}^h)} + \delta^{-1} \|\mathbf{u}_{k,h} - \mathbf{u}_k\|_{L^p(\Omega_{k,\delta}^h)} \\ &\lesssim \delta^{-2} h \|\nabla s_k\|_{L^\infty(\Omega)} + \delta^{-1} h \|\nabla \mathbf{u}_k\|_{L^\infty(\Omega)} \end{aligned}$$

for all $1 \leq p < \infty$. This shows that $\|\tilde{\mathbf{n}}_k - \mathbf{n}_k\|_{L^p(\Omega_{k,\delta}^h)} \rightarrow 0$ if $\delta^{-2}h \rightarrow 0$ as $h, \delta \rightarrow 0$. To deal with the gradient part, we resort to available expressions of $\nabla \mathbf{n}_k$ and $\nabla \tilde{\mathbf{n}}_k$ to write

$$\begin{aligned} \int_{\Omega_{k,\delta}^h} |\nabla \mathbf{n}_k - \nabla \tilde{\mathbf{n}}_k|^2 &= \int_{\Omega_{k,\delta}^h} |s_k^{-1} (\nabla \mathbf{u}_k - \mathbf{n}_k \otimes \nabla s_k) - s_{k,h}^{-1} (\nabla \mathbf{u}_{k,h} - \tilde{\mathbf{n}}_k \otimes \nabla s_{k,h})|^2 \\ &\leq T_1 + T_2 + T_3, \end{aligned}$$

where

$$\begin{aligned} T_1 &:= \int_{\Omega_{k,\delta}^h} |s_{k,h}^{-1} (\nabla \mathbf{u}_k - \nabla \mathbf{u}_{k,h})|^2, \quad T_2 := \int_{\Omega_{k,\delta}^h} |s_{k,h}^{-1} (\tilde{\mathbf{n}}_k \otimes \nabla s_{k,h} - \mathbf{n}_k \otimes \nabla s_k)|^2, \\ T_3 &:= \int_{\Omega_{k,\delta}^h} |(s_k^{-1} - s_{k,h}^{-1}) (\nabla \mathbf{u}_k - \mathbf{n}_k \otimes \nabla s_k)|^2. \end{aligned}$$

Recalling again $|s_k|, |s_{k,h}| > \delta/2$ in $\Omega_{k,\delta}^h$, as well as (6.2)–(6.3), the asserted convergence follows from

$$\begin{aligned} T_1 &\lesssim \delta^{-2} \|\nabla (\mathbf{u}_k - I_h[\mathbf{u}_k])\|_{L^2(\Omega)}^2, \\ T_2 &\lesssim \delta^{-2} (\|\nabla s_{k,h}\|_{L^4(\Omega_{k,\delta}^h)}^2 \|\tilde{\mathbf{n}}_k - \mathbf{n}_k\|_{L^4(\Omega_{k,\delta}^h)}^2 + \|\mathbf{n}_k\|_{L^4(\Omega_{k,\delta}^h)}^2 \|\nabla (s_k - I_h[s_k])\|_{L^4(\Omega_{k,\delta}^h)}^2), \\ T_3 &\lesssim \delta^{-4} \|s_k - I_h[s_k]\|_{L^4(\Omega)}^2 \|\nabla \mathbf{u}_k - \mathbf{n}_k \otimes \nabla s_k\|_{L^4(\Omega)}^2 \end{aligned}$$

and the fact that the right-hand sides of these estimates converge to 0 for a suitably faster convergence of h to 0 relative to that of δ .

Step 5: Proof of $\lim_{h,\delta \rightarrow 0} \int_{\Omega} s_{k,h}^2 |\nabla \mathbf{n}_{k,h}|^2 = \int_{\Omega \setminus \Sigma_k} s_k^2 |\nabla \mathbf{n}_k|^2$. Let us assume that $h, \delta \rightarrow 0$ in such a way (h sufficiently faster than δ) that the definitions and the convergence results established in Steps 2–4 are valid. Combining Steps 3–4 gives

$$(6.5) \quad \lim_{h,\delta \rightarrow 0} \int_{\Omega_{k,\delta}^h} |\nabla \mathbf{n}_k - \nabla \mathbf{n}_{k,h}|^2 = 0.$$

In order to exploit this property, we split the integral under consideration as

$$(6.6) \quad \int_{\Omega} s_{k,h}^2 |\nabla \mathbf{n}_{k,h}|^2 = \int_{\Omega_{k,\delta}^h} s_{k,h}^2 |\nabla \mathbf{n}_{k,h}|^2 + \int_{\Omega \setminus \Omega_{k,\delta}^h} s_{k,h}^2 |\nabla \mathbf{n}_{k,h}|^2.$$

The fact that $s_{k,h} \rightarrow s_k$ strongly in $L^p(\Omega)$ as $h \rightarrow 0$ for $d < p < \infty$, according to (6.3), together with $s_{k,h} \in L^\infty(\Omega)$ uniformly in h , $\nabla \mathbf{n}_k \in \mathbf{L}^\infty(\Omega \setminus \Sigma_{k,\delta})$, and (6.5), yields

$$\lim_{h,\delta \rightarrow 0} \left| \int_{\Omega_{k,\delta}^h} s_{k,h}^2 |\nabla \mathbf{n}_{k,h}|^2 - \int_{\Omega_{k,\delta}^h} s_k^2 |\nabla \mathbf{n}_k|^2 \right| = 0.$$

Since $\Omega \setminus \Sigma_{k,2\delta} \subset \Omega_{k,\delta}^h \subset \Omega \setminus \Sigma_{k,\delta}$ (recall Figure 9), we deduce

$$\lim_{h,\delta \rightarrow 0} \int_{\Omega_{k,\delta}^h} s_{k,h}^2 |\nabla \mathbf{n}_{k,h}|^2 = \int_{\Omega \setminus \Sigma_k} s_k^2 |\nabla \mathbf{n}_k|^2.$$

Now, we consider the second term on the right-hand side of (6.6). Since $\Omega \setminus \Omega_{k,\delta}^h \subset \Sigma_{k,2\delta}$ and $s_{k,h} \nabla \mathbf{n}_{k,h} = \nabla(s_{k,h} \mathbf{n}_{k,h}) - \mathbf{n}_{k,h} \otimes \nabla s_{k,h}$, using $\mathbf{u}_{k,h} = I_h[s_{k,h} \mathbf{n}_{k,h}]$, we see that

$$\begin{aligned} & \int_{\Omega \setminus \Omega_{k,\delta}^h} s_{k,h}^2 |\nabla \mathbf{n}_{k,h}|^2 \\ & \lesssim \int_{\Sigma_{k,2\delta}} |\nabla(s_{k,h} \mathbf{n}_{k,h})|^2 + \int_{\Sigma_{k,2\delta}} |\mathbf{n}_{k,h} \otimes \nabla s_{k,h}|^2 \\ & \lesssim \int_{\Sigma_{k,2\delta}} |\nabla(s_{k,h} \mathbf{n}_{k,h}) - \nabla I_h[s_{k,h} \mathbf{n}_{k,h}]|^2 + \int_{\Sigma_{k,2\delta}} |\nabla \mathbf{u}_{k,h}|^2 + \int_{\Sigma_{k,2\delta}} |\nabla s_{k,h}|^2. \end{aligned}$$

Combining an interpolation estimate with the fact that $s_{k,h}$ and $\mathbf{n}_{k,h}$ are piecewise affine, and exploiting an inverse estimate to bound $\|\nabla \mathbf{n}_{k,h}\|_{\mathbf{L}^\infty(K)}$ in terms of $\|\mathbf{n}_{k,h}\|_{\mathbf{L}^\infty(K)} = 1$, yields

$$\begin{aligned} & \int_{\Sigma_{k,2\delta}} |\nabla(s_{k,h} \mathbf{n}_{k,h}) - \nabla I_h[s_{k,h} \mathbf{n}_{k,h}]|^2 \lesssim \sum_{\substack{K \in \mathcal{T}_h \\ K \cap \Sigma_{k,2\delta} \neq \emptyset}} h_K^2 \|D^2(s_{k,h} \mathbf{n}_{k,h})\|_{\mathbf{L}^2(K)}^2 \\ & \lesssim \sum_{\substack{K \in \mathcal{T}_h \\ K \cap \Sigma_{k,2\delta} \neq \emptyset}} h_K^2 \|\nabla s_{k,h}\|_{\mathbf{L}^2(K)}^2 \|\nabla \mathbf{n}_{k,h}\|_{\mathbf{L}^\infty(K)}^2 \lesssim \sum_{\substack{K \in \mathcal{T}_h \\ K \cap \Sigma_{k,2\delta} \neq \emptyset}} \|\nabla s_{k,h}\|_{\mathbf{L}^2(K)}^2. \end{aligned}$$

Using the $W^{1,p}$ -stability (6.2) of the nodal interpolant with $p > d$ for all elements $K \in \mathcal{T}_h$ with $K \cap \Sigma_{k,2\delta} \neq \emptyset$, we end up with, as $h, \delta \rightarrow 0$,

$$\begin{aligned} & \int_{\Omega \setminus \Omega_{k,\delta}^h} s_{k,h}^2 |\nabla \mathbf{n}_{k,h}|^2 \lesssim \|\nabla \mathbf{u}_k\|_{\mathbf{L}^p(\Sigma_{k,3\delta})}^2 + \|\nabla s_k\|_{\mathbf{L}^p(\Sigma_{k,3\delta})}^2 \\ & \rightarrow \|\nabla \mathbf{u}_k\|_{\mathbf{L}^p(\Sigma_k)}^2 + \|\nabla s_k\|_{\mathbf{L}^p(\Sigma_k)}^2 = 0 \end{aligned}$$

(cf. [18, Theorem 4.4 (iv)]), because the Lipschitz continuity of s_k implies that all elements $K \in \mathcal{T}_h$ with $K \cap \Sigma_{k,2\delta} \neq \emptyset$ are contained in $\Sigma_{k,3\delta}$ provided $h\delta^{-1}$ is sufficiently small (cf. Step 2).

Step 6: Proof of $\lim_{h,\delta \rightarrow 0} \int_{\Omega} |\mathbf{n}_{k,h} \otimes \nabla s_{k,h}|^2 = \int_{\Omega} |\nabla s_k|^2$. We split the integral as

$$\int_{\Omega} |\mathbf{n}_{k,h} \otimes \nabla s_{k,h}|^2 = \int_{\Omega_{k,\delta}^h} |\mathbf{n}_{k,h} \otimes \nabla s_{k,h}|^2 + \int_{\Omega \setminus \Omega_{k,\delta}^h} |\mathbf{n}_{k,h} \otimes \nabla s_{k,h}|^2.$$

Exploiting the identity $\mathbf{n}_{k,h} \otimes \nabla s_{k,h} - \mathbf{n}_k \otimes \nabla s_k = (\mathbf{n}_{k,h} - \mathbf{n}_k) \otimes \nabla s_k + \mathbf{n}_{k,h} \otimes (\nabla s_{k,h} - \nabla s_k)$, and using the convergence results for $s_{k,h}$ and $\mathbf{n}_{k,h}$ in $\Omega_{k,\delta}^h$ from Steps 2–4, we readily see that

$$\lim_{h,\delta \rightarrow 0} \int_{\Omega_{k,\delta}^h} |\mathbf{n}_{k,h} \otimes \nabla s_{k,h}|^2 = \int_{\Omega \setminus \Sigma_k} |\mathbf{n}_k \otimes \nabla s_k|^2 = \int_{\Omega} |\nabla s_k|^2.$$

Moreover, employing $\Omega \setminus \Omega_{k,\delta} \subset \Sigma_{k,2\delta}$ together with (6.2) implies

$$\int_{\Omega \setminus \Omega_{k,\delta}^h} |\mathbf{n}_{k,h} \otimes \nabla s_{k,h}|^2 \lesssim \|\nabla s_{k,h}\|_{L^2(\Sigma_{k,2\delta})}^2 \lesssim \|\nabla s_{k,h}\|_{L^p(\Sigma_{k,2\delta})}^2 \lesssim \|\nabla s_k\|_{L^p(\Sigma_{k,3\delta})}^2.$$

Finally, taking $h, \delta \rightarrow 0$ yields $\|\nabla s_k\|_{L^p(\Sigma_{k,3\delta})} \rightarrow \|\nabla s_k\|_{L^p(\Sigma_k)} = 0$, which leads to the desired limit.

Step 7: End of the proof. Let $\epsilon > 0$ be the arbitrary value fixed in Step 1. The triangle inequality yields that

$$\|\mathbf{n}_{k,h} - \mathbf{n}_k\|_{L^2(\Omega \setminus \Sigma)} \lesssim \|\mathbf{n}_{k,h} - \mathbf{n}_k\|_{L^2(\Omega_{k,\delta}^h)} + \|\mathbf{n}_{k,h} - \mathbf{n}_k\|_{L^2(\Sigma_{k,2\delta} \setminus \Sigma)}.$$

For the first term, the convergences in Steps 3–4 guarantee that $\|\mathbf{n}_{k,h} - \mathbf{n}_k\|_{L^2(\Omega_{k,\delta}^h)} < \epsilon$ if h, δ are chosen properly and sufficiently small. For the second term, we have

$$\|\mathbf{n}_{k,h} - \mathbf{n}_k\|_{L^2(\Sigma_{k,2\delta} \setminus \Sigma)} \leq 2|\Sigma_{k,2\delta} \setminus \Sigma|^{1/2} \rightarrow 2|\Sigma_k \setminus \Sigma|^{1/2} \quad \text{as } \delta \rightarrow 0.$$

Since $|\Sigma_k \setminus \Sigma|^{1/2} < \epsilon^{1/2}$, we deduce that $\|\mathbf{n}_{k,h} - \mathbf{n}_k\|_{L^2(\Omega \setminus \Sigma)} \lesssim \epsilon + \epsilon^{1/2}$ provided h, δ are sufficiently small.

Next, recall that $(s_{k,h}, \mathbf{u}_{k,h}) \rightarrow (s_k, \mathbf{u}_k)$ in $H^1(\Omega)^{1+d}$ as $h \rightarrow 0$ (thanks to (6.3)) as well as $E_1^h[s_{k,h}, \mathbf{n}_{k,h}] \rightarrow E_1[s_k, \mathbf{n}_k]$ as $h, \delta \rightarrow 0$ (thanks to Steps 5–6). Furthermore, since $-1/(d-1) + 1/k \leq s_k \leq 1 - 1/k$ in Ω , assumption (2.14) guarantees that $0 \leq \psi(s_{k,h}) \leq \max\{\psi(-1/(d-1) + 1/k), \psi(1 - 1/k)\}$. Hence, the dominated convergence theorem implies that

$$\lim_{h \rightarrow 0} E_2^h[s_{k,h}] = \lim_{h \rightarrow 0} \int_{\Omega} \psi(I_h[s_k]) = \int_{\Omega} \lim_{h \rightarrow 0} \psi(I_h[s_k]) = \int_{\Omega} \psi(s_k) = E_2[s_k].$$

Altogether, we can therefore find sufficiently small $\delta > 0$ and $h > 0$ such that $\|s_{k,h} - s_k\|_{H^1(\Omega)} < \epsilon$, $\|\mathbf{u}_{k,h} - \mathbf{u}_k\|_{H^1(\Omega)} < \epsilon$, $\|\mathbf{n}_{k,h} - \mathbf{n}_k\|_{L^2(\Omega \setminus \Sigma)} \lesssim \epsilon + \epsilon^{1/2}$, and $|E^h[s_{k,h}, \mathbf{n}_{k,h}] - E[s_k, \mathbf{n}_k]| < \epsilon$. Combining these inequalities with those established in Step 1, we conclude that $\|s_{k,h} - s\|_{H^1(\Omega)} < 2\epsilon$, $\|\mathbf{u}_{k,h} - \mathbf{u}\|_{H^1(\Omega)} < 2\epsilon$, $\|\mathbf{n}_{k,h} - \mathbf{n}\|_{L^2(\Omega \setminus \Sigma)} \lesssim \epsilon + \epsilon^{1/2}$, and $|E^h[s_{k,h}, \mathbf{n}_{k,h}] - E[s, \mathbf{n}]| < 2\epsilon$. Since ϵ was arbitrary, this shows that the sequence $(s_h, \mathbf{n}_h, \mathbf{u}_h) := (s_{k,h}, \mathbf{n}_{k,h}, \mathbf{u}_{k,h}) \in \mathcal{A}_{h,\epsilon}(g_h, \mathbf{r}_h)$ satisfies the desired convergence toward $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}(g, \mathbf{r})$ as well as $\lim_{h \rightarrow 0} E^h[s_h, \mathbf{n}_h] = E[s, \mathbf{n}]$. This implies the lim-sup inequality (3.5) and concludes the proof. \square

6.3. Lim-inf inequality: Stability. To show the lim-inf inequality, we first prove that admissible discrete pairs (s_h, \mathbf{n}_h) with uniformly bounded energy are uniformly bounded in H^1 . In contrast to [23], we do not need to assume that \mathcal{T}_h is weakly acute.

LEMMA 6.4 (coercivity). *Let $\{(s_h, \mathbf{n}_h, \mathbf{u}_h)\} \subset V_h \times \mathbf{V}_h \times \mathbf{V}_h$ satisfy $\mathbf{u}_h = I_h[s_h \mathbf{n}_h]$ and $|\mathbf{n}_h(z)| \geq 1$ for all $z \in \mathcal{N}_h$. Then, there exists a constant $C > 0$ depending only on the shape-regularity of $\{\mathcal{T}_h\}$ and κ such that*

$$C \max \left\{ \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)}^2, \|\nabla(s_h \mathbf{n}_h)\|_{\mathbf{L}^2(\Omega)}^2, \|\nabla s_h\|_{\mathbf{L}^2(\Omega)}^2 \right\} \leq E_1^h[s_h, \mathbf{n}_h].$$

Proof. Since $\|\mathbf{n}_h\|_{\mathbf{L}^\infty(K)} \geq 1$ for all $K \in \mathcal{T}_h$ and ∇s_h is piecewise constant, it holds that

$$\begin{aligned} \|\nabla s_h\|_{\mathbf{L}^2(\Omega)}^2 &\leq \sum_{K \in \mathcal{T}_h} \|\mathbf{n}_h\|_{\mathbf{L}^\infty(K)}^2 \|\nabla s_h\|_{\mathbf{L}^2(K)}^2 = \sum_{K \in \mathcal{T}_h} |K| \|\mathbf{n}_h\|_{\mathbf{L}^\infty(K)}^2 |\nabla s_h|_K|^2 \\ &\lesssim \sum_{K \in \mathcal{T}_h} |\nabla s_h|_K|^2 \|\mathbf{n}_h\|_{\mathbf{L}^2(K)}^2 = \|\mathbf{n}_h \otimes \nabla s_h\|_{\mathbf{L}^2(\Omega)}^2 \leq \frac{2}{\kappa} E_1^h[s_h, \mathbf{n}_h], \end{aligned}$$

where the hidden multiplicative constant depends only on the shape-regularity of $\{\mathcal{T}_h\}$. Let $\tilde{\mathbf{u}}_h = s_h \mathbf{n}_h$ and use the Hölder inequality in conjunction with (6.2) for $p > d$ and an inverse estimate to obtain

$$\|\nabla I_h[\tilde{\mathbf{u}}_h]\|_{\mathbf{L}^2(K)} \lesssim |K|^{\frac{p-2}{2p}} \|\nabla I_h[\tilde{\mathbf{u}}_h]\|_{\mathbf{L}^p(K)} \lesssim |K|^{\frac{p-2}{2p}} \|\nabla \tilde{\mathbf{u}}_h\|_{\mathbf{L}^p(K)} \lesssim \|\nabla \tilde{\mathbf{u}}_h\|_{\mathbf{L}^2(K)}$$

for all $K \in \mathcal{T}_h$. Consequently, for $\mathbf{u}_h = I_h[\tilde{\mathbf{u}}_h]$ we deduce

$$\|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)}^2 \lesssim \|\nabla \tilde{\mathbf{u}}_h\|_{\mathbf{L}^2(\Omega)}^2 \lesssim \|\mathbf{n}_h \otimes \nabla s_h\|_{\mathbf{L}^2(\Omega)}^2 + \|s_h \nabla \mathbf{n}_h\|_{\mathbf{L}^2(\Omega)}^2 \lesssim E_1^h[s_h, \mathbf{n}_h].$$

This completes the proof. \square

We are now ready to extract convergent subsequences and characterize their limits.

LEMMA 6.5 (characterization of limits). *Let $\{(s_h, \mathbf{n}_h, \mathbf{u}_h)\} \subset \mathcal{A}_{h,\varepsilon}(g_h, \mathbf{r}_h)$ be a sequence such that $E_1^h[s_h, \mathbf{n}_h] \leq C$ and $\|\mathbf{n}_h\|_{\mathbf{L}^\infty(\Omega)} \leq C$, where $C > 0$ is a constant independent of h . Then, there exist a triple $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}(g, \mathbf{r})$ and a subsequence (not relabeled) of $\{(s_h, \mathbf{n}_h, \mathbf{u}_h)\}$ satisfying the following properties:*

- as $h \rightarrow 0$, $(s_h, \mathbf{u}_h, s_h \mathbf{n}_h)$ converges toward $(s, \mathbf{u}, \mathbf{u})$ weakly in $H^1(\Omega) \times \mathbf{H}^1(\Omega) \times \mathbf{H}^1(\Omega)$, strongly in $L^2(\Omega) \times \mathbf{L}^2(\Omega) \times \mathbf{L}^2(\Omega)$, and pointwise a.e. in Ω ;
- \mathbf{n}_h converges toward \mathbf{n} strongly in $\mathbf{L}^2(\Omega \setminus \Sigma)$ and pointwise a.e. in $\Omega \setminus \Sigma$ as $h \rightarrow 0$ and $\varepsilon \rightarrow 0$;
- \mathbf{n} is L^2 -differentiable a.e. in $\Omega \setminus \Sigma$ and the orthogonal decomposition $|\nabla \mathbf{u}|^2 = |\nabla s|^2 + s^2 |\nabla \mathbf{n}|^2$ is valid a.e. in $\Omega \setminus \Sigma$,

where $\Sigma \subset \Omega$ is given by (2.2).

Proof. For the sake of clarity, we divide the proof into three steps.

Step 1: Convergence of $\{s_h\}$, $\{\mathbf{u}_h\}$, and $\{s_h \mathbf{n}_h\}$. Since the energy $E_1^h[s_h, \mathbf{n}_h]$ is uniformly bounded, Lemma 6.4 (coercivity) gives uniform bounds in $H^1(\Omega) \times \mathbf{H}^1(\Omega) \times \mathbf{H}^1(\Omega)$ for the sequence $\{(s_h, \mathbf{u}_h, s_h \mathbf{n}_h)\}$. With successive extractions of subsequences (not relabeled), one can show that there exists a limit $(s, \mathbf{u}, \tilde{\mathbf{u}}) \in H^1(\Omega) \times \mathbf{H}^1(\Omega) \times \mathbf{H}^1(\Omega)$ such that $(s_h, \mathbf{u}_h, s_h \mathbf{n}_h)$ converges to $(s, \mathbf{u}, \tilde{\mathbf{u}})$ weakly in $H^1(\Omega) \times \mathbf{H}^1(\Omega) \times \mathbf{H}^1(\Omega)$, strongly in $L^2(\Omega) \times \mathbf{L}^2(\Omega) \times \mathbf{L}^2(\Omega)$, and pointwise a.e. in

Ω . Moreover, weak H^1 -convergence guarantees attainment of traces, namely $s = g$ and $\mathbf{u} = \tilde{\mathbf{u}} = \mathbf{r}$ on Γ_D . To see this, note that $g_h = I_h[g] \rightarrow g$ in $W^{1,p}(\Omega)$ for $p > d$, according to (6.3), and so in $H^1(\Omega)$. Therefore $s_h - g_h \in H_0^1(\Omega)$ satisfies

$$s_h - g_h \rightharpoonup s - g \in H_0^1(\Omega),$$

because $H_0^1(\Omega)$ is closed under weak convergence. Hence $s = g$ on Γ_D in the sense of traces, as asserted. Dealing with \mathbf{u}_h and $\tilde{\mathbf{u}}_h$ is identical. Since $\mathbf{u}_h = I_h[s_h \mathbf{n}_h]$, interpolation and inverse estimates yield

$$\begin{aligned} \|\mathbf{u}_h - s_h \mathbf{n}_h\|_{\mathbf{L}^2(\Omega)}^2 &\lesssim \sum_{K \in \mathcal{T}_h} h_K^4 \|D^2(s_h \mathbf{n}_h)\|_{\mathbf{L}^2(K)}^2 \\ &\lesssim \sum_{K \in \mathcal{T}_h} h_K^2 \|\nabla(s_h \mathbf{n}_h)\|_{\mathbf{L}^2(K)}^2 \lesssim h^2 E_1^h[s_h, \mathbf{n}_h] \leq Ch^2. \end{aligned}$$

This shows that $s_h \mathbf{n}_h$ and \mathbf{u}_h converge strongly in $\mathbf{L}^2(\Omega)$ toward the same limit, i.e., $\tilde{\mathbf{u}} = \mathbf{u}$. Moreover, $s_h \mathbf{n}_h$ converges to \mathbf{u} weakly in $\mathbf{H}^1(\Omega)$ and pointwise a.e. in Ω .

Step 2: $|s| = |\mathbf{u}|$ a.e. in Ω . The triangle inequality yields

$$\begin{aligned} \||\mathbf{u}_h|^2 - |s_h|^2\|_{L^1(\Omega)} &\leq \||\mathbf{u}_h|^2 - I_h[|\mathbf{u}_h|^2]\|_{L^1(\Omega)} \\ &\quad + \|I_h[|\mathbf{u}_h|^2] - |s_h|^2\|_{L^1(\Omega)} + \||s_h|^2 - I_h[|s_h|^2]\|_{L^1(\Omega)}. \end{aligned}$$

For the first and third terms on the right-hand side, standard interpolation estimates yield

$$\||s_h|^2 - I_h[|s_h|^2]\|_{L^1(\Omega)} \lesssim h^2 \|\nabla s_h\|_{L^2(\Omega)}^2, \quad \||\mathbf{u}_h|^2 - I_h[|\mathbf{u}_h|^2]\|_{L^1(\Omega)} \lesssim h^2 \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)}^2.$$

On the other hand, since $\{s_h\}$ is uniformly bounded in $L^\infty(\Omega)$, we infer that

$$\begin{aligned} \|I_h[|\mathbf{u}_h|^2] - |s_h|^2\|_{L^1(\Omega)} &= \|I_h[|s_h|^2(|\mathbf{n}_h|^2 - 1)]\|_{L^1(\Omega)} \\ &\leq \|s_h\|_{L^\infty(\Omega)}^2 \|I_h[|\mathbf{n}_h|^2 - 1]\|_{L^1(\Omega)} \leq \varepsilon \|s_h\|_{L^\infty(\Omega)}^2 \rightarrow 0 \end{aligned}$$

as $\varepsilon \rightarrow 0$. As $|s_h| \rightarrow |s|$ and $|\mathbf{u}_h| \rightarrow |\mathbf{u}|$ a.e. in Ω , we conclude that $|s| = |\mathbf{u}|$ a.e. in Ω .

Step 3: Convergence of $\{\mathbf{n}_h\}$. We now define $\mathbf{n} : \Omega \rightarrow \mathbb{R}^3$ as $\mathbf{n} := s^{-1} \mathbf{u}$ in $\Omega \setminus \Sigma$ and as an arbitrary unit vector in Σ . Step 2 implies, by construction, that $|\mathbf{n}| = 1$ a.e. in Ω . This shows that $(s, \mathbf{n}, \mathbf{u})$ satisfies the structural condition (2.6), i.e., $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}$.

We now observe that $s(x) \neq 0$ for a.e. $x \in \Omega \setminus \Sigma$ by definition of Σ . Since $s_h(x) \rightarrow s(x)$ as $h \rightarrow 0$, if h is sufficiently small (depending on x), then $s_h(x) \neq 0$ is valid. Consequently,

$$\mathbf{n}_h(x) = \frac{s_h(x) \mathbf{n}_h(x)}{s_h(x)} \rightarrow \frac{\mathbf{u}(x)}{s(x)} = \mathbf{n}(x),$$

i.e., $\mathbf{n}_h \rightarrow \mathbf{n}$ pointwise a.e. in $\Omega \setminus \Sigma$. Since $\{\mathbf{n}_h\}$ is uniformly bounded in $\mathbf{L}^\infty(\Omega)$, the Lebesgue dominated convergence theorem yields $\mathbf{n}_h \rightarrow \mathbf{n}$ strongly in $\mathbf{L}^2(\Omega \setminus \Sigma)$.

Finally, the L^2 -differentiability of \mathbf{n} and the orthogonal decomposition of $\nabla \mathbf{u}$, both valid a.e. in $\Omega \setminus \Sigma$, follow from Proposition 2.1 (orthogonal decomposition). This concludes the proof. \square

We are now in position to prove the lim-inf inequality.

Proof of Theorem 3.1(ii). The sequence $\{(s_h, \mathbf{n}_h, \mathbf{u}_h)\} \subset \mathcal{A}_{h,\varepsilon}(g_h, \mathbf{r}_h)$ satisfies the assumptions of Lemma 6.5 (characterization of limits). Hence, we can apply it to obtain subsequences (not relabeled) converging to the respective limits $(s, \mathbf{n}, \mathbf{u}) \in \mathcal{A}(g, \mathbf{r})$. Moreover, since also the sequences $\{\mathbf{n}_h \otimes \nabla s_h\}$ and $\{s_h \nabla \mathbf{n}_h\}$ are uniformly bounded in $\mathbf{L}^2(\Omega)$, there exist subsequences (not relabeled) and functions \mathbf{M}, \mathbf{N} in $\mathbf{L}^2(\Omega)$ such that $\mathbf{n}_h \otimes \nabla s_h \rightharpoonup \mathbf{M}$ and $s_h \nabla \mathbf{n}_h \rightharpoonup \mathbf{N}$ weakly in $\mathbf{L}^2(\Omega)$. Combining the equality $s_h \nabla \mathbf{n}_h = \nabla(s_h \mathbf{n}_h) - \mathbf{n}_h \otimes \nabla s_h$, which is valid in every element of \mathcal{T}_h , with $s_h \mathbf{n}_h \rightharpoonup \mathbf{u}$ weakly in $\mathbf{H}^1(\Omega)$, helps identify the limits $\mathbf{N} = \nabla \mathbf{u} - \mathbf{M}$.

Let $\Phi \in C_c^\infty(\Omega \setminus \Sigma)$ be an arbitrary $d \times d$ tensor field. We can thus write

$$\langle \mathbf{n}_h \otimes \nabla s_h - \mathbf{n} \otimes \nabla s, \Phi \rangle_{\Omega \setminus \Sigma} = \langle (\mathbf{n}_h - \mathbf{n}) \otimes \nabla s_h, \Phi \rangle_{\Omega \setminus \Sigma} + \langle \mathbf{n} \otimes (\nabla s_h - \nabla s), \Phi \rangle_{\Omega \setminus \Sigma}.$$

We note that $\mathbf{n}_h \rightarrow \mathbf{n}$ strongly in $L^2(\Omega \setminus \Sigma)$ implies

$$\langle (\mathbf{n}_h - \mathbf{n}) \otimes \nabla s_h, \Phi \rangle_{\Omega \setminus \Sigma} \leq \|\mathbf{n}_h - \mathbf{n}\|_{\mathbf{L}^2(\Omega \setminus \Sigma)} \|\nabla s_h\|_{\mathbf{L}^2(\Omega)} \|\Phi\|_{\mathbf{L}^\infty(\Omega \setminus \Sigma)} \rightarrow 0,$$

whereas $s_h \rightarrow s$ weakly in $H^1(\Omega)$ yields $\langle \mathbf{n} \otimes (\nabla s_h - \nabla s), \Phi \rangle_{\Omega \setminus \Sigma} \rightarrow 0$. Hence, we infer that $\langle \mathbf{n}_h \otimes \nabla s_h - \mathbf{n} \otimes \nabla s, \Phi \rangle_{\Omega \setminus \Sigma} \rightarrow 0$, whence $\mathbf{n}_h \otimes \nabla s_h \rightharpoonup \mathbf{n} \otimes \nabla s$ weakly in $\mathbf{L}^2(\Omega \setminus \Sigma)$. This in turn identifies the limit $\mathbf{M} = \mathbf{n} \otimes \nabla s$ and gives thus the identity $\mathbf{N} = \nabla \mathbf{u} - \mathbf{n} \otimes \nabla s$ a.e. in $\Omega \setminus \Sigma$. We deduce that $\nabla \mathbf{n} = \mathbf{N}/s$, where $\nabla \mathbf{n}$ is understood in the L^2 -sense according to Proposition 2.1. Exploiting the fact that norms are weakly lower semicontinuous, along with $|\mathbf{n} \otimes \nabla s|^2 = |\nabla s|^2$ a.e. in $\Omega \setminus \Sigma$, and $\nabla s = \mathbf{0}$ a.e. in Σ , it holds that

$$\begin{aligned} \liminf_{h \rightarrow 0} E_1^h[s_h, \mathbf{n}_h] &= \liminf_{h \rightarrow 0} \left\{ \frac{\kappa}{2} \|\mathbf{n}_h \otimes \nabla s_h\|_{\mathbf{L}^2(\Omega)}^2 + \frac{1}{2} \|s_h \nabla \mathbf{n}_h\|_{\mathbf{L}^2(\Omega)}^2 \right\} \\ &\geq \liminf_{h \rightarrow 0} \left\{ \frac{\kappa}{2} \|\mathbf{n}_h \otimes \nabla s_h\|_{\mathbf{L}^2(\Omega \setminus \Sigma)}^2 + \frac{1}{2} \|s_h \nabla \mathbf{n}_h\|_{\mathbf{L}^2(\Omega \setminus \Sigma)}^2 \right\} \\ &\geq \frac{\kappa}{2} \|\mathbf{n} \otimes \nabla s\|_{\mathbf{L}^2(\Omega \setminus \Sigma)}^2 + \frac{1}{2} \|s \nabla \mathbf{n}\|_{\mathbf{L}^2(\Omega \setminus \Sigma)}^2 = E_1[s, \mathbf{n}]. \end{aligned}$$

Since $s_h \rightarrow s$ a.e. in Ω and ψ is continuous, $\psi(s_h) \rightarrow \psi(s)$ a.e. in Ω . The Fatou lemma yields

$$E_2[s] = \int_{\Omega} \psi(s) = \int_{\Omega} \lim_{h \rightarrow 0} \psi(s_h) \leq \liminf_{h \rightarrow 0} \int_{\Omega} \psi(s_h) = \liminf_{h \rightarrow 0} E_2^h[s].$$

Altogether, we thus obtain the lim-inf inequality (3.6). This finishes the proof. \square

6.4. Properties of the numerical scheme. To start with, we prove the well-posedness and stability of Algorithm 4.1.

Proof of Proposition 4.1. Let $i \in \mathbb{N}_0$ and $\ell \in \mathbb{N}_0$. For fixed $s_h^i \in V_h$ (resp., $\mathbf{n}_h^{i+1} \in \mathbf{V}_h$), the left-hand side of (4.1) (resp., of (4.3)) is a coercive and continuous bilinear form on $\mathbf{V}_{h,D}$ (resp., on $V_{h,D}$). Therefore, the variational problem admits a unique solution $\mathbf{t}_h^{i,\ell} \in \mathcal{K}_h[\mathbf{n}_h^{i,\ell}]$ (resp., $s_h^{i+1} \in V_h$) by the Lax–Milgram theorem. This shows parts (i) and (iii) of Proposition 4.1.

Choosing the test function $\phi_h = \tau_n \mathbf{t}_h^{i,\ell} = \mathbf{n}_h^{i,\ell+1} - \mathbf{n}_h^{i,\ell} \in \mathcal{K}_h[\mathbf{n}_h^{i,\ell}]$ in (4.1) yields

$$\begin{aligned} \tau_n \|\mathbf{t}_h^{i,\ell}\|_*^2 + \kappa \langle \mathbf{n}_h^{i,\ell+1} \otimes \nabla s_h^i, (\mathbf{n}_h^{i,\ell+1} - \mathbf{n}_h^{i,\ell}) \otimes \nabla s_h^i \rangle_{\Omega} \\ + \langle s_h^i \nabla \mathbf{n}_h^{i,\ell+1}, s_h^i \nabla (\mathbf{n}_h^{i,\ell+1} - \mathbf{n}_h^{i,\ell}) \rangle_{\Omega} = 0. \end{aligned}$$

Using the identity $2a(a-b) = a^2 - b^2 + (a-b)^2$, valid for all $a, b \in \mathbb{R}$, we obtain

$$(6.7) \quad E_1^h[s_h^i, \mathbf{n}_h^{i,\ell+1}] - E_1^h[s_h^i, \mathbf{n}_h^{i,\ell}] + \tau_n \|\mathbf{t}_h^{i,\ell}\|_*^2 + \tau_n^2 E_1^h[s_h^i, \mathbf{t}_h^{i,\ell}] = 0.$$

In particular, $E_1^h[s_h^i, \mathbf{n}_h^{i,\ell+1}] \leq E_1^h[s_h^i, \mathbf{n}_h^{i,\ell}]$ is valid. Since $E_1^h[s_h^i, \mathbf{n}_h^{i,\ell}] \geq 0$ for all $i \in \mathbb{N}_0$, the sequence $\{E_1^h[s_h^i, \mathbf{n}_h^{i,\ell}]\}_{\ell \in \mathbb{N}_0}$ is convergent (as it is monotonically decreasing and bounded from below). In particular, it is a Cauchy sequence, which entails that the stopping criterion (4.2) is met in a finite number of iterations. This shows part (ii) of the proposition.

Let $\ell_i \in \mathbb{N}_0$ be the smallest integer for which the stopping criterion (4.2) is satisfied. Recall that $\mathbf{n}_h^{i+1} = \mathbf{n}_h^{i,\ell_i+1}$ and $\mathbf{n}_h^i = \mathbf{n}_h^{i,0}$. Summation of (6.7) over $\ell = 0, \dots, \ell_i$ yields

$$(6.8) \quad E_1^h[s_h^i, \mathbf{n}_h^{i+1}] - E_1^h[s_h^i, \mathbf{n}_h^i] + \tau_n \sum_{\ell=0}^{\ell_i} \|\mathbf{t}_h^{i,\ell}\|_*^2 + \tau_n^2 \sum_{\ell=0}^{\ell_i} E_1^h[s_h^i, \mathbf{t}_h^{i,\ell}] = 0.$$

Choosing the test function $w_h = \tau_s d_t s_h^{i+1} = s_h^{i+1} - s_h^i \in V_{h,D}$ in (4.3) and performing the same algebraic computation as above, we arrive at

$$\begin{aligned} E_1^h[s_h^{i+1}, \mathbf{n}_h^{i+1}] - E_1^h[s_h^i, \mathbf{n}_h^{i+1}] + \tau_s \|d_t s_h^{i+1}\|_{L^2(\Omega)}^2 + \tau_s^2 E_1^h[d_t s_h^{i+1}, \mathbf{n}_h^{i+1}] \\ + \langle \psi'_c(s_h^{i+1}) - \psi'_e(s_h^i), s_h^{i+1} - s_h^i \rangle_\Omega = 0. \end{aligned}$$

Applying [23, Lemma 4.1], which yields the inequality

$$E_2^h[s_h^{i+1}] - E_2^h[s_h^i] \leq \langle \psi'_c(s_h^{i+1}) - \psi'_e(s_h^i), s_h^{i+1} - s_h^i \rangle_\Omega,$$

we obtain

$$\begin{aligned} E_1^h[s_h^{i+1}, \mathbf{n}_h^{i+1}] - E_1^h[s_h^i, \mathbf{n}_h^{i+1}] + \tau_s \|d_t s_h^{i+1}\|_{L^2(\Omega)}^2 + \tau_s^2 E_1^h[d_t s_h^{i+1}, \mathbf{n}_h^{i+1}] \\ + E_2^h[s_h^{i+1}] - E_2^h[s_h^i] \leq 0. \end{aligned}$$

Adding the latter with (6.8), and exploiting cancellation of $E_1^h[s_h^i, \mathbf{n}_h^{i+1}]$, we deduce

$$(6.9) \quad \begin{aligned} E^h[s_h^{i+1}, \mathbf{n}_h^{i+1}] - E^h[s_h^i, \mathbf{n}_h^i] &\leq -\tau_n \sum_{\ell=0}^{\ell_i} \|\mathbf{t}_h^{i,\ell}\|_*^2 - \tau_n^2 \sum_{\ell=0}^{\ell_i} E_1^h[s_h^i, \mathbf{t}_h^{i,\ell}] \\ &\quad - \tau_s \|d_t s_h^{i+1}\|_{L^2(\Omega)}^2 - \tau_s^2 E_1^h[d_t s_h^{i+1}, \mathbf{n}_h^{i+1}] \leq 0. \end{aligned}$$

This shows (4.6) and concludes the proof. \square

We recall that Algorithm 4.1 does not enforce the unit-length constraint of the director field \mathbf{n}_h^j . We finish this paper with a proof that violation of such constraint is controlled by τ_n and that $\|\mathbf{n}_h^j\|_{L^\infty(\Omega)}$ is uniformly bounded provided the parameters h and τ_n are suitably chosen.

Proof of Proposition 4.3. Let $j \geq 1$. Summation of (6.9) over $i = 0, \dots, j-1$ yields

$$(6.10) \quad E^h[s_h^j, \mathbf{n}_h^j] + \tau_n \sum_{i=0}^{j-1} \sum_{\ell=0}^{\ell_i} \|\mathbf{t}_h^{i,\ell}\|_*^2 \leq E^h[s_h^0, \mathbf{n}_h^0].$$

Moreover, the tangential update $\mathbf{t}_h^{i,\ell}(z)$ is perpendicular to $\mathbf{n}_h^{i,\ell}(z)$ for all $z \in \mathcal{N}_h$, whence $\mathbf{n}_h^{i,\ell+1}(z) = \mathbf{n}_h^{i,\ell}(z) + \tau_{\mathbf{n}} \mathbf{t}_h^{i,\ell}(z)$ satisfies $|\mathbf{n}_h^{i,\ell+1}(z)|^2 = |\mathbf{n}_h^{i,\ell}(z)|^2 + \tau_{\mathbf{n}}^2 |\mathbf{t}_h^{i,\ell}(z)|^2$. Iterating in ℓ and i gives

$$|\mathbf{n}_h^j(z)|^2 = |\mathbf{n}_h^0(z)|^2 + \tau_{\mathbf{n}}^2 \sum_{i=0}^{j-1} \sum_{\ell=0}^{\ell_i} |\mathbf{t}_h^{i,\ell}(z)|^2 = 1 + \tau_{\mathbf{n}}^2 \sum_{i=0}^{j-1} \sum_{\ell=0}^{\ell_i} |\mathbf{t}_h^{i,\ell}(z)|^2 \geq 1.$$

Then, using the equivalence of the L^p -norm of a discrete function with the weighted ℓ^p -norm of the vector collecting its nodal values (see, e.g., [8, Lemma 3.4]), for h_z being the diameter of the nodal patch associated with $z \in \mathcal{N}_h$, we see that

$$\begin{aligned} \|I_h[|\mathbf{n}_h^j|^2] - 1\|_{L^1(\Omega)} &\lesssim \sum_{z \in \mathcal{N}_h} h_z^d (|\mathbf{n}_h^j(z)|^2 - 1) \leq \tau_{\mathbf{n}}^2 \sum_{z \in \mathcal{N}_h} h_z^d \sum_{i=0}^{j-1} \sum_{\ell=0}^{\ell_i} |\mathbf{t}_h^{i,\ell}(z)|^2 \\ &\lesssim \tau_{\mathbf{n}}^2 \sum_{i=0}^{j-1} \sum_{\ell=0}^{\ell_i} \|\mathbf{t}_h^{i,\ell}\|_{L^2(\Omega)}^2. \end{aligned}$$

Combining (4.8) with (6.10) leads to

$$\|I_h[|\mathbf{n}_h^j|^2] - 1\|_{L^1(\Omega)} \lesssim C_* \tau_{\mathbf{n}}^2 \sum_{i=0}^{j-1} \sum_{\ell=0}^{\ell_i} \|\mathbf{t}_h^{i,\ell}\|_*^2 \leq C_* \tau_{\mathbf{n}} E^h[s_h^0, \mathbf{n}_h^0],$$

which turns out to be (4.9).

It remains to estimate $\|\mathbf{n}_h^j\|_{L^\infty(\Omega)}$. Let us consider first the weighted H^1 -metric in (4.5). Using a global inverse estimate (see, e.g., [8, Remark 3.8]) and the Poincaré inequality, we obtain

$$\begin{aligned} \|\mathbf{n}_h^j\|_{L^\infty(\Omega)}^2 - 1 &= \max_{z \in \mathcal{N}_h} |\mathbf{n}_h^j(z)|^2 - 1 \leq \tau_{\mathbf{n}}^2 \sum_{i=0}^{j-1} \sum_{\ell=0}^{\ell_i} \max_{z \in \mathcal{N}_h} |\mathbf{t}_h^{i,\ell}(z)|^2 \\ &\lesssim \tau_{\mathbf{n}}^2 \sum_{i=0}^{j-1} \sum_{\ell=0}^{\ell_i} \|\mathbf{t}_h^{i,\ell}\|_{L^\infty(\Omega)}^2 \lesssim \tau_{\mathbf{n}}^2 h_{\min}^{2-d} |\log h_{\min}|^2 \sum_{i=0}^{j-1} \sum_{\ell=0}^{\ell_i} \|\mathbf{t}_h^{i,\ell}\|_{H^1(\Omega)}^2 \\ &\lesssim \tau_{\mathbf{n}}^2 h_{\min}^{2-d-\alpha} |\log h_{\min}|^2 \sum_{i=0}^{j-1} \sum_{\ell=0}^{\ell_i} \|h^{\alpha/2} \nabla \mathbf{t}_h^{i,\ell}\|_{L^2(\Omega)}^2 \\ &\leq \tau_{\mathbf{n}} h_{\min}^{2-d-\alpha} |\log h_{\min}|^2 E^h[s_h^0, \mathbf{n}_h^0]. \end{aligned}$$

Therefore, (4.11) is satisfied if $\tau_{\mathbf{n}} h_{\min}^{2-d-\alpha} |\log h_{\min}|^2 \leq C^*$ with C^* arbitrary. For the L^2 -metric (4.4), the result follows analogously, provided that $\tau_{\mathbf{n}} h_{\min}^{-d} \leq C^*$. \square

REFERENCES

- [1] P. J. ACKERMAN, J. VAN DE LAGEMAAT, AND I. I. SMALYUKH, *Self-assembly and electrostriction of arrays and chains of hopfion particles in chiral liquid crystals*, Nat. Commun., 6 (2015), 6012, <https://www.doi.org/10.1038/ncomms7012>.
- [2] J. AHRENS, B. GEVECI, AND C. LAW, *ParaView: An end-user tool for large data visualization*, in Visualization Handbook, C. D. Hansen and C. R. Johnson, eds., Elsevier, New York, 2005, pp. 717–731, <https://doi.org/10.1016/B978-012387582-2/50038-1>.
- [3] F. ALOUGES, *A new algorithm for computing liquid crystal stable configurations: The harmonic mapping case*, SIAM J. Numer. Anal., 34 (1997), pp. 1708–1726, <https://doi.org/10.1137/S0036142994264249>.

- [4] L. AMBROSIO, *Existence of minimal energy configurations of nematic liquid crystals with variable degree of orientation*, Manuscripta Math., 68 (1990), pp. 215–228, <https://doi.org/10.1007/BF02568761>.
- [5] T. ARAKI AND H. TANAKA, *Colloidal aggregation in a nematic liquid crystal: Topological arrest of particles by a single-stroke disclination line*, Phys. Rev. Lett., 97 (2006), 127801, <https://doi.org/10.1103/PhysRevLett.97.127801>.
- [6] J. W. BARRETT, X. FENG, AND A. PROHL, *Convergence of a fully discrete finite element method for a degenerate parabolic system modelling nematic liquid crystals with variable degree of orientation*, ESAIM Math. Model. Numer. Anal., 40 (2006), pp. 175–199, <https://doi.org/10.1051/m2an:2006005>.
- [7] S. BARTELS, *Numerical analysis of a finite element scheme for the approximation of harmonic maps into surfaces*, Math. Comp., 79 (2010), pp. 1263–1301, <https://doi.org/10.1090/S0025-5718-09-02300-X>.
- [8] S. BARTELS, *Numerical Methods for Nonlinear Partial Differential Equations*, Springer Ser. Comput. Math. 47, Springer, New York, 2015, <https://doi.org/10.1007/978-3-319-13797-1>.
- [9] S. BARTELS, *Projection-free approximation of geometrically constrained partial differential equations*, Math. Comp., 85 (2016), pp. 1033–1049, <https://doi.org/10.1090/mcom/3008>.
- [10] C. BLANC, *Colloidal crystal ordering in a liquid crystal*, Science, 352 (2016), pp. 40–41, <https://doi.org/10.1126/science.aaf4260>.
- [11] J. P. BORTHAGARAY, R. H. NOCHETTO, AND S. W. WALKER, *A structure-preserving FEM for the uniaxially constrained Q-tensor model of nematic liquid crystals*, Numer. Math., 145 (2020), pp. 837–881, <https://doi.org/10.1007/s00211-020-01133-z>.
- [12] J. P. BORTHAGARAY AND S. W. WALKER, *Chapter 5 – The Q-tensor model with uniaxial constraint*, in Geometric Partial Differential Equations, Part II, A. Bonito and R. H. Nochetto, eds., Handb. Numer. Anal. 22, Elsevier, New York, 2021, pp. 313–382, <https://doi.org/10.1016/bs.hna.2020.09.001>.
- [13] A. BRAIDES, *Γ -Convergence for Beginners*, Oxford Lecture Ser. Math. Appl. 22, Oxford University Press, Oxford, UK, 2002, <https://doi.org/10.1093/acprof:oso/9780198507840.001.0001>.
- [14] H. BREZIS, J.-M. CORON, AND E. H. LIEB, *Harmonic maps with defects*, Comm. Math. Phys., 107 (1986), pp. 649–705.
- [15] S. CARTER, A. ROTEM, AND S. W. WALKER, *A domain decomposition approach to accelerate simulations of structure preserving nematic liquid crystal models*, J. Non-Newton. Fluid Mech., 283 (2020), 104335, <https://doi.org/10.1016/j.jnnfm.2020.104335>.
- [16] P.-G. DE GENNES AND J. PROST, *The Physics of Liquid Crystals*, 2nd ed., Internat. Ser. Monogr. Phys., Oxford University Press, Oxford, UK, 1993.
- [17] J. L. ERICKSEN, *Liquid crystals with variable degree of orientation*, Arch. Ration. Mech. Anal., 113 (1991), pp. 97–120, <https://doi.org/10.1007/BF00380413>.
- [18] L. C. EVANS AND R. F. GARIEPY, *Measure Theory and Fine Properties of Functions*, Textb. Math., CRC Press, Boca Raton, FL, 2015, <https://doi.org/10.1201/b18333>.
- [19] Y. GU AND N. L. ABBOTT, *Observation of Saturn-ring defects around solid microspheres in nematic liquid crystals*, Phys. Rev. Lett., 85 (2000), 4719, <https://doi.org/10.1103/PhysRevLett.85.4719>.
- [20] J. KRAUS, C.-M. PFEILER, D. PRAETORIUS, M. RUGGERI, AND B. STIFTNER, *Iterative solution and preconditioning for the tangent plane scheme in computational micromagnetics*, J. Comput. Phys., 398 (2019), 108866, <https://doi.org/10.1016/j.jcp.2019.108866>.
- [21] F.-H. LIN, *On nematic liquid crystals with variable degree of orientation*, Comm. Pure Appl. Math., 44 (1991), pp. 453–468, <https://doi.org/10.1002/cpa.3160440404>.
- [22] S. Y. LIN AND M. LUSKIN, *Relaxation methods for liquid crystal problems*, SIAM J. Numer. Anal., 26 (1989), pp. 1310–1324, <https://doi.org/10.1137/0726076>.
- [23] R. H. NOCHETTO, S. W. WALKER, AND W. ZHANG, *A finite element method for nematic liquid crystals with variable degree of orientation*, SIAM J. Numer. Anal., 55 (2017), pp. 1357–1386, <https://doi.org/10.1137/15M103844X>.
- [24] R. H. NOCHETTO, S. W. WALKER, AND W. ZHANG, *The Ericksen model of liquid crystals with colloidal and electric effects*, J. Comput. Phys., 352 (2018), pp. 568–601, <https://doi.org/10.1016/j.jcp.2017.09.035>.
- [25] A. RAMAGE AND E. C. GARTLAND, JR., *A preconditioned nullspace method for liquid crystal director modeling*, SIAM J. Sci. Comput., 35 (2013), pp. B226–B247, <https://doi.org/10.1137/120870219>.
- [26] J. SCHÖBERL, *Netgen/NGSolve*, 2021, <https://ngsolve.org>.
- [27] R. SCHOEN AND K. UHLENBECK, *A regularity theory for harmonic maps*, J. Differential Geom., 17 (1982), pp. 307–335, <https://www.doi.org/10.4310/jdg/1214436923>.
- [28] H. STARK, *Director field configurations around a spherical particle in a nematic liquid crystal*, Eur. Phys. J. B, 10 (1999), pp. 311–321, <https://doi.org/10.1007/s100510050860>.

- [29] E. G. VIRGA, *Variational Theories for Liquid Crystals*, Appl. Math. Math. Comput. 8, Chapman & Hall, London, 1994, <https://doi.org/10.1201/9780203734421>.
- [30] S. W. WALKER, *A finite element method for the generalized Ericksen model of nematic liquid crystals*, ESAIM Math. Model. Numer. Anal., 54 (2020), pp. 1181–1220, <https://doi.org/10.1051/m2an/2019092>.
- [31] S. M. WISE, C. WANG, AND J. S. LOWENGRUB, *An energy-stable and convergent finite-difference scheme for the phase field crystal equation*, SIAM J. Numer. Anal., 47 (2009), pp. 2269–2288, <https://doi.org/10.1137/080738143>.