PERFECT MATCHINGS IN THE SEMI-RANDOM GRAPH PROCESS

PU GAO, CALUM MACRURY, AND PAWEŁ PRAŁAT

ABSTRACT. The semi-random graph process is a single player game in which the player is initially presented an empty graph on n vertices. In each round, a vertex u is presented to the player independently and uniformly at random. The player then adaptively selects a vertex v, and adds the edge uv to the graph. For a fixed monotone graph property, the objective of the player is to force the graph to satisfy this property with high probability in as few rounds as possible.

We focus on the problem of constructing a perfect matching in as few rounds as possible. In particular, we present an adaptive strategy for the player which achieves a perfect matching in βn rounds, where the value of $\beta < 1.206$ is derived from a solution to some system of differential equations. This improves upon the previously best known upper bound of $(1+2/e+o(1))\,n < 1.736\,n$ rounds. We also improve the previously best lower bound of $(\ln 2+o(1))\,n > 0.693\,n$ and show that the player cannot achieve the desired property in less than αn rounds, where the value of $\alpha > 0.932$ is derived from a solution to another system of differential equations. As a result, the gap between the upper and lower bounds is decreased roughly four times.

1. Introduction and Main Results

1.1. **Definitions.** In this paper, we consider the **semi-random process** suggested by Peleg Michaeli and studied recently in [3, 2, 5] that can be viewed as a "one player game". The process starts from G_0 , the empty graph on the vertex set $[n] := \{1, \ldots, n\}$ where $n \geq 1$. In each step t, a vertex u_t is chosen uniformly at random from [n]. Then, the player (who is aware of graph G_t and vertex u_t) must select a vertex v_t and add the edge $u_t v_t$ to G_t to form G_{t+1} . The goal of the player is to build a (multi)graph satisfying a given property \mathcal{P} as quickly as possible. It is convenient to refer to u_t as a **square**, and v_t as a **circle** so every edge in G_t joins a square with a circle. We say that vertex $j \in [n]$ is **covered** by the square u_t arriving at round t, provided $u_t = j$. The analogous definition extends to the circle v_t . Equivalently, we may view G_t as a directed graph where each arc directs from u_t to v_t . For this paper, it is easier to consider squares and circles for counting arguments.

A strategy S is defined by specifying for each $n \geq 1$, a sequence of functions $(f_t)_{t=1}^{\infty}$, where for each $t \in \mathbb{N}$, $f_t(u_1, v_1, \dots, u_{t-1}, v_{t-1}, u_t)$ is a distribution over [n] which depends on the vertex u_t , and the history of the process up until step t-1. Then, v_t is chosen according to this distribution. If f_t is an atomic distribution, then v_t is determined by $u_1, v_1, \dots, u_{t-1}, v_{t-1}, u_t$. Observe that this means that the player needs to select her strategy in advance, before the game actually starts. We then denote $(G_t^S(n))_{i=0}^t$ as the sequence of random (multi)graphs obtained by following the strategy S for t rounds; where we shorten $G_t^S(n)$ to G_t or $G_t(n)$ when clear.

Suppose \mathcal{P} is a monotonely increasing property. Given a strategy \mathcal{S} and a constant 0 < q < 1, let $\tau_{\mathcal{P}}(\mathcal{S}, q, n)$ be the minimum $t \geq 0$ for which $\mathbb{P}[G_t \in \mathcal{P}] \geq q$, where $\tau_{\mathcal{P}}(\mathcal{S}, q, n) := \infty$

if no such t exists. Define

$$\tau_{\mathcal{P}}(q,n) = \inf_{\mathcal{S}} \tau_{\mathcal{P}}(\mathcal{S}, q, n),$$

where the infimum is over all strategies on [n]. Observe that for each $n \geq 1$, if $0 \leq q_1 \leq q_2 \leq 1$, then $\tau_{\mathcal{P}}(q_1, n) \leq \tau_{\mathcal{P}}(q_2, n)$ as \mathcal{P} is increasing. Thus, the function $q \to \limsup_{n \to \infty} \tau_{\mathcal{P}}(q, n)$ is non-decreasing, and so the limit,

$$\tau_{\mathcal{P}} := \lim_{q \to 1^{-}} \limsup_{n \to \infty} \frac{\tau_{\mathcal{P}}(q, n)}{n},$$

is guaranteed to exist. The goal is typically to compute upper and lower bounds on $\tau_{\mathcal{P}}$ for various properties \mathcal{P} .

1.2. Main Results. In this work, we focus on the property of having a perfect matching, which we denote by PM. We remark that by convention, we say that a graph on an odd number of vertices has a perfect matching, if the matching saturates all but one vertex. Hence, if n is odd, then $\tau_{PM}(q,n) \leq \tau_{PM}(q,n+1)$ by the following natural coupling between the two corresponding processes. Indeed, the first strategy on n vertices may steal the second strategy on n+1 vertices if the square and the circle are both on [n]. Otherwise, the square and the circle are placed uniformly at random on [n]. Immediately the subgraph induced by [n] by the second strategy is a subgraph of the graph constructed by the first strategy and the inequality follows. Since τ_{PM} is an asymptotic definition in n, it suffices to consider only even n. Note also that since we focus on creating perfect matchings, we shall hereby restrict our attention to strategies which do not create self-loops.

Our first result is an improvement on the upper bound of τ_{PM} from 1+2/e<1.73576 to 1.20524. The current upper bound of 1+2/e follows from the fact, first observed in [3], that one may couple the semi-random process with the following process that generates a random bipartite graph. This process is known to have a perfect matching with probability tending to 1 as $n \to \infty$ (a.a.s.) [7]. The graph has two bipartite parts of size n/2 and is generated in two rounds. The first round is the bipartite version of the 1-out process. That is, each vertex chooses an out-neighbour independently and uniformly at random from the other vertex part. A vertex is classified as unpopular if it has been chosen by at most one vertex. In the second round, each unpopular vertex chooses another out-neighbour independently and uniformly at random from the other vertex part. The final bipartite graph is obtained by ignoring the directions of the arcs.

We propose a fully adaptive algorithm to construct a semi-random graph with a perfect matching, which gives the following upper bound on τ_{PM} .

Theorem 1.1. $\tau_{PM} \leq \beta + 10^{-5} \leq 1.20524$, where β is derived from a system of differential equations.

Our second result is an improvement on the lower bound of τ_{PM} from $\ln(2) \geq 0.69314$ to 0.93261. First observe that trivially $\tau_{PM} \geq 1/2$, as any strategy of the player must wait at least n/2 rounds in order to build a perfect matching. In fact, this lower bound can be improved to $\ln(2) \geq 0.69314$, as was first observed in [3]. There are two obvious necessary conditions for the existence of a perfect matching, both giving exactly the same lower bound. If there exists a perfect matching in G_t , then G_t has the minimum degree at least 1 and there are at least n/2 vertices in G_t with at least one square. In order to warm-up we re-prove this result in Subsection 3.1.

Proposition 1.2 ([3]). $\tau_{PM} \ge \ln(2) \ge 0.69314$.

The precise statement of our lower bound is in terms of a root of a function. Specifically, define

$$\alpha = \inf\{b \ge 0 : g(b) \ge 1/2\},\$$

where

$$g(b) := 1 + \frac{1-2b}{2} \exp(-b) - (b+1) \exp(-2b) - \frac{1}{2} \exp(-3b)$$

We prove the following:

Theorem 1.3. $\tau_{PM} \ge \alpha \ge 0.93261$.

1.3. **Previous Results.** Let us briefly describe a few known results on the semi-random process. In the very first paper [3], it was shown that the process is general enough to approximate (using suitable strategies) several well-studied random graph models. In the same paper, the process was studied for various natural properties such having minimum degree $k \in \mathbb{N}$ or having a fixed graph H as a subgraph. In particular, it was proven that a.a.s. one can construct H in less than $\omega n^{(d-1)/d}$ rounds where $d \geq 2$ is the degeneracy of H, and $\omega = \omega(n)$ is any function tending to infinity as $n \to \infty$. It was conjectured that their general lower bound is sharp. This conjecture was recently proven in [1].

The property of having a Hamilton cycle, which we denote by HAM, was also studied for the semi-random process. As observed in [3], if G_t has a Hamilton cycle, then G_t has the minimum degree at least 2 yielding $\tau_{\text{HAM}} \geq \ln 2 + \ln(1 + \ln 2) \geq 1.21973$. On the other hand, it is known that the famous 3-out process is a.a.s. Hamiltonian [4]. As the semi-random process can be coupled with the 3-out process, we get that $\tau_{\text{HAM}} \leq 3$. A new upper bound was obtained in [5] in terms of an optimal solution of an optimization problem, whose value is believed to be 2.61135 by numerical support. In the same paper, the lower bound mentioned above was shown to not be tight. Recently, the authors of this paper managed to analyze a fully adaptive algorithm further improving the upper bound and managed to substantially improve the lower bound. These results will be included in the forthcoming paper [6].

Finally, let us mention about the property of containing a given spanning graph H as a subgraph. It was asked by Noga Alon whether for any bounded-degree H, one can construct a copy of H a.a.s. in O(n) rounds. This question was answered positively in a strong sense in [2], in which is was shown that any graph with maximum degree Δ can be constructed a.a.s. in $(3\Delta/2 + o(\Delta))n$ rounds. They also proved that if $\Delta = \omega(\log(n))$, then this upper bound improves to $(\Delta/2 + o(\Delta))n$ rounds. Note that both of these upper bounds are asymptotic in Δ . When Δ is constant in n, such as in both the perfect matching and Hamiltonian cycle setting, determining the optimal dependence on Δ for the number of rounds needed to construct H remains open.

2. Upper Bound

In this section, we describe an adaptive algorithm which builds a perfect matching in around 1.2n rounds a.a.s. In order to warm-up, we start with a slightly simpler algorithm to analyze and then discuss the adjustments needed to claim the desired upper bound. Note that throughout the paper, floor is automatically applied to any number that needs to be an integer (typically a function of n).

2.1. Warming-up. In this section we analyse a simplified algorithm which constructs a semi-random graph with a perfect matching in less than 1.28n steps. The algorithm operates in two stages. In the first stage, the algorithm keeps building the matching greedily whenever possible, but will keep u_tv_t for future augmentation if u_t lands on a saturated vertex. In each step of the algorithm we will keep track of a matching M_t that is currently built and the set U_t of unsaturated vertices. We will also maintain a set of red vertices which is a subset of vertices that are saturated by M_t . If a red vertex is hit by a square, then an augmenting path will be created and we can update M_t to a larger matching. In order to briefly explain the idea, imagine uv is an edge in the matching. If a square lands on u then we will place its corresponding circle on an unsaturated vertex x, and then colour v red. If in the future a square is placed on v, then we will place its corresponding circle on an unsaturated vertex y. Now the path xuvy is an augmenting path. When the red vertex v receives a square, the algorithm needs to know the neighbour of u that is unsaturated, and there might be several of them. In this simple algorithm, we keep track of only the first square that landed on u.

We now formally describe the first stage of the algorithm. Suppose that at step t, u_t lands on a saturated vertex u and v_t is the corresponding circle which is necessarily placed on an unsaturated vertex. Colour vertex u as well as the edge $u_t v_t$ green, and colour the mate of u in the matching red. Below is a formal description of the t-th step of the algorithm.

- (A_1) If $u_t \in U_{t-1}$ then let v_t be a uniformly random vertex in U_{t-1} . Let $M_t = M_{t-1} \cup \{u_t v_t\}$ and $U_t = U_{t-1} \setminus \{u_t, v_t\}$. For every green vertex x, if it is adjacent to either u_t or v_t by a green edge then uncolour this green edge and uncolour x (from green) and uncolour the mate of x in M_t (from red).
- (A_2) If u_t lands on a red vertex, then let v_t be a uniformly random vertex in U_{t-1} . Let x be the mate of u_t in M_{t-1} . Let y be the vertex in U_{t-1} which is adjacent to x by a green edge. Let M_t be the matching obtained by augmenting along the path yxu_tv_t , and let $U_t = U_{t-1} \setminus \{y, v_t\}$. Update the green vertices and edges and the red vertices accordingly as in (A_1) .
- (A_3) If u_t lands on an uncoloured saturated vertex, then choose v_t uniformly at random from U_{t-1} . Colour the edge $u_t v_t$ and the vertex where u_t lands on green and colour the mate of u_t in M_{t-1} red. Let $M_t = M_{t-1}$ and $U_t = U_{t-1}$.
- (A_4) If u_t lands on a green vertex, then let v_t be an arbitrary vertex in [n]. Let $M_t = M_{t-1}$ and $U_t = U_{t-1}$. The edge $u_t v_t$ will not be used in the matching construction in the future.

The first stage terminates at the step when $|U_t|$ becomes at most εn where $\varepsilon = 10^{-14}$. In the second stage, we design the algorithm which saturates the remaining unsaturated vertices in at most $100\sqrt{\varepsilon}n = 10^{-5}n$ steps. We will define and analyse the second stage after analysing the first.

Let $X(t) = 2|M_t|$ and let R(t) denote the number of red vertices. By the algorithm, R(t) is also equal to the number of green vertices, and thus equal to the number of green edges. Let $H_t := (X(i), R(i))_{0 \le i \le t}$. Note that H_t does not encompass the full history of random graph process at time t (i.e., G_0, \ldots, G_t – the first t+1 graphs constructed by the randomized algorithm). We condition on less information so that the circle placements amongst the unsaturated vertices remain distributed uniformly at random. This allows us to claim the

following expected difference equations:

$$\mathbb{E}[X(t+1) - X(t) \mid H_t] = \frac{2(n - X(t) + R(t))}{n} + O(1/n),$$

$$\mathbb{E}[R(t+1) - R(t) \mid H_t] = \frac{n - X(t)}{n} \cdot \frac{-2R(t)}{n - X(t)} + \frac{R(t)}{n} \left(-1 - \frac{2(R(t) - 1)}{n - X(t)}\right) + \frac{X(t) - 2R(t)}{n} + O(1/n).$$

The first equation above is obvious, as the probability that u_{t+1} lands on a vertex in U_t or a red vertex is (n - X(t) + R(t))/n. The probability that v_{t+1} is on the same vertex as u_{t+1} in (A_1) , or is on vertex y in (A_2) , is O(1/n). In order to justify the second equation, note that if u_{t+1} lands on a vertex in U_t , then two vertices z_1 and z_2 in U_t become saturated after the augmentation. Since the ends of the set of green edges in U_t are uniformly distributed, the expected number of green edges incident with z_1 or z_2 is equal to 2R(t)/(n-X(t)). The other ends of these green edges become uncoloured from green after the augmentation which, in turn, forces their mates to become uncoloured from red. If u_{t+1} lands on a red vertex, the situation is similar, except that the mate of u_{t+1} is first uncoloured from green. If u_{t+1} lands on a green vertex, then there is no change to R(t). Finally, if u_{t+1} lands on a vertex that is neither unsaturated nor coloured, then a new green vertex is created. This explains the second equation above.

By writing x(s) = X(sn)/n and r(s) = R(sn)/n, we have that

$$x' = 2(1 - x + z), (1)$$

$$r' = \frac{-2r}{1-x}(1-x+r) - r + x - 2r, \tag{2}$$

with the initial conditions x(0) = r(0) = 0.

The right hand sides of (1) and (2) are continuous and Lipschitz in the connected open set $D = \{(s,x,r) : -\varepsilon < s < 3, |r| < 2, -\varepsilon < x < 1 - \varepsilon/4\}$ which contains the point (0,x(0),z(0)) = (0,0,0). Let T_D be the first step t where $(t/n,X(t)/n,R(t)/n) \notin D$. Then, $|X(t+1)-X(t)| \le 1$ for every $t < T_D$, and with probability $O(n^{-2})$, $|R(t+1)-R(t)| \le \log^2 n$ for every $t < T_D$ (in order to see the second assertion, note that, in expectation, every unsaturated vertex is adjacent to $R(t)/|U_t| \le 4/\varepsilon$ green vertices, and thus the claim easily follows by standard bounds on the tail probability of a binomial variable). By the differential equation method of Wormald (see, for example, [9, Theorem 5.1] or [8]) (applied with $\gamma = O(n^{-2})$, $\beta = \log^2 n$ and $\lambda = n^{-1/4}$), the differential equations (1) and (2) with the given initial conditions have a unique solution that can extend arbitrarily close to the boundary of D, and a.a.s. $X(t) = x(t/n)n + O(\lambda n)$ for every t where $t/n < \sigma$ where σ is the supremum of s where $x(s) \le 1 - \varepsilon/3$ and s < 3. Numerical calculations show that x reaches $1 - \varepsilon/3$ before s = 1.2769497. Hence, a.a.s. after 1.2769497n steps there are at most εn unsaturated vertices.

Finally, we describe the second stage of the algorithm, which we refer to as the **clean-up algorithm**. Let j_0 be the number of unsaturated vertices after the first stage. Thus, $j_0 \leq \varepsilon n$. Let $j_k = \lfloor (3/4)^k j_0 \rfloor$, for $k = 1, 2, \ldots$ For each $k \geq 1$, the algorithm saturates $j_{k-1} - j_k$ unsaturated vertices by doing the following.

(i) Uncolour all vertices and squares in the whole graph.

¹Maple worksheet may be accessed at https://math.ryerson.ca/~pralat/.

- (ii) Sprinkle $\lfloor \sqrt{3j_{k-1}n}/4 \rfloor$ semi-random edges u_tv_t as follows. If u_t lands on a saturated vertex x, then let v_t be an unsaturated vertex selected uniformly at random, and colour the mate of x in M_t by red (which is possibly already red). If u_t lands on an unsaturated vertex, then let v_t be an arbitrary vertex, and the algorithm will not use this edge for constructing the matching.
- (iii) Let \mathcal{R} be the set of red vertices after (ii). At each step t, if u_t lands on a red vertex or an unsaturated vertex then let v_t be a uniformly random unsaturated vertex, and then augment M_t as in (A_1) and (A_2) as in the first stage, and update \mathcal{R} accordingly. Stop when $|U_t| = j_k$. Let T_k be the total number of steps in (ii) and (iii).

The total number of steps in the final stage of constructing a perfect matching is then $\sum_{k\geq 1} T_k$. Next we bound T_k for every k. Of course, the number of steps in (ii) is at most $\sqrt{3j_{k-1}n}/4$. It is easy to show by standard concentration arguments that with probability $1 - O(n^{-2})$, $|\mathcal{R}| \geq \sqrt{3j_{k-1}n}/5$ after (ii). (Indeed, since $j_{k-1} \leq \varepsilon n$ and the number of rounds is $\sqrt{3j_{k-1}n}/4 = o(n)$, $\mathbb{E}[|\mathcal{R}|] \geq (1 - \varepsilon + o(1))\sqrt{3j_{k-1}n}/4$.) During the execution of (iii), for each augmentation that is performed, the expected number of red vertices that become uncoloured is at most

$$1 + 2 \cdot \frac{\sqrt{3j_{k-1}n}/4}{j_k},$$

since the total number of red vertices is at most $\sqrt{3j_{k-1}n}/4$, and the total number of unsaturated vertices in each round is at least j_k . There are $\frac{1}{2}(j_{k-1}-j_k)$ total augmentations in (iii). Thus, by standard concentration arguments, with probability $1 - O(n^{-2})$, at the end of (iii),

$$|\mathcal{R}| \ge \frac{\sqrt{3j_{k-1}n}}{5} - 2 \cdot \left(1 + 2 \cdot \frac{\sqrt{3j_{k-1}n}/4}{j_k}\right) \cdot \frac{j_{k-1} - j_k}{2} = \frac{\sqrt{3j_{k-1}n}}{30} - \frac{j_{k-1}}{4} \ge \frac{\sqrt{2j_{k-1}n}}{30},$$

where the last inequality holds since $j_{k-1} \leq \varepsilon n$ where $\varepsilon = 10^{-14}$ by our choice. Consequently, the probability that a square lands on a red vertex at each step in (iii) is at least $\sqrt{2j_{k-1}}/30\sqrt{n}$. Hence,

$$\mathbb{E}T_k \le \frac{1}{4}\sqrt{3j_{k-1}n} + \left(\frac{\sqrt{2j_{k-1}}}{30\sqrt{n}}\right)^{-1} \frac{j_{k-1} - j_k}{2} < 3.1\sqrt{j_{k-1}n}.$$

Then, again by standard concentration arguments, a.a.s.

$$T_k \le \max\{4\sqrt{j_{k-1}n}, \log^2 n\} = 4\sqrt{j_{k-1}n} \le 4\sqrt{(3/4)^{k-1}\varepsilon} \cdot n$$
, for every $k \ge 1$.

Thus, a.a.s. the total number of steps consumed in the second stage is at most

$$\sum_{k>1} T_k \le \sum_{k>1} 4\sqrt{(3/4)^{k-1}\varepsilon} \cdot n < 100\sqrt{\varepsilon}n.$$

By our choice of ε , $100\sqrt{\varepsilon} = 10^{-5}$, a.a.s. the total number of steps the algorithm spend on constructing a perfect matching is at most $(1.2769497 + 10^{-5})n < 1.27696n$.

2.2. Improving the Upper Bound. In this section, we describe a modified algorithm which allows us to get the claimed upper bound of Theorem 1.1. As before, the algorithm keeps building the matching greedily whenever possible. However, instead of making random choices, it will choose its circles both deterministically and in a strategic way. Specifically, suppose that u_t lands on a saturated vertex. In this case, we choose v_t such that v_t has the minimum number of circles on it amongst the vertices of U_{t-1} . We make the analogous

decision when u_t lands on an unsaturated vertex, as well as when u_t lands on a red vertex and we wish to extend the matching via an augmenting path. We now formally describe our algorithm for round t:

- (B_1) If $u_t \in U_{t-1}$, then let v_t be a vertex of U_{t-1} with a minimum number of circles among those in $U_{t-1} \setminus \{u_t\}$. Let $M_t = M_{t-1} \cup \{u_tv_t\}$ and $U_t = U_{t-1} \setminus \{u_t, v_t\}$. For every green vertex x, if it is adjacent to either u_t or v_t by a green edge then uncolour this green edge and x (from green) and uncolour the mate of x in M_t (from red).
- (B_2) If u_t lands on a red vertex, then let x be the mate of u_t in M_{t-1} . Let y be the vertex in U_t which is adjacent to x by a green edge. Let v_t be a vertex of U_{t-1} with a minimum number of circles among those in $U_{t-1} \setminus \{y\}$. Let M_t be the matching obtained by augmenting along the path yxu_tv_t , and let $U_t = U_{t-1} \setminus \{y, v_t\}$. Update the green vertices and edges and the red vertices accordingly as in (B_1) .
- (B_3) If u_t lands on an uncoloured saturated vertex, then let v_t be a vertex of U_{t-1} with a minimum number of circles. Colour the vertex where u_t lands on green and colour the mate of u_t in M_{t-1} red. Colour the edge $u_t v_t$ green. Let $M_t = M_{t-1}$ and $U_t = U_{t-1}$.
- (B_4) If u_t lands on a green vertex, then let v_t be an arbitrary vertex in $[n] \setminus U_{t-1}$. (We restrict to $[n] \setminus U_{t-1}$ so that we can conveniently keep track of the number of circles on our unsaturated vertices.) Let $M_t = M_{t-1}$ and $U_t = U_{t-1}$. The edge $u_t v_t$ will not be used in the matching construction in the future.

As before, we define X(t) to be the number of saturated vertices after $t \geq 0$ rounds. Given $q \geq 0$, we define $\widetilde{R}_q(t)$ to be the number of unsaturated vertices with precisely q circles after t rounds. Note that $\widetilde{R}_q(0) = 0$ for all $q \geq 1$, and $\widetilde{R}_0(0) = n$. Now, for $q \geq 1$, we say that a red vertex is of **type** q at round t, provided its mate in M_t is adjacent to an unsaturated vertex via a green edge with precisely q circles on it. We denote $R_q(t)$ as the number of red vertices of type q at time $t \geq 0$. Observe that $R_q(t) = q\widetilde{R}_q(t)$ for $q \geq 1$ and $t \geq 0$. Observe also that for every $t \geq 0$, there is q where all red vertices are either of type q or of type q + 1. For each $q \geq 1$, let us define the stopping time τ_q to be the smallest $t \geq 0$ such that

For each $q \geq 1$, let us define the stopping time τ_q to be the smallest $t \geq 0$ such that $\widetilde{R}_j(t) = 0$ for all j < q. It is obvious that τ_q is well-defined and is non-decreasing in q. Moreover, after the step τ_q , all unsaturated vertices in the graph have precisely q circles. This is obvious if what happened in step τ_q is in case (B_3) . Case (B_4) cannot happen in step τ_q , as the number of circles on the unsaturated vertices does not change in this case. If case (B_1) or (B_2) occurs in step τ_q , it is possible that v_t receives more than q circles in some rare situations, but v_t becomes saturated after the augmentation and so the number of circles on every other unsaturated vertex is equal to q. It follows then that $\widetilde{R}_q(t) = |U_t| = n - X(t)$. By definition, $\tau_0 = 0$. Let us refer to **phase** q as those $\tau_{q-1} \leq t < \tau_q$.

We shall analyze our algorithm for the first k phases using the differential equation method. For the case when k=1100, we show that a.a.s. $X(\tau_k) \geq (1-10^{-6})n$ and $\tau_k \leq 1.20365n$. We complete the perfect matching by running the randomized algorithm from Subsection 2.1.² The only difference is that the initial conditions are $x(0) = 1 - 10^{-6}$ (instead of x(0) = 0) and z(0) = 0. The conclusion is that a.a.s. after additional 0.00158n steps there are at most $10^{-14}n$ unsaturated vertices. As before, the clean-up algorithm for the remaining unsaturated vertices takes at most 0.00001n rounds, yielding the upper bound of (1.20365 + 0.00158 + 0.00001)n = 1.20524n.

²Maple worksheet may be accessed at https://math.ryerson.ca/~pralat/.

2.2.1. Analyzing the k phases. Unlike in the analysis of the randomized algorithm, we can safely condition on the full history of the random graph process in each phase. If G_0, \ldots, G_t correspond to the first t+1 graphs constructed by the deterministic algorithm, then let $H_t := (G_0, \ldots, G_t)$.

In phase 1, we place circles on the unsaturated vertices, thus creating red vertices of type 1. This leads to the following equations regarding the expected changes of X(t) and $R_1(t)$, for $0 \le t < \tau_1$:

$$\mathbb{E}[X(t+1) - X(t) \mid H_t] = 2\left(\frac{n - X(t) + R_1(t)}{n}\right),$$

$$\mathbb{E}[R_1(t+1) - R_1(t) \mid H_t] = \left(\frac{X(t) - 2R_1(t)}{n}\right) - \frac{2R_1(t)}{n} + O(1/n).$$

Observe that the first equation follows, since 2 vertices become saturated whenever a square lands on a red vertex, or an unsaturated vertex. The second equation follows since we create an additional red vertex whenever a square lands on an uncoloured saturated vertex (of which there exist $X(t) - 2R_1(t)$). We destroy a red vertex whenever a square lands on a type 1 red vertex, or an unsaturated vertex with one circle on it. There are $R_1(t)$ of each of these. Note that the O(1/n) term accounts for contributions from the following two cases. In the first case, u_t lands on an unsaturated vertex which is the unique one in U_t which is not covered by a circle. In that case, $R_1(t)$ decreases by one, but the probability that this case occurs is O(1/n). In the other case, u_t lands on a red vertex as in case (B_2) where y happens to be the only unsaturated vertex in U_t that is not covered by any circle. Again, it is easy to see that the expected contribution from this case is O(1/n).

Suppose that the functions $x_1, y_1 : [0, \infty) \to \mathbb{R}$ are the unique solution to the following system of differential equations:

$$x_1' = 2(1 - x_1 + y_1)$$

 $y_1' = x - 4y_1,$

with initial conditions $x_1(0) = y_1(0) = 0$. Let $c_1 > 0$ be the unique value such that $y_1(c_1) = 1 - x_1(c_1)$. By applying the differential equation method, we can derive the following:

Proposition 2.1. There exists a function $\varepsilon_1(n) = o(1)$, such that a.a.s for all $0 \le t \le \tau_1$ it holds that

$$\max\{|X(t) - x_1(t/n)n|, |R_1(t) - y_1(t/n)n|\} \le \varepsilon_1 n.$$

Moreover, $|\tau_1 - c_1 n| \le \varepsilon_1 n \ a.a.s.$

Let us now consider phase $q \geq 2$. Observe that there exist red vertices only of types q-1 and q during phase q. Thus, we can track the total number of red vertices at time t by just focusing on the random variables $R_{q-1}(t)$ and $R_q(t)$. Specifically, observe the following for $\tau_{q-1} \leq t < \tau_q$:

$$\mathbb{E}[X(t+1) - X(t) \mid H_t] = 2\left(\frac{n - X(t)}{n} + \frac{R_{q-1}(t)}{n} + \frac{R_q(t)}{n}\right),$$

$$\mathbb{E}[R_q(t+1) - R_q(t) \mid H_t] = q\left(\frac{(X(t) - 2R_q(t) - 2R_{q-1}(t)) - R_q(t)}{n}\right)$$

$$-\frac{R_q(t)}{n} + O(1/n),$$

where

$$R_{q-1}(t) = (q-1)\left(n - X(t) - \frac{R_q(t)}{q}\right).$$

The first equation follows since 2 vertices become saturated whenever a square lands on either an unsaturated vertex, or a red vertex. The second equation describes the expected change in the red vertices of type q. Note that q such vertices are created when a square lands on a saturated and uncoloured vertex, of which there are precisely $X(t) - 2R_q(t) - 2R_{q-1}(t)$. Similarly, q red vertices of type q are destroyed when the square lands on a red vertex of type q, as well as when the square lands on an unsaturated vertex with precisely q circles. Observe that there are $R_q(t)$ of the former, and $\widetilde{R}_q(t) = R_q(t)/q$ of the latter. Finally, the O(1/n) term accounts for the rare case when v_t is forced to be placed on a vertex with q circles for the same reason as in phase 1.

We define a system of differential equations for each phase $q \geq 2$, whose initial conditions are determined inductively. The differential equation method implies that the trajectories of the random variables X(t) and $R_q(t)$ are concentrated around the solutions of these differential equations in every phase. Let us assume that we have defined a system for phase q-1, whose solution has functions x_{q-1} and y_{q-1} , and a value $c_{q-1} \geq 0$ such that $y_{q-1}(c_{q-1}) = 1 - x_{q-1}(c_{q-1})$ (observe that the base case holds by Proposition 2.1). Note that we may interpret $c_{q-1}n$ as approximately the point in the process at which all the unsaturated vertices have q-1 circles, thus corresponding to the end of phase q-1. Define $x_q, y_q, z_q : [c_{q-1}, \infty) \to \mathbb{R}$ to be the unique solution to the following system of differential equations:

$$x'_q = 2(1 - x_q + y_q + z_q),$$

 $y'_q = q(x_q - 3y_q - 2z_q) - y_q,$

with initial conditions $x_q(c_{q-1}) = x_{q-1}(c_{q-1})$, $z_q(c_{q-1}) = y_{q-1}(c_{q-1})$, and $y_q(c_{q-1}) = 0$, where $z_q = (q-1)(1-x-y_q/q)$. Let $c_q \ge 0$ be the unique value such that $y_q(c_q) = 1-x_q(c_q)$ observe that this is precisely the value such that $z_q(c_q) = 0$, for all $q \ge 2$. By applying the differential equation method inductively for each phase q, we get the following:

Proposition 2.2. For each $q \ge 1$, there exists a function $\varepsilon_q(n) = o(1)$, such that a.a.s, for all $0 \le t \le \tau_q$ it holds that

$$\max\{|X(t) - x_q(t/n)n|, |R_q(t) - y_q(t/n)n|\} \le \varepsilon_q n.$$

Moreover, $|\tau_q - c_q n| \le \varepsilon_q n \ a.a.s.$

Proposition 2.3. If k = 1100, then a.a.s. $\tau_k \le 1.20365$ and $X(\tau_k) \ge (1 - 10^{-6})n$.

After k = 1100 phases, we have constructed a matching M_{τ_k} which a.a.s. has at least $(1-10^{-6})n/2$ edges. To complete the perfect matching, we apply the 2-stage randomised algorithm from Section 2.1. First, we uncolour all vertices and edges of the graph. Then, we keep exactly $(1-10^{-6})n/2$ edges in M_{τ_k} , and "artificially" delete the other edges in the matching, and put the incident vertices into U_{τ_k} . Starting again from t=0, the argument from Section 2.1 can be generalized to show that X(t) and R(t) follow the solutions of the differential equations (1) and (2) with initial conditions $x(0) = 1 - 10^{-6}$ and r(0) = 0. This is because we can reproduce the analysis of the algorithm from Section 2.1 with $(1-10^{-6})n$ initial edges in the matching, opposed to none. Numerical solutions show that $x(0.00158) > 1 - \varepsilon$, where $\varepsilon = 10^{-14}$. The clean-up algorithm from Section 2.1 completes the construction

in $100\sqrt{\varepsilon}n=10^{-5}n$ rounds. This yields an upper bound of $1.20365+0.00158+10^{-5}=1.20524$ on $\tau_{\rm PM}$, as in Theorem 1.1.

3. Lower Bound

Throughout the section, we will repeatedly apply the following claim when deriving bounds on $\tau_{\mathcal{P}}$ for a property \mathcal{P} :

Proposition 3.1. Given $\alpha > 0$, suppose that for any strategy S and $0 < \delta < \alpha$,

$$\mathbb{P}[G_T^{\mathcal{S}}(n) \in \mathcal{P}] \to 0$$

as
$$n \to \infty$$
, where $T = T(n) = (\alpha - \delta)n$. Then $\tau_{\mathcal{P}} \ge \alpha$.

3.1. Warming-up. We start with the proof of Proposition 1.2 as a warm-up for our improved result. Suppose that we consider some strategy S of the player which begins on the empty graph on vertex set [n]. Recall that we say the vertex $j \in [n]$ is **covered** by the square u_t arriving at round t, provided $u_t = j$. The analogous definition extends to the circle v_t . Let $S_j(t)$ denote the number of squares covering j, and let $C_j(t)$ denote the number of circles covering j, after t steps of the strategy.

Suppose that X(t) is the number of vertices of G_t which are covered by at least one square. Let us fix a constant $c < \ln(2)$, and consider T = T(n) = cn. Observe that if G_t has a perfect matching at time $t \ge 0$, then at least n/2 vertices of [n] must have been covered by at least one square; that is, $X(t) \ge n/2$. On the other hand, $X(t) = n - \sum_{i \in [n]} \mathbf{1}_{\{S_i(t)=0\}}$, and so

$$\mathbb{E}[X(t)] = \sum_{j \in [n]} (1 - \mathbb{P}[S_j(t) = 0]) = n \left(1 - \left(1 - \frac{1}{n} \right)^t \right)$$
$$= (1 + o(1))n(1 - \exp(-t/n)).$$

Thus, combined with a second moment computation, one can conclude that for any fixed $t \geq 0$, a.a.s.

$$X(t) = (1 + o(1))n(1 - \exp(-t/n)).$$

Now, $T/n \le c$, and $c < \ln(2)$, so a.a.s.

$$X(T) \le n(1 + o(1))(1 - \exp(-c)) < n/2.$$

Therefore, a.a.s. G_T does not have a perfect matching. Since this holds for each $c < \ln(2)$, Proposition 3.1 implies the desired property and the proof is finished.

3.2. Reducing to an Approximate Perfect Matching. Given a constant c > 0 and a function $\omega = \omega(n)$, we say that a strategy \mathcal{S} is ω -well-behaved for cn rounds, if for all $0 \le t \le \lfloor cn \rfloor$, each vertex of $G_t(n)$ is covered by at most ω circles. In order to prove Theorem 1.3, we work with strategies which are well-behaved for an appropriate choice of ω . We will also define a new property \mathcal{P} which satisfies $\tau_{\mathcal{P}} \le \tau_{\text{PM}}$. Moreover, we will show that general strategies do not perform significantly better than well-behaved strategies for building a graph in \mathcal{P} . This allows us to restrict to analysing well-behaved strategies, and the various random variables yielded by these strategies. Since the strategies are well-behaved, the set of random variables under concern have the nice "bounded-Lipschitz" property that is essential for the application of the differential equation method. We start with the definition of the new property \mathcal{P} on the family of directed graphs, and we use D[S] to denote the directed graph induced by set $S \subseteq V(D)$ for a directed graph D.

Definition 3.2. Suppose that D is a directed graph with m arcs on vertex set [n]. We say that D has an (μ, δ) -approximate perfect matching, denoted by $D \in PM(\mu, \delta)$, if there exists a subset $S \subseteq [n]$ such that:

- (A_1) D[S] has a perfect matching.
- (A_2) All the vertices of S have in-degree at most μ in D.
- $(A_3) |S| \geq n 2m\delta.$

A perfect matching of D[S] is then called a (μ, δ) -approximate perfect matching of D.

We first observe the following relation between PM and PM(μ, δ).

Proposition 3.3. For any function $\mu = \mu(n)$, it holds that

$$\tau_{PM(\mu, \mu^{-1})} \leq \tau_{PM}$$

Proof. Suppose that D is a directed graph which has a perfect matching M. Let \mathcal{V}_1 be the set of vertices whose in-degree is greater than $\mu(n)$. Let \mathcal{V}_2 be the set of isolated vertices obtained once vertices in \mathcal{V}_1 are removed from M. Clearly, $|\mathcal{V}_2| \leq |\mathcal{V}_1| \leq m/\mu$. Let $S = [n] \setminus (\mathcal{V}_1 \cup \mathcal{V}_2)$. Obviously, D[S] has a perfect matching, and $|S| \geq n - |\mathcal{V}_1| - |\mathcal{V}_2| \geq n - 2|\mathcal{V}_1| \geq n - 2m\mu^{-1}$. It follows that it is at least as easy to create an approximate perfect matching than to create a perfect one. The proof is completed.

Next, we confirm that general strategies do not behave significantly better than well-behaved strategies for constructing a directed graph in $PM(\mu, \mu^{-1})$, where we view every edge in G_t as an arc from the square to the circle. Moreover we may safely restrict to strategies that construct only simple graphs, as intuitively relocating v_t when u_tv_t is already an edge would perform no worse, if not better. In the next proposition, we obtain a strategy S_2 from a given strategy S_1 by copying the moves of S_1 whenever the move does not create a multiple edge, or create a vertex with too many circles. Otherwise, S_2 relocates the circle to a different vertex (for counting purpose we colour relocated circles blue). We show that S_2 performs at least as well as S_1 quantitatively.

Proposition 3.4. For any constant 0 < c < 1 and function $\mu = \mu(n) = \omega(1)$, if S_1 is a strategy of the player, then there exists another strategy S_2 which is 2μ -well-behaved for cn rounds, and satisfies

$$\mathbb{P}[G_{cn}^{\mathcal{S}_1}(n) \text{ satisfies PM}(\mu,\mu^{-1})] \leq \mathbb{P}[G_{cn}^{\mathcal{S}_2}(n) \text{ satisfies PM}(2\mu,\mu^{-1})].$$

Moreover, S_2 does not create any multi-edges, and so $G_{cn}^{S_2}(n)$ is a simple graph.

Proof. In order to prove the claim, we couple the execution of S_1 with another strategy S_2 which is 2μ -well-behaved, and which constructs a directed graph in $PM(2\mu, \mu^{-1})$ no later than when S_1 constructs one in $PM(\mu, \mu^{-1})$. We define S_2 by *stealing* the strategy of S_1 , while making a slight modification to avoid multi-edges and to ensure that no vertex of [n] is covered by more than 2μ circles.

Suppose that the squares presented to strategy S_1 are u_1, \ldots, u_{cn} , and whose corresponding circles are v_1^1, \ldots, v_{cn}^1 . We denote the graph formed by S_1 after $0 \le t \le cn$ rounds by G_t^1 . Let us define another strategy S_2 with the same set of squares as presented to S_1 , whose circles we denote by v_1^2, \ldots, v_{cn}^2 . For each $1 \le t \le cn$, suppose that u_t is presented to S_2 , and G_{t-1}^2 is the current graph constructed by S_2 (where G_0^2 is the empty graph):

(1) If adding $u_t v_t^1$ to G_{t-1}^1 does not cause the number of circles on v_t^1 to exceed μ , and $u_t v_t^1$ is not an edge in G_{t-1}^2 (note here that we consider G_{t-1}^2 as an undirected graph), then set $v_t^2 := v_t^1$.

- (2) Otherwise, choose $v_t^2 \in [n] \setminus N_{G_{t-1}^2}(u_t)$ amongst those vertices which have at most c/(1-c) blue circles, and then colour the circle v_t^2 blue.
- (3) Add (u_t, v_t^2) to G_{t-1}^2 to get G_t^2 .

We first observe that the strategy S_2 is well-defined in that there always exists a choice of v_t^2 which satisfies the requirements of (2). To see this, notice that trivially $|N_{G_{t-1}^2}(u_t)| \leq cn$, and so there are at least (1-c)n > 0 vertices which are not adjacent to u_t in G_{t-1}^2 , as c < 1 by assumption. Thus, since there are at most cn blue circles, a simple averaging argument ensures that there exists a vertex of $[n] \setminus N_{G_{t-1}^2}(u_t)$ which is covered by at most c/(1-c) blue circles.

We now argue that the strategy S_2 satisfies all the desired properties of Proposition 3.4. It is clear that S_2 does not create any multi-edges by definition, so it suffices to verify that each vertex of G_{cn}^2 is covered by at most 2μ circles, and that G_{cn}^2 is in $PM(2\mu, \mu^{-1})$ if G_{cn}^1 is in $PM(\mu, \mu^{-1})$. Observe that if we fix $j \in [n]$, then step (1) places at most μ (uncoloured) circles on j, and step (2) places at most $c/(1-c)+1 < \mu$ blue vertices of j. Thus, in total there are at most 2μ circles are placed on j, and so the strategy S_2 is 2μ -well-behaved. As a result, if G_{cn}^1 has a (μ, μ^{-1}) -approximate perfect matching, then G_{cn}^2 must have a $(2\mu, \mu^{-1})$ -approximate perfect matching. Thus,

$$\mathbb{P}[G^1_{cn} \text{ satisfies } \mathrm{PM}(\mu,\mu^{-1})] \leq \mathbb{P}[G^2_{cn} \text{ satisfies } \mathrm{PM}(2\mu,\mu^{-1})].$$

and so the proof is complete.

By combining the above propositions with Proposition 3.1, we get the following lemma:

Lemma 3.5. Given $0 < \alpha \le 1$ and $\mu = \mu(n) = \omega(1)$, suppose that for each $c < \alpha$ and each strategy S which is 2μ -well behaved for cn rounds, it holds that

$$\mathbb{P}[G_{cn}^{\mathcal{S}}(n) \text{ satisfies PM}(2\mu,\mu^{-1})] \to 0, \quad \text{as } n \to \infty.$$

Then, $\tau_{PM} \geq \alpha$.

3.3. **Proving Theorem 1.3.** In this section, we complete the proof of Theorem 1.3. Recall that $g:[0,\infty)\to\mathbb{R}$ is defined such that for $b\geq 0$,

$$g(b) := 1 + \frac{1 - 2b}{2} \exp(-b) - (b + 1) \exp(-2b) - \frac{1}{2} \exp(-3b), \tag{3}$$

and in viewing g as monotonely increasing on $[0, \infty)$, we have that

$$\alpha = \min\{\alpha \ge 0 : g(b) \ge 1/2\}.$$

Let $\mu = \sqrt{n}$ (indeed any function of n that grows with n in a sub-linear rate would work) and let $0 < c < \alpha \le 1$ be an arbitrary constant. Define ω to be 2μ . This specifies the parameters for the set of ω -well behaved strategies for cn rounds that are considered below.

Suppose that S is an ω -well behaved strategy for cn rounds. Since $c < \alpha \le 1$, we may apply Lemma 3.5 in order to prove Theorem 1.3. Thus, it suffices to show that

$$\mathbb{P}[G_{cn}^{\mathcal{S}}(n) \text{ satisfies } PM(\omega, \mu^{-1})] \to 0, \quad \text{as } n \to \infty.$$
 (4)

For simplicity and without confusion, we remove S from the superscript. In order to prove (4), we prove that a certain set of squares cannot be simultaneously contained in any (ω, μ^{-1}) -approximate perfect matching of G_{cn} and there are many such squares to exclude.

Given a vertex $j \in [n]$, we say that j is **redundant** at time $t \ge 0$, provided the following conditions hold:

- (B_1) Vertex j is covered by precisely one square, say u_s for $s \leq t$.
- (B_2) The circle v_s connected to u_s by the player is covered by at least one square, which arrives after round s.

We denote by U(t) the number of redundant vertices at time t. Observe that (B_2) ensures that U(t) depends only on the placement of the squares, not the strategy \mathcal{S} . Observe also that the redundant square on j and any square on v_s cannot be contained simultaneously in any (ω, μ^{-1}) -approximate perfect matching. Recall that X(t) denotes the number of vertices covered by at least one square after step t.

Suppose that j is redundant at time t thanks to the arrival of the square u_s at time $s \leq t$. We then say that j is **well-positioned**, provided the vertex v_s is also redundant at time t. Denote W(t) as the number of well-positioned redundant vertices at time t. Clearly, $W(t) \leq U(t)$ by definition. We observe then the following inequality:

Proposition 3.6. If G_t has an (ω, μ^{-1}) -approximate perfect matching at time $t \geq 0$, then

$$X(t) - U(t) + W(t) \ge \frac{n}{2} - \frac{3t}{\mu}.$$
 (5)

Proof. Since G_t has an (ω, μ^{-1}) -approximate perfect matching, there exists some subset $S \subseteq [n]$, such that $|S| \ge n - 2t/\mu$, for which $G_t[S]$ has a perfect matching. Now, if \mathcal{M} is a perfect matching of $G_t[S]$, then clearly $|\mathcal{M}| = |S|/2 \ge n/2 - t/\mu$.

On the other hand, suppose \mathcal{X}_S is the collection of vertices within S covered by at least one square, \mathcal{U}_S is the collection of vertices of S which are redundant, and \mathcal{W}_S is the number of well-positioned vertices of $G_t[S]$. Let us denote $X_S := |\mathcal{X}_S|$, $U_S := |\mathcal{U}_S|$ and $W_S := |\mathcal{W}_S|$ for convenience. We claim the following inequality:

$$X_S - U_S + W_S \ge |\mathcal{M}| \ge n/2 - t/\mu. \tag{6}$$

To see the left-hand side of this inequality, we first partition S into $\mathcal{X}_S \setminus \mathcal{U}_S$, \mathcal{U}_S and $\mathcal{Z}_S := S \setminus \mathcal{X}_S$, thus ensuring that

$$|\mathcal{M}| = e_{\mathcal{M}}(\mathcal{X}_S \setminus \mathcal{U}_S, S) + e_{\mathcal{M}}(\mathcal{U}_S, \mathcal{U}_S) + e_{\mathcal{M}}(\mathcal{U}_S, \mathcal{Z}_S). \tag{7}$$

Now, clearly $e_{\mathcal{M}}(\mathcal{X}_S \setminus \mathcal{U}_S, S) \leq |\mathcal{X}_S \setminus \mathcal{U}_S|$. On the other hand, we claim that $e_{\mathcal{M}}(\mathcal{U}_S, \mathcal{Z}_S) = 0$. This is because each vertex of \mathcal{U}_S is redundant, and so its single out-neighbour is a vertex covered by at least one square, and thus cannot be a member of \mathcal{Z}_S . Similarly, its inneighbours are covered by squares by definition, and thus are also not in \mathcal{Z}_S .

It remains to upper bound $e_{\mathcal{M}}(\mathcal{U}_S, \mathcal{U}_S)$. In order to do so, we partition \mathcal{U}_S into $\mathcal{U}_S \setminus \mathcal{W}_S$ and \mathcal{W}_S , which ensures that

$$e_{\mathcal{M}}(\mathcal{U}_S, \mathcal{U}_S) = e_{\mathcal{M}}(\mathcal{U}_S \setminus \mathcal{W}_S, \mathcal{U}_S \setminus \mathcal{W}_S) + e_{\mathcal{M}}(\mathcal{W}_S, \mathcal{U}_S).$$

Now, by definition each vertex $j \in \mathcal{U}_S \setminus \mathcal{W}_S$ has precisely one square on it, and the circle corresponding to this square lies in $\mathcal{X}_S \setminus \mathcal{U}_S$. Thus, $e_{\mathcal{M}}(\mathcal{U}_S \setminus \mathcal{W}_S) = 0$. Moreover, we can upper bound $e(\mathcal{W}_S, \mathcal{U}_S)$ by $|\mathcal{W}_S|$. By applying these bounds to (7), (6) follows.

We can complete the proof by observing that $X_S \leq X(t)$, $W_S \leq W(t)$, and $U_S \geq U(t) - (n - |S|) \geq U(t) - t/\mu$. Thus, after rearranging (6), we get that

$$X(t) - U(t) + W(t) \ge \frac{n}{2} - \frac{3t}{\mu},$$

as claimed. \Box

As we saw in Proposition 1.2, it is easy to control the behaviour of X(t) for $t \geq 0$. Controlling the behaviour of U(t) requires more care, and so it is useful to define Y(t) as the number of vertices of G_t which are covered by precisely one square. Let $H_t = (G_0, \ldots, G_t)$ be the history of the random graph process up to step t. We then have that for each $t \geq 1$:

$$\mathbb{E}[X(t+1) - X(t) \mid H_t] = 1 - \frac{X(t)}{n},\tag{8}$$

$$\mathbb{E}[Y(t+1) - Y(t) \mid H_t] = \frac{n - X(t)}{n} - \frac{Y(t)}{n},\tag{9}$$

and

$$\mathbb{E}[U(t+1) - U(t) \mid H_t] = \frac{Y(t) - U(t)}{n} - \frac{U(t)}{n}.$$
 (10)

Verification of (8)–(10) is straightforward. We briefly verify (10). The number of pairs of vertices (u, v) such that u is covered by exactly one square whose corresponding circle is on v and v has no squares arriving after that circle, is exactly Y(t) - U(t) by step t. A redundant vertex is created if a square lands on v. On the other hand, a redundant vertex can be destroyed if it receives another square. This explains (10).

In order to control the behaviour of W(t), we introduce another definition. Let us say that a vertex j_1 is **dangerous** at time $t \geq 0$, provided there exist times $t_1 < t_2 \leq t$, such that the following hold:

- (C_1) At time t_1 , the square u_{t_1} lands on j_1 , and connects its circle v_{t_1} to some vertex $j_2 \neq j_1$.
- (C_2) At time t_2 , the vertex j_2 is covered by the square u_{t_2} , at which point the circle v_{j_2} is placed on some vertex $j_3 \notin \{j_1, j_2\}$.
- (C_3) No squares land on j_3 between times t_2 and t, and u_{t_1} and u_{t_2} are the only squares on j_1 and j_2 , respectively.

Let us denote D(t) as the number of dangerous vertices at time t. Observe then that

$$\mathbb{E}[D(t+1) - D(t) \mid H_t] = \frac{Y(t) - U(t)}{n} - \frac{3D(t)}{n},\tag{11}$$

where we recall that Y(t) is the number of vertices covered by exactly one square, and U(t) is the number of redundant vertices. Note that the equality in (11) follows since the strategy we are analyzing does not create multi-edges by assumption. We can use D(t) to help us track the behaviour of W(t) at time t; that is, the well-positioned vertices. Specifically, observe that

$$\mathbb{E}[W(t+1) - W(t) | H_t] = \frac{D(t)}{n} - \frac{2W(t)}{n}.$$

Now, by assumption S is ω -well behaved—that is, there are at most $\omega = \omega(n)$ circles on any vertex of G_t . We can thus bound the worst case one-step changes of our random variables by $\omega(n)$. More explicitly, observe that for each $t \geq 0$

$$|X(t+1) - X(t)| \le \omega(n),$$

and the analogous upper bound is also true for the other random variables.

Suppose now that the functions $x, y, u, d, w : [0, \infty) \to \mathbb{R}$ are the unique solution to the following system of differential equations:

$$x' = 1 - x, y' = 1 - x - y, u' = y - 2u, d' = y - u - 3d, w' = d - 2w,$$

with initial conditions x(0) = y(0) = u(0) = d(0) = w(0) = 0. These functions have the following closed form solution:

- $x(b) = 1 \exp(-b)$,
- $y(b) = b \exp(-b)$,
- $u(b) = (b-1)\exp(-b) + \exp(-2b)$.
- $d(b) = \frac{1}{2} \exp(-b) + \frac{1}{2} \exp(-3b) \exp(-2b),$ $w(b) = \frac{1}{2} \exp(-b) b \exp(-2b) \frac{1}{2} \exp(-3b).$

The following lemma follows immediately by Nick Wormald's differential equation method (see, for example, [9, Theorem 5.1] or [8]).

Lemma 3.7. There exists a function $\varepsilon = \varepsilon(n) = o(1)$ such that a.a.s., for all $0 \le t \le cn$, $\max\{|X(t) - x(t/n)n|, |U(t) - u(t/n)n|, |W(t) - w(t/n)n|\} < \varepsilon n.$

By applying Lemma 3.7, we are guaranteed the existence of a function $\varepsilon = \varepsilon(n) = o(1)$, such that a.a.s. for all $t \geq 0$,

- $$\begin{split} \bullet \ & X(t)/n \leq 1 \exp(-t/n) + \varepsilon, \\ \bullet \ & U(t)/n \geq \left(\frac{t}{n} 1\right) \exp(-t/n) + \exp(-2t/n) \varepsilon, \\ \bullet \ & W(t)/n \leq \left(\frac{1}{2} \exp(-t/n) \frac{1}{2} \exp(-3t/n) \frac{t}{n} \exp(-2t/n)\right) + \varepsilon. \end{split}$$

Let us now assume that $\alpha \geq 0$ is the infimum over those $b \geq 0$ such that $g(b) \geq 1/2$, where g(b) is defined in (3). Now, $c < \alpha$ and $T(n) := \lfloor cn \rfloor$ satisfies T = (1 - o(1))cn. Thus,

$$\frac{X(T)}{n} + \frac{W(T)}{n} - \frac{U(T)}{n} \le g(c) + 3\varepsilon.$$

Moreover, g(c) < 1/2 as $c < \alpha$, and so since $\varepsilon, \mu^{-1} = o(1)$, it follows that a.a.s.

$$X(T) + W(T) - U(T) < n/2 - 3T/\mu$$
.

Thus, G_T does not have an (ω, μ^{-1}) -approximate perfect matching a.a.s. by Proposition 3.6. As $c < \alpha$ was arbitrary, the proof of Theorem 1.3 is complete after applying Lemma 3.5. \square

4. Conclusion

We have reduced the gap between the previous best upper and lower bounds on τ_{PM} by roughly a factor of four. That being said, we do not believe that any of our new bounds are tight. For instance, in the case of our upper bound, our strategy does not make use of subsequent squares which land on a green vertex. One way to improve the algorithm's performance would be to control these additional squares. The other way is to consider longer augmenting paths. However, there are not many saturated vertices with more than one square and thus the numerical improvement is not so significant. Tracing longer augmenting paths is harder to analyse and we did not attempt this. However, algorithmic simulations³ suggest that this approach will lead to an improved upper bound. In the case of our lower bound, we have identified certain subgraphs which preclude the strategy from building a perfect matching too quickly, and which no strategy can avoid creating. To improve our lower bound, one could track more complicated subgraphs which also prevent a perfect matching from being created too quickly. For instance, one could track the number of vertices which are covered by two squares, and whose neighbours are themselves *not* redundant. However, such subgraphs can be avoided by the strategy, and so they are harder to control. More importantly, these subgraphs are much rarer than redundant vertices, and so their contribution to the improvement of the lower bound is rather small.

We believe it is an interesting open question to resolve the remaining gap between our bounds on τ_{PM} . More generally, we believe it is an interesting direction to better understand the value of τ_{H} , when H is a graph with absolutely bounded maximum degree Δ which spans all the vertices in [n]. For instance, improving upon the upper bound of $3/2(\Delta + o(\Delta))$ of [2] remains an intriguing open question when Δ is small.

References

- [1] Natalie C Behague, Trent G Marbach, Pawel Pralat, and Andrzej Rucinski. Subgraph games in the semi-random graph process and its generalization to hypergraphs. arXiv preprint arXiv:2105.07034, 2021.
- [2] Omri Ben-Eliezer, Lior Gishboliner, Dan Hefetz, and Michael Krivelevich. Very fast construction of bounded-degree spanning graphs via the semi-random graph process. *Proceedings of the 31st Symposium on Discrete Algorithms (SODA'20)*, pages 728–737, 2020.
- [3] Omri Ben-Eliezer, Dan Hefetz, Gal Kronenberg, Olaf Parczyk, Clara Shikhelman, and Miloš Stojaković. Semi-random graph process. *Random Structures & Algorithms*, 56(3):648–675, 2020.
- [4] Tom Bohman and Alan Frieze. Hamilton cycles in 3-out. Random Structures & Algorithms, 35(4):393–417, 2009.
- [5] Pu Gao, Bogumił Kamiński, Calum MacRury, and Paweł Prałat. Hamilton cycles in the semi-random graph process. *European Journal of Combinatorics*, 99:103423, 2022.
- [6] Pu Gao, Calum MacRury, and Paweł Prałat. A fully adaptive strategy for hamiltonian cycles in the semi-random graph process. *draft*, 2022.
- [7] Michal Karoński, Ed Overman, and Boris Pittel. On a perfect matching in a random digraph with average out-degree below two. *Journal of Combinatorial Theory, Series B*, 143, 03 2020.
- [8] Lutz Warnke. On wormald's differential equation method. CoRR, abs/1905.08928, 2019.
- [9] Nicholas C Wormald et al. The differential equation method for random graph processes and greedy algorithms. Lectures on approximation and randomized algorithms, 73:155, 1999.

DEPARTMENT OF COMBINATORICS AND OPTIMIZATION, UNIVERSITY OF WATERLOO, WATERLOO, CANADA *Email address*: pu.gao@uwaterloo.ca

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF TORONTO, TORONTO, CANADA *Email address*: cmacrury@cs.toronto.edu

DEPARTMENT OF MATHEMATICS, RYERSON UNIVERSITY, TORONTO, CANADA *Email address*: pralat@ryerson.ca

³We did not simulate a longer augmenting path algorithm ourselves, however we thank an anonymous reviewer for informing us that their simulation suggests an upper bound of 1.09.