ON RESTRICTING SUBSETS OF BASES IN RELATIVELY FREE GROUPS

LUCAS SABALKA AND DMYTRO SAVCHUK

ABSTRACT. Let G be a finitely generated free, free abelian of arbitrary exponent, free nilpotent, or free solvable group, or a free group in the variety $\mathbf{A}_m \mathbf{A}_n$, and let $A = \{a_1, \ldots, a_r\}$ be a basis for G. We prove that, in most cases, if S is a subset of a basis for G which may be expressed as a word in A without using elements from $\{a_{l+1}, \ldots, a_r\}$, then S is a subset of a basis for the relatively free group on $\{a_1, \ldots, a_l\}$.

1. INTRODUCTION

Let F = F(A) be a free group of rank $r \ge 2$ with basis $A = \{a_1, \ldots, a_r\}$. A subgroup V of F is fully invariant if, for all endomorphisms $f: F \to F$, $f(V) \subseteq V$. Fully invariant subgroups of F include the trivial subgroup, the derived subgroups, the subgroups in the lower central series of F, and, more generally, verbal subgroups of F. Every fully invariant subgroup is normal in F, and the quotients of the form F/V include the free group, free nilpotent groups, and free solvable groups. A subset $S \subset F$ is primitive in F mod V if the corresponding set of cosets SV of V can be extended to a basis of F/V (i.e. to a generating set for F/V such that each mapping of this set to F/V extends to an endomorphisms of F/V).

We wish to study the following situation. Let V be a fully invariant subgroup of F and let S be a primitive set mod V. Assume that there exists some index $l \in \{1, \ldots, r-1\}$ such that, for each $s \in S$, the generators $a_t, t = l + 1, \ldots, r$ of F(A) are not used to express s as a reduced word in the alphabet A. In short, $S \subset \hat{F} \subset F$, where $\hat{F} := F(\{a_1, \ldots, a_l\})$. Then $\hat{V} = V \cap \hat{F}$ is a fully invariant subgroup of \hat{F} (as every endomorphism of \hat{F} can be extended to an endomorphism of F). Our main question is:

Main Question. Under what conditions is it true that S is also primitive in \hat{F} mod \hat{V} ?

If V is trivial – that is, if we are considering S to be primitive in F – and |S| = r - 1 (where |S| denotes the size of set S), then it is not difficult to show that S must be primitive in \hat{F} . The question becomes more interesting when |S| is allowed to vary, and when V is allowed to vary. Our main results answer this question for different choices of V, |S| and l. Let \mathbf{A}_m denote the variety of all abelian groups whose exponent divides m. In particular, $\mathbf{A} := \mathbf{A}_0$ is the variety of all abelian groups. Then the product variety $\mathbf{A}_m \mathbf{A}_n$ consists of extensions of abelian groups of exponent dividing m by abelian groups of exponent dividing n.

Theorem 1.1. Let $S \subset \hat{F}$ be primitive in $F \mod V$. Then S is primitive in \hat{F} mod \hat{V} if F/V is:

(1) free,

- (2) free abelian of arbitrary exponent (including 0),
- (3) free nilpotent,
- (4) free in the abelian-by-abelian variety $\mathbf{A}_m \mathbf{A}_n$ and at least one of the following conditions fails: (a) m = 0, (b) n > 0, or (c) either |S| = r-1 or |S| = l-1; or
- (5) free solvable and |S| = r 1.

The motivation for this theorem came from the statement in the free case, which was initially proved by the authors as a lemma pertaining to the work in [SS13]. The result was a generalization of one case of a result describing the structure of subwords of primitive elements in F that do not involve one of the generators of F. Though it was eventually realized that this lemma was unnecessary for the paper [SS13], the authors felt the lemma and proof interesting enough to generalize and publish.

We conclude the introduction with the following natural question.

Question 1.2. Does there exist a variety of groups for which the Main Question has a negative answer? In particular, are there varieties where a stably primitive element is not necessarily primitive?

The structure of the paper is as follows. We begin by examining the case where V is trivial in Section 2. Then we proceed to cases of the free abelian groups of arbitrary exponent in Section 3 and the free nilpotent and free solvable groups in Section 4. We conclude the paper by the cases of free abelian-by-abelian of arbitrary exponents in Section 5 and free solvable groups in Section 6.

The authors would like to thank Mark Sapir for helpful advice on this paper, and Michael Handel, Ilya Kapovich, Olga Kharlampovich, Marcin Mazur, and Benjamin Steinberg for useful conversations on this material. And we are especially grateful to Martin Kassabov for providing a short proof of the free nilpotent case that replaced a longer argument involving 'lifting primitivity' and to Vitaliĭ Roman'kov for pointing out an error in an earlier version of the paper. Finally, we greatly appreciate the comments and suggestions of the referee that enhanced the paper.

2. The Free Case

In the free case, the answer to our Main Question is: always. Our proof uses an interpretation of primitivity using Fox derivatives via a theorem of Umirbaev.

We begin by recalling the definition of free Fox derivatives and establish notation for them. Let $\mathbb{Z}F_r$ denote the integral ring over the free group F_r . For each $j = 1, 2, \ldots, r$ define a linear map $\partial_j : \mathbb{Z}F_r \to \mathbb{Z}F_r$ recursively by

$$\partial_j(a_j) = 1, \qquad \partial_j(a_i) = 0, i \neq j$$

and

$$\partial_j(uv) = \partial_j(u) + u\partial_j(v)$$
 for all $u, v \in F_i$

The map ∂_j is called the (left) free Fox derivative associated with a_j .

Theorem 2.1. [Umi94] A subset $\{x_1, \ldots, x_k\}$ is primitive in F_r if and only if the $k \times r$ Jacobian matrix $J = [\partial_j(x_i)]_{1 \le i \le k, 1 \le j \le r}$, is right invertible in the integral ring $\mathbb{Z}F_r$.

We note that the "only if" part in this theorem was proven by Birman in [Bir73].

Theorem 2.2. Let $S = \{x_1, x_2, \dots, x_k\} \subset \hat{F}$ be primitive in F. Then S is primitive in \hat{F} .

Proof. By Theorem 2.1, since the set S is primitive in F, the associated Jacobian J is such that there exists an $r \times k$ matrix $P = [p_{il}]$ with $p_{il} \in \mathbb{Z}F_r$ satisfying

where I_k is the $k \times k$ identity matrix over $\mathbb{Z}F_r$.

For m > l, since elements of S do not involve a_m , the m^{th} column of the matrix J consists of zeros and all other entries of J do not involve a_m . Each entry of matrix P can be uniquely written in the form $p_{ij} = q_{ij} + r_{ij}$, where q_{ij} represents the sum of all terms in p_{ij} involving some a_m with m > l, and r_{ij} is an element of $\mathbb{Z}F_l$ not involving any a_m with m > l. Then for matrices $Q = [q_{ij}]$ and $R = [r_{ij}]$ we have P = Q + R, and by Equation (1), $JQ + JR = I_k$. But each entry of matrix JQ is either 0 or involves a_m for some m > l. Since neither I_k nor JR involves a_m for m > l we must have that JQ = 0, yielding

$$(2) JR = I_k$$

Now let \hat{J} and \hat{R} be the matrices obtained from J and R, respectively, by deleting last r-l columns and rows, respectively. Then \hat{J} is the Jacobian matrix of the set S seen as a subset of $\hat{F} = F_l$ and \hat{R} is a matrix over $\mathbb{Z}\hat{F} = \mathbb{Z}F_l$. Also, equation (2) implies that

$$\hat{J}\hat{R} = I_k$$

Thus by Umirbaev's criterion the set S is a subset of the basis of F.

Via private communication, Ilya Kapovich has detailed an alternative proof of Theorem 2.2, using Gersten's characterization of Whitehead's algorithm for subgroups [Ger84]. Olga Kharlampovich has also suggested a proof using Bass-Serre theory. The benefit of the above proof is that it generalizes to the case of free abelian-by-abelian groups of arbitrary exponents in Section 5 (and also to the case of free metabelian groups and partially to the case of free solvable groups, though shorter proofs in these cases are given here).

3. The Abelian Case

The answer to our Main Question in the free abelian case is also: always. Our proof works for free abelian groups of arbitrary exponent and uses linear algebra.

For any n > 0, Let $\mathbb{Z}_n := \mathbb{Z}/n\mathbb{Z}$ denote the free group of rank 1 in the variety \mathbf{A}_n of all abelian groups of exponent dividing n, and let F^n denote the subgroup of F generated by all n^{th} powers and similarly define \hat{F}^n . Let F' denote the commutator subgroup of F and similarly define \hat{F}' . Note $(\hat{F})' = (\widehat{F'})$, so \hat{F}' is well defined. Further, for each $x \in F$ we denote by \bar{x} the image of x in F/F' under the canonical epimorphism. For a subset A of F define $\bar{A} = \{\bar{a} : a \in A\}$.

We start with the proof for free abelian groups in the case of exponent 0.

Proposition 3.1. Let $S = \{x_1, x_2, \ldots, x_k\} \subset \hat{F}$ be primitive in $F \mod F'$. Then S is primitive in $\hat{F} \mod \hat{F}'$.

Before we proceed to our proof, we note that this Proposition follows from a wellknown fact (see, for example, [KM79]) that the basis of any subgroup of a finitely generated free abelian group can be extended to a basis of the whole group. Indeed, since \bar{S} can be extended to a basis in F/F', we get that \bar{S} is linearly independent. Hence, \bar{S} is a basis of a subgroup $\langle \bar{S} \rangle < \hat{F}/\hat{F'}$. We give a proof of Proposition 3.1 that extends to the case of free abelian groups of finite exponent in Theorem 3.2.

Proof. There is a canonical embedding $\mathbb{Z}^l \cong \hat{F}/\hat{F}' \hookrightarrow F/F' \cong \mathbb{Z}^r$ and with a slight abuse of notation we will sometimes identify \hat{F}/\hat{F}' with its image under this embedding.

The standard basis of $\mathbb{Z}^r \cong F/F'$ is $\{\bar{a}_1, \ldots, \bar{a}_r\}$, whereas $\{\bar{a}_1, \ldots, \bar{a}_l\}$ is the standard basis of $\mathbb{Z}^l \cong \hat{F}/\hat{F}'$. Extend \bar{S} to a basis B of \mathbb{Z}^r . Express B with respect to the standard basis as a matrix M, where the elements of \bar{S} correspond to the first k columns of M. Let \hat{M} denote the submatrix of M obtained by deleting the last r-l rows and columns. Then \hat{M} corresponds to a set \hat{B} of l elements of \mathbb{Z}^l . Since the elements of \bar{S} do not involve generators $\bar{a}_{l+1}, \ldots, \bar{a}_r$, the matrix M has a block triangular shape:

$$M = \left[\begin{array}{cc} \hat{M} & P \\ 0 & Q \end{array} \right]$$

for some $l \times (r-l)$ -matrix P and $(r-l) \times (r-l)$ -matrix Q. Then since M is in $SL_r(\mathbb{Z})$ we have

$$\det M \cdot \det Q = \det M = \pm 1.$$

This is possible only if det $\hat{M} = \pm 1$, and \hat{M} is invertible. This shows \hat{B} is a basis of $\mathbb{Z}^l \cong \hat{F}/\hat{F}'$. But $\bar{S} \subset \hat{B}$, so \bar{S} is primitive in \hat{F}/\hat{F}' and thus S is primitive in \hat{F} mod \hat{F}' .

Now we proceed to the case of free abelian groups with exponent n > 0. First we note that the free group of rank r in this variety is isomorphic to $F/(F'F^n)$.

Theorem 3.2. Let $S = \{x_1, x_2, \ldots, x_k\} \subset \hat{F}$ be primitive in $F \mod F'F^n$. Then S is primitive in $\hat{F} \mod \hat{F}'\hat{F}^n$.

Proof. The proof is almost identical to that of Proposition 3.1. We replace F' with $F'F^n$, \hat{F}' with $\hat{F}'\hat{F}^n$, and \mathbb{Z} with \mathbb{Z}_n , but otherwise define all other objects in the same way. The only other change to make is the following argument to show that \hat{M} is invertible. The matrix M is invertible in $SL_r(\mathbb{Z}_n)$ if and only if det M is a unit in \mathbb{Z}_n . But a product of elements in \mathbb{Z}_n is a unit if and only if each of the elements is also a unit. Thus, det $M = \det \hat{M} \cdot \det Q$ is a unit in \mathbb{Z}_n if and only if det \hat{M} is a unit in \mathbb{Z}_n , if and only if \hat{M} is invertible in $SL_r(\mathbb{Z}_n)$.

4. The Free Nilpotent Case

We now turn to the free nilpotent and free solvable cases. A commutator of weight c is an expression recursively defined by $[y_1, \ldots, y_{c-1}, y_c] := [[y_1, \ldots, y_{c-1}], y_c]$ and $[y_1, y_2] := y_1^{-1} y_2^{-1} y_1 y_2$. Fix a value for $c \ge 2$, and let $\gamma_c(F)$ be the normal closure of subset of F consisting of all commutators of weight c (i.e. the c^{th} term in the lower central series of F). Then $F/\gamma_{c+1}(F)$ is the free nilpotent group of class c. Also define $\gamma_1(F) = F$. Further, we write $F^{(t)}$ for the t^{th} derived subgroup of F. Then $F/F^{(t)}$ is a free solvable group of derived length t. Note that $F' = \gamma_2(F) = F^{(1)}$, so the free abelian group is the free nilpotent group of class 1 and the free solvable group of derived length 1.

Theorem 4.1.

(1) Let $S = \{x_1, x_2, \dots, x_k\} \subset \hat{F}$ be primitive in $F \mod \gamma_{t+1}(F)$ for some $t \geq 1$. Then S is primitive in $\hat{F} \mod \gamma_{t+1}(\hat{F})$.

The following proof is based on suggestions and references from Martin Kassabov and the referee. In the nilpotent case, our original proof used a primitivity lifting technique [GG92]. In the solvable case, our original proof was based on a generalization of our approach in the free case, and used Fox derivatives and criteria for primitivity by Roman'kov and Timoshenko in the free metabelian case [Rom91a, Tim92], and by Krasnikov in the free solvable case [Kra79]. However, our original proofs were longer and less general.

Proof. It is a well-known fact (see, for example, [MKS04, Lemma 5.9]) that a set of elements generates a nilpotent group N if and only if the projection of these elements to the abelianization N/N' generates the abelianization. Assume a subset $S = \{x_1, \ldots, x_k\}$ is primitive in $F \mod \gamma_{t+1}(F)$ and S does not involve generators a_{l+1}, \ldots, a_r . Then S is also primitive over F', which implies by Proposition 3.1 that S is primitive in $\hat{F} \mod \hat{F}'$. Now by the above fact S must be primitive in $\hat{F} \mod \gamma_{t+1}(\hat{F})$.

5. The Abelian-by-Abelian Case

For integers $m, n \geq 0$, we consider the case of free groups in the variety $\mathbf{A}_m \mathbf{A}_n$ of all extensions of abelian groups of exponent dividing m by abelian groups of exponent dividing n. Let $V_{m,n}$ and $\hat{V}_{m,n}$ denote the corresponding fully invariant subgroups of F and \hat{F} , respectively. The free Fox derivatives ∂_i induce Fox derivatives $\partial_i^0 : \mathbb{Z}F \to \mathbb{Z}_m(F/F'F^n)$ (see Gupta and Timoshenko [GT96]) by

$$\mathbb{Z}F \xrightarrow{\partial_i} \mathbb{Z}F \xrightarrow{\alpha^*} \mathbb{Z}(F/F'F^n) \xrightarrow{\gamma^*} \mathbb{Z}_m(F/F'F^n),$$

where $\alpha^* \colon \mathbb{Z}F \to \mathbb{Z}(F/F'F^n)$ and $\gamma^* \colon \mathbb{Z}(F/F'F^n) \to \mathbb{Z}_m(F/F'F^n)$ are linear extensions of the canonical epimorphisms $\alpha \colon F \to F/F'F^n$ and $\gamma \colon \mathbb{Z} \to \mathbb{Z}_m$ respectively.

In [GT99], Gupta and Timoshenko prove the following theorem.

Theorem 5.1. [GT99] Let $S = \{x_1, x_2, \ldots, x_k\} \subset F$. Let $J = (\partial_j^0 x_i), i = 1, \ldots, k, j = 1, \ldots, r$ denote the $k \times r$ Jacobian matrix of S over $\mathbb{Z}_m(F/F'F^n)$. Assume that at least one of the conditions m = 0, n > 0, or k = r - 1 fails. Then S is primitive mod $V_{m,n}$ if and only if:

- (i) the ideal in the ring $\mathbb{Z}_m(F/F'F^n)$ generated by k^{th} order minors of J is the whole ring, and
- (ii) S is primitive mod $F'F^n$.

Using the above theorem we deduce:

Theorem 5.2. Let $S = \{x_1, x_2, \ldots, x_k\} \subset \hat{F}$ be primitive in $F \mod V_{m,n}$ and assume at least one of the following conditions fails: (a) m = 0, (b) n > 0, or (c) either k = r - 1 or k = l - 1. Then S is primitive in $\hat{F} \mod \hat{V}_{m,n}$.

Proof. Let m_1, \ldots, m_t denote the k^{th} order minors of J in $\mathbb{Z}_m(F/F'F^n)$, and let J_m denote the ideal of $\mathbb{Z}_m(F/F'F^n)$ generated by m_1, \ldots, m_t . As $S \subset \hat{F}$, J is a matrix over $\mathbb{Z}_m(\hat{F}/\hat{F}'\hat{F}^n)$ whose last r-l columns are zero columns. Therefore the $l \times k$ Jacobian matrix $\hat{J} = (\partial_j x_i)$ formed by viewing S as a subset of \hat{F} is obtained from J by simply removing last r-l columns. But removing columns of zeros does not change the k^{th} order minors of a matrix, so the k^{th} order minors of \hat{J} are also m_1, \ldots, m_t , considered now as elements in $\mathbb{Z}_m(\hat{F}/\hat{F}'\hat{F}^n)$ (via the canonical embedding of $\mathbb{Z}_m(\hat{F}/\hat{F}'\hat{F}^n)$ into $\mathbb{Z}_m(F/F'F^n)$). Let \hat{J}_m denote the ideal generated by m_1, \ldots, m_t in the ring $\mathbb{Z}_m(\hat{F}/\hat{F}'\hat{F}^n)$.

Since S is primitive in F mod $V_{m,n}$, by Theorem 5.1 we know that: (i) $J_m = \mathbb{Z}_m(F/F'F^n)$, and (ii) S is primitive mod $F'F^n$. It follows from (i) that J_m contains the identity in $\mathbb{Z}_m(F/F'F^n)$. Therefore for some $p_i \in \mathbb{Z}_m(F/F'F^n)$ we have

(3)
$$1 = \sum_{i=1}^{t} m_i p_i$$

We can decompose each p_i as

$$p_i = q_i + r_i,$$

where q_i is the sum of terms in p_i that do not involve a_{l+1}, \ldots, a_r , and r_i is the sum of terms in p_i that do involve at least one of the a_{l+1}, \ldots, a_r in non-zero power. Then equation (3) can be rewritten as

$$1 - \sum_{i=1}^{t} m_i q_i = \sum_{i=1}^{t} m_i r_i$$

The left-hand side of the above equation does not have terms involving a_{l+1}, \ldots, a_r . However, since every term of each m_i does not involve any of a_{l+1}, \ldots, a_r but every term of each r_i does, every term of $m_i r_i$ involves at least one of the elements a_{l+1}, \ldots, a_r . Some of the terms of $m_i r_i$ might cancel, but it follows that every surviving term of $m_i r_i$ must involve at least one of the elements a_{l+1}, \ldots, a_r . Further, when we take the sum of $m_i r_i$ we cannot create terms that do not involve a_{l+1}, \ldots, a_r as the support of $\sum_{i=1}^t m_i r_i$ viewed as a function from $F/F'F^n$ to \mathbb{Z}_m is included into the union of supports of $m_i r_i$, which, in turn, are included in the set of all elements of $F/F'F^n$ that involve a_{l+1}, \ldots, a_r . Thus, for the two sides to be equal, the only possibility is if both sides are equal to zero. But this yields that

$$1 = \sum_{i=1}^{t} m_i q_i$$

with $m_i, q_i \in \mathbb{Z}_m(\hat{F}/\hat{F}'\hat{F}^n)$. Thus the identity in $\mathbb{Z}_m(\hat{F}/\hat{F}'\hat{F}^n)$ belongs to the ideal \hat{J}_m , and so condition (i) holds for S with respect to the ring $\mathbb{Z}_m(\hat{F}/\hat{F}'\hat{F}^n)$.

It follows from (*ii*), the fact that $S \subset \hat{F}$, and Proposition 3.1 and Theorem 3.2 that S is primitive mod $\hat{F}'\hat{F}^n$. Thus, applying Theorem 5.1 to S thought of as a subset of \hat{F} , we see that S is primitive in $\hat{F} \mod \hat{V}_{m,n}$. Note that Theorem 5.1 applies in this situation if at least one of the conditions m = 0, n > 0, or $k = rank(\hat{F}) - 1 = l - 1$ fails, which is true by assumption.

Vitalii Roman'kov suggested that one could potentially use the basis cofinality property of free groups of countable rank in subvarieties **B** of $\mathbf{N}_c \mathbf{A}$ (where \mathbf{N}_c denotes the variety of nilpotent groups of class c) to generalize our results in this section. This property was proved in this case by Bryant and Roman'kov in [BR99] (see also [BE97] for definitions). To do so, one must show that primitive systems in a finitely generated relatively free group in **B** that miss certain generators can be lifted to primitive systems in F with the same property. For arbitrary varieties this statement is false, but it could be true for the varieties above.

6. The Free Solvable Case

The final case we consider is the case of free solvable groups. First of all, the free metabelian case is covered by Section 5. Alternatively, for the free metabelian case one can use the primitivity criteria developed by Timoshenko [Tim88, Tim98] and Roman'kov [Rom91b, Rom91a], which use Fox derivatives similar in spirit to Umirbaev's criterion [Umi94] stated in Theorem 2.1. Unfortunately, there is no such criterion for free solvable groups of arbitrary derived length. However, Krasnikov [Kra78] has obtained a result for groups of the form F/[N, N] where N is a normal subgroup of F, which works only in the case k = r. The Fox derivatives used here are the free Fox derivatives of Section 2.

Theorem 6.1. [Kra78] A set $\{x_1, \ldots, x_r\}$ in F/[N, N] is a generating set of this group if and only if the $r \times r$ Jacobian matrix $J = [D_j(x_i)]_{1 \le i,j \le r}$ of free Fox derivatives, projected to Z(F/N), is left invertible in the integral ring $\mathbb{Z}(F/N)$.

With this criterion we obtain an analog of Theorem 2.2 in the case of free solvable groups with a restriction on the size of S.

Theorem 6.2. Let $S = \{x_1, x_2, \ldots, x_{r-1}\} \subset \hat{F}$ be primitive in $F \mod F^{(t)}$, where $F^{(t)}$ is the t-th derived subgroup of F, t > 0. Then S is a basis for $\hat{F}/\hat{F}^{(t)}$.

Proof. Since S is primitive in $F \mod F^{(t)}$, we can extend S to a set $\tilde{S} = \{x_1, x_2, \ldots, x_r\} \subset F$ such that the images of elements of \tilde{S} in $F/F^{(t)}$ under the canonical epimorphism form a basis. By Theorem 6.1 we have that the associated to \tilde{S} Jacobian matrix $J = [D_j(x_i)]_{1 \leq i,j \leq r}$ is left invertible over $\mathbb{Z}(F/F^{(t-1)})$, so there exists an $r \times r$ matrix $P = [p_{ij}]_{1 \leq i,j < r}$ with entries over $\mathbb{Z}(F/F^{(t-1)})$ such that

$$PJ = I_r$$

where I_r is an $r \times r$ identity matrix over $\mathbb{Z}(F/F^{(t-1)})$.

Since elements of S do not involve the generator a_r , we have

$$D_r(x_j) = 0, \quad j = 1, \dots, r-1$$

and $D_i(x_j)$ does not involve a_r for $1 \leq j \leq r-1$. Therefore, matrix J can be written in the form

$$J = \begin{bmatrix} 0\\ \hat{J} & \vdots\\ 0\\ \hline D_1(x_r) \cdots D_{r-1}(x_r) & D_r(x_r) \end{bmatrix},$$

where $\hat{J} = [D_i(x_j)]_{1 \le i,j \le r-1}$ is the Jacobian martix of S in $\hat{F}/\hat{F}^{(t-1)}$.

Then

$$I_r = PJ = \begin{bmatrix} p_{1r}D_r(x_r) \\ * & \vdots \\ p_{r-1,r}D_r(x_r) \\ \hline & & p_{rr}D_r(x_r) \end{bmatrix}$$

Hence, $p_{rr}D_r(x_r) = 1$ in $\mathbb{Z}(F/F^{(t-1)})$ and, in particular, $D_r(x_r) \neq 0$. On the other hand,

$$p_{in}D_r(x_r) = 0, \quad i = 1, \dots, r-1,$$

so we must have $p_{in} = 0$ in $\mathbb{Z}(F/F^{(t-1)})$. This follows from the fact that the free solvable group $F/F^{(t-1)}$ is a torsion-free elementary amenable group, and for such groups the Kaplansky conjecture on zero divisors holds true [Lin91]. In particular, this implies that the ring $\mathbb{Z}(F/F^{(t-1)})$ does not have zero divisors.

Now we have

$$I_{r} = PJ = \begin{bmatrix} \hat{P} & 0 \\ \vdots \\ 0 \\ \hline p_{r1} \cdots p_{r,r-1} & p_{rr} \end{bmatrix} \begin{bmatrix} \hat{J} & 0 \\ \vdots \\ 0 \\ \hline D_{1}(x_{r}) \cdots D_{r-1}(x_{r}) & D_{r}(x_{r}) \end{bmatrix}$$
$$= \begin{bmatrix} \hat{P}\hat{J} & \vdots \\ 0 \\ \hline \vdots \\ \hline p_{r}D_{r}(x_{r}) \\ \hline p_{r}D_{r}(x_{r}) \end{bmatrix}.$$

Therefore we must have $\hat{P}\hat{J} = I_{r-1}$, where I_{r-1} is the $(r-1) \times (r-1)$ identity matrix over $\mathbb{Z}(F/F^{(t-1)})$. Similarly to the proof in the free case, it follows that \hat{P} can be chosen to be in $\mathbb{Z}(\hat{F}/\hat{F}^{(t-1)})$. Finally, applying Theorem 6.1 again, we obtain that S is a basis for $\hat{F}/\hat{F}^{(t)}$.

References

- [BE97] Roger M. Bryant and David M. Evans. The small index property for free groups and relatively free groups. J. London Math. Soc. (2), 55(2):363–369, 1997.
- [Bir73] Joan S. Birman. An inverse function theorem for free groups. Proc. Amer. Math. Soc., 41:634–638, 1973.
- [BR99] R. M. Bryant and V. A. Roman'kov. Automorphism groups of relatively free groups. Math. Proc. Cambridge Philos. Soc., 127(3):411-424, 1999.
- [Ger84] S. M. Gersten. On Whitehead's algorithm. Bull. Amer. Math. Soc. (N.S.), 10(2):281– 284, 1984.
- [GG92] C. K. Gupta and N. D. Gupta. Lifting primitivity of free nilpotent groups. Proc. Amer. Math. Soc., 114(3):617–621, 1992.
- [GT96] C. K. Gupta and E. I. Timoshenko. Primitivity in the free groups of the variety $\mathfrak{A}_m\mathfrak{A}_n$. Comm. Algebra, 24(9):2859–2876, 1996.
- [GT99] Ch. K. Gupta and E. I. Timoshenko. Primitive systems of elements in the varieties $A_m A_n$: a criterion and induction. Algebra Log., 38(5):513–530, 638, 1999.
- [KM79] M. I. Kargapolov and Ju. I. Merzljakov. Fundamentals of the theory of groups, volume 62 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1979. Translated from the second Russian edition by Robert G. Burns.
- [Kra78] A. F. Krasnikov. Generating elements of the group F/[N, N]. Mat. Zametki, 24(2):167– 173, 301, 1978.

8

- [Kra79] A. F. Krasnikov. Generating elements of the group F/[N, N]. Math. Notes, 24(2):591–594, 1979.
- [Lin91] P. A. Linnell. Zero divisors and group von Neumann algebras. Pacific J. Math., 149(2):349–363, 1991.
- [MKS04] Wilhelm Magnus, Abraham Karrass, and Donald Solitar. Combinatorial group theory. Dover Publications Inc., Mineola, NY, second edition, 2004. Presentations of groups in terms of generators and relations.
- [Rom91a] V. A. Roman'kov. Criteria for the primitivity of a system of elements of a free metabelian group. Ukrain. Mat. Zh., 43(7-8):996-1002, 1991.
- [Rom91b] V. A. Roman'kov. Primitive elements of free groups of rank 3. Mat. Sb., 182(7):1074– 1085, 1991.
- [SS13] Lucas Sabalka and Dmytro Savchuk. On the geometry of the edge splitting complex. to appear in Groups, Geometry, and Dynamics, 2013. Preprint: arXiv:1007.1998.
- [Tim88] E. I. Timoshenko. On the inclusion of given elements into a basis of a free metabelian group. VINITI, (No. 2699-V), 1988. (In Russian).
- [Tim92] E. I. Timoshenko. Algorithmic solvability of the inclusion problem in a basis for a free metabelian group. Mat. Zametki, 51(3):117–121, 1992.
- [Tim98] E. I. Timoshenko. Primitive systems of elements in the variety \mathfrak{AN}_2 and some locally finite varieties of groups. Algebra i Logika, 37(6):687–699, 746, 1998.
- [Umi94] U. U. Umirbaev. Primitive elements of free groups. Uspekhi Mat. Nauk, 49(2(296)):175– 176, 1994.

Department of Mathematics and Computer Science, Saint Louis University $E\text{-}mail\ address:\ lsabalka@slu.edu$

Department of Mathematics and Statistics, University of South Florida $E\text{-}mail \ address: savchuk@usf.edu$