

EVERY GROUP IS A MAXIMAL SUBGROUP OF THE FREE IDEMPOTENT GENERATED SEMIGROUP OVER A BAND

IGOR DOLINKA AND NIK RUŠKUC

Dedicated to Stuart W. Margolis on the occasion of his 60th birthday

ABSTRACT. Given an arbitrary group G we construct a semigroup of idempotents (band) B_G with the property that the free idempotent generated semigroup over B_G has a maximal subgroup isomorphic to G . If G is finitely presented then B_G is finite. This answers several questions from recent papers in the area.

1. INTRODUCTION

Let S be a semigroup. The set $E = E(S)$ of all idempotents of S carries a structure of a partial algebra, called the *biordered set* of S , by retaining the products of the so-called *basic pairs*: these are pairs of idempotents $\{e, f\}$ such that $\{ef, fe\} \cap \{e, f\} \neq \emptyset$. It should be noted that if $ef \in \{e, f\}$ then fe is also an idempotent, possibly different from e, f and ef . Also, if S is an idempotent semigroup (i.e. a *band*) then its biordered set is in general different from S itself, since not every pair is necessarily basic. The term ‘biordered set’ comes from an alternative (but equivalent) approach, where one considers $E(S)$ as a relational structure equipped with two partial pre-orders; here we shall not pursue this approach, directing instead to [4, 5, 6, 11, 16] for further background.

The class of idempotent generated semigroups is of prime importance in semigroup theory, with a host of natural examples, such as the semigroups of singular (non-bijective) transformations of a finite set (Howie [12]) or singular $n \times n$ matrices over a field (Erdős [7]). It is not difficult to show that the category of all idempotent generated semigroups with a fixed biordered set E has an initial object $\text{IG}(E)$, called the *free idempotent generated semigroup* over E (we shall also say ‘over S ’ when $E = E(S)$). This semigroup is defined by the presentation

$$\text{IG}(E) = \langle E \mid e \cdot f = ef \text{ } (\{e, f\} \text{ is a basic pair}) \rangle.$$

Here $e \cdot f$ stands for a word of length 2 in the free semigroup E^+ , while ef is the element of E to which the product equals in S . Unsurprisingly, $\text{IG}(E)$ plays a crucial role in understanding the structure of semigroups with a prescribed biordered set of idempotents.

For reasons that are intrinsic to basic structure theory of semigroups [11, 13], this in turn depends upon the knowledge of maximal subgroups of $\text{IG}(E)$. It was conjectured for a long time that the maximal subgroups of $\text{IG}(E)$ are always free; this conjecture was widely circulated back in the 1980s, and was explicitly recorded in [15]. The conjecture was proved in a number of particular cases, see e.g. [15, 17, 19]. In 2009, Brittenham, Margolis and Meakin [1] disproved the conjecture by means of

2010 *Mathematics Subject Classification.* 20M05, 20F05.

Key words and phrases. Free idempotent generated semigroup, maximal subgroup, band.

an explicit 72-element semigroup S such that $\text{IG}(E(S))$ has a maximal subgroup isomorphic to $\mathbb{Z} \oplus \mathbb{Z}$, the free abelian group of rank 2. This was followed by Gray and Ruškuc [9] who proved that *every* group arises as a maximal subgroup of $\text{IG}(E(S))$ for a suitably chosen semigroup S ; if the group in question is finitely presented then a finite S will suffice. Further ensuing work such as [10, 3, 8] investigates maximal subgroups of $\text{IG}(S)$ for some specific natural semigroups S , and the first author [2] initiates the study of $\text{IG}(B)$, where B is a band.

The aim of the present note is to prove the result announced in the title:

Theorem 1. *Let G be a group. Then there exists a band B_G such that $\text{IG}(B_G)$ has a maximal subgroup isomorphic to G . Furthermore, if G is finitely presented, then B_G can be constructed to be finite.*

This single construction provides an alternative, simpler proof of *all* the main results of [9] (Theorems 1–4), resolves [9, Problem 1] which asks whether every finitely presented group is a maximal subgroup of $\text{IG}(S)$ for some finite *regular* semigroup S , and solves [2, Problem 2] which calls for a characterisation of maximal semigroups of free idempotent generated semigroups over bands.

2. PRESENTATION FOR MAXIMAL SUBGROUPS

A general presentation for maximal subgroups of $\text{IG}(S)$ in terms of parameters that depend only on the structure of S has been exhibited in [9, Theorem 5]. Since we are interested here only in the case of bands, we utilise the particular form of this theorem, deduced in [2, Corollary 5].

First of all, recall [13, Theorem 4.4.1] that any band B decomposes into a semilattice of rectangular bands, which are the \mathcal{D} -classes of B . Thus a \mathcal{D} -class D of S can be viewed as an $I \times J$ ‘table’ of idempotents e_{ij} ($i \in I, j \in J$), where $\{R_i : i \in I\}$ and $\{L_j : j \in J\}$ are the \mathcal{R} - and \mathcal{L} -classes in D respectively. For $i, k \in I$ and $j, l \in J$ we refer to the tuple $(e_{ij}, e_{il}, e_{kj}, e_{kl})$ as the $(i, k; j, l)$ square.

Suppose now we have an element $f \in B$ belonging to a \mathcal{D} -class above D . From the basic theory of bands (see, for example, [13, Section 4.4]) we know that f induces idempotent mappings $\sigma : I \rightarrow I, i \mapsto \sigma(i)$, and $\tau : J \rightarrow J, j \mapsto (j)\tau$, such that for all $i \in I, j \in J$ we have

$$fe_{ij} = e_{\sigma(i),j}, \quad e_{ij}f = e_{i,(j)\tau}.$$

We say that the square $(i, k; j, l)$ is *singular* induced by f if one of the following holds:

- (a) $\sigma(i) = i, \sigma(k) = k$ and $(j)\tau = (l)\tau \in \{j, l\}$; or
- (b) $\sigma(i) = \sigma(k) \in \{i, k\}$ and $(j)\tau = j, (l)\tau = l$.

We talk of a *left-right* or *up-down* singular square depending on whether (a) or (b) applies.

With the above conventions the general presentation we need is as follows:

Proposition 2 ([9, 2]). *The maximal subgroup H of $\text{IG}(B)$ containing $e_{11} \in D$ is presented by*

$$\langle f_{ij} \ (i \in I, j \in J) \mid \quad f_{i1} = f_{1j} = 1 \quad (i \in I, j \in J), \quad (1)$$

$$f_{ij}^{-1}f_{il} = f_{kj}^{-1}f_{kl} \quad ((i, k; j, l) \text{ a singular square in } D) \rangle. \quad (2)$$

3. CONSTRUCTION OF B_G

Let G be any group. Let us choose and fix a presentation $\langle A \mid R \rangle$ for G in which every relation has the form $ab = c$ for some $a, b, c \in A$. It is clear that G has such a presentation – for instance the Cayley table would do. What is less obvious, but nonetheless still true, is that if G is finitely presented then it has a *finite* presentation of this form. One way of seeing this is as follows: A relation $a_1 \dots a_k = b_1 \dots b_l$ can be replaced by two relations of the form $a_1 \dots a_k = c, b_1 \dots b_l = c$, at the expense of introducing a new generator c . Furthermore, the relation $a_1 \dots a_k = c$ can be replaced by $k - 1$ relations $a_1 a_2 = d_2, d_2 a_3 = d_3, \dots, d_{k-2} a_{k-1} = d_{k-1}, d_{k-1} a_k = c$ of the desired form, with new generators d_2, \dots, d_{k-1} .

Define sets

$$A_0 = A \cup \{0\}, A'_0 = \{a' : a \in A_0\}, I = A_0 \cup A'_0, J = A_0 \cup \{\infty\},$$

where $0, \infty$ and a' ($a \in A_0$) are symbols distinct from each other and those already in A . Consider the semigroup $\mathcal{T} = \mathcal{T}_I^{(l)} \times \mathcal{T}_J^{(r)}$, where $\mathcal{T}_I^{(l)}$ (respectively $\mathcal{T}_J^{(r)}$) is the semigroup of all mappings $I \rightarrow I$ (resp. $J \rightarrow J$) written on the left (resp. right). The semigroup \mathcal{T} has a unique minimal ideal K consisting of all (σ, τ) with both σ and τ constant. This ideal is naturally isomorphic to the rectangular band $I \times J$, and we will identify the two. We will visualise K as in Figure 1.




FIGURE 1. A visual representation of $K = I \times J$, highlighting the partition $I = A_0 \cup A'_0$, as well as the four distinguished rows and columns.

We now define a set $L \subseteq \mathcal{T}$. All elements $(\sigma, \tau) \in L$ will have

$$\sigma^2 = \sigma, \tau^2 = \tau, \ker(\sigma) = \{A_0, A'_0\}, \text{im}(\tau) = A_0. \quad (3)$$

Recall that $\ker(\sigma)$ is the equivalence on I defined by $(i, i') \in \ker(\sigma)$ if and only if $\sigma(i) = \sigma(i')$, and that it can be identified with the resulting partition of I into equivalence classes. Therefore, each (σ, τ) will be uniquely determined by $\text{im}(\sigma)$ which must be a two-element set that is a cross-section of $\{A_0, A'_0\}$, and the value $(\infty)\tau \in A_0$. The elements of L come in four groups: Z – the initial pair; G, \overline{G} – the elements arising from the generators A ; R – the elements arising from the relations R :

Type	Notation	Indexing	$\text{im}(\sigma)$	$(\infty)\tau$
Z	(σ_0, τ_0)	–	$\{0, 0'\}$	0
G	(σ_a, τ_a)	$a \in A$	$\{0, a'\}$	a
\overline{G}	$(\overline{\sigma}_a, \overline{\tau}_a)$	$a \in A$	$\{a, a'\}$	0
R	(σ_r, τ_r)	$r = (ab, c) \in R$	$\{b, c'\}$	a

These elements can be visualised as shown in Figure 2.




FIGURE 2. The elements (σ_0, τ_0) , (σ_a, τ_a) , $(\overline{\sigma}_a, \overline{\tau}_a)$ ($a \in A$), (σ_r, τ_r) ($r = (ab, c) \in R$) of L . For each (σ, τ) shaded are the two rows corresponding to $\text{im}(\sigma)$ and one column corresponding to $(\infty)\tau$. They all have $\ker(\sigma) = \{A_0, A'_0\}$ and $\text{im}(\tau) = A_0$.

Because $\ker(\sigma)$ and $\text{im}(\tau)$ are the same for all $(\sigma, \tau) \in L$ it follows that L is a left zero semigroup (i.e. $xy = x$ for all $x, y \in L$). Furthermore, since K is an ideal in \mathcal{T} (i.e. $xy, yx \in K$ for all $x \in K, y \in \mathcal{T}$), the set $B_G = K \cup L$ is a subsemigroup of \mathcal{T} . We remark that, strictly speaking, B_G depends not only on G , but crucially on the chosen presentation for G .

4. PROOF OF THEOREM 1

We will now use the presentation given in Proposition 2 to compute the maximal subgroup H of $\text{IG}(B_G)$ containing the idempotent $e_0 = (0, 0) \in K$. Relations (1) in our context read

$$f_{0j} = f_{i0} = 1 \quad (i \in I, j \in J). \quad (4)$$

The remaining relations (2) arise from the singular squares induced by the elements of L acting on K . Each up-down singular square is of one of the following forms:

$$(a_1, a_2; c_1, c_2), (a'_1, a'_2, c_1, c_2) \quad (a_1, a_2, c_1, c_2 \in A_0).$$

The square $(a_1, a_2; c_1, c_2)$ yields the relation

$$f_{a_1, c_1}^{-1} f_{a_1, c_2} = f_{a_2, c_1}^{-1} f_{a_2, c_2} \quad (a_1, a_2, c_1, c_2 \in A_0). \quad (5)$$

Putting $a_1 = c_1 = 0, a_2 = a, c_2 = c$ and using (4) yields

$$f_{a, c} = 1 \quad (a, c \in A_0); \quad (6)$$

clearly, all the remaining relations (5) are consequences of (6). Similarly, the squares (a'_1, a'_2, c_1, c_2) yield the relations

$$f_{a',c} = f_{0',c} \quad (a, c \in A_0). \quad (7)$$

(Note that we do not necessarily have $f_{0',c} = 1$, and so cannot deduce $f_{a',c} = 1$.)

Turning to the left-right singular squares, each $(\sigma, \tau) \in L$ induces precisely one. Below we list respectively the squares introduced by (σ_0, τ_0) of type **Z**, (σ_a, τ_a) of type **G**, $(\bar{\sigma}_a, \bar{\tau}_a)$ of type $\bar{\mathbf{G}}$, and (σ_r, τ_r) of type **R**, together with the relations they yield:

$$(0, 0'; 0, \infty) : \quad f_{0,0}^{-1} f_{0,\infty} = f_{0',0}^{-1} f_{0',\infty} \quad (8)$$

$$(0, a'; a, \infty) : \quad f_{0,a}^{-1} f_{0,\infty} = f_{a',a}^{-1} f_{a',\infty} \quad (a \in A) \quad (9)$$

$$(a, a'; 0, \infty) : \quad f_{a,0}^{-1} f_{a,\infty} = f_{a',0}^{-1} f_{a',\infty} \quad (a \in A) \quad (10)$$

$$(b, c'; a, \infty) : \quad f_{b,a}^{-1} f_{b,\infty} = f_{c',a}^{-1} f_{c',\infty} \quad (r = (ab, c) \in R). \quad (11)$$

Using the relations (4), (6), (7), we can transform (8)–(11) into:

$$f_{0',\infty} = 1 \quad (12)$$

$$f_{a',\infty} = f_{0',a} \quad (a \in A) \quad (13)$$

$$f_{a,\infty} = f_{a',\infty} = f_{0',a} \quad (a \in A) \quad (14)$$

$$f_{0',b} = f_{0',a}^{-1} f_{0',c} \quad (r = (ab, c) \in R). \quad (15)$$

So, the group H is defined by the generators $f_{i,j}$ ($i \in I, j \in J$) and relations (4), (6), (7), (12)–(15). The relations (4), (6), (7), (12)–(14) can be used simply to eliminate all the generators except $f_{0',a}$ ($a \in A$). Replacing each symbol $f_{0',a}$ by the symbol a , the remaining relations (15) become

$$ab = c \quad (r = (ab, c) \in R).$$

In other words, we obtain the original presentation for G . This proves that $H \cong G$.

Finally note that if $\langle A \mid R \rangle$ is a finite presentation, the semigroup B_G is also finite, with

$$|B_G| = (2|A| + 2)(|A| + 2) + 1 + 2|A| + |R|,$$

and this completes the proof of our theorem.

5. AN EXAMPLE, TWO REMARKS AND AN OPEN PROBLEM

It may be instructive to follow in a specific example the sequence of Tietze transformations constituting the brunt of the above proof. Let us take $G = Q_8$, the quaternion group, with the well known Fibonacci $F(2, 3)$ presentation (see [14, Section 7.3]):

$$\langle a, b, c \mid ab = c, bc = a, ca = b \rangle.$$

The dimension of K in this case is 8×5 , and Proposition 2 gives a presentation in terms of 40 generators. This is then simplified by a sequence of generator eliminations, using relations (1), up-down singular squares, and left-right singular squares induced by the elements of L of types **Z**, **G**, $\bar{\mathbf{G}}$. In the final step further singular squares are revealed, giving back the original presentation.

If we record the original generators in a natural 8×5 grid, this process may be encapsulated as shown in Figure 3.




FIGURE 3. The sequence of Tietze transformations constituting the proof of Theorem 1.

Remark 3. It is possible to describe completely the structure of the free idempotent generated semigroup $\text{IG}(B_G)$. By known results (see e.g. [9, (IG1)–(IG4)]) $\text{IG}(B_G)$ has precisely two regular \mathcal{D} -classes. The ‘upper’ one is a left zero semigroup \bar{L} isomorphic to L (as all products in L are basic), while the ‘lower’ one \bar{K} , the completely simple minimal ideal, has a Rees matrix representation with structure group G and (normalised) sandwich matrix (a_{ji}^{-1}) , where (a_{ij}) is the $|I| \times |J|$ table that is the end-product of Tietze transformations performed in the proof of Theorem 1 (in our example this is the last table in Figure 3). We claim that in fact $\text{IG}(B_G) = \bar{L} \cup \bar{K}$. To confirm this, and see that the structure is completely determined, we need to show how to write products ef and fe with $e \in L, f \in K$ as products of idempotents from K in $\text{IG}(B)$. For the product ef note that there exists $g \in K$ such that $f\mathcal{R}g$ and $ge = g$; both pairs $\{e, g\}$ and $\{f, g\}$ are critical and we have $ef = egf = hf$, where $h = eg \in K$. The product fe can be treated similarly.

Remark 4. Associated to the biorder E of idempotents of a regular semigroup S there is another free idempotent generated object $\text{RIG}(E)$, the free regular idempotent generated semigroup on E . It is the largest regular semigroup with the biorder of idempotents E , and its presentation can be obtained by adding further relations to the defining presentation for $\text{IG}(E)$. For definition and references we refer the

reader to [9]. In particular, from (IG1)–(IG4), (RIG1), (RIG2) in [9] it follows that $\text{IG}(B_G) = \text{RIG}(B_G)$.

One way of interpreting Remarks 3, 4 is to say that the word problem for $\text{IG}(B_G)$ is decidable if and only if the word problem for G is decidable. It is the authors' belief that the next stage in the ongoing exploration of free idempotent generated semigroups is precisely an analysis of the word problem for $\text{IG}(S)$. This at present seems a daunting task, even in the case where S is finite. Nonetheless, we propose the following problem which may just be within reach at this stage:

Question 1. Let B be a finite band such that all maximal subgroups of $\text{IG}(B)$ have recursively soluble word problems. Is the word problem of $\text{IG}(B)$ necessarily recursively soluble?

Acknowledgement. The research of the first author is supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia through Grant No.174019, and by a grant (Contract 114–451–2675/2012) of the Secretariat of Science and Technological Development of the Autonomous Province of Vojvodina. Also, the first author gratefully acknowledges the hospitality of the School of Mathematics and Statistics of the University of St Andrews, where this research was carried out.

REFERENCES

- [1] M. Brittenham, S. W. Margolis and J. Meakin, Subgroups of free idempotent generated semigroups need not be free, *J. Algebra* **321** (2009), 3026–3042.
- [2] I. Dolinka, A note on maximal subgroups of free idempotent generated semigroups over bands, *Periodica Math. Hungar.* **65** (2012), 97–105.
- [3] I. Dolinka and R. Gray, Maximal subgroups of free idempotent generated semigroups over the full linear monoid, *Trans. Amer. Math. Soc.*, to appear. [arXiv:1112.0893](#)
- [4] D. Easdown, Biordered sets of bands, *Semigroup Forum* **29** (1984), 241–246.
- [5] D. Easdown, Biordered sets are biordered subsets of idempotents of semigroups, *J. Austral. Math. Soc. Ser. A* **37** (1984), 258–268.
- [6] D. Easdown, Biordered sets come from semigroups, *J. Algebra* **96** (1985), 581–591.
- [7] J. A. Erdos, On products of idempotent matrices, *Glasgow Math. J.* **8** (1967), 118–122.
- [8] V. Gould, D. Yang, Every group is the maximal subgroup of a naturally occurring free idempotent generated semigroup, [arXiv:1209.1242](#)
- [9] R. Gray and N. Ruškuc, On maximal subgroups of free idempotent generated semigroups, *Israel J. Math.* **189** (2012), 147–176.
- [10] R. Gray and N. Ruškuc, Maximal subgroups of free idempotent generated semigroups over the full transformation monoid, *Proc. London Math. Soc.* **105** (2012), 997–1018.
- [11] P. M. Higgins, *Techniques of Semigroup Theory*, Oxford University Press, New York, 1992.
- [12] J. M. Howie, The subsemigroup generated by the idempotents of a full transformation semigroup, *J. London Math. Soc.* **41** (1966), 707–716.
- [13] J. M. Howie, *Fundamentals of Semigroup Theory*, Oxford University Press, New York, 1995.
- [14] D. L. Johnson, *Presentations of Groups*, LMS Student Texts Vol. 15, Cambridge University Press, Cambridge, 1990.
- [15] B. McElwee, Subgroups of the free semigroup on a biordered set in which principal ideals are singletons, *Comm. Algebra* **30** (2002), 5513–5519.
- [16] K. S. S. Nambooripad, Structure of regular semigroups. I, *Mem. Amer. Math. Soc.* **22** (1979), no. 224, vii+119 pp.
- [17] K. S. S. Nambooripad and F. Pastijn, Subgroups of free idempotent generated regular semigroups, *Semigroup Forum* **21** (1980), 1–7.
- [18] N. Ruškuc, Presentations for subgroups of monoids, *J. Algebra* **220** (1999), 365–380.
- [19] F. Pastijn, The biorder on the partial groupoid of idempotents of a semigroup, *J. Algebra* **65** (1980), 147–187.

[20] M. Petrich, *Introduction to Semigroups*, Merrill, Columbus, 1973.

DEPARTMENT OF MATHEMATICS AND INFORMATICS, UNIVERSITY OF NOVI SAD, TRG DOSITEJA
OBRADOVIĆA 4, 21101 NOVI SAD, SERBIA

E-mail address: dockie@dmf.uns.ac.rs

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF ST ANDREWS, ST ANDREWS KY16
9SS, SCOTLAND, UK

E-mail address: nik@mcs.st-and.ac.uk