# LOCAL ANALYSIS OF GRAUERT–REMMERT-TYPE NORMALIZATION ALGORITHMS

### JANKO BÖHM, WOLFRAM DECKER, AND MATHIAS SCHULZE

ABSTRACT. Normalization is a fundamental ring-theoretic operation; geometrically it resolves singularities in codimension one. Existing algorithmic methods for computing the normalization rely on a common recipe: successively enlarge the given ring in form an endomorphism ring of a certain (fractional) ideal until the process becomes stationary. While Vasconcelos' method uses the dual Jacobian ideal, Grauert–Remmert-type algorithms rely on so-called test ideals.

For algebraic varieties, one can apply such normalization algorithms globally, locally, or formal analytically at all points of the variety. In this paper, we relate the number of iterations for global Grauert–Remmert-type normalization algorithms to that of its local descendants.

We complement our results by an explicit study of ADE singularities. This includes the description of the normalization process in terms of value semigroups of curves. It turns out that the intermediate steps produce only ADE singularities and simple space curve singularities from the list of Frühbis-Krüger.

## INTRODUCTION

Normalization of rings is an important concept in commutative algebra, with applications in algebraic geometry and singularity theory. In fact, geometrically, normalization leads to a desingularization in codimension one. In particular, in the case of curves, it is the same as desingularization.

Given a reduced Noetherian ring A for which the normalization  $\overline{A}$  is a finite A-module, an obvious approach for finding  $\overline{A}$  is to successively enlarge A in form of an endomorphism ring of a fractional ideal. Specific instances of such algorithms require a recipe for choosing the fractional ideal, supported by a suitable normality criterion. The latter must ensure that the endomorphism ring is strictly larger than A exactly if A is not normal. For a reduced affine K-algebra A, where K is a perfect field, two approaches of this type have proven to work:

Vasconcelos' algorithm [Vas91] (see also [Vas98, §6]) is based on a regularity criterion of Lipman (see [Lip69]). It first replaces A by its  $(S_2)$ -ification  $\operatorname{End}_A(\omega_A)$ , which is computed via a Noether normalization, and then iteratively by  $\operatorname{End}_A(J_A^{\vee})$ , where  $J_A$  is the Jacobian ideal of A.

De Jong's algorithm [dJ98, DdJGP99] is based on the normality criterion of Grauert and Remmert [GR71, Anhang, §3.3, Satz 7], see [GLS10] for a historical account. Here, the idea is to choose a so-called *test ideal* J (for example, the radical of the Jacobian ideal), and to replace A by End<sub>A</sub>(J), repeating the procedure if necessary. A variant of this algorithm with improved efficiency is described in [GLS10];

Date: April 22, 2022.

<sup>2010</sup> Mathematics Subject Classification. 13B22 (Primary) 13H10, 14H20, 13P10 (Secondary). Key words and phrases. Normalization, integral closure, Grauert–Remmert criterion, curve singularity, simple singularity.

The research leading to these results has received funding from the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme (FP7/2007-2013) under REA grant agreement n° PCIG12-GA-2012-334355.

we refer to this version as the *GLS normalization algorithm*. The performance of the GLS algorithm can be further enhanced by applying it to suitable strata of the singular locus and combining the individual results, see  $[BDL^+13]$ . This technique is particularly useful for parallel computing. The parallel GLS algorithm is the fastest normalization algorithm known to date. An implementation is available in the computer algebra system SINGULAR [DGPS13].

For further progress in algorithmic normalization, a deeper theoretical understanding of endomorphism rings of fractional ideals will be useful. The following natural questions arise:

How do properties of and relations between fractional ideals affect the associated endomorphism ring? The aspect of products of ideals is briefly discussed in [dJ98, p. 275].

More specifically, how does the choice of fractional ideal affect the number of iterations to reach the normalization? In [PV08], an analysis of the complexity of general normalization algorithms is given in the graded case. In [BEGvB09], examples of free discriminants of versal families are given where the normalization is obtained in one step.

How to measure and bound the progress of each iteration? In [GS12], a minimal step size is related to quasihomogeneity for Gorenstein curves.

What kind of endomorphism rings occur in the process of the algorithms? In the case of finite Coxeter arrangements and their discriminants, this question is studied in [GMS12].

In the curve case, how does the normalization algorithm relate to the resolution of singularities by blowups?

From a more practical point of view, which choice of fractional ideal leads to the best overall performance? Is it better to normalize by many small steps or by a few large steps?

In this paper, we focus on the GLS algorithm and study the number of iterations in terms of localization and completion of the given ring and with respect to the inclusion of test ideals.

In Section 1, we introduce the objects and operations relevant to algorithmic normalization and notice that they are compatible with localization. To obtain similar results for completion, we consider the class of excellent semilocal rings. For such rings, normalization is finite and commutes with completion.

Based on these preparations, in Section 2, we show that the GLS algorithm behaves well with respect to localization and completion. That is, these operations preserve the property of being a test ideal and commute with forming endomorphism rings. By considering test ideals relative to a subset of the spectrum of A, we prepare the ground for the study of stratified normalization in the next section.

In Section 3, we stratify the singular locus of A and consider test ideals relative to the individual strata. We show that the number of steps of the global algorithm (using the radical ideal of the singular locus) is at most the maximal number of steps among the strata. We prove equality in the case where A is equidimensional and satisfies Serre's condition  $(S_2)$ . Being Cohen–Macaulay, local complete intersections or curves satisfy these conditions. As a side result, in the equidimensional case, we show that the GLS algorithm preserves  $(S_2)$  at each step.

In Section 4, we prove formulas for the number of steps of the GLS algorithm for plane curve singularities of type ADE<sup>1</sup>. Even more, we explicitly determine the singularity types that occur in the process, describing the respective value semigroups on our way. Besides other instances of ADE singularities, we find simple space curve singularities from the list of Frühbis-Krüger [FK99].

<sup>&</sup>lt;sup>1</sup>Based on experiments using SINGULAR, these formulas are also given in [Mor13].

Acknowledgements. We thank Gert-Martin Greuel and Gerhard Pfister for helpful discussions.

## 1. NORMALIZATION AND COMPLETION

In the following, all rings are commutative with one and  $\mathbb{N}$  includes 0.

Let A be a reduced Noetherian ring. We write Q(A) for the total ring of fractions of A, which is again a reduced Noetherian ring.

**Definition 1.1.** The normalization of A, written  $\overline{A}$ , is the integral closure of A in Q(A). We call A normalization-finite if  $\overline{A}$  is a finite A-module, and we call A normal if  $A = \overline{A}$ . We denote by

$$N(A) = \{ P \in \text{Spec}(A) \mid A_P \text{ is not normal} \}$$

the non-normal locus of A, and by

$$\operatorname{Sing}(A) = \{ P \in \operatorname{Spec}(A) \mid A_P \text{ is not regular} \}$$

the singular locus of A.

Remark 1.2. Note that  $N(A) \subset \text{Sing}(A)$ . Equality holds if A is of pure dimension one. Indeed, a Noetherian local ring of dimension one is normal if and only if it is regular (see [dJP00, Thm. 4.4.9]).

*Remark* 1.3. Recall that a ring is reduced if and only if it satisfies Serre's conditions  $(R_0)$  and  $(S_1)$ ; it is normal if and only if satisfies  $(R_1)$  and  $(S_2)$  (see [HS06, §4.5]).

**Definition 1.4.** The conductor of A in  $\overline{A}$  is  $C_A = \operatorname{Ann}_A(\overline{A}/A)$ .

Remark 1.5. Note that  $C_A$  is the largest ideal of A which is also an ideal of  $\overline{A}$ . In particular, A is normal if and only if  $C_A = A$ .

**Definition 1.6.** If M is an A-module, we write  $M^{-1} = \text{Hom}_A(M, A)$  for the *dual* module, and call M reflexive if the canonical map  $M \to (M^{-1})^{-1}$  is an isomorphism.

Remark 1.7. Any map  $\varphi \in \overline{A}^{-1} = \operatorname{Hom}_A(\overline{A}, A)$  is multiplication by  $\varphi(1) \in A$ . We may, thus, identify  $\overline{A}^{-1} = \mathcal{C}_A$  by taking  $\varphi$  to  $\varphi(1)$ .

If A is normalization-finite, satisfies  $(S_2)$ , and is Gorenstein in codimension 1, then  $C_A$  is reflexive (see [GS11, Lem. 2.8]).

Notation 1.8. If I is an ideal of a ring R, we write  $V(I) = \{P \in \text{Spec}(R) \mid P \supseteq I\}$  for the vanishing locus of I in Spec(R).

Proposition 1.9. As above, let A be a reduced Noetherian ring. Then:

- (1) A is normalization-finite if and only if  $C_A$  contains a non-zerodivisor.
- (2)  $N(A) \subseteq V(\mathcal{C}_A)$ , with equality if A is normalization-finite.
- (3) If A is normalization-finite, then  $(\mathcal{C}_A)_P = \mathcal{C}_{A_P}$  for all  $P \in \operatorname{Spec}(A)$ .

*Proof.* (1) Note that a common denominator of a finite set of generators for  $\overline{A}$  over A is a non-zerodivisor in  $\mathcal{C}_A$ .

(2) See [GP08, Lem. 3.6.3].

(3) See [ZS75, Ch. V, §5].

By part (1) of the preceding proposition, if A is normalization-finite, then A is a fractional ideal in the following sense:

**Definition 1.10.** A *fractional ideal* of A is a finite A-submodule of Q(A) containing a non-zerodivisor of A.

**Lemma 1.11.** Let M and N be fractional ideals of A. Then, independently of the choice of a non-zerodivisor  $g \in M$  of A, we may identify  $\operatorname{Hom}_A(M, N)$  with a fractional ideal of A by means of

$$\operatorname{Hom}_A(M, N) \hookrightarrow Q(A), \quad \varphi \mapsto \varphi(g)/g.$$

In particular, any  $\varphi \in \operatorname{Hom}_A(M, N)$  is multiplication by  $\varphi(g)/g$ .

Proof. See, for example, [GLS10, Lem. 3.1].

Our applications will be geometric in nature: The rings under discussion will be finite K-algebras, where K is a field, localizations of such algebras, the semi-local rings appearing in the normalization process of a localization, or the rings obtained by completing one of the semi-local rings. As we recall in what follows, all these rings are normalization-finite.

Notation 1.12. The Jacobson radical of a ring R will be denoted by  $\mathfrak{m}_R$ .

**Theorem 1.13.** Let R be a Noetherian ring, and let  $\hat{}$  be completion at an ideal I of R. Then  $\hat{R}$  is Noetherian, and we have:

- (1) If M is a finite R-module, then the natural map  $M \otimes_R \widehat{R} \to \widehat{M}$  is an isomorphism, and  $\widehat{R}$  is flat over R.
- (2) If I is contained in  $\mathfrak{m}_R$ , then the natural map  $R \to \widehat{R}$  is an inclusion, and  $\widehat{R}$  is faithfully flat over R.

*Proof.* See [Bou98, Ch. III, §3.4 Thm. 3, Props. 8, 9]. Note that in case (2), R together with the *I*-adic topology is a Zariski ring.

Remark 1.14. If R is a semi-local Noetherian ring, then  $\widehat{}$  will always stand for the completion at  $\mathfrak{m}_R$ . In this case,  $\widehat{R}$  is again a semi-local Noetherian ring, and if  $R \subset R'$  is a finite ring extension, then R' is semi-local and Noetherian as well, and the completion of any R'-module M' at  $\mathfrak{m}_{R'}$  coincides with that at  $\mathfrak{m}_R$  if M' is regarded as an R-module. See [Bou98, Ch. III, §3.1, Cor.; Ch. IV, §2.5, Cor. 3] for details.

Grothendiecks's notion of an *excellent ring* provides the general framework for the rings considered here. Referring to [Mat80] for the defining properties of an excellent ring (which include that the ring is Noetherian), we recall a number of consequences of these properties.

**Theorem 1.15** (Grothendieck). Let R be an excellent ring. Then all localizations of R and all finitely generated R-algebras are excellent.

*Proof.* See [Mat80, Thms. 73, 77].

Remark 1.16. Complete semi-local Noetherian rings are excellent (for a proof see [Mat80, (28.P); Thms. 68, 74]). In particular, any field K and, hence, any localization of any finitely generated K-algebra are excellent.

**Theorem 1.17** (Grothendieck). Let A be a reduced excellent ring. Then:

- (1) The completion  $\widehat{A}$  of A at any ideal of A is reduced. If A is normal, then  $\widehat{A}$  is normal.
- (2) A is normalization-finite.
- (3) If A is semi-local, then  $\overline{\widehat{A}} = \overline{\widehat{A}}$ . In particular,
  - (a)  $\widehat{A} = \overline{A} = \overline{A} \otimes_A \widehat{A}$  is a finite  $\widehat{A}$ -module, and
  - (b) if A is complete, then  $\overline{A}$  is complete.

*Proof.* For (1), see [Mat80, Thm. 79].

For (2), note that excellent rings are Nagata rings (see [Mat80, Thm. 78]). Hence, if P is any prime ideal of A, then A/P is normalization-finite, so that  $\overline{A/P}$  is in particular finite over A. The result follows from the *splitting of normalization* (see [dJP00, Thm. 1.5.20]): If  $P_1, \ldots, P_s$  are the minimal primes of R, then

$$\overline{A} \cong \prod_{i=1}^{s} \overline{A/P_i}.$$

For (3), note that  $\overline{A}$  is finite over A by (2) and, hence, an excellent ring by Theorem 1.15. We conclude from (1) that  $\widehat{\overline{A}}$  is normal. Now consider the inclusions  $A \subseteq \overline{A} \subseteq Q(A)$  which by the flatness of completion give rise to the inclusions  $\widehat{A} \subseteq \widehat{\overline{A}} \subseteq Q(A) \otimes_A \widehat{A}$ , where  $\widehat{\overline{A}}$  is finite over  $\widehat{A}$ . Since every non-zerodivisor of A is a non-zerodivisor of  $\widehat{A}$ , we may regard  $Q(A) \otimes_A \widehat{A}$  as a subring of  $Q(\widehat{A})$ . Since  $\widehat{\overline{A}}$ is normal, we conclude that  $\overline{\widehat{A}} = \widehat{\overline{A}}$ .

**Corollary 1.18.** If A is a Noetherian complete local domain, then  $\overline{A}$  is a Noetherian complete local domain.

*Proof.* It is clear from Remark 1.16 that A is excellent. Hence, by Theorem 1.17, A is normalization-finite and  $\overline{A}$  is complete. Taking Remark 1.14 into account, we conclude that  $\overline{A}$  is a Noetherian complete semi-local ring and, thus, a product of local rings (see [Bou98, Ch. III, §2.13, Cor.]). Since  $\overline{A}$  is also a domain, it must be a local ring.

**Corollary 1.19.** If J is an ideal of an excellent ring R, and  $\hat{}$  is completion at an arbitrary ideal of R, then

$$\sqrt{\widehat{J}} = \widehat{\sqrt{J}}.$$

In particular, if J is radical, then  $\widehat{J}$  is radical as well.

Proof. By the flatness of completion, we first deduce from  $J \subseteq \sqrt{J}$  that  $\widehat{J} \subseteq \sqrt{\overline{J}}$ , and second from Theorem 1.17.(1) that  $\sqrt{J}$  is radical. Thus,  $\sqrt{\widehat{J}} \subseteq \sqrt{\overline{J}}$ . On the other hand, there is an m with  $(\sqrt{J})^m \subseteq J$ . Again by flatness, this implies that  $(\sqrt{J})^m \subseteq \widehat{J}$ , that is,  $\sqrt{\overline{J}} \subseteq \sqrt{\widehat{J}}$ .

**Proposition 1.20.** Suppose  $R \to S$  is a flat homomorphism of rings, M and N are R-modules, and M is finitely generated. Then there is a unique S-module isomorphism

$$\operatorname{Hom}_R(M, N) \otimes_R S \cong \operatorname{Hom}_S(M \otimes_R S, N \otimes_R S)$$

which takes  $\varphi \otimes 1$  to  $\varphi \otimes \mathrm{id}_S$ .

Proof. See [Eis95, Prop. 2.10].

**Corollary 1.21.** If A is a reduced excellent semi-local ring, then

$$\widehat{\mathcal{C}_A} = \mathcal{C}_{\widehat{A}}$$

*Proof.* Use Remark 1.7, Theorem 1.13.(1), Proposition 1.20, and Theorem 1.17.(3).  $\Box$ 

#### 2. Algorithmic Normalization

Throughout this section, A denotes a reduced excellent ring. In particular, A is normalization-finite.

**Definition 2.1.** A *test ideal* at a subset  $W \subseteq \text{Spec}(A)$  is an ideal  $J \subseteq A$  satisfying the following conditions:

(1) J contains a non-zerodivisor of A,

(2) J is a radical ideal, and

(3)  $V(\mathcal{C}_{A_P}) \subseteq V(J_P)$  for all  $P \in W$ .

A test ideal for A is an ideal  $J \subseteq A$  satisfying (1), (2), and

(3')  $V(\mathcal{C}_A) \subseteq V(J).$ 

Remark 2.2. Since we assume that A is normalization-finite,

$$V(\mathcal{C}_A) = N(A) \subseteq \operatorname{Sing}(A).$$

Hence the vanishing ideal J of Sing(A) is a valid test ideal for A. Indeed, J contains a non-zerodivisor since A is reduced and, hence, regular in codimension zero, so that  $J \otimes_A Q(A) = Q(A)$ .

In what follows, if  $J \subseteq A$  is any ideal containing a non-zerodivisor of A, we regard A as a subring of  $\operatorname{End}_A(J) = \operatorname{Hom}_A(J, J)$  by sending  $a \in A$  to multiplication by a, and  $\operatorname{End}_A(J)$  as a fractional ideal of A as in Lemma 1.11. Then

$$A \subseteq \operatorname{End}_A(J) \subseteq \overline{A} \tag{2.1}$$

(see [GP08, Lem. 3.6.1]).

**Proposition 2.3** (Grauert and Remmert Criterion). Let J be a test ideal for A. Then A is normal if and only if  $A = \text{End}_A(J)$ .

*Proof.* See [GR71], [GP08, Prop. 3.6.5].

Before describing the normalization algorithm arising from the Grauert and Remmert criterion, we discuss how the criterion behaves with respect to localization and completion. We first address the test ideals:

**Lemma 2.4.** Let  $J \subset A$  be an ideal, and let  $W \subset \text{Spec}(A)$ . Then:

- (1) If J is a test ideal at W, and  $P \in W$ , then  $J_P$  is a test ideal for  $A_P$ .
- (2) If  $W \supseteq N(A)$ , then J is a test ideal for A if and only if it is a test ideal at W.

Proof. Condition (3) of Definition 2.1 means that condition (3') of the definition holds for the rings  $A_P$  together with the ideals  $J_P$ ,  $P \in W$ . The first statement of the lemma follows since conditions (1) and (2) of Definition 2.1 carry over from Jto  $J_P$ : Use the flatness of localization and that localization commutes with passing to radicals, respectively. Taking into account that  $(\mathcal{C}_A)_P = \mathcal{C}_{A_P}$  since A is assumed normalization-finite (see Proposition 1.9.(3)), the same reasoning shows the second statement of the lemma. Indeed,  $J \subseteq \sqrt{\mathcal{C}_A}$  if and only if  $J_P \subseteq (\sqrt{\mathcal{C}_A})_P$  for all  $P \in \text{Spec}(A)$ , and  $\mathcal{C}_P = A_P$  if  $P \notin N(A)$ .

**Lemma 2.5.** If A is semi-local and J is a test ideal for A, then  $\widehat{J}$  is a test ideal for  $\widehat{A}$ .

*Proof.* If  $g \in J$  is a non-zerodivisor of A, then  $g \otimes_A 1 \in J \otimes_A \widehat{A} = \widehat{J}$  is a non-zerodivisor of  $\widehat{A}$  by the flatness of completion. Moreover, by Corollary 1.19,  $\widehat{J}$  is radical and  $\widehat{J} \subseteq \widehat{\sqrt{\mathcal{C}_A}} = \sqrt{\widehat{\mathcal{C}_A}} = \sqrt{\mathcal{C}_{\widehat{A}}}$ , where we use the assumption  $J \subseteq \sqrt{\mathcal{C}_A}$ , and where the last equality holds by Corollary 1.21.

Next, in the two corollaries of Proposition 1.20 below, we treat the endomorphism rings appearing in the Grauert and Remmert criterion.

Remark 2.6. As usual, if P is a prime of a ring R, and M is an R-module, we write  $M_P$  for the localization of M at  $R \setminus P$ . Recall that if  $R \subseteq R'$  is a ring extension, and M' is an R'-module, then the localization of M' at  $R \setminus P \subset R'$  coincides with  $M'_P$  if M' is considered as an R-module.

**Corollary 2.7.** Let J be an ideal of A. Then, for all  $P \in \text{Spec}(A)$ ,

$$(\operatorname{End}_A(J))_P = \operatorname{End}_{A_P}(J_P).$$

Further,  $A = \operatorname{End}_A(J)$  if and only if  $A_P = \operatorname{End}_{A_P}(J_P)$  for all  $P \in \operatorname{Spec}(A)$ .

*Proof.* Apply Proposition 1.20 and use that equality is a local property.

Corollary 2.8. If A is semi-local, and J is an ideal of A, then

$$\operatorname{End}_{A}(J) = \operatorname{End}_{\widehat{A}}(\widehat{J}),$$

and  $A = \operatorname{End}_A(J)$  if and only if  $\widehat{A} = \operatorname{End}_{\widehat{A}}(\widehat{J})$ .

*Proof.* Apply Proposition 1.20 and use that A is faithfully flat over A.

The Grauert and Remmert criterion allows us to compute  $\overline{A}$  by successively enlarging the given ring by an endomorphism ring. Here, we need:

**Lemma 2.9.** Let  $A \subseteq A' \subseteq \overline{A}$  be an intermediate ring, and let J be a test ideal for A. Then  $\sqrt{JA'}$  is a test ideal for A'.

*Proof.* Write  $J' := \sqrt{JA'}$ . If  $g \in J$  is a non-zerodivisor of A, then  $g \in J \subseteq JA' \subseteq J'$  is a non-zerodivisor of Q(A) and, hence, of A'. Since  $J \subseteq \sqrt{\mathcal{C}_A}$  by assumption, and  $\mathcal{C}_A \subseteq \mathcal{C}_{A'}$ , we have  $J' \subseteq \sqrt{\mathcal{C}_A A'} \subseteq \sqrt{\mathcal{C}_{A'}}$ .

Given any radical ideal  $J \subseteq A$  containing a non-zerodivisor, we inductively define radical ideals and intermediate rings by setting  $A_0 = A$ ,

$$J_i = \sqrt{JA_i}$$
, and  $A_{i+1} = \operatorname{End}_{A_i}(J_i) \subseteq Q(A)$ .

Here, with A, also Q(A) and, hence, all the  $A_i$  are reduced. Since we assume that A is Noetherian and normalization-finite, we get, thus, a finite chain of extensions of reduced Noetherian rings

$$A = A_0 \subsetneqq \cdots \subsetneqq A_{i-1} \gneqq A_i \gneqq \cdots \gneqq A_n = A_{n+1} \subseteq \overline{A}.$$

Notation 2.10. We write n(A, J) = n for the number of steps above.

Note that if J is a test ideal for A, then each  $J_i$  is a test ideal for  $A_i$ , so that  $A_n = \overline{A}$  by the Grauert and Remmert criterion. More generally, by the proof of [BDL<sup>+</sup>13, Prop. 3.3], we have:

**Proposition 2.11.** Let  $A \subseteq A' \subseteq \overline{A}$  be an intermediate ring. Let  $W \subseteq \text{Spec}(A)$ , let J be a test ideal at W, and let  $J' = \sqrt{JA'}$ . If

$$A' \cong \operatorname{End}_{A'}(J'),$$

then  $A'_P$  is normal for each  $P \in W$ .

We now relate the behaviour of the global version of the normalization algorithm to that of its local version:

**Proposition 2.12.** Let  $J \subseteq A$  be a radical ideal containing a non-zerodivisor of A, and let  $W \subseteq \text{Spec}(A)$ . Then

$$n(A, J) \ge \max_{P \in W} n(A_P, J_P)$$
$$= \max_{P \in W} n\left(\widehat{A_P}, \widehat{J_P}\right).$$

Equality holds if either  $V(J) \subseteq W$  or  $N(A) \subseteq W$ .

*Proof.* Let  $P \in \text{Spec}(A)$ . If  $P \notin V(J)$ , then  $J_P = A_P$ , so  $n(A_P, J_P) = 0$ . Furthermore, as is clear from (2.1), this number is also zero if  $P \notin N(A)$ . Hence, all statements of the proposition will follow once we show that

- $n(A, J) = \max_{P \in \operatorname{Spec}(A)} n(A_P, J_P)$ , and
- $n(A_P, J_P) = n\left(\widehat{A_P}, \widehat{J_P}\right)$  for all  $P \in \operatorname{Spec}(A)$ .

To establish these equalities, given  $P \in \text{Spec}(A)$ , we inductively set  $B_0 = A_P$ ,  $C_0 = \widehat{A_P}$ ,

$$B_{i+1} = \operatorname{End}_{B_i}\left(\sqrt{J_P B_i}\right)$$
, and  $C_{i+1} = \operatorname{End}_{C_i}\left(\sqrt{\widehat{J_P C_i}}\right)$ 

Since equality is a local property, and completion is faithfully flat in the semi-local case, it is enough to show that  $B_i = (A_i)_P$  and  $C_i = \widehat{B_i}$  for all i and P. For this, we do induction, assuming that our claim is true for i: We first note that

$$J_P(A_i)_P = (JA_i)_P$$
 and  $\widehat{J_PB_i} = \widehat{J_PB_i}$ .

Then, applying Corollary 2.7, we get

$$B_{i+1} = \operatorname{End}_{(A_i)_P} \left( \sqrt{J_P(A_i)_P} \right) = \operatorname{End}_{(A_i)_P} \left( \left( \sqrt{JA_i} \right)_P \right) = (A_{i+1})_P,$$

and Corollaries 1.19 and 2.8 give

$$C_{i+1} = \operatorname{End}_{\widehat{B_i}}\left(\sqrt{\widehat{J_P}\widehat{B_i}}\right) = \operatorname{End}_{\widehat{B_i}}\left(\widehat{\sqrt{J_P}B_i}\right) = \widehat{B_{i+1}}.$$

In the proposition, the maximum exists even though W may be infinite. On the other hand, if  $\operatorname{Sing}(A)$  is finite, then n(A, J) can be read off from just finitely many values  $n(A_P, J_P) = n(\widehat{A_P}, \widehat{J_P})$ . To obtain some sort of general analogue of this fact, we discuss a convenient stratification of  $\operatorname{Sing}(A)$ .

#### 3. Bounds for Stratified Normalization

Let again A be a reduced excellent ring. If  $P \in \text{Sing}(A)$ , set

$$L_P = \bigcap_{P \supseteq \widetilde{P} \in \operatorname{Sing}(A)} \widetilde{P}.$$

We stratify  $\operatorname{Sing}(A)$  according to the values of the function  $P \mapsto L_P$ . That is, if

 $\mathcal{L} = \{ L_P \mid P \in \operatorname{Sing}(A) \}$ 

denotes the set of all possible values, then the strata are the sets

$$W_L = \{ P \in \operatorname{Sing}(A) \mid L_P = L \}, \ L \in \mathcal{L}.$$

We write  $\text{Strata}(A) = \{W_L \mid L \in \mathcal{L}\}\$  for the set of all strata. This is a finite set. By construction, the singular locus is the disjoint union of the strata. For  $W \in \text{Strata}(A)$ , write  $L_W$  for the constant value of  $P \mapsto L_P$  on W.

**Lemma 3.1.** If  $W \in \text{Strata}(A)$ , then  $L_W$  is a test ideal at W.

*Proof.* By construction,  $L_W$  is radical. If  $J = \bigcap_{P \in \text{Sing}(A)} P$  is the vanishing ideal of Sing(A), then  $(L_W)_P = J_P \subseteq \sqrt{\mathcal{C}_{A_P}}$  for all  $P \in W$ : the equality holds by construction of  $L_W$ , and the inclusion since J is a test ideal for A (see Remark 2.2). From the latter, we also get that  $L_W$  contains a non-zerodivisor since  $J \subseteq L_W$ .  $\Box$ 

Considering the ideal  $J = L_W$  and proceeding as in the previous section, we obtain a chain of rings

$$A = A_0 \subsetneqq \cdots \subsetneqq A_{i-1} \gneqq A_i \gneqq \cdots \gneqq A_n = A_{n+1} \subseteq \overline{A},$$

where, by Lemma 3.1 and Proposition 2.11,  $(A_n)_P$  is normal and, hence, equal to  $\overline{(A_n)_P} = \overline{A_P}$  for all  $P \in W$ .

By considering all strata and combining the resulting rings, we get  $\overline{A}$ :

**Proposition 3.2** ([BDL<sup>+</sup>13]). Suppose  $\operatorname{Sing}(A) = \bigcup_{i=1}^{s} W_i$ . For  $i = 1, \ldots, s$ , let an intermediate ring  $A \subseteq A^{(i)} \subseteq \overline{A}$  be given such that  $(A^{(i)})_P = \overline{A_P}$  for each  $P \in W_i$ . Then

$$\sum_{i=1}^{s} A^{(i)} = \overline{A}.$$

We know from Remark 1.3 how the Serre conditions characterize reduced and normal rings, respectively. Since we will make explicit use of these conditions in what follows, we recall their definition:

**Definition 3.3.** Let R be a Noetherian ring, and let  $i \ge 0$  be an integer.

We say that R satisfies Serre's condition  $(R_i)$  if for all  $P \in \text{Spec}(R)$  with  $\dim R_P = \text{height } P \leq i, R_P$  is a regular local ring.

We say that R satisfies Serre's condition  $(S_i)$  if for all  $P \in \text{Spec}(R)$ ,

$$\operatorname{lepth} R_P \ge \min\{i, \dim R_P\}.$$

Notation 3.4. If R is a ring, and I is an ideal of R or an R-module, then Ass(I) denotes the set of associated ideals of I.

The following is well-known:

**Lemma 3.5.** Let R be a local ring with maximal ideal  $\mathfrak{m}_R$ . Then  $\operatorname{End}_R(\mathfrak{m}_R) = \operatorname{Hom}_R(\mathfrak{m}_R, R)$ .

*Proof.* Assuming the contrary, there is a surjection  $\mathfrak{m}_R \twoheadrightarrow R$  which splits as R is trivially projective. But then  $\mathfrak{m}_R = Rx \oplus I$  for a non-zerodivisor x of R and an ideal  $I \subset R$ , and  $xI \subseteq Rx \cap I = 0$  implies I = 0, a contradiction.

**Lemma 3.6.** Let  $J \subseteq A$  be any radical ideal containing a non-zerodivisor and assume that A is  $(S_2)$ . Then:

- (1) The ring  $\operatorname{End}_A(J)$  is  $(S_2)$ .
- (2) If  $A_P$  is regular for all  $P \in Ass(J)$  with height P = 1, then  $End_A(J) = A$ .

*Proof.* (1) Fix an arbitrary  $Q \in \text{Spec}(\text{End}_A(J))$  and let  $P = Q \cap A$ . Then, by [BH93, Prop. 1.2.10.(a)] and the proof of [AK70, III, Prop. 3.16]),

$$depth(End_A(J)_Q) \ge grade(P(End_A(J))_P, (End_A(J))_P)$$
$$= depth_{A_P}(End_{A_P}(J_P)).$$

On the other hand, since  $A_P \subseteq \operatorname{End}_A(J)_P$  is an integral ring extension,

$$\dim A_P = \dim(\operatorname{End}_A(J)_P) \ge \dim(\operatorname{End}_A(J)_Q).$$

Hence, by  $(S_2)$  for A, it is enough to show that

$$\operatorname{depth}_{A_P}(\operatorname{End}_{A_P}(J_P)) \ge \min\{2, \operatorname{depth} A_P\}.$$
(3.1)

We distinguish two cases.

If dim $(A_P/J_P) = 0$ , then  $J_P = \mathfrak{m}_P$ . Hence (3.1) follows from Lemma 3.5 and [BH93, Exc. 1.4.19].

If  $\dim(A_P/J_P) = 1$ , then

$$\operatorname{depth}(A_P/J_P) \ge 1 \tag{3.2}$$

by  $(S_1)$  for the reduced ring  $A_P/J_P$ . On the other hand, using again [BH93, Exc. 1.4.19], we get

$$\operatorname{depth}_{A_P}(\operatorname{End}_A(J_P)) \ge \min\{2, \operatorname{depth}_{J_P}\}.$$

To estimate depth  $J_P$ , we apply the Depth Lemma (see [BH93, Prop. 1.2.9]) to the short exact sequence

$$0 \to J_P \to A_P \to A_P / J_P \to 0$$

and obtain

$$\operatorname{depth} J_P \ge \min \{\operatorname{depth} A_P, \operatorname{depth} (A_P/J_P) + 1\}.$$

Combined with (3.2), this proves (3.1).

(2) Consider the exact sequence

$$0 \to A \to \operatorname{End}_A(J) \to B \to 0 \tag{3.3}$$

with cokernel B. We show that  $Ass(B) = \emptyset$  and, hence, that B = 0. For this, let  $P \in Spec(A)$ .

If  $P \notin V(J)$ , then  $J_P = A_P$ , so  $B_P = 0$ . If  $P \notin \operatorname{Sing}(A)$ , then  $A_P$  is normal and  $J_P \subseteq A_P = \mathcal{C}_{A_P}$  is a test ideal for  $A_P$ , so again  $B_P = 0$  by the Grauert and Remmert criterion. We conclude that if  $P \notin V(J) \cap \operatorname{Sing}(A)$ , then  $P \notin \operatorname{Ass}(B)$ .

If  $P \in V(J) \cap \text{Sing}(A)$ , then dim  $A_P \geq 2$  (otherwise, height  $P = \dim A_P = 1$ , so  $P \in \text{Ass}(J)$ ; by assumption,  $A_P$  would be regular, a contradiction). By (3.1) and  $(S_2)$  for A, this implies

$$\operatorname{depth}_{A_P}(\operatorname{End}_{A_P}(J_P)) \ge 2.$$

Localizing (3.3) at P and applying the Depth Lemma, this gives

$$\operatorname{depth}_{A_P}(B_P) \ge \min\{\operatorname{depth}_{A_P}(\operatorname{End}_{A_P}(J_P)), \operatorname{depth}(A_P) - 1\} \ge 1$$

using once more  $(S_2)$  for A. We conclude again that  $P \notin Ass(B)$ .

**Proposition 3.7.** Suppose A is equidimensional and satisfies  $(S_2)$ , and let J be a test ideal for A. If  $J' \subseteq A$  is a radical ideal with  $Ass(J') \subseteq Ass(J)$ , then

$$n(A, J') \le n(A, J).$$

Proof. If J' = A, then  $n(A, J') = 0 \le n(A, J)$ .

Now let  $J' \subsetneq A$ . Inductively, set  $A_0 = B_0 = A$ ,

$$A_{i+1} = \operatorname{End}_{A_i}\left(\sqrt{JA_i}\right)$$
, and  $B_{i+1} = \operatorname{End}_{B_i}\left(\sqrt{J'B_i}\right)$ .

By assumption and Lemma 3.6.(1), all rings  $A_i$ ,  $B_i$  satisfy  $(S_2)$ . Let n = n(A, J). Then  $A_n = A_{n+1} = \overline{A}$  by the Grauert and Remmert criterion, and we must show that  $B_n = B_{n+1}$ .

We use Lemma 3.6.(2). Since A is equidimensional and  $A \subset B_n$  is an integral extension, also  $B_n$  is equidimensional. Fix  $Q \in \operatorname{Ass}(\sqrt{J'B_n})$  with height Q = 1, and set  $P = Q \cap A$ . Then  $J' \subseteq P$  and height P = 1 by equidimensionality. We conclude that  $P \in \operatorname{Ass}(J') \subseteq \operatorname{Ass}(J)$  and, hence, that  $J'_P = J_P$  since J and J' are radical. From the proof of Proposition 2.12, it follows that

$$(A_i)_P = (B_i)_P$$
, and  $\sqrt{J_P(A_i)_P} = \sqrt{J'_P(B_i)_P}$ ,

for i = 0, ..., n. In particular,  $(B_n)_P = (A_n)_P = \overline{A_P}$  and, hence,  $(B_n)_Q$  are regular. This proves that the hypothesis of Lemma 3.6.(2) applied to  $\sqrt{J'B_n}$  and  $B_n$  is satisfied. Hence, as desired,

$$B_{n+1} = \operatorname{End}_{B_n}(\sqrt{J'B_n}) = B_n.$$

**Corollary 3.8.** Let J be the vanishing ideal of Sing(A). Then

$$n(A, J) \le \max_{W \in \operatorname{Strata}(A)} n(A, L_W).$$

If A is equidimensional and satisfies  $(S_2)$ , then equality holds.

*Proof.* Recall from the proof of Lemma 3.1 that  $J_P = (L_W)_P$  for all  $P \in W$ . Hence, by Proposition 2.12,

$$n(A, J) = \max_{\substack{P \in \operatorname{Sing}(A)}} n(A_P, J_P)$$
  
= 
$$\max_{\substack{W \in \operatorname{Strata}(A)}} \max_{\substack{P \in W}} n(A_P, (L_W)_P)$$
  
$$\leq \max_{\substack{W \in \operatorname{Strata}(A)}} n(A, L_W).$$

This gives the first statement of the corollary. The second statement follows from the first one and Proposition 3.7: If A satisfies  $(S_2)$ , then

$$n(A, L_W) \le n(A, J).$$

*Remark* 3.9. It would be interesting to know whether the assumptions of the second statement of the corollary can be weakened. On the other hand, in its present form, the statement applies already to interesting classes of examples such as local complete intersections.

## 4. Plane curves

In this section, by a curve we mean a reduced excellent ring A of pure dimension one. In particular, A is Noetherian and normalization-finite, and it has a finite singular locus, which coincides with its non-normal locus by Remark 1.2.

Remark 4.1. If A is local and non-normal, then there is a unique test ideal J for A, namely  $J = \mathfrak{m}_A$ . In this case, we write  $n(A) = n(A, \mathfrak{m}_A)$ .

With this notation, Proposition 2.12 reduces to:

**Corollary 4.2.** Let A be a non-normal curve. Then, if J is any test ideal for A, we have

$$n(A, J) = \max_{P \in \operatorname{Sing}(A)} n(\widehat{A_P}).$$

In particular, n(A) := n(A, J) does not depend on the choice of J.

If A is regular, we write n(A) = 0.

**Corollary 4.3.** Let A be a non-normal curve with  $\operatorname{Sing}(A) = \{P_1, \ldots, P_r\}$ . For each i, let a curve  $B_i$  and a prime  $Q_i \in \operatorname{Sing}(B_i)$  be given such that

$$\widehat{A_{P_i}} \cong \widehat{(B_i)_{Q_i}}.$$

Then

$$n(A) = \max_{i=1,\dots,r} n\left(B_i, Q_i\right).$$

*Proof.* We fix an index i and write  $B = B_i$  and  $Q = Q_i$ . If  $Q' \in \text{Sing}(B)$  is different from Q, then  $Q_{Q'} = B_{Q'}$ . Hence

$$n(B,Q) = \max_{Q' \in \text{Sing}(B)} n(B_{Q'}, Q_{Q'}) = n(B_Q, Q_Q) = n(\widehat{B_Q}),$$

and the claim follows by Corollary 4.2.

In the following, A will be an algebroid curve over  $K = \mathbb{C}$ , that is, a reduced complete local Noetherian K-algebra A of dimension one. Then A is excellent by Remark 1.16.

Notation 4.4. We write  $\mathfrak{m}_A$  for the maximal ideal of A and

$$A' = \operatorname{End}_A(\mathfrak{m}_A).$$

Then  $\mathfrak{m}_A$  is a test ideal for A by Remark 4.1 and  $A \subset A' = \mathfrak{m}_A^{-1}$  by Lemma 3.5.

Remark 4.5. Recall that an algebroid curve A is Gorenstein if and only if every fractional ideal of A is reflexive or, equivalently, if  $\dim_K(A'/A) = 1$  (see [HK71, p. 19, Bsp. (b)] and [Ber62, Satz 1]). It follows that each local ring of a plane curve singularity is Gorenstein (see [HK71, Satz 1.46.(d)]).

Let  $P_1, \ldots, P_s$  be the associated primes of A, and let  $A_i = A/P_i$ ,  $i = 1, \ldots, s$ , be the branches of A. We have inclusions

$$A \hookrightarrow \prod_{i=1}^{s} A_i \hookrightarrow \prod_{i=1}^{s} \overline{A_i} \cong \overline{A},$$

where, by Corollary 1.18 and Remark 1.2, each  $\overline{A_i}$  is a complete discrete valuation ring, say with valuation  $\nu_i \colon \overline{A_i} \to \mathbb{N} \cup \{\infty\}$  and uniformizing parameter  $t_i$ . Then  $\overline{A_i} \cong K[[t_i]]$ , and evaluating  $\nu_i$  at a power series means to take its order.

We write the elements of  $\overline{A}$  as  $a = (a_1, \ldots, a_s)$ , with all the  $a_i \in \overline{A_i}$ , and consider the valuation map

$$\nu \colon \overline{A} \to (\mathbb{N} \cup \{\infty\})^s, \ a \mapsto (\nu_1(a_1), \dots, \nu_s(a_s)).$$

For monomials in  $K[[t_1]] \times \cdots \times K[[t_s]] \cong \overline{A}$ , we use multi-index notation: If  $\alpha \in \mathbb{N}^s$ , we write  $\alpha_i$  for the *i*th component of  $\alpha$ , and  $t^{\alpha} = (t_1^{\alpha_1}, \ldots, t_s^{\alpha_s})$ .

**Definition 4.6.** The *semigroup* of a fractional ideal I of A is defined as

$$\Gamma_I = \{ (\nu_1(a_1), \dots, \nu_s(a_s)) \mid a = (a_1, \dots, a_s) \in I, \ a_i \neq 0 \text{ for all } i \} \subseteq \mathbb{N}^s.$$

We call I multigraded if it is invariant under the action

$$(K^*)^s \times \overline{A} \to \overline{A}, \ (\lambda, t^{\alpha}) \mapsto (\lambda_1^{\alpha_1} t_1^{\alpha_1}, \dots, \lambda_s^{\alpha_s} t_s^{\alpha_s})$$

corresponding to the choice of coordinates  $t_1, \ldots, t_s$ .

**Lemma 4.7.** If I is multigraded, then  $\alpha \in \Gamma_I$  implies that  $K^s \cdot t^{\alpha} \subseteq I$ .

*Proof.* Let  $\alpha \in \Gamma_I$ . Then there is an  $a = (a_1, \ldots, a_s) \in I$  with  $\nu(a) = \alpha$ . Using a standard Vandermonde determinant argument, we obtain

$$(0,\ldots,0,t_i^{\beta_i},0,\ldots,0) \in I + \mathfrak{m}_{\overline{A}}^N,$$

for all *i* and *N*, and for all monomials  $t_i^{\beta_i}$  occuring in  $a_i$ . Since  $\overline{A}$  is complete,  $\bigcap_N (I + \mathfrak{m}_{\overline{A}}^N) = I$ , so the claim follows.

Remark 4.8. Let I be a fractional ideal. Suppose that  $(0, \ldots, 0, t_i^{\beta_i}, 0, \ldots, 0) \in I$  for all i. Then  $K^s \cdot t^{\beta} \subseteq I$ .

*Remark* 4.9 (Properties of the Conductor). Recall that the conductor  $C_A$  is the largest ideal of A which is also an ideal of  $\overline{A} \cong \prod_{i=1}^{s} \overline{A_i}$ . It is, hence, generated by a monomial, say  $C_A = \langle t^{\gamma} \rangle = \langle t_1^{\gamma_1} \rangle \times \cdots \times \langle t_s^{\gamma_s} \rangle$ , where for each i,

$$\gamma_i = \min \left\{ \alpha_i \mid \alpha + \mathbb{N}^s \subseteq \Gamma_A \right\}.$$

In particular,  $C_A$  is multigraded, and it follows from Lemma 4.7 that if  $\alpha \in \Gamma_{C_A}$ , then  $K^s \cdot t^{\alpha} \subseteq C_A$ .

Notation 4.10. Set  $\tau = \gamma - \mathbf{1}$ , and for any  $\alpha \in \mathbb{Z}^s$ , write

$$\Delta(\alpha) = \bigcup_{i=1}^{s} \Delta_i(\alpha), \text{ where}$$
$$\Delta_i(\alpha) = \{\beta \in \mathbb{N}^s \mid \alpha_i = \beta_i \text{ and } \alpha_j < \beta_j \text{ if } j \neq i\}.$$

Note that for s = 1, we have  $\Delta(\alpha) = \{\alpha\}$ .

The following theorem generalizes a result of Kunz [Kun70] from irreducible to reducible algebroid curves:

**Theorem 4.11** ([DdlM88]). The algebroid curve A is Gorenstein if and only if for all  $\alpha \in \mathbb{Z}^s$ , the following symmetry condition is satisfied:

$$\alpha \in \Gamma_A \Leftrightarrow \Delta(\tau - \alpha) \cap \Gamma_A = \emptyset.$$

Lemma 4.12. Let A be any algebroid curve. Then:

- (1)  $\Delta(\tau) \cap \Gamma_A = \emptyset$ .
- (2)  $\Gamma_A \subset \{0\} \cup \Gamma_{\mathfrak{m}_{\overline{A}}}.$
- (3) Let A be Gorenstein with s branches. If s = 2, then  $\tau \in \Gamma_A$ . If  $s \ge 3$ , then  $\tau + \bigcup_{i=1}^s \mathbb{N}e_i \subseteq \Gamma_A$ .

Proof.

- (1) See [DdlM88, (1.9) Cor. (i)].
- (2) This follows from  $\mathfrak{m}_A \subset \mathfrak{m}_{\overline{A}}$ .
- (3) This follows from Theorem 4.11 using (2).

*Remark* 4.13. Let A be Gorenstein with  $s \ge 2$  branches. Taking  $\alpha = \tau \in \Gamma_A$  in Theorem 4.11, we get  $\Delta(\mathbf{0}) \cap \Gamma_A = \emptyset$ .

**Lemma 4.14.** If A is any algebroid curve, and  $I \subseteq J$  are fractional ideals of A, then I = J if and only if  $\Gamma_I = \Gamma_J$ .

Proof. There is only one direction to show: Suppose that  $\Gamma_I = \Gamma_J$ . Let  $g \in I$  be a non-zerodivisor of A. Since I = J if and only if  $(t^{\delta}/g)I = (t^{\delta}/g)J$ , we may assume that  $g = t^{\delta}$ . Now let  $b \in J$ . By adding a K-multiple of  $t^{\delta}$  to b, we may assume that  $\nu(b) \in \Gamma_J = \Gamma_I$ . Hence, there is an  $a \in I$  with  $\nu(a) = \nu(b)$ . If  $b \in C_A$ , then  $b \in I$ , and we are done. Otherwise, there is some j with  $\nu(b_j) < \delta_j$ . Choose a scalar  $c \in K$  with  $\nu(b_j - ca_j) > \nu(b_j)$ . Setting  $b^{(1)} = b - (c, \ldots, c) \cdot a$ , we have  $\nu(b^{(1)}) > \nu(b)$  with respect to the natural partial ordering on  $\mathbb{N}^s$ . Continuing in this way, after finitely many steps, we arrive at an element  $b^{(m)} \in C_A$ , and conclude that  $b \in I$ .

Recall that we write  $A' = \text{End}_A(\mathfrak{m}_A)$ , and that we think of this as a fractional ideal of A as in Lemma 1.11.

Lemma 4.15. 
$$\Gamma_{A'} \subseteq \bigcap_{m \in \mathfrak{m}_A} (\Gamma_A - \nu(m)).$$
  
*Proof.* If  $a' \in A'$  and  $m \in \mathfrak{m}_A$ , then  $a'm \in \mathfrak{m}_A \subseteq A.$ 

Proof. We have

$$(\overline{A} \cdot t^{\tau}) \cdot \mathfrak{m}_A \subseteq (\overline{A} \cdot t^{\gamma-1}) \cdot (\overline{A} \cdot t^1) = \overline{A} \cdot t^{\gamma} = \mathcal{C}_A \subseteq A,$$

so the result follows from Lemma 3.5.

In the following,  $A = K[[x, y]] = K[[X, Y]] / \langle f \rangle$  will be a plane algebroid curve with a singularity of type ADE.

**Proposition 4.17.** If A is of type  $A_n$  then A' is of type  $A_{n-2}$ . In particular,  $n(A) = \lfloor \frac{n+1}{2} \rfloor$ .

*Proof.* We may assume that  $f = X^2 - Y^{n+1}$ .

(1) Suppose n = 2k is even. Then f is irreducible,  $A \to K[[t]] = \overline{A}, x \mapsto t^{n+1}, y \mapsto t^2$  is the normalization, and  $\gamma = 2k$ . In accordance with Lemma 4.12.(1),  $t^{\tau} \notin A$ . On the other hand, by Lemma 4.16,  $t^{\tau} \in A'$ , and by Lemma 4.15,  $\Gamma_{A'} \subseteq \Gamma_A \cup \{\tau\} = \Gamma_{A+K\cdot t^{\tau}}$ . Hence, by Lemma 4.14,

$$A' = A + K \cdot t^{\tau} = K[[x', y]] \cong K[[X, Y]] / \langle X^2 - Y^{n-1} \rangle,$$

where  $x' = t^{\tau}$ . See Figure 1.





(2) Suppose n = 2k - 1 is odd. Then there are two branches,

$$f = (X - Y^k) \cdot (X + Y^k)$$

and  $A \to K[[t_1]] \times K[[t_2]] = \overline{A}, x \mapsto (t_1^k, -t_2^k), y \mapsto (t_1, t_2)$  is the normalization. The properties of the conductor discussed in Remark 4.9 allow us to determine  $\gamma$ : First,  $x = (t_1^k, -t_2^k) \in \mathcal{C}_A = \langle t^{\gamma} \rangle$  since, for all  $i, j \ge 0$ ,

$$2t_1^{k+i}, -2t_2^{k+j}) = y^i \left( x + y^k \right) + y^j \left( x - y^k \right) \in A;$$

then  $\gamma = (k, k)$  since  $\mathcal{C}_A$  is multigraded and  $(t_1^{k-1}, 0) \notin A$  and  $(0, t_2^{k-1}) \notin A$ .

Considering the powers  $y^j \in A$ , j = 0, ..., k - 1, we deduce from Theorem 4.11 that  $\Gamma_A$  is as shown in Figure 2 (cf. Lemma 4.12 and Remark 4.13 for  $\tau = \nu(y^{k-1})$  and  $\mathbf{0} = \nu(y^0)$ , respectively).

Applying Lemmas 4.16, 4.15, and 4.14 as in part (1) of the proof, we get  $C_{A'} = \langle t^{\gamma'} \rangle$ , where  $\gamma' = \tau = (k - 1, k - 1)$ , and

$$A' = A + \overline{A} \cdot t^{\tau}.$$

See again Figure 2. Setting  $x' = (t_1^{k-1}, -t_2^{k-1})$ , we have  $x' \in \mathcal{C}_{A'} \setminus A$  by Remark 4.9. As  $\dim_K(A'/A) = 1$  by Remark 4.5, it follows that

$$A' = A + K \cdot x' = K[[x', y]] \cong K[[X, Y]] / \langle X^2 - Y^{n-1} \rangle.$$

**Proposition 4.18.** Suppose A is of type  $D_n$ , where  $n \ge 4$ . Then A' is a simple, non-Gorenstein space curve singularity of type  $A_{n-3} \lor L$ , that is, a transversal union of an  $A_{n-3}$ -singularity and a line L as in [FK99, Tab. 2a]. Moreover, A'' is the disjoint union of an  $A_{n-5}$ -singularity and a line for  $n \ge 6$ , and it is smooth for n = 4, 5. In particular,  $n(A) = \lfloor \frac{n}{2} \rfloor$ .

14

	0—	-0-	-0-	-0-	-•-		•	•	•		-•			0—	-0-	-0-	•		•	•	•	•	••••	-•
А	Å	0 0 0	0 0 0	0 0 0	:	:	÷	÷	÷		ł			Å	0 0 0	0 0 0	:	:	:	:	:	:		ł
	6	0	0	0	٠	•	•	٠	٠	•••	•			þ	0	0	•	٠	•	٠	•	•	•••	•
	þ	0	0	0	•	•	•	٠	•	•••	•			þ	0	0	•	•	•	•	•	•	•••	•
	þ	0	0	0	•	•	•	٠	•	•••	•			þ	0	0	•	•	•	٠	•	•	•••	•
	þ	0	0	0	٠	•	•	٠	•	•••	•		$\Delta'$	þ	0	0	٠	٠	•	٠	•	•	•••	•
	þ	0	0	0	γ•	•	•	٠	٠	•••	•			þ	0	0	٠	٠	٠	•	•	•	•••	•
	þ	0	0 /	τ•	0	0	0	0	0		þ			þ	0	0	٠	٠	٠	٠	•	•	•••	•
	þ	0	•	0	0	0	0	0	0		¢			þ	0	•	0	0	0	0	0	0	000	þ
	þ	•	0	0	0	0	0	0	0		þ			þ	٠	0	0	0	0	0	0	0		þ
	-	_0_	_0_	_0_	_0_	_0_	_0_	_0_	_0_	•••	6			•	_0_	_0_	_0_	_0_	_0_	-0	-0	-0-	<b>&gt;</b> •••	9

FIGURE 2. Normalization steps for a singularity of type  $A_7$ .

*Proof.* We may assume that  $f = X (Y^2 - X^{n-2})$ .

(1) Suppose  $n = 2k + 3 \ge 5$  is odd. Then  $A \to K[[t_1]] \times K[[t_2]] = \overline{A}, x \mapsto (0, t_2^2), y \mapsto (t_1, t_2^{2k+1})$  is the normalization.

(A) Considering the elements  $x^j y^i = (0, t_2^{i(2k+1)+2j}) \in A$  for i = 0, 1 and  $j \ge 1$ , and  $y^i (y^2 - x^{2k+1}) = (t_1^{2+i}, 0) \in A$  for  $i \ge 0$ , we deduce from Remarks 4.8 and 4.9 that  $(t_1^2, t_2^{2k+2}) \in \mathcal{C}_A = \langle t^\gamma \rangle$ . Then  $\gamma = (2, 2k+2)$  by Remark 4.9 since  $(t_1, 0) \notin A$  and  $(0, t_2^{2k+1}) \notin A$ . Hence,  $\tau = (1, 2k+1) = \nu(y)$  and  $\Gamma_A \cap (\tau + \mathbb{N}^2)$  is of the form shown in Figure 3 by Lemma 4.12.(1).

Next, for  $i \ge 1$  and j = 1, ..., k, considering the elements  $x^j + y^i = (t_1^i, t_2^{2j} + t_2^{i(n-2)}) \in A$ , we get  $(i, 2j) \in \Gamma_A$ . Since  $y \equiv (t_1, 0) \mod \overline{A} \cdot (0, t_2^{2k+1})$ , we conclude from Remark 4.13 that  $\Gamma_A$  is as depicted in Figure 3.

(A') To determine A', we argue as in part (2) of the proof of the previous Proposition 4.17. This gives  $C_{A'} = \langle t^{\gamma'} \rangle$ , where  $\gamma' = (1, 2k)$ , and

 $A' = A + K \cdot z = K[[x, y, z]] \cong K[[X, Y, Z]]/I',$ 

where  $z = (t_1, -t_2^{2k+1})$  and

$$I' = \langle X, Y - Z \rangle \cap \langle Y + Z, Z^2 - X^{n-2} \rangle.$$

Indeed, for the latter, note that x, y, z satisfy the relations given by the generators of I', and that I' is a radical ideal with the right number of components of the right codimension. The coordinate change  $(X, Y, Z) \mapsto (X, Y+Z, -Y+Z)$  turns I' into the ideal of maximal minors of the matrix

$$egin{pmatrix} Z & Y & X^{n-3} \ 0 & X & Y \end{pmatrix}.$$

Hence, A' is of the claimed type. Note that  $(t_1, 0) = (y + z)/2 \in A'$  and  $(0, t_2^{2j}) \in A', j = 1, \ldots, k - 1$ , and hence  $\mathfrak{m}_{A'}$  is multigraded by Remark 4.8.

(A'') By Lemma 4.16,  $\overline{A} \cdot t^{\tau'} \subseteq A''$ . In particular,  $(1,0) \in A''$ . Applying again Remark 4.8, we see that  $\sum_{j=1}^{k-1} K^2 \cdot (1, t_2^{2j}) \subseteq A''$ . Furthermore, by the very definition of  $A'' = \operatorname{End}_{A'}(\mathfrak{m}_{A'}) \subset \overline{A}$ , and since  $\mathfrak{m}_{A'}$  is multigraded,  $\sum_{i=0}^{\infty} K^2 \cdot (t_1^i, 1) \subseteq A''$ . It follows that A'' is multigraded, with  $\mathcal{C}_{A''} = \langle t^{\gamma''} \rangle$ , where  $\gamma'' = (0, 2k - 2)$ , and with  $\Gamma_{A''}$  as in Figure 3. We conclude that A''admits a product decomposition

$$A'' = K[[t_1]] \times K[[x', y']],$$



FIGURE 3. Normalization steps for a singularity of type  $D_7$ .

where  $x' = t_2^2$  and  $y' = t_2^{2k-1}$ . Moreover,

$$K[[x',y']] \cong K[[X,Y]] / \langle Y^2 - X^{2k-1} \rangle$$

is of type  $A_{n-5}$  for  $n \ge 7$ , and it is smooth for n = 5. (2) Suppose n = 2k+2 is even. Then  $f = X(Y^2 - X^{n-2}) = X(Y - X^k)(Y + X^k)$ and  $A \to K[[t_1]] \times K[[t_2]] \times K[[t_3]] = \overline{A}, x \mapsto (0, t_2, t_3), y \mapsto (t_1, t_2^k, -t_3^k)$  is the normalization.

(A) For j = 1, ..., k-1 and  $i \ge 1$ , we have  $x^j + y^i = (t_1^i, t_2^j + t_2^{ki}, t_3^j + (-1)^i t_3^{ki})$ . Since  $y \equiv (t_1, 0, 0) \mod \overline{A} \cdot (0, t_2^k, t_3^k)$ , it follows that

$$\{\alpha \in \Gamma_A \mid \alpha_2 < k, \ \alpha_3 < k\} = \{\mathbf{0}\} \cup \bigcup_{i \ge 1, \ j=1,...,k-1} (i,j,j).$$

Next, set

$$a = y - x^{k} = (t_{1}, 0, -2t_{3}^{k})$$
  
$$b = y + x^{k} = (t_{1}, 2t_{2}^{k}, 0).$$

Then, for  $j \ge 0$ , we have

$$y^{2} - x^{2k} = (t_{1}^{2}, 0, 0),$$
  

$$x^{j+1} \cdot a = (0, 0, -2t_{3}^{k+j+1}),$$
  

$$x^{j+1} \cdot b = (0, 2t_{2}^{k+j+1}, 0).$$

Hence,  $(t_1^2, t_2^{k+1}, t_3^{k+1}) \in \mathcal{C}_A = \langle t^{\gamma} \rangle$  by Remarks 4.8 and 4.9. An easy argument using parts (2) and (3) of Lemma 4.12 shows that  $\gamma = (2, k+1, k+1)$ . We conclude from Theorem 4.11 that  $\Gamma_A$  is of the form shown in Figure 4.



FIGURE 4. Normalization steps for a singularity of type  $D_{10}$ .

(A') Determining A' as before, we get  $C_{A'} = \langle t^{\gamma'} \rangle$ , where  $\gamma' = \tau = (1, k, k)$ , and

 $A' = A + K \cdot z = K[[x, y, z]] \cong K[[X, Y, Z]]/I',$ 

where  $z = (0, t_2^k, -t_3^k)$  and

 $I' = \langle X, Z \rangle \cap \left\langle Y - Z, Z + X^k \right\rangle \cap \left\langle Y - Z, Z - Y^k \right\rangle.$ 

Indeed, x, y, z satisfy the relations given by the generators of I', and I' is a radical ideal with the right number of components of the right codimension. The coordinate change  $(X, Y, Z) \mapsto (X, Y - Z, Y)$  then leads to the claimed ideal of minors (see part (1) of the proof).

(A'') Multiplying (1,0,0) with the generators x, y, z of  $\mathfrak{m}_{A'}$ , we get either zero or  $(1,0,0) \cdot y = (t_1,0,0) = y - z \in \mathfrak{m}_{A'}$ . Hence,  $(1,0,0) \in A'' = \operatorname{End}_{A'}(\mathfrak{m}_{A'}) \subset \overline{A}$ . It follows from Lemmas 4.16 and 4.14 that

$$\mathbf{A}'' = A' + \overline{A} \cdot (1,0,0) + \overline{A} \cdot t^{\gamma''},$$

where  $\gamma'' = \tau' = (0, k - 1, k - 1)$  (for the inclusion from left to right, use again that  $A'' = \operatorname{End}_{A'}(\mathfrak{m}_{A'})$ ). Using once more Lemma 4.14, and that  $(1, 0, 0) \in A''$ , we see that A'' admits a product decomposition

$$A'' = K[[t_1]] \times K[[x', y']],$$

where  $x' = (t_2, t_3)$  and  $y' = (t_2^{k-1}, -t_3^{k-1})$ . Moreover,

$$K[[x',y']] \cong K[[X,Y]] / \langle Y^2 - X^{2k-2} \rangle$$

is of type  $A_{n-5}$  for  $n \ge 6$ , and it is smooth for n = 4.

**Proposition 4.19.** Suppose A is of type  $E_n$ , where n = 6, 7, 8. Then A' is a simple, non-Gorenstein space curve singularity of type  $E_n(1)$ , see [FK99, Tab. 2a]. Moreover, A'' is of type  $A_{n-6}$  for n = 7, 8, and it is smooth for n = 6. In particular,  $n(A) = \lfloor \frac{n-1}{2} \rfloor$ .

*Proof.* (1) Let n = 6. We may assume  $f = X^3 - Y^4$ . Then  $A \to K[[t]] = \overline{A}$ ,  $x \mapsto t^4, y \mapsto t^3$  is the normalization, and we have

$$\mathbf{A} = \left\langle 1, t^3, t^4 \right\rangle_K \oplus K[t] \cdot t^6.$$

Arguing as in the previous proofs, we get

$$A' = A + K \cdot z,$$

where  $z = \tau = t^5$ . Then

$$A' = K[[x, y, z]] \cong K[[X, Y, Z]]/I'$$

where I' is the ideal of minors of the matrix

$$\begin{pmatrix} Y & Z & X^2 \\ X & Y & Z \end{pmatrix}$$
.

This means that A' is of type  $E_6(1)$ . Since  $A' = K \cdot 1 \oplus K[t] \cdot t^3 = K \cdot 1 \oplus \mathfrak{m}_{A'}$ , we obtain that  $A'' = \operatorname{End}_{A'}(\mathfrak{m}_{A'}) = K[t]$ . See Figure 5.



FIGURE 5. Normalization steps for a singularity of type  $E_6$ .

(2) Let n = 7. We may assume  $f = X(X^2 - Y^3)$ . Then  $A \to K[[t_1, t_2]] = \overline{A}$ ,

Let n = t. We may assume  $f = X(X^2 - Y^2)$ . Then  $A \to K[[t_1, t_2]] = A$ ,  $x \mapsto (0, t_2^3), y \mapsto (t_1, t_2^2)$  is the normalization. Since  $x^i y^j = (0, t_2^{3i+2j}) \in A$  for  $i \ge 1$  and  $j \ge 0$ , and  $(y^3 - x^2)y^i = (t_1^{3+i}, 0) \in A$  for  $i \ge 0$ , we have  $(t_1^3, t_2^5) \in \mathcal{C}_A = \langle t^\gamma \rangle$  by Remarks 4.8 and 4.9. Then  $\gamma = (3, 5)$  by Remark 4.9 since  $(t_1^2, 0) \notin A$  and  $(0, t_2^4) \notin A$ . Considering the elements  $x + y^i = (t_1^i, t_2^3 + t_2^{2+i}) \in A$  for  $i \ge 2$  and  $y^2 \in A$ , parts (1) and (3) of Lemma 4.12 imply that  $\Gamma_A \cap ((2, 3) + \mathbb{N}^2)$  is of the form shown in Figure 6. Using Theorem 4.11 and  $y \in A$ .

shown in Figure 6. Using Theorem 4.11 and  $y \in A$ , we see that  $\Gamma_A \cap ((1,2) + \mathbb{N}^2)$ 



FIGURE 6. Normalization steps for a singularity of type  $E_7$ .

and, hence,  $\Gamma_A$  are as in the figure.

Now, as before, we get

$$A' = \langle 1, y \rangle_K \oplus \overline{A} \cdot (t_1^2, t_2^3) = A + K \cdot z,$$

where  $z = (t_1^2, 0)$  (see again Figure 6). So

$$A' = K[[x, y, z]] \cong K[[X, Y, Z]]/I',$$

where

$$I' = \langle XZ, Y^2Z - Z^2, Y^3 - X^2 - YZ \rangle.$$

After the coordinate change  $(X, Y, Z) \mapsto (Y, -X, Z + X^2)$ , this becomes the ideal of maximal minors of the matrix

$$\begin{pmatrix} Z+X^2 & Y & X \\ 0 & Z & Y \end{pmatrix}.$$

It follows that A' is of type  $E_7(1)$ . Since  $A' = \langle 1, y \rangle_K \oplus \overline{A} \cdot (t_1^2, t_2^3)$ , we must have  $A'' = K \cdot 1 \oplus \overline{A} \cdot (t_1, t_2)$ , so that A'' is of type  $A_1$ . (3) Let n = 8. We may assume  $f = X^3 - Y^5$ . Then  $A \to K[[t]] = \overline{A}, x \mapsto t^5$ ,

 $y \mapsto t^3$  is the normalization, and we have

$$A = \left\langle 1, t^3, t^5, t^6 \right\rangle_K \oplus K[t] \cdot t^8.$$

As before,

$$A' = A + K \cdot z,$$

where  $z = \tau = t^7$ . Then

$$A' = K[[x, y, z]] \cong K[[X, Y, Z]]/I'$$

where I' is the ideal of maximal minors of the matrix

$$\begin{pmatrix} X & Z & Y^3 \\ Y & X & Z \end{pmatrix}.$$

This means that A' is of type  $E_8(1)$ . Since  $A' = \langle 1, t^3 \rangle_K \oplus K[t] \cdot t^5$ , it follows that  $A'' = K \cdot 1 \oplus K[t] \cdot t^2$ . See Figure 7.



FIGURE 7. Normalization steps for a singularity of type  $E_8$ .

### References

- [AK70] Allen Altman and Steven Kleiman, Introduction to Grothendieck duality theory, Lecture Notes in Mathematics, Vol. 146, Springer-Verlag, Berlin, 1970. MR 0274461 (43 #224)
- Janko Böhm, Wolfram Decker, Santiago Laplagne, Gerhard Pfister, Andreas Steen- $[BDL^+13]$ paß, and Stefan Steidel, Parallel algorithms for normalization, J. Symbolic Comput. 51 (2013), 99–114. MR 3005784
- [BEGvB09] Ragnar-Olaf Buchweitz, Wolfgang Ebeling, and Hans-Christian Graf von Bothmer, Low-dimensional singularities with free divisors as discriminants, J. Algebraic Geom. 18 (2009), no. 2, 371-406. MR 2475818

- [Ber62] Robert Berger, Über eine Klasse unvergabelter lokaler Ringe, Math. Ann. 146 (1962), 98–102. MR 0150172 (27 #175)
- [BH93] Winfried Bruns and Jürgen Herzog, Cohen-Macaulay rings, Cambridge Studies in Advanced Mathematics, vol. 39, Cambridge University Press, Cambridge, 1993. MR MR1251956 (95h:13020)
- [Bou98] Nicolas Bourbaki, Commutative algebra. Chapters 1–7, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 1998, Translated from the French, Reprint of the 1989 English translation. MR 1727221 (2001g:13001)
- [DdJGP99] Wolfram Decker, Theo de Jong, Gert-Martin Greuel, and Gerhard Pfister, The normalization: a new algorithm, implementation and comparisons, Computational methods for representations of groups and algebras (Essen, 1997), Progr. Math., vol. 173, Birkhäuser, Basel, 1999, pp. 177–185. MR 1714609 (2001a:13047)
- [DdlM88] Félix Delgado de la Mata, Gorenstein curves and symmetry of the semigroup of values, Manuscripta Math. 61 (1988), no. 3, 285–296. MR 949819 (89h:14024)
- [DGPS13] Wolfram Decker, Gert-Martin Greuel, Gerhard Pfister, and Hans Schönemann, SIN-GULAR 3-1-6 — A computer algebra system for polynomial computations, 2013.
- [dJ98] Theo de Jong, An algorithm for computing the integral closure, J. Symbolic Comput. 26 (1998), no. 3, 273–277. MR 1633931 (99d:13007)
- [dJP00] Theo de Jong and Gerhard Pfister, Local analytic geometry, Advanced Lectures in Mathematics, Friedr. Vieweg & Sohn, Braunschweig, 2000, Basic theory and applications. MR 1760953 (2001c:32001)
- [Eis95] David Eisenbud, Commutative algebra, Graduate Texts in Mathematics, vol. 150, Springer-Verlag, New York, 1995, With a view toward algebraic geometry. MR 1322960 (97a:13001)
- [FK99] Anne Frühbis-Krüger, Classification of simple space curve singularities, Comm. Algebra 27 (1999), no. 8, 3993–4013. MR 1700205 (2000f:32037)
- [GLS10] Gert-Martin Greuel, Santiago Laplagne, and Frank Seelisch, Normalization of rings,
   J. Symbolic Comput. 45 (2010), no. 9, 887–901. MR 2661161 (2011m:13012)
- [GMS12] Michel Granger, David Mond, and Mathias Schulze, Partial normalizations of Coxeter arrangements and discriminants, Mosc. Math. J. 12 (2012), no. 2, 335–367, 460–461. MR 2978760
- [GP08] Gert-Martin Greuel and Gerhard Pfister, A SINGULAR introduction to commutative algebra, extended ed., Springer, Berlin, 2008, With contributions by Olaf Bachmann, Christoph Lossen and Hans Schönemann, With 1 CD-ROM (Windows, Macintosh and UNIX). MR 2363237 (2008j:13001)
- [GR71] H. Grauert and R. Remmert, Analytische Stellenalgebren, Springer-Verlag, Berlin, 1971, Unter Mitarbeit von O. Riemenschneider, Die Grundlehren der mathematischen Wissenschaften, Band 176. MR 0316742 (47 #5290)
- [GS11] Michel Granger and Mathias Schulze, Normal crossing properties of complex hypersurface singularities and logarithmic residues, arXiv.org 1109.2612 (2011), Submitted.
- [GS12] \_\_\_\_\_, Quasihomogeneity of curves and the Jacobian endomorphism ring, Submitted, 2012.
- [HK71] Jürgen Herzog and Ernst Kunz (eds.), Der kanonische Modul eines Cohen-Macaulay-Rings, Lecture Notes in Mathematics, Vol. 238, Springer-Verlag, Berlin, 1971, Seminar über die lokale Kohomologietheorie von Grothendieck, Universität Regensburg, Wintersemester 1970/1971. MR 0412177 (54 #304)
- [HS06] Craig Huneke and Irena Swanson, Integral closure of ideals, rings, and modules, London Mathematical Society Lecture Note Series, vol. 336, Cambridge University Press, Cambridge, 2006. MR 2266432 (2008m:13013)
- [Kun70] Ernst Kunz, The value-semigroup of a one-dimensional Gorenstein ring, Proc. Amer. Math. Soc. 25 (1970), 748–751. MR 0265353 (42 #263)
- [Lip69] Joseph Lipman, On the Jacobian ideal of the module of differentials, Proc. Amer. Math. Soc. 21 (1969), 422–426. MR 0237511 (38 #5793)
- [Mat80] Hideyuki Matsumura, Commutative algebra, second ed., Mathematics Lecture Note Series, vol. 56, Benjamin/Cummings Publishing Co., Inc., Reading, Mass., 1980. MR 575344 (82i:13003)
- [Mor13] Robert Moritz, The behaviour of the normalization algorithm for curves, 2013, TU Kaiserslautern.
- [PV08] Thuy Pham and Wolmer V. Vasconcelos, Complexity of the normalization of algebras, Math. Z. 258 (2008), no. 4, 729–743. MR 2369053 (2009):13005)
- [Vas91] Wolmer V. Vasconcelos, Computing the integral closure of an affine domain, Proc. Amer. Math. Soc. 113 (1991), no. 3, 633–638. MR 1055780 (92b:13013)

#### JANKO BÖHM, WOLFRAM DECKER, AND MATHIAS SCHULZE

- [Vas98] \_, Computational methods in commutative algebra and algebraic geometry, Algorithms and Computation in Mathematics, vol. 2, Springer-Verlag, Berlin, 1998, With chapters by David Eisenbud, Daniel R. Grayson, Jürgen Herzog and Michael Stillman. MR 1484973 (99c:13048)
- Oscar Zariski and Pierre Samuel, Commutative algebra. Vol. 1, Springer-Verlag, New [ZS75] York, 1975, With the cooperation of I. S. Cohen, Corrected reprinting of the 1958 edition, Graduate Texts in Mathematics, No. 28. MR 0384768 (52 #5641)

JANKO BÖHM, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KAISERSLAUTERN, ERWIN-Schrödinger-Str., 67663 Kaiserslautern, Germany  $E\text{-}mail\ address:$  boehm@mathematik.uni-kl.de

WOLFRAM DECKER, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KAISERSLAUTERN, ERWIN-Schrödinger-Str., 67663 Kaiserslautern, Germany  $E\text{-}mail\ address: \texttt{deckerQmathematik.uni-kl.de}$ 

MATHIAS SCHULZE, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KAISERSLAUTERN, ERWIN-Schrödinger-Str., 67663 Kaiserslautern, Germany

 $E\text{-}mail\ address: \texttt{mschulze@mathematik.uni-kl.de}$