LOCALLY FINITE *p*-GROUPS WITH A LEFT 3-ENGEL ELEMENT WHOSE NORMAL CLOSURE IS NOT NILPOTENT

ANASTASIA HADJIEVANGELOU, MARIALAURA NOCE, AND GUNNAR TRAUSTASON

ABSTRACT. For any odd prime p, we give an example of a locally finite p-group G containing a left 3-Engel element x where $\langle x \rangle^G$ is not nilpotent.

1. INTRODUCTION

Let G be a group. An element $a \in G$ is a left Engel element in G, if for each $x \in G$ there exists a non-negative integer n(x) such that

$$[x_{n(x)} a] = [[[x, \underline{a}], a], \dots, a]] = 1.$$

If n(x) is bounded above by n then we say that a is a left n-Engel element in G. It is straightforward to see that any element of the Hirsch-Plotkin radical HP(G) of G is a left Engel element and the converse is known to be true for some classes of groups, including solvable groups and finite groups (more generally groups satisfying the maximal condition on subgroups) [4,1]. The converse is however not true in general and this is the case even for bounded left Engel elements. In fact whereas one sees readily that a left 2-Engel element is always in the Hirsch-Plotkin radical this is still an open question for left 3-Engel elements. Recently there has been a breakthrough and in [7] it is shown that any left 3-Engel element of odd order is contained in HP(G). From [12] one also knows that in order to generalise this to left 3-Engel elements of any finite order it suffices to deal with elements of order 2.

It was observed by William Burnside [3] that every element in a group of exponent 3 is a left 2-Engel element and so the fact that every left 2-Engel element lies in the Hirsch-Plotkin radical can be seen as the underlying reason why groups of exponent 3 are locally finite. For groups of 2-power exponent there is a close link with left Engel elements. If G is a group of exponent 2^n then it is not difficult to see that any element a in G of order 2 is a left (n+1)-Engel element of G (see the introduction of [13] for details). For sufficiently large n we know that the variety of groups of exponent 2^n is not locally finite [6,8]. As a result one can see (for example in [13]) that it follows that for sufficiently large n we do not have in general that a left n-Engel element is contained in the Hirsch-Plotkin radical. Using the fact that groups of exponent 4 are locally finite [11], one can also see

²⁰¹⁰ Mathematics Subject Classification. 20F45, 20F40.

Key words and phrases. Engel elements, Lie algebras.

that if all left 4-Engel elements of a group G of exponent 8 are in HP(G) then G is locally finite.

Swapping the role of a and x in the definition of a left Engel element we get the notion of a right Engel element. Thus an element $a \in G$ is a right Engel element, if for each $x \in G$ there exists a non-negative integer n(x) such that

$$\left[a_{n(x)} x\right] = 1.$$

If n(x) is bounded above by n, we say that a is a right n-Engel element. By a classical result of Heineken [5] one knows that if a is a right n-Engel element in G then a^{-1} is a left (n + 1)-Engel element.

In [9] M. Newell proved that if a is a right 3-Engel element in G then $a \in HP(G)$ and in fact he proved the stronger result that $\langle a \rangle^G$ is nilpotent of class at most 3. The natural question arises whether the analogous result holds for left 3-Engel elements. In [10] it has been shown that this is not the case by giving an example of a locally finite 2-group with a left 3-Engel element a such that $\langle a \rangle^G$ is not nilpotent. In this paper we extend this result to include any odd prime. The odd case turns out to be more involved and the construction in many ways quite different from the p = 2 case.

The structure of the paper is as follows. In Section 2 we describe the pair (G, x) that will provide our example and we show directly that x is a left 3-Engel element in G. In order to show that $\langle x \rangle^G$ is not nilpotent we need more work. In Sections 3 and 4 we construct a pair (L, z) where L is a Lie algebra over \mathbb{F}_p , the field of p elements, and where $\mathrm{Id}(z)$ is not nilpotent. The pair (L, z)can be seen as the Lie algebra version of our group construction and in Section 5 we build a group H within $\mathrm{End}(L)$ containing $1 + \mathrm{ad}(z)$ where $(1 + \mathrm{ad}(z))^H$ is not nilpotent and where $(H, 1 + \mathrm{ad}(z))$ is a homomorphic image of (G, x).

2. The group G

Let x, a_2, a_2, \ldots be group variables. We recall that a simple commutator in x, a_1, a_2, \ldots is a group word defined recursively as follows: x, a_1, a_2, \ldots are simple commutators and if u, v are simple commutators then [u, v] is a simple commutator. We say that a simple commutator s has multi-weight (m, e_1, e_2, \ldots) in x, a_1, a_2, \ldots , if x occurs m times and a_i occurs e_i times in s. The weight of s is then $m + e_1 + e_2 + \cdots$. The following definition will also play a crucial role.

Definition. Let s be a simple commutator of multi-weight $(m, e_1, e_2, ...)$ in $x, a_1, a_2, ...$ The type of s is $t(s) = e_1 + e_2 + ... - 2m$.

Remark. If u, v are simple commutators in x, a_1, a_2, \ldots then t([u, v]) = t(u) + t(v). In particular $t([u, a_j]) = t(u) + 1$ and t([u, x]) = t(u) - 2.

For a fixed odd prime p, let $G = \langle x, a_1, a_2, \dots \rangle$ be the largest group satisfying the following relations:

(1) $\langle a_i \rangle^G$ is abelian for all $i \geq 1$;

 $\mathbf{2}$

- (2) $\langle x \rangle^G$ is metabelian;
- (3) $x^p = a_1^p = a_2^p = \dots = 1;$
- (4) if $s \neq x$ is a simple commutator in x, a_1, a_2, \ldots , where $t(s) \geq 2$ or $t(s) \leq -2$, then s = 1.

Remark. Notice that G is a locally finite p-group. Also, from (4) it follows immediately that if s is a simple commutator in x, a_1, a_2, \ldots , then [s, x] = 1 whenever $t(s) \neq 1$. This is something we will make use of later.

Let s be a left-normed commutator of weight 3m + 1 in x, a_1, a_2, \ldots , starting in x. The reader can convince himself/herself that from (4) it follows that such a commutator will be trivial if it is not of the form $[x, a_{j_1}, a_{j_2}, a_{j_3}, x, a_{j_4}, a_{j_5}, \ldots, x, a_{j_{2m}}, a_{j_{2m+1}}]$. We will establish later that there are such non-trivial commutators for any m that implies that $\langle x \rangle^G$ is not nilpotent. However the structure of G is not transparent enough at this stage for us to see this directly. Instead will will use Lie ring methods to lift up this property from an analogous Lie ring setting that we will deal with in the next two sections. On the other hand we can establish directly that x is a left 3-Engel element and this we will do now for the rest of this section. We start with a useful lemma.

Lemma 2.1. Let $g \in G$. We can write $g = \alpha \beta \gamma \delta$ where $\alpha, \beta, \gamma, \delta$ are products of simple commutators in x, a_1, a_2, \ldots of types 1, 2, -1, 0.

Proof. We prove first by induction on $k \geq 1$ that we can write $g = \alpha_k \beta_k \gamma_k \delta_k \epsilon(k+1)$ 1), where $\alpha_k, \beta_k, \gamma_k, \delta_k$ are products of simple commutators in x, a_1, a_2, \ldots of types 1, 2, -1, 0 and $\epsilon(k+1)$ is a product of commutators of weight k+1 or higher. For the induction basis observe that we can write $q = x_1 x_2 \cdots x_m$ where $x_i \in \{x, a_1, a_2, \ldots\}$ and modulo commutators of weight 2 or higher we can rearrange the product so that $g = \alpha_1 \beta_1 \epsilon(2)$ and where α_1 is a product of elements from $\{a_1, a_2, \ldots\}$, β_1 is a power of x and $\epsilon(2)$ is a product of simple commutators of weight 2 or higher (here $\gamma_1 = \delta_1 = 1$). For the induction step suppose that $k \geq 2$ and that the inductive claim holds for smaller values of k. Thus we know that we can write $g = \alpha_{k-1}\beta_{k-1}\gamma_{k-1}\delta_{k-1}\epsilon(k)$ where $\alpha_{k-1}, \beta_{k-1}, \gamma_{k-1}, \delta_{k-1}$ are products of simple commutators in x, a_1, a_2, \ldots of types 1, 2, -1, 0 and where $\epsilon(k)$ is a product of simple commutators of weight k or higher. Working modulo commutators of weight k + 1 and higher, we can now collect the commutators of weight k so that they appear with the commutator of smaller weight that are of the same type. This gives us the expression $g = \alpha_k \beta_k \gamma_k \delta_k \epsilon(k+1)$ that we wanted. This finishes the proof of the inductive hypothesis. Notice now that we are working with simple commutators in x_1, x_2, \ldots, x_m and as $H = \langle x_1, \ldots, x_m \rangle$ is nilpotent we have that for a large enough k we have $\epsilon(k+1) = 1$.

Let $g \in G$. We want to show that [g, x, x, x] = 1. Let $g = \alpha \beta \gamma \delta$ be an expression for g as in Lemma 2.1. As we remarked above, all simple commutators of type different from 1 commute with x. This means that β, γ, δ commute with x and without loss of generality we can assume that

$$g = b_1 \cdots b_m$$

where b_1, \ldots, b_m are simple commutators in x, a_1, a_2, \ldots of type 1.

Let $H = \langle x, b_1, \ldots, b_m \rangle$. Observe that relations (1) and (2) imply that $\langle b_i \rangle^H$ is abelian and $\langle x \rangle^H$ is metabelian. Notice also that relations (1) and (3) imply that $b_1^p = \cdots = b_m^p = 1$. Finally, suppose that we have a simple commutator cin x, b_1, \ldots, b_m of multi-weight $(l, f_1, f_2, \ldots, f_m)$ in x, b_1, \ldots, b_m then the type of c as a simple commutator in x, a_1, a_2, \ldots is $-2l + f_1t(b_1) + \cdots + f_mt(b_m) =$ $-2l + f_1 + \cdots + f_m$ that is equal to the type of c as a simple commutator in x, b_1, \ldots, b_m . Hence relations (4) hold for H with x, a_1, a_2, \ldots replaced by x, b_1, \ldots, b_m . It follows from this that there is a homomorphism $\alpha : G \to H$ that maps x, a_1, \ldots, a_m to x, b_1, \ldots, b_m and a_k to 1 for $k \ge m + 1$. In particular $[b_1 \cdots b_m, x, x, x] = 1$ would follow from $[a_1 \cdots a_m, x, x, x] = 1$. Thus the problem of showing that x is a left 3-Engel element reduces to showing that $[a_1 \cdots a_r, x, x, x] = 1$ for all integers $r \ge 1$. Before turning to this we need to introduce some more standard notation.

Definition. Let $y, c_1, c_2, ..., c_m$ be group elements. For any non-trivial subset $I = \{i_1, ..., i_k\}$ of $\{1, 2, ..., m\}$, where $i_1 < ... < i_k$ we denote by $[y, c_I]$ the commutator $[y, c_{i_1}, ..., c_{i_k}]$.

Proposition 2.2. We have $[a_1 \cdots a_r, x, x, x] = 1$ for all integers $r \ge 1$.

Proof. We prove this by induction on r. For the induction basis simply observe that $t([a_1, x, x, x]) = -5$ and thus $[a_1, x, x, x] = 1$ by relations (4). Now suppose $r \ge 2$ and that the result holds for all smaller values for r. First we see from standard commutator properties that

$$[a_1 \cdots a_r, x] = \left(\prod_{I \subseteq \{2, \dots, r\}} [a_1, x, a_I]\right) \cdot \left(\prod_{I \subseteq \{3, \dots, r\}} [a_2, x, a_I]\right) \dots [a_r, x].$$

Notice that relations (1) imply that the order of factors within each sub-product above does not matter. The RHS has $s = 2^r - 1$ factors. Suppose these are b_1, \ldots, b_s and thus $[a_1 \cdots a_r, x] = b_1 \cdots b_s$. We want to show that $[b_1 \cdots b_s, x, x] = 1$ or equivalently that $[x, b_1 \cdots b_s]$ commutes with x. Expanding again we see that

$$[x, b_1 \cdots b_s] = \prod_{\emptyset \neq I \subseteq \{1, \dots, s\}} [x, b_I].$$

Notice that each factor is in $(\langle x \rangle^G)'$ and as $\langle x \rangle^G$ is metabelian the order of the factors does not matter. Notice also that every non-trivial factor is of the form $[x, [x, a_{I_1}]^{-1}, \ldots, [x, a_{I_l}]^{-1}]$ where, because of relations (1), the sets I_1, \ldots, I_l are pairwise disjoint. We also have that $\min(I_1) < \min(I_2) < \ldots < \min(I_l)$. Let A be the product of such factors where not all a_1, a_2, \ldots, a_r are included and let B be the product of those factors where each a_1, a_2, \ldots, a_r occurs once. Then

$$[x, b_1 \cdots b_s] = AB.$$

It is not difficult to see by standard arguments that the induction hypothesis implies that A commutes with x. We are thus left with showing that B commutes with x. Now let

$$[x, [x, a_{I_1}], \dots, [x, a_{I_m}]]^{(-1)}$$

be one of the factors of *B*. This will commute with *x* if the type is not 1. As $I_1 \cup \cdots \cup I_m = \{1, 2, \ldots, r\}$ we have that the $t([x, [x, a_{I_1}], \ldots, [x, a_{I_m}]]) =$ r-2(m+1) that is equal to 1 if and only if r=2m+3. We are thus left with showing that e(m) commutes with x where

$$e(m) = \prod_{(I_1, I_2, \dots, I_m)} [x, [x, a_{I_1}], \dots, [x, a_{I_m}]].$$

Here the product is taken over all partitions of $\{1, 2, ..., 2m + 3\}$ into a disjoint union of m non-empty subsets where

$$1 = \min(I_1) < \min(I_2) < \ldots < \min(I_m).$$

As $\langle x \rangle^G$ is metabelian the value of a factor does not change if we permute the terms $[x, a_{I_2}], \ldots, [x, a_{I_m}]$. Notice also that for $[x, [x, a_{I_1}]]$ to be non-trivial we need $|I_1| = 3$. If some I_j has size less than 2 where $2 \leq j \leq m$ then (because we can permute the terms $[x, a_{I_2}], \ldots, [x, a_{I_m}]$ with out changing the value) we see that the factor is trivial as $[x, [x, a_{I_1}], [x, a_{I_j}]]$ has type less than -1. For a factor of e(m) to be non-trivial we thus need all of I_2, I_3, \ldots, I_m to have size 2 or 3. Suppose s are of size 2 and t of size 3. Then the we get the equations

$$s+t = m-1$$
$$2s+3t = 2m.$$

That has the solution t = 2 and s = m - 3 (in particular $m \ge 3$). This shows that

$$e(m) = \prod_{(I_1, I_2, \dots, I_m)} [x, [x, a_{I_1}], \dots, [x, a_{I_m}]]$$

where the product is taken over all partitions of $\{1, 2, \ldots, 2m+3\}$ where $|I_1| = |I_2| = |I_3| = 3$ and $|I_4| = \ldots = |I_m| = 2$ and $1 \in I_1$.

Consider any factor of this product. As $[x, a_{I_4}], \ldots, [x, a_{I_m}]$ commute with x, we have

$$[e(m), x] = \prod_{(I_1, I_2, \dots, I_m)} [x, [x, a_{I_1}], [x, a_{I_2}], [x, a_{I_3}], x, [x, a_{I_4}], \dots, [x, a_{I_m}]].$$

It follows from this that in order to show that [e(m), x] = 1 for all $m \ge 1$, it suffices to show that [e(3), x] = 1.

To see this take any $2 \le i < j < k < r \le 9$. We have

$$1 = [x, a_1, [x, a_i, a_j, a_k, a_r]] = \prod_{(I_1, I_2)} [x, a_{I_1}, x, a_{I_2}].$$

where the product is taken over all partitions of $\{1, i, j, k, r\}$ where $|I_1| = 3$, $|I_2| = 2$ and $1 \in I_1$. From this it follows that

$$e(3) = \prod_{(I_1, I_2, I_3)} [x, [x, a_{I_1}], [x, a_{I_2}], [x, a_{I_3}]]$$

is trivial and thus in particular [e(3), x] = 1. This finishes the proof.

As a consequence we have the following theorem.

Theorem 2.3. The element x is a left 3-Engel element in G.

3. The Lie Algebra L

In this section we will construct a pair (L, z) that will provide us with the Lie algebra analogue of (G, x). Most of this section will be about describing the Lie algebra L and getting some close transparent information about $\mathrm{Id}(z)$, the ideal in L generated by z. This will then be used in Section 4 to prove that $\mathrm{Id}(z)$ is not nilpotent. Throughout the paper p is a fixed odd prime and $\mathbb{F} = \mathbb{F}_p$ is the field of p elements.

We will start with a result that is probably well known although we do not have a reference. For that reason and for the convenience of the reader we will include a proof. We first need some notation. Let z, c_1, c_2, \ldots be some Lie algebra variables. For any non-trivial multi-set $\{c_{i_1}, c_{i_2}, \ldots, c_{i_k}\}$ with $i_1 \leq i_2 \cdots \leq i_k$ we denote by $[z, c_I]$ the left normed Lie commutator $[z, c_{i_1}, \ldots, c_{i_k}]$.

Proposition 3.1. Let *E* be the largest Lie algebra over \mathbb{F} generated z, c_1, c_2, \ldots where c_1, c_2, \ldots commute. Then $\mathrm{Id}(z)$ is freely generated by $\langle [z, c_I], I \subset_{fm} \mathbb{N}^m \rangle$, where *I* runs through all finite multi-subsets of \mathbb{N}^m . Here \mathbb{N}^m is the multi-set variant of the set of natural numbers namely $\{1, 1, \ldots, 2, 2, \ldots, \ldots\}$.

Proof. Let A be the enveloping algebra of E, that is the largest associative algebra generated by z, c_1, c_2, \ldots where c_1, c_2, \ldots commute. Thus E is the Lie sub-algebra generated by z, c_1, c_2, \ldots where the Lie product of $u, v \in A$ is [u, v] = uv - vu. In order to prove the proposition it suffices to show that the associative sub-algebra C generated by the set $\{[z, c_I], I \subset_{fm} \mathbb{N}^m\}$ is freely generated as an associative algebra by these elements. To see this, pick a variable y_I for each finite multi sub-set of \mathbb{N}^m and consider the free associative algebra B, freely generated by these. Order the generators of A with $c_1 < c_2 < \cdots < z$ and use these to define a lexicographical order on words in these. We also order the generators y_I by

$$y_I < y_J$$
 if and only if $zc_{i_1} \cdots c_{i_r} < zc_{j_1} \cdots c_{j_s}$

where $I = \{i_1, \ldots, i_r\}, J = \{j_1, \ldots, j_s\}, i_1 \leq \cdots \leq i_r \text{ and } j_1 \leq \cdots \leq j_s$. We are here using the lexicographical order above for the displayed inequalities. Notice that if we expand $[z, c_I]$ then the leading term with respect to that lexicographical order is $zc_{i_1}\cdots c_{i_r}$. We will denote the latter by zc_I . We then use this order do define a lexiographical order on words in the y_I 's. Now consider the algebra homomorphism from B to A that maps y_I to $[z, c_I]$. In order to show that C is freely generated by the $[z, c_I]$'s it suffices to show that this homomorphism is injective. Thus let y be a non-zero element in B say

$$y = \alpha y_{I_1} \cdots y_{I_r} + \cdots + \beta y_{J_1} \cdots y_{J_s}$$

where $\alpha y_{I_1} \cdots y_{I_r}$ is the leading term of y with respect to the lexicographical ordering on B described above. From the definition of this lexicographical order, we see that

$$\bar{y} = \alpha z c_{I_1} \cdots z c_{I_r} + \cdots + \beta z c_{J_1} \cdots z c_{J_s}$$

has leading term $\alpha z c_{I_1} \cdots z c_{I_r}$ with respect to the lexicographical order on A. But notice that this is also the leading term of the image of y under the homomorphism, namely

$$\tilde{y} = \alpha[z, c_{I_1}] \cdots [z, c_{I_r}] + \cdots + \beta[z, c_{J_1}] \cdots [z, c_{J_s}].$$

This shows that the image of y is non-zero and thus the homomorphism is injective. This finishes the proof.

Thus $\mathrm{Id}_E(z)$ is a free Lie algebra, freely generated by the $[z, c_I]$'s. For the following we will use a slightly different total order on this set of free generators. If I and J are both non-empty, then $[z, c_I] < [z, c_J]$ as above. However if one of the multi-sets is empty we define the order differently so that z is the largest element. That is $[z, c_I] < z$ whenever I is non-empty. Now consider the largest metabelian quotient, M(z), of $\mathrm{Id}(z)$. This is the free metabelian Lie algebra over \mathbb{F} with free generators the $[z, c_I]$'s. Now it is well known (see for example [2]), that M(z) has a basis consisting of the left-normed Lie commutators

$$[[z, c_{I_1}], [z, c_{I_2}], \dots, [z, c_{I_m}]]$$

where $m \ge 1$ and $[z, c_{I_1}] > [z, c_{I_2}] \le \cdots \le [z, c_{I_m}].$

We are now getting ready to describe our Lie algebra L. First we consider the largest Lie algebra $F = \langle z, c_1, c_2, \ldots \rangle$ over \mathbb{F}_p , such that

- (1) $Id(c_i)$ is abelian for i = 1, 2, ...;
- (2) Id(z) is metabelian.

From the analysis above, the following is a basis for $Id_F(z)$

(*)
$$[[z, c_{I_1}], [z, c_{I_2}], \dots, [z, c_{I_m}]]$$

where $m \geq 1, I_1, \ldots, I_m$ are pairwise disjoint and $[z, c_{I_1}] > [z, c_{I_2}] \leq \cdots \leq [z, c_{I_m}]$. We define a type of a Lie commutator in a similar way as we did for group commutators in Section 1. This term will play an important role in this section.

Definition. Let s be a simple commutator of multi-weight (m, e_1, \ldots, e_r) in z, c_1, \ldots, c_r . The type of s is $t(s) = e_1 + \cdots + e_r - 2m$.

Remark. If c and d are simple commutators then t([c,d]) = t(c) + t(d). In particular t([c,x]) = t(c) - 2 and $t([c,a_1]) = t(c) + 1$.

We will now construct a Lie algebra L that will be a quotient of F by a certain multi-homogeneous ideal J. Note that since F is multi-graded, L will also be multi-graded. Our multi-homogeneous ideal will be the smallest ideal of F containing the following elements from our basis for $\mathrm{Id}_F(z)$ described above. Let $c = [[z, c_{I_1}], [z, c_{I_2}], \ldots, [z, c_{I_m}]]$ be one of these basis elements.

- (3) c is in J if $c \neq z$ is a commutator of type either less than -1 or more than 1;
- (4) c is in J if one of I_1, I_3, \ldots, I_m has size greater than 2.

In this and following section we will determine the structure of J and prove that $Id_L(z)$ is not nilpotent.

3.1. Some detailed analysis about which basis elements are in J. One can see directly that a number of the commutators in (*) will be in J as a consequence of conditions (3) and (4). In this section, we will now look more closely at these commutators and see that a number of elements are in J as a less direct consequence. Consider one of the basis elements, $[[z, c_{I_1}], [z, c_{I_2}], \ldots, [z, c_{I_m}]]$, from (*) above. Note that, in particular, by condition (3) we have that $[z, c_I]$ is in J when $|I| \ge 4$. In the following we will denote with α the number of I_i of size 1 and β those of size 2, for $i \ge 3$.

Remark. Notice that, since Id(z) is metabelian, it does not matter how we order I_3, \ldots, I_m as they can be permuted without changing the value of the Lie commutator. In fact, the only constraint is that the smallest i in $I_1 \cup \cdots \cup I_m$ should be in I_2 , since $[z, c_{I_1}] > [z, c_{I_2}] \leq \cdots \leq [z, c_{I_m}]$.

Commutators of type -1 where $|I_1| + |I_2| = 3$. In this case we have $t([z, c_{I_1}], [z, c_{I_2}]) = -1$ and $|I_3| + \cdots + |I_m| = 2m - 4$. Notice that none of I_3, \ldots, I_m is empty or of size 1 since otherwise the commutator would be in J by condition (3). Then all commutators are in J except possibly those where

$$|I_1| + |I_2| = 3, |I_3| = \dots = |I_m| = 2, |I_1| \le 2.$$

Commutators of type -1 where $|I_1| + |I_2| = 4$. Here $t([z, c_{I_1}], [z, c_{I_2}]) = 0$ and $|I_3| + \cdots + |I_m| = 2m - 5$. As before, none of I_3, \ldots, I_m is empty since otherwise the commutator would be in J by condition (3). Furthermore none of the sets is of size greater than 2 by condition (4). We thus obtain the following system of equations

$$\begin{cases} + \beta = m - 2 \\ + 2\beta = 2m - 5 \end{cases}$$

whose solution is $\alpha = 1$ and $\beta = m - 3$. Therefore all commutators are in J except possibly those where

$$|I_1| + |I_2| = 4, |I_3| = 1, |I_4| = \dots = |I_m| = 2, |I_1| \le 2.$$

Commutators of type -1 where $|I_1| + |I_2| = 5$. We have $t([z, c_{I_1}], [z, c_{I_2}]) = 1$ and $|I_3| + \cdots + |I_m| = 2m - 6$. First, we consider the case when I_3 is empty. In this case $t([z, c_{I_1}], [z, c_{I_2}], [z, c_{I_3}]) = -1$ and none of I_4, \ldots, I_m are of size 1. Thus they are all of size 2, and all commutators are in J except possibly the cases

$$|I_1| = 2, |I_2| = 3, |I_3| = 0, |I_4| + \dots + |I_m| = 2.$$

Now it remains to deal with the case for which none of I_3, \ldots, I_m is empty. Using α and β as before, we have

$$\begin{cases} + \beta = m - 2 \\ + 2\beta = 2m - 6 \end{cases}$$

whose solution is $\alpha = 2$ and $\beta = m - 4$. Thus all commutators are in J except possibly the cases where

$$|I_1| = 2, |I_2| = 3, |I_3| = |I_4| = 1, |I_5| = \dots = |I_m| = 2.$$

Commutators of type 0 where $|I_1| + |I_2| = 3$. In this case none of I_3, \ldots, I_m are empty or of size 1, as otherwise the commutator is in J by condition (3). Also, none are of size 3 by condition (4). Therefore, they must all be of size 2, which implies that the type is -1, that is a contradiction since we supposed that the type is 0.

Commutators of type 0 where $|I_1| + |I_2| = 4$. We have that none of I_3, \ldots, I_m is empty. Using α and β as before, we have

$$\begin{cases} + \beta = m - 2 \\ + 2\beta = 2m - 4 \end{cases}$$

whose solution is $\alpha = 0$ and $\beta = m - 2$. Thus all commutators are in J except possibly those where

$$|I_1| + |I_2| = 4, |I_3| = \dots = |I_m| = 2, |I_1| \le 2.$$

Commutators of type 0 where $|I_1| + |I_2| = 5$. First, notice that none of I_3, \ldots, I_m is empty. Indeed, if I_3 is empty then $t([z, c_{I_1}], [z, c_{I_2}], [z, c_{I_3}]) = -1$ and the commutator has type at most -1 if it is non-trivial. We then get a contradiction since we are considering the case when the type is 0. Thus none of I_3, \ldots, I_m is empty. Therefore we have

$$\begin{cases} + \beta = m - 2 \\ + 2\beta = 2m - 5 \end{cases}$$

whose solution is $\alpha = 1$ and $\beta = m - 3$. Therefore all commutators are in J except possibly those where

$$|I_1| = 2, |I_2| = 3, |I_3| = 1, |I_4| = \dots = |I_m| = 2.$$

Commutators of type 1. Since $|I_3| + ... |I_m| \le 2m - 4$, we need $|I_1| + |I_2| = 5$. Then all commutators are in J except possibly those where

$$|I_1| = 2, |I_2| = 3, |I_3| = \dots = |I_m| = 2.$$

In view of the analysis above we come up with the following decomposition of $Id_F(z)$ as a direct sum of sub spaces.

$$\mathrm{Id}_F(z) = \langle X \setminus Z \rangle \oplus \langle Z \rangle$$

where Z is the list of basis elements with the following partitions (I_1, \ldots, I_m)

$[z, c_I], \text{ for } I \leq 3$	
$ I_1 + I_2 = 3, I_1 \le 2, I_3 = \dots = I_m = 2$	(ζ_1)
$ I_1 + I_2 = 4, I_1 \le 2, I_3 = 1, I_4 = \dots = I_m = 2$	(ζ_2)
$ I_1 = 2, I_2 = 3, I_3 = 0, I_4 = \dots = I_m = 2$	(ζ_3)
$ I_1 = 2, I_2 = 3, I_3 = I_4 = 1, I_5 = \dots = I_m = 2$	(ζ_4)
$ I_1 = 2, I_2 = 3, I_3 = 1, I_4 = \dots = I_m = 2$	(ξ_1)
$ I_1 + I_2 = 4, I_1 \le 2, I_3 = \dots = I_m = 2$	(ξ_2)
$ I_1 = 2, I_2 = 3, I_3 = \dots = I_m = 2.$	(au_1)

We have seen above that $\langle X \setminus Z \rangle \subseteq J$. Now J is the smallest subspace of F containing $X \setminus Z$ that is invariant under taking Lie commutators with the c_j 's and z. In the following we go systematically through all possible scenarios where we need to add new elements to $\langle X \setminus Z \rangle$ in order to get a subspace $Z_0 + \langle X \setminus Z \rangle$, where $Z_0 \leq Z$, that is invariant under Lie commutators with the c_j 's. All the new elements will be relations amongst the ζ 's, ξ 's, and τ 's. We will distinguish several cases considering the cardinality of the sets I_1, \ldots, I_m . We will then later ensure furthermore that we obtain a subspace that is also invariant under taking Lie commutators with z.

3.2. Relations among $\zeta_1, \zeta_2, \zeta_3, \zeta_4$.

We now begin systematically adding elements to $\langle X \setminus Z \rangle$ to ensure that the resulting subspace is invariant under taking Lie commutator with c_k . For this we only need to consider elements of types -2, -1 and 0. We start with a basis element u from $X \setminus Z$ that is of type -2. We write $[u, c_k]$ as a linear combination of basis elements in X. If all the components are in $X \setminus Z$, then nothing needs to be added. We thus only need to consider the cases when some of the components are among $\zeta_1, \zeta_2, \zeta_3$ or ζ_4 . This means reducing the size of one of I_1, \ldots, I_m for some of ζ_1, \ldots, ζ_4 by one, to get a basis element u of type -2 and then expand $[u, c_k]$ to get a linear combination of ζ_1, \ldots, ζ_4 modulo J. This may then give us a new relator to add to $\langle X \setminus Z \rangle$. Below we exhaust all possibilities.

Case A1:
$$|I_1| = 2, |I_2| = 3, |I_3| = 0, |I_4| = 1, |I_5| = \dots = |I_m| = 2.$$

Without loss of generality, we can assume that $I_1 \cup I_2 \cup \cdots \cup I_m \cup \{k\} = \{1, \ldots, 2m-1\}$. Recall that for the basis elements for $\mathrm{Id}_F(z)$ we should have $1 \in I_2$. For this reason we need to deal separately with the cases $k \geq 2$ and k = 1.

Consider first the case when $k \ge 2$. Suppose $I_4 = \{i\}$. Here as elsewhere in this section we will be calculating modulo J. We have

$$0 = [[z, c_{I_1}], [z, c_{I_2}], z, [z, c_i], [z, c_{I_5}], \dots, [z, c_{I_m}], c_k] = \zeta_4(I_1, I_2, \{k\}, \{i\}, I_5, \dots, I_m) + \zeta_3(I_1, I_2, \emptyset, \{i, k\}, I_5, \dots, I_m).$$

Therefore

(R1) $\zeta_4(I_1, I_2, \{k\}, \{i\}, I_5, \dots, I_m) = -\zeta_3(I_1, I_2, \emptyset, \{i, k\}, I_5, \dots, I_m).$ This means that we need to add the relator $\zeta_4(I_1, I_2, \{k\}, \{i\}, I_5, \dots, I_m) + \zeta_3(I_1, I_2, \emptyset, \{i, k\}, I_5, \dots, I_m)$ to $\langle X \setminus Z \rangle.$

The reader can convince himself that the case k = 1 does not give us a new relation.

Case A2: $|I_1| = 1, |I_2| = 3, |I_3| = 0, |I_4| = \cdots = |I_m| = 2.$

First consider the case when $k \ge 2$ and suppose that $I_1 = \{i\}$. We have

Therefore,

(R2) $\zeta_2(\{i\}, I_2, \{k\}, I_4, \dots, I_m) = -\zeta_3(\{i, k\}, I_2, \emptyset, I_4, \dots, I_m).$

Again one sees that for k = 1, one does not get a new relation.

Case A3: $|I_1| = |I_2| = 2, |I_3| = 0, |I_4| = \dots = |I_m| = 2.$

Consider the case when $k \ge 2$. We have

Therefore, we get

(R3) $\zeta_2(I_1, I_2, \{k\}, I_4, \dots, I_m) = -\zeta_3(I_1, I_2 \cup \{k\}, \emptyset, I_4, \dots, I_m).$ We will handle k = 1 later.

Case A4: $|I_1| = 2, |I_2| = 1, |I_3| = 1, |I_4| = \cdots = |I_m| = 2.$

Let $I_3 = \{i\}$. For the case $k \ge 2$ we get

$$0 = [[z, c_{I_1}], [z, c_{\{1,k\}}], [z, c_i], [z, c_{I_4}], \dots, [z, c_{I_m}]] + [[z, c_{I_1}], [z, c_1], [z, c_{\{i,k\}}], [z, c_{I_4}], \dots, [z, c_{I_m}]] = \zeta_2(I_1, \{1, k\}, \{i\}, I_4, \dots, I_m) + \zeta_1(I_1, \{1\}, \{i, k\}, I_4, \dots, I_m) = -\zeta_3(I_1, \{1, i, k\}, \emptyset, I_4, \dots, I_m) + \zeta_1(I_1, \{1\}, \{i, k\}, I_4, \dots, I_m),$$

where the last equality follows from R3. Therefore,

(R4) $\zeta_1(I_1, \{1\}, \{i, k\}, I_4, \dots, I_m) = \zeta_3(I_1, \{1, i, k\}, \emptyset, I_4, \dots, I_m).$

We will deal with the case k = 1 later.

Case A5:
$$|I_1| = 1, |I_2| = 2, |I_3| = 1, |I_4| = \dots = |I_m| = 2.$$

Consider first the case $k \ge 2$. Let $I_1 = \{i\}, I_3 = \{j\}$ and $k \ge 2$. We have

$$\begin{array}{lll} 0 &=& [[z,c_{\{i,k\}}], [z,c_{I_2}], [z,c_j], [z,c_{I_4}], \dots, [z,c_{I_m}]] \\ &+& [[z,c_i], [z,c_{I_2\cup\{k\}}], [z,c_j], [z,c_{I_4}], \dots, [z,c_{I_m}]] \\ &+& [[z,c_i], [z,c_{I_2}], [z,c_{\{j,k\}}], [z,c_{I_4}], \dots, [z,c_{I_m}]] \\ &=& \zeta_2(\{i,k\}, I_2, \{j\}, I_4, \dots, I_m) + \zeta_2(\{i\}, I_2 \cup \{k\}, \{j\}, I_4, \dots, I_m) \\ &+& \zeta_1(\{i\}, I_2, \{j,k\}, I_4, \dots, I_m) \\ &=& -\zeta_3(\{i,k\}, I_2 \cup \{j\}, \varnothing, I_4, \dots, I_m) - \zeta_3(\{i,j\}, I_2 \cup \{k\}, \varnothing, I_4, \dots, I_m) \\ &+& \zeta_1(\{i\}, I_2, \{j,k\}, I_4, \dots, I_m), \end{array}$$

where the last equality follows from R2 and R3. Therefore,

(R5)
$$\zeta_1(\{i\}, I_2, \{j, k\}, I_4, \dots, I_m) = \zeta_3(\{i, k\}, I_2 \cup \{j\}, \emptyset, I_4, \dots, I_m) + \zeta_3(\{i, j\}, I_2 \cup \{k\}, \emptyset, I_4, \dots, I_m).$$

We deal with the case k = 1 later.

Case A6:
$$|I_1| = 0, |I_2| = 3, |I_3| = 1, |I_4| = \dots = |I_m| = 2$$
.

We first look at the case $k \ge 2$. Let $|I_3| = \{i\}$. We have

Therefore,

(R6)
$$\zeta_1(\emptyset, I_2, \{i, k\}, I_4, \dots, I_m) = \zeta_3(\{i, k\}, I_2, \emptyset, I_4, \dots, I_m).$$

Again the case k = 1 does not add anything new.

Notice that it follows from the relations above that modulo J we can write the basis elements of types ζ_1, ζ_2 and ζ_4 as a linear combination of basis elements of type ζ_3 . Notice also that from (R6) and the fact that $\mathrm{Id}_F(z)$ is metabelian, it follows that

$$\begin{aligned} \zeta_3(I_1, I_2, \emptyset, I_4, I_5, \dots, I_m) &= \zeta_1(\emptyset, I_2, I_1, I_4, I_5, \dots, I_m) \\ &= \zeta_1(\emptyset, I_2, I_4, I_1, I_5, \dots, I_m) \\ &= \zeta_3(I_4, I_2, \emptyset, I_1, I_5, \dots, I_m). \end{aligned}$$

Therefore we get

(R7)
$$\zeta_3(I_1, I_2, \emptyset, I_4, I_5, \dots, I_m) = \zeta_3(I_4, I_2, \emptyset, I_1, I_5, \dots, I_m).$$

Case A7: $|I_1| = |I_2| = 1, |I_3| = \cdots = |I_m| = 2.$

Let $I_1 = \{i\}, I_3 = \{j, r\}$ and consider the case where $k \ge 2$. We have

Therefore,

(R8)
$$\zeta_3(\{i,k\},\{1,j,r\},\emptyset,I_4,\ldots,I_m) = -\zeta_3(\{i,j\},\{1,k,r\},\emptyset,I_4,\ldots,I_m) - \zeta_3(\{i,r\},\{1,j,k\},\emptyset,I_4,\ldots,I_m).$$

For k = 1 one can check that there is not a new relation.

Case A8: $|I_1| = 0, |I_2| = 2, |I_3| = \cdots = |I_m| = 2.$

Let $I_2 = \{1, i\}, I_3 = \{j, r\}$ and $k \ge 2$. We have

$$\begin{aligned} 0 &= [z, [z, c_{\{1,i,k\}}], [z, c_{I_3}], \dots, [z, c_m]] + [[z, c_k], [z, c_{\{1,i\}}], [z, c_{I_3}], \dots, [z, c_{I_m}]] \\ &= \zeta_1(\varnothing, \{1, i, k\}, I_3, \dots, I_m) + \zeta_1(\{k\}, \{1, i\}, I_3, \dots, I_m) \\ &= \zeta_3(\{j, r\}, \{1, i, k\}, \varnothing, I_4, \dots, I_m) + \zeta_3(\{k, j\}, \{1, i, r\}, \varnothing, I_4, \dots, I_m) \\ &+ \zeta_3(\{k, r\}, \{1, i, j\}, \varnothing, I_4, \dots, I_m). \end{aligned}$$

Modulo relation R8, we get

(R9)
$$\zeta_3(\{j,r\},\{1,i,k\},\emptyset,I_4,\ldots,I_m) = \zeta_3(\{i,k\},\{1,j,r\},\emptyset,I_4,\ldots,I_m).$$

The reader can check that here fas well as for A3, A4 and A5 with k = 1 one does not obtain any new relations. The same is true for the following remaining cases.

Case A9.
$$|I_1| = |I_2| = 2, |I_3| = |I_4| = 1, |I_5| = \dots = |I_m| = 2.$$

Case A10: $|I_1| = 2, |I_2| = 3, |I_3| = |I_4| = |I_5| = 1, |I_6| = \dots = |I_m| = 2.$
Case A11: $|I_1| = 1, |I_2| = 3, |I_3| = |I_4| = 1, |I_5| = \dots = |I_m| = 2.$
Case A12: $|I_1| = 2, |I_2| = 0, |I_3| = \dots = |I_m| = 2.$

To summarize we get the following set of equivalent relations for elements of type -1 modulo J.

$$\zeta_4(I_1, I_2, \{i\}, \{j\}, I_5, \dots, I_m) = -\zeta_3(I_1, I_2, \emptyset, \{i, j\}, I_5, \dots, I_m) \tag{E1}$$

$$\zeta_2(I_1, I_2, \{i\}, I_4, \dots, I_m) = -\zeta_3(I_1, I_2 \cup \{i\}, \emptyset, I_4, \dots, I_m) \ (|I_1| = |I_2| = 2) \quad (E2)$$

$$\zeta_2(\{i\}, I_2, \{j\}, I_4, \dots, I_m) = -\zeta_3(\{i, j\}, I_2, \emptyset, I_4, \dots, I_m)$$
(E3)

(E4)

$$\zeta_1(\emptyset, I_2, I_3, I_4, \dots, I_m) = \zeta_3(I_3, I_2, \emptyset, I_4, \dots, I_m)$$

$$\zeta_1(I_1, \{1\}, \{i, j\}, I_4, \dots, I_m) = \zeta_3(I_1, \{1, i, j\}, \emptyset, I_4, \dots, I_m)$$
(E5)

$$\zeta_1(\{i\}, I_2, \{j, k\}, I_4, \dots, I_m) = \zeta_3(\{i, j\}, I_2 \cup \{k\}, \emptyset, I_4, \dots, I_m)$$

$$+ \zeta_n(\{i, k\}, I_2 + \{i\}, \emptyset, I_1, \dots, I_m)$$
(E6)

$$+\zeta_3(\{i,k\}, I_2 \cup \{j\}, \emptyset, I_4, \dots, I_m) \tag{E6}$$

$$\zeta_3(I_1, I_2, \emptyset, I_4, I_5, \dots, I_m) = \zeta_3(I_4, I_2, \emptyset, I_1, I_5, \dots, I_m)$$
(E1)

$$\zeta_{3}(I_{1}, I_{2} \cup \{1\}, \emptyset, I_{4}, \dots, I_{m}) = \zeta_{3}(I_{2}, I_{1} \cup \{1\}, \emptyset, I_{4}, \dots, I_{m})$$
(E8)

$$\begin{aligned} \zeta_{3}(\{i,k\},\{1,j,r\},\emptyset,I_{4},\dots,I_{m}) + \zeta_{3}(\{i,j\},\{1,k,r\},\emptyset,I_{4},\dots,I_{m}) \\ + \zeta_{3}(\{i,r\},\{1,j,k\},\emptyset,I_{4},\dots,I_{m}) = 0. \end{aligned}$$
(E9)

Notice that relations E1-E6 tell us that, modulo J, all the basis elements of type -1 in X can be written as linear combinations of elements of type ζ_3 . Then E7 and E8 imply that

$$\zeta_3(I_1, I_2, I_4, \dots, I_m) = \zeta_3(I_1, I_2 \cup \{1\}, \emptyset, I_4, \dots, I_m)$$

is symmetric in $I_1, I_2, I_4, \ldots, I_m$. Apart from this we have one extra relation E9.

3.3. Relations among ξ_1, ξ_2 .

We continue the analysis and deal here with $[u, c_k]$ were u is of type -1. On readily sees that if $u \in X \setminus Z$, then $[u, c_k] \in \langle X \setminus Z \rangle$ and no new relators needed. We therefore only need to consider u where u is one of the extra relators that we obtained in Section 3.2.

B1. Consequences of **E1.** For $k \ge 2$ we have

$$[\zeta_4(I_1, I_2, \{i\}, \{j\}, I_5, \dots, I_m), c_k] = -[\zeta_3(I_1, I_2, \emptyset, \{i, j\}, I_5, \dots, I_m), c_k]$$
 which implies that

$$\begin{aligned} \xi_2(I_1, I_2, \{j\}, \{i, k\}, I_5, \dots, I_m) &= -\xi_2(I_1, I_2, \{i\}, \{j, k\}, I_5, \dots, I_m) \\ &- \xi_2(I_1, I_2, \{k\}, \{i, j\}, I_5, \dots, I_m). \end{aligned}$$

Therefore,

(F1)
$$\xi_2(I_1, I_2, \{j\}, \{i, k\}, I_5, \dots, I_m) = -\xi_2(I_1, I_2, \{i\}, \{j, k\}, I_5, \dots, I_m) - \xi_2(I_1, I_2, \{k\}, \{i, j\}, I_5, \dots, I_m).$$

For k = 1 one does not get a new relation.

B2. Consequences of E8. For the case $k \ge 2$ we get

$$[\zeta_3(I_1, I_2 \cup \{1\}, \emptyset, I_4, \dots, I_m), c_k] = [\zeta_3(I_2, I_1, \{1\}, \emptyset, I_4, \dots, I_m), c_k].$$

Therefore,

(F2)
$$\xi_2(I_1, I_2 \cup \{1\}, \{k\}, I_4, \dots, I_m) = \xi_2(I_2, I_1 \cup \{1\}, \{k\}, I_4, \dots, I_m).$$

Again for k = 1 we do not get any new relation.

B3. Consequences of E7. For the case $k \ge 2$ we get

 $[\zeta_3(I_1, I_2, \emptyset, I_4, I_5, \dots, I_m), c_k] = [\zeta_3(I_4, I_2, \emptyset, I_1, I_5, \dots, I_m), c_k].$

Therefore,

(F3)
$$\xi_2(I_1, I_2, \{k\}, I_4, I_5, \dots, I_m) = \xi_2(I_4, I_2, \{k\}, I_1, I_5, \dots, I_m).$$

For k = 1 we do not get anything new.

B4. Consequences of E9. Commuting E_9 with a_s , for the case $s \ge 2$ we get (F4) $\xi_2(\{i,k\},\{1,j,r\},\{s\},I_4,\ldots,I_m) = -\xi_2(\{i,j\},\{1,k,r\},\{s\},I_4,\ldots,I_m) -\xi_2(\{i,r\},\{1,j,k\},\{s\},I_4,\ldots,I_m).$

For k = 1 we get nothing new.

B5. Consequences of E4. For the case $k \ge 2$ we get

$$[\zeta_1(\emptyset, I_2, I_3, I_4, \dots, I_m), c_k] = [\zeta_3(I_3, I_2, \{k\}, I_4, \dots, I_m), c_k].$$

As a consequence we obtain that

$$\xi_2(\{k\}, I_2, I_3, I_4, \dots, I_m) = \xi_2(I_3, I_2, \{k\}, I_4, \dots, I_m).$$

Therefore,

(F5) $\xi_1(\{i\}, I_2, \{j, k\}, I_4, \dots, I_m) = \xi_2(\{j, k\}, I_2, \{i\}, I_4, \dots, I_m).$ Again there is no relation for k = 1.

B6. Consequences of E5. For the case $k \ge 2$ we have

 $[\zeta_1(I_1,\{1\},\{i,j\},I_4,\ldots,I_m),c_k] = [\zeta_3(I_1,\{1,i,j\},\emptyset,I_4,\ldots,I_m),c_k].$

Then

$$\xi_1(I_1, \{1, k\}, \{i, j\}, I_4, \dots, I_m) = \xi_2(I_1, \{1, i, j\}, \{k\}, I_4, \dots, I_m).$$

Therefore,

(F6)
$$\xi_1(I_1, \{1, k\}, \{i, j\}, I_4, \dots, I_m) = \xi_2(I_1, \{1, i, j\}, \{k\}, I_4, \dots, I_m).$$

For k = 1 we do not get anything new and the same is true when consider consequences of E2,E3 and E6. From this analysis we get the following set of equivalent relations.

$$\xi_1(\{i\}, I_2, \{j, k\}, I_4, \dots, I_m) = \xi_2(\{j, k\}, I_2, \{i\}, I_4, \dots, I_m) \tag{G1}$$

$$\xi_1(I_1, \{1, k\}, \{i, j\}, I_4, \dots, I_m) = \xi_2(I_1, \{1, i, j\}, \{k\}, I_4, \dots, I_m)$$
(G2)

$$\xi_2(I_1, I_2 \cup \{1\}, \{k\}, I_4, \dots, I_m) = \xi_2(I_2, I_1 \cup \{1\}, \{k\}, I_4, \dots, I_m)$$
(G3)

$$\xi_2(I_1, I_2, \{k\}, I_4, I_5, \dots, I_m) = \xi_2(I_4, I_2, \{k\}, I_1, I_5, \dots, I_m) \tag{G4}$$

$$\xi_{2}(\{i,k\},\{1,j,r\},\{s\},I_{4},\ldots,I_{m}) + \xi_{2}(\{i,j\},\{1,k,r\},\{s\},I_{4},\ldots,I_{m}) + \xi_{2}(\{i,r\},\{1,j,k\},\{s\},I_{4},\ldots,I_{m}) = 0$$
(G5)

$$\xi_{2}(I_{1}, \{1, i, k\}, \{j\}, I_{4}, \dots, I_{m}) + \xi_{2}(I_{1}, \{1, k, j\}, \{i\}, I_{4}, \dots, I_{m}) + \xi_{2}(I_{1}, \{1, i, j\}, \{k\}, I_{4}, \dots, I_{m}) = 0.$$
(G6)

3.4. Relations in τ_1 .

Here we deal with $[u, c_k]$ where u is of type 0. Again we only get a new relation when u is one of the relators we obtained in Section 3.3.

C1. Consequences of G3. For $k \ge 2$ we have

$$[\xi_2(I_1, I_2 \cup \{1\}, \{i\}, I_4, \dots, I_m), c_k] = [\xi_2(I_2, I_1 \cup \{1\}, \{i\}, I_4, \dots, I_m), c_k].$$

which gives

$$\tau_1(I_1, I_2 \cup \{1\}, \{i, k\}, I_4, \dots, I_m) = \tau_1(I_2, I_1 \cup \{1\}, \{i, k\}, I_4, \dots, I_m).$$

Therefore, we get

(H1)
$$\tau_1(I_1, I_2 \cup \{1\}, I_3, \dots, I_m) = \tau_1(I_2, I_1 \cup \{1\}, I_3, \dots, I_m).$$

For k = 1 we get nothing new.

C2. Consequences of G4. For $k \ge 2$ we have

For k = 1 we get nothing new.

C3. Consequences of G5. For $k \ge 2$ we have

(H3)
$$\tau_1(\{i,k\},\{1,j,r\},I_3,\ldots,I_m) = -\tau_1(\{i,j\},\{1,k,r\},I_3,\ldots,I_m) -\tau_1(\{i,r\},\{1,k,j\},I_3,\ldots,I_m).$$

Again for k = 1 we get nothing new and the same is true for the consequence of G1, G2 and G6. Thus the only extra relators we get are H1, H2 and H3. We will now see that if we add to $\langle X \setminus Z \rangle$ the subspace generated by the extra relators we have obtained in Sections 3.2, 3.3 and 3.4, then we get an ideal in F.

Lemma 3.2. The subspace generated by $X \setminus Z$ and the relators given in E1-E9, G1-G6 and H1-H3 is an ideal in F and thus equal to J.

Proof. Let V be this subspace. We found the relations E1-E9, G1-G6 and H1-H3 by systematically ensuring that we obtained a subspace that is invariant under taking Lie commutators with the c_j 's. Notice that if we take a basis element $u = [[c, c_{i_1}], [z, c_{i_2}], \ldots, [z, c_{I_m}]]$ from X where $[u, z] \notin V$ then we must have $|I_1|, |I_3|, |I_4|, \ldots, |I_m| \leq 2$ and $|I_2| \leq 3$. Thus u has type at most 1. On the other hand if the type of u is less than 1, then [u, z] has type less than -1 and is thus in V. This means that we only need to consider $u \in V$ that is linear combination of elements of type τ_1 . In other words we have to commute each relation H1, H2, H3 with z and check that the resulting element is in V. If we commute relation H1 with z, we get

$$[\tau_1(I_1, I_2 \cup \{1\}, I_3, \dots, I_m), z] = [\tau_1(I_2, I_1 \cup \{1\}, I_3, \dots, I_m), z]$$

which implies that

$$\zeta_3(I_1, I_2 \cup \{1\}, \emptyset, I_4, \dots, I_m) = \zeta_3(I_2, I_1 \cup \{1\}, \emptyset, I_4, \dots, I_m),$$

16

which is relation E8 from Section 3.2. Using similar arguments we commute relation H2 with z, in which case we have

$$au_1(I_1, I_2, I_3, I_4, \dots, I_m), z] = [au_1(I_4, I_2, I_3, I_1, \dots, I_m), z]$$

which implies

$$\zeta_3(I_1, I_2, \emptyset, I_4, I_5, \dots, I_m) = \zeta_3(I_4, I_2, \emptyset, I_1, I_5, \dots, I_m)$$

which is relation E7 from Section 3.2. Lastly, commuting relation H3 with z gives

$$\begin{bmatrix} \tau_1(\{i,k\},\{1,j,r\},I_3,\ldots,I_m),z \end{bmatrix} = -[\tau_1(\{i,j\},\{1,k,r\},I_3,\ldots,I_m),z] \\ -[\tau_1(\{i,r\},\{1,k,j\},I_3,\ldots,I_m),z]$$

which implies

$$\zeta_{3}(\{i,k\},\{1,j,r\},\emptyset,I_{4},\ldots,I_{m}) = -\zeta_{3}(\{i,j\},\{1,k,r\},\emptyset,I_{4},\ldots,I_{m}) -\zeta_{3}(\{i,r\},\{1,j,k\},\emptyset,I_{4},\ldots,I_{m})$$

which is relation E9. This completes the proof.

4. The ideal
$$\operatorname{Id}_L(z)$$

Let $W = W(m, 1, \stackrel{2m+1}{\dots}, 1)$ be the subspace of F generated by all

$$E(I_1, \dots, I_m) = [[z, c_{I_1}], [z, c_{\{1\} \cup I_2}], [z, c_{I_3}], \dots, [z, c_{I_m}]]$$

with $I_1 \cup \cdots \cup I_m = \{2, \ldots, 2m+1\}$ and $|I_1| = \cdots = |I_m| = 2$. We know that these elements are linearly independent and we have also seen that

 $N = N(m, 1, \stackrel{2m+1}{\dots}, 1) = W(m, 1, \stackrel{2m+1}{\dots}, 1) \cap J$

is generated by two sets of relations

$$\begin{aligned} &(\mathcal{R}_1) & E(I_{\sigma(1)}, \dots, I_{\sigma(m)}) - E(I_1, \dots, I_m), \text{ where } \sigma \in \operatorname{Sym}\{1, \dots, m\} \\ &(\mathcal{R}_2) \\ & E(\{i_1, i_2\}, \{j_1, j_2\}, I_3, \dots, I_m) = - E(\{i_1, j_1\}, \{i_2, j_2\}, I_3, \dots, I_m) \\ & - E(\{i_1, j_2\}, \{i_2, j_1\}, I_3, \dots, I_m). \end{aligned}$$

Our aim is to show that $\frac{W}{N} \neq 0$, for all $m \geq 1$. Then it will follow that $L = \frac{F}{J}$ has multihomogenous elements of arbitrary weight m in z that are non-zero. As a consequence, we see that $\mathrm{Id}_L(z)$ is not nilpotent.

From now on, we work modulo (\mathcal{R}_1) and thus the order of I_1, \ldots, I_m does not matter for the value of $E(I_1, \ldots, I_m)$. From now on we write $E(\{I_1, \ldots, I_m\})$ for $E(I_1, \ldots, I_m)$ modulo (\mathcal{R}_1) . In this way we get an element for each partition $\{I_1, \ldots, I_m\}$ and these are a basis for W modulo (\mathcal{R}_1) .

Definition. We say that a basis element $E(\{I_1, \ldots, I_m\})$ has norm k if there are exactly k of the I_j 's, where $I_j \subseteq \{2, \ldots, m+1\}$. Notice that we then also have exactly k of the I_j 's, where $I_j \subseteq \{m+2, \ldots, 2m+1\}$.

Suppose that $k \geq 1$. We can assume $I_1, I_3, \ldots, I_{2k-1} \subseteq \{2, \ldots, m+1\}$ and

 $I_2, I_4, \ldots, I_{2k} \subseteq \{m + 2, \ldots, 2m + 1\}$. If $I_1 = \{i_1, i_2\}$ and $I_2 = \{j_1, j_2\}$, then modulo (\mathcal{R}_2) we have

$$E(\{\{i_1, i_2\}, \{j_1, j_2\}, I_3, \dots, I_m\}) = -E(\{\{i_1, j_1\}, \{i_2, j_2\}, I_3, \dots, I_m\}) - E(\{\{i_1, j_2\}, \{i_2, j_1\}, I_3, \dots, I_m\}).$$

Notice that the terms on the right hand side have norm k-1. We will refer to this as a 1-step decomposition of E through the pair (I_1, I_2) and we denote it $\mathcal{D}_{\{(I_1, I_2)\}}(E(\{I_1, \ldots, I_m\}))$. Suppose now $k \geq 2$. We can continue and apply the 1-step decomposition to the two terms through the pair (I_3, I_4) to get a 2-step decomposition of $E(I_1, \ldots, I_m)$ through $\{(I_1, I_2), (I_3, I_4)\}$ that we denote

$$\mathcal{D}_{\{(I_1,I_2),(I_3,I_4)\}}(E(\{I_1,\ldots,I_m\}))$$

We can continue in this manner through the pairs $(I_1, I_2), \ldots, (I_{2k-1}, I_{2k})$ to get a decomposition into 2^k terms of norm 0, denoted

$$\mathcal{D}_{\{(I_1,I_2),\ldots,(I_{2k-1},I_{2k})\}}(E(\{I_1,\ldots,I_m\})).$$

Remark. To say that an element $E({I_1, I_2, \ldots, I_m})$ has norm 0 is to say that ${I_1, \ldots, I_m} = {(2, \sigma(m+2)), (3, \sigma(m+3)), \ldots, (m+1, \sigma(2m+1))}$ for some permutation of σ of $m+2, \ldots, 2m+1$. We have seen that, modulo (\mathcal{R}_1) and (\mathcal{R}_2) , W is generated by basis elements of norm 0. Let W_0 be the subspace of W generated by these. Thus $W/W \cap J \cong (W_0 + J)/J \cong W_0/W_0 \cap J$.

Notice that the decomposition $\mathcal{D}_{\{(I_1,I_2),\ldots,(I_{2k-1},I_{2k})\}}(E(\{I_1,\ldots,I_m\}))$, does not depend on the order of the pairs in $\{(I_1,I_2),\ldots,(I_{2k-1},I_{2k})\}$, but it may depend on what pairings are chosen. For each $E(\{I_1,\ldots,I_m\})$ of norm $k \geq 2$, pick one of the decompositions according to some pairing and denote this $E^0(\{I_1,\ldots,I_m\})$. Thus for every element $E(\{I_1,\ldots,I_m\})$ of norm $k \geq 2$, there is a defining relation

$$E(\{I_1, \dots, I_m\}) = E^0(\{I_1, \dots, I_m\})$$

modulo (\mathcal{R}_2) , where $E^0(\{I_1, \ldots, I_m\})$ is a linear combination of terms of norm 0. Using these defining relations, all relations in (\mathcal{R}_2) become relations in W_0 , the subspace of W generated by $E(\{I_1, \ldots, I_m\})$ of norm 0. We next want to understand what these relations in W_0 are. Let $(\mathcal{R}_2)_2$ be the collections of all relations in (\mathcal{R}_2) where none of the I_5, \ldots, I_m is contained in $\{2, \ldots, m+1\}$ or $\{m+2, \ldots, 2m+1\}$. Next lemma simplifies our task.

Lemma 4.1. Let $E = E(\{I_1, ..., I_m\})$ be a basis element of norm k where $I_1, I_3, ..., I_{2k-1} \subseteq \{2, ..., m+1\}$ and $I_2, I_4, ..., I_{2k} \subseteq \{m+2, ..., 2m+1\}$ modulo $(\mathcal{R}_2)_2$. We have

$$\mathcal{D}_{\{(I_1, I_2), \dots, (I_{2k-1}, I_{2k})\}}(E) = \mathcal{D}_{\{(I_1, I_{\sigma(2)}), \dots, (I_{2k-1}, I_{\sigma(2k)})\}}(E)$$

for any $\sigma \in \text{Sym}\{2, 4, \dots, 2k\}$.

Proof. We prove the result by induction on k. If k = 1 then it is clear, since in this case σ is the identity. For k = 2 this is a direct consequence of $(\mathcal{R}_2)_2$. Now suppose that $k \geq 3$ and that the result is true for smaller values of k. Without

loss of generality, we can suppose $\sigma(2) = 4$. Using the induction hypothesis twice we have

$$\mathcal{D}_{\{(I_1, I_4), (I_3, I_{\sigma(4)}), (I_5, I_{\sigma(6)}), \dots, (I_{2k-1}, I_{\sigma(2k)})\}}(E)$$

= $\mathcal{D}_{\{(I_1, I_4), (I_3, I_2), (I_5, I_6), \dots, (I_{2k-1}, I_{2k})\}}(E)$
= $\mathcal{D}_{\{(I_1, I_2), (I_3, I_4), (I_5, I_6), \dots, (I_{2k-1}, I_{2k})\}}(E),$

as required.

Now take any relation in (\mathcal{R}_2) . Without loss of generality, we can assume (after reordering I_1, \ldots, I_m) that half of the elements of $I_5 \cup \cdots \cup I_m$ are in $\{2, \ldots, m+1\}$ and half in $\{m+2, \ldots, 2m+1\}$. We are using here the fact that the size of $(I_1 \cup I_2) \cap \{2, \ldots, m+1\}$ is between 0 and 4). We thus have now a relation of the form:

(1)
$$E(\{\{i_1, i_2\}, \{j_1, j_2\}, I_3, \dots, I_m\}) = -E(\{\{i_1, j_1\}, \{i_2, j_2\}, I_3, \dots, I_m\}) - E(\{\{i_1, j_2\}, \{i_2, j_1\}, I_3, \dots, I_m\}).$$

Suppose exactly k of I_5, \ldots, I_m are subsets of $\{2, \ldots, m+1\}$. Without loss of generality we can suppose that

$$I_5, I_7, \ldots, I_{2k+3} \subseteq \{2, \ldots, m+1\}$$
 and $I_6, I_8, \ldots, I_{2k+4} \subseteq \{m+2, \ldots, 2m+1\}$.

By Lemma 4.1, we know that modulo $(\mathcal{R}_2)_2$, relation (1) is equivalent to a sum of relations of the type

$$E(\{\{i_1, i_2\}, \{j_1, j_2\}, I_3, I_4, J_5, \dots, J_m\}) = -E(\{\{i_1, j_1\}, \{i_2, j_1\}, I_3, I_4, J_5, \dots, J_m\})$$

$$(2) -E(\{\{i_1, j_2\}, \{i_2, j_1\}, I_3, I_4, J_5, \dots, J_m\}).$$

where no J_5, \ldots, J_m is a subset of $\{2, \ldots, m+1\}$ or $\{m+2, \ldots, 2m+1\}$. Notice that all the relations (2) are in $(\mathcal{R}_2)_2$.

The conclusion is that all the relations in W_0 we are looking for, will be consequence of $(\mathcal{R}_2)_2$. Let J_5, \ldots, J_m be fixed and consider the collection of all the relations in $(\mathcal{R}_2)_2$ where $I_5 = J_5 \ldots I_m = J_m$.

Let $\overline{E}(\{I_1, I_2, I_3, I_4\}) = E(\{I_1, I_2, I_3, I_4, J_5, \dots, J_m\})$. Suppose $\{2, \dots, m+1\} \setminus (J_5 \cup \dots \cup J_m) = \{i_1, i_2, i_3, i_4, j_1, j_2, j_3, j_4\}$, where $i_1, i_2, i_3, i_4 \in \{2, \dots, m+1\}$ and $j_1, j_2, j_3, j_4 \in \{m+2, \dots, 2m+1\}$. We now go systematically through all possible types of $(\mathcal{R}_2)_2$ relations.

Case 1: $I_3 = \{i_3, j_3\}, I_4 = \{i_4, j_4\}$. We have

$$\bar{E}(\{\{i_1, i_2\}, \{j_1, j_2\}, \{i_3, j_3\}, \{i_4, j_4\}\}) = -\bar{E}(\{\{i_1, j_1\}, \{i_2, j_2\}, \{i_3, j_3\}, \{i_4, j_4\}\}) \\ -\bar{E}(\{\{i_1, j_2\}, \{i_2, j_1\}, \{i_3, j_3\}, \{i_4, j_4\}\})$$

These are defining relations.

Case 2: $I_3 = \{i_3, i_4\}, I_4 = \{j_3, j_4\}$. We have

$$\begin{split} \bar{E}(\{\{i_1, i_2\}, \{j_1, j_2\}, \{i_3, i_4\}, \{j_3, j_4\}\}) &= -\bar{E}(\{\{i_1, j_1\}, \{i_2, j_2\}, \{i_3, i_4\}, \{j_3, j_4\}\}) \\ &- \bar{E}(\{\{i_1, j_2\}, \{i_2, j_1\}, \{i_3, i_4\}, \{j_3, j_4\}\})) \\ &= \bar{E}(\{\{i_1, j_1\}, \{i_2, j_2\}, \{i_3, j_3\}, \{i_4, j_4\}\}) \\ &+ \bar{E}(\{\{i_1, j_1\}, \{i_2, j_2\}, \{i_3, j_4\}, \{i_4, j_3\}\}) \\ &+ \bar{E}(\{\{i_1, j_2\}, \{i_2, j_1\}, \{i_3, j_4\}, \{i_4, j_3\}\})) \\ &+ \bar{E}(\{\{i_1, j_2\}, \{i_2, j_1\}, \{i_3, j_4\}, \{i_4, j_3\}\})) \end{split}$$

If we instead pair $\{i_1,i_2\}$ and $\{j_3,j_4\}$ we get

$$\begin{split} \bar{E}(\{\{i_1, i_2\}, \{j_1, j_2\}, \{i_3, i_4\}, \{j_3, j_4\}\}) &= -\bar{E}(\{\{i_1, j_3\}, \{j_1, j_2\}, \{i_3, i_4\}, \{i_2, j_4\}\}) \\ &- \bar{E}(\{\{i_1, j_4\}, \{j_1, j_2\}, \{i_3, i_4\}, \{i_2, j_4\}\}) \\ &= \bar{E}(\{\{i_1, j_3\}, \{i_2, j_4\}, \{i_3, j_1\}, \{i_4, j_2\}\}) \\ &+ \bar{E}(\{\{i_1, j_4\}, \{i_2, j_3\}, \{i_3, j_1\}, \{i_4, j_2\}\}) \\ &+ \bar{E}(\{\{i_1, j_4\}, \{i_2, j_3\}, \{i_3, j_2\}, \{i_4, j_1\}\}) \\ &+ \bar{E}(\{\{i_1, j_4\}, \{i_2, j_3\}, \{i_3, j_2\}, \{i_4, j_1\}\})) \end{split}$$

From these two pairings we get

$$\sum_{\sigma \in \text{Sym}\{1,2\}, \tau \in \text{Sym}\{3,4\}} \bar{E}(\{\{i_1, j_{\sigma(1)}\}, \{i_2, j_{\sigma(2)}\}, \{i_3, j_{\tau(3)}\}, \{i_4, j_{\tau(4)}\}\})$$
$$= \sum_{\sigma \in \text{Sym}\{1,2\}, \tau \in \text{Sym}\{3,4\}} \bar{E}(\{\{i_1, j_{\tau(3)}\}, \{i_2, j_{\tau(4)}\}, \{i_3, j_{\sigma(1)}\}, \{i_4, j_{\sigma(2)}\}\}).$$

Case 3: $I_3 = \{j_2, j_3\}, I_4 = \{i_4, j_4\}$. We have

$$\begin{array}{lll} 0 &=& \bar{E}(\{\{i_1,i_2\},\{i_3,j_1\},\{j_2,j_3\},\{i_4,j_4\}\}) + \bar{E}(\{\{i_1,i_3\},\{i_2,j_1\},\{j_2,j_3\},\{i_4,j_4\}\}) \\ &\quad + \bar{E}(\{\{i_2,i_3\},\{i_1,j_1\},\{j_2,j_3\},\{i_4,j_4\}\}) \\ &=& -\bar{E}(\{\{i_1,j_2\},\{i_2,j_1\},\{i_3,j_3\},\{i_4,j_4\}\}) - \bar{E}(\{\{i_1,j_3\},\{i_2,j_2\},\{i_3,j_1\},\{i_4,j_4\}\}) \\ &\quad - \bar{E}(\{\{i_1,j_2\},\{i_2,j_1\},\{i_3,j_3\},\{i_4,j_4\}\}) - \bar{E}(\{\{i_1,j_3\},\{i_2,j_1\},\{i_3,j_2\},\{i_4,j_4\}\}) \\ &\quad - \bar{E}(\{\{i_1,j_1\},\{i_2,j_2\},\{i_3,j_3\},\{i_4,j_4\}\}) - \bar{E}(\{\{i_1,j_2\},\{i_2,j_3\},\{i_3,j_2\},\{i_4,j_4\}\}) \\ &\quad - \bar{E}(\{\{i_1,j_1\},\{i_2,j_2\},\{i_3,j_3\},\{i_4,j_4\}\}) - \bar{E}(\{\{i_1,j_2\},\{i_2,j_3\},\{i_3,j_2\},\{i_4,j_4\}\}) \\ \end{array}$$

This gives us the relation

$$\sum_{\sigma \in \text{Sym}\{1,2,3\}} \bar{E}(\{\{i_1, j_{\sigma(1)}\}, \{i_2, j_{\sigma(2)}\}, \{i_3, j_{\sigma(3)}\}, \{i_4, j_4\}\}) = 0$$

By symmetry the case of having $I_3 = \{i_2, i_3\}, I_4 = \{i_4, j_4\}$ works in a similar manner.

Case 4: $I_3 = \{j_1, j_2\}, I_4 = \{j_3, j_4\}$. We have

Modulo Case 2, the choice of pairings does not matter, therefore we get

Cancelling the signs gives

$$\sum_{\sigma \in \text{Sym}\{2,3,4\}} \bar{E}(\{\{i_1, j_1\}, \{i_2, j_{\sigma(2)}\}, \{i_3, j_{\sigma(3)}\}, \{i_4, j_{\sigma(4)}\}\}) + \sum_{\sigma \in \text{Sym}\{1,3,4\}} \bar{E}(\{\{i_1, j_2\}, \{i_2, j_{\sigma(1)}\}, \{i_3, j_{\sigma(3)}\}, \{i_4, j_{\sigma(4)}\}\}) = 0,$$

which follows from Case 3. We have seen that $W/W \cap J \cong W_0/W_0 \cap J$, as vector spaces, and $W_0 \cap J$ is generated by relations

(3)
$$\sum_{\sigma \in \text{Sym}\{1,2,3\}} E(\{\{i_1, j_{\sigma(1)}\}, \{i_2, j_{\sigma(2)}\}, \{i_3, j_{\sigma(3)}\}, \{i_4, j_4\}, \dots, \{i_m, j_m\}\}) = 0$$

and

$$(4) \sum_{\sigma \in \text{Sym}\{1,2\}, \tau \in \text{Sym}\{3,4\}} E(\{\{i_1, j_{\sigma(1)}\}, \{i_2, j_{\sigma(2)}\}, \{i_3, j_{\tau(3)}\}, \{i_4, j_{\tau(4)}\}, \{i_5, j_5\}, \dots, \{i_m, j_m\}\})$$

$$= \sum_{\sigma \in \text{Sym}\{1,2\}, \tau \in \text{Sym}\{3,4\}} \bar{E}(\{\{i_1, j_{\tau(3)}\}, \{i_2, j_{\tau(4)}\}, \{i_3, j_{\sigma(1)}\}, \{i_4, j_{\sigma(2)}\}, \{i_5, j_5\}, \dots, \{i_m, j_m\}\});$$

where $\{i_1, \ldots, i_m\} = \{2, 3, \ldots, m+1\}$ and $\{j_1, \ldots, j_m\} = \{m+2, \ldots, 2m+1\}$. We show that $W_0 \cap J$ is a proper subspace of W_0 by showing that all the relators in (3) and (4) lie in a subspace of co-dimension 1 in W_0 .

Let M_0 be the subspace of W_0 generated by all

$$E(\{\{2,\sigma(j_1)\},\ldots,\{m+1,\sigma(j_m)\}\})+E(\{\{2,j_1\},\ldots,\{m+1,j_m\}\})$$

with $\{j_1, \ldots, j_m\} = \{m + 2, \ldots, 2m + 1\}$ and where σ is a transposition in Sym $(\{m+2, \ldots, 2m+1\})$. Thus modulo M_0 we have $E(\{\{2, \sigma(m+2)\}, \ldots, \{m+1, \sigma(2m+1)\}\}) = \operatorname{sign}(\sigma)E(\{\{2, m+2\}, \ldots, \{m+1, 2m+1\}\})$. Notice that M_0 is of codimension 1 in W_0 and contains $W_0 \cap J$. For (3) this is because S_3 has equally many even and odd permutations and for (4) we can transform the LHS to the RHS using even permutations. Thus $W_0/W_0 \cap J \neq 0$ that implies that $W/W \cap J \neq 0$.

As a consequence we get the following main result of this section.

Theorem 4.2. The ideal generated by z in L is non-nilpotent.

5. The normal subgroup $\langle x \rangle^G$

In Section 2 we considered the largest locally finite *p*-group $G = \langle x, a_1, a_2, ... \rangle$ satisfying the following relations:

- (1) $\langle a_i \rangle^G$ is abelian for all $i \geq 1$;
- (2) $\langle x \rangle^G$ is metabelian;
- (3) $x^p = a_1^p = a_2^p = \dots = 1;$
- (4) if $s \neq x$ is a simple commutator in x, a_1, a_2, \ldots , where $t(s) \geq 2$ or $t(s) \leq -2$, then s = 1.

Now we use the Lie algebra L constructed in the Section 3 to get a group $H = \langle 1 + \operatorname{ad}(z), 1 + \operatorname{ad}(c_1), 1 + \operatorname{ad}(c_2), \ldots \rangle$. In this section we will show that H is a homomorphic image of G by proving that the relations (1)-(4) hold in H where x, a_1, a_2, \ldots are replaced by $1 + \operatorname{ad}(z), 1 + \operatorname{ad}(c_1), 1 + \operatorname{ad}(c_2), \ldots$. We will also use the work in Section 2 and Section 3 to show that $\langle 1 + \operatorname{ad}(z) \rangle^H$ is non-nilpotent from which follows that $\langle x \rangle^G$ is non-nilpotent.

Lemma 5.1. We have $ad(z)^2 = 0$ and ad(z) ad(w) ad(z) = 0 for any simple Lie product w in z, c_1, c_2, \ldots .

Proof. Let u be a simple Lie product in z, c_1, c_2, \ldots . Notice first that if u is non-trivial then the type of [u, z, z] is less than or equal to -3 and therefore [u, z, z] = 0. Turning to the second claim we know that [u, z] = 0 if u is of type different from 1. We can therefore assume that u is of type 1. Then if w is non-zero, we have that [u, z, w, z] is of type at most -2 and [u, z, w, z] = 0. \Box

Lemma 5.2. Let w_H be a simple group commutator in $1 + \operatorname{ad}(z)$, $1 + \operatorname{ad}(c_1)$, $1 + \operatorname{ad}(c_2)$,... and let w_L be the corresponding Lie commutator in z, c_1, c_2, \ldots . Then

$$w_H = 1 + \operatorname{ad}(w_L).$$

Proof. We show this by using induction on the weight of w_H . If the weight is 1 then this is obvious. Now suppose the weight is $k \ge 2$ and the result holds whenever the weight is smaller. We have $w_H = [u_H, v_H]$, where u_H, v_H are simple commutators of smaller weight in $1 + \operatorname{ad}(z), 1 + \operatorname{ad}(c_1), 1 + \operatorname{ad}(c_2), \ldots$. Let u_L, v_L be the corresponding simple commutators in z, c_1, c_2, \ldots . Using the induction hypothesis we have

$$w_{H} = [u_{H}, v_{H}] = (1 + \operatorname{ad}(u_{L}))^{-1}(1 + \operatorname{ad}(v_{L}))^{-1}(1 + \operatorname{ad}(u_{L}))(1 + \operatorname{ad}(v_{L}))$$

= $(1 - \operatorname{ad}(u_{L}))(1 - \operatorname{ad}(v_{L}))(1 + \operatorname{ad}(u_{L}))(1 + \operatorname{ad}(v_{L}))$ (using Lemma 5.1)
= $1 + \operatorname{ad}(u_{L}) \operatorname{ad}(v_{L}) - \operatorname{ad}(v_{L}) \operatorname{ad}(u_{L})$
= $1 + \operatorname{ad}([u_{L}, v_{L}])$
= $1 + \operatorname{ad}(w_{L}).$

Now we show that H satisfies the relations (1)-(4) for G where x, a_1, a_2, \ldots are replaced by $1 + \operatorname{ad}(z), 1 + \operatorname{ad}(c_1), 1 + \operatorname{ad}(c_2), \ldots$ More precisely, we have the following.

Proposition 5.3. Let H be as before. Then

- (1) $\langle 1 + \operatorname{ad}(c_i) \rangle^H$ is abelian for all $i \ge 1$;
- (2) $\langle 1 + \operatorname{ad}(z) \rangle^H$ is metabelian;
- (3) $(1 + \operatorname{ad}(z))^p = (1 + \operatorname{ad}(c_1))^p = (1 + \operatorname{ad}(c_2))^p = \dots = 1;$
- (4) if $s \neq 1 + ad(z)$ is a simple commutator in $1 + ad(z), 1 + ad(c_1), 1 + ad(c_2), \dots$, where $t(s) \geq 2$ or $t(s) \leq -2$, then s = 1.

Proof. (1) Let w_H be a simple group commutator in $1 + \operatorname{ad}(z)$, $1 + \operatorname{ad}(c_1)$, $1 + \operatorname{ad}(c_2)$,... with a repeated occurrence of $1 + \operatorname{ad}(c_i)$. Then $w_H = 1 + \operatorname{ad}(w_L)$ and as $\operatorname{Id}(c_i)$ is abelian, $\operatorname{ad}(w_L) = 0$ and thus $w_H = 1$. This implies that $\langle 1 + \operatorname{ad}(c_i) \rangle^H$ is abelian for all $i \geq 1$.

(2) Let u_H, v_H be simple commutators in $1 + \operatorname{ad}(z), 1 + \operatorname{ad}(c_1), 1 + \operatorname{ad}(c_2), \ldots$ where both have at least two occurrences of $1 + \operatorname{ad}(z)$. Then

$$[u_H, v_H] = 1 + \mathrm{ad}([u_L, v_L])$$

and since $\mathrm{Id}(z)$ is metabelian, then $[u_L, v_L] = 0$ and $[u_H, v_H] = 1$. It follows then that $\langle 1 + \mathrm{ad}(z) \rangle^H$ is metabelian.

(3) Clearly $(1 + \operatorname{ad}(z))^p = 1 + p \operatorname{ad}(z) = 1$ and $(1 + \operatorname{ad}(c_i))^p = 1 + p \operatorname{ad}(c_i) = 1$. (4) Let $w_H \neq 1 + \operatorname{ad}(z)$ be a simple commutator in $1 + \operatorname{ad}(z), 1 + \operatorname{ad}(c_1), 1 + \operatorname{ad}(c_2), \ldots$, where $t(w_H) \geq 2$ or $t(w_H) \leqslant -2$. Then, $w_H = 1 + \operatorname{ad}(w_L)$, where w_L is a Lie commutator in z, c_1, c_2, \ldots , where $t(w_L) \geq 2$ or $t(w_L) \leqslant -2$. Here $w_L = 0$ and so $w_H = 1$. This completes the proof. \Box

Theorem 5.4. The normal closure of 1 + ad(z) is not nilpotent.

Proof. Consider

$$\begin{split} & [[1 + \mathrm{ad}(z), 1 + \mathrm{ad}(c_1), 1 + \mathrm{ad}(c_2), 1 + \mathrm{ad}(c_3)], [1 + \mathrm{ad}(z), 1 + \mathrm{ad}(c_4), 1 + \mathrm{ad}(c_5)], \\ & \dots, [1 + \mathrm{ad}(z), 1 + \mathrm{ad}(c_{2m}), 1 + \mathrm{ad}(c_{2m+1})]] \\ & = 1 + \mathrm{ad}([[z, c_1, c_2, c_3], [z, c_4, c_5], \dots, [z, c_{2m}, c_{2m+1}]]). \end{split}$$
Notice that

$$1 + \mathrm{ad}([[z, c_1, c_2, c_3], [z, c_4, c_5], \dots, [z, c_{2m}, c_{2m+1}]]) \neq 1,$$

since, for example $[[z, c_1, c_2, c_3], [z, c_4, c_5], \ldots, [z, c_{2m}, c_{2m+1}], z]) \neq 0$. Thus, we have shown that there is a simple commutator of arbitrary weight m in (1+ad(z)) that is non-trivial. This finishes the proof.

Acknowledgement. We acknowledge the EPSRC (grant number 16523160) for support. We also thank Gareth Tracey and James Williams for participating in discussions at an early stage of this project. In particular we would like to thank the latter for some useful GAP calculations.

References

- [1] R. Baer, Engelsche Elemente Noetherscher Gruppen, Math. Ann. 133 (1957), 256-270.
- [2] Y. Bakhturin, Identical relations in Lie algebras, VNU Science Press, Utrecht, 1987.
- [3] W. Burnside, On an unsettled question in the theory of discontinous groups, Quart. J. Pure Appl. Math. 37 (1901), 230-238.
- [4] K. W. Gruenberg, The Engel elements of a soluble group, Illinois J. Math. 3 (1959), 151-169.
- [5] H. Heineken, Eine Bemerkung über engelshe Elemente, Arch. Math. 12 (1960), 176-178.

- [6] S. V. Ivanov, The free Burnside groups of sufficiently large exponents, Int. J. Algebra and Comp. 4 (1994), 1-308.
- [7] E. Jabara and G. Traustason, Left 3-Engel elements of odd order in groups, Proc. Am. Soc, to appear.
- [8] I. G. Lysenok, Infinite Burnside groups of even period, Izv. Math. 60 (1996), 453-654.
- [9] M. Newell, On right-Engel elements of length three, Proc. Royal Irish Ac. 96A, No.1 (1996), 17-24.
- [10] M. Noce, G. Tracey, and G. Traustason, A left 3-Engel element whose normal closure is not nilpotent, J Pure Appl. Algebra 224(3) (2020), 1092-1101.
- [11] I. N. Sanov, Solution of Burnside's Problem for Exponent Four, Leningrad State Univ., Ann. Maths. Ser. 10 (1940), 166-170.
- [12] G. Tracey and G. Traustason, Left 3-Engel elements in groups of exponent 60, Int. J. Algebra and Comp. 28 No. 4 (2018), 673-695.
- [13] G. Traustason, Left 3-Engel elements in groups of exponent 5, J. Algebra, 414 (2014), 41-71.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF BATH, CLAVERTON DOWN, BATH BA2 7AY, UNITED KINGDOM

E-mail address: ah926@bath.ac.uk

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF BATH, CLAVERTON DOWN, BATH BA2 7AY, UNITED KINGDOM

E-mail address: mn670@bath.ac.uk

Department of Mathematical Sciences, University of Bath, Claverton Down, Bath BA2 7AY, United Kingdom

E-mail address: gt223@bath.ac.uk