

Uniform and Partially Uniform Redistribution Rules

Florentin Smarandache
Department of Mathematics
University of New Mexico
Gallup, NM 87301, U.S.A.
smarand@unm.edu

Jean Dezert
ONERA
The French Aerospace Lab,
F-91761 Palaiseau, France.
Jean.Dezert@onera.fr

Abstract - *This short paper introduces two new fusion rules for combining quantitative basic belief assignments. These rules although very simple have not been proposed in literature so far and could serve as useful alternatives because of their low computation cost with respect to the recent advanced Proportional Conflict Redistribution rules developed in the DSmT framework.*

Keywords: Uniform Redistribution Rule, Partially Uniform Redistribution Rule, information fusion, belief functions, Dezert-Smarandache Theory (DSmT).

1 Introduction

Since the development of DSmT (Dezert-Smarandache Theory) in 2002 [4, 5], a new look for information fusion in the framework of belief has been proposed which covers many aspects related to the fusion of uncertain and conflicting beliefs. Mainly, the fusion of quantitative or qualitative belief functions of highly uncertain and conflicting sources of evidence with theoretical advances in belief conditioning rules. The Shafer's milestone book [3] introducing the concept of belief functions and Dempster's rule of combination of beliefs has been the important step towards non probabilistic reasoning approach, aside Zadeh's fuzzy logic [6, 8]. Since Shafer's seminal work, many alternatives have been proposed to circumvent limitations of Dempster's rule pointed out first by Zadeh in [7] (see [2] and [5] for a review). The Proportional Conflict Redistribution rule number 5 (PCR5) [5] is one of the most efficient alternative to Dempster's rule which can be used both in Dempster-Shafer Theory (DST) as well as in DSmT. The simple idea behind PCR5 is to redistribute every partial conflict only onto propositions which are truly involved in the partial conflict and proportionally to the corresponding belief mass assignment of each source generating this conflict. Although very efficient and appealing, the PCR5 rule suffers of its relative complexity in implementation and in some cases, it is required to use simpler (but less precise) rule of combination which requires only a low complexity. For this purpose, we herein present two new cheap alternatives for combination of basic belief assignments (bba's): the Uniform Redistribution Rule (URR) and the Partially Uniform Redistribution Rule (PURR). In the sequel, we assume the reader familiar with the basics of DSmT, mainly with the definition and notation of hyper-power set G^Θ and also bba's defined over hyper-power set. Basics of DSmT can be found in chapter 1 of [4] which is freely downloadable on internet.

2 Uniform Redistribution Rule

Let's consider a finite and discrete frame of discernment Θ , its hyper-power set G^Θ (i.e. Dedekind's lattice) and two quantitative basic belief assignments $m_1(\cdot)$ and $m_2(\cdot)$ defined on G^Θ expressed by two independent sources of evidence.

The Uniform Redistribution Rule (URR) consists in redistributing the total conflicting mass k_{12} to all focal elements of G^Θ generated by the consensus operator. This way of redistributing mass is very simple and URR is different from Dempster's rule of combination [3], because Dempster's rule redistributes the total conflict proportionally with respect to the masses resulted from the conjunctive rule of non-empty sets. PCR5 and PCR4 [5] do proportional redistributions of partial conflicting masses to the sets involved in the conflict. Here it is the URR formula for two sources: $\forall A \neq \emptyset$, one has

$$m_{12URR}(A) = m_{12}(A) + \frac{1}{n_{12}} \sum_{\substack{X_1, X_2 \in G^\Theta \\ X_1 \cap X_2 = \emptyset}} m_1(X_1)m_2(X_2) \quad (1)$$

where $m_{12}(A)$ is the result of the conjunctive rule applied to belief assignments $m_1(\cdot)$ and $m_2(\cdot)$, and $n_{12} = \text{Card}\{Z \in G^\Theta, m_1(Z) \neq 0 \text{ or } m_2(Z) \neq 0\}$.

For $s \geq 2$ sources to combine: $\forall A \neq \emptyset$, one has

$$m_{12\dots sURR}(A) = m_{12\dots s}(A) + \frac{1}{n_{12\dots s}} \sum_{\substack{X_1, X_2, \dots, X_s \in G^\Theta \\ X_1 \cap X_2 \cap \dots \cap X_s = \emptyset}} \prod_{i=1}^s m_i(X_i) \quad (2)$$

where $m_{12\dots s}(A)$ is the result of the conjunctive rule applied to $m_i(\cdot)$, for all $i \in \{1, 2, \dots, s\}$ and

$$n_{12\dots s} = \text{Card}\{Z \in G^\Theta, m_1(Z) \neq 0 \text{ or } m_2(Z) \neq 0 \text{ or } \dots \text{ or } m_s(Z) \neq 0\}$$

As alternative, we can also consider the cardinal of the ensemble of sets whose masses resulted from the conjunctive rule are non-null, i.e. the cardinality of the core of conjunctive consensus:

$$n_{12\dots s}^c = \text{Card}\{Z \in G^\Theta, m_{12\dots s}(Z) \neq 0\}$$

We denote this modified version of URR as MURR in the sequel.

3 Example for URR and MURR

Example for URR: Let's consider $\Theta = \{A, B, C\}$ with the DSm hybrid model as shown on the Figure 1. In




Fig. 1: Hybrid model for $\Theta = \{A, B, C\}$.

this hybrid model $C \cap (A \cup B) = \emptyset$ (therefore $A \cap C = \emptyset$ and $B \cap C = \emptyset$). We consider also the following two belief assignments

$$\begin{aligned} m_1(A) &= 0.4 & m_1(B) &= 0.2 & m_1(A \cup B) &= 0.4 \\ m_2(A) &= 0.2 & m_2(C) &= 0.3 & m_2(A \cup B) &= 0.5 \end{aligned}$$

then the conjunctive operator provides for this DSm hybrid model a consensus on A , B , C , $A \cup B$, and $A \cap B$ with supporting masses

$$m_{12}(A) = 0.36 \quad m_{12}(B) = 0.10 \quad m_{12}(A \cup B) = 0.20 \quad m_{12}(A \cap B) = 0.04$$

and partial conflicts between two sources on $A \cap C$, $B \cap C$ and $C \cap (A \cup B)$ with

$$m_{12}(A \cap C) = 0.12 \quad m_{12}(B \cap C) = 0.06 \quad m_{12}(C \cap (A \cup B)) = 0.12$$

Then with URR, the total conflicting mass

$$m_{12}(A \cap C) + m_{12}(B \cap C) + m_{12}(C \cap (A \cup B)) = 0.12 + 0.06 + 0.12 = 0.30$$

is uniformly (i.e. equally) redistributed to A , B , C and $A \cup B$ because the sources support only these propositions. That is $n_{12} = 4$ and thus $0.30/n_{12} = 0.075$ is added to $m_{12}(A)$, $m_{12}(B)$, $m_{12}(C)$ and $m_{12}(A \cup B)$ with URR. One finally gets:

$$\begin{aligned} m_{12URR}(A) &= m_{12}(A) + \frac{0.30}{n_{12}} = 0.36 + 0.075 = 0.435 \\ m_{12URR}(B) &= m_{12}(B) + \frac{0.30}{n_{12}} = 0.10 + 0.075 = 0.175 \\ m_{12URR}(C) &= m_{12}(C) + \frac{0.30}{n_{12}} = 0.00 + 0.075 = 0.075 \end{aligned}$$

$$m_{12URR}(A \cup B) = m_{12}(A \cup B) + \frac{0.30}{n_{12}} = 0.20 + 0.075 = 0.275$$

while the others remain the same. That is $m_{12URR}(A \cap B) = 0.04$. Of course, one has also

$$m_{12URR}(A \cap C) = m_{12URR}(B \cap C) = m_{12URR}(C \cap (A \cup B)) = 0$$

Example for MURR: Let's consider the same frame, same model and same bba as in previous example. In this case the total conflicting mass 0.30 is uniformly redistributed to the sets A , B , $A \cup B$, and $A \cap B$ only, i.e. to the sets whose masses, after applying the conjunctive rule to the given sources, are non-zero. Thus $n_{12} = 4$, and $0.30/4 = 0.075$. Hence:

$$\begin{aligned} m_{12MURR}(A) &= 0.36 + 0.075 = 0.435 \\ m_{12MURR}(B) &= 0.10 + 0.075 = 0.175 \\ m_{12MURR}(A \cup B) &= 0.20 + 0.075 = 0.275 \\ m_{12MURR}(A \cap B) &= 0.04 + 0.075 = 0.115 \end{aligned}$$

4 Partially Uniform Redistribution Rule

It is also possible to do a uniformly partial redistribution, i.e. to uniformly redistribute the conflicting mass only to the sets involved in the conflict. For example, if $m_{12}(A \cap B) = 0.08$ and $A \cap B = \emptyset$, then 0.08 is equally redistributed to A and B only, supposing A and B are both non-empty, so 0.04 assigned to A and 0.04 to B .

$\forall A \neq \emptyset$, one has the Partially Uniform Redistribution Rule (PURR) for two sources

$$m_{12PURR}(A) = m_{12}(A) + \frac{1}{2} \sum_{\substack{X_1, X_2 \in G^\Theta \\ X_1 \cap X_2 = \emptyset \\ X_1 = A \text{ or } X_2 = A}} m_1(X_1)m_2(X_2) \quad (3)$$

where $m_{12}(A)$ is the result of the conjunctive rule applied to belief assignments $m_1(\cdot)$ and $m_2(\cdot)$.

For $s \geq 2$ sources to combine: $\forall A \neq \emptyset$, one has

$$m_{12\dots sPURR}(A) = m_{12\dots s}(A) + \frac{1}{s} \sum_{\substack{X_1, X_2, \dots, X_s \in G^\Theta \\ X_1 \cap X_2 \cap \dots \cap X_s = \emptyset \\ \text{at least one } X_j = A, j \in \{1, \dots, s\}}} \text{Card}_A(\{X_1, \dots, X_s\}) \prod_{i=1}^s m_i(X_i) \quad (4)$$

where $\text{Card}_A(\{X_1, \dots, X_s\})$ is the number of A 's occurring in $\{X_1, X_2, \dots, X_s\}$.

If $A = \emptyset$, $m_{12PURR}(A) = 0$ and $m_{12\dots sPURR}(A) = 0$.

5 Example for PURR

Let's take back the example of section 3. Based on PURR, $m_{12}(A \cap C) = 0.12$ is redistributed as follows: 0.06 to A and 0.06 to C ; $m_{12}(B \cap C) = 0.06$ is redistributed as follows: 0.03 to B and 0.03 to C ; and $m_{12}(C \cap (A \cup B)) = 0.12$ is redistributed in this way: 0.06 to C and 0.06 to $A \cup B$. Therefore we finally get

$$\begin{aligned} m_{12PURR}(A) &= m_{12}(A) + \frac{0.12}{2} = 0.36 + 0.06 = 0.42 \\ m_{12PURR}(B) &= m_{12}(B) + \frac{0.06}{2} = 0.10 + 0.03 = 0.13 \\ m_{12PURR}(C) &= m_{12}(C) + \frac{0.12}{2} + \frac{0.06}{2} + \frac{0.12}{2} = 0.15 \\ m_{12PURR}(A \cup B) &= m_{12}(A \cup B) + \frac{0.12}{2} = 0.20 + 0.06 = 0.26 \end{aligned}$$

while the others remain the same. That is $m_{12PURR}(A \cap B) = 0.04$. Of course, one has also

$$m_{12PURR}(A \cap C) = m_{12PURR}(B \cap C) = m_{12PURR}(C \cap (A \cup B)) = 0$$

6 Neutrality of vacuous belief assignment

Both URR (with MURR included) and PURR are commutative and quasi-associative, and they verify the neutrality of Vacuous Belief Assignment (VBA): since any bba $m_1(\cdot)$ combined with the VBA defined on any frame $\Theta = \{\theta_1, \dots, \theta_n\}$ by $m_{VBA}(\theta_1 \cup \dots \cup \theta_n) = 1$, using the conjunctive rule, gives $m_1(\cdot)$, so no conflicting mass is needed to transfer.

7 Conclusion

Two new simple rules of combination have been presented in the framework of DSMT which have a lower complexity than PCR5. These rules are very easy to implement but from a theoretical point of view remain less precise in their transfer of conflicting beliefs since they do not take into account the proportional redistribution with respect to the mass of each set involved in the conflict. So we cannot reasonably expect that URR or PURR outperforms PCR5 but they may hopefully appear as good enough in some specific fusion problems when the level of total conflict is not important. PURR does a more refined redistribution than URR and MURR but it requires a little more calculation.

References

- [1] Dezert J., Smarandache F., *DSMT: A new paradigm shift for information fusion*, Proceedings of Cogis 06 Conference, Paris, March 15-17, 2006.
- [2] Sentz K., Ferson S., *Combination of evidence in Dempster-Shafer Theory*, SANDIA Tech. Report, SAND2002-0835, 96 pages, April 2002.
<http://www.sandia.gov/epistemic/Reports/SAND2002-0835.pdf>
- [3] Shafer G., *A Mathematical Theory of Evidence*, Princeton University Press, Princeton, NJ, 1976.
- [4] Smarandache F., Dezert J. (Editors), *Applications and Advances of DSMT for Information Fusion (Collected works)*, Amer. Res. Press, Rehoboth, 2004,
<http://www.gallup.unm.edu/~smarandache/DSMT-book1.pdf>.
- [5] Smarandache F., Dezert J. (Editors), *Applications and Advances of DSMT for Information Fusion (Collected works)*, Vol. 2, Amer. Res. Press, Rehoboth, 2006.
<http://www.gallup.unm.edu/~smarandache/DSMT-book2.pdf>.
- [6] Zadeh L., *Concept of a linguistic variable and its application to approximate reasoning*, Information Sciences, Part 1 & 2, Vol. 8, pp. 199–249, 301–357, 1975, Part 3, Vol.9 pp. 43-80, 1976.
- [7] Zadeh L., *On the validity of Dempster's rule of combination*, Memo M 79/24, Univ. of California, Berkeley, 1979.
- [8] Zadeh L., *A Theory of Approximate Reasoning*, Machine Intelligence, Vol. 9, pp. 149-194, 1979.