Thermodynamic formalism for continuous-time quantum Markov semigroups: the detailed balance condition, entropy, pressure and equilibrium quantum processes

Jader E. Brasil, Josue Knorst and Artur O. Lopes

January 14, 2022

Abstract

 $M_n(\mathbb{C})$ denotes the set of n by n complex matrices. Consider continuous time quantum semigroups $\mathcal{P}_t = e^{t\mathcal{L}}, t \geq 0$, where \mathcal{L} : $M_n(\mathbb{C}) \to M_n(\mathbb{C})$ is the infinitesimal generator. If we assume that $\mathcal{L}(I) = 0$, we will call $e^{t\mathcal{L}}, t \geq 0$ a quantum Markov semigroup. Given a stationary density matrix $\rho = \rho_{\mathcal{L}}$, for the quantum Markov semigroup $\mathcal{P}_t, t \geq 0$, we can define a continuous time stationary quantum Markov process, denoted by $X_t, t \ge 0$. Given an *a priori* Laplacian operator $\mathcal{L}_0: M_n(\mathbb{C}) \to M_n(\mathbb{C})$, we will present a natural concept of entropy for a class of density matrices on $M_n(\mathbb{C})$. Given an Hermitian operator $A: \mathbb{C}^n \to \mathbb{C}^n$ (which plays the role of an Hamiltonian), we will study a version of the variational principle of pressure for A. A density matrix ρ_A maximizing pressure will be called an equilibrium density matrix. From ρ_A we will derive a new infinitesimal generator \mathcal{L}_A . Finally, the continuous time quantum Markov process defined by the semigroup $\mathcal{P}_t = e^{t\mathcal{L}_A}, t \geq 0$, and an initial stationary density matrix, will be called the continuous time equilibrium quantum Markov process for the Hamiltonian A. It corresponds to the quantum thermodynamical equilibrium for the action of the Hamiltonian A.

Key words: continuous time quantum Markov process, Lindbladian, detailed balance condition, entropy, pressure, equilibrium quantum processes MSC2010: 81P45; 81p47; 81p16; 81P17

1 Introduction

We are interested in continuous time stationary quantum Markov process which corresponds to equilibrium for a quantum bath (interacting with a quantum system) under the action of a certain given Hamiltonian. Therefore, our results concern continuous-time quantum channels.

In [5] the authors present a detailed study of a nice version of the detailed balance condition for a continuous-time quantum Markov semigroup on $M_n(\mathbb{C})$, $n \in \mathbb{N}$ (see also [9], [8] and [16] for related results). In Theorem 4.2 in [5] it is explained that the detailed balance condition for a classical continuous-time Markov Chain, with values on a finite state space, corresponds to the commutative part of the dynamical evolution of the continuoustime quantum Markov semigroup. Results of [5] are used here in an essential way.

On page 75 in Section 9 in [10], and also on page 114 in Section 5 in [17], for a classical continuous-time Markov Chain satisfying the detailed balance condition, a deviation function (which is a form of entropy) is introduced and a variational principle (in some sense a form of maximizing pressure) is considered (see expression 9.18 in page 76 in [10]). We would like to extend the results obtained for the classical commutative setting to the non-commutative setting of quantum Markov semigroups satisfying the detailed balance condition as described in [5].

We will present a natural concept of entropy for a class of density matrices (see Section 3). We point out that the dynamics of the flow in the set of matrices is encapsulated on the infinitesimal generator and the entropy we consider here is at the level of this linear operator. In this sense, this concept of entropy has no *direct* dynamical content. Our setting is the quantum channel version of the classical ones considered in [10] and [17].

After introducing entropy we will study a version of the variational principle of pressure and its relation to an eigenvalue problem for a certain type of transfer operator (see Section 5 and expression (32) in Section 6). In classical Thermodynamic Formalism, the Ruelle operator plays this role. The Ruelle Theorem describes a relation of equilibrium states with a corresponding eigenvalue problem (see [15]). The Ruelle operator is an infinite-dimensional version of the Perron-Frobenius operator. The transfer operator we consider here is not exactly an extension of the concept of Ruelle operator. A density matrix maximizing pressure will be called an equilibrium density matrix. We will provide examples in Section 5.

Our results are in some sense the quantum analogous of the reasoning delineated in [2], [13] and [11], which considered the dynamics of continuous-time dynamics (a flow) in the Skorohod space.

Our definition of entropy is also different from the ones in [3] and [4] which considered quantum channels in discrete-time dynamics.

Taking into account the concept and the notation described in section 5 in [5] we will denote by \mathcal{L}_0 (the Laplacian) the generator of the heat semigroup. We will choose a special \mathcal{L}_0 (see Definition 4) which will play the role of the *a priori* Laplacian. Our special choice of \mathcal{L}_0 is the analogous of taking the normalized counting probability as the *a priori* probability in the classical definition of Kolmogorov-Shannon entropy (see discussion in [14]).

From this \mathcal{L}_0 (which is fixed from now on) we will be able to define the detailed balanced condition (as described in [5]) and the Laplacian-entropy.

Definition 1. Given a density operator ρ define the Laplacian-entropy (entropy for short) by

$$h(\rho) = Tr \left[\rho^{1/2} \mathcal{L}_0^{\dagger}(\rho^{1/2}) \right], \tag{1}$$

where we set \mathcal{L}_0 by Definition 4.

Our definitions of entropy and pressure are quite natural. They are the non commutative extension of the concepts considered in the classical setting of continuous-time Markov chains as described by M. Kac in expressions (9.16) and (9.18) in Section 9 in [10] and by D. W. Strook in Section 5 in [17].

Expression (5.12) in [17] defines the so-called *rate function* I in the setting of classical continuous-time Markov Processes taking values on the compact metric space E. If L is the infinitesimal generator, then (5.12) means

$$I(\nu) = -\inf_{u>0, u\in C(E)} \int \frac{Lu}{u} d\nu,$$

where C(E) is the set of real continuous functions defined on E.

Later, for reversible processes, the above formula simplifies to expression (5.18) in [17], which claims

$$I(\nu) = -\int \phi^{1/2} L \phi^{1/2} d\mu, \quad \phi = \frac{d\nu}{d\mu}$$

In [17] it is used the term symmetric operator but in other contexts, this would correspond to conditions like reversibility or the detailed balanced condition.

Under the detailed balanced condition, in the quantum channel context, one should replace the role of L by the generator of a QMS, which is usually denoted by \mathcal{L} (the Lindbladian). Probabilities are replaced by densities ρ (states). In this case, (5.12) in [17] corresponds here to

$$I(\rho) = -\inf_{U>0} \operatorname{Tr}(\rho \, U^{-1} \, \mathcal{L}_0 \, U),$$

where the infimum is taken over the positive matrices $U \in M_n$.

In [17] (see also [10]) the variational principle is taken as

$$\lambda(V) = \sup_{\text{prob }\nu} \left(\int V d\nu - I(\nu) \right)$$

where $\lambda(V)$ is the main eigenvalue of a certain operator.

Denote $D_n = \{ \rho \ge 0 : \text{Tr}(\rho) = 1 \}.$

In section 5 we consider an analogous problem: given an Hermitian operator $A : \mathbb{C}^n \to \mathbb{C}^n$, we consider the variational problem:

$$P_A = \sup_{\rho \in D_n} \{ h(\rho) + \operatorname{Tr}(A\rho) \}.$$
(2)

A matrix ρ_A maximizing P_A will be called an equilibrium density the operator for A. We call P(A) the Laplacian-pressure (pressure for short) of the Hermitian operator $A : \mathbb{C}^n \to \mathbb{C}^n$.

A connection of P_A of expression (2) with the eigenvalue of a certain linear operator is described in expression (32) in Section 6 (see also (33)). In this way we get all elements for establishing a continuous time quantum channel version of the classical Ruelle operator (see [15], [14], [2], [13]).

Our definition of entropy has a difference of sign when compared with the setting of [17], so we wonder if there exists a connection between $h(\rho)$ and $-I(\rho)$. In section 7 we will show this connection in the special case of the heat-semigroup with the *a priori* generator \mathcal{L}_0 defined on section 3. We will show that:

Theorem 2. Given the density matrix ρ , then

$$h(\rho) = \inf_{A>0} Tr(\rho A^{-1} \mathcal{L}_0(A)).$$

Following [5], we will present in Section 8 the classical Markov Chain associated with a continuous-time quantum channel and we will provide examples.

The present work is part of the PhD. thesis of J. Brasil and J. Knorst in Graduate Program in Math. in UFRGS.

2 An outline of the main prerequisites

Given a linear operator $A : \mathbb{C} \to \mathbb{C}$, its dual (with respect to the canonical inner product), is denoted by $A^* : \mathbb{C} \to \mathbb{C}$.

Denote by $M_n(\mathbb{C}) = M_n$ the set of n by n complex matrices with the GNS inner product $\langle A, B \rangle = \text{Tr}(A^*B)$. Given a linear operator $T: M_n \to M_n$,

its dual with respect to this inner product is denoted by $T^{\dagger}: M_n \to M_n$. That is, for all matrices A, B we get

$$\langle T(A), B \rangle = \langle A, T^{\dagger}(B) \rangle$$
.

We denote by 1 the diagonal matrix with entries $\frac{1}{n}$. Then, 1 is a density matrix and also the unity of the C^* -algebra $M_n(\mathbb{C})$. We denote by \mathfrak{G}_+ the set of invertible density matrices (operators) $\rho: M_n \to M_n$.

We will consider **continuous time quantum semigroups** (QS) the ones given by $\mathcal{P}_t = e^{t\mathcal{L}}$, $t \geq 0$, where $\mathcal{L} : M_n(\mathbb{C}) \to M_n(\mathbb{C})$ is the infinitesimal generator (see Definition 5.5.1 and also Section 9.3.2 in [6]). The linear operator \mathcal{L} should satisfy the conditional complete positivity property (see section 5 in [6] or section 6 in [18]). We assume that $\mathcal{L}(A^*) = (\mathcal{L}(A))^*$, for all $A \in M_n(\mathbb{C})$. Given a selfadjoint matrix $A \in M_n(\mathbb{C})$, the dynamical evolution $t \to e^{t\mathcal{L}}(A)$ is called the Heisenberg dynamical evolution. Given a density matrix $\rho \in M_n(\mathbb{C})$, the dynamical evolution $t \to e^{t\mathcal{L}^{\dagger}}(\rho)$ is called the Schrödinger dynamical evolution.

If we assume that $\mathcal{L}(I) = 0$, we will call $\mathcal{P}_t = e^{t\mathcal{L}}$, $t \ge 0$, the continuous time **quantum Markov semigroup** (QMS) associated to \mathcal{L} (see Definition 5.5.2 and also section 7 in [6]). It is known that in this case $e^{t\mathcal{L}}(\mathbf{1}) = \mathbf{1}$, for all $t \ge 0$. Continuous time quantum Markov semigroups provide a convenient mathematical description of the irreversible dynamics of an open quantum system.

If $\rho = \rho_{\mathcal{L}}$ is such that $\mathcal{L}^{\dagger}(\rho) = 0$, then for all $t \ge 0$, we get $e^{t\mathcal{L}^{\dagger}}(\rho) = \rho$ and we say that ρ is the **stationary density matrix for the continuous time quantum Markov semigroup** with infinitesimal generator \mathcal{L} . Section 9.4 in [6] presents a discussion on the uniquenes of the stationary matrix ρ .

We call $X_t, t \ge 0$, the **continuous time quantum Markov process** (QMP) associated to the infinitesimal generator \mathcal{L} , the process associated to the pair $(e^{t\mathcal{L}}, \rho_{\mathcal{L}}), t \ge 0$. We can ask questions about ergodicity for such process (see Section 11 in [6]).

Given the (QMP) associated to \mathcal{L} and the stationary density operator ρ , take an observable (a self-adjoint matrix) $A \in M_n$. Then, we get that

$$t \to \operatorname{Tr}(\rho e^{t\mathcal{L}}(A))$$

describes the time evolution of the expected value of the observable A.

We say that \mathcal{L} is irreducible if for every non-zero matrix $A \geq 0$, and every strictly positive t > 0, we have $e^{t\mathcal{L}}(A) > 0$. We will also assume that \mathcal{L} is irreducible (see Sections 10 and 11 in [6]).

Given $\sigma \in \mathfrak{G}_+$, consider the inner product $\langle , \rangle_{\sigma}$ in the set of matrices in M_n given by $\langle A, B \rangle_{\sigma} = \text{Tr} (A^* B \sigma) = \langle A, B \sigma \rangle$. **Definition 3.** Given $\sigma \in \mathfrak{G}_+$, we say that the QMS $e^{t\mathcal{L}}, t \geq 0$, satisfies the σ -detailed balance condition if \mathcal{L} is symmetric with respect to $\langle , \rangle_{\sigma}$. That is, for all matrices $A, B \in M_n$ we get

$$< \mathcal{L}(A), B >_{\sigma} = < A, \mathcal{L}(B) >_{\sigma}$$
.

Given $\sigma \in \mathfrak{G}_+$, if the QMS $e^{t\mathcal{L}}, t \geq 0$, satisfies the σ -detailed balance condition, then, σ is stationary for the evolution of the semigroup $e^{t\mathcal{L}^{\dagger}}, t \geq 0$ (see Lemma 10).

The explicit form of the infinitesimal generator of a continuous-time quantum Markov semigroups satisfying the detailed balance condition is described by expression (3.4) in [5] (see our expression (13)).

Definition 4. We denote \mathcal{L}_0 the infinitesimal generator satisfying d.b.c. where we take $\sigma = \mathbf{1}$. \mathcal{L}_0 will be called the Laplacian (see Section 3 in [5]).

The semigroup $\mathcal{P}_t = e^{t\mathcal{L}_0}, t \geq 0$, describes the unperturbed continuous time quantum channel.

Given the *a priori* Laplacian operator $\mathcal{L}_0 : M_n(\mathbb{C}) \to M_n(\mathbb{C})$, we will present a natural concept of entropy for a class of density matrices ρ on $M_n(\mathbb{C})$ (see Definition 5).

Given a Hermitian operator $A : \mathbb{C}^n \to \mathbb{C}^n$ (which plays the role of minus the Hamiltonian), we will consider in Section 5 a variational principle of pressure for A, which is given by Definition 12.

A density matrix ρ_A maximizing pressure will be called an equilibrium density matrix for A. This matrix (in fact $\rho_A^{1/2}$) will satisfy an eigenvalue property for a certain linear operator \mathfrak{L}_A to be described in Section 6. From ρ_A we will derive a new infinitesimal generator \mathcal{L}_A . Finally, the continuoustime quantum Markov Process X_t , $t \geq 0$, associated to $\mathcal{P}_t = e^{t\mathcal{L}_A}$, $t \geq$ 0, and ρ_A , will be called the continuous-time equilibrium quantum Markov semigroup for the Hamiltonian A. This new process describes a continuoustime quantum channel after the perturbation by the selfadjoint operator A.

3 The heat semigroup and entropy of density operators

In this section the inner product in M_n is $\langle A, B \rangle = \text{Tr} (A^* B)$. Denote by $e_j, j = 1, ..., n$, the canonical base in \mathbb{C}^n , and by

$$\mathfrak{I}_{i,j} = |e_i\rangle\langle e_j|: \mathbb{C}^n \to \mathbb{C}^n,$$

where i, j = 1, ..., n. Note that $\mathfrak{I}_{i,j}^* = |e_j\rangle\langle e_i|$.

We denote by $\mathfrak{I}_{i,j}$ the matrix which is zero in all entries, up to the entry i, j, where it has the value 1.

We denote by $\mathbf{1}$ the operator identity I times $\frac{1}{n}$. The matrix $\mathbf{1}$ describes an invertible density operator.

 \mathcal{L}_0 denotes the infinitesimal generator satisfying d.b.c. for $\sigma = 1$.

One can show that $\mathfrak{I}_{i,j}$, i, j = 1, ..., n, is an orthonormal basis for \mathcal{L}_0 associated to the eigenvalue 0.

Following section 5 in [5] we call \mathcal{L}_0 (the Laplacian) the generator of the heat semigroup (the Laplacian)

$$A \to \mathcal{L}_0(A) = \sum_{i,j=1}^n \left(V_{i,j}^* \left[A, V_{i,j} \right] + \left[V_{i,j}, A \right] V_{i,j}^* \right)$$
(3)

This operator is negative-semi definite. (see page 1827 in [5] and also (13) and (15) of next section. Note that $\mathcal{L}_0(I) = 0$.

One can show that $\mathcal{L}_0^{\dagger} = \mathcal{L}_0$ and Tr $(\mathbf{1}\mathcal{L}_0(A)) = 0$, for all $A \in M_n$. Note that

$$\mathcal{L}_{0}^{\dagger}(\rho) = \sum_{i,j=1}^{n} \left(\left[V_{i,j} \, \rho, V_{i,j}^{*} \right] + \left[V_{i,j}, \rho \, V_{i,j}^{*} \right] \right). \tag{4}$$

 $\sigma = \mathbf{1}$ is invariant for the flow $e^{t\mathcal{L}_0^{\dagger}}$.

Definition 5. Given a density operator ρ define the Laplacian-entropy

$$h(\rho) = Tr[\rho^{1/2} \mathcal{L}_0^{\dagger}(\rho^{1/2})]$$
(5)

This definition is consistent with expression (5.18) on page 113 in [17].

Our main result in this section is the explicit expression for entropy to be described by Proposition 7.

First, we want to show the following Lemma:

Lemma 6. h(1) = 0.

Proof. From (4)

$$\mathcal{L}_{0}^{\dagger}(\rho^{1/2}) = \sum_{i,j=1}^{n} \left(\left[V_{i,j} \, \rho^{1/2}, V_{i,j}^{*} \right] + \left[V_{i,j}, \rho^{1/2} \, V_{i,j}^{*} \right] \right).$$

Then,

$$\mathcal{L}_{0}^{\dagger}(\mathbf{1}^{1/2}) = \sum_{i,j=1}^{n} \left(\left[V_{i,j} \, \mathbf{1}^{1/2}, V_{i,j}^{*} \right] + \left[V_{i,j}, \mathbf{1}^{1/2} \, V_{i,j}^{*} \right] \right) = \mathbf{1}^{1/2} \, 2 \sum_{i,j=1}^{n} \left(\left[V_{i,j}, V_{i,j}^{*} \right] \right)$$

Note that

$$\begin{split} [V_{i,j}, V_{i,j}^*] = \\ | \, e_i \, \rangle \langle \, e_i \, | - | \, e_j \, \rangle \langle \, e_j \, | \, . \end{split}$$

For each pair (i, j) there is a correspondent (j, i). From this follows that

$$\mathcal{L}_{0}^{\dagger}(\mathbf{1}^{1/2}) = \mathbf{1}^{1/2} 2 \sum_{i,j=1}^{n} |e_{i}\rangle \langle e_{i}| - |e_{j}\rangle \langle e_{j}| = 0.$$

Then, $h(\mathbf{1}) = 0$.

Consider now a general density operator $\rho \in \mathfrak{S}_+$. We want to estimate $h(\rho)$.

 $\begin{array}{l} h(\rho).\\ \text{Denote } \rho^{1/2} = \sum_{r,s=1}^n c_{rs} \mid e_r \rangle \langle \, e_s \, |.\\ \text{For } i,j \text{ fixed} \end{array}$

$$[V_{i,j} \rho^{1/2}, V_{i,j}^*] = [V_{i,j} \left(\sum_{r,s=1}^n c_{rs} | e_r \rangle \langle e_s | \right), V_{i,j}^*] =$$

$$V_{i,j} \left(\sum_{r,s=1}^n c_{rs} | e_r \rangle \langle e_s | \right) V_{i,j}^* - V_{i,j}^* V_{i,j} \left(\sum_{r,s=1}^n c_{rs} | e_r \rangle \langle e_s | \right) =$$

$$V_{i,j} \left(\sum_{r,s=1}^n c_{rs} | e_r \rangle \langle e_s | \right) V_{i,j}^* - | e_j \rangle \langle e_j | \left(\sum_{r,s=1}^n c_{rs} | e_r \rangle \langle e_s | \right) =$$

$$V_{i,j} \left(\sum_{r,s=1}^n c_{rs} | e_r \rangle \langle e_s | \right) | e_j \rangle \langle e_i | - \sum_{s=1}^n c_{js} | e_j \rangle \langle e_s | =$$

$$| e_i \rangle \langle e_j | \sum_{r=1}^n c_{rj} | e_r \rangle \langle e_i | - \sum_{s=1}^n c_{js} | e_j \rangle \langle e_s | =$$

$$c_{jj} | e_i \rangle \langle e_i | - \sum_{s=1}^n c_{js} | e_j \rangle \langle e_s |.$$

On the other hand, for i,j fixed

$$[V_{i,j}, \rho^{1/2} V_{i,j}^*] = \left[V_{i,j}, \left(\sum_{r,s=1}^n c_{rs} \,|\, e_r \,\rangle \langle \, e_s \,|\, \right) V_{i,j}^* \right] =$$

$$V_{i,j} \left(\sum_{r,s=1}^{n} c_{rs} |e_r\rangle \langle e_s| \right) V_{i,j}^* - \left(\sum_{r,s=1}^{n} c_{rs} |e_r\rangle \langle e_s| \right) V_{i,j}^* V_{ij} = c_{jj} |e_i\rangle \langle e_i| - \left(\sum_{r,s=1}^{n} c_{rs} |e_r\rangle \langle e_s| \right) |e_j\rangle \langle e_j| = c_{jj} |e_i\rangle \langle e_i| - \sum_{r=1}^{n} c_{rj} |e_r\rangle \langle e_j|.$$

Finally,

$$\mathcal{L}_{0}^{\dagger}(\rho^{1/2}) = \sum_{i,j=1}^{n} \left([V_{i,j} \rho^{1/2}, V_{i,j}^{*}] + [V_{i,j}, \rho^{1/2} V_{i,j}^{*}] \right) = \sum_{i,j=1}^{n} \left(c_{jj} | e_{i} \rangle \langle e_{i} | - \sum_{s=1}^{n} c_{js} | e_{j} \rangle \langle e_{s} | + c_{jj} | e_{i} \rangle \langle e_{i} | - \sum_{r=1}^{n} c_{rj} | e_{r} \rangle \langle e_{j} | \right) = 2\sum_{j=1}^{n} c_{jj} I - \sum_{i,j=1}^{n} \left(\sum_{s=1}^{n} c_{js} | e_{j} \rangle \langle e_{s} | + \sum_{r=1}^{n} c_{rj} | e_{r} \rangle \langle e_{j} | \right) = 2\sum_{j=1}^{n} c_{jj} I - n \sum_{j=1}^{n} \left(\sum_{s=1}^{n} c_{js} | e_{j} \rangle \langle e_{s} | + \sum_{r=1}^{n} c_{rj} | e_{r} \rangle \langle e_{j} | \right).$$

Then,

$$\rho^{1/2} \mathcal{L}_{0}^{\dagger}(\rho^{1/2}) =$$

$$\sum_{u,v=1}^{n} c_{uv} |e_{u}\rangle \langle e_{v}| 2 \sum_{j=1}^{n} c_{jj} I -$$

$$n \sum_{u,v=1}^{n} c_{uv} |e_{u}\rangle \langle e_{v}| \left(\sum_{j=1}^{n} \sum_{s=1}^{n} c_{js} |e_{j}\rangle \langle e_{s}| + \sum_{j=1}^{n} \sum_{r=1}^{n} c_{rj} |e_{r}\rangle \langle e_{j}|\right) =$$

$$\sum_{u,v=1}^{n} c_{uv} |e_{u}\rangle \langle e_{v}| 2 \sum_{j=1}^{n} c_{jj} I -$$

$$n \sum_{u,v=1}^{n} c_{uv} |e_{u}\rangle \left(\sum_{s=1}^{n} c_{vs} \langle e_{s}|\right) + n \sum_{u,v=1}^{n} c_{uv} |e_{u}\rangle \left(\sum_{j=1}^{n} c_{vj} \langle e_{j}|\right) =$$

$$\sum_{u,v=1}^{n} c_{uv} |e_{u}\rangle \langle e_{v}| 2 \sum_{j=1}^{n} c_{jj} I - 2n \sum_{u,v=1}^{n} c_{uv} |e_{u}\rangle \left(\sum_{s=1}^{n} c_{vs} \langle e_{s}|\right) = 2\sum_{j=1}^{n} c_{jj} \sum_{u,v=1}^{n} c_{uv} |e_{u}\rangle \langle e_{v}| - 2n \sum_{u,v=1}^{n} \sum_{s=1}^{n} c_{uv} c_{vs} |e_{u}\rangle \langle e_{s}|.$$

The diagonal of ρ is $\rho_{11}, \rho_{22}, ..., \rho_{nn}$ and $\sum_{j=1}^{n} c_{jj} = \sum_{j=1}^{n} \sqrt{\lambda_j}$, where λ_j , j = 1, ..., d, are the eigenvalues of ρ . Note that $\sup(\sum_{j=1}^{n} \sqrt{\lambda_j})^2 = n$ and $\inf(\sum_{j=1}^n \sqrt{\lambda_j})^2 = 1.$

Therefore, for fixed n and a given ρ

$$h(\rho) = \operatorname{Tr}\left[\rho^{1/2}\mathcal{L}_{0}^{\dagger}(\rho^{1/2})\right] = 2\sum_{j=1}^{n} c_{jj}\sum_{u=1}^{n} c_{uu} - 2n\sum_{v=1}^{n}\sum_{s=1}^{n} c_{sv}c_{vs} = 2\left(\sum_{j=1}^{n} c_{jj}\right)^{2} - 2n\sum_{s=1}^{n} \rho_{ss} = 2\left(\sum_{j=1}^{n}\sqrt{\lambda_{j}}\right)^{2} - 2n \le 0.$$

Note $h(\rho)$ can be very negative if n is large. We get the following proposition by looking at this last inequality:

Proposition 7. The entropy h depends only on $\{\lambda_i\}$ the eigenvalues of ρ and

$$h(\rho) = 2 \operatorname{Tr}(\rho^{1/2})^2 - 2n = 2 \left(\sum_{j=1}^n \sqrt{\lambda_i}\right)^2 - 2n$$

Note that as $\sum_{j=1}^{n} \lambda_j = 1$, the maximal value of $h(\rho)$ is zero, and this happens when all eigenvalues $\lambda_j = 1$ are equal to $\frac{1}{n}$. The maximal value of entropy is attained by the density matrix 1.

Remark 1. For fixed A we denote $\partial_{i,j}(A) = [V_{i,j}, A]$ and $\partial_{i,j}^{\dagger}(A) = [V_{i,j}^*, A]$. $\partial_{i,j}$ is a version of the momentum operator $\frac{1}{i} \frac{\partial}{\partial x}$ acting on the set L^2 of functions for the Lebesgue probability on the circle. Indeed, denote by D the operator $g \to D(g) = \frac{1}{i}g'$. For fixed $a: [0,1) \to \mathbb{R}$, take the multiplication operator $g \rightarrow a g$ acting on functions g. Then, the operator

$$g \to D(ag) - aD(g) = \frac{1}{i}a'g,$$

describes multiplication by $\frac{1}{i}a' = \frac{1}{i}\frac{\partial a}{\partial x}$.

We point out that $\sum_{i,j} \partial_{i,j} \partial_{i,j}^{\dagger}$ corresponds to second derivative (Lapla-cian). On the other hand $\sum_{i,j} \partial_{i,j} \partial_{i,j}$ corresponds to minus second derivative (minus Laplacian).

4 The general setting for detailed balanced condition

Before we begin the study of the quantum case we will state results for the detailed balanced condition when the continuous time Markov Chain takes values on $\{1, 2.., k\}$. Denote by $\mathfrak{s} = (\mathfrak{s}_1, ..., \mathfrak{s}_k)$ the initial invariant probability for the line sum zero matrix $W = (W_{i,j})_{i,j=1,...,k}$.

The detailed balance condition for W is: for all i, j = 1, ..., k

$$\mathfrak{s}_i W_{i,j} = \mathfrak{s}_j W_{j,i}.$$

Consider the inner product in \mathbb{R}^k

$$\langle x, y \rangle_{\mathfrak{s}} = \sum_{j=1}^{k} \mathfrak{s}_j x_j y_j.$$

It is easy to see that W satisfies the detailed balance condition, if and only if, W is self-adjoint for the inner product $\langle ., . \rangle_{\mathfrak{s}}$.

The above is the classical (commutative) setting for presenting the detailed balance condition. We are interested in presenting the non-commutative version of the concept.

We will be interested here in the C^* -Algebra $\mathcal{A} = M_n$ of complex n by n matrices. The inner product in M_n is $\langle A, B \rangle = \text{Tr} [A^* B]$. Following the notation of [5] the associated Hilbert space will be denoted by $\mathfrak{h}_{\mathcal{A}}$.

We will fix from now on an element $\sigma : \mathbb{C}^n \to \mathbb{C}^n$ in \mathfrak{S}_+ . A hypothesis that can be helpful for ergodic properties is to assume that all eigenvalues of σ are simple.

Now, we present some preliminary definitions and properties taken from [5].

Once we fix the Hamiltonian H we fix the density state σ via $\sigma = e^{-h}$ (in some sense we are considering a "normalized" Hamiltonian). In fact, $\sigma = \frac{e^{-H}}{Tr(e^{-H})}$ and $h = H + \log Tr(e^{-H})$.

The linear transformation, $\Delta_{\sigma} : M_n \to M_n$, is given by $A \to \Delta_{\sigma}(A) = \sigma A \sigma^{-1}$. Note that $\Delta_{\sigma}(\sigma) = \sigma$ and $\Delta_{\sigma}(\mathbf{1}) = \mathbf{1}$.

Assume each eigenvalue of σ is simple.

 $\Delta_{\sigma}^{-1}: M_n \to M_n$, is given by $B \to \tilde{\Delta}_{\sigma}(B) = \sigma^{-1} B \sigma$. Note that $(\Delta_{\sigma}(A))^* = \Delta_{\sigma}^{-1}(A^*)$.

 $\mathcal{K}: M_n \to M_n$ is positive preserving, if $\mathcal{K}(A) \ge 0$, in the case $A \ge 0$. Δ_{σ} is positive but not positive preserving. We say that $\mathcal{K}: M_n \to M_n$ is self-adjointness preserving, if $(\mathcal{K}(A))^* = \mathcal{K}(A^*)$.

Remember that by definition $\mathcal{K}^{\dagger}: M_n \to M_n$ is the one such that for all A, B

$$Tr[A^*\mathcal{K}(B)] = \langle A, \mathcal{K}(B) \rangle = \langle \mathcal{K}^{\dagger}(A), B \rangle = Tr[(\mathcal{K}^{\dagger}(A))^* B].$$

 $\mathcal{K}: M_n \to M_n$ is self-adjoint if $\mathcal{K} = \mathcal{K}^{\dagger}$.

 Δ_{σ} is self-adjoint.

Note that if $\Delta_{\sigma}(E) = e^{-w}E$, then $\Delta_{\sigma}(E^*) = e^{w}E^*$.

Assume that $\eta_j \in \mathbb{C}^n$, j = 1, 2, ..., n, is an orthonormal basis for $h = -\log \sigma$,

$$h(\eta_j) = \lambda_j \eta_j, \ \forall j.$$
(6)

Then, $\eta_j \in \mathbb{C}^n$, j = 1, 2, ..., n, is also an orthonormal basis for σ . $h = -\log \sigma$ plays the role of the Hamiltonian.

Then, we get that $\eta_j \in \mathbb{C}^n$, j = 1, 2, ..., n, is an orthonormal basis for σ ,

$$\sigma(\eta_j) = e^{-\lambda_j} \eta_j, \quad \forall j.$$
(7)

As $\sigma : \mathbb{C}^n \to \mathbb{C}^n$ is in \mathfrak{S}_+ , we get that

$$\sum_{j=1}^{n} e^{-\lambda_j} = 1.$$
 (8)

For $\alpha = (\alpha_1, \alpha_2)$, where $\alpha_1, \alpha_2 \in \{1, 2, ..., n\}$, denote

$$w_{\alpha_1,\alpha_2} = w_\alpha = \lambda_{\alpha_1} - \lambda_{\alpha_2}.$$
 (9)

If $\sigma = \mathbf{1}$, then, all $w_{\alpha_1,\alpha_2} = 0$. For each pair $\alpha \in \{1, 2, ..., n\}^2$, denote

$$F_{\alpha} = |\eta_{\alpha_1}\rangle \langle \eta_{\alpha_2}|.$$

Note that

$$F_{(\alpha_1,\alpha_2)}^* = F_{(\alpha_2,\alpha_1)}.$$
 (10)

Moreover,

$$\Delta_{\sigma}(F_{\alpha}) = e^{-\lambda_{\alpha_1} + \lambda_{\alpha_2}} F_{\alpha}.$$
 (11)

Indeed,

$$\Delta_{\sigma}(F_{\alpha}) = \sigma |\eta_{\alpha_{1}}\rangle \langle \eta_{\alpha_{2}} | \sigma^{-1} = e^{-\lambda_{\alpha_{1}}} |\eta_{\alpha_{1}}\rangle \langle \eta_{\alpha_{2}} | \sigma^{-1} = e^{-\lambda_{\alpha_{1}} + \lambda_{\alpha_{2}}} |\eta_{\alpha_{1}}\rangle \langle \eta_{\alpha_{2}} | = e^{-\lambda_{\alpha_{1}} + \lambda_{\alpha_{2}}} F_{\alpha}.$$

This shows that:

Lemma 8. The operators $F_{\alpha} = |\eta_{\alpha_1}\rangle\langle \eta_{\alpha_2}|, \alpha_1, \alpha_2 \in \{1, 2, ..., n\}$, describe a natural orthonormal basis of Δ_{σ} . The corresponding eigenvalues are $e^{-\lambda_{\alpha_1}+\lambda_{\alpha_2}}$.

Note that if τ represents the normalized trace and $\alpha = (\alpha_1, \alpha_2)$ is such that $\alpha_1 \neq \alpha_2$, then,

$$\tau(F_{\alpha}) = 0$$

and, for $\alpha, \tilde{\alpha} \in \{1, 2, ..., n\}^2$

$$\tau(F_{\alpha}^* F_{\tilde{\alpha}}) = \delta_{\alpha, \tilde{\alpha}} := \delta_{\alpha_1, \tilde{\alpha}_1} \delta_{\alpha_2, \tilde{\alpha}_2}.$$

Now we denote the different F_{α} , $\alpha \in \{1, 2, ..., n\}^2$, by V_k , $k = 1, ..., n^2$ (in order to use the same notation as in [5]). In this identification, we also denote for each $k = 1, ..., n^2$, the value $w_k = \lambda_{\alpha_1} - \lambda_{\alpha_2}$, for the corresponding $\alpha = (\lambda_{\alpha_2}, \lambda_{\alpha_1})$.

Then, the family $V_1, ..., V_{n^2}$ represent the different eigenmatrices (an orthonormal basis) for Δ_{σ} associated to the eigenvalues $e^{-w_1}, ..., e^{-w_{n^2}}$, where $w_k \in \mathbb{R}, k = 1, ..., n^2$. 1 and σ are eigenmatrices associated to the eigenvalue 1. The matrices V_k do not have to be self-adjoint, but from (10) we get

$$\{V_1, ..., V_{n^2}\} = \{V_1^*, ..., V_{n^2}^*\}.$$

Therefore, if w_k is in the above list, there exists a j such that $w_j = -w_k$. Given the Hamiltonian $h = -\log \sigma$, the modular automorphism $\alpha_t : M_n \to M_n, t \ge 0$, is defined by

 $A \to \alpha_t(A) = e^{ith} A e^{-ith} \Leftarrow$ The Heisenberg point of view.

A Quantum Markov Semigroup (QMS) is a continuous one-parameter semigroup of linear transformations $\mathcal{P}_t : M_n \to M_n, t \ge 0$, such that for each $t \ge 0, \mathcal{P}_t$ is completely positive and $\mathcal{P}_t(\mathbf{1}) = \mathbf{1}$.

It is natural to focus on quantum Markov semigroups that commute with the modular operator Δ_{σ} associated to their invariant states σ .

Consider a QMS $\mathcal{P}_t: M_n \to M_n, t \ge 0$, of the form

$$\mathcal{P}_t = e^{t\mathcal{L}},$$

for some linear operator $\mathcal{L}: M_n \to M_n$.

The operator \mathcal{L} acts on observables (self-adjoint matrices). Note that $\mathcal{L}(\mathbf{1}) = 0$. The dual operator \mathcal{L}^{\dagger} acts on density matrices.

A state σ is invariant if Tr $[\sigma \mathcal{L}(A)] = 0$, for all $A \in M_n$.

In terms of the possible inner products described on Definition 2.2 in [5] we will choose s = 1.

Remember that given σ we consider the inner product $<, >_{\sigma} = <, >_{1}$ in M_{n} , where

$$\langle A, B \rangle_{\sigma} = \operatorname{Tr} [\sigma A^* B].$$

From [5] we get:

Proposition 9. Given the density operator σ , the the QMS $\mathcal{P}_t : M_n \to M_n$, $t \geq 0$, of the form $\mathcal{P}_t = e^{t\mathcal{L}}$, satisfies the σ - detailed balance condition, if and only if,

$$\mathcal{L} \circ \Delta_{\sigma} = \Delta_{\sigma} \circ \mathcal{L}. \tag{12}$$

If $\mathcal{P}_t = e^{t\mathcal{L}}$, satisfies the σ -detailed balanced condition, then, for all $t \geq 0$,

$$\mathcal{P}_t \circ \Delta_\sigma = \Delta_\sigma \circ \mathcal{P}_t.$$

Moreover, for any t, t' and matrix A we get that

$$(\alpha_{t'} \circ \mathcal{P}_t)(A) = (\mathcal{P}_t \circ \alpha_{t'})(A).$$

It follows from (12) that $V_1, ..., V_{n^2}$ is an orthonormal basis for \mathcal{L} associated to the eigenvalues $e^{-v_1}, ..., e^{-v_{n^2}}$, where $v_j \in \mathbb{R}, j = 1, ..., n^2$.

Lemma 10. σ is a stationary density matrix for the semigroup with infinitesimal generator \mathcal{L} .

Proof. Note that

$$\begin{split} [V_{i,j}\,\sigma, V_{i,j}^*] &= |\,\eta_i\,\rangle\langle\,\eta_j\,|\sigma|\,\eta_j\,\rangle\langle\,\eta_i\,| - |\,\eta_j\,\rangle\langle\,\eta_i\,||\,\eta_i\,\rangle\langle\,\eta_j\,|\sigma\\ &= e^{-\lambda_j}|\,\eta_i\,\rangle\langle\,\eta_i\,| - e^{\lambda_j}|\,\eta_j\,\rangle\langle\,\eta_j\,|\\ &= e^{-\lambda_j}\,(|\,\eta_i\,\rangle\langle\,\eta_i\,| - |\,\eta_j\,\rangle\langle\,\eta_j\,|) \end{split}$$

and

$$[V_{i,j}, \sigma V_{i,j}^*] = |\eta_i\rangle \langle \eta_j |\sigma| \eta_j\rangle \langle \eta_i | - \sigma| \eta_j\rangle \langle \eta_i || \eta_i\rangle \langle \eta_j |$$

= $e^{-\lambda_j} |\eta_i\rangle \langle \eta_i | - e^{\lambda_j} |\eta_j\rangle \langle \eta_j |$
= $e^{-\lambda_j} (|\eta_i\rangle \langle \eta_i | - |\eta_j\rangle \langle \eta_j |).$

Using expression (15) we get

$$\mathcal{L}^{\dagger}(\sigma) = 2 \sum_{i,j} e^{(\lambda_j - \lambda_i)/2} e^{-\lambda_j} \left(|\eta_i\rangle \langle \eta_i| - |\eta_j\rangle \langle \eta_j| \right)$$

= $2 \left[\sum_j e^{-\lambda_j/2} \sum_i e^{-\lambda_i/2} |\eta_i\rangle \langle \eta_i| - \sum_i e^{-\lambda_i/2} \sum_j e^{-\lambda_j/2} |\eta_j\rangle \langle \eta_j| \right]$
= 0.

Remember that for each pair $i, j \in \{1, 2, ..., n\}$, we denote

$$V_{i,j} = |\eta_i\rangle \langle \eta_j|,$$

where η_i , $i \in \{1, 2, ..., n\}$, is the orthonormal basis of eigenvectors for $h = -\log \sigma$ associated to the eigenvalues λ_j (according to (6)). As we mention before $w_{i,j} = \lambda_i - \lambda_j$, $i, j \in \{1, 2, ..., n\}$.

Theorem 11. If $\mathcal{P}_t = e^{t\mathcal{L}}$, satisfies the σ -detailed balanced condition, then \mathcal{L} is of the form

$$A \to \mathcal{L}(A) = \sum_{i,j=1}^{n} e^{-w_{i,j}/2} \left(V_{i,j}^* \left[A, V_{i,j} \right] + \left[V_{i,j}^*, A \right] V_{i,j} \right), \tag{13}$$

where $V_{i,j} = |\eta_i\rangle\langle \eta_j|$ and $w_{i,j} = \lambda_i - \lambda_j$, $i, j \in \{1, 2, ..., n\}$, are real numbers such that (7), (8) and (9) are true (which also means $\Delta_{\sigma}(V_{i,j}) = e^{-\lambda_i + \lambda_j} V_{i,j}$).

Note that given $\sigma \in \mathfrak{G}_+$, the eigenvetors $|\eta_i\rangle$ and eigenvalues λ_j , $j \in \{1, 2, ..., n\}$, are determined. Therefore, if $\mathcal{P}_t = e^{t\mathcal{L}}$ satisfies the σ -detailed balanced condition, then, \mathcal{L} is uniquely determined.

Remark 2. Conversely, given σ in \mathfrak{G}_+ and V_j , $j = 1, 2, ..., n^2$, such that, 1. $\Delta_{\sigma} V_j = e^{-w_j} V_j$, 2. $\{V_j, j = 1, 2, ..., n^2\} = \{V_j^*, j = 1, 2, ..., n^2\}$, then, $A \to \mathcal{L}(A) = \sum_{i=1}^{n^2} e^{-w_j/2} (V_j^* [A, V_j] + [V_j^*, A] V_j),$ (14)

is the infinitesimal generator of a QMS $e^{t\mathcal{L}}$, $t \geq 0$, which satisfies de d.b.c. for the given σ . Therefore, σ is stationary for $e^{t\mathcal{L}^{\dagger}}$, $t \geq 0$.

The dual operator \mathcal{L}^{\dagger} satisfies

$$\rho \to \mathcal{L}^{\dagger}(\rho) = \sum_{i,j=1}^{n} e^{-w_{i,j}/2} \left([V_{i,j} \, \rho, V_{i,j}^*] + [V_{i,j}, \rho \, V_{i,j}^*] \right)$$
(15)

Remark 3. If $\mathcal{P}_t = e^{t\mathcal{L}}$, satisfies the detailed balanced condition for the $\sigma = \mathbf{1}$, then from (12) we get that $V_{i,j} = \mathfrak{I}_{i,j}$, i, j = 1, ..., n. This is the case when $\mathcal{L} = \mathcal{L}_0$.

5 The Pressure problem

Definition 12. Given an Hermitian operator $A : \mathbb{C}^n \to \mathbb{C}^n$, $D_n = \{\rho \ge 0 : Tr(\rho) = 1\}$, consider

$$P_A(\rho) = h(\rho) + Tr(A\rho) \tag{16}$$

and

$$P_A = \sup_{\rho \in D_n} P_A(\rho). \tag{17}$$

A matrix ρ_A maximizing P_A is called an equilibrium density operator for A.

Question: Is there \mathcal{L} such that ρ_A is stationary for \mathcal{L} ? The converse in Theorem 3.1 in [5] may be useful.

The expression of entropy we found in Proposition 7 suggests us to look at the matrices of the form $\xi = \rho^{1/2}$, where ρ is a density. In order to study the problem of who maximizes P_A , we then define

$$\Xi_n = \{\xi \ge 0 : \xi^2 \in D_n\},\tag{18}$$

the set of square roots of density operators and

$$p_A(\xi) = h_{1/2}(\xi) + Tr(A\xi^2),$$
 (19)

where $h_{1/2}(\xi) := 2\text{Tr}(\xi)^2 - 2n = h(\xi^2)$ by Proposition 7. Notice that $p_A(\xi) = P_A(\xi^2)$.

Proposition 13. If ξ maximizes the functional p_A , then there exists a κ satisfying

$$2\kappa\xi = 4\operatorname{Tr}(\xi)I + A\xi + \xi A. \tag{20}$$

Proof. We will use Lagrange Multipliers. Let $g: M^n \to \mathbb{C}, g(\zeta) = \text{Tr}(\zeta^2) - 1$. The maximal ξ then satisfies, for all $h \in M_n(\mathbb{C})$ and some $\kappa \in \mathbb{R}$,

$$\begin{cases} \mathrm{Dp}_A(\xi)(h) = \kappa \mathrm{D}g(\xi)(h) \\ g(\xi^2) = 0, \end{cases}$$
(21)

with $Dp_A(\xi)(h) = 4Tr(\xi)Tr(h) + Tr(A\{\xi, h\})$ and $Dg(\xi)(h) = 2Tr(h\xi)$.

Taking $h = |e_j\rangle\langle e_i|$ we have

$$2\kappa\xi_{ij} = 4\mathrm{Tr}(\xi)\delta_{i,j} + \{A,\xi\}_{ij}.$$

Since the above is true for every i, j, it corresponds to the coordinate equations of the matrix equation

$$2\kappa\xi = 4\mathrm{Tr}(\xi)I + A\xi + \xi A. \tag{22}$$

Note that if $\xi = \rho_A^{1/2}$, then it follows from (20)

$$\kappa \rho_A = 2 \operatorname{Tr}(\rho_A^{1/2}) \rho_A^{1/2} + \frac{1}{2} \left(A \rho_A + \rho_A^{1/2} A \rho_A^{1/2} \right).$$
(23)

Indeed,

$$2\kappa\rho_A = 2\kappa\xi\,\xi = 4\mathrm{Tr}(\xi)\,\xi + A\xi\xi + \xi A\xi = 4\mathrm{Tr}(\xi)\,\xi + \rho_A + \xi A\xi.$$
(24)

Proposition 14. If ξ maximizes p_A , then the following statements are true

1. $Tr(A\xi) = (\kappa - 2n) Tr(\xi);$ 2. $P_A(\rho_A) = P_A(\xi^2) = p_A(\xi) = \kappa - 2n;$

Proof. In (21) take h = I and $h = \xi$, respectively.

Remark 4. The problem of finding the maximizing density $\rho = \xi^2$ for a general Hermitian A can be reduced to the study of the diagonal case. In fact, since A is hermitian, it is diagonalizable. We have $UAU^* = \Lambda$, the later a diagonal matrix, for some unitary matrix U. Multiplying on the left by U and on the right by U^{*} in the above matrix equation gives us

$$2\kappa U\xi U^* = 4 \operatorname{Tr}(\xi) UU^* + UAU^* U\xi U^* + U\xi U^* UAU^*$$
$$\iff 2\kappa \eta = 4 \operatorname{Tr}(\eta) + \Lambda \eta + \eta \Lambda,$$

where $\eta := U\xi U^*$. We arrive at a particular version of (22) on which the matrix is diagonal.

Theorem 15. If $A = Diag(a_1, \ldots, a_n)$, the ξ that maximizes p_A is also diagonal, with

$$\xi_{ii} = \frac{c}{\kappa - a_i},$$

where κ is given implicitly on the data $a_1, ..., a_n$ and c is such that $Tr(\xi^2) = 1$. Consequently, the density that maximizes the pressure P_A is

$$\rho_A = \frac{1}{Tr(\rho_A)} \begin{pmatrix} \frac{1}{(\kappa - a_1)^2} & & \\ & \frac{1}{(\kappa - a_2)^2} & \\ & & \ddots & \\ & & & \frac{1}{(\kappa - a_n)^2} \end{pmatrix}$$

Proof. For A diagonal, the expression (22) becomes

$$2\kappa\xi_{ij} = 4\mathrm{Tr}(\xi)\delta_{i,j} + \xi_{ij}(a_i + a_j).$$
⁽²⁵⁾

If we take i = j in the expression above,

$$\kappa \xi_{ii} = 2 \operatorname{Tr}(\xi) + \xi_{ii} a_i \iff (\kappa - a_i) \xi_{ii} = 2 \operatorname{Tr}(\xi)$$
(26)

We know that $\operatorname{Tr}(\xi) > 0$, because $\operatorname{Tr}(\xi) = 0$ leads to $\xi = 0$. Then $\xi_{ii} \neq 0$ and $\kappa > a_i$ for all *i*. So,

$$\frac{1}{\kappa - a_i} = \frac{\xi_{ii}}{2\text{Tr}(\xi)} \tag{27}$$

and

$$\sum_{i=1}^{n} \frac{1}{\kappa - a_i} = \frac{1}{2}.$$
(28)

We need to find κ to completely characterize the maximal ξ . Suppose that $a_1 = \max_i a_i$. Notice that $f(x) = \sum_{i=1}^n \frac{1}{x-a_i}$, for $x \neq a_i$ has a vertical asymptote at a_1 , $\lim_{x\to a_1^+} f(x) = \infty$, and it decreases to $\lim_{x\to\infty} f(x) = 0$. By the intermediate value theorem, we have a $\kappa > a_1$ s.t. $f(\kappa) = 1/2$.

Alternatively, one can find it as one of the roots of the following polynomial, which is the expression (27) rewritten.

$$\frac{1}{2}\det(\kappa \mathbf{I} - A) - \sum_{i=1}^{n} \det\left(\kappa \mathbf{I} - A + (a_i - \kappa + 1)|e_i\rangle\langle e_i|\right) = 0.$$
(29)

There is just one root that is bigger than all a_i , therefore it is κ .

Finally we prove the elements out of the diagonal are null. For $i \neq j$, the expression (28) gives us $2\kappa\xi_{ij} = \xi_{ij}(a_i + a_j)$, or equivalently,

$$\xi_{ij}(2\kappa - (a_i + a_j)) = 0.$$

We know that $\kappa > a_i, \forall i$. Thus $2\kappa > a_i + a_j$. This leaves us with $\xi_{ij} = 0$.

To conclude the pressure problem, we write

$$\xi = \text{Diag}(\xi_{11}, ..., \xi_{nn}), \quad \xi_{ii} = \frac{c}{\kappa - a_i},$$
(30)

where c is the constant that makes $\text{Tr}(\xi^2) = 1$, i.e.,

$$c = \left(\sum_{i=1}^{n} \frac{1}{(\kappa - a_i)^2}\right)^{-1/2}$$

This way, given $a_1, ..., a_n$, we find κ , then c and finally ξ .

Corollary 16. If A is diagonalizable, with UAU^{*} diagonal, then the maximizing density ρ_A for P_A is such that $U\rho_A U^*$ is diagonal.

Proof. If A was not diagonal at first, we proceed as in Remark 4 and use the last theorem to find a maximal $\eta = U\xi U^*$ which is diagonal. Then $\eta^2 = U\xi^2 U^* = U\rho_A U^*$ is diagonal.

Remark 5. Using (27), we know that if $A = a_1 I$ then $\xi_{ii} = \xi_{11}$ for all *i*. Since A is diagonal and $Tr(\xi^2) = \sum_i \xi_{ii}^2 = n\xi_{11}^2 = 1$, it follows that $\xi = \frac{1}{\sqrt{n}}I$ is the only ξ that maximizes p_A .

Example 1. Let

$$A = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 2 \end{pmatrix}.$$

What is the equilibrium density ρ_A , i.e., the density that maximizes P_A ? Let's diagonalize A.

$$U = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 & 0\\ 1 & 1 & 0\\ 0 & 0 & \sqrt{2} \end{pmatrix}, \quad UAU^* = \begin{pmatrix} -1 & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & 2 \end{pmatrix}.$$

Now we apply Theorem 15. κ satisfies

$$\frac{1}{\kappa+1} + \frac{1}{\kappa-1} + \frac{1}{k-2} = \frac{1}{2},$$

and $\kappa > -1, 1, 2$. We get $\kappa \approx 6.902$. Thus, the maximizing density for UAU^* is

$$\rho_{UAU^*} = \frac{1}{\text{Tr}(\rho_{UAU^*})} \begin{pmatrix} \frac{1}{(6.902+1)^2} & & \\ & \frac{1}{(6.902-1)^2} & \\ & & \frac{1}{(6.902-2)^2} \end{pmatrix}$$
$$= \begin{pmatrix} 0.186 & & \\ & 0.332 & \\ & & 0.482 \end{pmatrix}.$$

Finally,

$$\rho_A = U^* \rho_{UAU^*} U$$

$$= \begin{pmatrix} 0.259 & 0.073 & 0\\ 0.073 & 0.259 & 0\\ 0 & 0 & 0.482 \end{pmatrix}.$$

Notice that

$$P(\rho_A) = h(\rho_A) + \operatorname{Tr}(A\rho_A)$$

= 2 Tr $(\rho_A^{1/2})^2 - 2n + \operatorname{Tr}(UAU^*U\rho_A U^*)$
= 2 Tr $((U\rho_A U^*)^{1/2})^2 - 2n + \operatorname{Tr}(UAU^*U\rho_A U^*)$
= 2 $(\sqrt{0.186} + \sqrt{0.332} + \sqrt{0.482})^2 - 6 + (-1 \cdot 0.186 + 1 \cdot 0.332 + 2 \cdot 0.482)$
 $\approx 0.902 \approx \kappa - 6.$

in accordance with Proposition 14.

6 The pressure P_A as an eigenvalue problem

Consider the linear operator \mathfrak{L}_A

$$\xi \to \mathfrak{L}_A(\xi) = 2\mathrm{Tr}(\xi)I + \frac{1}{2}(A\xi + \xi A).$$
(31)

Suppose ρ_A is an equilibrium density operator for the selfadjoint matrix A.

From (20) we get that $\xi = \rho_A^{1/2}$ is an eigenmatrix for the linear operator \mathfrak{L}_A associated to the eigenvalue κ , that is

$$\mathfrak{L}_A(\rho_A^{1/2}) = \kappa \,\rho_A^{1/2} \tag{32}$$

From item 2. in Theorem 14 we get that $P_A(\rho_A) = \kappa - 2n$.

In this way, the equilibrium density operator is related to an eigenvalue problem in a similar fashion as in classical Thermodynamic Formalism.

The equilibrium matrix ρ_A satisfies

$$\kappa \rho_A = 2 \operatorname{Tr}(\rho_A^{1/2}) \, \rho_A^{1/2} + \frac{1}{2} \, (A \, \rho_A + \rho_A^{1/2} A \rho_A^{1/2}), \tag{33}$$

but this is not exactly a linear relation.

7 A connection between $h(\rho)$ and $I(\nu)$

We will present a connection between the concepts of $h(\rho)$ and $-I(\rho)$.

Recall that $\mathbf{1} = Id/n$ satisfies the detailed balance condition, which is the quantum equivalent of reversibility.

We are going to establish a connection of the Laplacian-entropy of Definition 5 with the one in (5.18) of [17]. The notion of Radon-Nikodym derivative is not clear in the quantum setting, but we can consider a natural analogy in our reasoning and we write $\frac{d\nu}{d\mu} = A$, if

$$\operatorname{Tr}(\nu U) = \operatorname{Tr}(\mu A U).$$

This corresponds of writing A in the form $A = \mu^{-1}\nu$. When looking at (5.18), L is simmetric in $L^2(\mu)$, and our operator \mathcal{L}_0 satisfies d.b.c. for **1**. Therefore, here we will address the computation of the corresponding expression $\frac{d\nu}{d\mathbf{1}}$. Then, it is natural to consider $A = \mathbf{1}^{-1}\nu = n\nu$. Therefore, (5.18) in our setting corresponds to

$$I(\nu) = -\int (n\nu)^{1/2} \mathcal{L}_0((n\nu)^{1/2}) \, d\mathbf{1}$$

= $-n \operatorname{Tr}(\mathbf{1} \nu^{1/2} \mathcal{L}_0(\nu^{1/2}))$
= $-\operatorname{Tr}(\nu^{1/2} \mathcal{L}_0(\nu^{1/2})),$

and then, $-h(\nu)$ come up.

Remark 6. Recall that for an $A = \sum_{kl} a_{kl} |k\rangle \langle l|$,

$$\mathcal{L}_{0}(A) = \sum_{i,j=1}^{n} (V_{ij}^{*}[A, V_{ij}] + [V_{ij}, A]V_{ij}^{*}), \text{ for } V_{ij} = |i\rangle\langle j|$$
$$= \sum_{i,j=1}^{n} V_{ji}AV_{ij} - V_{ji}V_{ij}A + V_{ij}AV_{ji} - AV_{ij}V_{ji}$$

$$= 2\sum_{i,j=1}^{n} V_{ji}AV_{ij} - \sum_{i,j=1}^{n} |j\rangle\langle i||i\rangle\langle j|A - \sum_{i,j=1}^{n} A|i\rangle\langle j||j\rangle\langle i|$$

$$= 2\sum_{i,j=1}^{n} V_{ji}AV_{ij} - n\sum_{j=1}^{n} |j\rangle\langle j|A - n\sum_{i=1}^{n} A|i\rangle\langle i|$$

$$= 2\sum_{i,j=1}^{n} V_{ji}AV_{ij} - 2nA$$

$$= 2\sum_{i,j=1}^{n} |j\rangle\langle i|A|i\rangle\langle j| - 2nA$$

$$= 2\sum_{i,j=1}^{n} a_{kl}|j\rangle\langle i||k\rangle\langle l||i\rangle\langle j| - 2nA$$

$$= 2\sum_{i,j=1}^{n} a_{ii}|j\rangle\langle j| - 2nA$$

$$= 2\sum_{i,j=1}^{n} a_{ii}\sum_{j=1}^{n} |j\rangle\langle j| - 2nA$$

$$= 2\sum_{i,j=1}^{n} a_{ii}\sum_{j=1}^{n} |j\rangle\langle j| - 2nA$$

$$= 2Tr(A)I - 2nA.$$

The next step is to write the entropy as an infimum. We will show that: **Theorem 17.** Given the density matrix ρ

$$h(\rho) = \inf_{A>0} Tr(\rho A^{-1} \mathcal{L}_0(A))$$

For the proof, we will need the following result:

Lemma 18. In the space M_n of matrices $n \times n$, is true that

$$\inf_{B>0} Tr(BU) Tr(UB^{-1}) = Tr(U)^2$$

Proof. Let B > 0 be a general positive matrix. Let $|i\rangle$ be the orthonormal basis of eigenvectors of B. Then, we can write $B = \sum_{i=1}^{n} b_i |i\rangle\langle i|$, and in this basis, U can be written as $U = \sum_{j,k=1}^{n} u_{jk} |j\rangle\langle k|$. Thus,

$$BU = \sum_{ijk} b_i u_{jk} |i\rangle \langle i||j\rangle \langle k| = \sum_{ik} b_i u_{ik} |i\rangle \langle k|$$

$$\implies \operatorname{Tr}(BU) = \sum_{i=1}^{n} b_{i} u_{ii}.$$
$$UB^{-1} = \sum_{ijk} \frac{u_{jk}}{b_{i}} |j\rangle \langle k||i\rangle \langle i| = \sum_{ij} \frac{u_{ji}}{b_{i}} |j\rangle \langle i|$$
$$\implies \operatorname{Tr}(UB^{-1}) = \sum_{i=1}^{n} \frac{u_{ii}}{b_{i}}.$$

Therefore,

$$\operatorname{Tr}(UB)\operatorname{Tr}(BU^{-1}) = \sum_{i,j=1}^{n} \frac{b_j}{b_i} u_{ii} u_{jj} = \frac{1}{2} \sum_{i,j=1}^{n} \left(\frac{b_j}{b_i} + \frac{b_i}{b_j}\right) u_{ii} u_{jj}$$
$$\geq \sum_{i,j=1}^{n} u_{ii} u_{jj} = \left(\sum_{i=1}^{n} u_{ii}\right)^2 = \operatorname{Tr}(U)^2.$$

Notice that we used the fact that $x + 1/x \ge 2$, for all x > 0. By now, we have the lower bound $Tr(U)^2$. To finish, notice that B = Id achieves this bound, so we conclude that

$$\inf_{B>0} \operatorname{Tr}(BU) \operatorname{Tr}(UB^{-1}) = \operatorname{Tr}(U)^2.$$

Now we proceed to prove the theorem.

Proof. Using Remark 6, we have

$$A^{-1}\mathcal{L}_0(A) = 2A^{-1}\mathrm{Tr}(A) - 2nI$$

$$\Rightarrow \mathrm{Tr}(\rho A^{-1}\mathcal{L}_0(A)) = 2\mathrm{Tr}(\rho A^{-1}).\mathrm{Tr}(A) - 2n\mathrm{Tr}(\rho)$$

$$= 2\mathrm{Tr}(\rho A^{-1})\mathrm{Tr}(A) - 2n.$$

Writing A in the form $A = B\rho^{1/2}$, it will not change the infimum, which will be now taken over B > 0. This means

$$\inf_{A>0} 2\operatorname{Tr}(\rho A^{-1})\operatorname{Tr}(A) - 2n = 2 \inf_{B>0} \operatorname{Tr}(\rho^{1/2}B^{-1})\operatorname{Tr}(B\rho^{1/2}) - 2n$$
$$= 2\operatorname{Tr}(\rho^{1/2})^2 - 2n = h(\rho).$$

As the infimum was computed by Lemma 18 we proved the claim.

8 From quantum to classical

Definition 19. Given σ and an infinitesimal generator \mathcal{L} of the form (13), we say that the matrix Q is the matrix associated to \mathcal{L} , if Q is $n \times n$ real matrix with entries $Q_{i,j} = Tr[F_{i,i}\mathcal{L}F_{j,j}]$, where $F_{i,i} = |\eta_i\rangle\langle\eta_i|$.

This matrix is line sum zero with positive values outside the diagonal (see [5]). The matrix Q^{\dagger} , the transpose of Q, has a stationay eigenvector probability $\overrightarrow{\sigma} \in (0, 1)^n$ associated to the eigenvalue 0.

Lemma 20. Given l, k, the entry $Q_{l,k} = Tr[F_{l,l} \mathcal{L}F_{k,k}]$ is given by

$$Q_{l,k} = 2 e^{-w_{k,l}/2} - 2\delta_{l,k} \sum_{i=1}^{n} e^{-w_{i,l}/2}.$$
(34)

Proof. Indeed, when, $A = V_{k,k} = |\eta_k\rangle \langle \eta_k|$ we get

$$V_{i,j}^*[A, V_{i,j}] = |\eta_j\rangle \langle \eta_i | [A, |\eta_i\rangle \langle \eta_j |] =$$
$$|\eta_j\rangle \langle \eta_i | (|\eta_k\rangle \langle \eta_k | |\eta_i\rangle \langle \eta_j | - |\eta_i\rangle \langle \eta_j | |\eta_k\rangle \langle \eta_k |) =$$
$$\delta_{i,k} |\eta_j\rangle \langle \eta_j | - \delta_{j,k} |\eta_j\rangle \langle \eta_k |$$

Moreover, when $A = V_{k,k} = |\eta_k\rangle \langle \eta_k |$

$$[V_{i,j}^*, A] V_{i,j} = \left[|\eta_j\rangle \langle \eta_i|, A \right] |\eta_i\rangle \langle \eta_j| = \\ \left(|\eta_j\rangle \langle \eta_i| |\eta_k\rangle \langle \eta_k| - |\eta_k\rangle \langle \eta_k| |\eta_j\rangle \langle \eta_i| \right) |\eta_i\rangle \langle \eta_j| = \\ \delta_{i,k} |\eta_j\rangle \langle \eta_j| - \delta_{j,k} |\eta_k\rangle \langle \eta_j|.$$

Then, when $A = V_{k,k} = |\eta_k\rangle \langle \eta_k |$

$$e^{-w_{i,j}/2} \left(V_{i,j}^*[A, V_{i,j}] + [V_{i,j}^*, A] V_{i,j} \right) =$$

$$e^{-w_{i,j}/2} \left(2\delta_{i,k} | \eta_j \rangle \langle \eta_j | -\delta_{j,k} | \eta_j \rangle \langle \eta_k | -\delta_{j,k} | \eta_k \rangle \langle \eta_j | \right)$$

$$= e^{-w_{i,j}/2} \left(2\delta_{i,k} | \eta_j \rangle \langle \eta_j | -2\delta_{j,k} | \eta_k \rangle \langle \eta_k | \right),$$

and finally

$$\mathcal{L}(A) = \sum_{i,j=1}^{n} e^{-w_{i,j}/2} \left(V_{i,j}^{*}[A, V_{i,j}] + V_{i,j}^{*}, A] V_{i,j} \right) = \sum_{i=1}^{n} \sum_{j=1}^{n} e^{-w_{i,j}/2} \left(2\delta_{i,k} | \eta_{j} \rangle \langle \eta_{j} | - 2\delta_{j,k} | \eta_{k} \rangle \langle \eta_{k} | \right) =$$

$$2\sum_{j=1}^{n} e^{-w_{k,j}/2} |\eta_{j}\rangle \langle \eta_{j}| - 2\sum_{i=1}^{n} e^{-w_{i,k}/2} |\eta_{k}\rangle \langle \eta_{k}|$$

Therefore, when $A = V_{k,k} = |\eta_k\rangle \langle \eta_k|$, given l

$$|\eta_l\rangle\langle\eta_l|\mathcal{L}(A) = 2|\eta_l\rangle\langle\eta_l|\sum_{j=1}^n e^{-w_{k,j}/2}|\eta_j\rangle\langle\eta_j|-2|\eta_l\rangle\langle\eta_l|\sum_{i=1}^n e^{-w_{i,k}/2}|\eta_k\rangle\langle\eta_k|$$

$$= 2 \sum_{j=1}^{n} e^{-w_{k,j}/2} |\eta_l\rangle \langle \eta_l || \eta_j\rangle \langle \eta_j |-2 \sum_{i=1}^{n} e^{-w_{i,k}/2} |\eta_l\rangle \langle \eta_l || \eta_k\rangle \langle \eta_k |$$
$$= 2 e^{-w_{k,l}/2} |\eta_l\rangle \langle \eta_l |-2\delta_{l,k} \sum_{i=1}^{n} e^{-w_{i,l}/2} |\eta_l\rangle \langle \eta_l |.$$

From this,

=

$$Q_{lk} = 2 e^{-w_{k,l}/2} - 2\delta_{l,k} \sum_{i=1}^{n} e^{-w_{i,l}/2}$$
(35)
$$\begin{cases} 2 e^{-w_{k,l}/2} = 2 e^{(\lambda_l - \lambda_k)/2} & \text{if } l \neq k \\ 2 e^{-w_{l,l}/2} - 2 \sum_{i=1}^{n} e^{-w_{i,l}/2} = 2 - 2 \sum_{i=1}^{n} e^{(\lambda_l - \lambda_i)/2} & \text{if } l = k \end{cases}$$

Notice that:

$$\sum_{k=1}^{n} Q_{lk} = Q_{ll} + \sum_{k:k \neq l} Q_{lk}$$
$$= 2 e^{-w_{l,l}/2} - 2 \sum_{i=1}^{n} e^{-w_{i,l}/2} + 2 \sum_{k:k \neq l} e^{-w_{k,l}/2}$$
$$= -2 \sum_{i=1}^{n} e^{-w_{i,l}/2} + 2 \sum_{k=1}^{n} e^{-w_{k,l}/2} = 0.$$

Note that the expression (34) for the matrix Q depends on the eigenvalues $e^{-\lambda_i}$, $i \in \{1, 2.., n\}$, and not the specific eigenfunctions η_i , $i \in \{1, 2.., n\}$, of σ . This means that many density matrices σ can determine the same matrix Q.

Theorem 4.2 in [5] claims:

Theorem 21. Assume that \mathcal{L} is of the form (13) for σ . The matrix Q, given by $Q_{i,j} = Tr [F_{i,i}\mathcal{L}F_{j,j}]$ is line sum zero. The invariant probability for the classical continuous time Markov chain with infinitesimal generator Q is

$$\overrightarrow{\sigma} = (\sigma_1, \sigma_2, \dots, \sigma_n) = (Tr[\sigma F_{1,1}], Tr[\sigma F_{2,2}], \dots Tr[\sigma F_{n,n}]).$$
(36)

The classical detailed balance condition

$$\sigma_i Q_{i,k} = \sigma_k Q_{k,i} \tag{37}$$

is satisfied.

Consider the Chapman-Kolmogorov linear differential equation on $\overrightarrow{\rho}(t) = (\rho_1(t), \rho_2(t), ..., \rho_n(t)) \in \mathbb{R}^n$,

$$\frac{d}{dt}\rho_l(t) = \sum_{k=1}^n (\rho_k(t)Q_{k,l} - \rho_k(t)Q_{l,k}).$$
(38)

This is equivalent to

$$\overrightarrow{\rho}(t) = e^{tQ^{\dagger}}(\overrightarrow{\rho}(0)). \tag{39}$$

The occupation time probability in $\{1, 2.., n\}$ of the continuous time Markov Chain is described by $\overrightarrow{\rho}(t)$.

 $\overrightarrow{\rho}(t)$ satisfies (38), if and only if, the quantum continuous time evolution $\rho(t)$ in \mathcal{A} satisfies

$$\rho(t) = \sum_{k=1}^{n} \frac{\rho_k(t)}{Tr(F_{k,k})} F_{k,k}.$$
(40)

Remember that from (7)

$$\sigma = \sum_{k=1}^{n} e^{-\lambda_k} |\eta_k\rangle \langle \eta_k|.$$
(41)

Then, from (36), given j

$$\sigma_j = \operatorname{Tr}\left[\sigma \mid \eta_j \rangle \langle \eta_j \mid \right] = \operatorname{Tr}\left[\sum_{k=1}^n e^{-\lambda_k} \mid \eta_k \rangle \langle \eta_k \mid \mid \eta_j \rangle \langle \eta_j \mid \right] = e^{-\lambda_j}.$$
(42)

Expression (37) means for $k \neq l$

$$e^{-\lambda_k} e^{\lambda_k/2 - \lambda_l/2} = e^{-\lambda_k/2 - \lambda_l/2} = e^{-\lambda_l} e^{\lambda_l/2 - \lambda_k/2}.$$
(43)

From (35) and (36) we get

$$\overrightarrow{\sigma}Q = (\sigma_1, \sigma_2, ..., \sigma_n)Q = (0, 0, ..., 0).$$
 (44)

Example 2. Let

$$\sigma = \begin{pmatrix} \frac{1}{2} & 0 & 0\\ 0 & \frac{1}{3} & 0\\ 0 & 0 & \frac{1}{6} \end{pmatrix}.$$

Then

$$h = -\log \sigma = \begin{pmatrix} \log 2 & 0 & 0\\ 0 & \log 3 & 0\\ 0 & 0 & \log 6 \end{pmatrix},$$

so $\lambda_1 = \log 2$, $\lambda_2 = \log 3$ and $\lambda_3 = \log 6$. The Q matrix given by the expression (34) has entries

$$Q_{12} = 2e^{(\log 2 - \log 3)/2} = 2\left(\frac{2}{3}\right)^{1/2} = 2\frac{\sqrt{2}}{\sqrt{3}}$$

$$Q_{13} = 2e^{(\log 2 - \log 6)/2} = 2\left(\frac{2}{6}\right)^{1/2} = 2\frac{1}{\sqrt{3}}$$

$$Q_{11} = 2\left(-\frac{\sqrt{2}}{\sqrt{3}} - \frac{1}{\sqrt{3}}\right)$$

$$Q_{21} = 2e^{(\log 3 - \log 2)/2} = 2\left(\frac{3}{2}\right)^{1/2} = 2\frac{\sqrt{3}}{\sqrt{2}}$$

$$Q_{23} = 2e^{(\log 3 - \log 6)/2} = 2\left(\frac{3}{6}\right)^{1/2} = 2\frac{\sqrt{3}}{\sqrt{6}}$$

$$Q_{22} = 2\left(-\frac{\sqrt{3}}{\sqrt{2}} - \frac{1}{\frac{\sqrt{3}}{\sqrt{6}}}\right)$$

$$Q_{31} = 2e^{(\log 6 - \log 2)/2} = 2\left(\frac{6}{2}\right)^{1/2} = 2\sqrt{3}$$

$$Q_{32} = 2e^{(\log 6 - \log 3)/2} = 2\left(\frac{6}{3}\right)^{1/2} = 2\sqrt{2}$$

$$Q_{33} = 2(-\sqrt{3} - \sqrt{2})$$

Thus

$$Q = 2 \begin{pmatrix} -\frac{\sqrt{2}}{\sqrt{3}} - \frac{1}{\sqrt{3}} & \frac{\sqrt{2}}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ \frac{\sqrt{3}}{\sqrt{2}} & -\frac{\sqrt{3}}{\sqrt{2}} - \frac{\sqrt{3}}{\sqrt{6}} & \frac{\sqrt{3}}{\sqrt{6}} \\ \sqrt{3} & \sqrt{2} & -\sqrt{3} - \sqrt{2} \end{pmatrix}.$$

We should have that $\vec{\sigma} = (\frac{1}{2}, \frac{1}{3}, \frac{1}{6})$ is the invariant vector. In fact,

$$\frac{1}{2}(\vec{\sigma}Q)_1 = -\frac{(\sqrt{2}+1)}{2\sqrt{3}} + \frac{\sqrt{3}}{3\sqrt{2}} + \frac{\sqrt{3}}{6}$$
$$= \frac{-\sqrt{3}\sqrt{2} - \sqrt{3} + \sqrt{2}\sqrt{3} + \sqrt{3}}{6} = 0.$$

$$\begin{aligned} \frac{1}{2}(\vec{\sigma}Q)_2 &= \frac{\sqrt{2}}{2\sqrt{3}} - \frac{(1+\sqrt{3})}{3\sqrt{2}} + \frac{\sqrt{2}}{6} \\ &= \frac{\sqrt{2}\sqrt{3} - \sqrt{2} - \sqrt{2}\sqrt{3} + \sqrt{2}}{6} = 0. \end{aligned}$$

$$\frac{1}{2}(\vec{\sigma}Q)_3 = \frac{1}{2\sqrt{3}} + \frac{1}{3\sqrt{2}} - \frac{(\sqrt{2} + \sqrt{3})}{6}$$
$$= \frac{\sqrt{3} + \sqrt{2} - \sqrt{2} - \sqrt{3}}{6} = 0.$$

Example 3. Let

$$\sigma = \begin{pmatrix} \frac{1}{4} & 0 & \frac{i}{8} \\ 0 & \frac{1}{2} & 0 \\ -\frac{i}{8} & 0 & \frac{1}{4} \end{pmatrix}.$$

The eigenvalues of σ are $\frac{1}{8}$, $\frac{3}{8}$ and $\frac{1}{2}$. Then $h = -\log \sigma$ has eigenvalues $\lambda_1 = \log 8$, $\lambda_2 = \log \frac{8}{3}$ and $\lambda_3 = \log 2$. So, the Q matrix given by the expression (34) has entries

$$Q_{12} = 2e^{(\log 8 - \log \frac{8}{3})/2} = 2(3)^{1/2} = 2\sqrt{3}.$$
$$Q_{13} = 2e^{(\log 8 - \log 2)/2} = 2\left(\frac{8}{2}\right)^{1/2} = 4.$$
$$Q_{11} = -2\left(\sqrt{3} + 2\right).$$
$$Q_{21} = 2e^{(\log \frac{8}{3} - \log 8)/2} = 2\left(\frac{1}{3}\right)^{1/2} = \frac{2}{\sqrt{3}}.$$

$$Q_{23} = 2e^{(\log \frac{8}{3} - \log 2)/2} = 2\left(\frac{4}{3}\right)^{1/2} = \frac{4}{\sqrt{3}}.$$
$$Q_{22} = -\frac{6}{\sqrt{3}}.$$
$$Q_{31} = 2e^{(\log 2 - \log 8)/2} = 2\left(\frac{1}{4}\right)^{1/2} = 1.$$
$$Q_{32} = 2e^{(\log 2 - \log \frac{8}{3})/2} = 2\left(\frac{3}{4}\right)^{1/2} = \sqrt{3}.$$
$$Q_{33} = -\left(1 + \sqrt{3}\right).$$

Thus

$$Q = \begin{pmatrix} -2(\sqrt{3}+2) & 2\sqrt{3} & 4\\ \frac{2}{\sqrt{3}} & -\frac{6}{\sqrt{3}} & \frac{4}{\sqrt{3}}\\ 1 & \sqrt{3} & -(1+\sqrt{3}) \end{pmatrix}.$$

We should have that $\vec{\sigma} = (\frac{1}{8}, \frac{3}{8}, \frac{1}{2})$ is the invariant vector. In fact,

$$(\vec{\sigma}Q)_1 = \frac{1}{8} \left(-2\sqrt{3} - 4 \right) + \frac{3}{8} \frac{2}{\sqrt{3}} + \frac{1}{2}$$
$$= -\frac{\sqrt{3}}{4} - \frac{1}{2} + \frac{\sqrt{3}}{4} + \frac{1}{2} = 0.$$
$$(\vec{\sigma}Q)_2 = \frac{\sqrt{3}}{4} - \frac{3\sqrt{3}}{4} + \frac{2\sqrt{3}}{4} = 0.$$
$$1 - \sqrt{3} - \frac{1}{4} \left(-\frac{\sqrt{3}}{4} - \frac{\sqrt{3}}{4} - \frac{\sqrt{3}}{4$$

$$(\vec{\sigma}Q)_3 = \frac{1}{2} + \frac{\sqrt{3}}{2} - \frac{1}{2}\left(1 + \sqrt{3}\right) = 0.$$

Related results are described in (3) and (4) in [12].

Inst. Mat. UFRGS, Av. Bento Gonvalves 9500 - 91.500 Porto Alegre, Brazil

Email of J. E. Brasil is jaderebrasil@gmail.com Email of J. Knorst is jojoknorst@gmail.com Email of A. O. Lopes is arturoscar.lopes@gmail.com

References

- R. Alicki and M. Fannes. Quantum Dynamical Systems. Oxford University Press (2000).
- [2] A. Baraviera, R. Exel and A. O. Lopes, A Ruelle Operator for continuous time Markov Chains, Sao Paulo Journal of Mathematical Sciences, vol 4 n. 1, pp 1-16 (2010)
- [3] J. E. Brasil, and J. Knorst and A. O. Lopes, Thermodynamic Formalism for Quantum Channels: Entropy, Pressure, Gibbs Channels and generic properties, on line - Communications in Contemporary Mathematics.
- [4] J. E. Brasil, and J. Knorst and A. O. Lopes, Lyapunov exponents for Quantum Channels: an entropy formula and generic properties, Journal of Dynamical Systems and Geometric Theories, 21(2) 155-187 (2021)
- [5] E. A. Carlen and J. Maas, Gradient flow and entropy inequalities for quantum Markov semigroups with detailed balance, Journal of Functional Analysis, 273. 1810–1869 (2017)
- [6] M.-H. Chang, Quantum Stochastics. Cambridge University Press, (2015)
- [7] F. Cipriani, Dirichlet Forms and Markovian Semigroups on Standard Forms of von Neumann Algebras, Journal of functional analysis 147, 259–300 (1997)
- [8] F. Fagnola and R. Rebolledo, From classical to quantum entropy production, In Quantum Probability and Infinite Dimensional Analysis, 245-261 (2010)
- [9] V. Jaksic, C.-A. Pillet and M. Westrich, Entropic fluctuations of quantum dynamical semigroups, J. Stat. Phys., 154(1-2), 153–187, (2014)
- [10] M. Kac, Integration in function spaces and some of its applications, Acad Naz dei Lincei Scuola Superiore Normale Superiore, Piza, Italy (1980).
- [11] J. Knorst, A. O. Lopes, G. Muller and A. Neumann, Thermodynamic Formalism on the Skhorohd space: the continuous time Ruelle operator, entropy, pressure, entropy production and expansiveness. preprint UFRGS (2022)
- [12] M. de Leeuw, C. Paletta and Balazs Pozsgay, Constructing Integrable Lindblad Superoperators, arXiv (2021)

- [13] A. O. Lopes, A. Neumann and Ph. Thieullen, A thermodynamic formalism for continuous time Markov chains with values on the Bernoulli Space: entropy, pressure and large deviations, Journ. of Statist. Phys. Volume 152, Issue 5, Page 894-933 (2013)
- [14] A. O. Lopes, J. K. Mengue, J. Mohr, and R. R. Souza, Entropy and variational principle for one-dimensional lattice systems with a general *a priori* probability: positive and zero temperature. Ergodic Theory Dynam. Systems, 35(6):1925–1961 (2015)
- [15] W. Parry and M. Pollicott. Zeta functions and the periodic orbit strucuture of hyperbolic dynamics. Asterisque, Vol 187–188, pp 1–268 (1990).
- [16] C. A. Pillet, Quantum dynamical systems, In Open Quantum Systems I. The Hamiltonian Approach. S. Attal, A. Joye and C.-A. Pillet editors. Lecture Notes in Mathematics 1880. Springer, Berlin (2006)
- [17] D. W. Stroock, An Introduction to the Theory of Large Deviations, Springer Verlag (1984)
- [18] M. M. Wolf, Quantum Channels and Operations Guided Tour. 2010. https://www-m5.ma.tum.de/foswiki/pub/M5/
 Allgemeines/MichaelWolf/ QChannelLecture.pdf