REALIZABLE RANKS OF JOINS AND INTERSECTIONS OF SUBGROUPS IN FREE GROUPS

IGNAT SOROKO

ABSTRACT. The famous Hanna Neumann Conjecture (now the Friedman–Mineyev theorem) gives an upper bound for the ranks of the intersection of arbitrary subgroups H and K of a non-abelian free group. It is an interesting question to 'quantify' this bound with respect to the rank of $H \vee K$, the subgroup generated by H and K. We describe a set of realizable values $(rk(H \vee K), rk(H \cap K))$ for arbitrary H, K, and conjecture that this locus is complete. We study the combinatorial structure of the topological pushout of the core graphs for H and K with the help of graphs introduced by Dicks in the context of his Amalgamated Graph Conjecture. This allows us to show that certain conditions on ranks of $H \vee K$, $H \cap K$ are not realizable, thus resolving the remaining open case m = 4 of Guzman's "Group-Theoretic Conjecture" in the affirmative. This in turn implies the validity of the corresponding "Geometric Conjecture" on hyperbolic 3–manifolds with a 6–free fundamental group. Finally, we prove the main conjecture describing the locus of realizable values for the case when rk(H) = 2.

1. INTRODUCTION

Let F be a free group and $H, K \leq F$ finitely generated subgroups. Denote $H \vee K$ the subgroup of F generated by H and K. Define the *reduced rank* of H by

$$\operatorname{rr}(H) = \max(0, \operatorname{rk}(H) - 1).$$

The famous Hanna Neumann Conjecture (now the Friedman–Mineyev theorem [24], [8], [11], [7]; see also a recent proof of Jaikin-Zapirain [19]) states that

$$\operatorname{rr}(H \cap K) \le \operatorname{rr}(H) \operatorname{rr}(K).$$

It is an interesting problem to try to 'quantify' the possible ranks of $H \cap K$ with respect to the rank of the join $H \vee K$ of H and K (i.e. the subgroup generated by H and K). Ideally, one wishes to determine the set of all realizable values for tuples

$$(\operatorname{rk}(H \lor K), \operatorname{rk}(H \cap K))$$

for any given values of $\operatorname{rk}(H)$ and $\operatorname{rk}(K)$. It seems plausible, by the analogy with the linear algebra identity for vector spaces, $\dim(U \cap V) + \dim(U + V) = \dim(U) + \dim(V)$, that the bigger $\operatorname{rk}(H \cap K)$ is, the smaller $\operatorname{rk}(H \vee K)$ should be. Several partial results and conjectures have been made in this direction.

In [16] Imrich and Müller have proved the following:

Theorem (Imrich–Müller, 1994). If H, K are finitely generated subgroups of F and either H or K is of finite index in $H \vee K$ then

$$\operatorname{rr}(H \cap K)\operatorname{rr}(H \vee K) \le \operatorname{rr}(H)\operatorname{rr}(K).$$
(1)

In general, without the finite index assumption, this inequality does not hold, see [16, Ex. 3]. Moreover, Hunt [15] has shown that the ratio of the left-hand side of (1) to its right-hand side can be made arbitrarily large. Recently, Sergei Ivanov [17, p. 826] has posed the following open question:

Question (Ivanov, 2017). Does inequality (1) hold true if $\operatorname{rr}(H \cap K)$ is the maximal possible in the Friedman-Mineyev theorem, i.e. if $\operatorname{rr}(H \cap K) = \operatorname{rr}(H)\operatorname{rr}(K) > 0$? Equivalently, does this assumption imply that $\operatorname{rr}(H \vee K) = 1$?

Another circle of questions about the relationship between $\operatorname{rk}(H \cap K)$ and $\operatorname{rk}(H \vee K)$ was motivated by the study of hyperbolic 3-manifolds. Continuing the program started by Agol, Culler and Shalen [1], [5], Guzman [13] formulated the following "Group-Theoretic Conjecture" (GTC):

Date: October 31, 2019.

²⁰¹⁰ Mathematics Subject Classification. Primary 20E05, 20E07, 20F65, 57M07.

Conjecture (GTC, Guzman, 2014). If two subgroups $H, K \leq F$ both have ranks equal to $m \geq 2$, and $\operatorname{rk}(H \cap K) \geq m$, then $\operatorname{rk}(H \vee K) \leq m$.

She proved that this conjecture, if true, implies the following "Geometric Conjecture" (GC), with k = m + 2 (recall that a group is called *k*-free if all of its *k*-generator subgroups are free):

Conjecture (GC, Guzman, 2014). Let M be a closed, orientable, hyperbolic 3-manifold. If $\pi_1(M)$ is k-free for $k \geq 3$ then there exists a point P in M such that the set of all elements of $\pi_1(M, P)$ represented by loops of length less than $\log(2k-1)$ is contained in a (free) subgroup of $\pi_1(M)$ of rank $\leq k-3$.

The case m = 2 of the GTC appeared earlier as a question in a preprint of Culler and Shalen and was subsequently resolved in the affirmative by Kent [22] and, independently, by Louder–McReynolds [23]. This was used in [5, Th. 1.4] to prove a statement equivalent to the GC for k = 4 and to obtain a lower bound on the volume of a closed orientable hyperbolic 3–manifold with a 4–free fundamental group. (The case k = 3of the GC and the corresponding lower bound on the volume for the case of 3–free fundamental groups was proved earlier in [1, Cor. 9.3].) Using results and techniques from [22], Guzman proves the GTC for m = 3and hence the GC for k = 5. However, Hunt [15] has shown by example that the GTC is no longer true for m = 5. Below we will show that the GTC is false for all values $m \ge 6$, but holds true for m = 4.

It must be noted that, very recently, Guzman and Shalen [14] proved the Geometric Conjecture in full generality, without dependence on the Group-Theoretic Conjecture.

It is an easy consequence of the Hopfian property of finite rank free groups that the only possibility for $\operatorname{rk}(H \lor K)$ to equal the maximal possible value $\operatorname{rk}(H) + \operatorname{rk}(K)$ is to have $H \lor K \cong H * K$. Thus, in this case $\operatorname{rk}(H \cap K)$ must equal 0.

In [22], Kent proved the following inequality:

Theorem (Kent, 2009). Let H and K be nontrivial finitely generated subgroups of F with reduced ranks h = rr(H), k = rr(K), and $k \ge h$. Assume also that $H \cap K \ne 1$. Then

$$\operatorname{rr}(H \cap K) \le 2hk - h\operatorname{rr}(H \vee K).$$

Recently, Sergei Ivanov has improved the above estimate of Kent, see [18, (4.2)] (where he mentions that inequality (2) was also obtained independently by Dicks):

Theorem (Ivanov, 2018). Let H and K be nontrivial finitely generated subgroups of F with reduced ranks h = rr(H), k = rr(K). Then

$$\operatorname{rr}(H \cap K) \leq \frac{1}{2} \big(h + k - \operatorname{rr}(H \vee K) \big) \big(h + k - \operatorname{rr}(H \vee K) + 1 \big).$$

$$\tag{2}$$

The last result suggests that it may be convenient to describe the locus of possible values $(\operatorname{rk}(H \lor K), \operatorname{rk}(H \cap K))$ in terms of the difference between $\operatorname{rr}(H \lor K)$ and its largest possible value h+k+1. If we denote $i = h + k + 1 - \operatorname{rr}(H \lor K)$ then inequality (2) reads: $\operatorname{rr}(H \cap K) \leq \frac{i(i-1)}{2}$.

The results mentioned so far do not guarantee that if certain numbers (v, c) satisfy the respective inequalities, then there exist subgroups H, K realizing them as $(v, c) = (\operatorname{rk}(H \lor K), \operatorname{rk}(H \cap K))$. Hence these results are, in effect, describing the regions of tuples which are *non-realizable*. By contrast, in [21], Kent exhibited for arbitrary $h, k \ge 2$ a family of subgroups H = H(h, k, m) and K = K(k) such that $\operatorname{rk}(H) = h$, $\operatorname{rk}(K) = k$, $\operatorname{rk}(H \lor K) = 2$ and $\operatorname{rk}(H \cap K)$ takes on all possible ranks $m = 0, \ldots, (h-1)(k-1) + 1$ allowed by the Friedman–Mineyev theorem. This answered a question of Myasnikov [3, (AUX1)].

Our first contribution to this theme is the following addition to the region of known realizable values of $(\operatorname{rk}(H \lor K), \operatorname{rk}(H \cap K))$.

Theorem 1.1. Let F be a free group and let integers h, k, c, v satisfy $2 \le h \le k$, $2 \le v \le h + k$, $0 \le c \le (h-1)(k-1) + 1$. Define a sequence a_i as follows:

$$a_0 = 0;$$

 $a_i = \left\lfloor \frac{i^2}{4} \right\rfloor + 1, \quad for \quad i = 1, \dots, 2(h-1);$
 $a_i = (h-1)(i-h+1) + 1, \quad for \quad i = 2(h-1), \dots, h+k-2.$





FIGURE 1. The locus of realizable values for rk(H) = rk(K) = 6. Dark gray area: known realizable values. White area: proved non-realizable values. Light gray area: conjecturally non-realizable values (Conjecture 1.3). The cells marked with an asterisk are non-realizable by Theorem 1.5. Guzman's GTC for m = 6 claims that the red rectangle consists entirely of non-realizable values.

If we denote i = h + k - v, then for any $c \leq a_i$ there exist subgroups $H, K \leq F$ such that rk(H) = h, $\operatorname{rk}(K) = k$, $\operatorname{rk}(H \cap K) = c$ and $\operatorname{rk}(H \vee K) = v$.

The sequence a_i from this theorem is a splicing of a discrete quadratic function and a linear function. The linear part exists only if h < k. Written in terms of reduced ranks, the quadratic part implies: $\operatorname{rr}(H \cap K) \leq \left|\frac{i^2}{4}\right|$, which is smaller than Ivanov's upper bound $\frac{i(i-1)}{2}$ above, and the gap between the two becomes unbounded as i grows. If h = 2, the quadratic part trivializes and the sequence a_i becomes especially simple: $a_i = i$ for all i, see Figure 2.

The realizable values from Theorem 1.1 allow us to establish the following.

(i = 10) 2

3

 $\mathbf{4}$

 $\mathbf{5}$

6

Corollary 1.2. Guzman's "Group-Theoretic Conjecture" does not hold for any m > 5.

Figure 1 depicts all the regions described above for the case rk(H) = rk(K) = 6.

We conjecture that the set of realizable values from Theorem 1.1 is complete.

Conjecture 1.3. Let F be a free group and let integers h, k, v, c satisfy: $2 \le h \le k, 2 \le v \le h + k$ and $0 \leq c \leq (h-1)(k-1) + 1$. Then there exist subgroups $H, K \leq F$ such that $\operatorname{rk}(H) = h$, $\operatorname{rk}(K) = k$, $\operatorname{rk}(H \lor K) = v$, and $\operatorname{rk}(H \cap K) = c$ if and only if $c \leq a_i$ for i = h + k - v, where a_i is the sequence defined in Theorem 1.1.

The author tested this conjecture on a computer (searching for a possible counterexample) using a Monte-Carlo type algorithm of Bassino, Nicaud and Weil [2], which randomly generates core graphs on a given number of vertices with the uniform distribution for subgroups of the given size (of their core graph) in a free group. The author learned about this algorithm from the preprint of Hunt [15], and also used the computer algebra system GAP [12] (with the package FGA [25] for methods dealing with free groups) to implement it. Testing about $5 \cdot 10^8$ pairs of random core graphs did not produce any values outside of the conjectured locus.

Note that Conjecture 1.3 subsumes the open question of Ivanov above.

We prove this conjecture for the special case of h = rk(H) = 2 (when the sequence a_i from Theorem 1.1 becomes linear: $a_i = i$ for all i):

Theorem 1.4. Let F be a free group and let integers k, v, c satisfy: $k \ge 2, 2 \le v \le k+2$ and $0 \le c \le k$. Then there exist subgroups $H, K \leq F$ such that $\operatorname{rk}(H) = 2$, $\operatorname{rk}(K) = k$, $\operatorname{rk}(H \vee K) = v$, and $\operatorname{rk}(H \cap K) = c$ if and only if $c + v \leq k + 2$.

The diagram in Figure 2 shows all realizable values for rk(H) = 2, rk(K) = 10.



FIGURE 2. All realizable values for rk(H) = 2, rk(K) = 10.

Our next contribution is the resolution of a few extremal cases which are not covered by the realizable values from Theorem 1.1 and are not eliminated by Ivanov's inequality (2) (see the cells marked with an asterisk in Figure 1).

Theorem 1.5. Let F be a free group. Then there do not exist subgroups $H, K \leq F$ such that $\operatorname{rk}(H)$, $\operatorname{rk}(K) \geq 2$, $\operatorname{rk}(H \vee K) = \operatorname{rk}(H) + \operatorname{rk}(K) - i$ for some $i \geq 3$, and $\operatorname{rk}(H \cap K) = \frac{i(i-1)}{2} + 1$.

As an immediate consequence we conclude that Guzman's "Group-Theoretic Conjecture" holds true for the remaining unresolved case of m = 4:

Corollary 1.6. Let F be a free group. If two subgroups $H, K \leq F$ both have ranks equal to 4, and $\operatorname{rk}(H \cap K) \geq 4$, then $\operatorname{rk}(H \vee K) \leq 4$.

Invoking the implication theorem from [13], we obtain a proof of the "Geometric Conjecture" for k = 6:

Corollary 1.7. Let M be a closed, orientable, hyperbolic 3-manifold. If $\pi_1(M)$ is 6-free then there exists a point P in M such that the set of all elements of $\pi_1(M, P)$ represented by loops of length less than $\log(11)$ is contained in a free subgroup of $\pi_1(M)$ of rank at most 3.

Our paper is organized as follows.

Section 2 contains the basic definitions of graph related concepts necessary for our needs. We introduce Stallings' core graphs and describe the construction of the topological pushout of two core graphs, following Kent [22]. The topological pushout is an intermediate object between the join of two core graphs Γ_H , Γ_K and the core graph $\Gamma_{H\vee K}$ of the join of the two subgroups, and the rank of the topological pushout serves as an upper bound for the rank of the join.

In Section 3 we show how to obtain all the values of $(\operatorname{rk}(H \lor K), \operatorname{rk}(H \cap K))$ from Theorem 1.1 by properly adding new generators to the family of examples exhibited by Kent in [21].

In Section 4 we study the combinatorial structure of the topological pushout in terms of graphs introduced by Dicks in [6] in the context of the Amalgamated Graph Conjecture. Our approach is motivated by the construction of graphs Υ and Z in section 6 of [6]. In Section 5 we establish a technical condition on Dicks graphs which specifies when the rank of the topological pushout is the maximal possible.

Finally, in Section 6, we use the results obtained so far to prove Theorems 1.4, 1.5 and the consequences of the latter, the Guzman's GTC for the remaining case of m = 4, and hence the GC for k = 6. The key observation is that in the situation described in Theorems 1.4, 1.5, the components of the corresponding Dicks graphs Ω_{abc} are 'incompatible' in the sense that one component contains a highly connected subgraph (a complete bipartite graph $K_{i,i-1}$ in Theorem 1.5 and $K_{2,m}$ in Theorem 1.4), while others are singleton vertices. Dicks' duality implies the existence of an isomorphic copy of the highly connected subgraph in Ω , which must be 'spread' along two or more subgraphs Ω_{ab} , Ω_{ac} , Ω_{bc} . This forces the rank of the join to be less than required, which makes the corresponding tuples from Theorems 1.4, 1.5 non-realizable. Acknowledgments. The author is very grateful to Jing Tao for bringing Guzman's "Group-Theoretic Conjecture" to his attention and for supporting him with the research assistantship from her grants (NSF grant DMS 1611758 and NSF Career grant DMS 1651963). The author extends his gratitude to Noel Brady for his interest in this project and numerous constructive discussions and to Sergei Ivanov who suggested to the author that Warren Dicks' methods may prove useful for this project. The author would also like to thank the anonymous referee for numerous suggestions which greatly improved the clarity of the exposition, and Pallavi Dani, Warren Dicks, Max Forester, Autumn Kent, and Lars Louder for their valuable remarks. Last but not least the author would like to thank Till Tantau and Matthew Fayers for creating IATEX packages TikZ [27] and genyoungtabtikz [10], respectively, which proved to be extremely useful in typesetting this article.

2. Graphs

In this paper we will deal with two types of graphs: directed labeled graphs and undirected ones. Among the former are Stallings' core graphs that are used to represent finitely generated subgroups of a free group, see [26], [20]. Among the latter ones, are bipartite graphs introduced by Dicks [6] to study the structure of the intersection of two subgroups in a free group; they will be useful for the description of the topological pushout in the sense of Kent [22]. We now remind the reader of the relevant definitions adapted to our needs from [4], [20], [6].

2.1. **Basic definitions.** A graph Γ is a pair of sets $V(\Gamma)$, $E(\Gamma)$, where $V(\Gamma)$ is a nonempty set of vertices of Γ and $E(\Gamma)$ is a set of *(directed) edges* of Γ equipped with the three maps: $o: E(\Gamma) \to V(\Gamma)$, $t: E(\Gamma) \to V(\Gamma)$ and $\overline{-}: E(\Gamma) \to E(\Gamma)$ called the *origin* map, the *terminus* map and the map of taking the *inverse* of an edge, respectively, with the following properties: for each $e \in E(\Gamma)$, $\overline{\overline{e}} = e, \overline{e} \neq e$ and $o(\overline{e}) = t(e)$.

A morphism between two graphs Γ and Δ is a map $\pi \colon \Gamma \to \Delta$ that sends vertices to vertices and edges to edges and has the property that $o(\pi(e)) = \pi(o(e)), t(\pi(e)) = \pi(t(e))$ and $\pi(\overline{e}) = \overline{\pi(e)}$ for any edge $e \in E(\Gamma)$.

Each graph Γ admits a geometric realization as a 1-dimensional CW complex X_{Γ} , with vertices of Γ being the 0-cells of X_{Γ} , and each pair of mutually inverse edges e, \overline{e} of Γ corresponding to the two opposite orientations of the same open 1-cell of X_{Γ} .

A graph Γ is called *directed* (or *oriented*) if in each pair of its mutually inverse edges e, \overline{e} one edge is chosen, which is called *positively oriented*, and the other is called *negatively oriented*. The set of all positively (negatively) oriented edges is denoted $E^+(\Gamma)$ (respectively, $E^-(\Gamma)$). A morphism of directed graphs $\pi: \Gamma \to \Delta$ is required to send $E^+(\Gamma)$ to $E^+(\Delta)$.

Let \mathcal{A} be a finite alphabet, and \mathcal{A}^{-1} be the set of formal inverses of \mathcal{A} . A directed \mathcal{A} -labeled graph (or just a directed labeled graph, if \mathcal{A} is obvious from the context) is a directed graph Γ with a labeling $\mu: E(\Gamma) \to \mathcal{A} \sqcup \mathcal{A}^{-1}$ such that $\mu(E^+(\Gamma)) \subseteq \mathcal{A}$ and $\mu(\overline{e}) = \mu(e)^{-1}$ for each $e \in E(\Gamma)$. A morphism of directed \mathcal{A} -labeled graphs $\pi: \Gamma \to \Delta$ is required to preserve the labeling, i.e. $\mu(\pi(e)) = \mu(e)$ for each $e \in E(\Gamma)$.

The star of a vertex v of $V(\Gamma)$ is the set of all edges e in $E(\Gamma)$ such that o(e) = v. The star of v can be thought of as the link of v in the geometric realization of Γ in our context. The valence val(v) of v is the cardinality of the star of v. If k = val(v) we call the vertex v k-valent.

A morphism $\pi: \Gamma \to \Delta$ is called an *immersion* if its restriction to the star of each vertex of Γ is injective. A path p in Γ is a sequence of edges $p = e_1, \ldots, e_k$ of $E(\Gamma)$ such that for each $i = 2, \ldots, k$, we have $o(e_i) = t(e_{i-1})$. The length of p is set to be k (with the case k = 0 possible). In this situation we call vertex $x = o(e_1)$ the origin of p and $y = t(e_k)$ the terminus of p. We also say that p is a path from x to y, and use notation x-y to denote any such path.

If Γ is a directed labeled graph, then any path $p = e_1, \ldots, e_k$ has a naturally defined label $\mu(p) = \mu(e_1) \ldots \mu(e_k)$, which is a word in the alphabet $\mathcal{A} \sqcup \mathcal{A}^{-1}$. (If k = 0 then $\mu(p) = 1$, the empty word.)

The notion of the fundamental group of a graph is a combinatorial analog of the notion of the fundamental group of the geometric realization of the graph, see [4, Ch. 2.4]. In what follows, we will not distinguish graphs and their geometric realizations and will use these notions interchangeably. All directed labeled graphs will be labeled by the set $\mathcal{A} = \{a, b, c\}$ of free generators of a rank 3 free group F(a, b, c).

With each graph Γ we can associate an *undirected graph* Γ_u which has the same set of vertices $V(\Gamma)$ but whose set of *undirected edges* is obtained by identifying each pair $\{e, \overline{e}\}$ of mutually inverse directed edges of Γ into a single equivalence class. Such an undirected edge has two vertices that are *incident* to it, namely $\{o(e), t(e)\} = \{o(\overline{e}), t(\overline{e})\}$, and these two vertices may coincide if e is a loop. We also say that such vertices



FIGURE 3. The graph X with $\pi_1(X, u)$ equal to $F(ca^{-1}, cb^{-1}) \leq F(a, b, c)$.

are *adjacent* to each other. We will abuse the notation and denote an undirected edge $\{e, \overline{e}\}$ simply by e, and the set of all undirected edges of Γ_u also by $E(\Gamma_u)$. A *path* in an undirected graph is a sequence $p = x_1e_1x_2e_2\ldots x_ke_kx_{k+1}$ of pairwise distinct vertices x_i and undirected edges e_i such that for all i, the vertices x_i, x_{i+1} are incident to the edge e_i . We say that p is a path from x_1 to x_{k+1} , and we will also denote it as x_1-x_{k+1} . The length of it is k, as above, with k = 0 possible. If we denote a subpath $e_1x_2e_2\ldots x_ke_k$ as q, we can also write: $p = x_1qx_{k+1}$. A cycle in an undirected graph Γ is a union of a path $x_0e_0x_1e_1\ldots x_k$ with an edge e_k which is incident to both x_k and x_0 . We denote a cycle also as $x_0e_0x_1e_1\ldots x_ke_kx_0$ and consider its length to be equal k.

2.2. Core graphs represent subgroups. Let F = F(a, b) be a free group of rank 2 and let X be a finite graph viewed as a 1-dimensional CW complex such that $\pi_1(X)$ is isomorphic to F. Traditionally X is identified with a wedge of two circles, but we will implement Dicks' approach [6] and take X to be the graph with two 0-cells u, v and three 1-cells a, b, c all originating at u and terminating at v, see Figure 3. If we take u as the basepoint for X, this realizes F as a subgroup of a rank 3 free group F(a, b, c) on free generators $\{a, b, c\}$ with the inclusion $\theta: F(a, b) \hookrightarrow F(a, b, c)$ given by $a \mapsto ca^{-1}, b \mapsto cb^{-1}$. (Choosing v as the basepoint of X yields an inclusion given by $a \mapsto a^{-1}c$ and $b \mapsto b^{-1}c$.)

For any subgroup $H \leq F$ there is a covering $\tilde{X}_H \to X$ corresponding to H. If we fix the vertex u (or v, for that matter) as the basepoint of X, there is a choice of the basepoint x_H in \tilde{X}_H such that $\pi_1(\tilde{X}_H, x_H)$ is identical to H. (Such choice is not unique if \tilde{X}_H has a nontrivial deck transformation.) Let Γ_H be the smallest subgraph of \tilde{X}_H containing x_H that carries $\pi_1(\tilde{X}_H, x_H) = H$. We call (Γ_H, x_H) the core graph for H. The vertices of Γ_H fall into two classes: 'sources' (preimages of $u \in X$) and 'sinks' (preimages of $v \in X$). Every edge of Γ_H is oriented from a source to a sink and inherits a unique label a, b, or c induced by the covering map $\tilde{X}_H \to X$. Notice that this labeling is proper in the sense that for every vertex x of Γ_H and each letter $\eta \in \{a, b, c\}$ there is at most one edge in Γ_H with the origin x labeled η and there is at most one edge in Γ_H with the terminus x labeled η . Notice also that any vertex of Γ_H is at most 3-valent, and the only vertex that may have valence 1 is the basepoint x_H .

In what follows, we will call the edges of Γ_H which map to the edge *a* (respectively, *b*, *c*) of *X*, as *a*-edges (resp., *b*-edges, *c*-edges), and paint them in diagrams with red (resp., blue, green) color. (We also depict *a*-edges with a single arrow, *b*-edges with a double arrow, and *c*-edges with a solid triangular arrow in all diagrams.)

2.3. Intersection of subgroups is represented by pullback. Let $H, K \leq F$ be two finitely generated subgroups of F and (Γ_H, x_H) , (Γ_K, x_K) be the corresponding core graphs, with the natural maps $p_H: (\Gamma_H, x_H) \to (X, u), p_K: (\Gamma_K, x_K) \to (X, u)$, which are injective on links of vertices, i.e. are immersions. Let $G_{H\cap K}$ be the pullback of these maps, defined as follows. The vertex set of $G_{H\cap K}$ is $V(\Gamma_H) \times V(\Gamma_K)$ and there is an oriented edge labeled η ($\eta \in \{a, b, c\}$) from the vertex (p, q) to the vertex (r, s) in $G_{H\cap K}$ if and only if there is an edge labeled η from p to r in Γ_H and an edge labeled η from q to s in Γ_K . The natural projections $V(\Gamma_H) \times V(\Gamma_K) \to V(\Gamma_H)$ and $V(\Gamma_H) \times V(\Gamma_K) \to V(\Gamma_K)$ give rise to immersions $\Pi_H: G_{H\cap K} \to \Gamma_H, \Pi_K: G_{H\cap K} \to \Gamma_K$, and the fundamental group of the component of $G_{H\cap K}$ containing the basepoint (x_H, x_K) is equal to $H \cap K$, see [26, Th. 5.5]. Denote by $\Gamma_{H\cap K}$ the minimal subgraph of $G_{H\cap K}$ that contains (x_H, x_K) and carries the fundamental group of the connected component of (x_H, x_K) . Then $(\Gamma_{H\cap K}, (x_H, x_K))$ is the core graph for $H \cap K$. The construction guarantees that the two compositions of immersions $(\Gamma_{H\cap K}, (x_H, x_K)) \xrightarrow{\Pi_H} (\Gamma_H, x_H) \xrightarrow{p_H} (X, u)$ and $(\Gamma_{H\cap K}, (x_H, x_K)) \xrightarrow{\Pi_K} (\Gamma_K, x_K) \xrightarrow{p_K} (X, u)$ commute and thus define the canonical immersion $(\Gamma_{H\cap K}, (x_H, x_K)) \to (X, u)$.

Example 2.1. Figure 4 shows the core graphs Γ_H , Γ_K , $\Gamma_{H\cap K}$ for the subgroups $H, K \leq \theta(F) \leq F(a, b, c)$ given by $H = \theta(\langle a, bab^{-1} \rangle) = \langle ca^{-1}, cb^{-1}ca^{-1}bc^{-1} \rangle$, $K = \theta(\langle b^{-1}a, ba \rangle) = \langle ba^{-1}, cb^{-1}ca^{-1} \rangle$, and for their intersection $H \cap K = \theta(\langle bab^{-1}a \rangle) = \langle cb^{-1}ca^{-1}bc^{-1} \rangle$.



FIGURE 4. The core graphs for the subgroups $H, K, H \cap K$ of Example 2.1 and their topological pushout \mathcal{T} . The thin dashed *c*-edges belong to $G_{H\cap K}$ but not to $\Gamma_{H\cap K}$. The set of all *a*-edges of $\Gamma_H \sqcup \Gamma_K$ and the set of all *b*-edges of $\Gamma_H \sqcup \Gamma_K$ each form their own single equivalence class of edges in \mathcal{T} , whereas the *c*-edges of $\Gamma_H \sqcup \Gamma_K$ form two 2-element classes of edges in \mathcal{T} : {(1,2), (7,8)} and {(3,4), (5,6)}.

Now we show that without loss of generality we can assume that the core graphs Γ_H , Γ_K and $\Gamma_{H\cap K}$ do not have vertices of valence 1.

Lemma 2.2. For any $H, K \leq \theta(F) \leq F(a, b, c)$ such that $H \cap K \neq 1$, we can find an element $g \in F(a, b, c)$ such that all the core graphs $\Gamma_{H^g}, \Gamma_{K^g}, \Gamma_{H^g \cap K^g}$ for the conjugated subgroups $H^g, K^g, H^g \cap K^g$, respectively, do not have vertices of valence 1.

Proof. The only vertices in Γ_H , Γ_K , $\Gamma_{H\cap K}$ that may have valence 1 are the basepoints x_H , x_K , (x_H, x_K) , respectively. If (x_H, x_K) has valence 2 or more, then its projections x_H and x_K also have valence 2 or more, and there is nothing to prove.

Now suppose that the basepoint $p = (x_H, x_K)$ of $\Gamma_{H\cap K}$ has valence 1. Since $H \cap K \neq 1$, the graph $\Gamma_{H\cap K}$ must have a closed circuit, and the vertex p does not belong to it. Therefore there exists a vertex of valence 3 in $\Gamma_{H\cap K}$. Let p-q be the shortest path in $\Gamma_{H\cap K}$ to a valence 3 vertex q, and let $w \in F(a, b, c)$ be the label on this path. Then projections $\Pi_H(p-q)$ and $\Pi_K(p-q)$ are immersed paths in Γ_H , Γ_K , respectively, with the same label w on them. Since q is a valence 3 vertex in $\Gamma_{H\cap K}$, the vertices $q_H = \Pi_H(q)$ and $q_K = \Pi_K(q)$ also have valence 3. However, one may have other vertices of valence 3 on the paths $\Pi_H(p-q) = x_H-q_H$ and $\Pi_K(p-q) = x_K - q_K$. Let q'_H be a vertex on the path $x_H - q_H$ defined as follows: $q'_H = x_H$, if x_H has valence greater than 1 in Γ_H , and q'_H is the vertex of valence 3 on the path $x_H - q_H$ with the least distance along this path from x_H , otherwise. Define q'_K similarly in Γ_K . Then conjugating by $g = w^{-1}$ inside F(a, b, c) yields a triple of subgroups $H^g, K^g, H^g \cap K^g$ such that their core graphs $\Gamma_{H^g}, \Gamma_{K^g}, \Gamma_{H^g \cap K^g}$ differ from Γ_H , $\Gamma_K, \Gamma_{H \cap K}$ by moving their basepoint to vertices q_H, q_K , and q, respectively, and deleting the hanging trees p-q from $\Gamma_{H \cap K}$ and $x_H - q'_H, x_K - q'_K$ from Γ_H, Γ_K , respectively. (Vertices q, q'_H, q'_K themselves are not deleted.) This gives us a triple of the core graphs $\Gamma_{H^g}, \Gamma_{K^g}, \Gamma_{H^g \cap K^g}$ for the subgroups $H^g, K^g, H^g \cap K^g$ with no vertices of valence 1. See Figure 5 for an illustration.

Remark 2.3. Since $\operatorname{rk} H^g = \operatorname{rk} H$, $\operatorname{rk} K^g = \operatorname{rk} K$, $\operatorname{rk} H^g \cap K^g = \operatorname{rk} H \cap K$, for the purposes of this paper we may assume (and will do so from now on) without loss of generality that the groups $H, K, H \cap K$ have the core graphs which do not have vertices of valence 1. It may happen that after the procedure described in Lemma 2.2 the basepoints of Γ_H, Γ_K and $\Gamma_{H \cap K}$ all map to the vertex $v \in X$, instead of u, but that does not limit generality, since we may have chosen vertex $v \in X$ as the basepoint of X from the very beginning.



FIGURE 5. Eliminating vertices of valence 1 (for some generic H and K).

2.4. Join of subgroups and the topological pushout. As was shown by Stallings [26], the core graph $\Gamma_{H\vee K}$ for the join of two subgroups is obtained by joining the core graphs for Γ_H and Γ_K at their respective basepoints and performing a sequence of identifications of edges with the same labels called *foldings*:

$$\Gamma_H \vee \Gamma_K \xrightarrow{\text{foldings}} \Gamma_{H \vee K}$$

In general, the number of foldings required to produce $\Gamma_{H\vee K}$ and the rank of $\Gamma_{H\vee K}$ are hard to estimate directly from the information about Γ_H , Γ_K , without actually performing the required sequence of foldings. In [22], Kent works with an intermediate object, the *topological pushout* \mathcal{T} of Γ_H and Γ_K , which fits into the diagram:

$$\Gamma_H \vee \Gamma_K \xrightarrow{\text{foldings}} \mathcal{T} \xrightarrow{\text{foldings}} \Gamma_{H \vee K}$$

and whose rank is much easier to estimate than the rank of $\Gamma_{H \vee K}$. Since the folding operation is surjective at the level of fundamental groups [26, Cor. 4.4], we also have

$$\operatorname{rk} \mathcal{T} \geq \operatorname{rk} \Gamma_{H \vee K}.$$

 \mathcal{T} is defined as follows.

Let $x \in \Gamma_H$ and $y \in \Gamma_K$ be either two vertices or two edges of Γ_H and Γ_K . The graph \mathcal{T} is the quotient of the disjoint union $\Gamma_H \sqcup \Gamma_K$ by the equivalence relation generated by the following relation: $x \sim y$ if $x \in \Pi_H((\Pi_K|_{\Gamma_{H\cap K}})^{-1}(y))$ or $y \in \Pi_K((\Pi_H|_{\Gamma_{H\cap K}})^{-1}(x))$. In other words, $x \sim y$ if and only if there is an element z of $\Gamma_{H\cap K}$ such that x and y are the images under Π_H , Π_K , respectively, of z. Recall that, by construction, the vertices of $\Gamma_{H\cap K}$ can be identified with a certain subset of $V(\Gamma_H) \times V(\Gamma_K)$ and the same is true for edges. Thus two elements (i.e. two vertices or two edges) a, b of $\Gamma_H \sqcup \Gamma_K$ map to the same element in \mathcal{T} if and only if there is a sequence of elements $(x_1, y_1), \ldots, (x_n, y_n)$ in $\Gamma_{H\cap K}$ with $x_i \in \Gamma_H$, $y_i \in \Gamma_K$ such that a is either x_1 or y_1 , b is either x_n or y_n and for each i either $x_i = x_{i+1}$ or $y_i = y_{i+1}$. Equivalently, \mathcal{T} can be obtained from the join of Γ_H and Γ_K over their respective basepoints x_H and x_K , followed by a sequence of foldings along the edges of $\Gamma_{H\cap K}$ only. I.e. we may choose a circuit (i.e. a closed path) γ in $\Gamma_{H\cap K}$ that starts at the basepoint (x_H, x_K) and traverses each edge of $\Gamma_{H\cap K}$ at least once, and perform a sequence of foldings, identifying $\Pi_H(z) \in \Gamma_H$ with $\Pi_K(z) \in \Gamma_K$ for z running consecutively through all vertices and edges along γ . Since $\Gamma_{H\cap K}$ is connected, a simple inductive argument shows that the result of this sequence of foldings is exactly the topological pushout \mathcal{T} of Γ_H and Γ_K .

Figure 4 shows the topological pushout for the groups H, K of Example 2.1. We see that the topological pushout \mathcal{T} may be different from $\Gamma_{H\vee K}$. In Figure 4, the graph $\Gamma_{H\vee K}$ is equal to X and it is obtained from \mathcal{T} by identifying (folding) two *c*-edges.

3. Proof of Theorem 1.1

For the purposes of this section it will be convenient to assume that F is a free group of countable rank so that we have a freedom to add new generators if necessary, without the need of explicitly embedding them into the free group of rank 2. Also, for that purpose, we fix the wedge of countably many circles as the base CW complex for F.

We will call a tuple of values (h, k; v, c) realizable, if there exist finitely generated subgroups H, K of F with $\operatorname{rk} H = h$, $\operatorname{rk} K = k$, $\operatorname{rk}(H \lor K) = v$, and $\operatorname{rk}(H \cap K) = c$. If the values h, k (and sometimes also v) are clear from the context, we will also call the tuple (v, c) (respectively, the number c) realizable, and say that it belongs to page (h, k).

Excluding trivial cases, we may assume that $\operatorname{rk} H \geq 2$ and $\operatorname{rk} K \geq 2$, so that $\operatorname{rk}(H \vee K) \geq 2$. The upper bound for $\operatorname{rk}(H \vee K)$ is obviously h + k. On the other hand, the limits for $\operatorname{rk}(H \cap K)$ are 0 and (h-1)(k-1) + 1, as is stipulated by the Friedman–Mineyev theorem.

It turns out that the set of all known realizable values (h, k; v, c) can be described for any fixed (h, k) by a finite sequence of nonnegative integers (a_i) , such that for any given h, k, v all (known) realizable values of c are described as the range $0 \le c \le a_i$, where i = h + k - v:

Theorem 1.1. Let F be a free group and let integers h, k, c, v satisfy $2 \le h \le k$, $2 \le v \le h + k$, $0 \le c \le (h-1)(k-1) + 1$. Define a sequence (a_i) as follows:

$$a_0 = 0;$$

 $a_i = \left\lfloor \frac{i^2}{4} \right\rfloor + 1, \quad for \quad i = 1, \dots, 2(h-1);$
 $a_i = (h-1)(i-h+1) + 1, \quad for \quad i = 2(h-1), \dots, h+k-2.$

If we denote i = h + k - v, then for any $c \leq a_i$ there exist subgroups $H, K \leq F$ such that $\operatorname{rk}(H) = h$, $\operatorname{rk}(K) = k$, $\operatorname{rk}(H \cap K) = c$, and $\operatorname{rk}(H \vee K) = v$.

Example 3.1. The diagram in Figure 6 shows the realizable values from Theorem 1.1 for h = rk H = 5, k = rk K = 7. They correspond to the sequence

$$(a_i) = (0, 1, 2, 3, 5, 7, 10, 13, 17, 21, 25).$$

We see that the sequence (a_i) from the Theorem 1.1 is a union of a discrete quadratic function and a linear function, with the linear part present only when h < k. The value $a_0 = 0$ reflects the fact that if $\operatorname{rk}(H \lor K) = \operatorname{rk} H + \operatorname{rk} K$ then $\operatorname{rk}(H \cap K) = 0$. (This is a consequence of the property of finitely generated free groups being Hopfian.) On the other hand, the value v = 2 corresponds to i = h + k - 2, and a_{h+k-2} equals (h-1)(k-1)+1. This reflects the fact that all possible values for $\operatorname{rk}(H \cap K) = 0, \ldots, (h-1)(k-1)+1$ are realizable when $\operatorname{rk}(H \lor K) = 2$, as was shown by Kent in [21].

Proof of Theorem 1.1. In what follows, we will call a finite sequence (a_i) , (i = 0, ..., n), greater than a sequence (b_i) , (i = 0, ..., n), if for each *i*, we have $a_i \ge b_i$. In this case we also say that the sequence (b_i) is smaller than the sequence (a_i) .

We will obtain the required set of realizable values for page (h', k') inductively from the realizable values for page (h, k) with $h \leq h', k \leq k'$, by using the following operations:

Ia. Adding a new generator to H. This operation copies all realizable values from page (h, k) to page (h + 1, k) as follows:

$$(h, k; v, c) \longmapsto (h+1, k; v+1, c)$$

IGNAT SOROKO



FIGURE 6. The realizable values for (h, k) = (5, 7) from Theorem 1.1.



FIGURE 7. The realizable values for (h, k) = (2, 2).

Ib. Adding a new generator to K. This operation copies all realizable values from page (h, k) to page (h, k + 1) as follows:

$$(h,k;v,c) \longmapsto (h,k+1;v+1,c)$$

II. Adding the same new generator to both H and K. We get:

$$(h, k; v, c) \longmapsto (h+1, k+1; v+1, c+1)$$

III. Populating the first row of any page (h, k) with values

 $v = 2, \quad c = 0, \dots, (h-1)(k-1) + 1$

corresponding to the explicit examples produced by Kent [21].

To prove the claimed effect on ranks under operations Ia, Ib and II, we notice that adding a new generator to a subgroup H amounts to attaching the loop corresponding to this generator to the core graph Γ_H , at its basepoint. (We can do that in view of the assumption in the opening paragraph of the current section.) This makes the effect of operations Ia and Ib obvious, while for operation II we recall the construction of the core graph for $H \cap K$ from Section 2. It is clear that if we attach a loop labeled with the same new generator to both core graphs Γ_H , Γ_K at their respective basepoints, then the core graph of their intersection $\Gamma_{H \cap K}$ also gets the loop labeled with the same generator attached to its basepoint.

We start with page (h, k) = (2, 2), which corresponds to the sequence $(a_n) = (0, 1, 2)$, (as was shown in [22, p. 307], see Figure 7), and determine a sequence of operations Ia, Ib, II, III that leads to the greatest sequence (a_n) , for the required values (h, k).

We first show that if h < k then applying (Ia + III) followed by (Ib + III) produces a greater sequence (a_n) of realizable *c*-values for page (h + 1, k + 1) than applying first (Ib + III) and then (Ia + III). Indeed, let (a_n) be the sequence for (h, k) with *n* ranging from 0 to h + k - 2. Denote (a'_n) the result of applying (Ia + III) to (a_n) and (a''_n) the result of applying (Ib + III) to (a'_n) . Similarly, denote (b'_n) the result of applying (Ib + III) to (a_n) and (b''_n) the result of applying (Ia + III) to (b'_n) :

$$\begin{array}{c} (a_n) \xrightarrow{\text{Ia}+\text{III}} (a'_n) \xrightarrow{\text{Ib}+\text{III}} (a''_n) \\ (a_n) \xrightarrow{\text{Ib}+\text{III}} (b'_n) \xrightarrow{\text{Ia}+\text{III}} (b''_n) \end{array}$$



FIGURE 8. The effect of operations Ia and II followed by III. The horizontal axes of the tables correspond to the values of $c = \operatorname{rk}(H \cap K)$, the vertical axes to the values of $v = \operatorname{rk}(H \vee K)$. Operations Ia and II are shown in dark gray, operation III in light gray. Operation Ib has the same effect as Ia, but for the target page (h, k + 1).

We want to show that $a''_n \ge b''_n$ for all n. Since operation Ia copies all values of (a_i) to the new page, we have: $a'_i = a_i$ if i < (h + 1) + k - 2, and since operation III populates the first row (corresponding to the last value of a_i) with a sequence of h(k-1) + 1 values, we get $a'_{(h+1)+k-2} = h(k-1) + 1$. Similarly, $a''_i = a_i$ for i < (h + 1) + (k + 1) - 3, $a''_{(h+1)+(k+1)-3} = h(k-1) + 1$, and $a''_{(h+1)+(k+1)-2} = hk + 1$. Performing these operations in the opposite order, i.e. applying (Ib + III) to (a_n) first followed by (Ia + III), we get in a similar fashion: $b''_i = a_i$ for i < (h+1) + (k+1) - 3, $b''_{(h+1)+(k+1)-3} = hk - k + 1$, and $b''_{(h+1)+(k+1)-2} = hk + 1$. The sequences (a''_n) and (b''_n) agree for all values of n except the penultimate one, n = (h+1) + (k+1) - 3. Comparing them and taking into account that h < k, we see that hk - h + 1 > hk - k + 1, i.e. $a''_n \ge b''_n$ for all n. This means that if h < k, applying (Ia + III) followed by (Ib + III) produces a greater sequence (a_n) than if applying these operations in the opposite order.

The above formulas also show that in the case when h = k the result of applying operations (Ia + III) and (Ib + III) does not depend on their order.

Now let's examine operation (II + III). It copies realizable values from page (h, k) to page (h + 1, k + 1)as shown in the lower part of Figure 8. Let (a_n) and (a'''_n) be the corresponding sequences of realizable values for pages (h, k) and (h + 1, k + 1), respectively, and let (a''_n) be the sequence obtained from (a_n) by the composition of operations (Ia + III) and (Ib + III), as before. We claim that (a'''_n) is smaller than (a''_n) . Indeed, we saw in the previous paragraph that $a''_i = a_i$ for i < h + k - 1, $a''_{h+k-1} = hk - h + 1$ and $a''_{h+k} = hk + 1$. From Figure 8 we see that $a''_i = a_{i-1} + 1$ for $i = 1, \ldots, h + k - 1$ and $a''_{h+k} = hk + 1$. Thus to prove that $a''_i \ge a''_i$ for all i, we claim the following:

- (i) $a_i \ge a_{i-1} + 1$ for $i = 1, \dots, h + k 2$, and
- (ii) $hk h + 1 \ge a_{h+k-2} + 1$.

We will prove these using the following fact: for any page (h, k) the last value of the sequence (a_n) , i.e. the term a_{h+k-2} , equals (h-1)(k-1)+1, which is the content of operation III. (Recall that operation III is applied every time we apply Ia, Ib or II.) Thus, inequality (ii) is established: $a_{h+k-2} = (h-1)(k-1)+1 \leq hk-h$ since $k \geq 2$. To prove (i) we observe that all operations Ia, Ib, II preserve the difference between consecutive elements $a_j - a_{j-1}$, so all that needs to be proved is that the top value $a_{h+k-2} = (h-1)(k-1)+1$ is always at least 1 bigger than the previous value of a_{h+k-3} . Assuming that (i) holds true for page (h, k), we examine how it changes under the application of operations (Ia + III), (Ib + III) and (II + III). In the first case, operation Ia makes the next-to-last value of the sequence (a'_n) to be (h-1)(k-1)+1, while the last one is h(k-1)+1, with the difference k-1 between the two. In the second case, arguing in a similar fashion, we

(2,2)	(2,3)	(2,4)	(2,5)	(2,6)	(2,7)	(2,8)	(2,9)	(2,10)
(3,2)	(3,3)	(3,4)	(3,5)	(3,6)	(3,7)	(3,8)	(3,9)	(3,10)
(4,2)	(4,3)	(4,4)	(4,5)	(4,6)	(4,7)	(4,8)	(4,9)	(4,10)
(5,2)	(5,3)	(5,4)	(5,5)	(5,6)	(5,7)	(5,8)	(5,9)	(5,10)

FIGURE 9. The ways to reach page (h, k) = (5, 10) from page $(h_0, k_0) = (2, 2)$ are given by broken lines composed of horizontal and vertical arrows, which correspond to operations (Ib+III) and (Ia+III), respectively. The thick line corresponds to an optimal sequence of operations, giving the maximal sequence (a_n) .

get that the difference between the last two values of (b'_n) equals h - 1. And in the case of operation (II + III), we get that the difference equals (hk + 1) - ((h - 1)(k - 1) + 2) = h + k - 2. We see that in all three cases this difference is at least 1, which proves claim (i). Thus operation (II + III) creates a sequence (a''_n) which is smaller than the sequence (a''_n) created by (Ia + III) followed by (Ib + III).

The above analysis shows that to obtain the greatest sequence (a_n) of realizable values for page (h, k), $h \leq k$, one can discard operation (II + III) completely and apply only operations (Ia + III) and (Ib + III), starting with page $(h_0, k_0) = (2, 2)$. All the ways to get from page (2, 2) to page (h, k) by applying the said operations can be encoded by broken lines running in a rectangular table from entry (2, 2) to entry (h, k) with horizontal and vertical segments corresponding to operations (Ib + III) and (Ia + III), respectively, see Figure 9. Every time when operation (Ib + III) followed by (Ia + III) is applied to a page (h', k') with h' < k', we can interchange the order of these operations thus producing a bigger sequence (a_n) . By repeatedly doing this interchange, we obtain an optimal sequence which can be described as follows:

Start with $(h_0, k_0) = (2, 2)$. Alternate operations (Ib + III) and (Ia + III) to reach page (h, h). If h < k, keep applying (Ib + III) to reach page (h, k):

$$(2,2) \xrightarrow{\text{Ib}+\text{III}} (2,3) \xrightarrow{\text{Ia}+\text{III}} (3,3) \xrightarrow{\text{Ib}+\text{III}} \dots \xrightarrow{\text{Ia}+\text{III}} (h,h) \xrightarrow{\text{Ib}+\text{III}} (h,h+1) \xrightarrow{\text{Ib}+\text{III}} (h,h+2) \xrightarrow{\text{Ib}+\text{III}} \dots \xrightarrow{\text{Ib}+\text{III}} (h,k).$$

Note that $a_0 = 0, a_1 = 1, a_2 = 2$ for $(h_0, k_0) = (2, 2)$, and each of operations (Ia + III), (Ib + III) augments the existing sequence a_0, \ldots, a_n with a new value a_{n+1} , which is computed according to operation III as follows:

$$(\ell, \ell) \xrightarrow{\text{Ib+III}} (\ell, \ell+1): \qquad a_{n+1} = \ell(\ell-1) + 1, \text{ with } n = 2(\ell-1); \\ (\ell, \ell+1) \xrightarrow{\text{Ia+III}} (\ell+1, \ell+1): \qquad a_{n+1} = \ell^2 + 1, \text{ with } n = 2\ell - 1,$$

for $\ell = 2, \ldots, h - 1$, and

$$(h, h+j) \xrightarrow{\text{Ib+III}} (h, h+j+1): \quad a_{n+1} = (h-1)(h+j) + 1, \text{ with } n = 2(h-1) + j,$$

for j = 0, ..., k - h - 1, if h < k.

Now we observe that the values a_{n+1} for $0 \le n \le 2h-3$ can be written concisely as $a_{n+1} = \lfloor \left(\frac{n+1}{2}\right)^2 \rfloor + 1$. Indeed, if $n = 2(\ell - 1)$, then $\lfloor \left(\frac{n+1}{2}\right)^2 \rfloor = \lfloor (\ell - \frac{1}{2})^2 \rfloor = \lfloor \ell^2 - \ell + \frac{1}{4} \rfloor = \ell(\ell - 1)$, and if $n = 2\ell - 1$, then $\lfloor \left(\frac{n+1}{2}\right)^2 \rfloor = \ell^2$. This proves that the above sequence of operations produces the sequence a_i described in the Theorem, which finishes the proof.

Now we have counterexamples to GTC for all $m \geq 5$.

Corollary 3.2. Guzman's "Group-Theoretic Conjecture" does not hold for any $m \geq 5$.

Proof. Theorem 1.1 guarantees the existence of subgroups $H, K \leq F$ such that $h = \operatorname{rk}(H) = m, k = \operatorname{rk}(K) = m, c = \operatorname{rk}(H \cap K) = m$ and $v = \operatorname{rk}(H \vee K) = m + 1$, if $m \geq 5$. Indeed, i = h + k - v = m - 1 in this case,

which is less than 2(h-1) = 2(m-1). So the value of a_i in Theorem 1.1 is equal to $\lfloor \frac{i^2}{4} \rfloor + 1 = \lfloor \frac{(m-1)^2}{4} \rfloor + 1$, which is bigger than or equal to c = m for all $m \ge 5$. See Figure 1 for an illustration of the case m = 6, which shows the existence of rank 6 subgroups H, K with $\operatorname{rk}(H \lor K) = 7$ and $\operatorname{rk}(H \cap K) = 6, 7$.

4. The structure of the topological pushout

In this section we study the combinatorial structure of the topological pushout using graphs introduced by Dicks in [6]. This will allow us to obtain an upper bound on the rank of the topological pushout of two core graphs and hence on the rank of the join of the corresponding subgroups.

4.1. The Dicks graphs. Let Γ_H , Γ_K be the core graphs for subgroups $H, K \leq F$ and the core graph $\Gamma_{H \cap K}$ is constructed as the pullback of graph immersions, as in Section 2:

$$\begin{array}{ccc} \Gamma_{H\cap K} & \stackrel{\Pi_K}{\longrightarrow} & \Gamma_K \\ & & & \downarrow^{\pi_H} & & \downarrow^{p_K} \\ & & \Gamma_H & \stackrel{p_H}{\longrightarrow} & X \end{array}$$

Each element z (a vertex or an edge) of Γ_H and Γ_K inherits its type from the mapping to X, that is, an element of $V(X) \sqcup E(X) = \{u, v, a, b, c\}$ to which z maps.

We are now going to define five bipartite undirected graphs $\Omega_u, \Omega_v, \Omega_a, \Omega_b, \Omega_c$, adapting the construction of [6] to our needs.

First, we define Ω_u as follows.

$$V(\Omega_u) = \{ z \in V(\Gamma_H) \mid p_H(z) = u \} \sqcup \{ z' \in V(\Gamma_K) \mid p_K(z') = u \},\$$

$$E(\Omega_u) = \{ (z, z') \in V(\Gamma_{H \cap K}) \mid p_H(z) = u \text{ and } p_K(z') = u \}.$$

In other words, two vertices $z \in V(\Gamma_H)$, $z' \in V(\Gamma_K)$ from $V(\Omega_u)$ are connected with a single undirected edge if the vertex (z, z') of $G_{H \cap K}$ actually belongs to $\Gamma_{H \cap K}$.

The graph Ω_v is defined analogously to Ω_u with the obvious modification $(u \rightsquigarrow v)$. Denote also $\Omega = \Omega_u \sqcup \Omega_v$.

We define Ω_a similarly, by dealing with edges instead of vertices:

$$V(\Omega_a) = \{ e \in E(\Gamma_H) \mid p_H(e) = a \} \sqcup \{ e' \in E(\Gamma_K) \mid p_K(e') = a \},\$$

$$E(\Omega_a) = \{ (e, e') \in E(\Gamma_{H \cap K}) \mid p_H(e) = a \text{ and } p_K(e') = a \}$$

The graphs Ω_b , Ω_c are defined analogously with the obvious modifications.

The bipartite structure on the defined graphs is given by grouping all vertices/edges of graph Γ_H in one part, and those of Γ_K in the other.

Figure 10 shows the graphs $\Omega_u, \Omega_v, \Omega_a, \Omega_b, \Omega_c$ for the core graphs of the subgroups H, K from Example 2.1. Notice that the operations of taking the origin and the terminus of an edge induce embeddings \tilde{o}, \tilde{t} of the graph $\Omega_a \sqcup \Omega_b \sqcup \Omega_c$ into Ω_u, Ω_v , respectively. Let's show that $\tilde{o}|_{\Omega_a} : \Omega_a \to \Omega_u$ is an embedding. If $e, e' \in V(\Omega_a)$, with $e \neq e'$, then they correspond to a-edges of $\Gamma_H \sqcup \Gamma_K$. If $e \in E(\Gamma_H)$, $e' \in E(\Gamma_K)$, their origins are different. If they both belong to the same graph then their origins are also different, since Γ_H and Γ_K are folded, i.e. at every vertex of Γ_H, Γ_K there is at most one a-edge having this vertex as the origin. This proves that the map $\tilde{o}|_{\Omega_a} : V(\Omega_a) \to V(\Omega_u)$ is injective. Now, if $e, e' \in V(\Omega_a)$ are connected with an edge, this means that there is an a-edge (e, e') in $E(\Gamma_{H \cap K})$ whose projections under Π_H, Π_K are e, e'. In particular, the origin of (e, e') projects to the origins of e, e', and we see that the images of e, e' in $V(\Omega_u)$ are connected with an edge as well. This proves that $\tilde{o}|_{\Omega_a} : \Omega_a \to \Omega_u$ is an injective graph homomorphism, i.e. is an isomorphism onto its image. Denote

$$\Omega_{u,a} = \operatorname{im}(\Omega_a \stackrel{\tilde{o}}{\longrightarrow} \Omega_u), \qquad \Omega_{v,a} = \operatorname{im}(\Omega_a \stackrel{\tilde{t}}{\longrightarrow} \Omega_v),
\Omega_{u,b} = \operatorname{im}(\Omega_b \stackrel{\tilde{o}}{\longrightarrow} \Omega_u), \qquad \Omega_{v,b} = \operatorname{im}(\Omega_b \stackrel{\tilde{t}}{\longrightarrow} \Omega_v),
\Omega_{u,c} = \operatorname{im}(\Omega_c \stackrel{\tilde{o}}{\longrightarrow} \Omega_u), \qquad \Omega_{v,c} = \operatorname{im}(\Omega_c \stackrel{\tilde{t}}{\longrightarrow} \Omega_v)$$
(3)



FIGURE 10. The Dicks graphs $\Omega_u, \Omega_v, \Omega_a, \Omega_b, \Omega_c$ for the subgroups in Example 2.1. A pair (x, y) denotes the unique a-, b-, or c-edge of $E(\Gamma_H) \sqcup E(\Gamma_K)$ originating at a vertex x and terminating at a vertex y.

the images of these embeddings, and set

$$\begin{array}{ll} \Omega_{u,ab} = \Omega_{u,a} \cap \Omega_{u,b}, & \Omega_{v,ab} = \Omega_{v,a} \cap \Omega_{v,b}, & \Omega_{ab} = \Omega_{u,ab} \sqcup \Omega_{v,ab}, \\ \Omega_{u,bc} = \Omega_{u,b} \cap \Omega_{u,c}, & \Omega_{v,bc} = \Omega_{v,b} \cap \Omega_{v,c}, & \Omega_{bc} = \Omega_{u,bc} \sqcup \Omega_{v,bc}, \\ \Omega_{u,ac} = \Omega_{u,a} \cap \Omega_{u,c}, & \Omega_{v,ac} = \Omega_{v,a} \cap \Omega_{v,c}, & \Omega_{ac} = \Omega_{u,ac} \sqcup \Omega_{v,ac}, \\ \Omega_{u,abc} = \Omega_{u,a} \cap \Omega_{u,b} \cap \Omega_{u,c}, & \Omega_{v,abc} = \Omega_{v,a} \cap \Omega_{v,b} \cap \Omega_{v,c}, & \Omega_{abc} = \Omega_{u,abc} \sqcup \Omega_{v,abc}. \end{array}$$

Since by Remark 2.3 we assume that neither of graphs Γ_H , Γ_K , $\Gamma_{H \cap K}$ can have a vertex of valence 1, we observe that each vertex in $\Omega_{u,a}$ is also the origin of either another *b*-edge, or *c*-edge, or both. Thus,

$$\Omega_{u,a} = \Omega_{u,ab} \bigvee_{\Omega_{u,abc}} \Omega_{u,ac},$$

where $K = L \bigvee_N M$ means that $K = L \cup M$ and $L \cap M = N$. Similarly, each vertex in $\Omega_{v,a}$ is also the terminus of either another *b*-edge, or *c*-edge, or both. Thus,

$$\Omega_{v,a} = \Omega_{v,ab} \bigvee_{\Omega_{v,abc}} \Omega_{v,ac},$$

Of course, a completely similar statement holds for all other graphs (3) in place of $\Omega_{u,a}$, $\Omega_{v,a}$.

Clearly, $\Omega_{ab} \cap \Omega_{bc} = \Omega_{bc} \cap \Omega_{ab} = \Omega_{ac} \cap \Omega_{ab} = \Omega_{abc}$. Thus we can denote

$$A = \Omega_{ab} \bigvee_{\Omega_{abc}} \Omega_{ac}, \qquad B = \Omega_{ab} \bigvee_{\Omega_{abc}} \Omega_{bc}, \qquad C = \Omega_{ac} \bigvee_{\Omega_{abc}} \Omega_{bc}.$$

Notice that, by construction, all graphs A, B, C are subgraphs of $\Omega = \Omega_u \sqcup \Omega_v$.

In what follows we will depict connected components of Ω against the following 'trefoil' Venn diagram, illustrating the relation

$$\Omega_{ab} \cap \Omega_{bc} = \Omega_{ab} \cap \Omega_{ac} = \Omega_{bc} \cap \Omega_{ac} = \Omega_{abc},$$

see Figure 11.

Now, for A, we have:

$$A = \Omega_{ab} \cup \Omega_{ac} = (\Omega_{u,ab} \sqcup \Omega_{v,ab}) \cup (\Omega_{u,ac} \sqcup \Omega_{v,ac}) = (\Omega_{u,ab} \cup \Omega_{u,ac}) \sqcup (\Omega_{v,ab} \cup \Omega_{v,ac}) = \Omega_{u,a} \sqcup \Omega_{v,a} \cong \Omega_a \sqcup \Omega_a,$$
(4)

and similarly for B, C.

Thus we established the following duality, discovered by Dicks [6]:

Proposition 4.1 (Dicks' duality). Each of the graphs A, B, C defined above consists of an even number of connected components, which are isomorphic in pairs. If $\{Z, Z'\}$ is such a pair of components of A, then



FIGURE 11. Venn diagram for the Dicks graphs.

one of Z, Z' is a component of $\Omega_{u,a}$ and the other is a component of $\Omega_{v,a}$, and the isomorphism between them preserves the bipartite structure. Similar statements are true for B and C in place of A. \square

4.2. Modeling the topological pushout on the Dicks graphs. Notice that the Dicks graphs Ω_u , Ω_v , $\Omega_a, \Omega_b, \Omega_c$ express the equivalence relation ~ used to define the topological pushout in subsection 2.4: two vertices z, z' are connected by an edge in the graph Ω_x (where x stands for any of u, v, a, b, c) if and only if they are the two projections under Π_H , Π_K of the same element of $\Gamma_{H\cap K}$. Thus, the connected components of $\Omega_u \sqcup \Omega_v$ are exactly the vertices of the topological pushout, and the connected components of $\Omega_a \sqcup \Omega_b \sqcup \Omega_c$ are the edges of the topological pushout, with inclusions \tilde{o}, \tilde{t} defined above being the origin and the terminus maps. In particular, we see from Figure 10 that the topological pushout \mathcal{T} from Example 2.1 (see Figure 4) has exactly two vertices corresponding to the connected graphs Ω_u and Ω_v and four directed edges: one a-edge for the connected graph Ω_a , one b-edge for the connected graph Ω_b and two c-edges for the two connected components of Ω_c .

It will be useful for us to recast the topological pushout in terms of graphs A, B, C defined above:

Proposition 4.2. The topological pushout \mathcal{T} admits the following description:

Vertices: connected components of $\Omega = \Omega_u \sqcup \Omega_v = \Omega_{ab} \bigvee_{\Omega_{abc}} \Omega_{bc} \bigvee_{\Omega_{abc}} \Omega_{ac} = A \cup B \cup C$. **Edges:** pairs of connected components of A, B, C from the pairing in Proposition 4.1.

If $e = \{Z, Z'\}$ is such a pair, viewed as a directed edge, it inherits its type from the corresponding graph it belongs to: if $e \subset A$, then e is an a-edge, if $e \subset B$, then e is a b-edge, and if $e \subset C$ then e is a c-edge. The origin and terminus maps are defined as follows. If $e \subset A$, then one of Z, Z' is a connected component of $\Omega_{u,a}$, and the other, of $\Omega_{v,a}$. Let $Z \subset \Omega_{u,a}$, $Z' \subset \Omega_{v,a}$, say. Then the origin of e is the connected component of $(A \cup B \cup C) \cap \Omega_u$ in which Z lies, and the terminus of e is the connected component of $(A \cup B \cup C) \cap \Omega_v$ in which Z' lies. Similar definitions apply to B and C in place of A.

Proof. By the definition of \mathcal{T} , its vertices are the connected components of $\Omega = \Omega_u \sqcup \Omega_v$. Since every vertex of Γ_H , Γ_K is incident to either an *a*-edge, a *b*-edge, or a *c*-edge, the same is true for \mathcal{T} . Hence $\Omega_u = \Omega_{u,a} \cup \Omega_{u,b} \cup \Omega_{u,c} \text{ and } \Omega_v = \Omega_{v,a} \cup \Omega_{v,b} \cup \Omega_{v,c}. \text{ From (4) we see that } \Omega_{u,a} \sqcup \Omega_{v,a} = A, \Omega_{u,b} \sqcup \Omega_{v,b} = B,$ and $\Omega_{u,c} \sqcup \Omega_{v,c} = C$. It follows that $\Omega_u \sqcup \Omega_v = A \cup B \cup C$.

The edges of \mathcal{T} , by definition, are the connected components of $\Omega_a \sqcup \Omega_b \sqcup \Omega_c$, with the attaching maps \tilde{o} , t, defined above. If Z_0 is a connected component of Ω_a , say, it defines two isomorphic connected subgraphs $Z = \tilde{o}(Z_0) \subset \Omega_{u,a}$ and $Z' = \tilde{t}(Z_0) \subset \Omega_{v,a}$. Since $\Omega_{u,a}$ and $\Omega_{v,a}$ are, by definition, isomorphic copies of Ω_a , the subgraphs Z, Z' are the whole connected components of $\Omega_{u,a}, \Omega_{v,a}$, respectively. By (4), $A = \Omega_{u,a} \sqcup \Omega_{v,a}$, therefore the pair $\{Z, Z'\}$ is a pair of connected components of A determined by Z_0 . The attaching maps for $\{Z, Z'\}$ described in the Proposition are induced by \tilde{o} and \tilde{t} applied to Z_0 . \square

Figure 12 shows the topological pushout for the subgroups of Example 2.1, modeled by subsets Ω_{ab} , $\Omega_{bc}, \Omega_{ac}, \Omega_{abc}$, in accordance with Proposition 4.2. Notice that the connected components of $\Omega_a, \Omega_b, \Omega_c$ establish bijections between parts A, B, C of Ω_u (the left 'trefoil') and the corresponding parts of Ω_v (the right 'trefoil'). In particular, the connected graph Ω_a from Figure 10 acts as the *a*-edge of \mathcal{T} and establishes a bijection between $\tilde{o}(\Omega_a) = \{1-7-3\}$ with $\tilde{t}(\Omega_a) = \{2-6-4\}$. Also, the connected graph Ω_b from Figure 10 is the b-edge of \mathcal{T} and it establishes a bijection between $\tilde{o}(\Omega_b) = \{5-3-7\}$ and $\tilde{t}(\Omega_b) = \{8-2-6\}$. Finally, each connected component of Ω_c from Figure 10 serves as a *c*-edge of \mathcal{T} and they establish bijections of the two connected components 1–7, 3–5 of $\Omega_{u,c}$ with 2–8, 4–6 of $\Omega_{v,c}$. Notice also that this bijection does not preserve subsets Ω_{ac} and Ω_{bc} individually but leaves invariant their union $C = \Omega_{ac} \bigvee_{\Omega_{abc}} \Omega_{bc}$: the subgraph 1–7 of $\Omega_{u,ac}$ gets paired with the subgraph 2–8 of $\Omega_{v,bc}$, and also 3–5 of $\Omega_{u,bc}$ gets paired with 4–6 of $\Omega_{v,ac}$. The curved edges 3–7 and 2–6 reflect the fact that vertices 3 and 7 (and also 2 and 6) are adjacent in Ω_{ab} , but not in Ω_{abc} .



FIGURE 12. The topological pushout \mathcal{T} from Figure 4 modeled on the Dicks graphs. Vertices 3 and 7 (and also 2 and 6) are adjacent in Ω_{ab} , but not in Ω_{abc} , which is depicted by a curved edge lying in the 'petal' for Ω_{ab} .

Now we are ready to relate ranks of H, K, $H \cap K$ and \mathcal{T} with the structure of the Dicks graph $\Omega = \Omega_{ab} \bigvee_{\Omega_{abc}} \Omega_{bc} \bigvee_{\Omega_{abc}} \Omega_{ac}$. We are going to prove the following Theorem 4.3 assuming the validity of Proposition 5.1, which will be proved in Section 5. Recall that a *cycle* in an undirected graph is a path $x_0e_0x_1e_1\ldots x_ke_kx_0$ such that all vertices x_i are pairwise different. We denote by #W the cardinality of a finite set W and, as before, by r(H) the reduced rank of group H, i.e. the quantity $\max(0, \operatorname{rk}(H) - 1)$.

Theorem 4.3. Let F be a free group and $H, K \leq F$ be finitely generated subgroups. Let Γ_H, Γ_K and $\Gamma_{H\cap K}$ be the core graphs of the corresponding subgroups, and let \mathcal{T} be their topological pushout. Let also Ω and Ω_{abc} be the Dicks graphs defined above. Then

- (1) Ω_{abc} is a bipartite graph with $2 \operatorname{rr}(H)$ vertices in one part and $2 \operatorname{rr}(K)$ vertices in the other;
- (2) Ω_{abc} has $2 \operatorname{rr}(H \cap K)$ edges;
- (3) (# connected components of Ω_{abc}) $\geq 2 \operatorname{rr}(\mathcal{T})$, with the equality taking place if and only if every cycle of Ω lies entirely in one of the subgraphs Ω_{ab} , Ω_{bc} , Ω_{ac} (with different cycles possibly lying in different subgraphs).

Proof. The graph Ω_{abc} has as its vertices all vertices of valence 3 of Γ_H and Γ_K . Since graphs Γ_H , Γ_K , $\Gamma_{H\cap K}$ are normalized as in Remark 2.3, all their vertices have valence either 2 or 3. Computing Euler characteristic of Γ (where Γ stands for any of Γ_H , Γ_K , $\Gamma_{H\cap K}$) gives:

$$1 - \operatorname{rk}(\Gamma) = \#V(\Gamma) - \#(E^+(\Gamma)) = \sum_{v \in V(\Gamma)} \left(1 - \frac{\operatorname{val}(v)}{2}\right),$$

which is equivalent to

$$2\operatorname{rr}(\Gamma) = \sum_{v \in V(\Gamma)} (\operatorname{val}(v) - 2) = (\# \text{ vertices of valence 3 in } \Gamma),$$

the last equality being true since vertices of valence 2 contribute 0 to the sum. This proves part (1).

Edges of Ω_{abc} are exactly the vertices of valence 3 in $\Gamma_{H\cap K}$, hence the computation above establishes part (2) as well.

For part (3), exactly as above, we have

$$2\operatorname{rr}(\mathcal{T}) = \sum_{v \in V(\mathcal{T})} (\operatorname{val}(v) - 2).$$
(5)

According to Proposition 4.2, the valence of a vertex v of \mathcal{T} (represented by some connected component D of $A \cup B \cup C$) is the sum of numbers of connected components of $A \cap D$, $B \cap D$ and $C \cap D$. Our goal is to understand the relationship between components of Ω_{abc} , components of $A \cup B \cup C$, and components of A, B and C, taken separately. The difficulty lies in an observation that two components P, Q of Ω_{abc} may be connected by a path outside Ω_{abc} , i.e. by a path all edges of which lie in one of the graphs $\Omega_{ab} \setminus \Omega_{abc}$, $\Omega_{bc} \setminus \Omega_{abc}$, $\Omega_{ac} \setminus \Omega_{abc}$. If this is the case, then P and Q actually correspond to the same connected component D of $A \cup B \cup C$, i.e. to the same vertex of \mathcal{T} , and their contribution to the valence of \mathcal{T} may be different from the value $2 \cdot 3$ expected otherwise. A careful treatment of this situation is given in Section 5, where we take an abstract approach and study a certain class C_n of graphs Γ with a function Σ associated to them, which encode the connectedness of components of Ω_{abc} to each other through the three graphs Ω_{ab} , Ω_{ac} , Ω_{ac} , and their joint contribution to the right-hand side of (5). To get the input for the main result of Section 5, Proposition 5.1, we form the following undirected graph Γ , which we will call the *component connectivity graph* (CCG) of Ω_{abc} .

Vertices of Γ are connected components of Ω_{abc} .

If $p, q \in V(\Gamma)$, they are some components P, Q of Ω_{abc} . If there exists a path between P and Q which does not contain any vertices of Ω_{abc} (except the first vertex of the path and the last one), and all edges of which lie in $\Omega_{ab} \setminus \Omega_{abc}$, we connect vertices p, q in $E(\Gamma)$ with an undirected edge and assign the color magenta to it. Similarly, if a Ω_{abc} -avoidant path between P and Q lies in $\Omega_{ac} \setminus \Omega_{abc}$, we add an edge to $E(\Gamma)$ connecting p and q and assign the color yellow to it. Lastly, if such path lies in $\Omega_{bc} \setminus \Omega_{abc}$, we add an edge to $E(\Gamma)$ connecting p and q and assign the color cyan to it. Thus every two vertices of Γ may be connected by up to three undirected edges, each having a different color. (The choice of names for the colors is suggested by mixing the basic colors red, blue and green, which we used to depict a-edges, b-edges and c-edges, respectively. Hence, edges from $\Omega_{ab} \setminus \Omega_{abc}$ get color red-blue, i.e. magenta, edges from $\Omega_{ac} \setminus \Omega_{abc}$ get color red-green, i.e. yellow, and edges from $\Omega_{bc} \setminus \Omega_{abc}$ get color blue-green, i.e. cyan.)

Thus, the graph Γ encodes the connectedness information (within $A \cup B \cup C$) between different connected components of Ω_{abc} . The contribution of vertices of \mathcal{T} to the sum in (5) (i.e. the right-hand side of (5)) is equal to the function $\Sigma(\Gamma)$, defined in equation (6) of Section 5.

Having formed the input for Proposition 5.1, we can use its conclusion, which reads: $\Sigma(\Gamma) \leq n$. Here *n* is the number of vertices of Γ , i.e. the number of connected components of Ω_{abc} , and $\Sigma(\Gamma)$ is the right-hand side of (5). This proves the inequality in part (3) of the Theorem. Proposition 5.1 also specifies when we have the equality in $\Sigma(\Gamma) \leq n$: this happens if and only if all cycles of Γ are *monochromatic* in the terminology of Section 5. In terms of the Dicks graphs, this means that every cycle of $\Omega_{ab} \bigvee_{\Omega_{abc}} \Omega_{bc} \bigvee_{\Omega_{abc}} \Omega_{ac}$ lies entirely in either Ω_{ab} , or Ω_{bc} , or Ω_{ac} . This finishes the proof of part (3) of the Theorem.

Situations when the inequality in part (3) of the Theorem 4.3 is strict appear quite frequently, see for example Figure 20 in Section 6.

5. CLASS \mathcal{C}_n

The goal of this technical section is to prove Proposition 5.1 needed for the proof of part (3) of Theorem 4.3. Consider a class C_n of pairs (Γ, c) where Γ is an undirected graph with multiple edges allowed (but not loops), and $c: E(\Gamma) \to \{magenta, yellow, cyan\}$ is an edge-coloring map, with the following properties:

- (1) Each graph Γ from \mathcal{C}_n has exactly *n* vertices.
- (2) Let $E(\Gamma, p, q) = E(\Gamma, q, p)$ denote the set of all undirected edges between two different vertices $p, q \in V(\Gamma)$. Then the edge-coloring map c is injective on each set $E(\Gamma, p, q)$.

In other words, any two different vertices p, q of Γ may be joined by up to three undirected edges of different colors from the set {magenta, yellow, cyan}.

For any $\Gamma \in \mathcal{C}_n$ we define three subgraphs Γ_{my} , Γ_{yc} , Γ_{mc} of Γ as follows:

• $V(\Gamma_{my}) = V(\Gamma_{yc}) = V(\Gamma_{mc}) = V(\Gamma);$

- $E(\Gamma_{my}) = \{ \text{all magenta and yellow edges of } \Gamma \};$
- $E(\Gamma_{yc}) = \{ \text{all yellow and } cyan \text{ edges of } \Gamma \};$
- $E(\Gamma_{mc}) = \{ \text{all magenta and cyan edges of } \Gamma \};$



FIGURE 13. Venn diagram for graphs Γ_{my} , Γ_{yc} , Γ_{mc} .

We define a function $\Sigma \colon \mathcal{C}_n \to \mathbb{Z}_{>0}$ as follows:

$$\Sigma(\Gamma) = \sum_{C \in CC(\Gamma)} \left(\operatorname{val}_{my}(C) + \operatorname{val}_{yc}(C) + \operatorname{val}_{mc}(C) - 2 \right), \tag{6}$$

where $CC(\Gamma)$ denotes the set of all connected components of Γ , and

 $\operatorname{val}_{colors}(C) = \# \text{ connected components of } (C \cap \Gamma_{colors})$

for $colors \in \{my, yc, mc\}$.

Recall that a cycle in an undirected graph Γ is a sequence $x_0e_0x_1e_1\ldots x_ke_kx_0$ of pairwise different vertices x_i and edges $e_i \in E(\Gamma, x_i, x_{i+1}), e_k \in E(\Gamma, x_k, x_0)$. A cycle is called *monochromatic* if all its edges are of the same color (from the set {magenta, yellow, cyan}).

Proposition 5.1. For any $\Gamma \in C_n$, we have: $\Sigma(\Gamma) \leq n$, with the equality taking place if and only if all cycles of Γ are monochromatic (with different cycles possibly having different colors).

Proof. Our first observation is:

If $\Gamma \in \mathcal{C}_n$ is edgeless then $\Sigma(\Gamma) = n$.

Indeed, in that case Γ has *n* connected components which are singleton vertices and $\Gamma_{my} = \Gamma_{yc} = \Gamma_{mc} = \Gamma$, so $\Sigma(\Gamma) = n \cdot (1 + 1 + 1 - 2) = n$.

We are going to prove that adding an edge to an arbitrary graph $\Gamma \in C_n$ may only decrease Σ , and we identify all cases when the decrease does not happen.

Choose two vertices $p, q \in V(\Gamma)$ such that $|E(\Gamma, p, q)| < 3$ and consider

$$\Gamma' = \Gamma \cup e,$$

where e is a new edge between p and q of a color that is not present in $E(\Gamma, p, q)$. Without loss of generality, we may assume that the color of edge e is magenta.

Let [p], [q] denote the connected components of Γ containing p, q, respectively, and let $[p]_{colors}, [q]_{colors}$ be the connected components of Γ_{colors} containing p, q, respectively, for $colors \in \{my, yc, mc\}$.

Also, let [p]', [q]' denote the connected components of Γ' containing p, q, respectively, and let $[p]'_{colors}$, $[q]'_{colors}$ be the connected components of Γ'_{colors} containing p, q, respectively, for $colors \in \{my, yc, mc\}$. Let also

$$\operatorname{val}_{colors}'(C') = \# \text{ connected components of } (C' \cap \Gamma_{colors}')$$

for C' a connected component of Γ' and $colors \in \{my, yc, mc\}$.

We look at several cases.

Case I: $[p] \neq [q]$. Then $[p]_{colors} \neq [q]_{colors}$ for any $colors \in \{my, yc, mc\}$, and adding magenta edge e to p, q makes $[p]' = [q]', [p]'_{my} = [q]'_{my}$ and $[p]'_{mc} = [q]'_{mc}$ while $[p]'_{yc} \neq [q]'_{yc}$ (since Γ_{yc} by definition has only

yellow and cyan edges, so if $[p]'_{yc} = [q]'_{yc}$ then $[p]_{yc} = [q]_{yc}$). Thus

$$\Sigma(\Gamma) = \left(\operatorname{val}_{my}[p] + \operatorname{val}_{yc}[p] + \operatorname{val}_{mc}[p] - 2\right) + \left(\operatorname{val}_{my}[q] + \operatorname{val}_{yc}[q] + \operatorname{val}_{mc}[q] - 2\right) + \sum_{\substack{C \neq [p], [q] \\ C \in CC(\Gamma)}} (\dots),$$

and

$$\Sigma(\Gamma') = \left(\operatorname{val}'_{my}[p]' + \operatorname{val}'_{yc}[p]' + \operatorname{val}'_{mc}[p]' - 2 \right) + \sum_{\substack{C' \neq [p]'\\C' \in CC(\Gamma')}} (\dots),$$

Now,

 $\operatorname{val}'_{my}[p]' = (\# \text{ components of } [p]' \cap \Gamma'_{my}) =$ (# components of $[p] \cap \Gamma_{my}$) + (# components of $[q] \cap \Gamma_{my}$) - 1 =

 $\operatorname{val}_{my}[p] + \operatorname{val}_{my}[q] - 1.$



FIGURE 14. Case I.

Similarly,

$$\operatorname{val}_{mc}'[p]' = (\# \text{ components of } [p]' \cap \Gamma_{mc}') = (\# \text{ components of } [p] \cap \Gamma_{mc}) + (\# \text{ components of } [q] \cap \Gamma_{mc}) - 1 = \operatorname{val}_{mc}[p] + \operatorname{val}_{mc}[q] - 1.$$

And

$$\operatorname{val}_{yc}'[p]' = (\# \text{ components of } [p]' \cap \Gamma_{yc}') = (\# \text{ components of } [p]' \cap \Gamma_{yc}) = (\# \text{ components of } ([p] \cup [q]) \cap \Gamma_{yc}) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([q] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components of } ([p] \cap \Gamma_{yc}) \cup ([p] \cap \Gamma_{yc})) = (\# \text{ components o$$

 $\operatorname{val}_{yc}[p] + \operatorname{val}_{yc}[q].$

Comparing the contributions of the left- and right-hand sides to the function Σ , and noticing that the components $C \in CC(\Gamma) \setminus \{[p], [q]\}$ and $C' \in CC(\Gamma') \setminus \{[p]'\}$ pairwise coincide, we conclude that the two sums are equal: $\Sigma(\Gamma') = \Sigma(\Gamma)$ in Case I. This proves, in particular, that

If $\Gamma \in \mathcal{C}_n$ is a forest then $\Sigma(\Gamma) = n$.

Indeed, every forest can be obtained from an edgeless graph by adding edges which connect disjoint components.

Case II: [p] = [q]. Thus [p]' = [q]' and the four subcases are possible:

- (1) $[p]_{my} \neq [q]_{my}, [p]_{mc} \neq [q]_{mc};$ (2) $[p]_{my} = [q]_{my}, [p]_{mc} \neq [q]_{mc};$ (2') $[p]_{my} \neq [q]_{my}, [p]_{mc} = [q]_{mc};$ (3) $[p]_{my} = [q]_{my}, [p]_{mc} = [q]_{mc}.$

Subcase (1): $[p] = [q], [p]_{my} \neq [q]_{my}, [p]_{mc} \neq [q]_{mc}.$

These conditions mean that there exists a path p-q in Γ , but not in Γ_{my} or Γ_{mc} . This means that some edge of the path p-q must be *cyan*, and some other *yellow*. Thus adding a *magenta* edge *e* between *p* and *q* does create a non-monochromatic cycle.



FIGURE 15. Case II, subcase (1).

Computing the contribution of
$$[p] = [q]$$
 to $\Sigma(\Gamma)$ and of $[p]' = [q]'$ to $\Sigma(\Gamma')$ we see that:

$$\operatorname{val}_{my}'[p]' = \operatorname{val}_{my}[p] - 1,$$
$$\operatorname{val}_{mc}'[p]' = \operatorname{val}_{mc}[p] - 1,$$
$$\operatorname{val}_{yc}'[p]' = \operatorname{val}_{yc}[p],$$

and

$$\Sigma(\Gamma) = \sum_{[p] \in CC(\Gamma)} \left(\operatorname{val}_{my}[p] + \operatorname{val}_{yc}[p] + \operatorname{val}_{mc}[p] - 2 \right)$$

$$\Sigma(\Gamma') = \sum_{[p]' \in CC(\Gamma')} \left(\operatorname{val}'_{my}[p]' + \operatorname{val}'_{yc}[p]' + \operatorname{val}'_{mc}[p]' - 2 \right).$$

Hence, $\Sigma(\Gamma') = \Sigma(\Gamma) - 2$.

Subcases (2) and (2') are symmetric, so we consider only subcase (2). Subcase (2): $[p] = [q], [p]_{my} = [q]_{my}, [p]_{mc} \neq [q]_{mc}$.

This means that p and q are connected by a path p-q in Γ_{my} , but not in Γ_{mc} . In particular, every such path p-q must contain a *yellow* edge (otherwise all edges of p-q would be *magenta* and and p-q would lie in Γ_{mc}). Thus adding a new *magenta* edge e between p and q does create a non-monochromatic cycle.



FIGURE 16. Case II, subcase (2).

$$\operatorname{val}_{my}'[p]' = \operatorname{val}_{my}[p],$$
$$\operatorname{val}_{mc}'[p]' = \operatorname{val}_{mc}[p] - 1,$$
$$\operatorname{val}_{uc}'[p]' = \operatorname{val}_{uc}[p],$$

hence $\Sigma(\Gamma') = \Sigma(\Gamma) - 1$.

<u>Subcase (3):</u> $[p] = [q], [p]_{my} = [q]_{my}, [p]_{mc} = [q]_{mc}.$

Situation (3a): Every path p-q consists entirely of magenta edges. In this case, adding a new magenta edge does not create a non-monochromatic cycle. And in this case,

$$\operatorname{val}_{my}'[p]' = \operatorname{val}_{my}[p]$$
$$\operatorname{val}_{mc}'[p]' = \operatorname{val}_{mc}[p],$$
$$\operatorname{val}_{yc}'[p]' = \operatorname{val}_{yc}[p],$$

so that $\Sigma(\Gamma') = \Sigma(\Gamma)$.

Situation (3b): There exists a path p-q having a non-magenta edge, let it be yellow. Then there is a path pPq in Γ_{my} with at least one edge yellow and a path pQq in Γ_{mc} . Let p-p' be the maximal by inclusion common initial subpath of pPq and pQq consisting entirely of magenta edges. Similarly, let q'-q be the maximal by inclusion common terminal subpath of pPq and pQq consisting entirely of magenta edges. Similarly, let q'-q be the maximal by inclusion common terminal subpath of pPq and pQq consisting entirely of magenta edges. Thus, we can denote pPq = p-p'P'q'-q and pQq = p-p'Q'q'-q for some subpaths P', Q'. We observe at once that the union of the paths p'P'q' and p'Q'q' is a non-monochromatic cycle already existing in Γ .



FIGURE 17. Case II, subcase (3), situation (3b).

We notice that, in this situation, adding a magenta edge e to p, q does not change $\Sigma(\Gamma)$ since, as before,

$$\operatorname{val}_{my}'[p]' = \operatorname{val}_{my}[p],$$
$$\operatorname{val}_{mc}'[p]' = \operatorname{val}_{mc}[p],$$
$$\operatorname{val}_{yc}'[p]' = \operatorname{val}_{yc}[p],$$

so that $\Sigma(\Gamma') = \Sigma(\Gamma)$.

However, we can show by an inductive reasoning on the number of edges that in the situation (3b) we have $\Sigma(\Gamma)$ already less than n.

Indeed, we can construct Γ from the edgeless graph on *n* vertices Γ_0 by adding edges one at a time. We get a sequence of graphs:

$$\Gamma_0, \Gamma_1, \Gamma_2, \ldots, \Gamma_i, \ldots, \Gamma_m = \Gamma,$$

all belonging to C_n and such that for each i, $\Gamma_{i+1} = \Gamma_i \cup e_{i+1}$ for a new edge e_{i+1} .

If Γ_i is a forest, we showed above that $\Sigma(\Gamma_i) = \Sigma(\Gamma_0) = n$.

Suppose that we have already proved by induction on the number k of edges that if Γ_i has all cycles monochromatic and $|E(\Gamma_i)| \leq k$, then $\Sigma(\Gamma_i) = n$. Consider a new edge e_{i+1} such that $\Gamma_{i+1} = \Gamma_i \cup e_{i+1}$.

If e_{i+1} does not create a cycle, then e_{i+1} joins two components of Γ_i , and Γ_{i+1} is a forest. Hence, $\Sigma(\Gamma_{i+1}) = \Sigma(\Gamma_i)$.

If e_{i+1} creates a non-monochromatic cycle then we are in the subcase (1), (2) or (2') above, and we see that in this case $\Sigma(\Gamma_{i+1}) < \Sigma(\Gamma_i)$.

IGNAT SOROKO

If e_{i+1} creates a monochromatic cycle, i.e. e_{i+1} joins vertices $p, q \in V(\Gamma)$, and there exists a path p-q in Γ_i of the same color as e_{i+1} , then we are in the situation (3a), and $\Sigma(\Gamma_{i+1}) = \Sigma(\Gamma_i)$.

The last two cases are mutually exclusive, since, by the inductive hypothesis, Γ_i does not have a nonmonochromatic cycle.

This inductive reasoning, together with the consideration of the cases and all the subcases above, shows the following:

- adding an edge to a graph can only decrease the value of Σ ;
- graphs Γ with all cycles monochromatic have $\Sigma(\Gamma) = n$;
- creating a non-monochromatic cycle (when there were none) decreases the value of Σ by 1 or 2.

This proves the Proposition.

6. Proof of Theorems 1.4 and 1.5.

We will prove Theorem 1.5 first and Theorem 1.4 at the end of the section. We start with translating Theorem 1.5 into a statement about the Dicks graphs.

Proposition 6.1. Let F be a free group, and suppose there exist subgroups $H, K \leq F$ such that $\operatorname{rk}(H)$, $\operatorname{rk}(K) \geq 2$, $\operatorname{rk}(H \vee K) = \operatorname{rk}(H) + \operatorname{rk}(K) - i$, and $\operatorname{rk}(H \cap K) = \frac{i(i-1)}{2} + 1$, for some $i \geq 3$. Then the corresponding Dicks graphs have the following properties:

- (1) One component of Ω_{abc} is isomorphic to the complete bipartite graph $K_{i,i-1}$, while all others are singleton vertices.
- (2) Every cycle of $\Omega = \Omega_{ab} \bigvee_{\Omega_{abc}} \Omega_{bc} \bigvee_{\Omega_{abc}} \Omega_{ac}$ lies entirely in one of the subgraphs Ω_{ab} , Ω_{bc} , or Ω_{ac} (with different cycles possibly lying in different subgraphs).

Proof. Denote $h = \operatorname{rr}(H)$, $k = \operatorname{rr}(K)$. Then $\operatorname{rr}(H \cap K) = \frac{i(i-1)}{2}$ and from Theorem 4.3 we conclude that Ω_{abc} is a bipartite graph with 2h vertices in one part and 2k vertices in the other, and that Ω_{abc} has exactly i(i-1) edges. The condition $\operatorname{rk}(H \vee K) = \operatorname{rk}(H) + \operatorname{rk}(K) - i$ can be written in terms of the reduced ranks as $\operatorname{rr}(H \vee K) = h + k - (i-1)$. Also, from the discussion in subsection 2.4 we know that $\operatorname{rr}(\mathcal{T}) \geq \operatorname{rr}(H \vee K)$. Thus, part (3) of Theorem 4.3 gives us:

(# connected components of
$$\Omega_{abc}$$
) $\geq 2 \operatorname{rr}(\mathcal{T}) \geq 2h + 2k - 2(i-1).$ (7)

Let Ω_{abc} have p single-vertex components in the part corresponding to vertices from $V(\Gamma_H)$, q single-vertex components in the part corresponding to vertices from $V(\Gamma_K)$, and ℓ components C_j , $(j = 1, ..., \ell)$ each of which has at least one edge. Then inequality (7) implies:

$$p + q + \ell \ge 2h + 2k - 2(i - 1),$$

or

$$(2h-p) + (2k-q) - \ell \le 2(i-1).$$
(8)

Let s_j , t_j denote the number of vertices of C_j in each of the two parts of the bipartite graph Ω_{abc} . Then we have the following equalities: $\sum_{j=1}^{\ell} s_j = 2h - p$, $\sum_{j=1}^{\ell} t_j = 2k - q$, and (8), together with the condition $2 \operatorname{rr}(H \cap K) = i(i-1)$, becomes:

$$\sum_{j=1}^{\ell} (s_j + t_j - 1) \le 2(i-1), \tag{9}$$

$$\sum_{j=1}^{\ell} (\# \text{ edges of } C_j) = i(i-1).$$
(10)

Notice that the quantity $s_j + t_j - 1$ is the number of edges in a spanning tree of the component C_j . The following Lemma shows that the total number of edges in the left-hand side of (10) attains its maximum i(i-1) under the constraint (9) if and only if $\ell = 1$ and $C_1 = K_{i,i-1}$.

Let's call a component of a bipartite graph Γ having at least one edge, *nontrivial*.

Lemma 6.2. Let Γ be a bipartite graph having ℓ nontrivial connected components C_1, \ldots, C_{ℓ} . Under the constraint:

$$\sum_{j=1}^{\ell} (\# edges of a spanning tree of C_j) \le 2m,$$

for some $m \geq 2$, the maximal possible number of edges for Γ is achieved when $\ell = 1$ and $C_1 = K_{m,m+1}$.

Proof. (This is essentially the reasoning of Ivanov [18, (4.2)] with few more details provided.) Let's show first that for any two components C_1 , C_2 of Γ with the number of edges in their spanning trees m_1 , m_2 respectively, a single component C_0 with a spanning tree having $m_1 + m_2$ edges can have a bigger total number of edges than C_1 and C_2 together. Indeed, let s_1, t_1 be numbers of vertices in the two parts of C_1 and s_2, t_2 be numbers of vertices in the two parts of C_2 (we refer to the bipartite structure of Γ here). Then the maximal total number of edges in C_1 is s_1t_1 which is achieved when $C_1 = K_{s_1,t_1}$ and, similarly, the maximal total number of edges in C_2 is s_2t_2 . Now let's consider a join C_0 of C_1 and C_2 at a pair of vertices either in one part or in the other. In the first case we will get a bipartite graph C_0 on $s_1 + s_2 - 1$ vertices in one part and $t_1 + t_2$ vertices in the other, and in the second case C_0 will have $m_1 + m_2$ edges, so that the total number of edges in all spanning trees remains invariant. Notice that the number of edges of the complete bipartite graph $K_{s_1+s_2-1,t_1+t_2}$ is

$$(s_1 + s_2 - 1)(t_1 + t_2) = (s_1t_1 + s_2t_2) + (s_1 - 1)t_2 + (s_2 - 1)t_1 > s_1t_1 + s_2t_2, \text{ if either } s_1 \ge 2 \text{ or } s_2 \ge 2.$$

Similarly, the number of edges of the complete bipartite graph $K_{s_1+s_2,t_1+t_2-1}$ is

$$(s_1 + s_2)(t_1 + t_2 - 1) = (s_1t_1 + s_2t_2) + (t_1 - 1)s_2 + (t_2 - 1)s_1 > s_1t_1 + s_2t_2, \text{ if either } t_1 \ge 2 \text{ or } t_2 \ge 2.$$

We conclude that joining two components C_1 and C_2 allows us to have a bigger total number of edges in C_0 than the sum of edges in C_1 and C_2 , unless $s_1 = s_2 = t_1 = t_2 = 1$, when joining two $K_{1,1}$'s produces a $K_{1,2}$ with the same number of edges. But since 2m > 2, we are going to deal with components having more than two edges, and we can proceed by induction, joining components together, and each time (after possibly joining two $K_{1,1}$'s the very first time) we increase the maximal possible number of edges while preserving the total number of edges of all spanning trees. Hence we prove by induction that the maximal number of edges is achieved when there is only one nontrivial component, and it should be a complete bipartite graph $K_{s,t}$ with s+t-1=2m. Clearly, the number st of edges of $K_{s,t}$ is maximized under the constraint s+t-1=2mif and only if s and t are closest to being equal. Since s and t have opposite parity, we conclude that s=mand t=m+1, or vice versa.

Hence, part (1) is established. Notice also that the only solution to (9) and (10) implies the equality in (9), and this is equivalent to having two equalities in (7). Hence we may apply the last clause of part (3) of Theorem 4.3, which proves part (2).

We will need the following graph theoretic construction.

Let $\Omega = \Omega_{ab} \bigvee_{\Omega_{abc}} \Omega_{bc} \bigvee_{\Omega_{abc}} \Omega_{ac} = \Omega_u \sqcup \Omega_v$, Ω_a , Ω_b , Ω_c be the the Dicks graphs defined for given core graphs Γ_H , Γ_K , $\Gamma_{H\cap K}$, and let \tilde{o} , \tilde{t} be the embeddings of $\Omega_x \hookrightarrow \Omega_y$ ($x \in \{a, b, c\}$, $y \in \{u, v\}$) defined in Section 4. Let, as before, $A = \Omega_{ab} \bigvee_{\Omega_{abc}} \Omega_{ac}$, $B = \Omega_{ab} \bigvee_{\Omega_{abc}} \Omega_{bc}$, $C = \Omega_{ac} \bigvee_{\Omega_{abc}} \Omega_{bc}$. For any finite connected bipartite undirected graph Δ (with a fixed bipartite structure) define the following directed graph, which we will call the subgraph isomorphism graph for Δ , and denote it SIG(Δ):

Vertices of $SIG(\Delta)$:

 $V(SIG(\Delta)) = \{subgraphs \ \Gamma \subset \Omega, such that \ \Gamma \subset A, or \ \Gamma \subset B, or \ \Gamma \subset C, and \ \Gamma \cong \Delta \},\$

where \cong is the isomorphism of bipartite graphs, i.e. it is required to send parts of bipartite structure of Γ (induced by that of Ω) into the corresponding parts of Δ .

Since $\Omega = \Omega_u \sqcup \Omega_v$, we may define the set of directed edges of SIG(Δ) by specifying the stars of the 'source' vertices $\Gamma \subset \Omega_u$. Let Γ be a vertex of SIG(Δ) such that Γ , viewed as a subgraph of Ω , lies in $A \cap \Omega_u$. Then Γ lies in the image under \tilde{o} of some connected component $Q \subset \Omega_a$. Hence $\Gamma' = \tilde{t} \circ (\tilde{o}|_Q)^{-1}(\Gamma)$ is another vertex of SIG(Δ), with $\Gamma' \subset A \cap \Omega_v$. We connect Γ and Γ' in SIG(Δ) with a directed edge labeled a with the origin Γ and the terminus Γ' . Similarly, if $\Gamma \subset B \cap \Omega_u$, the star of Γ in SIG(Δ) will have an outgoing b-edge, and if $\Gamma \subset C \cap \Omega_u$, the star of Γ will have an outgoing c-edge, with their termini defined correspondingly. IGNAT SOROKO

Thus, a vertex Γ of SIG(Δ) may have valence 1, 2, or 3, if Γ lies in only one of the subsets A, B, C, or in only two of them, or in all three, respectively. Clearly, the same is true for vertices $\Gamma' \subset \Omega_v$.

In other words, edges of SIG(Δ) are in 1–1 correspondence with the subgraphs of Ω_a , Ω_b , Ω_c which are isomorphic to Δ , with the restrictions of \tilde{o} , \tilde{t} as the origin and the terminus maps.

Notice also that $SIG(\Delta)$ admits a natural immersion into the topological pushout \mathcal{T} , since the vertices and edges of $SIG(\Delta)$ are naturally mapped into the vertices and edges of \mathcal{T} , and this mapping is injective on stars.

Figure 18 shows the graph $SIG(K_{1,1})$ for the Dicks graphs in Figure 12.



FIGURE 18. The graph $SIG(K_{1,1})$ for the Dicks graphs in Figure 12.

An undirected graph Γ is called *k*-connected, for $k \in \mathbb{N}$, if $\#V(\Gamma) > k$ and $\Gamma \setminus Y$ is connected for every $Y \subset V(\Gamma)$ with #Y < k. We will make use of the following global version of Menger's theorem, see [9, Th. 3.3.6 (i)].

Menger's Theorem. A graph is k-connected if and only if it contains k independent paths between any two vertices. \Box

Recall that a *path* in an undirected graph is a sequence $x_1e_1x_2e_2...x_ke_kx_{k+1}$ of pairwise distinct vertices x_i and undirected edges e_i such that for all i, vertices x_i, x_{i+1} are incident to edge e_i . Two paths from x to y are *independent* if they share no other vertices except x and y.

Finally, we are ready to prove Theorem 1.5.

Proof of Theorem 1.5. Construct all the Dicks graphs Ω_{\bullet} for the core graphs Γ_H , Γ_K , $\Gamma_{H\cap K}$. Then graphs Ω and Ω_{abc} satisfy conditions (1) and (2) of Proposition 6.1. In particular, one connected component of Ω_{abc} is isomorphic to the complete bipartite graph $K_{i,i-1}$, and all other components are singleton vertices. Clearly, graph $K_{i,i-1}$ is 2–connected, if $i \geq 3$. For the rest of the proof, let Δ denote $K_{i,i-1}$. (The proof will be valid for an arbitrary 2–connected graph Δ .)

Consider the graph SIG(Δ) built for the Dicks graphs constructed above. It has a single vertex of valence 3, since, by part (1) of Proposition 6.1, only one subgraph isomorphic to Δ exists in $\Omega_{abc} = A \cap B \cap C$. We claim that all other vertices of SIG(Δ) have valence 2.

Suppose the contrary, that there exists a vertex Γ of SIG(Δ) which has valence 1. This means that subgraph $\Gamma \subset \Omega$ lies in only one of subgraphs A, B, C of Ω , let's say $\Gamma \subset A = \Omega_{ab} \bigvee_{\Omega_{abc}} \Omega_{ac}$. Since $\Gamma \not\subset B$ and $\Gamma \not\subset C$, we conclude that $\Gamma \cap (\Omega_{ab} \setminus \Omega_{abc}) \neq \emptyset$ and $\Gamma \cap (\Omega_{ac} \setminus \Omega_{abc}) \neq \emptyset$.

If there exist vertices p, q such that $p \in V(\Gamma) \cap (\Omega_{ab} \setminus \Omega_{abc})$ and $q \in V(\Gamma) \cap (\Omega_{ac} \setminus \Omega_{abc})$, then by Menger's theorem above, there exist two independent paths from p to q, and their union is a cycle which does not lie entirely in either of the subgraphs Ω_{ab} , Ω_{ac} , Ω_{bc} , thus contradicting condition (2) of Proposition 6.1.

Assume now that $V(\Gamma) \cap (\Omega_{ab} \setminus \Omega_{abc}) = \emptyset$ but there exists an edge $e \in E(\Gamma) \cap (\Omega_{ab} \setminus \Omega_{abc})$. Let s, t denote the vertices incident to the edge e. The last two conditions imply that $s, t \in V(\Omega_{abc})$. Form a new graph Γ' by subdividing the edge e into a sequence e_1pe_2 of two edges e_1, e_2 and a new vertex p such that

the vertices p and s are incident to e_1 and the vertices p and t are incident to e_2 . We claim that the graph Γ' obtained this way is also 2-connected. First, observe that $s \neq t$ since Ω has no loops (being bipartite). If we remove the vertex p from Γ' the result is the same as if we remove the edge e from Γ . Since Γ is 2-connected, Menger's theorem guarantees the existence of another path s-t in Γ which doesn't contain the edge e. Hence, $\Gamma \setminus e$ is still connected, and so is $\Gamma' \setminus p$. Also, the removal of any other vertex $p' \neq p$ from Γ' doesn't make the resulting graph disconnected. Indeed, for any two vertices p_1 , p_2 of Γ there exist at least two independent paths between them, by Menger's theorem, and only one of them may contain p'. This means that p_1 , p_2 are still connected via the other path in $\Gamma \setminus p'$ and hence in $\Gamma' \setminus p'$. Also, the vertex p is connected to any other vertex of $\Gamma' \setminus p'$ since $s \neq t$.

If $V(\Gamma) \cap (\Omega_{ac} \setminus \Omega_{abc}) \neq \emptyset$, pick a vertex q in that subset. Otherwise, as before, there is some edge $e' \in E(\Gamma) \cap (\Omega_{ac} \setminus \Omega_{abc})$, with the endpoints $s' \neq t' \in V(\Omega_{abc})$, and we perform the above operation of subdivision of edge again, applied to e', thus obtaining another 2-connected graph with e' changed into $e'_1qe'_2$. (For simplicity, we will still denote this graph by Γ' .)

Applying Menger's theorem again, we see that there exist two independent paths from p to q in Γ' . Their union is a cycle in Γ' , which has subpaths se_1pe_2t and $s'e'_1qe'_2t'$, see Figure 19. Going back to the original graph Γ and replacing these subpaths with the subpaths $s \ e \ t$ and s'e't', respectively, (the latter only if we performed the subdivision of edges twice), we get a cycle in the original graph Γ which does not lie entirely in either of the subgraphs Ω_{ab} , Ω_{ac} , Ω_{bc} , thus contradicting condition (2) of Proposition 6.1.



FIGURE 19. A cycle in Γ' from the proof of Theorem 1.5.

Therefore, all vertices of the graph $\operatorname{SIG}(\Delta)$ have valence 2, except for a single vertex of valence 3. But this is impossible, since in any graph the number of vertices of odd valence must be even (otherwise the count for the number of edges, $\#E^+(\Gamma) = \frac{1}{2} \sum_{v \in V(\Gamma)} \operatorname{val}(v)$, would be a half-integer). The obtained contradiction proves that the values of the ranks of $H, K, H \lor K$, and $H \cap K$ in Theorem 1.5

The obtained contradiction proves that the values of the ranks of $H, K, H \lor K$, and $H \cap K$ in Theorem 1.5 are non-realizable.

Example 6.3. Interestingly, condition (1) alone in Proposition 6.1 does not make the Dicks graphs non-realizable, as the following example shows. Let subgroups $H, K \leq \theta(F) \leq F(a, b, c)$ be given by $H = \theta(\langle b, a^3, ab^{-1}a, ab^2a^{-1}\rangle) = \langle cb^{-1}, (ca^{-1})^3, ca^{-1}ba^{-1}, ca^{-1}(cb^{-1})^2ac^{-1}\rangle, K = \theta(\langle a^{-1}b, b^{-2}ab^2\rangle) = \langle ab^{-1}, bc^{-1}ba^{-1}cb^{-1}cb^{-1}\rangle$. Then $\Omega_{abc} = K_{2,3} \cup \{$ three vertices $\}$, see Figure 20. The graph SIG($K_{2,3}$) has one vertex of valence 3 and three vertices of valence 1. Note also that (# connected components of Ω_{abc}) = $4 > 2 = 2 \operatorname{rr}(\mathcal{T})$, cf. part (3) of Theorem 4.3.

The case i = 3 of Theorem 1.5 resolves the remaining open case m = 4 of Guzman's "Group-Theoretic Conjecture" in the affirmative:

Corollary 1.6. Let F be a free group. If two subgroups $H, K \leq F$ both have ranks equal to 4, and $\operatorname{rk}(H \cap K) \geq 4$, then $\operatorname{rk}(H \vee K) \leq 4$.

Proof. Indeed, looking at the locus of known realizable values and the region of proved non-realizable values for $\operatorname{rk}(H) = \operatorname{rk}(K) = 4$, see Figure 21, we conclude that the GTC for m = 4 holds true if and only if the tuple $(\operatorname{rk}(H \lor K), \operatorname{rk}(H \cap K)) = (5, 4)$ is not realizable. But this is exactly what Theorem 1.5 says for $\operatorname{rk}(H) = \operatorname{rk}(K) = 4$ and i = 3.



FIGURE 20. The core graphs, the Dicks graphs and the topological pushout for the subgroups H, K from Example 6.3. The graph Ω_{abc} is the union of $\Omega_{v,abc} = K_{2,3}$ and $\Omega_{u,abc} = \{$ three singleton vertices 1, 3, 5 $\}$.

Invoking the implication theorem from [13], we obtain a proof of the "Geometric Conjecture" for k = 6: Corollary 1.7. Let M be a closed, orientable, hyperbolic 3-manifold. If $\pi_1(M)$ is 6-free then there exists a point P in M such that the set of all elements of $\pi_1(M, P)$ represented by loops of length less than $\log(11)$



FIGURE 21. Guzman's GTC for $m = \operatorname{rk}(H) = \operatorname{rk}(K) = 4$.

Now we are going to prove Theorem 1.4.

is contained in a free subgroup of $\pi_1(M)$ of rank at most 3.

Proof of Theorem 1.4. Let k, v and c be as in the statement of Theorem 1.4. The existence of subgroups $H, K \leq F$ with $\operatorname{rk}(H) = 2$, $\operatorname{rk}(K) = k$, $\operatorname{rk}(H \vee K) = v$, and $\operatorname{rk}(H \cap K) = c$, such that $c \leq k + 2 - v$, follows from Theorem 1.1. Let's prove that if c > k + 2 - v, such subgroups do not exist.

Suppose the contrary, that subgroups H, K with $\operatorname{rk}(H) = 2$, $\operatorname{rk}(K) = k$, $\operatorname{rk}(H \lor K) = v$, $\operatorname{rk}(H \cap K) = c$, and c > k + 2 - v do exist. Denote i = k + 2 - v and let $d \ge 1$ be such that c = i + d. From Theorem 4.3 we see that Ω_{abc} is a bipartite graph with 2 vertices in the $V(\Gamma_H)$ -part of Ω_{abc} and 2(k-1) vertices in the $V(\Gamma_K)$ -part (note that now k denotes $\operatorname{rk}(K)$ so that $\operatorname{rr}(K) = k - 1$), and that Ω_{abc} has 2(c-1) = 2(i-1)+2dedges. Denote nontrivial components of Ω_{abc} as C_1, C_2, \ldots, C_ℓ and call the edges of a spanning tree of C_j the spanning edges of C_j . Arguing as in the proof of Proposition 6.1, we get:

$$\sum_{j=1}^{\ell} \left(\# \text{ spanning edges of } C_j \right) \le 2(i-1)$$
(11)

and

$$\sum_{j=1}^{\ell} \left(\# \text{ edges of } C_j \right) = 2(i-1) + 2d.$$
(12)

Denote the two vertices in the $V(\Gamma_H)$ -part of Ω_{abc} as z and w. We claim that z and w belong to the same nontrivial component C_j . Indeed, if z and w belong to different components, then these components are trees, and, in particular, the count in (12) is equal to that of (11), which contradicts the condition $d \ge 1$. Hence, the component C_j containing z and w is the only nontrivial component of Ω_{abc} , i.e. $j = \ell = 1$.

Denote

$$s = (\# \text{ spanning edges of } C_1),$$
 (13)

 $m = (\# \text{ vertices of valence 2 in the } V(\Gamma_K) - \text{part of } \Omega_{abc}),$ (14)

$$q = (\# \text{ singleton vertices in the } V(\Gamma_K) \text{-part of } \Omega_{abc}), \tag{15}$$

see Figure 22.

In particular, we see that $m \ge 2d+1$ (indeed, m-1 is the rank of C_1 , i.e. the number of edges of C_1 minus the number of spanning edges of C_1 , hence is at least 2d), and that Ω_{abc} contains a subgraph Δ isomorphic to the complete bipartite graph $K_{2,m}$.

Consider the graph $SIG(K_{2,m})$, as defined after the proof of Proposition 6.1. Arguing as in the proof of Theorem 1.5, we observe that $SIG(K_{2,m})$ has a unique vertex of valence 3, and hence must have another vertex of odd valence, that is of valence 1. Let's call the subgraph of Ω , corresponding to this valence 1

 \square



FIGURE 22. The graph Ω_{abc} when $\operatorname{rk}(H) = 2$.

vertex, Δ' . Without loss of generality we can assume that Δ' lies in $A = \Omega_{ab} \bigvee_{\Omega_{abc}} \Omega_{ac}$, and hence that $\Delta' \cap (\Omega_{ab} \setminus \Omega_{abc}) \neq \emptyset$ and $\Delta' \cap (\Omega_{ac} \setminus \Omega_{abc}) \neq \emptyset$. We conclude that $\Delta' \cap \Omega_{abc} \neq \emptyset$, and, since the isomorphisms involved in the definition of SIG preserve the bipartite structure, the image of m vertices of $K_{2,m}$ in Δ' (call this subset M) is a subset of the q singleton vertices of the $V(\Gamma_K)$ -part of Ω_{abc} . In particular, $m \leq q$, see Figure 23.



FIGURE 23. The graph $\Delta' \cong K_{2,m}$ corresponding to the valence 1 vertex of SIG $(K_{2,m})$.

To estimate $rr(\mathcal{T})$ we recall equation (5) from Section 4:

$$2\operatorname{rr}(\mathcal{T}) = \sum_{v \in V(\mathcal{T})} (\operatorname{val}(v) - 2).$$
(5)

From Proposition 4.2 we know that the vertices of \mathcal{T} are connected components of Ω . Vertices of \mathcal{T} of valence ≥ 3 correspond to certain subgraphs of Ω_{abc} , with the exact relation between components of Ω_{abc} and the valence of the corresponding vertex of \mathcal{T} given by the component connectivity graph Γ , as constructed in the proof of Theorem 4.3. The vertices of Γ are connected components of Ω_{abc} . Two vertices p, q of Γ may be connected by up to three undirected edges (colored magenta, yellow, and cyan) in Γ , if there exists an Ω_{abc} -avoidant path connecting components p and q which lies entirely in Ω_{ab} , Ω_{ac} , or Ω_{bc} , respectively. The right-hand sum of equation (5) equals the value of the function Σ on Γ , as defined in (6) of Section 5. The main conclusion of Proposition 5.1 is that this value, and hence, the value $2 \operatorname{rr}(\mathcal{T})$, is bounded above by the number of vertices of Γ .

The component connectivity graph Γ in the situation we are considering will have one vertex for the only nontrivial component C_1 of Ω_{abc} and q vertices for the remaining singleton components. However, if we identify the m vertices of M in Δ' into a single vertex, thus forming a new graph Γ' , we observe that the value of function Σ on Γ and Γ' is the same. Indeed, these m vertices belong to the same component in each of the subgraphs A, B, C of Ω , and hence their contribution to the number of a-edges, b-edges, and c-edges of \mathcal{T} is the same as if they were a single vertex of Ω_{abc} . Hence, we can use the graph Γ' for computing the quantity $2 \operatorname{rr}(\mathcal{T})$, and we conclude that the latter is bounded above by $n = \#V(\Gamma')$. We now estimate n.

We have: n = 1 + 1 + (q - m), where the ones correspond to the component C_1 and the subset M, which is one vertex of Γ' . Since a spanning tree of C_1 contains only one vertex of valence 2 in the $V(\Gamma_K)$ -part (for otherwise there would be a cycle in it), all other vertices of the spanning tree that lie in the $V(\Gamma_K)$ -part have valence 1. Now from (15) and (13) above we deduce that:

$$q = 2(k-1) - (s-1).$$

Also from (14) and (12) it follows that:

 $m = (\# \text{ edges of } C_1) - (\# \text{ spanning edges of } C_1) + 1 = 2(i-1) + 2d - s + 1.$

Hence, we get:

$$n = 2 + q - m = 2 + 2(k - 1) - (s - 1) - 2(i - 1) - 2d + s - 1 = 2(k - i - d + 1),$$

and we deduce from formula (5) above (and the discussion following it) that

$$\operatorname{rr}(\mathcal{T}) \le k - i - d + 1.$$

Now recall that $v = \operatorname{rk}(H \lor K) = k + 2 - i$ and that $\operatorname{rk}(H \lor K) \leq \operatorname{rk}(\mathcal{T}) = \operatorname{rr}(\mathcal{T}) + 1$. We get:

$$v = \operatorname{rk}(H \lor K) \le \operatorname{rk}(\mathcal{T}) = (k+2-i) - d = v - d,$$

which yields a contradiction with the condition that $d \ge 1$. Since d was defined as d = c - (k + 2 - v), this proves that $c + v \le k + 2$.

References

- Ian Agol, Marc Culler, Peter B. Shalen, Singular surfaces, mod 2 homology, and hyperbolic volume. I. Trans. Amer. Math. Soc. 362 (2010), no. 7, 3463–3498.
- [2] Frédérique Bassino, Cyril Nicaud, Pascal Weil, Random generation of finitely generated subgroups of a free group. Internat. J. Algebra Comput. 18 (2008), no. 2, 375–405.
- [3] Gilbert Baumslag, Alexei G. Myasnikov, Vladimir Shpilrain, Open problems in combinatorial and geometric group theory. http://www.sci.ccny.cuny.edu/~shpil/gworld/problems/oproblems.html
- [4] Oleg Bogopolski, Introduction to group theory. (EMS Textbooks in Mathematics.) European Mathematical Society (EMS), Zürich, 2008. x+177 pp.
- [5] Marc Culler, Peter B. Shalen, 4–free groups and hyperbolic geometry. J. Topol. 5 (2012), no. 1, 81–136.
- [6] Warren Dicks, Equivalence of the strengthened Hanna Neumann conjecture and the amalgamated graph conjecture. Invent. Math. 117 (1994), 373–389.
- [7] Warren Dicks, Joel Friedman's proof of the strengthened Hanna Neumann conjecture, pp. 91–101 in [11].
- [8] Warren Dicks, Simplified Mineyev. Preprint at http://mat.uab.cat/~dicks/SimplifiedMineyev.pdf
- [9] Reinhard Diestel, Graph Theory, Fifth edition. Graduate Texts in Mathematics, 173. Springer, Berlin, 2017. xviii+428 pp.
- [10] Matthew Fayers, The genyoungtabtikz package, version 1.14, 2016-10-05. http://www.maths.qmul.ac.uk/~mf/genyoungtabtikz.html.
- [11] Joel Friedman, Sheaves on graphs, their homological invariants, and a proof of the Hanna Neumann conjecture: with an appendix by Warren Dicks. Mem. Amer. Math. Soc. 233 (2015), no. 1100, xii+106 pp.
- [12] The GAP Group, GAP Groups, Algorithms, and Programming, Version 4.10.0, 2018, https://www.gap-system.org.
- [13] Rosemary K. Guzman, Hyperbolic 3–manifolds with k–free fundamental group. Topology Appl. 173 (2014), 142–156.
- [14] Rosemary K. Guzman, Peter B. Shalen, The geometry of k-free hyperbolic 3-manifolds. J. Topol. Anal. https://doi. org/10.1142/S1793525320500016.
- [15] Joshua E. Hunt, The Hanna Neumann Conjecture and the rank of the join. arXiv:1509.04449
- [16] Wilfried Imrich, Thomas Müller, On Howson's theorem. Arch. Math. (Basel) 62 (1994), no. 3, 193–198.
- [17] Sergei V. Ivanov, On a conjecture of Imrich and Müller. J. Group Theory 20 (2017), no. 4, 823–828.
- [18] Sergei V. Ivanov, On joins and intersections of subgroups in free groups. J. Comb. Algebra 2 (2018), 1-18.
- [19] Andrei Jaikin-Zapirain, Approximation by subgroups of finite index and the Hanna Neumann conjecture. Duke Math. J. 166 (2017), 1955–1987.
- [20] Ilya Kapovich, Alexei Myasnikov, Stallings foldings and subgroups of free groups. J. Algebra 248 (2002), no. 2, 608–668.
- [21] Richard P. Kent IV, Achievable ranks of intersections of finitely generated free groups. Internat. J. Algebra Comput. 15 (2005), no. 2, 339–341.
- [22] Richard P. Kent IV, Intersections and joins of free groups. Algebr. Geom. Topol. 9 (2009), no. 1, 305–325.
- [23] Larsen Louder, D.B. McReynolds, Graphs of subgroups of free groups. Algebr. Geom. Topol. 9 (1) (2009), pp. 327–355.
- [24] Igor Mineyev, Groups, graphs, and the Hanna Neumann conjecture. J. Topol. Anal. 4 (2012), no. 1, 1–12.
- [25] Christian Sievers, FGA (Free Group Algorithms) a GAP package, Version 1.4.0, 2018, http://www.icm.tu-bs.de/ag_algebra/software/FGA.
- [26] John R. Stallings, Topology of finite graphs. Invent. Math. 71 (1983), no. 3, 551-565.
- [27] Till Tantau, The TikZ and PGF Packages, manual for version 3.1, http://sourceforge.net/projects/pgf, 2019-01-05.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LA 70803, USA *E-mail address*: ignatsoroko@lsu.edu