Fractal structures in Adversarial Prediction

Rina Panigrahy¹ and Preyas Popat²

 ¹ Microsoft Corp. Mountain View, CA rina@microsoft.com
 ² New York University and University of Chicago New York, NY and Chicago, IL popat@cs.nyu.edu

Abstract. Fractals are self-similar recursive structures that have been used in modeling several real world processes. In this work we study how "fractal-like" processes arise in a prediction game where an adversary is generating a sequence of bits and an algorithm is trying to predict them. We will see that under a certain formalization of the predictive payoff for the algorithm it is most optimal for the adversary to produce a fractal-like sequence to minimize the algorithm's ability to predict. Indeed it has been suggested before that financial markets exhibit a fractal-like behavior [1,2]. We prove that a fractal-like distribution arises naturally out of an optimization from the adversary's perspective.

In addition, we give optimal trade-offs between predictability and expected deviation (i.e. sum of bits) for our formalization of predictive payoff. This result is motivated by the observation that several time series data exhibit higher deviations than expected for a completely random walk.

1 Introduction

Consider an adversary who is producing a sequence of bits (each bit is +1 or -1) and an algorithm having seen a certain number of bits is interested in predicting the next x bits. Say the algorithm gets a payoff of 1 for every bit that it predicts correctly and -1 for every bit where it is wrong. This is like an idealized stock market where each day the price changes by +1 or -1 percent and the algorithm is required to make a bet on the daily direction. We ask what is the most adversarial distribution on sequence of bits so as to minimize the algorithm's payoff. Clearly the uniform distribution where every bit is chosen independently and uniformly at random is the most adversarial, since the expected payoff of any algorithm is always exactly 0.

Given a sequence s of bits, let h(s) be the sum of the bits in s i.e. the *height* of the sequence when plotted cumulatively. We will refer to the magnitude of height as deviation. For $s \in \{-1, 1\}^T$ chosen uniformly at random the typical deviation s is $\Theta(\sqrt{T})$.

The question we study here is: what is the most adversarial distribution on sequences if the distribution is required to be heavy-tailed, say the typical height should be $k\sqrt{T}$ where k > 1. Indeed it has been observed in several studies that the distribution of financial time series is heavy-tailed [3,4]. A natural heavytailed distribution is to pick a random string conditioned on its height being at least $k\sqrt{T}$. This is essentially the highest entropy distribution with the property that the typical height is around $k\sqrt{T}$. However the highest entropy distribution is not the least predictable. Indeed for large k, it tends to rise/drop rather linearly to its final height. Thus by observing the initial segment of bits, the algorithm can easily infer the direction of the remaining bits to get a large payoff.

One distribution that has been suggested for financial markets is the Fractional Brownian Motion (FBM) [5,6] which is a generalization of the Brownian motion. For our purposes, the Brownian motion can be thought of as a continuous variant of the uniform distribution on bits. FBM is characterized by a single parameter Hwhich is called the Hurst parameter, and the typical height achieved by sequences drawn from FBM(H) is around T^H . For H > 1/2, the increments of FBM are positively correlated while the case H = 1/2 corresponds to Brownian motion.

To make our question precise we introduce a measure of unpredictability for a distribution which is motivated by the notion that the expected payoff of an algorithm on an interval I having observed the previous bits should be small compared to the standard deviation of height in I. Intuitively, we are enforcing a low signal-to-noise ratio.

Definition 1 Let D be a distribution which produces bits in an online fashion and s be the sequence of bits that have been produced immediately preceding an interval I. Let $\mathbb{E}[A_s(I)]$ denote the expected payoff of an algorithm A on interval I (where the bits in I are produced according to D conditioned on having produced s immediately before I). Note that A must fix its prediction for I based solely on s and before looking at any bits within I.

We say that D is δ -unpredictable if for all A, s and I, $\mathbb{E}[A_s(I)] \leq \delta \cdot \sqrt{|I|}$.




Fig. 1: Growth charts for two different types of adversarial sequences. The first is the cumulative plot of a random i.i.d sequence with a constant upward bias. The second is an α -inverting sequence as in Definition 2. Note that the latter plots seems to change direction more significantly than the former.

For example an algorithm may notice a high density of +1's and may decide to predict +1 for the next few bits (this would correspond to a "buying" a stock) for the next x bits. Note that \sqrt{x} is the standard deviation in the payoff of an algorithm for the uniform distribution on x bit sequences and thus we are asking that the payoff of the algorithm for a δ -unpredictable distribution is negligible compared to this standard deviation (we will in fact construct distributions where the standard deviation is much higher than \sqrt{x}). Roughly, this is equivalent to saying that the signal to noise ratio in any interval is negligible.

We ask what is the maximum deviation that can be achieved by a δ -unpredictable distribution D. We will look at maximizing measures such as median deviation or mean deviation: $\mathbb{E}_{s\sim D}[|h(s)|]$ (we will show that our claims hold with respect to any of these measures).

We show that there is a δ -unpredictable distribution which achieves a deviation of $\sqrt{T}(1 + \Omega(\delta \log T))$. Thus, the deviation can be $\omega(\sqrt{T})$ for $\delta = o(1)$. The distribution we construct is a variant of a discretization of FBM. We also show that the highest deviation that can be achieved by a δ -unpredictable distribution is $\sqrt{T}(1 + O(\delta \log T))$. In addition, we construct a distribution which is a simple discretization of FBM and show that the deviation achieved by this distribution is $T^{1/2+\Theta(\delta)}$. Though this distribution is not δ -unpredictable, it satisfies a similar but weaker property.

A nice property of δ -unpredictable distributions is that they are "fractal-like" in some sense. We use the terms fractal-like and fractal somewhat interchangeably. Normally fractal is considered to be a self-similar recursive structure in Euclidean space (usually with non-integer dimension to exclude trivial patterns). Traditionally this has not been applied to bit sequences. Therefore we refrain from calling such sequences strictly a fractal. To formalize our "fractal-like" property, we first define a notion of inversion for a deterministic sequence. The property essentially says that if in any interval there is a huge rise, then there must be a sub-interval where there must be a proportionally big fall and vice versa.

Definition 2 (α -Inversion) Given a sequence $s \in \{-1, 1\}^T$, it is said to be α -inverting if for every interval X within [1,T] (of at least some constant length) there is a sub interval Y such that $h(s_X)$ and $h(s_Y)$ are of opposite sign and $|h(s_Y)|/|h(s_X)| \ge \alpha$. Here by s_I we mean the sequence s restricted to interval I. We refer to the largest feasible α as the inversion ratio of s.

Observe that an α -inverting sequence resembles a fractal in a certain sense. To see this, note that in a sequence s such that h(s) > 0, if we locate the biggest contiguous rise, it may be divided into three parts $s_1s_2s_3$ where s_2 has a net downward slope and s_1, s_3 have a positive slope each. But one can recurse and divide each of the three substrings further into three parts each and thus the sequence has a recursive, self-similar structure.

We show that any δ -unpredictable distribution is α -inverting in a certain sense. Since we are dealing with a distribution rather than a deterministic sequence we need an appropriate generalization of Definition 2 which is stated in Section 1.1. It will be clear from the definition that the highest entropy sequence we discussed earlier has a very small inversion ratio compared to δ -unpredictable distributions.

1.1 Main results

In this section we describe our main results in more detail. As we mentioned earlier, the adversarial distributions we construct are closely related to and inspired from FBM.

FBM with parameter H is the unique continuous time, Gaussian process $B_H(t)$ which satisfies B(0) = 0, $\mathbb{E}[B_H(t)] = 0$ for all t and has covariance function:

$$\mathbb{E}[B_H(t)B_H(s)] = \frac{1}{2}(|t|^{2H} + |s|^{2H} - |t-s|^{2H})$$

The process B_H is translation invariant and is self-similar in the sense that $\{B_H(at) : t \in \mathbb{R}\}$ is identical in distribution to $\{a^H B_H(t) : t \in \mathbb{R}\}$ for all a > 0. Furthermore, $B_H(t)$ is normally distributed with variance t^H . Thus any interval of length t has deviation about t^H . The case H = 0.5 corresponds to the standard Brownian motion.

The analysis of the FBM usually requires an understanding of integrated Wiener processes. The first adversarial distribution we construct is a discrete variant of the FBM that produces bits instead of real numbers. We denote this distribution as FRACTAL RANDOM WALK (FRW).

The sequence is constructed recursively in lengths that are powers of 2. To produce a sequence of length 2n, we concatenate two recursively constructed sequences of length n each, and change the height of the second sequence by a factor proportional to the height h of the first sequence. This is done by flipping approximately δh (-1)'s to +1's if h > 0 (and +1's to -1 otherwise.) A formal description of the construction appears in Section 3.

While this lacks the translation invariance and the exact self-similarity properties of the FBM, it still has the property that any interval of size t has deviation $t^{1/2+\Theta(\delta)}$.

To see this, note that if h_1, h_2 denote the heights of the two sequences that are concatenated to produce the sequence of length 2n after altering the second string then $E[h_1h_2] = 2\delta \mathbb{E}[h_1^2] = 2\delta \mathbb{E}[H(n)^2]$ where H(n) is a random variable that denotes the height of a random sequence of length n drawn from FRW. So $\mathbb{E}[H(2n)^2] = E[(h_1 + h_2)^2] = E[h_1^2] + E[h_2^2] + 2E[h_1h_2] = (2 + \delta)E[H(n)^2]$. The recurrence works out to a root mean square deviation $(\sqrt{E[H(n)^2]})$ of about $n^{1/2+\Theta(\delta)}$.

This informal description skips over technical issues such as discretization. Furthermore, extending this argument to show that the high deviation is achieved with constant probability is more complicated and is done in Theorem 7. Note that a constant probability bound for achieving a particular deviation is stronger than showing a high deviation in expectation (using Markov's inequality). We note that this distribution is not δ -unpredictable but satisfies a weaker property (Theorem 20). For completeness, we show that the FBM (continuous version)

with $H = 1/2 + \delta$ is also not δ -unpredictable in the strict sense (Claim D). We also note that the highest entropy distribution is very poor in terms of δ unpredictability (Claim 25).

We construct another distribution, which we call OPTIMAL FRACTAL RAN-DOM WALK (OPT-FRW) which has optimal trade-offs between deviation and predictability. The distribution OPT-FRW is a simple but important twist on the above process where instead of flipping $\delta \cdot h$ bits, we flip $\delta \cdot \sqrt{n}$ bits in the direction of h.

Theorem 3. (Theorems 12, 18 and 13) The distribution OPT-FRW is $O(\delta)$ unpredictable and achieves a deviation of $\sqrt{T}(1 + \Omega(\delta \log T))$ with constant probability. Further, no δ -unpredictable distribution can achieve an expected deviation higher than $\sqrt{T}(1 + O(\delta \log T))$.

We now turn to formalizing the relationship between δ -unpredictability and "fractal-like" property of a distribution.

For a deterministic sequence we show that an α -inverting sequence with the highest deviation is a fractal.

Theorem 4. (Claim C)

Let s be an α -inverting sequence of length t (Definition 2), where α is bounded above by a constant. Then the highest deviation that can be achieved by s for large t is t^{θ} where θ is the solution to the equation $1 = 2((1 + \alpha)/2)^{1/\theta} + \alpha^{1/\theta}$. Furthermore, this deviation is actually achieved by an appropriately designed fractal.

For distributions D over sequences we define the following variant of the earlier inversion rule.

Definition 5 ((α, q) -**Inversion**) A distribution D is said to be (α, q) -inverting if for any interval X of at least some constant length) with median deviation $\Delta = \Omega(\delta \sqrt{|X|})$, with probability at least q there is a sub interval Y such that $h(s_X)$ and $h(s_Y)$ are of opposite sign and $|h(s_Y)| \ge \alpha \cdot \Delta$. Here by s_I we mean the sequence s restricted to interval I. This should hold even if one conditions on a given history of bits seen before the interval X.

We note (see Observation 24) that a uniform random sequence is (α, q) inverting for some constants α, q . Further the probability parameter q can be made as high as $1 - \varepsilon$ by reducing the inversion ratio α to $\Theta(1/\log(1/\varepsilon))$.

The following theorem establishes that every δ -unpredictable distribution must be fractal-like in the sense that it is $(\Omega(1), \Omega(1))$ inverting.

Theorem 6. (Theorems 16 and 17) For δ small enough, any δ -unpredictable distribution is also (α, q) -inverting for some constants α, q . Further by dropping the inversion ratio α to $\Theta(1/\log T)$ the probability q can be made as high as $1-1/T^{\Omega(1)}$ for all intervals of length at least $\Omega(\log T)$. Thus the condition holds with high probability simultaneously for all such intervals.

1.2 Related Work

Many studies support the thesis that fractals occur naturally in several real world processes in diverse fields such as physics, finance and geography [7,8,9]. Ralph Elliot [1], a professional accountant, suggested the use of fractal like "waves" in understanding financial markets. Fractal models for finance have also been studied widely in the academic community. Fractional Brownian Motion (FBM) was introduced as a variant to the well known Brownian Motion by Mandelbrot and van Ness in [5]. In addition to financial time series modeling, FBM has also found applications in the study of network traffic and fluid turbulence [10,6].

The reason for considering FBM rather than the standard Brownian motion for financial modelling was the observation that the distribution of financial time series is heavy-tailed [3,4]. This means that the deviations achieved are a bit higher than those expected for Brownian motion. It has been argued that modeling S&P500 price data according to FBM produces an estimated value of the Hurst parameter H to be slightly over the 0.5 value that corresponds to the standard Brownian Motion [11]. Values of H > 0.5 allow for long range (positive) correlations in the time series that results in a higher than normal deviation. Besides FBM other models such as *p*-stable distributions and levy distributions [12,4,13] provide an alternate explanation for the heavy tailed nature of time series data by allowing heavier tails for the price changes *in each unit time* that are *independent* across time. In contrast, the FBM uses normally distributed price changes in each unit time, and the high deviations are achieved by correlations across time.

Works such as [14,15] have analyzed the level of arbitrage present in FBM. The authors in [16] have analyzed the predictability of the FBM using a different loss function from ours. Other researchers [17,18] have studied the prediction problem as a game between an algorithm and an adversary, and derived that the optimal strategy for the adversary resembles a Brownian Motion. The work in [18] was inspired by [19] where the authors provide robust upper and lower bounds for pricing European call options, under the no-arbitrage assumption when the price process is assumed to be *discrete and discontinuous* as opposed to the Black Scholes model [20] where the price process is taken to be continuous.

1.3 Discussion and Future work

Note that our notion of δ -unpredictable requires the algorithm to fix its prediction for an entire interval I before looking at any of its bits. A stronger notion of unpredictability is to allow the algorithm to change its prediction for the interval after looking at bits within I. In other words, at every point the algorithm tries to simply predict the next bit, based on the bits it has seen so far. One could ask what is most adversarial distribution in this setting which achieves a high deviation. In this setting, for any sequence s, a bounded regret algorithm such as Weighted Majority can achieve a payoff of $|h(s)| - c\sqrt{|s|}$ where $c := \sqrt{2/\pi}$ [21,22]. So for a distribution D which achieves typical deviation $k\sqrt{T}$, it is always possible to get a payoff of $(k - c)\sqrt{T}$. It is also fairly straightforward to construct a distribution D such that no algorithm can achieve an expected payoff better than $(k-c)\sqrt{T}$ even when it predicts one bit at a time. We also note that while the distributions inspired by FBM have some guarantees in terms of δ -unpredictability, they perform poorly in this model when one is allowed to predict based on all previous bits (see Claim 26).

One possible justification for our notion of δ -unpredictability is that changing predictions very often may have a cost associated with it. Although this may be a reasonable assumption (at least for financial markets), it is only a conjecture at this point and we invite further comments on this issue.

An interesting direction for further research is to look for natural constraints on real world processes which provably result in the formation of fractal-like processes.

2 Preliminaries

Here is some common notation we use throughout the paper. For a sequence of bits $s \in \{-1, 1\}^T$, h(s) denotes the sum of bits in s i.e. the *height* of s. We refer to the magnitude of height as deviation.

We will be working with several aggregate measures of deviation for a distribution such as median deviation (or generalized median), mean deviation and root-mean-squared deviation $(\sqrt{\mathbb{E}_{s\sim D}[h(s)^2]})$. Note that mean deviation is no more than root-mean-squared deviation and the generalized median is bounded by mean deviation up to constant factors using Markov's inequality (as long as the probability in generalized median is at least a constant). We will prove our upper bounds for root-mean-squared deviation and lower bounds for generalized median and so they will hold for all measures up to constants.

We will typically denote random variables by capital letters and fixed sequences by small letters.

3 Construction of Adversarial distributions

In this section we formally construct our adversarial distributions. Each of these distributions has two parameters, l which is the length of the sequence in the base case and $\delta > 0$.

We will construct the distributions inductively: having constructed $D_{\delta}(n)$ we will show how to construct $D_{\delta}(2n)$ (the base case for n = l is simply a random sequence in $\{-1, 1\}^l$). In both cases below, we describe the distribution $D_{\delta}(2n)$ in terms of how to generate a sequence $s \sim D_{\delta}(2n)$ given access to distribution $D_{\delta}(n)$.

FRACTAL RANDOM WALK (FRW_{l,δ}) (2n)

- 1. Generate sequences s_1, s_2 independently according to $\text{FRW}_{l,\delta}(n)$
- 2. If height of s_1 is positive, change exactly $\delta \cdot h(s_1) 1$'s in s_2 to 1 (if they exist, otherwise change as many as possible). Similarly, if height of s_1 is negative, change exactly $\delta \cdot h(s_1)$ 1's in s_2 to -1 (if they exist). Call the resulting sequence s'_2 .

3. Set $s = s_1 \cdot s'_2$ i.e. the concatenation of s_1 and s'_2

Optimum Fractal Random Walk (Opt-FRW_{l,δ})(2n)

- 1. Generate sequences s_1, s_2 independently according to OPT-FRW_{l,δ}(n)
- 2. If height of s_1 is positive, change exactly $\delta\sqrt{n} 1$'s in s_2 to 1 (if they exist, otherwise change as many as possible). Similarly, if height of s_1 is negative, change exactly $\delta\sqrt{n}$ 1's in s_2 to -1 (if they exist). Call the resulting sequence s'_2 .
- 3. Set $s = s_1 \cdot s'_2$ i.e. the concatenation of s_1 and s'_2

Note: Note that both distributions involve changing exactly r bits in s_2 where r is a real number. Intuitively, we want to change each bit of the appropriate sign in s_2 with probability r/n. However, it is simpler to analyze the deviation of the distributions when we change exactly r bits. The fact that r is a real number and not an integer will not make much difference since our base case l will be an increasing function of T (total number of bits to be produced) and so the discretization errors can be safely ignored.

3.1 High deviation

In this section we show that the distributions we constructed achieve high deviation with constant probability. What follows is a proof sketch for high deviation of distribution $FRW_{i,\delta}$. Due to space constraints, the proof for OPT- $FRW_{i,\delta}$ and for the intermediate claims appears in the appendix (Section B).

Theorem 7. The distribution $\operatorname{FRW}_{l,\delta}(T)$ achieves a deviation of $T^{1/2+\Theta(\delta)}$ with probability at least $1/2 - \varepsilon$ where $\varepsilon \leq T^{-10}$.

Proof: To analyze the height distribution of $\text{FRW}_{l,\delta}$ it will be more convenient to define another process which is similar to $\text{FRW}_{l,\delta}$ but which can assume integer values instead of bits.

Augmented Fractal Random Walk (AFRW_{l,δ}) (2n)

- 1. Generate sequences s_1, s_2 independently according to AFRW_{l,δ}(n)
- 2. If height of s_1 is positive, change exactly $\delta \cdot h(s_1) 1$'s in s_2 to 1 (if they exist). Similarly, if height of s_1 is negative, change exactly $\delta \cdot h(s_2)$ 1's in s_2 to -1 (if they exist). Call the resulting sequence s'_2 .
- 3. Augment: If there aren't enough -1's to flip in s_2 , then add 2 to some of the numbers so that the increase in height is exactly $\delta \cdot h(s_1)$. Similarly for 1's.
- 4. Set $s = s_1 \cdot s'_2$ i.e. the concatenation of s_1 and s'_2

For the random variable $S \sim AFRW_{l,\delta}$, we can exactly characterize the distribution of h(S).

Claim 8 (Claim 21) For $n = 2^i \cdot l$, $S \sim AFRW_{l,\delta}(n)$,

$$h(S) = \sum_{U \subseteq [i]} r^{|U|} h(X_U)$$
(3.1)

where $r = (1 + \delta)$ and each X_U is independently and uniformly distributed in $\{-1, 1\}^l$.

We then apply the Berry-Esseen theorem (Theorem 28) to show that the deviation of $|h(AFRW_{l,\delta})|$ is high.

Lemma 1. (Lemma 3) Median of $|h(AFRW_{l,\delta})|$ is $n^{1+\Omega(\delta)}$.

Next we show that the probability of executing step **Augment** in $AFRW_{l,\delta}$ is exponentially small. Note that when constructing a sequence of size T, the inductive steps of distribution $AFRW_{l,\delta}$ are executed at most 2T times. We show that when starting with sequences of size l where $l = 100 \log T$, the probability that sequence s_2 doesn't have enough 1's or -1's to flip at a particular stage is at most T^{-10} . Thus, taking a union bound over all inductive steps, we get the desired result.

Claim 9 (Claim 22) The probability that step **Augment** is executed at a particular step is at most T^{-10} .

When the step **Augment** is not executed, the distributions AFRW and FRW are identical. Thus, the probability that the distribution $\text{FRW}_{l,\delta}(T)$ achieves a deviation of $T^{1/2+\Theta(\delta)}$ is at least $1/2 - T^{-10}$.

3.2 Unpredictability

In this section we show that the distribution OPT-FRW_{l,δ} is δ -unpredictable.

We first observe that it suffices to work with *aligned* intervals i.e. intervals which start and end at appropriate powers of 2.

Definition 10 (Aligned interval)

We assume here that T is a power of 2. An aligned interval is one which is obtained by breaking [1,T] into 2^i equal parts for $i \in [0, \log T]$ and picking one of the parts. So for instance the first part is always $[1,2^i]$.

In other words, an interval [p+1, p+x] given by $p \in [0,T]$, $x \in [1, T-p]$ is said to be an aligned interval if $p = j \cdot 2^i$ and $x = 2^i$ for some $i \in [0, \log T]$ and $j \in [0, T-2^i]$.

Claim 11 If distribution D(T) is ε -unpredictable with respect to all aligned intervals then it is $c \cdot \varepsilon$ -unpredictable with respect to all intervals, where $c := \frac{\sqrt{2}}{\sqrt{2-1}}$.

The proof of Claim 11 is fairly straightforward and is moved to the appendix (Claim 23).

Theorem 12. The distribution OPT-FRW_{l,δ} is $O(\delta)$ -unpredictable.

Proof: [Sketch]

It can be shown that the process OPT-FRW_{l,δ} has very similar properties if in Step 2 of the construction, instead of changing exactly $\delta \cdot \sqrt{n}$ bits in s_2 we change each bit (of appropriate sign) in s_2 with probability $\frac{\delta}{\sqrt{n}}$. Here we assume this fact without proving it.

We need to show that for every A, s and I, $\mathbb{E}[A_s(I)] \leq O(\delta) \cdot \sqrt{|I|}$ where s and I are as in Definition 1. We may assume that I is an aligned interval (Claim 11).

From the construction it is clear that $\mathbb{E}[A_s(I)]$ is largest when h(s) = |s|or h(s) = -|s| i.e. all the bits before I are of the same sign. Without loss of generality assume $h_s = s$. Also, if there were no prefix (i.e. |s| = 0) then $\mathbb{E}[A_s(I)] = 0$ since the construction is symmetric. To provide an upper bound on $\mathbb{E}[A_s(I)]$ we simply need to bound the expected number of -1's which are changed to +1's due to the existence of s. We will use a simple union bound on the total probability of changing a -1 to a 1 according to the construction. This probability can be split into 2 parts, the first which occurs because of bit sequences immediately preceding I of length less than I and the second because of bit sequences immediately preceding I of length more than I. For sequences of the first kind, the number of bits changed in I is exactly $\delta \cdot \sqrt{l}$ while for sequences of the second kind we may assume that the expected number of bits changed in I is $\frac{\delta \cdot |I|}{\sqrt{l}}$ where l is the length of the bit sequence under discussion. Thus, the total probability is bounded by:-

$$\sum_{i=1}^{\infty} (\min(|I|, 2^i) \cdot \delta) / \sqrt{2^i} = \sum_{i=1}^{\log |I|} \delta \cdot \sqrt{2^i} + \sum_{i=\log |I|+1}^{\infty} \delta \cdot \frac{|I|}{\sqrt{2^i}}$$

Both terms can be bounded by $\delta \cdot \sqrt{|I|} \cdot \sum_{i=0}^{\infty} 1/\sqrt{2^i}$ and so the combined sum is at most $O(\delta) \cdot \sqrt{|I|}$.

4 Deviation upper bound for Adversarial Distributions

In this section we prove that the deviation achieved by OPT-FRW is essentially the best possible for a δ -unpredictable distribution up to a constant factor.

Theorem 13. The highest Root-Mean-Square deviation that can be achieved by a δ -unpredictable distribution on sequences of length T is $\sqrt{T}(1 + O(\delta)) \log T$.

Proof:

Let $\mathcal{D}_{\delta}(T)$ be the set of all δ -unpredictable distributions over sequences of length T, and let $h_n = \max_{D \in \mathcal{D}_{\delta}(n)} \mathbb{E}_{s \sim D}[h(s)^2]$. Clearly, $h_1 = 1$. We need to show that $\sqrt{h_T} = \sqrt{T}(1 + O(\delta)) \log T$.

Let D(T) be a δ -unpredictable distribution which maximizes $\mathbb{E}_{s\sim D}[h(s)^2]$. Given a sequence $s \sim D$, we write $s = s_1 s_2$ where s_1 and s_2 are of length n/2each. Then we have,

$$\begin{aligned} h_n &= \mathbb{E}_{s \sim D}[h(s)^2] = \mathbb{E}[(h(s_1) + h(s_2))^2] \\ &= \mathbb{E}[h(s_1)^2] + \mathbb{E}[h(s_2)^2] + 2\mathbb{E}[h(s_1)h(s_2)] \\ &\leq 2h_{n/2} + 2\sum_{x=0}^{n/2} \Pr[h(s_1) = x] \cdot x \cdot \mathbb{E}[h(s_2) \mid h(s_1) = x] \\ &\leq 2h_{n/2} + 2\delta\sqrt{n/2}\sum_{x=0}^{n/2} \Pr[h(s_1) = x] \cdot |x| \\ &= 2h_{n/2} + \delta \cdot \sqrt{2n} \cdot \mathbb{E}[|h(s_1)|] \\ &\leq 2h_{n/2} + \delta \cdot \sqrt{2n} \cdot \sqrt{\mathbb{E}[h(s_1)^2]} \\ &\leq 2h_{n/2} + \delta \cdot \sqrt{2n} \cdot \sqrt{h_{n/2}} \end{aligned}$$

The first inequality follows from the definition of $h_{n/2}$. The second inequality

follows from the fact that the distribution of s_2 is also δ -unpredictable. Let's substitute, $g_n^2 := h_n/n$. Then $h_{n/2} = (ng_{n/2}^2)/2$ and $\sqrt{h_{n/2}} = \sqrt{n/2}\sqrt{g_n^2/2}$. Thus, we get

$$\begin{array}{rcl} ng_n^2 \leq & ng_{n/2}^2 + \delta n \sqrt{g_{n/2}^2} \\ \Longrightarrow & g_n^2 \leq & g_{n/2}^2 + \delta \sqrt{g_{n/2}^2} & \leq & \left(\sqrt{g_{n/2}^2} + \delta/2\right)^2 \\ \Longrightarrow & g_n \leq & g_{n/2} + \delta/2 \end{array}$$

Since $g_1 = 1$, this gives the upper bound $g_n \leq 1 + (\delta/2) \log n$. This implies $\sqrt{h_n} = \sqrt{\mathbb{E}_{A \sim D}[h_A^2]} \leq \sqrt{n}(1 + \delta/2) \log n$.

$\mathbf{5}$ Acknowledgements

We thank Alex Andoni and Samuel Ieong for useful discussions.

References

1. Frost, A., Prechter, R.: Elliott wave principle: key to market behavior. Bookworld Services (1998)

- Mandelbrot, B.: The inescapable need for fractal tools in finance. Annals of Finance 1(2) (2005) 193–195
- Bradley, B., Taqqu, M.: Financial risk and heavy tails. Handbook of Heavy-Tailed Distributions in Finance (2003) 35–103
- 4. Rachev, S., Menn, C., Fabozzi, F., et al.: Fat-tailed and skewed asset return distributions: Implications for risk management, portfolio selection, and option pricing. Volume 139. Wiley (2005)
- Mandelbrot, B., Van Ness, J.: Fractional brownian motions, fractional noises and applications. SIAM review 10(4) (1968) 422–437
- Nualart Rodón, D.: Fractional brownian motion: stochastic calculus and applications. In: Proceedings on the International Congress of Mathematicians: Madrid, August 22-30, 2006: invited lectures. (2006) 1541–1562
- Mandelbrot, B., Passoja, D., Paullay, A.: Fractal character of fracture surfaces of metals. (1984)
- Davy, P., Sornette, A., Sornette, D.: Some consequences of a proposed fractal nature of continental faulting. Nature 348(6296) (1990) 56–58
- Mandelbrot, B.: Fractals and Chaos: the Mandelbrot set and beyond. Volume 3. Springer (2004)
- Norros, I.: On the use of fractional brownian motion in the theory of connectionless networks. Selected Areas in Communications, IEEE Journal on 13(6) (1995) 953– 962
- Bayraktar, E., Poor, H., Sircar, K.: Estimating the fractal dimension of the s&p 500 index using wavelet analysis. International Journal of Theoretical and Applied Finance 7(05) (2004) 615–643
- 12. Voit, J.: The statistical mechanics of financial markets. Springer (2005)
- 13. Nolan, J.: Stable distributions: models for heavy-tailed data. Birkhauser (2003)
- 14. Rogers, L.: Arbitrage with fractional brownian motion. Mathematical Finance 7(1) (2002) 95–105
- Sottinen, T., Valkeila, E.: On arbitrage and replication in the fractional black– scholes pricing model. Statistics & Decisions/International mathematical Journal for stochastic methods and models 21(2/2003) (2003) 93–108
- Gripenberg, G., Norros, I.: On the prediction of fractional brownian motion. Journal of Applied Probability (1996) 400–410
- 17. Shafer, G., Vovk, V.: Probability and finance: it's only a game! Volume 373. Wiley-Interscience
- Abernethy, J., Frongillo, R., Wibisono, A.: Minimax option pricing meets blackscholes in the limit. In: Proceedings of the 44th symposium on Theory of Computing, ACM (2012) 1029–1040
- DeMarzo, P., Kremer, I., Mansour, Y.: Online trading algorithms and robust option pricing. (2006)
- 20. Black, F., Scholes, M.: The pricing of options and corporate liabilities. The journal of political economy (1973) 637–654
- 21. Littlestone, N., Warmuth, M.: The weighted majority algorithm. FOCS (1989)
- 22. Cover, T.: Behaviour of sequential predictors of binary sequences. Transactions of the Fourth Prague Conference on Information Theory, Statistical Decision Functions, Random Processes (1965)
- Hoeffding, W.: Probability inequalities for sums of bounded random variables. Journal of the American Statistical Association 58(301) (1963) 13–30
- 24. Berry, A.: The accuracy of the gaussian approximation to the sum of independent variates. Transactions of the american mathematical society (1941) 122–136

A Fractal nature of Adversarial Distributions

Here we show that any distribution which is δ -unpredictable must have a fractal like nature (Theorem 6). We will first show that δ -unpredictable distributions are also unpredictable in a slightly stronger sense.

Definition 14 (Adaptive interval algorithm) An interval prediction algorithm is said to be adaptive if it can choose to stop making predictions on interval I at any point within I based on the bits it has seen so far. Note that we do not allow the prediction of the algorithm to depend on the bits in I, the only decision the algorithm can make based on bits in I is to stop predicting earlier than the end point of I.

Definition 15 (Adaptively δ -unpredictable) A distribution D is said to be adaptively δ -unpredictable if for any adaptive algorithm A, sequence of bits s and interval I, $\mathbb{E}[A_s(I)] \leq \delta \cdot \sqrt{l}$ where l is the expected time for which A continues making a prediction in I.

Here the bits in I are produced according to D conditioned on having produced s immediately before I, similarly as in Definition 1.

Theorem 16. A δ -predictable distribution is also adaptively $O(\delta)$ -unpredictable.

Proof:

Let D be a δ -predictable distribution and A' an adaptive interval algorithm. We first show that $\mathbb{E}[A'_s(I)] \leq 2\delta \cdot \sqrt{|I|}$ i.e. we replace the *expected* time for which A' continues making a prediction in I by the *maximum* time for which it makes a prediction.

We will construct a non-adaptive algorithm A such that $\mathbb{E}[|A_s(I) - A'_s(I)|] \leq \delta \cdot \sqrt{|I|}$. Since $\mathbb{E}[A_s(I)] \leq \delta \cdot \sqrt{|I|}$ (D is δ -unpredictable) this implies that $\mathbb{E}[A'_s(I)] \leq 2\delta \cdot \sqrt{|I|}$

Let p_u be the probability of producing a sequence of bits u as a prefix in I according to distribution D. Let E be the set of sequences u such that the algorithm A' stops making predictions on seeing u. Then $\sum_{u \in E} p_u = 1$.

Let $P_u(A)$ denote the expected payoff of A on the remaining part of I conditioned on the event that A' has stopped making predictions. Then $P_u(A) \leq \delta \cdot \sqrt{|I| - |u|} \leq \delta \cdot \sqrt{|I|}$. Thus, $\mathbb{E}[|A_s(I) - A'_s(I)|] \leq \sum_{u \in E} p_u \cdot \delta \cdot \sqrt{|I|} = \delta \cdot \sqrt{|I|}$.

Now we extend the proof to the case where A' makes a prediction for expected time x rather than maximum time x.

Let q_i be the probability that A' makes a prediction for time more thant $2^i x$. By Markov's inequality, $q_i \leq 2^{-i}$. Also, $q_i = \sum_{u \in E: |u|=2^i} p_u$, where p_u is as defined above. We will bound the payoff of A' in phases where the i^{th} phase consists of bits between $2^i x$ to $2^{i+1} x$ from the start of I, and show that it is at most $2\delta \cdot q_i \cdot \sqrt{2^i x}$. For a fixed sequence u, the payoff of algorithm A' in phase i conditioned on having seen u is at most $2\delta \sqrt{2^i x}$ (proved above). Thus, the total payoff of A' in phase i is at most $2\delta \cdot q_i \cdot \sqrt{2^i x}$. Finally, the expected payoff of A' over all phases is at most: $\begin{array}{l} \sum_i 2\delta q_i \sqrt{2^i x} \leq 2\delta \cdot \sum_i \sqrt{2^i x}/2^i \leq O(\delta) \cdot \sqrt{x}) \\ \text{which proves that } D \text{ is adaptively } O(\delta)\text{-unpredictable.} \end{array}$

Now we turn to showing that any adaptively δ -unpredictable distribution has a fractal like nature.

Theorem 17. If a distribution over T bit sequences is adaptively δ -unpredictable (Definition 15) then it is (α, q) -inverting for some constants α, q . Further by dropping the inversion ratio α to $\Theta(1/\log T)$ the probability q can be made as high as $1-1/T^{\Omega(1)}$ for all intervals of length at least $\Omega(\log T)$. Thus the condition holds with high probability simultaneously for all such intervals.

Proof:

For a certain given history of bits consider the interval I. Let h(I) denote the random variable that denotes the height of this interval. Let θ, p be such that the deviation in I exceeds θ with constant probability p (this generalizes the case when θ is the median deviation.)

We will show that some prefixes of I must achieve height at least $\alpha\theta$ and $-\alpha\theta$ each with constant probability (where $\alpha < 1/2$ is a constant). To show this, note that either $h \ge \theta$ or $h \le -\theta$ with probability at least p/2. Assume it is the former without loss of generality. So we only need to prove that $h \le -\alpha\theta$ with probability at least p/4. Assume the contrary and we will see that the interval cannot be δ -unpredictable.

Consider a prediction algorithm that predicts +1 for the interval but adaptively terminates its betting if the height drops to $-\alpha\theta$ or if the height exceeds $2\alpha\theta$, whichever happens first. Since the algorithm hits the lower limit of $-\alpha\theta$ only with probability at most p/4, so with at least probability p/4 it must realize the upper limit (payoff) of $2\alpha\theta$ (since $2\alpha < 1$). In all remaining cases the payoff is at least $-\alpha\theta$. So the expected payoff is at least $(p/4)(2\alpha\theta) - (p/4)(\alpha\theta)$ which needs to be at most $\delta\sqrt{x}$. This is not possible if $\alpha \leq 1/2$ and $\theta = \Omega(\frac{\delta \cdot \sqrt{|I|}}{p})$. Thus if the height in an interval has high magnitude with constant probability, it must reach in either direction with constant probability.

To convert this into a high probability argument, we will use (at most) s iterations of the above prediction algorithm each with limits that depend on θ/s instead of θ . Each iteration has limits of $2\alpha\theta/s$ and $-\alpha\theta/s$ on the sum of bits seen during its execution. The next iteration is initiated only if either of the upper or lower limit is reached in the previous iteration and if not all |I| bits in the full interval are exhausted. From the previous argument, conditioned on the event that a certain iteration is initiated, if an iteration is executed for expected time O(|I|/s) and hits the upper limit with probability p/2 then it must also hit the lower limit with probability p/4. Since the final height exceeds $\alpha\theta$ with constant probability p, in such cases all s iterations have been initiated. Since there are at most s iterations and all are initiated with constant probability, at least half of them must have an expected length of O(|I|/s) conditioned on the event that they are initiated; otherwise the total expected time of all the s iterations will exceed x.

1

Conditioned on the event that the i^{th} iteration is initiated, with probability p it must hit at least one of its two limits; otherwise the total height will not reach $2\alpha\theta$ with probability p. So conditioned on the event that the i^{th} iteration is initiated, for at least half the iterations, it must hit the lower limit (and upper limit) with probability at least p/4. So conditioned on the event that all s iterations are initiated the probability that none of them hit the lower limit and also the upper limit is at most $(p/4)^{s/2}$.

Thus, it follows that by choosing $s = \Theta(1)$, we get an α inversion for constant α with constant probability. This proves the first part of the theorem.

For the second part, note that with probability at least $1 - (p/4)^{s/2}$ either the final height is less than $2\alpha\theta$ or some subinterval has height $-\alpha\theta/s$. For $s = \Theta(\log T)$ the probability that the final height exceeds θ and there is no inversion of height $\leq -\alpha/s\theta$ is negligible.

B Omitted Proofs

Theorem 18. The distribution OPT-FRW_{l,δ}(T) achieves a deviation of $\sqrt{T}(1 + \Omega(\delta \log T))$ with constant probability for $l := T^{-3/4}$.

Proof:

To prove the theorem it will be more convenient to define another process which is similar to OPT-FRW_{l,δ} but which can assume integer values instead of bits.

Augmented Optimum Fractal Random Walk (AOpt-FRW_{l,δ})(2n)

- 1. Generate sequences $s_1, s_2 \in \{-1, 1\}^n$ independently according to AOPT-FRW_{l,\delta}(n)
- 2. If height of s_1 is positive, change exactly $\delta \cdot \sqrt{n} 1$'s in s_2 to 1 (if they exist). Similarly, if height of s_1 is negative, change exactly $\delta \cdot \sqrt{n}$ 1's in s_2 to -1 (if they exist). Call the resulting sequence s'_2 .
- 3. Augment: If there aren't enough (-1)'s to flip in s_2 , then add 2 to some of the numbers so that the increase in height is exactly $\delta \cdot \sqrt{n}$. Similarly for 1's.
- 4. Set $s = s_1 \cdot s'_2$ i.e. the concatenation of s_1 and s'_2

First we observe that when $l = T^{-3/4}$, the probability of executing step **Aug**ment is exponentially small in T. To see this note that if all the base sequences of length l have at least $c(T) := \delta \sqrt{T} \log T$ (-1)'s and at least c(T) 1's then the step **Augment** is never called. This is because every inductive step removes at most $\delta \sqrt{T}$ 1's or -1's at each stage and the number of times a base sequence is modified is at most $\log(T/l) \leq \log T$. Now note that by Chernoff bound, probability that a given base sequence does not have c(T) 1's or (-1)'s is exponentially small in T. Finally note that the number of base sequences is at most T/l, so we can simply take a union bound over all of them.

For brevity, let $D := \text{OPT-FRW}_{l,\delta}$ and $D' := \text{AOPT-FRW}_{l,\delta}$. The next observation is that it suffices to prove that $\mathbb{E}[|h(D')|]$ is $\sqrt{T}(1 + \Omega(\delta \log T))$ and

 $\mathbb{E}[h(D')^2] = O(\mathbb{E}[|h(D')|]^2)$ to prove the theorem. To see this, let NA be the event that the step **Augment** is never executed at any point in the construction, then we have:-

$$\mathbb{E}_{s \sim D}[|h(s)|] \ge \mathbb{E}_{s \sim D}[|h(s)| \mid NA]$$

= $\mathbb{E}_{s \sim D'}[|h(s)|] \mid NA]$
= $\mathbb{E}_{s \sim D'}[|h(s)|] - \Pr[NA] \cdot \max_{s \in D'}|h(s)$

We already saw that $\Pr[NA]$ is exponentially small in T. Note that the maximum value of |h(s)| is at most $T + \delta(\sqrt{T/2} + 2\sqrt{T/4}) + 4\sqrt{T/8} + \ldots + 2^{T/l-1}\sqrt{l}$ which is bounded by a polynomial in T. Thus, if $\mathbb{E}[|h(D')|]$ is $\sqrt{T}(1 + \Omega(\delta \log T))$ then so is $\mathbb{E}[|h(D)|]$. It is also easy to see that the maximum value of $h(s)^2$ is polynomial in T. This fact combined with our assumption about D', $\mathbb{E}[h(D')^2] = O(\mathbb{E}[|h(D')|]^2)$ implies that $\mathbb{E}[h(D)^2] = O(\mathbb{E}[|h(D)|]^2)$. Applying Lemma D to distribution D we get that deviation $\sqrt{T}(1 + \Omega(\delta \log T))$ is achieved with constant probability as required.

So to reiterate, we need to prove two things:-

$$- \mathbb{E}[|h(D')|] \text{ is } \sqrt{T}(1 + \Omega(\delta \log T)) \\ - \mathbb{E}[h(D')^2] = O(\mathbb{E}[|h(D')|]^2)$$

From now on, we denote by S_T a random sequence S drawn from the distribution D'(T). The random variable $h(S_T)$ can be written as $h(A_{T/2}) + h(B_{T/2}) + R$ where R is $\delta\sqrt{T/2}$ if $h(A_{T/2}) > 0$ and $-\delta\sqrt{T/2}$ otherwise. Here the pairs of variables $(A_{T/2}, B_{T/2})$ and $(B_{T/2}, R)$ are independent. Now define $h_T := h(S_T)$. We see that,

$$\begin{split} \mathbb{E}[h_T^2] &= \mathbb{E}[h(S_T)^2] \\ &= \mathbb{E}[(h(A_{T/2}) + h(B_{T/2}) + R)^2] \\ &= \mathbb{E}[h(A_{T/2})^2] + \mathbb{E}[h(B_{T/2})^2] + \mathbb{E}[R^2] + 2\mathbb{E}[h_A R] \\ &= 2\mathbb{E}[h_{T/2}^2] + \delta^2 \sqrt{T/2} + 2\delta \sqrt{T/2}\mathbb{E}[|h_A|] \\ &\geq 2\mathbb{E}[h_{T/2}^2] + \delta \sqrt{2T}\mathbb{E}[|h_{T/2}|] \end{split}$$

The following claim gives a lower bound for $\mathbb{E}[|h_T|]$.

Claim 19

$$\mathbb{E}[|h_T|] \ge \frac{\mathbb{E}[h_T^2]^2}{\mathbb{E}[h_T^4]^{3/4}}$$

Proof: Let the random variables X, Y be defined as $X := |h_T|^{1/2}, Y := |h_T|^{3/2}$. By Cauchy-Schwartz,

$\mathbb{E}[XY]^2 \le$	$\mathbb{E}[X^2] \cdot \mathbb{E}[Y^2]$
$\implies \mathbb{E}[h_T^2]^2 \le$	$\mathbb{E}[h_T] \cdot \mathbb{E}[h_T ^3]$
\leq	$\mathbb{E}[h_T] \cdot \mathbb{E}[h_T^4]^{3/4}$
$\implies \mathbb{E}[h_T] \ge$	$\frac{\mathbb{E}[h_T^2]^2}{\mathbb{E}[h_T^4]^{3/4}}$

Thus, we can say that

$$\begin{split} \mathbb{E}[h_T^2] &\geq 2\mathbb{E}[h_{T/2}^2] + \delta\sqrt{2T}\mathbb{E}[|h_{T/2}|] \\ &\geq 2\mathbb{E}[h_{T/2}^2] + \delta\sqrt{2T}\frac{\mathbb{E}[h_{T/2}^2]^2}{\mathbb{E}[h_{T/2}^4]^{3/4}} \\ &= 2\mathbb{E}[h_{T/2}^2] + \delta\sqrt{2T}\sqrt{\mathbb{E}[h_{T/2}^2]} \left(\frac{\mathbb{E}[h_{T/2}^2]}{\mathbb{E}[h_{T/2}^4]}\right)^{3/4} \end{split}$$

First, let's complete the proof assuming that $\frac{\mathbb{E}[h_T^2]^2}{\mathbb{E}[h_T^4]} \ge C$ for all T where C is an absolute constant. Let's substitute $g_T^2 := \mathbb{E}[h_T^2]/T$. Then,

$$\begin{split} \mathbb{E}[h_T^2] \geq & 2\mathbb{E}[h_{T/2}^2] + \Omega(\delta)\sqrt{2T}\sqrt{\mathbb{E}[h_{T/2}^2]} \\ \Longrightarrow & T \cdot g_T^2 \geq & 2 \cdot (T/2) \cdot g_{T/2}^2 + \Omega(\delta) \cdot \sqrt{2T} \cdot \sqrt{T/2} \cdot \sqrt{g_{T/2}^2} \\ \implies & g_T^2 \geq & g_{T/2}^2 + \Omega(\delta) \cdot g_{T/2} \\ &= & (g_{T/2} + \Omega(\delta))^2 - O(\delta^2) \\ \implies & g_T \geq & g_{T/2} + \Omega(\delta) \end{split}$$

For the base case, we have $\mathbb{E}[h_l^2] = l$, thus $g_l^2 = 1$. Thus,

$$g_T \ge 1 + \Omega(\delta) \cdot \log(T/l) = 1 + \Omega(\delta) \cdot \log T^{1/4} = 1 + \Omega(\delta \log T)$$

Thus, $\mathbb{E}[h_T^2] \ge T \cdot g_T^2 = T(1 + \Omega(\delta \log T)^2)$. By Lemma 19 and Lemma 2, this implies $\mathbb{E}[|h_T|] \ge \sqrt{T}(1 + \Omega(\delta \log T))$. These statements together imply both the guarantees we set out to prove about D'.

It remains to prove the following lemma.

Lemma 2. $\frac{\mathbb{E}[h_T^2]^2}{\mathbb{E}[h_T^4]} \ge C$ for all T where C is an absolute constant. **Proof:** Recall that for S_T drawn according to AOPT-FRW_{l,δ}(T), we have $h(S_T) = h(A_{T/2}) + h(B_{T/2}) + R$. We already saw that $\mathbb{E}[h_T^2] \ge 2\mathbb{E}[h_{T/2}^2]$. Let $r_T := \frac{\mathbb{E}[h_T^4]}{\mathbb{E}[h_T^2]^2}$. We need to show that $r_T \le C$. We have,

$$r_T = \frac{\mathbb{E}[h_T^4]}{\mathbb{E}[h_T^2]^2} \ge \frac{\mathbb{E}[h_T^4]}{4\mathbb{E}[h_{T/2}^2]^2}$$

Now, let's write a recurrence for $\mathbb{E}[h_T^4]$.

$$\begin{split} \mathbb{E}[h_T^4] &= \mathbb{E}[h(S_T^4)] \\ &= \mathbb{E}[(h_A + h(s_1) + R)^4] \\ &= \mathbb{E}[h_A^4] + \mathbb{E}[h(s_1)^4] + \mathbb{E}[R^4] + 6\mathbb{E}[h_A^2h(s_1)^2] + 6\mathbb{E}[h_A^2R^2] + \\ 6\mathbb{E}[h(s_1)^2R^2] + 4\mathbb{E}[h_AR^3] + 4\mathbb{E}[h_A^3R] \\ &= 2\mathbb{E}[h_{T/2}^4] + \delta^4(T/2)^2 + 6\mathbb{E}[h_{T/2}^2]^2 + 12\delta^2(T/2)\mathbb{E}[h_{T/2}^2] + \\ 4\delta^3(T/2)^{3/2}\mathbb{E}[|h_{T/2}|] + 4\delta\sqrt{T/2}\mathbb{E}[|h_{T/2}|^3] \\ &\leq 2\mathbb{E}[h_{T/2}^4] + O(\delta^4T^2) + 6\mathbb{E}[h_{T/2}^2]^2 + O(\delta^2T\mathbb{E}[h_{T/2}^2]) + \\ O(\delta^3T^{3/2})\sqrt{\mathbb{E}[h_{T/2}^2]} + O(\delta\sqrt{T})\mathbb{E}[h_{T/2}^4]^{3/4} \end{split}$$

Dividing both sides by $4\mathbb{E}[h_{T/2}^2]^2$ and using the fact that $\mathbb{E}[h_T^4] \ge \mathbb{E}[h_T^2]^2 \ge T^2$, we get:-

$$r_T \leq \frac{\mathbb{E}[h_T^4]}{4\mathbb{E}[h_{T/2}^2]^2} \leq (1/2) \cdot r_{T/2} + O(\delta^4) + (3/2) + O(\delta^2) + O(\delta^3) + O(\delta)r_{T/2}^{3/4} \leq (3/4) \cdot r_{T/2} + O(1)$$

which is clearly bounded above by an absolute constant for all T.

Thus, the theorem is proved.

-

Theorem 20. The distribution $D := \operatorname{FRW}_{l,\delta}$ is $O(\delta)$ -unpredictable in a weak sense i.e. $\mathbb{E}_{s,I}[A_s(I)] \leq O(\delta) \cdot h_{|I|}$ where $h_n := \mathbb{E}_{s \sim D(n)}[|h(s)|]$. Here s, I and A are as in Definition 1. Note that the expectation on the left is taken over I as well as the prefix s as opposed to Definition 1 where s is fixed and the expectation is over I only.

Proof: [Sketch]

It can be shown that the process $\operatorname{FRW}_{l,\delta}$ has very similar properties if in Step 2 of the construction, instead of changing exactly $\delta \cdot h(s_1)$ bits in s_2 we change each bit (of appropriate sign) in s_2 with probability $\frac{\delta \cdot h(s_1)}{n}$. Here we assume this fact without proving it.

We need to show that $\mathbb{E}_{s,I}[A_s(I)] \leq O(\delta) \cdot h_{|I|}$. We may assume that I is an aligned interval (Claim 11). Let s(i) be the suffix of length i in s. Then,

$$\begin{split} & \mathbb{E}_{s}[\mathbb{E}_{I}[A_{s}(I)] \\ & \leq \sum_{i=0}^{\log|I|} \delta \cdot \mathbb{E}_{s}[|h(s(2^{i}))|] + \sum_{i=\log|I|+1}^{\infty} \frac{\delta \cdot \mathbb{E}_{s}[|h(s(2^{i})|] \cdot |I|]}{2^{i}} \\ & \leq O(\delta) \cdot \mathbb{E}_{s}[|h(s(|I|))|] + \\ & O(\delta) \cdot \mathbb{E}_{s}[|h(s(|I|))|](1/2^{1/4} + 1/4^{1/4} + \ldots) \\ & \leq O(\delta) \cdot \mathbb{E}_{s}[|h(s(|I|))|] \\ & = O(\delta) \cdot h_{|I|} \end{split}$$

where the second inequality uses $h_T = T^{1/2 + \Theta(\delta)} \leq T^{3/4}$ (Theorem 7).

Claim 21 For $n = 2^i \cdot l$, $S \sim AFRW_{l,\delta}(n)$,

$$h(S) = \sum_{U \subseteq [i]} r^{|U|} h(X_U)$$
(B.1)

-

where $r = (1 + \delta)$ and each X_U is independently and uniformly distributed in $\{-1, 1\}^l$.

Proof: We will prove the claim by induction on i. For i = 0, the claim clearly holds.

Assume that the claim holds for i = k, and let $n := 2^{i+1} \cdot l$. Let $S = S_1 \cdot S'_2$ be the sequence produced by the distribution as described above where S_1 and S'_2 are random sequences of length n/2 each. Because of step **Augment**, it is clear that $h(S'_2) = h(S_2) + \delta h(S_1)$ which means $h(S) = (1 + \delta)h(S_1) + h(S_2)$. Thus,

$$h(S) = rh(S_1) + h(S_2)$$

= $\sum_{U \subseteq [k]} r(r^{|U|}h(X_U)) + \sum_{V \subseteq [k]} r(r^{|V|}h(X_V))$
= $\sum_{U \subseteq [k]} r^{|U \cup \{k+1\}|}h(X_{U \cup \{k+1\}}) + \sum_{V \subseteq [k]} r(r^{|V|}h(X_V))$
= $\sum_{U \subseteq [k+1]} (r^{|U|}h(X_U))$

Lemma 3. Median of $|h(AFRW_{l,\delta})|$ is $n^{1+\Omega(\delta)}$.

Proof: In the notation of Theorem 28 we think of each term in Equation B.1 as a random variable. There are exactly 2^i terms. It is clear that $\mathbb{E}[X_S] = 0$ for all $S \subseteq [i]$. Also, $\mathbb{E}[X_S^2] = r^{2|S|} \cdot l$ and $\mathbb{E}[|X_S^3|] = r^{3|S|} \cdot l$,

$$\sigma^2 := \sum_S r^{2|S|} \cdot l = (r^2 + 1)^i \cdot l \approx (2 + 2\delta)^i \cdot l \approx n^{1 + \Theta(\delta)}$$

Also, $\max_S \rho_S / \sigma_S = \max_S r^{|S|} = r^i \approx n^{\Theta(\delta)}$. Thus, we have

$$|S - N(0, \sigma^2)| \le n^{-1/2} \cdot n^{O(\delta)} \le n^{-\Omega(1)}$$

Thus, the distribution of $|h(\text{AFRW}_{l,\delta})|$ is very close to a half-normal distribution with variance σ^2 and thus the median of $|h(\text{AFRW}_{l,\delta})|$ is $\Omega(\sigma) = n^{1/2+\Omega(\delta)}$.

Claim 22 The probability that step Augment is executed at a particular step is at most T^{-10} .

Proof: This is a simple application of Theorem 27.

Let's say we are at the step where the length of the sequences is $n := 2^i \cdot l$. We consider random variables $Y_S := r^{|}S| \cdot X_S$ where X_S is as in Lemma 3. Observe that a single random variable Y_S is actually a sum of l independent random variables each of which take values in $\{-r^{|S|}, r^{|S|}\}$. Let us denote these random variables as $Y_{S,i}$ so that $X_S = \sum_{i=1}^{l} Y_{S,i}$. Thus, in the notation of Theorem 27,

$$\sum_{S,i} (b_{S,i} - a_{S,i})^2 = 4 \sum_{S,i} r^{2|S|} = 4l \sum_S r^{2|S|} = n^{1+O(\delta)}$$

where σ^2 is as in Lemma 3.

The step **Augment** is executed only when $h_S \ge n - \delta n$ or $h_S \le -(n - \delta n)$. Thus, we have the bound

$$\Pr[|h_S| \ge n - \delta n] \le \Pr[|h_S| \ge n/2] \le 2 \cdot \exp\left(-\frac{n^2}{n^{1+O(\delta)}}\right) \le 2 \cdot \exp\left(-\frac{l^2}{l^{1+O(\delta)}}\right) \le T^{-10}$$

as desired.

Claim 23 If distribution D(T) is ε -unpredictable with respect to all aligned intervals then it is $c \cdot \varepsilon$ -unpredictable with respect to all intervals, where $c := \frac{\sqrt{2}}{\sqrt{2-1}}$.

Proof: Consider an interval I of size x. If I is an aligned interval we are done, otherwise we write it as the minimal union of aligned intervals (take out the largest aligned interval in I and repeat). There are three possibilities:-

1. $I = I_1 \cup I_2$ is a union of two intervals of size x/2 each (eg. the interval [T/4 + 1, 3T/4])

- 2. $I = I_1 \cup I_2 \cup \ldots \cup I_k$, where each I_j is of a different size. Note that all interval sizes on the right are powers of 2 and strictly less than x
- 3. $I = J \cup J'$ where each J can be written as a union of intervals as in 1 or 2 above

In the first case,

$$|\mathbb{E}[h_I]| \le |\mathbb{E}[h_{I_1}]| + |\mathbb{E}[h_{I_1}]| \le 2 \cdot \varepsilon \cdot \sqrt{x/2} = \sqrt{2} \cdot \varepsilon \cdot \sqrt{x}$$

In the second case,

$$|\mathbb{E}[h_I]| \le \sum_{j=1}^k |\mathbb{E}[h_{I_j}]| \le \varepsilon \cdot \sqrt{x} \cdot \sum_{j=1}^\infty \sqrt{1/2^j} = \frac{1}{\sqrt{2}-1} \cdot \varepsilon \cdot \sqrt{x}$$

In the third case,

$$\begin{split} \mathbb{E}[h_I]| &\leq |\mathbb{E}[h_J]| + |\mathbb{E}[h'_J]| \leq \frac{1}{\sqrt{2} - 1} \cdot \varepsilon \cdot \sqrt{|J|} + \\ &\frac{1}{\sqrt{2} - 1} \cdot \varepsilon \cdot \sqrt{|J'|} \leq \frac{\sqrt{2}}{\sqrt{2} - 1} \cdot \varepsilon \cdot \sqrt{|I|} \end{split}$$

C Fractal nature of deterministic inverting sequences

We will argue that the optimal sequence with height h and inversion ratio α is achieved by the following fractal-like recursive process. To construct a sequence of height h, recursively generate a sequence s_1 of height $(1 + \alpha/2) \cdot h$ and s_2 of height $\alpha \cdot h$ respectively. Concatenate s_1 , an inverted copy of s_2 followed by another copy of s_1 . For simplicity for explanation we will ignore rounding errors from the discretization.

It turns out that for large h, the ratio of lengths of s_1 and s_2 is fixed to $\left(\frac{1+\alpha}{2}\right)^{1/\theta} : \alpha^{1/\theta}$ where θ is a constant defined below.

Claim. The above process produces an α -inverting sequence for α smaller than some constant.

Proof: Observe that by recurrence any interval that is contained within s_1 or s_2 is α -inverting. The full interval consisting of the three concatenated strings also has an α -inversion; and so are the intervals that span the first two and the last two strings. So we only need to argue about intervals that span parts of multiple of these pieces. Consider for example an interval that spans across some suffix of s_1 and some prefix of the inverted copy of s_2 . Now for small enough α , the two parts of the interval have heights of opposite signs. So the α -inversion in the piece with the larger absolute height suffices to produce an α -inversion in the interval. The same argument can be applied for intervals that span part of the first and the third sequence.

Claim. Let s be an α -inverting sequence of length t (Definition 2), where α is bounded above by a constant. Then the highest deviation that can be achieved by s for large t is t^{θ} where θ is the solution to the equation $1 = 2((1+\alpha)/2)^{1/\theta} + (\alpha)^{1/\theta}$. Furthermore, this deviation is actually achieved by the above process.

Proof: [Sketch] We will compute the amount of time t(h) when the process described above first achieves a height h > 0. By the construction, t(h) satisfies the recurrence $t(h) = 2t \cdot ((1 + \alpha)h/2) + t(\alpha h)$. In the limit, if this recurrence has a solution of the form $h^{1/\theta}$ then note that $h^{1/\theta} = 2((1 + \alpha)h/2)^{1/\theta} + (\alpha h)^{1/\theta}$ which means that $1 = 2((1 + \alpha)/2)^{1/\theta} + (\alpha)^{1/\theta}$. The proof can be formalized by sandwiching the solution to the recurrence in the limit between the functions h^{1/θ_1} and h^{1/θ_2} where θ_1 and θ_2 approach θ from above and below.

To prove the lower bound, let t(h) denote the required time to produce a height of absolute value h for any α -inverting sequence. We will prove that for large h, t(h) approaches $h^{1/\theta}$. We know that for large enough t there must be an inversion with ratio α . So to achieve height h in time t there must be a sub-interval with height less than $-\alpha h$. So t can be broken into three segments of lengths t_1 , t_2 , t_3 with heights h_1 , h_2 , h_3 such that $h = h_1 + h_2 + h_3$ where $h_2 \leq -\alpha h$. We wish to minimize $t(h) = t_1 + t_2 + t_3 \geq t(h_1) + t(h_2) + t(h_3)$. Since t(h) is non-decreasing in h, we may set $h_2 = -\alpha h$ and $h_1 + h_3 = h - h_2 = (1+\alpha)h$ giving $t(h) = \min t(h_1) + t(h_2) + t(\alpha h)$ where $h_1 + h_3 = (1 + \alpha)h$.

Note that if t(h) is of the form $h^{1/\theta}$ then it is convex and so $t(h_1) + t(h_2)$ is minimized when $h_1 = h_3 = \frac{(1+\alpha)\cdot h}{2}$ giving $t(h) = 2t(\frac{(1+\alpha)\cdot h}{2}) + t(\alpha h)$ whose solution approaches $h^{1/\theta}$ in the limit. That the solution must approach $h^{1/\theta}$, by looking at the behavior of $\log_t h$ in the limit and sandwiching it between θ_1 and θ_2 that approach θ from above and below.

D Miscellaneous Observations

Observation 24 A uniform random sequence is (α, q) inverting (Definition 5) for some constants α, q . Further the probability parameter q can be made as high as $1 - \varepsilon$ by reducing the inversion ratio α to $\Theta(1/\log(1/\varepsilon))$.

Proof: Let us divide the interval of length x into two halves of length x/2 each. With probability 1/2 the two parts have opposite heights and with constant probability both heights have magnitude $\Theta(\sqrt{x})$. Thus it has an α -inversion with some constant probability for some constant α . The higher probability statement is obtained similarly by dividing it into $\log(1/\varepsilon)$ intervals of equal length.

Observation 25 If a string is sampled from the highest entropy distribution with deviation $k\sqrt{T}$, then it is possible to get an expected payoff of $\Omega(1) \cdot k\sqrt{T}$ for $k = \Omega(1)$.

Proof: [Sketch] The algorithm simply predicts the sign of h(s) where s is the sequence seen in the first half i.e. |s| = T/2. A simple computation proves the observation.

The following theorem shows that the FBM with $H = 1/2 + \delta$ is not $O(\delta)$ unpredictable. In fact, an algorithm can get an expected payoff of $\Theta(x^H)$ on an interval of size x by predicting the sign of the height of the preceding interval of length x. (It can also be shown that one cannot do better than this if one is only allowed to use the sign of the height of some preceding interval.)

Claim. The algorithm that predicts an interval of length x using the sign of the height of the preceding interval of length x gets an expected payoff of $\Theta(x^H)$ where the expectation is taken over all values in the preceding interval. Further it is optimal to use a preceding interval of length x if one is using the sign of its height.

Proof. $E[B_H(sx)|B_H(x)]/B_H(x) = (1/2)(s^{2H} + 1 - |s - 1|^{2H})$ (See [5], Section 5.3)

Let us compute the expected payoff if one uses the height of the preceding interval of length x to predict the following interval of length x.

 $E[B_H((s+1)x)|B_H(sx)]/B_H(sx) = (1/2)((1+1/s)^{2H} + 1 - (1/s)^{2H}). E[B_H((s+1)x)]/B_H(sx)] = (1/2)((1+1/s)^{2H} + 1 - (1/s)^{2H}).$ $1(x) - B_H(sx)[B_H(sx)] = (1/2)((1+1/s)^{2H} - 1 - (1/s)^{2H})B_H(sx)$. So by predicting the sign of $B_H(sx)$ to predict the following interval of length x the expected payoff is $E[sign(B_H(sx))B_H(sx)] = (1/2)((1+1/s)^{2H} - 1 - (1/s)^{2H})E[|B_H(sx)|] = (1/2)((1+1/s)^{2H})E[|B_H(sx)|] = (1/2)((1+1/s)^{2H} - 1 - (1/s)^{2H})E[|B_H(sx)|] = (1/2)((1+1/s)^{2H})E[|B_H(sx)|] = (1/2)((1+1/s)^{2H} - 1 - (1/s)^{2H})E[|B_H(sx)|] = (1/2)((1+1/s)^{2H})E[|B_H(sx)|] = (1$ $\Theta(sx)^H(1/2)((1+1/s)^{2H}-1-(1/s)^{2H})$

Note that for s = 1, this is $\Theta(x^H)$. Further this is the best possible value of the above expression.

Observation 26 With continuous prediction the FBM and its binary (discretized) variants have a payoff of $\Omega(\delta T)$

Proof: Observe that if we take a sequence of length 2 the second bit is correlated to the first by $\Theta(\delta)$. This is true of every even bit. The observation follows for the binary variants. For the true FBM the statement holds since if B_1 and B_2 are the heights in two adjacent unit intervals of the FBM process with hurst coefficient $H = 1/2 + \Theta(\delta)$ then

 $E[B_1 + B_2 | B_1]/B_1 = (1/2)(2^{2H} + 1 - 1^{2H}) = 2^{\Theta(\delta)}$ (See [5], Section 5.3)

Therefore $E[B_2 \mid B_1] = (2^{\Theta(\delta)} - 1) \cdot B_1 = \Theta(\delta) \cdot B_1$ for $\delta \leq 1$. So again by predicting the sign of B_1 one can get a payoff of $\Theta(\delta) \cdot B_1 \cdot sign(B_1) =$ $\Theta(\delta) \cdot |B_1|$. This in expectation is $\Theta(\delta)$ as B_1 is normally distributed with constant variance.

Claim. For any random variable X that only takes non negative values and $E[X^2] = O((E[X])^2, Pr[X \ge \Omega(E[X]) = \Omega(1))$

Proof: Let $\mu = E[X]$. The the standard deviation $\sigma = O(\mu) = c\mu$ (say) where c is at most some constant. We will bound $E[X|X > \mu + rc\mu]$ for any $r \in \mathbb{N}$. Note that $Pr[X \ge \mu + rc\mu] \le 1/r^2$.

So $E[X|X \ge \mu + rc\mu] \le (\mu + rc\mu) + \mu \sum_{i>r} 1/i^2 \le (\mu + rc\mu) + c\mu/r$. Now $\mu = E[X] = Pr[X < \mu + rc\mu]E[X|X < \mu + rc\mu] + Pr[X \ge \mu + rc\mu]$ $rc\mu E[X|X \ge \mu + rc\mu] \le (1 - 1/r^2)E[X|X < \mu + rc\mu] + (1/r^2)(\mu + rc\mu + c\mu/r).$ By setting r to be a constant that is at least some large multiple of c, we can conclude that $E[X|X < \mu + rc\mu] = \Omega(\mu)$. So this conditioned random variable X has maximum value and mean value that are the same up to constant factors. Thus it must exceed $\Omega(\mu)$ with constant probability. So the unconditioned random variable X must also exceed $\Omega(\mu)$ with a smaller constant probability.

E Basic tools

Theorem 27 (Hoeffding's bound). [23]

Let X_1, X_2, \ldots, X_n be independent random variables such that $\mathbb{E}[X_i] = 0$ and $\Pr[X_i \in [a_i, b_i]] = 1$. Let $S := \sum_i X_i$. Then,

$$\Pr[|S| \ge y] \le 2 \cdot \exp\left(-\frac{2y^2}{\sum_{i=1}^n (b_i - a_i)^2}\right)$$

Theorem 28 (Berry-Esseen Theorem). [24]

Let X_1, X_2, \ldots, X_n be independent random variables such that $\mathbb{E}[X_i] = 0$, $\mathbb{E}[X_i^2] = \sigma_i^2 > 0$, and $\mathbb{E}[|X_i^3|] = \rho_i < \infty$. Let $\sigma^2 := \sum_i \sigma_i^2$ and $S := \frac{1}{\sigma} \sum_i X_i$. Then there is an absolute constant C such that

$$|S - N(0, \sigma^2)| \le \frac{C}{\sigma} \cdot \max_i \frac{\rho_i}{\sigma_i}$$

Here $|D-D'| := \max_x |\Pr[D \ge x] - \Pr[D' \ge x]|$ denotes the statistical distance between distributions D and D' and $N(\mu, \sigma^2)$ denotes the normal distribution with mean μ and variance σ^2 .