Sum-of-squares lower bounds for planted clique

Raghu Meka Department of Computer Science University of California, Los Angeles Aaron Potechin^{*} Massachusetts Institute of Technology

Avi Wigderson[†] Institute for Advanced Study, Princeton

Abstract

Finding cliques in random graphs and the closely related "planted" clique variant, where a clique of size k is planted in a random G(n, 1/2) graph, have been the focus of substantial study in algorithm design. Despite much effort, the best known polynomial-time algorithms only solve the problem for $k = \Theta(\sqrt{n})$.

In this paper we study the complexity of the *planted clique* problem under algorithms from the Sum-Of-Squares hierarchy. We prove the first average case lower bound for this model: for almost all graphs in G(n, 1/2), r rounds of the SOS hierarchy cannot find a planted k-clique unless $k \ge (\sqrt{n}/\log n)^{1/r}/C^r$. Thus, for any constant number of rounds planted cliques of size $n^{o(1)}$ cannot be found by this powerful class of algorithms. This is shown via an integrability gap for the natural formulation of maximum clique problem on random graphs for SOS and Lasserre hierarchies, which in turn follow from degree lower bounds for the Positivestellensatz proof system.

We follow the usual recipe for such proofs. First, we introduce a natural "dual certificate" (also known as a "vector-solution" or "pseudo-expectation") for the given system of polynomial equations representing the problem for every fixed input graph. Then we show that the matrix associated with this dual certificate is PSD (positive semi-definite) with high probability over the choice of the input graph. This requires the use of certain tools. One is the theory of association schemes, and in particular the eigenspaces and eigenvalues of the *Johnson scheme*. Another is a combinatorial method we develop to compute (via traces) norm bounds for certain random matrices whose entries are highly dependent; we hope this method will be useful elsewhere.

1 Introduction

1.1 The problem and main result

Finding cliques in random graphs has been the focus of substantial study in algorithm design. Let G(n, p) denote Erdös-Renyi random graphs on n vertices where each edge is kept in the graph with probability p. It is easy to check that in a random graph $G \leftarrow G(n, 1/2)$, the largest clique has size $(2 + o(1)) \log_2 n$ with high probability. On the other hand, the best known polynomial-time algorithms can only find cliques of size $(1 + o(1)) \log_2 n$ and obtaining better algorithms remains a

*Supported in part by an NSF Graduate Research Fellowship under grant No. 0645960 and by Microsoft Research

[†]Supported in part by NSF Expeditions grant CCF-0832797

longstanding open problem: Karp [Kar76] suggested that even finding cliques of size $(1 + \varepsilon) \log_2 n$ could require superpolynomial time.

Motivated by this, much attention has been given to the related *planted clique problem* or hidden clique problem introduced by Jerrum [Jer92] and Kucera [Kuc95]. Here, we are given a graph $G \leftarrow G(n, 1/2, k)$ generated by first choosing a G(n, 1/2) random graph and placing a clique of size k in the random graph for $t \gg \log_2 n$. The goal is to recover the hidden clique for as small a k as possible given G. The study of the planted clique problem and its variations (like finding planted dense subgraphs) is motivated from several other more recent directions. Its potential as being hard on average has lead to proposals to base crypto systems on variants of it [ABW10]. It was used to argue that testing k-wise independence is hard near the information theoretic limit by [AAK⁺07]. It is used in [ABBG10] to argue that evaluating some financial derivatives is hard. It was also used to justify the hardness of sparse principal component detection by Bethet and Rigollet [BR13]. Another source of interest comes from the related algorithmic problem of finding large communities in social networks. The best known polynomial-time algorithms can solve the problem for $k = \Theta(\sqrt{n})$ [AKS98] (see [DGGP14] for a near linear-time algorithm) and improving on this bound has received significant attention. The algorithmic problem has also been of much interest in the context of signal finding in molecular biology (pattern discovery in DNA sequences) as modeled in the work of $[PS^+00]$.

In this work we exhibit a lower bound for the problem in the powerful Lasserre [Las01] and "sum-of-squares" (SOS) [Par00] semi-definite programming hierarchies¹. As it happens, proving such lower bounds for the planted clique problem reduces easily to proving an integrality gap of value k for the natural formulation of the maximum clique problem in these hierarchies on G(n, 1/2)graphs. Our main result then is the following average-case lower bound for maximum clique. We defer the formal definition of the semi-definite relaxation and hierarchies for now, and only note a few facts. First, that implementing the rth level of the SOS hierarchy (namely, r rounds), takes roughly $n^{O(r)}$ time, which is polynomial for constant r. Second, the above algorithm for $k = \Theta(\sqrt{n})$ may be viewed as implementing only one round. Third, that $r = \log n$ suffices for exact solution of the problem, namely finding the maximum clique. Our lower bound implies that polynomial time (when the number of rounds r is constant) cannot handle even $k = n^{o(1)}$, and that as many as $(\log n)^{1/2}$ rounds cannot handle $k = (\log n)^{O(1)}$. Here are more precise statements².

Theorem 1.1. With high probability, for $G \leftarrow G(n, 1/2)$ the natural r-round SOS relaxation of the maximum clique problem has an integrality gap of at least $n^{1/2r}/C^r(\log n)^2$.

As a corollary we obtain the following lower bound for the planted clique problem.

Corollary 1.2. With high probability, for $G \leftarrow G(n, 1/2, t)$ the natural r-round SOS relaxation of the planted clique problem has an integrality gap of at least $n^{1/2r}/tC^r(\log n)^2$.

1.2 Background and related work

Linear and semi-definite hierarchies are one of the most powerful and well-studied techniques in algorithm design. The most prominent of these are the Sherali-Adams hierarchy (SA) [SA90], Lovasz-Schrijver hierarchy (LS) [LS91], their semi-definite versions SA_+ , LS_+ and Lasserre and SOS

¹For brevity, in the following, we will use SOS hierarchy as a common term for the formulations of Lasserre [Las01] and Parrilo [Par00] which are essentially the same in our context.

²Throughout, c, C denote constants.

hierarchies. The hierarchies present progressively stronger convex relaxations for combinatorial optimization problems parametrized by the number of rounds r, where the r-round relaxation can be solved in $n^{O(r)}$ time on instances of size n in all of them. In terms of relative power (barring some minor technicalities about how the numbering of rounds starts), it is known that $LS_+(r) < SA_+(r) < SOS(r)$. Because they capture most powerful techniques for combinatorial optimization, lower bounds for hierarchies serve as strong unconditional evidence for computational hardness. Such lower bounds are even more relevant and compelling in situations where we do not have NP-hardness results, as is the case for typical average-case optimization problems.

Broadly speaking, our understanding of the SOS hierarchy is more limited than those of LS_+ and SA_+ hierarchies and in fact the SOS hierarchy appears to be much more powerful. A particularly striking example of this phenomenon was provided by a recent work of Barak et al. [BBH⁺12]. They showed that a constant number of rounds of the SOS hierarchy can solve the much studied *unique games problem* on instances which need super constant number of LS_+ , SA_+ rounds. It was also shown by the works of [BRS11, GS11] that the SOS hierarchy captures the sub-exponential algorithm for unique games of [ABS10]. These results emphasize the need for a better understanding of the power and limitations of the SOS hierarchy.

From the perspective of proving limitations, all known lower bounds for the SOS hierarchy essentially have their origins in the works of Grigoriev [Gri01b, Gri01a], some of which were later independently rediscovered by Schoenebeck [Sch08]. These works show that even $\Omega(n)$ rounds of SOS hierarchy cannot solve random 3XOR or 3SAT instances, implying a strong unconditional average-case lower bound for a natural distribution.

Most subsequent lower bounds for SOS hierarchy such as those of [Tul09], [BCV⁺12] rely on [Gri01b] and [Sch08] and gadget reductions. For example, Tulsiani [Tul09] shows that $2^{O(\sqrt{\log n})}$ rounds of SOS has an integrality gap of $n/2^{O(\sqrt{\log n})}$ for maximum clique in worst-case. This is in stark contrast to the average-case setting: even a single round of SOS gets an integrality gap of at most $O(\sqrt{n})$ for maximum clique on G(n, 1/2) [FK00]. Thus, the worst-case and average-case problems have very different complexities. Finally, using reductions tend to induce distributions that are far from uniform and definitely not as natural as G(n, 1/2).

For max-clique on random G(n, 1/2) graphs, Feige and Krauthgamer [FK00] showed that $\mathsf{LS}_+(r)$, and hence $\mathsf{SOS}(r)$, has an integrality gap of at most $\sqrt{n}/2^{\Omega(r)}$ with high probability. Complementing this, they also showed [FK03] that the gap remains $\sqrt{n}/2^r$ for $\mathsf{LS}_+(r)$ with high probability. However, there were no non-trivial lower bounds known for the stronger SOS hierarchy.

For the planted clique problem, other algorithmic techniques were studied. Jerrum [Jer92] showed that a broad class of Markov chain Monte-Carlo (MCMC) based methods cannot solve the problem when the planted clique has size $O(n^{1/2-\delta})$ for any constant $\delta > 0$. Another approach for the planted clique problem based on optimizing a third order tensor was suggested by Frieze and Kannan [FK08]. However, the corresponding optimization problem is NP-hard in the worst-case.

In a recent work, Feldman et al. [FGR⁺13] introduced the framework of statistical algorithms which generalizes many algorithmic approaches like MCMC methods and showed that such algorithms cannot find large cliques when the planted clique has size $O(n^{1/2-\delta})$ in less than $n^{\Omega(\log n)}$ time³. However, their framework seems quite different from hierarchy based algorithms. In particular, the statistical algorithms framework is not applicable to algorithms which first pick a sample, fix it, and then perform various operations (such as convex relaxations) on it, as is the case for the

³The results of $[FGR^{+13}]$ actually apply to the harder *bipartite* planted clique problem, but this assumption is not too critical.

hierarchies above.

Meka and Wigderson [MW13] addressed SOS lower bounds for planted clique and claimed a stronger bound than Thm 1.1. While there was a fatal error in their proof, many of the techniques introduced there are used in the present paper.

Independent of our work, Deshpande and Montanari [DM15] recently gave a degree 4 SOS lower bound for planted clique; while they are only able to handle the degree 4 case (i.e., r = 2), they obtain a better bound for this case than us (roughly $n^{1/3}$ vs $n^{1/4}$ as we do).

1.3 Proof systems and SDP hierarchies

A potentially simpler problem than deciding is a large clique exists is the problem of producing short certificates to the non-existence of such cliques. This puts the problem in the realm of proof complexity. Indeed, we approach the problem of SOS lower bounds from this viewpoint, via the *positivstellensatz* proof system perspective of Grigoriev and Volobjov [GV01]. We explain this proof system next in general, and then specialize to Boolean problems and specifically to planted clique.

Suppose we are given a system of polynomial equations or "axioms"

$$f_1(x) = 0, \ f_2(x) = 0, \ \dots, \ f_m(x) = 0,$$

where each $f_i : \mathbb{R}^n \to \mathbb{R}$ is a *n*-variate polynomial. A positivstellensatz refutation of the system $\mathcal{F} = ((f_i))$ is an identity of the form

$$\sum_{i=1}^{m} f_i g_i \equiv 1 + \sum_{i=1}^{N} h_i^2,$$

where $\{g_1, \ldots, g_m\}$ and $\{h_1, \ldots, h_N\}$ are arbitrary *n*-variate polynomials. Clearly, if there exists an identity as above, then the system \mathcal{F} has no solution over reals. Starting with the seminal work of Artin on Hilbert's seventeenth problem [Art27], a long line of important results in real algebraic geometry – [Kri64, Ste73, Put93, Sch91]; cf. [BCR98] and references therein – showed that, under some (important) technical conditions⁴, such certifying identities *always exist* for an infeasible system. This motivates the following notion of complexity for refuting systems of polynomial equations.

Definition 1.3 (Positivstellensatz Refutation, [GV01]). Let $\mathcal{F} \equiv \{f_1, \ldots, f_n : \mathbb{R}^n \to \mathbb{R}\}$, be a system of axioms, where each f_i is a real n-variate polynomial. A positivstellensatz refutation of degree r (PS(r) refutation, henceforth) for \mathcal{F} is an identity of the form

$$\sum_{i=1}^{m} f_i g_i \equiv 1 + \sum_{i=1}^{N} h_i^2, \tag{1.1}$$

where $g_1, \ldots, g_m, h_1, \ldots, h_N$ are n-variate polynomials such that $deg(f_ig_i) \leq 2r$ for all $i \in [m]$ and $deg(h_j) \leq r$ for all $j \in [N]$.

⁴We avoid going into the details here as the conditions are easily met in the presence of Boolean axioms.

Our interest in positivstellensatz refutations as above comes from the known relations between such identities and SOS hierarchy. Informally (and under appropriate technical conditions), identities as above of degree r show that SOS hierarchy can *certify* infeasibility of the axioms in $2r + \Theta(1)$ rounds and vice versa. We will focus on showing degree lower bounds for identities as above and use them to get integrality gaps for the the SOS hierarchy. We formalize this in Section 12. For a brief history of the different formulations from [GV01], [Las01], [Par00] and the relations between them and results in real algebraic geometry we refer the reader to [OZ13].

Given the above setup, we shall consider the following set of natural axioms to test if a graph G has a clique of size k.

Definition 1.4. Given a graph G, let Clique(G, k) denote the following set of polynomial axioms:

(Max-Clique):
$$x_i^2 - x_i, \forall i \in [n]$$

 $x_i \cdot x_j, \forall \text{ pairs } \{i, j\} \notin G$
 $\sum_i x_i - k.$
(1.2)

Here, the equations on the first line are Boolean axioms restricting feasible solutions to be in $\{0,1\}^n$. The equations on the second line constrain the support of any feasible x to define a clique in G. Finally, the equation on the third line specifies the size of support of x. Thus, for any graph G, Clique(G,k) is feasible if and only if G has a clique of size k. Our core result is to show lower bounds on positivstellensatz refutations for Clique(G,k).

Theorem 1.5 (Main). With high probability over $G \leftarrow G(n, 1/2)$, the system Clique(G, k) defined by Equation 1.2 has no $\mathsf{PS}(r)$ refutation for $k \leq n^{1/2r}/C^r(\log n)^{1/r}$

Given the above theorem it is easy to deduce the integrality gap for the SOS hierarchy, Theorem 1.1: see Section 12. We next highlight the outline of the proof, and some of our techniques which may be of broader interest.

1.4 Outline

We now give an outline of our arguments. As in most previous works (cf. [Gri01a], [Gri01b], [Sch08]) on showing lower bounds for $\mathsf{PS}(r)$ refutations, our main tool will be a *dual certificate*. We note that in the context of hierarchies above, this object is called either a *vector solution*⁵, or *pseudo-expectation*⁶. We now turn to define this important notion, which arises naturally from using duality to prove that a degree r refutation like 1.1 does *not* exist. Let $\mathcal{P}(n, 2r) : \mathbb{R}^n \to \mathbb{R}$ be the set of *n*-variate real polynomials of total degree at most 2r.

Definition 1.6 (PSD Mappings). A linear mapping $\mathcal{M} : \mathcal{P}(n, 2r) \to \mathbb{R}$ is said to be positive semi-definite (PSD) if $\mathcal{M}(P^2) \geq 0$ for all n-variate polynomials P of degree at most r.

Definition 1.7 (Dual Certificates). Given a set of axioms f_1, \ldots, f_m , a dual certificate for the axioms is a PSD mapping $\mathcal{M} : \mathcal{P}(n, 2r) \to \mathbb{R}$ such that $\mathcal{M}(f_ig) = 0$ for all $i \in [m]$ and all polynomials g such that $deg(f_ig) \leq 2r$.

⁵ in which numerical values to variables are replaced by vector values

⁶reflecting the view of these values as moments of a (possibly nonexistent) probability distribution

Under reasonable technical conditions which ensure strong duality, the converse also holds. For the clique axioms from Equation 1.2, a dual certificate would correspond to a feasible vector solution for the r-round SOS relaxation for maximum clique (see Figure 12 for the exact formulation) with value k.

The following elementary lemma will be crucial.

Lemma 1.8 (Dual Certificate). Given a system of axioms $((f_i))$, there does not exist a $\mathsf{PS}(r)$ refutation of the system if there exists a dual certificate $\mathcal{M} : \mathcal{P}(n, 2r) \to \mathbb{R}$ for the axioms.

The existence of such a mapping trivially implies a lower bound for $\mathsf{PS}(r)$ refutations: apply \mathcal{M} to both sides of a purported $\mathsf{PS}(r)$ identity as in Equation 1.1 to arrive at a contradiction.

The lemma suggests a general recipe for proving PS(r) refutation lower bounds:

- Design a dual certificate \mathcal{M} : For the clique axioms we care about, it is easy to figure out what the right dual certificate \mathcal{M} "should be" by working backwards from the axioms. The same happens also for the $\mathsf{PS}(r)$ refutation lower bounds of [Gri01a, Gri01b]. The main hurdle then is to show that the obtained mapping \mathcal{M} is indeed PSD. At a high level, this reduces to proving a certain random matrix $M \in \mathbb{R}^{\binom{n}{r} \times \binom{n}{r}}$ is PSD. We show that M is PSD in three steps.
- Reduction to PSDness of another matrix M': The matrix M has many zero rows and columns which makes it difficult to work with. In Section 5 we fix this by filling in the zero rows and columns of M to obtain a new matrix M'. We then argue that to show M is PSD it is sufficient to show that M' is PSD.
- (Deterministic) Matrix analysis: E = E[M'] is PSD with a large minimum eigenvalue $\lambda_{min}(E)$. We show this statement in Section Section 7 by using the theory of association schemes described below.
- Large deviation: with high probability, $||M' E|| \leq \lambda_{min}(E)$. This is done by using the structure of our matrix M' along-with a careful application of the trace method to bound the norms of certain random matrices with dependent entries.

We note here the main techniques used.

Techniques: Association schemes As discussed, the essence of proving Theorem 1.5 involves showing that a certain random matrix is positive semi-definite (PSD) with high probability. In our case, this calls for showing a relation of the form $A \prec B^7$ for two matrices A, B whose rows and columns are indexed by subsets of [n] of size r. This in turn leads us to matrices which though complicated to describe, will be *set-symmetric* - the entry defined by any two (row and column) sets I, J depends solely on the size of the intersection $I \cap J$. The set of all such matrices, called the Johnson scheme, is quite well studied in combinatorics as a special case of association schemes. In particular, all such matrices commute with one another and their common eigenspaces are completely understood. This theory allows us to estimate the eigenvalues and norms of various matrices that arise in the analysis.

⁷Here and henceforth \prec denotes PSD ordering: $A \prec B$ if and only if B - A is positive definite.

Techniques: Trace bounds for locally random matrices After various simplifications and reductions, a central problem we have to deal with is upper bounding the spectral norm of certain random matrices, defined by the underlying random graph $G \leftarrow G(n, 1/2)$. As above, these matrices have rows and columns indexed by subsets of vertices. The entry (I, J) of the matrix will be a random variable of expectation zero, which depends only on the edges and non-edges of G in the subgraph induced by $I \cup J$ (hence we name such matrices *local*). In the simple case when r = 1 (so rows and columns are indexed by singletons), which is the one studied in the analysis of the \sqrt{n} approximation algorithm, the random variables in all entries are mutually independent, and a norm bound is easy to obtain by a straightforward use of the trace method. However, for r > 1 as we need to handle, the entries of the matrix are dependent whenever the edge sets of their entries intersect. This significantly complicates the trace calculation, and we develop some combinatorial tools to bound the trace of high powers of such local matrices.

2 Dual certificate for PS(r) refutations of max-clique

We will specify the dual certificate \mathcal{M} by defining it for polynomials where each individual variable has degree at most 1 and extend \mathcal{M} multi-linearly to all polynomials: for any polynomial P, $\mathcal{M}(P) = \mathcal{M}(\tilde{P})$ where \tilde{P} is obtained from P by reducing the individual degrees of all variables to 1. We can do this without loss of generality because of the Boolean axioms.

As mentioned in the introduction, we can often work out what the dual certificate should be from the axioms and basic linear algebra. As an example, we first work out the case where the graph G is the complete graph; this will also help us draw a concrete connection to the work of [Gri01a].

2.1 Complete graph and knapsack

For complete graph, the clique axioms simplify to

(Max-Clique):
$$x_i^2 - x_i, \ \forall i \in [n]$$

 $\sum_i x_i - k.$

These incidentally also correspond to proving lower bounds for knapsack as studied by Grigoriev [Gri01a] (and was what lead us to the specific dual certificate we study). However, in the context of lower bounds for knapsack, the axioms are mainly interesting for non-integer k and Grigoriev shows that for non-integer $k \leq n/2$, the above system has no $\mathsf{PS}(r)$ refutation for r < k.

The above axioms tell us that any candidate dual certificate $\mathcal{M}_{Gr} \equiv \mathcal{P}(n, 2r) \to \mathbb{R}$ should satisfy:

$$\mathcal{M}_{Gr}\left(\left(\sum_{i=1}^{n} x_i - k\right) \left(\prod_{i \in I} x_i\right)\right) = 0, \ \forall I, \ |I| < 2r.$$

For $I \subseteq [n]$, let $X_I = \prod_{i \in I} x_i$. Now, as the above equation is symmetric, it is natural to assume that \mathcal{M}_{Gr} is also symmetric in the sense that $\mathcal{M}_{Gr}(X_I) = f(|I|)$ for some function $f : \{0, \ldots, 2r\} \to \mathbb{R}_+$. Working from this assumption, Grigoriev derives the following recurrence relation for $f : \{0, \ldots, 2r\} \to \{0, \ldots, 2r\} \to \mathbb{R}_+$,

$$f(i+1) = \frac{k-i}{n-i}f(i).$$

From the above it follows that we can define f and hence \mathcal{M} as follows:

$$\mathcal{M}_{Gr}(X_I) = f(|I|) = f(0) \cdot \frac{k(k-1)\cdots(k-|I|)}{n(n-1)\cdots(n-|I|)}$$

Grigoriev takes f(0) = 1. Here we set $f(0) = \binom{n}{2r}$ with a view towards what is to come. Thus, the final certificate is

$$\mathcal{M}_{Gr}(X_I) = \binom{n}{2r} \cdot \frac{k(k-1)\cdots(k-|I|)}{n(n-1)\cdots(n-|I|)} = \binom{n-|I|}{2r-|I|} \cdot \frac{\binom{k}{|I|}}{\binom{2r}{|I|}}.$$
(2.1)

Grigoriev shows the following:

Theorem 2.1 ([Gri01a]). For k < n/2, the mapping \mathcal{M}_{Gr} defined above is PSD for r < k.

2.2 Certificate for clique axioms

Following a similar approach, we now derive the dual certificate for the clique axioms from Equations 1.2, which we restate below for convenience: given a graph G on n vertices, $k \leq n$, the axioms of Clique(G, k) are

(Max-Clique):
$$x_i^2 - x_i, \forall i \in [n]$$

 $x_i \cdot x_j, \forall \text{ pairs } \{i, j\} \notin G$
 $\sum_i x_i - k.$ (2.2)

The above axioms tell us that any candidate dual certificate $\mathcal{M} \equiv \mathcal{M}_G : \mathcal{P}(n, 2r) \to \mathbb{R}$ should satisfy:

$$\mathcal{M}(X_I) = 0, \ \forall I, \ |I| \le 2r, \ I \text{ is not a clique in } G,$$
$$\mathcal{M}\left(\left(\sum_{i=1}^n x_i - k\right) X_I\right) = 0, \ \forall I, \ |I| < 2r.$$
(2.3)

The above equations give us a system of linear equations that \mathcal{M} needs to satisfy. By working with the equations, it is easy to guess a natural solution for the system.

Given a graph G on [n], and $I \subseteq [n]$, $|I| \leq 2r$, let

 $deg_G(I) = |\{S \subseteq [n] : I \subseteq S, |S| = 2r, S \text{ is a clique in } G\}|.$

For instance, if r = 1 and $v \in G$, then $deg_G(\{v\})$ is the degree of vertex v.

We define $\mathcal{M} \equiv \mathcal{M}_G : \mathcal{P}(n, 2r) \to \mathbb{R}$ for monomials as follows: for $I \subseteq [n], |I| \leq 2r$, let

$$\mathcal{M}\left(\prod_{i\in I} x_i\right) = \deg_G(I) \cdot \frac{k(k-1)\cdots(k-|I|+1)}{2r(2r-1)\cdots(2r-|I|+1)} = \deg_G(I) \cdot \frac{\binom{k}{|I|}}{\binom{2r}{|I|}}.$$
(2.4)

It is easy to check the following claim:

Claim 2.2. For any graph G, $\mathcal{M} \equiv \mathcal{M}_G$ defined by Equation 2.4 satisfies Equations 2.3.

Proof. The first equation in Equation 2.3 follows immediately from the definition of \mathcal{M} . Now, for $I \subseteq [n], |I| < 2r$,

$$\begin{split} \mathcal{M}\left(\left(\sum_{i} x_{i} - k\right) X(I)\right) &= (|I| - k) \mathcal{M}(X(I)) + \sum_{j \notin I} \mathcal{M}(X(I \cup \{j\})) \\ &= (|I| - k) \cdot \deg_{G}(I) \cdot \frac{\binom{k}{|I|}}{\binom{2r}{|I|}} + \sum_{j \notin I} \deg_{G}(I \cup \{j\}) \cdot \frac{\binom{k}{|I|+1}}{\binom{2r}{|I|+1}} \\ &= \frac{\binom{k}{|I|+1}}{\binom{2r}{|I|+1}} \cdot \left(-(2r - |I|) \cdot \deg_{G}(I) + \sum_{j \notin I} \deg_{G}(I \cup \{j\})\right). \end{split}$$

Observe that our notion of degree, deg_G , satisfies the following recurrence: for |I| < 2r,

$$deg_G(I) = \frac{1}{2r - |I|} \cdot \sum_{j \notin I, \text{ j adjacent to all of } I} deg_G(I \cup \{j\}) = \frac{1}{2r - |I|} \sum_{j \notin I} deg_G(I \cup \{j\}).$$

The above two equations imply that \mathcal{M} satisfies the second equation in 2.3.

Thus, to prove our main theorem Theorem 1.5, it suffices to show that \mathcal{M} as defined above is PSD with high probability. We now argue that in fact, to show that \mathcal{M} is PSD we do not need to consider all polynomials P of degree at most r. Rather, it is sufficient to show that $\mathcal{M}(P_1^2) \geq 0$ whenever P_1 is multilinear and homogeneous of degree r.

Lemma 2.3. For any P of degree at most r we may write $P = P_1 + \sum_i P_{2i}(x_i^2 - x_i) + P_3(\sum_i x_i - k)$ where P_1 is multilinear and homogeneous of degree r, P_3 has degree at most r - 1, and all P_{2i} have degree at most r - 2.

Proof. We first make P multilinear by removing any terms which are not multilinear from P as follows. If P has a term of the form $x_i^2 f$ where f has degree at most r-2, write $x_i^2 f = (x_i^2 - x_i)f + x_i f$. Iteratively applying this procedure, we may write P = P' plus terms of the form $(x_i^2 - x_i)f$ where P' is multilinear of degree at most r and f has degree at most r-2.

We now make P' multilinear and homogeneous of degree r by removing any terms which have lower degree as follows. If P' has a term of the form X_I where |I| < r, write

$$X_{I} = \frac{1}{|I| - k} \left(\sum_{i} x_{i} - k \right) X_{I} + \frac{1}{k - |I|} \sum_{i \in I} (x_{i}^{2} - x_{i}) X_{I \setminus \{i\}} + \frac{1}{k - |I|} \sum_{i \notin I} X_{I \cup \{i\}}$$

Iteratively applying this procedure, we may write $P = P_1$ plus terms of the form $(x_i^2 - x_i)f$ and terms of the forms $(\sum_i x_i - k)g$ where P_1 is multilinear and homogeneous of degree r, all such f have degree at most r - 2 and all such g have degree at most r - 1. Putting everything together, the result follows.

Corollary 2.4. If $\mathcal{M}(P_1^2) \geq 0$ for all multilinear homogeneous P_1 of degree r then \mathcal{M} is PSD.

Proof. Assume $\mathcal{M}(P_1^2) \geq 0$ for all multilinear homogeneous P_1 of degree r and $\mathcal{M}(P^2) < 0$ for some $P \in \mathcal{P}(n, r)$. Using Lemma 2.3, we may write $P = P_1 + \sum_i P_{2i}(x_i^2 - x_i) + P_3(\sum_i x_i - k)$ where P_1 is multilinear and homogeneous of degree r. $\mathcal{M}(P^2) = \mathcal{M}(P_1^2)$ so $\mathcal{M}(P_1^2) < 0$. Contradiction. \Box

Thus, showing that \mathcal{M} is PSD with high probability is equivalent to showing that the following matrix $M \equiv M_G \in \mathbb{R}^{\binom{[n]}{r} \times \binom{[n]}{r}}$ is PSD with high probability for $G \leftarrow G(n, 1/2)$: for $I, J \in \binom{[n]}{r}$,

$$M(I,J) = \deg_G(I \cup J) \cdot \frac{\binom{k}{|I \cup J|}}{\binom{2r}{|I \cup J|}}.$$
(2.5)

In the remainder of the paper, we show that M is PSD with high probability for $k \leq \Omega_r (n^{1/2r}/(\log n)^{1/r})$.

Theorem 2.5 (Main Technical Theorem). There exists a constant c > 0 such that, with high probability over $G \leftarrow G(n, 1/2)$, the matrix M_G defined by Equation 2.5 is PSD for $k \leq 2^{-cr} \cdot (\sqrt{n}/\log n)^{1/r}$.

3 Overview of proof of Theorem **2.5**

The proof of Theorem 2.5 is quite technical, and is broken into two parts, where the second part is further broken down into smaller parts. While we gave a sketch of the proof of Theorem 2.5 in the inroduction, we give a more detailed overview of the proof here. Recall that all matrices mentioned below are random matrices which are specified by the choice of the random graph G.

As mentioned in the introduction, the matrix $M = M_G$ has many zero rows and columns which makes it difficult to work with. The first part is to fill in the zero rows and columns of M to obtain a new matrix, M', which is nonsingular and has no high variance entries. In Section 5 we define this matrix M' and show that if M' is PSD, so is M. The idea is that M and M' are symmetric and the nonzero part of M is a principal submatrix of M', so the smallest nonzero eigenvalue of M is at least as large as the smallest eigenvalue of M'.

The second part is to prove that M' is PSD (indeed we prove that it has a high positive smallest eigenvalue). This is stated in the main technical lemma Lemma 8.1. For the proof of Lemma 8.1 we decompose the matrix M' as $M' = E + L + \Delta$, where (a) $E = \mathbb{E}[M']$ is the expectation matrix; (b) L will be a "local" random matrix such that for sets I, J, L(I, J) only depends on the edges among the vertices of $I \cup J$ and (c) Δ is a "global" error matrix whose entries are small in magnitude.

Having defined E (which is set-symmetric), let us spell out what the other matrices are. The "local" random matrix L is defined in a simple way as follows:

$$L(I,J) = \begin{cases} -E(I,J) & \text{if some edge in } \mathcal{E}(I\cup J) \setminus (\mathcal{E}(I)\cup \mathcal{E}(J)) \text{ is missing from } G \\ \beta(|I\cap J|) & \text{otherwise} \end{cases},$$

where $\mathcal{E}(I)$ denotes the set of possible edges between vertices of I and $\beta : \{0, \ldots, r\} \to \mathbb{R}_+$ is suitably chosen so that each individual entry of L has expectation zero.

Finally, define the last matrix $\Delta = M' - E - L$.

The proof that M' is PSD proceeds in three modular steps:

- 1. We use the results about Johnson scheme to show that $E \succ 0$ and has a large least eigenvalue (roughly $\Omega_r(k^r n^r)$); see Section 7.
- 2. We next show that $||L|| < Ck^{2r}n^{r-1/2}\log n$ by exploiting the recursive structure of the matrix L and some careful trace calculations. This is the most technically intensive part of the proof, and requires the development of some combinatorial tools to estimate the trace of high powers of L; see Section 8.2.

3. We then show that $\|\Delta\| < Ck^{2r}n^{r-1/2}\log n$. This is done by first showing that *every* entry of Δ is small in magnitude, via concentration bounds on the number of cliques in random graphs, and bounding its norm using Gershgorin's circle theorem (Lemma 4.1); see Section 8.3.

4 Preliminaries

We shall use the following notations⁸:

- 1. $\mathcal{P}(n, 2r)$ denotes the set of *n*-variate polynomials of degree at most 2r.
- 2. PS(r) denotes positivstellensatz refutations of degree at most r as defined in Definition 1.3.
- 3. A linear mapping $\mathcal{M} : \mathcal{P}(n, 2r) \to \mathbb{R}$ is said to be positive semi-definite (PSD) if $\mathcal{M}(P^2) \ge 0$ for all $P \in \mathcal{P}(n, r)$.
- 4. For $0 \le r \le n$, let $\binom{[n]}{r}$, $\binom{[n]}{\le r}$ denote all subsets of size exactly and at most r, respectively.
- 5. For $0 \le r \le n$, $\mathbb{R}^{\binom{[n]}{r} \times \binom{[n]}{r}}$ denotes matrices with rows and columns indexed by subsets of [n] of size exactly r. Similarly, $\mathbb{R}^{\binom{[n]}{\le r} \times \binom{[n]}{\le r}}$ denotes matrices with rows and columns indexed by subsets of [n] of size at most r.
- 6. We will view linear functionals $\mathcal{M} : \mathcal{P}(n, 2r) \to \mathbb{R}$ as matrices $M \in \mathbb{R}^{\binom{[n]}{\leq r} \times \binom{[n]}{\leq r}}$, where for $I, J \in \binom{[n]}{\leq r}, M_{IJ} = \mathcal{M}(\prod_{s \in I \cup J} x_s)$. In general, this correspondence is not bijective. However, as we only deal with mappings which are constant under multi-linear extensions throughout, the correspondence is one-to-one. It is a standard fact that a mapping \mathcal{M} is PSD if and only if the matrix M is PSD.
- 7. For $I \subseteq [n]$, let $X_I = \prod_{i \in I} x_i$.
- 8. By default all vectors are column vectors. For a set I, $\mathbb{1}(I)$ denotes the indicator vector of the set I.
- 9. For a matrix $A \in \mathbb{R}^{m \times n}$, $A^{\dagger} \in \mathbb{R}^{n \times m}$ denotes its conjugate matrix.

We will also need the following standard fact from matrix theory (see [GVL96] for instance).

Lemma 4.1 (special case of Gershgorin circle theorem). For any square matrix $M \in \mathbb{R}^{N \times N}$,

$$\|M\| \le \max_{i \in [N]} \left(\sum_{j=1}^{N} |M_{ij}| \right).$$

Finally, we need McDiarmid's inequality for obtaining tail bounds for functions of independent random variables (see [?] for instance)

⁸Some are repeated from the introduction so as to have them at one place.

Theorem 4.2 (McDiarmid's inequality). Let X_1, \ldots, X_n be independent random variables and let f be a function over the domain space of (X_1, \ldots, X_n) . Let $c_1, \ldots, c_n > 0$ be such that for all i, x_1, \ldots, x_n, x'_i ,

$$|f(x_1,\ldots,x_{i-1},x_i,x_{i+1},\ldots,x_n) - f(x_1,\ldots,x_{i-1},x'_i,x_{i+1},\ldots,x_n)| \le c_i.$$

Then, for all t > 0,

$$\Pr\left[|f(X_1,\ldots,X_n) - \mathbb{E}[f]| > t\right] \le 2\exp\left(\frac{-2t^2}{\sum_{i=1}^n c_i^2}\right)$$

5 Reduction to PSDness of M'

In this section, we define the matrix M' and show that if M' is PSD then so is M. We use the following notations for brevity: For any set $I \subseteq [n]$, let $\mathcal{E}(I) = \{\{i, j\} : i \neq j \in I\}$. For $0 \leq i \leq r$, let

$$\beta(i) = \binom{k}{2r-i} / \binom{2r}{2r-i}.$$
(5.1)

For every $T \subseteq [n]$, let $M_T \in \mathbb{R}^{\binom{[n]}{r} \times \binom{[n]}{r}}$, with $M_T(I, J) = \beta(|I \cap J|)$ if $I \cup J \subseteq T$, and G contains every edge in $\mathcal{E}(T) \setminus \mathcal{E}(I) \cup \mathcal{E}(J)$ (i.e., the only edges in T missing in G are those with both end points in one of I or J). We will study the matrix

$$M' = \sum_{T:|T|=2r} M_T.$$
 (5.2)

Intuitively, for every I, J, M'(I, J) is what M(I, J) would be had we added cliques on the subsets I, J to the graph. The above definition avoids the problem of the whole row and column corresponding to I or J becoming zero if either was not a clique and controls the variance of the entries. We now show that to show M is PSD, it is sufficient to show that M' is PSD.

Lemma 5.1. If M' is PSD then M is PSD.

Proof. The reason this lemma is true is because as shown below, the nonzero part of M is a principal submatrix of M'.

Proposition 5.2. Whenever I and J are cliques of size r in G, M'(I, J) = M(I, J)

Proof. Suppose that I and J are cliques in G. Then, $M_T(I, J) = \beta(|I \cap J|)$ if $I \cup J \subseteq T$ and T is a clique and 0 otherwise. Therefore,

$$M'(I,J) = \sum_{T} M_T(I,J) = \beta(|I \cap J|) \cdot |\{T : I \cup J \subseteq T, T \text{ clique}\}| = M(I,J).$$

Corollary 5.3. The nonzero part of M is a principal submatrix of M'.

We now use the following elementary fact about matrices.

Proposition 5.4. If A is a principal submatrix of a symmetric matrix B then the smallest eigenvalue of A is at least as large as the smallest eigenvalue of B.

Proof. Without loss of generality, A is an $l \times l$ matrix and B is an $m \times m$ matrix where $l \leq m$. Let $v \in \mathbb{R}^l$ be a unit eigenvector of A with minimal eigenvalue λ_{min} . If we let $w \in \mathbb{R}^m$ be the extension of v to \mathbb{R}^m with zeros in the other coordinates, $w^T B w = v^T A v = \lambda_{min}$. This implies that the smallest eigenvalue of B is at most λ_{min} and the result follows.

Combining Corollary 5.3 and Proposition 5.4, if M' is PSD then M is PSD, as needed.

6 Johnson scheme

Association schemes is a classical area in combinatorics and coding theory (cf. for instance [vLW01]). We shall use a few classical results (lemmas 6.6, 6.7 below), about the eigenspaces and eigenvalues of association schemes and the *Johnson scheme* in particular. We also introduce two bases for the Johnson scheme, which will play a key role in bounding the eigenvalues of various matrices later.

We start with some basics about the Johnson scheme - some of our notations are non-standard but they fit better with the rest of the manuscript.

Definition 6.1 (Set-Symmetry). A matrix $M \in \mathbb{R}^{\binom{[n]}{r} \times \binom{[n]}{r}}$ is set-symmetric if for every $I, J \in \binom{[n]}{r}$, M(I, J) depends only on the size of $|I \cap J|$.

Definition 6.2 (Johnson Scheme). For $n, r \leq n/2$, let $\mathcal{J} \equiv \mathcal{J}_{n,r} \subseteq \mathbb{R}^{\binom{[n]}{r} \times \binom{[n]}{r}}$ be the subspace of all set-symmetric matrices. \mathcal{J} is called the Johnson scheme.

As we will soon see, \mathcal{J} is also a commutative algebra. There is a natural basis for the subspace \mathcal{J} :

Definition 6.3 (D-Basis). For $0 \le \ell \le r \le n$, let $D_{\ell} \equiv D_{n,r,\ell} \in \mathbb{R}^{\binom{[n]}{r} \times \binom{[n]}{r}}$ be defined by⁹

$$D_{\ell}(I,J) = \begin{cases} 1 & |I \cap J| = \ell \\ 0 & otherwise. \end{cases}$$
(6.1)

For example, D_0 is the well-studied disjointness matrix. Clearly, $\{D_{\ell} : 0 \leq \ell \leq r\}$ span the subspace \mathcal{J} . Also, it is easy to check that the D_{ℓ} 's and hence all the matrices in \mathcal{J} , commute with one another.

Another important collection of matrices that come up naturally while studying PSD'ness of set-symmetric matrices is the following which gives a basis of PSD matrices for the Johnson scheme.

Definition 6.4 (P-Basis). For $0 \le t \le r$, let $P_t \equiv P_{n,r,t} \in \mathbb{R}^{\binom{[n]}{r} \times \binom{[n]}{r}}$ be defined by¹⁰

$$P_t(I,J) = \binom{|I \cap J|}{t}.$$

Equivalently, for $T \subseteq [n]$, if we let P_T be the PSD rank one matrix

$$P_T = \mathbb{1}\left(\{I : I \subseteq [n], I \supseteq T\}\right) \cdot \mathbb{1}\left(\{I : I \subseteq [n], I \supseteq T\}\right)^{\dagger}$$

⁹We will often omit the subscripts n, r.

¹⁰We will often omit the subscripts n, r.

then

$$P_t = \sum_{T:T \subseteq [n], |T|=t} P_T.$$
 (6.2)

The equivalence of the above two definitions follows from a simple calculation: there is a non-zero contribution to (I, J)'th entry from the T'th summand from Equation 6.2 if and only if $T \subseteq I \cap J$. Clearly, $P_t \succeq 0$ for $0 \leq t \leq r$. We will exploit this relation repeatedly by expressing matrices in \mathcal{J} as linear combinations of P_t 's. The following elementary claim relates the two bases $((D_\ell))$ and $((P_t))$ for fixed n, r.

Claim 6.5. For fixed n, r, the following relations hold:

- 1. For $0 \le t \le r$, $P_t = \sum_{\ell=t}^r \binom{\ell}{t} D_{\ell}$.
- 2. For $0 \le \ell \le r$, $D_{\ell} = \sum_{t=\ell}^{r} (-1)^{t-\ell} {t \choose \ell} P_t$.

Proof. The first relation follows immediately from the definition of P_t . The second relation follows from inverting the set of equations given in (1).

The main nontrivial result from the theory of association schemes we use is the following characterization of the eigenspaces of matrices in \mathcal{J} . The starting point for these characterizations is the fact that matrices in \mathcal{J} commute with one another and hence are simultaneously diagonalizable. We refer the reader to Section 7.4 in [God] (the matrices P_t in our notation correspond to matrices C_t in [God]) for the proofs of these results.

Lemma 6.6. Fix $n, r \leq n/2$ and let $\mathcal{J} \equiv \mathcal{J}(n, r)$ be the Johnson scheme. Then, for P_t as defined by Equation 6.2, there exist subspaces $V_0, V_1, \ldots, V_r \in \mathbb{R}^{\binom{[n]}{r}}$ that are orthogonal to one another such that:

- 1. V_0, \ldots, V_r are eigenspaces for $\{P_t : 0 \le t \le r\}$ and consequently for all matrices in \mathcal{J} .
- 2. For $0 \le j \le r$, $dim(V_j) = \binom{n}{j} \binom{n}{j-1}$.
- 3. For any matrix $Q \in \mathcal{J}$, let $\lambda_i(Q)$ denote the eigenvalue of Q within the eigenspace V_i . Then,

$$\lambda_j(P_t) = \begin{cases} \binom{n-t-j}{r-t} \cdot \binom{r-j}{t-j} & j \le t\\ 0 & j > t \end{cases}.$$
(6.3)

The above lemma helps us estimate the eigenvalues of any matrix in $Q \in \mathcal{J}$ if we can write Q as a linear combination of the P_t 's or D_ℓ 's. To this end, we shall also use the following estimate on the eigenvalues of such linear combinations.

Lemma 6.7. Let $Q = \sum_{\ell} \alpha_{\ell} D_{\ell} \in \mathcal{J}(n, r)$, and $\beta_t = \sum_{\ell \leq t} {t \choose \ell} \alpha_{\ell}$, where $\alpha_{\ell} \geq 0$. Then, for $0 \leq j \leq r$, $\lambda_j(Q) \leq \sum_{t \geq i} \beta_t \cdot {n-t-j \choose r-t} \cdot {r-j \choose t-j}$. *Proof.* By Claim 6.5,

$$\sum_{\ell} \alpha_{\ell} D_{\ell} = \sum_{\ell} \alpha_{\ell} \left(\sum_{t \ge \ell} (-1)^{t-\ell} {t \choose \ell} P_t \right) = \sum_{t} P_t \left(\sum_{\ell \le t} (-1)^{t-\ell} {t \choose \ell} \alpha_{\ell} \right)$$
$$\preceq \sum_{t} P_t \left(\sum_{\ell \le t} {t \choose \ell} \alpha_{\ell} \right) = \sum_{t} \beta_t P_t.$$

Therefore, as Q and P_t 's have common eigenspaces, by Lemma 6.6,

$$\lambda_j(Q) \le \lambda_j\left(\sum_t \beta_t P_t\right) \le \sum_t \beta_t \lambda_j(P_t) = \sum_{t \ge j} \beta_t \cdot \binom{n-t-j}{r-t} \cdot \binom{r-j}{t-j}.$$

7 PSD'ness of the expectation matrices

In the section we show that if r is not too large then the expectation matrix $E = \mathbb{E}[M']$ is PSD with high minimal eigenvalue. As a warmup, we first show that the expectation matrix $E_M = \mathbb{E}[M]$ is PSD. We start by writing down E_M .

Claim 7.1. For $I, J \in \binom{n}{r}$, and $E_M = \mathbb{E}[M]$,

$$E_M(I,J) = \binom{n-|I\cup J|}{2r-|I\cup J|} \cdot \frac{\binom{k}{|I\cup J|}}{\binom{2r}{|I\cup J|}} \cdot 2^{-\binom{2r}{2}}.$$
(7.1)

Proof. The claim follows from observing that for all I and J, $\mathbb{E}[deg_G(I \cup J)] = \binom{n-|I\cup J|}{2r-|I\cup J|} \cdot 2^{-\binom{2r}{2}}$. To see this, note that for all I and J there are $\binom{n-|I\cup J|}{2r-|I\cup J|}$ sets of size 2r containing $I \cup J$ and each is a clique with probability $2^{-\binom{2r}{2}}$.

The expectation matrix above is just a scalar multiple of \mathcal{M}_{Gr} (viewed as a matrix) as defined in Equation 2.1. Therefore, by Theorem 2.1, E_M as defined above is PSD for $r < \min(k, n-k)$. We give a simpler proof of this claim here for the case when $r \leq \min(\frac{k}{2}, n-k)$.

Theorem 7.2. The matrix E_M is positive definite for $r \leq \min(\frac{k}{2}, n-k)$.

Proof. We will show this by writing E_M as a suitable positive linear combination of the PSD matrices P_t 's from Section 6. More concretely, for any $\alpha_0, \ldots, \alpha_t > 0$, we have

$$0 \prec \sum_{t} \alpha_t P_t = \sum_{\ell=0}^r \left(\sum_{t=0}^\ell \alpha_t \binom{\ell}{t} \right) D_\ell$$

Now, let $e_{\ell} = E_M(I, J)$ for any I and J with $|I \cup J| = 2r - \ell$, i.e.,

$$e_{\ell} = 2^{-\binom{2r}{2}} \cdot \binom{n-2r+\ell}{\ell} \cdot \frac{\binom{k}{2r-\ell}}{\binom{2r}{2r-\ell}}$$

Then, $E_M = \sum_{\ell=0}^r e_\ell D_\ell$. Therefore, we will be done if we can find α_t 's such that for every $0 \le \ell \le r$, $e_\ell = \sum_{t=0}^\ell \alpha_t {\ell \choose t}$. By examining the first values of ℓ , it is easy to guess what the α_t should be. First observe that $e_\ell = e_0 \cdot {\binom{n-2r+\ell}{\ell}}/{\binom{k-2r+\ell}{\ell}}$ and let $\alpha_t = e_0 {\binom{n-k}{t}}/{\binom{k-2r+t}{t}}$. Then,

$$e_0\binom{n-2r+\ell}{\ell} = e_0 \sum_{t=0}^{\ell} \binom{n-k}{t} \cdot \binom{k-2r+\ell}{\ell-t}$$
$$= \sum_{t=0}^{\ell} \alpha_t \cdot \binom{k-2r+t}{t} \binom{k-2r+\ell}{\ell-t}$$
$$= \sum_{t=0}^{\ell} \alpha_t \cdot \binom{\ell}{t} \cdot \binom{k-2r+\ell}{\ell}.$$

Therefore, $e_{\ell} = \sum_{t} {\ell \choose t} \alpha_{t}$ and the lemma now follows:

$$E_M = \sum_{\ell=0}^r e_\ell D_\ell = \sum_{\ell=0}^r \left(\sum_{t=0}^\ell \alpha_t \binom{\ell}{t} \right) D_\ell \succeq \alpha_r \mathbb{I}.$$

7.1 **PSD**'ness of E

Now that we have shown that E_M is positive definite when r is not too large, we use similar ideas to analyze $E = \mathbb{E}[M']$, the expectation matrix we will actually be using. We begin by writing down E.

Claim 7.3. For $I, J \in {n \choose r}$, and $E = \mathbb{E}[M']$,

$$E(I,J) = \binom{n - |I \cup J|}{2r - |I \cup J|} \cdot \frac{\binom{k}{|I \cup J|}}{\binom{2r}{|I \cup J|}} \cdot 2^{-r^2 - \binom{|I \cap J|}{2}}.$$
(7.2)

Proof. The claim follows from observing that for all I and J, conditioned on the edges in $\mathcal{E}(I)$ and $\mathcal{E}(J)$ being present, $\mathbb{E}[deg_G(I \cup J)] = \binom{n-|I\cup J|}{2r-|I\cup J|} \cdot 2^{-\binom{2r}{2} + \binom{|I|}{2} + \binom{|J|}{2} - \binom{|I\cap J|}{2}} = \binom{n-|I\cup J|}{2r-|I\cup J|} \cdot 2^{-r^2 - \binom{|I\cap J|}{2}}$. To see this, note that for all I and J there are $\binom{n-|I\cup J|}{2r-|I\cup J|}$ sets of size 2r containing $I \cup J$, and conditioned on the edges in $\mathcal{E}(I)$ and $\mathcal{E}(J)$ being present, each is a clique with probability $2^{-\binom{2r}{2} + \binom{|I|}{2} + \binom{|J|}{2} - \binom{|I\cap J|}{2}}$. Now note that |I| = |J| = r and $-\binom{2r}{2} + 2\binom{r}{2} = -(2r^2 - r) + (r^2 - r) = -r^2$ so $-\binom{2r}{2} + \binom{|I|}{2} + \binom{|J|}{2} - \binom{|I\cap J|}{2} = -r^2 - \binom{|I\cap J|}{2}$.

Lemma 7.4. If $k < \frac{n-2r}{3r \cdot 2^{r-1}}$ and $r \leq \frac{k}{2}$ then E is PSD with minimal eigenvalue $2^{-O(r^2)}k^r n^r$

Proof. By Equation 7.2, $E = \sum_{\ell} e_{\ell} D_{\ell}$, where $e_{\ell} = \binom{n-2r+l}{l} \cdot \frac{\binom{k}{2r-\ell}}{\binom{2r}{2r-\ell}} \cdot 2^{-r^2 - \binom{\ell}{2}}$. We next express E as a linear combination of P_t 's: $E = \sum_t \alpha_t P_t$. By Claim 6.5, $D_{\ell} = \sum_{t=\ell}^r (-1)^{t-\ell} \binom{t}{\ell} P_t$ so

$$\alpha_t = \sum_{\ell=0}^t (-1)^{t-\ell} \binom{t}{\ell} e_\ell.$$

Now note that for all $l \ge 1$, $e_{\ell} = \frac{n-2r+\ell}{\ell} \cdot \frac{\ell}{k-2r+\ell} \cdot 2^{\ell-l} \cdot e_{\ell-1} = \frac{n-2r+\ell}{2^{1-l}(k-2r+\ell)} \cdot e_{\ell-1}$. If $k < \frac{n-2r}{3r\cdot 2^{r-1}}$ then the terms in the sum for α_t increase geometrically by a factor of at least 3 and the sum will therefore be dominated by the last term. In particular, $\alpha_t \ge \frac{e_t}{2}$. Thus, $\alpha_t > 0$ for all $t \in [0, r]$ and

$$\alpha_r \ge \frac{e_r}{2} = \frac{1}{2} \cdot \binom{n-r}{r} \cdot \frac{\binom{k}{r}}{\binom{2r}{r}} \cdot 2^{-r^2 - \binom{r}{2}} = 2^{-O(r^2)} k^r n^r$$

Since the P_t 's are PSD and $P_r = I$, E is PSD with minimal eigenvalue $2^{-O(r^2)}k^r n^r$, as needed.

8 PSD'ness of dual certificate

We are now ready to prove our main result, Theorem 1.5, with the aid of several technical results whose proof is deferred to Section 9 and Section 10. We prove Theorem 1.5 by showing that the matrix M will be PSD with high probability (Theorem 2.5). In turn, we show that M is PSD with high probability with our main technical lemma, which says that M' is PSD with high probability (this is sufficient by Lemma 5.1).

Lemma 8.1 (Main Technical Lemma). For c a sufficiently large constant the following holds. The matrix $M' \in \mathbb{R}^{\binom{[n]}{r} \times \binom{[n]}{r}}$ defined by Equation 5.2 is positive definite with high probability, for $k < 2^{-cr} (\sqrt{n}/\log n)^{1/r}$.

To prove Lemma 8.1, we first decompose M' as $M' = E + L + \Delta$ in Section 8.1. We then analyze L and Δ in Section 8.2 and Section 8.3 respectively. We put all the pieces together to show the PSD'ness of M' in Section 8.4.

For the remainder of this section, we shall use the following additional notations:

• For $0 \le i \le r$, let

$$\alpha(i) = \frac{\binom{k}{2r-i}}{\binom{2r}{2r-i}} \cdot \binom{n-2r+i}{i} \cdot 2^{-r^2 - \binom{i}{2}}.$$
(8.1)

- For $0 \le i \le r$, let $p(i) = 2^{-(r-i)^2}$. Then, for $I, J \in {\binom{[n]}{r}}$ with $|I \cap J| = i, p(i)$ is the probability that $\mathcal{E}(I \cup J) \setminus (\mathcal{E}(I) \cup \mathcal{E}(J)) \subseteq G$.
- In the following we will adopt the convention that I, J, K denote elements of $\binom{[n]}{r}$ and T, T' denote elements of $\binom{[n]}{2r}$.
- All matrices considered below will be over $\mathbb{R}^{\binom{[n]}{r} \times \binom{[n]}{r}}$ unless otherwise specified.
- We write $A \approx_r B$ if there exist constants c, C such that $c^{r^2}B \leq A \leq C^{r^2}B$.

8.1 Decomposition of M'

For the proof of Lemma 8.1 we decompose the matrix M' as $M' = E + L + \Delta$, where (a) $E = \mathbb{E}[M']$ is the expectation matrix; (b) L will be a "local" random matrix such that for sets I, J, L(I, J) only depends on the edges between the vertices of $I \cup J$ and (c) Δ is a "global" error matrix whose entries are small in magnitude.

To this end, first observe that by Equation 7.2, for $E \equiv \mathbb{E}[M']$,

$$E(I,J) = \alpha(|I \cap J|). \tag{8.2}$$

Now, define $L \in \mathbb{R}^{\binom{[n]}{r} \times \binom{[n]}{r}}$ as follows: for $I, J \in \binom{[n]}{r}$,

$$L(I,J) = \begin{cases} \alpha(|I \cap J|) \cdot \frac{1-p(|I \cap J|)}{p(|I \cap J|)} & \text{if } \mathcal{E}(I \cup J) \setminus (\mathcal{E}(I) \cup \mathcal{E}(J)) \subseteq G \\ -\alpha(|I \cap J|) & \text{otherwise} \end{cases}.$$
(8.3)

Finally, define $\Delta = M' - E - L$. We have already shown in Section 7 that E is PSD with minimal eigenvalue $2^{-O(r^2)}k^r n^r$. There are now two remaining modular steps in the proof:

- 1. We show that ||L|| is $2^{O(r^2)}k^{2r}n^{r-1/2}\log n$ by exploiting the recursive structure of the matrix L and some careful trace calculations. This is the most technically intensive part of the proof.
- 2. We then show that $\|\Delta\|$ is $2^{O(r^2)}k^{2r}n^{r-1/2}\log n$. This is done by first showing that each entry of Δ is small in magnitude and using Lemma 4.1.

The next two subsections address these two steps with the corresponding technical elements dealt with in Section 9 and Section 10 respectively.

8.2 Bounding the norm of the locally random matrix L

In this subsection, we bound the norm of the matrix L.

Lemma 8.2. For some constant C > 0, with probability at least 1 - 1/n over the random graph G,

$$||L|| \le O(1) \cdot 2^{Cr^2} \cdot k^{2r} \cdot n^r \cdot \frac{\log n}{\sqrt{n}}$$

We will prove the lemma by further decomposing L according to the intersection sizes of the indexing sets and using the recursive structure of the matrix M'. To this end, we define the following closely related locally-random matrix. For $a \in [r]$, let $R_a \in \mathbb{R}^{\binom{n}{a} \times \binom{n}{a}}$ be the matrix supported only on disjoint sets and defined as follows: for $V, W \in \binom{[n]}{a}$,

$$R_a(V,W) = \begin{cases} 2^{a^2} - 1 & \text{if } V \cap W = \emptyset \text{ and } \{\{v,w\} : v \in V, w \in W\} \subseteq G \\ -1 & \text{if } V \cap W = \emptyset \text{ and } \{\{v,w\} : v \in V, w \in W\} \not\subseteq G \\ 0 & \text{if } V \cap W \neq \emptyset \end{cases}$$
(8.4)

In other words, for disjoint $V, W \in {\binom{[n]}{a}}$ the $R_a(V, W)$ 'th entry is essentially (up to a constant multiple) a shift of the indicator random variable which is 1 if all edges in $V \times W$ are in G and 0 otherwise.

Note that $\mathbb{E}[R_a] = 0$. The following technical claim proved in Section 9 bounds the norm of R_a . The proof relies on computing the trace of powers of R_a .

Claim 8.3 (See Section 9). If $n \ge 100$, for all $\varepsilon \in (0,1)$, $\Pr\left[||R_a|| > 2^{a^2+2a+2} \ln\left(\frac{n}{\varepsilon}\right)n^{a-\frac{1}{2}}\right] < \varepsilon$.

Note that $2^{a^2}n^a$ is an easy bound for $||R_a||$ (each entry of the matrix is at most 2^{a^2} in magnitude); the main advantage of the claim is the multiplicative $n^{-1/2}$ factor.

In the remainder of this section we use the recursive structure of the matrix L to prove Claim 8.2 assuming the above claim. We first introduce some notation:

- For a matrix $X \in \mathbb{R}^{\binom{[n]}{r_1} \times \binom{[n]}{r_2}}$, and $0 \le i \le \min\{r_1, r_2\}$, let $X^i \in \mathbb{R}^{\binom{[n]}{r_1} \times \binom{[n]}{r_2}}$ be the matrix such that $X^i(I, J) = X(I, J)$ if $|I \cap J| = i$ and 0 otherwise¹¹.
- For a matrix $X \in \mathbb{R}^{\binom{[n]}{r_1-i} \times \binom{[n]}{r_2-i}}$, let $X^{(i)} \in \mathbb{R}^{\binom{[n]}{r_1} \times \binom{[n]}{r_2}}$, be defined as follows:

$$X^{(i)}(I,J) = \begin{cases} X(I \setminus (I \cap J), J \setminus (I \cap J)) & \text{if } |I \cap J| = i \\ 0 & \text{otherwise} \end{cases}.$$
(8.5)

The next claim relates the norms of "lifts" of matrix R, $R^{(i)}$. Conceptually, bounding the norms of matrices with non-zero entries on intersecting indexing sets are reduced to that of the disjoint case. Note that the requirement $R = R^0$ exactly captures the latter.

Lemma 8.4. For $0 \le i \le \min\{r_1, r_2\}$ and $R \in \mathbb{R}^{\binom{[n]}{r_1-i} \times \binom{[n]}{r_2-i}}$, if $R = R^0$ then $||R^{(i)}|| \le \binom{r_1}{i}\binom{r_2}{i} \cdot ||R||$.

Proof. We partition the entries of $R^{(i)}$ as follows.

Definition 8.5. For any X, Y, K such that $X \subseteq [1, r_1], Y \subseteq [1, r_2]$, and $K \subseteq V(G)$ where |K| = |X| = |Y| = i, let $R_{X,Y,K}^{(i)}$ be the matrix such that the following is true:

- 1. $R_{X,Y,K}^{(i)}(I,J) = R^{(i)}(I,J) = R(I \setminus K, J \setminus K)$ if $K = \{i_x : x \in X\} = \{j_y : y \in Y\}$ where i_1, \dots, i_{r_1} are the elements of I in increasing order and j_1, \dots, j_{r_2} are the elements of J in increasing order.
- 2. $R_{X,Y,K}^{(i)}(I,J) = 0$ otherwise.

Proposition 8.6. For all X, Y, K, $||R_{X,Y,K}^{(i)}|| \le ||R||$.

Proof. The nonzero part of $R_{X,Y,K}^{(i)}$ can be viewed as a submatrix of R, so it cannot have larger induced norm than R.

Proposition 8.7. $R^{(i)} = \sum_{X,Y,K} R^{(i)}_{X,Y,K}$.

Proof. If $R^{(i)}(I,J) = 0$ then $\sum_{X,Y,K} R^{(i)}_{X,Y,K}(I,J) = 0$. If $R^{(i)}(I,J) \neq 0$ then $|I \cap J| = i$. This implies that $K = \{i_x : x \in X\} = \{j_y : y \in Y\}$ if and only if $K = I \cap J$, X is the set of indices of K in I, and Y is the set of indices of K in J, which happens for precisely one X, Y, K. Thus, $R^{(i)}_{X,Y,K}(I,J) = R^{(i)}(I,J)$ for precisely one I, J, X and is 0 otherwise, so $R^{(i)} = \sum_{X,Y,K} R^{(i)}_{X,Y,K}$, as needed.

Proposition 8.8. $||R^{(i)}|| \le \sum_{X,Y} ||\sum_{K} R^{(i)}_{X,Y,K}||$.

¹¹For this paper, we will only use the case where $r_1 = r_2 = r$. We put in this extra generality with an eye towards future work.

Proposition 8.9. If K_1, K_2 are distinct subsets of V(G) of size x, $R_{X,Y,K_1}^{(i)}(I_1, J_1) \neq 0$, and $R_{X,Y,K_2}^{(i)}(I_2, J_2) \neq 0$ then $I_1 \neq I_2$ and $J_1 \neq J_2$.

Proof. Assume that $I_1 = I_2 = I$ and let i_1, \dots, i_{r_1} be the elements of I in increasing order. Then $K_1 = \{i_x : x \in X\} = K_2$. Contradiction. Following similar logic, we cannot have that $J_1 = J_2$ either.

Proposition 8.10. For any $X, Y \subseteq [1, n], || \sum_{K} R_{X,Y,K}^{(i)}|| \le ||R||.$

Proof. Note that we can permute the rows and columns of a matrix without affecting its induced norm. By Proposition 8.9, we can permute the rows and columns of $\sum_{K} R_{X,Y,K}^{(i)}$ to put it into block form where each block is the nonzero part of $R_{X,Y,K}^{(i)}$ for some K. For a matrix in block form, its norm is the maximum of the norms of the individual blocks, which by Proposition 8.6 is at most ||R||, as needed.

With these results, Lemma 8.4 follows immediately. Plugging in Proposition 8.10 to Proposition 8.8 gives $||R^{(i)}|| \leq \sum_{X,Y} ||\sum_{K} R^{(i)}_{X,Y,K}|| \leq \sum_{X,Y} ||R|| \leq {r_1 \choose i} {r_2 \choose i} ||R||$, as needed.

We now use the above statements to prove Lemma 8.2.

Proof of Lemma 8.2. We claim that for $0 \le i \le r$, and α_i as in Equation 8.1

$$L^i = \alpha_i \cdot R_{r-i}^{(i)}. \tag{8.6}$$

To see the above, fix $I, J \in {[n] \choose r}$ with $|I \cap J| = i$ and let $V = I \setminus (I \cap J), W = J \setminus (I \cap J)$. Observe that

$$\mathcal{E}(I \cup J) \setminus (\mathcal{E}(I) \cup \mathcal{E}(J)) = \{\{v, w\} : v \in V, w \in W\}.$$

We cosider two cases as in the definition of L.

Case 1. $\mathcal{E}(I \cup J) \setminus (\mathcal{E}(I) \cup \mathcal{E}(J)) \subseteq G$. Then, $R_{r-i}^{(i)}(I,J) = R_{r-i}(V,W) = 2^{(r-i)^2-1} = (1 - p(i))/p(i)$. Equation 8.6 now follows from the first case of the definition of L.

Case 2. $\mathcal{E}(I \cup J) \setminus (\mathcal{E}(I) \cup \mathcal{E}(J)) \not\subseteq G$. Then, $R_{r-i}^{(i)}(I,J) = R_{r-i}(V,W) = -1$. Equation 8.6 now follows from the second case of the definition of L.

Therefore, by Claim 8.3, Lemma 8.4 and Equation 8.6,

$$||L^i|| \le O(1) \cdot 2^{Cr^2} \cdot k^{2r-i} \cdot n^r \cdot \frac{\log n}{\sqrt{n}}.$$

The lemma now follows as $L = \sum_{i=0}^{r} L^{i}$.

8.3 Bounding the norm of the global error matrix Δ

The main claim of this subsection is the following bound on the spectral norm of Δ .

Lemma 8.11. For $n > C2^{4r^2}$, with probability at least 1 - 1/n over the random graph G,

$$\|\Delta\| \le 2^{Cr^2} \cdot k^{2r} \cdot n^r \cdot \frac{\log n}{\sqrt{n}}$$

The proof relies on the following bound on the individual entries of Δ .

Lemma 8.12. For some universal constant C, and $n > C2^{4r^2}$, with probability at least 1 - 1/n over the random graph G, for all $I, J \in {[n] \choose r}$, with $i = |I \cap J|$,

$$|\Delta(I,J)| \le 2^{Cr^2} \cdot k^{2r-i} \cdot n^i \cdot \frac{\log n}{\sqrt{n}}.$$

Before proving the lemma, we first use it to bound $\|\Delta\|$.

Proof of Lemma 8.11. Suppose that the conclusion of the previous lemma holds. Then, for any $I \in {\binom{[n]}{r}}$,

$$\begin{split} \|\Delta\| &\leq \sum_{J} |\Delta(I,J)| = \sum_{i=0}^{r} \sum_{J:|I \cap J|=i} |\Delta(I,J)| \\ &\leq \frac{2^{Cr^{2}k^{2r}(\log n)}}{\sqrt{n}} \sum_{i=0}^{r} \sum_{J:|I \cap J|=i} (n/k)^{i} \\ &\leq \frac{2^{Cr^{2}k^{2r}(\log n)}}{\sqrt{n}} \sum_{i=0}^{r} (n/k)^{i} 2^{r} n^{r-i} \\ &\leq \frac{2^{Cr^{2}k^{2r}(\log n)n^{r}}}{\sqrt{n}}. \end{split}$$

The lemma now follows from the above bound and Lemma 4.1.

Proof of Lemma 8.12. Fix sets I, J with $|I \cap J| = i$. Let \mathcal{A} be the event that $\mathcal{E}(I \cup J) \setminus (\mathcal{E}(I) \cup \mathcal{E}(J)) \subseteq G$.

Then, by the second case of Equation 8.3, conditioned on $\neg \mathcal{A}$ we have $\Delta(I, J) = 0$. Thus, the claim holds trivially in this case. In the following we condition on \mathcal{A} . Observe that

$$\mathbb{E}[M'(I,J) \mid \mathcal{A}] = E(I,J)/\Pr[\mathcal{A}] = E(I,J)/p(i).$$

We next use the following claim that $\deg_G(I \cup J)$ is concentrated around its mean when conditioned on $I \cup J$ being a clique. At a high level, this follows from the fact that conditioned on $I \cup J$ being a clique, $\deg_G(I \cup J)$ can be written as a (structured) low-degree polynomial in the indicator variables of the edges not in $I \cup J$ with small variance. We defer the proof to the appendix.

Claim 8.13 (See Theorem 10.1 of the appendix). For some constant C > 0,

$$\Pr\left[\left|\deg_{G}(I\cup J) - 2^{-\binom{2r}{2} + \binom{2r-i}{2}} \cdot \binom{n-2r+i}{i}\right| > 2(\ln(C/\varepsilon))^{2}n^{i-1/2} \mid (I\cup J \ a \ clique)\right] < \varepsilon.$$

As a consequence of the above claim we also get concentration for $M'(I, J) \mid \mathcal{A}$. This is because $M'(I, J) \mid \mathcal{A}$ is identically distributed as $M(I, J) \mid (I \cup J \text{ a clique})$. Therefore, taking $\varepsilon = 1/n^{2r+1}$ and applying a union bound over all sets I, J we get that with probability at least 1 - 1/n, for all I, J such that $\mathcal{E}(I \cup J) \setminus (\mathcal{E}(I) \cup \mathcal{E}(J)) \subseteq G$, and $|I \cap J| = i$,

$$\left| M'_{r}(I,J) - \beta(i)2^{-\binom{2r}{2} + \binom{2r-i}{2}} \cdot \binom{n-2r+i}{i} \right| < Cr2^{2r^{2}} \cdot k^{2r-i} \cdot (\log n) \cdot n^{i-1/2}.$$

Finally, observe that

$$\beta(i)2^{-\binom{2r}{2}+\binom{2r-i}{2}} \cdot \binom{n-2r+i}{i} = \alpha(i)/p(i),$$

and conditioned on \mathcal{A} , $\Delta(I, J) = M'(I, J) - \alpha(|I \cap J|)/p(|I \cap J|)$. The lemma now follows by combining the above two bounds.

8.4 Putting things together

We now prove Lemma 8.1 and use it to prove our main results.

Proof of Lemma 8.1. By Lemma 7.4, we have that $E \succeq 2^{-Cr^2} k^r n^r \mathbb{I}$. Therefore, by Lemma 8.2 and Lemma 8.11, with probability at least 1 - 2/n, the least eigenvalue of M' is at least

$$2^{-Cr^2}k^r n^r - 2^{O(r^2)}k^{2r} n^r \frac{\log n}{\sqrt{n}} = k^r n^r \left(2^{-O(r^2)} - \frac{2^{O(r^2)}k^r (\log n)}{\sqrt{n}}\right) \ge 0,$$

for k as in the statement of the lemma for a sufficiently big constant c.

We bring the arguments from previous sections together to prove our main results Theorem 2.5 and Theorem 1.5.

Proof of Theorem 2.5. Follows immediately from Lemma 5.1 and Lemma 8.1. \Box

Proof of Theorem 1.5. Follows immediately from Lemma 1.8, Claim 2.2 and Theorem 2.5.

Theorems 1.1 and 1.2 follow immediately from our $\mathsf{PS}(r)$ -refutation lower bound using standard arguments. We defer these to the appendix.

9 Bounding norms of locally random matrices

In this section we shall develop tools for bounding the norms of *locally random* matrices (recall their informal definition from Section 1.4 and more formal one in Section 8.2) associated with random graphs $G \leftarrow G(n, 1/2)$, proving Claim 8.3. The idea behind our bounds is to use the *trace method*. Recall the trace method: for any matrix M, for any positive integer q, $||M|| \leq \sqrt[2q]{tr((M^T M)^q)}$ so we can probabilistically bound ||M|| by bounding $\mathbb{E}\left[tr((M^T M)^q)\right]$.

Going back to Claim 8.3 let us first look at the special case of a = 1 to gain some intuition. In this case, the entries of R_1 are (essentially) independent, and so the trace method is easy to apply. More precisely, R_1 is a symmetric random matrix with zeros on the diagonal and the entries in the upper diagonal taking independent uniformly random ± 1 values. It is well known that $||R_1|| = O(\sqrt{n})$ in this case (see [Ver] for instance). One can also prove the bound by the trace method as follows. We have that

$$\mathbb{E}\left[tr((R_1^T R_1)^q)\right] = \mathbb{E}\left[tr(R_1^{2q})\right] = \sum_{i_1,\cdots,i_{2q}} \mathbb{E}\left[\prod_{j=1}^{2q} R_1(i_j, i_{j+1})\right],$$

where $i_{2q+1} = i_1$. We can then look at which products $\prod_{j=1}^{2q} R_1(i_j, i_{j+1})$ have expectation 0.

Since each individual $R_1(i_j, i_{j+1})$ is an independent ± 1 random variable with expectation 0, a term in the summation $\mathbb{E}\left[\prod_{j=1}^{2q} R_1(i_j, i_{j+1})\right] = 0$ unless every $R_1(i_j, i_{j+1})$ appears an even number of times in the product. Thus, the vast majority of the terms $\mathbb{E}\left[\prod_{j=1}^{2q} R_1(i_j, i_{j+1})\right]$ are 0 and we can count the remaining terms to bound $\mathbb{E}\left[tr(R_1^{2q})\right]$.

One way to implement the above argument is to first look at terms $\mathbb{E}\left[\prod_{j=1}^{2q} R_1(i_j, i_{j+1})\right]$ which have non-zero expectation and observe that in all such terms, the number of distinct entries in $\{i_j : j = 1, \ldots, 2q\}$ is at most q + 1. We can then bound the number of terms with non-zero expected value by the number of possible terms which contain at most q+1 distinct elements. This number can be easily bounded by $O((nq)^{q+1})$, and picking q optimally results in showing that with high probability $||R_1|| = O(\sqrt{n} \log n)$, a near-optimal bound.

To handle higher a's we first generalize the above argument based on constraint graphs to work with general locally-random matrices. However, unlike for a = 1, distinct entries of the matrix are now dependent, which significantly complicates the structure of the terms and the associated count of the terms which have non-zero expectation. The rest of the section is devoted to this. While we apply our arguments to the particular locally-random matrices arising in our proof, these techniques should apply more generally to other locally-random matrices.

9.1 Constraint graphs

We next state our main technical result which gives us a way to bound traces of high powers of locally random matrices based on the structure of the individual terms. The advantage being that the conditions on the terms will be easier to ascertain in our applications.

Here we use V rather than I for subsets because we will be viewing the individual elements of each V as vertices.

Theorem 9.1. Assume that we have values a, B > 0 and for every positive q, we have a function p(G, 2q) such that $p(G, 2q) \ge 0$ and p(G, 2q) can be written in the form

$$p(G, 2q) = \sum_{\{V_1, \dots, V_{2q}\}} f(G, \{V_1, \dots, V_{2q}\})$$

where the following are true:

- 1. $V_j \subseteq V(G)$ and $|V_j| = a$.
- 2. For every term $f(G, \{V_1, \ldots, V_{2q}\})$ with non-zero expected value, $|\cup_j V_j| \leq 2aq qy + z$ for some integers y and z where $1 \leq y \leq 2a$ and $z \geq 0$.
- 3. $\mathbb{E}[f(G, \{V_1, \dots, V_{2q}\})] \le B^{2q}$.

Then, if $n \ge 10$, for all $\varepsilon \in (0, 1)$,

$$\Pr\left[|\min_{q\in\mathbb{Z}^+}\left\{\sqrt[2q]{p(G,2q)}\right\}| > \frac{B}{a!} \cdot \left(2ea\left(\frac{\ln(n^z/\varepsilon)}{2y} + 1\right)\right)^y \cdot n^{a-y/2}\right] < \varepsilon.$$

Remark 9.2. We will use this theorem with two types of functions p. When $p(G, 2q) = tr((M^T M)^q)$ for some matrix M depending on G, $||M|| \leq \sqrt[2q]{p(G, 2q)}$ for all q > 0 so this theorem gives us a probabilistic bound on ||M||. When $p(G, 2q) = h(G)^{2q}$ for some function h, then $h(G) = \sqrt[2q]{p(G, 2q)}$ for all q > 0 so this theorem gives us a probabilistic bound on h(G).

Example 9.3. In the case when $p(G, 2q) = tr(R_1^{2q})$, $p(G, 2q) = \sum_{i_1, \dots, i_{2q}} \prod_{j=1}^{2q} R_1(i_j, i_{j+1})$. Each term here has expected value at most 1 and it is easy to argue that for any term with non-zero expected value, the number of distinct elements is at most q + 1. Applying Theorem 9.1 with y = z = 1, and B = 1 we have that for all $n \ge 10$, and $\varepsilon \in (0, 1)$,

$$\Pr\left[||R_1|| > 2e\sqrt{n}(\ln\left(n/\varepsilon\right) + 2)\right] < \varepsilon$$

This bound is weaker (by a logarithmic factor) than the bounds in e.g. [Ver], but is sufficient for our purposes.

Before proving the theorem we introduce the concept of *constraint graphs* which are a useful way to visualize our calculations. While the statement of the above theorem does not involve constraint graphs, thinking in terms of constraint graphs is helpful in proving the conditions required to apply the theorem.

Definition 9.4. Given a family of sets of vertices $\{V_i\}$, we define a corresponding constraint graph C whose vertices are the sets $\{V_i\}$ and there is an edge between $V_i, V_j, i \neq j$, if $V_i \cap V_j \neq \emptyset$.

The above definition is useful because of the following elementary lemma.

Lemma 9.5. For any collection of sets $\{V_1, \ldots, V_\ell\}$, if the corresponding constraint graph C has t connected components, then $|\cup_i V_i| < \sum_i |V_i| - \ell + t$.

Proof. Let V_{i_1}, \ldots, V_{i_t} belong to the *t* different connected components of *C*. Now add the remaining elements of $\{V_1, \ldots, V_\ell\}$ so that each new set is adjacent (in *C*) to at least one of the previously added sets (we can do this as the number of connected components is *t*). Then, each such step adding a set V_i can increase the size of the union by at most $|V_i| - 1$. Therefore, the size of the union is at most $\sum_i |V_i| - \ell + t$.

Proof of Theorem 9.1. In the following we use $\{V_i\}$ as a short form for $\{V_1, \ldots, V_{2q}\}$. We prove this result by obtaining an upper bound on the number of terms in $p(G, 2q) = \sum_{\{V_i\}} f(G, \{V_i\})$ with nonzero expected value. This gives us a probabilistic upper bound for p(G, 2q), implying the upper bound on $\min_q \{ \sqrt[2q]{p(G, 2q)} \}$.

Definition 9.6. Define N(n, a, q, m) to be the number of ways to choose subsets $\{V_i : i \in [2q]\}$ of [n] such that $|\cup_i V_i| \leq m$ and for all $i, |V_i| = a$.

Lemma 9.7. If $m \leq 2aq$, then

$$N(n, a, q, m) \le \left(\frac{1}{a!}\right)^{2q} \binom{2aq}{2aq-m} n^m m^{2aq-m}$$

Proof. We can choose each ordered 2aq-tuple (v_1, \dots, v_{2aq}) of elements in [n] which contains at most m distinct elements as follows. There must be at least 2aq - m elements which are duplicates of other elements, so we can first choose a set I of 2aq - m indices such that for all $i \in I$, $v_i = v_j$ for some $j \notin I$. There are $\binom{2aq}{2aq-m}$ choices for I. We then choose the elements $\{v_j : j \notin I\}$. There are no restrictions on these elements so there are n^m choices for these elements. Finally, we choose the elements $\{v_i : i \in I\}$. To determine each v_i it is sufficient to specify the $j \notin I$ such that $v_i = v_j$. For each i there are m choices for the corresponding j, so the number of choices for these elements is at most m^{2aq-m} . Putting everything together, the total number of choices is at most

 $\binom{2aq}{2aq-m}n^mm^{2aq-m}$. Now note that since we are choosing subsets $\{V_i : i \in [2q]\}$ of [n] rather than one big ordered tuple, the order within each subset does not matter. Thus, there are $(a!)^{2q}$ different ordered tuples which give the same subsets of elements, so the total number of possibilities for the subsets $\{V_i\}$ is at most $(a!)^{-2q} \binom{2aq}{2aq-m} n^mm^{2aq-m}$, as needed.

Now $\mathbb{E}[p(G, 2q)] = \sum_{\{V_i\}} \mathbb{E}[f(G, \{V_i\})]$. For every nonzero term $\mathbb{E}[f(G, \{V_i\})]$, we have that $|\cup_i V_i| \leq 2aq - qy + z$. If $q > \frac{z}{y}$ then applying Lemma 9.7 with m = 2aq - qy + z, the number of non-zero terms $\mathbb{E}[f(G, \{V_i\})]$ is at most

$$\left(\frac{1}{a!}\right)^{2q} \binom{2aq}{2aq-m} n^m m^{2aq-m} \le \left(\frac{1}{a!}\right)^{2q} (2aqm)^{2aq-m} n^m$$

Moreover, by our assumptions, each of these nonzero terms $E[f(G, \{V_i\})]$ has value at most B^{2q} , so

$$\mathbb{E}[p(G,2q)] \le \left(\frac{1}{a!}\right)^{2q} (2aqm)^{2aq-m} n^m B^{2q}.$$

Now, by Markov's inequality applied to p(G, 2q),

$$\Pr\left[\sqrt[2q]{p(G,2q)} > \sqrt[2q]{\mathbb{E}[p(G,2q)]/\varepsilon}\right] < \varepsilon.$$

We next choose a value q so as to minimize our estimate on $\sqrt[2q]{\mathbb{E}[p(G,2q)]/\varepsilon}$. Specifically, we set $q = \lceil \ln(n^z/\varepsilon)/2y \rceil$ (we arrive at this value by minimizing the general estimate as a function of q by setting the derivative to 0 - we spare the reader the details). As long as $n \ge 10$, this guarantees that q > z/y so that

$$\begin{split} {}^{2q} \sqrt{\mathbb{E}[p(G,2q)]/\varepsilon} &\leq \frac{B}{a!} \cdot \frac{1}{\varepsilon^{1/2q}} \cdot (2aqm)^{a-m/2q} \cdot n^{m/2q} \\ &= \frac{B}{a!} \cdot n^{a-y/2} \cdot \left(\frac{n^z}{\varepsilon}\right)^{1/2q} (2aqm)^{a-m/2q} \\ &\leq \frac{B}{a!} \cdot n^{a-y/2} \cdot \left(\frac{n^z}{\varepsilon}\right)^{1/2q} \cdot (2aq)^y \\ &\leq \frac{B}{a!} \cdot n^{a-y/2} \cdot e^y \cdot (2a)^y \cdot \left(\frac{\ln(n^z/\varepsilon)}{2y} + 1\right)^y \end{split}$$

The claim now follows by rearranging the above bound.

9.2 Bounds on $||R_a||$

In this subsection, we prove Claim 8.3 using Theorem 9.1. For convenience, we restate Claim 8.3 here with more precise constants.

Theorem 9.8. If
$$n \ge 100$$
, for all $\varepsilon \in (0,1)$, $\Pr\left[||R_a|| > 2^{a^2+2a+2} \ln\left(\frac{n}{\varepsilon}\right)n^{a-\frac{1}{2}}\right] < \varepsilon$.

The core of the proof will be to bound $|\bigcup_{j=1}^{2q} V_{i_j}|$ for any term $\prod_{j=1}^{2q} R_a(V_{i_j}, V_{i_{j+1}})$ with non-zero expectation which appear in the expansion of $tr((R_a^T R_a)^q)$. We will do so by arguing that the constraint graph associated with the term has at most 2aq - q + 1 connected components, which we do by inductively decomposing R_a as follows.

Definition 9.9. Given a partition (A, B) of [1, n], define $R_{a,A,B}(V_1, V_2) = R_a(V_1, V_2)$ if $V_1 \subseteq A$ and $V_2 \subseteq B$ and 0 otherwise.

Proposition 9.10. $\sum_{A,B} R_{a,A,B} = 2^{n-2a} R_a$

Proof. $R_{a,A,B}(V_1, V_2) = R_a(V_1, V_2) = 0$ whenever V_1 and V_2 are not disjoint. For all disjoint V_1 and V_2 , $R_{a,A,B}(V_1, V_2) = R_a(V_1, V_2)$ for 2^{n-2a} choices of A and B and is 0 for the rest.

Corollary 9.11. $||R_a|| \le 2^{2a} \max_{A,B} \{||R_{a,A,B}||\}$

Proof. Since $R_a = 2^{2a-n} \sum_{A,B} R_{a,A,B}, ||R_a|| \le 2^{2a-n} \sum_{A,B} ||R_{a,A,B}|| \le 2^{2a} \max_{A,B} \{||R_{a,A,B}||\}$

Now given A and B, take

$$p(G,2q) = tr((R_{a,A,B}^T R_{a,A,B})^q) = \sum_{\substack{\{V_{ij}:i \in [1,q], j \in [1,2]\}\\ i=1}} \prod_{i=1}^q R_{a,A,B}^T(V_{i1}, V_{i2}) R_{a,A,B}(V_{i2}, V_{(i+1)1}) = \sum_{\substack{\{V_{ij}:i \in [1,q], j \in [1,2]\}:\\ \forall i, V_{i1} \subseteq B\\ \forall i, V_{i2} \subseteq A}} \prod_{i=1}^q R_a(V_{i1}, V_{i2}) R_a(V_{i2}, V_{(i+1)1})$$

where we take $V_{(q+1)1} = V_{11}$.

To simplify this expression, rename the sets of vertices as follows.

Definition 9.12.

- 1. If $i \in [1, 2q]$ and i is odd then take $W_i = V_{(\frac{i+1}{2})1}$
- 2. If $i \in [1, 2q]$ and i is even then take $W_i = V_{(\frac{i}{2})2}$

We now have that

$$p(G, 2q) = \sum_{\substack{\{W_i: i \in [1, 2q]\}: \\ \forall \text{ odd } i, W_i \subseteq B \\ \forall \text{ even } i, W_i \subseteq A}} \prod_{i=1}^{2q} R_a(W_i, W_{i+1}),$$
(9.1)

where we take $W_{2q+1} = W_1$. To study which of these terms may have non-zero expectation, we first define a graph related to the corresponding constraint graph.

Definition 9.13. Given a constraint graph C, let H be a graph with two types of edges, product edges and constraint edges, such that

- 1. $V(H) = \{W_i : i \in [1, 2q]\}$
- 2. $E_P(H) = \{(W_i, W_{i+1}) : i \in [1, 2q]\}$
- 3. $E_C(H) = \{(W_i, W_j) : i \neq j, W_i \cap W_j \neq \emptyset\}$

Now, each $R_a(W_i, W_{i+1})$ is a random variable with expectation 0, so if any $R_a(W_i, W_{i+1})$ is independent from everything else, the product will have expectation 0. Such dependencies arise due to the presence of edges from G occurring in (at least two) different "elements" (say (W_i, W_{i+1}) , (W_j, W_{j+1}) for $i \neq j$) of the term. Such repeated occurrences manifest in our constraint graphs (and the graph H defined above) as (three or four) cycles in the graph, which we call independence breaking. For a term to have non-zero expectation it must be that every element (W_i, W_{i+1}) is on some such cycle. This implies that each product $\prod_{i=1}^{2q} R_a(W_i, W_{i+1})$ has zero expected value unless all of the product edges in the corresponding H are part of independence-breaking cycles. This places restrictions on H (see Lemma 9.17) which in turn places restrictions on the constraint graph C, allowing us to use Theorem 9.1. We make these ideas precise below.

Definition 9.14. Given q and $\{W_1, \dots, W_{2q}\}$, we define $W_{i\pm 2q} = W_i$ for all $i \in [1, 2q]$.

Definition 9.15. If $q \ge 2$,

- 1. Define an independence breaking 3-cycle in H to consist of product edges $(W_i, W_{i+1}), (W_{i+1}, W_{i+2})$ and a constraint edge $((W_i, j), (W_{i+2}, j))$.
- 2. Define an independence breaking 4-cycle to consist of product edges $e_1 = (W_{i_1}, W_{i_1+1}), e_2 = (W_{i_2}, W_{i_2\pm 1})$ and constraint edges (W_{i_1}, W_{i_2}) and $(W_{i_1+1}, W_{i_2\pm 1})$.

Proposition 9.16. For all W_1, \dots, W_{2q} such that $W_i \subseteq B$ whenever *i* is odd and $W_i \subseteq A$ whenever *i* is even, if the corresponding *H* has a product edge (W_i, W_{i+1}) which is not contained in any independence-breaking cycle then $E[\prod_{i=1}^{2q} R_a(W_i, W_{i+1})] = 0$

Proof. If (W_i, W_{i+1}) is not contained in any independence-breaking cycle then no edge between W_i and W_{i+1} appears anywhere else so $R_a(W_i, W_{i+1})$ is a random variable with expectation 0 which is independent from everything else and thus $E[\prod_{i=1}^{2q} R_a(W_i, W_{i+1})] = 0.$

We now bound the number of connected components in H with the following lemma.

Lemma 9.17. Let $q \ge 2$ and H be a graph such that

- 1. Every product edge of H is contained in an independence-breaking cycle.
- 2. Every constraint edge of H is of the form (W_i, W_{i+j}) where j is even.

Then, the number of connected components in the graph defined by only the constraint edges of H is at most q + 1.

The intuitive idea behind this lemma is that if we add the constraint edges in the right order, every new constraint edge can put two product edges into independence breaking cycles. For example, a constraint edge between W_{i-1} and W_{i+1} puts the product edges (W_{i-1}, W_i) and (W_i, W_{i+1}) into an independence breaking 3-cycle. If we then add a constraint edge between W_{i-2} and W_{i+2} , this puts the product edges (W_{i-2}, W_{i-1}) and (W_{i+1}, W_{i+2}) into an independence breaking 4-cycle. The final constraint edge can put 4 product edges into independence breaking cycles, so the number of constraint edges needed is q - 1.

To make this argument work, we use an inductive proof. We note that if there is no W_i which is isolated in H, we must have at least q constraint edges. On the other hand, if there a W_i which is isolated, there must be a constraint edge between W_{i-1} and W_{i+1} . As noted above, this constraint edge puts the product edges (W_{i-1}, W_i) and (W_i, W_{i+1}) into an independence breaking 3-cycle. We take this to be the first constraint edge. We then argue that we can essentially delete W_i and merge W_{i-1} and W_{i+1} which allows us to use the inductive hypothesis. We make these ideas rigorous below.

Proof of Lemma 9.17. We prove Lemma 9.17 by induction on q. The base case q = 2 is trivial, as we clearly need at least one constraint edge, so the number of connected components in H is at most 3. Now assume that $q = k \ge 3$ and the result is true for q = k - 1.

First note that if there is no W_i which is isolated (when looking only at constraint edges), then there are at most q connected components in H. Thus, we may assume that W_i is isolated for some i. Now note that for the product edge (W_{i-1}, W_i) , since W_i is isolated, there are no independence breaking 3-cycles or 4-cycles where W_i is the endpoint of a constraint edge. Thus, we must have that (W_{i-1}, W_i) is part of an independence breaking 3-cycle consisting of (W_{i-1}, W_i) , (W_i, W_{i+1}) , and a constraint edge (W_{i-1}, W_{i+1}) .

Now form a new graph H' as follows. Delete W_i and contract the constraint edge between W_{i-1} and W_{i+1} . More precisely,

1. Take
$$V(H') = V(H) \setminus \{W_{i-1}, W_i, W_{i+1}\} \cup \{U\}$$

- 2. Take $E_{product}(H') = E_{product}(H) \setminus \{(W_j, W_{j+1}) : j \in [i-2, i+1]\} \cup \{(W_{i-2}, U), (U, W_{i+2})\}$
- 3. Take

$$E_{constraint}(H') = E_{constraint}(H) \setminus \{(W_{i-1}, W_j) : (W_{i-1}, W_j) \in E_{constraint}(H)\}$$
$$\setminus \{(W_{i+1}, W_j) : (W_{i+1}, W_j) \in E_{constraint}(H)\}$$
$$\cup \{(U, W_j) : (W_{i-1}, W_j) \in E_{constraint}(H) \text{ or } (W_{i+1}, W_j) \in E_{constraint}(H)\}$$

After doing this, rename U as W_{i-1} and rename each W_j where j > i + 1 as W_{j-2} . In going from H to H', we have effectively reduced both q and the number of connected components by 1. To complete the proof, we need to check that H' satisfies the inductive hypotheses. Based on the reduction from H to H', we still have that every constraint edge is of the form (W_i, W_{i+j}) where j is even. We check that every product edge is still part of an independence-breaking cycle case by case.

- 1. Every independence-breaking cycle which did not contain the constraint edge (W_{i-1}, W_{i+1}) in H is preserved in H' except that the vertices may have been renamed. The reason for this is that such an independence breaking cycle in H cannot contain W_i and can contain at most one of $\{W_{i-1}, W_{i+1}\}$.
- 2. The independence-breaking 3-cycle in H consisting of the product edges (W_{i-1}, W_i) , (W_i, W_{i+1}) and the constraint edge (W_{i-1}, W_{i+1}) is removed, but so are the product edges (W_{i-1}, W_i) and (W_i, W_{i+1}) , so this is fine.
- 3. If we have an independence breaking 4-cycle in H consisting of the product edges (W_{i-2}, W_{i-1}), (W_{i+1}, W_{i+2}) and the constraint edges (W_{i-1}, W_{i+1}), (W_{i-2}, W_{i+2}), this becomes an independence-breaking 3-cycle in H' with product edges (W_{i-2}, W_{i-1}), (W_{i-1}, W_i) and a constraint edge (W_{i-2}, W_i) (note that W_{i-1} and W_{i+1} are merged into W_{i-1} in H' and W_{i+2} is renamed as W_i in H').

H' satisfies the inductive hypotheses, so looking only at the constraint edges, H' has at most (q-1)+1=q connected components. H has one more connected component than H' (the vertex W_i in H), so H has at most q+1 connected components, as needed.

The above lemma combined with Lemma 9.5 gives the following corollary.

Corollary 9.18. For all terms $\prod_{i=1}^{2q} R_a(W_i, W_{i+1})$ occurring in Equation 9.1 with nonzero expectation, $|\bigcup_{i=1}^{2q} W_i| \leq 2aq - (2q) + q + 1$.

We can now prove Theorem 9.8

Proof of Theorem 9.8. We can now apply Theorem 9.1 with y = 1, and z = 1 by the above corollary. Every entry of $R_{a,A,B}$ has magnitude at most 2^{a^2} so we can take $B = 2^{a^2}$. By Theorem 9.1, if $n \ge 10$, for all A and B, for every $\varepsilon \in (0, 1)$,

$$\Pr\left[||R_{a,A,B}|| > \frac{2^{a^2}}{a!} \left(2ea\left(\frac{\ln n - \ln \varepsilon}{2} + 1\right)\right) n^{a-1/2}\right] = \Pr\left[||R_{a,A,B}|| > \frac{2^{a^2}ea}{a!} (\ln n - \ln \varepsilon + 2) n^{a-\frac{1}{2}}\right] < \varepsilon$$

Since $4 \ln n \ge e(\ln n + 2)$ for all $n \ge 100$ and $a! \ge a$, we have that for all $n \ge 100$, for all A and B and all $\varepsilon \in (0, 1)$,

$$\Pr\left[||R_{a,A,B}|| > 2^{a^2+2} \ln\left(\frac{n}{\varepsilon}\right) n^{a-\frac{1}{2}}\right] < \varepsilon$$

Now by Corollary 9.11, $||R_a|| \le 2^{2a} \max_{A,B} \{||R_{a,A,B}||\}$ so

$$\Pr\left[||R_a|| > 2^{a^2 + 2a + 2} \ln\left(\frac{n}{\varepsilon}\right) n^{a - \frac{1}{2}}\right] < \varepsilon$$

10 Concentration bounds for number of cliques and $deg_G(I)$

We now prove large deviation bounds for $deg_G()$ leading to Claim 8.13 which we state below in a more precise form.

Theorem 10.1. If $n \ge 10$, and $\varepsilon \in (0,1)$, then for all $I \subseteq [n]$, with $|I| = i \le 2r$,

$$\Pr\left[\left|\deg_{G}(I) - 2^{-\binom{2r}{2} + \binom{i}{2}} \cdot \binom{n-i}{2r-i}\right| > 2(\ln(128/\varepsilon))^{2}n^{2r-i-1/2} \mid (I \text{ is a clique})\right] \le \varepsilon.$$

To prove the claim we first show a similar concentration bound for the number of cliques of a certain size in G. While similar results appear in the literature, see for instance [Ruc88, Vu01, JLR11], we give a short direct proof based on Theorem 9.1.

Definition 10.2. For a graph G, define $N_a(G)$ to be the number of a-cliques in G.

Theorem 10.3. For all a, for all $n \ge 10$ and $\varepsilon \in (0,1)$, $E[N_a(G)] = 2^{-\binom{a}{2}}\binom{n}{a}$ and

$$\Pr\left[|N_a(G) - E[N_a(G)]| > (\ln(64/\varepsilon))^2 \cdot n^{a-1}\right] < \varepsilon.$$

Proof. The first part of the theorem is trivial so we focus on the second part. Given a set of vertices V of size a, define c_V to be $1 - 2^{-\binom{a}{2}}$ if V is a clique and $-2^{-\binom{a}{2}}$ otherwise. Then,

$$N_a(G) - E[N_a(G)] = \sum_{V:|V|=a} c_V$$

Now let's consider the function $p(G, 2q) = (\sum_{V:|V|=a} c_V)^{2q} = \sum_{W_1, \dots, W_{2q}} \prod_{i=1}^{2q} c_{W_i}.$

Note that $E[\prod_{i=1}^{2q} c_{W_i}] = 0$ unless each set of vertices W_i has two vertices in common with a different set of vertices W_j . Now consider a graph C_2 where the vertices are $\{W_1, \ldots, W_{2q}\}$ and an edge between W_i, W_j if $|W_i \cap W_j| \ge 2$. Let t be the number of connected components in C_2 . We claim that $|\cup_i W_i| \le 2aq - 4q + 2t$. For, as in the proof of Lemma 9.5, first consider elements W_{i_1}, \ldots, W_{i_t} belonging to the t different connected components. Now, add the remaining elements of $\{W_1, \ldots, W_{2q}\}$ so that each new element is adjacent to at least one of the previously added sets. When doing so, each step can increase the size of the union by at most a - 2. Therefore, the size of the union is at most at + (a - 2)(2q - t) = 2aq - 4q + 2t. On the other hand, each connected component in C_2 must have at least two vertices, so $t \le q$. Therefore, $|\cup_i W_i| \le 2aq - 2q$.

We can now apply Theorem 9.1 with y = 2, z = 0 and B = 1 so that for $n \ge 10$, and $\varepsilon \in (0, 1)$,

$$\Pr\left[|N_a(G) - E[N_a(G)]| > \frac{1}{a!} \cdot \left(ea\left(\frac{-\ln\varepsilon}{4} + 1\right)\right)^2 \cdot n^{a-1}\right] < \varepsilon.$$

Using the facts that $e^2 < 8$ and $\frac{m^2}{m!} \leq 2$ for all nonnegative integers m, we have that

$$\Pr\left[|N_a(G) - E[N_a(G)]| > (\ln(64/\varepsilon))^2 \cdot n^{a-1}\right] < \varepsilon.$$

We are now ready to prove Theorem 10.1. The idea is as follows. Let A_I be the collection of vertices which are adjacent to all the vertices in I. Then, conditioned on I being a clique, $deg_G(I)$ is just the number of cliques of size 2r - i in the vertices A_I which is primarily determined by $|A_I|$. This is because the edges between vertices of A_I are independent of the edges involving vertices in I so that we can apply Theorem 10.3 to A_I .

Proof of Theorem 10.1. Let A_I be as above and let us condition on I being a clique. Then, $deg_G(I)$ is just the number of cliques of size 2r - i among the vertices in A_I . Therefore, by Theorem 10.3, with probability at least $1 - \varepsilon/2$,

$$\left| deg_G(I) - 2^{-\binom{2r-i}{2}} \binom{|A_I|}{2r-i} \right| \le (\ln(128/\varepsilon))^2 \cdot n^{2r-i-1}.$$

We next argue that $\binom{|A_I|}{2r-i}$ is concentrated around its mean. For $j \notin I$, let X_j be the indicator random variable that is 1 if the *j*'th vertex is adjacent to all the vertices in *I* and 0 otherwise. Then, $|A_I| = \sum_{j \notin I} X_j$ and

$$\binom{|A_I|}{2r-i} = \sum_{J \subseteq [n] \setminus I, |J| = 2r-i j \in J} \prod_{X_j \in J} X_j \equiv f(\{X_j : j \notin I\}).$$

Observe that the random variables X_i are independent of each other and that

$$\mathbb{E}[f(\{X_j : J \notin I\})] = 2^{-i(2r-i)} \binom{n-i}{2r-i}.$$

We next apply McDiarmid's inequality to the function f. Note that changing any single coordinate of the inputs to f can change its value by at most n^{2r-i-1} . Therefore, by Theorem 4.2, with probability at least $1 - \varepsilon/2$,

$$\left| \binom{|A_I|}{2r-i} - 2^{-i(2r-i)} \binom{n-i}{2r-i} \right| \le \sqrt{\ln(4/\varepsilon)} \cdot n^{2r-i-.5}.$$

Combining the above equations, we get that with probability at least $1 - \varepsilon$,

$$\left| deg_G(I) - 2^{-\binom{2r-i}{2} - i(2r-i)} \binom{n-i}{2r-i} \right| \le (\ln(128/\varepsilon))^2 \cdot n^{2r-i-1} + 2^{-\binom{2r-i}{2}} \cdot \sqrt{\ln(4/\varepsilon)} \cdot n^{2r-i-.5} \le 2\ln((128/\varepsilon)^2) \cdot n^{2r-i-.5}.$$

The theorem now follows as $\binom{2r-i}{2} + i(2r-i) = \binom{2r}{2} - \binom{i}{2}$.

11 Conclusion and future work

In this work we showed a lower bound for the maximum clique problem on random G(n, 1/2) graphs in the SOS hierarchy and positivstellensatz proof system. Besides the specific application to clique lower bounds, the PSD'ness of the matrix M from Equation 2.5 seems to carry further information that could be potentially useful elsewhere, perhaps for studying various sub-graph statistics. Further, the arguments related to association schemes and bounding the norm of locally random matrices could also be useful elsewhere, especially for other SOS hierarchy lower bounds. One natural and interesting candidate is the densest subgraph problem.

For planted clique itself, the most obvious open problem is to tighten the gap between the current upper bound of $O(\sqrt{n}/2^r)$ and our lower bound of $2^{-O(r)}(\sqrt{n}/\log n)^{1/r}$ for r rounds of the SOS hierarchy. In particular, can a constant number of rounds of SOS beat the square-root barrier and identify planted cliques of size $o(n^{1/2})$? Kelner¹² showed that our dual certificate M actually is not PSD for k roughly $O(n^{1/(r+1)})$. Thus one needs to come up with a different dual certificate to approach the upper bound of \sqrt{n} even for r = 2.

Acknowledgements We thank Boaz Barak, Siu-on Chan, Jonathan Kelner, Robert Krauthgamer, James Lee, Nati Linial, David Steurer, Madhu Sudan and Amir Yehudayoff for several useful comments.

References

[AAK⁺07] Noga Alon, Alexandr Andoni, Tali Kaufman, Kevin Matulef, Ronitt Rubinfeld, and Ning Xie. Testing k-wise and almost k-wise independence. In Proceedings of the 39th Annual ACM Symposium on Theory of Computing, San Diego, California, USA, June 11-13, 2007, pages 496–505, 2007.

¹²Personal comminication

- [ABBG10] Sanjeev Arora, Boaz Barak, Markus Brunnermeier, and Rong Ge. Computational complexity and information asymmetry in financial products. In *ICS*, pages 49–65, 2010.
- [ABS10] Sanjeev Arora, Boaz Barak, and David Steurer. Subexponential algorithms for unique games and related problems. In *FOCS*, pages 563–572, 2010.
- [ABW10] Benny Applebaum, Boaz Barak, and Avi Wigderson. Public-key cryptography from different assumptions. In Proceedings of the forty-second ACM symposium on Theory of computing, pages 171–180. ACM, 2010.
- [AKS98] Noga Alon, Michael Krivelevich, and Benny Sudakov. Finding a large hidden clique in a random graph. *Random Struct. Algorithms*, 13(3-4):457–466, 1998.
- [Art27] Emil Artin. Über die zerlegung deniter funktionen in quadrate. Abhandlungen aus dem Mathematischen Seminar der Universitat Hamburg, 5(1):100–115, 1927.
- [BBH⁺12] Boaz Barak, Fernando G. S. L. Brandão, Aram Wettroth Harrow, Jonathan A. Kelner, David Steurer, and Yuan Zhou. Hypercontractivity, sum-of-squares proofs, and their applications. In STOC, pages 307–326, 2012.
- [BCR98] Jacek Bochnak, Michel Coste, and Marie-Francoise Roy. *Real algebraic geometry*. Springer, 1998.
- [BCV⁺12] Aditya Bhaskara, Moses Charikar, Aravindan Vijayaraghavan, Venkatesan Guruswami, and Yuan Zhou. Polynomial integrality gaps for strong SDP relaxations of densest k-subgraph. In SODA, pages 388–405, 2012.
- [BR13] Q. Berthet and P. Rigollet. Complexity theoretic lower bounds for sparse principal component detection. J. Mach. Learn. Res., W and CP, 30:1046–1066 (electronic), 2013.
- [BRS11] Boaz Barak, Prasad Raghavendra, and David Steurer. Rounding semidefinite programming hierarchies via global correlation. In *FOCS*, pages 472–481, 2011.
- [DGGP14] YAEL DEKEL, ORI GUREL-GUREVICH, and YUVAL PERES. Finding hidden cliques in linear time with high probability. *Combinatorics, Probability and Computing*, 23:29–49, 1 2014.
- [DM15] Yash Deshpande and Andrea Montanari. Improved sum-of-squares lower bounds for hidden clique and hidden submatrix problems. *CoRR*, abs/1502.06590, 2015.
- [FGR⁺13] Vitaly Feldman, Elena Grigorescu, Lev Reyzin, Santosh Vempala, and Ying Xiao. Statistical algorithms and a lower bound for detecting planted cliques. In STOC, pages 655–664, 2013.
- [FK00] Uriel Feige and Robert Krauthgamer. Finding and certifying a large hidden clique in a semirandom graph. *Random Struct. Algorithms*, 16(2):195–208, 2000.
- [FK03] Uriel Feige and Robert Krauthgamer. The probable value of the Lovász–Schrijver relaxations for maximum independent set. *SIAM J. Comput.*, 32(2):345–370, 2003.

- [FK08] Alan M. Frieze and Ravi Kannan. A new approach to the planted clique problem. In IARCS Annual Conference on Foundations of Software Technology and Theoretical Computer Science, FSTTCS 2008, December 9-11, 2008, Bangalore, India, pages 187– 198, 2008.
- [God] Chris Godsil. Association schemes. Lecture Notes available at http://quoll.uwaterloo.ca/mine/Notes/assoc1.pdf.
- [Gri01a] Dima Grigoriev. Complexity of positivstellensatz proofs for the knapsack. Computational Complexity, 10(2):139–154, 2001.
- [Gri01b] Dima Grigoriev. Linear lower bound on degrees of positivstellensatz calculus proofs for the parity. *Theor. Comput. Sci.*, 259(1-2):613–622, 2001.
- [GS11] Venkatesan Guruswami and Ali Kemal Sinop. Lasserre hierarchy, higher eigenvalues, and approximation schemes for graph partitioning and quadratic integer programming with PSD objectives. In *FOCS*, pages 482–491, 2011.
- [GV01] Dima Grigoriev and Nicolai Vorobjov. Complexity of null-and positivstellensatz proofs. Ann. Pure Appl. Logic, 113(1-3):153–160, 2001.
- [GVL96] G.H. Golub and C.F. Van Loan. *Matrix Computations*. Johns Hopkins Studies in the Mathematical Sciences. Johns Hopkins University Press, 1996.
- [Jer92] Mark Jerrum. Large cliques elude the metropolis process. *Random Struct. Algorithms*, 3(4):347–360, 1992.
- [JLR11] S. Janson, T. Luczak, and A. Rucinski. *Random Graphs*. Wiley Series in Discrete Mathematics and Optimization. Wiley, 2011.
- [Kar76] R. M. Karp. Probabilistic analysis of some combinatorial search problems. In: Algorithms and Complexity: New Directions and Recent Results, pages 1–19, 1976.
- [Kri64] Jean-Louis Krivine. Anneaux preordonnes. Journal d'Analyse Mathematique, 12(1):307–326, 1964.
- [Kuc95] Ludek Kucera. Expected complexity of graph partitioning problems. *Discrete Applied Mathematics*, 57(2-3):193–212, 1995.
- [Las01] Jean B. Lasserre. Global optimization with polynomials and the problem of moments. SIAM Journal on Optimization, 11(3):796–817, 2001.
- [LS91] L. Lovász and A. Schrijver. Cones of matrices and set-functions and 0-1 optimization. SIAM Journal on Optimization, 1(2):166–190, 1991.
- [MW13] Raghu Meka and Avi Wigderson. Association schemes, non-commutative polynomial concentration, and sum-of-squares lower bounds for planted clique. *CoRR*, abs/1307.7615, 2013.
- [OZ13] Ryan O'Donnell and Yuan Zhou. Approximability and proof complexity. In SODA, pages 1537–1556, 2013.

- [Par00] Pablo Parrilo. Structured semidefinite programs and semialgebraic geometry methods in robustness and optimization, 2000. PhD thesis, California Institute of Technology.
- [PS⁺00] Pavel A Pevzner, Sing-Hoi Sze, et al. Combinatorial approaches to finding subtle signals in dna sequences. In *ISMB*, volume 8, pages 269–278, 2000.
- [Put93] Mihai Putinar. Positive polynomials on compact semi-algebraic sets. Indiana University Mathematics Journal, 42(3):969–984, 1993.
- [Ruc88] Andrzej Ruciski. When are small subgraphs of a random graph normally distributed? *Probability Theory and Related Fields*, 78(1):1–10, 1988.
- [SA90] H. Sherali and W. Adams. A hierarchy of relaxations between the continuous and convex hull representations for zero-one programming problems. SIAM Journal on Discrete Mathematics, 3(3):411–430, 1990.
- [Sch91] Konrad Schmudgen. The k-moment problem for compact semi-algebraic sets. *Mathe*matische Annalen, 289(1):203–206, 1991.
- [Sch08] Grant Schoenebeck. Linear level Lasserre lower bounds for certain k-CSPs. In *FOCS*, pages 593–602, 2008.
- [Ste73] Gilbert Stengle. A nullstellensatz and a positivstellensatz in semialgebraic geometry. Mathematische Annalen, 207(2):87–97, 1973.
- [Tul09] Madhur Tulsiani. CSP gaps and reductions in the Lasserre hierarchy. In *STOC*, pages 303–312, 2009.
- [Ver] Roman Vershynin. Lecture 6: Norm of a random matrix. Lecture Notes on Non-Asymptotic Random Matrix Theory - Available online.
- [vLW01] J.H. van Lint and R.M. Wilson. A Course in Combinatorics. Cambridge University Press, 2001.
- [Vu01] Van Vu. A large deviation result on the number of small subgraphs of a random graph. Combinatorics, Probability and Computing, 10:79–94, 1 2001.

12 Hierarchy Gaps and Positivstellensatz Refutations

For a detailed discussion of the hierarchies and PS(r)-refutations we refer the reader to the discussions in [OZ13]. The basic principle is that, typically, PS(r)-refutations are more robust and stronger than the hierarchy formulations.

The SOS (or Lasserre) relaxation for maximum clique is stated in Figure 12 (cf. [Tul09]). Although, the formulation itself is not in terms of an SDP, it is a standard fact that as the program only involves inner products of vectors, the optimization can be done by semi-definite programming. The connection between Figure 12 and PS(r)-refutations comes from the following straightforward lemma stating that a certificate for PS(r)-refutations is simply a primal solution to the standard *r*-round SOS-relaxation of the problem.

SOS-relaxation for Max-Clique. Input: Graph G = (V, E), r - number of rounds. Variables of the SDP are vectors U_S , where $S \subseteq [n], |S| \leq r$.

maximize
$$\sum_{i \in V} \| \boldsymbol{U}_{\{i\}} \|_2^2$$
.

such that

 $\langle \boldsymbol{U}_{\{i\}}, \boldsymbol{U}_{\{j\}} \rangle = 0,$ $\forall i, j, \{i, j\} \notin E$ $\langle \boldsymbol{U}_{S_1}, \boldsymbol{U}_{S_2} \rangle = \langle \boldsymbol{U}_{S_3}, \boldsymbol{U}_{S_4} \rangle, \qquad S_1 \cup S_2 = S_3 \cup S_4, |S_1 \cup S_2| \le r$ $\langle \boldsymbol{U}_{S_1}, \boldsymbol{U}_{S_2} \rangle \in [0,1],$ $|S_1|, |S_2| \leq r$ $\|U_{\emptyset}\|_{2}^{2} = 1$

Figure 1: r-round SOS-relaxation for Maximum Clique

Lemma 12.1. Let G = (V, E) be a graph and let Clique(G, k) denote the clique axioms as defined by Equations 1.2. Suppose that there exists a dual certificate $\mathcal{M}: \mathcal{P}(n,2r) \to \mathbb{R}$ for $\operatorname{Clique}(G,k)$ as defined in Definition 1.7. Then, the value of the r-round SOS-relaxation for maximum clique given by Figure 12 is at least k.

Proof. Let $\mathcal{M} : \mathcal{P}(n, 2r) \to \mathbb{R}$ be the dual certificate and $M \in \mathbb{R}^{\binom{[n]}{\leq r} \times \binom{[n]}{\leq r}}$ be the corresponding PSD matrix. Without loss of generality suppose that $M(\emptyset, \emptyset) = 1$. Let $M = UU^{\dagger}$, where $U = \mathbb{R}^{\binom{[n]}{\leq r} \times N}$ for some N. Finally, for $S \in {[n] \choose \leq r}$, let U_S be the S'th row of U. We claim that the collection $(U_S,$ $|S| \leq r$ gives a feasible solution for the SDP in Figure 12.

Observe that for any two subsets $S_1, S_2 \in {\binom{[n]}{< r}}$,

$$\langle \boldsymbol{U}_{S_1}, \boldsymbol{U}_{S_2} \rangle = M(S_1, S_2) = \mathcal{M}(X_{S_1 \cup S_2}).$$

Therefore, the vectors $(U_S : |S| \le r)$ satisfy the first two constraints of Figure 12 as \mathcal{M} is a dual certificate. Further, $\|\boldsymbol{U}_{\boldsymbol{\emptyset}}\|^2 = M(\boldsymbol{\emptyset}, \boldsymbol{\emptyset}) = 1$ and for any set S,

$$\|\boldsymbol{U}_S\|_2^2 = \langle \boldsymbol{U}_S, \boldsymbol{U}_S \rangle = \langle \boldsymbol{U}_S, \boldsymbol{U}_{\emptyset} \rangle \le \|\boldsymbol{U}_S\|_2,$$

so that $||U_S|| \leq 1$. Thus, $(U_S; |S| \leq r)$ give a feasible solution for the program in Figure 12. Finally, the value of the solution is

$$\sum_{i \in V} \|\boldsymbol{U}_{\{i\}}\|_2^2 = \sum_{i \in V} \mathcal{M}(X_{\{i\}}) = k.$$

This proves the lemma.

Our main theorems now follow.

Proof of Theorem 1.1. Let $G \leftarrow G(n, 1/2)$. Then, from the above lemma and the proof of Theorem 1.5 (where we showed the existence of a dual certificate for the clique axioms), the value of the r-round SOS-relaxation for max-clique on G is at least $n^{1/2r}/C^r(\log n)^{1/r}$ with high probability. The claim follows as the integral value is $(2 + o(1)) \log_2 n$ with high probability.

Proof of Corollary 1.2. The value of the relaxation in Figure 12 is clearly monotone with respect to adding edges. Therefore, from the above argument, for $G \leftarrow G(n, 1/2, t)$ the value of the *r*-round SOS-relaxation for max-clique on G is at least $n^{1/2r}/C^r(\log n)^{1/r}$ with high probability. The claim follows as the integral value is t with high probability.