Differential elimination by differential specialization of Sylvester style matrices

Sonia L. Rueda

Dpto. de Matemática Aplicada, E.T.S. Arquitectura. Universidad Politécnica de Madrid. Avda. Juan de Herrera 4, 28040-Madrid, Spain. sonialuisa.rueda@upm.es

Abstract

Differential resultant formulas are defined, for a system \mathcal{P} of n ordinary Laurent differential polynomials in n-1 differential variables. These are determinants of coefficient matrices of an extended system of polynomials obtained from \mathcal{P} through derivations and multiplications by Laurent monomials. To start, through derivations, a system $ps(\mathcal{P})$ of L polynomials in L-1 algebraic variables is obtained, which is non sparse in the order of derivation. This enables the use of existing formulas for the computation of algebraic resultants, of the multivariate sparse algebraic polynomials in $ps(\mathcal{P})$, to obtain polynomials in the differential elimination ideal generated by \mathcal{P} . The formulas obtained are multiples of the sparse differential resultant defined by Li, Yuan and Gao, and provide order and degree bounds in terms of mixed volumes in the generic case.

Keywords

differential elimination, Laurent differential polynomial, sparse resultant, differential specialization, sparse differential resultant

1 Introduction

The algebraic treatment via symbolic computation of differential equations has gained importance in the last years [34], [23], [27]. In addition, algebraic and differential elimination techniques have proven to be relevant tools in constructive, algorithmic algebra and symbolic computation [24], [9], [10], [26], [20]. This work establishes a bridge between the differential elimination problem for systems of ordinary differential polynomials and the use of sparse algebraic resultants. Let us consider a system of two ordinary differential polynomials in the differential indeterminates x and y,

$$f_1(x) = y' + yx + x' + xx' + yx^2 + y'(x')^2,$$

$$f_2(x) = y + y'x + yx' + y^2xx' + x^2 + (x')^2.$$
(1)

To eliminate the differential indeterminate x (and all its derivatives), they can be seen as two differential polynomials in the differential indeterminate x, whose coefficients are polynomials in the differential indeterminate y.

Differential elimination for differential polynomials can be achieved by characteristic set methods via symbolic computation algorithms [19], [4] (implemented in the Maple package diffalg, [3] and in the BLAD libraries [2] respectively), see also [15], [26]. These methods do not have an elementary complexity bound [16] and, the development of algorithms based on order and degree bounds, of the output elimination polynomials, would contribute to improve the complexity. Searching for order and degree bounds of the elimination polynomials is a problem closely related to the study of differential resultants.

For a system of sparse algebraic multivariate polynomials Canny and Emiris defined in [5] a Sylvester type matrix, whose determinant is a multiple of the sparse algebraic resultant, in the generic case (defined in [17]). Furthermore, the sparse multivariate algebraic resultant can be represented as the quotient of two determinants, as proved in [13]. These so called Macaulay style formulas provide degree bounds and furthermore methods to predict the support of the sparse algebraic resultant. While the studies and achievements on algebraic resultants are quite numerous, the differential case is at an initial state of development. A rigorous definition of the differential resultant $\partial \text{Res}(\mathfrak{P})$, of a set \mathfrak{P} of *n* sparse generic ordinary Laurent differential polynomials in n-1 differential variables, has been recently presented in [21] (and in [15], for the non sparse nonhomogeneous polynomial case), together with a single exponential algorithm in terms of bounds for degree and order of derivation. A matrix representation of the sparse differential resultant does not exist even for the simplest cases and, as noted in [21], having Macaulay style formulas in the differential case would improve the existing bounds for degree and order. The study of such formulas is the basis for efficient computation algorithms and, it promises to have a great contribution to the development and applicability of differential elimination techniques.

The first attempt to give Macaulay style formulas for a system \mathcal{P} of n ordinary differential polynomials, in n-1 differential variables, was made by G. Carrà-Ferro in [7]. Previous definitions of differential resultants were given for two ordinary differential operators, [1], [8] (refer to [7], [21] for an extended history of these developments). The differential resultant CFRes(\mathcal{P}) of \mathcal{P} defined by Carrà-Ferro is the algebraic resultant of Macaulay [22], of a set of derivatives of the differential polynomials in \mathcal{P} . For two non sparse differential polynomials of order 1 and degree 2, say (1), CFRes(\mathcal{P}) is the Macaulay algebraic resultant of the polynomial set ps = { f_1, f'_1, f_2, f'_2 }. This is the greatest common divisor of the determinant of all the minors of maximal order of a matrix \mathcal{M} , whose columns are indexed by all the monomials in x, x' and x'' of degree less than or equal to 5. The rows of \mathcal{M} are the coefficients of polynomials obtained by multiplying the polynomials in ps by certain monomials in x, x' and x'', see [6] and [28] for details. Observe that, even if f_1 and f_2 are nonsparse in x and x', the extended system ps is sparse. The polynomials in ps do not contain the monomial $(x'')^2$, thus the columns indexed by $(x'')^i, i = 2, \ldots, 5$ are all zero and CFRes(f_1, f_2) = 0.

Carrà-Ferro's construction is not taking into consideration the sparsity of differential polynomials and therefore it is zero in many cases, giving thus no further information. Contemporary of Carrà-Ferro's construction is the definition of the sparse algebraic resultant in [17] and [32]. Later on, methods to compute sparse algebraic resultants were developed in [5], [13] via Sylvester style matrices. Therefore, an alternative natural approach to treat example (1) (using Carrà-Ferro's philosophy) would be to consider the sparse algebraic resultant formula of ps given in [13]. A determinantal formula for 2 generic differential polynomials of arbitrary degree and order 1 has been recently presented in [33].

The system (1) can only be sparse in the degree but if we considered the elimination of two or more differential variables, the system can be also sparse in the order of derivation of such variables. This fact motivated the works in [29], [30] and [31] where the linear case is considered, to focus on the study of the sparsity with respect to the order of derivation, as defined in Section 3. An easy example is given by the next system of 3 polynomials

$$\mathcal{P} = \{f_1 = z + x + y + y', f_2 = z + tx' + y'', f_3 = z + x + y'\}$$

in 3 differential variables x, y and z w.r.t. the derivation $\partial/\partial t$. The differential resultant of Carra'Ferro is the determinant of the next coefficient matrix, whose columns are indexed by y^v , $x^v, \ldots, y', x', y, x, 1$,

1	0	1	1	0	0	0	0	0	0	$z^{\prime\prime\prime}$
0	0	1	0	1	1	0	0	0	0	$z^{\prime\prime}$
0	0	0	0	1	0	1	1	0	0	z'
0	0	0	0	0	0	1	0	1	1	z
1	0	0	t	0	2	0	0	0	0	z''
0	0	1	0	0	t	0	1	0	0	z'
0	0	0	0	1	0	0	t	0	0	z
1	0	0	1	0	0	0	0	0	0	$z^{\prime\prime\prime}$
0	0	1	0	0	1	0	0	0	0	$z^{\prime\prime}$
0	0	0	0	1	0	0	1	0	0	z'
0	0	0	0	0	0	1	0	0	1	z

Thus $CFRes(\mathcal{P}) = 0$ and the reason is the sparsity in the order of derivation of the variable x (the column indexed by x^v is zero).

In Section 2, differential resultant formulas are defined for a system \mathcal{P} of n ordinary Laurent differential polynomials in n-1 differential variables. These are determinants of coefficient matrices

of an extended system of polynomials obtained from \mathcal{P} through derivations and multiplications by Laurent monomials (Carrà-Ferro's construction is a particular case). To built such formulas, in Section 3 the results in [31] are extended to the nonlinear case, namely, an extended system $ps(\mathcal{P})$ of L polynomials in L-1 algebraic variables is obtained through the appropriate number of derivations of the elements of \mathcal{P} , which is non sparse in the order of derivation. As explained in Section 3, this is only possible for systems \mathcal{P} that verify the "super essential" condition, but it is there proved that every system contains such a subsystem. For $n \geq 3$, this is a necessary step to be able to use the existing formulas for sparse algebraic polynomials in [5] applied to the system $ps(\mathcal{P})$. An algebraic generic sparse system $ags(\mathcal{P})$ of L polynomials in L-1 algebraic variables associated to \mathcal{P} is defined, as explained in Section 4, from which a Sylvester style matrix $S(\mathcal{P})$ can be constructed using the results in [5]. The specialization of $S(\mathcal{P})$, to the differential coefficients of $p_{S}(\mathcal{P})$, gives a determinantal formula $\partial FRes(\mathcal{P})$, as explained in Section 5. In [5], $det(S(\mathcal{P}))$ is guaranteed to be nonzero (under some conditions) so, if $\partial FRes(\mathcal{P}) = 0$ then we know that it is not because of sparsity reasons, but due to the specialization final step. In Section 6 the generic case is treated. It is shown how these formulas provide order and degree bounds for the sparse differential resultant $\partial \operatorname{Res}(\mathcal{P})$ of \mathcal{P} defined by Li, Yuan and Gao in [21]. To achieve this goal, conditions for $\partial \operatorname{Res}(\mathcal{P})$ to be a factor of the given differential resultant formulas are explored, providing degree bounds of $\partial \operatorname{Res}(\mathcal{P})$ in terms of mixed volumes, under the appropriate conditions.

2 Differential resultant formulas

Let \mathcal{D} be an ordinary differential domain with derivation ∂ . Let $U = \{u_1, \ldots, u_{n-1}\}$ be a set of differential indeterminates over \mathcal{D} . By \mathbb{N} we mean the natural numbers including 0. For $k \in \mathbb{N}$, we denote by $u_{j,k}$ the kth derivative of u_j and for $u_{j,0}$ we simply write u_j . We denote by $\{U\}$ the set of derivatives of the elements of U, $\{U\} = \{\partial^k u \mid u \in U, k \in \mathbb{N}\}$, and by $\mathcal{D}\{U\}$ the ring of differential polynomials in the differential indeterminates U, which is a differential ring with derivation ∂ . For definitions in differential algebra we refer to [26] and [20].

As introduced in [21], the ring of Laurent differential polynomials generated by U is defined to be

$$\mathcal{D}\{U^{\pm}\} := \mathcal{D}[u_{j,k}, u_{j,k}^{-1} \mid j = 1, \dots, n-1, k \in \mathbb{N}]\},\$$

which is a differential ring under the derivation ∂ , (emphasize that $\mathcal{D}\{U^{\pm}\}$ is just notation). Given a subset $\mathcal{U} \subset \{U\}$, we denote by $\mathcal{D}[\mathcal{U}]$ the ring of polynomials in the indeterminates \mathcal{U} and by $\mathcal{D}[\mathcal{U}^{\pm}]$ the ring of Laurent polynomials in the variables \mathcal{U} , that is

$$\mathcal{D}[\mathcal{U}^{\pm}] := \mathcal{D}[u, u^{-1} \mid u \in \mathcal{U}].$$

Given $f \in \mathcal{D}\{U^{\pm}\}$, $f = \sum_{\iota=1}^{m} \theta_{\iota} \omega_{\iota}$, where $\theta_{\iota} \in \mathcal{D}$ and ω_{ι} is a Laurent differential monomial in $\mathcal{D}\{U^{\pm}\}$. Let us denote the differential support in u_j of f by

$$\mathfrak{S}_{j}(f) = \{k \in \mathbb{N} \mid \partial \omega_{\iota} / \partial u_{j,k} \neq 0 \text{ for some } \iota \in \{1, \dots, m\}\},\$$

to define $\operatorname{ord}(f, u_j) := \max \mathfrak{S}_j(f)$ and $\operatorname{lord}(f, u_j) := \min \mathfrak{S}_j(f)$ if $\mathfrak{S}_j(f) \neq \emptyset$, otherwise $\operatorname{ord}(f, u_j) = \operatorname{lord}(f, u_j) = -\infty$. Thus, the order of f is the maximum of $\{\operatorname{ord}(f, u) \mid u \in U\}$.

Let $\mathcal{P} := \{f_1, \ldots, f_n\}$ be a system of differential polynomials in $\mathcal{D}\{U^{\pm}\}$. We assume that:

 $(\mathcal{P}1)$ The order of f_i is $o_i \geq 0, i = 1, \ldots, n$. So that no f_i belongs to \mathcal{D} .

 $(\mathcal{P}2)$ \mathcal{P} contains *n* distinct polynomials.

 $(\mathcal{P}3)$ For every $j \in \{1, \ldots, n-1\}$ there exists $i \in \{1, \ldots, n\}$ such that $\mathfrak{S}_j(f_i) \neq \emptyset$.

Let $[\mathcal{P}]$ denote the differential ideal generated by \mathcal{P} in $\mathcal{D}\{U^{\pm}\}$. Our goal is to obtain elements of the differential elimination ideal $[\mathcal{P}] \cap \mathcal{D}$, using differential resultant formulas.

Let us denote by $\partial \mathcal{P} := \{\partial^k f_i \mid i = 1, \dots, n, k \in \mathbb{N}\}$ and $f_i^{[L_i]} := \{\partial^k f_i \mid k \in [0, L_i] \cap \mathbb{N}\}$, for $L_i \in \mathbb{N}$. For this purpose, we consider a polynomial subset ps of $\partial \mathcal{P}$, a set of differential indeterminates $\mathcal{U} \subset \{U\}$ and sets of Laurent differential monomials Ω_f , $f \in ps$, Ω , in $\mathcal{D}[\mathcal{U}^{\pm}]$, verifying:

$$(ps1) ps = \bigcup_{i=1}^{n} f_i^{\lfloor L_i \rfloor}, \ L_i \in \mathbb{N},$$

(ps2) ps $\subset \mathcal{D}[\mathcal{U}^{\pm}]$ and $|\mathcal{U}| = |ps| - 1$,

(ps3) $\sum_{f \in ps} |\Omega_f| = |\Omega|$ and $\bigcup_{f \in ps} \Omega_f f \in \bigoplus_{\omega \in \Omega} \mathcal{D}\omega$, ($|\Omega|$ denotes de number of elements of Ω).

Under assumptions (ps1), (ps2) and (ps3), we consider a total set of polynomials PS := $\cup_{f \in ps} \Omega_f f$ whose elements are

$$p = \sum_{\omega \in \Omega} \theta_{p,\omega} \omega$$
, with $\theta_{p,\omega} \in \mathcal{D}$.

The coefficient matrix of the elements in PS as polynomials in the monomials Ω , $\mathcal{M}(PS, \Omega) = (\theta_{p,\omega})$, $p \in PS$, $\omega \in \Omega$, is an $|\Omega| \times |\Omega|$ matrix. We call

$$\det(\mathcal{M}(\mathrm{PS},\Omega)) \tag{2}$$

a differential resultant formula for \mathcal{P} .

Example 2.1. A differential resultant formula was defined by Carrà-Ferro in [7] for a system \mathcal{P} of nonhomogeneous differential polynomials in $\mathcal{D}\{U\}$. In [7], $L_i = N - o_i$, i = 1, ..., n, $N := \sum_{i=1}^n o_i$ and $\mathcal{U} = \{u_{j,k} \mid k \in [0, N] \cap \mathbb{N}, j = 1, ..., n-1\}$. The sets of monomials Ω_f , $f \in ps$ and Ω are taken so that $\mathcal{M}(PS, \Omega)$ is the specialization of the numerator matrix of the Macaulay algebraic resultant [22] of generic algebraic polynomials P_f , $f \in ps$ of degree $\deg(P_f) = \deg(f)$ in the variables \mathcal{U} . See [7] and [28] for a detailed construction and examples.

If (ps2) holds, the set

$$\nu(\mathrm{ps}) := \{ u_{j,k} \in \mathcal{U} \mid k \in \mathfrak{S}_j(f) \text{ for some } f \in \mathrm{ps}, j \in \{1, \dots, n-1\} \} \subseteq \mathcal{U}\}$$

verifies $|\nu(ps)| \le |ps| - 1$. Observe that, if $|\nu(ps)| > |ps| - 1$ we cannot guarantee the elimination of the variables in $\nu(ps)$.

3 A system $ps(\mathcal{P})$ of L polynomials in L-1 algebraic variables

In this section, we construct $ps(\mathcal{P}) \subset \partial \mathcal{P}$ and $\mathcal{V}(\mathcal{P}) \subset \{U\}$ verifying (ps1), (ps2) and give conditions on \mathcal{P} so that $\mathcal{V}(\mathcal{P}) = \nu(ps(\mathcal{P}))$. In particular, it is precisely stated what it means for the system \mathcal{P} to be sparse in the order and under what conditions can this phenomenon be avoided.

Let us denote $o_{i,j} := \operatorname{ord}(f_i, u_j)$, which equals $-\infty$ if $\mathfrak{S}_j(f_i) = \emptyset$ and belongs to N otherwise. Let us define the order matrix of \mathcal{P} by $\mathcal{O}(\mathcal{P}) = (o_{i,j})$. Given $\mathcal{P}_i := \mathcal{P} \setminus \{f_i\}, i = 1, \ldots, n$, the diagonals of the matrix $\mathcal{O}(\mathcal{P}_i)$ are indexed by the set Γ_i of all possible bijections between $\{1, \ldots, n\} \setminus \{i\}$ and $\{1, \ldots, n-1\}$. The Jacobi number $J_i(\mathcal{P})$ of the matrix $\mathcal{O}(\mathcal{P}_i)$ (see [21], Section 5.2) equals

$$J_i(\mathcal{P}) := \operatorname{Jac}(\mathcal{O}(\mathcal{P}_i)) := \max\left\{\sum_{j \in \{1,\dots,n\} \setminus \{i\}} o_{j,\mu(j)} \mid \mu \in \Gamma_i\right\}.$$

Throughout the paper, if there is no need to specify, we will simply write J_i . Observe that J_i is either $-\infty$ or it belongs to \mathbb{N} . There exists $\mu_i \in \Gamma_i$ such that $J_i = \sum_{j \in \{1,...,n\} \setminus \{i\}} o_{j,\mu_i(j)}$ but μ_i may not be unique.

The situation where $J_i \ge 0$, i = 1, ..., n is of special interest. Let $x_{i,j}$, i = 1, ..., n, j = 1, ..., n - 1 be algebraic indeterminates over \mathbb{Q} , the field of rational numbers. Let $X(\mathcal{P}) = (X_{i,j})$ be the $n \times (n-1)$ matrix, such that

$$X_{i,j} := \begin{cases} x_{i,j}, & \mathfrak{S}_j(f_i) \neq \emptyset, \\ 0, & \mathfrak{S}_j(f_i) = \emptyset. \end{cases}$$
(3)

Let $X(\mathcal{P}_i)$, i = 1, ..., n, be the submatrix of $X(\mathcal{P})$ obtained by removing its *i*th row. It follows easily that

Lemma 3.1. $J_i \ge 0 \Leftrightarrow \det(X(\mathcal{P}_i)) \ne 0, i = 1, \dots, n.$

Proof. If det $(X(\mathcal{P}_i)) \neq 0$ then the matrix $X(\mathcal{P}_i)$ has a nonzero diagonal. Thus, there exists $\mu \in \Gamma_i$ such that

$$J_i \ge \sum_{j \in \{1,\dots,n\} \setminus \{i\}} o_{j,\mu(j)} \ge 0.$$

Conversely if $J_i \ge 0$, there exists $\mu \in \Gamma_i$ such that $J_i = \sum_{j \in \{1,...,n\} \setminus \{i\}} o_{j,\mu(j)} \ge 0$. Thus $\prod_{j \in \{1,...,n\} \setminus \{i\}} x_{j,\mu(j)} \ge 0$ and $\det(X(\mathcal{P}_i)) \ne 0$.

The notion of super essential system of differential polynomials was introduced in [31], for systems of linear differential polynomials and it is extended here to the nonlinear case.

Definition 3.2. The system \mathcal{P} is called super essential if $det(X(\mathcal{P}_i)) \neq 0$, i = 1, ..., n. Equivalently, by Lemma 3.1, \mathcal{P} is super essential if $J_i \geq 0$, i = 1, ..., n.

For $j = 1, \ldots, n-1$ let us define integers in \mathbb{N}

$$\gamma_j(\mathcal{P}) := \min\{ \operatorname{lord}(f_i, u_j) \mid \mathfrak{S}_j(f_i) \neq \emptyset, \ i = 1, \dots, n \},$$
$$\gamma(\mathcal{P}) := \sum_{j=1}^{n-1} \gamma_j(\mathcal{P}).$$

We write just γ_j and γ when there is no room for confusion. If $J_i \ge 0$, i = 1, ..., n then $J_i - \gamma \ge 0$ and the sets of lattice points $[0, J_i - \gamma] \cap \mathbb{N}$ are non empty. For i = 1, ..., n, we define the set of differential polynomials

$$ps(\mathcal{P}) := \bigcup_{i=1}^{n} f_i^{[J_i - \gamma]},\tag{4}$$

containing $L := \sum_{i=1}^{n} (J_i - \gamma + 1)$ differential polynomials, whose variables belong to the set $\mathcal{V}(\mathcal{P})$ of differential indeterminates

$$\mathcal{V}(\mathcal{P}) := \{ u_{j,k} \mid k \in [\gamma_j, M_j] \cap \mathbb{N}, \, j = 1, \dots, n-1 \},\$$

with $M_j := m_j - \gamma$ and $m_j := \max\{o_{i,j} + J_i - \gamma \mid i = 1, \dots, n\}$. By [21], Lemma 5.6, if $J_i \ge 0$, $i = 1, \dots, n$ then $\sum_{i=1}^n J_i = \sum_{j=1}^{n-1} m_j$. Thus the number of elements of $\mathcal{V}(\mathcal{P})$ equals

$$\sum_{j=1}^{n-1} (M_j - \gamma_j + 1) = \sum_{j=1}^{n-1} (m_j - \gamma_j - \gamma + 1) = \sum_{i=1}^n J_i - n\gamma + n - 1 = L - 1.$$

Observe that $\nu(ps(\mathcal{P})) \subseteq \mathcal{V}(\mathcal{P})$ and given $j \in \{1, \ldots, n-1\}$ we have

$$\cup_{f \in \mathrm{ps}(\mathcal{P})} \mathfrak{S}_j(f) \subseteq [\gamma_j, M_j] \cap \mathbb{N},\tag{5}$$

but we cannot guarantee that the equality holds.

Definition 3.3. If there exists j such that (5) is not an equality, we will say that the system \mathcal{P} is sparse in the order.

It can be proved as in [31], Section 4, that every system \mathcal{P} contains a super essential subsystem \mathcal{P}^* and if rank $(X(\mathcal{P})) = n - 1$ then \mathcal{P}^* is unique. Namely, the system \mathcal{P}^* can be obtained as follows:

- 1. Consider the system $\mathbb{P} = \{p_i = c_i + \sum_{j=1}^{n-1} X_{i,j} u_j \mid l = 1, \dots, m, \}$ of algebraic polynomials in $\mathbb{K}[c_1, \dots, c_m][U], \mathbb{K} := \mathbb{Q}(X_{i,j} \mid X_{i,j} \neq 0).$
- 2. Compute a reduced Gröbner basis $\mathcal{B} = \{e_0, e_1, \ldots, e_{m-1}\}$ of the algebraic ideal (\mathbb{P}) generated by \mathbb{P} in $\mathbb{K}[c_1, \ldots, c_m][U]$, with respect to lex monomial order with $u_1 > \cdots > u_p > c_1 > \cdots > c_m$. We assume that $e_0 < e_1 < \cdots < e_{m-1}$. By [9], p. 95, Exercise 10, this can be computed through an echelon form of the coefficient matrix of the system \mathbb{P} .
- 3. Observe that at least $e_0 \in \mathcal{B}_0 := \mathcal{B} \cap \mathbb{K}[c_1, \dots, c_m], e_0 = \sum_{l=1}^m \chi_l c_l, \ \chi_l \in \mathbb{K}$. Let $\Delta(e_0) := \{l \in \{1, \dots, m\} \mid \chi_l \neq 0\}.$

4.
$$\mathcal{P}^* := \{ f_l \mid l \in \Delta(e_0) \}.$$

Example 3.4. Let us consider the systems $\mathcal{P} = \{f_1, f_2, f_3, f_4\}$ and $\mathcal{P}' = \{f_1, f_2, f_3, f_5\}$ with

$$f_1 = 2 + u_1 u_{1,1} + u_{1,2}, f_2 = u_1 u_{1,2}, f_3 = u_2 u_{3,1}, f_4 = u_{1,1} u_2, f_5 = u_{1,2}$$

$$X(\mathcal{P}) = \begin{pmatrix} x_{1,1} & 0 & 0 \\ x_{2,1} & 0 & 0 \\ 0 & x_{3,2} & x_{3,3} \\ x_{4,1} & x_{4,2} & 0 \end{pmatrix} \text{ and } X(\mathcal{P}') = \begin{pmatrix} x_{1,1} & 0 & 0 \\ x_{2,1} & 0 & 0 \\ 0 & x_{3,2} & x_{3,3} \\ x_{4,1} & 0 & 0 \end{pmatrix}.$$

 \mathcal{P} is not super essential but since $\operatorname{rank}(X(P)) = 3$, it has a unique super essential subsystem, which is $\{f_1, f_2\}$. \mathcal{P}' is not super essential and $\operatorname{rank}(X(P')) < 3$, super essential subsystems are $\{f_1, f_2\}$, $\{f_1, f_5\}$ and $\{f_2, f_5\}$.

We prove next that if \mathcal{P} is super essential then \mathcal{P} is not sparse in the order (Theorem 3.7). For this purpose we need two preparatory lemmas.

Given $j \in \{1, \ldots, n-1\}$, the set $\mathcal{I}(j) := \{i \in \{1, \ldots, n\} \mid \mathfrak{S}_j(f_i) \neq \emptyset\}$ is not empty, because of assumption ($\mathcal{P}3$). If \mathcal{P} is super essential, the next lemma shows, in particular, that $|\mathcal{I}(j)| \geq 2$. Given $I, I' \in \{1, \ldots, n\}$, let $\mathcal{P}_{I,I'} := \mathcal{P} \setminus \{f_I, f_{I'}\}$. Let us denote by $X(\mathcal{P}_{I,I'})^j$ and $\mathcal{O}(\mathcal{P}_{I,I'})^j$ the submatrices of $X(\mathcal{P}_{I,I'})$ and $\mathcal{O}(\mathcal{P}_{I,I'})$ respectively, obtained by removing their *j*th column. Observe that det $(X(\mathcal{P}_{I,I'})^j) = 0$ if and only if $\operatorname{Jac}(\mathcal{O}(\mathcal{P}_{I,I'})^j) = -\infty$.

Lemma 3.5. Let \mathcal{P} be super essential and $j \in \{1, \ldots, n-1\}$.

- 1. Given $I \in \mathcal{I}(j)$, there exists $I' \in \mathcal{I}(j) \setminus \{I\}$ such that $o_{I,j} \gamma_j \leq J_{I'} \gamma$.
- 2. Given distinct $I, \overline{I} \in \mathcal{I}(j)$ such that $\operatorname{Jac}(\mathcal{O}(\mathcal{P}_{I,\overline{I}})^j) = -\infty$, there exists $I' \in \mathcal{I}(j) \setminus \{\overline{I}\}$ such that $o_{\overline{I},j} \gamma_j \leq J_{I'} \gamma$ and, if $I \neq I'$ then $o_{I',j} \gamma_j \leq J_I \gamma$.

Proof. We denote $X(\mathcal{P}_{I,I'})^j$ and $\mathcal{O}(\mathcal{P}_{I,I'})^j$ simply by $\mathcal{X}_{I,I'}$ and $\mathcal{O}_{I,I'}$ in this proof. Let $X(P)^j$ be the submatrix of $X(\mathcal{P})$ obtained by removing its *j*th column. By r_I we denote the row corresponding to f_I in $X(\mathcal{P})^j$ and by $\mathcal{X}_{\overline{I},I,I'}$ the matrix $X(\mathcal{P}_{\overline{I},I,I'})^j$ with $\mathcal{P}_{\overline{I},I,I'} := \mathcal{P} \setminus \{f_{\overline{I}}, f_I, f_{I'}\}.$

1. By definition of super essential system, $X(\mathcal{P}_I)$ contains a nonzero diagonal. That is, there exists $I' \in \mathcal{I}(j) \setminus \{I\}$ and an $(n-2) \times (n-2)$ non singular submatrix $\mathcal{X}_{I,I'}$ of $X(\mathcal{P}_I)$. That is $Jac(\mathcal{O}_{I,I'}) \neq -\infty$ and

$$o_{I,j} - \gamma_j \leq \operatorname{Jac}(\mathcal{O}_{I,I'}) + o_{I,j} - \gamma \leq J_{I'} - \gamma_j$$

2. By 1, there exists $I' \in \mathcal{I}(j) \setminus \{\overline{I}\}$ such that $\operatorname{Jac}(\mathcal{O}_{\overline{I},I'}) \neq -\infty$ and $o_{\overline{I},j} - \gamma_j \leq J_{I'} - \gamma$. If $I \neq I'$, let us assume that $\operatorname{Jac}(\mathcal{O}_{I,I'}) = -\infty$ to get a contradiction. Thus we have $\det(\mathcal{X}_{\overline{I},I}) = 0$ and $\det(\mathcal{X}_{I,I'}) = 0$.

The $(n-2) \times (n-2)$ matrices $\mathcal{X}_{\overline{I},I}$, $\mathcal{X}_{\overline{I},I'}$ and $\mathcal{X}_{I,I'}$ have n-3 rows in common, namely $\mathcal{X}_{\overline{I},I,I'}$. Since $\det(\mathcal{X}_{\overline{I},I'}) \neq 0$, the rows of $\mathcal{X}_{\overline{I},I,I'}$ are linearly independent. This proves that

$$\operatorname{rank}(\mathcal{X}_{\overline{I},I}) = \operatorname{rank}(\mathcal{X}_{I,I'}) = \operatorname{rank}(\mathcal{X}_{\overline{I},I,I'}) = n - 3.$$

Thus row I' of $\mathcal{X}_{\overline{I},I}$ and row \overline{I} of $\mathcal{X}_{I,I'}$ are a linear combination of the rows of $\mathcal{X}_{I,I',\overline{I}}$. Therefore both rows I' and \overline{I} of $X(\mathcal{P}_I)$ can be reduced to the form $(0,\ldots,0,\star_j,0,\ldots,0)$. Thus $\det(X(\mathcal{P}_I)) = 0$ contradicting that \mathcal{P} is super essential. This proves that $\operatorname{Jac}(\mathcal{O}_{I,I'}) \neq -\infty$ and

$$o_{I',j} - \gamma_j \leq \operatorname{Jac}(\mathcal{O}_{I,I'}) + o_{I',j} - \gamma \leq J_I - \gamma$$

Lemma 3.6. Let $f \in \mathcal{D}\{U^{\pm}\}$. If $k \in \mathfrak{S}_j(f)$ but $k+1 \notin \mathfrak{S}_j(f)$ then $k+1 \in \mathfrak{S}_j(\partial f)$.

Proof. Observe that $f = A_{-l}u_{j,k}^{-l} + \dots + A_{-1}u_{j,k}^{-1} + A_0 + A_1u_{j,k} + \dots + A_mu_{j,k}^m$ with $A_t \in \mathcal{D}\{U^{\pm}\}, t = -l, \dots, -1, 0, 1, \dots, m$, such that $k, k+1 \notin \mathfrak{S}_j(A_t), A_{-l} \neq 0$ or $A_m \neq 0$ and $l \geq 1$ or $m \geq 1$. The claim follows since

$$\partial f = \partial A_{-l} u_{j,k}^{-l} + \dots + \partial A_0 + \dots + \partial A_m u_{j,k}^m + \left(\sum_{h \neq 0, h = -l}^m h A_h u_{j,k}^{h-1}\right) u_{j,k+1}.$$

Theorem 3.7. If \mathcal{P} is super essential then

$$\cup_{f \in \mathrm{ps}(\mathcal{P})} \mathfrak{S}_j(f) = [\gamma_j, M_j] \cap \mathbb{N}, \ j = 1, \dots, n-1.$$

That is, $ps(\mathcal{P})$ is a system of L polynomials in L-1 algebraic indeterminates.

Proof. Given $j \in \{1, ..., n-1\}$, there exists $\overline{I} \in \{1, ..., n\}$ such that $m_j = o_{\overline{I}, j} + J_{\overline{I}}$. Recall $M_j = m_j - \gamma$. We can write

$$[\gamma_j, M_j] = [\gamma_j, o_{\overline{I},j} - 1] \cup [o_{\overline{I},j}, M_j].$$

- 1. For every $k \in [o_{\overline{I},j}, M_j] \cap \mathbb{N}, \ k o_{\overline{I},j} \leq M_j o_{\overline{I},j} = J_{\overline{I}} \gamma$. By Lemma 3.6, $k \in \mathfrak{S}_j(\partial^{k-o_{\overline{I},j}} f_{\overline{I}})$.
- 2. If $o_{\overline{I},j} = \gamma_j$ then the first interval is empty. If $o_{\overline{I},j} \neq \gamma_j$, there exists $\underline{I} \in \mathcal{I}(j)$ such that $\operatorname{ldeg}(f_{\underline{I}}, u_j) = \gamma_j$ and a bijection $\mu_{\underline{I}} : \{1, \ldots, n\} \setminus \{\underline{I}\} \longrightarrow \{1, \ldots, n-1\}$, with $I := \mu_{\underline{I}}^{-1}(j)$ such that

$$o_{I,j} - \gamma_j \leq J_{\underline{I}} - \gamma = \sum_{l \in \{1,\dots,n\} \setminus \{\underline{I}\}} o_{l,\mu_{\underline{I}}(l)} - \gamma.$$

If there exists $k \in ([\gamma_j, o_{I,j}] \cap \mathbb{N}) \setminus \mathfrak{S}_j(f_I)$ then let us consider

$$k' := \max([\gamma_j, k-1] \cap \mathbb{N}) \cap \mathfrak{S}_j(f_{\underline{I}}).$$

Since $k - k' \leq o_{I,j} - k' \leq o_{I,j} - \gamma_j \leq J_{\underline{I}} - \gamma$, by Lemma 3.6, $k \in \mathfrak{S}_j(\partial^{k-k'} f_{\underline{I}})$. Thus, for $ps(f_{\underline{I}})$ as in (4), it holds

$$[\gamma_j, o_{I,j}] \cap \mathbb{N} \subseteq \cup_{f \in \mathrm{ps}(f_{\underline{I}})} \mathfrak{S}_j(f).$$
(6)

2.1. If $o_{I,j} \ge o_{\overline{I},j} - 1$ then, by (6) $[\gamma_j, o_{\overline{I},j} - 1] \cap \mathbb{N} \subseteq \bigcup_{f \in \mathrm{ps}(f_{\underline{I}})} \mathfrak{S}_j(f)$.

2.2. If $o_{I,j} < o_{\overline{I},j} - 1$ then $[\gamma_j, o_{\overline{I},j} - 1] = [\gamma_j, o_{I,j}] \cup [o_{I,j} + 1, o_{\overline{I},j} - 1]$. Consequently, if $o_{\overline{I},j} \le o_{I,j} + J_I - \gamma$ then $[o_{I,j} + 1, o_{\overline{I},j} - 1] \subset [o_{I,j} + 1, o_{I,j} + J_I - \gamma]$. If $o_{I,j} + J_I - \gamma < o_{\overline{I},j}$ then $\operatorname{Jac}(\mathcal{O}(\mathcal{P}_{I,\overline{I}})^j) = -\infty$ since otherwise

$$o_{\overline{I},j} \leq \gamma_j + o_{\overline{I},j} + \operatorname{Jac}(\mathcal{O}(\mathcal{P}_{I,\overline{I}})^j) - \gamma \leq o_{I,j} + J_I - \gamma$$

By Lemma 3.5 (2), there exists $I' \in \mathcal{I}(j) \setminus \{\overline{I}\}$ such that $o_{\overline{I},j} - \gamma_j \leq J_{I'} - \gamma$ and, if $I \neq I'$ then $o_{I',j} - \gamma_j \leq J_I - \gamma$. Note this implies

$$o_{\overline{I},j} \leq o_{I',j} + J_{I'} - \gamma$$
 and $o_{I',j} \leq o_{I,j} + J_I - \gamma$

If $o_{I',j} \le o_{I,j}$ then $[o_{I,j} + 1, o_{\overline{I},j} - 1] \subset [o_{I',j}, o_{I',j} + J_{I'} - \gamma]$, otherwise $o_{I,j} < o_{I',j}$ and

$$\begin{split} [o_{I,j}+1, o_{\overline{I},j}-1] &= [o_{I,j}+1, o_{I',j}-1] \cup [o_{I',j}, o_{\overline{I},j}-1] \\ &\subset [o_{I,j}+1, o_{I,j}+J_I-\gamma] \cup [o_{I',j}, o_{I',j}+J_{I'}-\gamma]. \end{split}$$

Thus given $k \in [o_{I,j}+1, o_{\overline{I},j}-1] \cap \mathbb{N}$, if $k \in [o_{I,j}+1, o_{I,j}+J_I-\gamma]$ then $k - o_{I,j} \leq J_I - \gamma$ and, by Lemma 3.6, $k \in \mathfrak{S}_j(\partial^{k-o_{I,j}}f_I)$. Analogously, if $k \in [o_{I',j}, o_{I',j}+J_{I'}-\gamma]$ then $k \in \mathfrak{S}_j(\partial^{k-o_{I',j}}f_{I'})$.

Example 3.8. Let \mathcal{P} be a system with $\gamma = 0$,

$$\mathcal{O}(\mathcal{P}) = \begin{pmatrix} 2 & 0 \\ -\infty & 1 \\ 2 & 0 \end{pmatrix}, \text{ thus } X(\mathcal{P}) = \begin{pmatrix} x_{1,1} & x_{1,2} \\ 0 & x_{2,3} \\ x_{3,1} & x_{3,2} \end{pmatrix}.$$

Then $J_1 = 3$, $J_2 = 2$ and $J_3 = 3$ and \mathcal{P} is super essential. By Theorem 3.7, $ps(\mathcal{P})$ is a system with 11 polynomials in 10 algebraic variables $\mathcal{V} = \{u_1, u_{1,1}, \ldots, u_{1,5}, u_2, u_{2,1}, \ldots, u_{2,3}\}.$

If we consider, for instance, the system ps, with $L_1 = 2 < J_1$, $L_2 = J_2$ and $L_3 = J_3$. We have 10 polynomials in 10 algebraic variables $\mathcal{V}(\mathcal{P})$, in this case we cannot guarantee the elimination of the algebraic variables $\mathcal{V}(\mathcal{P})$.

4 Sparse algebraic resultant associated to \mathcal{P}

The result in Theorem 3.7, allows the construction of a Sylvester matrix associated to the system $ps(\mathcal{P})$, choosing orderings on the sets $\mathcal{V}(\mathcal{P})$ and $ps(\mathcal{P})$, as it is next explained.

Through a bijection $\beta : \mathcal{V}(\mathcal{P}) \to \{1, \dots, L-1\}$ we establish an ordering of the set of variables $\mathcal{V}(\mathcal{P})$. Let $\mathcal{Y} = \{y_1, \dots, y_{L-1}\}$ be a set of L-1 algebraic indeterminates over \mathbb{Q} . A natural bijection $v : \mathcal{Y} \to \mathcal{V}(\mathcal{P})$ is defined by $v(y_l) = \beta^{-1}(l)$. Given the Laurent polynomial ring $\mathcal{D}[\mathcal{Y}^{\pm}], v$ extends to a ring isomorphism

$$v: \mathcal{D}[\mathcal{Y}^{\pm}] \to \mathcal{D}[\mathcal{V}(\mathcal{P})^{\pm}].$$

Monomials in $\mathcal{D}[\mathcal{Y}^{\pm}]$ are $y^{\alpha} = y_1^{\alpha_1} \cdots y_{L-1}^{\alpha_{L-1}}$, with $\alpha = (\alpha_1, \ldots, \alpha_{L-1}) \in \mathbb{Z}^{L-1}$ and $v(y^{\alpha}) = v(y_1)^{\alpha_1} \cdots v(y_{L-1})^{\alpha_{L-1}}$. Now given $f = \sum_{\alpha \in \mathbb{Z}^{L-1}} a_{\alpha} v(y^{\alpha})$ in $\mathcal{D}[\mathcal{V}(\mathcal{P})^{\pm}]$, we define the algebraic support $\mathcal{A}(f)$ of f as

$$\mathcal{A}(f) := \left\{ \alpha \in \mathbb{Z}^{L-1} \mid a_{\alpha} \neq 0 \right\}$$

A bijection $\lambda : ps(\mathcal{P}) \to \{1, \dots, L\}$ defines an ordering in the set $ps(\mathcal{P})$. Let us call its inverse ρ . We define the algebraic generic system associated to \mathcal{P} as

$$\operatorname{ags}(\mathcal{P}) := \left\{ \sum_{\alpha \in \mathcal{A}(f)} c_{\alpha}^{\lambda(f)} y^{\alpha} \mid f \in \operatorname{ps}(\mathcal{P}) \right\},\$$

where $c_{\alpha}^{\lambda(f)}$ are algebraic indeterminates over \mathbb{Q} . Thus we have

$$\operatorname{ags}(\mathcal{P}) = \left\{ P_l := \sum_{\alpha \in \mathcal{A}(\rho(l))} c_{\alpha}^l y^{\alpha} \mid l = 1, \dots, L \right\}.$$

Given $l \in \{1, \ldots, L\}$, let us consider sets of algebraic indeterminates over \mathbb{Q}

$$C_l := \{ c_{\alpha}^l \mid \alpha \in \mathcal{A}(\rho(l)) \} \text{ and } C := \cup_{l=1}^L C_l.$$

The system of algebraic generic polynomials $\operatorname{ags}(\mathcal{P})$ is included in $\mathbb{E}[\mathcal{Y}^{\pm}]$, for $\mathbb{E} := \mathbb{Q}(\mathcal{C})$. Given a subsystem $\mathcal{S} \subseteq \operatorname{ags}(\mathcal{P})$, its elements are polynomials $P = \sum_t a_t M_{P,t}$, with $M_{P,t}$ monomials in $\mathbb{E}[\mathcal{Y}^{\pm}]$. We denote by

$$\mathcal{Y}(\mathcal{S}) := \{ y \in \mathcal{Y} \mid \partial M_{P,t} / \partial y \neq 0 \text{ for some monomial of } P \in \mathcal{S} \}.$$
(7)

If \mathcal{P} is super essential, Theorem 3.7 implies that $\mathcal{Y}(ags(\mathcal{P})) = \mathcal{Y}$, so $ags(\mathcal{P})$ is a set of L polynomials in L-1 indeterminates \mathcal{Y} .

A Sylvester matrix $Syl(ags(\mathcal{P}))$ for $ags(\mathcal{P})$ can be constructed as in [5] (see also [11], [32] and [13]), where finite sets of monomials $\Lambda_1, \ldots, \Lambda_L, \Lambda$ in $\mathbb{E}[\mathcal{Y}^{\pm}]$ are determined. Let $\langle \Lambda \rangle_{\mathbb{E}}$ denote the \mathbb{E} -vector space generated by Λ . The matrix in the monomial bases of the linear map

$$\langle \Lambda_1 \rangle_{\mathbb{E}} \oplus \cdots \oplus \langle \Lambda_L \rangle_{\mathbb{E}} \to \langle \Lambda \rangle_{\mathbb{E}} : (g_1, \ldots, g_L) \mapsto \sum g_l P_l,$$

is $Syl(ags(\mathcal{P}))$. In [5] and [32], it is assumed without loss of generality that

(A1) the affine lattice generated by the Minkowski sum $\sum_{f \in ps(\mathcal{P})} \mathcal{A}(f)$ has dimension L-1.

This technical hypothesis is removed in [11], thus a Sylvester matrix $Syl(ags(\mathcal{P}))$ for $ags(\mathcal{P})$ can be constructed without any additional assumption.

Let $(ags(\mathcal{P}))$ be the algebraic ideal generated by $ags(\mathcal{P})$ in $\mathbb{Q}[\mathcal{C}][\mathcal{Y}^{\pm}]$. No reference was found for the next result, which is proved for the sake of completeness, although it seems natural that it should exist in the sparse algebraic resultant literature.

Proposition 4.1. det $(Syl(ags(\mathcal{P}))) \in (ags(\mathcal{P})) \cap \mathbb{Q}[\mathcal{C}].$

Proof. Let us denote $D = \det(Syl(\operatorname{ags}(\mathcal{P})))$ and $S = Syl(\operatorname{ags}(\mathcal{P}))$. Assume $\Lambda_l = \{y^{\sigma_{l,h}} \mid h = 1, \ldots, \tau_l\}, l = 1, \ldots, L$. Let us choose $y^{\alpha} \in \Lambda$ and define $C_{\alpha - \sigma_{l,h}}^l$ equal to $c_{\alpha - \sigma_{l,h}}^l$ if $\alpha - \sigma_{l,h} \in \mathcal{A}(\rho(l))$ and zero otherwise. Let us define the linear map

$$\Psi: \mathbb{E}^{\tau_1} \oplus \cdots \oplus \mathbb{E}^{\tau_L} \to \langle \Lambda \setminus \{y^{\alpha}\} \rangle_{\mathbb{E}}$$
$$g = (g_{1,1}, \dots, g_{1,\tau_1}, \dots, g_{L,1}, \dots, g_{L,\tau_L}) \mapsto \sum_{l=1}^L \sum_{h=1}^{\tau_l} g_{l,h}(y^{\sigma_{l,h}} P_l - C^l_{\alpha - \sigma_{l,h}} y^{\alpha}).$$

The columns of S are indexed by the elements of Λ . The matrix $M(\Psi)$ of Ψ , in the monomial bases, is the submatrix of S obtained by removing the column indexed by y^{α} . Observe that S and $M(\Psi)$ are matrices with elements in $\mathbb{Q}[\mathcal{C}]$.

There exists a nonzero $g \in \operatorname{Ker}(\Psi) \cap \mathbb{Q}[\mathcal{C}]^{\sum_{l=1}^{L} \tau_L}$. We can assume w.l.o.g. that $g_{1,1} \neq 0$ There exists a nonsingular matrix E such that $\det(E) = g_{1,1}$ and the first row of $E \cdot S$ has all its entries equal to zero, except for the entry in the column indexed by y^{α} , which equals $\sum_{l=1}^{L} \sum_{h=1}^{\tau_l} g_{l,h} C_{\alpha-\sigma_{l,h}}^l$. Thus

$$g_{1,1}D = \det(E \cdot S) = \gamma \sum_{l=1}^{L} \sum_{h=1}^{\tau_l} g_{l,h} C_{\alpha - \sigma_{l,h}}^l,$$

for some $\gamma \in \mathbb{Q}[\mathcal{C}]$. If we develop D by the column of S indexed by y^{α} we obtain

$$D = \sum_{l=1}^{L} \sum_{h=1}^{\tau_l} r_{l,h} C^l_{\alpha - \sigma_{l,h}}, \text{ with } r_{l,h} \in \mathbb{Q}[\mathcal{C}],$$

which implies $g_{1,1}r_{l,h} = \gamma g_{l,h}$, $l = 1, \ldots, L$, $h = 1, \ldots, \tau_l$. This proves

$$g_{1,1}\sum_{l=1}^{L}\sum_{h=1}^{\tau_l} r_{l,h}(y^{\sigma_{l,h}}P_l - C_{\alpha-\sigma_{l,h}}^l y^{\alpha}) = \gamma \sum_{l=1}^{L}\sum_{h=1}^{\tau_l} g_{l,h}(y^{\sigma_{l,h}}P_l - C_{\alpha-\sigma_{l,h}}^l y^{\alpha}) = 0,$$

and $g_{1,1} \neq 0$ in the domain $\mathbb{Q}[\mathcal{C}][\mathcal{Y}^{\pm}]$ implies

$$\sum_{l=1}^{L} \sum_{h=1}^{\tau_l} r_{l,h} (y^{\sigma_{l,h}} P_l - C_{\alpha - \sigma_{l,h}}^l y^{\alpha}) = 0.$$

Furthermore

$$\sum_{l=1}^{L} \sum_{h=1}^{\tau_l} r_{l,h} y^{\sigma_{l,h}} P_l = y^{\alpha} \sum_{l=1}^{L} \sum_{h=1}^{\tau_l} r_{l,h} C_{\alpha-\sigma_{l,h}}^l = y^{\alpha} D.$$

Since $D \in \mathbb{Q}[\mathcal{C}]$, we have

$$D = y^{-\alpha} \sum_{l=1}^{L} \sum_{h=1}^{\tau_l} r_{l,h} y^{\sigma_{l,h}} P_l \in (\operatorname{ags}(\mathcal{P})) \cap \mathbb{Q}[\mathcal{C}].$$

By [20], Chapter 0, §11, an ideal \mathcal{I} in a polynomial algebra $\mathbb{Q}[\mathcal{C}]$ is prime if and only if it has a generic zero ϵ in $\mathbb{E}^{|\mathcal{C}|}$, for a natural field extension \mathbb{E} of \mathbb{Q} . That is, a polynomial in $\mathbb{Q}[\mathcal{C}]$ belongs to \mathcal{I} if and only if it vanishes at the generic zero ϵ . In the next proof, concepts as autoreduced set and pseudo remainder will be used in the algebraic case, we refer to [24].

Given $l \in \{1, ..., L\}$, let us suppose that $C_l = \{c_l, c_{l,h} \mid h = 1, ..., h_l\}$ and $\{T_l, T_{l,h} \mid h = 1, ..., h_l\} = \{y^{\alpha} \mid \alpha \in \mathcal{A}(\rho(l))\}$, then

$$P_l = c_l T_l + \sum_{h=1}^{h_l} c_{l,h} T_{l,h}, \text{ with } h_l := |\mathcal{A}(\rho(l))| - 1.$$
(8)

Let $\overline{\mathcal{C}} := \mathcal{C} \setminus \{c_1, \ldots, c_L\}$ and define,

$$\epsilon_l := -\sum_{h=1}^{h_l} c_{l,h} \frac{T_{l,h}}{T_l} \text{ and } \epsilon := (\overline{\mathcal{C}}; \epsilon_1, \dots, \epsilon_L).$$
(9)

Lemma 4.2. The elimination ideal $(ags(\mathcal{P})) \cap \mathbb{Q}[\mathcal{C}]$ is a prime ideal with ϵ as a generic zero.

Proof. We only need to prove that ϵ is a generic zero of $\mathcal{I} = (\operatorname{ags}(\mathcal{P})) \cap \mathbb{Q}[\mathcal{C}]$. Given $G \in \mathcal{I}$, $G = \sum_{l} \alpha_{l} P_{l}$, with $\alpha_{l} \in \mathbb{Q}[\mathcal{C}][\mathcal{Y}^{\pm}]$. Since $P_{l}(\epsilon) = \epsilon_{l} T_{l} + \sum_{h=1}^{h_{l}} c_{l,h} T_{l,h} = 0$ we have $G(\epsilon) = 0$. Conversely, let $G \in \mathbb{Q}[\mathcal{C}]$ with $G(\epsilon) = 0$. For each $l \in \{1, \ldots, L\}$, there exists a monomial N_{l} in the variables \mathcal{Y} such that $N_{l}P_{l} \in \mathbb{Q}[\mathcal{C}][\mathcal{Y}]$. Furthermore, $\mathcal{A} = \{N_{1}P_{1}, \ldots, N_{L}P_{L}\}$ is an autoreduced set with c_{l} as leaders. Let G_{0} be the pseudo remainder of G w.r.t. \mathcal{A} , that is $MG = \sum_{l} \beta_{l} N_{l} P_{l} + G_{0}$, for some monomial M in \mathcal{Y} . Observe that $G_{0} \in \mathbb{Q}[\overline{\mathcal{C}}][\mathcal{Y}]$ because each $N_{l}P_{l}$ is linear in c_{l} . Hence

$$G_0 = G_0(\epsilon) = MG(\epsilon) - \sum_l \beta_l(\epsilon) N_l P_l(\epsilon) = 0$$

and

$$G = \sum_{l} \beta_l \frac{N_l}{M} P_l = \sum_{l} \gamma_l P_l, \ \gamma_l \in \mathbb{Q}[\mathcal{C}][\mathcal{Y}^{\pm}].$$

Thus $G \in \mathcal{I}$ and the result is proved.

Let us denote the system of generic algebraic polynomials $\operatorname{ags}(\mathcal{P})$ by \mathcal{S} . The dimension of $(\mathcal{S}) \cap \mathbb{Q}[\mathcal{C}]$ is by definition the transcendence degree of $\mathbb{Q}(\epsilon)$ over \mathbb{Q} ([20], Chapter 0, §11), let us denote it by $\operatorname{trdeg}(\mathbb{Q}(\epsilon)/\mathbb{Q})$.

Remark 4.3. If $\operatorname{trdeg}(\mathbb{Q}(\epsilon)/\mathbb{Q}) = L - 1$ then $(S) \cap \mathbb{Q}[\mathcal{C}]$ is a prime ideal of codimension one, which implies it is a principal ideal. Namely, there exists an irreducible polynomial denoted by $\mathsf{R}(S)$ in $\mathbb{Z}[\mathcal{C}]$ such that $(S) \cap \mathbb{Q}[\mathcal{C}] = (\mathsf{R}(S))$. If $\operatorname{trdeg}(\mathbb{Q}(\epsilon)/\mathbb{Q}) < L - 1$ we define $\mathsf{R}(S)$ to be equal to 1.

A vector of coefficients for the system $S = \{P_1, \ldots, P_L\}$ defines a point **c** of the product of complex projective spaces $\mathbf{P}^{h_1} \times \cdots \times \mathbf{P}^{h_L}$, namely

$$\mathbf{c} = (\mathbf{c}_1, \mathbf{c}_{1,1}, \dots, \mathbf{c}_{1,h_1}, \dots, \mathbf{c}_L, \mathbf{c}_{L,1}, \dots, \mathbf{c}_{L,h_L}).$$

Let us denote $P_l^{\mathbf{c}} := \mathbf{c}_l T_l + \sum_{h=1}^{h_l} \mathbf{c}_{l,h} T_{l,h}$, $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$ and define

$$Z_0 := \{ \mathbf{c} \in \mathbf{P}^{h_1} \times \cdots \times \mathbf{P}^{h_L} \mid P_l^{\mathbf{c}} = 0, l = 1, \dots, L \text{ have a common solution in } (\mathbb{C}^*)^{L-1} \}.$$

By [25], the Zariski closure Z of Z_0 in $\mathbf{P}^{h_1} \times \cdots \times \mathbf{P}^{h_L}$ is an irreducible variety. As defined in [25], if the codimension of Z is one then the sparse resultant $\operatorname{Res}(\mathcal{S})$ of the system $\mathcal{S} = \operatorname{ags}(\mathcal{P})$ is the irreducible polynomial in $\mathbb{Z}[\mathcal{C}]$ defining the hypersurface Z. If the codimension of Z is greater than one then $\operatorname{Res}(\mathcal{S})$ is defined to be the constant 1. Observe that $(\mathcal{S}) \cap \mathbb{Q}[\mathcal{C}]$ is included in the ideal of the variety Z, thus if Z has codimension one then

$$(\operatorname{Res}(\mathcal{S})) = (\mathcal{S}) \cap \mathbb{Q}[\mathcal{C}] = (\mathsf{R}(\mathcal{S})).$$
(10)

It is proved in [5], Section 6 (see also [32]) that $det(Syl(ags(\mathcal{P})))$ is a nonzero multiple of (the nontrivial) $\operatorname{Res}(\mathcal{S})$, thus $det(Syl(ags(\mathcal{P}))) \in (\mathcal{S}) \cap \mathbb{Q}[\mathcal{C}]$. Note that the proof of Proposition 4.1 is needed only in the case $\operatorname{Res}(\mathcal{S}) = 1$.

Given $J \subseteq \{1, \ldots, L\}$, let us consider the affine lattice \mathcal{L}_J generated by $\sum_{l \in J} \mathcal{A}(\rho(l))$,

$$\mathcal{L}_J = \left\{ \sum_{l \in J} \lambda_l \alpha_l \mid \alpha_l \in \mathcal{A}(\rho(l)), \lambda_l \in \mathbb{Z}, \sum_{l \in J} \lambda_l = 1 \right\},\$$

with $\mathcal{L} := \mathcal{L}_{\{1,...,L\}}$. Let rank (\mathcal{L}_J) denote the rank of \mathcal{L}_J . In [32], the system $\mathcal{S}_J = \{P_l \mid l \in J\}$ is said to be algebraically essential if rank $(\mathcal{L}_J) = |J| - 1$ and rank $(\mathcal{L}_{J'}) \ge |J'|$, for each proper subset J' of J. The condition,

(A2) there exists a unique algebraically essential subsystem S_I of S.

is proved, in [25] and [32], to be a necessary and sufficient for Z to have codimension one (see also [17]). In such case, $\operatorname{Res}(\mathcal{S})$ coincides with $\operatorname{Res}(\mathcal{S}_I)$, considered w.r.t. the lattice \mathcal{L}_I , and hence

$$(\mathcal{S}_I) \cap \mathbb{Q}[\mathcal{C}_I] = (\mathsf{R}(\mathcal{S}_I)) = (\operatorname{Res}(\mathcal{S}_I)), \tag{11}$$

with $C_I = \bigcup_{l \in I} C_l$ and (S_I) the ideal generated by S_I in $\mathbb{Q}[C_I][\mathcal{Y}(S_I)^{\pm}]$, with $\mathcal{Y}(S_I)$ as in (7).

In [25], if S is essential, the degree of $\operatorname{Res}(S)$ in C_l , $l = 1, \ldots, L$ was proved to equal the normalized mixed volume

$$MV_{-l}(\mathcal{S}) := \mathcal{M}(\mathcal{Q}_h \mid h \in \{1, \dots, L\} \setminus \{l\}) = \frac{\sum_{J \subset \{1, \dots, L\} \setminus \{l\}} (-1)^{L-|J|} \operatorname{vol}\left(\sum_{j \in J} \mathcal{Q}_j\right)}{\operatorname{vol}(\mathcal{Q})}, \quad (12)$$

where \mathcal{Q}_l is the convex hull of $\mathcal{A}(\rho(l))$ in $\mathcal{L} \otimes \mathbb{R}$, $\operatorname{vol}(\mathcal{Q}_l)$ its L-1 dimensional volume, $\sum_{j \in J} \mathcal{Q}_j$ is the Minkowski sum of \mathcal{Q}_j , $j \in J$ and \mathcal{Q} a fundamental lattice parallelotope in \mathcal{L} .

The Sylvester matrix $Syl(ags(\mathcal{P}))$ constructed in [5] and [13] assigns a special role to P_1 , let us denote $S_1(\mathcal{P}) := Syl(ags(\mathcal{P}))$. The same construction can be done choosing P_l , $l = 2, \ldots, L$ as a distinguished polynomial, obtaining a matrix denoted by $S_l(\mathcal{P})$. As noted in [5], Section 9 and [13], Section 4.3, $S_l(\mathcal{P})$ has the minimum number of rows containing coefficients of P_l , its degree in the coefficients of P_l coincides with the degree of $\operatorname{Res}(\mathcal{S})$ in the coefficients of P_l . Furthermore, $\operatorname{Res}(\mathcal{S})$ can be computed as the gcd in $\mathbb{Q}[\mathcal{C}]$ of the determinants

$$D_l(\mathcal{P}) := \det(S_l(\mathcal{P})), \ l = 1, \dots, L.$$
(13)

Example 4.4. The next system $\mathcal{P} = \{f_1, f_2\}$ in $\mathcal{D}\{u_1\}$, is a simplified version of a predator-prey model studied in [12] that we take as a toy example,

$$f_1 = a_2 x + (a_1 + a_4 x)u_1 + u_{1,1} + (a_3 + a_6 x)u_1^2 + a_5 u_1^3,$$

$$f_2 = x' + (b_1 + b_3 x)u_1 + (b_2 + b_5 x)u_1^2 + b_4 u_1^3,$$

with a_i, b_j algebraic indeterminates over \mathbb{Q} , $\mathcal{D} = \mathbb{Q}(t)[a_i, b_j]\{x\}$ and $\partial = \frac{\partial}{\partial t}$. The first attempt to eliminate the differential variable u_1 was done using the Maple package diffalg, [3] (using characteristic set methods). The computation was interrupted with no answer after two hours. We carry the example to show the elimination of u_1 . Computations were done with Maple 15.

Since $ps(\mathcal{P}) = \{f_1, f_2, \partial f_2\}$, with $\partial f_2 = x'' + b_3 x' u_1 + (b_3 x + b_1) u_{1,1} + b_5 x' u_1^2 + (2b_5 x + 2b_2) u_1 u_{1,1} + 3b_4 u_1^2 u_{1,1}$ and $\mathcal{V}(\mathcal{P}) = \{u_1, u_{1,1}\}$, we have the following associated system of algebraic generic polynomials in y_1, y_2

$$\operatorname{ags}(\mathcal{P}) = \left\{ \begin{array}{l} P_1 = c_1 + c_{11}y_1 + c_{12}y_2 + c_{13}y_1^2 + c_{14}y_1^3, \\ P_2 = c_2 + c_{21}y_1 + c_{22}y_1^2 + c_{23}y_1^3, \\ P_3 = c_3 + c_{31}y_1 + c_{32}y_2 + c_{33}y_1^2 + c_{34}y_1y_2 + c_{35}y_1^2y_2 \end{array} \right\}.$$

Observe that $ags(\mathcal{P})$ is algebraically essential because the linear part of the polynomials in $ags(\mathcal{P})$, $\{c_1 + c_{11}y_1 + c_{12}y_2, c_2 + c_{21}y_1, c_3 + c_{31}y_1 + c_{32}y_2\}$ is an algebraically essential system that verifies (A1). Thus the algebraic resultant $Res(\mathcal{P})$ is nontrivial. Using "toricres04", Maple 9 code for sparse (toric) resultant matrices by I.Z. Emiris, [5], we obtain a 12×12 matrix $S_1(\mathcal{P})$ whose rows contain the coefficients of the polynomials

$$y_1P_1, y_1y_2P_1, y_1y_2^2P_1, y_1^2P_2, y_1y_2P_2, y_1^2y_2P_2, y_1y_2^2P_2, y_1^2y_2^2P_2, y_1P_3, y_1y_2P_3, y_1y_2^2P_3, y_1y_2^3P_3$$

 $in \ the \ monomials$

$$y_1, y_1^2, y_1y_2, y_1^2y_2, y_1y_2^2, y_1^2y_2^2, y_1y_2^3, y_1^2y_2^3, y_1y_2^4, y_1^2y_2^4, y_1y_2^5, y_1^2y_2^5$$

If the order of the input polynomials is P_2 , P_3 , P_1 , we get a 13×13 matrix $S_2(\mathcal{P})$ and if the order is P_3 , P_1 , P_2 , the matrix $S_3(\mathcal{P})$ obtained is 11×11 , namely

c_1	c_{12}	c_{11}	0	c_{13}	0	c_{14}	0	0	0	0]
c_3	c_{32}	c_{31}	c_{34}	c_{33}	c_{35}	0	0	0	0	0	
0	0	c_1	c_{12}	c_{11}	0	c_{13}	0	c_{14}	0	0	
0	0	c_3	c_{32}	c_{31}	c_{34}	c_{33}	c_{35}	0	0	0	
0	0	0	0	c_1	c_{12}	c_{11}	0	c_{13}	0	c_{14}	
0	0	0	0	c_3	c_{32}	c_{31}	c_{34}	c_{33}	c_{35}	0	
c_2	0	c_{21}	0	c_{22}	0	c_{23}	0	0	0	0	
0	c_2	0	c_{21}	0	c_{22}	0	c_{23}	0	0	0	
0	0	c_2	0	c_{21}	0	c_{22}	0	c_{23}	0	0	
0	0	0	c_2	0	c_{21}	0	c_{22}	0	c_{23}	0	
0	0	0	0	c_2	0	c_{21}	0	c_{22}	0	c_{23}	

The determinants of these matrices are

$$D_1(\mathcal{P}) = -c_3 \operatorname{Res}(\mathcal{P}), D_2(\mathcal{P}) = c_1^2 \operatorname{Res}(\mathcal{P}) \text{ and } D_3(\mathcal{P}) = \operatorname{Res}(\mathcal{P}).$$

5 Differential specialization

We are ready to define differential resultant formulas for \mathcal{P} , through the specialization of the previously defined Sylvester matrices.

Given $f \in ps(\mathcal{P})$, with $f = \sum_{\alpha \in \mathcal{A}(f)} a_{\alpha}^{f} v(y^{\alpha})$, let us denote by $A_{f} := \{a_{\alpha}^{f} \mid \alpha \in \mathcal{A}(f)\}$ its coefficient set and $A(\mathcal{P}) := \bigcup_{f \in ps(\mathcal{P})} A_{f}.$ (14)

Given
$$l \in \{1, \ldots, L\}$$
, such that $\rho(l) = f$, and $c_{\alpha}^{l} \in C_{l}$, it holds that $a_{\alpha}^{\rho(l)} \in A_{f}$. Thus we can define the specialization map

$$\Xi: \mathcal{C} \to A(\mathcal{P}), \text{ by } \Xi(c^l_\alpha) = a^{\rho(l)}_\alpha,$$

which naturally extends to a ring epimorphism, defining $\Xi(y_l) = \upsilon(y_l)$,

$$\Xi: \mathbb{Q}[\mathcal{C}][\mathcal{Y}^{\pm}] \to \mathbb{Q}[A(\mathcal{P})][\mathcal{V}(\mathcal{P})^{\pm}].$$

 $\mathbb{Q}[A(\mathcal{P})][\mathcal{V}(\mathcal{P})^{\pm}]$ is included in the differential ring $\mathbb{Q}\{A(\mathcal{P})\}\{U^{\pm}\} \subseteq \mathcal{D}\{U^{\pm}\}$ and obviously

$$\Xi(P_l) = \rho(l) \in \operatorname{ps}(\mathcal{P}), l = 1, \dots, L$$

Let us assume that \mathcal{P} is supper essential to define the determinants $D_l(\mathcal{P})$, $l = 1, \ldots, L$ in (13). By Proposition 4.1, each $D_l(\mathcal{P})$ belongs to the ideal $(ags(\mathcal{P})) \cap \mathbb{Q}[\mathcal{C}]$ and

$$\Xi(D_l(\mathcal{P})) \in [\mathcal{P}] \cap \mathcal{D}.$$
(15)

As defined in (2), $\Xi(D_l(\mathcal{P}))$ is a differential resultant formula for \mathcal{P} with

$$L_i = J_i(\mathcal{P}) - \gamma(\mathcal{P}), \ \mathcal{U} = \mathcal{V}(\mathcal{P}) \text{ and } \Omega_f = \Xi(\Lambda_{\lambda(f)}), \Omega = \Xi(\Lambda),$$

 $f \in ps(\mathcal{P}).$

Example 5.1. To finish Example 4.4. The specializations $\Xi(D_1(\mathcal{P}))$, $\Xi(D_2(\mathcal{P}))$ and $\Xi(D_3(\mathcal{P}))$ are nonzero polynomials in the differential elimination ideal $[\mathcal{P}] \cap \mathcal{D}$ (they are not included due to their size), in particular $\Xi(D_3(\mathcal{P})) = \Xi(\text{Res}(\mathcal{P}))$.

Observe that, even for nonzero $D_l(\mathcal{P})$, $\Xi(D_l(\mathcal{P}))$ could be zero, in which case the perturbation methods in [14] could be used to obtain a nonzero differential polynomial in $[\mathcal{P}] \cap \mathcal{D}$. Alternatively, an algorithm to specialize step by step and obtain a factor of the specialization, which is a nonzero differential polynomial in $[\mathcal{P}] \cap \mathcal{D}$, is proposed next. A similar argument is used in other specialization results as [18], p. 168-169 and it was used in the proof of [21], Theorem 6.5 for a system of non sparse generic non homogeneous differential polynomials.

For i = 1, ..., n, let us assume that $A_{f_i} = \{a_i, a_{i,k} \mid k = 1, ..., l_i\}$ and $\{v(y^{\alpha}) \mid \alpha \in \mathcal{A}(f_i)\} = \{M_i, M_{i,k} \mid k = 1, ..., l_i\}$, then

$$f_i = a_i M_i + a_{i,1} M_{i,1} + \dots + a_{i,l_i} M_{i,l_i}.$$

In the remaining parts of this section, we consider differential indeterminates a_i , i = 1, ..., n over \mathbb{Q} and the system

$$\tilde{\mathcal{P}} := \{ F_i := \mathfrak{a}_i M_i + a_{i,1} M_{i,1} + \dots + a_{i,l_i} M_{i,l_i} \mid i = 1, \dots, n \},\$$

of sparse Laurent differential polynomials in $\tilde{\mathcal{D}}\{U^{\pm}\}$, with differential domain $\tilde{\mathcal{D}} := \mathcal{D}\{\mathfrak{a}_1, \ldots, \mathfrak{a}_n\}$. Observe that $\operatorname{ags}(\tilde{\mathcal{P}}) = \operatorname{ags}(\mathcal{P}) = \{P_1, \ldots, P_L\}$, thus $D_l(\mathcal{P}) = D_l(\tilde{\mathcal{P}})$. Let us assume that specialization map $\Xi : \mathcal{C} \to A(\tilde{\mathcal{P}})$ verifies

$$\Xi(c_l) = \partial^k \mathfrak{a}_i, \text{ if } \rho(l) = \partial^k F_i, \tag{16}$$

given $l \in \{1, \ldots, L\}$ and as in (8)

$$P_l = c_l T_l + \sum_{h=1}^{h_l} c_{l,h} T_{l,h}$$
, with $h_l := |\mathcal{A}(\rho(l))| - 1$.

Thus $\Xi(\{c_l, c_{l,h} \mid h = 1, \dots, h_l\}) = A_{\rho(l)}$ and

$$A(\tilde{\mathcal{P}}) = \Xi(\overline{\mathcal{C}}) \cup \Xi(\mathbf{c}) = \Xi(\overline{\mathcal{C}}) \cup \{\mathfrak{a}_1, \dots, \partial^{J_1 - \gamma} \mathfrak{a}_1, \dots, \mathfrak{a}_n, \dots, \partial^{J_n - \gamma} \mathfrak{a}_n\},$$

with $C = \overline{C} \cup \mathbf{c}$ and $\mathbf{c} = \{c_1, \ldots, c_L\}.$

The idea is that, to study the specialization of $D_l(\mathcal{P})$ to the coefficients $A(\mathcal{P})$, one can first study the specialization to the coefficients $A(\tilde{\mathcal{P}})$ and then specialize $\{\mathfrak{a}_i \mid i = 1, \ldots, n\}$ to $\{a_i \mid i = 1, \ldots, n\}$. We dedicate the rest of the section to the first part of this specialization. The results obtained will be used in Section 6 to study the case of sparse generic differential systems. The behavior of the specialization of $\{\mathfrak{a}_i \mid i = 1, \ldots, n\}$ would depend on the specific domain \mathcal{D} to be considered.

Given a nonzero differential polynomial $Q \in (\operatorname{ags}(\tilde{\mathcal{P}})) \cap \mathbb{Q}[\mathcal{C}]$, note that $\Xi(Q) \in [\tilde{\mathcal{P}}] \cap \tilde{\mathcal{D}}$ but we cannot guarantee that $\Xi(Q)$ is nonzero. For a subset Δ of \mathcal{C} , define partial specializations

$$\Xi_{\Delta}: \mathcal{C} \to (\mathcal{C} \setminus \Delta) \cup \Xi(\Delta), \text{ by } \Xi_{\Delta}(c) = \begin{cases} \Xi(c), & c \in \Delta, \\ c, & c \notin \Delta, \end{cases}$$

which naturally extend to ring epimorphisms

$$\Xi_{\Delta}: \mathbb{Q}[\mathcal{C}][\mathcal{Y}^{\pm}] \to \mathbb{Q}[(\mathcal{C} \setminus \Delta) \cup \Xi(\Delta)][\mathcal{Y}^{\pm}].$$

Observe that we leave the monomials in $\mathbb{Q}[\mathcal{Y}^{\pm}]$ fixed for the moment and, if $\Delta = \mathcal{C}$ then $\Xi_{\Delta}(\mathbb{Q}[\mathcal{C}]) = \mathbb{Q}[A(\tilde{\mathcal{P}})]$. Let

$$\Xi_{\mathcal{Y}^{\pm}}: \mathbb{Q}[A(\tilde{\mathcal{P}})][\mathcal{Y}^{\pm}] \to \mathbb{Q}[A(\tilde{\mathcal{P}})][\mathcal{V}(\tilde{\mathcal{P}})^{\pm}],$$

be defined by $\Xi_{\mathcal{Y}^{\pm}}(y_l) = \upsilon(y_l).$

Algorithm 5.2. • <u>Given</u> a nonzero polynomial Q in $(ags(\tilde{\mathcal{P}})) \cap \mathbb{Q}[\mathcal{C}]$.

- <u>Return</u> a nonzero differential polynomial H in $[\tilde{\mathcal{P}}] \cap \tilde{\mathcal{D}}$.
- 1. Let $\Delta := \emptyset$ and H := Q.
- 2. If $\mathcal{C} \setminus \Delta = \emptyset$, return $\Xi_{\mathcal{Y}^{\pm}}(H)$.
- 3. Choose $c \in \mathcal{C} \setminus \Delta$ and define $\Delta := \Delta \cup \{c\}$.
- 4. If $\Xi_{\Delta}(H) \neq 0$ then $H := \Xi_{\Delta}(H)$, go to step 2.
- 5. $H = (c \Xi(c))^s \overline{H}, s \in \mathbb{N} \setminus \{0\}, set H := \Xi_{\Delta}(\overline{H}) \neq 0 and go to step 2.$

We prove next that the output of the previous algorithm is a nonzero differential polynomial in $[\tilde{\mathcal{P}}] \cap \tilde{\mathcal{D}}$. Given $\emptyset \neq \Delta \subset \mathcal{C}$, observe that $\Xi_{\Delta}(c_l)$ equals $\partial^k \mathfrak{a}_i$, if $c_l \in \Delta$, and c_l otherwise. Let us consider ideals

$$\mathcal{I}_{\Delta,\mathcal{Y}^{\pm}} := (\Xi_{\Delta}(P_1), \dots, \Xi_{\Delta}(P_L))_{\mathbb{D}_{\Delta}[\mathcal{Y}^{\pm}]}$$

generated by $\Xi_{\Delta}(ags(\tilde{\mathcal{P}}))$ in $\mathbb{D}_{\Delta}[\mathcal{Y}^{\pm}]$, with

$$\mathbb{D}_{\Delta} := \begin{cases} \mathbb{Q}[\mathcal{C}] \text{ if } \Delta = \emptyset, \\ \mathbb{Q}[\Xi(\overline{\mathcal{C}} \cap \Delta)][\overline{\mathcal{C}} \setminus \Delta \cup \Xi_{\Delta}(\mathbf{c})] \text{ if } \Delta \neq \emptyset \end{cases}$$

Observe that $\mathcal{I}_{\mathcal{C},\mathcal{Y}^{\pm}}$ is the ideal generated by $\Xi_{\mathcal{C}}(ags(\tilde{\mathcal{P}}))$ in

$$\mathbb{D}_{\mathcal{C}}[\mathcal{Y}^{\pm}] = \mathbb{Q}[\Xi(\overline{\mathcal{C}})][\mathfrak{a}_1, \dots, \partial^{J_1 - \gamma}\mathfrak{a}_1, \dots, \mathfrak{a}_n, \dots, \partial^{J_n - \gamma}\mathfrak{a}_n][\mathcal{Y}^{\pm}].$$

Let $\mathbb{K}_{\Delta} := \mathbb{Q}(\Xi(\overline{\mathcal{C}} \cap \Delta))$ if $\Delta \neq \emptyset$ and $\mathbb{K}_{\emptyset} := \mathbb{Q}$. Observe that

$$\mathbb{D}_{\Delta} \subset \mathbb{K}_{\Delta}[\overline{\mathcal{C}} \setminus \Delta \cup \Xi_{\Delta}(\mathbf{c})]. \tag{17}$$

Lemma 5.3. $\mathcal{I}_{\Delta,\mathcal{Y}^{\pm}} \cap \mathbb{K}_{\Delta}[\overline{\mathcal{C}} \setminus \Delta \cup \Xi_{\Delta}(\mathbf{c})]$ is a prime ideal.

Proof. As in the proof of Lemma 4.2, we prove that $\mathcal{I} = \mathcal{I}_{\Delta,\mathcal{Y}^{\pm}} \cap \mathbb{K}_{\Delta}[\overline{\mathcal{C}} \setminus \Delta \cup \Xi_{\Delta}(\mathbf{c})]$ has a generic zero. Let us define

$$\epsilon_l^{\Delta} := -\sum_{h=1}^{n_l} \Xi_{\Delta}(c_{l,h}) \frac{T_{l,h}}{T_l}.$$

We can adapt the proof of Lemma 4.2 to show that $\epsilon^{\Delta} := (\overline{\mathcal{C}} \setminus \Delta; \epsilon_1^{\Delta}, \ldots, \epsilon_L^{\Delta})$ is a generic zero of \mathcal{I} . Observe that $\Xi_{\Delta}(P_l) = \Xi_{\Delta}(c_l)T_l + \sum_h \Xi_{\Delta}(c_{l,h})T_{l,h}$ and $\Xi_{\Delta}(c_l)$ is replaced by ϵ_l^{Δ} . By [20], Chapter 0, Section 11, \mathcal{I} is a prime ideal.

Theorem 5.4. Given a nonzero differential polynomial in $(\operatorname{ags}(\tilde{\mathcal{P}})) \cap \mathbb{Q}[\mathcal{C}]$, the output of Algorithm 5.2 is a nonzero differential polynomial in $[\tilde{\mathcal{P}}] \cap \tilde{\mathcal{D}}$.

Proof. Let $c \in \mathcal{C} \setminus \Delta$, if $H \in \mathcal{I}_{\Delta,\mathcal{Y}^{\pm}} \cap \mathbb{D}_{\Delta}$ verifies $\Xi_{\Delta}(H) = 0$ then $H = (c - \Xi(c))^{s}\overline{H}$ with $\overline{H} \in \mathbb{D}_{\Delta}$. By Lemma 5.3 and (17), $\overline{H} \in \mathcal{I}_{\Delta,\mathcal{Y}^{\pm}} \cap \mathbb{D}_{\Delta}$. For $\Delta' = \Delta \cup \{c\}$, if $H \in \mathcal{I}_{\Delta,\mathcal{Y}^{\pm}} \cap \mathbb{D}_{\Delta}$ then

$$\Xi_{\Delta'}(H) \in \mathcal{I}_{\Delta',\mathcal{Y}^{\pm}} \cap \mathbb{D}_{\Delta'}.$$

Thus steps 4 and 5 of Algorithm 5.2 return polynomials in $\mathcal{I}_{\Delta',\mathcal{V}^{\pm}}$. If $\Delta' = \mathcal{C}$ then step 2 returns $\Xi_{\mathcal{V}^{\pm}}(H)$ that belongs to $(\mathrm{ps}(\tilde{\mathcal{P}}))$ the ideal generated by $\mathrm{ps}(\tilde{\mathcal{P}})$ in $\mathbb{Q}[A(\tilde{\mathcal{P}})][\mathcal{V}(\tilde{\mathcal{P}})^{\pm}]$, thus $\Xi_{\mathcal{V}^{\pm}}(H) \in [\tilde{\mathcal{P}}] \cap \tilde{\mathcal{D}}$.

Therefore, if $D_l(\tilde{P})$ is nonzero, by Proposition 4.1 it can be taken as the input of Algorithm 5.2 and, by Theorem 5.4, we obtain a nonzero differential polynomial in $[\tilde{\mathcal{P}}] \cap \tilde{\mathcal{D}}$.

6 Order and degree bounds

In this section, we prove that the formulas obtained are multiples of the differential resultant of a system of generic sparse differential polynomials, defined by Li, Gao and Yuan in [21] and whose definition we include below. This fact is used to give order and degree bounds of the sparse differential resultant.

Let us consider sets of differential indeterminates over \mathbb{Q} , $A_i := \{\mathfrak{a}^i_\alpha \mid \alpha \in \mathcal{A}(f_i)\}, i = 1, ..., n, A := \bigcup_{i=1}^n A_i$ and a differential domain $\mathfrak{D} := \mathbb{Q}\{A\}$, to define the system $\mathfrak{P} := \{\mathbb{F}_1, \ldots, \mathbb{F}_n\}$ of sparse generic differential polynomials in $\mathfrak{D}\{U^{\pm}\}$. The sparse generic polynomial \mathbb{F}_i in $\mathfrak{D}\{U^{\pm}\}$ with algebraic support $\mathcal{A}(f_i)$ is

$$\mathbb{F}_i := \sum_{\alpha \in \mathcal{A}(f_i)} \mathfrak{a}^i_{\alpha} \upsilon(y^{\alpha}),$$

which has order o_i . The ideal generated by \mathfrak{P} in $\mathfrak{D}\{U^{\pm}\}$ is denoted by $[\mathfrak{P}]$. In this section J_i and γ denote $J_i(\mathfrak{P})$ and $\gamma(\mathfrak{P})$ respectively. Let $A(\mathfrak{P})$ be as in (14) and $\partial^k A_i := \{\partial^k \mathfrak{a} \mid \mathfrak{a} \in A_i\}, k \in \mathbb{N}$. It holds

$$A \subset A(\mathfrak{P}) \text{ and } \mathbb{Q}[A(\mathfrak{P})] = \mathbb{Q}[\cup_{i=1}^{n} A_i^{[J_i - \gamma]}],$$

with

$$A_i^{[\tau_i]} := \begin{cases} \bigcup_{i=1}^n \bigcup_{k=0}^{\tau_i} \partial^k A_i \text{ if } \tau_i \in \mathbb{N}, \\ \emptyset \text{ if } \tau_i = -\infty. \end{cases}$$
(18)

If the differential elimination ideal $[\mathfrak{P}] \cap \mathfrak{D}$ has dimension n-1 then $[\mathfrak{P}] \cap \mathfrak{D} = \operatorname{sat}(\partial \operatorname{Res}(\mathfrak{P}))$, the saturated ideal determined by an irreducible differential polynomial $\partial \operatorname{Res}(\mathfrak{P})$, which is called the sparse differential resultant of \mathfrak{P} , [21], Definition 3.10. The saturated ideal of $\partial \operatorname{Res}(\mathfrak{P})$ is the set of all differential polynomials in \mathfrak{D} whose differential remainder (under any elimination ranking) w.r.t. $\partial \operatorname{Res}(\mathfrak{P})$ is zero, see [26] and [20].

Lemma 6.1. For every $H \in [\mathfrak{P}] \cap \mathfrak{D}$

$$\omega_i := \operatorname{ord}(\partial \operatorname{Res}(\mathfrak{P}), A_i) \le \operatorname{ord}(H, A_i), \ i = 1, \dots, n$$

Proof. If $\operatorname{ord}(H, A_k) = -\infty$ then $\operatorname{ord}(\partial \operatorname{Res}(\mathfrak{P}, A_k)) = -\infty$, since otherwise H cannot be reduced to zero by an elimination ranking with elementes in A_k greater than the elements of A_i , $i \neq k$. Thus the result follows easily from $H \in \operatorname{sat}(\partial \operatorname{Res}(\mathfrak{P}))$.

For i = 1, ..., n, let us assume that $A_i = \{\mathfrak{a}_i, \mathfrak{a}_{i,k} \mid k = 1, ..., l_i\}$ and $\{\upsilon(y^{\alpha}) \mid \alpha \in \mathcal{A}(f_i)\} = \{M_i, M_{i,k} \mid k = 1, ..., l_i\}$, then

$$\mathbb{F}_i = \mathfrak{a}_i M_i + \mathfrak{a}_{i,1} M_{i,1} + \dots + \mathfrak{a}_{i,l_i} M_{i,l_i}.$$
(19)

By [21], Definition 3.6, \mathfrak{P} is Laurent differentially essential if for each $i \in \{1, \ldots, n\}$ there exists $k_i \in \{1, \ldots, l_i\}$ such that the differential transcendence degree of the set of monomials $\{M_{i,k_i}M_i^{-1} \mid i = 1, \ldots, n\}$ over \mathbb{Q} is n-1.

Let \mathcal{I} be a differential ideal in $\mathfrak{D} = \mathbb{Q}\{A\}$. Given a differential field extension \mathcal{E} of \mathbb{Q} , ζ in $\mathcal{E}^{|A|}$ is called a generic zero of \mathcal{I} if, a differential polynomial $F \in \mathfrak{D}$ belongs to \mathcal{I} if and only if $F(\zeta) = 0$, [26], p. 27. Furthermore, \mathcal{I} is prime if and only if it has a generic zero. Given $\overline{A} := A \setminus \{\mathfrak{a}_1, \ldots, \mathfrak{a}_n\}$, the differential field extension $\mathbb{Q}\langle \overline{A}, \mathcal{Y}^{\pm} \rangle$ of \mathbb{Q} contains

$$\zeta_i := -\sum_{k=1}^{l_i} \mathfrak{a}_{i,k} \frac{M_{i,k}}{M_i}, i = 1, \dots, n.$$
$$\zeta := (\overline{A}; \zeta_1, \dots, \zeta_n)$$
(20)

By [21], Corollary 3.12,

is a generic zero of the differential prime ideal $[\mathfrak{P}] \cap \mathfrak{D}$, which has codimension one if and only if \mathfrak{P} is Laurent differentially essential. To prove this last claim, in [21], Theorem 3.9, it is proved that the differential transcendence degree (see [20]) of $\mathbb{Q}\langle\zeta\rangle$ over \mathbb{Q} is $|A| - 1 = |\overline{A}| + n - 1$, using the next result.

Lemma 6.2 ([21], proof of Theorem 3.9). For $\mathcal{F} = \mathbb{Q}\langle \overline{A} \rangle$, \mathfrak{P} is Laurent differentially essential if and only if the differential transcendence degree of $\mathcal{F}\langle \zeta_1, \ldots, \zeta_n \rangle$ over \mathcal{F} is n-1.

In the remaining parts of this section, let us assume that \mathfrak{P} is Laurent differentially essential and therefore that $\partial \operatorname{Res}(\mathfrak{P})$ exists. We will use the next result about the order ω_i of $\partial \operatorname{Res}(\mathfrak{P})$ in A_i .

Lemma 6.3 ([21], Lemma 5.4). For i = 1, ..., n, if $\omega_i \ge 0$ then $\omega_i = \operatorname{ord}(\partial \operatorname{Res}(\mathfrak{P}), \mathfrak{a}_i)$ and $\omega_i = \operatorname{ord}(\partial \operatorname{Res}(\mathfrak{P}), \mathfrak{a}_{i,k}), k = 1, ..., l_i$.

In [21], order and degree bounds for $\partial \text{Res}(\mathfrak{P})$ were given. Recall that if $J_i \geq 0$, $i = 1, \ldots, n$ the system \mathfrak{P} is called super essential in Definition 3.2. The next theorem follows from this fact and [21], Theorem 5.13. Emphasize that if \mathfrak{P} is super essential then $J_i - \gamma \geq 0$, $i = 1, \ldots, n$ but $\omega_i \geq 0$ is not guaranted.

Theorem 6.4. Let \mathfrak{P} be a Laurent differentially essential system. If \mathfrak{P} is super essential then

$$\omega_i = \operatorname{ord}(\partial \operatorname{Res}(\mathfrak{P}), A_i) \le J_i - \gamma, \ i = 1, \dots, n.$$
(21)

Observe that to obtain the same conclusion, in [21], Corollary 5.11 a much stronger condition on \mathfrak{P} than super essential was demanded (namely rank essential, see [21], Definition 4.20). The same conclusion is obtained later in this section by Remmarks 6.8 and 6.15.

In this section, we revise the order bounds and we provide degree bounds for the sparse differential resultant of \mathfrak{P} in terms of normalized mixed volumes. The goal of the next results is to prove Theorem 6.11 and its corollaries. They explain under which conditions the nonzero specialization of certain polynomials in the algebraic elimination ideal $(\operatorname{ags}(\mathfrak{P})) \cap \mathbb{Q}[\mathcal{C}]$ are multiples of the sparse differential resultant $\partial \operatorname{Res}(\mathfrak{P})$. Those specializations can then be used to give order and degree bounds of $\partial \operatorname{Res}(\mathfrak{P})$.

Given $\tau = (\tau_1, \dots, \tau_n) \in \mathbb{N}_{-\infty}^n$ with $\mathbb{N}_{-\infty} := \mathbb{N} \cup \{-\infty\}$, let us define $\mathbb{F}_i^{[\tau_i]} := \{\mathbb{F}_i, \partial \mathbb{F}_i, \dots, \partial^{\omega_i} \mathbb{F}_i\}$ if $\tau_i \in \mathbb{N}$ and $\mathbb{F}_i^{[\tau_i]} := \emptyset$ if $\tau_i = -\infty$, $i = 1, \dots, n$, $\kappa_j := \max\{o_{i,j} + \tau_i \mid i = 1, \dots, n\} - \gamma$, $j = 1, \dots, n-1$ and

$$PS(\tau) := \bigcup_{i=1}^{n} \mathbb{F}_{i}^{[\tau_{i}]},$$

$$\mathcal{V}(\tau) := \{u_{j,k} \mid k \in [\gamma_{j}, \kappa_{j}] \cap \mathbb{N}, j = 1, \dots, n-1\}.$$

Observe that, if $\tau_i \leq J_i - \gamma$ then $PS(\tau) \subseteq ps(\mathfrak{P})$. Let us consider the algebraic generic system associated to $PS(\tau)$

$$\mathcal{S}_{\tau} := \left\{ \sum_{\alpha \in \mathcal{A}(\mathbb{F})} c_{\alpha}^{\lambda(\mathbb{F})} y^{\alpha} \mid \mathbb{F} \in \mathrm{PS}(\tau) \right\} \subseteq \mathrm{ags}(\mathfrak{P}),$$

and assume that $S_{\tau} = \{P_{l_1}, \dots, P_{l_{|\mathrm{PS}(\tau)|}}\}$ with, as in (8),

$$P_t := c_t T_t + \sum_{h=1}^{h_t} c_{t,h} T_{t,h}, \ t \in \Lambda(\tau) := \{l_1, \dots, l_{|\mathcal{S}_\tau|}\} \subseteq \{1, \dots, L\}.$$
(22)

The sets of algebraic indeterminates over \mathbb{Q}

 $\mathcal{C}_{\tau} := \bigcup_{t \in \Lambda(\tau)} \mathcal{C}_t = \bigcup_{t \in \Lambda(\tau)} \{c_t, c_{t,h} \mid h = 1, \dots, h_t\} \text{ and } \overline{\mathcal{C}}_{\tau} := \mathcal{C}_{\tau} \setminus \{c_t \mid t \in \Lambda(\tau)\}$

together with $\mathcal{Y}(\mathcal{S}_{\tau}) := \{ y \in \mathcal{Y} \mid v(y) \in \mathcal{V}(\tau) \}$ describe the ring $\mathbb{Q}[\mathcal{C}_{\tau}][\mathcal{Y}(\mathcal{S}_{\tau})^{\pm}]$ that contains \mathcal{S}_{τ} . As in (9)

$$\epsilon_t := -\sum_{h=1}^{h_t} c_{t,h} \frac{T_{t,h}}{T_t}, \ t \in \Lambda(\tau).$$

We assume that the specialization map $\Xi : \mathbb{Q}[\mathcal{C}][\mathcal{Y}^{\pm}] \to \mathbb{Q}[A(\mathfrak{P})][\mathcal{V}(\mathfrak{P})^{\pm}]$ defined in Section 5 verifies (16).

Lemma 6.5. Let (S_{τ}) be the ideal generated by S_{τ} in $\mathbb{Q}[\mathcal{C}_{\tau}][\mathcal{Y}(S_{\tau})^{\pm}]$.

- 1. $(S_{\tau}) \cap \mathbb{Q}[\mathcal{C}_{\tau}]$ is a prime ideal with generic zero $\epsilon_{\tau} := (\overline{\mathcal{C}}_{\tau}; \epsilon_{l_1}, \ldots, \epsilon_{l_{|S_{\tau}|}}).$
- 2. Let $\omega = (\omega_1, \ldots, \omega_n)$, with $\omega_i = \deg(\partial \operatorname{Res}(\mathfrak{P}), A_i)$. The prime ideal $(\mathcal{S}_{\omega}) \cap \mathbb{Q}[\mathcal{C}_{\omega}]$ has codimension one.

Furthermore, there exists an irreducible polynomial $\mathsf{R}(\mathcal{S}_{\omega})$ in $\mathbb{Q}[\mathcal{C}_{\omega}]$ such that $(\mathcal{S}_{\omega}) \cap \mathbb{Q}[\mathcal{C}_{\omega}] = (\mathsf{R}(\mathcal{S}_{\omega})).$

Proof. 1. Analogously to Lemma 4.2 we can prove that ϵ_{τ} is a generic point of $(S_{\tau}) \cap \mathbb{Q}[\mathcal{C}_{\tau}]$.

2. Let us assume w.l.o.g. that $\omega_n \geq 0$ and $\Xi(P_{l_{|S\omega|}}) = \partial^{\omega_n} F_n$. By (16) $\Xi(c_{l_{|S\omega|}}) = \partial^{\omega_n} \mathfrak{a}_n$. We will prove that $\epsilon_{l_1}, \ldots, \epsilon_{l_{|S\omega|-1}}$ are algebraically independent over $\mathbb{Q}(\overline{\mathcal{C}}_{\omega})$. Otherwise, there exists a nonzero polynomial $Q \in \mathbb{Q}[\overline{\mathcal{C}}_{\omega}][c_{l_1}, \ldots, c_{l_{|S\omega|-1}}]$ such that $Q(\epsilon_{l_1}, \ldots, \epsilon_{l_{|S\omega|-1}}) = 0$. By definition of generic zero, $Q \in (S_{\omega}) \cap \mathbb{Q}[\mathcal{C}_{\omega}]$ and, by Theorem 5.4, there exists a nonzero differential polynomial $H \in [\mathfrak{P}] \cap \mathfrak{D} = \operatorname{sat}(\partial \operatorname{Res}(\mathfrak{P}))$ given by Algorithm 5.2. Observe that $\partial^{\omega_n}\mathfrak{a}_n$ cannot appear in H, by definition of Q. Thus $\operatorname{ord}(H,\mathfrak{a}_n) \leq \omega_n - 1 < \operatorname{ord}(\partial \operatorname{Res}(\mathfrak{P}),\mathfrak{a}_n) = \omega_n$, contradicting that $\partial \operatorname{Res}(\mathfrak{P})$ is the differential resultant, by Lemma 6.3. This proves that

$$\operatorname{trdeg}(\mathbb{Q}(\epsilon_{\omega})/\mathbb{Q}) = |\overline{\mathcal{C}}_{\omega}| + |\mathcal{S}_{\omega}| - 1 = |\mathcal{C}_{\omega}| - 1$$

and hence $(\mathcal{S}_{\omega}) \cap \mathbb{Q}[\mathcal{C}_{\omega}]$ has codimension one. The conclusion follows as in Remark 4.3.

The second part of the next lemma is part of the proof of [21], Theorem 6.5 and it is used there to bound the degree of $\partial \text{Res}(\mathfrak{P})$.

Lemma 6.6. Let $(PS(\tau))$ be the ideal generated by $PS(\tau)$ in $\mathbb{Q}[\Xi(\mathcal{C}_{\tau})][\mathcal{V}(\tau)^{\pm}]$.

- 1. $(PS(\tau)) \cap \mathbb{Q}[\Xi(\mathcal{C}_{\tau})]$ is a prime ideal with generic zero ζ_{τ} given by (24).
- 2. Let $\omega = (\omega_1, \ldots, \omega_n)$, with $\omega_i = \deg(\partial \operatorname{Res}(\mathfrak{P}), A_i)$. The prime ideal $(\operatorname{PS}(\omega)) \cap \mathbb{Q}[\Xi(\mathcal{C}_{\omega})]$ has codimension one and is equal to $(\partial \operatorname{Res}(\mathfrak{P}))$.

Proof. 1. Let ζ be as in (20) and \mathfrak{a}_i as in (19). If $\tau_i \in \mathbb{N}$, let us define the sets

$$\zeta_i^{[\tau_i]} := \{\zeta_i, \partial \zeta_i, \dots, \partial^{\tau_i} \zeta_i\} \text{ and } \mathfrak{a}_i^{[\tau_i]} := \{\mathfrak{a}_i, \dots, \partial^{\tau_i} \mathfrak{a}_i\},\$$

otherwise these sets are defined to be empty. $A_i^{[\tau_i]}$ was defined in (18). Let $\mathfrak{a} := \bigcup_{i=1}^n \mathfrak{a}^{[\tau_i]}$ and observe that

$$\mathbb{Q}[\Xi(\mathcal{C}_{\omega})] = \mathbb{Q}[\bigcup_{i=1}^{n} A_{i}^{[\tau_{i}]}].$$
(23)

For $\partial^k \mathbb{F}_i$ in $\mathrm{PS}(\omega)$ it holds

$$\partial^k \mathbb{F}_i(\zeta) = \partial^k \mathbb{F}_i(\zeta_i, \partial \zeta_i, \dots, \partial^k \zeta_i) = 0.$$

Analogously to Lemma 4.2, it can be proved that $(PS(\omega)) \cap \mathbb{Q}[\Xi(\mathcal{C}_{\omega})]$ is a prime ideal with

$$\zeta_{\tau} := \left(\bigcup_{i=1}^{n} A_{i}^{\lfloor \tau_{i} \rfloor} \backslash \mathfrak{a}; \bigcup_{i=1}^{n} \zeta_{i}^{\lfloor \tau_{i} \rfloor} \right)$$
(24)

as a generic zero.

2. Similarly to Lemma 6.5, (2) it follows that $|PS(\omega)| - 1$ of the elements in $\bigcup_{i=1}^{n} \zeta_i^{[\omega_i]}$ are algebraically independent and therefore this ideal has codimension one.

By Lemma 6.3 and (23), $\partial \operatorname{Res}(\mathfrak{P}) \in \mathbb{Q}[\Xi(\mathcal{C}_{\omega})]$. Since ζ in (20) is a generic zero of $[\mathfrak{P}] \cap \mathfrak{D} = \operatorname{sat}(\partial \operatorname{Res}(\mathfrak{P}))$, it holds $\partial \operatorname{Res}(\mathfrak{P})(\zeta) = 0$, which means that replacing $\partial^k \mathfrak{a}_i$ by $\partial^k \zeta_i$, $k = 0, \ldots, \omega_i$, $\partial \operatorname{Res}(\mathfrak{P})$ becomes zero (for $i \in \{1, \ldots, n\}$ with $\omega_i \geq 0$). This implies that $\partial \operatorname{Res}(\mathfrak{P})$ is an irreducible polynomial in $(\operatorname{PS}(\omega)) \cap \mathbb{Q}[\Xi(\mathcal{C}_{\omega})]$ and proves the result.

Observe that

$$\Xi((\mathcal{S}_{\tau}) \cap \mathbb{Q}[\mathcal{C}_{\tau}]) \subset (\mathrm{PS}(\tau)) \cap \Xi(\mathbb{Q}[\mathcal{C}_{\tau}]).$$
(25)

The next result shows that the irreducible polynomials $\mathsf{R}(\mathcal{S}_{\omega})$ and $\partial \operatorname{Res}(\mathfrak{P})$, defining respectively algebraic and differential elimination ideals of codimension one, are related through the specialization process.

Proposition 6.7. Let \mathfrak{P} be a Laurent differentially essential system and $\omega = (\omega_1, \ldots, \omega_n)$, with $\omega_i = \operatorname{ord}(\partial \operatorname{Res}(\mathfrak{P}), A_i)$. There exists $\mathfrak{E} \in \mathbb{Q}[\Xi(\mathfrak{C})]$ such that $\Xi(\mathsf{R}(\mathcal{S}_\omega)) = \mathfrak{E}\partial \operatorname{Res}(\mathfrak{P})$ and

$$\deg(\partial \operatorname{Res}(\mathfrak{P}), A_i^{[\omega_i]}) \le \sum_{k=0}^{\omega_i} \deg(\mathsf{R}(\mathcal{S}_{\omega}), C_{\lambda(\partial^k \mathbb{F}_i)}), \ i = 1, \dots, n.$$

Proof. By Lemmas 6.5, 6.6 and (25)

$$\Xi((\mathsf{R}(\mathcal{S}_{\omega}))) = \Xi((\mathcal{S}_{\omega}) \cap \mathbb{Q}[\mathcal{C}_{\omega}]) \subset (\partial \operatorname{Res}(\mathfrak{P})) = (\operatorname{PS}(\omega)) \cap \mathbb{Q}[\Xi(\mathcal{C}_{\omega})].$$

Thus $\Xi(\mathsf{R}(\mathcal{S}_{\omega})) = \mathfrak{E}\partial \operatorname{Res}(\mathfrak{P})$, with $\mathfrak{E} \in \mathbb{Q}[\Xi(\mathcal{C}_{\omega})]$, which implies the inequality.

Since a priori we do not know the value of the orders ω_i , we can use the differential resultant formulas $D_l(\mathfrak{P})$, $l = 1, \ldots, L$ and Algorithm 5.2 to get new upper bounds of ω_i .

Remark 6.8. Assuming \mathfrak{P} to be super esential, to compute $D_l(\mathfrak{P})$. If $\Xi(D_l(\mathfrak{P})) \neq 0$ then, by (15), it belongs to $[\mathfrak{P}] \cap \mathfrak{D}$. Thus, by Lemma 6.1 and construction of $D_l(\mathfrak{P})$

$$\omega_i \leq \operatorname{ord}(\partial \operatorname{Res}(\mathfrak{P}), A_i) \leq \operatorname{ord}(\Xi(D_l(\mathfrak{P})), A_i) \leq J_i - \gamma, \ i = 1, \dots, n,$$

which proves Theorem 6.4 if $\Xi(D_l(\mathfrak{P})) \neq 0$.

In the remaining parts of this section, let us assume that \mathfrak{P} is super essential, to construct $D_l(\mathfrak{P}), l = 1, \ldots, L$. Given $l \in \{1, \ldots, L\}$, let us assume that $D_l(\mathfrak{P}) \neq 0$ and $D_l(\mathfrak{P}) = Q_1 \cdot \ldots \cdot Q_r$ as a product of irreducible factors in $\mathbb{Q}[\mathcal{C}]$. By Proposition 4.1, $D_l(\mathfrak{P}) \in (\mathrm{ags}(\mathfrak{P})) \cap \mathbb{Q}[\mathcal{C}]$, which by Lemma 4.2 has ϵ as a generic zero. Let

$$\mathcal{Q} = \{ Q \in \{Q_1, \dots, Q_r\} \mid Q(\epsilon) = 0 \},\$$

which is nonempty because $(ags(\mathfrak{P})) \cap \mathbb{Q}[\mathcal{C}]$ is prime by Lemma 4.2.

Lemma 6.9. Given $Q \in Q$, there exists a unique $\tau^Q \in \mathbb{N}^n_{-\infty}$ such that $Q \in \mathbb{Q}[\mathcal{C}_{\tau^Q}]$ and $\mathcal{C}_{\tau^Q} \subset \mathcal{C}_{\tau}$, for each $\tau \in \mathbb{N}^n_{-\infty}$ such that $Q \in \mathbb{Q}[\mathcal{C}_{\tau}]$.

Proof. Given $J = (J_1 - \gamma, \ldots, J_n - \gamma) \in \mathbb{N}^n$, $\mathcal{C} = \mathcal{C}_J$. Thus the next set is nonempty

$$\Gamma = \{ \gamma = (\gamma_1, \dots, \gamma_n) \in \mathbb{N}^n_{-\infty} \mid Q \in \mathbb{Q}[\mathcal{C}_{\gamma}] \}.$$

Then $\tau^Q = (\tau_1, \ldots, \tau_n)$, with $\tau_i := \min\{\gamma_i \mid \gamma \in \Gamma\}$.

By Lemma 6.5, $(\mathcal{S}_{\tau^Q}) \cap \mathbb{Q}[\mathcal{C}_{\tau^Q}]$ is a prime ideal with generic zero ϵ_{τ^Q} . The facts that $Q \in \mathbb{Q}[\mathcal{C}_{\tau^Q}]$ and $Q(\epsilon) = 0$, imply that $Q(\epsilon_{\tau^Q}) = Q(\epsilon) = 0$. Hence Q is an irreducible polynomial in $(\mathcal{S}_{\tau^Q}) \cap \mathbb{Q}[\mathcal{C}_{\tau^Q}]$ and if this ideal has codimension one, it holds

$$(\mathcal{S}_{\tau^Q}) \cap \mathbb{Q}[\mathcal{C}_{\tau^Q}] = (\mathsf{R}(\mathcal{S}_{\tau^Q})) = (Q).$$
(26)

Lemma 6.10. Given $Q \in Q$, with $\tau^Q = (\tau_1, \ldots, \tau_n)$ if $\Xi(Q) \neq 0$ then

$$\omega_i \leq \operatorname{ord}(\Xi(Q), A_i) \leq \tau_i \leq J_i - \gamma, \ i = 1, \dots, n.$$

In particular, if $\tau_i = -\infty$ then $\omega_i = -\infty$.

Proof. Observe that $Q \in \mathbb{Q}[\mathcal{C}]$ implies $\tau_i \leq J_i - \gamma$. It is also easy to see that $Q \in (\mathcal{S}_{\tau^Q}) \cap \mathbb{Q}[\mathcal{C}_{\tau^Q}]$ implies $\Xi(Q) \in (\mathrm{PS}(\tau^Q)) \cap \mathbb{Q}[\Xi(\mathcal{C}_{\tau^Q})]$ and if $\Xi(Q) \neq 0$ then $\mathrm{ord}(\Xi(Q), A_i) \leq \tau_i$. The first inequality follows by Lemma 6.1.

We are ready now to prove the main result of this section.

Theorem 6.11. Let \mathfrak{P} be a Laurent differentially essential and super essential system. Let us suppose that there exists $Q \in \mathcal{Q}$ such that $(\mathcal{S}_{\tau^Q}) \cap \mathbb{Q}[\mathcal{C}_{\tau^Q}]$ has codimension one. If $\Xi(Q) \neq 0$ then $\Xi(Q) = \mathfrak{E}\partial \operatorname{Res}(\mathfrak{P})$, with $\mathfrak{E} \in \mathfrak{D}$.

Proof. Let $\tau^Q = (\tau_1, \ldots, \tau_n)$, by Lemma 6.10, $\omega_i \leq \tau_i, i = 1, \ldots, n$ implies

$$(\mathsf{R}(\mathcal{S}_{\omega})) = (\mathcal{S}_{\omega}) \cap \mathbb{Q}[\mathcal{C}_{\omega}] \subseteq (\mathcal{S}_{\tau^{Q}}) \cap \mathbb{Q}[\mathcal{C}_{\tau^{Q}}] = (Q).$$

Since Q is irreducible, $Q = \alpha \mathsf{R}(\mathcal{S}_{\omega}), \alpha \in \mathbb{Q}$. By Proposition 6.7, $\Xi(\mathsf{R}(\mathcal{S}_{\omega})) = \mathfrak{E}_1 \partial \operatorname{Res}(\mathfrak{P})$, for some $\mathfrak{E}_1 \in \mathfrak{D}$. Thus

$$\Xi(Q) = \alpha \Xi(\mathsf{R}(\mathcal{S}_{\omega})) = \alpha \mathfrak{E}_1 \partial \operatorname{Res}(\mathfrak{P}),$$

with $\mathfrak{E} = \alpha \mathfrak{E}_1$ in \mathfrak{D} .

It would be stronger to replace the assumption $\Xi(Q) \neq 0$ by $\Xi(D_l(\mathfrak{P})) \neq 0$.

Corollary 6.12. Let \mathfrak{P} be a Laurent differentially essential and super essential system. Let us suppose that there exists $Q \in \mathcal{Q}$ such that $(\mathcal{S}_{\tau^Q}) \cap \mathbb{Q}[\mathcal{C}_{\tau^Q}]$ has codimension one. If $\Xi(D_l(\mathfrak{P})) \neq 0$ then $\Xi(D_l(\mathfrak{P})) = \mathfrak{E}\partial \operatorname{Res}(\mathfrak{P})$, with $\mathfrak{E} \in \mathfrak{D}$.

Proof. Since $D_l(\mathfrak{P}) = Q' \cdot Q$, with $Q' \in \mathbb{Q}[\mathcal{C}]$, by Theorem 6.11, we get

$$\Xi(D_l(\mathfrak{P})) = \Xi(Q')\Xi(Q) = \Xi(Q')\mathfrak{E}_1\partial\operatorname{Res}(\mathfrak{P})$$

with $\mathfrak{E} = \Xi(Q')\mathfrak{E}_1$ in \mathfrak{D} .

As in Section 4, let $\operatorname{Res}(S)$ be the sparse algebraic resultant of the sparse algebraic generic system $S = \operatorname{ags}(\mathfrak{P})$. Even stronger than the assumption of the previous corollary is to assume that $\operatorname{Res}(S)$ is nontrivial and $\Xi(\operatorname{Res}(S)) \neq 0$.

Corollary 6.13. Let \mathfrak{P} be a Laurent differentially essential and super essential system. Let us suppose that $\operatorname{Res}(S)$ is nontrivial. If $\Xi(\operatorname{Res}(S)) \neq 0$ then $\Xi(\operatorname{Res}(S)) = \mathfrak{E}\partial\operatorname{Res}(\mathfrak{P})$, with $\mathfrak{E} \in \mathfrak{D}$.

Proof. As explained in Section 4, $Q = \text{Res}(S) \in Q$ and

$$\operatorname{Res}(\mathcal{S}) \in (\mathcal{S}_{\tau^Q}) \cap \mathbb{Q}[\mathcal{C}_{\tau^Q}] \subset (\mathcal{S}) \cap \mathbb{Q}[\mathcal{S}] = (\operatorname{Res}(\mathcal{S})).$$

Thus $(\mathcal{S}_{\tau^Q}) \cap \mathbb{Q}[\mathcal{C}_{\tau^Q}]$ has codimension one and by Theorem 6.11 the result follows.

It is natural to wonder what are the conditions on \mathfrak{P} to guarantee $\Xi(D_l(\mathfrak{P})) \neq 0$ (or $\Xi(Q) \neq 0$ for some $Q \in \mathcal{Q}$) but these are not even available so far in the linear case, see [31]. This question is left as a future research direction.

Example 6.14. Let us consider the generic sparse differential system $\mathfrak{P} = \{\mathbb{F}_1 = \mathfrak{a}_1 + \mathfrak{a}_{11}u_1u_2, \mathbb{F}_2 = \mathfrak{a}_2 + \mathfrak{a}_{21}u_1u_{22}, \mathbb{F}_3 = \mathfrak{a}_3 + \mathfrak{a}_{31}u_{21}\}$, which is easily Laurent differentially essential and super essential. The modified Jacobi numbers are $J_1 - \gamma = 1, J_2 - \gamma = 1, J_3 - \gamma = 2$ and

$$\begin{split} \mathrm{ps}(\mathfrak{P}) &= \{ \partial \mathbb{F}_1 = \partial \mathfrak{a}_1 + \partial \mathfrak{a}_{11} u_1 u_2 + \mathfrak{a}_{11} u_{11} u_2 + \mathfrak{a}_{11} u_1 u_{21}, \mathbb{F}_1 = \mathfrak{a}_1 + \mathfrak{a}_{11} u_1 u_2, \\ \partial \mathbb{F}_2 &= \partial \mathfrak{a}_2 + \partial \mathfrak{a}_{21} u_1 u_{22} + \mathfrak{a}_{21} u_{11} u_{22} + \mathfrak{a}_{21} u_1 u_{23}, \mathbb{F}_2 = \mathfrak{a}_2 + \mathfrak{a}_{21} u_1 u_{22}, \\ \partial^2 \mathbb{F}_3 &= \partial^2 \mathfrak{a}_3 + \partial^2 \mathfrak{a}_{31} u_{21} + 2 \partial \mathfrak{a}_{31} u_{22} + \mathfrak{a}_{31} u_{23}, \partial \mathbb{F}_3 = \partial \mathfrak{a}_3 + \partial \mathfrak{a}_{31} u_{21} + \mathfrak{a}_{31} u_{22}, \mathbb{F}_3 = \mathfrak{a}_3 + \mathfrak{a}_{31} u_{21} \}. \end{split}$$

The generic algebraic system associated to \mathfrak{P} is

$$S = \operatorname{ags}(\mathfrak{P}) = \{P_1 = c_{10} + c_{11}y_2y_1 + c_{12}y_2y_3 + c_{13}y_4y_1, P_2 = c_{20} + c_{21}y_2y_1, P_3 = c_{30} + c_{31}y_5y_1 + c_{32}y_5y_3 + c_{33}y_6y_1, P_4 = c_{40} + c_{41}y_5y_1, P_5 = c_{50} + c_{51}y_4 + c_{52}y_5 + c_{53}y_6, P_6 = c_{60} + c_{61}y_4 + c_{62}y_5, P_7 = c_{70} + c_{71}y_4\}.$$

We compute $D_1(\mathfrak{P})$ (using "toricres04", [5], with Maple 15) which has the next irreducible factors

$$Q_1 = c_{62}, Q_2 = c_{40}, Q_3 = -c_{70}c_{62}c_{51} + c_{70}c_{61}c_{52} - c_{60}c_{71}c_{52} + c_{62}c_{50}c_{71}$$

$$Q_4 = -c_{61}c_{70} + c_{71}c_{60}, Q_5 = c_{70}, Q_6 = c_{20}c_{40}c_{12}c_{41}c_{33}c_{71}^2c_{62}c_{50}$$

- $-c_{62}c_{40}c_{70}c_{53}c_{32}c_{13}c_{21} c_{62}c_{40}c_{70}c_{20}c_{51}c_{12}c_{41}c_{33} c_{71}c_{40}c_{20}c_{12}c_{41}c_{60}c_{33}c_{52}$
- $+ c_{71}c_{60}c_{40}c_{53}c_{10}c_{32}c_{41}c_{21} + c_{71}c_{40}c_{20}c_{12}c_{41}c_{60}c_{53}c_{31} c_{71}c_{20}c_{60}c_{30}c_{12}c_{41}^2c_{53}$
- $-c_{71}c_{40}c_{20}c_{32}c_{41}c_{60}c_{53}c_{11} + c_{40}c_{70}c_{20}c_{61}c_{52}c_{12}c_{41}c_{33} c_{40}c_{70}c_{53}c_{10}c_{61}c_{32}c_{41}c_{21}$
- $-c_{40}c_{70}c_{20}c_{12}c_{41}c_{53}c_{31}c_{61}+c_{70}c_{20}c_{30}c_{61}c_{12}c_{41}^2c_{53}+c_{40}c_{70}c_{20}c_{32}c_{41}c_{53}c_{11}c_{61}.$

Only $Q_6(\epsilon) = 0$, thus $\mathcal{Q} = \{Q_6\}$ and its specialization $\Xi(Q_6) \neq 0$,

$$\begin{split} \Xi(Q_6) &= -\mathfrak{a}_{21}(-\mathfrak{a}_{21}\mathfrak{a}_1\mathfrak{a}_{2}\mathfrak{a}_{11}\mathfrak{a}_{31}^2\partial^2\mathfrak{a}_3 + 2\mathfrak{a}_{21}\mathfrak{a}_{31}\mathfrak{a}_2\mathfrak{a}_1\mathfrak{a}_{11}\partial\mathfrak{a}_3\partial\mathfrak{a}_{31} - \mathfrak{a}_{21}\mathfrak{a}_{31}^2\partial\mathfrak{a}_3\mathfrak{a}_2\partial\mathfrak{a}_1\mathfrak{a}_{11} \\ &+ \mathfrak{a}_{21}\mathfrak{a}_{31}\mathfrak{a}_2\mathfrak{a}_3\mathfrak{a}_1\partial^2\mathfrak{a}_{31}\mathfrak{a}_{11} + \mathfrak{a}_{21}\mathfrak{a}_{31}^2\mathfrak{a}_2\mathfrak{a}_1\partial\mathfrak{a}_3\mathfrak{a}_1 - 2\mathfrak{a}_{21}\mathfrak{a}_2\mathfrak{a}_3\mathfrak{a}_1\partial\mathfrak{a}_{31}^2\mathfrak{a}_{11} \\ &+ \mathfrak{a}_{21}\mathfrak{a}_2\mathfrak{a}_3\mathfrak{a}_{31}\partial\mathfrak{a}_1\partial\mathfrak{a}_{31}\mathfrak{a}_{11} + \mathfrak{a}_{21}\mathfrak{a}_{31}^2\mathfrak{a}_2\mathfrak{a}_1\partial\mathfrak{a}_3\partial\mathfrak{a}_2\mathfrak{a}_{11} - \mathfrak{a}_{21}\mathfrak{a}_3\mathfrak{a}_1\partial\mathfrak{a}_2\partial\mathfrak{a}_{31}\mathfrak{a}_{11} \\ &- \mathfrak{a}_{21}\mathfrak{a}_2\mathfrak{a}_3\mathfrak{a}_1\mathfrak{a}_{31}\partial\mathfrak{a}_{11}\partial\mathfrak{a}_{31} + \mathfrak{a}_{31}^2\mathfrak{a}_2^2\mathfrak{a}_3\mathfrak{a}_{11}^2 + \mathfrak{a}_2\mathfrak{a}_3\mathfrak{a}_1\mathfrak{a}_{11}\mathfrak{a}_{31}\partial\mathfrak{a}_{21}\partial\mathfrak{a}_{31} - \mathfrak{a}_{31}^2\mathfrak{a}_2\mathfrak{a}_1\mathfrak{a}_{11}\partial\mathfrak{a}_3\partial\mathfrak{a}_{21}). \end{split}$$

Observe that $\epsilon_2, \ldots, \epsilon_7$ are algebraically independent, since we can choose monomials y_1y_2 , y_1y_5 , y_3y_5 , y_4 , y_5 , y_6 respectively in each of them that are algebraically independent. Therefore $(S) \cap \mathbb{Q}[\mathcal{C}] = (Q_6)$ and $\Xi(Q_6) = \mathfrak{E}\partial \operatorname{Res}(\mathfrak{P})$. We can see that $\Xi(Q_6) = -\mathfrak{a}_{21}H$, with $H(\zeta) = 0$. Thus $H = \partial \operatorname{Res}(\mathfrak{P})$, which illustrates Theorem 6.11 and in particular Corollary 6.12. With a bit more work, we can prove that S is algebraically essential so Q_6 is in fact the sparse algebraic resultant $\operatorname{Res}(S)$. Therefore this example illustrates Corollary 6.13 as well.

Another question for future investigation is, to give conditions on \mathfrak{P} so that $D_l(\mathfrak{P}) \neq 0$. Thus far we assume $D_l(\mathfrak{P}) \neq 0$ and remove the assumption $\Xi(D_l(\mathfrak{P})) \neq 0$ (or $\Xi(Q) \neq 0$), making use of Algorithm 5.2. If $\Xi(D_l(\mathfrak{P})) = 0$, using Algorithm 5.2 with $D_l(\mathfrak{P})$ as an input, by Theorem 5.4, we obtain H_0 in $[\mathfrak{P}] \cap \mathfrak{D}$.

Remark 6.15. Assuming \mathfrak{P} is a Laurent differentially essential and super essential system with $D_l(\mathfrak{P}) \neq 0$, by Lemma 6.1 and construction of $D_l(\mathfrak{P})$ and H_0

$$\omega_i \leq \operatorname{ord}(H_0, A_i) \leq J_i - \gamma, \ i = 1, \dots, n.$$

This proves Theorem 6.4 under the assumption $D_l(\mathfrak{P}) \neq 0$.

Let us assume that $H_0 = H_1 \cdot \ldots \cdot H_s$ as a product of irreducible factors. Since $[\mathfrak{P}] \cap \mathfrak{D}$ is prime with ζ as a generic zero, the next set is nonempty

$$\mathcal{H} := \{ H \in \{ H_1, \dots, H_s \} \mid H(\zeta) = 0 \}.$$

Lemma 6.16. Given $H \in \mathcal{H}$ and $\sigma = (\sigma_1, \ldots, \sigma_n)$, with $\sigma_i := \operatorname{ord}(H, A_i)$, it holds

$$\omega_i \leq \sigma_i \leq J_i - \gamma, \ i = 1, \dots, n.$$

Proof. Since $H(\zeta) = 0$ and ζ is a generic zero of $[\mathfrak{P}] \cap \mathfrak{D}$ then $H \in [\mathfrak{P}] \cap \mathfrak{D}$. By Lemma 6.1 and construction of $D_l(\mathfrak{P})$ and H, the result follows.

Given $H \in \mathcal{H}$ and $\sigma = (\sigma_1, \ldots, \sigma_n)$, with $\sigma_i := \operatorname{ord}(H, A_i)$, by Lemma 6.6 $(\operatorname{PS}(\sigma)) \cap \mathbb{Q}[\Xi(\mathcal{C}_{\sigma})]$ is a prime ideal with ζ_{σ} as a generic zero. We have $H \in \mathbb{Q}[\Xi(\mathcal{C}_{\sigma})]$ and $H(\zeta) = 0$, thus $H(\zeta_{\sigma}) = 0$ and $H \in (\operatorname{PS}(\sigma)) \cap \mathbb{Q}[\Xi(\mathcal{C}_{\sigma})]$. If $\mathcal{I}(H) := (\operatorname{PS}(\sigma)) \cap \mathbb{Q}[\Xi(\mathcal{C}_{\sigma})]$ has codimension one then $\mathcal{I}(H) = (H)$ and, by Lemma 6.16,

$$(\partial \operatorname{Res}(\mathfrak{P})) = (\operatorname{PS}(\omega)) \cap \mathbb{Q}[\Xi(\mathcal{C}_{\omega})] \subseteq (H).$$
(27)

Hence $H = \alpha \partial \text{Res}(\mathfrak{P}), \alpha \in \mathbb{Q}$ because H is irreducible. The previous construction proves the next result.

Theorem 6.17. Let \mathfrak{P} be Laurent differentially essential and super essential system. Let H_0 be the output of Algorithm 5.2 with $D_l(\mathfrak{P}) \neq 0$ as an input. If there exists $H \in \mathcal{H}$ such that $\mathcal{I}(H)$ has codimension one then $H_0 = \mathfrak{E}\partial \operatorname{Res}(\mathfrak{P})$, with $\mathfrak{E} \in \mathfrak{D}$. To finish, we give degree bounds of $\partial \operatorname{Res}(\mathfrak{P})$ in terms of normalized mixed volumes. As mentioned in Section 4, if $\operatorname{Res}(\mathcal{S})$ is nontrivial, for $\mathcal{S} = \operatorname{ags}(\mathfrak{P})$ then $D_l(\mathfrak{P}) \neq 0$ has $\operatorname{Res}(\mathcal{S})$ as an irreducible factor. Furthermore, if \mathcal{S} is algebraically essential (and hence $\operatorname{Res}(\mathcal{S})$ non trivial)

$$\deg(\operatorname{Res}(\mathcal{S}), C_{\lambda(f)}) = MV_{-\lambda(f)}(\mathcal{S}), \ f \in \operatorname{ps}(\mathfrak{P}),$$
(28)

as in (12).

Theorem 6.18. Let \mathfrak{P} be a Laurent differentially essential and super essential system such that $\mathcal{S} = \operatorname{ags}(\mathfrak{P})$ is algebraically essential. If $\Xi(\operatorname{Res}(\mathcal{S})) \neq 0$ then

$$\deg(\partial \operatorname{Res}(\mathfrak{P}), A_i^{[\tau_i]}) \le \deg(\Xi(\operatorname{Res}(\mathcal{S})), A_i^{[\tau_i]}) \le \sum_{k=0}^{\tau_i} \deg(\operatorname{Res}(\mathcal{S}), C_{\lambda(\partial^k f_i)}) = \sum_{k=0}^{\tau_i} MV_{-\lambda(\partial^k f_i)}(\mathcal{S}),$$

with $\tau^{\operatorname{Res}(\mathcal{S})} = (\tau_1, \ldots, \tau_n)$ given by Lemma 6.9.

Proof. By Corollary 6.13, $\Xi(\text{Res}(S)) = \mathfrak{E}\partial \text{Res}(\mathfrak{P})$. The result follows from Lemma 6.10 and (28).

Example 6.19. If $\mathfrak{P} = {\mathbb{G}_1, \ldots, \mathbb{G}_n}$ is a non sparse system, with \mathbb{G}_i a nonhomogeneous generic polynomial of order o_i and degree d_i , it was proven in [21], Theorem 6.18 that $S = \operatorname{ags}(\mathfrak{P})$ is algebraically essential and degree bounds for $\partial \operatorname{Res}(\mathfrak{P})$ in terms of mixed volumes are given in this case.

If $\Xi(\operatorname{Res}(S)) = 0$, we can use $\operatorname{Res}(S)$ as an input of Algorithm 5.2, that returns a nonzero polynomial $H_0 \in [\mathfrak{P}] \cap \mathfrak{D}$. Assuming $H_0 = H_1 \cdot \ldots \cdot H_s$ as a product of irreducible factors, the set $\mathcal{H}(\operatorname{Res}(S)) = \{H \in \{H_1, \ldots, H_s\} \mid H(\zeta) = 0\}$ is nonempty. Given $H \in \mathcal{H}(\operatorname{Res}(S))$, we have $\sigma_i = \operatorname{ord}(H, A_i) \leq \tau_i$, and by construction of H_0

$$\deg(H, A_i^{[\sigma_i]}) \le \deg(H_0, A_i^{[\tau_i]}) \le \sum_{k=0}^{\tau_i} \deg(\operatorname{Res}(\mathcal{S}), C_{\lambda(\partial^k f_i)}).$$
⁽²⁹⁾

Furthermore, if $\mathcal{I}(H)$ has codimension one then $\partial \operatorname{Res}(\mathfrak{P}) = \alpha H$, $\alpha \in \mathbb{Q}$. If \mathcal{S} is algebraically essential then $\operatorname{Res}(\mathcal{S})$ is nontrivial and the next result follows from (29) and (28).

Theorem 6.20. Let \mathfrak{P} be a Laurent differentially essential and super essential system such that $\mathcal{S} = \operatorname{ags}(\mathfrak{P})$ is algebraically essential. If there exists $H \in \mathcal{H}(\partial \operatorname{Res}(\mathfrak{P}))$ such that $\mathcal{I}(H)$ has codimension one then

$$\deg(\partial \operatorname{Res}(\mathfrak{P}), A_i^{[\sigma_i]}) = \deg(H, A_i^{[\sigma_i]}) \le \sum_{k=0}^{\tau_i} \deg(\operatorname{Res}(\mathcal{S}), C_{\lambda(\partial^k f_i)}) = \sum_{k=0}^{\tau_i} MV_{-\lambda(\partial^k f_i)}(\mathcal{S}),$$

with $\tau^{\operatorname{Res}(\mathcal{S})} = (\tau_1, \ldots, \tau_n)$ given by Lemma 6.9 and $\sigma_i = \operatorname{ord}(H, A_i)$.

7 Conclusions

Given a system \mathcal{P} of n Laurent sparse differential polynomials in n-1 differential variables U $(\mathcal{P} \in \mathcal{D}\{U\})$, a method has been designed to compute differential resultant formulas, which provide differential polynomials where the variables U have been eliminated, elements of the differential elimination ideal $[\mathcal{P}] \cap \mathcal{D}$. The steps of this method are:

- 1. Through derivation obtain an extended system $ps(\mathcal{P})$ of \mathcal{P} with L polynomials in L-1 variables.
- 2. Compute determinants of Sylvester matrices $D_l(\mathcal{P})$, l = 1, ..., L of an sparse algebraic generic system \mathcal{S} associated to \mathcal{P} .
- 3. The specialization $\Xi(D_l(\mathcal{P}))$ of $D_l(\mathcal{P})$ to the coefficients of $ps(\mathcal{P})$ is a differential resultant formula for \mathcal{P} and it belongs to the differential elimination ideal $[\mathcal{P}] \cap \mathcal{D}$.

For a generic system \mathfrak{P} , if $\Xi(D_l(\mathfrak{P}))$ is nonzero then it was shown that, under appropriate conditions on S, it is a multiple of the sparse differential resultant $\partial \operatorname{Res}(\mathfrak{P})$ defined by Li, Yuan and Gao in [21]. If $\Xi(D_l(\mathfrak{P})) = 0$, for $D_l(\mathfrak{P}) \neq 0$ an algorithm is given to still obtain an element H of the differential elimination ideal, which is proved to be a multiple of the sparse differential resultant, in the appropriate situation. If the sparse algebraic resultant $\operatorname{Res}(S)$ is nontrivial, its degree can be computed in terms of normalized mixed volumes, we use those to give bounds of the degree of $\partial \operatorname{Res}(\mathfrak{P})$.

It would be interesting to study the appropriate conditions on \mathfrak{P} that guarantee to have nonzero determinats $D_l(\mathfrak{P}), l = 1, \ldots, L$, or furthermore $\Xi(D_l(\mathfrak{P})) \neq 0$. If $\Xi(D_l(\mathfrak{P})) \neq 0$, it still remains to give close formulas to describe $\partial \operatorname{Res}(\mathfrak{P})$ as a quotient of two determinants, as it was done by D'Andrea in the algebraic case, [13]. If $\Xi(D_l(\mathfrak{P})) = 0$, one would need to have more control on the method to obtain the multiple H of $\partial \operatorname{Res}(\mathfrak{P})$, to give closed formulas to compute H and the extraneous factor.

Acknowledgments: The author kindly thanks Alfonso C. Casal and Scott MacCallum for interesting discussions on these topics, and specially Carlos D'Andrea for helpful comments on this manuscript. This work was developed, and partially supported, under the research project MTM2011-25816-C02-01. The author belongs to the Research Group ASYNACS (Ref. CCEE2011/R34).

References

- Berkovich, L.M. and Tsirulik, V.G., 1986. Differential resultants and some of their applications. Differential Equations, Plenum Publ. Corp., 22, 750-757.
- [2] Boulier, F., 2012. The BLAD libraries. http://www.lifl.fr/~boulier/BLAD.
- [3] Boulier, F., Hubert, E., 2004. DIFFALG: description, help pages and examples of use. Now available at http://www-sop.inria.fr/cafe/Evelyne.Hubert/diffalg.
- [4] Boulier, F., Lemaire, F., Maza, M. M., 2010. Computing Differential Characteristic Sets by Change of Ordering, Journal of Symbolic Computation, 45(1), 124-149.
- [5] Canny, J.F. and Emiris, I.Z., 2000. A Subdivision-Based Algorithm for the Sparse Resultant, J. ACM 47,417-451.
- [6] Carra'Ferro, G., 1997. A resultant theory for systems of two ordinary algebraic differential equations. Applicable Algebra in Engineering, Communication and Computing, 8, 539-560.
- [7] Carrà-Ferro, G., 1997. A resultant theory for ordinary algebraic differential equations. Lecture Notes in Computer Science, 1255. Applied Algebra, Algebraic Algorithms and Error-Correcting Codes. Proceedings.
- [8] Chardin, M., 1991. Differential Resultants and Subresultants. Proc. FCT'91, Lecture Notes in Computer Science, 529, Springer-Verlag.
- [9] Cox, D., Little, J., O'Shea, D., 1997. Ideals, Varieties and Algorithms, second ed. Springer, New York.
- [10] Cox, D., Little, J., O'Shea, D., 1998. Using Algebraic Geometry. Springer-Verlag, New York.
- [11] Canny, J., Pedersen, P., 1993. An algorithm for the Newton resultant. Tech. Rep. 1394. Comput. Science Dept., Cornell Univ., Ithaca, N.Y.
- [12] Casal, A.C., Sotomolinos, A.S., 2004. Parametric excitation in a predator-prey model. In the first 60 years of Jean Mawhin, World Scientific, River Edge N.J., 41-54.
- [13] D'Andrea, C., 2002. Macaulay Style Formulas for Sparse Resultants. Trans. of AMS, 354(7), 2595-2629.
- [14] D'Andrea, C. and Emiris, I.Z., 2001. Computing Sparse Projection Operators. Symbolic computation: solving equations in algebra, geometry, and engineering (South Hadley, MA, 2000), 121139, Contemp. Math., 286, Amer. Math. Soc., Providence, RI.

- [15] Gao, X. S., Li, W., Yuan, C. M., 2013. Intersection Theory in Differential Algebraic Geometry: Generic Intersections and the Differential Chow Form. Trans. of AMS, 365(9), 4575-4632.
- [16] Golubitsky, O., Kondratieva, M., Ovchinnikov, A., Szanto, A., 2009. A Bound for Orders in Differential Nullstellensatz. Journal of Algebra, 322, 3852-3877.
- [17] Gelfand, I.M., Kapranov, M., Zelevinsky, A. V., 1994. Discriminants, Resultants and Multidimensional Determinants. Boston, Birkhäuser.
- [18] Hodge, W.V.D. and Pedoe, D., 1968. Methods of Algebraic Geometry, Volume I. Cambridge University Press.
- [19] Hubert, E., 2000. Factorization-free decomposition algorithms in differential algebra. Journal of Symbolic Computation 29, 641-662.
- [20] Kolchin, E.R., 1973. Differential Algebra and Algebraic Groups. Academic Press., London.
- [21] Li, W., Yuan, C.M., Gao, X.S., 2012. Sparse Differential Resultant for Laurent Differential polynomials. In arXiv:1111.1084v3, 1-70.
- [22] Macaulay, F.S., 1916. The Algebraic Theory of Modular Systems. Proc. Cambridge Univ. Press., Cambridge.
- [23] MacCallum, M.A.H., Mikhailov, A., 2009. Algebraic Theory of Differential Equations. London Mathematical Society Lecture Note Series (Book 357). Cambridge University Press.
- [24] Mishra, B., 1993. Algorithmic algebra. Springer.
- [25] Pedersen, P., Sturmfels, B., 1993. Product formulas for resultants and Chorw forms. Math. Z., 214, 337-396.
- [26] Ritt, J.F., 1950. Differential Algebra. Amer. Math. Sco. Colloquium, New York.
- [27] Rosenkranz, M., Wang, D., 2007. Gröbner Bases in Symbolic Analysis. Radon Series on Computational and Applied Mathematics (Book 2). Walter de Gruyter.
- [28] Rueda, S.L., 2011. On the computation of differential resultants. Modern Mathematical Tools and Techniques in Capturing Complexity. Understanding Complex Systems, Springer, 91-103.
- [29] Rueda, S.L. and Sendra, J.F., 2010. Linear complete differential resultants and the implicitization of linear DPPEs. Journal of Symbolic Computation, 45, 324-341.
- [30] Rueda, S.L., 2011. A perturbed differential resultant based implicitization algorithm for linear DPPEs. Journal of Symbolic Computation, 46, 977-996.
- [31] Rueda, S.L., 2013. Linear sparse differential resultant formulas. Linear Algebra and Its Applications 438, 4296 - 4321.
- [32] Sturmfels, B., 1994. On The Newton Polytope of the Resultant. Journal of Algebraic Combinatorics, 3, 207-236.
- [33] Zhang, Z.Y., Yuan, C.M., Gao, X.S., 2012. Matrix Formula of Differential Resultant for First Order Generic Ordinary Differential Polynomials. In arXiv: 1204.3773v1, 1-22.
- [34] Wang, D.M., Zheng, Z., 2005. Differential equations with symbolic computation. Birkhuser Basel.