



Simple Pricing Schemes for the Cloud

IAN A. KASH, University of Illinois at Chicago

PETER KEY, Cambridge

WARUT SUKSOMPONG, University of Oxford

The problem of pricing the cloud has attracted much recent attention due to the widespread use of cloud computing and cloud services. From a theoretical perspective, several mechanisms that provide strong efficiency or fairness guarantees and desirable incentive properties have been designed. However, these mechanisms often rely on a rigid model, with several parameters needing to be precisely known for the guarantees to hold. In this article, we consider a stochastic model and show that it is possible to obtain good welfare and revenue guarantees with simple mechanisms that do not make use of the information on some of these parameters. In particular, we prove that a mechanism that sets the same price per timestep for jobs of any length achieves at least 50% of the welfare and revenue obtained by a mechanism that can set different prices for jobs of different lengths, and the ratio can be improved if we have more specific knowledge of some parameters. Similarly, a mechanism that sets the same price for all servers even though the servers may receive different kinds of jobs can provide a reasonable welfare and revenue approximation compared to a mechanism that is allowed to set different prices for different servers.

CCS Concepts: • **Theory of computation** → **Computational pricing and auctions**; *Algorithmic mechanism design*; • **Software and its engineering** → **Cloud computing**;

Additional Key Words and Phrases: Cloud computing, pricing, simple auctions, mechanism design

ACM Reference format:

Ian A. Kash, Peter Key, and Warut Suksompong. 2019. Simple Pricing Schemes for the Cloud. *ACM Trans. Econ. Comput.* 7, 2, Article 7 (June 2019), 27 pages.
<https://doi.org/10.1145/3327973>

1 INTRODUCTION

With cloud computing generating billions of dollars per year and forming a significant portion of the revenue of large software companies (Columbus 2016), the problem of how to price cloud resources and services is of great importance. On the one hand, for a pricing scheme to be used, it is necessary that the scheme provide strong welfare and revenue guarantees. On the other hand,

The project was started at Microsoft Research Cambridge, UK. Preliminary versions of the article appeared in *Proceedings of the 13th Conference on Web and Internet Economics*, December 2017, and *Proceedings of the 12th Workshop on the Economics of Networks, Systems, and Computation*, June 2017.

This work was partially supported by NSF Award CCF-1813188, by the European Research Council (ERC) under grant number 639945 (ACCORD), and by a Stanford Graduate Fellowship.

Authors' addresses: I. A. Kash, 851 S. Morgan (M/C 152), Chicago, IL 60607, USA; email: iankash@uic.edu; P. Key, 44 Main Street, Hardwick, Cambridge CB23 7QS, UK; email: peter.key@outlook.com; W. Suksompong, Wolfson Building, Parks Road, Oxford OX1 3QD, UK; email: warut.suksompong@cs.ox.ac.uk.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than the author(s) must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

© 2019 Copyright held by the owner/author(s). Publication rights licensed to ACM.

2167-8375/2019/06-ART7 \$15.00

<https://doi.org/10.1145/3327973>

Microsoft Azure

SALES 0800 090 8435 MY ACCOUNT PORTAL Search

Why Azure Solutions Products Documentation Pricing Partners Blog Resources Support FREE ACCOUNT

+ Add items Clear all

VPN Gateway SQL Database App Service

Virtual Machines

REGION: Australia East TYPE: Red Hat Enterprise Linux

PRICING TIER: Standard LICENSE: On Demand

INSTANCE SIZE: D1 SSD 1 cores 3.5 GB RAM 50 GB disk \$0.153/hr

NOTE: The displayed price includes the Virtual Machine and the license prices.

1 Virtual Machines × 100 Hours = \$15.30/MO

Your estimate

US Dollar (\$)

Virtual Machin... \$15.30

Support Options \$0.00

\$15.30
Estimated monthly cost

Purchase options

Export estimate

Prices are estimates and are not intended as actual price quotes.

Fig. 1. Pricing scheme for virtual machines on Microsoft Azure (Azure 2016).

it is also often desirable that the scheme be simple. We combine the two objectives in this article and show that simple pricing schemes perform almost as well as more complex ones with respect to welfare and revenue guarantees. In particular, consider the pricing scheme for virtual machines on Microsoft Azure shown in Figure 1. Once the user chooses the basic parameters such as region, type, and instance size, the price is calculated by simply multiplying an hourly base price by the number of virtual machines and number of hours desired. The question that we study can be phrased in this setting as follows: How much more welfare or revenue could be created if instead of this simple multiplication formula, a complex table specifying the price for each number of hours were to be used? Our main result is that the former offers at worst a two approximation to the latter, both in terms of welfare and revenue. Similarly, we demonstrate that setting a single price for a group of servers, even though the servers may receive different kinds of jobs, can provide a reasonable welfare and revenue approximation compared to setting different prices for different servers.

In much of the prior work in this space, which focuses more explicitly on scheduling, prices depend in a complex way on a number of parameters (typically including job length, arrival time, deadline, and value) and the current state of the system (Azar et al. 2015; Dehghani et al. 2016; Jain et al. 2011, 2012; Lucier et al. 2013). A weakness of such schemes is that they require these parameters to be known up front for the desirable properties of the mechanisms, such as their approximation ratios, to hold. The availability of such information is not always realistic in practice. Even when it is in principle possible to provide this information, there is a cost to participants in both time and resources to figure it out. In this work, we show that good results are possible with no up-front information.

For our initial results, we assume that there is a single server that receives jobs of various lengths whose value per timestep is drawn from the same probability distribution regardless of length. We compare the welfare and revenue that can be obtained by setting a price per timestep that is

Spot Instances	Defined Duration for Linux	Defined Duration for Windows
Region: US East (Ohio)		
	1 hour	6 hours
General Purpose - Current Generation		
m4.large	\$0.142 per Hour	\$0.157 per Hour
m4.xlarge	\$0.284 per Hour	\$0.314 per Hour
m4.2xlarge	\$0.568 per Hour	\$0.628 per Hour
m4.4xlarge	\$1.136 per Hour	\$1.256 per Hour
m4.10xlarge	\$2.84 per Hour	\$3.14 per Hour
m4.16xlarge	\$4.544 per Hour	\$5.024 per Hour

Fig. 2. Pricing scheme for defined duration spot instances on Amazon (Amazon 2017).

independent of the job length against the corresponding objective obtained by setting an individual price for each job length. When we are allowed the freedom of setting different prices for different job lengths, intuitively we want to set a higher price per timestep for longer jobs as a premium for reserving the server for a longer period of time.¹ However, as we show, we do not lose more than 50% of the welfare or revenue (or any convex combination of the two) if we are only allowed to set one price. We would like to emphasize that this is a worst-case bound over a wide range of parameters, including the number of job lengths, the distribution over job lengths, and the distribution over job values. Indeed, as we show, we can obtain improved bounds if we know the value of some of these parameters. The price that we use in the single-price setting can be chosen from one of the prices used in the multi-price setting, meaning that we do not have to calculate a price from scratch. Moreover, all of our approximation guarantees hold generally for arbitrary prices, meaning that for any prices that we may set in a multi-price setting (i.e., not necessarily optimal ones), we can obtain an approximation of the welfare or revenue by setting one of those prices alone. Finally, we emphasize that these results put no restrictions on the form of the distribution; it can be discrete, continuous, or mixed. The only substantive constraint is that jobs of all lengths share the same distribution of value per timestep. However, in an extension, we show that a version of our results continues to hold even if this constraint is relaxed.

We then generalize our results to a setting where there are multiple servers, each of which receives jobs of various lengths. The distribution over job lengths can be different for different servers. This is conceivable, for instance, if the servers are in various geographic locations or are utilized by various groups of users. We compare the welfare and revenue obtained by a simple pricing scheme that sets the same price for all servers against the corresponding objective achieved by a scheme that can set a different (single) price for each server. Roughly speaking, we show that as long as the parameters are not too extreme (e.g., the number of servers or the job lengths are not too large), then we do not lose too much of the welfare or revenue by setting a single price. Combining this with our initial results, we obtain an approximation of a very restricted

¹Amazon recently started offering a product called *defined duration spot instances*, where users can specify a duration in hourly increments up to 6 hours. Indeed, the price *per hour* of this product increases as the number of hours increases. (See Figure 2).

pricing scheme where we must set the same price for all servers and all job lengths against one where we can set an individual price for each job length of each server. These results require an assumption that all servers have the same probability of not receiving a job at a timestep. Using similar techniques, we also obtain approximation bounds when this assumption does not hold but there is only one job length across all servers.

1.1 Related Work

Much recent work has focused on designing online scheduling mechanisms with good welfare guarantees and incentive properties. Jain et al. (2011) exhibited a truthful mechanism for batch jobs on cloud systems where jobs are allocated non-preemptively, and the same group of authors came up with mechanisms for deadline-sensitive jobs in large computing clusters (Jain et al. 2012). Lucier et al. (2013) also considered the problem of scheduling deadline-sensitive jobs; they circumvented known lower bounds by assuming that jobs could be delayed and still finish by their deadline. Zhang et al. (2013) developed a framework for truthful online cloud auctions where users with heterogeneous demands can come and leave on the fly. More recently, Azar et al. (2015) constructed a truthful mechanism that achieves a constant competitive ratio given that slackness is allowed, whereas Dehghani et al. (2016) assumed a stochastic model and developed a truthful mechanism that approximates the expected maximum welfare up to a constant factor. Wang et al. (2015) designed mechanisms for selling reserved instances where users are allowed to reserve resources of any length and from any time point in the future. Their mechanisms determine the acceptance and payment immediately when the job arrives, and achieve a competitive ratio that is optimal to within a constant factor with regard to welfare.

Other work in this space has dealt with comparing pricing mechanisms such as the on-demand market and the spot market (Abhishek et al. 2012; Dierks and Seuken 2016; Hoy et al. 2016), achieving fairness in job allocation (Friedman et al. 2014), and studying models of real-time pricing with budget constraints (Friedman et al. 2015). Kash and Key (2016) gave a survey of the current state of research in economics and computer science with respect to cloud pricing.

From a technical perspective, our work bears a resemblance to the work of Dütting et al. (2018) on discriminatory and anonymous posted pricing and of Disser et al. (2016) on hiring secretaries. In particular, Dütting et al. (2018) considered the problem of selling homogeneous items to buyers who arrive sequentially with values drawn independently from identical distributions. They showed that by posting discriminatory prices, one can obtain at most $2 - k/n$ times as much revenue as that obtained by posting the same anonymous price, where n is the number of buyers and k the number of items. As is the case in our work, their anonymous price can always be chosen from one of the discriminatory prices, and their proof of tightness is also based on a discrete bimodal distribution. Interestingly, their proof of the $2 - k/n$ bound relies on first comparing the revenue from discriminatory prices to that from a lottery over two anonymous prices, then estimating this latter revenue in relation to setting a single anonymous price. Unlike in our work, however, their techniques only yield guarantees for revenue and not welfare. Disser et al. (2016) provided a competitive online algorithm for a variant of the stochastic secretary problem, where applicants need to be hired over time. Upon the arrival of each applicant, the cost per timestep of the applicant is revealed, and we have to decide on the duration of the employment, starting immediately. Once an applicant is accepted, we cannot terminate the contract until the duration of the job is over.

Our work falls into the broader area of the design and analysis of simple mechanisms, particularly posted price mechanisms. One of the motivations for studying simple mechanisms is that in practice, designers are often willing to partially give up optimality in return for simplicity. Mechanisms that simply post prices on goods have received significant attention because they reflect perhaps the most common way of selling goods in the real world, and moreover they leave no room

for strategizing, making them easy for agents to participate in. A long line of work has investigated how well such mechanisms can approximate optimal mechanisms with respect to various objectives, including welfare (Cohen-Addad et al. 2016; Ezra et al. 2018; Feldman et al. 2015), revenue (Babaioff et al. 2011; Blumrosen and Holenstein 2008; Chawla et al. 2010), and social costs (Cohen et al. 2015). In Section 3.4, we show that techniques from this literature can recover some of our results under relaxed assumptions.

2 PRELIMINARIES

We consider a system with a number of servers and discrete timesteps. Each job takes an integer number of timesteps to complete and yields a value upon completion. The value *per timestep* of a job is drawn from a distribution that is independent of the length of the job. Let F be the cumulative distribution function of this distribution and f the probability density function with respect to a base measure μ , and define $\ell(x) = xf(x)$.² We do not make any assumption on our distribution; in particular, it need not be continuous or discrete, which is why we allow flexibility in terms of the base measure.

When a job request is made for a job to be served by a server, there is a price p per timestep that may depend on the job length and/or the server. If the value per timestep of the job is at least p , the server accepts and executes the job to completion. Otherwise, the server rejects the job. The objectives in our model are the steady-state welfare and revenue for each pricing scheme. In particular, we will be interested in the expected welfare and revenue per timestep, given that the job values are drawn from a probability distribution. This can also be thought of as the average welfare and revenue per timestep that result from a pricing scheme over a long period of time.

In Section 3, we assume that there is a single server. Each timestep, either zero or one job appears. A job with length a_i appears with probability $0 < r_i \leq 1$, where $\sum_{i=1}^n r_i \leq 1$ and n denotes the number of job lengths. We are allowed to set a price p_i for jobs of length a_i . If a server accepts a job of length a_i , it is busy and cannot accept other jobs for a_i timesteps, including the current one. We compare the setting where we are forced to set the same price per timestep p for all job lengths against the setting where we can set a different price per timestep p_i for each job length a_i . Note that if we could set different prices for different job lengths, then to optimize welfare or revenue, intuitively we would set a higher price per timestep for longer jobs as a premium for reserving the server for a longer period. Put differently, once we accept a longer job, we are stuck with it for a longer period, during which we miss the opportunity to accept other jobs. Consequently, we should set a higher standard for accepting longer jobs. (See also Footnote 1.)

In Section 4, we assume that there are multiple servers. Each timestep, either zero or one job appears for each server $1 \leq j \leq n$. For server j , a job with length a_{ji} appears with probability $0 < r_{ji} \leq 1$ for $1 \leq i \leq n_j$, where n_j denotes the number of job lengths for server j . We do not assume that the set of job lengths or the number of job lengths is identical across servers. However, we assume that the probability of no job appearing at a timestep is the same for all servers (i.e., $\sum_{i=1}^{n_j} r_{ji}$ is constant for any j). In Section 4.1, we assume that we can set one price per server, and we compare the setting where we are forced to set the same price per timestep p for all servers against that where we can set a different price per timestep p_j for each server j . In Section 4.2, we assume that we can set a different price per timestep p_{ji} for each server j and each of its job lengths a_{ji} , and we compare that setting against the one where we are forced to set the same price per timestep p for all servers and all job lengths.

²For technical reasons, we will deviate slightly from the usual notion of cumulative distribution function. In particular, if y is a random variable drawn from a distribution, then we define its cumulative distribution function $F(x)$ as $\Pr[y < x]$ instead of the usual $\Pr[y \leq x]$. This will only be important when we deal with discrete distributions.

3 ONE SERVER

In this section, we assume that there is a single server, which receives jobs of various lengths. After presenting an introductory example in Section 3.1, we consider the general setting with an arbitrary number of job lengths in Section 3.2. In this setting, we show a 50% approximation for both welfare and revenue (or any convex combination of the two) of setting one price for all job lengths compared to setting an individual price for each job length, for any realization of the parameters. Moreover, we show in Section 3.3 that our techniques provide a template for deriving tighter bounds if we have more specific information on the parameters. In particular, when there are two job lengths, we show for each setting of the parameters a tight approximation bound for welfare and revenue. Our approximation results hold for arbitrary (i.e., not necessarily optimal) pricing schemes, and the price we use in the single-price setting can be drawn from one of the prices in the multi-price setting. Finally, in Section 3.4, we consider an extension that does not assume independence between the job length and the value per timestep.

3.1 Warm-Up: Uniform Distribution

As a warm-up example, assume that at any timestep a job with length 1 or 2 appears with probability 50% each. The value per timestep of a job is drawn from the uniform distribution over $[0, 1]$. Suppose that we set a price per timestep p_1 for jobs of length 1 and p_2 for jobs of length 2.

Consider an arbitrary timestep when the server is free. If the job drawn at that timestep has length 1, then with probability p_1 it has value below p_1 and is rejected. In this case, the server passes one timestep without a job. Otherwise, the job has value at least p_1 and is accepted. In this case, the expected welfare from executing the job is $\frac{1+p_1}{2}$. Similarly, if the job has length 2, then with probability p_2 it is rejected, and with probability $1 - p_2$ it is accepted and yields expected welfare $2 \cdot \frac{1+p_2}{2} = 1 + p_2$ over two timesteps. Letting c_w denote the expected welfare per timestep assuming that the server is free at the current timestep, we have

$$0 = \frac{1}{2} \left(-p_1 c_w + (1 - p_1) \left(\frac{1 + p_1}{2} - c_w \right) \right) + \frac{1}{2} \left(-p_2 c_w + (1 - p_2) (1 + p_2 - 2c_w) \right).$$

The two terms on the right-hand side correspond to jobs of length 1 and 2, which are drawn with probability 1/2 each. In the case that a job of length 2 is drawn, with probability p_2 it is rejected and the server is idle for one timestep, during which it would otherwise have produced expected welfare c_w . With the remaining probability $1 - p_2$ the job is accepted, yielding expected welfare $1 + p_2$ over two timesteps, during which the server would otherwise have produced expected welfare $2c_w$. The derivation for the term corresponding to jobs of length 1 is similar. By equating the expected welfare with the variable denoting this quantity, we arrive at the preceding equation.

Solving for c_w , we get

$$c_w(p_1, p_2) = \frac{\frac{(1-p_1)(1+p_1)}{2} + (1-p_2)(1+p_2)}{3 - p_2}.$$

To maximize $c_w(p_1, p_2)$ over all values of p_1, p_2 , we should set $p_1 = 0$. (Indeed, to maximize welfare, we should always accept jobs of length 1 because they do not interfere with future jobs.) Then the value of p_2 that maximizes $c_w(p_1, p_2)$ is $p_2 = 3 - \sqrt{\frac{15}{2}} \approx 0.261$, yielding $c_w(p_1, p_2) = 6 - \sqrt{30} \approx 0.522$.

However, if we set the same price $p = p_1 = p_2$ for jobs with different lengths, our welfare per timestep becomes

$$c_w(p) = \frac{\frac{(1-p)(1+p)}{2} + (1-p)(1+p)}{3 - p} = \frac{3(1-p)(1+p)}{2(3 - p)}.$$

This is maximized at $p = 3 - 2\sqrt{2} \approx 0.172$, yielding $c_w(p) = 9 - 6\sqrt{2} \approx 0.515$. Moreover, if we use either of the prices in the optimal price combination for the two-price setting as the single price, we get $c_w(0) = 0.5$ and $c_w(3 - \sqrt{\frac{15}{2}}) \approx 0.510$.

Next, we repeat the same exercise for revenue. We can derive the equations in the same way, with the only difference being that the revenue from accepting a job at price p is simply p . Letting c_r denote the revenue per timestep, we have

$$0 = \frac{1}{2} (-p_1 c_r + (1 - p_1)(p_1 - c_r)) + \frac{1}{2} (-p_2 c_r + (1 - p_2)(2p_2 - 2c_r)).$$

Solving for c_r , we get

$$c_r(p_1, p_2) = \frac{(1 - p_1)p_1 + 2(1 - p_2)p_2}{3 - p_2}.$$

To maximize c_r over all values of p_1, p_2 , we should set $p_1 = 0.5$. (Indeed, to maximize revenue, we should always set the monopoly price for jobs of length 1 because they do not interfere with future jobs.) Then the value of p_2 that maximizes $c_r(p_1, p_2)$ is $p_2 = 3 - \sqrt{\frac{47}{8}} \approx 0.576$, yielding $c_r(p_1, p_2) = 10 - \sqrt{94} \approx 0.304$.

However, if we set the same price $p = p_1 = p_2$ for jobs with different lengths, our revenue per timestep becomes

$$c_r(p) = \frac{(1 - p)p + 2(1 - p)p}{3 - p} = \frac{3(1 - p)p}{3 - p}.$$

This is maximized at $p = 3 - \sqrt{6} \approx 0.551$, yielding $c_r(p) = 15 - 6\sqrt{6} \approx 0.303$. Moreover, if we use either of the prices in the optimal price combination for the two-price setting as the single price, we get $c_r(0.5) = 0.3$ and $c_r(3 - \sqrt{\frac{47}{8}}) \approx 0.302$.

Observe that for both welfare and revenue, the maximum in the one-price setting is not far from that in the two-price setting. In addition, in both cases, at least one of the two prices in the optimal price combination for the two-price setting, when used alone as a single price, performs almost as well as the maximum in the two-price setting. In the remainder of Section 3, we will show that this is not a coincidence but rather a phenomenon that occurs for any set of job lengths, any probability distribution over job lengths, and any probability distribution over job values.

3.2 General 50% Approximation

In this section, we consider a general setting with an arbitrary number of job lengths. We show that even at this level of generality, it is always possible to obtain 50% of the welfare and revenue of setting an individual price for each job length by setting just one price. Although the optimal price in the one-price setting might be different from any of the prices in the multi-price setting, we show that at least one of the prices in the latter setting can be used alone to achieve the 50% guarantee.

Assume that there are jobs of lengths $a_1 \leq a_2 \leq \dots \leq a_n$ that appear at each timestep with probability r_1, r_2, \dots, r_n , respectively. Suppose that we set a price per timestep p_i for jobs of length a_i . Recall that the value per timestep of a job is drawn from a distribution with cumulative distribution function F and probability density function f .

The following lemma gives the formulas for the expected welfare and revenue per timestep.

LEMMA 3.1. Let $S = a_1r_1 + \dots + a_nr_n$ and $R = r_1 + \dots + r_n$, and let c_w and c_r denote the expected welfare and revenue per timestep, respectively. We have

$$c_w(p_1, p_2, \dots, p_n) = \frac{a_1r_1 \int_{x \geq p_1} \ell d\mu + \dots + a_nr_n \int_{x \geq p_n} \ell d\mu}{S - ((a_1 - 1)r_1F(p_1) + \dots + (a_n - 1)r_nF(p_n)) + (1 - R)} \quad (1)$$

and

$$c_r(p_1, p_2, \dots, p_n) = \frac{a_1r_1(1 - F(p_1))p_1 + \dots + a_nr_n(1 - F(p_n))p_n}{S - ((a_1 - 1)r_1F(p_1) + \dots + (a_n - 1)r_nF(p_n)) + (1 - R)}. \quad (2)$$

In particular, if $p_1 = \dots = p_n = p$, then

$$c_w(p) = \frac{S \int_{x \geq p} \ell d\mu}{S - (S - R)F(p) + (1 - R)}$$

and

$$c_r(p) = \frac{S(1 - F(p))p}{S - (S - R)F(p) + (1 - R)}.$$

PROOF. We represent the states of the server by a Markov chain. The states are categorized into types T_0, T_1, \dots, T_n . Type T_0 consists of only one state, which is an idle state corresponding to when the server is free. For $1 \leq i \leq n$, type T_i consists of $a_i - 1$ states. When the server accepts a job of length a_i , it deterministically steps through each of the $a_i - 1$ states of type T_i before returning to the idle state. (If $a_i = 1$, there is no state of type T_i ; when the server accepts a job of length a_i , it simply transitions from the idle state back to itself.) The rewards, which can be either welfare or revenue, are collected at each transition. If the threshold for accepting a job of a certain length is p , then the revenue collected for each transition with that job length is always p , whereas the expected welfare gained during the transition is given by $\frac{\int_{x \geq p} \ell d\mu}{1 - F(p)}$.

By virtue of local balance, for each type T_i with $a_i \neq 1$, the proportion of time spent in each of the states of that type is the same for all $a_i - 1$ states. Denote this proportion by q_i . From the idle state, the probability of transitioning to the first state of type T_i is $r_i(1 - F(p_i))$, so $q_i = r_i(1 - F(p_i))q_0$. Since the probabilities across all states sum to 1, we have

$$q_0 + \sum_{i=1}^n (a_i - 1)r_i(1 - F(p_i))q_0 = 1.$$

This yields

$$q_0 = \frac{1}{1 + \sum_{i=1}^n (a_i - 1)r_i(1 - F(p_i))} = \frac{1}{S - \sum_{i=1}^n (a_i - 1)r_iF(p_i) + (1 - R)}.$$

We are now in a position to compute the expected welfare and revenue per timestep. For states of type T_i , the expected revenue collected in the transitions through these states is $q_i p_i a_i = q_0 r_i (1 - F(p_i)) p_i a_i$. The expected revenue per timestep is therefore

$$\sum_{i=1}^n q_0 r_i (1 - F(p_i)) p_i a_i = \frac{\sum_{i=1}^n a_i r_i (1 - F(p_i)) p_i}{S - \sum_{i=1}^n (a_i - 1) r_i F(p_i) + (1 - R)}.$$

Similarly, for states of type T_i , the expected revenue collected in the transitions through these states is $q_i \cdot \frac{\int_{x \geq p_i} \ell d\mu}{1 - F(p_i)} \cdot a_i = q_0 r_i (1 - F(p_i)) \cdot \frac{\int_{x \geq p_i} \ell d\mu}{1 - F(p_i)} \cdot a_i$. The expected welfare per timestep is therefore

$$\sum_{i=1}^n q_0 r_i (1 - F(p_i)) \cdot \frac{\int_{x \geq p_i} \ell d\mu}{1 - F(p_i)} \cdot a_i = \frac{\sum_{i=1}^n a_i r_i \int_{x \geq p_i} \ell d\mu}{S - \sum_{i=1}^n (a_i - 1) r_i F(p_i) + (1 - R)},$$

as claimed. \square

With the formulas for welfare and revenue in hand, we are ready to show the main result of this section, which exhibits that the worst-case approximation ratio for welfare or revenue between the single-price setting and the multi-price setting is at least 50%. As we will see later in Section 3.3, this bound is in fact tight, and it remains tight even when there are only two job lengths. Note that the bound holds for any number of job lengths, any distribution over job lengths, and any distribution over job values.

THEOREM 3.2. *For any prices p_1, p_2, \dots, p_n that we set in the multi-price setting, there exists a price p_i such that we can achieve a welfare (respectively, revenue, or any convex combination of welfare and revenue) approximation of at least 50% in the one-price setting by using p_i as the single price.*

To prove Theorem 3.2, we work with the ratio

$$\frac{\max(c_w(p_1), \dots, c_w(p_n))}{c_w(p_1, \dots, p_n)}$$

and show that it is at least $1/2$ for any p_1, \dots, p_n (and similarly for revenue or any convex combination of welfare and revenue). Using the formula (1) for c_w given in Lemma 3.1, we can write the ratio in terms of the variables $A_i = \frac{\int_{x \geq p_i} \ell d\mu}{\int_{x \geq p_1} \ell d\mu}$ and $B_i = F(p_i)$ for $1 \leq i \leq n$. For any fixed values of B_i , we then deduce the values of A_i that minimize the ratio of interest. Finally, we show that the remaining expression is always at least $1/2$ no matter the values of B_i .

PROOF. We first consider welfare. We wish to show that setting one of the prices p_i alone achieves an approximation of $1/2$ of setting all n prices. In other words,

$$\max(c_w(p_1), \dots, c_w(p_n)) \geq \frac{1}{2} \cdot c_w(p_1, \dots, p_n).$$

To establish this inequality, we will work with the ratio

$$\frac{\max(c_w(p_1), \dots, c_w(p_n))}{c_w(p_1, \dots, p_n)}$$

and show that its minimum is at least $1/2$.

Writing $A_i = \frac{\int_{x \geq p_i} \ell d\mu}{\int_{x \geq p_1} \ell d\mu}$ (in particular, $A_1 = 1$) and $B_i = F(p_i)$ for $1 \leq i \leq n$, the ratio to minimize becomes

$$g(A_1, \dots, A_n, B_1, \dots, B_n) := \max_{i=1}^n \left(\frac{SA_i}{S - (S - R)B_i + (1 - R)} \right) \cdot \frac{S - ((a_1 - 1)r_1B_1 + \dots + (a_n - 1)r_nB_n) + (1 - R)}{a_1r_1A_1 + \dots + a_nr_nA_n}.$$

By dividing into cases according to the term that the max function outputs, we can deduce that we only need to minimize the function

$$h(B_1, \dots, B_n) := \frac{S^2 - ((a_1 - 1)r_1B_1 + \dots + (a_n - 1)r_nB_n)S + S(1 - R)}{S^2 - (a_1r_1B_1 + \dots + a_nr_nB_n)(S - R) + S(1 - R)};$$

the details of this casework can be found in Appendix A. In particular, we want to show that $h(B_1, \dots, B_n) \geq \frac{1}{2}$ for any choice of B_1, \dots, B_n . This is equivalent to

$$S^2 + S(1 - R) \geq (2S(a_1 - 1) - a_1(S - R))r_1B_1 + \dots + (2S(a_n - 1) - a_n(S - R))r_nB_n$$

or

$$S^2 + S(1 - R) \geq ((a_1 - 2)S + a_1R)r_1B_1 + \dots + ((a_n - 2)S + a_nR)r_nB_n.$$

We consider two cases:

- (1) $a_1 \geq 2$ (and hence $a_2, \dots, a_n \geq 2$). All coefficients of $r_i B_i$ on the right-hand side are positive, so we only need to verify the inequality for $B_1 = \dots = B_n = 1$. We have

$$\begin{aligned} h(B_1 = 1, \dots, B_n = 1) &= \frac{S^2 - ((a_1 - 1)r_1 + \dots + (a_n - 1)r_n)S + S(1 - R)}{S^2 - (a_1 r_1 + \dots + a_n r_n)(S - R) + S(1 - R)} \\ &= \frac{S^2 - (S - R)S + S(1 - R)}{S^2 - S(S - R) + S(1 - R)} = 1 > \frac{1}{2}. \end{aligned}$$

- (2) $a_1 = 1$.³ All coefficients of $r_i B_i$ on the right-hand side except the first one are positive, so we only need to verify the inequality for $B_1 = 0$ and $B_2 = \dots = B_n = 1$. We have

$$h(B_1 = 0, B_2 = 1, \dots, B_n = 1) = \frac{S}{S + r_1(S - R)} > \frac{1}{2},$$

where the last line follows from $r_1(S - R) < 1 \cdot S = S$.

Thus, the inequality holds in both cases, and the approximation ratio is at least $\frac{1}{2}$, as claimed.

Finally, we can obtain analogous results for revenue by essentially repeating the same argument but instead writing $A_i = \frac{(1-F(p_i))p_i}{(1-F(p_1))p_1}$ for $1 \leq i \leq n$, and for any convex combination of welfare and revenue by writing A_i as the appropriate convex combination of the two corresponding terms. \square

3.3 Tighter Bounds for Specific Parameters

Assume in this section that there are jobs of two lengths $a < b$ that appear at each timestep with probability r_1 and r_2 , respectively, where $r_1 + r_2 \leq 1$. Suppose that we set a price per timestep p_1 for jobs of length a and p_2 for jobs of length b . Recall that the value per timestep of a job is drawn from a distribution with cumulative distribution function F and probability density function f .

Our next result exhibits a tight approximation bound for any fixed setting of the job lengths and their distribution.

THEOREM 3.3. *For any prices p_1 and p_2 that we set in the two-price setting, we can achieve a welfare (respectively, revenue, or any convex combination of welfare and revenue) approximation of at least*

$$\rho(a, b, r_1, r_2) := \frac{(ar_1 + br_2)(ar_1 + 1 - r_1)}{a(a - 1)r_1^2 + a(b - 1)r_1 r_2 + ar_1 + br_2}$$

in the one-price setting by setting either p_1 or p_2 alone. Moreover, this bound is the best possible even if we are allowed to set a price different from p_1 or p_2 in the one-price setting.

Note that although the approximation ratios for welfare and revenue can be different for a particular distribution (see Section 3.1), Theorem 3.3 shows that the worst-case approximation over all distributions is the same for both benchmarks.

To prove this theorem, we work with the expression in terms of $B_i = F(p_i)$ that we have from the proof of Theorem 3.2. We then show that the expression is minimized when we take $B_1 = 0$ and $B_2 = 1$, meaning that the distribution on job values is bimodal. The proof method readily yields an example showing that our bound is tight, where the bimodal distribution on job values puts a large probability on a low value and a small probability on a high value.

PROOF. Using the proof of Theorem 3.2, we are left with minimizing the function

$$h(B_1, B_2) := \frac{S^2 - ((a - 1)r_1 B_1 + (b - 1)r_2 B_2)S + S(1 - R)}{S^2 - ar_1(S - R)B_1 - br_2(S - R)B_2 + S(1 - R)}.$$

³The proof proceeds similarly if there are other a_i 's equal to 1.

Note that the numerator and the denominator are positive for all $0 \leq B_1, B_2 \leq 1$. When $B_1 = B_2 = B$, both terms are equal to

$$S^2 - S(S - R)B + S(1 - R),$$

and so $h(B_1, B_2) = 1$. Moreover, using the assumption $a < b$, we find that

- $h(B_1, B_2) < 1$ when $B_1 = 0$ and $B_2 > 0$, and
- $h(B_1, B_2) > 1$ when $B_1 > 0$ and $B_2 = 0$.

Hence, the function $h(B_1, B_2)$ is increasing in B_1 for fixed B_2 and decreasing in B_2 for fixed B_1 .⁴ This implies that the function is minimized when $B_1 = 0$ and $B_2 = 1$, where its value is

$$\begin{aligned} h(B_1 = 0, B_2 = 1) &= \frac{S^2 - (b - 1)r_2S + S(1 - R)}{S^2 - br_2(S - R) + S(1 - R)} \\ &= \frac{(ar_1 + br_2)(ar_1 + 1 - r_1)}{a(a - 1)r_1^2 + a(b - 1)r_1r_2 + ar_1 + br_2} \\ &= \rho(a, b, r_1, r_2), \end{aligned}$$

as claimed. The proof that this approximation ratio is tight is left to Appendix B.

Finally, we can obtain analogous results for revenue by essentially repeating the same argument but instead writing $A_2 = \frac{(1-F(p_2))p_2}{(1-F(p_1))p_1}$ in the proof of Theorem 3.2, and for any convex combination of welfare and revenue by writing A_2 as the appropriate convex combination of the two corresponding terms. \square

Theorem 3.3 allows us to obtain the worst-case approximation ratio in arbitrary settings of the parameters. Some examples follow:

- Suppose that $r_1 + r_2 = 1$ (i.e., a job appears at every timestep). This assumption can in fact be made without loss of generality, because we can convert jobs not arriving to jobs arriving with a value of 0 as long as all prices are non-zero. This only changes F and thus is irrelevant for the calculation of ρ . In this case, the approximation ratio is

$$\rho(a, b, r, 1 - r) = \frac{(ar + b - br)(ar + 1 - r)}{a(a - b)r^2 + b(a - 1)r + b}.$$

- Next, we look at how the bound behaves when the job lengths are close together or far apart. Suppose for convenience that $r_1 = r_2 = 1/2$ (i.e., a job appears at every timestep and is of length a or b with equal probability). Then the approximation ratio is

$$\rho\left(a, b, \frac{1}{2}, \frac{1}{2}\right) = \frac{(a + b)(a + 1)}{a^2 + ab + 2b}.$$

The approximation ratio for $a = 1, 2, 3$ and different values of b can be found in Figure 3. In particular, the approximation ratio is

- $\frac{6}{7}$ if $a = 1$ and $b = 2$,
- $\frac{4}{5}$ if $a = 1$ and $b = 3$,
- $\frac{15}{16}$ if $a = 2$ and $b = 3$, and
- $\frac{a+1}{a+2}$ if $b \rightarrow \infty$.

⁴To see this, observe that an arbitrary function of the form $f(x) = \frac{p+qx}{r+sx}$ for real constants p, q, r, s is either increasing, decreasing, or constant in any interval where the denominator is non-zero, depending on whether $qr - ps$ is positive, negative, or zero, respectively.




Fig. 3. The approximation ratio $\rho(a, b, 1/2, 1/2)$ for $a = 1, 2, 3$ and different values of b .




Fig. 4. The approximation ratio $\rho(1, b, r, 1-r)$ for $b = 2, 3, 4$ and different values of $r \in [0, 1]$.

Note that if $b \rightarrow \infty$, then the ratio approaches 1 as a grows. This makes sense because when the two job lengths are large and close to each other, there is little difference between accepting one or the other. In addition, if we take $b = a + 1$, then the ratio becomes

$$\rho\left(a, a+1, \frac{1}{2}, \frac{1}{2}\right) = \frac{2a^2 + 3a + 1}{2a^2 + 3a + 2}.$$

This also converges to 1 as $a \rightarrow \infty$.

- We now consider the behavior of the bound when the shorter job length is fixed. Suppose for convenience that $a = 1$ (i.e., the shorter job length is 1). Then the approximation ratio is

$$\rho(1, b, r_1, r_2) = \frac{r_1 + br_2}{(b-1)r_1r_2 + r_1 + br_2}.$$

The approximation ratio for $b = 2, 3, 4$ and different values of r_1, r_2 with $r_1 + r_2 = 1$ can be found in Figure 4. In particular, as the longer job length grows, this ratio decreases and approaches $\frac{1}{1+r_1}$. This is consistent with the intuition that the approximation gets worse as the job lengths are farther apart.




Fig. 5. The approximation ratio $\rho(a, \infty, r_1, r_2)$ for $a = 1, 2, 3$ and different values of $r_1 \in [0, 1]$. The ratio is independent of r_2 .

- We next look at the “opposite” of the previous case and assume that the longer job length is extremely large. Suppose that $b \rightarrow \infty$. Then the approximation ratio is

$$\rho(a, \infty, r_1, r_2) = \frac{ar_1 + 1 - r_1}{ar_1 + 1}.$$

Note that this ratio does not depend on r_2 . The ratio for $a = 1, 2, 3$ and different values of r_1 can be found in Figure 5. In particular, the ratio increases as the shorter job length grows. Again, this is consistent with the intuition that the approximation gets worse as the job lengths are farther apart.

If we fix the probabilities r_1, r_2 , we can derive a tight worst-case bound over all possible job lengths a, b .

THEOREM 3.4. *For fixed r_1, r_2 , we have*

$$\rho(a, b, r_1, r_2) \geq \frac{1}{1 + r_1}$$

for arbitrary a, b . Moreover, this bound is the best possible.

PROOF. From Theorem 3.3, we need to show the inequality

$$\frac{(ar_1 + br_2)(ar_1 + 1 - r_1)}{a(a-1)r_1^2 + a(b-1)r_1r_2 + ar_1 + br_2} \geq \frac{1}{1 + r_1}.$$

This simplifies to

$$r_1(a(a-1)r_1^2 + (a-1)br_1r_2 + ar_1 + ar_2) \geq 0.$$

Since each term on the left-hand side is non-negative, the inequality holds.

For tightness of the bound, let $a = 1$ and $b \rightarrow \infty$. The approximation ratio is

$$\begin{aligned} \rho(1, \infty, r_1, r_2) &= \frac{r_1 + br_2}{r_1(1 - r_2) + br_2(1 + r_1)} \\ &= \frac{1}{1 + r_1}, \end{aligned}$$

as desired. □

Note that the fact that the bound is tight at $a = 1$ and $b \rightarrow \infty$ is consistent with the intuition that the further apart the job lengths are, the more welfare and revenue there is to be gained by setting different prices for different job lengths, and consequently the worse the approximation ratio.

Finally, we show that we can obtain at least 50% of the welfare or revenue from setting two prices by using one of those prices.

THEOREM 3.5. *For arbitrary a, b, r_1, r_2 , we have*

$$\rho(a, b, r_1, r_2) \geq \frac{1}{2}.$$

Moreover, this bound is the best possible.

PROOF. Using Theorem 3.4, we find that

$$\rho(a, b, r_1, r_2) \geq \frac{1}{1 + r_1} \geq \frac{1}{2}$$

since $r_1 \leq 1$.

For tightness of the bound, let $a = 1$ and $r_2 = 1 - r_1$. The approximation ratio is

$$\rho(1, b, r, 1 - r) = \frac{r + (1 - r)b}{r^2 + (1 - r^2)b}.$$

Taking $b = \frac{1}{(1-r)^2}$, the ratio becomes

$$\frac{r + \frac{1}{1-r}}{r^2 + \frac{1+r}{1-r}} = \frac{r - r^2 + 1}{r^2 - r^3 + 1 + r},$$

which approaches $1/2$ as r approaches 1 from below.⁵ □

Although we do not have a general formula for the worst-case approximation ratio for each choice of the parameters $a_1, \dots, a_n, r_1, \dots, r_n$ as we do for the case of two job lengths, the function h in the proof of Theorem 3.2 still allows us to derive a tighter bound for each specific case. Note that to find the minimum of h , it suffices to check $B_i = 0$ or 1 (see Footnote 4), so we only have a finite number of cases to check.

In fact, one can show that the minimum is always attained when $B_1 = 0$ and $B_n = 1$. However, as we show next, the remaining B_i 's may be 0 or 1 at the minimum, depending on the job lengths and the probability distribution over them. We take $n = 3$ and assume for convenience that $r_1 = r_2 = r_3 = 1/3$ and $(a_1, a_2) = (2, 3)$. The following examples show that B_2 can be 0 or 1 at the minimum depending on how far the longest job length a_3 is from a_1 and a_2 :

- Suppose that $a_3 = 6$. Then we have

$$h(B_1, B_2, B_3) = \frac{121 - 11B_1 - 22B_2 - 55B_3}{121 - 16B_1 - 24B_2 - 48B_3}.$$

This is minimized at $(B_1, B_2, B_3) = (0, 1, 1)$, where its value is $44/49$.

- Suppose that $a_3 = 7$. Then we have

$$h(B_1, B_2, B_3) = \frac{48 - 4B_1 - 8B_2 - 24B_3}{48 - 6B_1 - 9B_2 - 21B_3}.$$

This is minimized at $(B_1, B_2, B_3) = (0, B_2, 1)$ for any $0 \leq B_2 \leq 1$, where its value is $8/9$.

⁵The approximation ratio does not necessarily converge to $1/2$ for an arbitrary direction as $b \rightarrow \infty$ and r approaches 1 from below. For instance, if we take $b = \frac{1}{1-r}$ and r approaches 1 from below, then the ratio converges to $2/3$.

- Suppose that $a_3 = 8$. Then we have

$$h(B_1, B_2, B_3) = \frac{169 - 13B_1 - 26B_2 - 91B_3}{169 - 20B_1 - 30B_2 - 80B_3}.$$

This is minimized at $(B_1, B_2, B_3) = (0, 0, 1)$, where its value is $78/89$.

The preceding examples show that the transition point where the optimal value of B_2 goes from 0 to 1 for $r_1 = r_2 = r_3 = 1/3$ and $(a_1, a_2) = (2, 3)$ is at $a_3 = 7$, where $h(B_1, B_2, B_3)$ takes on the same value for any $0 \leq B_2 \leq 1$.

3.4 Extension

In this section, we show that by using a single price, we can obtain 50% of the welfare not only compared to using multiple prices but also compared to the offline optimal welfare.⁶ In fact, we will also not need the assumption that the job length and the value per timestep are independent. However, the result only works for particular prices rather than arbitrary ones, and we cannot obtain tighter results for specific parameters using this method. The proof of this result is similar to prophet inequality proofs of Feldman et al. (2015) and Dütting et al. (2017).

THEOREM 3.6. *Assume that the job length and the value per timestep are not necessarily independent. There exists a price p such that we can achieve a 50% approximation of the offline optimal welfare by using p as the single price.*

PROOF. We prove the result for a distribution with discrete support and extend it to continuous and mixed support later. Our approach follows that of Feldman et al. (2015). As job length and value are not independent, assume that there are jobs of classes $1, 2, \dots, n$ and that a job of class j arrives with probability r_j , has length a_j and value per timestep v_j . Note that a_j and v_j need not be different for different classes of jobs.

We can write an “expected LP” to upper bound the maximum welfare per timestep as follows:

$$\begin{aligned} Opt &= \max \sum_j x_j v_j a_j \text{ subject to} \\ x_j &\leq r_j \\ \sum_j x_j a_j &\leq 1. \end{aligned}$$

Here, Opt is the optimal long-run average welfare per timestep, and the ratio x_j/r_j can be thought of as the probability that an arriving job of class j is accepted. The constraints then say that this probability is at most 1 and that we cannot accept more jobs in expectation than, if they arrived perfectly, would saturate the machine.

Take $p = Opt/2$, and consider some time t . Let y_t be the probability that the server is occupied at time t (possibly by the job arriving at time t). Then the expected revenue at time t is $py_t = (Opt/2) \cdot y_t$. The expected consumer surplus of the job arriving at time t is at least $\sum_j r_j (v_j - p) a_j (1 - y_t)$, because the probability that the server is unoccupied at time t (either by the job arriving at time t or by an earlier job) is a lower bound on the probability that the server was not occupied when the job at time t arrived. By the constraints of the LP, the consumer surplus is at least $(Opt/2) \cdot (1 - y_t)$. Welfare is the sum of revenue and consumer surplus, so summing over all t shows that the welfare per timestep is at least $Opt/2$.

⁶For the offline optimal welfare, we compute the limit of the expected average offline optimal welfare per timestep as the time horizon grows.

If the support of the distribution is not discrete, a similar argument still works. To determine Opt , we order the jobs in decreasing order of value per timestep and take the jobs until the machine is saturated. The rest of the proof then follows in the same way as before, with sums replaced by integrals.

As a further extension, this argument can also be adapted to the multiple server case. We can solve the expected LP for each server individually and compute a price $p_i = Opt_i/2$. We can then select whichever of these prices maximizes welfare to recover the $1/H_n$ bound from Theorem 4.2. (For some i , the welfare from the i servers with the largest values of Opt_i at price p_i , which we can lower bound as $iOpt_i/2$, must be at least $\frac{1}{2H_n} \sum_i Opt_i$.) \square

4 MULTIPLE SERVERS

In this section, we assume that there are multiple servers, each of which receives jobs of various lengths. Under the assumption that the servers have the same probability of receiving no job at a timestep, we show in Section 4.1 an approximation bound of the welfare and revenue of setting one price for all servers compared to setting an individual price for each server. This yields a strong bound when at least one of the dimensions of the parameters is not too extreme (e.g., the number of servers or the job lengths are not too large). In Section 4.2, we combine the newly obtained results with those from Section 3. Using a composition technique, we derive a general result that compares the welfare and revenue obtained by a restricted mechanism that sets the same price for all servers and all job lengths against those obtained by a mechanism that can set a different price for each job length of each particular server. We show that even with the heavy restrictions, the former mechanism still provides a reasonable approximation to the latter in a wide range of situations. Using similar techniques, we also obtain approximation bounds when this assumption does not hold but there is only one job length across all servers. The analysis of the latter setting is deferred to Appendix C.

As in Section 3, our approximation results hold for arbitrary (i.e., not necessarily optimal) pricing schemes, and the price we use in the single-price setting can be drawn from one of the prices in the multi-price setting.

4.1 One Price per Server

Assume that at each timestep, either zero or one job appears for each server $1 \leq j \leq n$. Server j receives jobs of length $a_{j1} \leq a_{j2} \leq \dots \leq a_{jn_j}$ with probability $r_{j1}, r_{j2}, \dots, r_{jn_j}$, respectively. Suppose that we set a price per timestep p_j for all jobs on server j . Recall that the value per timestep of a job is drawn from a distribution with cumulative distribution function F and probability density function f , and that we assume that $\sum_{i=1}^{n_j} r_{ji}$ is constant. Let $S_j = a_{j1}r_{j1} + \dots + a_{jn_j}r_{jn_j}$ and $R = r_{j1} + \dots + r_{jn_j}$.

Using the formula (1) for c_w given in Lemma 3.1, we find that the welfare per timestep is

$$\begin{aligned} d_w(p_1, p_2, \dots, p_n) &= \sum_{j=1}^n \frac{S_j \int_{x \geq p_j} \ell d\mu}{S_j - (S_j - R)F(p_j) + (1 - R)} \\ &= \sum_{j=1}^n \frac{\int_{x \geq p_j} \ell d\mu}{1 - \left(1 - \frac{R}{S_j}\right)F(p_j) + \frac{1-R}{S_j}}. \end{aligned}$$

If we set the same price $p = p_1 = \dots = p_n$ for different servers, our welfare per timestep becomes

$$d_w(p) = \sum_{j=1}^n \frac{\int_{x \geq p} \ell d\mu}{1 - \left(1 - \frac{R}{S_j}\right) F(p) + \frac{1-R}{S_j}}.$$

Similarly, we have the formulas for revenue per timestep

$$d_r(p_1, p_2, \dots, p_n) = \sum_{j=1}^n \frac{(1 - F(p_j))p_j}{1 - \left(1 - \frac{R}{S_j}\right) F(p_j) + \frac{1-R}{S_j}}$$

and

$$d_r(p) = \sum_{j=1}^n \frac{(1 - F(p))p}{1 - \left(1 - \frac{R}{S_j}\right) F(p) + \frac{1-R}{S_j}}.$$

We show that if at least one dimension of the parameters is not too extreme (e.g., the number of servers or the job lengths are bounded), then we can obtain a reasonable approximation of the welfare and revenue in the multi-price setting by setting just one price.

THEOREM 4.1. *For any prices p_1, p_2, \dots, p_n that we set in the multi-price setting, we can achieve a welfare (respectively, revenue, or any convex combination of welfare and revenue) approximation of at least*

$$\max\left(\frac{1}{H_n}, \frac{M-1}{M \ln M}\right)$$

in the one-price setting, where $H_n = 1 + \frac{1}{2} + \dots + \frac{1}{n} \approx \ln n$ is the n th harmonic number and $M = \max_{i,j} \frac{S_i}{S_j}$.

In particular, if all job lengths are bounded above by c , then $R \leq S_j \leq cR$ for all $1 \leq j \leq n$, and so $\max_{i,j} \frac{S_i}{S_j} \leq c$. The theorem then implies that the approximation ratio is at least $\frac{c-1}{c \ln c}$.

The proof follows a similar outline to that of Theorem 3.2, but the details are more involved.

PROOF. We first consider welfare. We will work with the ratio

$$\frac{\max(d_w(p_1), \dots, d_w(p_n))}{d_w(p_1, \dots, p_n)}$$

and try to minimize it.

Writing $A_j = \frac{\int_{x \geq p_j} \ell d\mu}{\int_{x \geq p_1} \ell d\mu}$ (in particular, $A_1 = 1$), $B_j = F(p_j)$, and $s_j = \frac{1}{S_j}$ for $1 \leq j \leq n$, the ratio to minimize becomes

$$g(A_1, \dots, A_n, B_1, \dots, B_n) := \max_{j=1}^n \left(\sum_{i=1}^n \frac{A_j}{1 - (1 - R s_i) B_j + (1 - R) s_i} \right) \cdot \frac{1}{\sum_{i=1}^n \frac{A_i}{1 - (1 - R s_i) B_i + (1 - R) s_i}}.$$

Let $U_j = \sum_{i=1}^n \frac{1}{1 - (1 - R s_i) B_j + (1 - R) s_i}$ for $1 \leq j \leq n$, and $T = \sum_{i=1}^n \frac{A_i}{1 - (1 - R s_i) B_i + (1 - R) s_i}$. We have

$$g(A_1, \dots, A_n, B_1, \dots, B_n) = \frac{\max_{j=1}^n (A_j U_j)}{T}.$$

By dividing into cases according to the term that the max function outputs, we can deduce that we only need to minimize the function

$$h(B_1, \dots, B_n) := \frac{1}{\sum_{j=1}^n \frac{1}{\sum_{i=1}^n \frac{1}{1 - (1 - R s_j) B_j + (1 - R) s_j}}};$$

the details of this casework can be found in Appendix D. In other words, the reciprocal of this function, which we want to maximize, is

$$\begin{aligned} h_0(B_1, \dots, B_n) &:= \sum_{j=1}^n \frac{1}{\sum_{i=1}^n \frac{1-(1-Rs_j)B_j+(1-R)s_j}{1-(1-Rs_i)B_j+(1-R)s_i}} \\ &= \sum_{j=1}^n \frac{1}{\sum_{i=1}^n \frac{1-B_j+s_j(1-R+RB_j)}{1-B_j+s_i(1-R+RB_j)}}. \end{aligned}$$

Assume without loss of generality that $s_1 \geq s_2 \geq \dots \geq s_n$. For $1 \leq j \leq n$, write $h_j = \frac{1}{\sum_{i=1}^n \frac{1-B_j+s_j(1-R+RB_j)}{1-B_j+s_i(1-R+RB_j)}}$ (i.e., the j th term of the sum that constitutes h_0). The last $n+1-j$ terms of the sum in the denominator of h_j are at least 1 since $s_j \geq s_i$ for $i \geq j$. This implies that $h_j \leq \frac{1}{n+1-j}$, and therefore

$$\begin{aligned} h_0(B_1, \dots, B_n) &= h_1 + h_2 + \dots + h_n \\ &\leq \frac{1}{n} + \frac{1}{n-1} + \dots + 1 = H_n. \end{aligned}$$

Equivalently, $h(B_1, \dots, B_n) \geq \frac{1}{H_n}$.

Next, since $M = \max_{i,j} \frac{s_i}{s_j}$, we have $\frac{s_i}{s_j} \leq M$ for all i, j . Note that

$$\frac{1-(1-Rs_j)B_j+(1-R)s_j}{1-(1-Rs_i)B_j+(1-R)s_i} \geq \frac{1}{M}$$

for all i, j , as both the numerator and the denominator are positive, and when seen as a function of B_j , the expression is monotonic and takes on values at least $\frac{1}{M}$ at $B_j = 0$ and $B_j = 1$. This implies that

$$h_j \leq \frac{1}{n+1-j + \frac{j-1}{M}}$$

for all $1 \leq j \leq n$, and therefore

$$\begin{aligned} h_0(B_1, \dots, B_n) &= \sum_{j=1}^n \frac{1}{n-(j-1)\left(1-\frac{1}{M}\right)} \\ &= \sum_{j=1}^n \frac{1}{n} \cdot \frac{1}{1-\frac{j-1}{n}\left(1-\frac{1}{M}\right)}. \end{aligned}$$

The last sum is the left Riemann sum of the function $\frac{1}{1-x\left(1-\frac{1}{M}\right)}$. This function is increasing in x , so its integral between 0 and 1 is an upper bound for our sum. Hence, we have

$$h_0(B_1, \dots, B_n) \leq \int_0^1 \frac{1}{1-x\left(1-\frac{1}{M}\right)} dx = \frac{M \ln M}{M-1}.$$

Equivalently, $h(B_1, \dots, B_n) \geq \frac{M-1}{M \ln M}$.

Combining the two bounds, we have

$$h(B_1, \dots, B_n) \geq \max\left(\frac{1}{H_n}, \frac{M-1}{M \ln M}\right),$$

as desired.

Finally, we can obtain analogous results for revenue by essentially repeating the same argument but instead writing $A_j = \frac{(1-F(p_j))p_j}{(1-F(p_1))p_1}$ for $1 \leq j \leq n$, and for any convex combination of welfare and revenue by writing A_j as the appropriate convex combination of the two corresponding terms. \square

Done with the proof of the theorem, we now address the tightness of the approximation ratio. The upper bound H_n for h_0 is the best possible in the sense that there exist values $B_1, \dots, B_n, s_1, \dots, s_n, R$ such that h_0 gets arbitrarily close to H_n . In particular, take $s_j = c^{-2j}$ and $B_j = 1 - c^{-2j+1}$ for some large constant c , and $R = 1$. We have

$$\begin{aligned} h_0(B_1, \dots, B_n) &= \sum_{j=1}^n \frac{1}{\sum_{i=1}^n \frac{1-B_j+B_j s_j}{1-B_j+B_j s_i}} \\ &= \sum_{j=1}^n \frac{1}{\sum_{i=1}^n \frac{c^{-2j+1}+c^{-2j}-c^{-4j+1}}{c^{-2j+1}+c^{-2i}-c^{-2j-2i+1}}} \\ &= \sum_{j=1}^n \frac{1}{\sum_{i=1}^n \frac{c+1-c^{-2j+1}}{c+c^{2(j-i)}-c^{-2i+1}}}. \end{aligned}$$

Taking $c \rightarrow \infty$, we find that for $j \leq i$, the fraction $\frac{c+1-c^{-2j+1}}{c+c^{2(j-i)}-c^{-2i+1}}$ converges to 1, whereas for $j > i$, it converges to 0. Hence, $h_j \rightarrow \frac{1}{n+1-j}$, and consequently $h_0 \rightarrow H_n$.

Although this argument does not directly imply the tightness of the approximation ratio, we see it as strong evidence for that claim.

4.2 Multiple Prices per Server

Assume as in Section 4.1 that at each timestep, server j receives jobs of length $a_{j1} \leq a_{j2} \leq \dots \leq a_{jn_j}$ with probability $r_{j1}, r_{j2}, \dots, r_{jn_j}$, respectively. In this section, we consider setting an individual price not only for each server but also for each job length of that server. In particular, suppose that we set a price per timestep p_{ji} for jobs of length a_{ji} on server j . Recall that the value per timestep of a job is drawn from a distribution with cumulative distribution function F and probability density function f , and that we assume that $\sum_{i=1}^{n_j} r_{ji}$ is constant. Let $S_j = a_{j1}r_{j1} + \dots + a_{jn_j}r_{jn_j}$.

We will compare a setting where we have considerable freedom with our pricing scheme and can set a different price p_{ji} for each job length a_{ji} on each server j with a setting where we have limited freedom and must set the same price p for all job lengths and all servers. We show that by “composing” our results on the two dimensions, we can obtain an approximation of the welfare and revenue of setting different prices by setting a single price.

THEOREM 4.2. *For any prices p_{ji} , where $1 \leq j \leq n$ and $1 \leq i \leq n_j$ for each j , that we set in the multi-price setting, we can achieve a welfare (respectively, revenue, or any convex combination of welfare and revenue) approximation of at least*

$$\frac{1}{2} \cdot \max\left(\frac{1}{H_n}, \frac{M-1}{M \ln M}\right)$$

in the one-price setting, where $H_n = 1 + \frac{1}{2} + \dots + \frac{1}{n} \approx \ln n$ is the n th harmonic number and $M = \max_{i,j} \frac{S_i}{S_j}$.

PROOF. Consider welfare. By Theorem 3.2, for each server j we can achieve a $\frac{1}{2}$ -approximation of the welfare in the multi-price setting by setting a single price p_j for all job lengths. However, using Theorem 4.1, we can approximate the latter welfare by a factor of $\max(\frac{1}{H_n}, \frac{M-1}{M \ln M})$ by setting

a single price p for all servers. Therefore, setting a single price p also yields a $\frac{1}{2} \cdot \max(\frac{1}{H_n}, \frac{M-1}{M \ln M})$ -approximation of the original welfare.

The same argument holds for revenue and for any convex combination of welfare and revenue. \square

If we have tighter approximations for either the “different prices for different job lengths” or the “different prices for different servers” dimension, for instance, by knowing the values of some of the parameters, then the same composition argument yields a correspondingly tighter bound.

5 CONCLUSION

In this article, we study how well simple pricing schemes that are oblivious to certain parameters can approximate optimal schemes with respect to welfare and revenue, and prove several results when the simple schemes are restricted to setting the same price for all servers or all job lengths. Our results provide an explanation of the efficacy of such schemes in practice, including the one shown in Figure 1 for virtual machines on Microsoft Azure. Since simple schemes do not require agents to spend time and resources to determine their specific parameter values, our results also serve as an argument in favor of using these schemes in a range of applications. It is worth noting that as all of our results are of worst-case nature, we can expect the guarantees on welfare and revenue to be significantly better than these pessimistic bounds in practical instances where the parameters are not adversarially tailored.

We believe that there is still much interesting work to be done in the study of simple pricing schemes for the cloud. We conclude our article by listing some intriguing future directions:

- The example in Section 3.1 shows that the approximations for the uniform distribution are significantly better than 50%. How does the shape of the distribution affect the approximation ratio that can be achieved?
- In many scheduling applications, a job can be scheduled online to any server that is not occupied at the time. Does a good welfare or revenue approximation hold in such a model?
- Can our results be extended to models with more fluid job arrivals, such as one where several jobs can arrive at each timestep?
- Can we approximate welfare and revenue simultaneously? A trivial randomized approach would be to choose with equal probability whether to approximate welfare or revenue. According to Theorem 3.2, this yields a 1/4-approximation for both expected welfare and expected revenue of the single-price setting in comparison to the multi-price setting for job lengths.
- As we mentioned in Section 3.4, the proof of Theorem 3.6 resembles the prophet inequality proofs of Feldman et al. (2015) and Dütting et al. (2017). Does there exist a reduction from one setting to the other?

APPENDICES

A MISSING DETAILS IN THE PROOF OF THEOREM 3.2

We divide into cases according to the term that the max function outputs.

Case 1. The max function outputs the first term, $\frac{SA_1}{S-(S-R)B_1+(1-R)}$.

Taking into account that $A_1 = 1$, we want to minimize the ratio

$$\frac{S}{S - (S - R)B_1 + (1 - R)} \cdot \frac{S - ((a_1 - 1)r_1B_1 + \dots + (a_n - 1)r_nB_n) + (1 - R)}{a_1r_1 + a_2r_2A_2 + \dots + a_nr_nA_n},$$

where $A_i \leq \frac{S-(S-R)B_i+(1-R)}{S-(S-R)B_1+(1-R)}$ for $i \geq 2$.

For any A_i , if we fix the remaining A_j and all B_j , this is a decreasing function in A_i . To minimize it, we should set $A_i = \frac{S-(S-R)B_i+(1-R)}{S-(S-R)B_i+(1-R)}$ for all $i \geq 2$. The ratio becomes

$$\frac{S^2 - ((a_1 - 1)r_1B_1 + \dots + (a_n - 1)r_nB_n)S + S(1 - R)}{S^2 - (a_1r_1B_1 + \dots + a_nr_nB_n)(S - R) + S(1 - R)}.$$

Case 2. The max function outputs the i th term, $\frac{SA_i}{S-(S-R)B_i+(1-R)}$, for some $i \geq 2$. This means that $A_j \leq \frac{S-(S-R)B_j+(1-R)}{S-(S-R)B_i+(1-R)} \cdot A_i$ for all $j \geq 2$ with $j \neq i$, and $A_i \geq \frac{S-(S-R)B_i+(1-R)}{S-(S-R)B_i+(1-R)}$. We want to minimize the ratio

$$\frac{SA_i}{S - (S - R)B_i + (1 - R)} \cdot \frac{S - ((a_1 - 1)r_1B_1 + \dots + (a_n - 1)r_nB_n) + (1 - R)}{a_1r_1 + a_2r_2A_2 + \dots + a_nr_nA_n},$$

or equivalently,

$$\frac{SA_i}{a_1r_1A_1 + \dots + a_nr_nA_n} \cdot \frac{S - ((a_1 - 1)r_1B_1 + \dots + (a_n - 1)r_nB_n) + (1 - R)}{S - (S - R)B_i + (1 - R)}.$$

For any $j \geq 2$ with $j \neq i$, if we fix all terms A_k except A_j and fix all B_k , then this function is decreasing in A_j . To minimize it, we should set $A_j = \frac{S-(S-R)B_j+(1-R)}{S-(S-R)B_i+(1-R)} \cdot A_i$. The resulting function is increasing in A_i if we fix all B_k , so we should set $A_i = \frac{S-(S-R)B_i+(1-R)}{S-(S-R)B_i+(1-R)}$. We obtain the same ratio as in Case 1.

Hence, in either case, we are left with minimizing the function

$$h(B_1, \dots, B_n) := \frac{S^2 - ((a_1 - 1)r_1B_1 + \dots + (a_n - 1)r_nB_n)S + S(1 - R)}{S^2 - (a_1r_1B_1 + \dots + a_nr_nB_n)(S - R) + S(1 - R)},$$

as claimed.

B MISSING DETAILS IN THE PROOF OF THEOREM 3.3

We show that the approximation ratio is tight even if we are allowed to set an arbitrary price (i.e., not necessarily p_1 or p_2) in the one-price setting. To this end, consider a discrete bimodal distribution where a high probability $q_1 \approx 1$ is put on a value $v_1 \approx 0$ and a small probability $q_2 \approx 0$ is put on a value $v_2 \approx 1$.⁷ The values v_1, v_2 and the probabilities are chosen arbitrarily close to 0 and 1 and so that the relation

$$\frac{q_2v_2}{q_1v_1} = \frac{1}{S - R}$$

is satisfied.

In the two-price setting, we can set prices $p_1 = v_1$ and $p_2 = v_2$ and obtain welfare

$$\begin{aligned} c_w(p_1 = v_1, p_2 = v_2) &= \frac{ar_1(q_1v_1 + q_2v_2) + br_2(q_2v_2)}{S - ((a - 1)r_1F(v_1) + (b - 1)r_2F(v_2)) + (1 - R)} \\ &\approx \frac{ar_1(q_1v_1 + q_2v_2) + br_2(q_2v_2)}{S - (b - 1)r_2 + (1 - R)} \\ &= \frac{q_2v_2(ar_1(S + 1 - R) + br_2)}{S - (b - 1)r_2 + (1 - R)}. \end{aligned}$$

However, in the one-price setting, there are three price ranges that we can pick: $[0, v_1]$, $(v_1, v_2]$, and (v_2, ∞) . If we set a price in the range (v_2, ∞) , no job is accepted and the welfare is zero. For

⁷Intuitively, we want to accept all short jobs but only high-valued long jobs. We can do this in the two-price setting; however, in the one-price setting, we are forced to either accept low-valued long jobs or reject low-valued short jobs.

each of the other two ranges, setting any price in the range yields the same set of accepted jobs and thus the same welfare. Hence, it suffices to consider setting prices v_1 and v_2 . We have

$$c_w(p = v_1) = \frac{S(q_1v_1 + q_2v_2)}{S + 1 - R} = Sq_2v_2$$

and

$$c_w(p = v_2) \approx \frac{Sq_2v_2}{S - (S - R) + (1 - R)} = Sq_2v_2.$$

We obtain the same welfare per timestep in either case. It follows that the approximation ratio is at most

$$\begin{aligned} \frac{c_w(p = v_1)}{c_w(p_1 = v_1, p_2 = v_2)} &= \frac{Sq_2v_2(S - (b - 1)r_2 + (1 - R))}{q_2v_2(ar_1(S + 1 - R) + br_2)} \\ &= \frac{S(ar_1 + 1 - r_1)}{a(a - 1)r_1^2 + a(b - 1)r_1r_2 + ar_1 + br_2} \\ &= \rho(a, b, r_1, r_2), \end{aligned}$$

as desired.

C MULTIPLE SERVERS, ONE JOB LENGTH

Assume that at each timestep, either zero or one job appears for each server. Server j receives a job of length a with probability r_j . Suppose that we set a price per timestep p_j for jobs on server j . Recall that the value per timestep of a job is drawn from a distribution with cumulative distribution function F and probability density function f .

Using the formula (1) for c_w given in Lemma 3.1, we find that the welfare per timestep is

$$\begin{aligned} e_w(p_1, p_2, \dots, p_n) &= \sum_{j=1}^n \frac{ar_j \int_{x \geq p_j} \ell d\mu}{ar_j - (a - 1)r_jF(p_j) + (1 - r_j)} \\ &= \sum_{j=1}^n \frac{a \int_{x \geq p_j} \ell d\mu}{(a - 1)(1 - F(p_j)) + \frac{1}{r_j}}. \end{aligned}$$

If we set the same price $p = p_1 = \dots = p_n$ for different servers, our welfare per timestep becomes

$$e_w(p) = \sum_{j=1}^n \frac{a \int_{x \geq p} \ell d\mu}{\frac{1}{r_j} + (a - 1)(1 - F(p))}.$$

Similarly, we have the formulas for revenue per timestep

$$e_r(p_1, p_2, \dots, p_n) = \sum_{j=1}^n \frac{a(1 - F(p_j))p_j}{(a - 1)(1 - F(p_j)) + \frac{1}{r_j}}$$

and

$$e_r(p) = \sum_{j=1}^n \frac{a(1 - F(p))p}{(a - 1)(1 - F(p)) + \frac{1}{r_j}}.$$

We will compare the welfare and revenue that can be obtained by setting a single price against setting several prices. Similarly to Section 4, we will show that if at least one dimension of the parameters is not too extreme (e.g., the number of servers, the job length, or the probabilities of jobs occurrence are bounded), then we can obtain a reasonable approximation of the welfare and revenue in the multi-price setting by setting just one price.

THEOREM C.1. *For any prices p_1, p_2, \dots, p_n that we set in the multi-price setting, we can achieve a welfare (respectively, revenue, or any convex combination of welfare and revenue) approximation of at least*

$$\max\left(\frac{1}{H_n}, \frac{M-1}{M \ln M}, \frac{1}{a}\right)$$

in the one-price setting, where $H_n = 1 + \frac{1}{2} + \dots + \frac{1}{n} \approx \ln n$ is the n th harmonic number and $M = \max_{i,j} \frac{r_i}{r_j}$.

In particular, if all probabilities of job occurrence are bounded below by $c \leq 1$, then $\max_{i,j} \frac{S_i}{S_j} \leq \frac{1}{c}$. The theorem then implies that the approximation ratio is also at least $\frac{\frac{1}{c}-1}{\frac{1}{c} \ln(\frac{1}{c})} = \frac{c-1}{\ln c}$.

PROOF. We first consider welfare. We will work with the ratio

$$\frac{\max(e_w(p_1), \dots, e_w(p_n))}{e_w(p_1, \dots, p_n)}$$

and try to minimize it.

Writing $A_j = \frac{\int_{x \geq p_j} \ell d\mu}{\int_{x \geq p_1} \ell d\mu}$ (in particular, $A_1 = 1$), $B_j = F(p_j)$, and $R_j = 1/r_j$ for $1 \leq j \leq n$, the ratio to minimize becomes

$$g(A_1, \dots, A_n, B_1, \dots, B_n) := \max_{j=1}^n \left(\sum_{i=1}^n \frac{A_j}{R_i + (a-1)(1-B_j)} \right) \cdot \frac{1}{\sum_{j=1}^n \frac{A_i}{R_i + (a-1)(1-B_i)}}.$$

Let $U_j = \sum_{i=1}^n \frac{1}{R_i + (a-1)(1-B_j)}$ for $1 \leq j \leq n$, and $T = \sum_{i=1}^n \frac{A_i}{R_i + (a-1)(1-B_i)}$. We have

$$g(A_1, \dots, A_n, B_1, \dots, B_n) = \frac{\max_{j=1}^n (A_j U_j)}{T}.$$

Case 1. The max function outputs the first term, $A_1 U_1$.

Taking into account that $A_1 = 1$, we want to minimize the ratio

$$\frac{U_1}{T} = \left(\sum_{i=1}^n \frac{1}{R_i + (a-1)(1-B_1)} \right) \cdot \frac{1}{\sum_{i=1}^n \frac{A_i}{R_i + (a-1)(1-B_i)}},$$

where $A_i \leq \frac{U_1}{U_i}$ for all $i \geq 2$.

For any A_j , if we fix the remaining A_i and all B_i , this is a decreasing function in A_j . To minimize it, we should set $A_i = \frac{U_1}{U_i}$ for all $i \geq 2$. The ratio becomes

$$\frac{1}{\sum_{j=1}^n \frac{1}{R_j + (a-1)(1-B_j)}}.$$

Case 2. The max function outputs the j th term, $A_j U_j$, for some $j \geq 2$.

This means that $A_i \leq \frac{A_j U_j}{U_i}$ for all $i \geq 2$ with $i \neq j$, and $A_j \geq \frac{U_1}{U_j}$. We want to minimize the ratio

$$\frac{A_j U_j}{T} = \left(\sum_{i=1}^n \frac{A_j}{R_i + (a-1)(1-B_j)} \right) \cdot \frac{1}{\sum_{i=1}^n \frac{A_i}{R_i + (a-1)(1-B_i)}}.$$

For any $i \geq 2$ with $i \neq j$, if we fix all terms A_k except A_i and fix all B_k , then this function is decreasing in A_i . To minimize it, we should set $A_i = \frac{A_j U_j}{U_i}$. The resulting function is increasing in A_j if we fix all B_k , so we should set $A_j = \frac{U_1}{U_j}$. We obtain the same ratio as in Case 1.

Hence, in either case, we are left with minimizing the function

$$h(B_1, \dots, B_n) := \frac{1}{\sum_{j=1}^n \frac{1}{\frac{R_j + (a-1)(1-B_j)}{R_i + (a-1)(1-B_j)}}}.$$

In other words, the reciprocal of this function, which we want to maximize, is

$$h_0(B_1, \dots, B_n) := \sum_{j=1}^n \frac{1}{\sum_{i=1}^n \frac{R_j + (a-1)(1-B_j)}{R_i + (a-1)(1-B_j)}}.$$

Assume without loss of generality that $R_1 \geq R_2 \geq \dots \geq R_n$. For $1 \leq j \leq n$, write $h_j = \frac{1}{\sum_{i=1}^n \frac{R_j + (a-1)(1-B_j)}{R_i + (a-1)(1-B_j)}}$ (i.e., the j th term of the sum that constitutes h_0). The last $n+1-j$ terms of the sum in the denominator of h_j are at least 1 since $R_j \geq R_i$ for $i \geq j$. This implies that $h_j \leq \frac{1}{n+1-j}$, and therefore

$$\begin{aligned} h_0(B_1, \dots, B_n) &= h_1 + h_2 + \dots + h_n \\ &\leq \frac{1}{n} + \frac{1}{n-1} + \dots + 1 = H_n. \end{aligned}$$

Equivalently, $h(B_1, \dots, B_n) \geq \frac{1}{H_n}$.

Next, since $M = \max_{i,j} \frac{r_i}{r_j}$, we have $\frac{R_i}{R_j} \leq M$ for all i, j . Note that

$$\frac{R_i + (a-1)(1-B_i)}{R_j + (a-1)(1-B_i)} \geq \frac{1}{M}$$

for all i, j . This implies that

$$h_j \leq \frac{1}{n+1-j + \frac{j-1}{M}}$$

for all $1 \leq j \leq n$, and therefore

$$\begin{aligned} h_0(B_1, \dots, B_n) &= \sum_{j=1}^n \frac{1}{n - (j-1) \left(1 - \frac{1}{M}\right)} \\ &= \sum_{j=1}^n \frac{1}{n} \cdot \frac{1}{1 - \frac{j-1}{n} \left(1 - \frac{1}{M}\right)}. \end{aligned}$$

The last sum is the left Riemann sum of the function $\frac{1}{1-x(1-\frac{1}{M})}$. This function is increasing in x , so its integral between 0 and 1 is an upper bound for our sum. Hence, we have

$$h_0(B_1, \dots, B_n) \leq \int_0^1 \frac{1}{1-x\left(1-\frac{1}{M}\right)} dx = \frac{M \ln M}{M-1}.$$

Equivalently, $h(B_1, \dots, B_n) \geq \frac{M-1}{M \ln M}$.

Finally, one can check that since $R_i \geq 1$ and $1 - B_i \leq 1$ for all i ,

$$\frac{R_i + (a-1)(1-B_i)}{R_j + (a-1)(1-B_i)} \geq \frac{1}{a} \cdot \frac{R_i}{R_j}$$

for all i, j . It follows that

$$\begin{aligned} h_0(B_1, \dots, B_n) &\leq \sum_{i=1}^n \frac{1}{\sum_{j=1}^n \frac{1}{a} \cdot \frac{R_i}{R_j}} \\ &= a \sum_{i=1}^n \frac{1}{\sum_{j=1}^n \frac{R_i}{R_j}} \\ &= a. \end{aligned}$$

Equivalently, $h(B_1, \dots, B_n) \geq \frac{1}{a}$.

Combining the three bounds, we have

$$h(B_1, \dots, B_n) \geq \max \left(\frac{1}{H_n}, \frac{M-1}{M \ln M}, \frac{1}{a} \right),$$

as desired.

Finally, we can obtain analogous results for revenue by essentially repeating the same argument but instead writing $A_j = \frac{(1-F(p_j))p_j}{(1-F(p_1))p_1}$ for $1 \leq j \leq n$, and for any convex combination of welfare and revenue by writing A_j as the appropriate convex combination of the two corresponding terms. \square

We now address the tightness of the approximation ratio. The upper bound H_n for h_0 is the best possible in the sense that there exist values $B_1, \dots, B_n, R_1, \dots, R_n, a$ such that h_0 gets arbitrarily close to H_n . In particular, take $R_j = c^{2(n-j)}$ and $B_j = 1 - \frac{c^{2(n-j)+1}}{a-1}$ for some large constant c . We have

$$\begin{aligned} h_0(B_1, \dots, B_n) &= \sum_{j=1}^n \frac{1}{\sum_{i=1}^n \frac{R_j + (a-1)(1-B_j)}{R_i + (a-1)(1-B_j)}} \\ &= \sum_{j=1}^n \frac{1}{\sum_{i=1}^n \frac{c^{2(n-j)} + c^{2(n-j)+1}}{c^{2(n-i)} + c^{2(n-j)+1}}} \\ &= \sum_{j=1}^n \frac{1}{\sum_{i=1}^n \frac{1+c}{c^{2(j-i)}+c}}. \end{aligned}$$

Taking $c \rightarrow \infty$, we find that for $j \leq i$, the fraction $\frac{1+c}{c^{2(j-i)}+c}$ converges to 1, whereas for $j > i$, it converges to 0. Hence, $h_j \rightarrow \frac{1}{n+1-j}$, and consequently $h_0 \rightarrow H_n$.

Although this argument does not directly imply the tightness of the approximation ratio, we see it as strong evidence for that claim.

D MISSING DETAILS IN THE PROOF OF THEOREM 4.1

We divide into cases according to the term that the max function outputs.

Case 1. The max function outputs the first term, $A_1 U_1$.

Taking into account that $A_1 = 1$, we want to minimize the ratio

$$\frac{U_1}{T} = \left(\sum_{i=1}^n \frac{1}{1 - (1 - R s_i) B_1 + (1 - R) s_i} \right) \cdot \frac{1}{\sum_{i=1}^n \frac{A_i}{1 - (1 - R s_i) B_1 + (1 - R) s_i}},$$

where $A_i \leq \frac{U_i}{U_1}$ for all $i \geq 2$.

For any A_j , if we fix the remaining A_i and all B_i , this is a decreasing function in A_j . To minimize it, we should set $A_i = \frac{U_i}{U_j}$ for all $i \geq 2$. The ratio becomes

$$\frac{1}{\sum_{j=1}^n \frac{1}{\sum_{i=1}^n \frac{1}{1-(1-Rs_j)B_j+(1-R)s_j}}}$$

Case 2. The max function outputs the j th term, $A_j U_j$, for some $j \geq 2$.

This means that $A_i \leq \frac{A_j U_j}{U_i}$ for all $i \geq 2$ with $i \neq j$, and $A_j \geq \frac{U_i}{U_j}$. We want to minimize the ratio

$$\frac{A_j U_j}{T} = \left(\sum_{i=1}^n \frac{A_j}{1 - (1 - R s_i) B_j + (1 - R) s_i} \right) \cdot \frac{1}{\sum_{i=1}^n \frac{A_i}{1 - (1 - R s_i) B_j + (1 - R) s_i}}.$$

For any $i \geq 2$ with $i \neq j$, if we fix all terms A_k except A_i and fix all B_k , then this function is decreasing in A_i . To minimize it, we should set $A_i = \frac{A_j U_j}{U_i}$. The resulting function is increasing in A_j if we fix all B_k , so we should set $A_j = \frac{U_i}{U_j}$. We obtain the same ratio as in Case 1.

Hence, in either case, we are left with minimizing the function

$$h(B_1, \dots, B_n) := \frac{1}{\sum_{j=1}^n \frac{1}{\sum_{i=1}^n \frac{1}{1-(1-Rs_j)B_j+(1-R)s_j}}},$$

as claimed.

ACKNOWLEDGMENTS

We thank the anonymous reviewers for helpful comments.

REFERENCES

- Vineet Abhishek, Ian A. Kash, and Peter Key. 2012. Fixed and market pricing for cloud services. In *Proceedings of the 7th Workshop on the Economics of Networks, Systems, and Computation*.
- Amazon. 2017. Amazon EC2 Spot Instances Pricing. Retrieved May 1, 2019 from <http://aws.amazon.com/ec2/spot/pricing>.
- Yossi Azar, Inna Kalp-Shaltiel, Brendan Lucier, Ishai Menache, Joseph (Seffi) Naor, and Jonathan Yaniv. 2015. Truthful online scheduling with commitments. In *Proceedings of the 16th ACM Conference on Economics and Computation*. 715–732.
- Azure. 2016. Microsoft Azure Pricing Calculator. Retrieved May 1, 2019 from <http://azure.microsoft.com/en-us/pricing/calculator>.
- Moshe Babaioff, Liad Blumrosen, Shaddin Dughmi, and Yaron Singer. 2011. Posting prices with unknown distributions. In *Proceedings of the 1st Innovations in Computer Science Conference*. 166–178.
- Liad Blumrosen and Thomas Holenstein. 2008. Posted prices vs. negotiations: An asymptotic analysis. In *Proceedings of the 9th ACM Conference on Electronic Commerce*. 49.
- Shuchi Chawla, Jason D. Hartline, David L. Malec, and Balasubramanian Sivan. 2010. Multi-parameter mechanism design and sequential posted pricing. In *Proceedings of the 42nd ACM Symposium on Theory of Computing*. 311–320.
- Ilan Reuven Cohen, Alon Eden, Amos Fiat, and Lukasz Jez. 2015. Pricing online decisions: Beyond auctions. In *Proceedings of the 26th Annual ACM-SIAM Symposium on Discrete Algorithms*. 73–91.
- Vincent Cohen-Addad, Alon Eden, Michal Feldman, and Amos Fiat. 2016. The invisible hand of dynamic market pricing. In *Proceedings of the 17th ACM Conference on Economics and Computation*. 383–400.
- Louis Columbus. 2016. Roundup of Cloud Computing Forecasts and Market Estimates, 2016. Retrieved May 1, 2019 from <http://www.forbes.com/sites/louiscolumnbus/2016/03/13/roundup-of-cloud-computing-forecasts-and-market-estimates-2016>.
- Sina Dehghani, Ian A. Kash, and Peter Key. 2016. *Online Stochastic Scheduling and Pricing the Cloud*. Working Paper.
- Ludwig Dierks and Sven Seuken. 2016. Cloud Pricing: The Spot Market Strikes Back. In *Proceedings of the Workshop on Economics of Cloud Computing*.
- Yann Disser, John Fearnley, Martin Gairing, Oliver Göbel, Max Klimm, Daniel Schmand, Alexander Skopalik, et al. 2016. Hiring secretaries over time: The benefit of concurrent employment. arXiv:1604.08125.

- Paul Dütting, Michal Feldman, Thomas Kesselheim, and Brendan Lucier. 2017. Prophet inequalities made easy: Stochastic optimization by pricing non-stochastic inputs. In *Proceedings of the 58th IEEE Annual Symposium on Foundations of Computer Science*. 540–551.
- Paul Dütting, Felix Fischer, and Max Klimm. 2018. Revenue gaps for static and dynamic posted pricing of homogeneous goods. arXiv:1607.07105.
- Tomer Ezra, Michal Feldman, Tim Roughgarden, and Warut Suksompong. 2018. Pricing multi-unit markets. In *Proceedings of the 14th Conference on Web and Internet Economics*. 140–153.
- Michal Feldman, Nick Gravin, and Brendan Lucier. 2015. Combinatorial auctions via posted prices. In *Proceedings of the 26th Annual ACM-SIAM Symposium on Discrete Algorithms*. 123–135.
- Eric J. Friedman, Ali Ghodsi, and Christos-Alexandros Psomas. 2014. Strategyproof allocation of discrete jobs on multiple machines. In *Proceedings of the 15th ACM Conference on Economics and Computation*. 529–546.
- Eric J. Friedman, Miklós Z. Rácz, and Scott Shenker. 2015. Dynamic budget-constrained pricing in the cloud. In *Proceedings of the 28th Canadian Conference on Artificial Intelligence*. 114–121.
- Darrell Hoy, Nicole Immorlica, and Brendan Lucier. 2016. On-demand or spot? Selling the cloud to risk-averse customers. In *Proceedings of the 12th International Conference on Web and Internet Economics*. 73–86.
- Navendu Jain, Ishai Menache, Joseph (Seffi) Naor, and Jonathan Yaniv. 2011. A truthful mechanism for value-based scheduling in cloud computing. In *Proceedings of the 4th International Symposium on Algorithmic Game Theory*. 178–189.
- Navendu Jain, Ishai Menache, Joseph (Seffi) Naor, and Jonathan Yaniv. 2012. Near-optimal scheduling mechanisms for deadline-sensitive jobs in large computing clusters. In *Proceedings of the 24th ACM Symposium on Parallelism in Algorithms and Architectures*. 255–266.
- Ian A. Kash and Peter Key. 2016. Pricing the cloud. *IEEE Internet Computing* 20, 1 (2016), 36–43.
- Brendan Lucier, Ishai Menache, Joseph (Seffi) Naor, and Jonathan Yaniv. 2013. Efficient online scheduling for deadline-sensitive jobs. In *Proceedings of the 25th ACM Symposium on Parallelism in Algorithms and Architectures*. 305–314.
- Changjun Wang, Weidong Ma, Tao Qin, Xujin Chen, Xiaodong Hu, and Tie-Yan Liu. 2015. Selling reserved instances in cloud computing. In *Proceedings of the 24th International Conference on Artificial Intelligence*. 224–230.
- Hong Zhang, Bo Li, Hongbo Jiang, Fangming Liu, Athanasios V. Vasilakos, and Jiangchuan Liu. 2013. A framework for truthful online auctions in cloud computing with heterogeneous user demands. In *Proceedings of IEEE INFOCOM 2013*. 1510–1518.

Received January 2018; revised January 2019; accepted April 2019