Approximating Geometric Knapsack via L-packings*

Waldo Gálvez*, Fabrizio Grandoni*, Sandy Heydrich**, Salvatore Ingala*, Arindam Khan***, and Andreas Wiese****

- *IDSIA, USI-SUPSI, Switzerland, [waldo, fabrizio, salvatore]@idsia.ch
- **MPI for Informatics and Saarbrücken Graduate School of Computer Science, Germany, heydrich@mpi-inf.mpg.de
- ***Department of Computer Science, Technical University of Munich, Munich, Germany, arindam.khan@in.tum.de
- *****Department of Industrial Engineering and Center for Mathematical Modeling, Universidad de Chile, Chile, awiese@dii.uchile.cl

Abstract

We study the two-dimensional geometric knapsack problem (2DK) in which we are given a set of n axis-aligned rectangular items, each one with an associated profit, and an axis-aligned square knapsack. The goal is to find a (non-overlapping) packing of a maximum profit subset of items inside the knapsack (without rotating items). The best-known polynomial-time approximation factor for this problem (even just in the cardinality case) is $2+\varepsilon$ [Jansen and Zhang, SODA 2004]. In this paper we break the 2 approximation barrier, achieving a polynomial-time $\frac{17}{9}+\varepsilon<1.89$ approximation, which improves to $\frac{558}{325}+\varepsilon<1.72$ in the cardinality case.

Essentially all prior work on 2DK approximation packs items inside a constant number of rectangular containers, where items inside each container are packed using a simple greedy strategy. We deviate for the first time from this setting: we show that there exists a large profit solution where items are packed inside a constant number of containers plus one L-shaped region at the boundary of the knapsack which contains items that are high and narrow and items that are wide and thin. The items of these two types possibly interact in a complex manner at the corner of the L.

The above structural result is not enough however: the best-known approximation ratio for the subproblem in the L-shaped region is $2+\varepsilon$ (obtained via a trivial reduction to one-dimensional knapsack by considering tall or wide items only). Indeed this is one of the simplest special settings of the problem for which this is the best known approximation factor. As a second major, and the main algorithmic contribution of this paper, we present a PTAS for this case. We believe that this will turn out to be useful in future work in geometric packing problems.

We also consider the variant of the problem with rotations (2DKR), where items can be rotated by 90 degrees. Also in this case the best-known polynomial-time approximation factor (even for the cardinality case) is $2 + \varepsilon$ [Jansen and Zhang, SODA 2004]. Exploiting part of the machinery developed for 2DK plus a few additional ideas, we obtain a polynomial-time $3/2 + \varepsilon$ -approximation for 2DKR, which improves to $4/3 + \varepsilon$ in the cardinality case.

^{*}The authors from IDSIA are partially supported by ERC Starting Grant NEWNET 279352 and SNSF Grant APXNET 200021_159697/1. Arindam Khan is supported in part by the European Research Council, Grant Agreement No. 691672, the work was primarily done when the author was at IDSIA. Sandy Heydrich is in part supported by the Google Europe PhD Fellowship.

1 Introduction

The (two-dimensional) geometric knapsack problem (2DK) is the geometric variant of the classical (one-dimensional) knapsack problem. We are given a set of n items $I = \{1, \ldots, n\}$, where each item $i \in I$ is an axis-aligned open rectangle $(0, w(i)) \times (0, h(i))$ in the two-dimensional plane, and has an associated profit p(i). Furthermore, we are given an axis-aligned square knapsack $K = [0, N] \times [0, N]$. W.l.o.g. we next assume that all values w(i), h(i), p(i) and N are positive integers. Our goal is to select a subset of items $OPT \subseteq I$ of maximum total profit $opt = p(OPT) := \sum_{i \in OPT} p(i)$ and to place them so that the selected rectangles are pairwise disjoint and fully contained in the knapsack. More formally, for each $i \in OPT$ we have to define a pair of coordinates (left(i), bottom(i)) that specify the position of the bottom-left corner of i in the packing. In other words, i is mapped into a rectangle $R(i) := (left(i), right(i)) \times (bottom(i), top(i))$, with right(i) = left(i) + w(i) and top(i) = bottom(i) + h(i). For any two $i, j \in OPT$, we must have $R(i) \subseteq K$ and $R(i) \cap R(j) = \emptyset$.

Besides being a natural mathematical problem, 2DK is well-motivated by practical applications. For instance, one might want to place advertisements on a board or a website, or cut rectangular pieces from a sheet of some material. Also, it models a scheduling setting where each rectangle corresponds to a job that needs some "consecutive amount" of a given resource (memory storage, frequencies, etc.). In all these cases, dealing with rectangular shapes only is a reasonable simplification and often the developed techniques can be extended to deal with more general instances.

2DK is NP-hard [29], and it was intensively studied from the point of view of approximation algorithms. The best known polynomial time approximation algorithm for it is due to Jansen and Zhang and yields a $(2+\varepsilon)$ -approximation [25]. This is the best known result even in the *cardinality* case (with all profits being 1). However, there are reasons to believe that much better polynomial time approximation ratios are possible: there is a QPTAS under the assumption that $N=n^{\mathrm{poly}(\log n)}$ [3], and there are PTASs if the profit of each item equals its area [4], if the size of the knapsack can be slightly increased (resource augmentation) [14, 21], if all items are relatively small [13] and if all input items are squares [22, 19]. Note that, with no restriction on N, the current best approximation for 2DK is $2+\varepsilon$ even in quasi-polynomial time¹.

All prior polynomial-time approximation algorithms for 2DK implicitly or explicitly exploit a *container-based* packing approach. The idea is to partition the knapsack into a constant number of axis-aligned rectangular regions (*containers*). The sizes (and therefore positions) of these containers can be *guessed* in polynomial time. Then items are packed inside the containers in a simple way: either one next to the other from left to right or from bottom to top (similarly to the one-dimensional case), or by means of the simple greedy Next-Fit-Decreasing-Height algorithm. Indeed, also the QPTAS in [3] can be cast in this framework, with the relevant difference that the number of containers in this case is poly-logarithmic (leading to a quasi-polynomial running time).

One of the major bottlenecks to achieve approximation factors better than 2 (in polynomial-time) is that items that are high and narrow (*vertical* items) and items that are wide and thin (*horizontal* items) can interact in a very complicated way. Indeed, consider the following seemingly simple L-packing problem: we are given a set of items i with either w(i) > N/2 (horizontal items) or h(i) > N/2 (vertical items). Our goal is to pack a maximum profit subset of them inside an L-shaped region $L = ([0, N] \times [0, h_L]) \cup ([0, w_L] \times [0, N])$, so that horizontal (resp., vertical) items are packed in the bottom-right (resp., top-left) of L. To the best of our knowledge, the best-known approximation ratio for L-packing is $2 + \varepsilon$: Remove either all vertical or all horizontal items, and then pack the remaining items by a simple reduction to one-

¹The role of N in the running time is delicate, as shown by recent results on the related *strip packing* problem [1, 16, 18, 20, 30].

dimensional knapsack (for which an FPTAS is known). It is unclear whether a container-based packing can achieve a better approximation factor, and we conjecture that this is not the case. As we will see, a better understanding of L-packing will play a major role in the design of improved approximation algorithms for 2DK.

1.1 Our contribution

In this paper we break the 2-approximation barrier for 2DK. In order to do that, we substantially deviate for the first time from *pure* container-based packings, which are, either implicitly or explicitly, at the hearth of prior work. Namely, we consider L&C-packings that combine $O_{\varepsilon}(1)$ containers *plus* one L-packing of the above type (see Fig.1.(a)), and show that one such packing has large enough profit.

While it is easy to pack almost optimally items into containers, the mentioned $2 + \varepsilon$ approximation for L-packings is not sufficient to achieve altogether a better than 2 approximation factor: indeed, the items of the L-packing might carry all the profit! The main algorithmic contribution of this paper is a PTAS for the L-packing problem. It is easy to solve this problem optimally in pseudo-polynomial time $(Nn)^{O(1)}$ by means of dynamic programming. We show that a $1 + \varepsilon$ approximation can be obtained by restricting the top (resp., right) coordinates of horizontal (resp., vertical) items to a proper set that can be computed in polynomial time $n^{O_{\varepsilon}(1)}$. Given that, one can adapt the above dynamic program to run in polynomial time.

Theorem 1. There is a PTAS for the L-packing problem.

In order to illustrate the power of our approach, we next sketch a simple $\frac{16}{9} + O(\varepsilon)$ approximation for the cardinality case of 2DK (details in Section 3). By standard arguments² it is possible to discard *large* items with both sides longer than $\varepsilon \cdot N$. The remaining items have height or width smaller than $\varepsilon \cdot N$ (horizontal and vertical items, resp.). Let us delete all items intersecting a random vertical or horizontal strip of width $\varepsilon \cdot N$ inside the knapsack. We can pack the remaining items into $O_{\varepsilon}(1)$ containers by exploiting the PTAS under one-dimensional resource augmentation for 2DK in $[21]^3$. A vertical strip deletes vertical items with $O(\varepsilon)$ probability, and horizontal ones with probability roughly proportional to their width, and symmetrically for a horizontal strip. In particular, let us call *long* the items with longer side larger than N/2, and short the remaining items. Then the above argument gives in expectation roughly one half of the profit opt_{long} of long items, and three quarters of the profit opt_{short} of short ones. This is already good enough unless opt_{long} is large compared to opt_{short} .

At this point L-packings and our PTAS come into play. We shift long items such that they form 4 stacks at the sides of the knapsack in a ring-shaped region, see Fig.1.(b)-(c): this is possible since any vertical long item cannot have a horizontal long item both at its left and at its right, and vice versa. Next we delete the least profitable of these stacks and rearrange the remaining long items into an L-packing, see Fig.1.(d). Thus using our PTAS for L-packings, we can compute a solution of profit roughly three quarters of opt_{long} . The reader might check that the combination of these two algorithms gives the claimed approximation factor.

Above we used either $O_{\varepsilon}(1)$ containers or one L-packing: by combining the two approaches together and with a more sophisticated case analysis we achieve the following result (see Section B).

Theorem 2. There is a polynomial-time $\frac{558}{325} + \varepsilon < 1.72$ approximation algorithm for cardinality 2DK.

²There can be at most $O_{\varepsilon}(1)$ such items in any feasible solution, and if the optimum solution contains only $O_{\varepsilon}(1)$ items we can solve the problem optimally by brute force.

³Technically this PTAS is not container-based, however in Section F we show that it can be cast in that framework. Our version of the PTAS simplifies the algorithms and works also in the case with rotations: this might be a handy black-box tool.




Figure 1: (a) An L&C-packing with 4 containers, where the top-left container is packed by means of Next-Fit-Decreasing-Height. (b) A subset of long items. (c) Such items are shifted into 4 stacks at the sides of the knapsack, and the top stack is deleted. (d) The final packing into an L-shaped region.

For weighted 2DK we face severe technical complications for proving that there is a profitable L&C-packing. One key reason is that in the weighted case we cannot discard large items since even one such item might contribute a large fraction to the optimal profit. In order to circumvent these difficulties, we exploit the corridor-partition at the hearth of the QPTAS for 2DK in [3] (in turn inspired by prior work in [2]). Roughly speaking, there exists a partition of the knapsack into $O_{\varepsilon}(1)$ corridors, consisting of the concatenation of $O_{\varepsilon}(1)$ (partially overlapping) rectangular regions (subcorridors). In [3] the authors partition the corridors into a poly-logarithmic number of containers. Their main algorithm then guesses these containers in time $n^{\text{poly}(\log n)}$. However, we can only handle a constant number of containers in polynomial time. Therefore, we present a different way to partition the corridors into containers: here we lose the profit of a set of thin items, which in some sense play the role of long items in the previous discussion. These thin items fit in a very narrow ring at the boundary of the knapsack and we map them to an L-packing in the same way as in the cardinality case above. Some of the remaining non-thin items are then packed into $O_{\varepsilon}(1)$ containers that are placed in the (large) part of the knapsack not occupied by the L-packing. Our partition of the corridors is based on a somewhat intricate case analysis that exploits the fact that long consecutive subcorridors are arranged in the shape of rings or spirals: this is used to show the existence of a profitable L&C-packing.

Theorem 3. There is a polynomial-time $\frac{17}{9} + \varepsilon < 1.89$ approximation algorithm for (weighted) 2DK.

Rotation setting. In the variant of 2DK with rotations (2DKR), we are allowed to rotate any rectangle i by 90 degrees. This means that i can also be placed in the knapsack as a rectangle of the form $(left(i), left(i) + h(i)) \times (bottom(i), bottom(i) + w(i))$. The best known polynomial time approximation factor for 2DKR (even for the cardinality case) is again $2 + \varepsilon$ due to [25] and the mentioned QPTAS in [3] works also for this case.

By using the techniques described above and exploiting a few more ideas, we are also able to improve the approximation factor for 2DKR (see Sections C and D for the cardinality and general case, resp.). The basic idea is that any thin item can now be packed inside a narrow vertical strip (say at the right edge of the knapsack) by possibly rotating it. This way we do not lose one quarter of the profit due to the mapping to an L-packing and instead place all items from the ring into the mentioned strip (while we ensure that their total width is small). The remaining short items are packed by means of a novel resource contraction lemma: unless there is one huge item that occupies almost the whole knapsack (a case that we consider separately), we can pack almost one half of the profit of non-thin items in a reduced knapsack where one of the two sides is shortened by a factor $1 - \varepsilon$ (hence leaving enough space for the vertical strip). We remark that

here we heavily exploit the possibility to rotate items. Thus, roughly speaking, we obtain either all profit of non-thin items, or all profit of thin items plus one half of the profit of non-thin items: this gives a $3/2 + \varepsilon$ approximation. A further refinement of this approach yields a $4/3 + \varepsilon$ approximation in the cardinality case. We remark that, while resource augmentation is a well-established notion in approximation algorithms, resource contraction seems to be a rather novel direction to explore.

Theorem 4. For any constant $\varepsilon > 0$, there exists a polynomial-time $\frac{3}{2} + \varepsilon$ approximation algorithm for 2DKR. In the cardinality case the approximation factor can be improved to $\frac{4}{3} + \varepsilon$.

1.2 Other related work

The mentioned $(2+\varepsilon)$ -approximation for two-dimensional knapsack [25] works in the weighted case of the problem. However, in the unweighted case a simpler $(2+\varepsilon)$ -approximation is known [24]. If one can increase the size of the knapsack by a factor $1+\varepsilon$ in both dimensions then one can compute a solution of optimal weight, rather than an approximation, in time $f(1/\varepsilon) \cdot n^{O(1)}$ where the exponent of n does not depend on ε [19] (for some suitable function f). Similarly, for the case of squares there is a $(1+\varepsilon)$ -approximation algorithm known with such a running time, i.e., an EPTAS [19]. This improves previous results such as a $(5/4+\varepsilon)$ -approximation [17] and the mentioned PTAS [22]. Two-dimensional knapsack is the separation problem when we want to solve the configuration-LP for two-dimensional bin-packing. Even though we do not have a PTAS for the former problem, Bansal et al. [4] show how to solve the latter LP to an $(1+\varepsilon)$ -accuracy using their PTAS for two-dimensional knapsack for the special case where the profit of each item equals its area. The best known (asymptotic) result for two-dimensional bin packing is due to Bansal and Khan and it is based on this configuration-LP, achieving an approximation ratio of 1.405 [6] which improves a series of previous results [21, 5, 7, 26, 10]. See also the recent survey in [9] and [27].

2 A PTAS for L-packings

In this section we present a PTAS for the problem of finding an optimal L-packing. In this problem we are given a set of horizontal items I_{hor} with width larger than N/2, and a set of vertical items I_{ver} with height larger than N/2. Furthermore, we are given an L-shaped region $L = ([0,N] \times [0,h_L]) \cup ([0,w_L] \times [0,N])$. Our goal is to pack a subset $OPT \subseteq I := I_{hor} \cup I_{ver}$ of maximum total profit $opt = p(OPT) := \sum_{i \in OPT} p(i)$, such that $OPT_{hor} := OPT \cap I_{hor}$ is packed inside the $horizontal\ box\ [0,N] \times [0,h_L]$ and $OPT_{ver} := OPT \cap I_{ver}$ is packed inside the $vertical\ box\ [0,w_L] \times [0,N]$. We remark that packing horizontal and vertical items independently is not possible due to the possible overlaps in the intersection of the two boxes: this is what makes this problem non-trivial, in particular harder than standard (one-dimensional) knapsack.

Observe that in an optimal packing we can assume w.l.o.g. that items in OPT_{hor} are pushed as far to the right/bottom as possible. Furthermore, the items in OPT_{hor} are packed from bottom to top in non-increasing order of width. Indeed, it is possible to *permute* any two items violating this property while keeping the packing feasible. A symmetric claim holds for OPT_{ver} . See Fig. 1.(d) for an illustration.

Given the above structure, it is relatively easy to define a dynamic program (DP) that computes an optimal L-packing in pseudo-polynomial time $(Nn)^{O(1)}$. The basic idea is to scan items of I_{hor} (resp. I_{ver}) in decreasing order of width (resp., height), and each time *guess* if they are part of the optimal solution OPT. At each step either both the considered horizontal item i and vertical item j are not part of the

optimal solution, or there exist a guillotine cut^4 separating i or j from the rest of OPT. Depending on the cases, one can define a smaller L-packing sub-instance (among N^2 choices) for which the DP table already contains a solution.

In order to achieve a $(1+\varepsilon)$ -approximation in polynomial time $n^{O_\varepsilon(1)}$, we show that it is possible (with a small loss in the profit) to restrict the possible top coordinates of OPT_{hor} and right coordinates of OPT_{ver} to proper polynomial-size subsets $\mathcal T$ and $\mathcal R$, resp. We call such an L-packing $(\mathcal T,\mathcal R)$ -restricted. By adapting the above DP one obtains:

Lemma 5. An optimal $(\mathcal{T}, \mathcal{R})$ -restricted L-packing can be computed in time polynomial in $m := n + |\mathcal{T}| + |\mathcal{R}|$ using dynamic programming.

Proof. For notational convenience we assume $0 \in \mathcal{T}$ and $0 \in \mathcal{R}$. Let $h_1, \ldots, h_{n(h)}$ be the items in I_{hor} in decreasing order of width and $v_1, \ldots, v_{n(v)}$ be the items in I_{ver} in decreasing order of height (breaking ties arbitrarily). For $w \in [0, w_L]$ and $h \in [0, h_L]$, let $L(w, h) = ([0, w] \times [0, N]) \cup ([0, N] \times [0, h]) \subseteq L$. Let also $\Delta L(w, h) = ([w, w_L] \times [h, N]) \cup ([w, N] \times [h, h_L]) \subseteq L$. Note that $L = L(w, h) \cup \Delta L(w, h)$.

We define a dynamic program table DP indexed by $i \in [1, n(h)]$ and $j \in [1, n(v)]$, by a top coordinate $t \in \mathcal{T}$, and a right coordinate $r \in \mathcal{R}$. The value of DP(i, t, j, r) is the maximum profit of a $(\mathcal{T}, \mathcal{R})$ -restricted packing of a subset of $\{h_i, \ldots, h_{n(h)}\} \cup \{v_j, \ldots, v_{n(v)}\}$ inside $\Delta L(r, t)$. The value of DP(1, 0, 1, 0) is the value of the optimum solution we are searching for. Note that the number of table entries is upper bounded by m^4 .

We fill in DP according to the partial order induced by vectors (i, t, j, r), processing larger vectors first. The base cases are given by (i, j) = (n(h) + 1, n(v) + 1) and $(r, t) = (w_L, h_L)$, in which case the table entry has value 0.

In order to compute any other table entry DP(i, t, j, r), with optimal solution OPT', we take the maximum of the following few values:

- If $i \le n(h)$, the value DP(i+1,t,j,r). This covers the case that $h_i \notin OPT'$;
- If $j \leq n(v)$, the value DP(i, t, j + 1, r). This covers the case that $v_i \notin OPT'$;
- Assume that there exists $t' \in \mathcal{T}$ such that $t' h(h_i) \geq t$ and that $w(h_i) \leq N r$. Then for the minimum such t' we consider the value $p(h_i) + DP(i+1,t',j,r)$. This covers the case that $h_i \in OPT'$, and there exists a (horizontal) guillotine cut separating h_i from $OPT' \setminus \{h_i\}$.
- Assume that there exists $r' \in \mathcal{R}$ such that $r' w(v_j) \geq r$ and that $h(v_j) \leq N t$. Then for the minimum such r' we consider the value $p(v_j) + DP(i,t,j+1,r')$. This covers the case that $v_j \in OPT'$, and there exists a (vertical) guillotine cut separating v_j from $OPT' \setminus \{v_j\}$.

We observe that the above cases (which can be explored in polynomial time) cover all the possible configurations in OPT'. Indeed, if the first two cases do not apply, we have that $h_i, v_j \in OPT'$. Then either the line containing the right side of v_j does not intersect h_i (hence any other item in OPT') or the line containing the top side of h_i does not intersect v_j (hence any other item in OPT'). Indeed, the only remaining case is that v_j and h_i overlap, which is impossible since they both belong to OPT'.

We will show that there exists a $(\mathcal{T}, \mathcal{R})$ -restricted L-packing with the desired properties.

Lemma 6. There exists a $(\mathcal{T}, \mathcal{R})$ -restricted L-packing solution of profit at least $(1 - 2\varepsilon)$ opt, where the sets \mathcal{T} and \mathcal{R} have cardinality at most $n^{O(1/\varepsilon^{1/\varepsilon})}$ and can be computed in polynomial time based on the input (without knowing OPT).

 $^{^4}$ A guillotine cut is an infinite, axis-parallel line ℓ that partitions the items in a given packing in two subsets without intersecting any item.




Figure 2: Illustration of the delete&shift procedure with $r_{hor}=2$. The dashed lines indicate the value of the new baselines in the different stages of the algorithm. (Left) The starting packing. Dark and light grey items denote the growing sequences for the calls with r=2 and r=1, resp. (Middle) The shift of items at the end of the recursive calls with r=1. Note that light grey items are all deleted, and dark grey items are not shifted. (Right) The shift of items at the end of the process. Here we assume that the middle dark grey item is deleted.

Lemmas 5 and 6 together immediately imply a PTAS for L-packings (showing Theorem 1). The rest of this section is devoted to the proof of Lemma 6.

We will describe a way to delete a subset of items $D_{hor} \subseteq OPT_{hor}$ with $p(D_{hor}) \le 2\varepsilon p(OPT_{hor})$, and shift down the remaining items $OPT_{hor} \setminus D_{hor}$ so that their top coordinate belongs to a set \mathcal{T} with the desired properties. Symmetrically, we will delete a subset of items $D_{ver} \subseteq OPT_{ver}$ with $p(D_{ver}) \le 2\varepsilon p(OPT_{ver})$, and shift to the left the remaining items $OPT_{ver} \setminus D_{ver}$ so that their right coordinate belongs to a set \mathcal{R} with the desired properties. We remark that shifting down (resp. to the left) items of OPT_{hor} (resp., OPT_{ver}) cannot create any overlap with items of OPT_{ver} (resp., OPT_{hor}). This allows us to reason on each such set separately.

We next focus on OPT_{hor} only: the construction for OPT_{ver} is symmetric. For notational convenience we let $1, \ldots, n_{hor}$ be the items of OPT_{hor} in non-increasing order of width *and* from bottom to top in the starting optimal packing. We remark that this sequence is not necessarily sorted (increasingly or decreasingly) in terms of item heights: this makes our construction much more complicated.

Let us first introduce some useful notation. Consider any subsequence $B = \{b_{start}, \dots, b_{end}\}$ of consecutive items (interval). For any $i \in B$, we define $top_B(i) := \sum_{k \in B, k \le i} h(k)$ and $bottom_B(i) = top_B(i) - h(i)$. The growing subsequence $G = G(B) = \{g_1, \dots, g_h\}$ of B (with possibly non-contiguous items) is defined as follows. We initially set $g_1 = b_{start}$. Given the item g_i, g_{i+1} is the smallest-index (i.e., lowest) item in $\{g_i + 1, \dots, b_{end}\}$ such that $h(g_{i+1}) \ge h(g_i)$. We halt the construction of G when we cannot find a proper g_{i+1} . For notational convenience, define $g_{h+1} = b_{end} + 1$. We let $B_i^G := \{g_i + 1, \dots, g_{i+1} - 1\}$ for $i = 1, \dots, h$. Observe that the sets B_i^G partition $B \setminus G$. We will crucially exploit the following simple property.

Lemma 7. For any $g_i \in G$ and any $j \in \{b_{start}, \dots, g_{i+1} - 1\}$, $h(j) \leq h(g_i)$.

Proof. The items $j \in B_i^G = \{g_i + 1, \dots, g_{i+1} - 1\}$ have $h(j) < h(g_i)$. Indeed, any such j with $h(j) \ge h(g_i)$ would have been added to G, a contradiction.

Consider next any $j \in \{b_{start}, \dots g_i - 1\}$. If $j \in G$ the claim is trivially true by construction of G. Otherwise, one has $j \in B_k^G$ for some $g_k \in G$, $g_k < g_i$. Hence, by the previous argument and by construction of G, $h(j) < h(g_k) \le h(g_i)$.

The intuition behind our construction is as follows. Consider the growing sequence $G = G(OPT_{hor})$, and suppose that $p(G) \leq \varepsilon \cdot p(OPT_{hor})$. Then we might simply delete G, and shift the remaining items $OPT_{hor} \setminus G = \bigcup_i B_i^G$ as follows. Let $[x]_y$ denote the smallest multiple of y not smaller than x. We consider each set B_i^G separately. For each such set, we define a baseline vertical coordinate $base_j =$ $\lceil bottom(g_j) \rceil_{h(g_j)/2}$, where $bottom(g_j)$ is the bottom coordinate of g_j in the original packing. We next round up the height of $i \in B_i^G$ to $\hat{h}(i) := \lceil h(i) \rceil_{h(g_i)/(2n)}$, and pack the rounded items of B_i^G as low as possible above the baseline. The reader might check that the possible top coordinates for rounded items fall in a polynomial size set (using Lemma 7). It is also not hard to check that items are *not* shifted up.

We use recursion in order to handle the case $p(G) > \varepsilon \cdot p(OPT_{hor})$. Rather than deleting G, we consider each B_i^G and build a new growing subsequence for each such set. We repeat the process recursively for r_{hor} many rounds. Let \mathcal{G}^r be the union of all the growing subsequences in the recursive calls of level r. Since the sets \mathcal{G}^r are disjoint by construction, there must exist a value $r_{hor} \leq \frac{1}{\varepsilon}$ such that $p(\mathcal{G}^{r_{hor}}) \leq$ $\varepsilon \cdot p(OPT_{hor})$. Therefore we can apply the same shifting argument to all growing subsequences of level r_{hor} (in particular we delete all of them). In the remaining growing subsequences we can afford to delete 1 out of $1/\varepsilon$ consecutive items (with a small loss of the profit), and then apply a similar shifting argument.

We next describe our approach in more detail. We exploit a recursive procedure delete&shift. This procedure takes as input two parameters: an interval $B = \{b_{start}, \dots, b_{end}\}$, and an integer round parameter $r \geq 1$. Procedure delete& shift returns a set $D(B) \subseteq B$ of deleted items, and a shift function $shift: B \setminus D(B) \to \mathbb{N}$. Intuitively, shift(i) is the value of the top coordinate of i in the shifted packing w.r.t. a proper baseline value which is implicitly defined. We initially call delete&shift(OPT_{hor}, r_{hor}), for a proper $r_{hor} \in \{1, \dots, \frac{1}{\varepsilon}\}$ to be fixed later. Let (D, shift) be the output of this call. The desired set of deleted items is $D_{hor} = D$, and in the final packing top(i) = shift(i) for any $i \in OPT_{hor} \setminus D_{hor}$ (the right coordinate of any such i is N).

The procedure behaves differently in the cases r=1 and r>1. If r=1, we compute the growing sequence $G = G(B) = \{g_1 = b_{start}, \dots, g_h\}$, and set D(B) = G(B). Consider any set $B_i^G = \{g_j + g_j\}$ $1, \ldots, g_{j+1} - 1\}, j = 1, \ldots, h$. Let $base_j := \lceil bottom_B(g_j) \rceil_{h(g_j)/2}$. We define for any $i \in B_j^{G'}$,

$$shift(i) = base_j + \sum_{k \in B_j^G, k \le i} \lceil h(k) \rceil_{h(g_j)/(2n)}.$$

Observe that shift is fully defined since $\bigcup_{j=1}^h B_j^G = B \setminus D(B)$. If instead r > 1, we compute the growing sequence $G = G(B) = \{g_1 = b_{start}, \dots, g_h\}$. We next delete a subset of items $D' \subseteq G$. If $h < \frac{1}{\varepsilon}$, we let $D' = D'(B) = \emptyset$. Otherwise, let $G_k = \{g_j \in G : g_j \in G : g_j$ $j=k \pmod{1/\varepsilon}$ $\subseteq G$, for $k \in \{0,\ldots,1/\varepsilon-1\}$. We set $D'=D'(B)=\{d_1,\ldots,d_p\}=G_x$ where $x = \arg\min_{k \in \{0, \dots, 1/\varepsilon - 1\}} p(G_k).$

Proposition 8. One has $p(D') \leq \varepsilon \cdot p(G)$. Furthermore, any subsequence $\{g_x, g_{x+1}, \dots, g_y\}$ of G with at least $1/\varepsilon$ items contains at least one item from D'.

Consider each set $B_j^G = \{g_j+1,\ldots,g_{j+1}-1\}, j=1,\ldots,h$: We run delete&shift $(B_j^G,r-1)$. Let $(D_j,shift_j)$ be the output of the latter procedure, and $shift_j^{max}$ be the maximum value of $shift_j$. We set the output set of deleted items to $D(B) = D' \cup (\bigcup_{i=1}^{h} D_i)$.

It remains to define the function shift. Consider any set B_i^G , and let d_q be the deleted item in D' with largest index (hence in topmost position) in $\{b_{start}, \ldots, g_j\}$: define $base_q = \lceil bottom_B(d_q) \rceil_{h(d_q)/2}$. If there is no such d_q , we let $d_q=0$ and $base_q=0$. For any $i\in B_j^G$ we set:

$$shift(i) = base_q + \sum_{g_k \in G, d_q < g_k \le g_j} h(g_k) + \sum_{g_k \in G, d_q \le g_k < g_j} shift_k^{max} + shift_j(i).$$

Analogously, if $g_j \neq d_q$, we set

$$shift(g_j) = base_q + \sum_{g_k \in G, d_q < g_k \le g_j} h(g_k) + \sum_{g_k \in G, d_q \le g_k < g_j} shift_k^{max}.$$

This concludes the description of delete&shift. We next show that the final packing has the desired properties. Next lemma shows that the total profit of deleted items is small for a proper choice of the starting round parameter r_{hor} .

Lemma 9. There is a choice of $r_{hor} \in \{1, \dots, \frac{1}{\varepsilon}\}$ such that the final set D_{hor} of deleted items satisfies $p(D_{hor}) \leq 2\varepsilon \cdot p(OPT_{hor})$.

Proof. Let \mathcal{G}^r denote the union of the sets G(B) computed by all the recursive calls with input round parameter r. Observe that by construction these sets are disjoint. Let also \mathcal{D}^r be the union of the sets D'(B) on those calls (the union of sets D(B) for $r = r_{hor}$). By Proposition 8 and the disjointness of sets \mathcal{G}^r one has

$$p(D_{hor}) = \sum_{1 \le r \le r_{hor}} p(\mathcal{D}^r)$$

$$\le \varepsilon \cdot \sum_{r < r_{hor}} p(\mathcal{G}^r) + p(\mathcal{D}^{r_{hor}})$$

$$\le \varepsilon \cdot p(OPT_{hor}) + p(\mathcal{D}^{r_{hor}}).$$

Again by the disjointness of sets \mathcal{G}^r (hence \mathcal{D}^r), there must exists a value of $r_{hor} \in \{1, \dots, \frac{1}{\varepsilon}\}$ such that $p(\mathcal{D}^{r_{hor}}) \leq \varepsilon \cdot p(OPT_{hor})$. The claim follows.

Next lemma shows that, intuitively, items are only shifted down w.r.t. the initial packing.

Lemma 10. Let (D, shift) be the output of some execution of delete&shift(B, r). Then, for any $i \in B \setminus D$, $shift(i) \le top_B(i)$.

Proof. We prove the claim by induction on r. Consider first the case r=1. In this case, for any $i \in B_j^G$:

$$\begin{split} shift(i) \\ = \lceil bottom_B(g_j) \rceil_{h(g_j)/2} + \sum_{k \in B_j^G, k \le i} \lceil h(k) \rceil_{h(g_j)/(2n)} \\ \le top_B(g_j) - \frac{1}{2}h(g_j) + \sum_{k \in B_j^G, k \le i} h(k) + n \cdot \frac{h(g_j)}{2n} \\ = top_B(i). \end{split}$$

Assume next that the claim holds up to round parameter $r-1 \geq 1$, and consider round r. For any $i \in B_j^G$

with $base_q = \lceil bottom_B(d_q) \rceil_{h(d_q)/2}$, one has

$$shift(i)$$

$$=\lceil bottom_B(d_q) \rceil_{h(d_q)/2} + \sum_{g_k \in G, d_q < g_k \le g_j} h(g_k)$$

$$+ \sum_{g_k \in G, d_q \le g_k < g_j} shift_k^{max} + shift_j(i)$$

$$\le top_B(d_q) + \sum_{g_k \in G, d_q < g_k \le g_j} h(g_k)$$

$$+ \sum_{g_k \in G, d_q \le g_k < g_j} top_{B_k^G}(g_{k+1} - 1) + top_{B_j^G}(i)$$

$$= top_B(i).$$

An analogous chain of inequalities shows that $shift(g_j) \leq top_B(g_j)$ for any $g_j \in G \setminus D'$. A similar proof works for the special case $base_q = 0$.

It remains to show that the final set of values of top(i) = shift(i) has the desired properties. This is the most delicate part of our analysis. We define a set \mathcal{T}^r of candidate top coordinates recursively in r. Set \mathcal{T}^1 contains, for any item $j \in I_{hor}$, and any integer $1 \leq a \leq 4n^2$, the value $a \cdot \frac{h(j)}{2n}$. Set \mathcal{T}^r , for r>1 is defined recursively w.r.t. to \mathcal{T}^{r-1} . For any item j, any integer $0 \leq a \leq 2n-1$, any tuple of $b \leq 1/\varepsilon-1$ items $j(1),\ldots,j(b)$, and any tuple of $c \leq 1/\varepsilon$ values $s(1),\ldots,s(c) \in \mathcal{T}^{r-1}$, \mathcal{T}^r contains the sum $a \cdot \frac{h(j)}{2} + \sum_{k=1}^b h(j(k)) + \sum_{k=1}^c s(k)$. Note that sets \mathcal{T}^r can be computed based on the input only (without knowing OPT). It is easy to show that \mathcal{T}^r has polynomial size for $r = O_\varepsilon(1)$.

Lemma 11. For any integer
$$r \geq 1$$
, $|\mathcal{T}^r| \leq (2n)^{\frac{r+2+(r-1)\varepsilon}{\varepsilon^{r-1}}}$.

Proof. We prove the claim by induction on r. The claim is trivially true for r=1 since there are n choices for item j and $4n^2$ choices for the integer a, hence altogether at most $n \cdot 4n^2 < 8n^3$ choices. For r > 1, the number of possible values of \mathcal{T}^r is at most

$$n \cdot 2n \cdot \left(\sum_{b=0}^{1/\varepsilon - 1} n^b\right) \cdot \left(\sum_{c=0}^{1/\varepsilon} |\mathcal{T}^{r-1}|^c\right) \le 4n^2 \cdot n^{\frac{1}{\varepsilon} - 1} \cdot |\mathcal{T}^{r-1}|^{\frac{1}{\varepsilon}}$$
$$\le (2n)^{\frac{1}{\varepsilon} + 1} \left((2n)^{\frac{r+1+(r-2)\varepsilon}{\varepsilon^{r-2}}} \right)^{\frac{1}{\varepsilon}} \le (2n)^{\frac{r+2+(r-1)\varepsilon}{\varepsilon^{r-1}}}.$$

Next lemma shows that the values of shift returned by delete&shift for round parameter r belong to \mathcal{T}^r , hence the final top coordinates belong to $\mathcal{T}:=\mathcal{T}^{r_{hor}}$.

Lemma 12. Let (D, shift) be the output of some execution of delete&shift(B, r). Then, for any $i \in B \setminus D$, $shift(i) \in \mathcal{T}^r$.

Proof. We prove the claim by induction on r. For the case r=1, recall that for any $i \in B_i^G$ one has

$$shift(i) = \lceil bottom_B(g_j) \rceil_{h(g_j)/2}$$

$$+ \sum_{k \in B_j^G, k \le i} \lceil h(k) \rceil_{h(g_j)/(2n)}.$$

By Lemma 7, $bottom_B(g_j) = \sum_{k \in B, k < g_j} h(k) \le (n-1) \cdot h(g_j)$. By the same lemma, $\sum_{k \in B_j^G, k \le i} h(k) \le (n-1) \cdot h(g_j)$. It follows that

$$shift(i) \le 2(n-1) \cdot h(g_j) + \frac{h(g_j)}{2} + (n-1) \cdot \frac{h(g_j)}{2n}$$

 $\le 4n^2 \cdot \frac{h(g_j)}{2n}.$

Hence $shift(i) = a \cdot \frac{h(g_j)}{2n}$ for some integer $1 \le a \le 4n^2$, and $shift(i) \in \mathcal{T}^1$ for $j = g_j$ and for a proper choice of a.

Assume next that the claim is true up to $r-1 \ge 1$, and consider the case r. Consider any $i \in B_j^G$, and assume $0 < base_q = \lceil bottom_B(d_q) \rceil_{h(d_q)/2}$. One has:

$$shift(i) = \lceil bottom_B(d_q) \rceil_{h(d_q)/2} + \sum_{g_k \in G, d_q < g_k \le g_j} h(g_k)$$

$$+ \sum_{g_k \in G, d_q \le g_k < g_j} shift_k^{max} + shift_j(i).$$

By Lemma 7, $bottom_B(d_q) \leq (n-1)h(d_q)$, therefore $\lceil bottom_B(d_q) \rceil_{h(d_q)/2} = a \cdot \frac{h(d_q)}{2}$ for some integer $1 \leq a \leq 2(n-1)+1$. By Proposition 8, $|\{g_k \in G, d_q < g_k \leq g_j\}| \leq 1/\varepsilon-1$. Hence $\sum_{g_k \in G, d_q < g_k \leq g_j} h(g_k)$ is a value contained in the set of sums of $b \leq 1/\varepsilon-1$ item heights. By inductive hypothesis $shift_k^{max}$, $shift_j(i) \in \mathcal{T}^{r-1}$. Hence by a similar argument the value of $\sum_{g_k \in G, d_q \leq g_k < g_j} shift_k^{max} + shift_j(i)$ is contained in the set of sums of $c \leq 1/\varepsilon-1+1$ values taken from \mathcal{T}^{r-1} . Altogether, $shift(i) \in \mathcal{T}^r$. A similar argument, without the term $shift_j(i)$, shows that $shift(g_j) \in \mathcal{T}^r$ for any $g_j \in G \setminus D'$. The proof works similarly in the case $base_q = 0$ by setting a = 0. The claim follows. \square

Proof of Lemma 6. We apply the procedure delete&shift to OPT_{hor} as described before, and a symmetric procedure to OPT_{ver} . In particular the latter procedure computes a set $D_{ver} \subseteq OPT_{ver}$ of deleted items, and the remaining items are shifted to the left so that their right coordinate belongs to a set $\mathcal{R} := \mathcal{R}^{r_{ver}}$, defined analogously to the case of $\mathcal{T} := \mathcal{T}^{r_{hor}}$, for some integer $r_{ver} \in \{1, \dots, 1/\varepsilon\}$ (possibly different from r_{hor} , though by averaging this is not critical).

It is easy to see that the profit of non-deleted items satisfies the claim by Lemma 9 and its symmetric version. Similarly, the sets \mathcal{T} and \mathcal{R} satisfy the claim by Lemmas 11 and 12, and their symmetric versions. Finally, w.r.t. the original packing non-deleted items in OPT_{hor} and OPT_{ver} can be only shifted to the bottom and to the left, resp., by Lemma 10 and its symmetric version. This implies that the overall packing is feasible.

3 A Simple Improved Approximation for Cardinality 2DK

In this section we present a simple improved approximation for the cardinality case of 2DK. We can assume that the optimal solution $OPT \subseteq I$ satisfies that $|OPT| \ge 1/\varepsilon^3$ since otherwise we can solve the problem optimally by brute force in time $n^{O(1/\varepsilon^3)}$. Therefore, we can discard from the input all *large* items with both sides larger than $\varepsilon \cdot N$: any feasible solution can contain at most $1/\varepsilon^2$ such items, and discarding them decreases the cardinality of OPT at most by a factor $1 + \varepsilon$. Let OPT denote this slightly sub-optimal solution obtained by removing large items.

We will need the following technical lemma, that holds also in the weighted case (see also Fig.1.(b)-(d)).

Lemma 13. Let H and V be given subsets of items from some feasible solution with width and height strictly larger than N/2, resp. Let h_H and w_V be the total height and width of items of H and V, resp. Then there exists an L-packing of a set $APX \subseteq H \cup V$ with $p(APX) \ge \frac{3}{4}(p(H) + p(V))$ into the area $L = ([0, N] \times [0, h_H]) \cup ([0, w_V] \times [0, N])$.

Proof. Let us consider the packing of $H \cup V$. Consider each $i \in H$ that has no $j \in V$ to its top (resp., to its bottom) and shift it up (resp. down) until it hits another $i' \in H$ or the top (resp, bottom) side of the knapsack. Note that, since h(j) > N/2 for any $j \in V$, one of the two cases above always applies. We iterate this process as long as possible to move any such i. We perform a symmetric process on V. At the end of the process all items in $H \cup V$ are stacked on the 4 sides of the knapsack⁵.

Next we remove the least profitable of the 4 stacks: by a simple permutation argument we can guarantee that this is the top or right stack. We next discuss the case that it is the top one, the other case being symmetric. We show how to repack the remaining items in a boundary L of the desired size by permuting items in a proper order. In more detail, suppose that the items packed on the left (resp., right and bottom) have a total width of w_l (resp., total width of w_r and total height of h_b). We next show that there exists a packing into $L' = ([0, N] \times [0, h_b]) \cup ([0, w_l + w_r] \times [0, N])$. We prove the claim by induction. Suppose that we have proved it for all packings into left, right and bottom stacks with parameters w_l' , w_r' , and h' such that $h' < h_b$ or $w_l' + w_r' < w_l + w_r$ or $w_l' + w_r' = w_l + w_r$ and $w_r' < w_r$.

In the considered packing we can always find a guillotine cut ℓ , such that one side of the cut contains precisely one *lonely* item among the leftmost, rightmost and bottommost items. Let ℓ be such a cut. First assume that the lonely item j is the bottommost one. Then by induction the claim is true for the part above ℓ since the part of the packing above ℓ has parameters w_{ℓ} , w_{r} , and h - h(j). Thus, it is also true for the entire packing. A similar argument applies if the lonely item j is the leftmost one.

It remains to consider the case that the lonely item j is the rightmost one. We remove j temporarily and move *all* other items by w(j) to the right. Then we insert j at the left (in the space freed by the previous shifting). By induction, the claim is true for the resulting packing since it has parameters $w_l + w(j)$, $w_r - w(j)$, and h, resp.

For our algorithm, we consider the following three packings. The first uses an L that occupies the full knapsack, i.e., $w_L = h_L = N$. Let $OPT_{long} \subseteq OPT$ be the items in OPT with height or width strictly larger than N/2 and define $OPT_{short} = OPT \setminus OPT_{long}$. We apply Lemma 13 to OPT_{long} and hence obtain a packing for this L with a profit of at least $\frac{3}{4}p(OPT_{long})$. We run our PTAS for L-packings from Theorem 1 on this L, the input consisting of all items in I having one side longer than N/2. Hence we obtain a solution with profit at least $(\frac{3}{4} - O(\varepsilon))p(OPT_{long})$.

For the other two packings we employ the one-sided resource augmentation PTAS from [21]. We apply this algorithm to the slightly reduced knapsacks $[0,N] \times [0,N/(1+\varepsilon)]$ and $[0,N/(1+\varepsilon)] \times [0,N]$ such that in both cases it outputs a solution that fits in the full knapsack $[0,N] \times [0,N]$ and whose profit is by at most a factor $1+O(\varepsilon)$ worse than the optimal solution for the respective reduced knapsacks. We will prove in Theorem 14 that one of these solutions yields a profit of at least $(\frac{1}{2}-O(\varepsilon))p(OPT)+(\frac{1}{4}-O(\varepsilon))p(OPT_{short})$ and hence one of our packings yields a $(\frac{16}{9}+\varepsilon)$ -approximation.

Theorem 14. There is a $\frac{16}{9} + \varepsilon$ approximation for the cardinality case of 2DK.

Proof. Let OPT be the considered optimal solution with opt = p(OPT). Recall that there are no large items. Let also $OPT_{vert} \subseteq OPT$ be the (vertical) items with height more than $\varepsilon \cdot N$ (hence with width at

⁵It is possible to permute items in the left stack so that items appear from left to right in non-increasing order of height, and symmetrically for the other stacks. This is not crucial for this proof, but we implemented this permutation in Fig.1.(c).

most $\varepsilon \cdot N$), and $OPT_{hor} = OPT \setminus OPT_{ver}$ (horizontal items). Note that with this definition both sides of a horizontal item might have a length of at most $\varepsilon \cdot N$. We let $opt_{long} = p(OPT_{long})$ and $opt_{short} = p(OPT_{short})$.

As mentioned above, our L-packing PTAS achieves a profit of at least $(\frac{3}{4} - O(\varepsilon))opt_{long}$ which can be seen by applying Lemma 13 with $H = OPT_{long} \cap OPT_{hor}$ and $V = OPT_{long} \cap OPT_{ver}$. In order to show that the other two packings yield a good profit, consider a $random\ horizontal\ strip\ S = [0,N] \times [a,a+\varepsilon\cdot N]$ (fully contained in the knapsack) where $a\in [0,(1-\varepsilon)N)$ is chosen unformly at random. We remove all items of OPT intersecting S. Each item in OPT_{hor} and $OPT_{short} \cap OPT_{ver}$ is deleted with probability at most 3ε and $\frac{1}{2}+2\varepsilon$, resp. Therefore the total profit of the remaining items is in expectation at least $(1-3\varepsilon)p(OPT_{hor})+(\frac{1}{2}-2\varepsilon)p(OPT_{short}\cap OPT_{vert})$. Observe that the resulting solution can be packed into a restricted knapsack of size $[0,N]\times[0,N/(1+\varepsilon)]$ by shifting down the items above the horizontal strip. Therefore, when we apply the resource augmentation algorithm in [21] to the knapsack $[0,N]\times[0,N/(1+\varepsilon)]$, up to a factor $1-\varepsilon$, we will find a solution of (deterministically!) at least the same profit. In other terms, this profit is at least $(1-4\varepsilon)p(OPT_{hor})+(\frac{1}{2}-\frac{5}{2}\varepsilon)p(OPT_{short}\cap OPT_{vert})$. By a symmetric argument, we obtain a solution of profit at least $(1-4\varepsilon)p(OPT_{ver})+(\frac{1}{2}-\frac{5}{2}\varepsilon)p(OPT_{short}\cap OPT_{vert})$.

By a symmetric argument, we obtain a solution of profit at least $(1-4\varepsilon)p(OPT_{ver})+(\frac{1}{2}-\frac{5}{2}\varepsilon)p(OPT_{short}\cap OPT_{hor})$ when we apply the algorithm in [21] to the knapsack $[0,N/(1+\varepsilon)]\times[0,N]$. Thus the best of the latter two solutions has profit at least $(\frac{1}{2}-2\varepsilon)opt_{long}+(\frac{3}{4}-\frac{13}{4}\varepsilon)opt_{short}=(\frac{1}{2}-2\varepsilon)opt+(\frac{1}{4}-\frac{5}{4}\varepsilon)opt_{short}$. The best of our three solutions has therefore value at least $(\frac{9}{16}-O(\varepsilon))opt$ where the worst case is achieved for roughly $opt_{long}=3\cdot opt_{short}$.

In the above result we use either an L-packing or a container packing. The $\frac{558}{325} + \varepsilon$ approximation claimed in Theorem 2 is obtained by a careful combination of these two packings. In particular, we consider configurations where long items (or a subset of them) can be packed into a relatively small L, and pack part of the remaining short items in the complementary rectangular region (using container packings and Steinberg's algorithm [32]). See Section B for details.

4 Open Problems

The main problem that we left open is to find a PTAS, if any, for 2DK and 2DKR. This would be interesting even in the cardinality case. We believe that a better understanding of natural generalizations of L-packings might be useful. For example, is there are PTAS for *ring-packing* instances arising by shifting of long items? This would directly lead to an improved approximation factor for 2DK (though not to a PTAS). Is there a PTAS for L-packings *with rotations*? Our improved approximation algorithms for 2DKR are indeed based on a different approach. Is there a PTAS for O(1) instances of L-packing? This would also lead to an improved approximation factor for 2DK, and might be an important step towards a PTAS.

References

- [1] Anna Adamaszek, Tomasz Kociumaka, Marcin Pilipczuk, and Michal Pilipczuk. Hardness of approximation for strip packing. *CoRR*, abs/1610.07766, 2016.
- [2] Anna Adamaszek and Andreas Wiese. Approximation schemes for maximum weight independent set of rectangles. In *FOCS*, pages 400–409, 2013.
- [3] Anna Adamaszek and Andreas Wiese. A quasi-PTAS for the two-dimensional geometric knapsack problem. In *SODA*, pages 1491–1505, 2015.
- [4] Nikhil Bansal, Alberto Caprara, Klaus Jansen, Lars Prädel, and Maxim Sviridenko. A structural lemma in 2-dimensional packing, and its implications on approximability. In *ISAAC*, pages 77–86, 2009.
- [5] Nikhil Bansal, Alberto Caprara, and Maxim Sviridenko. A new approximation method for set covering problems, with applications to multidimensional bin packing. *SIAM Journal on Computing*, 39(4):1256–1278, 2009.
- [6] Nikhil Bansal and Arindam Khan. Improved approximation algorithm for two-dimensional bin packing. In *SODA*, pages 13–25, 2014.
- [7] Alberto Caprara. Packing 2-dimensional bins in harmony. In *FOCS*, pages 490–499, 2002.
- [8] Chandra Chekuri and Sanjeev Khanna. A polynomial time approximation scheme for the multiple knapsack problem. *SIAM Journal on Computing*, 35(3):713–728, 2005.
- [9] Henrik I. Christensen, Arindam Khan, Sebastian Pokutta, and Prasad Tetali. Approximation and online algorithms for multidimensional bin packing: A survey. *Computer Science Review*, 24:63–79, 2017.
- [10] Fan Chung, Michael R. Garey, and David S. Johnson. On packing two-dimensional bins. *SIAM Journal on Algebraic Discrete Methods*, 3:66–76, 1982.
- [11] Edward G. Coffman, Jr, Michael R. Garey, David S. Johnson, and Robert E. Tarjan. Performance bounds for level-oriented two-dimensional packing algorithms. *SIAM Journal on Computing*, 9:808–826, 1980.
- [12] Uriel Feige and Jan Vondrak. Approximation algorithms for allocation problems: Improving the factor of 1-1/e. In *Proceedings of the 47th Annual IEEE Symposium on Foundations of Computer Science (FOCS 2006)*, pages 667–676. IEEE, 2006.
- [13] Aleksei V. Fishkin, Olga Gerber, and Klaus Jansen. On efficient weighted rectangle packing with large resources. In *ISAAC*, pages 1039–1050, 2005.
- [14] Aleksei V. Fishkin, Olga Gerber, Klaus Jansen, and Roberto Solis-Oba. Packing weighted rectangles into a square. In *MFCS*, pages 352–363, 2005.
- [15] Lisa Fleischer, Michel X. Goemans, Vahab S. Mirrokni, and Maxim Sviridenko. Tight approximation algorithms for maximum separable assignment problems. *Mathematics of Operations Research*, 36(3):416–431, 2011.

- [16] Waldo Gálvez, Fabrizio Grandoni, Salvatore Ingala, and Arindam Khan. Improved pseudo-polynomial-time approximation for strip packing. In *FSTTCS*, pages 9:1–9:14, 2016.
- [17] Rolf Harren. Approximating the orthogonal knapsack problem for hypercubes. In *ICALP*, pages 238–249, 2006.
- [18] Rolf Harren, Klaus Jansen, Lars Prädel, and Rob van Stee. A $(5/3 + \epsilon)$ -approximation for strip packing. *Comput. Geom.*, 47(2):248–267, 2014.
- [19] Sandy Heydrich and Andreas Wiese. Faster approximation schemes for the two-dimensional knapsack problem. In *SODA*, pages 79–98, 2017.
- [20] Klaus Jansen and Malin Rau. Improved approximation for two dimensional strip packing with polynomial bounded width. *CoRR*, abs/1610.04430, 2016.
- [21] Klaus Jansen and Roberto Solis-Oba. New approximability results for 2-dimensional packing problems. In *MFCS*, pages 103–114, 2007.
- [22] Klaus Jansen and Roberto Solis-Oba. A polynomial time approximation scheme for the square packing problem. In *IPCO*, pages 184–198, 2008.
- [23] Klaus Jansen and Roberto Solis-Oba. Rectangle packing with one-dimensional resource augmentation. *Discrete Optimization*, 6(3):310 323, 2009.
- [24] Klaus Jansen and Guochuan Zhang. Maximizing the number of packed rectangles. In *SWAT*, pages 362–371, 2004.
- [25] Klaus Jansen and Guochuan Zhang. On rectangle packing: maximizing benefits. In *SODA*, pages 204–213, 2004.
- [26] Claire Kenyon and Eric Rémila. A near-optimal solution to a two-dimensional cutting stock problem. *Mathematics of Operations Research*, 25(4):645–656, 2000.
- [27] Arindam Khan. *Approximation algorithms for multidimensional bin packing*. PhD thesis, Georgia Institute of Technology, Atlanta, GA, USA, 2016.
- [28] Jan Karel Lenstra, David B. Shmoys, and Éva Tardos. Approximation algorithms for scheduling unrelated parallel machines. In *Proceedings of the 28th Annual Symposium on Foundations of Computer Science (FOCS 1987)*, pages 217–224, Washington, DC, USA, 1987. IEEE Computer Society.
- [29] Joseph YT Leung, Tommy W Tam, CS Wong, Gilbert H Young, and Francis YL Chin. Packing squares into a square. *Journal of Parallel and Distributed Computing*, 10(3):271–275, 1990.
- [30] Giorgi Nadiradze and Andreas Wiese. On approximating strip packing with a better ratio than 3/2. In *SODA*, pages 1491–1510, 2016.
- [31] David B. Shmoys and Éva Tardos. An approximation algorithm for the generalized assignment problem. *Mathematical Programming*, 62(1-3):461–474, 1993.
- [32] A. Steinberg. A strip-packing algorithm with absolute performance bound 2. *SIAM Journal on Computing*, 26(2):401–409, 1997.

A Weighted Case Without Rotations

In this section we show how to extend the reasoning of the unweighted case to the weighted case. This requires much more complicated technical machinery than the algorithm presented in Section 3.

Our strategy is to start with a partition of the knapsack into thin corridors as defined in [3]. Then, we partition these corridors into a set of rectangular boxes and an L-packing. We first present a simplified version of our argumentation in which we assume that we are allowed to drop $O_{\varepsilon}(1)$ many items at no cost, i.e., we pretend that we have the right to remove $O_{\varepsilon}(1)$ items from OPT and compare the profit of our computed solution with the remaining set. Building on this, we give an argumentation for the general case which will involve some additional shifting arguments.

A.1 Item classification

We start with a classification of the input items according to their heights and widths. For two given constants $1 \ge \varepsilon_{large} > \varepsilon_{small} > 0$, we classify an item i as:

- small if $h_i, w_i \leq \varepsilon_{small} N$;
- large if $h_i, w_i > \varepsilon_{large} N$;
- horizontal if $w_i > \varepsilon_{large} N$ and $h_i \leq \varepsilon_{small} N$;
- vertical if $h_i > \varepsilon_{large} N$ and $w_i \leq \varepsilon_{small} N$;
- intermediate otherwise (i.e., at least one side has length in $(\varepsilon_{small}N, \varepsilon_{large}N]$).

We also call *skewed* items that are horizontal or vertical. We let I_{small} , I_{large} , I_{hor} , I_{ver} , I_{skew} , and I_{int} be the items which are small, large, horizontal, vertical, skewed, and intermediate, respectively. The corresponding intersection with OPT defines the sets OPT_{small} , OPT_{large} , OPT_{hor} , OPT_{ver} , OPT_{skew} , OPT_{int} , respectively.

Observe that $|OPT_{large}| = O(1/\varepsilon_{large}^2)$ and since we are allowed to drop $O_{\varepsilon}(1)$ items from now on we ignore OPT_{large} . The next lemma shows that we can neglect also OPT_{int} .

Lemma 15. For any constant $\varepsilon > 0$ and positive increasing function $f(\cdot)$, f(x) > x, there exist constant values $\varepsilon_{large}, \varepsilon_{small}$, with $\varepsilon \geq \varepsilon_{large} \geq f(\varepsilon_{small}) \geq \Omega_{\varepsilon}(1)$ and $\varepsilon_{small} \in \Omega_{\varepsilon}(1)$ such that the total profit of intermediate rectangles is bounded by $\varepsilon p(OPT)$. The pair $(\varepsilon_{large}, \varepsilon_{small})$ is one pair from a set of $O_{\varepsilon}(1)$ pairs and this set can be computed in polynomial time.

Proof. Define $k+1=2/\varepsilon+1$ constants $\varepsilon_1,\ldots,\varepsilon_{k+1}$, such that $\varepsilon=f(\varepsilon_1)$ and $\varepsilon_i=f(\varepsilon_{i+1})$ for each i. Consider the k ranges of widths and heights of type $(\varepsilon_{i+1}N,\varepsilon_iN]$. By an averaging argument there exists one index j such that the total profit of items in OPT with at least one side length in the range $(\varepsilon_{j+1}N,\varepsilon_jN]$ is at most $2\frac{\varepsilon}{2}p(OPT)$. It is then sufficient to set $\varepsilon_{large}=\varepsilon_j$ and $\varepsilon_{small}=\varepsilon_{j+1}$.

We transform now the packing of the optimal solution OPT. To this end, we temporarily remove the small items OPT_{small} . We will add them back later. Thus, the reader may now assume that we need to pack only the skewed items from OPT_{skew} .




Figure 3: Illustration of two specific types of corridors: spirals (A) and rings (B)...

A.2 Corridors, Spirals and Rings

We build on a partition of the knapsack into corridors as used in [3]. We define an *open corridor* to be a face on the 2D-plane bounded by a simple rectilinear polygon with 2k edges e_0,\ldots,e_{2k-1} for some integer $k\geq 2$, such that for each pair of horizontal (resp., vertical) edges $e_i,e_{2k-i},i\in\{1,...,k-1\}$ there exists a vertical (resp., horizontal) line segment ℓ_i such that both e_i and e_{2k-i} intersect ℓ_i and ℓ_i does not intersect any other edge. Note that e_0 and e_k are not required to satisfy this property: we call them the *boundary edges* of the corridor. Similarly a *closed corridor* (or *cycle*) is a face on the 2D-plane bounded by two simple rectilinear polygons defined by edges e_0,\ldots,e_{k-1} and e'_0,\ldots,e'_{k-1} such that the second polygon is contained inside the first one, and for each pair of horizontal (resp., vertical) edges $e_i,e'_i,i\in\{0,...,k-1\}$, there exists a vertical (resp., horizontal) line segment ℓ_i such that both e_i and e'_i intersect ℓ_i and ℓ_i does not intersect any other edge. See Figures 3 and 4 for examples. Let us focus on minimum length such ℓ_i 's: then the *width* α of the corridor is the maximum length of any such ℓ_i . We say that an open (resp., closed) corridor of the above kind has k-2 (resp., k) bends. A corridor partition is a partition of the knapsack into corridors.

Lemma 16 (Corridor Packing Lemma [3]). *There exists a corridor partition and a set* $OPT_{corr} \subseteq OPT_{skew}$ *such that:*

- 1. there is a subset $OPT_{corr}^{cross} \subseteq OPT_{corr}$ with $|OPT_{corr}^{cross}| \leq O_{\varepsilon}(1)$ such that each item $i \in OPT_{corr} \setminus OPT_{corr}^{cross}$ is fully contained in some corridor,
- 2. $p(OPT_{corr}) \ge (1 O(\varepsilon))p(OPT_{skew}),$
- 3. the number of corridors is $O_{\varepsilon,\varepsilon_{large}}(1)$ and each corridor has width at most $\varepsilon_{large}N$ and has at most $1/\varepsilon$ bends.

Since we are allowed to drop $O_{\varepsilon}(1)$ items from now on we ignore OPT_{corr}^{cross} . We next identify some structural properties of the corridors that are later exploited in our analysis. Observe that an open (resp., closed) corridor of the above type is the union of k-1 (resp., k) boxes, that we next call *subcorridors* (see also Figure 4). Each such box is a maximally large rectangle that is contained in the corridor. The subcorridor S_i of an open (resp., closed) corridor of the above kind is the one containing edges e_i, e_{2k-i} (resp., $e_i, e_{i'}$) on its boundary. The length of S_i is the *length* of the shortest such edge. We say that a subcorridor is *long* if its length is more than N/2, and *short* otherwise. The partition of subcorridors into short and long will be crucial in our analysis.

We call a subcorridor horizontal (resp., vertical) if the corresponding edges are so. Note that each rectangle in OPT_{corr} is univocally associated with the only subcorridor that fully contains it: indeed, the longer side of a skewed rectangle is longer than the width of any corridor. Consider the sequence of consecutive subcorridors $S_1, \ldots, S_{k'}$ of an open or closed corridor. Consider two consecutive corridors S_i and $S_{i'}$, with i'=i+1 in the case of an open corridor and $i'=(i+1)\pmod{k'}$ otherwise. First assume that $S_{i'}$ is horizontal. We say that $S_{i'}$ is to the right (resp., left) of S_i if the right-most (left-most) boundary of $S_{i'}$ is to the right (left) of the right-most (left-most) boundary of S_i . If instead $S_{i'}$ is vertical, then S_i must be horizontal and we say that $S_{i'}$ is to the right (left) of S_i if S_i is to the left (right) of $S_{i'}$. Similarly, if $S_{i'}$ is vertical, we say that $S_{i'}$ is above (below) S_i if the top (bottom) boundary of $S_{i'}$ is above (below) the top (bottom) boundary of S_i . If $S_{i'}$ is horizontal, we say that it is above (below) S_i if S_i (which is vertical) is below (above) $S_{i'}$. We say that the pair $(S_i, S_{i'})$ forms a clockwise bend if S_i is horizontal and $S_{i'}$ is to its bottom-right or top-left, and the complementary cases if S_i is vertical. In all the other cases the pairs form a counter-clockwise bend. Consider a triple $(S_i, S_{i'}, S_{i''})$ of consecutive subcorridors in the above sense. It forms a *U*-bend if $(S_i, S_{i'})$ and $(S_{i'}, S_{i''})$ are both clockwise or counterclockwise bends. Otherwise it forms a Z-bend. In both cases $S_{i'}$ is the *center* of the bend, and $S_i, S_{i''}$ its *sides*. An open corridor whose bends are all clockwise (resp., counter-clockwise) is a *spiral*. A closed corridor with k=4 is a *ring*. Note that in a ring all bends are clockwise or counter-clockwise, hence in some sense it is the closed analogue of a spiral. We remark that a corridor whose subcorridors are all long is a spiral or a ring⁶. As we will see, spirals and rings play a crucial role in our analysis. In particular, we will exploit the following simple fact.

Lemma 17. The following properties hold:

- 1. The two sides of a Z-bend cannot be long. In particular, an open corridor whose subcorridors are all long is a spiral.
- 2. A closed corridor contains at least 4 distinct (possibly overlapping) U-bends.

Proof. (1) By definition of long subcorridors and Z-bend, the 3 subcorridors of the Z-bend would otherwise have total width or height larger than N. (2) Consider the left-most and right-most vertical subcorridords, and the top-most and bottom-most horizontal subcorridors. These 4 subcorridors exist, are distinct, and are centers of a U-bend.

A.3 Partitioning Corridors into Rectangular Boxes

We next describe a routine to partition the corridors into rectangular boxes such that each item is contained in one such box. We remark that to achieve this partitioning we sometimes have to sacrifice a large fraction of OPT_{corr} , hence we do not achieve a $1+\varepsilon$ approximation as in [2]. On the positive side, we generate only a constant (rather than polylogarithmic) number of boxes. This is crucial to obtain a polynomial time algorithm in the later steps.

Recall that each $i \in OPT_{corr}$ is univocally associated with the only subcorridor that fully contains it. We will say that we *delete* a sub-corridor, when we delete all rectangles univocally associated with the subcorridor. Note that in deletion of a sub-corridor we do not delete rectangles that are partially contained in that subcorridor but completely contained in a neighbor sub-corridor. Given a corridor, we sometimes *delete* some of its subcorridors, and consider the *residual* corridors (possibly more than one) given by the union of the remaining subcorridors. Note that removing any subcorridor from a closed corridor turns it into an open corridor. We implicitly assume that items associated with a deleted subcorridor are also removed (and consequently the corresponding area can be used to pack other items).

⁶We leave the simple proof for the ring case to the reader since we do not explicitly need this claim.




Figure 4: Left: The subcorridors S_1 and S_3 are vertical, S_2 and S_4 are horizontal. The subcorridor S_3 is on the top-right of S_2 . The curve on the bottom left shows the boundary curve between S_1 and S_2 . The pair (S_3, S_4) forms a clockwise bend and the pair (S_2, S_3) forms a counter-clockwise bend. The triple (S_1, S_2, S_3) forms a U-bend and the triple (S_2, S_3, S_4) forms a Z-bend. Right: Our operation that divides a corridor into $O_{\varepsilon}(1)$ boxes and $O_{\varepsilon}(1)$ shorter corridors. The dark gray items show thin items that are removed in this operation. The light gray items are fat items that are shifted to the box below their respective original box. The value α denotes the width of the depicted corridor.

Given two consecutive subcorridors S_i and $S_{i'}$, we define the *boundary curve* among them as follows (see also Figure 4). Suppose that $S_{i'}$ is to the top-right of S_i , the other cases being symmetric. Let $S_{i,i'} = S_i \cap S_{i'}$ be the rectangular region shared by the two subcorridors. Then the boundary curve among them is any simple rectilinear polygon inside $S_{i,i'}$ that decreases monotonically from its top-left corner to its bottom-right one and that does not cut any rectangle in these subcorridors. For a boundary horizontal (resp., vertical) subcorridor of an open corridor (i.e., a subcorridor containing e_0 or e_{2k-1}) we define a dummy boundary curve given by the vertical (resp., horizontal) side of the subcorridor that coincides with a (boundary) edge of the corridor.

Remark 1. Each subcorridor has two boundary curves (including possibly dummy ones). Furthermore, all its items are fully contained in the region delimited by such curves plus the two edges of the corridor associated with the subcorridor (private region).

Given a corridor, we partition its area into a constant number of boxes as follows (see also Figure 4, and [2] for a more detailed description of an analogous construction). Let S be one of its boundary subcorridors (if any), or the central subcorridor of a U-bend. Note that one such S must exist (trivially for an open corridor, otherwise by Lemma 17.2). In the corridor partition, there might be several subcorridors fulfilling the latter condition. We will explain later in which order to process the subcorridors, here we explain only how to apply our routine to *one* subcorridor, which we call *processing* of subcorridor.

Suppose that S is horizontal with height b, with the shorter horizontal associated edge being the top one. The other cases are symmetric. Let $\varepsilon_{box}>0$ be a sufficiently small constant to be defined later. If S is the only subcorridor in the considered corridor, S forms a box and all its items are marked as fat. Otherwise, we draw $1/\varepsilon_{box}$ horizontal lines that partition the private region of S into subregions of height $\varepsilon_{box}b$. We mark as thin the items of the bottom-most (i.e., the widest) such subregion, and as killed the items of the subcorridor cut by these horizontal lines. All the remaining items of the subcorridor are marked as fat.

For each such subregion, we define an associated (horizontal) box as the largest axis-aligned box that is contained in the subregion. Given these boxes, we partition the rest of the corridor into $1/\varepsilon_{box}$ corridors as follows. Let S' be a corridor next to S, say to its top-right. Let P be the set of corners of the boxes contained in the boundary curve between S and S'. We project P vertically on the boundary curve of S' not shared with S, hence getting a set P' of $1/\varepsilon_{box}$ points. We iterate the process on the pair (S', P'). At the end of the process, we obtain a set of $1/\varepsilon_{box}$ boxes from the starting subcorridor S, plus a collection of $1/\varepsilon_{box}$ new (open) corridors each one having one less bend with respect to the original corridor. Later, we will also apply this process on the latter corridors. Each newly created corridor will have one bend less than the

original corridor and thus this process eventually terminates. Note that, since initially there are $O_{\varepsilon,\varepsilon_{large}}(1)$ corridors each one with $O(1/\varepsilon)$ bends, the final number of boxes is $O_{\varepsilon,\varepsilon_{large},\varepsilon_{box}}(1)$. See Figure 4 for an illustration.

Remark 2. Assume that we execute the above procedure on the subcorridors until there is no subcorridor left on which we can apply it. Then we obtain a partition of OPT_{corr} into disjoint sets OPT_{thin} , OPT_{fat} , and OPT_{kill} of thin, fat, and killed items, respectively. Note that each order to process the subcorridors leads to different such partition. We will define this order carefully in our analysis.

Remark 3. By a simple shifting argument, there exists a packing of OPT_{fat} into the boxes. Intuitively, in the above construction each subregion is fully contained in the box associated with the subregion immediately below (when no lower subregion exists, the corresponding items are thin).

We will from now on assume that the shifting of items as described in Remark 3 has been done.

The following lemma summarises some of the properties of the boxes and of the associated partition of OPT_{corr} (independently from the way ties are broken). Let I_{hor} and I_{ver} denote the set of horizontal and vertical input items, respectively.

Lemma 18. The following properties hold:

- 1. $|OPT_{kill}| = O_{\varepsilon, \varepsilon_{large}, \varepsilon_{box}}(1);$
- 2. For any given constant $\varepsilon_{ring} > 0$ there is a sufficiently small $\varepsilon_{box} > 0$ such that the total height (resp., width) of items in $OPT_{thin} \cap I_{hor}$ (resp., $OPT_{thin} \cap I_{ver}$) is at most $\varepsilon_{ring}N$.

Proof. (1) Each horizontal (resp., vertical) line in the construction can kill at most $1/\varepsilon_{large}$ items, since those items must be horizontal (resp., vertical). Hence we kill $O_{\varepsilon,\varepsilon_{large},\varepsilon_{box}}(1)$ items in total.

(2) The mentioned total height/width is at most $\varepsilon_{box}N$ times the number of subcorridors, which is $O_{\varepsilon,\varepsilon_{large}}(1)$. The claim follows for ε_{box} small enough.

A.4 Containers

Assume that we applied the routine described in Section A.3 above until each corridor is partitioned into boxes. We explain how to partition each box into $O_{\varepsilon}(1)$ subboxes, to which we refer to as *containers* in the sequel. Hence, we apply the routine described below to each box.

Consider a box of size $a \times b$ coming from the above construction, and on the associated set OPT_{box} of items from OPT_{fat} . We will show how to pack a set $OPT'_{box} \subseteq OPT_{box}$ with $p(OPT'_{box}) \ge (1 - \varepsilon)p(OPT_{box})$ into $O_{\varepsilon}(1)$ containers packed inside the box, such that both the containers and the packing of OPT'_{box} inside them satisfy some extra properties that are useful in the design of an efficient algorithm. This part is similar in spirit to prior work, though here we present a refined analysis that simplifies the algorithm (in particular, we can avoid LP rounding).

A container is a box labelled as horizontal, vertical, or area. A container packing of a set of items I' into a collection of non-overlapping containers has to satisfy the following properties:

- Items in a horizontal (resp., vertical) container are stacked one on top of the other (resp., one next to the other).
- Each $i \in I'$ packed in an area container of size $a \times b$ must have $w_i \leq \varepsilon a$ and $h_i \leq \varepsilon b$.

Our main building block is the following resource augmentation packing lemma essentially taken from [23]⁷.

Lemma 19 (Resource Augmentation Packing Lemma). Let I' be a collection of items that can be packed into a box of size $a \times b$, and $\varepsilon_{ra} > 0$ be a given constant. Then there exists a container packing of $I'' \subseteq I'$ inside a box of size $a \times (1 + \varepsilon_{ra})b$ (resp., $(1 + \varepsilon_{ra})a \times b$) such that:

- 1. $p(I'') \ge (1 O(\varepsilon_{ra}))p(I');$
- 2. The number of containers is $O_{\varepsilon_{ra}}(1)$ and their sizes belong to a set of cardinality $n^{O_{\varepsilon_{ra}}(1)}$ that can be computed in polynomial time.

Applying Lemma 19 to each box yields the following lemma.

Lemma 20 (Container Packing Lemma). For a given constant $\varepsilon_{ra} > 0$, there exists a set $OPT_{fat}^{cont} \subseteq OPT_{fat}$ such that there is a container packing for all apart from $O_{\varepsilon}(1)$ items in OPT_{fat}^{cont} such that:

- 1. $p(OPT_{fat}^{cont}) \ge (1 O(\varepsilon))p(OPT_{fat});$
- 2. The number of containers is $O_{\varepsilon,\varepsilon_{large},\varepsilon_{box},\varepsilon_{ra}}(1)$ and their sizes belong to a set of cardinality $n^{O_{\varepsilon,\varepsilon_{large},\varepsilon_{box},\varepsilon_{ra}}(1)}$ that can be computed in polynomial time.

Proof. Let us focus on a specific box of size $a \times b$ from the previous construction in Section A.3, and on the items $OPT_{box} \subseteq OPT_{fat}$ inside it. If $|OPT_{box}| = O_{\varepsilon}(1)$ then we can simply create one container for each item and we are done. Otherwise, assume w.l.o.g. that this box (hence its items) is horizontal. We obtain a set $\overline{OPT_{box}}$ by removing from OPT_{box} all items intersecting a proper horizontal strip of height $3\varepsilon b$. Clearly these items can be repacked in a box of size $a \times (1 - 3\varepsilon)b$. By a simple averaging argument, it is possible to choose the strip so that the items fully contained in it have total profit at most $O(\varepsilon)p(OPT_{box})$. Furthermore, there can be at most $O(1/\varepsilon_{large})$ items that partially overlap with the strip (since items are skewed). We drop these items and do not pack them.

At this point we can use the Resource Augmentation Lemma 19 to pack a large profit subset $OPT'_{box} \subseteq \overline{OPT_{box}}$ into $O_{\varepsilon_{ra}}(1)$ containers that can be packed in a box of size $a \times (1-3\varepsilon)(1+\varepsilon_{ra})b \leq a \times (1-2\varepsilon)b$. We perform the above operation on each box of the previous construction and define OPT^{cont}_{fat} to be the union of the respective sets OPT'_{box} . The claim follows.

A.5 A Profitable Structured Packing

We next prove our main structural lemma which yields that there exists a structured packing which is partitioned into $O_{\varepsilon}(1)$ containers and an L. We will refer to such a packing as an L&C packing (formally defined below). Note that in the previous section we did not specify in which order we partition the subcorridors into boxes. In this section, we give several such orders which will then result in different packings. The last such packing is special since we will modify it a bit to gain some space and then reinsert the thin items that were removed in the process of partitioning the corridors into containers. Afterwards, we will show that one of the resulting packings will yield an approximation ratio of $17/9 + \varepsilon$.

A boundary ring of width N' is a ring having as external boundary the edges of the knapsack and as internal boundary the boundary of a square box of size $(N - N') \times (N - N')$ in the middle of the knapsack.

⁷In Appendix F we reprove this lemma in a *container-based* form, rather than using LP-based arguments, since this is more convenient for our final algorithm. Our version of the lemma might also be a handy tool for future work.

A boundary L of width N' is the region covered by two boxes of size $N' \times N$ and $N \times N'$ that are placed on the left and bottom boundaries of the knapsack.

An L&C packing is defined as follows. We are given two integer parameters $N' \in [0, N/2]$ and $\ell \in (N/2, N]$. We define a boundary L of width N', and a collection of non-overlapping containers contained in the space not occupied by the boundary L. The number of containers and their sizes are as in Lemma 20. We let $I_{\text{long}} \subseteq I$ be the items whose longer side has length longer than ℓ (hence longer than N/2), and $I_{\text{short}} = I \setminus I_{\text{long}}$ be the remaining items. We can pack only items from I_{long} in the boundary L, and only items from I_{short} in the containers (satisfying the usual container packing constraints). See also Figure 1.

Remark 4. In the analysis sometimes we will not need the boundary L. This case is captured by setting N'=0 and $\ell=N$ (degenerate L case).

Lemma 21. Let $OPT_{L\&C}$ be the most profitable solution that is packed by an L&C packing. Then $p(OPT_{L\&C}) \ge (\frac{9}{17} - O(\varepsilon))p(OPT)$.

In the remainder of this section we prove Lemma 21, assuming that we can drop $O_{\varepsilon}(1)$ items at no cost. Hence, formally we will prove that there is an L&C packing I' and a set of $O_{\varepsilon}(1)$ items I_{drop} such that $p(I') + p(I_{\text{drop}}) \geq (\frac{9}{17} - O(\varepsilon))p(OPT)$. Subsequently, we will prove Lemma 21 in full generality (without dropping any items).

The proof of Lemma 21 involves some case analysis. Recall that we classify subcorridors into short and long, and horizontal and vertical. We further partition short subcorridors as follows: let $S_1, \ldots, S_{k'}$ be the subcorridors of a given corridor, and let $S_1^s, \ldots, S_{k''}^s$ be the subsequence of short subcorridors (if any). Mark S_i^s as *even* if i is so, and *odd* otherwise. Note that corridors are subdivided into several other corridors during the box construction process (see the right side of Figure 4), and these new corridors might have fewer subcorridors than the initial corridor. However, the marking of the subcorridors (short, long, even, odd, horizontal, vertical) is inherited from the marking of the original subcorridor.

We will describe now 7 different ways to partition the subcorridors into boxes, for some of them we delete some of the subcorridors. Each of these different processing orders will give different sets OPT_{thin}, OPT_{kill} and OPT_{fat}^{cont} , and based on these, we will partition the items into three sets. We will then prove three different lower bounds on $p(OPT_{L\&C})$ w.r.t. the sizes of these three sets using averaging arguments about the seven cases.

Cases 1a, 1b, 2a, 2b: Short horizontal/short vertical subcorridors. We delete either all vertical short (case 1) or all horizontal short subcorridors (case 2). We first process all short subcorridors, then either all vertical (subcases a) or horizontal long ones (subcases b), and finally the remaining (horizontal or vertical, resp.) long ones. We can start by processing all short corridors. Indeed, any such corridor cannot be the center of a Z-bend by Lemma 17.1 since its two sides would be long, hence it must be boundary or the center of a U-bend. After processing short subcorridors, by the same argument the residual (long) subcorridors are the boundary or the center of a U-bend. So we can process the long subcorridors in any order. This gives in total four cases. See Fig. 5 for deletion/processing of subcorridors for these cases.

Cases 3a, 3b: Even/odd short subcorridors. We delete the odd (or even) short subcorridors and then process even (resp., odd) short subcorridors last. We exploit the fact that each residual corridor contains at most one short subcorridor. Then, if there is another (long) subcorridor, there is also one which is boundary (trivially for an open corridor) or the center of a *U*-bend (by Lemma 17, Property 2). Hence we can always process some long subcorridor leaving the unique short subcorridor as last one. This gives two cases.




Figure 5: Figure for Case 1 and 2. The knapsack on the left contains two corridors, where short subcorridors are marked light grey and long subcorridors are marked dark grey. In case 1, we delete vertical short subcorridors and then consider two processing orders in subcases a and b. In case 2, we delete horizontal short subcorridors and again consider two processing orders in subcases a and b.

Case 4: Fat only. Do not delete any short subcorridor. Process subcorridors in any feasible order.

In each of the cases, we apply the procedure described in Section A.4 to partition each box into $O_{\varepsilon}(1)$ containers. We next label items as follows. Consider the classification of items into OPT_{fat}^{cont} , OPT_{thin} , and OPT_{kill} in each one of the 7 cases above. Then:

- OPT_T is the set of items which are in OPT_{thin} in at least one case;
- OPT_K is the set of items which are in OPT_{kill} in at least one case;
- OPT_F is the set of items which are in OPT_{fat}^{cont} in all the cases.

Remark 5. Consider the subcorridor of a given corridor that is processed last in one of the above cases. None of its items are assigned to OPT_{thin} in that case and thus essentially all its items are packed in one of

the constructed containers. In particular, for an item in set OPT_T , in some of the above cases it might be in such a subcorridor and thus marked fat and packed into a container.

Lemma 22. One has
$$p(OPT_F \cup OPT_T) + p(OPT_K) + p(OPT_{large}) + p(OPT_{corr}^{cross}) \ge (1 - O(\varepsilon))p(OPT)$$
.

Proof. Let us initialize $OPT_F = OPT_{fat}^{cont}$, $OPT_T = OPT_{thin}$, and $OPT_K = OPT_{kill}$ by considering one of the above cases. Next we consider the aforementioned cases, hence moving some items in OPT_F to either OPT_T or OPT_K . Note that initially $p(OPT_F \cup OPT_T) + p(OPT_{kill}) + p(OPT_{large}) + p(OPT_{corr}^{cross}) \ge (1 - O(\varepsilon))p(OPT)$ by Lemma 16 and hence we keep this property.

Let I_{lc} and I_{sc} denote the items in long and short corridors, respectively. We also let $OPT_{LF} = I_{lc} \cap OPT_F$, and define analogously OPT_{SF} , OPT_{LT} , and OPT_{LF} . The next three lemmas provide a lower bound on the case of a degenerate L.

Lemma 23.
$$p(OPT_{L\&C}) \ge p(OPT_{LF}) + p(OPT_{SF})$$
.

Proof. Follows immediately since we pack a superset of OPT_F in case 4.

Lemma 24.
$$p(OPT_{L\&C}) \ge p(OPT_{LF}) + p(OPT_{LT})/2 + p(OPT_{SF})/2$$
.

Proof. Consider the sum of the profit of the packed items corresponding to the in total four subcases of cases 1 and 2. Each $i \in OPT_{LF}$ appears 4 times in the sum (as items in OPT_F are fat in all cases and all long subcorridors get processed), and each $i \in OPT_{LT}$ at least twice by Remark 5: If a long subcorridor $\mathfrak L$ neighbors a short subcorridor, the short subcorridor is either deleted or processed first. Further, all neighboring long subcorridors are processed first in case 1a and 2a (if $\mathfrak L$ is horizontal, then its neighbors are vertical) or 1b and 2b (if $\mathfrak L$ is vertical and its neighbors are horizontal). Thus, $\mathfrak L$ is the last processed subcorridor in at least two cases. Additionally, each item $i \in OPT_{SF}$ also appears twice in the sum, as it gets deleted either in case 1 (if it is vertical) or in case 2 (if it is horizontal) and is fat otherwise.

The claim follows by an averaging argument. \Box

Lemma 25.
$$p(OPT_{L\&C}) \ge p(OPT_{LF}) + p(OPT_{SF})/2 + p(OPT_{ST})/2$$
.

Proof. Consider the sum of the number of packed items corresponding to cases 3a and 3b. Each $i \in OPT_{LF}$ appears twice in the sum as it is fat and all long subcorridors get processed. Each $i \in OPT_{SF} \cup OPT_{ST}$ appears at least once in the sum by Remark 5: An item $i \in OPT_{SF}$ is deleted in one of the two cases (depending on whether it is in an even or odd subcorridor) and otherwise fat. An item $i \in OPT_{ST}$ is also deleted in one of the two cases and otherwise its subcorridor is processed last. The claim follows by an averaging argument.

There is one last (and slightly more involving) case to be considered, corresponding to a non-degenerate L.

Lemma 26.
$$p(OPT_{L\&C}) \ge \frac{3}{4}p(OPT_{LT}) + p(OPT_{ST}) + \frac{1 - O(\varepsilon)}{2}p(OPT_{SF}).$$

Proof. Recall that $\varepsilon_{large}N$ is the maximum width of a corridor. We consider an execution of the algorithm with boundary L width $N' = \varepsilon_{ring}N$, and threshold length $\ell = (\frac{1}{2} + 2\varepsilon_{large})N$. We remark that this length guarantees that items in I_{long} are not contained in short subcorridors.

By Lemma 13, we can pack a subset of $OPT_T \cap I_{long}$ of profit at least $\frac{3}{4}p(OPT_T \cap I_{long})$ in a boundary L of width $\varepsilon_{ring}N$. By Lemma 18 the remaining items in OPT_T can be packed in two containers of size $\ell \times \varepsilon_{ring}N$ and $\varepsilon_{ring}N \times \ell$ that we place on the two sides of the knapsack not occupied by the boundary L.

In the free area we can identify a square region K'' with side length $(1-\varepsilon)N$. We next show that there exists a feasible solution $OPT'_{SF}\subseteq OPT_{SF}$ with $p(OPT'_{SF})\geq (1-O(\varepsilon))p(OPT_{SF})/2$ that can be packed in a square of side length $(1-3\varepsilon)N$. We can then apply the Resource Augmentation Lemma 19 to pack $OPT''_{SF}\subseteq OPT'_{SF}$ of cardinality $p(OPT''_{SF})\geq (1-O(\varepsilon))p(OPT'_{SF})$ inside a central square region of side length $(1-3\varepsilon)(1+\varepsilon_{ra})N\leq (1-2\varepsilon)N$ using containers according to Lemma 20.

Consider the packing of OPT_{SF} as in the optimum solution. Choose a random vertical (resp., horizontal) strip in the knapsack of width (resp., height) $3\varepsilon N$. Delete from OPT_{SF} all the items intersecting the vertical and horizontal strips: clearly the remaining items OPT'_{SF} can be packed into a square of side length $(1-3\varepsilon)N$. Consider any $i\in OPT_{SF}$, and assume i is horizontal (the vertical case being symmetric). Recall that it has height at most $\varepsilon_{small}N\leq \varepsilon N$ and width at most $\ell\leq 1/2+2\varepsilon$. Therefore i intersects the horizontal strip with probability at most 5ε and the vertical strip with probability at most $1/2+8\varepsilon$. Thus by the union bound $i\in OPT'_{SF}$ with probability at least $1/2-13\varepsilon$. The claim follows by linearity of expectation.

Combining the above Lemmas 22, 23, 24, 25, and 26 we achieve the desired approximation factor, assuming that the (dropped) $O_{\varepsilon}(1)$ items in $OPT_{kill} \cup OPT_{large} \cup OPT_{corr}^{cross}$ have zero profit. The worst case is obtained, up to $1 - O(\varepsilon)$ factors, for $p(OPT_{LT}) = p(OPT_{SF}) = p(OPT_{ST})$ and $p(OPT_{LF}) = 5p(OPT_{LT})/4$. This gives $p(OPT_{LT}) = 4/17 \cdot p(OPT_{T} \cup OPT_{F})$ and a total profit of $9/17 \cdot p(OPT_{T} \cup OPT_{F})$.

A.6 Adding small items

Note that up to now we ignored the small items OPT_{small} . In this section, we explain how to pack a large fraction of these items.

We described above how to pack a large enough fraction of OPT_{skew} into containers. We next refine the mentioned analysis to bound the total area of such containers. It turns out that the residual area is sufficient to pack almost all the items of OPT_{small} into a constant number of area containers (not overlapping with the previous containers) for ε_{small} small enough.

To this aim we use a refined version of the Resource Augmentation Lemma 19, i.e., Lemma 55. Essentially, besides the other properties, we can guarantee that the total area of the containers is at most $a(I') + \varepsilon_{ra} \ a \cdot b$, where $a \times b$ and I' are the size of the box and the initial set of items in the claim of Lemma 19, respectively.

Lemma 27. In the packings due to Lemmas 23, 24, 25, and 26 the total area occupied by containers is at most $\min\{(1-2\varepsilon)N^2, a(OPT_{corr}) + \varepsilon_{ra}N^2\}$.

Proof. Consider the first upper bound on the area. We have to distinguish between the containers considered in Lemma 26 and the remaining cases. In the first case, there is a region not occupied by the boundary L nor by the containers of area at least $4\varepsilon N^2 - 4\varepsilon^2 N^2 - 4\varepsilon_{ring}N^2 \geq 2\varepsilon N^2$ for ε_{ring} small enough, e.g., $\varepsilon_{ring} \leq \epsilon^2$ suffices. The claim follows. For the remaining cases, recall that in each horizontal box of size $a \times b$ we remove a horizontal strip of height $3\varepsilon b$, and then use the Resource Augmentation Packing Lemma to pack the residual items in a box of size $a \times b(1-3\varepsilon)(1+\varepsilon_{ra}) \leq a \times b(1-2\varepsilon)$ for $\varepsilon_{ra} \leq \varepsilon$. Thus the total area of the containers is at most a fraction $1-2\varepsilon$ of the area of the original box. A symmetric argument applies to vertical boxes. Thus the total area of the containers is at most a fraction $1-2\varepsilon$ of the total area of the boxes, which in turn is at most N^2 . This gives the first upper bound in the claim.

For the second upper bound, we just apply the area bound in Lemma 55 to get that the total area of the containers is at most $a(OPT_{corr})$ plus $\varepsilon_{ra} \, a \cdot b$ for each box of size $a \times b$. Summing the latter terms over the boxes one obtains at most $\varepsilon_{ra} N^2$.

We are now ready to state a lemma that provides the desired packing of small items. By slightly adapting the analysis we can guarantee that the boundary L that we use to prove the claimed approximation ratio has width at most $\varepsilon^2 N$.

Lemma 28 (Small Items Packing Lemma). Suppose we are given a packing of non small items of the above type into k containers of total area A and, possibly, a boundary L of width at most $\varepsilon^2 N$. Then for ε_{small} small enough it is possible to define $O_{\varepsilon_{small}}(1)$ area containers of size $\frac{\varepsilon_{small}}{\varepsilon} N \times \frac{\varepsilon_{small}}{\varepsilon} N$ neither overlapping with the containers nor with the boundary L (if any) such that it is possible to pack $OPT'_{small} \subseteq OPT_{small}$ of profit $p(OPT'_{small}) \ge (1 - O(\varepsilon))p(OPT_{small})$ inside these new area containers.

Proof. Let us build a grid of width $\varepsilon'N = \frac{\varepsilon_{small}}{\varepsilon} \cdot N$ inside the knapsack. We delete any cell of the grid that overlaps with a container or with the boundary L, and call the remaining cells *free*. The new area containers are the free cells.

The total area of the deleted grid cells is, for ε_{small} and ε_{ra} small enough, at most

$$(\varepsilon^2 N^2 + A) + (2N + 4Nk) \frac{1}{\varepsilon' N} \cdot (\varepsilon' N)^2 \le A + 2\varepsilon^2 N^2 \stackrel{Lem.27}{\le} \min\{(1 - \varepsilon)N^2, a(OPT_{corr}) + 3\varepsilon^2 N^2\}$$

For the sake of simplicity, suppose that any empty space in the optimal packing of $OPT_{corr} \cup OPT_{small}$ is filled in with dummy small items of profit 0, so that $a(OPT_{corr} \cup OPT_{small}) = N^2$. We observe that the area of the free cells is at least $(1 - O(\varepsilon))a(OPT_{small})$: Either, $a(OPT_{small}) \geq \varepsilon N^2$ and then the area of the free cells is at least $a(OPT_{small}) - 3\varepsilon^2 N^2 \geq (1 - 3\varepsilon)a(OPT_{small})$; otherwise, we have that the area of the free cells is at least $\varepsilon N^2 > a(OPT_{small})$. Therefore we can select a subset of small items $OPT'_{small} \subseteq OPT_{small}$, with $p(OPT'_{small}) \geq (1 - O(\varepsilon))p(OPT_{small})$ and area $a(OPT_{small}) \leq (1 - O(\varepsilon))a(OPT_{small})$ that can be fully packed into free cells using classical Next Fit Decreasing Height algorithm (NFDH) according to Lemma 52 described later. The key argument for this is that each free cell is by a factor $1/\varepsilon$ larger in each dimension than each small item.

Thus, we have proven now that if the items in $OPT_{kill} \cup OPT_{large} \cup OPT_{corr}^{cross}$ had zero profit, then there is an L&C-packing for the skewed and small items with a profit of at least $9/17 \cdot p(OPT_{corr}) + (1 - O(\epsilon))(OPT_{small}) \ge (9/17 - O(\epsilon))p(OPT)$.

A.7 Shifting argumentation

We remove now the assumption that we can drop $O_{\varepsilon}(1)$ items from OPT. We will add a couple of shifting steps to the argumentation above to prove Lemma 21 without that assumption.

It is no longer true that we can neglect the large rectangles OPT_{large} since they might contribute a large amount towards the objective, even though their total number is guaranteed to be small. Also, in the process of constructing the boxes, we killed up to $O_{\varepsilon}(1)$ rectangles (the rectangles in OPT_{kill}). Similarly, we can no longer drop the constantly many items in OPT_{corr}^{cross} . Therefore, we apply some careful shifting arguments in order to ensure that we can still use a similar construction as above, while losing only a factor $1 + O(\varepsilon)$ due to some items that we will discard.

The general idea is as follows: For $t=0,\ldots,k$ (we will later argue that $k<1/\varepsilon$), we define disjoint sets K(t) recursively, each containing at most $O_\varepsilon(1)$ items. Each set $K(t)=\bigcup_{j=0}^t K(j)$ is used to define a grid G(t) in the knapsack. Based on an item classification that depends on this grid, we identify a set of skewed items and create a corridor partition w.r.t. these skewed items as described in Lemma 16. We then create a partition of the knapsack into corridors and a constant (depending on ε) number of containers (see Section A.7.1). Next, we decompose the corridors into boxes (Section A.7.2) and these boxes into containers (section A.7.3) similarly as we did in Sections A.3 and A.4 (but with some notable changes as we did not delete small items from the knapsack and thus need to handle those as well). In the last step, we add small items to the packing (Section A.7.4). During this whole process, we define the set K(i+1) of items that are somehow "in our way" during the decomposition process (e.g., items that are not fully contained in some corridor of the corridor partition), but which we cannot delete directly as they might have large profit. These items are similar to the killed items in the previous argumentation. However, using a shifting argument we can simply show that after at most $k < 1/\varepsilon$ steps of this process, we encounter a set K(k) of low total profit, that we can remove, at which point we can apply almost the same argumentation as in Lemmas 23, 24, and 25 to obtain lower bounds on the profit of an optimal L&C packing (Section A.7.5).

We initiate this iterative process as follows: Denote by K(0) a set containing all items that are killed in at least one of the cases arising in Section A.5 (the set OPT_K in that Section) and additionally the large items OPT_{large} and the $O_{\varepsilon}(1)$ items in OPT_{corr}^{cross} (in fact $OPT_{large} \subseteq OPT_{corr}^{cross}$). Note that $|K(0)| \leq O_{\varepsilon}(1)$. If $p(K(0)) \leq \varepsilon \cdot p(OPT)$ then we can simply remove these rectangles (losing only a factor of $1 + \varepsilon$) and then apply the remaining argumentation exactly as above and we are done. Therefore, from now on suppose that $p(K(0)) > \varepsilon \cdot p(OPT)$.

A.7.1 Definition of grid and corridor partition

Assume we are in round t of this process, i.e., we defined K(t) in the previous step (unless t=0, then K(t) is defined as specified above) and assume that $p(K(t)) > \varepsilon OPT$ (otherwise, see Section A.7.5). We are now going to define the non-uniform grid G(t) and the induced partition of the knapsack into a collection of cells \mathcal{C}_t . The x-coordinates (y-coordinates) of the grid cells are the x-coordinates (y-coordinates, respectively) of the items in $\mathcal{K}(t)$. This yields a partition of the knapsack into $O_{\varepsilon}(1)$ rectangular cells, such that each item of $\mathcal{K}(t)$ covers one or multiple cells. Note that an item might intersect many cells.

Similarly as above, we define constants $1 \geq \varepsilon_{large} \geq \varepsilon_{small} \geq \Omega_{\varepsilon}(1)$ and apply a shifting step such that we can assume that for each item $i \in OPT$ touching some cell C we have that $w(i \cap C) \in (0, \varepsilon_{small}w(C)] \cup (\varepsilon_{large}w(C), w(C)]$ and $h(i \cap C) \in (0, \varepsilon_{small}h(C)] \cup (\varepsilon_{large}h(C), h(C)]$, where h(C) and w(C) denote the height and the width of the cell C and $w(i \cap C)$ and $h(i \cap C)$ denote the height and the width of the intersection of the rectangle i with the cell C, respectively. For a cell C denote by OPT(C) the set of rectangles that intersect C in OPT. We obtain a partition of OPT(C) into $OPT_{small}(C)$, $OPT_{large}(C)$, $OPT_{hor}(C)$, and $OPT_{ver}(C)$:

- $OPT_{small}(C)$ contains all items $i \in OPT(C)$ with $h(i \cap C) \leq \varepsilon_{small}h(C)$ and $w(i \cap C) \leq \varepsilon_{small}w(C)$,
- $OPT_{large}(C)$ contains all items $i \in OPT(C)$ with $h(i \cap C) > \varepsilon_{large}h(C)$ and $w(i \cap C) > \varepsilon_{large}w(C)$,
- $OPT_{hor}(C)$ contains all items $i \in OPT(C)$ with $h(i \cap C) \le \varepsilon_{small}h(C)$ and $w(i \cap C) > \varepsilon_{large}w(C)$, and
- $OPT_{ver}(C)$ contains all items $i \in OPT(C)$ with $h(i \cap C) > \varepsilon_{large}h(C)$ and $w(i \cap C) \le \varepsilon_{small}w(C)$.

We call a rectangle *i intermediate* if there is a cell C such that $w(i \cap C) \in (\varepsilon_{small}w(C), \varepsilon_{large}w(C)]$ or $h(i \cap C) \in (\varepsilon_{small}w(C), \varepsilon_{large}w(C)]$. Note that a rectangle i is intermediate if and only if the last condition is satisfied for one of the at most four cells that contain a corner of i.

Lemma 29. For any constant $\varepsilon > 0$ and positive increasing function $f(\cdot)$, f(x) > x, there exist constant values $\varepsilon_{large}, \varepsilon_{small}$, with $\varepsilon \geq \varepsilon_{large} \geq f(\varepsilon_{small}) \geq \Omega_{\varepsilon}(1)$ and $\varepsilon_{small} \in \Omega_{\varepsilon}(1)$ such that the total profit of intermediate rectangles is bounded by $\varepsilon p(OPT)$.

For each cell C that is not entirely covered by some item in K(t) we add all rectangles in $OPT_{large}(C)$ that are not contained in K(t) to K(t+1). Note that here, in contrast to before, we do *not* remove small items from the packing but keep them.

Based on the items $OPT_{skew}(\mathcal{C}_t) := \cup_{C \in \mathcal{C}_t} OPT_{hor}(C) \cup OPT_{ver}(C)$ we create a corridor decomposition and consequently a box decomposition of the knapsack. To make this decomposition clearer, we assume that we first stretch the non-uniform grid into a uniform $[0,1] \times [0,1]$ grid. After this operation, for each cell C and for each element of $OPT_{hor}(C) \cup OPT_{ver}(C)$ we know that its height or width is at least $\varepsilon_{large} \cdot \frac{1}{1+2|\mathcal{K}(t)|}$. We apply Lemma 16 on the set $OPT_{skew}(\mathcal{C}_t)$ which yields a decomposition of the $[0,1] \times [0,1]$ square into at most $O_{\varepsilon,\varepsilon_{large},\mathcal{K}(t)}(1) = O_{\varepsilon,\varepsilon_{large}}(1)$ corridors. The decomposition for the stretched $[0,1] \times [0,1]$ square corresponds to the decomposition for the original knapsack, with the same items being intersected. Since we can assume that all items of OPT are placed within the knapsack so that they have integer coordinates, we can assume that the corridors of the decomposition also have integer coordinates. We can do that, because shifting the edges of the decomposition to the closest integral coordinate will not make the decomposition worse, i.e., no new items of OPT will be intersected.

We add all rectangles in $OPT_{skew}(\mathcal{C}_t)$ that are not contained in a corridor (at most $O_{\varepsilon}(1)$ many) and that are not contained in $\mathcal{K}(t)$ to K(t+1). The corridor partition has the following useful property: we started with a fixed (optimal) solution OPT for the overall problem with a fixed placement of the items in this solution. Then we considered the items in $OPT_{skew}(\mathcal{C}_t)$ and obtained the sets $OPT_{corr}\subseteq OPT_{skew}(\mathcal{C}_t)$ and $OPT_{corr}^{cross}\subseteq OPT_{corr}$. With the mentioned fixed placement, apart from the $O_{\varepsilon}(1)$ items in OPT_{corr}^{cross} , each item in OPT_{corr} is contained in one corridor. In particular, the items in OPT_{corr} do not overlap the items in $\mathcal{K}(t)$. We construct now a partition of the knapsack into $O_{\varepsilon}(1)$ corridors and $O_{\varepsilon}(1)$ containers where each container contains exactly one item from $\mathcal{K}(t)$. The main obstacle is that there can be an item $i \in \mathcal{K}(t)$ that overlaps a corridor (see Figure 6). We solve this problem by applying the following lemma on each such corridor.

Lemma 30. Let S be an open corridor with b(S) bends. Let $I' \subseteq OPT$ be a collection of items which intersect the boundary of S with $I' \cap OPT_{skew}(C_t) = \emptyset$. Then there is a collection of $|I'| \cdot b(S)$ line segments $\mathcal L$ within S which partition S into corridors with at most b(S) bends each such that no item from I' is intersected by $\mathcal L$ and there are at most $O_{\varepsilon}(|I'| \cdot b(S))$ items of $OPT_{skew}(C_t)$ intersected by line segments in $\mathcal L$.

Proof. Let $i \in I'$ and assume w.l.o.g. that i lies within a horizontal subcorridor S_i of the corridor S. If the top or bottom edge e of i lies within S_i , we define a horizontal line segment ℓ which contains the edge e and which is maximally long so that it does not intersect the interior of any item in I', and such that it does not cross the boundary curve between S_i and an adjacent subcorridor, or an edge of the boundary of S (we can assume w.l.o.g. that e does not intersect the boundary curve between S_i and some adjacent subcorridor). We say that ℓ crosses a boundary curve c (or an edge e of the boundary of S) if $c \setminus \ell$ (or $e \setminus \ell$) has two connected components.

We now "extend" each end-point of ℓ which does not lie at the boundary of some other item of I' or at the boundary of S, we call such an end point a *loose end*. For each loose end x of ℓ lying on the boundary




Figure 6: Circumventing the items in I', shown in black. The connected components between the dashed lines show the resulting new corridors.

curve c_{ij} partitioning the subcorridors S_i and S_j , we introduce a new line ℓ' perpendicular to ℓ , starting at x and crossing the subcorridor S_j such that the end point of ℓ' is maximally far away from x subject to the constraint that ℓ' does not cross an item in I', another boundary curve inside S, or the boundary of S. We continue iteratively. Since the corridor has b(S) bends, after at most b(S) iterations this operation will finish. We repeat the above operation for every item $i \in I'$, and we denote by \mathcal{L} the resulting set of line segments, see Figure 6 for a sketch. Notice that $|\mathcal{L}| = b(S) \cdot |I'|$. By construction, if an item $i \in OPT_{skew}(\mathcal{C}_t)$ is intersected by a line in \mathcal{L} then it is intersected parallel to its longer edge. Thus, each line segment in \mathcal{L} can intersect at most $O_{\mathcal{E}}(1)$ items of $OPT_{skew}(\mathcal{C}_t)$. Thus, in total there are at most $O_{\mathcal{E},\mathcal{E}_{large}}(|I'| \cdot b(S))$ items of $OPT_{skew}(\mathcal{C}_t)$ intersected by line segments in \mathcal{L} .

We apply Lemma 30 to each open corridor that intersects an item in K(t). We add all items of $I_{skew}(C_t)$ that are intersected by line segments in L to K(t+1). This adds $O_{\varepsilon}(1)$ items in total to K(t+1) since $|K(t)| \in O_{\varepsilon}(1)$ and $b(S) \leq 1/\varepsilon$ for each corridor S. For closed corridors we prove the following lemma.

Lemma 31. Let S be a closed corridor with b(S) bends. Let $OPT_{skew}(S)$ denote the items in $OPT_{skew}(\mathcal{C}_t)$ that are contained in S. Let $I' \subseteq OPT$ be a collection of items which intersect the boundary of S with $I' \cap OPT_{skew}(\mathcal{C}_t) = \emptyset$. Then there is a collection of $O_{\varepsilon}(|I'|^2/\varepsilon)$ line segments \mathcal{L} within S which partition S into a collection of closed corridors with at most $1/\varepsilon$ bends each and possibly an open corridor with b(S) bends such that no item from I' is intersected by \mathcal{L} and there is a set of items $OPT'_{skew}(S) \subseteq OPT_{skew}(S)$ with $|OPT'_{skew}(S)| \leq O_{\varepsilon}(|I'|^2)$ such that the items in $OPT_{skew}(S) \setminus OPT'_{skew}(S)$ intersected by line segments in \mathcal{L} have a total profit of at most $O(\varepsilon) \cdot p(OPT_{skew}(\mathcal{C}_t))$.

Proof. Similarly as for the case of open corridors, we take each item $i \in I'$ whose edge e is contained in S, and we create a path containing e that partitions S. Here the situation is a bit more complicated, as our newly created paths could extend over more than $\frac{1}{\varepsilon}$ bends inside S. In this case we will have to do some shortcutting, i.e., some items contained in S will be crossed parallel to their shorter edge and we cannot guarantee that their total number will be small. However, we will ensure that the total weight of such items is small. We proceed as follows (see Figure 6 for a sketch).

Consider any item $i \in I'$ and assume w.l.o.g. that i intersects a horizontal subcorridor S_i of the closed corridor S. Let e be the edge of i within S_i . For each endpoint of e we create a path p as for the case of closed corridors. If after at most $b(S) \leq 1/\varepsilon$ bends the path hits an item of I' (possibly the same item i), the boundary of S or another path created earlier, we stop the construction of the path. Otherwise, if p is the first path inside of S that did not finish after at most b(S) bends, we proceed with the construction of the path,

only now at each bend we check the total weight of the items of $OPT_{skew}(S)$ that would be crossed parallel to their shorter edge, if, instead of bending, the path would continue at the bend to hit itself. From the construction of the boundary curves in the intersection of two subcorridors we know that for two bends of the constructed path, the sets of items that would be crossed at these bends of the path are pairwise disjoint. Thus, after $O(|I'|/\varepsilon)$ bends we encounter a collection of items $OPT''_{skew}(S) \subseteq OPT_{skew}(S)$ such that $p(OPT''_{skew}(S)) \leq \frac{\varepsilon}{|I'|}p(OPT_{skew}(S))$, and we end the path p by crossing the items of $OPT''_{skew}(S)$. This operation creates an open corridor with up to $O(|I'|/\varepsilon)$ bends. We divide it into up to O(|I'|) corridors with up to $1/\varepsilon$ bends each. Via a shifting argument we can argue that this loses at most a factor of $1+\varepsilon$ in the profit due to these items. When we perform this operation for each item $i \in I'$ the total weight of items intersected parallel to their shorter edge (i.e., due to the above shortcutting) is bounded by $|I'| \cdot \frac{\varepsilon}{|I'|} p(OPT_{skew}(S)) = \varepsilon \cdot p(OPT_{skew}(S))$. This way, we introduce at most $O(|I'|^2/\varepsilon)$ line segments. Denote them by \mathcal{L} . They intersect at most $O_{\varepsilon}(|I'|^2)$ items parallel to their respective longer edge, denote them by $OPT'_{skew}(S)$. Thus, the set \mathcal{L} satisfies the claim of the lemma.

Similarly as for Lemma 30 we apply Lemma 31 to each closed corridor. We add all items in the respective set $OPT'_{skew}(S)$ to the set K(t+1) which yields $O_{\varepsilon}(1)$ many items. The items in $OPT_{skew}(S) \setminus OPT'_{skew}(S)$ are removed from the instance, as their total profit is small.

A.7.2 Partitioning corridors into boxes

Then we partition the resulting corridors into boxes according to the different cases described in Section A.5. There is one difference to the argumentation above: we define that the set OPT_{fat} contains not only skewed items contained in the respective subregions of a subcorridor, but all items contained in such a subregion. In particular, this includes items that might have been considered as small items above. Thus, when we move items from one subregion to the box associated to the subregion below (see Remark 3) then we move *every* item that is contained in that subregion. If an item is killed in one of the orderings of the subcorridors to apply the procedure from Section A.3 then we add it to K(t+1). Note that $|K(t+1)| \in O_{\varepsilon,\varepsilon_{large},\varepsilon_{box}}(1)$ and $K(t) \cap K(t+1) = \emptyset$. Also note here that we ignore for the moment small items that cross the boundary curves of the subcorridors; they will be taken care of in Section A.7.4.

A.7.3 Partitioning boxes into containers

Then we subdivide the boxes into containers. We apply Lemma 20 to each box with a slight modification. Assume that we apply it to a box of size $a \times b$ containing a set of items I_{box} . Like above we first remove the items in a thin strip of width $3\varepsilon b$ such that via a shifting argument the items (fully!) contained in this strip have a small profit of $O(\varepsilon)p(I_{box})$. However, in contrast to the setting above the set I_{box} contains not only skewed items but also small items. We call an item i small if there is no cell C such that $i \in OPT_{large}(C) \cup OPT_{hor}(C) \cup OPT_{ver}(C)$ and denote by $OPT_{small}(C_t)$ the set of small items. When we choose the strip to be removed we ensure that the profit of the removed skewed and small items is small. There are $O_{\varepsilon}(1)$ skewed items that partially (but not completely) overlap the strip whose items we remove. We add those $O_{\varepsilon}(1)$ items to K(t+1). Small items that partially overlap the strip are taken care of later in Section A.7.4, we ignore them for the moment. Then we apply Lemma 19. In contrast to the setting above, we do not only apply it to the skewed items but apply it also to small items that are contained in the box. Denote by $OPT'_{small}(C_t)$ the set of small items that are contained in some box of the box partition.

Thus, we obtain an L&C packing for the items in $\mathcal{K}(t)$, for a set of items $OPT'_{skew}(\mathcal{C}_t) \subseteq OPT_{skew}(\mathcal{C}_t)$, and for a set of items $OPT''_{small}(\mathcal{C}_t) \subseteq OPT'_{small}(\mathcal{C}_t)$ such that $p(OPT'_{skew}(\mathcal{C}_t)) + p(OPT''_{small}(\mathcal{C}_t)) + p(OPT''_{small}(\mathcal{C}_t))$

$$p(K(t+1)) \ge (1 - O(\varepsilon))p(OPT_{skew}(C_t) \cup OPT'_{small}(C_t)).$$

A.7.4 Handling small items

So far we ignored the small items in $OPT''_{small}(\mathcal{C}_t) := OPT_{small}(\mathcal{C}_t) \setminus OPT'_{small}(\mathcal{C}_t)$. This set consists of small items that in the original packing intersect a line segment of the corridor partition, the boundary of a box, or a boundary curve within a corridor. We describe now how to add them into the empty space of the so far computed packing. First, we assign each item in $OPT''_{small}(\mathcal{C}_t)$ to a grid cell. We assign each small item $i \in OPT''_{small}(\mathcal{C}_t)$ to the cell C such that in the original packing i intersects with C and the area of $i \cap C$ is not smaller than $i \cap C'$ for any cell C' ($i \cap C'$ denotes the part of i intersecting C' in the original packing for any grid cell C').

Consider a grid cell C and let $OPT''_{small}(C)$ denote the small items in $OPT''_{small}(C_t)$ assigned to C. Intuitively, we want to pack them into the empty space in the cell C that is not used by any of the containers, similarly as above. We first prove an analog of Lemma 27 of the setting above.

Lemma 32. Let C be a cell. The total area of C occupied by containers is at most $(1-2\varepsilon)a(C)$.

Proof. In our construction of the boxes we moved some of the items (within a corridor). In particular, it can happen that we moved some items into C that were originally in some other grid cell C'. This reduces the empty space in C for the items in $OPT''_{small}(C)$. Assume that there is a horizontal subcorridor H intersecting C such that some items or parts of items within H were moved into C that were not in C before. Then such items were moved vertically and the corridor containing H must intersect the upper or lower boundary of C. The part of this subcorridor lying within C has a height of at most $\varepsilon_{large} \cdot h(C)$. Thus, the total area of C lost in this way is bounded by $O(\varepsilon_{large}a(C))$ which includes analogous vertical subcorridors.

Like in Lemma 27 we argue that in each horizontal box of size $a \times b$ we remove a horizontal strip of height $3\varepsilon b$ and then the created containers lie in a box of height $(1-3\varepsilon)(1+\varepsilon_{ra})b$. In particular, if the box does not intersect the top or bottom edge of C then within C its containers use only a box of dimension $a' \times (1-3\varepsilon)(1+\varepsilon_{ra})b$ where a' denotes the width of the box within C, i.e., the width of the intersection of the box with C. If the box intersects the top or bottom edge of C then we cannot guarantee that the free space lies within C. However, the total area of such boxes is bounded by $O(\varepsilon_{large}a(C))$. We can apply a symmetric argument to vertical boxes. Then, the total area of C used by containers is at most $(1-3\varepsilon)(1+\varepsilon_{ra})a(C)+O(\varepsilon_{large}a(C))\leq (1-2\varepsilon)a(C)$. This gives the claim of the lemma. \Box

Next, we argue that the items in $OPT''_{small}(C)$ have very small total area. Recall that they are the items intersecting C that are not contained in a box. The total number of boxes and boundary curves intersecting C is $O_{\varepsilon,\varepsilon_{large}}(1)$ and in particular, this quantity does not depend on ε_{small} . Hence, by choosing ε_{small} sufficiently small, we can ensure that $a(OPT''_{small}(C)) \leq \varepsilon a(C)$. Then, similarly as in Lemma 28 we can argue that if ε_{small} is small enough then we can pack the items in $OPT''_{small}(C)$ using NFDH into the empty space within C.

A.7.5 L&C packings

We iterate the above construction, obtaining pairwise disjoint sets K(1), K(2), ... until we find a set K(k) such that $p(K(k)) \le \varepsilon \cdot OPT$. Since the sets K(0), K(1), ... are pairwise disjoint there must be such a value k with $k \le 1/\varepsilon$. Thus, $|\mathcal{K}(k-1)| \le O_{\varepsilon}(1)$. We build the grid given by the x- and y-coordinates of $\mathcal{K}(k-1)$, giving a set of cells \mathcal{C}_k . As described above we define the corridor partition, the partition of the

corridors into boxes (with the different orders to process the subcorridors as described in Section A.3) and finally into containers. Denote by $OPT_{small}(\mathcal{C}_k)$ the resulting set of small items.

We consider the candidate packings based on the grid C_k . For each of the six candidate packings with a degenerate L we can pack almost all small items of the original packing. We define $I_{\rm lc}$ and $I_{\rm sc}$ the sets of items in long and short subcorridors in the initial corridor partition, respectively. Exactly as in the cardinality case, a subcorridor is long if it is longer than N/2 and short otherwise. As before we divide the items into fat and thin items and define the sets OPT_{SF} , OPT_{LT} , and OPT_{ST} accordingly. Moreover, we define the set OPT_{LF} to contain all items in $I_{\rm lc}$ that are fat in all candidate packings plus the items in $\mathcal{K}(k-1)$.

Thus, we obtain the respective claims of Lemmas 23, 24, and 25 in the weighted setting. For the following lemma let $OPT_{small} := OPT_{small}(\mathcal{C}_k)$.

Lemma 33. Let $OPT_{L\&C}$ the most profitable solution that is packed by an L&C packing.

(a)
$$p(OPT_{L\&C}) \ge (1 - O(\varepsilon))(p(OPT_{LF}) + p(OPT_{SF}) + p(OPT_{small}))$$

(b)
$$p(OPT_{L\&C}) \ge (1 - O(\varepsilon))(p(OPT_{LF}) + p(OPT_{SF})/2 + p(OPT_{LT})/2 + p(OPT_{small}))$$

(c)
$$p(OPT_{L\&C}) \ge (1 - O(\varepsilon))(p(OPT_{LF}) + p(OPT_{SF})/2 + p(OPT_{ST})/2 + p(OPT_{small}))$$
.

For the candidate packing for the non-degenerate-L case (Lemma 26 in Section A.5) we first add the small items as described above. Then we remove the items in $\mathcal{K}(k-1)$. Then, like above, with a random shift we delete items touching a horizontal and a vertical strip of width $3\varepsilon N$. Like before, each item i is still contained in the resulting solution with probability $1/2-15\varepsilon$ (note that we cannot make such a claim for the items in $\mathcal{K}(k-1)$). For each small item we can even argue that it still contained in the resulting solution with probability $1-O(\varepsilon)$ (since it is that small in both dimensions). We proceed with the construction of the boundary L and the assignment of the items into it like in the unweighted case.

Lemma 34. For the solution
$$OPT_{L\&C}$$
 we have that $p(OPT_{L\&C}) \ge (1-O(\varepsilon))(\frac{3}{4}p(OPT_{LT})+p(OPT_{ST})+\frac{1-O(\varepsilon)}{2}p(OPT_{SF})+p(OPT_{small}))$.

When we combine Lemmas 33 and 34 we conclude that $p(OPT_{L\&C}) \ge (17/9 + O(\varepsilon))p(OPT)$. Similarly as before, the worst case is obtained, up to $1 - O(\varepsilon)$ factors, when we have that $p(OPT_{LT}) = p(OPT_{SF}) = p(OPT_{ST})$, $p(OPT_{LF}) = 5p(OPT_{LT})/4$, and $p(OPT_{small}) = 0$. This completes the proof of Lemma 21.

A.8 Main algorithm

In this Section we present our main algorithm for the weighted case of 2DK. It is in fact an approximation scheme for L&C packings. Its approximation factor therefore follows immediately from Lemma 21.

Given $\epsilon > 0$, we first guess the quantities ε_{large} , ε_{small} , the proof of Lemma 15 reveals that there are only $2/\epsilon + 1$ values we need to consider. We choose $\varepsilon_{ring} := \epsilon^2$ and subsequently define ε_{box} according to Lemma 18. Our algorithm combines two basic packing procedures. The first one is the following standard PTAS to pack items into a constant number of containers. The same basic approach works also with rotations. The basic idea is to reduce the problem to an instance of the Maximum Generalized Assignment Problem (GAP) with one bin per container, and then use a PTAS for the latter problem plus Next Fit Decreasing Height to repack items in area containers.

Lemma 35. There is a PTAS for the problem of computing a maximum profit packing of a subset of items of a given set I' into a given set of containers of constant cardinality, both with and without rotations.

The second packing procedure is the PTAS for the L-packing problem, see Theorem 1.

To use these packing procedures, we first guess whether the optimal L&C-packing due to Lemma 21 uses a non-degenerate boundary L. If yes, we guess a parameter ℓ which denotes the minimum height of the vertical items in the boundary L and the minimum width of the horizontal items in the boundary L. For ℓ we allow all heights and widths of the input items that are larger than N/2, i.e., at most 2n values. Let I_{long} be the items whose longer side has length at least ℓ (hence longer than N/2). We set the width of the boundary L to be $\epsilon^2 N$ and solve the resulting instance (L, I_{long}) optimally using the PTAS for L-packings due to Theorem 1. Then we enumerate all the possible subsets of non-overlapping containers in the space not occupied by the boundary L (or in the full knapsack, in the case of a degenerate L), where the number and sizes of the containers are defined as in Lemma 20. In particular, there are at most $O_{\varepsilon}(1)$ containers and there is a set of size $n^{O_{\varepsilon}(1)}$ that we can compute in polynomial time such that the height and the width of each container is contained in this set. We compute an approximate solution for the resulting container packing instance with items $I_{\rm short} = I \setminus I_{\rm long}$ using the PTAS from Lemma 35. Finally, we output the most profitable solution that we computed.

B Cardinality case without rotations

In this section, we present a refined algorithm with approximation factor of $\frac{558}{325} + \varepsilon < 1.717$ for the cardinality case when rotations are not allowed.

Theorem 36. There exists a polynomial-time $\frac{558}{325} + \varepsilon < 1.717$ -approximation algorithm for cardinality 2DK.

Along this section, since the profit of each item is equal to 1, instead of p(I) for a set of items I we will just write |I|. We will use most of the notation defined in Section A. Recall that for two given constants $0 < \varepsilon_{small} < \varepsilon_{large} \le 1$, we partition the instance into:

- I_{small} , the set of rectangles with $h_i, w_i \leq \varepsilon_{small} N$, and we denote them as *small* rectangles;
- I_{large} , the set of rectangles with $h_i, w_i > \varepsilon_{large}N$, and we denote them as large rectangles;
- I_{hor} , the set of rectangles with $w_i > \varepsilon_{large}N$ and $h_i \le \varepsilon_{small}N$, and we denote them as horizontal rectangles;
- I_{ver} , the set of rectangles with $h_i > \varepsilon_{large}N$ and $w_i \le \varepsilon_{small}N$, and we denote them as vertical rectangles;
- I_{int} , the set of remaining rectangles, and we denote them as *intermediate* rectangles.

The corresponding intersection with OPT defines the sets OPT_{small} , OPT_{large} , OPT_{hor} , OPT_{ver} and OPT_{int} , respectively. As discussed in Section 3, since any feasible solution contains at most $\frac{1}{\varepsilon_{large}^2}$ large rectangles, we can assume in this case that $OPT_{large} = \emptyset$. Furthermore, thanks to Lemma 15, ε_{small} and ε_{large} can be chosen in such a way that $\varepsilon_{small} \leq \varepsilon_{large} \leq \varepsilon$, ε_{small} differs from ε_{large} by a large factor and $|OPT_{int}| \leq \varepsilon |OPT|$. Building upon the corridors decomposition from [2], we will again consider OPT_T (thin rectangles), OPT_F (fat rectangles) and OPT_K (killed rectangles) as defined in Section A.5. Thanks to Lemma 18, $|OPT_K| = O_\varepsilon(1)$ and all the involved parameters can be fixed in such a way that the total height (resp. width) of $OPT_T \cap I_{hor}$ (resp. $OPT_T \cap I_{ver}$) is at most εN . Recall that a subcorridor is called long if its shortest edge has length at least $\frac{N}{2}$ and short otherwise. In the analysis of the algorithm we will again use

sets OPT_{LF} , OPT_{LT} , OPT_{SF} and OPT_{ST} as defined in Section A.3, corresponding to rectangles from OPT_F inside long corridors, rectangles from OPT_F inside long corridors, rectangles from OPT_F inside short corridors and rectangles from OPT_T inside short corridors respectively. For a given $\ell \in (\frac{N}{2}, N]$, we let $I_{long} \subseteq I$ be the rectangles whose longest side has length longer than ℓ and $I_{short} = I \setminus I_{long}$. We will assume as in the proof of Lemma 26 that $\ell = (\frac{1}{2} + 2\varepsilon_{large}) N$. That way we make sure that no long rectangle belongs to a short subcorridor (however it is worth remarking that long corridors may contain short rectangles).

Let us define $OPT_{\mathrm{long}} := I_{\mathrm{long}} \cap OPT$ and $OPT_{\mathrm{short}} := I_{\mathrm{short}} \cap OPT$. Let us define $\varepsilon_L = \sqrt{\varepsilon}$. Note that $\varepsilon_L \geq \varepsilon \geq \varepsilon_{large} \geq \varepsilon_{small}$. For simplicity and readability of the section, sometimes we will slightly abuse the notation and for any small constant depending on $\varepsilon, \varepsilon_{large}, \varepsilon_{small}$, we will just use $O(\varepsilon_L)$. Now we give a brief informal overview of the refinement and the cases before we go to the details.

Overview of the refined packing. For the refined packing we will consider several L&C packings. Some of the packings are just extensions of previous constructions (such as from Theorem 14 and Lemma 33). Then we consider several other new L&C packings where an L-region is packed with items from I_{long} and the remaining region is used for packing items from I_{short} using Steinberg's theorem (See Theorem 38). Note that in the definition of L&C packing in Section A, we assumed the height of the horizontal part of L-region and width of the vertical part of L-region to be the same. However, for these new packings we will consider L-regions where the height of the horizontal part and width of vertical part may differ. Now several cases arise depending on the structure and profit of the L-region. To pack items in OPT_{short} we have three options:

- 1. We can pack items in $I_{\rm short}$ using Steinberg's theorem into one rectangular region. Then we need both sides of the region to be greater than $\frac{1}{2} + 2\varepsilon_{large}$.
- 2. We can pack items in $I_{\rm short}$ using Steinberg's theorem such that vertical and horizontal items are packed separately into different vertical and horizontal rectangular regions inside the knapsack.
- 3. If $a(OPT_{\rm short})$ is large, we might pack only a small region with items in $OPT_{\rm long}$, and use the remaining larger space in the knapsack to pack a significant fraction of profit from $OPT_{\rm short}$.

Now depending on the structure of the L-packing and $a(OPT_{\rm short})$, we arrive at several different cases. If the L-region has very small width and height, we have case (1). Else if the L-region has very large width (or height), we have case (2B), where we pack nearly $\frac{1}{2}|OPT_{\rm long}|$ in the L-region and then pack items from $I_{\rm short}$ in one large rectangular region. Otherwise, we have case (2A), where either we pack only items from $OPT_{\rm long}\cap OPT_T$ (See Lemma 40, used in case: (2Ai)) or nearly $3/4|OPT_{\rm long}|$ (See Lemma 41, used in cases (2Aii), (2Aiiia)) or in another case, we pack the vertical and horizontal items in $OPT_{\rm short}$ in two different regions through a more complicated packing (See case (2Aiiib)). The details of these cases can be found in the proof of Theorem 36.

Now first, we start with some extensions of previous packings. Note that by using analogous arguments as in the proof of Theorem 14, we can derive the following inequalities leading to a $\left(\frac{16}{9} + O(\varepsilon_L)\right)$ -approximation algorithm.

$$|OPT_{L\&C}| \ge \frac{3}{4}|OPT_{\text{long}}|\tag{1}$$

$$|OPT_{L\&C}| \ge \left(\frac{1}{2} - O(\varepsilon_L)\right)|OPT_{long}| + \left(\frac{3}{4} - O(\varepsilon_L)\right)|OPT_{short}|$$
 (2)

Now from Lemma 18, items in $OPT_{\mathrm{short}} \cap OPT_T$ can be packed into two containers of size $\ell \times \varepsilon N$ and $\varepsilon N \times \ell$. We can adapt part of the results in Lemma 33 to obtain the following inequalities.

Proposition 37. The following inequalities hold:

$$|OPT_{L\&C}| \ge (1 - O(\varepsilon_L))(|OPT_{long} \setminus OPT_T| + |OPT_{short} \setminus OPT_T|).$$
 (3)

$$|OPT_{L\&C}| \ge (1 - O(\varepsilon_L))(|OPT_{\text{long}} \setminus OPT_T| + \frac{1}{2}(|OPT_{\text{short}} \setminus OPT_T| + |OPT_{\text{long}} \cap OPT_T|)). \quad (4)$$

Proof. Inequality (3) follows directly from Lemma 33 since $OPT_{LF} \cup OPT_{SF} \cup OPT_{small} = (OPT_{long} \setminus OPT_T) \cup (OPT_{short} \setminus OPT_T)$ and both sets are disjoint. Inequality (4) follows from Lemma 24: if we consider the sum of the number of packed rectangles corresponding to the 4 subcases associated with the case "short horizontal/short vertical", then every $i \in OPT_{long} \setminus OPT_T \subseteq OPT_{LF}$ appears four times, every $i \in OPT_{short} \cap OPT_{LF}$ appears four times, every $i \in OPT_{SF}$ appears twice and every $i \in OPT_{long} \cap OPT_T$ appears twice. After including a $(1 - O(\varepsilon_L))$ fraction of OPT_{small} , and since $(OPT_{short} \cap OPT_{LF}) \cup OPT_{SF} \cup OPT_{small} = OPT_{short} \setminus OPT_T$, the inequality follows by taking average of the four packings. □

The following theorem due to Steinberg [32] will be useful to pack items from $OPT_{\rm short}$ in order to obtain better packings.

Theorem 38 (A. Steinberg [32]). We are given a set of rectangles I' and a box Q of size $w \times h$. Let $w_{max} \leq w$ and $h_{max} \leq h$ be the maximum width and maximum height among the items in I' respectively. Also we denote $x_+ := max(x, 0)$. If

$$2a(I') \le wh - (2w_{max} - w)_{+}(2h_{max} - h)_{+}$$

then I' can be packed into Q.

Corollary 39. Let I' be a set of rectangles such that $\max_{i \in I'} h(i) \leq \left(\frac{1}{2} + 2\varepsilon_{large}\right) N$ and $\max_{i \in I'} w(i) \leq \left(\frac{1}{2} + 2\varepsilon_{large}\right) N$. Then for any $\alpha, \beta \leq \frac{1}{2} - 2\varepsilon_{large}$, all of I' can be packed into a knapsack of width $(1-\alpha)N$ and height $(1-\beta)N$ if

$$a(I') \leq \left(\frac{1}{2} - (\alpha + \beta) \left(\frac{1}{2} + 2\varepsilon_{large}\right) - 8\varepsilon_{large}^2\right) N^2.$$

Now we prove a more general version of Lemma 26 which holds for the cardinality case.

Lemma 40. If $a(OPT_{short} \setminus OPT_T) \leq \gamma N^2$ for any $\gamma \leq 1$, then

$$|OPT_{L\&C}| \ge \frac{3}{4}|OPT_{\operatorname{long}} \cap OPT_T| + |OPT_{\operatorname{short}} \cap OPT_T| + \min\left\{1, \frac{1 - O(\varepsilon_L)}{2\gamma}\right\}|OPT_{\operatorname{short}} \setminus OPT_T|.$$

Proof. As in Lemma 26, we can pack $\frac{3}{4}|OPT_{\log}\cap OPT_T| + |OPT_{\text{short}}\cap OPT_T|$ many rectangles in a boundary L-region plus two boxes on the other two sides of the knapsack and then a free square region with side length $(1-3\varepsilon)N$ can be used to pack items from $OPT_{\text{short}}\setminus OPT_T$. From Corollary 39, any subset of rectangles of $OPT_{\text{short}}\setminus OPT_T$ with total area at most $(1-O(\varepsilon_L))N^2/2$ can be packed into that square region of length $(1-3\varepsilon)N$. Thus we sort rectangles from $OPT_{\text{short}}\setminus OPT_T$ in the order of nondecreasing area and iteratively pick them until their total area reaches $(1-O(\varepsilon_L)-\varepsilon_{small})N^2/2$. Using Steinberg's theorem, there exists a packing of the selected rectangles. If $2\gamma \leq 1-O(\varepsilon_L)-\varepsilon_{small}$ then the profit of this packing is $|OPT_{\text{short}}\setminus OPT_T|$, and otherwise the total profit is at least $\frac{1-O(\varepsilon_L)}{2\gamma}|OPT_{\text{short}}\setminus OPT_T|$. The packing coming from Steinberg's theorem may not be container-based, but we can then use resource augmentation as in Lemma 26 to obtain an L&C packing.

Now the following lemma will be useful when $a(OPT_{short})$ is large.

Lemma 41. If $a(OPT_{\text{short}}) > \gamma N^2$ for any $\gamma \geq \frac{3}{4} + \varepsilon + \varepsilon_{large}$, then

$$|OPT_{L\&C}| \geq \frac{3}{4}|OPT_{\mathrm{long}}| + \frac{(3\gamma - 1 - O(\varepsilon_L))}{4\gamma}|OPT_{\mathrm{short}}|.$$

Proof. Similar to Lemma 13 in Section 3, we start from the optimal packing and move all rectangles in OPT_{long} to the boundary such that all of them are contained in a boundary ring. Note that unlike the case when we only pack $OPT_{long} \cap OPT_T$ in the boundary region, the boundary ring formed by OPT_{long} may have width or height $\gg \varepsilon N$. Let us call the 4 stacks in the ring to be subrings. Let us assume that left and right subrings have width $\alpha_{left}N$ and $\alpha_{right}N$ respectively and bottom and top subrings have height $\beta_{bottom}N$ and $\beta_{top}N$ respectively. By possibly killing one of the long rectangles, subrings can be arranged such that $\alpha_{left}, \alpha_{right}, \beta_{bottom}, \beta_{top} \leq 1/2$: If no vertical rectangle intersects the vertical line $x = \frac{N}{2}$ and no horizontal rectangle intersects the horizontal line $y = \frac{N}{2}$ this property holds directly. If one of the previous cases is not satisfied, by deleting such rectangle we can ensure the desired property at a negligible loss of profit, and notice that it is not possible that both cases happen at the same time since rectangles are long.

As $a(OPT_{\rm short}) > \gamma N^2$, then $a(OPT_{\rm long}) < (1-\gamma)N^2$. Let us define $\alpha = \alpha_{left} + \alpha_{right}$ and $\beta = \beta_{bottom} + \beta_{top}$. Then $(\alpha + \beta)N \cdot \frac{N}{2} \leq a(OPT_{\rm long})$, which implies that $\frac{\alpha+\beta}{2} < 1-\gamma$. Hence, we get the following two inequalities:

$$(\alpha + \beta) \le 2(1 - \gamma);\tag{5}$$

$$a(OPT_{\text{short}}) \le N^2 - a(OPT_{long}) \le \left(1 - \frac{(\alpha + \beta)}{2}\right)N^2.$$
 (6)

Now consider the case when we remove the top horizontal subring and construct a boundary L-region as in Lemma 13. We will assume that rectangles in the L-region are pushed to the left and bottom as much as possible. Then, the boundary L-region has width $(\alpha_{left} + \alpha_{right})N$ and height $\beta_{bottom}N$. We will use Steinberg's theorem to show the existence of a packing of rectangles from $OPT_{\rm short}$ in a subregion of the remaining space with width $N-(\alpha_{left}+\alpha_{right}+\varepsilon)N$ and height $N-(\beta_{bottom}+\varepsilon)N$, and use the rest of the area for resource augmentation to get an L&C-based packing. Since $\gamma \geq \frac{3}{4}+\varepsilon+\varepsilon_{large}$, we have from (5) that $\alpha+\beta+2\varepsilon\leq 2(1-\gamma)+2\varepsilon\leq 1/2-2\varepsilon_{large}$. So $\alpha+\varepsilon\leq 1/2-2\varepsilon_{large}$ and $\beta+\varepsilon\leq 1/2-2\varepsilon_{large}$. Thus from Corollary 39, in the region with width $N-(\alpha_{left}+\alpha_{right}+\varepsilon)N$ and height $N(1-\beta_{bottom}-\varepsilon)$ we can pack rectangles from $OPT_{\rm short}$ of total area at most $\left(\frac{1}{2}-\frac{(\alpha_{left}+\alpha_{right}+\beta_{bottom})}{2}-O(\varepsilon_L)\right)N^2$. Hence, we can take the rectangles in $OPT_{\rm short}$ in the order of nondecreasing area until their total area reaches $\left(\frac{1}{2}-\frac{(\alpha_{left}+\alpha_{right}+\beta_{bottom})}{2}-O(\varepsilon_L)-\varepsilon_{small}\right)N^2$ and pack at least $|OPT_{\rm short}|\cdot\frac{\left(\frac{1}{2}-\frac{(\alpha_{left}+\alpha_{right}+\beta_{bottom})}{2}-O(\varepsilon_L)-\varepsilon_{small}\right)}{(1-\frac{(\alpha+\beta)}{2})}$ using Steinberg's theorem. If we now consider all the four different cases corresponding to removal of the four different subrings and take the average of profits obtained in each case, we pack at least

$$\frac{3}{4}|OPT_{\text{long}}| + |OPT_{\text{short}}| \cdot \left(\frac{\left(\frac{1}{2} - \frac{3}{8}(\alpha_{left} + \alpha_{right} + \beta_{bottom} + \beta_{top}) - O(\varepsilon_L)}{\left(1 - \frac{(\alpha + \beta)}{2}\right)}\right)$$

$$= \frac{3}{4}|OPT_{\text{long}}| + |OPT_{\text{short}}| \cdot \left(\frac{\left(\frac{1}{2} - \frac{3}{8}(\alpha + \beta) - O(\varepsilon_L)\right)}{\left(1 - \frac{(\alpha + \beta)}{2}\right)}\right)$$

$$\geq \frac{3}{4}|OPT_{\text{long}}| + |OPT_{\text{short}}| \cdot \frac{3\gamma - 1 - O(\varepsilon_L)}{4\gamma},$$




Figure 7: Summary of the cases.

where the last inequality follows from (5) and the fact that the expression is decreasing as a function of $(\alpha + \beta)$.

Now we start with the proof of Theorem 36.

Proof of Theorem 36. In the refined analysis, we will consider different solutions and show that the best of them always achieves the claimed approximation guarantee. We will pack some rectangles in a boundary L-region (either a subset of only $OPT_{long} \cap OPT_{T}$ or a subset of OPT_{long}) using the PTAS for L-packings described in Section 2, and in the remaining area of the knapsack (outside of the boundary L-region), we will pack a subset of rectangles from OPT_{short} .

Consider the ring as constructed in the beginning of the proof of Lemma 41. Then we remove the least profitable subring and repack the remaining rectangles from OPT_{long} in a boundary L-region. W.l.o.g. assume that the horizontal top subring was the least profitable subring. The other cases are analogous. We will use the same notation as in Lemma 41, and also define $w_L = (\alpha_{left} + \alpha_{right}), h_L = \beta_{bottom}$. Now let us consider two cases (see Figure 7 for an overview of the subcases of case 2).

• Case 1. $w_L \leq \varepsilon_L, h_L \leq \varepsilon_L$.


In this case, following the proof of Lemma 40 (using $\gamma=1$), we can pack $\frac{3}{4}|OPT_{\rm long}|+|OPT_{\rm short}\cap OPT_T|+\frac{1-O(\varepsilon_L)}{2}|OPT_{\rm short}\setminus OPT_T|$. This along with inequalities (2), (3) and (4) will give us a solution with good enough approximation factor. Check Section B.1 and Table 1 for details.


• Case 2. $w_L > \varepsilon_L$ or $h_L > \varepsilon_L$.


Let $V_{1-2\varepsilon_L}$ be the set of vertical rectangles having height strictly larger than $(1-2\varepsilon_L)N$. Let us define $w(V_{1-2\varepsilon_L}) = \sum_{i \in V_{1-2\varepsilon_L}} w(i)$. Similarly, let $H_{1-2\varepsilon_L}$ be the set of horizontal rectangles of width strictly larger than $(1-2\varepsilon_L)N$ and $h(H_{1-2\varepsilon_L}) = \sum_{i \in H_{1-2\varepsilon_L}} h(i)$.

$$\Diamond$$
 Case 2A. $\left(w_L > \varepsilon_L \text{ and } h_L > \varepsilon_L\right)$ or $\left(w_L > \varepsilon_L, h_L \leq \varepsilon_L, \text{ and } w(V_{1-2\varepsilon_L}) \leq \left(\frac{1}{2} - 2\varepsilon_{large}\right)N\right)$ or $\left(h_L > \varepsilon_L, w_L \leq \varepsilon_L, \text{ and } h(H_{1-2\varepsilon_L}) \leq \left(\frac{1}{2} - 2\varepsilon_{large}\right)N\right)$.

We will show that if any of the above three conditions is met, then we can pack $\frac{3(1-O(\varepsilon))}{4}|OPT_{\text{long}}| + |OPT_{\text{short}} \cap OPT_T|$ in a boundary L-region of width close to w_LN and height close to h_LN , and then in









(b) Packing of rectangles in $OPT_{\mathrm{long}} \cup (OPT_{\mathrm{short}} \cap OPT_T)$. Dark gray rectangles are from $OPT_{\mathrm{short}} \cap OPT_T$.

Figure 8: The case for $w_L > \varepsilon_L$ and $h_L > \varepsilon_L$.



(a) Packing of L-region using rectangles from OPT_{long} . Striped strip is the cheapest εN -width strip.



(b) Packing of rectangles in $OPT_{long} \cup (OPT_{short} \cap OPT_T)$. Dark gray rectangles are from $OPT_{short} \cap OPT_T$.

Figure 9: The case for $w_L > \varepsilon_L$ and $h_L \le \varepsilon_L$.

the remaining area we will pack some rectangles from $OPT_{\mathrm{short}} \setminus OPT_T$ using Steinberg's theorem and resource augmentation.

Packing of subset of rectangles from $OPT_{long} \cup (OPT_{short} \cap OPT_T)$ into L-region.

If $(w_L > \varepsilon_L \text{ and } h_L > \varepsilon_L)$, we partition the vertical part of the L-region into consecutive strips of width εN . Consider the strip that intersects the least number of vertical rectangles from OPT_{long} among all strips, and we call it to be the *cheapest* εN -width vertical strip (See Figure 8a). Clearly the cheapest εN -width vertical strip intersects at most a $\frac{\varepsilon + 2\varepsilon_{small}}{\varepsilon_L} \leq 3\varepsilon_L$ fraction of the rectangles in the vertical part of the L-region, so we can remove all such vertical rectangles intersected by that strip at a small loss of profit. Similarly, we remove the horizontal rectangles intersected by the cheapest εN -height horizontal strip in the boundary L-region. We now pack the horizontal container for $OPT_{short} \cap OPT_T$ in the free region left by the removed horizontal strip, and the vertical container for $OPT_{short} \cap OPT_T$ in the free region left by the removed vertical strip. Similarly to the proof of Lemma 13 we can sort rectangles in the vertical (resp. horizontal)

pile of the L-region according to their height (resp. width), obtaining a feasible L&C-packing (See Figure 8b).

In the other case $(w_L > \varepsilon_L, h_L \le \varepsilon_L \text{ and } w(V_{1-2\varepsilon_L}) \le \left(\frac{1}{2} - 2\varepsilon_{large}\right) N)$, we can again remove the cheapest εN -width vertical strip in the boundary L-region and pack the vertical container for $OPT_{\mathrm{short}} \cap OPT_T$ there (See Figure 9a). Now we show how to pack horizontal items from $OPT_{\mathrm{short}} \cap OPT_T$. In the packing of the boundary L-region, we can assume that the vertical rectangles are sorted non-increasingly by height from left to right and pushed upwards until they touch the top boundary. Then, since $w(V_{1-2\varepsilon_L}) \le \left(\frac{1}{2} - 2\varepsilon_{large}\right) N$ and $(h_L \le \varepsilon_L)$, the region $\left[\left(\frac{1}{2} - 2\varepsilon_{large}\right) N, N\right] \times \left[\varepsilon_L N, 2\varepsilon_L N\right]$ will be completely empty and thus we will have enough space to pack the horizontal container for $OPT_{\mathrm{short}} \cap OPT_T$ on top of the horizontal part of the L-region (See Figure 9b). This leads to a packing in a boundary L-region of width at most $w_L N$ and height at most $w_L N$ with total profit at least $\frac{3(1-O(\varepsilon))}{4}|OPT_{\mathrm{long}}| + |OPT_{\mathrm{short}} \cap OPT_T|$. The last case, when $w_L \le \varepsilon_L$, is analogous, leading to a packing into a boundary L-region of width at most $w_L N$ and height at most $w_L N$ with at least the same profit. Thus,

$$|OPT_{L\&C}| \ge \frac{3(1 - O(\varepsilon_L))}{4}|OPT_{long}| + |OPT_{short} \cap OPT_T|$$
 (7)

Packing of a subset of rectangles from $OPT_{\mathrm{short}} \setminus OPT_T$ into the remaining region.

Note that after packing at least $\frac{3(1-O(\varepsilon))}{4}|OPT_{\mathrm{long}}| + |OPT_{\mathrm{short}} \cap OPT_{T}|$ many rectangles in the boundary L-region, the remaining rectangular region of width $(1-w_L-\varepsilon_L)N$ and height $(1-h_L-\varepsilon_L)N$ is completely empty. Now we will show the existence of a packing of some rectangles from $OPT_{\mathrm{short}} \setminus OPT_{T}$ in the remaining space of the packing (even using some space from the L-boundary region). Let $(OPT_{\mathrm{short}} \setminus OPT_{T})_{hor} := ((OPT_{\mathrm{short}} \setminus OPT_{T}) \cap I_{hor}) \cup ((OPT_{\mathrm{short}} \setminus OPT_{T}) \cap I_{small})$ and $(OPT_{\mathrm{short}} \setminus OPT_{T})_{ver} := (OPT_{\mathrm{short}} \setminus OPT_{T}) \cap I_{ver}$. Let us assume w.l.o.g. that vertical rectangles are shifted as much as possible to the left and top of the knapsack and horizontal ones are pushed as much as possible to the right and bottom. We divide the analysis in three subcases depending on $a(OPT_{\mathrm{short}} \setminus OPT_{T})$.

- Subcase (i). If $a(OPT_{\text{short}} \setminus OPT_T) \leq \frac{3}{5}N^2$, from inequalities (2), (3), (4), (7) and Lemma 40, we get a solution with good enough approximation factor. Check Section B.1 and Table 1 for details.
- Subcase (ii). If $a(OPT_{short} \setminus OPT_T) > (\frac{3}{4} + \varepsilon + \varepsilon_{large})N^2$, from inequalities (2), (3), (4), (7) and Lemma 41, we get a solution with good enough approximation factor. Check Section B.1 and Table 1 for details.
- Subcase (iii). Finally, if $\frac{3}{5}N^2 \le a(OPT_{\rm short} \setminus OPT_T) \le (\frac{3}{4} + \varepsilon + \varepsilon_{large})N^2$, from inequality (5) we get $\alpha + \beta \le 2(1 \frac{3}{5}) = \frac{4}{5}$. Now we consider two subcases.
- \odot Subcase (iii a): $w_L \leq \frac{1}{2}$ and $h_L \leq \frac{1}{2}$. Note that in this case if $w_L \geq \frac{1}{2} 2\varepsilon_{large} 2\varepsilon_L$ (resp., $h_L \geq \frac{1}{2} 2\varepsilon_{large} 2\varepsilon_L$), we can remove the cheapest $2(\varepsilon_L + \varepsilon_{large})N$ -width vertical (resp., horizontal) strip from the L-region by removing $O(\varepsilon_L)$ fraction of rectangles in OPT_{long} . Otherwise we have $w_L < \frac{1}{2} 2\varepsilon_{large} 2\varepsilon_L$ and $h_L < \frac{1}{2} 2\varepsilon_{large} 2\varepsilon_L$. So there is a free rectangular region that has both side lengths at least $N(\frac{1}{2} + 2\varepsilon_{large} + \varepsilon_L)$; we will keep $\varepsilon_L N$ width and $\varepsilon_L N$ height for resource augmentation and use the rest of the rectangular region (with both sides length at least $(\frac{1}{2} + 2\varepsilon_{large})N$) for showing existence of a packing using Steinberg's theorem.

Note that this free rectangular region has area at least $N^2(1-w_L-2\varepsilon_L)(1-h_L-2\varepsilon_L)$. Now consider rectangles from $(OPT_{\rm short}\setminus OPT_T)_{hor}$ (by sorting them non-decreasingly by area and picking them iteratively) until their total area becomes at least $min\{a((OPT_{\rm short}\setminus OPT_T)_{hor}), \frac{N^2(1-w_L-2\varepsilon_L)(1-h_L-2\varepsilon_L)}{2} - \varepsilon_{small}N^2\}$. Thus their total area is at most $\leq \frac{N^2(1-w_L-2\varepsilon_L)(1-h_L-2\varepsilon_L)}{2}$ as the area of any rectangle in $(OPT_{\rm short}\setminus OPT_T)_{hor}$ is at most $\varepsilon_{small}N^2$. Hence, from Steinberg's theorem, we can pack these rectangles in the free rectangular region. Similarly, we can pack there rectangles from $(OPT_{\rm short}\setminus OPT_T)_{ver}$

with total area at least $min\{a((OPT_{\mathrm{short}} \setminus OPT_T)_{ver}), \frac{N^2(1-w_L-2\varepsilon_L)(1-h_L-2\varepsilon_L)}{2} - \varepsilon_{small}N^2\}.$

Since rectangles are sorted non-decreasingly according to their areas, the total profit of the aforementioned packings is bounded below by $min\{1, \left(\frac{(1-w_L)(1-h_L)}{2a((OPT_{\rm short}\setminus OPT_T)_{hor})} - O(\varepsilon_L)\right)N^2\}|(OPT_{\rm short}\setminus OPT_T)_{hor}|$ and $min\{1, \left(\frac{(1-w_L)(1-h_L)}{2a((OPT_{\rm short}\setminus OPT_T)_{ver})} - O(\varepsilon_L)\right)N^2\}|(OPT_{\rm short}\setminus OPT_T)_{ver}|$ respectively. We claim that if we keep the best of the two packings, we can always pack at least $\left(\frac{7}{48} - O(\varepsilon_L)\right)|OPT_{\rm short}\setminus OPT_T|$ many rectangles. To show this we will consider the four possible cases:

- If $min\{1, \left(\frac{(1-w_L)(1-h_L)}{2a((OPT_{\rm short}\setminus OPT_T)_{hor})} O(\varepsilon_L)\right)N^2\} = min\{1, \left(\frac{(1-w_L)(1-h_L)}{2a((OPT_{\rm short}\setminus OPT_T)_{ver})} O(\varepsilon_L)\right)N^2\} = 1$, then, by an averaging argument, the best among the two packings has profit at least $\frac{1}{2}(|(OPT_{\rm short}\setminus OPT_T)_{ver}| + |(OPT_{\rm short}\setminus OPT_T)_{hor}|) = \frac{1}{2}|OPT_{\rm short}\setminus OPT_T|$.
- If $\left(\frac{(1-w_L)(1-h_L)}{2a((OPT_{\mathrm{short}}\setminus OPT_T)_{hor})} O(\varepsilon_L)\right)N^2 < 1$ and $\left(\frac{(1-w_L)(1-h_L)}{2a((OPT_{\mathrm{short}}\setminus OPT_T)_{ver})} O(\varepsilon_L)\right)N^2 < 1$, then by an averaging argument we pack at least

$$\begin{split} &\frac{N^2}{2} \left((\frac{(1-w_L)(1-h_L)}{2a((OPT_{\text{short}} \backslash OPT_T)_{hor})} - O(\varepsilon_L)) | (OPT_{\text{short}} \backslash OPT_T)_{hor}| + (\frac{(1-w_L)(1-h_L)}{2a((OPT_{\text{short}} \backslash OPT_T)_{ver})} - O(\varepsilon_L)) | (OPT_{\text{short}} \backslash OPT_T)_{ver}| \right) \\ &\geq &\frac{N^2}{2} \left(\frac{(1-w_L)(1-h_L)}{2a(OPT_{\text{short}} \backslash OPT_T)} - O(\varepsilon_L) \right) | OPT_{\text{short}} \backslash OPT_T| \end{split}$$

where the inequality follows from the fact that $\frac{a}{b} + \frac{c}{d} \ge \frac{(a+c)}{(b+d)}$ for $a,b,c,d \ge 0$. Since $a(OPT_{\rm short} \setminus OPT_T) \le (N^2 - a(OPT_{\rm long})) \le (1 - \frac{\alpha}{2} - \frac{\beta}{2})N^2 \le (1 - \frac{w_L}{2} - \frac{h_L}{2})N^2$ and $w_L + h_L \le \alpha + \beta \le \frac{4}{5}$, the amount of rectangles we are packing from $OPT_{\rm short} \setminus OPT_T$ is bounded below by the minimum of

$$f(h_L, w_L) = \left(\frac{(1 - w_L)(1 - h_L)}{(4 - 2w_L - 2h_L)} - O(\varepsilon_L)\right) N^2 |OPT_{\text{short}} \setminus OPT_T|$$

over the domain $\{w_L+h_L\leq \frac{4}{5},0\leq w_L\leq \frac{1}{2},0\leq h_L\leq \frac{1}{2}\}$. Since $\frac{\partial f(h_L,w_L)}{\partial h_L}=\frac{-2(1-w_L)^2}{(4-2w_L-2h_L)^2}\leq 0$ and $\frac{\partial f(h_L,w_L)}{\partial w_L}=\frac{-2(1-h_L)^2}{(4-2w_L-2h_L)^2}\leq 0$, the function is decreasing with respect to both its arguments, implying that the minimum value must be attained when $h_L+w_L=\frac{4}{5}$. This in turn implies that the amount of rectangles from $OPT_{\rm short}\setminus OPT_T$ we are packing is bounded below by the minimum of $f(h_L,\frac{4}{5}-h_L)$ over the interval $[\frac{3}{10},\frac{1}{2}]$. Since

$$f(h_L, \frac{4}{5} - h_L) = \left(\frac{5}{12}(1 - h_L)(\frac{1}{5} - h_L) - O(\varepsilon_L)\right) N^2 |OPT_{\text{short}} \setminus OPT_T|$$

describes a parabola centered at $h_L=\frac{2}{5}$, the minimum value on the aforementioned interval is attained at both limits $h_L=\frac{3}{10}$ and $h_L=\frac{1}{2}$ with a value of $\left(\frac{7}{48}-O(\varepsilon_L)\right)|OPT_{\rm short}\setminus OPT_T|$.

• If
$$min\{1, \left(\frac{(1-w_L)(1-h_L)}{2a((OPT_{\mathrm{short}} \backslash OPT_T)_{hor})} - O(\varepsilon_L)\right)N^2\} = 1$$
 and $\left(\frac{(1-w_L)(1-h_L)}{2a((OPT_{\mathrm{short}} \backslash OPT_T)_{ver})} - O(\varepsilon_L)\right)N^2 < 1$




Figure 10: Case 2A(iii)b in the proof of Theorem 36

(the remaining case being analogous), then we are packing at least

$$\frac{1}{2} \left(|(OPT_{\text{short}} \setminus OPT_{T})_{hor}| + \left(\frac{(1 - w_{L})(1 - h_{L})}{2a((OPT_{\text{short}} \setminus OPT_{T})_{ver})} - O(\varepsilon_{L}) \right) N^{2} |(OPT_{\text{short}} \setminus OPT_{T})_{ver}| \right) \\
\geq \frac{N^{2}}{2} \left(\frac{(1 - w_{L})(1 - h_{L})}{2a((OPT_{\text{short}} \setminus OPT_{T})_{ver})} - O(\varepsilon_{L}) \right) (|(OPT_{\text{short}} \setminus OPT_{T})_{hor}| + |(OPT_{\text{short}} \setminus OPT_{T})_{ver}|) \\
\geq \frac{N^{2}}{2} \left(\frac{(1 - w_{L})(1 - h_{L})}{2a(OPT_{\text{short}} \setminus OPT_{T})} - O(\varepsilon_{L}) \right) |OPT_{\text{short}} \setminus OPT_{T}| \\
\geq \left(\frac{7}{48} - O(\varepsilon_{L}) \right) |OPT_{\text{short}} \setminus OPT_{T}|,$$

where the last inequality comes from the analysis of the previous case.

From this we conclude that

$$|OPT_{L\&C}| \ge \frac{3(1 - O(\varepsilon_L))}{4}|OPT_{long}| + |OPT_{short} \cap OPT_T| + \left(\frac{7}{48} - O(\varepsilon_L)\right)|OPT_{short} \setminus OPT_T|.$$

This together with inequalities (2), (3), (4) and Lemma 40 gives us a solution with good enough approximation factor. Check Section B.1 and Table 1 for details.

$$\odot$$
 Subcase (iii b): $w_L > \frac{1}{2}$ (then from inequality (5), $h_L \leq \frac{3}{10}$). Note that $a(OPT_{long}) \leq (1 - \frac{3}{5})N^2 = \frac{2}{5}N^2$.

Let us define some parameters from the current packing to simplify the calculations. Let λN be the height of the rectangle in the packing that intersects or touches the vertical line $x=\left(\frac{1}{2}-2\varepsilon_{large}\right)N$ (if two rectangles touch such line, we choose that tallest one) and γN be the total width of vertical rectangles having height greater than $(1-h_L)N$. We define also the following three regions in the knapsack: \mathcal{A} , the rectangular region of width $w_L N$ and height $\frac{1}{2}N$ in the top left corner of the knapsack; \mathcal{B} , the rectangular region of width $w_L N$ and height $(\lambda - \frac{1}{2})N$ below \mathcal{A} and left-aligned with the knapsack; and \mathcal{C} , the rectangular region of width N and height N touching the bottom boundary of the knapsack. Notice that N is fully occupied by vertical rectangles, N is almost fully occupied by vertical rectangles except for the right region

of width $w_LN-\left(\frac{1}{2}-2\varepsilon_{large}\right)N$, and at least half of $\mathcal C$ is occupied by horizontal rectangles (some vertical rectangles may overlap with this region). Our goal is to pack some rectangles from $OPT_{\mathrm{short}}\setminus OPT_T$ in the \sqcup -shaped region outside $\mathcal A\cup\mathcal B\cup\mathcal C$. Let $\psi\in[\lambda,1-h_L]$ be a parameter to be fixed later. We will use, when possible, the following regions for packing items from $OPT_{\mathrm{short}}\setminus OPT_T$: Region 1 on the top right corner of the knapsack with width $N(1-w_L)$ and height ψN and Region 2 which is the rectangular region $[0,N]\times[h_LN,(1-\psi)\cdot N]$ (see Figure 10). Region 1 is completely free but Region 2 may overlap with vertical rectangles.

We will now divide Region 2 into a constant number of boxes such that: they do not overlap with vertical rectangles, the total area inside Region 2 which is neither overlapping with vertical rectangles nor covered by boxes is at most $O(\varepsilon_L)N^2$ and each box has width at least $(\frac{1}{2}+2\varepsilon_{large})N$ and height at least εN . That way we will be able to pack rectangles from $(OPT_{\rm short}\setminus OPT_T)_{ver}$ into the box defined by Region 1 and rectangles from $(OPT_{\rm short}\setminus OPT_T)_{hor}$ into the boxes defined inside Region 2 using almost completely its free space. In order to create the boxes inside Region 2 we first create a monotone chain by doing the following: Let $(x_1,y_1)=(\gamma N,h_L)$. Starting from position (x_1,y_1) , we draw an horizontal line of length $\varepsilon_L N$ and then a vertical line from bottom to top until it touches a vertical rectangle, reaching position (x_2,y_2) . From (x_2,y_2) we start again the same procedure and iterate until we reach the vertical line $x=(\frac{1}{2}-2\varepsilon_{large})N$ or the horizontal line $y=(1-\psi)N$. Notice that the area above the monotone chain and below $y=(1-\psi)N$ that is not occupied by vertical rectangles, is at most $\sum_i \varepsilon_L N(y_{i+1}-y_i) \le \varepsilon_L N^2$. The number of points (x_i,y_i) defined in the previous procedure is at most $1/\varepsilon_L$. By drawing an horizontal line starting from each (x_i,y_i) up to (N,y_i) , together with the drawn lines from the monotone chain and the right limit of the knapsack, we define $k \le 1/\varepsilon_L$ boxes. We discard the boxes having height less than εN , whose total area is at most $\frac{\varepsilon}{\varepsilon_L}N^2=\varepsilon_LN^2$, and have all the desired properties for the boxes. Note that the area that is occupied for sure by rectangles in $OPT_{\rm long}$ in regions \mathcal{A} , \mathcal{B} and \mathcal{C} by rectangles


Note that the area that is occupied for sure by rectangles in $OPT_{\rm long}$ in regions \mathcal{A},\mathcal{B} and \mathcal{C} by rectangles in $OPT_{\rm long}$ is at least $(\frac{1}{2}w_L + (\lambda - \frac{1}{2})(\frac{1}{2} - 2\varepsilon_{large}) + \frac{1}{2}h_L)N^2$. Since the total area of rectangles from $OPT_{\rm long}$ is at most $\frac{2}{5}N^2$, the total area occupied by rectangles in $OPT_{\rm long}$ in Region 2 is at most $N^2(\frac{2}{5} - \frac{1}{2}w_L - (\lambda - \frac{1}{2})(\frac{1}{2} - 2\varepsilon_{large}) - \frac{1}{2}h_L) \leq N^2(\frac{13}{20} - \frac{w_L}{2} - \frac{\lambda}{2} - \frac{h_L}{2} + \varepsilon_{large})$. This implies that the total area of the horizontal boxes is at least $N^2(1 - \psi - h_L) - N^2(\frac{13}{20} - \frac{w_L}{2} - \frac{\lambda}{2} - \frac{h_L}{2}) - O(\varepsilon_L)N^2$ and the area of the vertical box is $(1 - w_L)\psi N^2$. Ignoring the $O(\varepsilon_{large})$ -term, these two areas become equal if we set $\psi = \frac{7+10(w_L+\lambda-h_L)}{40-20w_L}$. It is not difficult to verify that in this case $\psi \leq 1 - h_L$. If $\frac{7+10(w_L+\lambda-h_L)}{40-20w_L} \geq \lambda$, then we set $\psi = \frac{7+10(w_L+\lambda-h_L)}{40-20w_L}$. Otherwise we set $\psi = \lambda$.


First, consider the case $\psi = \frac{7+10(w_L + \lambda - h_L)}{40-20w_L}$. Since $\psi \geq \lambda$, boxes inside Region 2 have width at least $(\frac{1}{2} + 2\varepsilon_{large}) N$ and height at least $\varepsilon N \gg \varepsilon_{small} N$ (recall that ε_{small} differs by a large factor from $\varepsilon_{large} \leq \varepsilon$), and the box in Region 1 has height at least $(\frac{1}{2} + 2\varepsilon_{large}) N$ and width at least $\frac{1}{5}N \gg \varepsilon_{small} N$. By using Steinberg's theorem, we can always pack in these boxes at least


$$\left(\min\left\{1,\frac{\frac{1}{2}(N-w_LN)\psi N}{a((OPT_{\mathrm{short}}\backslash OPT_T)_{hor})}-\varepsilon_{small}N^2\right\}\right)|(OPT_{\mathrm{short}}\backslash OPT_T)_{hor}| \\ + \left(\min\left\{1,\frac{\frac{1}{2}(N-w_LN)\psi N}{a((OPT_{\mathrm{short}}\backslash OPT_T)_{ver})}-\varepsilon_{small}N^2\right\}\right)|(OPT_{\mathrm{short}}\backslash OPT_T)_{ver}|.$$

Note that from each box B' of height $h(\geq \varepsilon N)$, we can remove the cheapest εh -horizontal strip and use resource augmentation to get a container based packing with nearly the same profit as B'. Thus by performing a similar analysis to the one done in Subcase (iii a), and using the fact that $a(OPT_{\rm short} \setminus OPT_T) \leq N^2 - (\frac{\alpha}{2} + (\lambda - \frac{1}{2})\frac{1}{2} + \frac{\beta}{2})N^2 \leq N^2 - N^2(\frac{w_L}{2} + (\lambda - \frac{1}{2})\frac{1}{2} + \frac{h_L}{2})$, we can minimize the whole expression over the domain $\{\frac{w_L}{2} + (\lambda - \frac{1}{2})\frac{1}{2} + \frac{h_L}{2} \leq \frac{2}{5}, \lambda \leq \psi, \frac{1}{2} \leq w_L \leq \frac{4}{5}, \frac{1}{2} \leq \lambda \leq 1, 0 \leq h_L \leq \frac{3}{10}\}$ and prove that this solution packs at least

$$\left(\frac{3 - O(\varepsilon_L)}{4}\right) |OPT_{\text{long}}| + |OPT_{\text{short}} \cap OPT_T| + \left(\frac{5}{36} - O(\varepsilon_L)\right) |OPT_{\text{short}} \setminus OPT_T|. \tag{8}$$







(b) Packing of rectangles in $OPT_{long} \cup (OPT_{short} \cap OPT_T)$. Dark gray rectangles are from $OPT_{short} \cap OPT_T$.

Figure 11: The case 2B.

Thus, using the above inequality along with (2), (3), (4) and Lemma 40, we get a solution with good enough approximation factor. Check Section B.1 and Table 1 for details.

Finally, if $\psi = \lambda > \frac{7+10(w_L + \lambda - h_L)}{40-20w_L}$, the bound for the area of horizontal boxes will not be equal to the area of the vertical box constructed to pack rectangles from isfat. In this case we change the width of the box inside Region 1 to be $w'_L < N(1-w_L)$ fixed in such a way that the area of this box is equal to the bound we have for the area of the boxes in Region 2, i.e., $N^2(1-\lambda-h_L)-(\frac{13}{20}-\frac{w_L}{2}-\frac{h_L}{2}-\frac{\lambda}{2}+O(\varepsilon_L))N^2$. Performing the same analysis as before, it can be shown that in this case we pack at least

$$\left(\frac{(1-\lambda-h_L)N^2-(\frac{13}{20}-\frac{w_L}{2}-\frac{h_L}{2}-\frac{\lambda}{2})N^2}{2a(OPT_{\text{short}}\setminus OPT_T)}-O(\varepsilon_L)N^2\right)|OPT_{\text{short}}\setminus OPT_T|,$$

which is at least $(\frac{1}{6} - O(\varepsilon_L))|OPT_{\mathrm{short}} \setminus OPT_T|$ over the domain $\{\frac{w_L}{2} + (\lambda - \frac{1}{2})\frac{1}{2} + \frac{h_L}{2} \leq \frac{2}{5}, \frac{1}{2} \leq w_L \leq \frac{4}{5}, \frac{7+10(w_L+\lambda-h_L)}{40-20w_L} < \lambda \leq 1, 0 \leq h_L \leq \frac{3}{10}\}$ (and this solution leads to a better bound than (8)).

$$\Diamond$$
 Case 2B. $\left(h_L \leq \varepsilon_L \text{ and } w_L N \geq w(V_{1-2\varepsilon_L}) > \left(\frac{1}{2} - 2\varepsilon_{large}\right) N\right)$ or $\left(w_L \leq \varepsilon_L \text{ and } h_L N \geq h(H_{1-2\varepsilon_L}) > \left(\frac{1}{2} - 2\varepsilon_{large}\right) N\right)$

In the first case, area of rectangles in $V_{1-2\varepsilon_L} > \left(\frac{1}{2} - 2\varepsilon_{large}\right)(1 - 2\varepsilon_L)N^2$. Remaining rectangles in OPT_{long} have area at least $(w_L - \frac{1}{2} + 2\varepsilon_{large})N \cdot \frac{N}{2}$. So, $a(OPT_{long}) > \left(\frac{1}{2} - 2\varepsilon_{large}\right)(1 - 2\varepsilon_L)N^2 + (w_L - \frac{1}{2})N \cdot \frac{N}{2} \ge \left(\frac{1}{4} + \frac{w_L}{2} - \varepsilon_L - 2\varepsilon_{large}\right)N^2$. Thus $a(OPT_{short} \setminus OPT_T) \le a(OPT_{short}) < \left(\frac{3}{4} - \frac{w_L}{2} + \varepsilon_L + 2\varepsilon_{large}\right)N^2$.

Now consider the vertical rectangles in the boundary L-region sorted non-increasingly by width and pick them iteratively until their total width crosses $(\frac{w_L}{2} + 3\varepsilon_L + 2\varepsilon_{large})N$. Remove these rectangles and push the remaining vertical rectangles in the L-region to the left as much as possible. This modified L-region will have profit at least $(\frac{1}{2} - O(\varepsilon_L))|OPT_{long}|$. Now we can put an εN -strip for the vertical items from $OPT_{short} \cap OPT_T$ next to the vertical part of L-region. On the other hand, the horizontal items of $OPT_{short} \cap OPT_T$ can be placed on top of the horizontal part of the L-region. The remaining space will be a free rectangular region of height at least $(1 - 2\varepsilon_L)N$ and width $(1 - \frac{w_L}{2} + 2\varepsilon_L + 2\varepsilon_L)$

 $2arepsilon_{large})N$. We will use a part of this rectangular region of height $(1-3arepsilon_L)N$ and width $(1-\frac{w_L}{2}+arepsilon_L)N$ to pack rectangles from $OPT_{\mathrm{short}}\setminus OPT_T$ and the rest of the region for resource augmentation. Since $\frac{w_L}{2}-arepsilon_L \leq \frac{1}{2}-arepsilon_{large}$, we can use Corollary 39 to pack short rectangles in this region with profit at least $\left(\frac{(1-\frac{w_L}{2})/2}{\frac{3}{4}-\frac{w_L}{2}}-O(arepsilon_L)\right)|OPT_{\mathrm{short}}\setminus OPT_T| \geq (\frac{3}{4}-O(arepsilon_L))|OPT_{\mathrm{short}}\setminus OPT_T|$ as the expression is increasing with respect to w_L and $w_L>\frac{1}{2}-2arepsilon_{large}$. Thus, we get,

$$|OPT_{L\&C}| \ge \left(\frac{1}{2} - O(\varepsilon_L)\right)|OPT_{long}| + |OPT_{short} \cap OPT_T| + \left(\frac{3}{4} - O(\varepsilon_L)\right)|OPT_{short} \setminus OPT_T|.$$
(9)

On the other hand, as $a(OPT_{\mathrm{short}} \setminus OPT_T) \leq (\frac{3}{4} - \frac{w_L}{2} + \varepsilon_L + 2\varepsilon_{large})N^2$ and $w_L > \frac{1}{2} - 2\varepsilon_{large}$, we get $a(OPT_{\mathrm{short}} \setminus OPT_T) \leq (\frac{1}{2} + 3\varepsilon_{large} + \varepsilon_L)N^2$ and thus from Lemma 40 we get

$$|OPT_{L\&C}| \geq \frac{3}{4}|OPT_{\text{long}} \cap OPT_{T}| + |OPT_{\text{short}} \cap OPT_{T}| + (1 - O(\varepsilon_{L}))|OPT_{\text{short}} \setminus OPT_{T}|$$

$$\geq \frac{3}{4}|OPT_{\text{long}} \cap OPT_{T}| + (1 - O(\varepsilon_{L}))|OPT_{\text{short}}|. \tag{10}$$

From inequalities (1), (3), (4), (9) and (10) we get a solution with good enough approximation factor. Check Section B.1 and Table 1 for details.

Now we consider the last case when $w_L \leq \varepsilon_L$ and $h_L N \geq h(H_{1-2\varepsilon_L}) > \left(\frac{1}{2} - 2\varepsilon_{large}\right) N$. Note that as we assumed the cheapest subring was the top subring, after removing it we might be left with only $|OPT_{long} \cap I_{hor}|/2$ profit in the horizontal part of L-region. So, further removal of items from the horizontal part might not give us a good solution. Thus we show an alternate good packing. We restart with the ring packing and delete the cheapest vertical subring instead of the cheapest subring (i.e., the top subring) and create a new boundary L-region. Here, consider the horizontal rectangles in the boundary L-region in non-increasing order of height and take them until their total height crosses $(\frac{\beta_{bottom} + \beta_{top}}{2} + \varepsilon_{small} + \varepsilon)N$. Remove these rectangles and push the remaining horizontal rectangles to the bottom as much as possible. Then, following similar arguments as before, we will obtain the same bounds for the constructed solution.

B.1 Bounding the approximation factor

In each one of the cases listed before we are developing a set of different solutions in order to achieve a good approximation factor. Let $z = |OPT_{L\&C}|/|OPT|$, $x_1 = |OPT_{long} \cap OPT_T|/|OPT|$, $x_2 = |OPT_{long} \setminus OPT_T|/|OPT|$, $x_3 = |OPT_{short} \cap OPT_T|/|OPT|$ and $x_4 = |OPT_{short} \setminus OPT_T|/|OPT|$. The following list enumerates all the obtained inequalities in this section, and it is worth remarking that not all of them hold simultaneously.

1.
$$z \ge \frac{3}{4}x_1 + \frac{3}{4}x_2 + x_3 + \left(\frac{1}{2} - O(\varepsilon_L)\right)x_4;$$

2.
$$z \ge \frac{3}{4}x_1 + \frac{3}{4}x_2$$
;

3.
$$z \ge \left(\frac{1}{2} - O(\varepsilon_L)\right)(x_1 + x_2) + \left(\frac{3}{4} - O(\varepsilon_L)\right)(x_3 + x_4);$$

4.
$$z \ge (1 - O(\varepsilon_L))(x_2 + x_4);$$

5.
$$z \ge (1 - O(\varepsilon_L)) \left(\frac{1}{2}x_1 + x_2 + \frac{1}{2}x_4\right);$$

6.
$$z \ge \left(\frac{3}{4} - O(\varepsilon_L)\right)(x_1 + x_2) + x_3;$$

7.
$$z \ge \frac{3}{4}x_1 + x_3 + \left(\frac{5}{6} - O(\varepsilon_L)\right)x_4;$$

8.
$$z \ge \frac{3}{4}(x_1 + x_2) + \left(\frac{5}{12} - O(\varepsilon_L)\right)(x_3 + x_4);$$

9.
$$z \ge \frac{3}{4}x_1 + x_3 + \left(\frac{2}{3} - O(\varepsilon_L)\right)x_4;$$

10.
$$z \ge \frac{3}{4}(x_1 + x_2) + x_3 + (\frac{7}{48} - O(\varepsilon_L))x_4;$$

11.
$$z \ge \left(\frac{3}{4} - O(\varepsilon_L)\right) (x_1 + x_2) + x_3 + \left(\frac{5}{36} - O(\varepsilon_L)\right) x_4;$$

12.
$$z \ge \left(\frac{1}{2} - O(\varepsilon_L)\right)(x_1 + x_2) + x_3 + \left(\frac{3}{4} - O(\varepsilon_L)\right)x_4;$$

13.
$$z \ge \frac{3}{4}x_1 + (1 - O(\varepsilon_L))(x_3 + x_4)$$
.

For each case i, let A_i be the set of indexes of valid inequalities for case i. Then we can write the following linear program to compute the obtained approximation factor in that case:

$$\begin{array}{ll} \min & z \\ s.t. & \text{Inequalities indexed by } \mathcal{A}_i \\ & \sum_{i=1}^4 x_i = 1 \\ & z, x_i \geq 0 & \text{for } i = 1, 2, 3, 4. \end{array}$$

Let $c_{j,k}$ be the coefficient accompanying x_k in the constraint $j \in \mathcal{A}_i$, k = 1, 2, 3, 4. The dual of the program for case i has the form

$$\max_{s.t.} \quad \sum_{j \in \mathcal{A}_i}^{-w} y_j \le 1$$

$$\sum_{j \in \mathcal{A}_i}^{-w} c_{j,k} y_j + w \ge 0 \quad \text{for } k = 1, 2, 3, 4$$

$$y_j \ge 0 \quad \text{for } j \in \mathcal{A}_i$$

$$w \in \mathbb{R}$$

Any feasible solution for the dual program of case i is a lower bound on the fraction of OPT packed in that case. Table 1 summarizes the analysis described along this section for all the cases, stating the valid inequalities and the approximation factor obtained in each one of them, together with a dual feasible solution. It is not difficult to see that the worst case is 2A(iii)b, implying that $|OPT_{L\&C}| \geq (\frac{325}{558} - O(\varepsilon_L))|OPT|$. Applying Lemma 35 concludes the proof of Theorem 36.

Case	Valid inequalities	Dual feasible solution	Fraction of OPT
	1		packed (w)
1	1, 3, 4, 5	$y_1 = \frac{1}{2}, y_3 = \frac{1}{2}, y_4 = 0, y_5 = 0$	$\frac{5}{8} - O(arepsilon_L)$
2A(i)	3, 4, 5, 6, 7	$y_3 = \frac{17}{54}, y_4 = 0, y_5 = \frac{1}{3}, y_6 = \frac{7}{54}, y_7 = \frac{2}{9}$	$\frac{127}{216} - O(arepsilon_L)$
2A(ii)	3, 4, 5, 6, 8	$y_3 = \frac{4}{7}, y_4 = 0, y_5 = 0, y_6 = 0, y_8 = \frac{3}{7}$	$\frac{17}{28} - O(\varepsilon_L)$
2A(iii)a	3, 4, 5, 9, 10	$y_3 = \frac{124}{369}, y_4 = 0, y_5 = \frac{1}{3}, y_9 = \frac{2}{9}, y_{10} = \frac{40}{369}$	$\frac{215}{369} - O(\varepsilon_L)$
2A(iii)b	3, 4, 5, 9, 11	$y_3 = \frac{94}{279}, y_4 = 0, y_5 = \frac{1}{3}, y_9 = \frac{2}{9}, y_{11} = \frac{10}{93}$	$\frac{325}{558} - O(\varepsilon_L)$
2B	2, 4, 5, 12, 13	$y_2 = \frac{8}{41}, y_4 = 0, y_5 = \frac{9}{41}, y_{12} = \frac{18}{41}, y_{13} = \frac{6}{41}$	$\frac{24}{41} - O(\varepsilon_L)$

Table 1: Summary of the case analysis in Theorem 36.

C Cardinality Case With Rotations

In this section we present a polynomial time $(4/3 + \varepsilon)$ -approximation algorithm for 2DKR for the cardinality case. We next assume w.l.o.g. that $\varepsilon > 0$ is a sufficiently small constant and also that $h(i) \ge w(i)$ for all items i in the input.

Consider some optimal solution OPT for 2DKR, with an associated packing in the knapsack. We crucially exploit the following resource contraction lemma, which is our main new idea in the rotation case.

Lemma 42. (Resource Contraction Lemma) For given positive constants $\varepsilon \leq 1/13$ and $\varepsilon_{small} < \varepsilon^{\frac{1}{2\varepsilon}+1}$, suppose that there exists a feasible packing of a set of items M, with $|M| \geq 1/\varepsilon_{small}^3$. Then it is possible to pack a subset $M' \subseteq M$ of cardinality at least $\frac{2}{3}(1 - O(\varepsilon))|M|$ into $[0, (1 - \varepsilon^{\frac{1}{2\varepsilon}+1})N] \times [0, N]$ if rotations are allowed.

We defer the proof of the lemma to the end of this section.

As in the case without rotations, we will first show the existence of a container packing that packs at least $(3/4 - O(\varepsilon))|OPT|$ many items. Let APX be the container packing with largest possible profit. As in Section A, we assume all items to be skewed. Note that the small items can be handled with the techniques used in Lemma 28. We start with the corridor partition as in Section A and define *thin*, *fat* and *killed* rectangles accordingly. Let T and F be the set of thin and fat rectangles respectively.

We will show that $|APX| \ge (3/4 - O(\varepsilon))|OPT|$.

Lemma 43.
$$|APX| \ge (1 - \varepsilon)|F|$$
.

Proof. After removal of T, we can get a container packing for almost all items in F as discussed in Lemma 23 in Section A.

Now we show that using Lemma 42 we can prove the following:

Lemma 44.
$$|APX| \ge (1 - O(\varepsilon))(|T| + \frac{2}{3}|F|).$$

Proof. Thanks to 18 we can ensure that the total height of rectangles in T is at most $\frac{\varepsilon^{\frac{1}{2\varepsilon}+1}N}{2}$. So we can pack them in a vertical container of width $\frac{\varepsilon^{\frac{1}{2\varepsilon}+1}N}{2}$.

Now if $|F| \geq \frac{1}{\varepsilon_{small}^3}$, then by Lemma 42 there exists $F' \subseteq F$ of cardinality at least $\frac{2}{3}(1-O(\varepsilon))|F|$ that can be packed inside $K' := [0, \left(1-\varepsilon^{\frac{1}{2\varepsilon}+1}\right)N] \times [0,N]$. Then we can use the resource augmentation PTAS in [21] to get a container packing of $(2/3-O(\varepsilon))|F|$ many rectangles in the area $K'' := [0, \left(1-\varepsilon^{\frac{1}{2\varepsilon}+1}/2\right)N] \times [0,N]$, and pack the vertical container for items in T to the right of K'' in the area $[\left(1-\varepsilon^{\frac{1}{2\varepsilon}+1}/2\right)N, 1] \times [0,N]$.

Otherwise, if $|F| < \frac{1}{\varepsilon_{small}^3}$, there are two cases. If $|T| < \frac{1}{\varepsilon_{small}^4}$, then $|F \cup T| < \frac{2}{\varepsilon_{small}^4}$ and we can find the packing just by brute-force. Otherwise if, $|T| \ge \frac{1}{\varepsilon_{small}^4} \ge |F|/\varepsilon_{small}$, then $APX \ge |T| \ge (1 - O(\varepsilon))(|T| + |F|)$.

Thus we get the following theorem:

Theorem 45. $|APX| \ge (3/4 - O(\varepsilon))|OPT|$.

Proof. The claim follows by combining Lemma 44 and 43. Up to a factor $(1 - O(\varepsilon))$, the worst case is obtained when $|F| = |T| + 2/3 \cdot |F|$, i.e., |F| = 3|T|. This gives a total profit of $3/4 \cdot |T \cup F|$.

It remains to prove Lemma 42. Let us remove from M all items that are larger than $\varepsilon_{small}N$ in both dimensions. Let M_2 be the resulting set: observe that $|M_2| \ge (1 - \varepsilon_{small})|M|$.

We next show how to remove from M_2 a set of cardinality at most $\varepsilon |M_2|$ such that the remaining items M_3 are either *very tall* or *not too tall*, where the exact meaning of this will be given next. We use the notation $[k] = \{1, \ldots, k\}$ for a positive integer k.

Lemma 46. Given any constant $1/2 > \varepsilon > 0$, there exists a value $i \in [\lceil 1/(2\varepsilon) \rceil]$ such that all items in M_2 having height in $((1-2\varepsilon^i)N, (1-\varepsilon^{i+1})N]$ have total cardinality at most $\varepsilon |M_2|$.


Proof. Let K_i be the set of items in M_2 with height in $((1-2\varepsilon^i)N, (1-\varepsilon^{i+1})N]$ for $i \in [\lceil 1/(2\varepsilon) \rceil]$. An item can belong to at most two such sets as $\varepsilon < 1/2$. Thus, the smallest such set has cardinality at most $\varepsilon |M_2|$.


We remove from M_2 the elements from the set K_i of minimum cardinality guaranteed by the above lemma, and let M_3 be the resulting set. We also define $\varepsilon_s = \varepsilon^i$ for the same i. Thus, $\varepsilon_s \geq \varepsilon^{1/2\varepsilon} > \varepsilon_{small}/\varepsilon$. Note that the items in M_3 have height either at most $(1 - 2\varepsilon_s)N$ or above $(1 - \varepsilon \cdot \varepsilon_s)N$.

For any $\delta>0$ denote the strips of width N and height δN at the top and bottom of the knapsack by $S_{T,\delta}:=[0,N]\times[(1-\delta)N,N]$ and $S_{B,\delta}:=[0,N]\times[0,\delta N]$, resp. Similarly, denote the strips of height N and width δN to the left and right of the knapsack by $S_{L,\delta}:=[0,\delta N]\times[0,N]$ and $S_{R,\delta}:=[(1-\delta)N,N]\times[0,N]$, resp. The set of items in M_3 intersected by and fully contained in strip $S_{K,\delta}$ are denoted by $E_{K,\delta}$ and $C_{K,\delta}$, resp. Obviously $C_{K,\delta}\subseteq E_{K,\delta}$, and we define, $D_{K,\delta}=E_{K,\delta}\setminus C_{K,\delta}$. Let a(I) denote the total area of items in I, i.e., $a(I)=\sum_{i\in I}w(i)\cdot h(i)$.

Lemma 47. Either
$$a(E_{L,\varepsilon_s} \cup E_{R,\varepsilon_s}) \leq \frac{(1+8\varepsilon_s)}{2} N^2$$
 or $a(E_{T,\varepsilon_s} \cup E_{B,\varepsilon_s}) \leq \frac{(1+8\varepsilon_s)}{2} N^2$.

Proof. Let us define $V:=E_{L,\varepsilon_s}\cup E_{R,\varepsilon_s}$ and $H:=E_{T,\varepsilon_s}\cup E_{B,\varepsilon_s}$. Note that, $a(V)+a(H)=a(V\cup H)+a(V\cap H)$. Clearly $a(V\cup H)\leq N^2$ since all items fit into the knapsack. On the other hand, except possibly four items (the ones that contain at least one of the points $(\varepsilon_sN,\varepsilon_sN),((1-\varepsilon_s)N,\varepsilon_sN),(\varepsilon_sN,(1-\varepsilon_s)N),((1-\varepsilon_s)N,(1-\varepsilon_s)N)$) all other items in $V\cap H$ lie entirely within the four strips $S_{L,\varepsilon_s}\cup S_{R,\varepsilon_s}\cup S_{T,\varepsilon_s}\cup S_{T,\varepsilon_s}\cup$






(a) Strips $S_{L,\alpha}, S_{R,\beta}, S_{B,\delta}, S_{T,\gamma}$

(b) $C_{L,\alpha}, D_{L,\alpha}$ are dark and light gray resp.

Figure 12: Definitions for cardinality 2DK with rotations.






Figure 13: Case A for cardinality 2DK with rotations. Dark gray rectangles are X, light gray rectangles are Z, gray (and hatched) rectangles are Y, hatched rectangles are $C_{T,\varepsilon^{i+1}}$ and $C_{B,\varepsilon^{i+1}}$. Figure (a): original packing in $N \times N$, Figure (b): modified packing leaving space for resource contraction on the right.


Now we use Steinberg's Theorem [32] to prove the following lemma.

Lemma 48. Let $0 < \varepsilon_a < 1/2$ be a constant and $\tilde{M} := \{1, \ldots, k\}$ a set of items satisfying $w(i) \le \varepsilon_{small} N$ for all $i \in \tilde{M}$. If $a(\tilde{M}) \le (1/2 + \varepsilon_a)N^2$. Then, a subset of \tilde{M} of cardinality at least $(1 - 2\varepsilon_s - 2\varepsilon_a)|\tilde{M}|$ can be packed into $[0, (1 - \varepsilon_s)N] \times [0, N]$.

Proof. W.l.o.g., assume the items in \tilde{M} are given in nondecreasing order according to their area. Note that $a(i) \leq \varepsilon_{small} N^2 \leq \frac{\varepsilon_s}{2} N^2$ for any $i \in \tilde{M}$. Let $S := \{1,\ldots,j\}$ be such that $\frac{(1-2\varepsilon_s)}{2} N^2 \leq \sum_{i=1}^j a(i) \leq \frac{(1-\varepsilon_s)}{2} N^2$ and $\sum_{i=1}^{j+1} a(i) > \frac{(1-\varepsilon_s)}{2} N^2$. Then from Theorem 38, S can be packed into $[0,(1-\varepsilon_s)N] \times [0,N]$.

As we considered items in the order of nondecreasing area, $\frac{|S|}{|\tilde{M}|} \geq \frac{\left(\frac{1}{2} - \varepsilon_s\right)}{\left(\frac{1}{2} + \varepsilon_a\right)}$. Thus, $|S| \geq \left(1 - \frac{(\varepsilon_a + \varepsilon_s)}{\left(\frac{1}{2} + \varepsilon_a\right)}\right) |\tilde{M}| > (1 - 2\varepsilon_a - 2\varepsilon_s)|\tilde{M}|$.

From Lemma 47, we can assume w.l.o.g. that $a(E_{T,\varepsilon_s} \cup E_{B,\varepsilon_s}) \leq \frac{(1+8\varepsilon_s)}{2}N^2$. Let X be the set of items in M_3 that intersect both S_{T,ε_s} and S_{B,ε_s} and $Y:=\{E_{T,\varepsilon_s} \cup E_{B,\varepsilon_s}\} \setminus X$. Define $Z:=M_3 \setminus \{X \cup Y\}$ to






Figure 14: Case B for cardinality 2DK with rotations. Dark gray rectangles are X, light gray rectangles are Z, gray (and hatched) rectangles are Y, hatched rectangles are $C_{T,\varepsilon^{i+1}}$ and $C_{T,\varepsilon^{i+1}}$. Figure (a): original packing, Figure (b): modified packing leaving space for resource contraction on the top.

be the rest of the items. Let us define $w(X) = \sum_{i \in X} w(i)$. Now there are two cases.

Case A. $w(X) \geq 12\varepsilon \cdot \varepsilon_s N$. From Lemma 46, all items in X intersect both $S_{T,\varepsilon\cdot\varepsilon_s}$ and $S_{B,\varepsilon\cdot\varepsilon_s}$. So the removal of $X \cup C_{T,\varepsilon\cdot\varepsilon_s} \cup C_{B,\varepsilon\cdot\varepsilon_s}$ creates a set of empty strips of height N and total width of w(X). By a simple permutation argument, all items in $Y \cup Z$ can be packed inside $[0,N-w(X)] \times [0,N]$, leaving an empty vertical strip of width w(X) on the right side of the knapsack. Next we rotate $C_{T,\varepsilon\cdot\varepsilon_s}$ and $C_{B,\varepsilon\cdot\varepsilon_s}$ and pack them in two vertical strips, each of width $\varepsilon \cdot \varepsilon_s N$. Note that $w(i) \leq \varepsilon \cdot \varepsilon_s N$ for all $i \in X$. Now take items in X by nondecreasing width, till their total width is in $[w(X) - 4\varepsilon \cdot \varepsilon_s N, w(X) - 3\varepsilon \cdot \varepsilon_s N]$ and pack them into another vertical strip. The cardinality of this set is at least $\frac{(w(X) - 4\varepsilon \cdot \varepsilon_s N)}{w(X)} |X| \geq \frac{2}{3} |X|$, where the last inequality follow by the Case A assumption. Hence, at least $\frac{2}{3} |X| + |Y| + |Z| \geq \frac{2}{3} (|X| + |Y| + |Z|)$ items can be packed into $[0, (1 - \varepsilon \cdot \varepsilon_s)N] \times [0, N]$.

Case B. $w(X) < 12\varepsilon \cdot \varepsilon_s N$. Observe that $Y = (E_{T,\varepsilon_s} \setminus X) \dot{\cup} (E_{B,\varepsilon_s} \setminus X)$, hence $|Y| = |E_{T,\varepsilon_s} \setminus X| + |E_{B,\varepsilon_s} \setminus X|$. Assume w.l.o.g. that $|E_{B,\varepsilon_s} \setminus X| \ge |Y|/2 \ge |E_{T,\varepsilon_s} \setminus X|$. Then remove E_{T,ε_s} . We can pack X on top of $M \setminus E_{T,\varepsilon_s}$ as $12\varepsilon \cdot \varepsilon_s \le \varepsilon_s - \varepsilon \cdot \varepsilon_s$ for $\varepsilon \le 1/13$. This gives a packing of $|X| + |Z| + \frac{|Y|}{2}$ many rectangles. On the other hand, as $a(X \cup Y) = a(E_{T,\varepsilon_s} \cup E_{B,\varepsilon_s}) \le \frac{(1+8\varepsilon_s)}{2}N^2$, from Lemma 48, it is possible to pack at least $(1-2\varepsilon_s-8\varepsilon_s)|X \cup Y| \ge (1-10\varepsilon_s)(|X|+|Y|)$ many items into $[0,(1-\varepsilon \cdot \varepsilon_s)N] \times [0,N]$.

Thus we can always pack a set of items of cardinality at least

$$\max\{(1 - 10\varepsilon_s)(|X| + |Y|), |X| + |Z| + \frac{|Y|}{2}\}$$

$$\geq \frac{1}{3}(1 - 10\varepsilon_s)(|X| + |Y|) + \frac{2}{3}(|X| + |Z| + \frac{|Y|}{2})$$

$$\geq \frac{2}{3}(1 - 10\varepsilon_s)(|X| + |Y| + |Z|)$$

$$= \frac{2}{3}(1 - 10\varepsilon_s)|M_3|.$$

This concludes the proof of Lemma 42.

D Weighted Case with Rotations

In this section we give a polynomial time $(3/2+\varepsilon)$ -approximation algorithm for the weighted 2-dimensional geometric knapsack problem when items are allowed to be rotated by 90 degrees. In contrary to the unweighted case, where it is possible to remove a constant number of *large* items, the same is not possible in the weighted case, where an item could have a big profit.

We call an item i massive if $w(i) \ge (1 - \varepsilon)N$ and $h(i) \ge (1 - \varepsilon)N$. The presence of such a big item in

the optimal solution requires a different analysis, that we present below. In both the cases, we can show that there exists a container packing with roughly 2/3 of the profit of the optimal solution.

Let us assume that $\varepsilon < 1/6$. We will prove the following result:

Theorem 49. Let $\varepsilon > 0$ and let R be a set of items that can be packed into the $N \times N$ knapsack. Then there exists a container packing with $O_{\varepsilon}(1)$ containers of a subset $R' \subseteq R$ into the $N \times N$ knapsack such that $p(R') \ge (2/3 - O(\varepsilon))p(R)$, if rotations are allowed.

We start by analyzing the case of a set R that has a massive item.

Lemma 50. Suppose that a set R of items can be packed into a $N \times N$ bin and there is a massive item $m \in R$. Then, there is a container packing with at most $O_{\varepsilon}(1)$ containers for a subset $R' \subseteq R$ such that $p(R') \ge \left(\frac{2}{3} - O(\varepsilon)\right) p(R)$.


Proof. Assume, without loss of generality, that $1/(3\varepsilon)$ is an integer. Consider the items in $R\setminus\{m\}$. Clearly, each of them has width or height at most ε ; moreover, $a(R\setminus\{m\}) \leq (1-(1-\varepsilon)^2)N^2 = (2\varepsilon-\varepsilon^2)N^2 \leq \frac{N^2}{2(1+\varepsilon)}$, as $\varepsilon < 1/6$; thus, by possibly rotating each element so that the height is smaller than ε , by Theorem 38 all the items in $R\setminus\{m\}$ can be packed in a $N\times\frac{N}{1+\varepsilon}$ bin; then, by Lemma 55, there is a container packing for a subset of $R\setminus\{m\}$ with $O_\varepsilon(1)$ containers that fits in the $N\times N$ bin and has profit at least $(1-O(\varepsilon))p(R\setminus\{m\})$

Consider now the packing of R. Clearly, the region $[\varepsilon N, (1-\varepsilon)N]^2$ is entirely contained within the boundaries of the massive item m. Partition the region with x-coordinate between εN and $(1-\varepsilon)N$ in $k=1/(3\varepsilon)$ strips of width $3\varepsilon(1-2\varepsilon)N\geq 2\varepsilon N$ and height N, let them be S_1,\ldots,S_k ; let $R(S_i)$ be the set of items in R such that their left or right edge (or both) are contained in the interior of strip S_i . Since each item belongs to at most two of these sets, there exists i such that $p(R(S_i))\leq 6\varepsilon p(R)$.


Symmetrically, we define k horizontal strips T_1,\ldots,T_k , obtaining an index j such that $p(R(T_j)) \leq 6\varepsilon p(R)$. Thus, no item in $\overline{R}:=R\setminus (R(S_i)\cup R(T_j))$ has a side contained in the interior of S_i or T_j , and $p(\overline{R})\geq (1-12\varepsilon)p(R)$. Let M_V be the set of items in $\overline{R}\setminus \{m\}$ that overlap T_j , and let M_H be the set of items in $\overline{R}\setminus \{m\}$ that overlap S_i . Clearly, the items in M_H can be packed in a horizontal container with width N and height N-h(m), and the items in M_V can be packed in a vertical container of width N-w(m) and height N.

Let H be the set of items of $\overline{R} \setminus M_H$ that are completely above the massive item m or completely below it; symmetrically, let V be the set of items of $\overline{R} \setminus M_V$ that are completely to the left or completely to the right of m. We will now show that there is a container packing for $M_H \cup V \cup \{m\}$. Since all the elements overlapping T_j have been removed, V can be packed in a bin of size $(N-w(m)) \times (1-2\varepsilon)N$ (see Figure 15). Since $(1-2\varepsilon)N \cdot (1+\varepsilon) < (1-\varepsilon)N \le h(m)$, Lemma 55 implies that there is a container packing of a subset of V with profit at least $(1-O(\varepsilon))p(V)$ in a bin of size $(N-w(m)) \times h(m)$ and using $O_\varepsilon(1)$ containers; thus, by adding a horizontal container of the same size as m and a horizontal container of size $N \times (N-h(m))$, we obtain a container packing for $M_H \cup V \cup \{m\}$ with $O_\varepsilon(1)$ containers and profit at least $(1-O(\varepsilon))p(M_H \cup V \cup \{m\})$. Symmetrically, there is a container packing for a subset of $M_V \cup H \cup \{m\}$ with profit at least $(1-O(\varepsilon))p(M_V \cup H \cup \{m\})$ and $O_\varepsilon(1)$ containers.

Let R_{MAX} be the set of maximum profit among the sets $R \setminus \{m\}$, $M_H \cup V \cup \{m\}$ and $M_V \cup H \cup \{m\}$. By the discussion above, there is a container packing for $R' \subseteq R_{MAX}$ with $O_{\varepsilon}(1)$ containers and profit at least $(1 - O(\varepsilon))p(R_{MAX})$. Since each element in \overline{R} is contained in at least two of the above three sets, it follows that:



(a) Massive item case. Items intersecting strips M_H and M_V (hatched rectangles) cross them completely.



(b) Bottom(i), Top(i), Left(i), Right(i) are represented by vertical, horizontal, north east and north west stripes respectively.

Figure 15

$$p(R') \ge (1 - O(\varepsilon))p(R_{MAX}) \ge (1 - O(\varepsilon))\left(\frac{2}{3}p(\overline{R})\right)$$

 $\ge \left(\frac{2}{3} - O(\varepsilon)\right)p(R)$

If there is no massive item, we will show existence of two *container packings* and show the maximum of them always packs items with total profit at least $(\frac{2}{3} - O(\varepsilon))$ fraction of the optimal profit.

First, we follow the corridor decomposition and the classification of items as in Section A to define sets $LF, SF, LT, ST, OPT_{small}$. Let $T := LT \cup ST$ be the set of thin items. Also let APX be the best container packing and OPT be the optimal solution. Then similar to Lemma 33, we can show $p(APX) \ge (1 - \varepsilon)(p(LF) + p(SF) + p(OPT_{small}))$. Thus,

$$p(APX) \ge (1 - \varepsilon)p(OPT) - p(T).$$
 (11)

In the second case, we define the set T as above. Then in Resource Contraction Lemma (Lemma 51), we will show that one can pack 1/2 of the remaining profit in the optimal solution, i.e., $p(OPT \setminus T)/2$ in a knapsack of size $N \times (1 - \varepsilon/2)N$. Now, we can pack T in a horizontal container of height $\varepsilon/4$ and using Lemma 51 and resource augmentation we can pack $p(OPT \setminus T)/2$ in the remaining space $N \times (1 - \varepsilon/4)N$. Thus,

$$p(APX) \ge p(T) + (1 - \varepsilon)(p(OPT) - p(T))/2. \tag{12}$$

Hence, up to $(1 - O(\varepsilon))$ factor, we pack at least $\max\{(p(T) + p(OPT \setminus T)/2), p(OPT \setminus T)\} \ge 2/3 \cdot p(OPT)$, thus proving Theorem 49.

Note that using techniques similar to Appendix A, we can get a PTAS for the best container packing. Now to complete the proof of Theorem 4, it only remains to prove Lemma 51.




Figure 16: Cases for Resource Contraction Lemma (Lemma 51).

Lemma 51. (Resource Contraction Lemma) If a set of items M contains no massive item and can be packed into a $N \times N$ bin, then it is possible to pack a set M' of profit at least $p(M) \cdot \frac{1}{2}$ into a $N \times (1 - \frac{\varepsilon}{2})N$ bin (or a $(1 - \frac{\varepsilon}{2})N \times N$ bin), if rotations are allowed.

Proof. Let $\varepsilon_s = \varepsilon/2$. We will partition M into two sets $M_1, M \setminus M_1$ and show that both these sets can be packed into $N \times (1 - \varepsilon_s)N$ bin. If an item i is embedded in position (x_i, y_i) , we define $x_i' := x_i + w(i), y_i' := y_i + h(i)$.

In a packing of a set of items M, for item i we define $Left(i) := \{k \in M : x_k' \leq x_i\}$, $Right(i) := \{k \in M : x_k \geq x_i'\}$, $Top(i) := \{k \in M : y_k \geq y_i'\}$, $Bottom(i) := \{k \in M : y_k' \leq y_i\}$, i.e., the set of items that lie completely on left, right, top and bottom of i respectively. Now consider four strips $S_{T,3\varepsilon_s}, S_{B,\varepsilon_s}, S_{L,\varepsilon_s}, S_{R,\varepsilon_s}$ (see Figure 16).

Case 1. $D_{B,\varepsilon_s}\cap D_{T,3\varepsilon_s}=\emptyset$, i.e., no item intersecting S_{B,ε_s} intersects $S_{T,3\varepsilon_s}$. Define $M_1:=E_{T,3\varepsilon_s}$. As these items in M_1 do not intersect S_{B,ε_s} , M_1 can be packed into a $(N,N(1-\varepsilon_s))$ bin. For the remaining items, pack $M\setminus (M_1\cup C_{L,\varepsilon_s}\cup C_{R,\varepsilon_s})$ as it is. Now rotate C_{L,ε_s} and C_{R,ε_s} and pack on top of $M\setminus (M_1\cup C_{L,\varepsilon_s}\cup C_{R,\varepsilon_s})$ into two strips of height $\varepsilon_s N$ and width N. This packing will have total height $\le (1-3\varepsilon_s+2\varepsilon_s)N\le (1-\varepsilon_s)N$.

Case 2. $D_{B,\varepsilon_s} \cap D_{T,3\varepsilon_s} \neq \emptyset$, i.e., there is some item intersecting S_{B,ε_s} that also crosses $S_{T,3\varepsilon_s}$. Now, there

are three subcases:

Case 2A. There exists an item i that does neither intersect S_{L,ε_s} nor S_{R,ε_s} . Then item i partitions the items in $M\setminus (C_{T,3\varepsilon_s}\cup C_{B,\varepsilon_s}\cup \{i\})$ into two sets: Left(i) and Right(i). W.l.o.g., assume $x_i\leq 1/2$. Then remove $Right(i), i, C_{T,3\varepsilon_s}$ and C_{B,ε_s} from the packing. Now rotate $C_{T,3\varepsilon_s}$ and C_{B,ε_s} to pack right of Left(i). Define this set $M\setminus (Right(i)\cup \{i\})$ to be M_1 . Clearly packing of M_1 takes height N and width $x_i+4\varepsilon_sN\leq (\frac{1}{2}+4\varepsilon_s)N\leq (1-\varepsilon_s)N$ as $\varepsilon_s\leq \frac{1}{10}$. As the item i does not intersect the strip S_{L,ε_s} , $(Right(i)\cup \{i\})$ can be packed into height N and width $(1-\varepsilon_s)N$.

Case 2B. There exists an item i that intersects both S_{L,ε_s} and S_{R,ε_s} . Consider M_1 to be $M\setminus (C_{L,\varepsilon_s}\cup C_{R,\varepsilon_s}\cup Top(i))$. As there is no massive item, M_1 is packed in height $(1-\varepsilon_s)N$ and width N. Now, pack Top(i) and then rotate C_{L,ε_s} and C_{R,ε_s} to pack on top of it. These items can be packed into height $(1-y_i'+2\varepsilon_s)N\leq 5\varepsilon_sN\leq (1-\varepsilon_s)N$ as $\varepsilon_s\leq 1/10$.

Case 2C. If an item i intersects both S_{B,ε_s} and $S_{T,3\varepsilon_s}$, then the item i intersects exactly one of S_{L,ε_s} and S_{R,ε_s} . Consider the set of items in $D_{B,\varepsilon_s} \cap D_{T,3\varepsilon_s}$.

First, consider the case when the set $D_{B,\varepsilon_s}\cap D_{T,3\varepsilon_s}$ contains an item $i\in D_{L,\varepsilon_s}$ (similarly one can consider $i\in D_{R,\varepsilon_s}$). Now if $x_i'\leq N/2$, take $M_1:=Right(i)$. Then, we can rotate Right(i) and pack into height $(1-\varepsilon_s)N$ and width N. On the other hand, pack $M\setminus\{M_1\cup C_{T,3\varepsilon_s}\cup C_{B,\varepsilon_s}\}$ as it is. Then rotate $C_{T,3\varepsilon_s}\cup C_{B,\varepsilon_s}$ and pack on its side. Total width $\leq (1/2+3\varepsilon_s+\varepsilon_s)N\leq (1-\varepsilon_s)N$ as $\varepsilon_s\leq 1/6$. Otherwise if $x_i'>N/2$ take $M_1:=Left(i)\cup i$. Now, consider packing of $M\setminus\{M_1\cup C_{T,3\varepsilon_s}\cup C_{B,\varepsilon_s}\}$, rotate $C_{T,3\varepsilon_s}\cup C_{B,\varepsilon_s}$ and pack on its left. Total width $\leq (1/2+4\varepsilon_s)N\leq (1-\varepsilon_s)N$ as $\varepsilon_s\leq 1/10$.

Otherwise, no items in $S_{B,\varepsilon_s}\cap S_{T,3\varepsilon_s}$ are in $D_{L,\varepsilon_s}\cup D_{R,\varepsilon_s}$. So let us assume that $i\in C_{L,\varepsilon_s}$ (similarly one can consider $i\in C_{R,\varepsilon_s}$), then we take $M_1=E_{T,3\varepsilon_s}\setminus (C_{L,\varepsilon_s}\cup C_{R,\varepsilon_s})$. Then we can rotate C_{L,ε_s} and C_{R,ε_s} and pack them on top of $M\setminus (M_1\cup C_{L,\varepsilon_s}\cup C_{R,\varepsilon_s})$ as in Case 1.

E Some Tools

In this section, we review some standard building blocks that we rely on in our construction.

E.1 Next Fit Decreasing Height

One of the most recurring tools used as a subroutine in countless results on geometric packing problems is the Next Fit Decreasing Height (NFDH) algorithm, which was originally analyzed in [11] in the context of Strip Packing. We will use a variant of this algorithm to pack items inside a box, and analyze its properties. We provide a full proof for the sake of self-completeness.

Suppose you are given a box C of size $w \times h$, and a set of items I' each one fitting in the box (without rotations). NFDH computes in polynomial time a packing (without rotations) of $I'' \subseteq I'$ as follows. It sorts the items $i \in I'$ in non-increasing order of height h_i , and considers items in that order i_1, \ldots, i_n . Then the algorithm works in rounds $j \geq 1$. At the beginning of round j it is given an index n(j) and a horizontal segment L(j) going from the left to the right side of C. Initially n(1) = 1 and L(1) is the bottom side of C. In round j the algorithm packs a maximal set of items $i_{n(j)}, \ldots, i_{n(j+1)-1}$, with bottom side touching L(j) one next to the other from left to right (a shelf). The segment L(j+1) is the horizontal segment containing the top side of $i_{n(j)}$ and ranging from the left to the right side of C. The process halts at round r when either all items have being packed or $i_{n(r+1)}$ does not fit above $i_{n(r)}$.

We prove the following:

Lemma 52. Assume that, for some given parameter $\varepsilon \in (0,1)$, for each $i \in I'$ one has $w(i) \le \varepsilon w$ and $h(i) \le \varepsilon h$. Then NFDH is able to pack in C a subset $I'' \subseteq I'$ of area at least $a(I'') \ge \min\{a(I'), (1-\varepsilon)\}$

 $(2\varepsilon)w \cdot h$. In particular, if $a(I') \leq (1-2\varepsilon)w \cdot h$, all items in I' are packed.

Proof. The claim trivially holds if all items are packed. Thus suppose that this is not the case. Observe that $\sum_{j=1}^{r+1} h(i_{n(j)}) > h$, otherwise item $i_{n(r+1)}$ would fit in the next shelf above $i_{n(r)}$; hence $\sum_{i=2}^{r+1} h(i_{n(j)}) > h - h(i_{n(1)}) \ge (1-\varepsilon)h$. Observe also that the total width of items packed in each round j is at least $w - \varepsilon w = (1-\varepsilon)w$, since $i_{n(j+1)}$, of width at most εw , does not fit to the right of $i_{n(j+1)-1}$. It follows that the total area of items packed in round j is at least $(w - \varepsilon w)h(n(j+1) - 1)$, and thus

$$a(I'') \ge \sum_{j=1}^r (1-\varepsilon)w \cdot h(n(j+1)-1) \ge (1-\varepsilon)w \sum_{j=2}^{r+1} h(n(j)) \ge (1-\varepsilon)^2 w \cdot h \ge (1-2\varepsilon)w \cdot h.$$

E.2 Maximum Generalized Assignment Problem

In this section we show that there is a PTAS for the Maximum Generalized Assignment Problem (GAP) if the number of bins is constant. In GAP, we are given a set of k bins with capacity constraints and a set of n items that have a possibly different size and profit for each bin and the goal is to pack a maximum-profit subset of items into the bins. Let us assume that if item i is packed in bin j, then it requires size $s_{ij} \in \mathbb{Z}$ and profit $p_{ij} \in \mathbb{Z}$.

GAP is known to be APX-hard and the best known polynomial time approximation algorithm has ratio $(1-1/e+\varepsilon)$ [15, 12]. In fact, for arbitrarily small constant $\delta>0$ (which can even be a function of n) GAP remains APX-hard even on the following instances: bin capacities are identical, and for each item i and bin j, $p_{ij}=1$, and $s_{ij}=1$ or $s_{ij}=1+\delta$ [8]. The complementary case, where item sizes do not vary across bins but profits do, is also APX-hard [8]. However, when all profits and sizes are same across all bins (i.e., $p_{ij}=p_{ik}$ and $s_{ij}=s_{ik}$ for all bins j,k), the problem is known as multiple knapsack problem (MKP) and it admits PTAS [8].

On the other hand, for our purposes we only need instances where k = O(1). A PTAS for GAP for a constant number of bins follows from extending known techniques from the literature [28, 31]. However, we did not find an explicit proof in the literature and thus, for the sake of completeness, in this section we present a full, self-contained description of such an algorithm.

Let C_j be the capacity of bin j for $j \in [k]$. Let p(OPT) be the cost of the optimal assignment.

Lemma 53. There is a $O\left(\left(\frac{1+\varepsilon}{\varepsilon}\right)^k n^{k+1}\right)$ time algorithm for the maximum generalized assignment problem with k bins, which returns a solution with profit at least p(OPT) if we are allowed to augment the bin capacities by a $(1+\varepsilon)$ -factor for any fixed $\varepsilon>0$.

Proof. For each $i \in [n]$ and $c_j \in [C_j]$ for $j \in [k]$, let S_{i,c_1,c_2,\dots,c_k} denote a subset of the set of items $\{1,2,\dots,i\}$ packed into the bins such that the profit is maximized and capacity of bin j is at most c_j . Let $P[i,c_1,c_2,\dots,c_k]$ denote the profit of S_{i,c_1,c_2,\dots,c_k} . Clearly $P[1,c_1,c_2,\dots,c_k]$ is known for all $c_j \in [C_j]$ for $j \in [k]$. Moreover, we define $P[i,c_1,c_2,\dots,c_k]=0$ if $c_j<0$ for any $j\in [k]$. We can compute the value of $P[i,c_1,c_2,\dots,c_k]$ by using a dynamic program (DP), that exploits the following recurrence:

$$P[i, c_1, c_2, \dots, c_k] = \max\{P[i-1, c_1, c_2, \dots, c_k],$$

$$\max_j \{p_{ij} + P[i-1, c_1, \dots, c_j - s_{ij}, \dots, c_k]\}\}$$

With a similar recurrence, we can easily compute a corresponding set $S_{i,c_1,c_2,...,c_k}$.

The running time of the above program is $O\left(n\prod_{j=1}^k C_j\right)$. If each C_j is polynomially bounded, then this running time is polynomial. Therefore, we now create a modified instance where each bin size is polynomially bounded.

Let
$$\mu_j = \frac{\varepsilon C_j}{n}$$
. For item i and bin j , define the modified size $s'_{ij} = \left\lceil \frac{s_{ij}}{\mu_j} \right\rceil = \left\lceil \frac{ns_{ij}}{\varepsilon C_j} \right\rceil$ and $C'_j = \left\lfloor \frac{(1+\varepsilon)C_j}{\mu_j} \right\rfloor$. Note that $C'_j = \left\lfloor \frac{(1+\varepsilon)n}{\varepsilon} \right\rfloor \leq \frac{(1+\varepsilon)n}{\varepsilon}$, so the above DP requires time at most $O\left(n \cdot \left(\frac{(1+\varepsilon)n}{\varepsilon}\right)^k\right)$

The above DP finds the optimal solution $OPT_{modified}$ for the modified instance. Now consider the optimal solution for the original instance (i.e., with original item sizes and bin sizes) $OPT_{original}$. If we show the same assignment of items to the bins is a feasible solution (with modified bin sizes and item sizes) for the modified instance, we get $OPT_{modified} \geq OPT_{original}$ and that will conclude the proof.

Let S_j be the set of items packed in bin j in the $OPT_{original}$. So, $\sum_{i \in S_j} s_{ij} \leq C_j$. Hence,

$$\sum_{i \in S_j} s'_{ij} \le \left[\sum_{i \in S_j} \left(\frac{s_{ij}}{\mu_j} + 1 \right) \right] \le \left[\frac{1}{\mu_j} \left(\sum_{i \in S_j} s_{ij} + |S_j| \mu_j \right) \right] \le \left[\frac{1}{\mu_j} (C_j + n\mu_j) \right] \le \left[\frac{(1+\varepsilon)C_j}{\mu_j} \right] = C'_j$$

Thus $OPT_{original}$ is a feasible solution for the modified instance and the DP will return a packing with profit at least p(OPT) under ε -resource augmentation.

Now we can show how to employ this result to obtain a feasible solution with an almost optimal profit using the original bin capacities.

Lemma 54. There is an algorithm for maximum generalized assignment problem with k bins that runs in time $O\left(\left(\frac{1+\varepsilon}{\varepsilon}\right)^k n^{k/\varepsilon^2+k+1}\right)$ and returns a solution that has profit at least $(1-3\varepsilon)p(OPT)$, for any fixed $\varepsilon>0$.

Proof. First, we claim the following:

Claim 1. If a set of items R_j is packed in a bin B_j with capacity C_j , then there exists a set of at most $O(1/\varepsilon^2)$ items X_j , and a set of items Y_j with $p(Y_j) \leq \varepsilon p(R_j)$ such that all items in $R_j \setminus (X_j \cup Y_j)$ have size at most $\varepsilon(C_j - \sum_{i \in X_j} s_{ij})$.

Proof. Let Q_1 be the set of items i with $s_{ij} > \varepsilon C_j$. If $p(Q_1) \le \varepsilon p(R_j)$, we are done by taking $Y_j = Q_1$ and $X_j = \phi$. Otherwise, define $X_j := Q_1$ and we continue the next iteration with the remaining items. Let Q_2 be the items with size greater than $\varepsilon(C_j - \sum_{i \in X_j} s_{ij})$ in $R_j \setminus X_j$. If $p(Q_2) \le \varepsilon p(R_j)$, we are done by taking $Y_j = Q_2$. Otherwise define $X_j := Q_1 \cup Q_2$ and we continue with further iterations till we get a set Q_t with $p(Q_t) \le \varepsilon p(R_j)$. Note that we need at most $\frac{1}{\varepsilon}$ iterations since the sets Q_i are disjoint.

Otherwise,
$$p(R_j) \geq \sum_{i=1}^{1/\varepsilon} p(Q_i) > \sum_{i=1}^{1/\varepsilon} \varepsilon p(R_j) \geq p(R_j)$$
, which is a contradiction. Thus, consider $Y_j = Q_t$ and $X_j = \bigcup_{l=1}^{t-1} Q_l$. One has $|X_j| \leq 1/\varepsilon^2$ and $p(Y_j) \leq \varepsilon p(R_j)$. On the other hand, after removing Q_t , the remaining items have size $< \varepsilon (C_j - \sum_{i \in X_j} s_{ij})$.

Now consider a bin with bin capacity of $(C_j - \sum_{i \in X_j} s_{ij})$ where all packed items R'_j have sizes $< \varepsilon(C_j - \sum_{i \in X_j} s_{ij})$, then we can divide the bin into $1/\varepsilon$ equal sized intervals $S_{j,1}, S_{j,2}, \ldots, S_{j,1/\varepsilon}$ of lengths $\varepsilon(C_j - \sum_{i \in X_j} s_{ij})$. Let $R'_{j,l}$ be the set of items intersecting the interval $S_{j,l}$. As each packed item can belong

to at most two such intervals, the cheapest set R'' among $\{R'_{j,1},\ldots,R'_{j,1/\varepsilon}\}$ has profit at most $2\varepsilon p(R'_j)$. Thus we can remove this set R'' and reduce the bin size by a factor of $(1-\varepsilon)$.

Now consider the packing of k bins B_j 's in the optimal packing OPT. Let R_j be the set of items packed in bin B_j . Now the algorithm first guesses all X_j 's, a constant number of items, in all k bins. We assign them to corresponding bins in $O(n^{k/\varepsilon^2})$ time. Then for bin j we are left with capacity $r_j := C_j - \sum_{i \in X} s_{ij}$. From previous discussion, we know that there is packing of $R_j'' \subseteq R_j \setminus X_j$ of profit $(1-2\varepsilon)p(R_j \setminus X_j)$ in a bin with capacity $(1-\varepsilon)C_j$. Thus we can use resource augmentation algorithm for GAP in Lemma 53 to pack remaining items in k bins where for bin j we use original capacity to be $(1-\varepsilon)C_j$ for $j \in [k]$ before the resource augmentation. As Lemma 53 returns the optimal packing on this modified bin sizes we get total profit $\geq (1-3\varepsilon)p(OPT)$.

F Packing rectangles with resource augmentation

In this section we prove that it is possible to pack a high profit subset of rectangles into boxes, if we are allowed to augment one side of a knapsack by a small fraction.

The result is essentially proved in [23], although we introduced some modifications and extensions to obtain the additional properties relative to packing into containers and a guarantee on the area of the obtained packing. For the sake of completeness, we provide a full proof, which follows in spirit the proof of the original result, from which we also borrow several notations. We will prove the following stronger version of Lemma 19:

We say that a container C' is smaller than a container C if $w(C') \leq w(C)$ and $h(C') \leq h(C)$. Given a container C and a positive $\varepsilon < 1$, we say that a rectangle R_j is ε -small for C if $w_j \leq \varepsilon w(C)$ and $h_j \leq \varepsilon h(C)$.

Lemma 55 (Resource Augmentation Packing Lemma). Let I' be a collection of rectangles that can be packed into a box of size $a \times b$, and $\varepsilon_{ra} > 0$ be a given constant. Then there exists a container packing of $I'' \subseteq I'$ inside a box of size $a \times (1 + \varepsilon_{ra})b$ (resp., $(1 + \varepsilon_{ra})a \times b$) such that:

- 1. $p(I'') \ge (1 O(\varepsilon_{ra}))p(I');$
- 2. the number of containers is $O_{\varepsilon_{ra}}(1)$ and their sizes belong to a set of cardinality $n^{O_{\varepsilon_{ra}}(1)}$ that can be computed in polynomial time;
- 3. the total area of the containers is at most $a(I') + \varepsilon_{ra}ab$.

In this result, we assume that rectangles in an area container C are ε_{ra} -small for C.

Note that we do not allow rotations, that is, rectangles are packed with the same orientation as in the original packing. However, as an existential result we can apply it also to the case with rotations. Moreover, since Lemma 35 gives a PTAS for approximating container packings, this implies a simple algorithm that does not need to solve any LP to find the solution, in both the cases with and without rotations.

For simplicity, in this section we assume that widths and heights are positive real numbers in (0,1], and a=b=1: in fact, all elements, container and boxes can be rescaled without affecting the property of a packing of being a *container packing* with the above conditions. Thus, without loss of generality, we prove the statement for the augmented $1 \times (1 + \varepsilon_{ra})$ box.

Let $\varepsilon'_{ra} = \varepsilon_{ra}/2 < \varepsilon_{ra}$. We will first obtain a packing where all the elements of each area container C are ε'_{ra} -small for C, and in Section F.6 we will obtain the final packing, where the sizes of each container are taken from a polynomially sized set of choices.

We will use the following Lemma, that follows from the analysis in [26]:

Lemma 56 (Kenyon and Rémila [26]). Let $\overline{\varepsilon} > 0$, and let Q be a set of rectangles, each of height and width at most 1. Let $\mathcal{L} \subseteq Q$ be the set of rectangles of width at least $\overline{\varepsilon}$, and let $OPT_{SP}(\mathcal{L})$ be the minimum width such that the rectangles in \mathcal{L} can be packed in a box of size $OPT_{SP}(\mathcal{L}) \times 1$.

Then \mathcal{Q} can be packed in polynomial time into a box of height 1 and width $\tilde{w} \leq \max\{OPT_{SP}(\mathcal{L}) + \frac{18}{\overline{\varepsilon}^2}w_{\max}, a(\mathcal{Q})(1+\overline{\varepsilon}) + \frac{19}{\overline{\varepsilon}^2}w_{\max}\}$, where w_{\max} is the maximum width of rectangles in \mathcal{Q} . Furthermore, all the rectangles with both width and height less than $\overline{\varepsilon}$ are packed into at most $\frac{9}{\overline{\varepsilon}^2}$ boxes, and all the remaining rectangles into at most $\frac{27}{\overline{\varepsilon}^3}$ vertical containers.

Note that the boxes containing the rectangles that are smaller than $\overline{\varepsilon}$ are not necessarily packed as containers.

We need the following technical lemma:

Lemma 57. Let $\varepsilon > 0$ and let $f(\cdot)$ be any positive increasing function such that f(x) < x for all x. Then, there exist positive constant values $\delta, \mu \in \Omega_{\varepsilon}(1)$, with $f(\varepsilon) \geq \delta$ and $f(\delta) \geq \mu$ such that the total profit of all the rectangles whose width or height lies in $(\mu, \delta]$ is at most $\varepsilon \cdot p(I')$.

Proof. Define $k+1=2/\varepsilon+1$ constants $\varepsilon_1,\ldots,\varepsilon_{k+1}$, with $\varepsilon_1=f(\varepsilon)$ and $\varepsilon_i=f(\varepsilon_{i+1})$ for each i. Consider the k ranges of widths and heights of type $(\varepsilon_{i+1},\varepsilon_i]$. By an averaging argument there exists one index j such that the total profit of the rectangles in I' with at least one side length in the range $(\varepsilon_{j+1}N,\varepsilon_jN]$ is at most $2\frac{\varepsilon}{2}p(I')$. It is then sufficient to set $\delta=\varepsilon_j$ and $\mu=\varepsilon_{j+1}$.

We use this lemma with $\varepsilon = \varepsilon'_{ra}$, and we will specify the function f later. By properly choosing the function f, in fact, we can enforce constraints on the value of μ with respect to δ , which will be useful several times; the exact constraints will be clear from the analysis. Thus, we remove from I' the rectangles that have at least one side length in $(\mu, \delta]$.

We call a rectangle R_i wide if $w_i > \delta$, high if $h_i > \delta$, short if $w_i \le \mu$ and narrow if $h_i \le \mu$. From now on, we will assume that we start with the optimal packing of the rectangles in R', and we will modify it until we obtain a packing with the desired properties. We remove from R' all the short-narrow rectangles, initially obtaining a packing. We will show in section F.5 how to use the residual space to pack them, with a negligible loss of profit.

As a first step, we round up the widths of all the *wide* rectangles in R' to the nearest multiple of δ^2 ; moreover, we shift them horizontally so that their starting coordinate is an integer multiple of δ^2 (note that, in this process, we might have to shift also the other rectangles in order to make space). Since the width of each wide rectangle is at least δ and $\frac{1}{\delta} \cdot 2\delta^2 = 2\delta$, it is easy to see that it is sufficient to increase the width of the box to $1 + 2\delta$ to perform such a rounding.

F.1 Containers for short-high rectangles

We draw vertical lines across the $1 \times (1+2\delta)$ region spaced by δ^2 , splitting it into $M := \frac{1+2\delta}{\delta^2}$ vertical strips (see Figure 17). Consider each maximal rectangular region which is contained in one such strip and does not overlap any wide rectangle; we define a box for each such region that contains at least one short-high rectangle, and we denote the set of such boxes by \mathcal{B} .

Note that some short rectangles might intersect the vertical edges of the boxes, but in this case they overlap with exactly two boxes. Using a standard shifting technique, we can assume that no rectangle is cut by the boxes by losing profit at most $\varepsilon'_{ra}OPT$: first, we assume that the rectangles intersecting two boxes

⁸Note that the classification of the rectangles in this section is different from the ones used in the main results of this paper, although similar in spirit.




Figure 17: An example of a packing after the short-narrow rectangles have been removed, and the wide rectangles (in dark grey) have been aligned to the M vertical strips. Note that the short-high rectangles (in light gray) are much smaller than the vertical strips.




Figure 18: For each vertical box, we can remove a low profit subset of rectangles (red in the picture), to make space for short-high rectangles that cross the right edge of the box (blue).

belong to the leftmost of those boxes. For each box $B \in \mathcal{B}$ (which has width δ^2 by definition), we divide it into vertical strips of width μ . Since there are $\frac{\delta^2}{\mu} > 2/\varepsilon'_{ra}$ strips and each rectangle overlaps with at most 2 such strips, there must exist one of them such that the profit of the rectangles intersecting it is at most $2\mu p(B) \le \varepsilon'_{ra} p(B)$, where p(B) is the profit of all the rectangles that are contained in or belong to B. We can remove all the rectangles overlapping such strip, creating in B an empty vertical gap of width μ , and then we can move all the rectangles intersecting the right boundary of B to the empty space.

Proposition 58. The number of boxes in \mathcal{B} is at most $\frac{1+2\delta}{\delta^2} \cdot \frac{1}{\delta} \leq \frac{2}{\delta^3}$.

First, by a shifting argument similar to above, we can reduce the width of each box to $\delta^2-\delta^4$ while losing only an ε'_{ra} fraction of the profit of the rectangles in B. Then, for each $B\in\mathcal{B}$, since the maximum width of the rectangles in B is at most μ , by applying Lemma 56 with $\overline{\varepsilon}=\delta^2/2$ we obtain that the rectangles packed inside B can be repacked into a box B' of height h(B) and width at most $w'(B)\leq \max\{\delta^2-\delta^4+\frac{72}{\delta^4}\mu,(\delta^2-\delta^4)(1+\frac{\delta^2}{2})+\frac{76}{\delta^4}\mu\}\leq \delta^2$, which is true if we make sure that $\mu\leq \delta^{10}/76$. Furthermore, the short-high rectangles in B are packed into at most $\frac{216}{\delta^6}\leq \frac{1}{\delta^7}$ vertical containers, assuming without loss of generality that $\delta\leq 1/216$. Note that all the rectangles are packed into vertical containers, because rectangles that have both width and height smaller than $\overline{\varepsilon}$ are short-narrow and we already removed them. Summarizing:

Proposition 59. There is a set $I^+ \subseteq I'$ of rectangles with total profit at least $(1 - O(\varepsilon'_{ra})) \cdot p(I')$ and a corresponding packing for them in a $1 \times (1 + 2\delta)$ region such that:

- every wide rectangle in I^+ has its length rounded up to the nearest multiple of δ^2 and it is positioned so that its left side is at a position x which is a multiple of δ^2 , and
- each box $B \in \mathcal{B}$ storing at least one short-high rectangle has width δ^2 , and the rectangles inside are packed into at most $1/\delta^7$ vertical containers.

F.2 Fractional packing with O(1) containers

Let us consider now the set of rectangles I^+ and an almost optimal packing S^+ for them according to Proposition 59. We remove the rectangles assigned to boxes in $\mathcal B$ and consider each box $B \in \mathcal B$ as a single pseudoitem. Thus, in the new almost optimal solution there are just pseudoitems from $\mathcal B$ and wide rectangles with right and left coordinates that are multiples of δ^2 . We will now show that we can derive a fractional packing with the same profit, and such that the rectangles and pseudoitems can be (fractionally) assigned to a constant number of containers. By *fractional packing* we mean a packing where horizontal rectangles are allowed to be sliced horizontally (but not vertically); we can think of the profit as being split proportionally to the heights of the slices.

Let $\mathcal K$ be a subset of the horizontal rectangles of size K that will be specified later. By extending horizontally the top and bottom edges of the rectangles in $\mathcal K$ and the pseudoitems in $\mathcal B$, we partition the knapsack into at most $2(|K|+|\mathcal B|)+1\leq 2(K+\frac{2}{\delta^3})+1\leq 2(K+\frac{3}{\delta^3})$ horizontal stripes.

Let us focus on the (possibly sliced) rectangles contained in one such stripe of height h. For any vertical coordinate $y \in [0,h]$ we can define the *configuration* at coordinate y as the set of positions where the horizontal line at distance y from the bottom cuts a vertical edge of a horizontal rectangle which is not in \mathcal{K} . There are at most 2^{M-1} possible configurations in a stripe.

We can further partition the stripe in maximal contiguous regions with the same configuration. Note that the number of such regions is not bounded, since configurations can be repeated. But since the rectangles




Figure 19: Rearranging the rectangles in a horizontal stripe. On the right, rectangles are repacked so that regions with the same configuration appear next to each other. Note that the yellow rectangle has been sliced, since it partakes in two regions with different configurations.

are allowed to be sliced, we can rearrange the regions so that all the ones with the same configuration appear next to each other; see Figure 19 for an example. After this step is completed, we define up to M horizontal containers per each configuration, where we repack the sliced horizontal rectangles. Clearly, all sliced rectangles are repacked.

Thus, the number of horizontal containers that we defined per each stripe is bounded by $M2^{M-1}$, and the total number overall is at most

$$2\left(K+\frac{3}{\delta^3}\right)M2^{M-1}=\left(K+\frac{3}{\delta^3}\right)M2^M.$$

F.3 Existence of an integral packing

We will now show the existence of an integral packing, at a small loss of profit.

Consider a fractional packing in N containers. Since each rectangle slice is packed in a container of exactly the same width, it is possible to pack all but at most N rectangles integrally by a simple greedy algorithm: choose a container, and greedily pack in it rectangles of the same width, until either there are no rectangles left for that width, or the next rectangle does not fit in the current container. In this case, we discard this rectangle and close the container, meaning that we do not use it further. Clearly, only one rectangle per container is discarded, and no rectangle is left unpacked.

The only problem is that the total profit of the discarded rectangles can be large. To solve this problem, we use the following shifting argument. Let $\mathcal{K}_0 = \emptyset$ and $K_0 = 0$. For convenience, let us define $f(K) = (K + \frac{3}{\delta^3}) M2^M$.

First, consider the fractional packing obtained by choosing $\mathcal{K}=\mathcal{K}_0$, so that $K=K_0=0$. Let \mathcal{K}_1 be the set of discarded rectangles obtained by the greedy algorithm, and let $K_1=|\mathcal{K}_1|$. Clearly, by the above reasoning, the number of discarded rectangles is bounded by $f(K_0)$. If the profit $p(\mathcal{K}_1)$ of the discarded rectangles is at most $\varepsilon'_{ra}p(OPT)$, then we remove them and there is nothing else to prove. Otherwise, consider the fractional packing obtained by fixing $\mathcal{K}=\mathcal{K}_0\cup\mathcal{K}_1$. Again, we will obtain a set \mathcal{K}_2 of discarded rectangles such that $K_2:=|\mathcal{K}_2|\leq f(K_0+K_1)$. Since the sets $\mathcal{K}_1,\mathcal{K}_2,\ldots$ that we obtain are all disjoint, the process must stop after at most $1/\varepsilon'_{ra}$ iterations. Setting $p:=M2^M$ and $q:=\frac{3}{\delta^3}M2^M$, we have that $K_{i+1}\leq p(K_0+K_1+\ldots K_i)+q$ for each $i\geq 0$. Crudely bounding it as $K_{i+1}\leq i\cdot pq\cdot K_i$, we immediately obtain that $K_i\leq (pq)^i$. Thus, in the successful iteration, the size of \mathcal{K} is at most $K_{1/\varepsilon'_{ra}-1}$ and the number of containers is at most $K_{1/\varepsilon'_{ra}}\leq (pq)^{1/\varepsilon'_{ra}}=(\frac{3}{\delta^2}M^22^{2M})^{1/\varepsilon'_{ra}}=O_{\varepsilon'_{ra},\delta}(1)$.

F.4 Rounding down horizontal and vertical containers

As per the above analysis, the total number of horizontal containers is at most $(\frac{3}{\delta^2}M^22^{2M})^{\varepsilon'_{ra}}$ and the total number of vertical containers is at most $\frac{2}{\delta^3} \cdot \frac{1}{\delta^7} = \frac{2}{\delta^{10}}$.

We will now show that, at a small loss of profit, it is possible to replace each horizontal and each vertical container defined so far with a constant number of smaller containers, so that the total area of the new containers is at most as big as the total area of the rectangles originally packed in the container. Note that in each container we consider the rectangles with the original widths (not rounded up). We use the following lemma:

Lemma 60. Let C be a horizontal (resp. vertical) container defined above, and let I_C be the set of rectangles packed in C. Then, it is possible to pack a set $I'_C \subseteq I_C$ of profit at least $(1 - 3\varepsilon'_{ra})p(I_C)$ in a set of at most $\left\lceil \log_{1+\varepsilon'_{ra}}(\frac{1}{\delta}) \right\rceil / \varepsilon'^2_{ra}$ horizontal (resp. vertical) containers that can be packed inside C and such that their total area is at most $a(I_C)$.

Proof. Without loss of generality, we prove the result only for the case of a horizontal container.

Since $w_i \geq \delta$ for each rectangle $R_i \in I_C$, we can partition the rectangles in I_C into at most $\left|\log_{1+\varepsilon_{ra}'}(\frac{1}{\delta})\right|$ groups I_1, I_2, \ldots , so that in each I_j the widest rectangle has width bigger than the smallest by a factor at most $1+\varepsilon_{ra}'$; we can then define a container C_j for each group I_j that has the width of the widest rectangle it contains and height equal to the sum of the heights of the contained rectangles.

Consider now one such C_j and the set of rectangles I_j that it contains, and let $P:=p(I_j)$. Clearly, $w(C_j) \leq (1+\varepsilon'_{ra})w_i$ for each $R_i \in I_j$, and so $a(C_j) \leq (1+\varepsilon'_{ra})a(I_j)$. If all the rectangles in I_j have height at most $\varepsilon'_{ra}h(C_j)$, then we can remove a set of rectangles with total height at least $\varepsilon'_{ra}h(C)$ and profit at most $2\varepsilon'_{ra}p(I_j)$. Otherwise, let $\mathcal Q$ be the set of rectangles of height larger than $\varepsilon'_{ra}h(C_j)$, and note that $a(Q) \geq \varepsilon'_{ra}h(C_j)w(C_j)/(1+\varepsilon'_{ra})$. If the $p(\mathcal Q) \leq \varepsilon'_{ra}P$, then we remove the rectangles in $\mathcal Q$ from the container C_j and reduce its height as much as possible, obtaining a smaller container C'_j ; since $a(C'_j) \leq a(C_j) - \varepsilon'_{ra}a(C_j) = (1-\varepsilon'_{ra})a(C_j) \leq (1-\varepsilon'_{ra})(1+\varepsilon'_{ra})a(I_j) < a(I_j)$, then the proof is finished. Otherwise, we define one container for each of the rectangles in $\mathcal Q$ (which are at most $1/\varepsilon'_{ra}$) of exactly the same size, and we still shrink the container with the remaining rectangles as before; note that there is no lost area for each of the newly defined container. Since at every non-terminating iteration a set of rectangles with profit larger than $\varepsilon'_{ra}P$ is removed, the process must end within $1/\varepsilon'_{ra}$ iterations.

Note that the total number of containers that we produce for each initial container C_j is at most $1/\varepsilon_{ra}^{\prime 2}$, and this concludes the proof.

Thus, by applying the above lemma to each horizontal and each vertical container, we obtain a modified packing where the total area of the horizontal and vertical containers is at most the area of the rectangles of I' (without the short-narrow rectangles, which we will take into account in the next subsection), while the number of containers increases at most by a factor $\left\lceil \log_{1+\varepsilon'_{ra}}(\frac{1}{\delta}) \right\rceil / \varepsilon'^2_{ra}$.

F.5 Packing short-narrow rectangles

Consider the integral packing obtained from the previous subsection, which has at most $K':=\left(\frac{2}{\delta^{10}}+(\frac{3}{\delta^2}M^22^{2M})^{\varepsilon'_{ra}}\right)\left\lceil\log_{1+\varepsilon'_{ra}}(\frac{1}{\delta})\right\rceil/\varepsilon'^2_{ra}$ containers. We can create a non-uniform grid extending each side of the containers until they hit another container or the boundary of the knapsack. Moreover, we also add horizontal and vertical lines spaced at distance ε'_{ra} . We call *free cell* each face defined by the above lines that does not overlap a container of the packing; by construction, no free cell has a

side bigger than ε'_{ra} . The number of free cells in this grid plus the existing containers is bounded by $K_{TOTAL} = (2K' + 1/\varepsilon'_{ra})^2 = O_{\varepsilon'_{ra},\delta}(1)$. We crucially use the fact that this number does not depend on value of μ .

Note that the total area of the free cells is no less than the total area of the short-narrow rectangles, as a consequence of the guarantees on the area of the containers introduced so far. We will pack the short-narrow rectangles into the free cells of this grid using NFDH, but we only use cells that have width and height at least $\frac{8\mu}{\varepsilon_{ra}}$; thus, each short-narrow rectangle will be assigned to a cell whose width (resp. height) is larger by at least a factor $8/\varepsilon_{ra}'$ than the width (resp. height) of the rectangle. Each discarded cell has area at most $\frac{8\mu}{\varepsilon_{ra}'}$, which implies that the total area of discarded cells is at most $\frac{8\mu K_{TOTAL}}{\varepsilon_{ra}'}$. Now we consider the selected cells in an arbitrary order and pack short narrow rectangles into them using NFDH, defining a new area container for each cell that is used. Thanks to Lemma 52, we know that each new container C (except maybe the last one) that is used by NFDH contains rectangles for a total area of at least $(1 - \varepsilon_{ra}'/4)a(C)$. Thus, if all rectangles are packed, we remove the last container opened by NFDH, and we call S the set of rectangles inside, that we will repack elsewhere; note that $a(S) \le \varepsilon_{ra}'^2 \le \varepsilon_{ra}'/3$, since all the rectangles in S were packed in a free cell. Instead, if not all rectangles are packed by NFDH, let S be the residual rectangles. In this case, the area of the unpacked rectangles is $a(S) \le \frac{8\mu K_{TOTAL}}{\varepsilon_{ra}'} + \varepsilon_{ra}'/4 \le \varepsilon_{ra}'/3$, assuming that s

 $\mu \leq \frac{\varepsilon_{ra}'^2}{96K_{TOTAL}}.$ In order to repack the rectangles of S, we define a new area container C_S of height 1 and width $\varepsilon_{ra}'/2$. Since $a(C_S) = \varepsilon_{ra}'/2 \geq (\varepsilon_{ra}'/3)/(1-2\varepsilon_{ra}')$, all elements from S are packed in C_S by NFDH, and the container can be added to the knapsack by further enlarging its width from $1+2\delta$ to $1+2\delta+\varepsilon_{ra}'/2 < 1+\varepsilon_{ra}'$.

The last required step is to guarantee the necessary constraint on the total area of the area containers, similarly to what was done in Section F.4 for the horizontal and vertical containers.

Let D be any full area container (that is, any area container except for C_S). We know that the area of the rectangles R_D in D is $a(R_D) \geq (1 - \varepsilon'_{ra})a(D)$, since each rectangle R_i inside D has width less than $\varepsilon'_{ra}w(D)/2$ and height less than $\varepsilon'_{ra}h(D)/2$, by construction. We remove rectangles from R_D in non-decreasing order of profit/area ratio, until the total area of the residual rectangles is between $(1 - 4\varepsilon'_{ra})a(D)$ and $(1 - 3\varepsilon'_{ra})a(D)$ (this is possible, since each element has area at most $\varepsilon'^2_{ra}a(D)$); let R'_D be the resulting set. We have that $p(R'_D) \geq (1 - 4\varepsilon'_{ra})p(R_D)$, due to the greedy choice. Let us define a container D' of width w(D) and height $(1 - \varepsilon'_{ra})h(D)$. It is easy to verify that each rectangle in R_D has width (resp. height) at most $\varepsilon'_{ra}w(D')$ (resp. $\varepsilon'_{ra}h(D')$). Moreover, since $a(R'_D) \leq (1 - 3\varepsilon'_{ra})a(D) \leq (1 - 2\varepsilon'_{ra})(1 - \varepsilon'_{ra})a(C) \leq (1 - 2\varepsilon'_{ra})a(C')$, then all elements in R'_D are packed in D'. By applying this reasoning to each area container (except C_S), we obtain property (3) of Lemma 55.

Note that the constraints on μ and δ that we imposed are $\mu \leq \frac{\delta^{10}}{76}$ (from Section F.1), and $\mu \leq \frac{\varepsilon_{ra}'^2}{96K_{TOTAL}}$. It is easy to check that both of them are satisfied if we choose $f(x) = (\varepsilon_{ra}'x)^C$ for a big enough constant C that depends only on δ and ε_{ra}' .

F.6 Rounding containers to a polynomial set of sizes

In this subsection we show that it is possible to round down the size of each horizontal, vertical or area container so that the resulting sizes can be chosen from a polynomially sized set, while incurring in a marginal loss of profit.

For a set I of rectangles, we define $WIDTHS(I) = \{w_j \mid R_j \in I\}$ and $HEIGHTS(R) = \{h_j \mid R_j \in I\}$.

Given a finite set P of real numbers and a fixed natural number k, we define the set $P^{(k)} = \{(p_1 + p_2 + p_3) \in P \mid (p_1 + p_2) \in P \}$

 $\cdots + p_l$) + $ip_{l+1} \mid p_j \in P \; \forall \; j, l \leq k, 0 \leq i \leq n, i \; \text{integer}$ }; note that if |P| = O(n), then $|P^{(k)}| = O(n^{k+2})$. Moreover, if $P \subseteq Q$, then obviously $P^{(k)} \subseteq Q^{(k)}$, and if $k' \leq k''$, then $P^{(k')} \subseteq P^{(k'')}$.

Lemma 61. Let $\varepsilon > 0$, and let I be a set of rectangles packed in a horizontal or vertical container C. Then, for any $k \ge 1/\varepsilon$, there is a set $I' \subseteq I$ with profit $p(I') \ge (1-\varepsilon)p(I)$ that can be packed in a container C' smaller than C such that $w(C') \in WIDTHS(I)^{(k)}$ and $h(C') \in HEIGHTS(R)^{(k)}$.

Proof. Without loss of generality, we prove the thesis for an horizontal container C; the proof for vertical containers is symmetric. Clearly, the width of C can be reduced to $w_{max}(I)$, and $w_{max}(I) \in WIDTHS(I) \subseteq WIDTHS(I)^{(k)}$.

If $|I| \leq 1/\varepsilon$, then $\sum_{R_i \in I} h_i \in HEIGHTS(I)^{(k)}$ and there is no need to round the height of C down. Otherwise, let I_{TALL} be the set of the $1/\varepsilon$ rectangles in I with largest height (breaking ties arbitrarily), let R_j be the least profitable of them, and let $I' = I \setminus \{R_j\}$. Clearly, $p(I') \geq (1-\varepsilon)p(I)$. Since each element of $I' \setminus I_{TALL}$ has height at most h_j , it follows that $h(I \setminus I_{TALL}) \leq (n-1/\varepsilon)h_j$. Thus, letting $i = \lceil h(I' \setminus I_{TALL})/h_j \rceil \leq n$, all the rectangles in I' fit in a container C' of width $w_{max}(I)$ and height $h(C') := h(I_{TALL}) + ih_j \in HEIGHTS(R)^{(k)}$. Since $h(I_{TALL}) + ih_j \leq h(I_{TALL}) + h(I' \setminus I_{TALL}) + h_j = h(I) \leq h(C)$, this proves the result.

Lemma 62. Let $\varepsilon > 0$, and let I be a set of rectangles that are assigned to an area container C. Then there exists a subset $I' \subseteq I$ with profit $p(I') \ge (1 - 3\varepsilon)p(I)$ and a container C' smaller than C such that: $a(I') \le a(C)$, $w(C') \in WIDTHS(I)^{(0)}$, $h(C') \in HEIGHTS(I)^{(0)}$, and each $R_j \in I'$ is $\frac{\varepsilon}{1 - \varepsilon}$ -small for C'.

Proof. Without loss of generality, we can assume that $w(C) \leq nw_{max}(I)$ and $h(C) \leq nh_{max}(I)$: if not, we can first shrink C so that these conditions are satisfied, and all the rectangles still fit in C.

Define a container C' so that it has width $w(C') = w_{max}(I) \lfloor w(C)/w_{max}(I) \rfloor$ and height $h(C') = h_{max}(I) \lfloor h(C)/h_{max}(I) \rfloor$, that is, C' is obtained by shrinking C to the closest integer multiples of $w_{max}(I)$ and $h_{max}(I)$. Observe that $w(C') \in WIDTHS(I)^{(0)}$ and $h(C') \in HEIGHTS(I)^{(0)}$. Clearly, $w(C') \geq w(C) - w_{max}(I) \geq w(C) - \varepsilon w(C) = (1 - \varepsilon)w(C)$, and similarly $h(C') \geq (1 - \varepsilon)h(C')$. Hence $a(C') \geq (1 - \varepsilon)^2 a(C) \geq (1 - 2\varepsilon)a(C)$.

We now select a set $I'\subseteq I$ by greedily choosing elements from I in non-increasing order of profit/area ratio, adding as many elements as possible without exceeding a total area of $(1-2\varepsilon)a(C)$. Since each element of I has area at most $\varepsilon^2a(C)$, then either all elements are selected (and then p(I')=p(I)), or the total area of the selected elements is at least $(1-2\varepsilon-\varepsilon^2)a(C)\geq (1-3\varepsilon)a(C)$. By the greedy choice, we have that $p(I')\geq (1-3\varepsilon)p(I)$.

Since each rectangle in I is $\frac{\varepsilon}{1-\varepsilon}$ -small for C', this proves the thesis.

By applying Lemmas 61 and 62 with $\varepsilon = \varepsilon'_{ra}$ to all the containers and noting that $\frac{\varepsilon'_{ra}}{1 - \varepsilon'_{ra}} \le \varepsilon_{ra}$, we completed the proof of Lemma 55.

Remark 6. Note that in the above, the size of the container is rounded to a family of sizes that depends on the rectangles inside; of course, they are not known in advance in an algorithm that enumerates over all the container packings. On the other hand, if the instance is a set \mathcal{I} of n rectangles, then for any natural number k we have that $WIDTHS(I)^{(k)} \subseteq WIDTHS(I)^{(k)}$ and $HEIGHTS(I)^{(k)} \subseteq WIDTHS(\mathcal{I})^{(k)}$ for any $I \subseteq I$; clearly, $WIDTHS(\mathcal{I})^{(k)} \times HEIGHTS(I)^{(k)}$ has a polynomial size and can be computed explicitly.

Similarly, when finding container packings for the case with rotations, one can compute the set $SIZES(I) := WIDTHS(I) \cup HEIGHTS(I)$, and consider containers of width and height in $SIZES(I)^{(k)}$ for a sufficiently high constant k.