

The Role of Quantifier Alternations in Cut Elimination

Philipp Gerhardy

Abstract Extending previous results from work on the complexity of cut elimination for the sequent calculus **LK**, we discuss the role of quantifier alternations and develop a measure to describe the complexity of cut elimination in terms of quantifier alternations in cut formulas and contractions on such formulas.

1 Introduction

In this note we will present an extension of results on the complexity of cut elimination in the sequent calculus **LK** first developed in Gerhardy [1] and subsequently published in Gerhardy [2]. There it is shown how the complexity of cut elimination primarily depends on the nesting of quantifiers in cut formulas and contractions on ancestors of such cut formulas. A more complicated proof of the role of quantifier nestings was first given by Zhang in [4].

In this note we extend the analysis and develop a measure that describes with sufficient accuracy the role of quantifier alternations in cut elimination. The measure will be slightly more complicated than the notion of nested quantifier depth, *nqf*, described in [2], but will generalize with similar ease to incorporate the role of contractions. An earlier, more complicated treatment of the role of quantifier alternations by Zhang can be found in [5]. Though leading to comparable results, the measure of the cut complexity described in [5] is far more complicated than the one presented in this note. For another attempt at defining a measure capturing quantifier alternations, albeit without accompanying proof of cut elimination, see Visser [3]. Neither Zhang nor Visser treats the role of contractions in the complexity of cut elimination.

Received February 17, 2004; accepted May 17, 2004; printed May 25, 2005

2000 Mathematics Subject Classification: Primary, 03F05, 03F20

Keywords: proof complexity, cut elimination, quantifier alternations

©2005 University of Notre Dame

2 Previous Results

Let **LK** be the sequent calculus as defined in [2], that is, with multiplicative rules and with no implicit contractions. Let $|\cdot|$ denote the depth and $||\cdot||$ the size of formulas and proofs, in the latter case not counting weakenings and contractions. Let $nqf(\cdot)$, $dqf(\cdot)$, and $cnqf(\cdot)$ denote the nested quantifier depth, the deepest quantified formula (informally, the largest number of propositional connectives one has to “peel off” to get to a quantifier) and the contracted nested quantifier depth, respectively, as defined in [2].

In [2] the following results are proved.

Lemma 2.1 (Refined Reduction Lemma) *Let φ be an **LK**-proof of a sequent $\Gamma \vdash \Delta$ with the final inference a cut with cut formula A . Then, if for all other cut formulas B ,*

1. *$nqf(A) \geq nqf(B)$ and $dqf(\varphi) = dqf(A) > dqf(B)$, then there exists a proof φ' of the same sequent with $dqf(\varphi') \leq dqf(\varphi) - 1$ and $|\varphi'| \leq |\varphi| + 1$,*
2. *$nqf(\varphi) = nqf(A) > nqf(B)$ and $dqf(A) = 0$, then there exists a proof φ' of the same sequent with $nqf(\varphi') \leq nqf(\varphi) - 1$ and $|\varphi'| < 2 \cdot |\varphi|$.*

If the cut formula A is atomic and both subproofs are cut-free, then there is a cut-free proof φ' of the same sequent with $|\varphi'| < 2 \cdot |\varphi|$.

Lemma 2.2 *Let φ be an **LK**-proof of a sequent $\Gamma \vdash \Delta$. If $dqf(\varphi) = d > 0$, then there is a proof φ' of the same sequent with $dqf(\varphi') = 0$ and $|\varphi'| \leq 2^d \cdot |\varphi|$.*

Lemma 2.3 *Let φ be an **LK**-proof of a sequent $\Gamma \vdash \Delta$. If $dqf(\varphi) = 0$ and $nqf(\varphi) = d > 0$, then there is a proof φ' of the same sequent with $nqf(\varphi') \leq d - 1$ and $|\varphi'| < 2^{|\varphi|}$.*

Theorem 2.4 (First Refined Cut Elimination Theorem) *Let φ be an **LK**-proof of a sequent $\Gamma \vdash \Delta$. If $nqf(\varphi) = d > 0$, then there is a proof φ' of the same sequent and a constant c , depending only on the propositional nesting of the cut formulas so that $nqf(\varphi') \leq d - 1$ and $|\varphi'| \leq 2^{c \cdot |\varphi|}$.*

Corollary 2.5 *Let φ be an **LK**-proof of a sequent $\Gamma \vdash \Delta$ and let $nqf(\varphi) = d$. Then there is a constant c , depending only on the propositional nesting of the cut formulas, and a proof φ' of the same sequent where φ' is cut-free and $|\varphi'| \leq 2^{c \cdot |\varphi|}_{d+1}$.*

Lemma 2.6 (Contraction Lemma) *Let φ be an **LK**-proof of a sequent $\Gamma \vdash \Delta$, with $nqf(\varphi) > cnqf(\varphi)$. Then there is proof φ' of the same sequent with $nqf(\varphi') = cnqf(\varphi')$ and $||\varphi'|| \leq ||\varphi||$. As a consequence also $|\varphi'| \leq 2^{|\varphi|}$.*

Theorem 2.7 (Second Refined Cut Elimination Theorem) *Let φ be an **LK**-proof of a sequent $\Gamma \vdash \Delta$. Then there is a constant c depending only on the propositional nesting of the cut formulas and a cut-free proof φ' of the same sequent where $|\varphi'| \leq 2^{c \cdot |\varphi|}_{cnqf(\varphi)+2}$.*

The main work is to prove the Refined Reduction Lemma and the Contraction Lemma from which the remaining results follow easily.

To sum up, first it is shown that the complexity of cut elimination primarily depends on the nesting of quantifiers in cut formulas, whereas the elimination of the propositional connectives has a negligible contribution to the complexity of cut elimination. As mentioned above, this result was also shown by Zhang in [4]. Moreover,

if for a cut formula none of the direct ancestors has been contracted, then the cut can be eliminated with low complexity by a mere rearrangement of the proof that does not increase the size of the proof. Thus the nonelementary complexity of cut elimination was shown to depend only on the nested quantifier depth of cut formulas whose ancestors, of sufficient quantifier depth, also have been contracted.

3 Quantifier Alternations

In Section 3.4 of [2] it is discussed that blocks of \forall, \wedge -connectives, respectively, \exists, \vee , can be eliminated together, and it is shown that eliminating such a block from a cut formula at most doubles the depth of the proof. In [2] the following lemma is proved.

Lemma 3.1 *Let φ be a proof of a sequent $\Gamma \vdash \Delta$ with the last inference a cut. Let the cut formula be constructed from formulas B_1, \dots, B_n by the connectives \forall and \wedge only (respectively, \exists and \vee). Then we can replace that cut by a number of smaller cuts with cut formulas B_i . For the resulting proof φ' we have $|\varphi'| < 2 \cdot |\varphi|$.*

This lemma immediately suggests a bound on cut elimination based on the number of alternations between such blocks. We propose the following cut elimination strategy: first we eliminate as many outermost propositional connectives as possible, next we eliminate all outermost \forall, \wedge and \exists, \vee blocks. Repeating this we eventually arrive at a cut-free proof. By the Refined Reduction Lemma and subsequently Lemma 2.2, both of which can easily be adapted to some measure of the number of quantifier alternations instead of nqf , we see that the first step, eliminating propositional connectives, is not critical for the complexity of cut elimination. However, defining a new appropriate complexity measure for this cut elimination strategy is not trivial, as can also be seen by the complicated measure defined by Zhang [5] in order to prove a comparable result.

We want to define a measure aqf , the alternating quantifier depth. First consider the following very naïve approach: Let us restrict the logical connectives to \forall, \exists, \wedge , and \vee and let us count the propositional connectives \wedge, \vee as the quantifiers \forall, \exists . Defining the aqf as the number of alternations between quantifier blocks in cut formulas would not give the desired result. Alternations between propositional connectives \wedge, \vee , which can be eliminated easily, would be perceived as alternations between quantifiers \forall, \exists , which are expensive to eliminate. Thus this definition of aqf would lead to a bound on cut elimination much worse in such situations than the bound already achieved via the nested quantifier depth nqf .

In general it turns out to be difficult to define, inductively on the formula, a measure of cut complexity that correctly captures the role of quantifier alternations. The difficulty is to decide when to increase the alternating quantifier depth.

For example, when facing a formula composed of two subformulas and one of the propositional connectives, for example, the connective \vee , it is nontrivial to decide or predict whether the connective is part of a block of propositional connectives, and hence relatively harmless with respect to the alternations already present in the two subformulas, or marks the beginning of an \exists, \vee block and hence leads to an increase in the number of alternations.

Consider the formula $A := B \vee C$, where the subformula C is assumed to be purely \forall, \wedge :

1. if B is purely \exists, \vee then
 - (a) $aqf(A)$ should be 0—eliminate the \vee (simple), then two blocks each without alternations remain;
 - (b) $aqf(\exists x A)$ should be 1—all the \exists, \vee constitute one block, the \forall, \wedge block below constitutes the alternation;
 - (c) $aqf(\forall x A)$ should be 1—eliminate the \forall (expensive), eliminate the \vee (simple), eliminate the two alternation free blocks;
2. if B has an outermost \exists, \vee block and one \forall, \wedge block below
 - (a) $aqf(A)$ should be 1—eliminate the \vee (simple), then one alternation remains;
 - (b) $aqf(\exists x A)$ should be 1—all the \exists, \vee constitute one block, the \forall, \wedge blocks below constitute the alternation;
 - (c) $aqf(\forall x A)$ should be 2—eliminate the \forall (expensive), eliminate the \vee (simple), and still one alternation remains in the subformula B .

The example demonstrates the problem of deciding *inductively* on the formula when to increase the alternating quantifier depth. At the point of the propositional connective we might not yet have sufficient information to decide whether to increase or not. On the other hand postponing the decision until we meet the next quantifier requires information on the exact structure of the subformulas that may no longer be available.

The solution is to let the complexity measure mirror the intended cut elimination strategy. This leads to defining the measure aqf for the cut complexity *recursively* on the cut formula instead of inductively.

Definition 3.2 We define aqf as follows:

1. if A is atomic, purely \exists, \vee , or purely \forall, \wedge then $aqf(A) = 0$;
2. if A is composed of formulas B_1, \dots, B_n (each with outermost connective a quantifier) by propositional connectives only then $aqf(A) = \max\{aqf(B_i)\}$;
3. if $A := \forall x C$ for some C is composed of connectives \forall, \wedge and formulas B_1, \dots, B_n (each with outermost connective \exists, \vee) then $aqf(A) = \max\{aqf(B_i)\} + 1$;
4. if $A := \exists x C$ for some C is composed of connectives \exists, \vee and formulas B_1, \dots, B_n (each with outermost connective \forall, \wedge) then $aqf(A) = \max\{aqf(B_i)\} + 1$.

Moreover, we treat implication $B \rightarrow C$ as $\neg B \vee C$, and negation $\neg B$ simply flips the polarity of other connectives below, that is, $\exists, \vee \mapsto \forall, \wedge$ and vice versa.

With this definition of aqf for formulas, we define aqf for proofs.

Definition 3.3 $aqf(\varphi) := \sup\{aqf(A) : A \text{ is a cut formula in } \varphi\}$.

Also the notion of deepest quantified formula dqf defined in [2] can be adapted to aqf yielding a version of the Refined Reduction Lemma with aqf instead of nqf . Now it is easy to show the following theorem.

Theorem 3.4 *Let φ be an **LK**-proof of a sequent $\Gamma \vdash \Delta$ and let $aqf(\varphi) = d$. Then there is a constant c , depending only on the propositional blocks of the cut formulas, and a proof φ' of the same sequent where φ' is cut-free and $|\varphi'| \leq 2_{d+1}^{c \cdot |\varphi|}$.*

Proof As discussed the Refined Reduction Lemma can easily be adapted to the measure aqf instead of nqf . It then follows that one can adapt Lemma 2.2 and

Lemma 2.3 to the measure aqf . The theorem then follows easily from the cut elimination strategy sketched above and the adaptations of Lemma 2.2 and Lemma 2.3 to the measure aqf . \square

The definition of $caqf$, the contracted alternating quantifier depth, is defined from aqf in the same way $cnqf$ is defined from nqf (see [2]). Thus also taking the role of contractions into account we get the following theorem.

Theorem 3.5 *Let φ be an **LK**-proof of a sequent $\Gamma \vdash \Delta$. Then there is a constant c depending only on the propositional blocks of the cut formulas and a cut-free proof φ' of the same sequent where $|\varphi'| \leq 2^{c \cdot |\varphi|}_{caqf(\varphi)+2}$.*

Proof The theorem follows easily from the above theorem and the Contraction Lemma adapted to the measure aqf . \square

In conclusion both theorems follow easily from the analysis of cut elimination presented in [2], in particular Lemma 3.1, and the cut elimination strategy described above. As mentioned above a comparable result is proved in [5] but with a far more complicated complexity measure and a more complicated proof.

Furthermore, as with the upper bounds on cut elimination presented in [2], the bounds aqf and $caqf$ are optimal with regard to Statman's lower bound example, that is, the upper and the lower bound coincide. Conversely, one can say that every proof that yields an example of nonelementary cut elimination must use cut formulas with alternating quantifiers and contractions in a way similar to Statman's lower bound example.

Finally, the exponential bound on cut elimination in the case of pure \forall, \wedge -cuts, respectively \exists, \vee -cuts, that is stated in [2], follows as a special case from these bounds.

4 Comparison with the Literature

In this section we will briefly discuss the measures for the number of quantifier alternations proposed by, respectively, Zhang [5] and Visser [3].

The measure defined by Zhang uses two formula classes δ and δ' .

Definition 4.1 (Zhang [5]) A formula B is in $\delta(A)$ if and only if

1. $A = B$, or
2. $A = C \wedge D$ and $B \in \delta(C) \cup \delta(D)$, or
3. $A = \forall x C(x)$ and $B \in \delta(C(t))$ for any term t .

Definition 4.2 (Zhang [5]) A formula B is in $\delta'(A)$ if and only if

1. $B \in \delta(A)$, and
2. B is either a disjunction of two formulas, or
3. B is an \exists -formula such that all terms occurring in B also occur in A .

Definition 4.3 (Zhang [5]) The cut complexity $\rho(A)$ of a formula A is defined as a polynomial in w as follows:

1. if $\delta'(A)$ is empty, then $\rho(A) := w$,
2. if $\delta'(A) = \{B_i | 1 \leq i \leq n\}$ and there is a formula C such that $\forall x C(x) \in \delta(A)$ and $nqf(C) \geq nqf(B_i)$ for $1 \leq i \leq n$, then $\rho(A) := (\rho(B_1) \oplus \dots \oplus \rho(B_n)) \cdot w$,
3. if $A = B \wedge C$, $\delta'(A) = \{B_i | 1 \leq i \leq n\}$ and there is no formula C such that $\forall x C(x) \in \delta(A)$ and $nqf(C) \geq nqf(B_i)$ for $1 \leq i \leq n$, then $\rho(A) := (\rho(B) \oplus \rho(C)) + 1$,

4. if $\delta'(A) = \{A\}$, then $\rho(A) := \rho(\neg A)$,

where \oplus is the operation of summing two polynomials by raising them to the same degree and then taking the pointwise maximum over their coefficients.

The definition is somewhat similar to the definition of the alternating quantifier depth *aqf* presented in this note, as the degree of the cut complexity polynomial corresponds to our notion of *aqf*.

In detail, the first item in the definition covers the case when the formula is atomic, purely \forall, \wedge or (via the fourth item) purely \exists, \vee . The second item corresponds to eliminating a \forall, \wedge block (or via the fourth item an \exists, \vee block), and hence here the degree of the polynomial is increased. The third item corresponds to eliminating in-between propositional connectives, which only adds a constant to the polynomial.

The proof of cut elimination given the cut complexity polynomial above proceeds via several rather technical lemmas and uses an additional formula class δ^* .

Visser defines a measure “depth of quantifierchanges” via a three place function ϱ ([3], p. 281), where the first parameter is 0 when the formula under consideration occurs positively and 1 if it occurs negatively, the second parameter is 0 when we are in existential mode and 1 when we are in universal mode, while the last parameter is the formula under consideration.

The definition of ϱ is as follows.

Definition 4.4 (Visser [3]) Let $\varrho(A)$ of a formula A be $\varrho(0, 0, A)$ and let

1. $\varrho(i, j, A) := 0$ if A is atomic,
2. $\varrho(i, j, B \wedge C) = \varrho(i, j, B \vee C) := \max\{\varrho(i, j, B), \varrho(i, j, C)\}$,
3. $\varrho(i, j, \neg B) := \varrho(1 - i, 1 - j, B)$,
4. $\varrho(0, j, B \rightarrow C) := \max\{\varrho(1, 1 - j, B), \varrho(0, j, C)\}$,
5. $\varrho(1, j, B \rightarrow C) := \max\{\varrho(0, j, B), \varrho(1, 1 - j, C)\}$,
6. $\varrho(i, 0, \exists x B) := \varrho(i, 0, B)$,
7. $\varrho(i, 1, \exists x B) := \varrho(i, 0, B) + 1$,
8. $\varrho(i, 0, \forall x B) := \varrho(i, 1, B) + 1$,
9. $\varrho(i, 1, \forall x B) := \varrho(i, 1, B)$.

The merit of the measure defined by Visser is that it treats negation and implication directly. Contrary to the measure defined by Zhang and the measure *aqf* defined in this note, Visser’s ϱ makes no distinctions for the propositional connectives as to whether they appear in existential or universal mode, that is, below an existential or a universal quantifier. Thus the measure ϱ assigns the same “depth of quantifierchanges” to the formulas

$$\forall x (\forall y P(x, y) \vee \forall z Q(x, z))$$

and

$$\forall x (\forall y P(x, y) \wedge \forall z Q(x, z)).$$

However, one can show that the complexity of cut elimination for the two (cut) formulas is not the same, that is, cut elimination for the second formula, which is purely \forall, \wedge , is simpler than cut elimination for the first formula $\forall x (\forall y P(x, y) \vee \forall z Q(x, z))$, which contains a disjunction.

Thus, although capturing the main ideas, namely, that cut elimination mainly depends on quantifier alternations, the measure as it is defined in [3], is not optimal in all cases to estimate the complexity of cut elimination.

References

- [1] Gerhardy, P., “Improved Complexity Analysis of Cut Elimination and Herbrand’s Theorem,” Master’s thesis, Aarhus, 2003. [165](#)
- [2] Gerhardy, P., “Refined complexity analysis of cut elimination,” pp. 212–25 in *Proceedings of the 17th International Workshop CSL 2003*, edited by M. Baaz and J. Makovsky, vol. 2803 of *LNCS*, Springer-Verlag, Berlin, 2003. [165](#), [166](#), [167](#), [168](#), [169](#)
- [3] Visser, A., “The unprovability of small inconsistency. A study of local and global interpretability,” *Archive for Mathematical Logic*, vol. 32 (1993), pp. 275–98. [Zbl 0795.03080](#). [MR 94c:03073](#). [165](#), [169](#), [170](#)
- [4] Zhang, W., “Cut elimination and automatic proof procedures,” *Theoretical Computer Science*, vol. 91 (1991), pp. 265–84. [Zbl 0745.68098](#). [MR 93a:03064](#). [165](#), [166](#)
- [5] Zhang, W., “Depth of proofs, depth of cut-formulas and complexity of cut formulas,” *Theoretical Computer Science*, vol. 129 (1994), pp. 193–206. [Zbl 0807.03038](#). [MR 95h:03126](#). [165](#), [167](#), [169](#)

Acknowledgments

I am grateful to Lev Beklemishev, Ulrich Kohlenbach, and Albert Visser for many useful discussions on the subject of this note.

BRICS
Department of Computer Science
University of Aarhus
IT-parken, Aabogade 34
DK-8200 Aarhus N
DENMARK
peegee@brics.dk