

Internal consistency for embedding complexity

Sy-David Friedman and Katherine Thompson

December 18, 2007

Abstract

In a previous paper with M. Džamonja, class forcings were given which fixed the complexity (a universality covering number) for certain types of structures of size λ together with the value of 2^λ for every regular λ . As part of a programme for examining when such global results can be true in an inner model, we build generics for these class forcings.¹

1 Introduction

The *internal consistency programme* was introduced in [6]. This programme aims to determine which consistent statements of set theory are in fact *internally* consistent, i.e., true in an inner model, assuming the existence of large cardinals. One of the main advantages of internal consistency is that this rules out consistent statements, such as the nonexistence of transitive set models of ZFC, which can only hold in universes which are incompatible with V .

There are many consistency results achieved using forcing which are not yet known to be obtainable via inner models, since internal consistency results are harder to achieve. To obtain any consistency result, one typically makes

¹This research was supported by the Austrian Science Fund (FWF), Grants P 16334-NO5 and P 16790-NO4.

use of a generic filter. To obtain consistency results using set-forcing, one may assume that the starting model is countable and therefore the existence of a generic is trivial. To achieve internal consistency, one needs an inner model with all the ordinals of the ambient universe, therefore one cannot restrict to a countable submodel.

The question then is how to build a generic, if one even exists. The methods used to do this require new techniques which are mathematically of independent interest. Since there are strong connections between work done in large cardinal theory and in internal consistency (see e.g. [6], [2]), these techniques can be applied in the context of other questions involving large cardinals (see e.g. [8]).

For more about this programme, including a discussion of some of the results that have been obtained within it, we refer the reader to [6].

A number of the early internal consistency results below $0^\#$ are discussed in Chapter 3 of [5]. These results use either reverse Easton or forward Easton forcing methods. An example of the latter is due to Jensen [1], who showed that in $L[0^\#]$ there is a real which is class-generic but not set-generic over L . All of these early results however only produce models of GCH.

The first example of an internal consistency result where GCH fails appears in [7]. In that paper, Easton's theorem ([4]) is examined in the internal consistency context, and it is shown that any (parameter-free) L -definable Easton function can be realised as the generalised continuum function $\kappa \mapsto 2^\kappa$ in an inner model of $L[0^\#]$. Such a result cannot be obtained using the forcing that Easton introduced, the Easton product. Instead, [7] introduces a reverse Easton iteration of Easton products, and then shows how to obtain a generic over L for this class-forcing using the Silver indiscernibles. A generic for the forcing up to and including i , denoted $G(\leq i)$, is constructed by induction on the Silver indiscernible i ; the main task is to choose these to cohere in the sense that $G(\leq j)$ contains the image of $G(\leq i)$ under the natural embedding $\pi_{ij} : L \rightarrow L$ obtained by “shifting” the class I of indiscernibles onto $I \setminus [i, j)$. The technique of “generic modification” is used in order to modify a given generic $G'(\leq j)$ to a generic $G(\leq j)$ with the desired coherence property.

Our work in the present paper extends that of [7] in two new directions. First, we show how to construct generics for forcings for which the technique

of generic modification cannot be used. Instead, we construct our generics directly so as to cohere. Second, we establish the first internal consistency result (section 5) that handles a reverse Easton iteration of forcings whose building blocks are not products, as in [7], but which are themselves iterations. This presents some significant new challenges that we meet using special dense sets of conditions and elementary submodels.

The context of the present paper is that of embedding complexity. However, the techniques we introduce here for the construction of generics have the potential of application to many other contexts. We discuss this point further in the final section.

In order to explain our results on embedding complexity, we begin with the basic universality definitions. An embedding (also called weak embedding) for ordered sets is an injective order-preserving map. A strong embedding is an embedding which also preserves incomparability. The range of a strong embedding is an isomorphic copy of the order in the domain. For graphs, an embedding is an injective function which preserves edges and a strong embedding is an injective function whose range is an induced subgraph.

Given a set $\mathcal{A}(\lambda)$ of structures each of size λ , a (*strong*) *universal model* for $\mathcal{A}(\lambda)$ is an element of $\mathcal{A}(\lambda)$ which (strongly) embeds all other structures in $\mathcal{A}(\lambda)$. If there does not exist a universal model for $\mathcal{A}(\lambda)$, then we consider its *complexity*, or the smallest size of a subfamily of $\mathcal{A}(\lambda)$ which embeds the rest. This subfamily of structures is called a *universal family*. All of these notions have weak and strong counterparts depending on the type of embedding used.

In this paper, λ, κ will be regular cardinals such that $\lambda \geq \kappa \geq \aleph_0$. Let $\mathcal{C}(\lambda, \kappa)$ be the set (of representatives under isomorphism) of all posets of size λ which omit κ -chains. Denote by $G(\lambda, \kappa)$ the set of all graphs of size λ which omit κ -cliques.

In the paper [3], two class forcing results were given. One says that the weak complexity for $\mathcal{C}(\lambda, \kappa)$ (or $G(\lambda, \kappa)$) can be 2^λ for any “reasonable” value of 2^λ and κ , and that all these values can be fixed simultaneously for all regular λ . Reasonable here means that 2^λ is limited only by the usual restrictions on such values and κ is as above. The other states that the strong complexity for $G(\lambda, \kappa)$ can be any regular cardinal between λ^+ and 2^λ for reasonable

values of 2^λ and κ , and that these values can all be fixed simultaneously for all regular λ . These results will be stated in the sections 2 and 3, thus, the paper can be considered self-contained.

In sections 2 and 3, we take the global product forcings given in [3] and modify them so that they are reverse Easton iterations. In both cases we are fixing the value of $F(\lambda) = 2^\lambda$ for all regular λ simultaneously. This could lead to problems as the values that F takes can skip over cardinals. However, there is a club class of cardinals λ such that $F(\theta) < \lambda$ for all $\theta < \lambda$. We will form our iteration over these closure points. This is important in order to build generics.

In sections 4 and 5, we build generics for the global iterations given in Sections 2 and 3.

In section 6, we isolate the special properties of our forcings which are required for our construction. As our forcings lack the homogeneity properties of Cohen forcing which were heavily used in [7], we instead rely on other distinctive properties of the forcing in order to directly build the generic. We also discuss other types of forcings which may meet these criteria.

We will use the convention that for $p, q \in P$, a forcing notion, $q \leq p$ indicates that q is stronger than p . For a forcing condition p , we denote by $\text{Dom}(p)$ the non-trivial domain, or support, of p .

2 Iterating high complexity

In the paper [3] a forcing notion was given for any regular uncountable λ which fixed the complexity of $\mathcal{C}(\lambda, \kappa(\lambda))$ to be maximal (that is, 2^λ) and fixed the value of $2^\lambda > \lambda^+$. A globalisation of that result was also given as an Easton product, which proved the following result:

Theorem 2.1. Assume $V = L$. To each uncountable regular cardinal λ associate a regular cardinal $\kappa(\lambda) \leq \lambda$ and a cardinal $F(\lambda)$ such that $\text{cf}(F(\lambda)) > \lambda$ and $\lambda < \theta$ implies $F(\lambda) \leq F(\theta)$. Assume that the functions κ, F are L -definable. Then there exists an L -definable, ZFC-preserving, cardinal preserving class forcing notion P such that in L^P the complexity of $\mathcal{C}(\lambda, \kappa(\lambda))$

is $F(\lambda) = 2^\lambda$ for each regular uncountable λ .

Remark 2.2. In the theorem above, the functions κ and F are L -definable with or without parameters. When building the generic in Section 4, it will be required that these functions are definable without parameters.

Namely, P was defined as a product forcing notion of $P(\lambda, \kappa(\lambda), F(\lambda))$ (defined below) with Easton support. Recall that Easton support means that the support is bounded at inaccessibles.

The forcing notion $P(\lambda, \kappa(\lambda), F(\lambda))$ was defined to be the $(< \lambda)$ -support product of $F(\lambda)$ copies of $Q = Q(\lambda, \kappa(\lambda))$. We will repeat the definition of Q below, but first a remark is needed about the terminology.

The definition of Q requires a set of bounded subsets of λ which we call low sets. These are used to restrict the growth of chains when extending sequences of posets without creating large antichains in the forcing. This method was introduced in [9] and also used in [3]. The exact definition is not required for this paper, but can be found in both of the papers above.

Let the elements of Q be $q = (\delta, X, \mathcal{A})$, where

1. $\delta < \lambda$ is an ordinal,
2. $X \subseteq \delta$ is a poset (X, \leq_X) which omits $\kappa(\lambda)$ -chains and in which $\alpha \leq_X \beta$ implies that $\alpha \leq \beta$ (as ordinals),
3. the conditions on \mathcal{A} are as follows:
 - (a) \mathcal{A} is a family of size $< \lambda$ of low subsets A of λ such that $A \subseteq \delta$ and $|A| < \kappa(\lambda)$,
 - (b) if $A \in \mathcal{A}$ is such that $\sup(A) \leq x < \delta$, then $A \not\leq_X x$.

The ordering of Q is as follows. If $q = (\delta, X, \mathcal{A})$ and $q' = (\delta', X', \mathcal{A}')$ are in Q , then $q' \leq_Q q$ (i.e. q' is stronger than q) if and only if $\delta' \geq \delta$, with X a subposet of X' such that $X = X' \cap \delta$ and $\mathcal{A} = \mathcal{A}' \cap [\delta]^{<\kappa(\lambda)}$.

If a forcing notion has the property that one can build decreasing sequences of size $< \lambda$ only controlling what happens at limit stages, we say that the forcing is weakly λ -closed. More precisely:

Definition 2.3. A forcing notion P is *weakly λ -closed*, if and only if there exists a function $w : P^{<\lambda} \rightarrow P$ such that for all limit $\tau < \lambda$ if $\langle p_\alpha : \alpha < \tau \rangle$ is a descending chain and satisfies the following:

$$\text{for all limit } \tau' < \tau \text{ we have } p_{\tau'} \leq w(\langle p_\alpha : \alpha < \tau' \rangle)$$

then the sequence $\langle p_\alpha : \alpha < \tau \rangle$ has a lower bound.

Note that weakly λ -closed falls in between the notions of λ -closed and λ -strategically closed. The forcing $P(\lambda, \kappa(\lambda), F(\lambda))$ was shown to be weakly λ -closed and satisfy the λ^+ -cc.

In order to build a generic for the global result Theorem 2.1, we must first have an iteration.

Lemma 2.4. We can find a class forcing P such that it satisfies the conditions of Theorem 2.1 and in addition, P is given as a reverse Easton iteration $\langle (P_\alpha, Q_\alpha) : \alpha \in \text{Ord} \rangle$ where $P_\alpha \Vdash Q_\alpha$ is weakly α -closed for each $\alpha \in \text{Ord}$.

Proof To form an iteration out of this, we must iterate over the closure points of F and use a product in between.

Let $\langle \lambda_\beta : \beta \in \text{Ord} \rangle$ be the class of closure points of the the F function. That is, for all ordinals β and for all $\theta < \lambda_\beta$, a regular cardinal, $F(\theta) < \lambda_\beta$. Note that each λ_β is a limit cardinal.

Define $\langle (P_\alpha, Q_\alpha) : \alpha \in \text{Ord} \rangle$ as follows. Let Q_α be the Easton support product of $P(\beta, \kappa(\beta), F(\beta))$ for $\beta \in [\lambda_\alpha, \lambda_{\alpha+1})$ a regular, uncountable cardinal. Let P be the reverse Easton iteration of the Q_α .

As in [3], Q_α preserves cofinalities and forces that $\mathcal{C}(\beta, \kappa(\beta))$ has complexity $F(\beta)$ for all $\beta \in [\lambda_\alpha, \lambda_{\alpha+1})$.

We must only check that at closure points λ_β , the complexity has not been affected by the forcing below λ_β . Since λ_β is always a limit cardinal, we must check that the complexity of $\mathcal{C}(\lambda, \kappa(\lambda))$ is $F(\lambda)$ in M^{P_λ} if $\lambda = \lambda_\beta$ is inaccessible or $\lambda = \lambda_\beta^+$ and λ_β is singular.

When λ_β is inaccessible, we will have two cases, depending on whether or not it is Mahlo. If λ_β is a Mahlo cardinal, then the forcing has the λ_β -cc and the result follows as in [3].

So assume that λ_β is not Mahlo. We will use the elementary submodels argument as in [3] to check that we have the correct complexity. In this case, we only have the λ_β^+ -cc. We must ensure that the forcing below λ_β does not introduce inconvenient embeddings which would lower the complexity at λ_β . To that end, we can anticipate a generic for the forcing below λ_β and run the complexity argument based on the information in the anticipated generic. As we have the λ_β^+ -cc, we can anticipate a maximal antichain in λ_β steps and we use elementary submodels to ensure that this happens in order type λ_β .

If λ_β is singular, then the forcing at λ_β is trivial. However, we must check the complexity at λ_β^+ . Here we have the λ_β^{++} -cc. The argument then proceeds as in the non-Mahlo inaccessible case, replacing λ_β with λ_β^+ . \square

3 Iterating low complexity

We will also make an iteration out of the globalisation of the low complexity result in [9]. As before, we have already proved in [3] that the forcing below exists as a product.

Theorem 3.1. Assume $V = L$. Let $\lambda, \kappa(\lambda), F$ be as in Theorem 2.1 and let $\nu(\lambda) \in [\lambda^+, F(\lambda))$ be L -definable. Then there exists an L -definable, ZFC-preserving, cardinal preserving class forcing notion P such that in L^P the strong complexity of $G(\lambda, \kappa(\lambda))$ is $\nu(\lambda)$ and $F(\lambda) = 2^\lambda$ for each regular uncountable λ .

Remark 3.2. In the theorem above, the functions κ, ν and F are L -definable with or without parameters. When building the generic in Section 4, it will be required that these functions are definable without parameters.

The general construction of this global iteration is as before, except at each regular cardinal λ , the forcing is an iteration of length $\nu(\lambda)$ which produces a member of a universal family at every successor stage. When necessary in order to distinguish the local iteration at a particular λ and the global iteration, the local iteration will have a superscript l .

Here, the global forcing P will be defined as an iterated forcing notion with Easton support. At each regular stage λ of this global iteration, we force with

$P(\lambda, F(\lambda), \kappa(\lambda), \nu(\lambda)) = \langle P_\alpha^l, Q_\alpha^l : 0 \leq \alpha < \nu(\lambda) \rangle$ which is a $(< \lambda)$ -support iteration of length $\nu(\lambda)$. At singular stages, the forcing will be trivial. For a particular regular λ , we describe the forcing $P(\lambda, F(\lambda), \kappa(\lambda), \nu(\lambda))$ below.

The first step of the iteration, Q_0^l , is used to add $F(\lambda)$ -many Cohen subsets to λ , using the standard λ^+ -cc and λ -closed forcing.

In each step $\alpha \geq 1$ of the iteration, we will add a graph of size λ which omits $\kappa(\lambda)$ -cliques that strongly embeds every member of $G(\lambda, \kappa(\lambda))$ in $M^{P_\alpha^l}$. We will modify the conditions for Q_α^l slightly from the version in [9] in order to make the construction of the generic easier.

Namely, let $Q_\alpha^l = (Q, \leq)$ for some ordinal α such that $1 \leq \alpha < \nu(\lambda)$. Fix some “canonical” enumeration $\{Z_\gamma : \gamma < F(\lambda)\}$ of all graphs of size λ which omit $\kappa(\lambda)$ -cliques which exist at this stage. Let $q \in Q$ if and only if $q = (\delta, X, \mathcal{A}, \mathcal{Z}, \Phi)$ with the following properties:

1. $\delta < \lambda$ is an ordinal,
2. $X \subseteq \delta$ is a graph which omits $\kappa(\lambda)$ -cliques,
3. $\mathcal{A} \subseteq [\delta]^{<\kappa(\lambda)}$ is a family of low subsets of δ such that $|\mathcal{A}| < \lambda$,
4. $\mathcal{Z} \subseteq F(\lambda)$ such that $|\mathcal{Z}| < \lambda$,
5. $\Phi : \mathcal{Z} \times \delta \rightarrow \delta$ is a function such that if $\gamma \in \mathcal{Z}$, then the mapping $x \mapsto \Phi(\gamma, x)$ is a strong embedding from $Z_\gamma \upharpoonright \delta$ into X ,
6. if $A \in \mathcal{A}$ is such that $\sup(A) \leq x < \delta$, then $A \times \{x\} \not\subseteq X$ (i.e. there exists an element of A which is not connected to x in X),
7. if $A \in \mathcal{A}$ and $\gamma \in \mathcal{Z}$, then $A \not\subseteq \Phi''(\{Z_\gamma\} \times \delta) := \{\Phi(Z_\gamma, \beta) : \beta < \delta\}$.

If $q = (\delta, X, \mathcal{A}, \mathcal{Z}, \Phi)$ and $q' = (\delta', X', \mathcal{A}', \mathcal{Z}', \Phi')$ are in Q , then $q' \leq_Q q$ (i.e. q' is stronger than q) if $\delta' \geq \delta$, with $X = X' \cap \delta$ as graphs, $\mathcal{A} = \mathcal{A}' \cap [\delta]^{<\kappa(\lambda)}$ and $\mathcal{Z} \subseteq \mathcal{Z}'$. Additionally, the following requirements must be met:

- (a) if $(Z_\gamma, x) \in \text{Dom}(\Phi)$ then $\Phi'(Z_\gamma, x) = \Phi(Z_\gamma, x)$,
- (b) if $\gamma \in \mathcal{Z}$ and $\delta \leq x < \delta'$ then $\Phi'(Z_\gamma, x) \geq \delta$,

(c) if $\gamma \neq \gamma' \in \mathcal{Z}$ and $\delta \leq x, y < \delta'$, then $\Phi'(Z_\gamma, x) \neq \Phi'(Z_{\gamma'}, y)$.

The intuition for the definition of conditionhood is as follows. The graph X generically becomes a model which is universal for graphs in $G(\lambda, \kappa(\lambda))$ which exist at that stage of the iteration. The set \mathcal{A} prevents large cliques from forming in X via condition 6. The graphs indexed by \mathcal{Z} are the “ground model” graphs which exist at that stage of the iteration whose restriction to δ embed into X . The functions Φ are the appropriate partial embeddings.

The modification that we made to the conditions was to take the actual graphs of size λ out of the condition and replace them by their indices. This changes the effective size of the conditions to be $< \lambda$.

Forming the global iteration for the forcing P works exactly as in Section 2. Namely, let $\langle \lambda_\beta : \beta \in \text{Ord} \rangle$ be the class of closure points of the the F function as before. Define $P = \langle (P_\alpha, Q_\alpha) : \alpha \in \text{Ord} \rangle$ as follows. Let Q_α be the Easton support product of $P(\beta, \kappa(\beta), F(\beta), \nu(\beta))$ for $\beta \in [\lambda_\alpha, \lambda_{\alpha+1})$ a regular, uncountable cardinal. Let P be the reverse Easton iteration of the Q_α . Using similar techniques as in the case of high complexity, P forces the complexity of $G(\lambda, \kappa(\lambda))$ to be $\nu(\lambda)$ for all regular λ .

4 Generic for high complexity

In this section, we will build a generic for the forcing P such that in M^P the complexity of $\mathcal{C}(\lambda, \kappa(\lambda))$ is $F(\lambda) = 2^\lambda$ for all regular λ . Namely, this is the forcing P as defined in Lemma 2.4 above.

Let $I = \{i_\alpha : \alpha \in \text{Ord}\}$ be the Silver indiscernibles for L in increasing order. We will make use of the following facts:

1. For all formulas φ , it is the case that $L \models \varphi(\alpha, \vec{i}) \Leftrightarrow \varphi(\alpha, \vec{j})$ whenever $\alpha < \min(\vec{i} \cup \vec{j})$ where $\vec{i}, \vec{j} \in I^{<\omega}$.
2. If t is a Skolem term in L and $t(j_1, \dots, j_k, j_{k+1}, \dots, j_n) < j_{k+1}$ where $j_1 < \dots < j_k < j_{k+1} < \dots < j_n$, then this term has the same value as $t(j_1, \dots, j_k, j'_{k+1}, \dots, j'_n)$ where $j'_i > j_k$ for $i > k$.

Since the second fact indicates that the parameters above j_k are irrelevant (when t takes values below j_{k+1}), we often write $t(j_1, \dots, j_k, \vec{\infty})$. When we say that $\vec{j} < \alpha$ for some vector j , we mean that all elements of \vec{j} are less than α .

There are \aleph_0 many Skolem terms, so we will enumerate them as $\{t_n : n < \omega\}$.

Theorem 4.1 ($0^\#$). Let $P_\infty = \langle P(\alpha) : \alpha \in \text{Ord} \rangle$ be the iteration with Easton support definable in L without parameters as in Lemma 2.4 above. Also assume that the functions κ and F (defined in Theorem 2.1) are L -definable without parameters. Then there exists a G such that G is P_∞ -generic over L .

Proof First note that it suffices to meet all set maximal antichains since there are no cofinal antichains in P_∞ . We will define $G(\leq i)$ generic for $P(\leq i)$ by induction on $i \in I$. Since the function F is definable without parameters, the indiscernibles are closure points of F .

The following properties of P were proved in the previous paper.

- Lemma 4.2.**
1. $P(> \lambda)$ is weakly λ^+ -closed for all regular λ .
 2. $P(< \lambda)$ has the λ -cc for Mahlo (and hence, indiscernible) λ .

Finding a generic for $P(\leq i_0)$ is trivial since $\mathcal{P}(P(\leq i_0))^L$ is countable.

If α is a limit ordinal then we want $G(\leq i_\alpha)$ to be the “direct limit” of $G(\leq i_\beta)$ for $\beta < \alpha$. In order to achieve the compatibility needed to make this generic, we will use the shift map below.

Let $\beta < \beta'$. We will define $\pi_{i_\beta, i_{\beta'}}$ as follows:

$$\pi_{i_\beta, i_{\beta'}}(i_\gamma) = \begin{cases} i_\gamma & \gamma < \beta \\ i_{\beta' + (\gamma - \beta)} & \gamma \geq \beta. \end{cases}$$

Shifting one indiscernible up to another extends uniquely to an elementary embedding from L into L . We will abuse notation and denote this extension in the same way.

Let $G(\leq i_\beta) \rightarrow G(\leq i_{\beta'})$ denote that $G(\leq i_\beta)$ embeds into $G(\leq i_{\beta'})$ in the sense that $\pi_{i_\beta, i_{\beta'}}[G(\leq i_\beta)] \subseteq G(\leq i_{\beta'})$. Using the shift map we want to find $G(\leq i_{\beta'})$ such that $G(\leq i_\beta) \rightarrow G(\leq i_{\beta'})$ and $G(i_{\beta'})$ is generic for $P(i_{\beta'})$.

Lemma 4.3. If $G(\leq i_\beta) \rightarrow G(\leq i_{\beta'})$ for all $\beta < \beta' < \alpha$ for α limit, then the direct limit $G(\leq i_\alpha) = \bigcup_{\beta < \alpha} \pi_{i_\beta, i_\alpha}[G(\leq i_\beta)]$ is generic for $P(\leq i_\alpha)$.

Proof Let $\Delta = t(\vec{i}, i_\alpha, \vec{\infty})$ be a maximal antichain in $P(\leq i_\alpha)$. Then $\bar{\Delta} = t(\vec{i}, i_\beta, \vec{\infty})$ is a maximal antichain in $P(\leq i_\beta)$. If $p \in G(\leq i_\beta) \cap \bar{\Delta}$ and $G(\leq i_\beta) \rightarrow G(\leq i_\alpha)$ then $\pi_{i_\beta, i_\alpha}(p) \in \Delta \cap G(\leq i_\alpha)$. \square

The successor case, $i_{\alpha+1}$ is the interesting one. We wish to find $G(\leq i_{\alpha+1})$ assuming that we have already built $G(\leq i_\alpha)$. We first note that $P(< i_{\alpha+1}) = P(\leq i_\alpha) * P(i_\alpha, i_{\alpha+1})$ where $P(i_\alpha, i_{\alpha+1})$ is the forcing in the interval $(i_\alpha, i_{\alpha+1})$. By the induction hypothesis, we have $G(\leq i_\alpha)$ which is $P(\leq i_\alpha)$ -generic. We can split our task up into finding $G(i_\alpha, i_{\alpha+1})$, generic for $P(i_\alpha, i_{\alpha+1})$ and then finding $G(i_{\alpha+1})$ which is $P(i_{\alpha+1})$ -generic.

We start by building $G(i_\alpha, i_{\alpha+1})$. First note that by Lemma 4.2(2), every antichain in $P(i_\alpha, i_{\alpha+1})$ has size $< i_{\alpha+1}$. We will show how to group all open dense subsets of $P(i_\alpha, i_{\alpha+1})$ in $L_{i_{\alpha+1}}[G(\leq i_\alpha)]$ into \aleph_0 many families each of size i_α . Then we can use the weak i_α^+ -closure of $P(i_\alpha, i_{\alpha+1})$ to handle maximal antichains in each family.

Lemma 4.4. Let D be the collection of all open dense sets in $P(i_\alpha, i_{\alpha+1})$ which belong to $L_{i_{\alpha+1}}[G(\leq i_\alpha)]$. Then $D = \bigcup_{n < \omega} D_n$ such that each $D_n \in L_{i_{\alpha+1}}[G(\leq i_\alpha)]$ and $|D_n| = i_\alpha$.

Proof For every $x \in L_{i_{\alpha+1}}[G(\leq i_\alpha)]$, there exists an n such that $x = t_n(\vec{j}, \vec{k})$ for $\vec{j} < i_\alpha$ and $\vec{k} \geq i_{\alpha+1}$. As we are only considering subsets of $i_{\alpha+1}$, we know that \vec{k} is irrelevant to the value of t_n . We will arrange all elements of $L_{i_{\alpha+1}}[G(\leq i_\alpha)]$ according to the enumeration of the Skolem terms for which they take values.

Let D_n be the set of all open dense subsets $t_n^{L[G(\leq i_\alpha)]}(\vec{\beta}, \vec{\infty})$ of $P(i_\alpha, i_{\alpha+1})$ such that $\vec{\beta} \leq i_\alpha$ is a finite set of ordinals, $t_n^{L[G(\leq i_\alpha)]}(\vec{\beta}, \vec{\infty}) \in L_{i_{\alpha+1}}[G(\leq i_\alpha)]$ and $\vec{\infty}$ is a finite set of fixed indiscernibles of the appropriate length. L cannot see indiscernibles, so we take all ordinal vectors instead which will include the indiscernibles.

So, for each n , $D_n \in L[G(\leq i_\alpha)]$. Now $|D_n|^{L[G(i_\alpha)]} = i_\alpha$ and there are only countably many of them. \square

Thus, the sequence of the D_n 's is not in $L[G(\leq i_\alpha)]$, but the individual pieces are. We can also show that the shift of a D_n will also be in $L[G(\leq i_\alpha)]$.

Claim 4.5. 1. If $X \in L$ and $|X| \leq i_{\beta'}$ then $\text{Ran}(\pi_{i_\beta, i_{\beta'}}) \cap X \in L$.

2. If $X \in L$ and $|X| \leq i_{\beta'}$ then $\pi_{i_\beta, i_{\beta'}} \restriction X \in L$.

Proof The proof of (1) breaks down into two cases. In the first case, we have $X \in \text{Ran}(\pi_{i_\beta, i_{\beta'}})$. Let $s : X \rightarrow i_{\beta'}$ be a bijection which is in L such that $s \in \text{Ran}(\pi_{i_\beta, i_{\beta'}})$. Therefore, $\text{Ran}(\pi_{i_\beta, i_{\beta'}}) \cap X = s^{-1}[\text{Ran}(\pi_{i_\beta, i_{\beta'}}) \cap i_{\beta'}]$. Since the critical point of $\pi_{i_\beta, i_{\beta'}}$ is i_β , this is equal to $s^{-1}[i_\beta]$, which is in L .

If $X \notin \text{Ran}(\pi_{i_\beta, i_{\beta'}})$ we can show that there exists $Y \in L$ with $Y \in \text{Ran}(\pi_{i_\beta, i_{\beta'}})$ and $X \subseteq Y$.

First note that $\text{Ran}(\pi_{i_\beta, i_{\beta'}}) = \{t(\vec{\alpha}, i_{\beta'}, \vec{k}) : \vec{\alpha} < i_\beta, \vec{k} > i_{\beta'}\}$. Also there is some Skolem term t such that $X = t(\vec{\gamma}, i_\beta, i_{\beta'}, \vec{l})$ for some $\vec{\gamma} < i_\beta$ and $\vec{l} > i_{\beta'}$. Let

$$Y = \bigcup \{t(\vec{\alpha}, \delta, i_{\beta'}, \vec{j}) : \delta \in (\max(\vec{\alpha}), i_{\beta'}), \delta \in \text{Ord}, \\ |t(\vec{\alpha}, \delta, i_{\beta'}, \vec{j})| = i_{\beta'}\}$$

One can see that $Y \in L$ as all parameters are either ordinals or fixed. It is also the case that $Y \supseteq \text{Ran}(\pi_{i_\beta, i_{\beta'}}) \cap X$. Thus,

$$\text{Ran}(\pi_{i_\beta, i_{\beta'}}) \cap X = (\text{Ran}(\pi_{i_\beta, i_{\beta'}}) \cap Y) \cap X \in L.$$

To prove (2), let $f : i_{\beta'} \rightarrow X$ be a bijection in L . For all $x \in X$, we have

$$\begin{aligned} \pi_{i_\beta, i_{\beta'}}(x) &= \pi_{i_\beta, i_{\beta'}}(f(f^{-1}(x))) \\ &= \pi_{i_\beta, i_{\beta'}}(f)\pi_{i_\beta, i_{\beta'}}(f^{-1}(x)) \\ &= \pi_{i_\beta, i_{\beta'}}(f)(f^{-1}(x)). \end{aligned}$$

The conclusion is clear as f and $f^{-1}(x)$ are in L . \square

We will build $G(i_\alpha, i_{\alpha+1})$ by induction on n in the enumeration of the Skolem terms. By the weak i_α^+ -closure of $P(i_\alpha, i_{\alpha+1})$ we can find $p_0 \in P(i_\alpha, i_{\alpha+1})$ such that p_0 meets every dense set in D_0 . This can be done since there can only be i_α many of them. Assume that we have found $\langle p_m : m < n \rangle$ such that p_m is compatible with all $p_{m'}$ where $m' < m$ and meets all dense sets in D_m . Use the weak closure again to find p_n such that p_n is compatible with all p_m where $m < n$ and meets all dense sets in D_n . Let $G(i_\alpha, i_{\alpha+1}) = \{p_n : n < \omega\}$.

Now we must find $G(i_{\alpha+1})$, a generic for $P(i_{\alpha+1})$. We do this by first shifting the conditions in $G(i_\alpha)$ up to $i_{\alpha+1}$.

We will show that if we shift the conditions in the generic up to $i_{\alpha+1}$, they remain conditions in $P_{i_{\alpha+1}}$. So we need to see that each poset that we build generically at i_α omits $\kappa(i_{\alpha+1})$ -chains. Because the forcing is L -definable without parameters, $\kappa(i_\alpha)$ is either i_α or some $\theta < i_0$, the least indiscernible. In both cases, we have $\kappa(i_\alpha) \leq \kappa(i_{\alpha+1})$. Consider $\pi_{i_\alpha, i_{\alpha+1}}[G(i_\alpha)]$. This is a set of posets of size i_α which omit chains of size $\pi_{i_\alpha, i_{\alpha+1}}(\kappa(i_\alpha)) = \kappa(i_{\alpha+1})$.

Let $\pi_{i_\alpha, i_{\alpha+1}}[G(i_\alpha)] = \{p_\beta : \beta < F(i_\alpha)\}$. Since we are taking the pointwise image of the generic at i_α and the conditions have size $< i_\alpha$, this remains the case in the image. Our goal is to extend this image to a generic for $P(i_{\alpha+1})$.

The idea is to break up the product at i_α into ω -many pieces each of size i_α , call them M_n^α for $n < \omega$. The M_n^α 's will be increasing, that is, $M_n^\alpha \subseteq M_{n+1}^\alpha$. The sequence of pieces is not in $L[G(\leq i_\alpha)]$, but each piece is. For each n we will form the image of the generic $G(i_\alpha)$ restricted to M_n^α and then extend this to form a generic for the corresponding piece of the forcing $P(i_{\alpha+1})$.

For each $\alpha \in \text{Ord}$ and $n < \omega$ let M_n^α be the Σ_1 Hull($i_\alpha \cup \{i_\alpha, i_{\alpha+1}, \dots, i_{\alpha+n}\}$). These are Σ_1 -elementary submodels of L and are i_α -closed.

For $n = 1$, let $\{p : p \in M_1^\alpha \cap G(i_\alpha)\} =: A_1$. Then $\pi_{i_\alpha, i_{\alpha+1}}[A_1]$ is a set of conditions of $P(i_{\alpha+1})$ and there are only i_α -many of them.

Let $p_1^* = \bigcup \pi_{i_\alpha, i_{\alpha+1}}[A_1]$. To see that this is a condition, we must show that for each component in the support, no $\kappa_{\alpha+1}$ sequences have been added. One may write $p_1^* = \bigcup \{\pi_{i_\alpha, i_{\alpha+1}}(\langle p(\beta) : \beta \in \text{Dom}(p) \rangle) : p \in A_1\}$ which means that $p_1^*(\beta) = \bigcup \{\pi_{i_\alpha, i_{\alpha+1}}(p)(\beta) : p \in A_1\}$ for each $\beta \in \text{Dom}(p_1^*)$. We need to see

that $X_{p_1^*(\beta)}$ omits $\kappa(i_{\alpha+1})$ -chains. This has been proved by [3, Lemma 3.9], where it was shown that the generic poset $\bigcup\{X : (\delta, X, \mathcal{A}) \in G_Q \text{ generic for } Q(i_\alpha, \kappa(i_\alpha))\}$ (using the notation for the forcing from Section 2) has no $\kappa(i_\alpha)$ -chain.

The filtration $\{M_n^\alpha : n < \omega\}$ can be shifted up to a filtration of $P(i_{\alpha+1})$, namely $M_n^{\alpha+1} = \pi_{i_\alpha, i_{\alpha+1}}(M_n^\alpha)$. Note that each of the $M_n^{\alpha+1}$ has size $i_{\alpha+1}$, but there are still only ω -many of them. Thus, if i_α is a limit indiscernible, M_1^α is the direct limit of $\pi_{i_\beta, i_\alpha}(M_1^\beta)$ for $\beta < \alpha$.

Since $M_1^{\alpha+1}$ is $i_{\alpha+1}^+$ -closed and p_1^* is a lower bound for $\pi_{i_\alpha, i_{\alpha+1}}[A_1]$, we have $p_1^* \in M_1^{\alpha+1}$. We may extend below p_1^* to form a generic for $M_1^{\alpha+1}$ by breaking up the dense sets that we must meet into ω -many blocks as before. We call this restricted generic $G^1(i_{\alpha+1})$.

The case $n = 2$ is essentially the general case. Again form $A_2 := \{p_\beta \in G(i_\alpha) \cap M_2^\alpha : \beta < i_\alpha\}$. Extend below $\pi_{i_\alpha, i_{\alpha+1}}[A_2]$ to form $p_2^* \in M_2^{\alpha+1}$. Here we need to check that p_2^* is compatible with $G^1(i_{\alpha+1})$.

One can decompose $p_2^* = p_2^* \restriction M_1^{\alpha+1} \cup p_2^* \restriction (M_2^{\alpha+1} - M_1^{\alpha+1})$. Note that $p_2^* \restriction M_1^{\alpha+1} \in M_1^{\alpha+1}$ by the $i_{\alpha+1}$ -closure of $M_1^{\alpha+1}$. If there were a compatibility issue with $p_2^* \restriction M_1^{\alpha+1} = p_1^* \geq G^1(i_{\alpha+1})$, then this would have already been a problem in $M_1^{\alpha+1}$ which would contradict the genericity of $G^1(i_{\alpha+1})$. However, $p_2^* \restriction (M_2^{\alpha+1} - M_1^{\alpha+1})$ is irrelevant to $G^1(i_{\alpha+1})$, that is, the conditions of $G^1(i_{\alpha+1})$ are not defined in this domain.

Now, as p_2^* is compatible with $G^1(i_{\alpha+1})$, we may extend below p_2^* to form a generic $G^2(i_{\alpha+1})$ for $M_2^{\alpha+1}$ such that $G^1(i_{\alpha+1}) \subseteq G^2(i_{\alpha+1})$.

Continue this process for all $n < \omega$. Let $G(i_{\alpha+1}) = \bigcup_{n < \omega} G^n(i_{\alpha+1})$. We would like to show that $G(i_{\alpha+1})$ is generic for $P(i_{\alpha+1})$. Recall that $P(i_{\alpha+1})$ has the $i_{\alpha+1}^+$ -cc so any antichain has size at most $i_{\alpha+1}$. Any maximal antichain must be contained in $M_n^{\alpha+1}$ for some $n < \omega$, since each $M_n^{\alpha+1}$ is $i_{\alpha+1}$ -closed and thus intersects $G^n(i_{\alpha+1})$. \square

5 Generic for low complexity

Here, the underlying forcing for each regular cardinal is an iteration and we have globally iterated products of these iterations. Forming a generic for this global iteration will be similar to the construction for the high complexity case, but more complicated. In this case, we can find a filtration $\{M_n^i : n < \omega\}$ for the local forcing at i , but the restriction of the local forcing at i to these M_n^i is not simply a regular subforcing.

Theorem 5.1 ($0^\#$). Assume that $P_\infty = \langle P(\alpha) : \alpha \in \text{Ord} \rangle$ is the iteration with Easton support definable in L without parameters as in Section 3 above. Then there exists G which is P_∞ -generic over L .

Proof As before, we will define $G(\leq i)$ generic for $P(\leq i)$ by induction on $i \in I$. Let $i < j$ be adjacent indiscernibles in I . The only difference in the proof from the previous one will be the case of building a generic for $G(j)$ assuming that we have built $G(< j)$. Again, we will build $G(j)$ such that $\pi_{i,j}[G(i)] \subseteq G(j)$. (Note that, as before, we abuse the notation $\pi_{i,j}$ to mean the canonical extension of the indiscernible shift to the model $L[G(< i)]$.)

Define $\{M_n^i : n < \omega\}$ and $\{M_n^j : n < \omega\}$, filtrations at i and j respectively, in a canonical way as before such that $\pi_{i,j}(M_n^i) = M_n^j$ for each n . That is, let M_n^i be the Σ_1 Hull of i together with the next n indiscernibles. Recall that each M_n^i is in $L[G(\leq i)]$, but the sequence $\langle M_n^i : n < \omega \rangle$ is not.

As before, we would like to define the restriction of a condition to one of these models. Before we can do that, we will see that we can extend any condition such that all of its components have ground model names.

For each $p \in P(i)$, we can write $p = \langle p(\alpha) : \alpha \in \text{Dom}(p) \rangle$. We say that p is *self-determined* if it has the property that for every α in the support of p , $p \restriction \alpha$ forces $p(\alpha) = (\delta, X, \mathcal{A}, \mathcal{Z}, \Phi)$ as well as all $Z_\gamma \restriction \delta$, for $\gamma \in \mathcal{Z}$ to equal a name in the ground model $L[G(< i)]$.

It is dense for conditions in $P(i)$ to be self-determined since $p(\alpha)$ and $Z_\gamma \restriction \delta$ for γ mentioned in $p(\alpha)$ all have size $< i$ and therefore, we may extend any condition to be self-determined using the weak i -closure. Therefore, we will concentrate on conditions which have this property.

Let $P^*(i)$ denote the set of conditions in $P(i)$ which are self-determined. We will define the restriction of $p \in P^*(i)$ to the model M_n^i which we will denote as $p \restriction M_n^i$. For $p \in P^*(i)$ let $p \restriction M_n^i = \{p(\alpha) \restriction M_n^i : \alpha \in \text{Dom}(p) \cap M_n^i\}$ where $p(\alpha) \restriction M_n^i$ is the obvious restriction of $p(\alpha)$ to the ordinals of M_n^i . Note that this restriction is an element of M_n^i as M_n^i is i -closed. We must see that this is still a condition and that extensions of it inside M_n^i are compatible with p .

Claim 5.2. For all self-determined $p \in P(i)$, all $n < \omega$ and $\alpha \in \text{Dom}(p) \restriction M_n^i$, we have that

- $(p \restriction \alpha) \restriction M_n^i$ is a condition and
- if $q \in M_n^i$ is such that q extends $(p \restriction \alpha) \restriction M_n^i$, then $p \restriction \alpha$ and q are compatible.

Proof Let $\langle \alpha_k : k < \text{ordertype}(\text{Dom}(p) \cap M_n^i) \rangle$ be the enumeration of $\text{Dom}(p) \cap M_n^i$ in increasing order. We will prove the claim by induction on k , first at successor stages and then at limit k .

Assume that p is self-determined, that $(p \restriction \alpha_k) \restriction M_n^i$ is a condition and if $q \in M_n^i$ is such that q extends $(p \restriction \alpha_k) \restriction M_n^i$, then q is compatible with $p \restriction \alpha_k$. We will first show that $(p \restriction \alpha_{k+1}) \restriction M_n^i$ is a condition.

The only reason that $(p \restriction \alpha_{k+1}) \restriction M_n^i$ could fail to be a condition is if it fails to force $p(\alpha_k) \restriction M_n^i$ to be a condition. So extend $(p \restriction \alpha_k) \restriction M_n^i$ to a condition q which forces $p(\alpha_k) \restriction M_n^i$ to not be a condition. We may assume that q belongs to M_n^i . However, by the induction hypothesis, q is compatible with $p \restriction \alpha_k$ which forces $p(\alpha_k)$ to be a condition, contradiction.

To show the second part of the claim for α_{k+1} , suppose that $(p \restriction \alpha_{k+1}) \restriction M_n^i$ is extended to a condition $q \in M_n^i$. We will show that there exists a condition r which extends both q and $p \restriction \alpha_{k+1}$. Let $\text{Dom}(r) = \text{Dom}(p \restriction \alpha_{k+1}) \cup \text{Dom}(q)$. First we will consider $\alpha \in \text{Dom}(p \restriction \alpha_{k+1}) \cap \text{Dom}(q)$. Since $\text{Dom}(p \restriction \alpha_{k+1}) \cap \text{Dom}(q) = \text{Dom}((p \restriction \alpha_{k+1}) \restriction M_n^i)$, we know that $\delta_{q(\alpha)} \geq \delta_{p(\alpha)}$. We can take $r(\alpha) = q(\alpha) \cup p(\alpha)$ where

1. $\mathcal{A}_{r(\alpha)} = \mathcal{A}_{q(\alpha)} \cup \mathcal{A}_{p(\alpha)}$,
2. $\mathcal{Z}_{r(\alpha)} = \mathcal{Z}_{q(\alpha)} \cup \mathcal{Z}_{p(\alpha)}$,

3. $\delta_{r(\alpha)}$ is a large enough ordinal $< i$ such that ordinals in $[\delta_{q(\alpha)}, \delta_{r(\alpha)})$ can be written as the disjoint union of $|\mathcal{Z}_{r(\alpha)}|$ -many copies of $[\delta_{p(\alpha)}, \delta_{r(\alpha)})$,
4. $\Phi_{r(\alpha)}$ extends $\Phi_{q(\alpha)} \cup \Phi_{p(\alpha)} \restriction (\text{Dom}(\Phi_{p(\alpha)}) - M_n^i)$ such that for all $\gamma \in \mathcal{Z}_{q(\alpha)}$, $\Phi_{r(\alpha)}$ maps $Z_\gamma \restriction [\delta_{q(\alpha)}, \delta_{r(\alpha)})$ into one of the copies of $[\delta_{p(\alpha)}, \delta_{r(\alpha)})$ above $\delta_{q(\alpha)}$ and for all $\gamma \in \mathcal{Z}_{p(\alpha)}$, $\Phi_{r(\alpha)}$ maps $Z_\gamma \restriction [\delta_{p(\alpha)}, \delta_{r(\alpha)})$ into one of the copies of $[\delta_{p(\alpha)}, \delta_{r(\alpha)})$ above $\delta_{q(\alpha)}$,
5. $X_{r(\alpha)} = X_{q(\alpha)} \cup [\delta_{q(\alpha)}, \delta_{r(\alpha)})$. The extra relations on $X_{r(\alpha)}$ connect the images of $Z_\gamma \restriction \delta_{p(\alpha)}$ with those of $Z_\gamma \restriction [\delta_{p(\alpha)}, \delta_{r(\alpha)})$ in the same way as Z_γ for all $\gamma \in \mathcal{Z}_{p(\alpha)}$, and likewise for $\gamma \in \mathcal{Z}_{q(\alpha)}$.

If $\alpha \in \text{Dom}(r) \setminus \text{Dom}(p \restriction \alpha_{k+1})$, then let $r(\alpha) = q(\alpha)$ and if $\alpha \in \text{Dom}(r) \setminus \text{Dom}(q)$, then let $r(\alpha) = p(\alpha)$.

Since the components inside and outside the model M_n^i are disjoint and $X_q \restriction \delta_p = X_p$, the unions will cause no conflict. The rest of the argument that this is a condition follows from [9, proof of Theorem 4, Claims 2 and 3].

If k is a limit ordinal, then assume for all $l < k$, we have $(p \restriction \alpha_l) \restriction M_n^i$ is a condition and if $q \in M_n^i$ is such that q extends $(p \restriction \alpha_l) \restriction M_n^i$, then q is compatible with $p \restriction \alpha_l$. To see that $(p \restriction \alpha_k) \restriction M_n^i$ is a condition, we only need to note that the restriction operations are canonical, that is, for $l < m < k$ we have $((p \restriction \alpha_m) \restriction M_n^i) \restriction \alpha_l = (p \restriction \alpha_l) \restriction M_n^i$.

Finally, we need to show that if q extends $(p \restriction \alpha_k) \restriction M_n^i$, then q is compatible with $p \restriction \alpha_k$. We may construct r extending $(p \restriction \alpha_k) \restriction M_n^i$ and q as before letting $\text{Dom}(r) = \bigcup_{l < k} \text{Dom}(p \restriction \alpha_l) \cup \text{Dom}(q)$. \square

Claim 5.3. If $p \in P^*(i)$ and $q \in M_n^i$ such that q extends $p \restriction M_n^i$, then p and q are compatible.

Proof By Claim 5.2, we know that $p \restriction M_n^i$ is a condition. Construct r extending p and q exactly as in Claim 5.2, letting $\text{Dom}(r) = \text{Dom}(p) \cup \text{Dom}(q)$. \square

Let $G^*(i) = G(i) \cap P^*(i)$. We have sufficiently prepared the conditions in $G^*(i) \restriction M_0^i$ in order to shift them up to $P(j)$. Let $A_0 = \{\pi_{i,j}(p \restriction M_0^i) : p \in G^*(i) \cap M_0^i\}$.

Since each condition p has size $< i$, this is also true of each condition in A_0 . We know that $|A_0| \leq i$ since $|M_0^i| = i$. We may take the union of the conditions in A_0 , call this p_0^0 . By [9, proof of Theorem 4, Claim 5], this is a condition in $P(j)$. Using the weak j -closure of $P(j)$, we may extend p_0^0 to $p_0^1 \in P(j) \cap M_0^j$ hitting $< j$ -many dense sets in M_0^j .

We want to continue this at each M_n^j for $n < \omega$. We will do this by constructing the master condition at M_n^j and for each $m \leq n$ we will extend the conditions built before to hit even more (but less than j many) dense sets in M_m^j . Since in $L[0^\#]$ we have $\text{cf}(j) = \omega$, at the end of the construction for each m we will have a condition which hits all dense sets in M_m^j .

After step n we formed p_m^{n-m} for $m \leq n$ as above. Extend each p_m^{n-m} to $p_m^{(n-m)+1}$, hitting $< j$ -many dense sets in M_m^j . Form p_{n+1}^0 as the union of conditions in $A_{n+1} = \{\pi_{i,j}(p \restriction M_{n+1}^i) : p \in G^*(i) \cap M_{n+1}^i\}$. We know that p_{n+1}^0 is compatible with $p_m^{(n-m)+1}$ by Claim 5.3. Extend p_{n+1}^0 to $p_{n+1}^1 \leq p_m^{(n-m)+1}$ for all $m \leq n$ such that p_{n+1}^1 hits $< j$ -many dense sets in M_{n+1}^j .

The generic $G(j)$ will then be $\{p \in P(j) : p \geq p_n^m : \text{for some } m, n < \omega\}$.

□

6 Conclusion

As remarked in the introduction, in the inductive process of building generics for reverse Easton class forcings, it is often not possible to simply modify any generic to cohere with what was built before. The results in sections 4 and 5 give techniques for finding generics for reverse Easton iterations even when this modification is not possible.

To continue with the programme of deciding when such forcings can have generics, the question remains: what intrinsic properties of the given forcings are used to allow these techniques to work? Below we isolate the main attributes of a global forcing which are sufficient to use the techniques of sections 4 and 5.

Theorem 6.1. If $P_\infty = \langle P(\alpha) : \alpha \in \text{Ord} \rangle$ is a class forcing with Easton

support which is L -definable without parameters and the following properties hold, then there exist a generic for P_∞ over L .

1. For all $\alpha \in \text{Ord}$ we have $P(\alpha)$ has the α -weak closure.
2. For $i \in I$ let $G(i)$ be generic for $P(i)$ and let $\langle M_n^i : n < \omega \rangle$ be the sequence of elementary submodels of $L[G(< i)]$ defined by $M_n^i = \Sigma_1 \text{Hull}(i \cup \vec{\alpha}_n)$ where $\vec{\alpha}_n$ are the first n indiscernibles greater than i . (Each M_n^i is i -closed and has size i .) For an indiscernible $j > i$, let $\langle M_n^j : n < \omega \rangle$ be the corresponding sequence of elementary submodels of $L[G(< j)]$. Note that $\pi_{i,j}(M_n^i) = M_n^j$.

For any P_∞ -generic G there are sequences $\langle D_\alpha : \alpha \in \text{Ord} \rangle$, $\langle \upharpoonright_\alpha : \alpha \in \text{Ord} \rangle$, $\langle mc_\alpha : \alpha \in \text{Ord} \rangle$ where D_α , \upharpoonright_α and mc_α are definable in $L[G(< \alpha)]$ uniformly (and independently of G) such that for each $i \in I$:

- (a) D_i is dense in $P(i)$.
- (b) For $p \in D_i$ and M an i -closed elementary submodel of $L[G(< i)]$ of size i , $p \upharpoonright_i M$ is a condition in $P(i)$ such that for all q extending $p \upharpoonright_i M$, if $q \in M$ then q is compatible with p .
- (c) For $i < j$, $i, j \in I$ and for each $n < \omega$, $mc(\pi_{i,j}[G(i) \cap M_n^i])$ is a lower bound for the conditions in $\pi_{i,j}[G(i) \cap M_n^i]$; moreover, for any $n_0 < n$, $mc(\pi_{i,j}[G(i) \cap M_n^i]) \upharpoonright M_{n_0}^j = mc(\pi_{i,j}[G(i) \cap M_{n_0}^i])$.

Note that in the proofs in sections 4 and 5, property (1) was used to build generics between indiscernibles and property (2) was used to construct a generic at a successor indiscernible stage which coheres with the generics at previous indiscernible stages.

If we restrict our attention to a natural class of product forcings, we can get a result which is more easily verifiable.

Definition 6.2. A product forcing $P(\alpha)$ is *internally small* iff the support of P has size $< \alpha$ and each component of the product is a forcing which has size α .

Note that we do not say that $P(\alpha)$ is simply “small” as this would imply that the length of the product is also restricted. We do not require this.

Theorem 6.3. If $P_\infty = \langle P(\alpha) : \alpha \in \text{Ord} \rangle$ is a class forcing with Easton support which is L -definable without parameters and the following properties hold, then there exist a generic for P_∞ over L .

1. $P(\alpha)$ is an internally small product forcing.
2. For all $\alpha \in \text{Ord}$ we have $P(\alpha)$ has α -closure.

Note that here we require full α -closure whereas the product forcing notion introduced in section 2 only has weak closure. In general weak closure is not enough to shift up conditions in the generic restricted to a model from an indiscernible i to a greater indiscernible j and get a master condition as in 2(c) of Theorem 6.1. In particular, the lower bounds for the sequences in the generic which satisfy a weak closure operation may not be in the generic. In section 4, we do get such a master condition, but this requires a proof which is particular to our case.

References

- [1] A. Beller, R. Jensen, and P. Welch. *Coding the universe*, volume 47 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge-New York, 1982.
- [2] N. Dobrinen and S. Friedman. Internal consistency and co-stationarity of the ground model. accepted in the *Journal of Symbolic Logic*. <http://www.logic.univie.ac.at/~sdf/papers/>.
- [3] M. Džamonja, S. Friedman, and K. Thompson. Global complexity results. In A. Andretta, editor, *Set Theory: Recent Trends and Applications*, volume 17 of *Quaderni di Matematica*, pages 25–45. Seconda Università di Napoli, 2007.
- [4] W.B. Easton. Powers of regular cardinals. *Annals of Mathematical Logic*, 1, 1970.

- [5] S. Friedman. *Fine structure and class forcing*. de Gruyter, 2000.
- [6] S. Friedman. Internal consistency and the inner model hypothesis. *Bulletin of Symbolic Logic*, 12(4):591–600, 2006.
- [7] S. Friedman and P. Ondrejovič. Internal consistency of Easton’s theorem. submitted. <http://www.logic.univie.ac.at/~sdf/papers/>.
- [8] S. Friedman and K. Thompson. Perfect trees and generic modification. submitted. <http://www.logic.univie.ac.at/~thompson/pubs.html>.
- [9] P. Komjáth and S. Shelah. Universal graphs without large cliques. *Journal of Combinatorial Theory*, 63(1):125–135, 1995.

Sy-David Friedman
sdf@logic.univie.ac.at
Kurt Gödel Research Center for Mathematical Logic
University of Vienna
Austria

Katherine Thompson
aleph_nought@yahoo.com
Kurt Gödel Research Center for Mathematical Logic
University of Vienna
Austria