

THE PROOF-THEORETIC ANALYSIS OF TRANSFINITELY ITERATED QUASI LEAST FIXED POINTS

DIETER PROBST

Abstract. The starting point of this article is an old question asked by Feferman in his paper on Hancock's conjecture [6] about the strength of ID_1^* . This theory is obtained from the well-known theory ID_1 by restricting fixed point induction to formulas that contain fixed point constants only positively. The techniques used to perform the proof-theoretic analysis of ID_1^* also permit to analyze its transinitely iterated variants ID_α^* . Thus, we eventually know that $|ID_\alpha| = |ID_\alpha^*|$.

§1. Introduction. The theories ID_α of iterated inductive definitions formalize hierarchies of least (definable) fixed points. In the past years, these theories have been exhaustively studied and their proof-theoretic analysis has been carried out a long time ago, (cf. Buchholz et al. [3]). Also their metapredicative relatives \widehat{ID}_α , that speak about hierarchies of (not necessary least) fixed points are well understood by now. The proof-theoretic ordinal of \widehat{ID}_1 is due to Aczel [1], who used a recursion theoretic argument, nowadays known as Aczel's trick, to embed \widehat{ID}_1 into Σ_1^1 -AC. The theories \widehat{ID}_n of n -times iterated inductive definitions have been analyzed by Feferman in connection with Hancock's conjecture in [6]. The proof-theoretic analysis of \widehat{ID}_α has been carried out in all details by Jäger, Kahle, Setzer and Strahm [9].

Some problems however, have remained unsolved: In the theories \widehat{ID}_α , induction on fixed points is dropped completely. It is natural to study theories, where fixed point induction is only restricted. Kreisel pointed out in [11], that "an inductive definition tells you what is *in* $P^{\mathcal{A}}$ not what is *not in* $P^{\mathcal{A}}$ ". As mentioned in Feferman [6], this motivated to consider restricted versions of ID_1 such as ID_1^* , a theory credited to H. Friedman where the scheme for proof by induction on fixed points is restricted to formulas that contain fixed point constants only positively. The question for a sharp upper bound is raised loc. cit. No answer to this question has yet been published, although partial results have been attained: If the fixed point axioms of ID_1^* are restricted to so-called accessibility inductive definitions, then the resulting theory $ID_1^*(\mathcal{AEC})$ can be embedded in Σ_1^1 -DC as sketched by Feferman in [6]. There, it is also stated that Friedman [8] introduced the theory ID_1^* and showed that its ordinal is bounded by α_1 , where $\alpha_0 := \varepsilon_0$ and $\alpha_{n+1} := \varphi_{\alpha_n}0$. Further, upper bounds for

Received June 15, 2004.

Key words and phrases. Fixed points; Iteration; Pseudo-hierarchies.

The author is supported by the Swiss National Science Foundation.

© 2006, Association for Symbolic Logic
0022-4812/06/7103-0001/\$3.60

the theories ID_n^* are computed by Cantini [4]. Thereby the two common ordinal measures for a theory T of inductive definitions are considered: The proof-theoretic ordinal $|T|$, i.e., the least ordinal α such that for no primitive recursive well-ordering \prec of ordertype α the well-orderedness of \prec is provable within T , and alternatively, the least stage $\|T\|$ not provable in T . The ξ th stage $I_\xi^\mathcal{E}$ of an inductive definition \mathcal{E} is given by $\{x : \mathcal{E}(\bigcup_{\eta < \xi} I_\eta^\mathcal{E}, x)\}$, and a stage α is called provable if there is an inductive definition \mathcal{A} and an $m \in \mathbb{N}$ with $T \vdash m \in P^\mathcal{A}$, such that $\alpha \leq \min\{\beta : m \in I_\beta^\mathcal{A}\}$. In [4], Cantini proves that $|ID_n^*| \leq \alpha_{2^n}$ and $\|ID_n^*\| \leq \alpha_{2^n-1}$ for $n > 0$ and conjectures, like Feferman in [6], $|ID_n^*| = \alpha_n = \|ID_n^*\|$ to hold true. We point out that for impredicative theories these two ordinal assignments usually coincide, whereas in (meta-)predicative theories they usually differ: $|ID_n| = \|ID_n\|$ is the case, however $\|\widehat{ID}_n\| = \alpha_{n-1}$ only unfolds to $|\widehat{ID}_n| = \alpha_n$. In respect thereof, the theories ID_n^* take an exceptional position, as $|ID_n^*| = \|ID_n^*\|$ follows from our wellordering proof.

This article provides the proof-theoretic ordinals of the theories ID_α^* and $ID_\alpha^* \upharpoonright$, a variant where also induction on the natural numbers is restricted to formulas that contain fixed point constants only positively. In the first sections we treat the case ID_1^* . Thereto we present a new embedding of \widehat{ID}_1 into Σ_1^1 -AC that extends to an embedding of ID_1^* into Σ_1^1 -DC. This embedding relies on the following two observations: Already Σ_1^1 -AC proves that for an operator form $\mathcal{A}(P^+, p)$, the standard Π_1^1 definition of the fixed point, namely the intersection of all sets X satisfying $\forall x[\mathcal{A}(X, x) \rightarrow x \in X]$, is a fixed point of the operator defined by \mathcal{A} . Moreover, Σ_1^1 -DC proves that this is indeed the least Π_1^1 -definable fixed point.

To demonstrate the power of restricted fixed point induction, we give well-ordering proofs for ID_α^* and $ID_\alpha^* \upharpoonright$. The general idea is the same as in [9] or [7], but things are simpler and the proofs are carried out in ID_α^* and $ID_\alpha^* \upharpoonright$ themselves. Section 7 is devoted to the upper bounds. In [15], R\"uede has developed and analyzed semi-formal systems to treat theories M_α , formalizing transfinite hierarchies of models of Σ_1^1 -AC. To embed the theories ID_α^* and $ID_\alpha^* \upharpoonright$ into such systems, we require uniform hierarchies of models of Σ_1^1 -DC. Towards this, we extend M_α to M_α^+ , by an axiom claiming that transfinite induction for $|M_\alpha|$, the proof-theoretic ordinal of M_α , fails. According to J\"ager and Probst [10], this extension is conservative. In M_α^+ , the technique of pseudo-hierarchies can be applied to construct the required hierarchies. R\"uede's results then yield sharp upper bounds.

§2. The theories ID_1^* and $ID_1^* \upharpoonright$. Let L_1 be a standard language of first order arithmetic that includes number variables $a, b, c, d, e, u, v, w, x, y, z, \dots$ and function and relation symbols for all primitive recursive functions and relations. In particular, we have a unary relation symbol N for the natural numbers. Moreover, we have unary relation symbols U and V that are required for technical reasons. Since we consider Tait-style calculi in the sequel, we use the symbol \sim for forming negative literals, and define the negation $\neg A$ of a formula A of L_1 or some language containing L_1 by making use of De Morgan's laws and the law of double negation. For $U(t)$ and $\sim U(t)$ we write $t \in U$ and $t \notin U$.

Towards the formulation of ID_1^* , we extend L_1 by fresh unary relation symbols \bar{P}, \bar{Q} and a fresh number constant p , which serve as placeholders. Then, a P -positive formula of $L_1(P, \bar{Q}, p)$, the extension of L_1 by P, \bar{Q} and p , is called an inductive

operator form, and we let \mathcal{A} range over such forms. For sets \vec{Y} and numbers \vec{y} , an operator form $\mathcal{A}(P, \vec{Q}, q, \vec{u})$ defines an operator on the powerset of the natural numbers, namely

$$F_{\vec{Y}, \vec{y}}^{\mathcal{A}}(X) := \{x : \mathcal{A}(X, \vec{Y}, x, \vec{y})\}.$$

Next, we add to the first order language L_1 a fixed point constant $P^{\mathcal{A}}$ for each inductive operator form \mathcal{A} of $L_1(P, p)$ without free variables, and denote this new language by L_{Fix} . Technically, we treat fixed point constants as unary relation symbols, but write $t \in P^{\mathcal{A}}$ instead of $P^{\mathcal{A}}(t)$. The formulas A, B, C, \dots and the number terms r, s, t, \dots of L_{Fix} are defined in the expected way and the formulas of L_{Fix}^+ are the formulas of L_{Fix} that contain fixed point constants only positively.

The axioms of ID_1^* consist of the axioms of PA without induction, complete induction along the natural numbers for all formulas of L_{Fix} as well as the following two fixed point axioms: For all inductive operator forms $\mathcal{A}(P, p)$ without free variables, we have

$$(FIX) \quad \forall x[\mathcal{A}(P^{\mathcal{A}}, x) \leftrightarrow x \in P^{\mathcal{A}}],$$

and for all inductive operator forms $\mathcal{A}(P, p)$, $\mathcal{A}_1(P, p), \dots, \mathcal{A}_n(P, p)$ without free variables, and each \vec{P} -positive formula $B(\vec{P}, p, \vec{u})$ of $L_1(\vec{P}, p)$, we have

$$(IND_{\text{Fix}}^+) \quad \forall x[\mathcal{A}(\{z : B(P^{\vec{\mathcal{A}}}, z, \vec{y})\}, x) \rightarrow B(P^{\vec{\mathcal{A}}}, x, \vec{y})] \rightarrow \\ \forall x[x \in P^{\vec{\mathcal{A}}} \rightarrow B(P^{\vec{\mathcal{A}}}, x, \vec{y})].$$

Note that we wrote $P^{\vec{\mathcal{A}}}$ for the string $P^{\mathcal{A}_1}, \dots, P^{\mathcal{A}_n}$ and that \mathcal{A} may be syntactically identical to some \mathcal{A}_i . The axiom (FIX) asserts that $P^{\mathcal{A}}$ is indeed a fixed point of the operator $F^{\mathcal{A}}$ and (IND_{Fix}^+) is the scheme for proof by induction on $P^{\mathcal{A}}$ restricted to formulas of L_{Fix}^+ . Finally, $ID_1^+ \upharpoonright$ denotes the theory where also complete induction along the natural numbers is restricted to formulas of L_{Fix}^+ , and \widehat{ID}_1 is ID_1^* without (IND_{Fix}^+) .

In this article, we make use of the term *proof-theoretic ordinal*. For theories T that are formulated in a language comprising L_1 , the proof-theoretic ordinal of T can be defined in the following way: We set

$$\text{Prog}_{\prec}(Z) := \forall u(\forall v \prec u)(v \in Z \rightarrow u \in Z), \\ \text{TI}_{\prec}(Z, t) := \text{Prog}_{\prec}(Z) \rightarrow (\forall u \prec t)(u \in Z),$$

and call an ordinal α provable in T , if there exists a primitive recursive well-ordering \prec such that $T \vdash \text{TI}_{\prec}(U, \alpha)$. Any ordinal that is not provable in T is called an *upper bound* of T and the least ordinal that is not provable in T is then the *proof-theoretic ordinal* of T , denoted by $|T|$.

§3. A new embedding of \widehat{ID}_1 into Σ_1^1 -AC. The standard embedding of \widehat{ID}_1 into Σ_1^1 -AC is due to Aczel [1]. He makes use of a universal Σ_1^1 formula and a standard diagonalization argument to find a Σ_1^1 definable solution for each fixed point constant $P^{\mathcal{A}}$ respecting (FIX). Of course, there is no chance to prove that such a solution is minimal with respect to classes definable by L_{Fix}^+ formulas. Bearing such a minimality condition in mind, the most natural way to interpret a fixed point constant $P^{\mathcal{A}}$ is to take its Π_1^1 definition, i.e., the intersection of all sets satisfying

$F^{\mathcal{A}}(X) \subseteq X$. This is indeed in accord with axiom (FIX). Surprisingly enough, the compact proof of this fact has not yet been discovered. Prior to its presentation, we specify the language and axioms of the theories involved, and briefly recap Aczel's argument.

The theories $\Sigma_1^1\text{-AC}$ and $\Sigma_1^1\text{-AC}_0$ are formulated in the language L_2 that canonically extends our language L_1 to a language of second order arithmetic by set variables U, V, W, X, Y, Z, \dots , a symbol \in to denote elementhood and quantifiers for second order variables. Note, that we write $t \notin X$ for $\sim(t \in X)$. The number terms of L_2 are the number terms of L_1 . Formulas of L_2 that do not contain bounded set variables are called arithmetical. L_2 formulas of the form $\exists X A(X)$, where A is arithmetical, are called Σ_1^1 formulas, and formulas of the form $\neg B$, where B is Σ_1^1 , are called Π_1^1 formulas. The class of Π formulas of L_2 is the smallest class containing the arithmetical formulas of L_2 that is closed under conjunction, disjunction, number quantification and universal set quantification. If A is a Π formula of L_2 , then $\neg A$ is a Σ formula of L_2 . Arithmetical formulas of L_2 where all number quantifiers appear in the context $(\forall x < t)$ and $(\exists x < t)$ are called Δ_0^0 .

In the sequel, we make use of the usual coding machinery: $\langle \dots \rangle$ is a standard primitive recursive function for forming n -tuples $\langle t_0, \dots, t_{n-1} \rangle$, so-called sequence numbers; $(t)_i$ is the i th component of (the sequence coded by) t , if i is less than the length $\text{lh}(t)$ of t ; i.e., $(t)_i = t_i$ for all $0 \leq i \leq n-1$, provided that $t = \langle t_0, \dots, t_{n-1} \rangle$. Further, we write $s \in (X)_i$ for $\langle s, t \rangle \in X$, and $X = Y$ is to abbreviate the formula $\forall x [x \in X \leftrightarrow x \in Y]$.

Besides the usual axioms of classical logic with equality in the first sort and axioms for the primitive recursive functions and relations, the theory $\Sigma_1^1\text{-AC}$ comprises the schema of complete induction on the natural numbers for all formulas of L_2 , arithmetical comprehension (ACA), and for all Σ_1^1 formulas $A(U, u)$ an axiom

$$(\Sigma_1^1\text{-AC}) \quad \forall x \exists X A(X, x) \rightarrow \exists Y \forall x A((Y)_x, x).$$

$\Sigma_1^1\text{-AC}_0$ is $\Sigma_1^1\text{-AC}$ with the schema of complete induction on the natural numbers restricted to sets.

Below we observe that in a theory comprising $\Sigma_1^1\text{-AC}_0$, we do not have to distinguish between Π and Π_1^1 formulas of L_2 . Of course, this applies also to the dual classes of Σ_1^1 and Σ formulas.

LEMMA 3.1. *For each Π formula C of L_2 there is a Π_1^1 formula C' of L_2 containing the same free variables as C , such that $\Sigma_1^1\text{-AC}$ proves: $C \leftrightarrow C'$.*

Aczel's embedding of $\widehat{\text{ID}}_1$ into $\Sigma_1^1\text{-AC}$ relies on this observation and the fact, that there exists a universal Σ_1^1 formula $E(u, v, w)$ of L_2 : For each Σ formula $B(u, v)$ of L_2 , there exists an $e \in \mathbb{N}$ such that

$$\Sigma_1^1\text{-AC}_0 \vdash B(x, y) \leftrightarrow E(\bar{e}, x, y),$$

where \bar{e} denotes the constant for the natural number e . This means in particular, that for a given operator form \mathcal{A} of $L_1(P, p)$, there is an $e_{\mathcal{A}} \in \mathbb{N}$ such that

$$\Sigma_1^1\text{-AC}_0 \vdash \mathcal{A}(\{z : E(x, x, z)\}, y) \leftrightarrow E(\bar{e}_{\mathcal{A}}, x, y).$$

Letting $C(u)$ be the Σ_1^1 formula $E(\bar{e}_{\mathcal{A}}, \bar{e}_{\mathcal{A}}, u)$, then $\Sigma_1^1\text{-AC}_0$ proves:

$$\begin{aligned}\mathcal{A}(\{z : C(z)\}, x) &\leftrightarrow \mathcal{A}(\{z : E(\bar{e}_{\mathcal{A}}, \bar{e}_{\mathcal{A}}, z)\}, x) \\ &\leftrightarrow E(\bar{e}_{\mathcal{A}}, \bar{e}_{\mathcal{A}}, x) \\ &\leftrightarrow C(x).\end{aligned}$$

If we translate an L_{Fix} formula B to an L_2 formula \tilde{B} by substituting each subformula of B of the form $t \in P^{\mathcal{A}}$ by the Σ_1^1 formula $E(\bar{e}_{\mathcal{A}}, \bar{e}_{\mathcal{A}}, t)$, then we obtain the following theorem:

THEOREM 3.2 (Aczel). *For every L_{Fix} formula B the following holds:*

$$\widehat{\text{ID}}_1 \vdash B \implies \Sigma_1^1\text{-AC} \vdash \tilde{B}.$$

The canonic candidate to interpret the fixed point constant $P^{\mathcal{A}}$, however, is the intersection of all \mathcal{A} -closed sets, namely the Π_1^1 -definable class

$$\text{Fix}^{\mathcal{A}} := \bigcap \{X : F^{\mathcal{A}}(X) \subseteq X\}.$$

Of course, we cannot prove in $\Sigma_1^1\text{-AC}$ that $\text{Fix}^{\mathcal{A}}$ is a set, yet $F^{\mathcal{A}}(\text{Fix}^{\mathcal{A}}) \subseteq \text{Fix}^{\mathcal{A}}$ is still immediate: For all \mathcal{A} -closed sets X , the positivity of the operator form \mathcal{A} yields $F^{\mathcal{A}}(\text{Fix}^{\mathcal{A}}) \subseteq F^{\mathcal{A}}(X) \subseteq X$. For the other direction, though, we can no longer argue that $F^{\mathcal{A}}(\text{Fix}^{\mathcal{A}})$ is \mathcal{A} -closed, and therefore a superset of $\text{Fix}^{\mathcal{A}}$. To show that $\Sigma_1^1\text{-AC}_0$ proves $\text{Fix}^{\mathcal{A}} \subseteq F^{\mathcal{A}}(\text{Fix}^{\mathcal{A}})$, a more refined argument is required.

We prove $F^{\mathcal{A}}(\text{Fix}^{\mathcal{A}}) = \text{Fix}^{\mathcal{A}}$ in a slightly more general context. For an operator form $\mathcal{A}(P, \vec{Q}, p, \vec{u})$ of $L_1(P, \vec{Q}, p)$ we set

$$\begin{aligned}\text{Cl}_{\vec{Y}, \vec{y}}^{\mathcal{A}}(X) &:= \forall x (\mathcal{A}(X, \vec{Y}, x, \vec{y}) \rightarrow x \in X), \\ \text{Fix}_{\vec{Y}, \vec{y}}^{\mathcal{A}} &:= \{x : \forall X [\text{Cl}_{\vec{Y}, \vec{y}}^{\mathcal{A}}(X) \rightarrow x \in X]\}.\end{aligned}$$

Often, we do not explicitly indicate the parameters in the operator form \mathcal{A} , and write $\text{Cl}^{\mathcal{A}}(X)$, $\text{Fix}^{\mathcal{A}}$ and $F^{\mathcal{A}}$ instead of $\text{Cl}_{\vec{Y}, \vec{y}}^{\mathcal{A}}(X)$, $\text{Fix}_{\vec{Y}, \vec{y}}^{\mathcal{A}}$ and $F_{\vec{Y}, \vec{y}}^{\mathcal{A}}$. The context provides always enough information to identify the dropped parameters. Below, we prove within $\Sigma_1^1\text{-AC}_0$ that $\text{Fix}_{\vec{Y}, \vec{y}}^{\mathcal{A}}$ is a fixed point of the operator $F_{\vec{Y}, \vec{y}}^{\mathcal{A}}$. The direction from right to left is again immediate. For the other direction, the following lemma almost handles the job.

LEMMA 3.3 (Separation Lemma). *For all operator forms \mathcal{A} of $L_1(P, \vec{Q}, p)$ and each arithmetical, U -positive formula $B(U, u)$ of L_2 , $\Sigma_1^1\text{-AC}_0$ proves:*

$$\neg B(\text{Fix}_{\vec{Y}, \vec{y}}^{\mathcal{A}}, x) \rightarrow \exists X [\text{Cl}_{\vec{Y}, \vec{y}}^{\mathcal{A}}(X) \wedge \neg B(X, x)].$$

PROOF. We prove the lemma by induction on the build-up of the formula $B(U, u)$. If U does not occur in B there is nothing to prove, and if B is the formula $t \in U$, then the claim follows from the definition of $\text{Fix}^{\mathcal{A}}$. If B is a conjunction or a disjunction, a similar argument applies as in the cases treated below.

- (i) $B(U, u)$ is of the form $\exists y B_1(U, u, y)$. Assume $\forall y \neg B_1(\text{Fix}^{\mathcal{A}}, x, y)$. The I.H. reads

$$\neg B_1(\text{Fix}^{\mathcal{A}}, x, y) \rightarrow \exists X [\text{Cl}^{\mathcal{A}}(X) \wedge \neg B_1(X, x, y)],$$

hence our assumption yields that

$$\forall y \exists X [\text{Cl}^{\mathcal{A}}(X) \wedge \neg B_1(X, x, y)].$$

Applying $(\Sigma_1^1\text{-AC})$ gives us a set Y such that

$$\forall y[\text{Cl}^{\mathcal{A}}((Y)_y) \wedge \neg B_1((Y)_y, x, y)].$$

Now we set

$$Z := \{x : \forall y(x \in (Y)_y)\},$$

and observe that $\text{Cl}^{\mathcal{A}}(Z)$: From $\mathcal{A}(Z, x)$ we conclude that $\forall y\mathcal{A}((Y)_y, x)$, and so $\forall y\text{Cl}^{\mathcal{A}}((Y)_y)$ yields $\forall y(x \in (Y)_y)$. Hence, by the positivity of B_1 , we have

$$\text{Cl}^{\mathcal{A}}(Z) \wedge \forall y\neg B_1(Z, x, y).$$

- (ii) $B(U, u)$ is of the form $\forall yB_1(U, u, y)$. Assume $\exists y\neg B_1(\text{Fix}^{\mathcal{A}}, x, y)$. Now the I.H. yields $\exists y\exists X[\text{Cl}^{\mathcal{A}}(X) \wedge \neg B_1(X, x, y)]$, which implies $\exists X[\text{Cl}^{\mathcal{A}}(X) \wedge \neg B(X, x)]$.

⊥

Our claim is now obtained effortlessly.

LEMMA 3.4. *For all operator forms \mathcal{A} of $L_1(P, \vec{Q}, p)$, $\Sigma_1^1\text{-AC}_0$ proves:*

$$\forall x[x \in \text{Fix}_{\vec{Y}, \vec{y}}^{\mathcal{A}} \leftrightarrow \mathcal{A}(\text{Fix}_{\vec{Y}, \vec{y}}^{\mathcal{A}}, \vec{Y}, x, \vec{y})].$$

PROOF. It remains to show that $x \in \text{Fix}^{\mathcal{A}}$ implies $\mathcal{A}(\text{Fix}^{\mathcal{A}}, x)$. To show the contraposition, we assume that $x \notin F^{\mathcal{A}}(\text{Fix}^{\mathcal{A}})$. By Lemma 3.3 there is a \mathcal{A} -closed set Z with $x \notin F^{\mathcal{A}}(Z)$. Since also $F^{\mathcal{A}}(Z)$ is \mathcal{A} -closed, $x \notin \text{Fix}^{\mathcal{A}}$ follows. ⊥

Summing up, we have established that for each operator form \mathcal{A} of $L_1(P, p)$, the intersection of all \mathcal{A} -closed sets is a fixed point of the operator $F^{\mathcal{A}}$, provable in $\Sigma_1^1\text{-AC}_0$. This gives rise to the following embedding:

THEOREM 3.5. *If we translate an L_{Fix} formula B to a L_2 formula B^* by substituting each fixed point constant $P^{\mathcal{A}}$ by the Π_1^1 -definable class $\text{Fix}^{\mathcal{A}}$, the following holds:*

$$\widehat{\text{ID}}_1 \vdash B \implies \Sigma_1^1\text{-AC} \vdash B^*.$$

§4. Embedding ID_1^* into $\Sigma_1^1\text{-DC}_0$. The theory $\Sigma_1^1\text{-AC}_0$ proves that $\text{Fix}^{\mathcal{A}}$ is a subclass of every \mathcal{A} -closed set. When we move to the slightly stronger theory $\Sigma_1^1\text{-DC}_0$ by strengthening the choice principle $(\Sigma_1^1\text{-AC})$, we even can prove that $\text{Fix}^{\mathcal{A}}$ is contained in every \mathcal{A} -closed, Π_1^1 -definable class. As a consequence, we also obtain induction along the natural numbers for Π_1^1 formulas. Thus, the aforementioned embedding extends to an embedding of ID_1^* into $\Sigma_1^1\text{-DC}_0$.

Formally, the theories $\Sigma_1^1\text{-DC}$ and $\Sigma_1^1\text{-DC}_0$ are obtained from $\Sigma_1^1\text{-AC}$ and $\Sigma_1^1\text{-AC}_0$ by replacing the axiom schema $(\Sigma_1^1\text{-AC})$ by the schema $(\Sigma_1^1\text{-DC})$: For each Σ_1^1 formula $A(U, V)$ of L_2 we have

$$(\Sigma_1^1\text{-DC}) \quad \forall X\exists Y A(X, Y) \rightarrow \forall Q\exists Z[(Z)_0 = Q \wedge \forall x A((Z)_x, (Z)_{x+1})].$$

Note, that the theory $\Sigma_1^1\text{-DC}_0$ proves each instance of the axiom schema $(\Sigma_1^1\text{-AC})$.

Next, we proof within $\Sigma_1^1\text{-DC}_0$ that $\text{Fix}^{\mathcal{A}}$ is the least Π_1^1 -definable fixed point of the operator $F^{\mathcal{A}}$.

THEOREM 4.1. *For all operator forms \mathcal{A} of $L_1(P, \vec{Q}, p)$ and each Π_1^1 formula $C(u)$ of L_2 , the following is provable in $\Sigma_1^1\text{-DC}_0$:*

$$\text{Cl}_{\vec{Y}, \vec{y}}^{\mathcal{A}}(\{x : C(x)\}) \rightarrow \text{Fix}_{\vec{Y}, \vec{y}}^{\mathcal{A}} \subseteq \{x : C(x)\}.$$

Before we give the proof, we consider a simpler case to illustrate the proof idea: Suppose that \mathcal{A} and \mathcal{B} are operator forms and that $\text{Fix}^{\mathcal{B}}$ is \mathcal{A} -closed. We assume that there is an $x \in \text{Fix}^{\mathcal{A}}$ with $x \notin \text{Fix}^{\mathcal{B}}$, and argue for a contradiction. Thereto, we construct a sequence $V_0 \supseteq V_1 \supseteq \dots$ of \mathcal{B} -closed set, such that for all $n \in \mathbb{N}$, we have $x \notin V_n$ and $V_n \supseteq F^{\mathcal{A}}(V_{n+1})$. Then $W := \bigcap_{n \in \mathbb{N}} V_n$ is \mathcal{A} -closed, but $x \notin W$.

To apply this argument in the general case, we require that every Π_1^1 -definable class $\{x : C(x)\}$ is primitive recursive in a fixed point.

LEMMA 4.2 (Representation Lemma). *For each Π_1^1 formula $C(U, u)$ of L_2 there exists an operator form \mathcal{A} of $L_1(P, Q, p)$ and a U -positive Δ_0^0 formula $D(U, u)$ of L_2 , such that $\Sigma_1^1\text{-AC}_0$ proves: For all sets Y , there exists a set T , such that*

$$\forall x [D(\text{Fix}_T^{\mathcal{A}}, x) \leftrightarrow C(Y, x)].$$

PROOF. As follows e.g., from results in Simpson [18], $\Sigma_1^1\text{-AC}_0$ proves that there is a set T , depending on the number and set parameters occurring in C , such that for all n ,

$$(T)_n \text{ is a tree,} \quad \text{and} \quad C(n) \leftrightarrow [(T)_n \text{ is well-founded}].$$

As usual, a tree is a set of finite sequences that is closed under initial segments. Now we define an operator $F^{\mathcal{A}}$ that collects the leafs of the trees $(T)_n$. If the tree $(T)_n$ is well-founded, then the root $\langle \rangle$ of the tree $(T)_n$ is an element of $\text{Fix}^{\mathcal{A}}$, otherwise the infinite branches and therefore the root do not enter the fixed point.

Thus, we set

$$\mathcal{A}(P, Q, p) := \exists n [p = \langle y, n \rangle \wedge y \in (Q)_n \wedge (\forall z \in (Q)_n) (z \supset y \rightarrow z \in (P)_n)],$$

where $z \supset y$ states that z is a proper extension of the sequence y . It is now easy to see that

$$\forall n [\langle \langle \rangle, n \rangle \in \text{Fix}_T^{\mathcal{A}} \leftrightarrow C(n)]. \quad \dashv$$

Next we return to the proof of Theorem 4.1.

PROOF. Assume that \mathcal{A} is an operator form and $C(u)$ a Π_1^1 formula of L_2 such that $\text{Cl}^{\mathcal{A}}(\{x : C(x)\})$. We aim to prove that $x \in \text{Fix}^{\mathcal{A}}$ implies $C(x)$.

Lemma 4.2 provides a set T , an operator form \mathcal{B} of $L_1(P, Q, p)$ and a U -positive Δ_0^0 formula $D(U, u)$ of L_2 such that

$$\forall x [D(\text{Fix}_T^{\mathcal{B}}, x) \leftrightarrow C(x)].$$

Hence our assumption reads $\text{Cl}^{\mathcal{A}}(\{x : D(\text{Fix}_T^{\mathcal{B}}, x)\})$. We show that this implies

$$(1) \quad \forall X \exists Z [F^D(X) \neq N \wedge \text{Cl}_T^{\mathcal{B}}(X) \rightarrow \text{Cl}_T^{\mathcal{B}}(Z) \wedge Z \subseteq X \wedge F^{\mathcal{A}} \circ F^D(Z) \subseteq F^D(X)].$$

Fix an arbitrary X , such that $\text{Cl}_T^{\mathcal{B}}(X)$, and suppose that $F^D(X)$ does not contain all natural numbers. If $x \notin F^D(X)$, then $x \notin F^D(\text{Fix}_T^{\mathcal{B}})$, so our assumption yields $x \notin F^{\mathcal{A}} \circ F^D(\text{Fix}_T^{\mathcal{B}})$, and Lemma 3.3 provides a set Y that is \mathcal{B} -closed with respect to T , such that $x \notin F^{\mathcal{A}} \circ F^D(Y)$. If $\text{Cl}_T^{\mathcal{B}}(X)$ and $\text{Cl}_T^{\mathcal{B}}(Y)$ then also $\text{Cl}_T^{\mathcal{B}}(X \cap Y)$, thus we may assume that $Y \subseteq X$. Summarizing, we obtain

$$\forall x \exists Y [x \notin F^D(X) \rightarrow \text{Cl}_T^{\mathcal{B}}(Y) \wedge Y \subseteq X \wedge x \notin F^{\mathcal{A}} \circ F^D(Y)].$$

Now $(\Sigma_1^1\text{-AC})$ gives us a set Y such that for all $x \notin F^D(X)$

$$\text{Cl}_T^{\mathcal{B}}((Y)_x) \wedge (Y)_x \subseteq X \wedge x \notin F^{\mathcal{A}} \circ F^D((Y)_x).$$

Therefore, if we set

$$Z := \bigcap_{x \notin F^D(X)} (Y)_x,$$

we have $\text{Cl}_T^{\mathcal{B}}(Z)$ and $Z \subseteq X$ and

$$\forall x[x \notin F^D(X) \rightarrow x \notin F^{\mathcal{A}} \circ F^D(Z)],$$

which means $F^{\mathcal{A}} \circ F^D(Z) \subseteq F^D(X)$. Thus we have shown claim (1).

Now we suppose that there is an $x \in \text{Fix}^{\mathcal{A}}$ that is not an element of $x \notin F^D(\text{Fix}_T^{\mathcal{B}})$ and argue for a contradiction. Again, Lemma 3.3 provides a set Q that is \mathcal{B} -closed with respect to T and $x \notin F^D(Q)$. Applying $(\Sigma_1^1\text{-DC})$ to (1) gives us a set V such that $(V)_0 = Q$ and

$$\forall n[\text{Cl}_T^{\mathcal{B}}((V)_n) \rightarrow \text{Cl}_T^{\mathcal{B}}((V)_{n+1}) \wedge (V)_{n+1} \subseteq (V)_n \wedge F^{\mathcal{A}} \circ F^D((V)_{n+1}) \subseteq F^D((V)_n)].$$

One easily proves by induction that

$$\forall n[\text{Cl}_T^{\mathcal{B}}((V)_n) \wedge (V)_{n+1} \subseteq (V)_n \wedge F^{\mathcal{A}} \circ F^D((V)_{n+1}) \subseteq F^D((V)_n)].$$

Hence, for $W := \bigcap_{n \in \mathbb{N}} (V)_n$, we have that

$$F^{\mathcal{A}} \circ F^D(W) \subseteq \bigcap_{n \in \mathbb{N}} F^D((V)_n) = F^D\left(\bigcap_{n \in \mathbb{N}} (V)_n\right) = F^D(W).$$

The second but last equality follows from the fact that D is positive and Δ_0^0 , and that $(\forall n \in \mathbb{N})((V)_{n+1} \subseteq (V)_n)$. So $W \subseteq Q$ and $\text{Cl}^{\mathcal{A}}(F^D(W))$, i.e., $\text{Fix}^{\mathcal{A}} \subseteq F^D(W)$. Now $x \notin F^D(Q)$ yields $x \notin F^D(W)$, thus $x \notin \text{Fix}^{\mathcal{A}}$. A contradiction! \dashv

The following corollary is an immediate consequence of Theorem 4.1. To enhance readability, we let $\text{Fix}_{\vec{P}}^{\vec{\mathcal{A}}}$ stand for $\text{Fix}_{\vec{P}}^{\mathcal{A}_1}, \dots, \text{Fix}_{\vec{P}}^{\mathcal{A}_n}$.

COROLLARY 4.3. For all operator forms \mathcal{A} and $\vec{\mathcal{A}}$ of $L_1(P, \vec{Q}, p)$ and each \vec{U} -positive arithmetical formula $B(\vec{U}, \vec{u})$ of $L_2, \Sigma_1^1\text{-DC}_0$ proves:

$$\text{Cl}^{\mathcal{A}}(\{x : B(\text{Fix}_{\vec{P}}^{\vec{\mathcal{A}}}, x, \vec{z})\}) \rightarrow \text{Fix}_{\vec{P}}^{\mathcal{A}} \subseteq \{x : B(\text{Fix}_{\vec{P}}^{\vec{\mathcal{A}}}, x, \vec{z})\}.$$

PROOF. Note that $B(\text{Fix}_{\vec{P}}^{\vec{\mathcal{A}}}, x, \vec{z})$ is equivalent to a Π_1^1 formula of L_2 . \dashv

REMARK 4.4. We think of U as coding an ordering, and set

$$\text{Prog}(U, V) := \forall x[\forall y(\langle y, x \rangle \in U \rightarrow y \in V) \rightarrow x \in V],$$

$$\text{TI}(U, V) := \text{Prog}(U, V) \rightarrow \text{Field}(U) \subseteq V,$$

$$\text{Wo}(U) := \forall Y \text{TI}(U, Y).$$

Further, we consider the operator form

$$\mathcal{A}^{\text{Wo}}(P, Q, p) := \forall y[\langle y, p \rangle \in Q \rightarrow y \in P].$$

Observe, that $\text{Cl}_U^{\mathcal{A}^{\text{Wo}}}(V)$ is the formula $\text{Prog}(U, V)$ and $\text{Wo}(X)$ can be written as $\forall Y[\text{Cl}_X^{\mathcal{A}^{\text{Wo}}}(Y) \rightarrow \text{Field}(X) \subseteq Y]$. It is immediate, that $\text{Wo}(X)$ is equivalent to

$\text{Fix}_X^{\mathcal{A}} = \text{Field}(X)$. Due to Theorem 4.1, $\Sigma_1^1\text{-DC}_0$ proves for each Π_1^1 formula $C(u)$ of L_2 that

$$\text{Wo}(X) \rightarrow [\text{Cl}_X^{\mathcal{A}}(\{z : C(z)\}) \rightarrow \text{Field}(X) \subseteq \{z : C(z)\}],$$

which is normally written as

$$(\Pi_1^1\text{-TI}) \quad \text{Wo}(X) \rightarrow \text{TI}(X, \{z : C(z)\}).$$

It is shown, e.g., in [18], that $(\Pi_1^1\text{-TI})$ is provable in $\Sigma_1^1\text{-DC}_0$. In this sense, Corollary 4.1 is a generalization of this result.

Since $(\Pi_1^1\text{-TI})$ implies induction along the natural numbers for all Π_1^1 formulas of L_2 , the embedding given in the previous section extends to an embedding of $\text{ID}_1^* \upharpoonright$ and ID_1^* into $\Sigma_1^1\text{-DC}_0$ and $\Sigma_1^1\text{-DC}$, respectively.

THEOREM 4.5. *If we translate an L_{Fix} formula B to a L_2 formula B^* by substituting each fixed point constant $P^{\mathcal{A}}$ by the Π_1^1 -definable class $\text{Fix}^{\mathcal{A}}$, then the following holds:*

$$\text{ID}_1^* \upharpoonright \vdash B \implies \Sigma_1^1\text{-DC}_0 \vdash B^* \quad \text{and} \quad \text{ID}_1^* \vdash B \implies \Sigma_1^1\text{-DC} \vdash B^*.$$

Since $|\Sigma_1^1\text{-DC}_0| = \varphi\omega 0$ and $|\Sigma_1^1\text{-DC}| = \varphi\varepsilon_0 0$, this answers the question for a sharp upper bound of ID_1^* :

COROLLARY 4.6.

$$|\text{ID}_1^* \upharpoonright| \leq \varphi\omega 0, \quad \text{and} \quad |\text{ID}_1^*| \leq \varphi\varepsilon_0 0.$$

§5. The theories ID_α^* and $\text{ID}_\alpha^* \upharpoonright$. To formulate transfinite iterations of the theories ID_1^* and $\text{ID}_1^* \upharpoonright$, we follow the lines chosen by Jäger, Kahle, Setzer and Strahm [9] and we presuppose the same ordinal-theoretic facts. Again, $(\text{OT}, \triangleleft)$ is a standard notation system based on the ternary Veblen or φ -function. As usual, we write 0 for the least element of OT with respect to the primitive recursive ordering \triangleleft . Ordinals are often identified with their notations. If an ordinal α appears within a formal argument, the closed term representing its notation is meant instead. Also, we do not distinguish between operations on ordinals and the primitive recursive analogues on their codes. By Φ_0 we denote the least ordinal greater than 0 such that with $\alpha < \Phi_0$ also $\varphi 1 \alpha 0 < \Phi_0$. We restrict ourselves to ordinals below Φ_0 because we only bother to fix fundamental sequences for these ordinals in the subsequent well-ordering proof. However, it is straight forward to extend the following to all ordinals below Φ_1 , the least ordinal greater than 0 which is closed under all n -ary φ -functions.

The language L_1 and operator forms \mathcal{A} are defined as in Section 2, but this time, we extend the language L_1 by a unary relation symbol $P^{\mathcal{A}}$ for each operator form $\mathcal{A}(P, Q, p, u)$ of $L_1(P, Q, p)$ which contains at most the variable u free, and denote this new language again by L_{Fix} . To simplify the notation, $t \in P_s^{\mathcal{A}}$ stands for $P^{\mathcal{A}}(\langle t, s \rangle)$ and $t \in P_{\triangleleft s}^{\mathcal{A}}$ is to abbreviate $t = \langle (t)_0, (t)_1 \rangle \wedge (t)_1 \triangleleft s \wedge t \in P^{\mathcal{A}}$. For each ordinal α less than Φ_0 , the theory ID_α^* comprises the axioms of PA without induction, the axioms $\text{TI}_{\triangleleft}(\mathcal{A}, \max\{\alpha, \omega\})$ for all L_{Fix} formulas \mathcal{A} and the following fixed point axioms:

$$(\text{FIX}) \quad (\forall a \triangleleft \alpha) \forall x [x \in P_a^{\mathcal{A}} \leftrightarrow \mathcal{A}(P_a^{\mathcal{A}}, P_{\triangleleft a}^{\mathcal{A}}, x, a)],$$

and

$$(\text{IND}_{\text{FIX}}^+) \quad (\forall a \triangleleft \alpha)[\text{Cl}_{\text{P}_{\triangleleft a}, a}^{\vec{\mathcal{A}}}(\{x : B(\text{P}_a^{\vec{\mathcal{A}}}, \text{P}_{\triangleleft a}^{\vec{\mathcal{A}}}, x, \vec{y})\}) \rightarrow \\ \text{P}_a^{\vec{\mathcal{A}}} \subseteq \{x : B(\text{P}_a^{\vec{\mathcal{A}}}, \text{P}_{\triangleleft a}^{\vec{\mathcal{A}}}, x, \vec{y})\}],$$

for all operator forms $\vec{\mathcal{A}}(\text{P}, \text{Q}, \text{p}, u)$ containing at most the variable u free and each \vec{P} -positive formula $B(\vec{P}, \vec{Q}, \text{p}, \vec{u})$ of $\text{L}_1(\vec{P}, \vec{Q}, \text{p})$. In ID_α^* , only restricted (transfinite) induction is available, i.e., instead of $\text{TI}_{\triangleleft}(A, \max\{\alpha, \omega\})$, we only have

$$(\forall a \triangleleft \alpha)[\text{Prog}_{\triangleleft}(\lambda b. B(\text{P}_a^{\vec{\mathcal{A}}}, \text{P}_{\triangleleft a}^{\vec{\mathcal{A}}}, b, \vec{y})) \rightarrow (\forall c \triangleleft \max\{\alpha, \omega\}) B(\text{P}_a^{\vec{\mathcal{A}}}, \text{P}_{\triangleleft a}^{\vec{\mathcal{A}}}, c, \vec{y})]$$

for each \vec{P} -positive formula $B(\vec{P}, \vec{Q}, \text{p}, \vec{u})$ of $\text{L}_1(\vec{P}, \vec{Q}, \text{p})$. As usual, for a formula $A(u)$ of L_{FIX} , $\text{Prog}_{\triangleleft}(\lambda x. A(x))$ abbreviates $\forall u(\forall v \triangleleft u)(A(v) \rightarrow A(u))$. Note that the axioms concerning transfinite induction imply also induction along the natural numbers for the corresponding class of formulas. Again, $\widehat{\text{ID}}_\alpha$ is ID_α^* without $(\text{IND}_{\text{FIX}}^+)$.

§6. Wellordering proofs for ID_α^* and $\text{ID}_\alpha^* \upharpoonright$. To demonstrate the power of the axiom $(\text{IND}_{\text{FIX}}^+)$, we give wellordering proofs for the theories ID_α^* and $\text{ID}_\alpha^* \upharpoonright$. The proof idea is the same as in [9], where the wellordering proof of $\widehat{\text{ID}}_\alpha$ is carried out in the transfinitely iterated theory of self-reflecting truth SRT_α . However, things are easier in the present context and the wellordering proof is performed in ID_α^* itself. As Corollary 4.6 suggests, we obtain that also ID_α^* and $\widehat{\text{ID}}_\alpha$ prove the same ordinals.

For the wellordering proof, we fix fundamental sequences for the ordinals below Φ_0 . A fundamental sequence for α is a primitive recursive, increasing sequence $\alpha[n]$ on the corresponding notations such that for each $\beta < \alpha < \Phi_0$ there is an n with $\beta \leq \alpha[n]$. We set $(\alpha + 1)[n] := \alpha$ for all $n \in \mathbb{N}$, and if $\omega^{\alpha_k} + \dots + \omega^{\alpha_1}$ is the Cantor normal form of λ and $\lambda < \omega^\lambda$, then $\lambda[n] := \omega^{\alpha_k} + \dots + \omega^{\alpha_2} + \omega^{\alpha_1}[n]$. The remaining cases where $\lambda < \Phi_0$ is of the form $\varphi\alpha\beta\gamma$ for $\alpha \in \{0, 1\}$ and $\beta, \gamma < \lambda$ are given next: $\Gamma_0[0] := 0$ and $\Gamma_0[n+1] := \varphi(\Gamma_0[n])0$, $\varphi 00(\gamma+1)[n] = \omega^{\gamma+1}[n] := \omega^\gamma \cdot n$, and

- (i) $\varphi\alpha(\beta+1)0[0] = 0$ and $\varphi\alpha(\beta+1)0[n+1] = \varphi\alpha\beta(\varphi\alpha(\beta+1)0[n])$.
- (ii) For a limit λ : $\varphi\alpha\beta\lambda[n] = \varphi\alpha\beta(\lambda[n])$ and $\varphi\alpha\lambda 0[n] = \varphi\alpha(\lambda[n])0$.
- (iii) $\varphi\alpha\beta(\gamma+1)[0] = \varphi\alpha\beta\gamma + 1$, and

$$\begin{aligned} \varphi\alpha\beta(\gamma+1)[n+1] &= \varphi\alpha(\beta[n])(\varphi\alpha\beta(\gamma+1)[n]), \text{ if } \beta > 0, \\ &= \varphi 0(\varphi 10(\gamma+1)[n])0, \text{ if } \beta = 0 \text{ and } \alpha = 1. \end{aligned}$$

In the course of the wellordering proof, we let a, b, c, d, e range over the elements of OT and use l to denote limit notations. We start with the cases ID_1^* and $\text{ID}_1^* \upharpoonright$.

Let $\mathcal{ACC} := (\forall z \triangleleft x)(z \in \text{P})$ and denote the corresponding fixed point constant $\text{P}^{\mathcal{A}}$ by ACC. By means of the axiom $(\text{IND}_{\text{FIX}}^+)$ one immediately proves in $\text{ID}_1^* \upharpoonright$ that $a, b \in \text{ACC}$ implies $\text{ACC} \subseteq \{c : a + c \in \text{ACC}\}$ and $\text{ACC} \subseteq \{c : a \cdot c \in \text{ACC}\}$, hence $a, b \in \text{ACC}$ yields $a + b \in \text{ACC}$ and $a \cdot b \in \text{ACC}$.

LEMMA 6.1. *For each ordinal $k < \omega$, and each ordinal $\kappa < \varepsilon_0$, the following holds:*

$$\text{ID}_1^* \upharpoonright \vdash \text{Prog}_{\triangleleft}(\lambda a. \varphi k a \in \text{ACC}) \quad \text{and} \quad \text{ID}_1^* \vdash \text{Prog}_{\triangleleft}(\lambda a. \varphi \kappa a \in \text{ACC}).$$

PROOF. Note that $\text{Prog}_{\triangleleft}(\lambda x. A(x))$ is another way of writing $\text{Cl}_{\text{P}^{\mathcal{A}}}^{\mathcal{A}}(\{x : A(x)\})$. We prove the first claim by (meta-) induction on k . For $k = 0$, it is to show that

if $\omega^b \in \text{ACC}$ holds for all $b \triangleleft a$, then also $\omega^a \in \text{ACC}$. If a is a limit notation, this follows from $\text{Prog}_{\triangleleft}(\lambda b. b \in \text{ACC})$ and the continuity of the function $\lambda \xi. \omega^\xi$. If a is of the form $b + 1$, then we use restricted induction to show that $\forall n(\omega^b \cdot n \in \text{ACC})$, thus $\text{Prog}_{\triangleleft}(\lambda b. b \in \text{ACC})$ yields $\omega^{b+1} \in \text{ACC}$.

For the induction step, we assume that $\varphi(k+1)b \in \text{ACC}$ for all $b \triangleleft a$. Now the I.H. allows to prove by restricted induction that $\forall n(\varphi(k+1)a[n] \in \text{ACC})$. Thus, also $\varphi(k+1)a \in \text{ACC}$.

For the second claim, observe that in ID_1^* transfinite induction along ordinals $\kappa < \varepsilon_0$ is available for all formulas of L_{FIX} . Instead of meta-induction, transfinite induction within ID_1^* is used. If λ is a limit ordinal, the induction step is performed by showing $\forall n(\varphi \lambda a[n] \in \text{ACC})$. \dashv

The axiom $(\text{IND}_{\text{FIX}}^+)$ implies $\text{Prog}_{\triangleleft}(\text{U}) \rightarrow \text{ACC} \subseteq \text{U}$. Since the previous lemma yields $\text{ID}_1^* \upharpoonright \vdash \varphi k 0 \in \text{ACC}$ and $\text{ID}_1^* \vdash \varphi \kappa 0 \in \text{ACC}$ for $k < \omega$ and $\kappa < \varepsilon_0$, Theorem 4.6 gives rise to the following corollary.

COROLLARY 6.2.

$$|\text{ID}_1^*| = \varphi \omega 0, \quad \text{and} \quad |\text{ID}_1^*| = \varphi \varepsilon_0 0.$$

Next we consider $\text{ID}_\alpha^* \upharpoonright$ and ID_α^* . By mentioning $\text{ID}_\alpha^* \upharpoonright$ or ID_α^* , we implicitly imply $\alpha < \Phi_0$. This time, let $\mathcal{A} := (\forall z \triangleleft p)(z \in P_a)$ and denote the corresponding relation symbol $P^{\mathcal{A}}$ by ACC . In the sequel we write $c \in \text{ACC}^a$ for $(\forall b \triangleleft a)(c \in \text{ACC}_b)$. Note that this reads $(\forall b \triangleleft a)((c, b) \in \text{ACC}_{\triangleleft a})$. Therefore $\text{ID}_\alpha^* \upharpoonright$ proves for each formula $B(P, p)$ of $\text{L}_1(P, p)$, that $a \triangleleft \alpha$ and $\text{Prog}_{\triangleleft}(\lambda x. B(\text{ACC}^a, x))$ imply $\text{ACC}_a \subseteq \{x : B(\text{ACC}^a, x)\}$. Further, $a \triangleleft \alpha$ implies the progressivity of ACC^a and $b \triangleleft a \triangleleft \alpha$ implies $\text{ACC}_a \subseteq \text{ACC}_b$. Since $\text{Prog}_{\triangleleft}(\text{U})$ implies $\text{ACC}_0 \subseteq \text{U}$, proving an ordinal β in $\text{ID}_\alpha^* \upharpoonright$ breaks down to show $\beta \in \text{ACC}_0$.

LEMMA 6.3. Let $A(a, b) := \forall c(c \in \text{ACC}_a \rightarrow \varphi bc \in \text{ACC}_a)$. Then it is provable in $\text{ID}_\alpha^* \upharpoonright$ that $a \triangleleft \alpha \rightarrow \text{Prog}_{\triangleleft}(\lambda b. A(a, b))$.

PROOF. Let $a \triangleleft \alpha$. We assume $(\forall b' \triangleleft b)A(a, b')$ and show $A(a, b)$.

$(\text{IND}_{\text{FIX}}^+)$ tells us that $A(a, b)$ follows from $\text{Prog}_{\triangleleft}(\lambda c. \varphi bc \in \text{ACC}_a)$, which in turn follows from the assumption $(\forall b' \triangleleft b)A(a, b')$: Given $\varphi bc' \in \text{ACC}_a$ for all $c' \triangleleft c$, restricted induction yields that $\forall n(\varphi bc[n] \in \text{ACC}_a)$, thus $\varphi bc \in \text{ACC}_a$. \dashv

COROLLARY 6.4. For all limit notations $l \trianglelefteq \alpha$, $\text{ID}_\alpha^* \upharpoonright$ proves:

$$d \in \text{ACC}^l \rightarrow \varphi d 0 \in \text{ACC}^l.$$

PROOF. Pick an arbitrary $a \triangleleft l$. So $d \in \text{ACC}^l$ implies $d \in \text{ACC}_{a+1}$. Now $(\text{IND}_{\text{FIX}}^+)$ and $\text{Prog}_{\triangleleft}(\lambda b. \forall c(c \in \text{ACC}_a \rightarrow \varphi bc \in \text{ACC}_a))$ yield $\varphi d 0 \in \text{ACC}_a$. \dashv

COROLLARY 6.5. For all limit notations $l \triangleleft \alpha$, $\text{ID}_\alpha^* \upharpoonright$ proves:

$$\text{Prog}_{\triangleleft}(\lambda c. \varphi 10c \in \text{ACC}^l).$$

PROOF. Assume that $l \triangleleft \alpha$ and that $\varphi 10d \in \text{ACC}^l$ for all $d \triangleleft c$. Restricted induction and the previous corollary imply $\forall n(\varphi 10c[n] \in \text{ACC}^l)$. Thus $\varphi 10c \in \text{ACC}^l$. \dashv

COROLLARY 6.6. For all limits $l \leq \alpha$, ID_α^* proves:

$$\text{Prog}_{\triangleleft}(\lambda c. \varphi 10c \in \text{ACC}^l).$$

PROOF. In the case $l = \alpha$, full induction is needed to show $\forall n(\varphi 10c[n] \in \text{ACC}^l)$. \dashv

The following lemma corresponds to the Main Lemma in [9]. Again, the proof is simpler in the present context.

LEMMA 6.7. *Let*

$$A(a, b) := \forall d, c(d + \omega^{1+b} \leq a \wedge c \in \text{ACC}_{d+\omega^{1+b}} \rightarrow \varphi 1bc \in \text{ACC}^{d+\omega^{1+b}}).$$

Then it is provable in ID_α^ that $a \triangleleft \alpha \rightarrow \text{Prog}_\triangleleft(\lambda b.A(a, b))$.*

PROOF. Assume $a \triangleleft \alpha$ and that $A(a, b')$ holds for all $b' \triangleleft b$. We aim for $A(a, b)$. So suppose $d + \omega^{1+b} \leq a$. Now $c \in \text{ACC}_{d+\omega^{1+b}} \rightarrow \varphi 1bc \in \text{ACC}^{d+\omega^{1+b}}$ follows, if we can establish

$$(1) \quad \text{Prog}_\triangleleft(\lambda c.\varphi 1bc \in \text{ACC}^{d+\omega^{1+b}}).$$

There to we further suppose that $\varphi 1bc' \in \text{ACC}^{d+\omega^{1+b}}$ for all $c' \triangleleft c$, and use restricted induction to show $\forall n(\varphi 1bc[n] \in \text{ACC}^{d+\omega^{1+b}})$. We only consider the case where b is not 0 and c a successor: $\varphi 1bc[0] \in \text{ACC}^{d+\omega^{1+b}}$ follows immediately from our further supposition, and the induction step can be performed because we have for all $m \in \mathbb{N}$,

$$(2) \quad \forall e(e \in \text{ACC}^{d+\omega^{1+b}} \rightarrow \varphi 1(b[m])e \in \text{ACC}^{d+\omega^{1+b}}).$$

To see that (2) holds, fix an $m \in \mathbb{N}$ and suppose that $e \in \text{ACC}^{d+\omega^{1+b}}$. We argue that $\varphi 1(b[m])e \in \text{ACC}_{d+\omega^{1+b}[k]}^{d+\omega^{1+b}}$ for all $k \in \mathbb{N}$: So we fix an arbitrary $k \in \mathbb{N}$, and observe that $e \in \text{ACC}_{d+\omega^{1+b}[k]+\omega^{1+b}[m]}^{d+\omega^{1+b}}$. Thus, the assumption $(\forall b' \triangleleft b)A(a, b')$ forces $\varphi 1(b[m])e \in \text{ACC}_{d+\omega^{1+b}[k]}^{d+\omega^{1+b}}$.

The other cases are shown similarly or are easy. If $b = 0$, (1) becomes Corollary 6.5. \dashv

From the above proof we immediately extract the following corollaries:

COROLLARY 6.8. For all notations b and all d with $d + \omega^{1+b} \triangleleft \alpha$, ID_α^* proves:

$$\text{Prog}_\triangleleft(\lambda c.\varphi 1bc \in \text{ACC}^{d+\omega^{1+b}}).$$

COROLLARY 6.9. For all notations b and all d with $d + \omega^{1+b} \leq \alpha$, ID_α^* proves:

$$\text{Prog}_\triangleleft(\lambda c.\varphi 1bc \in \text{ACC}^{d+\omega^{1+b}}).$$

PROOF. Let $A(a, b)$ be as defined in Lemma 6.7. By transfinite induction we obtain $A(a, b)$ for all $a \triangleleft \alpha$ and $b \leq \alpha$. Using full induction, the claim is shown as in the proof of Lemma 6.7. \dashv

In order to speak about lower and upper bounds of ID_α^* and $\text{ID}_\alpha^* \upharpoonright$, we define for all $\alpha, \beta < \Phi_0$ a function $\sigma(\alpha, \beta)$.

DEFINITION 6.10. Let $\alpha = \omega^{1+\alpha_n} + \omega^{1+\alpha_{n-1}} + \dots + \omega^{1+\alpha_1} + m$, where $\alpha_n \geq \dots \geq \alpha_1$ and $m < \omega$, be an ordinal below Φ_0 in Cantor normal form. We set for all ordinals $\beta < \Phi_0$:

$$\sigma(\alpha, \beta) := \varphi 1\alpha_n(\varphi 1\alpha_{n-1}(\dots(\varphi 1\alpha_1\beta)\dots)), \text{ if } \alpha \geq \omega \quad \text{and} \quad \sigma(m, \beta) := \beta.$$

Moreover, $(\alpha|0) := \varepsilon(\alpha)$, i.e., the least fixed point of the function $\lambda \xi.\omega^\xi$ bigger than α , and $(\alpha|i+1) := \varphi(\alpha|i)0$.

$(\alpha \upharpoonright 0)$ is the least limit ordinal $\lambda > 0$ such that $\varphi 1 \alpha_1 \lambda > \alpha$, and $(\alpha \upharpoonright 1)$ is the least upper bound of $\{\varphi k(\alpha + 1) : k < \omega\}$. Eventually, $(\alpha \upharpoonright i + 2) := \varphi(\alpha \upharpoonright i + 1)0$.

Towards further simplifications, we write in the sequel $\sigma(\alpha)$ for $\sigma(\alpha, (\alpha \upharpoonright m))$ and $\sigma \upharpoonright (\alpha)$ for $\sigma(\alpha, (\alpha \upharpoonright m))$.

Note, that $\alpha \upharpoonright 0$ is of the form $\beta + \omega$, where β is a limit or zero, and if $\beta > 0$, then $\alpha \geq \varphi 1 \alpha_1 \beta$: Let $\beta_0 := \min\{\beta : \varphi 1 \alpha_1 \beta \geq \alpha\}$. Now if β_0 is zero or a successor, the claim is immediate, and if β_0 is a limit, then the continuity of the function $\lambda \xi. \varphi 1 \alpha_1 \xi$ yields $\varphi 1 \alpha_1 \beta_0 = \alpha$, thus $\alpha \upharpoonright 0 = \beta_0 + \omega$.

LEMMA 6.11. *Let $\lambda = \omega^{1+\alpha_n} + \omega^{1+\alpha_{n-1}} + \dots + \omega^{1+\alpha_1} < \Phi_0$, where $\alpha_n \geq \dots \geq \alpha_1$, and assume that $\lambda \upharpoonright 0 = \beta + \omega$. Then, for each $n \in \mathbb{N}$, $\text{ID}_\lambda^* \upharpoonright$ proves the following:*

$$\varphi 1 \alpha_1(\beta + n) \in \text{ACC}^\lambda.$$

PROOF. We choose δ such that $\lambda = \delta + \omega^{1+\alpha_1}$ and prove the claim by meta-induction on n . We start with the case $n = 0$: If $\beta > 0$, then $\varphi 1 \alpha_1 \beta \leq \lambda$ and the claim trivially follows by restricted transfinite induction up to λ . If $\beta = 0$, then the claim follows similar to the induction step, which we prove below. Thereby, we distinguish whether α_1 is zero, a successor or a limit. Exemplarily, we show the case $\alpha_1 = \alpha'_1 + 1$: To establish $\varphi 1 \alpha_1(\beta + n + 1) \in \text{ACC}^{\delta + \omega^{1+\alpha'_1}}$, we show that for all $k \in \mathbb{N}$, $\varphi 1 \alpha_1(\beta + n + 1)[k] \in \text{ACC}^{\delta + \omega^{1+\alpha'_1} \cdot k}$ by proving, using restricted induction on m , that for each $k \in \mathbb{N}$,

$$(\forall m \leq k)[\varphi 1 \alpha_1(\beta + n + 1)[m] \in \text{ACC}_{\delta + \omega^{1+\alpha'_1} \cdot (2k-m)}]:$$

If $m < k$, then the I.H. yields $\varphi 1 \alpha_1(\beta + n + 1)[m] \in \text{ACC}_{\delta + \omega^{1+\alpha'_1} \cdot (2k-m)}$ and Corollary 6.8 tells us

$$\text{Prog}_{\triangleleft}(\lambda c. \varphi 1 \alpha'_1 c \in \text{ACC}^{\delta + \omega^{1+\alpha'_1} \cdot (2k-m)}).$$

Thus, $\varphi 1 \alpha_1(\beta + n + 1)[m + 1] \in \text{ACC}^{\delta + \omega^{1+\alpha'_1} \cdot (2k - (m+1))} \subseteq \text{ACC}_{\delta + \omega^{1+\alpha'_1} \cdot (2k - (m+1))}$. \dashv

We conclude this section by presenting the lower bounds:

THEOREM 6.12. *For all $0 < \alpha < \Phi_0$ we have:*

$$|\text{ID}_\alpha^*| \geq \sigma(\alpha) \quad \text{and} \quad |\text{ID}_\alpha^* \upharpoonright| \geq \sigma \upharpoonright (\alpha).$$

PROOF. Assume that $\alpha = \omega^{1+\alpha_n} + \dots + \omega^{1+\alpha_1} + m$ for ordinals $\alpha_n \geq \dots \geq \alpha_1$ and $m < \omega$, and set $\delta_k := \omega^{1+\alpha_n} + \dots + \omega^{1+\alpha_k}$ for $k \leq n$, and $\sigma_k := \varphi 1 \alpha_k(\dots \varphi 1 \alpha_1(\alpha \upharpoonright m)) \dots$. By meta-induction on k we now show that for all $\beta < \sigma_k$, the theory $\text{ID}_\alpha^* \upharpoonright$ proves $\beta \in \text{ACC}^{\delta_k}$:

We first consider the case $k = 1$. If $m = 0$, then $\delta_1 = \alpha$ and $\sigma_1 = \varphi 1 \alpha_1(\alpha \upharpoonright 0)$. Hence the claim follows by Lemma 6.11. If $m = m' + 1$, then there exists for each $\beta < \sigma_1 = \varphi 1 \alpha_1(\alpha \upharpoonright 1)$ a $k < \omega$ and ordinals ξ_1, \dots, ξ_m such that $\xi_1 = \varphi k(\alpha + 1)$ and $\xi_{i+i} = \varphi \xi_i 0$ and $\beta < \varphi 1 \alpha_1 \xi_m$. It follows from the proof of Lemma 6.1 that $\text{ID}_\alpha^* \upharpoonright$ proves the progressivity of $\lambda a. \varphi k a \in \text{ACC}^{\delta_1 + m'}$, thus $\xi_1 \in \text{ACC}_{\delta_1 + m'}$. Applying m' -times Lemma 6.3 and $(\text{IND}_{\text{FIX}}^+)$ yields $\xi_m \in \text{ACC}_{\delta_1}$. Now Lemma 6.7 yields $\varphi 1 \alpha_1 \xi_m \in \text{ACC}^{\delta_1}$.

The induction step from k to $k + 1$ follows with Corollary 6.8.

The case ID_α^* is treated similarly. If $m = 0$, we use that for all formulas $A(u)$ of L_{FIX} , ID_α^* proves $\text{Prog}_{\triangleleft}(\lambda a. A(a)) \rightarrow A(\beta)$, for all $\beta < \varepsilon(\alpha)$. \dashv

That these bounds are sharp is established in the next section.

§7. Upper bounds for ID_α^* and $ID_\alpha^*\uparrow$. The aim of this section is to determine the upper bounds of the ID^* -theories ID_α^* and $ID_\alpha^*\uparrow$. In a first step, we introduce for each ordinal $\alpha < \Phi_0$ a theory M_α which formalizes an α -hierarchy of models of Σ_1^1 -AC. Then upper bounds of the theories $(M_\alpha + ACA)_0$, $(M_\alpha + \Sigma_1^1\text{-DC})_0$ and $M_\lambda\uparrow$, the so called M-theories, are identified by reducing them to semi-formal systems E_α^0 , presented and analyzed by R\"uede in [15]. Next we extend each M-theory to a corresponding M^\dagger -theory by adding the axiom $\neg T\downarrow(U, \xi)$, where ξ is the previously determined upper bound of the M-theory and argue that ξ is still an upper bound of the corresponding M^\dagger -theory. Finally, we give embeddings of the ID^* -theories into the M^\dagger -theories, namely ID_α^* into $(M_\alpha + ACA)_0^\dagger$, $ID_{\alpha+1}^*\uparrow$ into $(M_\alpha + \Sigma_1^1\text{-DC})_0^\dagger$ and $ID_\lambda^*\uparrow$ into $(M_\lambda\uparrow)^\dagger$ if λ is a limit.

7.1. Upper bounds for $(M_\alpha + ACA)_0$, $(M_\alpha + \Sigma_1^1\text{-DC})_0$ and $M_\lambda\uparrow$. For each $\alpha < \Phi_0$, the theory M_α is formulated in the language $L_2(D)$ which extends L_2 by the unary relation symbol D . Formulas of $L_2(D)$ that do not contain bound set variables are called elementary. To simplify the notation, we write $t \in D$ for $D(t)$, $t \in D_s$ for $\langle t, s \rangle \in D$ and $X \in D_s$ is to abbreviate the formula $\exists x(X = (D_s)_x)$, where again, $X = (D_s)_t$ is short for $\forall x(x \in X \leftrightarrow \langle x, t \rangle \in D_s)$. The expression $t \in X_{\triangleleft a}$ stands for $t = \langle (t)_0, (t)_1 \rangle \wedge (t)_1 \triangleleft a \wedge t \in X$ and $t \in D_{\triangleleft a}$ is defined accordingly. $X \in D_{\triangleleft a}$ is read as $(\exists b \triangleleft a)(X \in D_b)$. The relativization A^{D_a} of an $L_2(D)$ formula to D_a is A for an elementary A , $(\forall xA(X))^{D_a} := \forall xA^{D_a}((D_a)_x)$ and $(\exists xA(X))^{D_a}$ is $\exists xA^{D_a}((D_a)_x)$. Relativizations to $D_{\triangleleft a}$ are defined analogously. Observe that if A is an $L_2(D)$ formula without free set variables, then A^{D_a} is a formula of $L_1(D)$. Finally, $Ax_{\Sigma_1^1\text{-AC}}$ denotes the finite axiomatization of Σ_1^1 -AC given in [14], namely the conjunction of the formulas listed below:

- (i) $\forall X, Y \exists Z(Z = X \oplus Y)$,
- (ii) $\forall e, z, Z \exists Y \forall x[x \in Y \leftrightarrow \pi_1^0(Z, e, x, z)]$,
- (iii) $\forall e, z, Z[\forall x \exists X \pi_2^0(X, Z, e, x, z) \rightarrow \exists Y \forall x \pi_2^0((Y)_x, Z, e, x, z)]$,

where π_k^0 is a universal Π_k^0 formula of L_2 of the appropriate arity and $X \oplus Y$ denotes the set $\{\langle x, 0 \rangle : x \in X\} \cup \{\langle x, 1 \rangle : x \in Y\}$.

The idea is that D constitutes an α -hierarchy of models of Σ_1^1 -AC, i.e., for all ordinal notations $a \triangleleft \alpha$, we have that D_a is a model of Σ_1^1 -AC and $D_{\triangleleft a} \in D_a$. Note however, that $[\forall x \exists X A(X, x) \rightarrow \exists Y \forall x A((Y)_x, x)]^{D_a}$ holds only for Σ_1^1 formulas of L_2 , not for Σ_1^1 formulas of $L_2(D)$.

In order to have partial cut elimination at hand, we formulate the M-theories in a Tait-style calculus that extends the classical Tait-calculus (cf. [17]) by the non-logical axioms and rules of the M-theories. We let Γ, Δ, \dots range over finite sets of $L_2(D)$ formulas and write Γ, A for the union of Γ and $\{A\}$. For each $\alpha < \Phi_0$, the theory M_α consist of the following axioms and rules:

Basic axioms. For all finite sets Γ of $L_2(D)$ formulas, all elementary formulas A of $L_2(D)$ and all arithmetical formulas B of L_2 which are axioms for the primitive recursive functions and relations:

$$\Gamma, A, \neg A \quad \text{and} \quad \Gamma, B.$$

Propositional and quantifier rules. These include the usual Tait-style inference rules for the propositional connectives as well as number and set quantifiers.

D-axioms. For all finite sets Γ of $L_2(D)$ formulas:

$$\Gamma, a \triangleleft \alpha \rightarrow \exists X[X = D_a], \quad \Gamma, a \triangleleft \alpha \rightarrow (Ax_{\Sigma_1^1-AC})^{D_a} \quad \text{and} \quad \Gamma, a \triangleleft \alpha \rightarrow D_{\triangleleft a} \in D_a;$$

Transfinite induction. For all finite sets Γ of $L_2(D)$ formulas:

$$\Gamma, \text{TI}_{\triangleleft}(U, \max\{\alpha, \omega\}).$$

Cut rules. For all finite sets Γ of $L_2(D)$ formulas and all $L_2(D)$ formulas A :

$$\frac{\Gamma, A \quad \Gamma, \neg A}{\Gamma}$$

The formulas A and $\neg A$ are the cut formulas of this cut.

Note that the D-axioms imply that $U, V \in D_0$. For limit ordinals $\lambda < \Phi_0$, the theory M_λ is obtained by replacing the axioms for transfinite induction by the following restricted version:

Restricted transfinite induction. For all finite sets Γ of $L_2(D)$ formulas:

$$\Gamma, a \triangleleft \lambda \rightarrow (\forall X \in D_a) \text{TI}_{\triangleleft}(X, \lambda).$$

The theory $(M_\alpha + \text{ACA})_0$ extends M_α by axioms for arithmetical comprehension and $(M_\alpha + \Sigma_1^1\text{-DC})_0$ extends $(M_\alpha + \text{ACA})_0$ by rules that imply all instances of dependent choice for Σ_1^1 formulas of L_2 :

Arithmetical comprehension. For all finite sets Γ of $L_2(D)$ formulas and all arithmetical L_2 formulas A :

$$\Gamma, \exists X[\forall x(x \in X \leftrightarrow A(x))].$$

Dependent choice. For all finite sets Γ of $L_2(D)$ formulas and all arithmetical L_2 formulas A :

$$\frac{\Gamma, \forall X \exists Y A(X, Y)}{\Gamma, \exists Z[(Z)_0 = W \wedge \forall x A((Z)_x, (Z)_{x+1})]}.$$

The formulas mentioned beside Γ in an axiom or the conclusion of a rule are called main formulas. Note that due to the axiom about transfinite induction, induction along the natural numbers for sets is available in $(M_\alpha + \text{ACA})_0$ and $(M_\alpha + \Sigma_1^1\text{-DC})_0$.

To apply the machinery developed by R\"ude in [15], we aim to embed our M-theories into a semi-formal systems E_α^0 , that we introduce later. In a first step, we eliminate the comprehension and dependent choice part of $(M_\alpha + \Sigma_1^1\text{-DC})_0$ and the comprehension part of $(M_\alpha + \text{ACA})_0$. For that purpose we introduce for each $\alpha < \Phi_0$ a semi-formal system RA_α , which is essentially an extension of RA^* of Sch\"utte [16] by the D-axioms for M_α .

Also the system RA_α is formulated in a Tait-style calculus. The language $\text{L}_{\text{RA}_\alpha}$ of RA_α is the language $L_2(D)$, where the set variables X, Y, Z, \dots are replaced by set variables $X^\beta, Y^\beta, Z^\beta, \dots$ for each ordinal $\beta < \alpha$. In RA_α we have set terms, which we define inductively together with the formulas of $\text{L}_{\text{RA}_\alpha}$:

- (i) Each set variable X^β is a set term.
- (ii) If $A(u)$ is a formula of $\text{L}_{\text{RA}_\alpha}$, then $\{x : A(x)\}$ is a set term.
- (iii) $[\sim]D(t)$, $[\sim]U(t)$, $[\sim]V(t)$ and $[\sim]R(\vec{t})$ are formulas of $\text{L}_{\text{RA}_\alpha}$, where R is a primitive recursive relation symbol.
- (iv) If t is a number term and T a set term, then $[\sim](t \in T)$ is formula of $\text{L}_{\text{RA}_\alpha}$.
- (v) The formulas of $\text{L}_{\text{RA}_\alpha}$ are closed under $\wedge, \vee, \forall x, \exists x, \forall X^\beta, \exists X^\beta$ for $\beta > 0$.

The level of a set term T and the level of a formula A of L_{RA_α} is defined by

$$\text{lev}(T) := \max\{0, \beta : X^\beta \text{ occurs in } T\} \text{ and } \text{lev}(A) := \max\{0, \beta : X^\beta \text{ occurs in } A\}.$$

The rank $\text{rk}(A)$ of a formula A of L_{RA_α} is inductively defined as follows: If A contains no set terms, then $\text{rk}(A) := 0$. Otherwise, letting \mathcal{Q} range over $\{\forall, \exists\}$:

- (i) For each set variable X^β , $\text{rk}(t \in X^\beta) := \text{rk}(t \notin X^\beta) := \max\{1, \omega \cdot \beta\}$,
- (ii) $\text{rk}([\sim](s \in \{x : A(x)\})) := \text{rk}(A(0)) + 1$,
- (iii) $\text{rk}(A \vee B) = \text{rk}(A \wedge B) := \max\{\text{rk}(A), \text{rk}(B)\} + 1$,
- (iv) $\text{rk}(\mathcal{Q}x A(x)) := \text{rk}(A(0)) + 1$,
- (v) $\text{rk}(\mathcal{Q}X^\beta A(X^\beta)) := \max\{\omega \cdot \text{lev}(\forall X^\beta A(X^\beta)), \text{rk}(A(X^0)) + 1\}$.

Notice that $\text{rk}(A) = \text{rk}(\neg(A))$. Also, if $\text{lev}(A) = \gamma$ and $\text{lev}(T) < \gamma$, then we have $\omega \cdot \gamma \leq \text{rk}(A) < \omega(\gamma + 1)$ and $\text{rk}(A(T)) < \text{rk}(\mathcal{Q}X^\gamma A(X^\gamma))$. These properties lead to the partial cut elimination Lemma 7.1.

The semi-formal system RA_α is formulate in the language L_{RA_α} . The formulas of RA_α are the closed formulas of L_{RA_α} . Thereby we consider the variable x to occur bound in the set term $\{x : A(x)\}$ and the formula $t \in \{x : A(x)\}$. In order to state the axioms and rules of RA_α , we assign to each closed number term t of L_1 its value t^N in the standard model. The *true literals* of L_1 are the closed literals of L_1 that evaluate to true in the standard model. The axioms and rules of RA_α are listed below.

Logical axioms. For all finite sets Γ of RA_α formulas, all set variables X^β , all true literals A of L_1 and all closed number terms s, t with $s^N = t^N$:

$$\Gamma, A \quad \text{and} \quad \Gamma, t \in X^\beta, s \notin X^\beta \quad \text{and} \quad \Gamma, t \in D, s \notin D.$$

Set term rules. For all finite sets Γ of RA_α formulas, all formulas A of RA_α and all closed number terms t :

$$\frac{\Gamma, A(t)}{\Gamma, t \in \{x : A(x)\}}, \quad \frac{\Gamma, \neg A(t)}{\Gamma, t \notin \{x : A(x)\}}.$$

Quantifier rules. For all finite sets Γ of RA_α formulas, all formulas A of RA_α , all closed number terms t and all set terms T :

$$\frac{\Gamma, A(t)}{\Gamma, \exists x A(x)}, \quad \frac{\Gamma, A(s) \text{ for all closed number terms } s}{\Gamma, \forall x A(x)},$$

$$\frac{\Gamma, A(T) \text{ and } \text{lev}(T) < \beta}{\Gamma, \exists X^\beta A(X^\beta)}, \quad \frac{\Gamma, A(T) \text{ for all set terms with } \text{lev}(T) < \beta}{\Gamma, \forall X^\beta A(X^\beta)}.$$

D-axioms. For all finite sets Γ of RA_α formulas and all closed number terms $t \triangleleft \alpha$:

$$\Gamma, (Ax_{\Sigma_1} \text{-AC})^{D_t} \quad \text{and} \quad \Gamma, D_{\triangleleft t} \in D_t.$$

Rules for \wedge and \vee and cut rules. The usual Tait-style rules for \wedge and \vee as well as the cut rules.

Observe that the D-axioms imply the existence of closed number terms s and t , such that $U = (D_0)_t$ and $V = (D_0)_s$. Also partial cut elimination is available:

LEMMA 7.1. We have for all finite sets Γ of RA_α formulas and all ordinals $\rho > 0$:

- (i) $RA_\alpha \mid_{\rho+1}^\beta \Gamma \implies RA_\alpha \mid_\rho^{\omega^\beta} \Gamma$,
- (ii) $RA_\alpha \mid_{1+\gamma+\omega^\rho}^\beta \Gamma \implies RA_\alpha \mid_{1+\gamma}^{\varphi\rho\beta} \Gamma$.

If a finite set Γ of $L_2(D)$ formulas is provable in $(M_\alpha + ACA)_0$, then standard cut elimination techniques yield that it is already provable in $(M_\alpha + ACA)_0$ where the cut rule is restricted to cut formulas A that are either elementary or $[\neg]A$ is the main formula of an axiom for arithmetical comprehension. Such restricted derivations are denoted by $(M_\alpha + ACA)_0 \vdash_* \Gamma$.

LEMMA 7.2. *Let $\Gamma(\vec{X}, \vec{x})$ be a finite set of elementary formulas of $L_2(D)$ such that $(M_\alpha + ACA)_0 \vdash_* \Gamma(\vec{X}, \vec{x})$. Then there exists for all set terms \vec{S} of level 0 an ordinal $\beta < \omega \cdot \max\{\alpha, \omega\} + \omega$ such that for all closed number terms \vec{s} ,*

$$RA_\alpha \vdash_{<\omega}^\beta \Gamma(\vec{S}, \vec{s}).$$

PROOF. We proof the claim by induction on the length of the proof in $(M_\alpha + ACA)_0$. For a set term S of level 0, $RA_\alpha \vdash_{<\omega}^{\omega \cdot \max\{\alpha, \omega\} + \omega} TI(S, \max\{\alpha, \omega\})$, and similarly for the other elementary main formulas of an axiom of M_α . The only case that follows not directly from the I.H. is if the last inference was a cut with a cut formula of the form $\exists X \forall x [x \in X \leftrightarrow A(\vec{Y}, x)]$ for some arithmetical formula A , or $t \triangleleft \alpha \rightarrow \exists X [X = D_a]$. We just consider the first case: \forall -inversion and the I.H. yield that for arbitrary set terms \vec{S} of level 0 and $T := \{x : A(\vec{S}, x)\}$, there is a $\beta < \omega \cdot \max\{\alpha, \omega\} + \omega$ such that for all closed number terms \vec{s} ,

$$RA_\alpha \vdash_{<\omega}^\beta \Gamma(\vec{S}, \vec{s}), \neg \forall x [x \in T \leftrightarrow A(\vec{S}, x)].$$

Because of $RA_\alpha \vdash_{<\omega}^{\omega \cdot \max\{\alpha, \omega\} + \omega} \forall x [x \in T \leftrightarrow A(\vec{S}, x)]$, a cut yields the claim. \dashv

Now we move to the theory $(M_\alpha + \Sigma_1^1\text{-DC})_0$. Cantini has shown in [5] that there is an asymmetric interpretation of $\Sigma_1^1\text{-DC}_0$ into $\Pi_0^1\text{-CA}_{<\omega^\omega}$. The same proof allows to perform an asymmetric interpretation of the theory $(M_\alpha + \Sigma_1^1\text{-DC})_0$ into $(M_\alpha + \text{Hier}_{<\omega^\omega}^J)_0$, which extends M_α by an axiom asserting the existence of the jump-hierarchy above any set X along an initial segment of \triangleleft of ordertype less than ω^ω :

$$(\text{Hier}_{<\omega^\omega}^J) \quad \Gamma, \exists F \text{Hier}^J(F, X, \beta),$$

for each ordinal $\beta < \omega^\omega$ and finite sets Γ of $L_2(D)$ formulas, where $\text{Hier}^J(F, X, a)$ denotes the formula

$$(\forall b \triangleleft a)[(F)_b = \{\langle x, \langle c, e \rangle \rangle : \pi_1^0(((F)_{\triangleleft b})_c, X, e, x)\}],$$

expressing that F constitutes a jump-hierarchy above X along $a \in \text{Field}(\triangleleft)$. For more details on this particular definition of the jump-hierarchy, we refer to [14].

LEMMA 7.3. $(M_\alpha + \Sigma_1^1\text{-DC})_0$ and $(M_\alpha + \text{Hier}_{<\omega^\omega}^J)_0$ prove the same Π_2^1 formulas of L_2 .

If $(M_\alpha + \text{Hier}_{<\omega^\omega}^J)_0 \vdash \Gamma$, then there is already a derivation $(M_\alpha + \text{Hier}_{<\omega^\omega}^J)_0 \vdash_* \Gamma$ which only uses the cut rule for elementary formulas and main formulas of instances of arithmetical comprehension and $(\text{Hier}_{<\omega^\omega}^J)$. For each $\beta, \gamma < \omega^\omega$ and each set terms S with level $\text{lev}(S) = \gamma$, there is a set term T of level $< \omega^\omega$ such that

$$RA_\alpha \vdash_{<\omega^\omega}^{\omega^\omega} \text{Hier}^J(T, S, \beta).$$

Similar as before, we obtain the following lemma.

LEMMA 7.4. Let $\Gamma(\vec{X}, \vec{x})$ be a finite set of elementary formulas of $L_2(D)$ such that $M_\alpha + (\text{Hier}_{<\omega^\omega}^J) \vdash_* \Gamma(\vec{X}, \vec{x})$. Then there exists for all set terms \vec{S} of level less than ω^ω an ordinal $n < \omega$ such that for all closed number terms \vec{s}

$$RA_\alpha \vdash \frac{<\omega \cdot \alpha + \omega^n}{<\omega^n} \vdash \Gamma(\vec{S}, \vec{s}).$$

Collecting the previous results and applying partial cut elimination for RA_α yields the following:

LEMMA 7.5. Suppose that A is a sentence of L_1 . Then we have:

- (i) $(M_\alpha + \text{ACA})_0 \vdash A \implies RA_\alpha \vdash \frac{<\varepsilon_0(\alpha)}{1} A$,
- (ii) $(M_\alpha + \Sigma_1^1\text{-DC})_0 \vdash A \implies RA_\alpha \vdash \frac{<(\alpha \uparrow 1)}{1} A$.

Next, we want to reduce RA_α to the semi-formal system E_α^0 . Basically, E_α^0 corresponds to the first order part of RA_α . Due to Rüede's results in [15], a prove of an L_1 formula A in E_α^0 yields a cut-free derivation of A in E_α^0 , which corresponds to a derivation of A in PA^* , a Tait-style reformulation of Peano Arithmetic PA with ω -rule.

For the reader's convenience, we restate Rüede's system E_α^0 . The language of E_α^0 is the extension of L_1 by unary relation symbols D_β^0 and $D_{<\gamma}^0$ for each $\beta < \alpha$ and $\gamma \leq \alpha$. The formulas of E_α^0 are the formulas of the language of E_α^0 that do not contain free number variables.

The ontological axioms and rules of E_α^0 state that for $\beta < \alpha$, D_β^0 contains only pairs, i.e., if the closed number term t is an element of D_β^0 , then its value is $\langle m, n \rangle$ for some natural numbers m, n . This expresses that m is an element of the set with code n in D_β^0 . The closure axioms and rules express that for all $\beta < \alpha$, D_β^0 is a model of $\Sigma_1^1\text{-AC}$ and that $D_{<\beta}^0 \in D_\beta^0$.

Logical axioms of E_α^0 . For all finite sets Γ of E_α^0 formulas, all true literals A of L_1 and all closed number terms s, t with identical values and all ordinals $\beta < \alpha, \gamma \leq \alpha$:

$$\Gamma, A \quad \text{and} \quad \Gamma, t \in D_\beta^0, s \notin D_\beta^0 \quad \text{and} \quad \Gamma, t \in D_{<\gamma}^0, s \notin D_{<\gamma}^0.$$

Ontological axioms and rules of E_α^0 . For all finite sets Γ of formulas of E_α^0 , all $\beta \leq \alpha, \gamma < \beta$, all closed number terms r, s, t such that r is not a pair, s is a pair but $(s)_0$ is not a pair and $\beta \sqsubseteq (s)_1$, and t is a pair and $\gamma = (t)_1$,

$$\Gamma, r \notin D_{<\beta}^0, \quad \Gamma, s \notin D_{<\beta}^0, \quad \frac{\Gamma, (t)_0 \in D_\gamma^0}{\Gamma, t \in D_{<\beta}^0}, \quad \frac{\Gamma, (t)_0 \notin D_\gamma^0}{\Gamma, t \notin D_{<\beta}^0}.$$

D-axioms of E_α^0 . For all finite sets Γ of E_α^0 formulas, all closed number terms e, r, s, t and all ordinals $\beta < \alpha$:

$$\Gamma, \exists k[(D_\beta^0)_k = (D_\beta^0)_t \oplus (D_\beta^0)_s],$$

$$\Gamma, \exists k[\forall x(x \in (D_\beta^0)_k \leftrightarrow \pi_1^0((D_\beta^0)_t, D_{<\beta}^0, e, x, r))].$$

D-rules of E_α^0 . For all finite sets Γ of E_α^0 formulas, all closed number terms e, r, s, t and all ordinals $\beta < \alpha$:

$$\frac{\Gamma, \forall x \exists k \pi_2^0((D_\beta^0)_k, (D_\beta^0)_t, D_{<\beta}^0, e, x, r)}{\Gamma, \exists k \forall x \pi_2^0(((D_\beta^0)_k)_x, (D_\beta^0)_t, D_{<\beta}^0, e, x, r)}.$$

Propositional rule, rules for the first order quantifiers and cut rules. These are the rules for RA_α adapted to the language of E_α^0 .

For a precise definition of the rank of formulas of E_α^0 we refer to definition 11 and the subsequent paragraph in [15]. We just try to capture the general idea: For example, if $\beta < \alpha$ and $A(X)$ is a formula of L_2 with exactly the displayed set variable free, then $(\forall X \in D_\beta^0)A(X)$, $(\exists X \in D_{<\beta}^0)A(X)$, $(\forall X \in D_\beta^0)A(X)$ and $(\exists X \in D_{<\beta}^0)A(X)$ are formulas of the language of E_α^0 of rank zero. Further, formulas of the language of E_α^0 of rank zero are closed under number quantification and propositional connectives. If $\beta + 1 = \alpha$ and t is a closed term that is not a pair, then $t \in D_\beta^0$ has rank 1. Also $t \in D_{<\alpha}^0$ has rank 1. Moreover, the rank of an E_α^0 formula is always finite and the rank of all main formulas of axioms of E_α^0 is zero.

LEMMA 7.6. *For all natural numbers $n > 0$ we have:*

$$E_\alpha^0 \mid_{\frac{\beta}{n+1}} \Gamma \implies E_\alpha^0 \mid_{\frac{2^\beta}{n}} \Gamma.$$

A closed $L_1(D)$ formula A is translated to an formula $A^{D_{<\alpha}^0}$ of E_α^0 by simply replacing D by $D_{<\alpha}^0$ in A . Now the following lemma, which corresponds to c) of Theorem 20 in [15], yields an embedding of RA_α into E_α^0 .

LEMMA 7.7. *Let $\alpha < \Phi_0$ and Γ a finite set of closed formulas of $L_1(D)$. Then we have:*

$$RA_\alpha \mid_{\Gamma}^\delta \Gamma \implies E_\alpha^0 \mid_{\frac{<\omega \cdot \delta}{<\omega}} \Gamma^{D_{<\alpha}^0}.$$

PROOF. As in [15] this follows by an induction on δ . Since in our case, the D-axioms and D-rules of E_α^0 are syntactically different from the D-axioms of RA_α , we have to use that for a closed number term $t < \alpha$ with $t = \beta$, $E_\alpha^0 \mid_{\frac{<\omega}{0}} (D_{<\alpha}^0)_t = D_\beta^0$ and $E_\alpha^0 \mid_{\frac{<\omega}{<\omega}} (AX_{\Sigma_1} - AC)^{(D_{<\alpha}^0)_t}$. \dashv

For $M_\lambda \upharpoonright$, the detour over RA_λ is not necessary. We first embed $M_\lambda \upharpoonright$ into E_λ^+ , the extension of E_λ^0 by axioms

$$\Gamma, (\forall X \in D_\beta) TI_{<}(X, \lambda),$$

for each $\beta < \lambda$ and all finite sets Γ of $L_1(D)$ formulas.

LEMMA 7.8. *Let $\lambda < \Phi_0$ and $\Gamma(\vec{u})$ a finite set of formulas of $L_1(D)$. Then there exists for each $n \in \mathbb{N}$ an $n' \in \mathbb{N}$, such that we have for all closed number terms \vec{t} ,*

$$M_\lambda \upharpoonright \mid_{\Gamma}^n \Gamma(\vec{u}) \implies E_\lambda^+ \mid_{\Gamma}^{n'} \Gamma^{D_{<\lambda}^0}(\vec{t}).$$

PROOF. By induction on the proof length one first shows that $E_\lambda^+ \mid_{\frac{<\omega}{<\omega}} \Gamma^{D_{<\lambda}^0}(\vec{t})$. Then cut elimination in E_λ^+ yields the claim. \dashv

Theorem 26 in [15] tells us at what cost we can reduce $E_{\beta+\omega^{1+\rho}}^0$ to E_β^0 . Given this result, the reduction of $E_{\beta+\omega^{1+\rho}}^+$ to E_β^0 does not cause additional difficulties.

LEMMA 7.9. *Assume that $\lambda = \beta + \omega^{1+\rho} < \Phi_0$, $\lambda \upharpoonright 0 = \lambda_0 + \omega$ and k is the least natural number such that $\varphi_{1\rho}(\lambda_0 + k) > \lambda$. Further, suppose that $E_\lambda^+ \mid_{\Gamma}^n \Gamma$ for a finite set Γ of formulas of $E_{\beta+\omega^{1+\rho}}^0$ of rank 0. Then the following holds: If each formula in Γ is a formula of $E_{\beta+\xi}^0$, for some $\xi' < \xi < \omega^{1+\rho}$, then*

$$E_{\beta+\xi}^0 \mid_{\frac{\varphi_{1\rho}(\lambda_0+k+n)}{1}} \Gamma.$$

PROOF. The lemma is proved as Theorem 26 in [15] by main induction on ρ and side induction on n . For $n = 0$, observe that $\gamma < \xi < \omega^{1+\rho}$ yields

$$E_{\beta+\xi}^0 \mid \frac{<\varphi 1 \rho(\lambda_0+k)}{1} (\forall X \in D_\gamma^0) \text{TI}_\triangleleft(X, \lambda). \quad \dashv$$

Now we conclude that the theories $(M_\alpha + \text{ACA})_0$, $(M_\alpha + \Sigma_1^1\text{-DC})_0$ and $M_\lambda \upharpoonright$ have the desired upper bounds.

THEOREM 7.10 (Upper Bounds). *For $\alpha < \Phi_0$ and limit ordinals $\lambda < \Phi_0$, we have:*

$$|(M_\alpha + \text{ACA})_0| \leq \sigma(\alpha), \quad |(M_\alpha + \Sigma_1^1\text{-DC})_0| \leq \sigma \upharpoonright (\alpha + 1) \quad \text{and} \quad |M_\lambda \upharpoonright| \leq \sigma \upharpoonright (\lambda).$$

PROOF. Suppose that $\alpha = \omega^{1+\alpha_n} + \dots + \omega^{1+\alpha_1} + m$, where $\alpha_n \geq \dots \geq \alpha_1$ and $m < \omega$, and let δ and λ such that $\alpha = \lambda + m$ and $\lambda = \delta + \omega^{1+\alpha_1}$. Further, assume that A is a sentence of L_1 . If $(M_\alpha + \text{ACA})_0 \vdash A$, then the Lemmas 7.5, 7.6 and 7.7 yield $E_\alpha^0 \mid \frac{<(\alpha|0)}{1} A$, so that applying m -times Corollary 21 in [15] gives $E_\lambda^0 \mid \frac{<(\alpha|m)}{1} A$. Now n -fold application of Theorem 26 in [15] confirms $E_0^0 \mid \frac{<\sigma(\alpha)}{1} A$. Similarly, if $(M_\alpha + \Sigma_1^1\text{-DC})_0 \vdash A$, then $E_\alpha^0 \mid \frac{<(\alpha \upharpoonright 1)}{1} A$, thus we obtain $E_\lambda^0 \mid \frac{<(\alpha \upharpoonright m+1)}{1} A$ and $E_0^0 \mid \frac{<\sigma \upharpoonright (\alpha+1)}{1} A$. Finally, if $M_\lambda \upharpoonright \vdash A$, then $E_\lambda^+ \mid \frac{<\omega}{1} A$ due to Lemma 7.8, therefore $E_\delta^0 \mid \frac{<\varphi 1 \alpha_n(\lambda \upharpoonright 0)}{1} A$ and $E_0^0 \mid \frac{<\sigma \upharpoonright (\lambda)}{1} A$. Cut-elimination in PA^* yields the claim. \dashv

7.2. Embedding the ID^* -theories into the M^\dagger -theories. Let M denote one of the M -theories and let ξ be the upper bound according to Theorem 7.10. Note that for $\beta < \xi$ the ordinal $\omega \cdot \beta$ is still less than ξ . By choice of ξ , we have that $\text{TI}_\triangleleft(V, \xi)$ is not provable in the theory M . Therefore, the theory M^\dagger , the extension of M by the axiom $\neg \text{TI}_\triangleleft(V, \xi)$, is consistent. Moreover, ξ is still an upper bound of M^\dagger : Assume that M^\dagger proves $\text{TI}_\triangleleft(U, \alpha)$ for a primitive recursive well-ordering \triangleleft . Thus $M \vdash \text{TI}_\triangleleft(V, \xi) \vee \text{TI}_\triangleleft(U, \alpha)$. The proof of Theorem 7.10 yields that

$$\text{PA}^* \mid \frac{<\xi}{0} \neg \text{Prog}_\triangleleft(V), \xi \in V, \text{TI}_\triangleleft(U, \alpha).$$

With Lemma 4 in Jäger and Probst [10] we conclude that also $\text{PA}^* \mid \frac{<\xi}{0} \text{TI}_\triangleleft(U, \alpha)$. Hence, by Schütte's boundedness theorem (cf. [16] or [12]) we obtain $\alpha < \xi$. Embedding the ID^* -theory with lower bound ξ into M^\dagger , then yields $\xi \leq |\text{ID}^*| \leq |M^\dagger| \leq \xi$.

To embed the ID^* -theories into the M^\dagger -theories, we show that these theories prove the existence of α -hierarchies of models of $\Sigma_1^1\text{-DC}$. Thereby, we make use of so-called *pseudo-hierarchy arguments*. For second order arithmetic, this method is described in Simpson [18] in extenso and a typical application is given in Avigad [2]. In subsystems of second order arithmetic comprising (ACA), the existence of a pseudo-hierarchy follows from the fact that being a well-ordering is not expressible by a Σ_1^1 formula of L_2 . However, this method does not provide uniform pseudo-hierarchies.

We apply a more general method to obtain pseudo-hierarchies: Due to the axiom $\neg \text{TI}_\triangleleft(V, |M|)$ of the theory M_α^\dagger one can prove that $\{a \in \text{Field}(\triangleleft) : \forall Z \text{TI}_\triangleleft(Z, a)\}$ is not a set. The existence of [uniform] pseudo-hierarchies is then derived from this observation. Using this method, the application of pseudo-hierarchy arguments is no longer limited to second order analysis and can be applied in the context of explicit mathematics and admissible set theory as well; cf. [13, 14]. In the sequel,

$\text{Wo}_{\triangleleft}(a)$ is to abbreviate $\forall Z \text{TI}_{\triangleleft}(Z, a)$ and $\text{Hier}^J(U, V, u)$ is the formula defined above Lemma 7.3.

LEMMA 7.11. *The following is provable in $(M_{\alpha} + \text{ACA})_0^{\dagger}$:*

- (i) $\{a : \text{Wo}_{\triangleleft}(a)\}$ is not a set.
- (ii) If $b \triangleleft \alpha$, then $\{a : \text{Wo}_{\triangleleft}^{D_b}(a)\} \notin D_b$.
- (iii) If $b+1 \triangleleft \alpha$ and $X \in D_b$, then $\{a : \text{Wo}_{\triangleleft}^{D_{b+1}}(a)\} \subsetneq \{a : (\exists F \in D_b) \text{Hier}^J(F, X, a)\}$.

PROOF. Suppose for a moment that $S := \{a : \text{Wo}_{\triangleleft}(a)\}$ is a set. Then the set $S_0 := \{a : (\forall b \in S)(\varphi 0ab \in S)\}$ is easily shown to be progressive w.r.t. \triangleleft , which in turn yields the progressivity of the set $S_1 := \{a : (\forall b \in S)(\varphi 1ab \in S)\}$. If $a = 0$, this is due to the progressivity of S_0 , otherwise assume that $(\forall a' \triangleleft a)(a' \in S_1)$ and show that $\text{Prog}_{\triangleleft}(\{b \in S : \varphi 1ab \in S\})$, which yields $a \in S_1$. Hence $a, b \in S$ implies $\varphi 1ab \in S$. In particular, $\alpha \in S$ yields $\sigma(\alpha) \in S$. A contradiction!

If we relativize the above argument to D_b , we obtain that $\sigma(\alpha) \in \{a : \text{Wo}_{\triangleleft}^{D_b}(a)\}$. Since $V \in D_b$, this contradicts $\neg \text{TI}_{\triangleleft}(V, \sigma(\alpha))$. Thus (ii) holds.

Because D_b is a model of $\Sigma_1^1\text{-AC}$, $\text{Wo}_{\triangleleft}^{D_{b+1}}(a)$ implies the existence of an $F \in D_b$ such that $\text{Hier}^J(F, X, a)$. Because $\{a : \text{Wo}_{\triangleleft}^{D_{b+1}}(a)\}$ is not a set in D_{b+1} , the inclusion is proper. \dashv

LEMMA 7.12. *The following is provable in $(M_{\alpha+1} + \text{ACA})_0^{\dagger}$: For each set $X \in D_{\alpha}$ there exists a model M of $\Sigma_1^1\text{-DC}$ with $X \in M$.*

PROOF. Fix $X \in D_{\alpha}$. Since D_{α} is a model of $\Sigma_1^1\text{-AC}$, one easily proves that

$$\forall a[\text{Wo}_{\triangleleft}(a) \rightarrow (\exists F \in D_{\alpha}) \text{Hier}^J(F, X, a) \wedge \text{Wo}_{\triangleleft}^{D_{\alpha}}(a)].$$

Because $\{a : \text{Wo}_{\triangleleft}(a)\}$ is not a set, there exists a $b \in \text{Field}(\triangleleft)$ and an $F \in D_{\alpha}$ such that

$$\neg \text{Wo}_{\triangleleft}(b) \wedge \text{Hier}^J(F, X, b) \wedge \text{Wo}_{\triangleleft}^{D_{\alpha}}(b).$$

Thus, there exists a non-empty, upward closed $K \subseteq \text{Field}(\triangleleft)$ without a \triangleleft -least element and with $b \in K$. Surely, $\text{Wo}_{\triangleleft}(a)$ implies $(\forall x \in K)(a \triangleleft x)$, subsequently abbreviated by $a \triangleleft K$. Next we consider the sets

$$M := M_{\triangleleft K}^F := \{\langle x, \langle c, e \rangle \rangle : c \triangleleft K \wedge \langle x, \langle c, e \rangle \rangle \in (F)_{c+1}\},$$

$$M_{\triangleleft d}^F := \{\langle x, \langle c, e \rangle \rangle : c \triangleleft d \wedge \langle x, \langle c, e \rangle \rangle \in (F)_{c+1}\}, \text{ for each } d \in \text{Field}(\triangleleft),$$

and prove that M is a model of $\Sigma_1^1\text{-DC}$. We just show that M satisfies $(\Sigma_1^1\text{-DC})$, that M is a model of ACA follows from standard results concerning the jump-hierarchy. So, let $A(U, V)$ be an arithmetical formula of L_2 and assume that

$$(1) \quad (\forall X \in M)(\exists Y \in M)A(X, Y).$$

If $X \in M$, then there exists an index a such that $X = (M)_a$. The definition of M implies that a is of the form $\langle c, e \rangle$, where e is a natural number and c an element of the field of \triangleleft . Now, we set

$$I := \{\langle c, e \rangle : e \in \mathbb{N} \wedge c \in \text{Field}(\triangleleft)\},$$

and order I by $<_I$, letting $\langle c, e \rangle <_I \langle d, e' \rangle$ if $c \triangleleft d$, or $c = d$ and $e <_{\mathbb{N}} e'$. Note, that $\langle c, e \rangle \in I$ and $\neg(c \triangleleft K)$ implies $(M)_{\langle c, e \rangle} = \emptyset$. Therefore, (1) becomes equivalent

to the formula $(\forall y \in I)(\exists z \in I)A((M)_y, (M)_z)$. Moreover, for each $y \in I$, the set $\{z \in I : A((M)_y, (M)_z)\}$ has a $<_I$ -least element. To see this, observe that

$$S_1 := \{z \in I : A((M)_y, (M)_z)\} \subseteq \{z \in I : A((M_{<_I}^F)_y, (M_{<_I}^F)_z)\} =: S_2.$$

Since $S_2 \in D_\alpha$ and $\text{Wo}_{<_I}^{D_\alpha}(b)$, it has a $<_I$ -least element. This is also the minimum of the set S_1 , because $z \in S_1$, $y \in S_2$ and $y <_I z$ yields already $y \in S_1$. Therefore, we conclude that $(\forall y \in I)(\exists! z \in I)A'(M, y, z)$, where A' is an arithmetical formula of L_2 expressing that z is the least index w.r.t. our index ordering $<_I$, such that $A((M)_y, (M)_z)$ holds.

Next, we fix an index $w \in I$ with $(w)_0 < K$ and show that there exists a choice sequence $Z \in M$, such that $(Z)_0 = (M)_w$ and $\forall n A((Z)_n, (Z)_{n+1})$. First, we look for initial segments of such a choice sequence. In the present setting, this is a finite sequence s , (respectively a natural number of the form $\langle x_1, \dots, x_n \rangle$) of indices such that

$$\text{ChSeq}_{A'}(M, s, w, n) := \text{lh}(s) = n + 1 \wedge (s)_0 = w \wedge (\forall m < n) A'(M, (s)_m, (s)_{m+1}).$$

Assumption (1) allows us to prove by set induction that $\forall n \exists! s \text{ChSeq}_{A'}(M, s, w, n)$. Further, $c < K$ implies $(M)_{\langle c, e \rangle} = (M_{<_I}^F)_{\langle c, e \rangle}$ for each $a \in K$, thus the set

$$\{a < b : \forall n \exists s \text{ChSeq}_{A'}(M_{<_I}^F, s, w, n)\}$$

is not empty. Moreover, it is in D_α , so it has a least element a_0 . Since $a_0 < K$,

$$Z := \{\langle x, n \rangle : \exists s [\text{ChSeq}_{A'}(M_{<_{a_0}}^F, s, w, n) \wedge x \in (M_{<_{a_0}}^F)_{(s)_n}]\}$$

is a set in M and serves as a witness for our sought for choice sequence. \dashv

The model constructed in the previous proof is not uniform in the sense that we only know about the existence of a set K without a $<_I$ -least element, but cannot explicitly define it. However, if $X \in D_b$ and $b + 1 < \alpha$, then we can construct in $(M_\alpha + \text{ACA})_0^\dagger$ a uniform model M of Σ_1^1 -DC above X . More precisely: If $X \in D_b$, then we call the set

$$M_{\Sigma_1^1\text{-DC}}(X, b) := \{x : \text{Wo}_{<_I}^{D_{b+1}}(c) \wedge (\exists F \in D_b)[\text{Hier}^J(F, X, c + 1) \wedge x \in M_{<_I}^F]\},$$

the uniform model of Σ_1^1 -DC above $X \in D_b$. $M_{<_I}^F$ is as defined in the above proof. Because $\text{Wo}_{<_I}^{D_{b+1}}(c)$ and $X \in D_b$ imply that there is exactly one $F \in D_b$ satisfying $\text{Hier}^J(F, X, c + 1)$, the set $M_{\Sigma_1^1\text{-DC}}(X, b)$ is clearly unique.

LEMMA 7.13. *The following is provable in $(M_\alpha + \text{ACA})_0^\dagger$: If $b + 1 < \alpha$ and $X \in D_b$, then $M_{\Sigma_1^1\text{-DC}}(X, b)$ is a model of Σ_1^1 -DC with $X \in M_{\Sigma_1^1\text{-DC}}(X, b)$. If $b + 2 < \alpha$ then $M_{\Sigma_1^1\text{-DC}}(X, b) \in D_{b+2}$.*

PROOF. Assume that $X \in D_b$. Using (iii) of Lemma 7.11, we can find an $F \in D_b$ and a notation $d \in \text{Field}(<_I)$ such that $\neg \text{Wo}_{<_I}^{D_{b+1}}(d)$ and $\text{Hier}^J(F, X, d)$. Further,

$$K := \{a \in \text{Field}(<_I) : \neg \text{Wo}_{<_I}^{D_{b+1}}(a)\},$$

is a non-empty subset of $\text{Field}(<_I)$ without a $<_I$ -least element that contains d . Now $M_{\Sigma_1^1\text{-DC}}(X, b)$ becomes the set $M_{<_K}^F$ according to the definition in the previous proof, and we may continue as there. \dashv

Next we want to speak about [uniform] a -hierarchies of models of Σ_1^1 -DC. For this purpose, we set:

$$\begin{aligned}\text{Hier}_{\Sigma_1^1\text{-DC}}(F, a) &: \iff (\forall b \triangleleft a)[(F)_b = M_{\Sigma_1^1\text{-DC}}((F)_{\triangleleft b}, 2b)], \\ \text{Hier}_{\Sigma_1^1\text{-DC}}(F, a) &: \iff (\forall b \triangleleft a)[(F)_{\triangleleft b} \in (F)_b \wedge (Ax_{\Sigma_1^1\text{-DC}})^{(F)_b}].\end{aligned}$$

Such hierarchies indeed exist and have the intended properties:

LEMMA 7.14. *The following is provable in $(M_\lambda)^\dagger$: If $a \triangleleft \lambda < \Phi_0$, then*

- (i) $(\exists F \in D_{2a})\text{Hier}_{\Sigma_1^1\text{-DC}}(F, a)$,
- (ii) $\forall F, G[\text{Hier}_{\Sigma_1^1\text{-DC}}(F, a) \wedge \text{Hier}_{\Sigma_1^1\text{-DC}}(G, a) \rightarrow (\forall b \triangleleft a)(F)_b = (G)_b]$.

PROOF. For this proof, we denote claim (i) by $C_1(a)$ and claim (ii) by $C_2(a)$.

Assume that $a \triangleleft \lambda$. Since $\{b \triangleleft a + 1 : (\exists G \in D_{2b})\text{Hier}_{\Sigma_1^1\text{-DC}}(G, b)\}$ is a set in $D_{2(a+1)}$, (i) and (ii) can be shown simultaneously by restricted transfinite induction.

So suppose $c \triangleleft b \triangleleft a + 1$ and that $C_1(c)$ and $C_2(c)$ hold. If b is a successor, then the induction step follows easily from the previous lemma. If b is a limit, then $2b = b$ and the I.H. yields that

$$F := \{\langle x, d \rangle : d \triangleleft b \wedge (\exists G \in (D_{\triangleleft b})_{2(d+1)})(\text{Hier}_{\Sigma_1^1\text{-DC}}(G, d+1) \wedge x \in (G)_d)\}$$

is an element of D_b satisfying $\text{Hier}_{\Sigma_1^1\text{-DC}}(F, b)$, thus $C_1(b)$. $C_2(b)$ follows from $C_1(b)$ and the I.H. \dashv

COROLLARY 7.15. For $\alpha < \Phi_0$, the following is provable in $(M_\alpha + \text{ACA})_0^\dagger$:

$$\exists F \text{Hier}_{\Sigma_1^1\text{-DC}}(F, \alpha).$$

PROOF. α can be written in the form $\omega \cdot \beta + n$ for some $n < \omega$. The claim follows now by (meta-) induction on n : If $n = 0$, the previous lemma and arithmetical comprehension yield that

$$F := \{\langle x, a \rangle : a \triangleleft \omega \cdot \beta \wedge (\exists G \in D_{2(a+1)})(\text{Hier}_{\Sigma_1^1\text{-DC}}(G, a+1) \wedge x \in (G)_a)\}$$

constitutes the sought for hierarchy. If $(M_{\omega \cdot \beta + n} + \text{ACA})_0^\dagger$ proves the existence of an hierarchy F such that $\text{Hier}_{\Sigma_1^1\text{-DC}}(F, \omega \cdot \beta + n)$, then $(M_{\omega \cdot \beta + n + 1} + \text{ACA})_0^\dagger$ proves that such a F exists already in $D_{\omega \cdot \beta + n}$. Then also $(F)_{\triangleleft b} \in D_{\omega \cdot \beta + n}$, and Lemma 7.12 tells us that there exists a model of Σ_1^1 -DC above $(F)_{\triangleleft b}$ in $D_{\omega \cdot \beta + n + 1}$. The existence of a hierarchy G satisfying $\text{Hier}_{\Sigma_1^1\text{-DC}}(G, \omega \cdot \beta + n + 1)$ follows. \dashv

Next we introduce fixed point hierarchies. For each operator form $\mathcal{A}(P, Q, p, u)$ of $L_1(P, Q)$ we say that H constitutes a fixed point hierarchy for \mathcal{A} along a w.r.t. G , if

$$\text{Hier}_{\text{Fix}}^{\mathcal{A}}(G, H, a) := \text{Hier}_{\Sigma_1^1\text{-DC}}(G, a) \wedge (\forall b \triangleleft a)[(H)_b = (\text{Fix}_{(H)_{\triangleleft b}, b}^{\mathcal{A}})^{(G)_b}],$$

holds.

LEMMA 7.16. For $\alpha < \Phi_0$, $(M_\alpha + \text{ACA})_0^\dagger$ proves:

- (i) $a \trianglelefteq \alpha \wedge \text{Hier}_{\Sigma_1^1\text{-DC}}(G, \alpha) \rightarrow \exists! H \text{Hier}_{\text{Fix}}^{\mathcal{A}}(G, H, a)$,
- (ii) $\text{Hier}_{\text{Fix}}^{\mathcal{A}}(G, H, \alpha) \wedge \text{Hier}_{\text{Fix}}^{\mathcal{A}}(G, H', \alpha) \rightarrow (\forall a \triangleleft \alpha)[(H)_a = (H')_a]$,
- (iii) $\text{Hier}_{\text{Fix}}^{\mathcal{A}}(G, H, \alpha) \rightarrow (\forall a \triangleleft \alpha)[(H)_{\triangleleft a} \in (G)_a]$.

PROOF. Transfinite induction and the definition of the fixed point hierarchy yields immediately the uniqueness assertion (ii). For (i), we assume that G satisfies $\text{Hier}_{\text{Fix}}^{\mathcal{A}}(G, \alpha)$, and show by induction that for each $a \triangleleft \alpha$,

$$(\exists! H \in (G)_a) \text{Hier}_{\text{Fix}}^{\mathcal{A}}((G)_{\triangleleft a}, H, a).$$

We just consider the successor case, the limit case is similar. So assume that there exists a unique $H \in (G)_b$ such that $\text{Hier}_{\text{Fix}}^{\mathcal{A}}((G)_{\triangleleft b}, H, b)$. With $H \in (G)_b$ we have also $(H)_{\triangleleft b} \in (G)_b$, which implies that $H' := (\text{Fix}_{(H)_{\triangleleft b}, b}^{\mathcal{A}})^{(G)_b}$ is an element of $(G)_{b+1}$. Now we obtain a fixed point hierarchy for \mathcal{A} along $b+1$. Claim (iii) follows from (i) and the definition of $\text{Hier}_{\text{Fix}}^{\mathcal{A}}(G, H, a)$. \dashv

Lemma 7.14 gives way to the following corollary.

COROLLARY 7.17. For a limit ordinal $\lambda < \Phi_0$, $(M_\lambda \upharpoonright)^\dagger$ proves:

$$(\forall a \triangleleft \lambda) \exists! G \exists! H [\text{Hier}_{\Sigma_1\text{-DC}}^!(G, a) \wedge \text{Hier}_{\text{Fix}}^{\mathcal{A}}(G, H, a)].$$

Finally, we see that a fixed point hierarchy H w.r.t. G is indeed suitable for an interpretation of the relation symbols $P^{\mathcal{A}}$ of ID_α^* and $\text{ID}_\alpha^* \upharpoonright$:

LEMMA 7.18. Given $\alpha < \Phi_0$, then for each operator form \mathcal{A} and each Π_1^1 formula $D(U, u)$ of L_2 , the following is provable in $(M_\alpha + \text{ACA})_0^\dagger$. If $\text{Hier}_{\text{Fix}}^{\mathcal{A}}(G, H, \alpha)$ holds, then also

- (i) $(\forall a \triangleleft \alpha) \forall x [x \in (H)_a \leftrightarrow \mathcal{A}((H)_a, (H)_{\triangleleft a}, x, a)],$
- (ii) $(\forall a \triangleleft \alpha) [\text{Cl}^{\mathcal{A}}(\{x : D((G)_{\triangleleft a}, x)\}, a) \rightarrow (H)_a \subseteq \{x : D((G)_{\triangleleft a}, x)\}].$

PROOF. By Lemma 3.4 we know that ACA proves

$$\forall x [x \in \text{Fix}_{Y, Y}^{\mathcal{A}} \leftrightarrow \mathcal{A}(\text{Fix}_{Y, Y}^{\mathcal{A}}, Y, x, y)].$$

For each $a \triangleleft \alpha$ we have that $(H)_{\triangleleft a} \in (G)_a$ and $(G)_a$ is a model of $\Sigma_1^1\text{-AC}$. Thus we can relativize Lemma 3.4 to $(G)_a$ and obtain

$$\forall x [x \in (\text{Fix}_{(H)_{\triangleleft a}, a}^{\mathcal{A}})^{(G)_a} \leftrightarrow \mathcal{A}((\text{Fix}_{(H)_{\triangleleft a}, a}^{\mathcal{A}})^{(G)_a}, (H)_{\triangleleft a}, x, a)].$$

By the definition of $(H)_a$ this is equivalent to (i). Claim (ii) is shown analogously by relativizing Theorem 4.1 to $(G)_a$. \dashv

Now we are ready to present an embedding of the ID^* -theories into the M^\dagger -theories: If we have a list $\mathcal{A}_1, \dots, \mathcal{A}_n$ of operator forms, then we write $\text{Hier}_{\text{Fix}}^{\vec{\mathcal{A}}}(\vec{H}, G, a)$ for

$$\text{Hier}_{\text{Fix}}^{\mathcal{A}_1}(H_1, G, a) \wedge \dots \wedge \text{Hier}_{\text{Fix}}^{\mathcal{A}_n}(H_n, G, a).$$

An L_{Fix} formula A is translated to an $L_2(D)$ formula A^* , A^* or A° by replacing each subformula of the form $t \in P^{\mathcal{A}}$, by either the formula $(t \in P^{\mathcal{A}})^*$, $(t \in P^{\mathcal{A}})^*$ or $(t \in P^{\mathcal{A}})^\circ$, depending on whether we embed ID_α^* into $(M_\alpha + \text{ACA})_0^\dagger$, $\text{ID}_{\alpha+1}^* \upharpoonright$ into $(M_\alpha + \Sigma_1^1\text{-DC})_0^\dagger$ or $\text{ID}_\lambda^* \upharpoonright$ into $(M_\lambda \upharpoonright)^\dagger$:

THEOREM 7.19. Let A be an L_{Fix} formula that contains exactly the set constants $P^{\vec{\mathcal{A}}}$. Then the following holds for each $\alpha < \Phi_0$ and each limit $\lambda < \Phi_0$:

$$\begin{array}{lll} \text{ID}_\alpha^* \vdash A & \implies & (M_\alpha + \text{ACA})_0^\dagger \vdash \text{Hier}_{\text{Fix}}^{\vec{\mathcal{A}}}(G, \vec{H}, \alpha) \rightarrow A^*, \\ \text{ID}_{\alpha+1}^* \upharpoonright \vdash A & \implies & (M_\alpha + \Sigma_1^1\text{-DC})_0^\dagger \vdash \text{Hier}_{\text{Fix}}^{\vec{\mathcal{A}}}(G, \vec{H}, \alpha) \rightarrow A^*, \\ \text{ID}_\lambda^* \upharpoonright \vdash A & \implies & (M_\lambda \upharpoonright)^\dagger \vdash A^\circ, \end{array}$$

where for $1 \leq i \leq n$,

$$\begin{aligned}(t \in \mathcal{P}^{\mathcal{A}_i})^* &:= t = \langle s, a \rangle \wedge a \triangleleft \alpha \wedge t \in H_i, \\ (t \in \mathcal{P}^{\mathcal{A}_i})^* &:= [t = \langle s, a \rangle \wedge a \triangleleft \alpha \wedge s \in (H_i)_a] \vee [t = \langle s, \alpha \rangle \wedge s \in \text{Fix}_{(H_i) \triangleleft \alpha, \alpha}^{\mathcal{A}_i}], \\ (t \in \mathcal{P}^{\mathcal{A}_i})^\circ &:= t = \langle s, a \rangle \wedge a \triangleleft \lambda \wedge (\exists G, H \in D_{2(a+2)}) \\ &\quad [\text{Hier}_{\Sigma_1^1\text{-DC}}^!(G, a+1) \wedge (\text{Hier}_{\text{Fix}}^{\mathcal{A}_i}(G, H, a+1) \wedge s \in (H)_a)].\end{aligned}$$

Since the existence of the fixed point hierarchies follows from Corollary 7.15 Lemma 7.16 and Corollary 7.17, we also obtain that corresponding theories prove the same L_1 formulas.

COROLLARY 7.20. For each $\alpha < \Phi_0$, each limit $\lambda < \Phi_0$ and each L_1 formula A , the following holds:

$$\begin{aligned}\text{ID}_\alpha^* &\vdash A \implies (\text{M}_\alpha + \text{ACA})_0^\dagger \vdash A, \\ \text{ID}_{\alpha+1}^* \upharpoonright &\vdash A \implies (\text{M}_\alpha + \Sigma_1^1\text{-DC})_0^\dagger \vdash A, \\ \text{ID}_\lambda^* \upharpoonright &\vdash A \implies (\text{M}_\lambda \upharpoonright)^\dagger \vdash A.\end{aligned}$$

The circle closes: In Section 6 we computed lower bounds for the ID^* -theories. In Section 7.1 we proved that these lower bounds are upper bounds for the M -theories and thus also for the M^\dagger -theories. Eventually, we managed to embed the ID^* -theories into the M^\dagger -theories. Summing up, we can state the following theorem:

THEOREM 7.21. For each $\alpha < \Phi_0$ we have:

$$|\text{ID}_\alpha^*| = |\widehat{\text{ID}}_\alpha| = \sigma(\alpha) \quad \text{and} \quad |\text{ID}_\alpha^* \upharpoonright| = \sigma \upharpoonright(\alpha).$$

REFERENCES

- [1] PETER ACZEL, *The strength of Martin-Löf's type theory with one universe*, Technical report, Department of Philosophy, University of Helsinki, 1977.
- [2] JEREMY AVIGAD, *On the relationship between ATR_0 and $\widehat{\text{ID}}_{<\omega}$* , this JOURNAL, vol. 61 (1996), no. 3, pp. 768–779.
- [3] WILFRIED BUCHHOLZ, SOLOMON FEFERMAN, WOLFRAM POHLERS, and WILFRIED SIEG, *Iterated inductive definitions and subsystems of analysis: Recent proof-theoretical studies*, Lecture Notes in Mathematics, vol. 897, Springer, 1981.
- [4] ANDREA CANTINI, *A note on a predicatively reducible theory of iterated elementary induction*, *Bollettino Unione Matematica Italiana*, vol. 4-B (1985), no. 6, pp. 413–430.
- [5] ———, *On the relationship between choice and comprehension principles in second order arithmetic*, this JOURNAL, vol. 51 (1986), pp. 360–373.
- [6] SOLOMON FEFERMAN, *Iterated inductive fixed-point theories: Application to Hancock's conjecture*, *The Patras Symposium* (G. Metakides, editor), North Holland, 1982, pp. 171–196.
- [7] ———, *Reflecting on incompleteness*, this JOURNAL, vol. 56 (1991), no. 1, pp. 1–49.
- [8] HARVEY FRIEDMAN, *Theories of inductive definition*, 1969, Unpublished notes.
- [9] GERHARD JÄGER, REINHARD KAHLE, ANTON SETZER, and THOMAS STRAHM, *The proof-theoretic analysis of transfinutely iterated fixed point theories*, this JOURNAL, vol. 64 (1999), no. 1, pp. 53–67.
- [10] GERHARD JÄGER and DIETER PROBST, *Variation on a theme of Schütte*, *Mathematical Logic Quarterly*, vol. 50 (2004), no. 3, pp. 258–264.
- [11] G. KREISEL, *Mathematical Logic, Lectures on modern mathematics* (Thomas Lorie Saaty, editor), vol. III, Wiley, 1965, pp. 95–195.
- [12] WOLFRAM POHLERS, *Proof Theory: An Introduction*, Lecture Notes in Mathematics, vol. 1407, Springer, 1989.

- [13] DIETER PROBST, *On the relationship between fixed points and iteration in admissible set theory without foundation*, *Archive for Mathematical Logic*, (2005), no. 44, pp. 561–580.
- [14] ———, *Pseudo-hierarchies in admissible set theories without foundation and explicit mathematics*, Ph.D. thesis, Universität Bern, 2005.
- [15] CHRISTIAN RÜEDE, *The proof-theoretic analysis of Σ_1^1 transfinite dependent choice*, *Annals of Pure and Applied Logic*, vol. 121 (2003), no. 1, pp. 195–234.
- [16] KURT SCHÜTTE, *Proof Theory*, Springer, 1977.
- [17] HELMUT SCHWICHTENBERG, *Proof theory: Some applications of cut-elimination*, *Handbook of Mathematical Logic* (J. Barwise, editor), North Holland, 1977, pp. 867–895.
- [18] STEPHEN G. SIMPSON, *Subsystems of Second Order Arithmetic*, Perspectives in Mathematical Logic, Springer, 1998.

UNIVERSITÄT BERN

INSTITUT FÜR INFORMATIK UND ANGEWANDTE MATHEMATIK

NEUBRÜCKSTRASSE 10, CH-3012 BERN, SWITZERLAND

E-mail: probst@iam.unibe.ch