MODEL THEORY WITHOUT CHOICE? CATEGORICITY

SAHARON SHELAH

The Hebrew University of Jerusalem Einstein Institute of Mathematics Edmond J. Safra Campus, Givat Ram Jerusalem 91904, Israel

Department of Mathematics Hill Center-Busch Campus Rutgers, The State University of New Jersey 110 Frelinghuysen Road Piscataway, NJ 08854-8019 USA

ABSTRACT. We prove Los conjecture = Morley theorem in ZF, with the same characterization (of first order countable theories categorical in \aleph_{α} for some (equivalently for every ordinal) $\alpha > 0$. Another central result here is, in this context: the number of models of a countable first order T of cardinality \aleph_{α} is either $\geq |\alpha|$ for every α or it has a small upper bound (independent of α close to \beth_2).

The author would like to thank the Israel Science Foundation for partial support of this research (Grant No.242/03) and Alice Leonhardt for the beautiful typing. Publication 840.

Annotated Content

§0 Introduction, pg.3-8

Part I:

§1 Morley's proof revisited, pg.9-10

[We clarify when does Morley proof works - when there is an ω_1 -sequence of reals]

§2 Stability and categoricity, pg.11-18

[We prove the choiceless Łoś conjecture. This requires a different proof as possibly there is no well ordered uncountable set of reals. Note that it is harder to construct non-isomorphic models, as e.g. we do not know whether successor cardinals are regular and even so whether, e.g. they have a stationary/co-stationary subset. Also we have to use more of stability theory.]

§3 A dichotomy on $\dot{I}(\aleph_{\alpha}, T)$: either bounded or $\geq |\alpha|$, pg.19-37

[This shows that "the lower part of the family of functions $\{\dot{I}(\lambda, T) : T \text{ first order complete countable}\}$ " is nice.]

 $\S 4$ On T categorical in |T|, pg.38-43

[Here we get only partial results.]

Part II:

§5 Consistency results, pg.44-49

[We look for cases of "classes have few models" which do not occur in the ZFC context.]

- §6 Comments on model theory in ZF, pg.50-53
- §7 On powers which are not cardinals: categoricity, pg.55-60

[We deal with models of (first order) theories in so called reasonable powers (which are not cardinals), it is equivalent to the completeness theorem holds. We throw some light on "can a countable first order T be categorical in some reasonable power".]

References, pg.59

§0 Introduction

I have known for long that there is no interesting model theory without (the axiom of) choice, not an exciting question anyhow as we all know that AC is true. This work is dedicated to a try to refute this opinion, i.e., this work throws some light on this in the contrary direction: Theorem 0.2 seriously, Theorem 3.14, (the parallel of the ZFC theorem 0.3) in a stronger way.

Lately, I have continued my work on pcf without full choice (see [Sh 835], earlier [Sh 497], [Sh:E38], later [Sh:F728]) and saw that with suitable "reasonable" weak version of (the axiom of) choice essentially we can redo all [Sh:c] (for first order classes with well ordered vocabulary; see 6.3).

Then it seems reasonable to see if older established version suffices, say ZF + DC_{\aleph_0} . We first consider Loś conjecture which can be phrased (why only \aleph_{α} 's and not other powers? see below)

0.1 The choiceless Łoś Conjecture:

For a countable (first order theory) T:

- (*)₁ T is categorical in \aleph_{α} for (at least) one ordinal $\alpha > 0$ iff
- $(*)_2$ T is categorical in \aleph_{β} for every ordinal $\beta > 0$.

In §1 we shall show that the Morley's proof works exactly when there is an uncountable well ordered set of reals. In §2 we give a new proof which works always (under ZF); it used Hrushovski [Hr89d], so:

0.2 Theorem. (ZF) For any countable T we have: $(*)_1$ of 0.1 iff $(*)_2$ of 0.1 iff T is \aleph_0 -stable with no two cardinal models¹.

Note that though we have been ready enough to use $ZF + DC_{\aleph_0}$ in fact we solve the problem in ZF.

A theorem from [Sh:c] is

- **0.3 Theorem.** [ZFC] For a countable complete (first order theory) T, one of the following occurs:
 - (a) $\dot{I}(\aleph_{\alpha}, T) \ge |\alpha|$ for every ordinal α
 - (b) $\dot{I}(\aleph_{\alpha}, T) \leq \beth_2$ for every ordinal α (and can analyze this case: either $\dot{I}(\aleph_{\alpha}, T) = 1$ for every α or $\dot{I}(\aleph_{\alpha}, T) = \min\{\beth_2, 2^{\aleph_{\alpha}}\}$ for every $\alpha > 0$).

¹that is, for no model M of T, formula $\varphi(x,\bar{y}) \in \mathbb{L}_{\tau(T)}$ and sequence $\bar{a} \in {}^{\ell g(\bar{y})}M$, do we have $\aleph_0 \leq |\varphi(M,\bar{a})| < ||M||$.

We shall prove a similar theorem in ZF in 3.14.

Thirdly, we consider an old conjecture from Morley [Mo65]: if a complete (first order) T is categorical in the cardinal $\lambda, \lambda = |T| > \aleph_0$ then T is a definitional extension of some $T' \subseteq T$ of smaller cardinality. The conjecture actually says that T is not really of cardinality λ . This was proved in ZFC. Keisler [Ke71a] proved it when $|T| < 2^{\aleph_0}$. By [Sh 4] it holds if $|T|^{\aleph_0} = |T|$. It is fully proven in [Sh:c, IX,1.19,pg.491]). The old proof which goes by division to three cases is helpful but not sufficient. Without choice (but note that λ is an \aleph) the case T superstable (or just $\kappa_r(T) < \lambda$) has really a similar proof. The other two cases, T is unstable and T stable with large $\kappa_r(T)$, are not. Here in §4 it is partially confirmed, e.g., when λ is regular, the proofs are different though related.

In §7 we deal with power of non-well orderable sets, in §5 we deal with consistency results and in §6 we look what occurs to classical theorems of model theory.

We may consider isomorphism after appropriate forcing. Baldwin-Laskowski-Shelah [BLSh 464], Laskowski-Shelah [LwSh 518] deal with the question "does T or even $PC(T_1,T)$ have non-isomorphic models which become isomorphic after some c.c.c. forcing?" But this turns out to be very different and does not seem related to the work here.

However, the following definition 0.4 suggests a problem which is closely related but it may be easier to find examples of such objects, so called below "cardinal cases" with "not so nice behaviour" than to find forcing extension of \mathbf{V} which satisfies $\mathbf{ZF} + \mathbf{a}$ failure of some hopeful theorem.

- **0.4 Definition.** 1) A cardinal case is a pair (λ, \mathbf{P}) where λ is a cardinal and \mathbf{P} is a family of forcing notions.
- 2) A cardinal⁺ case is a triple $(\lambda, \mathbf{P}, <)$ such that λ is a cardinal, \mathbf{P} a family of forcing notions and < a partial order on \mathbf{P} such that $\mathbb{P}_1 \leq \mathbb{P}_2 \Rightarrow \mathbb{P}_1 \lessdot \mathbb{P}_2$ so if we omit < we mean \lessdot .
- 4) We say that a theory T or more generally a (definition, absolute enough, of a) class \mathfrak{K} of models is categorical in the cardinal case (λ, \mathbf{P}) when: for every $M_1, M_2 \in \mathfrak{K}_{\lambda}$ (i.e., $\in \mathfrak{K}$ of cardinality λ), for some $\mathbb{P} \in \mathbf{P}$ we have $\Vdash_{\mathbb{P}}$ " $M_1 \cong M_2$ ".
- 5) We say that a theory T or more generally a (definition, absolute enough, of a) class \mathfrak{K} of models is categorical in the cardinal⁺ case $(\lambda, \mathbf{P}, <)$ when for any $\mathbb{P} \in \mathbf{P}$, in $\mathbf{V}^{\mathbb{P}}$ we have: if $M_1, M_2 \in \mathfrak{K}_{\lambda}^{\mathbf{V}[\mathbb{P}]}$, then for some $\mathbb{P}' \in \mathbf{P}$ satisfying $\mathbb{P} \leq \mathbb{P}'$ we have $\Vdash_{\mathbb{P}'/\mathbb{P}}$ " $M_1 \cong M_2$ ".
- 4) Similarly uncategorical, has/does not have μ pairwise non-isomorphic models, etc.
- 5) We may replace cardinal by power.
- <u>0.5 Question</u>: Characterize countable (complete first order) T which may be categorical in some uncountable power (say in some forcing extension of $\mathbf{L}[T]$). See on

this $\S 7$.

This work may be continued in [Sh:F701].

We thank Udi Hrushovski for various comments and pointing out that Łoś conjecture proof is over after 2.12 as Kueker conjecture is known in the relevant case (in earlier versions the proof (of the choiceless Łoś conjecture) was more interesting and longer). We thank Moti Gitik for a discussion of the consistency results and for pointing out 5.5).

Lately, I have learned that Truss and his students were pursuing the connection between universes with restricted choice and model theory by a different guiding line: using model theory to throw light on the arithmetic of Dedekind finite powers, works in this direction are Agatha Walczak-Typke [WT05], [WT07]. Very interesting, does not interact with the present investigation, but may be relevant to Question 0.5.

* * *

Recall

0.6 Definition. A cardinal is the power of some well ordered set (so an \aleph or a natural number).

In [Sh:F701] we may deal with theories in a vocabulary which is not well ordered. 0.7 Convention: If not said otherwise

- (a) T is a first order theory in a vocabulary $\tau \subseteq \mathbf{L}$
- (b) T is complete
- (c) T is infinite
- (d) if T is countable for simplicity $\tau, T \subseteq \mathcal{H}(\aleph_0)$ (for notational simplicity).

This is justified by

- 0.8 Observation. Assume τ is a countable vocabulary and T is a first order theory in τ , i.e., $T \subseteq \mathbb{L}_{\tau}$.
- 1) There is a vocabulary $\tau' \subseteq \mathcal{H}(\aleph_0) \subseteq \mathbf{L}_{\omega}$ and first order theory T' in τ' (so $T' \subseteq \mathbb{L}_{\tau} \subseteq \mathcal{H}(\aleph_0)$) such that for every cardinal λ, T is categorical in λ iff T' is categorical in λ (and even $\dot{I}(\lambda, T) = \dot{I}(\lambda, T')$, similarly for power and the parallel of 0.9 below).
- 2) If T is categorical in some cardinal λ then $T \cup \{(\exists^{\geq n} x)(x = x) : n < \omega\}$ is complete.
- 0.9 Observation. Assume τ is a vocabulary which can be well ordered (i.e., $|\tau| \in \text{Card}$).

There is a vocabulary $\tau' \in \mathbf{L}$ (or even $\tau' \in \mathbf{L}_{|\tau|^+}$) and a function f from $\mathbb{L}(\tau')$ onto $\mathbb{L}(\tau)$ (note that $\mathbb{L}(\tau') \subseteq \mathbf{L}_{|\tau|^+}$) mapping predicates/functions symbols to predicate/function symbols respectively with the same arity such that:

- \boxtimes_1 f maps the set of (complete) first order theories in \mathbb{L}_{τ} onto the set of (complete) first order theories in $\mathbb{L}_{\tau'}$ (really this is a derived map, \hat{f})
- \boxtimes_2 for some definable class **F** which is a function, **F** maps the class of τ -models onto the class of τ' -models such that
 - (a) **F** is one to one onto and $Th(\mathbf{F}(M)) = \hat{f}(Th(M))$
 - (b) **F** preserves isomorphisms and non-isomorphisms
 - (c) **F** preserves $M \subseteq N, M \prec N$
 - (d) for some function f, if $\mathbf{F}(M) = M'$ then for every sentence $\psi \in \mathbb{L}(\tau'), M' \models \psi \Leftrightarrow M \models f(\psi)$ where $f(\psi)$ is defined naturally
 - (e) **F** preserves power, so equality and inequality of powers (hence for any theory $T \subseteq \mathbb{L}_{\tau}$, letting $T' = \hat{f}(T)$, for any set $X, (\{M/\cong: M \in \text{Mod}_T \text{ has power } |X|\}| = |\{M'/\cong: M' \in \text{Mod}_{T'} \text{ has power } |X|\}|$.

We shall use absoluteness freely recalling the main variant.

- **0.10 Definition.** 1) We say $\varphi(\bar{x})$ is upward ZFC-absolute when: if $\mathbf{V}_1 \subseteq \mathbf{V}_2$ (are transitive classes containing the class Ord of ordinals, both models of ZFC) and $\bar{a} \in \mathbf{V}_1$ then $\mathbf{V}_1 \models \varphi(\bar{a}) \Rightarrow \mathbf{V}_2 \models \varphi(\bar{a})$.
- 2) Replacing upward by downward mean we use \Leftarrow ; omitting upward mean we use \Leftrightarrow . Similarly for version ZFC' of ZFC (e.g. ZF + DC); but absolute means ZFC-absolute.
- 0.11 Convention. 1) If not said otherwise, for a theory T belonging to $\mathbf{L}[Y_0], Y_0 \subseteq \mathrm{Ord}$, saying "T satisfies Pr ", (" Pr " stands for " $\mathrm{Property}$ ") we mean "for some $Y_1 \subseteq \mathrm{Ord}$ for every $Y_2 \subseteq \mathrm{Ord}, T$ satisfies Pr in $\mathbf{L}[Y_0, Y_1, Y_2]$ ".
- 2) But "T categorical in λ " always means in \mathbf{V} .

Recall

- **0.12 Definition.** 1) $\theta(A) = \min\{\alpha : \text{ there is no function from } A \text{ onto } \alpha\}.$
- 2) $\Upsilon(A) = \text{Min}\{\alpha: \text{ there is no one-to-one function from } \alpha \text{ into } A\}.$

- **0.13 Definition.** 1) If $T \subseteq \mathbb{L}(\tau)$, Γ is a set of types in $\mathbb{L}(\tau)$, i.e., each is an m-type for some m, then $\mathrm{EC}(T,\Gamma)$ is the class of τ -models M of T which omits every $p(\bar{x}) \in \Gamma$.
- 2) If $T \subseteq \mathbb{L}(\tau)$ is complete, $T \subseteq T_1 \subseteq \mathbb{L}(\tau_1)$ and $\tau \subseteq \tau_1$ then $PC(T_1, T)$ is the class of τ -reducts of models M_1 of T_1 . Similarly for a set Γ of types in $\mathbb{L}(\tau_1)$ let $PC(T, T_1, \Gamma)$ be the class of τ -reducts of models $M \in EC(T_1, \Gamma)$.

We shall use Ehrenfuecht-Mostowski models.

0.14 Definition. 1) Φ is proper for linear orders when:

- (a) for some vocabulary $\tau = \tau_{\Phi} = \tau(\Phi)$, Φ is an ω -sequence, the *n*-th element a complete quantifier free *n*-type in the vocabulary τ
- (b) for every linear order I there is a τ -model M denoted by $\mathrm{EM}(I,\Phi)$, generated by $\{a_t: t\in I\}$ such that $s\neq t\Rightarrow a_s\neq a_t$ for $s,t\in I$ and $\langle a_{t_0},\ldots,a_{t_{n-1}}\rangle$ realizes the quantifier free n-type from clause (a) whenever $n<\omega$ and $t_0<_I\ldots<_I t_{n-1}$; so really M is determined only up to isomorphism but we may ignore this and use $I_1\subseteq J_1\Rightarrow \mathrm{EM}(I_1,\Phi)\subseteq \mathrm{EM}(I_2,\Phi)$. We call $\langle a_t: t\in I\rangle$ "the" skeleton of M; of course "the" is an abuse of notation as it is not necessarily unique.
- 2) If $\tau \subseteq \tau(\Phi)$ then we let $\mathrm{EM}_{\tau}(I,\Phi)$ be the τ -reduct of $\mathrm{EM}(I,\Phi)$.
- 3) For first order T, let $\Upsilon_{\kappa}^{\text{or}}[T]$ be the class of Φ proper for linear orders such that
 - (a) $\tau_T \subseteq \tau_{\Phi}$ and τ_{Φ} has cardinality $\leq \kappa$
 - (b) for any linear order I the model $\mathrm{EM}(I,\Phi)$ has cardinality $|\tau(\Phi)|+|I|$ and we have $\mathrm{EM}_{\tau(T)}(I,\Phi)\in K$
 - (c) for any linear orders $I \subseteq J$ we have $\mathrm{EM}_{\tau(T)}(I,\Phi) \prec \mathrm{EM}_{\tau(T)}(J,\Phi)$.
- 4) We may use Skeleton $\langle \bar{a}_t : t \in I \rangle$ with $\alpha = \ell g(\bar{a}_t)$ constant but in the definition of " $\Phi \in \Upsilon_{\kappa}^{\text{or}}[T]$; we add $\alpha < \kappa^+$. Alternatively $\bar{a}_t = \langle F_i^{\text{EM}(I,\Phi)}(a_t) : i < \alpha \rangle$, where $F_i \in \tau_{\Phi}$ are unary function symbols. We use Φ, Ψ only for such objects. Let $\Upsilon_T^{\text{or}} = \Upsilon_{|T| + \aleph_0}^{\text{or}}[T]$.

§1 Morley's proof revisited

The main theorem of this section is 1.1. The proof is just adapting Morley's proof in ZFC. We shall use 0.8(2) and convention 0.7 freely.

- **1.1 Theorem.** [ZF + there is an uncountable well ordered set of reals]. The following conditions on a countable (first order) T are equivalent:
 - (A) T is categorical in some cardinal $\aleph_{\alpha} > \aleph_0$, in V, of course
 - (B) T is categorical in every cardinal $\aleph_{\beta} > \aleph_0$, in V, of course
 - (C) T is (in $\mathbf{L}[T]$), totally transcendental (i.e. \aleph_0 -stable) with no two cardinal models (i.e., for no model M of T and formula $\varphi(x, \bar{y}) \in \mathbb{L}(\tau_T)$ and $\bar{a} \in \ell^{g(\bar{y})}M$ do we have $\aleph_0 \leq |\varphi(M, \bar{a})| < ||M||$ and ||M|| is a cardinal, i.e. the set of elements of M is well-orderable hence its power is a cardinal)
 - (D) if $\mathbf{V}' \subseteq \mathbf{V}$ is a transitive class extending $\mathbf{L}, T \in \mathbf{V}'$ and \mathbf{V}' satisfies ZFC then the conditions in (C) hold
 - (E) for some V' clause (D) holds.

Proof. By 0.7 or better 0.8(2) without loss of generality T is complete, $T \subseteq \mathcal{H}(\aleph_0)$. Trivially $(B) \Rightarrow (A)$. Next $(A) \Rightarrow (C)$ by claims 1.2, 1.3 below. Lastly, $(C) \Rightarrow (A)$ by 1.4 below and $(C) \Leftrightarrow (D) \Leftrightarrow (E)$ holds by absoluteness. $\square_{1.1}$

1.2 Claim. $/ZF + \exists \ a \ set \ of \aleph_1 \ reals/$

If T is (countable) and in $\mathbf{L}[T]$ the theory T is not \aleph_0 -stable and $\lambda > \aleph_0$ then T is not categorical in λ .

Proof. In $\mathbf{L}[T]$ we can find E.M. models, i.e. $\Phi \in \Upsilon_T^{\text{or}}$ such that $\tau(\Phi)$ is countable, extends $\tau = \tau_T$ and $\mathrm{EM}_{\tau}(I, \Phi)$ is a model of T (of cardinality λ) for every linear order I (of cardinality λ) and let $M_1 = \mathrm{EM}_{\tau}(\lambda, \Phi)$ and without loss of generality the universe of M_1 is λ .

In \mathbf{V} let $\bar{\eta} = \langle \eta_{\alpha} : \alpha < \omega_1 \rangle$ be a sequence of pairwise distinct reals. In $\mathbf{L}[T]$ there is a countable model M_0 of T with $\mathbf{S}(M_0)$ uncountable so containing a perfect set. Hence also in $\mathbf{L}[T, \bar{\eta}], M_0$ is a countable model of τ with $\mathbf{S}(M_0)$ containing a perfect set, hence there is (in $\mathbf{L}[T, \bar{\eta}]$) a model M_2 of T of cardinality λ (λ is still an uncountable cardinal in $\mathbf{L}[T, \bar{\eta}]$) such that $M_0 \prec M_2$ and there is a sequence $\langle a_i : i < \omega_1 \rangle, a_i \in M_2$ realizes $p_i \in \mathbf{S}(M_0)$ with $\langle p_i : i < \omega_1^{\mathbf{V}} \rangle$ pairwise distinct. Without loss of generality the universe of M_2 is λ .

Clearly even in V, the model M_1 satisfies "if $A \subseteq M_1$ is countable then the set $\{\operatorname{tp}(a,A,M_1): a\in M_1\}$ is countable" whereas M_2 fails this; hence the models M_1,M_2 have universe λ , are models of T and are not isomorphic, so we are done. $\square_{1,2}$

1.3 Claim. Assume T is countable \aleph_0 -stable and has a two cardinal model (in $\mathbf{L}[T]$, but both are absolute).

<u>Then</u> T is not categorical in λ , in fact, $I(\aleph_{\alpha}, T) \geq |\alpha|$ for every ordinal α .

Proof. So in $\mathbf{L}(T)$ it has a model M_1 and a finite sequence $\bar{a} \in {}^{\ell g(\bar{y})}(M_1)$ and a formula $\varphi(x,\bar{y}) \in \mathbb{L}(\tau_T)$ such that $\aleph_0 \leq |\varphi(M_1,\bar{a})| < ||M_1||$. If $\aleph_\beta < \lambda$, working in $\mathbf{L}[T]$ without loss of generality $|\varphi(M_1,\bar{a})| = \aleph_\beta$, $||M_1|| = \lambda$ (by [Sh 3]) and the universe of M_1 is λ . But \aleph_0 -stability T has (in $\mathbf{L}[T]$) a saturated model M_2 of cardinality λ , so without loss of generality the universe of M_2 is λ . So $\bar{a}' \in {}^{\ell g(\bar{y})}(M_2) \Rightarrow |\varphi(M_2,\bar{a}')| \notin [\aleph_0,\lambda)$. Clearly even in \mathbf{V},M_1,M_2 are models of T of cardinality λ and are not isomorphic. In fact for every $\aleph_\beta \leq \lambda$, T has an \aleph_β -saturated not $\aleph_{\beta+1}$ -saturated model M_β of cardinality λ such that $|\varphi(M_\beta,\bar{a}_\beta)| = \aleph_\beta$ for some $\bar{a} \in {}^{\ell g(\bar{y})}(M_\beta)$. So $\{|\varphi((M_\beta,\bar{b})| + \aleph_0 : \bar{b} \in {}^{\ell g(\bar{y})}(M_\beta)|\} = \{\aleph_\beta,\lambda\}$, hence $\mathbf{V} \models M_\beta \ncong M_\gamma$ when $\aleph_\beta < \aleph_\gamma \leq \lambda$ so also the second phrase in the conclusion of the claim holds and even $\dot{I}(\aleph_\alpha,T) \geq |\alpha+1|$.

1.4 Claim. Assume T is countable \aleph_0 -stable with no two cardinal models even just in $\mathbf{L}[T]$ and $\lambda > \aleph_0$. Then T is categorical in λ .

Proof. Let M_1, M_2 be models of T of cardinality λ , without loss of generality both have universe λ , clearly $\mathbf{L}[T, M_1, M_2]$ is a model of ZFC and by absoluteness T still satisfies the assumption of 1.4 in it, and M_1, M_2 are (also in it) uncountable models of T of the same uncountable cardinality in this universe. But by 1.1 being a theorem of ZFC clearly M_1, M_2 are isomorphic in $\mathbf{L}[T, M_1, M_2]$, hence in \mathbf{V} . $\square_{1.4}$

§2 Stability and categoricity

Our aim in this section is the categoricity spectrum for countable T (i.e. Th. 2.1), but in the claims leading to the proof we do not assume countability. Note that the absoluteness of various properties is easier for countable T.

- **2.1 Theorem.** [ZF] For countable T, clauses (A),(B),(C),(D),(E) of Theorem 1.1 are equivalent.
- 2.2 Observation. 1) If T is unstable so has the order property, say as witnessed by $\varphi(\bar{x}, \bar{y})$ and, of course, $\tau = \tau_T \subseteq \mathbf{L}$ then for some $\Phi \in \mathbf{L}[T]$
 - \circledast (a) Φ is proper for linear orders
 - (b) $\tau \subseteq \tau_{\Phi}$ and for every linear order I, $\mathrm{EM}_{\tau}(I, \Phi)$ is a model of T with skeleton $\langle \bar{a}_t : t \in I \rangle$, $\ell g(\bar{a}_{\tau}) = \ell g(\bar{x}) = \ell g(\bar{y})$
 - (c) $\mathrm{EM}_{\tau}(I, \Phi) \models \varphi[\bar{a}_s, \bar{a}_t]^{\mathrm{if}(s < t)}$
 - (d) $\tau(\Phi) \subseteq \mathbf{L}$ and $|\tau(\Phi)| = |T|$ (if $\tau(T) \in \mathbf{L}$, without loss of generality $\tau(\Phi) \in \mathbf{L}$) and without loss of generality $\mathbf{L}[T] \models |\tau(\Phi)| = |T|$.
- 2) It follows that if I is well orderable then the universe of $\text{EM}(I, \Phi)$ is well orderable so it is of cardinality $|I| + |T| + \aleph_0$ hence we can assume it has this cardinal as its universe.

Proof. 1) By [Sh:c]. $\square_{2.2}$ Follows. $\square_{2.2}$

Our first aim is to derive stability from categoricity, for diversion we give some versions.

2.3 Claim. Let Φ be as in 2.2. Then

 $M_1 \ncong M_2$ when κ_1, κ_2 are regular uncountable cardinals (> |T|) and for some $A \subseteq \text{Ord}$, in $\mathbf{L}[A]$

- \circledast (a) $M_{\ell} = \mathrm{EM}_{\tau}(I_{\ell}, \Phi)$ in $\mathbf{L}[A]$, (so $T, \Phi \in \mathbf{L}[A]$, $I_{\ell} \in \mathbf{L}[A]$) for $\ell = 1, 2$
 - (b) $\bar{s}^1 = \langle s^1_{\alpha} : \alpha < \kappa_1 \rangle$ is increasing in $I_1, \bar{t}^1 = \langle t^1_{\alpha} : \alpha < \kappa_2 \rangle$ is decreasing in I_1 (in $\mathbf{L}[A]$)
 - (c) $\alpha < \kappa_1 \wedge \beta < \kappa_2 \Rightarrow s^1_{\alpha} <_{I_1} t^1_{\beta} \text{ but } \neg (\exists s \in I_1) [(\forall \alpha < \kappa_1)(s^1_{\alpha} < r) \land (\forall \beta < \kappa_2)(s < t^1_{\beta})]$

- (d) in I_2 there is no pair of sequences like \bar{s}^1, \bar{t}^1
- (e) also in the inverse of I_2 , there is no such pair
- (f) [only for simplicity, implies (d)+(e)] I_2 is $\cong I_2 \times \mathbb{Q}$ ordered lexicographically.

Proof. Without loss of generality the universes of M_1 , M_2 are ordinals, and toward contradiction assume f is an isomorphism from M_1 onto M_2 . We can work in $\mathbf{L}[A, M_1, M_2, f]$ which is a model of ZFC, so easy to contradict (as in [Sh 12], see detailed proof showing more in 3.2).

 $\square_{2.3}$

2.4 Conclusion. [ZF + $|T|^+$ is regular] If T is categorical in some cardinal $\lambda > |T|$, then T is stable (in $\mathbf{L}[T]$).

A fuller version is

- **2.5 Claim.** $M_1 \ncong M_2$ when for some $\lambda > |T|$ we have:
 - \circledast (a) $M_{\ell} = \mathrm{EM}_{\tau}(I_{\ell}, \Phi)$ where T, Φ are as in 2.2 so $\Phi \in \mathbf{L}[T]$
 - (b) $I_1 = \lambda \times \mathbb{Q}$ ordered lexicographically
 - (c) $I_2 = \sum_{\alpha \leq \lambda} I_{\alpha}^2 \in \mathbf{L}[T]$ where I_{α}^2 is isomorphic to $\alpha + \alpha^*$ (α^* the inverse
 - of α) or just
 - (c)⁻ I_2 is a linear order of cardinality λ such that for every limit ordinal $\delta \leq |T|^+, I_2$ has an interval isomorphic to $\delta + \delta^*$
 - (d) I_1, I_2 has cardinality λ .

Proof. Let $\theta = |T|$ in $\mathbf{L}[T]$ and $\theta_1 = (\theta^+)^{\mathbf{V}}$. Without loss of generality M_ℓ has universe λ , assume toward contradiction that $M_1 \cong M_2$ let f be an isomorphism from M_1 onto M_2 and consider the universe $\mathbf{L}[T, M_1, M_2, f]$. In this universe $\theta_1^{\mathbf{V}}$ may be singular but is still a cardinal so $\delta =: (\theta^+)^{\mathbf{L}[T, M_1, M_2, f]}$ is necessarily $\leq (|\theta|^+)^{\mathbf{V}}$ hence I_2 has an interval isomorphic to $\delta + \delta^*$. Now we continue as in 2.3 (see details in 3.2).

2.6 Conclusion. If T is categorical in the cardinal $\lambda > |T|$, then T is stable.

<u>2.7 Discussion</u>: 1) We may like to have many models. So for T unstable if there are α regular cardinals $\leq \lambda$ we can get a set of pairwise non-isomorphic models of T of cardinality λ indexed by $|\mathscr{P}(\alpha)|$.

It is not clear what, e.g., we can get in \aleph_1 . As 2.5 indicate it is hard to have few models, i.e., to have such universe (see more in §3); but for our present purpose all this is peripheral, as we have gotten two.

On uni-dimensional see [Sh:c, V,Definition 2.2,pg.241] and [Sh:c, V.Theorem 2.10,p.246].

- **2.8 Definition.** A stable theory T is uni-dimensional <u>if</u> there are no $M \models T$ and two infinite indiscernible sets in M which are orthogonal.
- **2.9 Claim.** Assume T is stable (in $\mathbf{L}[T]$, anyhow this is Z^- -absolute). Then for every $\lambda > |T|$, T has a model $M \in \mathbf{L}[T]$ of cardinality λ such that:
 - \odot in M there are no two (infinite) indiscernible non-trivial sets each of cardinality $\geq |T|^+$ which are orthogonal.

Proof. We work in $\mathbf{L}[T]$ or $\mathbf{L}[T,Y],Y\subseteq \mathrm{Ord}$ and let $\kappa=|T|^{\mathbf{L}[T]}$ and $\partial=\theta^{\mathbf{V}}(\mathscr{P}(\kappa))$. Let μ be large enough (e.g., $\beth((2^{\partial})^+)$, i.e. the $(2^{\partial})^+$ -th beth), let \mathfrak{C} be a μ^+ -saturated model of T. Let $\mathbf{I}=\{a_i:i<\mu\}\subseteq\mathfrak{C}$ be an infinite indiscernible set of cardinality μ and minimal, i.e. $\mathrm{Av}(\mathbf{I},\cup\mathbf{I})$ is a minimal type. Let $M_1\prec\mathfrak{C}$ be κ^+ -prime over \mathbf{I} .

More specifically

 \otimes $\langle A_{\varepsilon} : \varepsilon \leq \kappa^{+} \rangle$ is an increasing sequence of subsets of $M_{1}, A_{\kappa^{+}} = M_{1}, A_{0} = \{a_{i} : i < \mu\}$ and $\bar{B} = \langle B_{a} : a \in M_{1} \rangle$, satisfies $[a \in A_{0} \Rightarrow B_{a} = \{a\}]$ and if $a \in A_{\varepsilon+1} \setminus A_{\varepsilon}$ then $B_{a} \subseteq A_{\varepsilon}$ and $\operatorname{tp}(a, B_{a}, \mathfrak{C}) \vdash \operatorname{tp}(a, A_{\varepsilon+i} \setminus \{a\}, \mathfrak{C})$ and $|B_{a}| \leq \kappa$ and without loss of generality $B_{a} = \{b_{a,j} : j < \kappa\}$ and $\zeta < \varepsilon \Rightarrow B_{a} \cap A_{\zeta+1} \not\subseteq A_{\zeta}$ and $a' \in B_{a} \Rightarrow B_{a'} \subseteq B_{a}$.

Expand M_1 to M_2 by adding $P^{M_2} = \mathbf{I}, <^{M_2} = \{(a_i, a_j) : i < j < \mu\}, E^{M_2} = \{(b_1, b_2): \text{ for some } \varepsilon < \kappa^+ \text{ we have } b_1 \in A_{\varepsilon} \land b_2 \in (A_{\varepsilon+1} \backslash A_{\varepsilon})\}, F_j^{M_2}(a) = b_{a,j} \text{ for } j < \kappa^+, \text{ (hence } a \in A_{\varepsilon+1} \backslash A_{\varepsilon} \land \zeta < \varepsilon \Rightarrow c\ell_{M_2}\{a\} \cap A_{\zeta+1} \backslash A_{\zeta} \neq \emptyset) \text{ and add Skolem functions, still } \tau(M_2) \in \mathbf{L}[T] \text{ has cardinality } \kappa = (|T| + \aleph_0)^{\mathbf{L}[T]}.$

Now (as in the proof of the omitting type theorem, see e.g., [Sh:c, VII,§5]) we can find $\langle \mathbf{I}_n : n < \omega \rangle, \mathbf{I}_n \subseteq \mathbf{I}$ is an *n*-indiscernible sequence in M_2 of cardinality $> 2^{\partial}$ and

 $(*)_1$ if for $n < \omega, M_2 \models a_0^n < \ldots < a_{n-1}^n$ and $\ell < n \Rightarrow a_\ell^n \in \mathbf{I}_n$ then $p_n = \operatorname{tp}(\langle a_0^n, \ldots, a_{n-1}^n \rangle, \emptyset, M_2) = \operatorname{tp}(\langle a_0^{n+1}, \ldots, a_{n-1}^{n+1} \rangle, \emptyset, M_2)$.

Let $\mathbf{I}_n = \{a_\alpha : \alpha \in \mathscr{U}_n\}$ and note that

- $\boxtimes_1 \langle \bar{\sigma}(a_{i(\alpha,0)},\ldots,a_{i(\alpha,m-1)}) : \alpha \in Z \rangle$ is an indiscernible set in M_1 (equivalently in \mathfrak{C}) when:
 - (a) $2m \leq n < \omega$
 - (b) $Z \subseteq \mathcal{U}_m$ is infinite
 - (c) $i(\alpha, \ell) \in \mathcal{U}_n$ is increasing with $\ell < m$ for $\alpha \in Z$
 - (d) for each $\ell, k < m$ and $\alpha_1 < \beta_1, \alpha_2 < \beta_2$ from Z we have $i(\alpha_1, \ell) < i(\beta_1, k) \Leftrightarrow i(\alpha_2, \ell) < i(\beta_2, k)$
 - (e) $\bar{\sigma}(x_0,\ldots,x_{m-1})$ is a finite sequence of $\tau(M_2)$ -term
 - (f) all $\langle a_{i(\alpha,0)}, \ldots, a_{i(\alpha,m-1)} \rangle$ for $\alpha \in \mathbb{Z}$ realize the same type (equivalently of quantifier-free type) in M_2 .

[Why? If $k \leq n$ and $j < \ell g(\bar{\sigma})$ then the truth values of $\sigma_j(a_{i_0}, \ldots, a_{i_{k-1}}) \in A_{\varepsilon}$ for $i_0 < \ldots < i_{k-1} < \mu$ such that $a_{i_0}, \ldots, a_{i_{k-1}} \in \mathbf{I}_n$ depend on σ only, we can prove this by induction on $\max\{\varepsilon_j : j < \ell g(\bar{\sigma})\}$, using the properties of the B_a 's. By the properties of $\mathbf{F}_{\kappa^+}^t$ -constructions² ([Sh:c, IV]) we are easily done³.]

Moreover

 \boxtimes_2 in \boxtimes_1 , the $\mathbb{L}(\tau_T)$ -type of $\langle \bar{\sigma}(a_{i(\alpha,0)},\ldots,a_{i(\alpha,m-1)}) : \alpha \in Z \rangle$ depends just on $Z, \bar{\sigma}$ and the truth values in (d) from \boxtimes_1 and the types in (f) of \boxtimes_1 over $\operatorname{acl}_{M_1}(\emptyset)$.

[Why: Note that $\operatorname{acl}_{M_2}(\emptyset) \subseteq (\text{the Skolem hull of } \emptyset \text{ in } M_2) \text{ and } \mathbf{I} \cap \operatorname{acl}_{M_2}(\emptyset) \text{ is infinite.}]$

So we can find a $\tau(M_2)$ -model M_3 generated by the indiscernible sequence $\langle b_{\alpha} : \alpha < \lambda \rangle$ such that for every $n < \omega$ and $\alpha_0 < \ldots < \alpha_{n-1} < \lambda$, recalling $(*)_1$ we have $p_n = \operatorname{tp}(\langle b_{\alpha_0}, \ldots, b_{\alpha_{n-1}} \rangle, \emptyset, M_3)$. Without loss of generality the Skolem hull of \emptyset in M_3 is the same as in M_2 . Let $M_4 = M_3 \upharpoonright \tau_T$. Clearly M_4 is a model of T of cardinality λ .

Now suppose that

(*)₂ in **V** we have $\mathbf{J} \subseteq M_4$ (or even $\mathbf{J} \subseteq {}^{\omega \geq}(M_4)$), is an indiscernible set of cardinality $\geq \kappa^+$ orthogonal to P^{M_3} which is an infinite indiscernible set in M_4 (this is absolute enough).

²instead we can use the conclusion derived in \boxtimes_2

³this is not the end of the proof, we still need to show another indiscernible set does not exist

Let $\mathbf{J} \supseteq \{c_{\alpha} : \alpha < (\kappa^{+})^{\mathbf{V}}\}$ with the c_{α} 's pairwise distinct. Let $c_{i} = \sigma_{i}(b_{\alpha(i,0)}, \ldots, b_{\alpha(i),n(\alpha)-1})$ where $\alpha(i,0) < \ldots < \alpha(i,n(i))$ (may be clearer in $\mathbf{L}[T,Y,\mathbf{J}]$).

So in $\mathbf{L}[T,Y,\mathbf{J}]$ for some $Z\subseteq (\kappa^+)^{\mathbf{V}}$ of cardinality $\geq (\kappa^+)^{\mathbf{L}[T,Y,\mathbf{J}]}$ (so maybe $\mathbf{V} \models |Z| < \kappa^+$) we have $i \in Z \Rightarrow \sigma_i = \sigma_* \wedge n(i) = n(*)$, and the truth value of $\alpha(i_1, \ell_1) < \alpha(i_2, \ell_2)$ for $i_1 < i_2$ depend just on (ℓ_1, ℓ_2) . In $\mathbf{L}[T, Y]$ for each $n \ge 2n(*)$, we can find $a_{i,\ell}^n \in \mathbf{I}_n$ for $i < \partial, \ell < n(*)$ such that $M_2 \models a_{i_1,\ell_1}^n < a_{i_2,\ell_2}^n \Leftrightarrow \alpha(0,\ell_1) < \alpha(1,\ell_2)$ for every $i_1 < i_2 < \partial$ and $M_2 \models a_{i,0}^n < a_{i,1}^n < \ldots < a_{i,n(*)-1}^n$ for $i < \partial$. By \boxtimes_1 we know that $\langle \sigma_*(a_{i,0}^n,\ldots,a_{i,n(*)-1}^n):i<\partial\rangle$ is an indiscernible set in M_1 hence an indiscernible set over $\operatorname{acl}_{M_2}(\emptyset)$. By \boxtimes_2 , its type over $\operatorname{acl}_{M_2}(\emptyset)$ does not depend on n. As $|\mathbf{I}_n| > \partial > (2^{|T|})^{\mathbf{L}[T,Y,\mathbf{J}]}$, we easily get, see [Sh.c, Ch.V,2.5,pg.244] that this indiscernible set is not orthogonal to the indiscernible set $\{a_i : i < \mu\}$. Also easily letting $f: Z \to \partial$ be one to one order preserving, the type which $\langle c_i : i \in Z \rangle$ realizes over $\operatorname{acl}_{M_2}(\emptyset)$ in M_3 is the same as the type of $\langle \sigma_*(a^n_{f(i),0},\ldots) : i \in Z \rangle$ realized in M_2 over $\operatorname{acl}_{M_2}(\emptyset)$ for $n \geq 2n(*) + 1$, as for formulas with $\leq m$ variable we consider n > m. As **J** was chosen to be indiscernible not orthogonal to $\operatorname{tp}(a_{0,0}^n,\operatorname{acl}_{M_2}(\emptyset),M_1),$ i.e., to the indiscernible set P^{M_4} , we get a contradiction. So there is no **J** as in $(*)_2$. As **I** is minimal, it follows that in **V**, if for $\ell = 1, 2$ the set $\mathbf{J}_{\ell} \subseteq n(\ell)(M_4)$ is indiscernible of cardinality $\geq \theta^+$ then $\mathbf{J}_1, \mathbf{J}_2$ are not orthogonal to I hence $\mathbf{J}_1, \mathbf{J}_2$ not orthogonal (e.g. works in $L[T, Y, J_1, J_2]$).

But this says that M_4 is a model as required in the conclusion of 2.9. $\square_{2.9}$

- 2.10 Remark. 1) By $\mathbf{F}_{\aleph_0}^f$ -constructions (see [Sh:c, IV]) we can get models with peculiar properties.
- 2) On absoluteness see 3.1.
- 3) In fact by [Sh 300f, §1], we can assume that $\langle \sigma(b_0, \ldots, b_{m-1}) : m < n, b_0 <^{M_2} \ldots <^{M_2} b_{m-1}$ are from $\mathbf{I}_n \rangle$ where $\sigma = \sigma(x_0, \ldots, x_{m-1})$ and σ is a $\tau(M_2)$ -term, is (fully) indiscernible in the model $M_2 \upharpoonright \tau_T$, i.e., in M_1 , see definition there. But the argument above is simpler.
- 2.11 Conclusion. If T is stable and categorical in $\lambda > |T|$ then (in $\mathbf{L}[T,Y]$ where $Y \subseteq \mathrm{Ord}$):
 - \boxtimes (a) T is uni-dimensional
 - (b) T is superstable
 - (c) T has no two cardinal models
 - (d) D(T) has cardinality $\leq |T|$; moreover $D(T) \in \mathbf{L}[T]$ and $\mathbf{L}[T] \models |D(T)| = \lambda$.

Proof. Assume clause (a) fails and we shall produce two models of cardinality (and universe) λ . The first N_1 is from 2.9. The second is a model N_2 such that there are indiscernible $\mathbf{I}, \mathbf{J} \subseteq N_1$ (or $\omega > (N_1)$) of cardinality λ which are orthogonal; this contradicts the categoricity hence clause (a).

The superstability, i.e., clause (b) follows from clause (a) by Hrushovski [Hr89d]. Clause (c), no two cardinal models follows from clause (a) by [Sh:c, V,§6].

Now $|D(T)| \leq |T|$ (clause (d)) is trivial as otherwise we have two models M_1, M_2 of T of cardinality λ such that some $p \in D(T)$ is realized in one but not the other (i.e., first choose $M_1 \in \mathbf{L}[T]$ realizing $\leq |T|$ types. Clearly $\{p \in D(T) : p \text{ is realized in } M\}$ is a well ordered set so by the assumption we can choose $p \in D(T)$ not realized in M_1 and lastly choose M_2 realizing p).

2.12 Claim. If clauses (b),(c),(d) of 2.11 hold and T is categorical in $\lambda > |T|$ then:

(e) any model M of T of cardinality μ , for any $\mu > |T|$ is \aleph_0 -saturated.

Proof. Assume clause (e) fails as exemplified by M and we shall get contradiction to clause (c) of 2.11, so without loss of generality the universe of M is μ .

For any $Y \subseteq$ Ord working in $\mathbf{L}[T,Y,M]$ we can find $\bar{a} \subseteq M$ and formula $\varphi(x,\bar{y}) \in \mathbb{L}_{\tau(T)}$ such that $\varphi(x,\bar{a})$ is a weakly minimal formula in M, existence as in [Sh 31]. Let $N_0 \prec M$ be of cardinality |T| such that $\bar{a} \subseteq N_0$ and $N_0 \in \mathbf{L}[T,M]$.

<u>Case 1</u>: $\{p \in \mathbf{S}(N_0) : p \text{ is realized in } M\}$ has power > |T| in \mathbf{V} .

But as |M| is an ordinal this set is well ordered so the proof of 1.2 applies contradicting categoricity in λ and we get more than needed.

<u>Case 2</u>: Not Case 1 but there is a finite $A \subseteq M$ such that $\bar{a} \subseteq A$ and $p \in \mathbf{S}(A), \varphi(x, \bar{a}) \in p$ and the type p is omitted by M.

As in [Sh 31] (using "not Case 1" here instead "T stable in |T|" there) we can find (in $\mathbf{L}[T,M]$) a model M' such that $M \prec M', M'$ omits the type p and $||M'|| \geq \lambda$, so by DLST (= the downward Lowenheim-Skolem-Tarksi) some $N_1 \prec M'$ has cardinality λ and is not \aleph_0 -saturated. Hence for some complete type $p(\bar{x},\bar{y}) \in D(T)^{\mathbf{L}[T,M]}$, for some $\bar{b} \in {}^{\ell g(\bar{y})}(N_1)$, the model N_1 omits the type $p(\bar{x},\bar{b})$ which is a type, i.e. finitely satisfiable in N_1 .

By clause (d) of \boxtimes of 2.11 we have $|D(T)| \leq |T|$ in $\mathbf{L}[T,Y]$ and D(T) is included in $\mathbf{L}[T,Y]$. So in $\mathbf{L}[T,Y]$, for every finite $A \subseteq N \models T$, $\mathbf{S}(A,N)^{\mathbf{V}}$ is the same as $\mathbf{S}(A,N)$ computed in $\mathbf{L}[T,Y]$ and is there of cardinality $\leq |D(T)|$ hence absolute. So in $\mathbf{L}[T,Y]$ we can find a model N_2 of T of cardinality λ which is \aleph_0 -saturated. [Alternatively to this, we can choose a model N_2 of cardinality λ such that: if

 $\bar{b}' \in {}^{\ell g(\bar{y})}N_2$ realizes $\operatorname{tp}(\bar{b}',\emptyset,M')$ then for some $\bar{a} \in {}^{\ell g(\bar{x})}N_2$ the sequence $\bar{a}' \hat{b}'$ realizes $p(\bar{x},\bar{y})$.

By the previous paragraphs this is a contradiction to categoricity.

Case 3: Neither Case 1 nor Case 2.

Subcase A: T countable.

Let N_1 be such that

- \circledast (a) $N_1 \prec M$ is countable
 - (b) $\bar{a} \subseteq N_1$
 - (c) if $\bar{a} \subseteq A \subseteq N_1$, A finite and p is a non-algebraic type satisfying $\varphi(x,\bar{a}) \in p \in \mathbf{S}(A,M)$ then p is realized in N_1

(possible as by clause (d) of \boxtimes of 2.11 the set D(T) is countable and "neither case 1 nor case 2").

Let N_2 be a countable saturated model of T such that $N_1 \prec N_2$. We can build an elementary embedding f (still working in $\mathbf{L}[T, M]$) from N_1 into N_2 such that $f(\varphi(N_1, \bar{a})) = (\varphi(N_2, \bar{a}))$. This contradicts clause (c) of \boxtimes of 2.11.

The last subcase is not needed for this section's main theorem 2.1, (but is needed for 2.11).

Subcase B: T uncountable.

So possibly increasing $Y \subseteq \operatorname{Ord}$, in $\mathbf{L}[T,Y]$ we have two models M_1, M_2 of T, M_1 is \aleph_0 -saturated, M_2 is not but $\varphi(x, \bar{a}), M_2$ fails cases 1 and 2; we work in $\mathbf{L}[T,Y]$. Let $\ell g(\bar{a}) = n$ and $T^+ \in \mathbf{L}[T,Y]$ be the first order theory in the vocabulary $\tau^+ = \tau_T \cup \{c_\ell : \ell < n\} \cup \{P\}$ where c_ℓ an individual constant, P a unary predicate such that $M^+ = (M, c_0^{M^+}, \dots, c_{n-1}^{M^+}, P^{M^+})$ is a model of T^+ iff $M = M^+ \upharpoonright \tau$ is a model of $T, \varphi(M, c_0^{M^+}, \dots, c_{n-1}^{M^+})$ is infinite and $\subseteq P^{M^+}$ and $M \upharpoonright P^{M^+} \prec M$. As T is uni-dimensional (more specifically clause (c) of \boxtimes of $2.11 + [\operatorname{Sh:c}, V, \S 6]$) T^+ is inconsistent, hence for some finite $\tau' \subseteq \tau, T^+ \cap \mathbb{L}(\tau' \cup \{c_0, \dots, c_{n-1}, P\})$ is inconsistent. Now choose $\bar{a}_1 \in {}^n(M_1)$ realizing $\operatorname{tp}(\bar{a}, \emptyset, M_2)$, let $\bar{a}_2 = \bar{a}$ and let χ be large enough, $\mathfrak{B} \prec (\mathscr{H}(\chi)^{\mathbf{L}[T,Y]}, \in)$ be countable such that $\{M_1, M_2, \tau', \bar{a}_1, \bar{a}_2\} \in \mathfrak{B}$; recall that we are working in $\mathbf{L}[T, Y]$. Now replacing M_1, M_2 by $(M_1 \upharpoonright \tau') \cap \mathfrak{B}, (M_2 \upharpoonright \tau') \cap \mathfrak{B}$ we get a contradiction as in Subcase A. $\square_{2.12}$

Proof of Theorem 2.1. By 0.8(2) and 0.9 without loss of generality T is complete, $T \subseteq \mathcal{H}(\aleph_0)$. Trivially $(B) \Rightarrow (A)$, by 1.4 we have $(C) \Rightarrow (B)$ and by absoluteness

 $(C) \Leftrightarrow (D) \Leftrightarrow (E)$, so it suffices to prove (C) assuming (A). By 2.6 the theory T is stable hence the assumption of 2.11 holds hence its conclusion, i.e., \boxtimes of 2.11 holds whenever $Y \subseteq \operatorname{Ord}$, in particular $D(T) \in \mathbf{L}[T]$. So by 2.12 we can conclude: every model of T of cardinality $\lambda > \aleph_0$ is \aleph_0 -saturated (in \mathbf{V} or, equivalently, in $\mathbf{L}[T, M]$ when M has universe λ). If T is \aleph_0 -stable use 1.3. So we can assume T is not \aleph_0 -stable but is superstable (recall clause (b) of \boxtimes of 2.11) hence T is not categorical in \aleph_0 (even has $\geq \aleph_0$ non-isomorphic models, by a theorem of Lachlan, see, e.g., $[\operatorname{Sh:c}]$), in any $\mathbf{L}[T,Y]$. So by Kueker conjecture (proved by Buechler [Be84] for T superstable and by Hrushovski $[\operatorname{Hr}89]$ for stable T), we get contradiction. $\square_{2.1}$

2.13 Remark. See more in [Sh:F701] about T which is categorical in the cardinal $\lambda > |T|, T$ not categorical in some $\mu > |T|$.

§3 A dichotomy for $\dot{I}(\aleph_{\alpha}, T)$: bounded or $\geq |\alpha|$

Our aim is to understand the lower part of the family of functions $\dot{I}(\lambda, T), T$ countable: either $(\forall \alpha)\dot{I}(\aleph_{\alpha}, T) \geq |\alpha|$ or $\dot{I}(\aleph_{\alpha}, T)$ is constant and not too large (for α not too small), see 3.14. For completeness we give a full proof of 3.2.

We need here absoluteness between models of the form $\mathbf{L}[Y]$ and this may fail for " $\kappa(T) > \kappa$ ", "T stable uni-dimensional". But usually more is true.

- 3.1 Observation. 1) "T is first order", " $\tau_T \subseteq \mathbf{L}_{\omega}$ ", " $\tau_T \subseteq \mathbf{L}$ ", "T is complete" are Z^- -absolute.
- 2) For T (not necessarily $\in \mathbf{L}$ but τ_T well orderable, our standard assumption) which is complete:
 - (a) "T is stable" is Z^- -absolute
 - (b) "T is superstable" is $(Z^- + DC)$ -absolute (and downward Z^- -absolute; Z^- -absolute if $T \subseteq \mathbf{L}$)
 - (c) "T totally transcendental" is $(Z^- + DC)$ -absolute (and downward Z^- -absolute; Z^- -absolute if $T \subseteq \mathbf{L}$); "T is \aleph_0 -stable, $\tau(T) \subseteq \mathbf{L}$ " is Z^- -absolute
 - (d) the appropriate ranks are $(Z^-+ DC)$ -absolute $(Z^-$ -absolute if $T \subseteq \mathbf{L})$ as the rank of $\{\varphi(x,\bar{a})\}$ in M depend just on $T, \varphi(x,\bar{y})$ and $\operatorname{tp}(\bar{a},\emptyset,M)$
 - (e) "M a model of T and $\mathbf{I}, \mathbf{J} \subseteq M$ (or $\omega > M$) are infinite indiscernible sets, \mathbf{I}, \mathbf{J} are orthogonal and where T is stable", is Z^- -absolute
 - (f) "T is stabe not uni-dimensional" is upward Z^- -absolute
 - (g) for countable T, "T is stable not uni-dimensional" is Z^- -absolute when $T \subseteq \mathbf{L}$,
 - (h) 'T is countable stable with the OTOP (omitting type order property, see 3.7 below)" is $(Z^- + DC)$ -absolute
 - (i) "M is primary over A, M a model of the (complete) stable theory T" is upward Z^- -absolute.

Proof. E.g.

2) <u>Clause (b)</u>:

This just asks if the tree \mathscr{T} has an ω -branch where the n-th level of \mathscr{T} is the set of sequence $\langle \varphi_{\ell}(x, \bar{y}_{\ell}) : \ell < n \rangle$ such that for every $m, \{ \varphi(x_{\eta}, \bar{y}_{\nu})^{\mathrm{if}(\nu \triangleleft \eta)} : \ell < n, \nu \in \ell m, \eta \in \ell^n m \}$ is consistent with T.

<u>Clause (e)</u>: Recall that this is equivalent to

 $(*)_1$ Av $(\mathbf{I}, \mathbf{I} \cup \mathbf{J})$ and Av $(\mathbf{J}, \mathbf{I} \cup \mathbf{J})$ are weakly orthogonal types which is equivalent to

(*)₂ for every $\varphi = \varphi(\bar{x}, \bar{y}, \bar{z}) \in \mathbb{L}(\tau_T)$ and $\bar{b} \in {}^{\ell g(\bar{z})}(\mathbf{I} \cup \mathbf{J})$ for some $\psi_{\ell}(x, \bar{z}_{\ell}) \in \mathbb{L}(\tau_T), \bar{c}_{\ell} \in {}^{\ell g(\bar{z}_{\ell})}(\mathbf{I} \cup \mathbf{J})$ such that $\psi_{\ell}(\bar{x}, \bar{c}_{\ell})$ is satisfied by infinitely many $\bar{a} \in \mathbf{I}$ if $\ell = 1, \bar{a} \in \mathbf{J}$ if $\ell = 2$ and truth value \mathbf{t} we have $M \models (\forall \bar{x}, \bar{y})[\psi_1(\bar{x}, \bar{c}_1) \land \psi_2(\bar{y}, \bar{c}_2) \to \varphi(\bar{x}, \bar{y}, \bar{c})^{\mathbf{t}}].$

Clause (h):

We just ask for the existence of the $\Phi \in \Upsilon_T^{\text{or}}$ so with τ_{Φ} countable $\supseteq \tau_T$ and type $p(\bar{x}, \bar{y}, \bar{z})$ from D(T) such that $(\ell g(\bar{y}) = \ell g(\bar{z})$ and) for any linear order I, which is well orderable $\text{EM}_{\tau}(I, \Phi)$ is a model of T of cardinality |T| + |I| and $p(\bar{x}, \bar{a}_s, \bar{a}_t)$ is realized in it iff $s <_I t$ (so O.K. for stable). $\square_{3.1}$

3.2 Claim. If T is unstable and $|T| = \aleph_{\beta_*} < \aleph_{\alpha} = \lambda$ then $\dot{I}(\lambda, T) \ge |\alpha - \beta_*|$.

Proof. In $\mathbf{L}[T]$, let Φ be as in 2.2 such that for every a linear order I we have $s, t \in I \Rightarrow \mathrm{EM}(I, \Phi) \models \varphi[\bar{a}_s, \bar{a}_t]^{\mathrm{if}(s < t)}$, where, of course, $\varphi(\bar{x}, \bar{z}) \in \mathbb{L}(\tau(T))$. First, we define for $\gamma \leq \aleph_{\alpha}$

$$J_{\gamma} =: \gamma + (\gamma)^*.$$

We can specify: the set of members of J_{γ} is $\{(\gamma, \ell, \zeta) : \ell \in \{0, 1\}, \zeta < \gamma\}$ and $(\gamma, \ell_1, \zeta_1) < (\gamma, \ell_2, \zeta_2)$ iff $\ell_1 = 0 \land \ell_2 = 1$ or $\ell_1 = \ell_2 = 0 \land \zeta_1 < \zeta_2$ or $\ell_1 = \ell_2 = 1 \land \zeta_1 > \zeta_2$.

Second, for $\beta \in [\beta_*, \alpha]$ let $J^{\beta} = \sum_{\gamma < \aleph_{\beta}} J_{\gamma} + J_{\infty}$ where $J_{\infty} = (\aleph_{\alpha} + 1) \times \mathbb{Q}$ ordered

lexicographically.

Third, let $M_1^{\beta} = \text{EM}(J^{\beta}, \Phi)$.

Lastly, $M^{\beta} := M_1^{\beta} \upharpoonright \tau_T$ - clearly a model of T of cardinality \aleph_{α} . We like to "recover", "define" \aleph_{β} from M^{β}/\cong at least when $\beta \geq \beta_*$. This is sufficient as the sequence $\langle M_{\beta} : \beta \in [\beta_*, \alpha] \rangle$ exists (in fact in $\mathbf{L}[T]$). We shall continue after stating 3.3.

<u>Discussion</u>: 1) In ZFC we could recover from the isomorphism types, stationary subsets modulo the club filter so as we get $2^{\aleph_{\alpha}}$, if, e.g., \aleph_{α} is regular and there are $2^{\aleph_{\alpha}}$ subsets of \aleph_{α} any two with a stationary difference so we get $\dot{I}(\aleph_{\alpha},T)=2^{\aleph_{\alpha}}$. But here (ZF) the stationary subsets of a regular uncountable λ may form an ultrafilter or all uncountable cardinals are singulars.

- 2) More than 3.3 is true in $\mathbf{L}[T,Y]$; $\mathrm{EM}(J,\Phi)$ satisfies \otimes_{θ} iff J has a (θ,θ) -cut (provided J has no $(1,\theta), (\theta,1), (0,\theta), (\theta,0)$ cuts), see below.
- 3) See more in 3.6 on OTOP.

- 4) Of course, we can prove theorems saying e.g.: if $\aleph_{\alpha} > |T|$ is regular, T unstable then $\dot{I}(\aleph_{\alpha}, T) \geq |\mathscr{P}(\aleph_{\alpha})/(\text{the club filter on }\aleph_{\alpha})|$.
- **3.3 Subclaim.** If $J = J^{\beta}$, $M = M^{\beta}$ are as above and $Y \subseteq \text{Ord satisfies } M \in \mathbf{L}[T,Y]$ then in $\mathbf{L}[T,Y]$ for any regular cardinal θ (of $\mathbf{L}[T,Y]$)
 - $(*) \otimes_{\theta} \Leftrightarrow \theta > \aleph_{\beta} \text{ where }$
 - \otimes_{θ} if p is a set of Δ -formulas with parameters from M of cardinality θ where

$$\Delta =: \{\varphi(\bar{x}, \bar{z}_1) \land \neg \varphi(\bar{x}, \bar{z}_1)\}$$

and any subset q of p of cardinality $< \theta$ is realized in M then some $q \subseteq p$, $|q| = \theta$ is realized in M.

Proof of Claim 3.2 from the subclaim 3.3. Why does this subclaim help us to prove the Theorem? Assume $\beta_* \leq \beta_1 < \beta_2 \leq \alpha$ and we consider M^{β_1}, M^{β_2} as above and toward a contradiction we assume that there is an isomorphism f from M^{β_1} onto M^{β_2} .

Let $Y \subseteq \text{Ord code } T, M^{\beta_1}, M^{\beta_2} \text{ and } f$. So $\mathbf{L}[T, M^{\beta_1}, Y] = \mathbf{L}[Y] = \mathbf{L}[T, M^{\beta_2}, Y]$. In this universe let θ the first cardinal greater than the ordinal $> \aleph_{\beta_1}^{\mathbf{V}}$ so $\aleph_{\beta_1}^{\mathbf{V}} < \theta \le \aleph_{\beta_1+1}^{\mathbf{V}} < \aleph_{\beta_2}$.

Question: Why we cannot prove that $\theta = \aleph_{\beta_1+1}^{\mathbf{V}}$? As possibly $\mathbf{L}[Y] \models \aleph_{\beta_1+1}^{\mathbf{V}}$ is singular or just a limit cardinal.

Note: Maybe every $\mathbf{L}[Y]$ -cardinal from $(\aleph_{\beta_1}^{\mathbf{V}}, \aleph_{\beta_1+1}^{\mathbf{V}})$ have cofinality \aleph_0 in \mathbf{V} !

But in $\mathbf{L}[Y], \aleph_{\beta}^{\mathbf{V}}, \aleph_{\beta+1}^{\mathbf{V}}$ are still cardinals so the successor of $\aleph_{\beta}^{\mathbf{V}}$ in $\mathbf{L}[Y]$ is $\leq \aleph_{\beta+1}^{\mathbf{V}}$ but in $\mathbf{L}[Y]$ this successor, θ is regular. (In \mathbf{V}, θ may not be a cardinal at all). In $\mathbf{L}[T]$ there are many possibilities for θ (it was defined from Y!) and we have built M_1^{β} before knowing who they will be in $\mathbf{L}[Y]$ so

$$\theta > \aleph_{\beta_1} \Leftrightarrow M^{\beta_1} \models \otimes_{\theta} \Leftrightarrow M^{\beta_2} \models \otimes_{\theta} \Leftrightarrow \theta > \aleph_{\beta_2}$$

(the first \Leftrightarrow by (*) of the subclaim and the second \Leftrightarrow as f is an isomorphism)

but $\aleph_{\beta_1} < \theta \leq \aleph_{\beta_2}$; contradiction.

Proof of the subclaim 3.3. I.e., in L[T, Y] we have to prove:

$$(*) \ [\otimes_{\theta} \Leftrightarrow \theta > \aleph_{\beta}]$$

First we will prove:

$$(*)_1 \ \theta \leq \aleph_{\beta} \Rightarrow \neg \otimes_{\theta}.$$

By the choice of $J = J^{\beta}$ clearly J_{θ} is an interval of J so let

$$p =: \{ \varphi(\bar{x}, \bar{a}_{(\theta,1,i)} \land \neg \varphi(\bar{x}, \bar{a}_{(\theta,0,i)}) : i < \theta \}.$$

Let $q \subseteq p, |q| < \theta$ now as θ is regular (in $\mathbf{L}[T, Y]$) for some $j < \theta$ we have

$$q \subseteq p_j = \{ \varphi(\bar{x}, \bar{a}_{(\theta,1,i)}) \land \neg \varphi(\bar{x}, \bar{a}_{(\theta,0,i)}) : i < j \}.$$

We have a natural candidate for a sequence realizing q: the sequence $\bar{a}_{(\theta,1,j)}$. Now

$$i < j \Rightarrow (\theta, 1, j) <_{J_{\theta}} (\theta, 1, i) \Rightarrow M \models \varphi[\bar{a}_{(\theta, 1, j)}, \bar{a}_{(\theta, 1, i)}]$$

$$i < j \Rightarrow (\theta, 0, i) <_{J_{\theta}} (\theta, 1, j) \Rightarrow M \models \neg \varphi[\bar{a}_{(\theta, 1, j)}, \bar{a}_{(\theta, 0, i)}].$$

So we have proved that every $q \subseteq p, |q| < \theta$ is realized in the model. Secondly, we need to show:

 \otimes no $\bar{a} \in M$ satisfies θ of formulas from p.

Assume toward contradiction that \bar{a} is a counterexample.

So we can find $n < \omega$, a finite sequence of terms $\bar{\sigma}(\bar{x}_0, \ldots, \bar{x}_{n-1})$ from $\tau(\Phi)$ and $t_0 <_J t_1 <_J \ldots <_J t_{n-1}$ such that $\bar{a} = \bar{\sigma}(\bar{a}_{t_0}, \ldots, \bar{a}_{t_{n-1}})$. Now for each ℓ for some $i_{\ell} < \theta, t_{\ell}$ is not in the interval $((\theta, 0, i_{\ell}), (\theta, 1, i_{\ell}))_J$.

Let:

$$j^* = \max[\{i_{\ell} + 1 : \ell < n\} \cup \{1\}].$$

Now consider $\varphi(\bar{x}, \bar{a}_{(\theta,1,j)} \land \neg \varphi(\bar{x}, \bar{a}_{(\theta,0,j)}))$ for $j \in [j^*, \theta)$. So $t_{\ell} <_J (\theta, 1, j) \equiv t_{\ell} <_J (\theta, 0, j)$ for $\ell = 0, \dots, n-1$ hence $M \models \varphi[\bar{\sigma}(\bar{a}_{t_0}, \dots, \bar{a}_{t_{n-1}}), \bar{a}_{(\theta,1,j)}] \Leftrightarrow M \models \varphi[\bar{\sigma}(\bar{a}_{t_0}, \dots, \bar{a}_{t_{n-1}}), \bar{a}_{(\theta,0,j)}]$. So $\bar{a} = \sigma(\bar{a}_{t_0}, \dots, \bar{a}_{t_{n-1}})$ fail the j-th formula from p for $j \in [j^*, \theta)$. So p really exemplifies the $\neg \otimes_{\theta}$. So we have proved $(*)_1$ which is one implication of the Subclaim.

Now we will prove:

$$(*)_2$$
 if $\mathbf{L}[T,Y] \models "\theta$ is regular $> \aleph_{\beta}$ " then \otimes_{θ} .

So let $p = \{ \varphi(\bar{x}, \bar{a}_i) \land \neg \varphi(\bar{x}, \bar{b}_i) : i < \theta \} \in \mathbf{L}[T, Y]$ be given. For $j < \theta$ let

$$p_j =: \{ \varphi(\bar{x}, \bar{a}_i) \land \neg \varphi(\bar{x}, \bar{b}_i) : i < j \}.$$

So some $c_j \in M$ realizes it and let $(\bar{a}_i, \bar{b}_i, \bar{c}_i) = \langle \bar{\sigma}_i^k(a_{t_0^i}, \dots, a_{t_{n_i-1}^i}) : k = 0, 1, 2 \rangle$ where $\bar{\sigma}_i^k$ is a finite sequence of terms from $\tau(\Phi)$ and $J \models t_0^i < t_1^i < \dots < t_{n_i-1}^i$; note that we can make $\langle t_\ell^i : \ell < n_i \rangle$ not to depend on k because we can add dummy variables.

As $\tau(\Phi)$ is of cardinality $\langle \theta = \operatorname{cf}(\theta) \text{ (in } \mathbf{L}[T,Y])$, for some σ_*^k, n_* the set $S = \{i : \sigma_i^k = \sigma_*^k \text{ for } k = 0, 1, 2 \text{ and } n_i = n_*\}$ is unbounded in θ . Recall

$$J^{\beta} = \sum_{\gamma < \aleph_{\beta}} J_{\gamma} + (\aleph_{\alpha} + 1) \times \mathbb{Q}.$$

So for some $m_i \leq n_*$

$$t_{\ell}^{i} \in \sum_{\gamma < \aleph_{\beta}} J_{\gamma} \Leftrightarrow \ell < m_{i}$$

shrinking S without loss of generality $i \in S \Rightarrow m_i = m_*$.

$$\underline{\text{Now}} \ \mathbf{L}[T,Y] \models \text{``}|\sum_{\gamma \leq \aleph_{\beta}} J_{\gamma}| \leq \sum_{\gamma \leq \aleph_{\beta}} |J_{\gamma}| = \sum_{\gamma \leq \aleph_{\beta}} (|\gamma| + \aleph_{0}) \leq \aleph_{\beta} < \theta = \text{cf}(\theta)\text{''}.$$

So without loss of generality

$$\circledast_1 \ell < m_* \Rightarrow t_\ell^i = t_\ell^* \text{ for } i \in S \text{ and for } \ell \in [m_*, n_*) \text{ let } t_\ell^i = (\varepsilon_\ell^i, q_\ell^i) \text{ where } q_\ell^i \in \mathbb{Q}.$$

Clearly for q_{ℓ}^{i} there are \aleph_{0} possibilities so without loss of generality, for each $\ell \in [m_{*}, n_{*})$

- $\circledast_2^{\ell} q_{\ell}^i = q_{\ell}^* \text{ for } i \in S,$
- \circledast_3^{ℓ} $\langle \varepsilon_{\ell}^i : i \in S \rangle$ is <u>constant</u> say ε_{ℓ}^* <u>or is strictly increasing</u> with limit ε_{ℓ}^* and is strictly increasing iff $\ell \in u$

so without loss of generality

- \circledast_4 (i) if $\ell_1 \neq \ell_2$ are in the interval $[m_*, n_*)$ and $\varepsilon_{\ell_1}^* < \varepsilon_{\ell_2}^*$ then $i, j \in S \Rightarrow \varepsilon_{\ell_1}^i \leq \varepsilon_{\ell_2}^* < \varepsilon_{\ell_2}^j$
 - (ii) if $\ell_1 \neq \ell_2 \in [m_*, n_*)$ and $\varepsilon_{\ell_1}^i = \varepsilon_{\ell_2}^* \wedge \ell_1 \in u \wedge \ell_2 \notin u$ and i < j are in S then $\varepsilon_{\ell_1}^i < \varepsilon_{\ell_2}^j$.

We choose $t_0 <_J t_1 <_J \ldots <_J t_{n-1}$ which satisfies

- \circledast_5 (a) if $\ell < m_*$ then $t_\ell = t_\ell^*$
 - (b) if $\ell \in [m_*, n_*)$ and $\langle t^i_\ell : i \in S \rangle$ is constant then $t_\ell = t^*_\ell$
 - (c) if $\ell \in [m_*, n_*), \langle t^i_{\ell} : i \in S \rangle$ is not constant (i.e. $\ell \in u$) then: (recall that $\langle q^i_{\ell} : i \in S \rangle$ is constantly $q^*_{\ell}, \langle \varepsilon^i_{\ell} : i \in S \rangle$ is strictly increasing with limit ε^*_{ℓ}) we choose $t_{\ell} = (\varepsilon_{\ell}, q_{\ell})$ such that $\varepsilon_{\ell} = \varepsilon^*_{\ell}, q_{\ell} = \min(\{0\} \cup \{q^*_k : k \in [m^*, n^*)\}) - n^* + \ell$ (the computation is in \mathbb{Q} !)

Hence

- \circledast_6 (α) q_ℓ is q_k^* for every $k \in [m^*, n_*)$ when $\ell \in u$
 - (β) if $\varepsilon_{\ell}^* = \varepsilon_k^*$ and $\ell, k \in u$ then $q_{\ell} < q_k \equiv \ell < k$.

Now note that:

 \circledast_7 for $\varepsilon < \zeta < \theta$ from S, in J the quantifier free types of $\langle t_\ell^{\varepsilon} : \ell < n_* \rangle \cap \langle t_\ell : \ell < n_* \rangle$ and $\langle t_\ell^{\varepsilon} : \ell < n_* \rangle \cap \langle t_\ell^{\zeta} : \ell < n_* \rangle$ are equal [all the shrinking was done for this].

Now for $\varepsilon < \zeta$ from S, by the original choice above $M^{\beta} \models \varphi[\bar{c}_{\zeta}, \bar{a}_{\varepsilon}] \land \neg \varphi[\bar{c}_{\zeta}, \bar{b}_{\varepsilon}]$ that is: $M_{1}^{\beta} \models \varphi[\bar{\sigma}_{*}^{0}(a_{t_{0}^{\zeta}}, \ldots), \bar{\sigma}_{\varepsilon}^{1}(a_{t_{0}^{\varepsilon}}, \ldots)] \land \neg \varphi[\bar{\sigma}_{*}^{0}(a_{t_{0}^{\zeta}}, \ldots), \bar{\sigma}_{\varepsilon}^{2}(a_{t_{0}^{\varepsilon}}, \ldots)].$

By the last sentence and \circledast_7+ indiscernibility of $\langle \bar{a}_t : t \in J \rangle$ in M_1^{β} we have $M \models \varphi[\bar{\sigma}^0_*(a_{t_0},\ldots),\bar{\sigma}^1_{\varepsilon}(a_{t_0^{\varepsilon}},\ldots)] \wedge \neg \varphi[\bar{\sigma}^0_*(a_{t_0},\ldots),\bar{\sigma}^2_{\varepsilon}(a_{t_0^{\varepsilon}},\ldots)].$

Let $\bar{c} = \bar{\sigma}^0_*(a_{t_0}, \dots)$ in M_1^{β} -sense, so $\varepsilon \in S \Rightarrow M \models \varphi[\bar{c}, \bar{a}_{\varepsilon}] \land \neg \varphi[\bar{c}, \bar{b}_{\varepsilon}]$. Hence $\{\varphi(\bar{x}, \bar{a}_{\varepsilon}) \land \neg \varphi(\bar{x}, \bar{b}_{\varepsilon}) : \varepsilon \in S\}$ is realized in M^{β} and $\mathbf{L}[T, Y] \models "|S| = \theta$ " as promised. $\Box_{3.2}$

3.4 Claim. If T is stable not uni-dimensional, $|T| = \aleph_{\beta} < \aleph_{\alpha} = \lambda \ \underline{then} \ \dot{I}(\lambda, T) \ge |\alpha - \beta|$.

Proof. As in 2.9; if $\gamma \in [\beta, \alpha]$ then there is a model M of T of cardinality λ such that M satisfies $(*)_{\gamma}$ but not $\beta \leq \gamma_1 < \gamma \Rightarrow \neg(*)_{\gamma_1}$ where

- (*) $_{\gamma}$ if $\mathbf{I}, \mathbf{J} \subseteq M$ are infinite orthogonal indiscernible sets and $|\mathbf{I}| = \lambda$ then $|\mathbf{J}| \leq \aleph_{\gamma}$.
- 3.5 Conclusion. If $\lambda = \aleph_{\alpha} > \aleph_{\beta} = |T|$ and $\dot{I}(\lambda, T) < |\alpha \beta|$ then (in $\mathbf{L}[T, Y]$ when $Y \subseteq \text{Ord}$)

- \boxtimes_T (a) T is stable and uni-dimensional
 - (b) T is superstable
 - (c) T has no two cardinal models
 - (d) D(T) has cardinality $\leq |T|$ or cardinality $< |\alpha \beta|$.

Proof. T is stable by 3.2 and uni-dimensional by 3.4 so clause (a) holds. This implies clause (c), see [Sh:c, V,§6]. Clause (d) is trivial by now and clause (b) follows from clause (a) by Hrushovski [Hr89d]. $\square_{3.5}$

- **3.6 Claim.** In 3.5 we can add to \boxtimes_T also clause (e) and if T is countable also clause (f) where
 - \boxtimes_T (e) T fails the OTOP (see [Sh:c, XII,Def.4.1,pg.608] or 3.7(1) below)
 - (f) T has the prime existence property (see [Sh:c, XII, Def.4.2, pg.608] or 3.7(2) below) hence $for \mathfrak{C}_T \text{ a model of } T \text{ with universe } |\mathfrak{C}_T| \subseteq \mathbf{L}:$

for any non-forking tree $\langle N_{\eta} : \eta \in \mathcal{T} \rangle$ of models $N_{\eta} \prec \mathfrak{C}_{T}$, there is a prime (even primary, i.e. $\mathbf{F}_{\aleph_{0}}^{t}$ -primary) model $N \prec \mathfrak{C}_{T}$ over $\cup \{N_{\eta} : \eta \in \mathcal{T}\}$, it is unique up to isomorphism over $\cup \{N_{\eta} : \eta \in \mathcal{T}\}$.

Proof. Clause (e) holds exactly as for stability, i.e., as in 3.2 only the formulas $\varphi(\bar{x}, \bar{y})$ are not first order but an infinite conjunction of such formulas. Clause (f) follows by [Sh:c, XII], i.e., it holds in any $\mathbf{L}[T, Y]$ which suffices.

- **3.7 Definition.** 1) T has OTOP if for some type $p = p(\bar{x}, \bar{y}, \bar{z})$ in $\mathbb{L}(\tau_T)$ the theory T has it for p, which means that for every λ for some model M of T with well ordered universe and $\bar{b}_{\alpha} \in {}^{\ell g(\bar{y})}M$, $\bar{c}_{\alpha} \in {}^{\ell g(\bar{z})}M$, for $\alpha, \beta < \lambda$ we have: for any $\alpha, \beta < \lambda$ the model M realizes the type $p(\bar{x}, \bar{b}_{\alpha}, \bar{c}_{\beta})$ iff $\alpha < \beta$.
- 2) T has the prime existence property when for every triple (M_0, M_1, M_2) in stable amalgamation in a model \mathfrak{C}_T of T such that $|\mathfrak{C}_T|$ is well orderable (so $M_\ell \prec \mathfrak{C}_T$), the set of isolated types is dense in $\mathbf{S}^m(M_1 \cup M_2)$ for every m.

- **3.8 Claim.** [T countable] We can add clause (g) below to \boxtimes_T from 3.5 + 3.6:
 - \boxtimes_T (g) if clause (A) <u>then</u> for some M' clause (B) below holds (both in $\mathbf{L}[T,Y]$) where
 - (A) (a) $M_{\emptyset} \prec M_{\{i\}} \prec M^*$ are countable models of T for $i < \omega \times 2$
 - (β) $(M_{\{i\}}, c)_{c \in M_{\emptyset}} \cong (M_{\{0\}}, c)_{c \in M_{\emptyset}} \text{ for } i < \omega \text{ that is } M_{\{i\}} \text{ is isomorphic to } M_{\{0\}} \text{ over } M_{\emptyset} \text{ for } i < \omega$
 - $(\gamma) \quad (M_{\{\omega+i\}}, c)_{c \in M_{\emptyset}} \cong (M_{\{\omega\}}, c)_{c \in M_{\emptyset}} \text{ that is } M_{\{\omega+i\}} \text{ is isomorphic}$ $\text{to } M_{\{\omega\}} \text{ over } M_{\emptyset} \text{ for } i < \omega$
 - (δ) $\{M_{\{i\}}: i < \omega \times 2\}$ is independent over M_{\emptyset} inside M^*
 - $(\varepsilon) \quad M^* \ \textit{is prime over} \cup \{M_{\{i\}}: i < \omega \times 2\}$
 - (B) (α) $M_{\emptyset} \prec M' \prec M^*$
 - $(\beta) \quad (M',c)_{c \in M_{\emptyset}} \cong (M_{\{0\}},c)_{c \in M_{\emptyset}}$
 - (γ) $\langle M_{\{i\}} : i < \omega \rangle^{\frown} \langle M' \rangle$ is independent over M_{\emptyset} .
- 3.9 Remark. 1) We can formulate (B) closer to \circledast_6 inside the proof of 3.10.
- 2) We can omit "T countable" but then have to change Y with the same proof.
- 3) We know more on T's satisfying \boxtimes_T of 3.5 by Laskowski [Las88] and Hart-Hrushovski-Laskowski [HHL00].

Proof. Note that $|T| = \aleph_0$ and choose the ordinals β_* , α_* such that $\beta_* = 0$, $\lambda = \aleph_{\alpha_*}$; most of the proof we do not use $\beta_* = 0$ but we use $\boxtimes_T (a) - (f)$.

We do more than is strictly necessary for the proof; we use \odot to denote definitions, working in $\mathbf{L}[T,Y]$ if not said otherwise and \mathfrak{C}_Y is a monster for T in $\mathbf{L}[T,Y]$:

- \odot_1 (a) for a model $M \prec \mathfrak{C}_Y$ let $\mathbf{S}_Y^{c,\theta}(M) = \{ \operatorname{tp}(\bar{a}, M, N) : M \prec N \prec \mathfrak{C}, \|N\| \leq \theta \text{ and } \bar{a} \text{ enumerates } N \}$, omitting θ means some θ
 - (b) in this case we say N realizes p
 - (c) if $p = \operatorname{tp}(\bar{a}, M, N)$ is as above, then we denote |p| = ||N||,
- \odot_2 for $\bar{\alpha} = \langle \alpha_{\varepsilon} : \varepsilon < \zeta \rangle$ and $\bar{p} = \langle p_{\varepsilon} : \varepsilon < \zeta \rangle, p_i \in \mathbf{S}_Y^c(M)$, we say N is $(\bar{p}, \bar{\alpha})$ -constructed over M when there is \bar{M} such that
 - (a) $\bar{M} = \langle M_{\{i\}} : i < \alpha^{\zeta} \rangle$, where $\alpha^{\varepsilon} = \sum_{\xi < \varepsilon} \alpha_{\xi}$ for $\varepsilon \le \zeta$

- (b) $M_{\{i\}}$ realizes p_{ε} if $i \in [\alpha^{\varepsilon}, \alpha^{\varepsilon} + \alpha_{\varepsilon})$
- (c) $\langle M_{\{i\}} : i < \alpha^{\zeta} \rangle$ is independent over M
- (d) N is primary over $\bigcup_{i < \alpha^{\zeta}} M_{\{i\}}$
- \odot_3 we say N is \bar{p} -constructed over M if this holds for some $\bar{\alpha}$
- \odot_4 if $M \prec N \prec \mathfrak{C}_Y, p \in \mathbf{S}_Y^c(M)$ then we say q lifts p or (p, M) to N when $q \in \mathbf{S}_Y^c(N)$ and for some M_1, N_1 realizing p, q respectively, $\operatorname{tp}(M_1, N)$ does not fork over M and N_1 is primary over $N \cup M_1$
- \odot_5 for $M \prec \mathfrak{C}$ and $p_1, p_2 \in \mathbf{S}_Y^c(M)$ we say p_2 pushes p_1 (in $\mathbf{L}[T, Y]$) when for some ordinals α_1, α_2 there are $M'_{\{i\}}$ for $i < \alpha_2 + \alpha_1$ and \bar{M}, M^*, M' satisfying
 - (a) M^* is $(\langle p_1, p_2 \rangle, \langle \alpha_1, \alpha_2 \rangle)$ -constructed over M as witnessed by $\bar{M}' = \langle M'_{\{i\}} : i < \alpha_1 + \alpha_2 \rangle$
 - (b) $M \prec M' \prec M^*$
 - (c) M' realizes p_1
 - (d) $\langle M_{\{i\}} : i < \alpha_1 \rangle^{\hat{}} \langle M' \rangle$ is independent over M
- \odot_6 (α) assume $p_{\varepsilon}, q_{\varepsilon} \in \mathbf{S}_Y^c(M)$ for $\varepsilon < \varepsilon(*)$; we say $(\bar{p}, \bar{\alpha})$ is equivalent to $(\bar{q}, \bar{\beta})$ when $\bar{\alpha} = \langle \alpha_{\varepsilon} : \varepsilon < \varepsilon(*) \rangle, \bar{\beta} = \langle \beta_{\varepsilon} : \varepsilon < \varepsilon(*) \rangle$ and there is M' which is both $(\bar{p}, \bar{\alpha})$ -constructed over M and $(\bar{q}, \bar{\beta})$ -constructed over M
 - (β) we may write p instead of $\langle p \rangle$, q instead of $\langle q \rangle$, and omitting $\bar{\alpha}, \bar{\beta}$ means "for some $\bar{\alpha}, \bar{\beta}$ ".
- \circledast_1 if $p_1, p_2 \in \mathbf{S}_Y^c(M)$ and q pushes p then in \odot_5 without loss of generality $\alpha, \beta \leq ||M|| + |T| + |p_1| + |p_2|$.

[Why? By the DLST argument.]

- \odot_7 (a) let $AP_Y^{\theta} = \{(M, p_1, q_1) : \text{ in } \mathbf{L}[T, Y], M \prec \mathfrak{C}_Y \text{ and } p_1, q_1 \in \mathbf{S}_Y^c(M) \text{ have cardinality } \leq \theta\}$
 - (b) $AP_Y^{\theta} \models \text{``}(M_1, p_1, q_1) \leq (M_2, p_2, q_2)\text{''}$ means that
 - (α) both triples are from AP_Y^{θ}
 - (β) M_2 is (p_1, q_1) -constructed over M_1
 - (γ) p_2, q_2 lift p_1, q_1 over M_2 respectively
- \circledast_2 if $AP_V^{\theta} \models "(M_1, p_1, q_1) \leq (M_2, p_2, q_2)"$ and q_2 pushes p_2 then q_1 pushes p_1 .

[Why? Straight.]

- \circledast_3 if $(M, p_1, q_1) \in AP_Y^{\aleph_{\beta(*)}}$ and p_1 does not push q_1 then we can find $\mu_0, \mu_1, M_*, p_2, q_2$ and r such that
 - (a) $AP_Y^{\aleph_0} \models \text{``}(M, p_1, q_1) \leq (M_*, p_2, q_2)\text{''}$ hence by \circledast_2 the type p_2 does not push q_2 , (this is the only point where we use " p_1 does not push q_1 ")
 - (b) $||M_*|| = \mu_0$
 - (c) $r \in \mathbf{S}_{V}^{c,\mu_0}(M_*)$ and $\aleph_{\beta(*)} \leq \mu_0 < \mu_1 < \lambda$
 - (d) $(\langle p_2, q_2 \rangle, \langle \lambda, \mu_1 \rangle)$ is equivalent to $(\langle r \rangle, \langle \lambda \rangle)$, see \odot_6 .

[Why? For every $\mu \in [\aleph_{\beta(*)}, \lambda)$ let N^{μ} be $(\langle p_1, q_1 \rangle, \langle \lambda, \mu \rangle)$ -constructed over M as witnessed by $\langle N_i : i < \lambda + \mu \rangle$.]

As we are assuming that $\dot{I}(\lambda, T) < |\alpha_* - \beta_*|$, there are μ_0, μ_1 such that $|T| = \aleph_{\beta(*)} \le \mu_0 < \mu_1 < \lambda$ and there is an isomorphism $f \in \mathbf{V}$ from N^{μ_0} onto N^{μ_1} ; of course f is not necessarily from $\mathbf{L}[T, Y]$. We now work in $\mathbf{L}[T, Y, f]$ and in the end we use absoluteness (here we use "T countable").

Now by the DLST argument and properties of $\mathbf{F}_{\aleph_0}^t$ -primary we can find (u_0, u_1, M^0, M^1) such that

- (*)₄ (a) u_{ℓ} is a subset of $\lambda + \mu_{\ell}$ of cardinality μ_0 satisfying $|u_{\ell} \cap \lambda| = \mu_0 = |u_{\ell} \setminus \lambda|$ and $[\lambda, \lambda + \mu_0) \subseteq u_{\ell}$ for $\ell = 0, 1$
 - (b) $M^{\ell} \prec N^{\mu_{\ell}}$ is primary over $M \cup \{N_i : i \in u_{\ell}\}$ for $\ell = 0, 1$
 - (c) $N^{\mu_{\ell}}$ is primary over $M^{\ell} \cup \{N_i : i \in (\lambda + \mu_{\ell}) \setminus u_{\ell}\}$ for $\ell = 0, 1$
 - (d) f maps M^0 onto M^1 .

For $i \in (\lambda + \mu_{\ell}) \setminus u_{\ell}$ let $N_{\ell,i} \prec N^{\mu_{\ell}}$ be primary over $M^{\ell} \cup N_{\{i\}}$ such that N^{ℓ} is primary over $\cup \{N_{\ell,i} : j \in (\lambda + \mu_{\ell}) \setminus u_{\ell}\}$; clearly

- $(*)_5$ for $\ell = 0, 1$
 - (a) $M^{\ell} \prec N_{\ell,i} \prec N^{\ell}$
 - (b) $(N_{\ell,i},c)_{c\in M^{\ell}} \cong (N_{\ell,j},c)_{c\in M^{\ell}}$ when $i,j\in \lambda\setminus u_{\ell}$ or $i,j\in (\lambda+\mu_{\ell})\setminus \lambda\setminus u_{\ell}$
 - (c) $\langle N_{\ell,i} : i \in (\lambda + \mu_{\ell}) \setminus u_{\ell} \rangle$ is independent over M^{ℓ} and N^{ℓ} is primary over their union.

Choose $\gamma_1 \in \lambda \setminus u_1, \gamma_2 \in [\lambda, \lambda + \mu_1) \setminus u_1$, so $(M^1, \operatorname{tp}(N_{1,\gamma_1}, M^1_{\emptyset}), \operatorname{tp}(N_{1,\gamma_2}, M^1_{\emptyset}))$ can serve as (M_*, p_1, q_1) and r is $f(\operatorname{tp}(M_{0,\gamma}, M^0))$ for any $\gamma \in \lambda \setminus u_0$.

So we have finished proving \circledast_3 .

 \circledast_4 assume \bar{p}, \bar{q} are sequences of members of $\mathbf{S}^c_Y(M)$ and $(\bar{p}, \bar{\alpha}), (\bar{q}, \bar{\beta})$ are equivalent and $M \prec N$ and $p'_{\varepsilon} \in \mathbf{S}^c_Y(N)$ lift p_{ε} for $\varepsilon < \ell g(\bar{p})$ and $q'_{\varepsilon} \in \mathbf{S}^c_Y(N)$ lift q_{ε} for $\varepsilon < \ell g(\bar{q})$ then $(\langle p'_{\varepsilon} : \varepsilon < \ell g(\bar{p}) \rangle, \bar{\alpha})$ and $(\langle q'_{\varepsilon} : \varepsilon < \ell g(q) \rangle, \bar{\beta})$ are equivalent.

[Why? By properties of "primary".]

 \circledast_5 if $p, q \in \mathbf{S}_Y^{c,\theta}(M)$ are equivalent then $(p, \theta), (q, \theta)$ are equivalent.

[Why? By DLST.] Note

 \circledast_6 in \circledast_3 we can conclude p_2, r are equivalent.

[Why? In clause (d) of \circledast_3 , let N be the model and let a witness for N being (r, λ) -constructed be $\langle N_i^r : i < \lambda \rangle$ and for N being $(\langle p_2, q_2 \rangle, \langle \lambda, \mu_1 \rangle)$ -constructed be $\langle N_i^* : i < \lambda + \mu_1 \rangle$. Let $u_0 \subseteq \lambda, u_1 \subseteq \lambda + \mu_1$ be of cardinality $\mu_1, [\lambda, \lambda + \mu_1) \subseteq u_1$ and M'_* be such that:

- $(*)_6$ (a) $M'* \prec N$
 - (b) M'_* is primary over $\cup \{N_i^2 : i \in u_1\}$
 - (c) M'_* is primary over $\cup \{N_i^* : i \in u_2\}$
 - (d) N is primary over $\cup \{N_i^2 : i \in \lambda \setminus u_1\} \cup M'_*$
 - (e) N is primary over $\cup \{N_i^* : i \in \lambda + \mu_1 \setminus u_2\}.$

The liftings r', p'_2 of r, p_2 to M'_* are equivalent, so we "collapse" to cardinality μ_0 getting M''_* so M''_* is (r, μ_0) -constructed over M_* and (p_2, μ_0) -constructed over M_* . Then find liftings $r'', p''_2 \in \mathbf{S}_Y^c(M'_*)$ of r, p respectively, so r'', p''_2 are equivalent naturally but M''_*, M_* are isomorphic over M by an isomorphism mapping r'', p''_2 to r, p_2 so we get that r, p_2 are equivalent as required.]

 \circledast_7 if M' is (p_2, μ_0) -constructed over M_* then q_2 is realized in M'.

[Why? Assume M' is a counterexample. Look again at the proof of \circledast_3 , so M_* is M^1 there, and so M' is (p_2, λ) -constructed over $M^1 = M_*$ and N^1 is (p_2, λ) -constructed over $M^1 = M_*$, so by uniqueness of primary also in N^1 we cannot find $N' \prec N^1$ realizing q_2 . But for any $\gamma \in [\lambda, \lambda + \mu_1] \setminus u_2$ the model $f^{-1}(N_{1,\gamma})$ contradict this.]

Now we can prove 3.8. Let $p, q \in \mathbf{S}_Y^{c,\aleph_{\beta(*)}}(M_{\emptyset})$ be types which $M_{\{0\}}, M_{\{\omega\}}$ respectively realizes. Let $(M, p_1, q_1) = (M_{\emptyset}, p, q)$.

So by \circledast_7 , there are $\mu_0 < \lambda$ and Y_1 and $M_2 \in \mathbf{L}[T, Y_1], M_2$ which is $(\langle p, q \rangle, \langle \mu_0, \mu_0 \rangle)$ constructed over M_\emptyset and p_2, q_2 lifting of p, q in $\mathbf{S}_{Y_1}^{c,\mu_0}(M_2)$ as there. So by DLST

we can find such M'_2, p'_2, q'_2 for the case $\mu_0 = \aleph_{\beta(*)}$, but this is absolute as $\beta(*) = 0$. Also it gives the required result.

3.10 Theorem. [ZF] If T is countable and \boxtimes_T below holds, then (recalling 0.12(2)) in every cardinal $\mu \geq \Upsilon(\mathscr{P}(\omega))$ we have $\dot{I}(\mu,T)$ is $\leq |\mathscr{F}^*/E|$ where $\mathscr{F}^* = \{f : f \text{ a function from } \mathscr{P}(\omega) \text{ to } \omega + 1\}$, for some equivalence relation E on the set of those functions where:

$$\boxtimes_T (a) - (d)$$
 from 3.5

- (e) (f) from 3.6
- (g) from 3.8.

Remark. 1) Countability of T is not used (if we write $\mathscr{P}(|T|)$ instead of $\mathscr{P}(\omega)$), but the gain is not substantial. This applies to 3.11, 3.13, too.

2) Fuller more accurate information is given in 3.13.

Proof. Let N be a model of T of cardinality μ so without loss of generality with universe μ , we work in $\mathbf{L}[T,N]$ and we shall analyze it. Now we first choose a countable $M_{\emptyset} \prec N_{\ell}$. As T is superstable, uni-dimensional we can find $\varphi(x,\bar{y}) \in \mathbb{L}(\tau_T)$ and $\bar{a} \in {}^{\ell g(\bar{y})}(M_{\emptyset})$ such that $\varphi(x,\bar{a})$ is weakly minimal.

We can find $\langle a_{\alpha} : \alpha < \mu \rangle$ such that:

- \circledast_1 (a) $a_{\alpha} \in \varphi(N, \bar{a}_{\ell}) \backslash M_{\emptyset}$
 - (b) $\{a_{\alpha} : \alpha < \mu\}$ is independent in N over M_{\emptyset} (in particular with no repetitions)
 - (c) modulo (a) + (b) the set $\{a_{\alpha} : \alpha < \mu\}$ is maximal hence
 - (d) $\varphi(N, \bar{a}) \subseteq \operatorname{acl}(M_{\emptyset} \cup \{a_{\alpha} : \alpha < \mu\}).$

Let $f \in \mathbf{L}[T, N]$ be a function from μ to μ such that $f(\alpha) \leq \alpha$ and $(\forall \beta < \mu)(\exists^{\mu} \alpha < \mu)(f(\alpha) = \beta)$. Now we try to choose $(M_{\{\alpha\}}, b_{\alpha})$ by induction on $\alpha < \mu$ such that

- \circledast_2 (a) $b_{\alpha} \in \varphi(N, \bar{a})$
 - (b) $b_{\alpha} \notin \operatorname{acl}(M_{\emptyset} \cup \{b_{\beta} : \beta < \alpha\})$
 - (c) $M_{\{\alpha\}} \prec N$ is $\mathbf{F}_{\aleph_0}^c$ -primary over $M_{\emptyset} \cup \{b_{\alpha}\}$, see [Sh:c, IV]
 - (d) if $\alpha = 2\beta + 1$ and we can find $(M_{\{\alpha\}}, b_{\alpha})$ satisfying (a) + (b) + (c) and $(M_{\{\alpha\}}^{N_{\alpha}}, c)_{c \in M_{\emptyset}} \cong (M_{\{f(\beta)\}}, c)_{c \in M_{\emptyset}}$ then $(M_{\{\alpha\}}, b_{\alpha})$ satisfies this
 - (e) if $\alpha = 2\beta$ and $\gamma_{\alpha} = \min\{\gamma : a_{\gamma} \notin \operatorname{acl}(M_{\emptyset} \cup \{b_{\varepsilon} : \varepsilon < 2\beta\})\}$ then $b_{\alpha} = a_{\gamma_{\alpha}}$.

[Why can we can carry the induction? We can ignore clause (d) as if its hypothesis hold, then clause (e) is irrelevant, and this hypothesis says that we can fulfil clause (a),(b),(c),(d). Also if $\alpha = 2\beta + 1$ and the further assumption of (d) fail then we can act as in clause (e). Also in all cases by cardinality considerations recalling $|\varphi(M,\bar{a})| = ||M||$ by (c) of \boxtimes_T of 3.5 there is b_α satisfying clauses (a) + (b) and if clause (e)'s assumption holds, without loss of generality also its conclusion.

Let $B_{\alpha} = \operatorname{acl}_{M}(M_{\emptyset} \cup \{b_{\alpha}\})$. By the choice of $\varphi(x, \bar{a})$ if $\varphi(x, \bar{a}) \in p \in \mathbf{S}(B_{\alpha}, M)$ then either p forks over \bar{a} hence is algebraic hence realized in B_{α} or p does not fork over \bar{a} hence is finitely satisfiable in M_{\emptyset} . Let $\langle b_{\alpha,i} : i < i_{\alpha} \rangle$ be a maximal sequence of members of M such that for each i for some formula $\varphi(x, \bar{c}_{\alpha,i}) \in \operatorname{tp}(b_{\alpha,i}, B_{\alpha} \cup \{b_{\alpha,j} : j < i\})$ hence no extension in $\mathbf{S}(B_{\alpha} \cup \{b_{\alpha,j} : j < i\})$ forking over $\bar{c}_{\alpha,i}$. By [Sh 31] there is $M_{\{\alpha\}} \prec N$ with universe $B_{\alpha} \cup \{b_{\alpha,i} : i < i_{\alpha}\}$.

So we are done.]

 $\circledast_3 \langle b_\alpha : \alpha < \mu \rangle$ satisfies the requirements on $\langle a_\alpha : \alpha < \mu \rangle$.

[Why? Easy to check.]

So $\langle M_{\{\alpha\}} : \alpha < \mu \rangle$ is independent over M_{\emptyset} inside N hence (by $\boxtimes_T(f)$) there is $N' \prec N$ primary over $\cup \{M_{\{\alpha\}} : \alpha < \mu\}$ and by $\circledast_1(d)$ include $\varphi(N, \bar{a})$ hence by $\boxtimes_T(c)$ we have N' = N.

We can find a set S and a partition $\langle I_t : t \in S \rangle$ of μ such that: for $\ell = 1, 2$ and $\alpha, \beta < \mu$ we have

$$\circledast_4 (M_{\{\alpha\}}, a_{\alpha}, c)_{c \in M_{\emptyset}}$$
 is isomorphic to $(M_{\{\beta\}}, a_{\beta}, c)_{c \in M_{\emptyset}}$ iff $\bigvee_{t \in S} \{\alpha, \beta\} \subseteq I_t$.

Now how large can |S| be? It is, in $\mathbf{V}, \leq |\mathscr{P}(\omega)|^{\mathbf{L}[T,N]} \leq |\mathscr{P}(\omega)|^{\mathbf{V}}$ (there is a function from a subset of $\mathscr{P}(\omega)$ onto this set). So $|S| < \theta(\mathscr{P}(\omega))$, but $\mathbf{L}[T,N] \models$ " $ZFC + |S| \leq 2^{\aleph_0} = |\mathscr{P}(\omega)|$ " so there is a well ordering of $\mathscr{P}(\omega) \cap \mathbf{L}[T,N] = \mathscr{P}(\omega)^{\mathbf{L}[T,N]}, |S| \leq |\mathscr{P}(\omega)^{\mathbf{L}[T,N]}| < |S| < \Upsilon(\mathscr{P}(\omega))$ and $\mathbf{L}[T,N] \models$ " $|S| \leq 2^{\aleph_0}$ ". Now we shall prove:

$$\circledast_5$$
 if $t \in S$ and $\mathbf{L}[T, N] \models "\aleph_0 \leq |I_t| < \mu"$ then for some $\alpha < \mu$ we have $(\forall s \in S)(|I_s \setminus \alpha| < \aleph_0)$ and $\mu < \Upsilon(\mathscr{P}(\omega))$.

Clearly \circledast_5 helps because " $\mu < \Upsilon(\mathscr{P}(\omega))$ " contradict an assumption on μ ".

Why \circledast_5 holds? Let $\alpha(*) = \operatorname{Min}(I_t)$, it is well defined as $I_t \neq \emptyset$ because " $\aleph_0 \leq |I_t|$ " was assumed. Let $J = \{2\beta + 1 : f(\beta) = \alpha(*)\}$. If $2\beta + 1 \in J \Rightarrow 2\beta + 1 \in I_t$, then we get $|I_t| \geq |\{2\beta + 1 : f(\beta) = \alpha(*)\}| = \mu$ hence the assumption " $|I_t| < \mu$ " is contradicted, so assume that $\alpha = 2\beta + 1 \in J \setminus I_t$. By clause (d) of \circledast_2 apply to $\alpha = 2\beta + 1$, we know that if (N', b) satisfies the demands on $(M_{\{\alpha\}}, b_{\alpha})$ in clauses (a),(b),(c) (i.e., $b_{\alpha} \in \varphi(N, \bar{a}) \setminus \operatorname{acl}(M_{\emptyset} \cup \{b_{\varepsilon} : \varepsilon < \alpha\})$ and $N' \prec N$ is $\mathbf{F}_{\aleph_0}^c$ -primary

over $M_{\emptyset} \cup \{b\}$) then $(N',c)_{c \in M_{\emptyset}} \ncong (M_{\{\alpha(*)\}},c)_{c \in M_{\emptyset}}$. This implies that $I_t \subseteq \alpha$. As it is infinite, by $\boxtimes_T(g)$ we get $(\forall s \in S)(|I_s \setminus \alpha| < \aleph_0)$ and recall $|\mu \setminus \alpha| = \mu$. So $\{\min(I_s \setminus \alpha) : s \in S \text{ and } I_s \nsubseteq \alpha\}$ is a subset of μ of cardinality μ (working in $\mathbf{L}[T,N]$) and there is a one-to-one mapping from it into $\mathscr{P}(\omega)$ (using the isomorphism types of $(M_{\{\alpha\}},c)_{c \in M_{\emptyset}}$). This gives $\mu < \Upsilon(\mathscr{P}(\omega))$. But this contradicts an assumption on μ .

So we know

 \circledast_6 if I_t is infinite then it has cardinality μ .

Let $f = f_N = f_{N,M_{\emptyset},\varphi(x,\bar{a})}$ be the partial function from $\mathscr{P}(\omega)$ into $\omega + 1$ defined as follows: if $t \in S$ and $\eta \in \mathscr{P}(\omega)$ codes⁴ a model isomorphic to $(M_{\alpha},c)_{c\in M_{\emptyset}}$ for $\alpha \in I_t$ then $f_N(\eta) = |I_t|$ if I_t is finite and $f_N(\eta) = \omega$ otherwise: of course, the choice of f_N is unique if we use the canonical well ordering of $\mathbf{L}[T,N]$ to make our choices in particular of $M_{\emptyset}, \varphi(x,\bar{a})$, but we could use "any such f" so increasing \mathscr{F}_{μ} below (and fix the coding).

Now in V for any model M of T of cardinality μ we define

 $\mathscr{F}_M = \{(f_N, N \upharpoonright \omega, \varphi(x, \bar{a})) : N \text{ is a model with universe } \mu \text{ isomorphic to } M \text{ such that } N \upharpoonright \omega \prec N \text{ so can serve as } M_\emptyset \text{ and } \bar{a} \in {}^{\omega >} N \text{ and } \varphi(x, \bar{a}) \text{ is weakly minimal} \}$

 $\mathbf{F}^* = \bigcup \{ \mathscr{F}_M : M \text{ a model of } T \text{ of cardinality } \mu \}.$

Clearly

- \circledast_7 (a) \mathscr{F}_M depends just on M/\cong
 - (b) if $\mathscr{F}_{M_1} \cap \mathscr{F}_{M_2} \neq \emptyset$ then $M_1 \approx M_2$ hence $\mathscr{F}_{M_1} = \mathscr{F}_{M_2}$ so there is an equivalence relation $E_{T,\mu}$ on a subset of \mathscr{F}^* such that the \mathscr{F}_M 's are its equivalence classes
 - (c) the number of models of T in μ up to isomorphism is equal to the number of $E_{T,\mu}$ -equivalence classes.

So we are done. $\square_{3.10}$

⁴see more details on this and similar points in the proof of 3.13

3.11 Claim. [T countable] The demand \boxtimes_T from 3.10 is absolute (property of T).

Proof. The new point is $\boxtimes_T(g)$ which should be clear.

 $\square_{3.11}$

- 3.12 Remark. 1) The proof of 3.10, 3.11 is really a particular case of "the number of special dimensions" from [Sh:c, XIII,§3] the number being here 1; see more on this Hrushovski Hart Laskowski [HHL00].
- 2) The "primary over $\cup \{N_{\{\alpha\}} : \alpha\}$ " is a special case of decompositions.
- **3.13 Theorem.** If T is countable and \boxtimes_T from 3.10 holds then:
 - (a) $\dot{I}(\mu, T)$ is the same whenever $\mu \geq \mu_* =: \theta(\mathscr{F}^*)$ recalling $\mathscr{F}^* = \{f : f \ a \ function from <math>^{\omega}2$ to ω with $\operatorname{supp}(f) = \{\eta : f(\eta) \neq 0\}$ well orderable $\}$.

Proof. We elaborate some parts done in passing in the proof of 3.10 (and add one point).

We can interpret $\eta \in {}^{\omega}2$ as a triple $(M_0, M_1, \varphi(x, \bar{a})) = (M_0^{\eta}, M_1^{\eta}, \varphi_{\eta}(x, \bar{a}_{\eta}))$ such that $M_0 \prec M_1$ are models of T, M_1 with universe ω, M_0 with universe $\{2n : n < \omega\}$ and $\varphi(x, \bar{a})$ a weakly minimal formula in M_0 . So the equivalence relation E_1 is Σ_1^1 where $\eta E_1 \nu \Leftrightarrow [M_0^{\eta} = M_0^{\nu}, \varphi_{\eta}(x, \bar{a}_{\eta}) = \varphi_{\nu}(x, \bar{a}_{\nu}) \text{ and } M_1^{\eta}, M_1^{\nu} \text{ are isomorphic over } M_0^{\eta} = M_0^{\nu}]$ and E_0 a Borel equivalent relation where $\eta E_0 \nu \Leftrightarrow M_0^{\eta} = M_{\eta}^0$.

 $\mathscr{P}_1 = \{A : A \subseteq {}^{\omega}2 \text{ is not empty, any two members are}$ E_0 -equivalent not E_1 equivalent and A is well orderable.

Let

 $\mathscr{F} = \{f : \text{ for some } A \in \mathscr{P}_1, f \text{ is a function from } A \text{ to } \omega + 1 \text{ such that } \omega \in \operatorname{Rang}(f)\}.$

Let

$$\theta_* = \theta(\mathscr{F}) (\leq \theta(\mathscr{F}^*)).$$

For N a model of T of cardinality $\geq \theta_*$ let $\mathscr{F}_N \subseteq \mathscr{F}$ be defined as in the proof of 3.10 but we can write f and not $(f, M_{\emptyset}, \varphi(x, \bar{a}))$ as $M_0, \varphi(x, \bar{a})$ are determined by $\mathrm{Dom}(f)$. Let

$$E_{\mu}^2 = E_{T,\mu}^2 = \{(f_1, f_2) : \text{ there is } N \in \text{Mod}_{T,\mu} \text{ for which } f_1, f_2 \in \mathscr{F}_N\}.$$

(Recalling $\operatorname{Mod}_{T,\mu} = \{M : M \text{ is a model of } T \text{ of cardinality } \mu\}$). Now

- $(*)_1$ if $\mu \geq \theta_*$ and $f \in \mathscr{F}$ then for some model N of T of cardinality μ we have $f \in \mathscr{F}_N$
- $(*)_2$ if $N_1 \cong N_2$ are from $\mathrm{Mod}_{T,\mu}$ and $\mu \geq \theta_*$ then $\mathscr{F}_{N_1} = \mathscr{F}_{N_2}$
- $(*)_3$ if $N_1, N_2 \in \operatorname{Mod}_{T,\mu}, \mu \geq \theta_*$ and $\mathscr{F}_{N_1} \cap \mathscr{F}_{N_2} \neq \emptyset$ then $N_1 \cong N_2$
- $(*)_4$ E_{μ}^2 is an equivalence relation on \mathscr{F}
- $(*)_5$ \mathscr{F}_N for N a model of T of cardinality $\geq \theta_*$ is an E_2 -equivalence class
- $(*)_6$ E_{μ}^2 is the same for all $\mu \geq \theta_*$.

[Why? Assume N^1, N^2 are models of T with universe $\mu, f_\ell \in \mathscr{F}_{N_\ell}$ and let $N_\emptyset^i, a_\alpha^\ell, N_{\{\alpha\}}^\ell$ ($\alpha < \mu$) be as in the proof of 3.10 exemplifying this. Let $\theta_* \leq \mu_1 < \mu_2$. If $\mu = \mu_1, f_1 E_{T,\mu_2}^2 f_2 \Rightarrow f_1 E_{T,\mu_1}^2 f_2$ by the LS argument. The other direction, i.e., if $\mu = \mu_2$ is similar to the proof of 3.10, i.e., we blow up $\langle a_\alpha : \alpha \in I_t \rangle$ for some t (or every t) such that $|I_t| = \mu$ and continue as in 3.8.]

- 3.14 Conclusion. For every countable complete first order theory T, one of the following occurs
 - (A) for every $\alpha, \dot{I}(\aleph_{\alpha}, T) \geq |\alpha|$, in fact there is a sequence $\langle M_{\beta} : \beta < \alpha \rangle$ of pairwise non-isomorphic models of T of cardinality \aleph_{α}
 - (B) for all $\mu \geq \mu_* =: \theta(\mathscr{F}^*)$ (which $\leq \theta(\mathscr{P}(\omega)\omega)$), $\dot{I}(\mu,T)$ is the same and has the form \mathscr{F}^*/E for some equivalence relation E (see more in 3.14 and its proof).
- <u>3.15 Problem</u> [ZF] Give complete classification of $\dot{I}(\lambda, T)$ for T countable by the model theoretic properties of T and the set theoretic properties of the universe.

But it maybe wiser to make less fine distinctions.

- **3.16 Definition.** 1) Let $|X| \lesssim |Y|$ mean that $X = \emptyset$ or there is a function from Y onto X (so $|X| \leq |Y|$ implies this).
- 2) Let $|X| \approx |Y|$ if $|X| \lesssim |Y| \lesssim |X|$ (so this weakens |X| = |Y| and is an equivalence relation) and $|X|/\approx$ is called the essential power.
- <u>3.17 Thesis</u>: It is most reasonable to interpret "determining $\dot{I}(\lambda, T)$ " as finding $\dot{I}(\lambda, T)/\approx$ which is the essential power $|\{M/\cong: M \text{ a model of } T \text{ with universe } \lambda\}|/\approx$.
- **3.18 Claim.** Assume \boxtimes_T of 3.10 and T is countable.
- 1) If T is \aleph_0 -stable then $I(\aleph_\alpha, T) = 1$ for every $\alpha > 0$.
- 2) If D(T) is uncountable and $\alpha > 0$ then:
 - (a) $|\{A \subseteq {}^{\omega}2 : |A| \leq \aleph_{\alpha}\}| \leq \dot{I}(\aleph_{\alpha}, T)\}$
 - (b) $\dot{I}(\aleph_{\alpha}, T)$ is < -below $|\{A \subseteq {}^{\omega}2 : |A| \leq \aleph_{\alpha}\}|$

(note: $|A| \leq \aleph_{\alpha} \Rightarrow A$ is well ordered).

3) If D(T) is countable, T is not \aleph_0 -stable and there is a set of \aleph_1 reals and $\alpha > 0$ then

$$\dot{I}(\aleph_{\alpha}, T) \approx |\{A \subset {}^{\omega}2 : |A| < \aleph_{\alpha}\}|.$$

Proof. As in [Sh:c]. (E.g. in (2) the first inequality holds as in $\mathbf{L}[T,Y]$ we can find countable complete $T_1 \supseteq T$ with Skolem functions $M_1 \models T_1, a_{\eta} \in {}^m M_1$ for $\eta \in {}^{\omega} 2$ and $b_n \in M_1$ for $n < \omega$ such that letting $\alpha = \omega, A = ({}^{\omega} 2)^{\mathbf{L}[T,Y]}$ we have

- $(*)_A^{\alpha}$ (a) $\langle b_n : n < \alpha \rangle$ is a non-trivial indiscernible sequence in M_1 over $\{\bar{a}_{\eta} : \eta \in A\}$
 - (b) $\langle \operatorname{tp}(\bar{a}_{\eta}, \emptyset, M_1 \upharpoonright \tau_T) : \eta \in {}^{\omega}2 \rangle$ are pairwise distinct
 - (c) $\langle \bar{a}_{\eta}: \eta \in {}^{\omega}2 \rangle$ is in discernible in $(M_1, b_n)_{n < \omega}$ in the weak sense of [Sh:c, VII,§2]
 - (d) $\operatorname{tp}(\bar{a}_{\eta}, \emptyset, M_1 \upharpoonright \tau_T)$ is not realized in $M_1 \upharpoonright \operatorname{acl}(\{\bar{a}_{\nu} : \nu \in {}^{\omega}2 \setminus \{\eta\}\}) \cup \{b_n : n < \omega\})$.

So in bigger universe this M_1 has a natural extension. So we can define $M_1^+, \langle a_\eta : \eta \in ({}^{\omega}2)^{\mathbf{V}} \rangle, \langle b_\alpha : \alpha \in [\omega, \mu) \rangle$ naturally such that $(*)^{\mu}_{\mathscr{P}(\omega)}$ and define M_A for $A \subseteq {}^{\omega}2$ as $\mathrm{Sk}(\{\bar{a}_\eta : \eta \in A\} \cup \{b_\alpha : \alpha < \mu\}, M_1)\} \upharpoonright \tau_T$; if A is well orderable then M_A has cardinality μ .

We now look at a well known example in our context.

- 3.19 Example: There is countable stable, not superstable T with D(T) countable such that: if there are no sets of \aleph_1 reals then $I(\aleph_\alpha, T)$ is "manageable"
 - (A) let G be an infinite abelian group, each element of order 2. So ${}^{\omega}G$ is also such a group. We define a model M:
 - (a) its universe: $G \cup {}^{\omega}G$ (assuming $G \cap {}^{\omega}G = \emptyset$)
 - (b) predicates $P^M = G, Q^M = {}^{\omega}G$
 - (c) the partial two-place function H_1^M which is the addition of G (you may add $x \notin G \land y \notin G \Rightarrow x + {}^M y = x$)
 - (d) H_2^M is the addition on ${}^\omega G$ (coordinatewise)
 - (e) a partial unary function F_n^M such that $\eta \in {}^{\omega}G \Rightarrow F_n^M(\eta) = \eta(n)$
 - (f) individual constants c_1, c_2 the zeroes of G and ${}^{\omega}G$ respectively
 - (B) let $T = \operatorname{Th}(M)$. Let $K^* = \{N : N \models T \text{ and } N \text{ omit } \{Q(x) \land Q(y) \land F_n(x) = 1\}$ $F_n(y) \land x \neq y : n < \omega$
 - (C) if $\langle M_t : t \in I \rangle$ is a sequence of models of T we can naturally define their sum $\bigoplus_{t \in I} M_t$. Clearly K^* is closed under sum (i.e.,

 $|M| = P^M \cup Q^M$

- $P^{M} = \{f : f \text{ is a function with domain } I \text{ such that } f(t) \in P^{M_t} \text{ and } f(t) \}$ is the zero $c_1^{M_t}$ of the abelian group $(P^{M_t}, H_1^{M_t})$ for all but finitely many t's},
- $Q^M = \{g : g \text{ a function with domain } I, f(t) \in Q^{M_t} \text{ and for all but finitely } \}$ many $t \in I$ we have $f(t) = c_1^{M_t}$ (we ignore that for I finite, formally $P^M \cap Q^M \neq \emptyset$), etc.)

- (D) (ZF) If M is a model from K^* of cardinality λ and λ is a ($<\lambda$)-free cardinal (see Definition 5.2 below) then $M = \bigoplus M_i$ for some sequence $\langle M_i : i < \lambda \rangle$ such that $i < \lambda \Rightarrow ||M_i|| < \lambda$
- (E) in (D) if the cardinal λ is ($<\mu$)-free we can add $||M_i|| < \mu$ (see Definition 5.2 below)
- (F) (a)define for $M \models T$ a two-place relation E_M on M: $aE_Mb \Leftrightarrow (a=b) \vee (Q(a) \wedge Q(b) \wedge \bigwedge F_n(x) = F_n(y)).$

It is an equivalence relation on M

- (b) define M/E_M naturally
- (G) (a) $M/E_M \in K^*$ for any model M of T
 - (b) $M_1 \cong M_2 \Rightarrow M_1/E_{M_1} \cong M_2/E_{M_1}$.

- (H) (a) if $M \models T$ and $a, b \in Q^M$ then $|a/E_M| = |b/E_M|$
 - (b) we can consider a/E_M (where $M \models T, a \in Q^M$) an abelian group $G_{a,M}$ with every element of order 2 except that the zero is not given
 - (c) if $a, b \in Q^M$ then $G_{a,M}, G_{b,M}$ as vector spaces of $\mathbb{Z}/2\mathbb{Z}$ without zero has the same dimension
 - (d) call this dimension $\lambda(M)$
- (I) if M_1, M_2 are models of T of cardinality \aleph_{α} then $M_1 \approx M_2$ iff $M_1/E_{M_2} \approx M_2/E_{M_2}$ and $\lambda(M_1) = \lambda(M_2)$ hence $\dot{I}(\aleph_{\alpha}, T) = \dot{I}(\leq \aleph_{\alpha}, K^*) \times |\omega + \alpha|$ where $\dot{I}(\leq \aleph_{\alpha}, K^*) = \Sigma\{\dot{I}(\aleph_{\beta}, K^*) : \beta \leq \alpha\} = \Sigma\{\dot{I}(\aleph_{\beta}, K^*) : \beta \leq \alpha, \aleph_{\beta} < \theta(\mathscr{P}, \omega)\}.$

§4 On T categorical in |T|

The ZFC parallel of 4.2 - 4.4 is the known "|D(T)| < |T| implies T is the definitional extension of some $T' \subseteq T, |T'| < |T|$ ", see Keisler [Ke71a], which in Boolean algebra terms say "the number of ultrafilters of an infinite Boolean algebra B is $\geq |B|$ ".

- 4.1 Convention. For 4.2-4.6, T is first order (with τ_T not necessarily well-orderable).
- **4.2 Definition.** For a first order T in the vocabulary $\tau = \tau_T$, usually for simplicity closed under deduction, we define the equivalence relation E_T on τ_T by
 - (a) for predicates $P_1, P_2 \in \tau$ P_1EP_2 iff: $P_1, P_2 \in \tau$ are predicates with the same arity and $(\forall \bar{x})(P_1(\bar{x}) \equiv P_2(\bar{x})) \in T$
 - (b) for function symbols $F_1, F_2 \in \tau$, e.g. individual constants F_1EF_2 iff: $P_1, P_2 \in \tau$ are function symbols with the same arity and $(\forall \bar{x})(F_1(\bar{x}) = F_2(\bar{x})) \in T$
 - (c) no predicate $P \in \tau$ is E-equivalent to a function symbol $F \in \tau$.
- **4.3 Definition.** Let $\tau = \tau_T, T$ a first order theory.
- 1) The theory T is called reduced if E_T is the equality.
- 2) Let τ/E_T be the vocabulary with predicates $P/E_T, P \in \tau$ a predicate with $\operatorname{arity}(P/E_T) = \operatorname{arity}_{\tau}(P)$ and similarly F/E_T .
- 3) For a τ -model M of T we define $M^{[E_T]}$ naturally, i.e.,
- $N = M^{[E_T]}$ iff they have the same universe, N is a (τ/E_T) -model, M is a τ -model $M \models T$ and $(R/E_T)^N = R^M$ for every predicate $R \in \tau(T)$ and $(F/E_T)^N = F^M$ for any function symbol $F \in \tau(T)$.
- 4) For N a (τ/E_T) -model, $M = {}^{[E_T]}N$ is the τ -model such that $N = M^{[E_T]}$ if one exists.
- 5) Let T/E_T be the set of $\psi \in \mathbb{L}(\tau(T)/E_T)$ such that if we replace any predicate R/E_T appearing in ψ by some $R' \in R/E_T$ and similarly for F/E_T , we get a sentence from $\{\psi \in \mathbb{L}(\tau_T) : T \vdash \psi\}$, see 4.2.
- 4.4 Observation. [ZF] For every first order T (as in 4.2) in $\mathbb{L}(\tau_T)$
 - (a) E_T is an equivalence relation on τ

- (b) if M is a τ -model of T then $M^{[E_T]}$ is a uniquely determined (τ/E_T) -model of T/E and $[E_t](M^{[E_T]}) = M$
- (c) for every (τ/E) -model M of T/E_T the τ -model $[E_t]M$ uniquely determined and is a model of T and $([E_T]M)^{[E_T]}=M$
- (d) T/E_T is a reduced first order theory

See hopefully more on such T's in [Sh:F701].

4.5 Hypothesis. $\tau(T) \subseteq \mathbf{L}$ as usual.

4.6 Claim. [ZF] If T is a complete first order theory in $\mathbb{L}(\tau)$ and T is reduced and $Y \subseteq \mathbf{L}$, then $T \in \mathbf{L}[Y] \Rightarrow |D(T)|^{\mathbf{L}[T]} \geq |T|$.

Proof. By the ZFC case (see Keisler [Ke71a]).

 $\square_{4.6}$

4.7 Claim. If $T \subseteq \mathbf{L}$ is categorical in λ and $Y \in \operatorname{Ord} \underline{then}$ in $\mathbf{L}[T, Y]$ the following is impossible

- \circledast (a) T stable, $\lambda \geq |T| + \aleph_1 + \mu$,
 - (b) $M \prec \mathfrak{C}$ is $\mathbf{F}_{\aleph_0}^f$ primary over \emptyset , see [Sh:c, IV]
 - (c) $\bar{a}_i \in {}^n M \text{ for } i < \mu$
 - (d) $\operatorname{tp}(\bar{a}_{\delta}, \cup \{\bar{a}_i : i < \delta\})$ forks over $\cup \{a_j : j < \alpha\}$ whenever $\alpha < \delta < \mu, \delta$ a limit ordinal from S
 - (e) every type over $\cup \{\bar{a}_i : i < \mu\}$ which is realized in M does not fork over some $\cup \{\bar{a}_i : i < \alpha\}$ for some $\alpha < \mu$
 - (f) in $\mathbf{L}[T, Y]$ we have: μ regular uncountable, $S \subseteq \mu$ stationary.

Proof. Work in $\mathbf{L}[T, Y]$; without loss of generality M has cardinality λ , and toward contradiction assume \circledast holds. By clause (b) there is $\bar{\mathbf{c}}$ such that

- $(*)_1 \ \bar{\mathbf{c}} = \langle \bar{c}_i : i < i^* \rangle$
- $(*)_2$ $M = \cup \{\bar{c}_i : i < i^*\}$ and $\operatorname{tp}(\bar{c}_i, \cup \{\bar{c}_j : j < i\})$ does not fork over some finite $B_i \subseteq \cup \{\bar{c}_j : j < i\}$ for each $i < i^*$

So by the properties of non-forking (or of $\mathbf{F}_{\aleph_0}^f$ -constructions, [Sh:c, IV]) without loss of generality we have $(i^* \geq \mu \text{ and}) \cup \{\bar{a}_i : i < \mu\} \subseteq \cup \{\bar{c}_j : j < \mu\}$. Hence for some club E of μ we have $\bar{a}_i \subseteq \bigcup_{j < \delta} \bar{c}_j \Leftrightarrow i < \delta \text{ for } i < \mu, \delta \in E$; clearly

 $\operatorname{tp}(\bar{a}_{\delta}, \cup \{\bar{c}_i : i < \delta\})$ does not fork over some finite $C_{\delta} \subseteq \cup \{\bar{c}_j : j < \delta\}$. Hence there is stationary $S_1 \subseteq S \cap C$ such that $\delta \in S \Rightarrow C_{\delta} = C_*$, and let \bar{c} list C_* .

By clause (e) of the assumption for some $\alpha_* < \mu$,

 $(*)_2 \operatorname{tp}(\bar{c}, \cup \{\bar{a}_i : i < \mu\})$ does not fork over $\cup \{\bar{a}_i : i < \alpha_*\}$

hence by the non-forking calculus

(*)₃ for $\delta \in S_1 \setminus (\alpha + 1)$ the type $\operatorname{tp}(\bar{a}_{\delta}, \cup \{\bar{a}_i : i < \delta\})$ does not fork over $\cup \{\bar{a}_i : i < \alpha_*\}$.

By this contradicts clause (d) of the assumption.

 $\square_{4.7}$

4.8 Claim. If T is stable, categorical in λ and $\lambda = |T| > \aleph_0$ then $\underline{Case}(\alpha)$: if $(\exists Y \subseteq \operatorname{Ord})(\aleph_1 = \aleph_1^{\mathbf{L}[T,Y]})$ then $\kappa_r(T) = \aleph_0$, i.e. T is superstable. $\underline{Case}(\beta)$: if $(\forall Y \subseteq \operatorname{Ord})(\aleph_1 > \aleph_1^{\mathbf{L}[T,Y]})$ then for every $Y \subseteq \operatorname{Ord}$ we have $\mathbf{L}[T,Y] \models \text{``}\kappa(T) < \aleph_1^{\mathbf{V}}$.

Proof. Case (α) :

Assume the conclusion fails. Fix $Y \subseteq \text{Ord}$ such that $T \in \mathbf{L}[Y], \aleph_1 = \aleph_1^{\mathbf{L}[Y]}$ and $\mathfrak{C} = \mathfrak{C}_T \in \mathbf{L}[Y]$ is a χ -saturated (in $\mathbf{L}[Y]$) model of T and $\mathbf{L}[Y] \models \kappa(T) \geq \aleph_1$ where χ is large enough and regular in $\mathbf{L}[Y]$; and we shall work inside $\mathbf{L}[Y]$.

Let $\mu = \aleph_1^{\mathbf{L}[Y]} = \aleph_1^{\mathbf{V}}$. We can find $\langle \bar{a}_n : n < \omega \rangle, \bar{a}_n \in {}^{\omega >} \mathfrak{C}_Y$ and a type $p = \{\varphi_n(x,\bar{a}_n) : n < \omega\}$ such that $\varphi_n(x,\bar{a}_n)$ forks over $\cup \{\bar{a}_m : m < \omega\}$. Let $\langle \eta_i : i < \mu \rangle$ list ${}^{\omega >}(\mu)$ such that $\eta_i \triangleleft \eta_j \Rightarrow i < j$ and for every limit ordinal $\delta < \mu$ we have ${}^{\omega >}\delta = \{\eta_i : i < \delta\}$. We choose $\bar{\nu} = \langle \nu_\delta : \delta < \mu \text{ limit} \rangle$ such that ν_δ is increasing with limit δ .

We choose $\langle \bar{a}_{\eta} : \eta \in {}^{\omega >} \mu \rangle$ such that: if $\ell g(\eta) = n$ then $\hat{a}_{\eta \upharpoonright 0} \hat{a}_{\eta \upharpoonright 1} \hat{\ldots} \hat{a}_{\eta}$ and $\bar{a}_{0} \hat{\ldots} \hat{a}_{n}$ realize the same type in \mathfrak{C} and $\operatorname{tp}(\bar{a}_{\eta}, \cup \{\bar{a}_{\nu} : \nu \in {}^{\omega >} \mu \text{ and } \neg (\eta \leq \nu)\})$ does not fork over $\cup \{\bar{a}_{\eta \upharpoonright k} : k < \ell g(\eta)\}$. For limit $\delta < \mu$ we choose b_{δ} which realizes $\{\varphi_{n}(x, \bar{a}_{\nu_{\delta} \upharpoonright n}) : n < \omega\}$ such that $\operatorname{tp}(b_{\delta}, \cup \{\bar{a}_{\nu} : \nu \in {}^{\omega >} \mu\} \cup \{b_{\delta'} : \delta' < \delta \text{ is limit}\})$ does not fork over $\cup \{\bar{a}_{\nu_{\delta} \upharpoonright n} : n < \omega\}$.

Lastly, let a_i' be b_δ if $i = \delta$ and be \bar{a}_{η_j} if i = j + 1, and be <> if i = 0.

Let $M_1 \prec \mathfrak{C}, M_1 \in \mathbf{L}[Y]$ be a model of cardinality λ which is $\mathbf{F}_{\aleph_0}^f$ -primary over \emptyset .

Let $M_2 \prec \mathfrak{C}_T, M_2 \in \mathbf{L}[Y]$ be $\mathbf{F}_{\aleph_0}^f$ -primary over $\cup \{\bar{a}_i' : i < \mu\}$ of cardinality λ (see [Sh:c, IV]). Now by 4.7 for $\mu = \aleph_1$, the models M_1, M_2 are not isomorphic even in $\mathbf{L}[Y, Y_1]$ for any $Y_2 \subseteq \mathrm{Ord}$ (as $\aleph_1^{\mathbf{L}[Y, Y_0]} = \aleph_1^{\mathbf{L}[Y]} = \aleph_1^{\mathbf{V}}$), contradiction.

Case (β) : Assume that the conclusion fails for Y. Clearly $\aleph_1^{\mathbf{V}}$ is a limit cardinal in $\mathbf{L}[T,Y']$ for every $Y'\subseteq \mathrm{Ord}$. So for every $\mu\in\mathrm{Card}^{\mathbf{L}[T,Y]}\cap\omega_1^{\mathbf{V}}$ we can find (in $\mathfrak{C}_T^{\mathbf{L}[T,Y]}\in\mathbf{L}[T,Y]$ chosen as above) a sequence $\bar{\mathbf{a}}_{\mu}=\langle\bar{a}_{\mu,i}:i<\mu\rangle$ such that $\bar{a}_{\mu,i}\in\omega^{\mathbf{L}[T,Y]}\in\mathbf{L}[T,Y]$ chosen as above) a sequence $\bar{\mathbf{a}}_{\mu}=\langle\bar{a}_{\mu,i}:i<\mu\rangle$ such that $\bar{a}_{\mu,i}\in\omega^{\mathbf{L}[T,Y]}\in\mathbf{L}[T,Y]$ and a type $p=\{\varphi_{\mu,i}(x,\bar{a}_{\mu,i}):i<\mu\}$ in \mathfrak{C} such that $\varphi_i(x,\bar{a}_{\mu}^{\mu})$ forks over $\cup\{\bar{a}_{\mu,j}:j< i\}$ for every i. Choose by induction on $i<\mu$ an element $b_i^{\mu}\in\mathfrak{C}$ which realizes $\{\varphi_{\mu,j}(x,\bar{a}_{\mu,j}):j< i\}$ but $\mathrm{tp}(b_i^{\mu},\cup\{\bar{a}_{\mu,j}:j<\omega_1\}\cup\{b_j^{\mu}:j< i\})$ does not fork over $\cup\{\bar{a}_{\mu,j}:j< i\}$. Let $\bar{a}_i^{\mu}=\bar{a}_{\mu,i}^{-}(b_i^{\mu})$ so $\langle\bar{a}_i^{\mu}:i<\mu\rangle$ is as in clauses (c) + (d) of 4.7. Note that the function $(\mu,i)\mapsto\bar{a}_i^{\mu}$ belongs to $\mathbf{L}[T,Y]$. Without loss of generality $\{\bar{\mathbf{a}}_{\mu}:\mu\in\mathrm{Card}^{\mathbf{L}[T,Y]}\cap\omega_1^{\mathbf{V}}\}$ is independent over \emptyset in $\mathfrak{C}_Y^{\mathbf{L}[T,Y]}$. In $\mathbf{L}[T,Y]$ let $M_1\prec\mathfrak{C}_T^{\mathbf{L}[T,Y]}$ be of cardinality λ , $\mathbf{F}_{\aleph_0}^f$ -primary over \emptyset . Let $M_2\prec\mathfrak{C}_T^{\mathbf{L}[T,Y]}$ be of cardinality λ and $\mathbf{F}_{\aleph_0}^f$ -primary over $\cup\{\bar{\mathbf{a}}_{\mu}:\mu\in\mathrm{Card}^{\mathbf{L}[T,Y]}\cap\omega_1^{\mathbf{V}}\}$. But T is categorical in λ so there is an isomorphism $\mathbf{f}\in\mathbf{V}$ from M_1 onto M_2 and now we shall work in $\mathbf{L}[T,Y,f]$ and let $\mu_*=\aleph_1^{\mathbf{L}[T,Y,f]}$, clearly $\mu_*\in\mathrm{Reg}^{\mathbf{L}[T,Y]}\cap\omega_1^{\mathbf{V}}$ so $\bar{\mathbf{a}}_{\mu_*}$ is well defined. By the non-forking calculus, the statement \circledast of 4.7 holds for μ_* so we are done.

4.9 Remark. Assume T is stable, (complete with infinite models of course), $\lambda = |T| \geq \aleph_{\alpha} > \aleph_0$ and for some $Y \subseteq \text{Ord}$ we have $\mathbf{L}[Y] \models \text{``}\kappa(T) > \aleph_{\alpha}$ or \aleph_{α} is a limit cardinal and $\kappa(T) \geq \aleph_{\alpha}$ ". Then $\dot{I}(\lambda, T) \geq |\alpha|$. The proof is similar.

4.10 Claim. T is not categorical in $\lambda = |T| > \aleph_0$ when for some $Y \subseteq \operatorname{Ord}$:

- \circledast (a) T is stable
 - (b) $\mathbf{L}[T,Y] \models \text{``}|D(T)| \ge \lambda = |T|\text{'`} \text{ (holds if } T \text{ is reduced, see 4.6)}$
 - (c) the conclusion of 4.8 holds, (or just for every $Y' \subseteq \text{Ord } we \ have$ $\lambda > \kappa(T)^{\mathbf{L}[Y',Y,T]}$).

Proof. Choose $Y \subseteq \text{Ord}$ which exemplify the assumption of case (α) of 4.8 if it holds. In $\mathbf{L}[T,Y]$ letting $\kappa = \kappa(T)^{\mathbf{L}[T,Y]}$ let:

- (*)₁ M_1 be \mathbf{F}_{κ}^a -constructible over \emptyset of cardinality λ , i.e., for some sequence $\langle a_i, B_i : i < \lambda \rangle$ we have $M_1 = \{a_i : i < \lambda\}$ and $B_i \subseteq \{a_j : j < i\}$ has cardinality $< \kappa$ and $\mathrm{stp}(a_i, B_i) \vdash \mathrm{stp}(a_i, \{a_j : j < i\})$ (not necessarily \mathbf{F}_{κ}^a -saturated!)
- (*)₂ M_2 be a model of T of cardinality λ with $\mathbf{I} \subseteq M_2$ indiscernible of cardinality λ .

[Why (*)₂ is possible? E.g., we can have $||M_1|| = \lambda$ because $\mathbf{L}[T, Y] \models "|D(T)| \ge \lambda$ ".]

So assume toward contradiction that M_1, M_2 are isomorphic, let $\mathbf{f}: M_1 \xrightarrow[\text{onto}]{\text{iso}} M_2$ be such an isomorphism and work in $\mathbf{L}[T,Y,\mathbf{f}]$. Now $\kappa(T)^{\mathbf{L}[T,Y,\mathbf{f}]}$ may be $> \kappa = \kappa(T)^{\mathbf{L}[T,Y]}$ and κ may be not a cardinality still the properties of M_1, M_2 from $(*)_1, (*)_2$ respectively holds in $\mathbf{L}[T,Y,\mathbf{f}]$ for $\kappa = \kappa(T)^{\mathbf{L}[T,Y,\mathbf{f}]}$. Now we can get a contradiction as in [Sh:c, IV].

Putting together Claims 4.8, 4.10.

4.11 Conclusion. If T is stable in $\lambda = |T| \leq |D(T)|$ then T is not categorical in λ .

* * *

4.12 Free Models. Let T be complete and stable. $\mathfrak{C} = \mathfrak{C}_{Y,T}$ a monster for T in $\mathbf{L}[T,Y]$.

The proofs above (and actually [Sh:c]) suggest that we look more into free models.

4.13 Definition. 1) A model M of T, (a stable theory) is free <u>when</u> we can find a sequence $\langle a_i : i < \alpha \rangle$ enumerating M such that for each $i < \alpha$ the type $\operatorname{tp}(a_i, \{a_j : j < i\}, M)$ does not fork over some finite subset say B_i .

2) We call $\langle (A_i, a_i, B_i) : i < \alpha \rangle$ is a free representation of M where $A_i = \{a_j : j < i\}$.

Remark. So free is the same as being $\mathbf{F}_{\aleph_0}^f$ -constructible over \emptyset .

4.14 Claim. If $A \subseteq \mathfrak{C}$, $\lambda = |A|$ is singular and every $A' \subseteq A$ of cardinality $< \lambda$ is free <u>then</u> A is free.

Proof. By compactness in singular ([Sh 54], [Sh:E18]). $\square_{4.14}$

§5 Consistency results

In spite of the evidence of §1,§4, without choice characterization for the number of non-isomorphic models is different without choice. We look for consistency results for "there are few models in cases impossible by ZFC", in particular we ask (and give a partial answer):

- 5.1 Question: 1) Is it consistent with ZF that for some/many $\kappa > \aleph_0$ we have: every two strongly \aleph_0 -homogeneous linear orders of cardinality κ , are isomorphic? (Add " κ singular or κ regular"; or add $cf(\kappa) = \aleph_0$.)
- 2) Similarly is it consistent with ZF that
- "if $M_1, M_2 \subseteq ({}^{\omega}\lambda, E_n)_{n < \omega}$ are strongly \aleph_0 -homogeneous of cardinality κ then they are isomorphic".
- 3) Instead categoricity proves the consistency of all models has nice descriptions, (see below):

Clearly 5.8 below proves that our use of elementary classes in the proof for stable, un-superstable T is necessary, that is we could not prove too good theorems on PC classes parallel to the ZFC case.

Toward 5.1(2) we consider:

- **5.2 Definition.** 1) A cardinality λ is free or ω -sequence-free when every subset of ${}^{\omega}\lambda$ of cardinality λ is free, where
- 2) A subset $A \subseteq {}^{\omega}\lambda$ is free when there is a one-to-one function $f: A \to {}^{\omega}{}^{>}\lambda$ such that $\eta \in A \Rightarrow f(\eta) \triangleleft \eta$.
- 3) A cardinal λ is $(< \mu)$ -free when every subset of ${}^{\omega}\lambda$ of cardinality $\leq \lambda$ is $(< \mu)$ -free where
- 4) We say " $A \subseteq {}^{\omega}\lambda$ is $(<\mu)$ -free if there is a function $f:A \to {}^{\omega}{}^{>}\lambda$ which in some $\mathbf{L}[Y]$ is $(<\mu)$ -to-one" and $\eta \in A \Rightarrow f(\eta) \triangleleft \eta$.
- <u>5.3 Question</u>: 1) Is it consistent (with ZF) that for arbitrarily large μ, μ^+ is μ^+ -free? (\aleph_0 always is).
- 2) Is it consistent with ZF that all cardinals are free?
- **5.4 Claim.** [ZF + DC] Let $\kappa = \aleph_1$. The following is a sufficient condition for λ being $(< \kappa)$ -free (equivalently free)

 $\Box_{\lambda,\mu}$ for every $A \subseteq \lambda$ for some $B \subseteq \lambda$ we have:

- (*)₁ if $\mathbf{L}[A] \models \text{``}\mu \text{ is a cardinal} < \lambda \text{ but } \geq \kappa \text{ such that } \mu < \mu^{\aleph_0}, \mu' = \min\{\lambda, \mu^{\aleph_0}\}\text{''} \text{ then } \mathbf{L}[A, B] \models \text{``}\mu' \text{ is an ordinal of cardinality} \leq \mu''$
- (*)₂ if $\mathbf{L}[A] \models "\mu \leq \lambda$ is regular uncountable $\geq \kappa$ and $S = \{\delta < \lambda : \mathrm{cf}(\delta) = \aleph_0\}$ " then $\mathbf{L}[A, B] \models "S$ is a non-stationary subset of μ ".

Remark. 1) A condition for $\kappa > \aleph_1$ will be more complicated. 2) ($< \aleph_1$)-free is equivalent to free (note that "in some $\mathbf{L}[Y]$ " in Definition 5.2).

Proof. So assume that A is a subset of ${}^{\omega}\lambda$ of cardinality $\leq \lambda$. Let

$$\Xi = \{(Y, f) : Y \subseteq \text{ Ord and } f \in \mathbf{L}[Y] \text{ is a function from } A \text{ to }^{\omega > \lambda} \}$$

such that $\eta \in A \Rightarrow f(\eta) \triangleleft \eta\}$

and let $\bar{\mu}_{Y,f} = \langle \mu_{\bar{\eta}}^{Y,f} : \eta \in A \rangle$ be defined by $\mu_{\eta}^{Y,f} = |\{\eta' \in A : f(\eta') = f(\eta)\}|^{\mathbf{L}[f,Y]}$. So the role of Y is in determining where we compute $\mu_{\eta}^{Y,f}$. Now it suffices to prove

 \circledast if $(Y, f) \in \Xi$ then there is $(Z, g) \in \Xi$ such that $\eta \in A \Rightarrow \bar{\mu}_{\eta}^{Y_1, f_1} < \mu_{\eta}^{Y, f} \vee \mu_{\eta}^{Y, f} < \kappa$.

[Why it suffices? If so by DC we can find $\langle (Y_n, f_n) : n < \omega \rangle \in V$ such that $(Y_n, f_n) \in \Xi$ and

$$(*) \ \eta \in A \Rightarrow (\mu_{\eta}^{Y_n, f_n} > \mu_{\eta}^{Y_{n+1}, f_{n+1}}) \lor (\mu_{\eta}^{Y_n, f_n} < \kappa).$$

Let $Y_* = \{\operatorname{cd}(\langle 1, n, \ell g(\eta) \rangle \hat{\ } \eta \hat{\ } \langle f_n(\eta) \rangle) : \eta \in {}^{\omega >} \lambda \text{ and } n < \omega\} \cup \{\operatorname{cd}(2, n, \alpha) : \alpha \in Y_n \text{ and } n < \omega\} \text{ where cd is a one-to-one definable function in } \mathbf{L} \text{ from } {}^{\omega >} \text{Ord into Ord.}$ Clearly $\langle f_n : n < \omega \rangle \in \mathbf{L}[Y_*]$ and define $h : A \to \omega$ by $h(\eta) = \operatorname{Min}\{n : \mu_{\eta}^{Y_n, f_n} < \kappa\}$, it clearly exists by \circledast .

Lastly, let $f: A \to {}^{\omega} > \lambda$ be defined by $f(\eta) = \eta \upharpoonright \operatorname{pr}(h(\eta), \ell g(f_{h(\eta)}(\eta)))$ where $\operatorname{pr}(n, m)$ is, e.g. $(n + m + 1)^2 + n$.

Now check.]

Proof of \circledast .: Let Z be like B is the claim's assumption with Y playing the roles of A; we work in $\mathbf{L}[Y,Z]$, without loss of generality $Y \in \mathbf{L}[Z]$. Let $\langle \eta_{\alpha} : \alpha < |A| \rangle$ list A with no repetitions. Let $\mathscr{U} = \{\alpha < |A| :$ for no $\beta < \alpha$ do we have $f(\eta_{\beta}) = f(\eta_{\alpha}) \}$ and let $\mu_{\alpha} = |\{\beta : f(\eta_{\beta}) = f(\eta_{\alpha})\}$ for $\alpha \in \mathscr{U}$ and let (so $\langle \mu_{\alpha} : \alpha \in \mathscr{U} \rangle \in \mathbf{L}[Y]$.

In $\mathbf{L}[Y]$ let $\langle \langle \eta_{\alpha,\varepsilon} : \varepsilon < \mu_{\alpha} \rangle : \alpha \in \mathscr{U} \rangle$ be such that for each $\alpha \in \mathscr{U}$ the sequence $\langle \eta_{\alpha,\varepsilon} : \varepsilon < \mu_{\alpha} \rangle$ list $A_{\alpha} := \{\beta : f(\beta) = f(\alpha)\}$. Now

 \boxtimes it suffices to prove that in $\mathbf{L}[Z]$, for every $\alpha \in \mathscr{U}$ there is $f_{\alpha} : A_{\alpha} \to {}^{\omega} > \lambda$ such that $\eta \in A_{\alpha} \Rightarrow |\{\nu \in A_{\alpha} : f_{\alpha}(\nu) = f_{\alpha}(\eta)\}| < \mu_{\alpha}$.

Note that in $\mathbf{L}[Z]$, μ_{α} is not necessary a cardinal, in this case $f_{\alpha} = f \upharpoonright A_{\alpha}$ can serve!

[Why? In $\mathbf{L}[Z]$ we can choose $\langle f_{\alpha} : \alpha \in \mathcal{U} \rangle$ in \circledast and then put together f and $\cup \{f_{\alpha} : \alpha \in \mathcal{U}\}$ as above.]

The proof of the condition in \oplus is by cases (on α):

Case 1: $\alpha \in \mathcal{U}$ and μ_{α} is not a cardinal in $\mathbf{L}[Z]$ or $\mu_{\alpha} < \kappa$. Trivial.

Hence by clause (a) of the assumption

 $(*)_2$ without loss of generality $L[Z] \models "\mu_{\alpha}$ is a cardinality".

<u>Case 2</u>: In L[Z], μ_{α} is regular $> \kappa$.

Let $B_{\alpha,\varepsilon} = \{\eta_{\alpha,\zeta}(n) : n < \omega \text{ and } \zeta < \varepsilon\}$, so in $\mathbf{L}[Z], \langle B_{\alpha,\varepsilon} : \varepsilon < \mu_{\alpha} \rangle$ is \subseteq -increasing continuous and let $C_{\alpha,0} = \{\delta < \lambda : \delta \text{ is a limit ordinal and for every } \varepsilon < \mu \text{ we have } \varepsilon < \delta \text{ iff for some } \zeta < \delta, \operatorname{Rang}(\eta_{\alpha,\varepsilon}) \subseteq B_{\alpha,\zeta}\}.$

In $\mathbf{L}[Z]$ there is a club $C_{\alpha} = \{\beta_{\xi} : \xi < \mu_{\alpha}\}$ of μ_{α} such that $\delta \in C = \mathrm{cf}(\delta)^{\mathbf{L}[Y]} > \aleph_0$ and $C_{\alpha} \subseteq C$ and $\beta_0 = 0$.

For $\varepsilon < \mu_{\alpha}$ let $\xi = \xi(\varepsilon)$ be maximal such that $\varepsilon \geq \beta_{\xi}$ and easily $\eta_{\alpha,\varepsilon} \notin {}^{\omega}(B_{\alpha,\beta_{\varepsilon}})$, and let $g(\eta_{\alpha,\varepsilon})$ be the shortest $\nu \leq \eta_{\alpha,\varepsilon}$ which $\notin B_{\alpha,\beta_{\xi(\varepsilon)}}$. Now check.

Case 3: $\operatorname{cf}^{\mathbf{L}[Z]}(\mu_{\alpha}) \geq \kappa$. Similarly.

Case 4: $\operatorname{cf}^{\mathbf{L}[Z]}(\mu_{\alpha}) = \aleph_0$.

Here we can find an increasing sequence $\langle B_n : n < \omega \rangle$ of subsets of λ of cardinality $< \mu$ such that $A_{\alpha} \subseteq \bigcup^{\omega} (B_n)$.

So we can proceed as above.

 $\square_{5,4}$

<u>Discussion</u>: Question 5.3 seems to me to call for iterating Radin forcing but for \aleph_2 there is a short cut. For this we quote.

5.5 Theorem. Assume ZF + DC + AD and $\kappa = \aleph_1$. Then

 $(*)_{\kappa}$ for every $A \subseteq \kappa$ for some $\eta \in {}^{\omega}2$ we have $A \in \mathbf{L}[\eta]$ and $\eta^{\#}$ (hence $A^{\#}$) exist.

Proof. Well known.

- **5.6 Claim.** [ZF] 1) If $DC + AD + \kappa = \aleph_1$ or just $(*)_{\kappa}$ from 5.5 holds, <u>then</u> \aleph_1 is free.
- 2) Also κ is Ord-free (see Definition 5.10 below).
- *Proof.* 1) We can easily check the criterion from 5.4 as for M a model with universe κ and vocabulary $\subseteq \mathbf{L}_{\omega}$, let $\eta \in {}^{\omega}2$ be such that $M \in \mathbf{L}[\eta]$ and can work in $\mathbf{L}[\eta, \eta^{\#}]$. 2) Easy, too.
- 5.7 Observation. [ZFC + DC] If $(*)_{\lambda,\partial}$ then $\Box_{\lambda,\partial}$ where
- $(*)_{\lambda,\theta}$ for every $A \subseteq \lambda$ there is $B \subseteq \partial$ such that $A \in \mathbf{L}[B]$ and $B^{\#}$ exists (so $(*)_{\kappa}$ is $(*)_{\kappa,\aleph_0}$)
- $\Box_{\lambda,\theta}$ every model M of cardinality λ with vocabulary of cardinality $\leq \partial$ (so τ_M well ordered) is isomorphic to a model of the form $\mathrm{EM}_{\tau}(\lambda,\Phi)$ for some template Φ with $|\tau_{\Phi}| \leq \theta$ (so τ_{Φ} well ordered).

Remark. This includes $(\lambda, <_{\alpha})$ where $<_{\alpha}$ is a well order of λ of order type $\alpha \in [\lambda, \lambda^{+}]$.

- **5.8 Claim.** Assume $T \subseteq T_1$ are countable complete first order theories.
- 1) If T is stable not superstable and $\lambda > \aleph_0 + |T_1|$ is not free (see Definition 5.2) then $PC(T_1, T)$ is not categorical in λ .
- 2) If T is unstable and $\lambda > \aleph_0$ then $PC(T_1, T)$ is not categorical in λ .

Proof. Without loss of generality $T, T_1 \subseteq \mathbf{L}_{\omega}$.

- 1) Working in $\mathbf{L}[T_1, T]$ we can find Φ proper for trees with $\omega + 1$ levels as in [Sh:c, VII], i.e., $\tau_{\Phi} \in \mathbf{L}[T_1, T]$, $\mathrm{EM}_{\tau(T_1)}(I, \Phi)$ a model of T_1 (e.g. for $I \subseteq {}^{\omega \geq} \lambda$) satisfying $\mathrm{EM}({}^{\omega \geq} \lambda, \Phi) \models \varphi_n(\bar{a}_n, \bar{a}_{\nu})^{\mathrm{if}(\nu = \eta \upharpoonright n)}$ when $\eta \in {}^{\omega} \lambda, \nu \in {}^{n} \lambda$.
- Let $F: \lambda \to {}^{\omega}\lambda$ exemplify that λ is not free, i.e., its range is not free. Working in $\mathbf{L}[T, T_1, F]$ (so without loss of generality F is one to one), let $M_1 = \mathrm{EM}_{\tau}({}^1\lambda, \Phi), M_2 = \mathrm{EM}_{\tau}({}^{\omega} \to \lambda \cup \mathrm{Rang}(F), \Phi)$ and assume toward contradiction that f is an isomorphism from M_1 onto M_2 and we shall work in $\mathbf{L}[T, T_1, F, f]$, in this universe let $\mathscr{U} \subseteq \lambda$ be of minimal cardinality such that $\{F(\alpha) : \alpha \in \mathscr{U}\}$ is not free (in the same sense). By [Sh 52] (or [Sh:E18]), $|\mathscr{U}|$ is a regular uncountable cardinal, so by renaming without loss of generality $\mathscr{U} = \mu = \mathrm{cf}(\mu) > \aleph_0$. Let $W \subseteq \lambda, |W| = \mu, \{f(a_{\alpha}) : \alpha \in \mathscr{U}\} \subseteq \mathrm{EM}({}^1W, \Phi)$ and let $\langle w_{\alpha} : \alpha < \mu \rangle$ be a filtration of W. Clearly M_1 satisfies $A \subseteq M_1 \land |A| < \mu \Rightarrow \mathbf{S}(A, M) = \{\mathrm{tp}(a, A, M) : a \in M_1\}$

has cardinality $\leq |A| + \aleph_0 < \mu$. This holds in M_2 hence $(\forall \alpha < \mu)(\exists \beta < \mu)[\forall \gamma \in w_{\alpha})[\{f(a_{F(\gamma)}|_n) : n \leq \omega\} \subseteq \mathrm{EM}(^1(w_{\beta}), \Phi)$. We continue as in [Sh:c, VIII,§2] and get contradiction.

2) As in 2.5. $\Box_{5.8}$

5.9 Claim. Assume $\lambda > \aleph_0$ is a free cardinal.

- 1) For $T = \text{Th}(^{\omega}\omega, E_n)_{n<\omega}$, $E_n = \{(\eta, \nu) : \eta, \nu \in {}^{\omega}\omega, \eta \upharpoonright n = \nu \upharpoonright n\}$ for some countable complete $T_1 \supseteq T$, $PC(T_1, T)$ is categorical in λ (T_1 does not depend on λ).
- 2) There is a countable complete stable not superstable T such that if $M \models T, ||M|| \le \lambda$, then the isomorphism type of M is determined by two dimensions.
- *Proof.* 1) As in [Sh 100], T_1 will guarantee that for any $M \in PC(T_1, T)$ we have:
 - $(*)_1$ if $a \in M$ then $\{b \in M : M \models bE_n a \text{ for every } n < \omega\}$ has cardinality ||M||
 - $(*)_2$ if $a \in M, n < \omega$ then $\{b/E_{n+1}^M : b \in a/E_n\}$ has cardinality ||M||.

So suppose $M_1, M_2 \in PC(T_1, T)$ has universe λ and we work in $\mathbf{L}[T, M_1, M_2]$. There is $M'_{\ell} \cong M_{\ell}$ of cardinality λ and $A_{\ell} \subseteq {}^{\omega}\lambda, |A_{\ell}| = \lambda$ for $\ell = 1, 2$ such that

(*)₃
$$|M'_{\ell}| = A_{\ell} \times \lambda$$
, $(\eta, \alpha) E_n(\nu, \beta)$ iff $(\eta, \nu \in A_{\alpha}, \alpha, \beta < \lambda \text{ and}) \eta \upharpoonright n = \nu \upharpoonright n$
(*)₄ $\nu \in {}^{\omega} \gt \lambda \Rightarrow (\exists^{\lambda} \eta) (\nu \triangleleft \eta \in A_{\ell})$.

By the assummption " λ is free" (see Definition 5.2) we can find $g_{\ell}: A_{\ell} \to \omega$ such that $\langle \eta \upharpoonright g_{\ell}(\eta) : \eta \in A_{\ell} \rangle$ is with no repetitions and we shall work in $\mathbf{L}[T_2, M_1, M_2, A_1, A_2, g_1, g_2]$. For $\kappa < \mu$ let \mathscr{F}_{κ} be the family of functions h such that

- $(*)_h^5$ (a) h is a partial one-to-one function from A_1 into A_2
 - (b) $|\mathrm{Dom}(h)| = \kappa$
 - (c) for $\eta_1, \eta_2 \in \text{Dom}(h)$ and $n < \omega$ we have $\eta_1 \upharpoonright n = \eta_2 \upharpoonright n \Leftrightarrow h(\eta_1) \upharpoonright n = h(\eta_2)$
 - (d) if $\ell \in \{1, 2\}$ and $\nu \in A_{\ell}$ and $(\forall n < \omega)(\exists \eta \in A_{\ell})(\nu \upharpoonright n = \eta \upharpoonright n)$ then $\nu \in A_{\ell}$.

Let $A_{\ell} = \{\eta_{\alpha}^{\ell} : \alpha < \lambda\}$. It is easy to choose $h_{\alpha} \in \mathscr{F}_{\aleph_0 + |\alpha|}$ by induction on α increasing continuous with α such that $\eta_{\alpha}^1 \in \text{Dom}(h_{\alpha+1}), \eta_{\alpha}^2 \in \text{Rang}(h_{\alpha+1})$. 2) As in Example 3.19 using ω -power. **5.10 Definition.** 1) We say λ is Ord- μ -free when:

for every linear order $M = (\lambda, <^M)$, λ for some $B \subseteq \lambda$ in $\mathbf{L}[A, B]$, M can be represented as $\cup \{M_i : i < \mu\}$, M_i embeddable into $({}^n\lambda, <_{\mathrm{even}})$ where $\eta <_{\mathrm{even}} \nu \Leftrightarrow (\exists m < n)(m = \ell g(\eta) = \ell g(\nu) \wedge (\eta(m) \neq \nu(m)) \wedge (\eta(m) < \nu(m) \equiv m \text{ even})$ (see Laver [Lv71], [Sh:e, XII,§2]).

- 2) If $\mu = \aleph_0$ we may omit it.
- **5.11 Claim.** If λ is Ord-free <u>then</u> any two strongly \aleph_0 -homogeneous linear orders (see below) of cardinality λ of the same cofinality are isomorphic.
- **5.12 Definition.** I is a strongly \aleph_0 -homogeneous if I is infinite dense isomorphic to any open interval and its interval.

Proof. See above.

§6 Comments on model theory in ZF

Before we comment on model theory without choice we write up the amount of absolute which holds.

- 6.1 Observation. Let T be countable complete first order theory, without loss of generality $\mathbb{L}_{\tau(T)} \subseteq \mathcal{H}(\aleph_0)$ (or if you like $\subseteq \omega$), so $T \subseteq \mathcal{H}(\aleph_0)$.
- 1) "T is stable" is a Borel relation.
- 2) "M is a countable model of $T, q(\bar{y}) \in \mathbf{S}^{<\omega}(M), p(\bar{x}) \in \mathbf{S}^{<\omega}(M)$ and $M_{\ell} \prec M$ for $\ell = 0, 1, 2$ and for stable $T, M_1 \bigcup_{M_0}^{M} M_2, p$ does not fork over M_0 , all coded naturally as a subset of ω " are Borel.
- 3) In part (2), " $p \perp q$ " is Borel as well as " $p \perp_{\text{wk}} q$ " is Borel, also " $p \perp M_0$ " by clause (e) of part (3A).
- 3A) Let T^{eq} be T when we add predicates naming the equivalence classes so have a predicate $P_{\varphi(\bar{x},\bar{y})}$ equivalent to every $\varphi(\bar{x}) \in \mathbb{L}(\tau_T)$, ([Sh:c, III]) and T_{\forall}^{eq} be the universal part (pedantically the consequences of T^{eq}), so $\tau(T), \mathbb{L}(\tau(T^{\text{eq}})), T^{\text{eq}}, T_{\forall}^{\text{eq}}$ are Borel definable from T. Also the following are Borel
 - (a) A is a model of T_{\forall}^{eq} in this observation with universe $\subseteq \omega$ and we use A, A_{ℓ} to denote such models
 - (b) $A_1 \subseteq A_2$ are models of T_{\forall}^{eq} , $A_2 = acl(A_1)$ (in any $M, A_2 \subseteq M \models T^{eq}$), and computing such A_2 naturally defined
 - (c) $p \in \mathbf{S}^m(A)$, A a model of T^{eq}_{\forall} ; i.e. $\{\text{tp}(\bar{a}, A, M) : A \subseteq M \models T^{\text{eq}}, \bar{a} \in {}^rM\}$ we may write acl(A)
 - (d) computing $p \upharpoonright A_2$ from $A_1 \subseteq A_2$ and $p \in \mathbf{S}^m(A_2)$
 - (e) computing $R^m(p, \Delta, 2), R^m(p, \Delta, \aleph_0)$, $\operatorname{rk}^m(p, \Delta, \aleph_0)$ for p an m-type over A, (a model of $T_{\forall}^{\operatorname{eq}}$), $\Delta \subseteq \mathbb{L}(\tau_T)$ finite)
 - (f) $A_1 \subseteq A_2, p(\bar{x})$ an m-type over A_2 (in clause (c)'s sense) and $p(\bar{x})$ does not fork over A_2
 - (g) $A_1 \subseteq A_2, p(\bar{x})$ an m-type over A_2 , the type $p(\bar{x})$ does not fork over A_2 and is stationary over A_1
 - (h) in (g) computing the unique extension $q \in \mathbf{S}^{\ell g(\bar{x})}(A_2)$ of $p(\bar{x})$ not forking over A_1 and $\operatorname{tp}(\bar{a}_0 \hat{a}_1 \hat{a}_1 \dots A_2, M)$ when $A_2 \subseteq M \models T^{\operatorname{eq}}, \bar{a}_n$ realizes $p(\bar{x})$ in M and $p_{A_2}^{\omega} = \operatorname{tp}((\bar{a}_n, A_2 \cup \{\bar{a}_0, \dots, \bar{a}_{n-1}\}, M)$ does not fork over A_1
 - (i) $p_{\ell}(\bar{x}_{\ell}) \in \mathbf{S}^{m(\ell)}(A)$ for $\ell = 1, 2$ are weakly orthogonal
 - (j) for $A_{\ell} \subseteq A$, the stationary types $p_{\ell}(\bar{x}) \in \mathbf{S}^{m(\ell)}(A_{\ell})$ for $\ell = 1, 2$ are orthogonal

- (k) from $A \subseteq A_{\ell}$, A_2 such that $\operatorname{tp}(A_{\ell}, A)$ is stationary for $\ell = 1, 2$ and computing $A', (f_{\ell}, A'_{\ell})$ for $\ell = 1, 2$ such that $A \subseteq A', f_{\ell}$ an isomorphism from A_{ℓ} onto A'_{ℓ} over $A, A'_{\ell} \subseteq A$ for $\ell = 1, 2$ and $A'_1 \bigcup_{A} A'_2$
- (l) $A_1 \subseteq A_2, A \subseteq A_2$ and $p(\bar{x}) \in \mathbf{S}^m(A)$ is orthogonal to A_1 .
- 4) "T has DOP" is a Σ^1_1 -relation (so NDOP is Π^1_2).
- 5) "T has DIDIP" is Σ_1^1 (so NDIDIP is Π_1^1).
- 6) "T has OTOP" in Σ_1^1 .

Proof. Sometimes we give equivalent formulations to prove. (1),(2) are obvious; for "dnf" see clause (f) of part (3A).

3)

- (a) $p \perp_{\text{wq}} q$ just says: for some $A \subseteq M, p, q \in \mathbf{S}^{<\omega}(A, M)$ (or even $p, q \in \mathbf{S}^{\leq\omega}(A, M)$) satisfying: if $\varphi(\bar{x}, \bar{y}, \bar{z}) \in \mathbb{L}(\tau_T)$ and \bar{a} from A we have $p(\bar{x}) \cup q(\bar{y}) \vdash \varphi(\bar{x}, \bar{y}, \bar{a})$ or $p(\bar{x}) \cup p(\bar{y}) \vdash \neg \varphi(\bar{x}, \bar{y}, \bar{a})$ and remember compactness
- (b) $p \perp q$, see clause (j) of part (3A)
- (c) $p \perp M_0$ by clause $(\ell)(\beta)$ of part (3A).
- 3A) E.g.

Clause (e): Because Δ is finite, the value is a natural number and for $\theta \leq \aleph_0, k < \omega$ we have $R^m(p(\bar{x}), \Delta, \theta) \geq k$ iff some Borel set of formulas, see [Sh:c, II,§2] is consistent. Similarly for $(R^m(p(\bar{x}), \Delta, \theta) > k_1) \vee (R^m(p(\bar{x}), \Delta, \theta) = k_1) \wedge \operatorname{Mlt}^m(p(\bar{x}), \Delta, \theta) \geq k_2$).

Clause (f): This is equivalent to "if $A_3 = acl(A_2)$, $\Delta \subseteq \mathbb{L}(\tau_{T^{eq}})$ is finite then there is $q \in \mathbf{S}^m_{\Delta}(A_2)$, i.e. a definition of such type which extend $p \upharpoonright \Delta$ and is definable over $acl(A_1)$.

Clause (g): We can use the definition: for every finite Δ there is $q \in \mathbf{S}_{\Delta}^{m}(A_3)$ definable over $acl(A_1, A_3)$ such that $R^{m}(p(\bar{x}), \Delta, 2) = R^{m}(p(\bar{x}) \cup q(\bar{x}), \Delta, 2)$.

Clause (j): This is equivalent to: $p_{\ell} \in \mathbf{S}^{<\omega}(A_{\ell})$ for $\ell = 1, 2, A_{\ell} \subseteq M$, and for every $n < \omega$ and finite $\Delta_1 \subseteq \mathbb{L}(\tau_T)$ for some finite $\Delta_2 \subseteq \mathbb{L}(\tau_T)$, if $\langle a_0^{\ell}, \dots, a_{n-1}^{\ell} \rangle$ is as in clause (h) with $(A_{\ell}, A_1 \cup A_2, p_{\ell})$ here standing for (A_1, A_2, p) but we have finitely many possibilities for each, then $\operatorname{tp}_{\Delta_2}(\bar{a}_0^1 \cap \dots \cap a_{n-1}^1, A_1 \cup A_2, M), \operatorname{tp}_{\Delta_1}(\bar{a}_0^2 \cap \dots \cap \bar{a}_{n-1}^2, A_1 \cup A_2, M)$ determine the A_1 -type of $\langle \bar{a}_0^1 \cap \dots \cap \bar{a}_{n-1}^1 \cap \bar{a}_0^2 \cap \dots \cap \bar{a}_{n-1}^2 \rangle$ over $A_1 \cup A_2$ in M.

<u>Clause</u> (ℓ): First assume A_1, A_2, A are algebraically closed. We know that $p \perp A_1$ iff there are f, M such that $A_2 \subseteq M \models T^{eq}, f \supseteq id_A(M, M)$ -elementary mapping

(i.e. an automorphism of M) and mapping A_2 to A_2' , $A_1 \bigcup_{A}^{M} A_2'$ such that $p \perp f(p)$.

In the general case as in clause (j) work with "for every finite Δ_1 ...".

4) Obvious by (3) and by the definition (there are countable models of $T, M_{\ell} (\ell \leq 3)$

such that $M_1 \bigcup_{M_1}^{M_3} M_2$, M_3 is $\mathbf{F}_{\aleph_0}^{\ell}$ -constructible over $M_1 \cup M_2$ and $p \in \mathbf{S}^{<\omega}(M_3)$ non-

algebraic such that $p \perp M_1, p \perp M_2$).

5) Obvious by (3) and the definition (equivalent to: there are countable models M_n of $T, M_n \prec M_{n+1}$, and countable N which is $\mathbf{F}_{\aleph_0}^{\ell}$ -atomic over $\cup \{M_n : n < \omega\}$ and non-algebraic $p \in \mathbf{S}^{<\omega}(N)$ such that $n < \omega \Rightarrow p_n \perp M_n$).

6.2 Claim. : 0) <u>Convention</u>:

- (a) T's vocabulary, $\tau = \tau_T$ is well orderable and for simplicity $\subseteq \mathbf{L}$
- (b) M, N denote models of T with universe a set of ordinals
- (c) T a theory in $\mathbb{L}(\tau)$ so |T| is a cardinal; without loss of generality $\tau \subseteq \mathbf{L}_{\lambda}$, $\lambda = |T| + \aleph_0$
- (d) "a model of T" means one with well ordered universe so without loss of generality a set of ordinals.
- 1) DLST and ULST holds (for models as in clause (b)), short for the downward Löwenheim-Skolem-Tarski and the upward Löwenheim-Skolem-Tarski theorems respectively. If T is categorical in $\lambda \geq |T|$ then $T \cup \{\exists \geq n x (x = x) : n < \omega\}$ is complete, etc., all that takes place in some $\mathbf{L}[Y]$ is fine.
- 2) [T complete] T has an \aleph_0 -saturated model iff every model of T has an \aleph_0 -saturated elementary extension iff D(T) can be well ordered.
- 3) Define $\kappa(T) =: \sup\{\kappa(T)^{\mathbf{L}[T,Y]} : Y \text{ a set of ordinals}\}\$ but probably better to use $\kappa^+(T) = \cup \{(\kappa(T)^+)^{\mathbf{L}(T,Y)} : Y \text{ a set of ordinals}\}.$
- 4) Assume T is complete. Every model M of T of cardinality $\leq \lambda$ has a κ -saturated elementary extension of cardinality $\leq \lambda$ iff $|D(T)| \leq \lambda$ and $(a) \vee (b)$ where
 - (a) $|\kappa\rangle \lambda = \lambda$, i.e. $[\lambda]^{\kappa}$ is well ordered
 - (b) $T = \operatorname{Th}(M)$ is stable, $|\lambda^{<\kappa(T)}| = \lambda$ and $|\mathscr{P}(\omega)|$ is a cardinal $\leq \lambda$ if some $p \in \mathbf{S}(B), B \subseteq M \models T, M$ well orderable, $|B| < \kappa(T)$ has a perfect set of stationarization and $\lambda > \aleph_0$.

[Why? As in [Sh:c, III], particularly section 5, hopefully see the proof of [Sh:F701, 1.1].] 5) [T complete]

(a) if $\neg(|D(T)| \leq |T|)$ then there is a family \mathscr{P} of subsets of D(T), each of cardinality $\leq |T|$ and $\cup \{\mathbf{P} : \mathbf{P} \in \mathscr{P}\} = D(T)$ and for each $\mathbf{P} \in \mathscr{P}$ there

is a Φ proper for linear orders, with $\tau(T), \tau(\Phi) \subseteq \mathbf{L}$ such that every model $\mathrm{EM}_{\tau(T)}(I,\Phi)$ satisfies: the model realizes $p \in D(T)$ iff $p \in \mathbf{P}$

- 6) Assume there is no set of \aleph_1 reals. If T is complete countable, D(T) uncountable, $M \models T$ and $\mathbf{P}_M = \{p \in D(T) : M \text{ realized } p\}$ then \mathbf{P}_M is countable.
- 6A) Of course, it is possible that $|D(T)|^{\mathbf{L}[T,\hat{Y}]}$ is large in $\mathbf{L}[T,Y]!$ (e.g. there is a set of |T| independent formulas see 12)(c)).
- 7) If T is complete not superstable, $T_1 \supseteq T$ complete, $\lambda = \operatorname{cf}(\lambda) > |T_1|, \lambda \ge \theta(\mathscr{P}(\omega))$ and axiom $\operatorname{Ax}^3_{\lambda}$ (see [Sh 835], i.e., $|[\lambda]^{\aleph_0}|$ a cardinal) then there is $\langle M_u : u \subseteq \lambda \rangle$ such that
 - (a) $M_u \in PC(T_1, T)$
 - (b) $||M_u|| = \lambda$
 - (c) $u \neq v \subseteq \lambda \Rightarrow M_u \ncong M_v$. [Why? There is a sequence $\langle C_{\delta} : \delta \in S \rangle$, $S \subseteq S_{\aleph_0}^{\lambda}$ stationary $C_{\delta} \subseteq \delta = \sup(C_{\delta})$, $\operatorname{otp}(C_{\delta}) = 0$ (hence we can partition S to λ stationary sets)].
- 8) Define $\beth'_{\alpha}(\lambda)$ by $\beth'_{0}(\lambda) = \lambda$, $\beth'_{\alpha+1}(\lambda) = \theta(\mathscr{P}(\beth'_{\alpha}(\lambda)), \beth'_{\delta}(\lambda) = \cup\{\beth'_{\alpha}(\lambda) : \alpha < \delta\}$, it is a cardinal. If T is countable, Γ is a countable set of $\mathbf{L}(\tau_{T})$ -types and for every $\alpha < \omega_{1}$ there is $M \in \mathrm{EC}_{\beth_{\alpha}}(T,\Gamma)$ so $|M| \subseteq \mathbf{L}$ of power \beth'_{α} or just of power $\geq \beth^{\mathbf{L}[M]}_{\alpha}$ (but we do not say that an ω_{1} -sequence of such models exists!), then there is an $\Phi \in \Upsilon^{\mathrm{or}}_{\aleph_{0}}[T]$ so $|\tau_{\Phi}| = \aleph_{0}$ such that $\mathrm{EM}_{\tau(T)}(I,\Phi) \in \mathrm{EC}(T,\Gamma)$ for every linear order I.

[Why? See proof of (9), but here the members of the tree are finite set of formulas hence the tree is $\subseteq \mathbf{L}[T]$ and we can define the rank in $\mathbf{L}[T]$ but: we let $\langle \Delta_n : n < \omega \rangle$ be an increasing sequence of finite sets of formulas, each $\varphi \in \Delta_n$ has a set of free variables $\subseteq \{x_0, \ldots, x_{n-1}\}$ and $\bigcup_n \Delta_n = \mathbb{L}(\tau_T), \Delta_n$ is closed under change of free variables (modulo the restriction above). We define \mathscr{T}_n as in the proof of part (9) by $p \in \mathscr{T}_n$ is a complete (Δ_n, n) -type. The tree is really $\subseteq \omega > \omega$.]

9) [DC] Assume T is an (infinite) theory with Skolem functions, Γ a set of $\mathbb{L}(\tau_T)$ types and for every $\alpha < \theta(\mathscr{P}(|T|))$ there is $M \in \mathrm{EC}(T,\Gamma)$ of power $\geq \beth_{\alpha}$ in $\mathbf{L}[T,M]$,
then there is Φ such that $\mathrm{EM}_{\tau(T)}(I,\Phi) \in \mathrm{EC}(T,\Gamma)$ for every linear order I.

[Why? A wrong way is to assume $\theta(\mathscr{P}(|T|))$ is regular and in stage n we have an n-indisernible sequence $\mathbf{I}_{\alpha}^{n} \subseteq M$ of cardinality \beth_{α}' for $\alpha < \theta(\mathscr{P}(|T|))$ with n-tuple from \mathbf{I}_{α}^{n} realizing p_{n} , as in the ZFC proof. The problem is that there may be no regular cardinal $\geq \theta(\mathscr{P}(|T|))$. But more carefully let \mathscr{T}_{n} be the set of complete types $p_{n}(x_{0},\ldots,x_{n-1})$ consistent with T, such that it is the type of a sequence of length n which is m-indiscernible for each $m \leq n$. The order on $\mathscr{T} = \bigcup \{\mathscr{T}_{n} : n < \omega\}$ is inclusion, so really \mathscr{T}_{n} is the n-th level. We need $\mathrm{DC}_{\aleph_{0}}$ to have a rank function on this set which has power $\leq \mathscr{P}(|T|)$. We prove by induction on the ordinal γ for each n, that if $p \in \mathscr{T}_{n}$ has rank γ that no indiscernible $\mathbf{I} \subseteq M, M \in \mathrm{EC}(T,\Gamma)$ of cardinality $\geq \beth_{\omega\gamma}^{\mathbf{L}[T]}$ exists.]

- 9A) Of course, if $EC(T,\Gamma)$ has a model $M, |M| \subseteq \mathbf{L}$ of cardinality $\geq \mathbf{Z}'_{\delta}$ where $\delta := \theta(\mathscr{P}(|T|))$ then we do not need DC.
- 9B) We can avoid "T has Skolem functions", see [Sh:F701], in both parts (9) and (9A). The point is that T needs not be complete, without loss of generality T has elimination of quantifiers and we can define T^{SK} which is T+ the axioms of Skolem functions; now for every $\alpha < \theta(\mathscr{P}(|T|))$, there is a model M of T of cardinality $\geq \beth_{\alpha}$ it can be expanded to M^+ , a model of T^{SK} and we can continue (with new function symbols).
- 10) Assume T is complete uncountable. Then all the proofs in $\S 2 + \S 3$ holds except that we do not have the dichotomy OTOP/existence of primes over stable amalgamation. We intend to return to it in [Sh:F701].
- 11) If $p(x_0, \ldots, x_{n-1})$ is a set of $\mathbb{L}(\tau_T)$ -formulas consistent with T then it is realized in some model M in some universe $\mathbf{L}[T, Y]$ hence can be extended to a complete type realized in such M, p hence $\in D_n(T)$ when T is complete. [Why? Work in $\mathbf{L}[T, p]$ O.K. as $p \subseteq \mathbf{L}$ as $T \subseteq L$.]
- **6.3 Lemma.** [Sh:c] can be done in ZF $+(\forall \alpha)([\alpha]^{\aleph_0}$ is well ordered), see [Sh 835] as long as
 - (a) the theory T is in a vocabulary which can be well ordered
 - (b) we deal only with models whose power is a cardinal
 - (c) all notions are in $\mathbf{L}[T,Y], Y \subseteq \text{Ord large enough (so } \mathfrak{C} \text{ is not constant it depends on the universe)}$
 - (d) in [Sh:c, VIII], the case $\lambda > |T_1|$ regular $(\tau(T_1)$ well orderable, too) is clear as using the well ordering $[\lambda]^{\aleph_0}$ we can find $\langle C_{\delta} : \delta \in S_{\aleph_0}^{\lambda} \rangle \in \mathbf{L}[T,Y]$ hence define a partition $\langle S_{\alpha} : \alpha < \lambda \rangle$ of $S_{\aleph_0}^{\lambda}$ such that $(\exists^{\lambda} \alpha)$ $(S_{\alpha}$ stationary (in \mathbf{V}), so increasing Y we are there but
 - (e) Ch VI on ultrapower should be considered separately.

§7 Powers which are not cardinals

We suggest to look at categoricity of countable theories in so-called reasonable cardinals. For them we have the completeness theorem in 7.7. We then uncharacteristically examine a classical example: Ehrenfuecht example (with 3 models in \aleph_0 , see 7.10).

We naturally ask

Question: Can an expansion of the theory of linear orders be categorical in some uncountable power?

We then deal with criterion, i.e. sufficient conditions for categoricity. We intend to continue this in [Sh:F701].

7.1 Convention. T not necessarily $\subseteq \mathbf{L}$.

We may consider

- **7.2 Definition.** 1) For a class \mathbf{C} of powers we say $T_1 \leq_{\mathbf{C}}^{\mathrm{ex}} T_2$ when: for every set X of power $\in \mathbf{C}$ if T_2 has a model with universe X then T_1 has a model with universe X.
- 2) For a class **C** of powers we say $T_1 \leq_{\mathbf{C}}^{\text{cat}} T_2$ when: for every set X of power \in **C** if T_2 is categorical in |X|, (i.e., has one and only one model with universe X up to isomorphism) then T_1 is categorical in |X|.
- 3) In both cases, if **C** is the class of all powers $\geq |T_2|$ we may omit it.
- 7.3 Observation. $\leq_{\mathbf{C}}^{\text{ex}}, \leq_{\mathbf{C}}^{\text{cat}}$ are partial orders.

We may also consider

- <u>7.4 Question</u>: 1) For which countable theories T is there a forcing extension $\mathbf{V}^{\mathbb{P}}$ of \mathbf{V} , model of ZF such that in $\mathbf{V}^{\mathbb{P}}$ the theory T is categorical in some uncountable power?
- 2) As in (1) for reasonable powers, see below.
- **7.5 Definition.** We say that X is a set of reasonable power (or |X| is a reasonable power) when:
 - (a) there is a linear order of X
 - (b) $|X| = |X \times X|$.

7.6 Claim. If T is countable theory and X a set of reasonable power <u>then</u> T has a model with universe X.

Proof. By 7.7. $\square_{7.6}$

- **7.7 Claim.** [ZF] 1) For some first order sentence ψ we have: for a set X the following are equivalent:
 - (a) X is a set of reasonable power
 - (b) if T is a countable theory then T has a model with universe X
 - (c) ψ has a model with universe ψ .
- 2) If T is categorical in |X|, a reasonable power then $T \cup \{(\exists^{\geq n} x)(x=x) : n < \omega\}$ is a complete theory.

Proof. 1) $(b) \Rightarrow (a)$.

First apply clause (b) to $T_1 =$ (the theory of dense linear order with neither first nor last elements), or just $T_1' = \{\psi_1\} \subseteq T_1$, where $\psi_1 \vdash T_1$ so it has a model $M = (X, <^M)$, so $<^M$ linearly ordered X.

Second, apply clause (b) to $T_2 = \text{Th}(\omega, F)$, F a one-to-one function from $\omega \times \omega$ onto ω , or just $T_2' = \{\psi_2\} \in T_2$ expresses this so there is a model $M = (X, F^M)$ of T, so F^M exemplifies $|X| = |X \times X|$.

Note that we have used (b) only for theories consisting of one sentence.

$(a) \rightarrow (b)$.

Use Ehrenfeuch-Mostoswki models.

That is it is enough to prove: using I = (X, <) a linear order

- \boxplus if T' is a countable complete theory with Skolem functions, every term $\sigma(x_0,\ldots,x_{n-1})$ is (by T') equal to a function symbol, $M' \models T$ and $\langle a_n : n < \omega \rangle$ is an indiscernible sequence in $M', p_n = \operatorname{tp}_{qf}(\langle a_0,\ldots,a_{n+1}\rangle,\emptyset,M)$ for $n < \omega$ then we can find $M, \langle a_t : t \in I \rangle$ such that
 - \circledast (a) M is a model of T'
 - (b) M^* has universe X
 - (c) $\langle a_t : t \in I \rangle$ is an indiscernible sequence in M
 - (d) $\langle a_{t_0}, \ldots, a_{t_{n-1}} \rangle$ realizes p_n in M when $t_0 <_I \ldots <_I t_{n-1}$.

Let $<^*$ be a well order $\tau(T)$.

Let $\langle (k_n, F_n) : n < \alpha \leq \omega \rangle$ list with no repetition the pairs (k, F) satisfying $(*)_{k,F}$ such that $k_0 = 1, M \models \forall x [F_0(x) = x]$ where

- $(*)_{k,F}$ (a) $F \in \tau(T)$ is a k-place function symbol
 - (b) there is no $u \subset \{0, \dots, k-1\}$ such that $F^{M'}(a_0, a_1, \dots, a_{k-1}) \in \operatorname{Sk}_M(\{a_\ell : \ell \in u\})$
 - (c) there is no k-place function symbol $F_1 \in \tau(T')$ such that $F_1 <^* F$ and $F_1^{M'}(a_0, \ldots, a_{k_1}) = F^{M'}(a_0, \ldots, a_{k-1})$.

Let
$$Y = \bigcup_{n < \omega} Y_n$$
 where $Y_n = \{(n, t_0, \dots, t_{k_n - 1}) : n < \alpha \text{ and } t_0 <_I \dots < t_{k_{n - 1}}\}.$

Let $g: X \times X \to X$ be one to one onto.

Clearly there is a model as required with universe Y, hence it is enough to prove |Y| = |X|. Clearly $|X| \le |Y|$ as $\{(0,t) : t \in I\} \subseteq Y$. Also $|Y_n| = |X|^{k_n}$ which is 1 if $k_n = 0$ and is |X| if $k_n \ge 1$ as we can prove by induction on n. Moreover, we can choose $\langle f_n : n < \alpha, k_n \ge 1 \rangle$ such that f_n is one-to-one from Y_n onto X as f_n is gotten by composition $k_n - 1$ times of g. This leads to $|Y| \le |X \times \omega| + |\omega|$. But trivially $\aleph_0 \le |X|$ by g hence $|X| \le |Y| \le |X| \times |X| + \aleph_0 = |X|$ hence we are done proving $(b) \Rightarrow (a)$.

Let ψ say "< is a linear order and F(x,y) is a one-to-one function onto.

 $(c) \Rightarrow (a)$: as in the proof of $(b) \Rightarrow (a)$ and also

 $(b) \Rightarrow (c)$: should be clear.

2) Easy, too. $\square_{7.7}$

- 7.8 Discusion We can use an \aleph_0 -saturated model M of T as a set of urelements, i.e. we use a Fraenkel-Mostowski model for the triple (M', a copy of M; finite support; finite partial automorphism of M). Is T categorical in |M'|? The problem is that maybe some $\psi \in \mathbb{L}_{(2^{\aleph_0})^+,\omega}$ define in M' with finitely many parameters, a model M'' of T with universe |M'| such that there is no permutation f of |M'| definable similarly such that f is an isomorphism from M' onto M''. But we may consider (D,\aleph_0) -homomogeneous models of some extension of T (in bigger vocabulary). This seems related to [Sh 199], [Sh 750].
- **7.9 Definition.** T_1 is the theory of dense linear order with neither first nor last element and $c_n < c_{n+1}$ for $n < \omega$ (so $\tau(T_1) = \{<\} \cup \{c_n : n < \omega\}$.

Remark. 1) This is the Eherenfeucht example for $I(\aleph_0, T) = 3$.

2) We can replace T_2 by $T_{i,n}$ with 3 below replaced by 3 + n.

- **7.10 Claim.** [ZF] 1) T_1 is a complete countable first order which is not categorical in any infinite power.
- 2) In fact if T_1 has a model with universe X then T_1 has at least three non-isomorphic models with this universe.
- 3) If in (2) the set X is uncountable (i.e. $|X| \neq |\omega|$) then T_1 has at least \aleph_0 non-isomorphic models with this universe.
- <u>7.11 Question</u>: 1) Consistently (with ZF), in some uncountable power, does $Th(\mathbb{Q}, <$) has exactly 3 models.

Proof. 1) Follows by (2).

2) Let X be a set. For $\ell = 0, 1, 2, 3$ let

 $K_{\ell} = \{N : (a) \mid N \text{ is a model of } T_1 \text{ with universe } X;$

- (b) if $\ell = 1$ then N omit $p(x) = \{c_n < x : n < \omega\}$
- (c) if $\ell = 2$ some $a \in N$ realizes p(x) but no $a \in N$ is the first such element;
- (d) if $\ell = 3$ some element $a \in N$ realizes p(x) and is the first such element ℓ .

Clearly

- \boxplus (a) K_0 is the class of models of T_1 with universe X
 - (b) K_0 is the disjoint union of K_1, K_2, K_3 .

By $(*)_1, (*)_2, (*)_3$ below the result follows:

 $(*)_1$ if $K_2 \neq \emptyset$ then $K_3 \neq \emptyset$.

[Why? Let $M \in K_3$ and we define a $\tau(T_1)$ -model N as follows:

- (i) the universe of N is X = |M|
- (ii) $c_n^N = c_{n+1}^M \text{ for } n < \omega$
- (iii) $N \models a < b \text{ iff } M \models \text{``}a < b \land a \neq c_0 \land b \neq c_0 \text{ or } a = c_0^M \land (b \text{ realizes} p(x) \text{ in } M) \text{ or } b = c_0^N \land a \neq c_0 \land \bigvee b < c_m$ "

Now check that $N \in K_3$ with c_0^M being the $<^N$ -first member of X realizing p(x)]

 $(*)_2$ if $K_3 \neq \emptyset$ then $K_1 \neq \emptyset$.

[Why? Let $M \in K_3$ and $c \in M$ realizes p(x) be the first such element. We define a $\tau(T_1)$ -model N by

- (i) the universe of N is X = |M|
- (ii) $c_n^N = c_n^M$
- (iii) $N \models a < b \text{ iff } M \models \text{``}a < b < c\text{''} \text{ or } M \models \text{``}c < b < a\text{''} \text{ or } M \models \text{``}c \leq a \land b < c\text{''}.$

Now check that $N \in K_1$.]

 $(*)_3$ if $K_1 \neq \emptyset$ then $K_2 \neq \emptyset$.

[Why? Let $M \in K_1$, let $Y = \{a \in X : M \models "c_{2n+1} \le a < c_{2n+2}" \text{ for some } n < \omega\}$ and we define a $\tau(T_1)$ -model N

- (i) the universe of N is X = |M|
- (ii) $c_n^N \equiv c_{2n}^M \text{ for } n < \omega$
- (iii) $N \models a < b \text{ iff } M \models \text{``}a < b \land (a \notin Y) \land (b \notin Y)\text{''} \text{ or } M \models \text{``}a < b \land a \in Y \land b \in Y\text{''} \text{ or } (b \in Y) \land (a \notin Y).$

Now check that $N \in K_2$.]

3) Let $M \in K_1$ have universe X and stipulate $c_{-1} = -\infty, X_n = \{a : M \models c_{n-1} < a \le c_n\}$ for $n < \omega$ so $\langle X_n : n < \omega \rangle$ is a partition of X. Let $S_* = \{n < \omega : X_n \text{ is uncountable}\}.$

Case 1: S_* is infinite.

For any partition $\langle S_n : n < \omega \rangle$ of ω to infinite sets we can define $N \in K_1$ with universe X such that $\{c_n^N : n < \omega\} = \{c_n^M : n \in S_0\}$, on this set $<^M, <^N$ agree, and the set $\{n < \omega : (c_n, c_{n+1})_N \text{ is uncountable}\}$ is any infinite co-infinite set.

Case 2: S_* is finite.

We can find $N \in K_0$ with universe X such that $\max\{n : (c_n, c_{n+1})_N \text{ is uncountable}\}$ is any natural number. $\square_{7.10}$

7.12 Definition. 1) Let N is $(\mathbb{L}_{\infty,\kappa}, \lambda)$ -interpretable in M means (without loss of generality τ_N consist of predicates only): there is $\bar{d} \in {}^{\lambda>}M$ and sequence $\langle \varphi_R(\bar{x}_R, \bar{d}) : R \in \tau_N \rangle$, including R being equality such that

$$\varphi_R(\bar{x}_R, \bar{y}) \in \mathbb{L}_{\infty,\kappa}$$

$$\ell g(\bar{x}_R) = \operatorname{arity}(R)$$

$$|N| = \{ a \in M : M \models \varphi_{=}(a, a, \bar{d}) \}$$

$$R^N = \{ \bar{a} \in {}^{\ell g(x_R)}|M| : M \models \varphi_R(\bar{a}, \bar{d}) \}.$$

- 2) We add "fully" if $\varphi_{=}(\bar{x}_R) = (x_0 = x_1)$ for R being the equality.
- **7.13 Claim.** 1) To prove the consistency of "a first order complete T is categorical in some power $\neq \aleph_0$ " it is enough
 - (*) find a model N of T and $\kappa > \aleph_0$ satisfying: if M is a model of T fully $\mathbb{L}_{\infty,\kappa}(\tau_M)$ -interpretable in N <u>then</u> $M \cong N$; moreover there is a function which is $\mathbb{L}_{\infty,\kappa}(\tau_M)$ -definable in N (with $< \kappa$ parameters) and is an isomorphism from N onto M.
- 2) We can replace $\mathbb{L}_{\infty,\kappa}(N)$ by: there is a set \mathscr{F} such that
 - (a) $\mathscr{F} \subseteq \{f : f \text{ a partial automorphism of } N \text{ with domain of cardinality} < \kappa\}$
 - (b) $(\forall A \subseteq N)(|A| < \kappa \Rightarrow (\exists f \in \mathscr{F})(A \subseteq Dom(f))$
 - (c) \mathscr{F} closed under inverse and composition
 - (d) if $f \in \mathscr{F}, A \in N$ then $(\exists g \in \mathscr{F})(f \subseteq g \cap a \in \text{Dom}(g))$.

Proof. Straight.

Remark. So this categoricity does not imply "not complicated". $\square_{7.13}$

REFERENCES.

- [BLSh 464] John T. Baldwin, Michael C. Laskowski, and Saharon Shelah. Forcing Isomorphism. *Journal of Symbolic Logic*, **58**:1291–1301, 1993. math.LO/9301208.
- [Be84] Steven Buechler. Kueker's conjecture for superstable theories. *Journal of Symbolic Logic*, **49**:930–934, 1984.
- [HHL00] Bradd Hart, Ehud Hrushovski, and Michael C. Laskowski. The uncountable spectra of countable theories. *Annals of Mathematics*, **152**:207–257, 2000.
- [Hr89] Ehud Hrushovski. Kueker's conjecture for stable theories. *Journal of Symbolic Logic*, **54**:207–220, 1989.
- [Hr89d] Ehud Hrushovski. Unidimensional theories. In *Logic Colloquium 88*. North–Holland, 1989.
- [Ke71a] Jerome H. Keisler. On theories categorical in their own power, volume 36. 1971.
- [Las88] Michael C. Laskowski. Uncountable theories that are categorical in a higher power. *The Journal of Symbolic Logic*, **53**:512–530, 1988.
- [LwSh 518] Michael C. Laskowski and Saharon Shelah. Forcing Isomorphism II. Journal of Symbolic Logic, **61**:1305–1320, 1996. math.LO/0011169.
- [Lv71] Richard Laver. On Fraissé's order type conjecture. Annals of Mathematics, 93:89–111, 1971.
- [Mo65] Michael Morley. Categoricity in power. Transaction of the American Mathematical Society, 114:514–538, 1965.
- [Sh 300f] Saharon Shelah. Chapter VI.
- [Sh:e] Saharon Shelah. *Non-structure theory*, accepted. Oxford University Press.
- [Sh 750] Saharon Shelah. On Weak Beth for Cofinality Logic. *Preprint*.
- [Sh:F728] Saharon Shelah. PCF arithmetic with little choice.
- [Sh 835] Saharon Shelah. PCF without choice. Archive for Mathematical Logic, submitted. math.LO/0510229.
- [Sh 3] Saharon Shelah. Finite diagrams stable in power. Annals of Mathematical Logic, **2**:69–118, 1970.

- [Sh 4] Saharon Shelah. On theories T categorical in |T|. Journal of Symbolic Logic, **35**:73–82, 1970.
- [Sh 12] Saharon Shelah. The number of non-isomorphic models of an unstable first-order theory. *Israel Journal of Mathematics*, **9**:473–487, 1971.
- [Sh 31] Saharon Shelah. Categoricity of uncountable theories. In *Proceedings of the Tarski Symposium (Univ. of California, Berkeley, Calif., 1971)*, volume XXV of *Proc. Sympos. Pure Math.*, pages 187–203. Amer. Math. Soc., Providence, R.I, 1974.
- [Sh 52] Saharon Shelah. A compactness theorem for singular cardinals, free algebras, Whitehead problem and transversals. *Israel Journal of Mathematics*, **21**:319–349, 1975.
- [Sh 54] Saharon Shelah. The lazy model-theoretician's guide to stability. Logique et Analyse, 18:241–308, 1975.
- [Sh:E18] Saharon Shelah. A combinatorial proof of the singular compactness theorem. Mineograph notes and lecture in a mini-conference, Berlin, August'77, 1977.
- [Sh 100] Saharon Shelah. Independence results. The Journal of Symbolic Logic, 45:563–573, 1980.
- [Sh 199] Saharon Shelah. Remarks in abstract model theory. Annals of Pure and Applied Logic, 29:255–288, 1985.
- [Sh:c] Saharon Shelah. Classification theory and the number of nonisomorphic models, volume 92 of Studies in Logic and the Foundations of Mathematics. North-Holland Publishing Co., Amsterdam, xxxiv+705 pp, 1990.
- [Sh 497] Saharon Shelah. Set Theory without choice: not everything on cofinality is possible. Archive for Mathematical Logic, **36**:81–125, 1997. A special volume dedicated to Prof. Azriel Levy. math.LO/9512227.
- [Sh:E38] Shelah, Saharon. Continuation of 497: Universes without Choice.
- [Sh:F701] Shelah, Saharon. More on model theory without choice.
- [WT05] Agatha Walczak-Typke. The first-order structure of weakly Dedekind-finite sets. *Journal of Symbolic Logic*, **70**:1161–1170, 2005.
- [WT07] Agatha Walczak-Typke. A model-theoretic approach to structures in set theory without the axiom of choice. In B. Loewe, editor, Algebra, Logic, Set Theory: Festschrift fur Ulrich Felgner zum 65 Geburtstag, Studies in Logic. College Publications at Kings College London. to appear.