

# $\gamma_k(n) = \max\{\lfloor n/(2k+1) \rfloor, 1\}$ for Maximal Outerplanar Graphs with $n \bmod (2k+1) \leq 6$

LIANG ZHAO<sup>1,a)</sup>

Received: November 7, 2016, Accepted: February 9, 2017

**Abstract:** Let  $G = (V, E)$  be an undirected graph with a set  $V$  of nodes and a set  $E$  of edges,  $|V| = n$ . A node  $v$  is said to *distance- $k$  dominate* a node  $w$  if  $w$  is reachable from  $v$  by a path consisting of at most  $k$  edges. A set  $D \subseteq V$  is said a *distance- $k$  dominating set* if every node can be distance- $k$  dominated by some  $v \in D$ . The size of a *minimum distance- $k$  dominating set*, denoted by  $\gamma_k(G)$ , is called the *distance- $k$  domination number* of  $G$ . The value  $\gamma_k(n)$  is defined by  $\gamma_k(n) = \max\{\gamma_k(G) : G \text{ has } n \text{ nodes}\}$ . This paper considers  $\gamma_k(n)$  for *maximal outerplanar graphs*. There is a conjecture  $\gamma_k(n) = \max\{\lfloor n/(2k+1) \rfloor, 1\}$ , which was proved for  $k = 1, 2$ . This paper gives a unified and simpler proof for  $k = 1, 2, 3$ . In fact, a *stronger* result is shown that for all  $n > 2k$  and  $r = n \bmod (2k+1) \leq 6$ , there exist at least  $2k+1-r$  distinct distance- $k$  dominating sets of size at most  $\lfloor n/(2k+1) \rfloor$ , which can be found in linear time.

**Keywords:** distance domination, maximal outerplanar graph, linear-time algorithm

## 1. Introduction

Let  $G = (V, E)$  be an undirected graph with a set  $V$  of nodes and a set  $E$  of edges, where  $|V| = n$ . A node  $v$  is said to *distance- $k$  dominate* a node  $w$  if  $w$  is reachable from  $v$  by a path consisting of at most  $k$  edges. A set  $D \subseteq V$  is said a *distance- $k$  dominating set* if every node can be distance- $k$  dominated by some node  $v \in D$ . The size of a minimum distance- $k$  dominating set, denoted by  $\gamma_k(G)$ , is called the *distance- $k$  domination number* of  $G$ . Let


$$\gamma_k(n) = \max\{\gamma_k(G) : G \text{ is a graph of } n \text{ nodes}\}.$$

In particular,  $\gamma_1(\cdot)$  is the well-known *domination number*.

Domination is one of the fundamental topics in graph theory, see Refs. [1], [5], [6], [10], [11], [12]. This paper considers  $\gamma_k(n)$  for *maximal outerplanar graphs* (MOG). A graph is said *outerplanar* if it can be drawn in the plane without crossing and the nodes belong to the unbounded outer face. It is *maximal* if adding an extra edge breaks this property. It is known that a graph is outerplanar if and only if it does not contain  $K_4$  or  $K_{2,3}$  as a minor (Ref. [3]), and a MOG is a *visibility graph*, i.e., a *triangulation* graph of a simple polygon of  $n$  nodes (Ref. [4]). See illustrations in Fig. 1.

In general, it is not trivial to determine  $\gamma_k(G)$  even for a MOG. Nevertheless, since the outer boundary  $C$  of a MOG is a Hamilton cycle in  $G$ , we see  $\gamma_k(G) \leq \gamma_k(C) = \lfloor \frac{n}{2k+1} \rfloor$ . Hence  $\gamma_k(G) = 1$  if  $n \leq 2k$ . Thus in the following we only consider for  $n > 2k$ .

The above argument shows  $\gamma_k(n) \leq \lfloor \frac{n}{2k+1} \rfloor$ . But in general it is not tight. Instead there is a conjecture  $\gamma_k(n) = \lfloor \frac{n}{2k+1} \rfloor$ , proved for  $k = 1, 2$  (Refs. [1], [10]). In this paper, we give a unified and simpler proof for  $k = 1, 2, 3$ . In fact, we show a stronger result that



**Fig. 1** An illustration of some graphs. Graph  $G_1$  is planar but not outerplanar since it has a  $K_{2,3}$  minor. On the other hand,  $G_2$  is outerplanar but not maximal.  $G_3$  is a maximal outerplanar graph and it is a triangulation of the outer polygon.

for all  $r = n \bmod (2k+1) \leq 6$  (hence for all  $k \leq 3$  and  $n > 2k$ ), there exist at least  $2k+1-r$  *distinct* distance- $k$  dominating sets of size at most  $\lfloor \frac{n}{2k+1} \rfloor$ , which can be found in linear time.

**Related works** Campos and Wakabayashi [2] showed  $\gamma_1(n) = \lfloor (n+t)/4 \rfloor$ , where  $t$  is the number of degree-2 nodes ( $t \geq 2$ ). This result was independently proved by Tokunaga [11] using a coloring-based and simpler proof.

## 2. Preliminaries

Let  $P = u - w - v$  denote a path with nodes  $u, w, v$  and edges  $(u, w), (w, v)$ . A *triangle ear* (simply an *ear* in the following) with respect to a graph  $G = (V, E)$  is such a path  $P = u - w - v$  that  $w \notin V$ ,  $u, v \in V$ , and  $(u, v) \in E$  (see an illustration in Fig. 2). We use  $G + P$  to denote the graph obtained by adding  $P$  to  $G$ , and similarly  $G + P_1 + \dots + P_i = (G + P_1 + \dots + P_{i-1}) + P_i$  for  $i \geq 2$ . In this paper, we prove the following theorem.

**Theorem 1** For any  $k \geq 1$ ,  $p \geq 1$ ,  $0 \leq r \leq \min\{6, 2k\}$  and  $n = p(2k+1) + r$ ,  $\gamma_k(G) \leq p = \lfloor \frac{n}{2k+1} \rfloor$  for any graph  $G = C + P_1 + \dots + P_r$ , where  $C$  is a simple cycle of  $p(2k+1) = n-r$  nodes,  $P_i$  are ears with respect to  $C + P_1 + \dots + P_{i-1}$ ,  $i \geq 2$ . Moreover, at least  $2k+1-r$  distinct distance- $k$  dominating set of  $G$  consisting of at most  $p$  nodes of  $C$  can be found in  $O(n)$  time.

<sup>1</sup> Graduate School of Advanced Integrated Studies in Human Survivability (Shishu-Kan), Kyoto University, Kyoto 606–8306, Japan

<sup>a)</sup> liang@gsais.kyoto-u.ac.jp




Fig. 2 An illustration of an ear  $P = u - w - v$  with respect to  $G$ .




Fig. 3 An illustration for Corollary 1: the  $k = 2$  case for graph  $G_3$  in Fig. 1.

**Corollary 1**  $\gamma_k(G) \leq \left\lfloor \frac{n}{2k+1} \right\rfloor$  for a MOG  $G$  of  $n \geq 2k+1$  nodes and  $k = 1, 2, 3$ .

*Proof.* Let  $p = \left\lfloor \frac{n}{2k+1} \right\rfloor \geq 1$  and  $r = n - p(2k+1)$ . We have  $0 \leq r \leq 2k \leq 6$  since  $k \leq 3$ .

It is well-known (and easy to see) that any MOG with four or more nodes must have an ear  $P = u - w - v$  on the outer boundary, where  $w$  is of degree two. Removing  $w$  we get a MOG with one fewer nodes. Repeating this procedure we can get an *ear decomposition*  $G = G_0 + P_1 + \dots + P_r$ , where  $G_0$  is a MOG of  $p(2k+1) = n - r$  nodes and  $P_i$  are ears on the outer boundary of  $G_{i-1} = G_0 + \dots + P_{i-1}$ ,  $i = 1, \dots, r$ .

Let  $C$  be the outer boundary of  $G_0$ . Clearly  $C$  is a Hamilton cycle of  $G_0$ . Thus  $P_1$  is an ear with respect to  $C$  too, and graph  $G'_1 = C + P_1$  has the same outer boundary as  $G_1$ . Repeating the argument, we see  $P_i$  is an ear with respect to graph  $G'_{i-1} = C + \dots + P_{i-1}$  too, and  $G'_i = G'_{i-1} + P_i$  has the same outer boundary as  $G_i$ ,  $i \geq 2$ . See an illustration in Fig. 3.

By Theorem 1, we have  $\gamma_k(G'_r) \leq p$ . Since graph  $G = G_r$  has the same node set as  $G'_r$  but with a superset of edges,  $\gamma_k(G) \leq \gamma_k(G'_r) \leq p = \left\lfloor \frac{n}{2k+1} \right\rfloor$ .  $\square$

Since the tight example in Ref. [10] for  $k = 1$  also serves as a tight example for any  $k \geq 2$ , we have the next corollary.

**Corollary 2**  $\gamma_k(n) = \left\lfloor \frac{n}{2k+1} \right\rfloor$  for MOGs of  $n \geq 2k+1$  nodes and  $k = 1, 2, 3$ .  $\square$

**Corollary 3** For a MOG with  $n \geq 2k+1$  nodes, at least  $2k+1-r$  distinct distance- $k$  dominating set of size at most  $\left\lfloor \frac{n}{2k+1} \right\rfloor$  can be found in  $O(n)$  time if  $k \leq 3$ , where  $r = n \bmod (2k+1)$ .

*Proof.* An ear decomposition  $G = G_0 + P_1 + \dots + P_r$  can be found by repeatedly finding and removing degree-2 nodes. For that purpose, we store the graph by an adjacency list and calculate the degrees in  $O(n)$  time (notice that the number of edges is  $2n-3$ ). We store the nodes using a bucket by their degrees. This can be done in  $O(n)$  time. Finding a node with (residual) degree two takes  $O(1)$  time. Then we set its degree to zero and for *all its neighbors in the adjacency list*, subtract their degrees by one unless it is zero (notice that we do not change the adjacency list). Then we update the bucket and continue. It is easy to see that the total time for updating the bucket is  $O(n)$  as there are  $O(n)$  edges.

On the other hand, determining the Hamilton cycle  $C$  for  $G_0$

requires  $O(n)$  time (Ref. [8]). Finding  $2k+1-r$  distinct distance- $k$  dominating set for  $G'_r$ , which is also a distance- $k$  dominating set for  $G$ , requires  $O(n)$  time by Theorem 1. Thus the total running time is  $O(n)$ .  $\square$

**Remark** We remark that Theorem 1 can be applied to non-MOGs. For example, it can be applied to graph  $G_1$  in Fig. 1, which is even not outerplanar.

### 3. Proof for Theorem 1

In this section, we prove Theorem 1. Let  $\text{dist}_G(u, v)$  denote the distance between nodes  $u$  and  $v$  in a graph  $G$ , i.e., the minimum number of edges required to connect  $u$  and  $v$  in  $G$ . Given a set  $D$  of nodes, let  $\text{dist}_G(D, v)$  denote the distance between  $D$  and a node  $v$ , i.e.,

$$\text{dist}_G(D, v) = \min_{u \in D} \{\text{dist}(u, v)\}.$$

Thus  $D$  is a distance- $k$  dominating set for graph  $G$  if and only if  $\text{dist}_G(D, v) \leq k$  for all nodes  $v$ . The next lemma is obvious by the definition of ear.

**Lemma 1** Let  $G = (V, E)$  be an undirected graph,  $D \subseteq V$  be a set of nodes, and  $P$  be an ear to  $G$ . Then  $\text{dist}_{G+P}(D, v) = \text{dist}_G(D, v)$  for all  $v \in V$ .  $\square$

#### 3.1 Case $r = 0$ (i.e., $G = C$ is a Cycle of $p(2k+1)$ Nodes)

Let the  $p(2k+1)$  nodes of cycle  $C$  be  $v_1, v_2, \dots, v_{p(2k+1)}$  such that  $v_{i+1}$  is a neighbor of  $v_i$ . Obviously  $\gamma_k(C) = p$ . In fact, there are exactly  $2k+1$  size- $p$  distinct distance- $k$  dominating sets on  $C$ :

$$S_i = \{v_i, v_{2k+1+i}, \dots, v_{(p-1)(2k+1)+i}\}, \quad i = 1, 2, \dots, 2k+1.$$

Finding and outputting all of them requires  $O(n)$  time, proving this case.

#### 3.2 Case $r \geq 1$ : a Glance of the Proof

Consider sets  $S_i$  defined in Section 3.1. First notice that for any *adjacent* nodes  $u$  and  $u'$  on  $C$ , their distances to an  $S_i$  can have exactly one of the next two relations:

- $\text{dist}_C(S_i, u) = \text{dist}_C(S_i, u') = k$ ,
- $\text{dist}_C(S_i, u) - \text{dist}_C(S_i, u') = \pm 1$ .

Moreover, knowing  $u, u', \text{dist}_C(S_i, u)$  and  $\text{dist}_C(S_i, u')$  (of valid values), we can *uniquely* determine  $S_i$  in  $O(1)$  time.




Observe that adding an ear  $P$  to  $C$  may make some  $S_i$  infeasible (i.e., cannot distance- $k$  dominate  $C + P$ ), but there can be *exactly one* of them. We conjecture that adding  $r$  ears can make at most  $r$  sets  $S_i$  infeasible, and thus Theorem 1 holds for all  $r$ . So far we can prove it for  $r \leq 6$ , as shown in the following subsections.

#### 3.3 Case $r = 1$ (One Ear)

Suppose that an ear  $P_1 = u_1 - w_1 - u'_1$  is added to  $C$  and node  $w_1$  is *not* distance- $k$  dominated by some  $S_i$  (see an illustration in Fig. 4). Clearly this can happen only if  $\text{dist}_{C+P_1}(S_i, u_1) = \text{dist}_{C+P_1}(S_i, u'_1) = k$ , thus  $\text{dist}_C(S_i, u_1) = \text{dist}_C(S_i, u'_1) = k$  by Lemma 1, hence  $S_i$  is unique and can be determined in  $O(1)$  time as shown in Section 3.2. This proved for case  $r = 1$ .

#### 3.4 Case $r = 2$ (Two Ears)

Let  $S_i$  be the unique set that cannot distance- $k$  dominate  $C + P_1$ .

Fig. 4 Illustration for Case  $r = 1$ .Fig. 5 Illustration for Case  $r = 2$ : In the first two sub-cases,  $P_2$  is an ear to  $C$  too (notice that the graph may not be outerplanar), hence the argument for Case  $r = 1$  can be applied again; In the last sub-case,  $P_2$  is an ear with respect to  $P_1$ .Fig. 6 Illustration for case  $r = 3$ : We only need to consider these two sub-cases.

Suppose an ear  $P_2 = u_2 - w_2 - u'_2$  is added to  $C + P_1$  (see Fig. 5). Clearly if  $P_2$  is an ear with respect to  $C$  too, then we are done, because, by applying the argument for  $r = 1$ ,  $P_2$  can make at most one more  $S_h$  infeasible to  $C + P_1 + P_2$ . Thus we only need to consider when  $P_2$  is not an ear with respect to  $C$  but to  $P_1$ . Without loss of generality, assume  $u_2 = w_1$  and  $u'_2 = u'_1$ .

Now suppose that  $w_2$  is not distance- $k$  dominated by some set  $S_h$  that distance- $k$  dominates  $C + P_1$ . This can happen only if  $\text{dist}_{C+P_1+P_2}(S_h, u_2) = \text{dist}_{C+P_1+P_2}(S_h, u'_2) = k$ . Hence  $\text{dist}_C(S_h, u'_1) = k$  and  $\text{dist}_C(S_h, u_1) = k - 1$  by easy calculation. Therefore  $S_h$  can be uniquely determined as shown in Section 3.2. Thus we have at least  $2k - 1$  sets  $S_j$  ( $j \notin \{i, h\}$ ) that distance- $k$  dominates both  $w_1$  and  $w_2$ , hence  $C + P_1 + P_2$ .

Determining the sub-case and, if necessary,  $S_h$ , requires  $O(1)$  time, hence the running time for case  $r = 2$  is  $O(n)$  too. For  $k = 1$ , since  $r = n \bmod (2k + 1) \leq 2$ , the proof finishes here.

### 3.5 Case $r = 3$ (Three Ears, Thus $k \geq 2$ )

So far we showed that adding two ears  $P_1$  and  $P_2$  can make at most two of the sets  $S_j$  infeasible to  $C + P_1 + P_2$ . Now we show that adding a third ear  $P_3 = u_3 - w_3 - u'_3$  to  $C + P_1 + P_2$  can make at most one more  $S_j$  infeasible.

It is easy to see that if  $P_2$  is an ear to  $C$ , or  $P_2$  is an ear to  $P_1$  but  $P_3$  is an ear to  $C$  or  $P_1$ , then we can apply the same argument for case  $r = 2$ . In fact, by Lemma 1, we only need to consider sub-cases in which every  $P_i$  is an ear to  $P_{i-1}$ . For  $r = 3$ , they are illustrated by sub-cases (a) and (b) in Fig. 6.

Assume that  $w_3$  is not distance- $k$  dominated by some  $S_\ell$  that distance- $k$  dominates  $C + P_1 + P_2$ . Sub-case (a) can happen only

if  $\text{dist}_C(S_\ell, u'_1) = k - 1$  and  $\text{dist}_C(S_\ell, u_1) = k$ . Hence we can determine the unique  $S_\ell$ . For sub-case (b), it can happen only if  $\text{dist}_{C+P_1+P_2+P_3}(S_\ell, u'_1) = \text{dist}_{C+P_1+P_2+P_3}(S_\ell, w_2) = k$ , implying  $\text{dist}_{C+P_1}(S_\ell, w_1) = k - 1$ , hence  $\text{dist}_C(S_\ell, u_1) = k - 2$ . This is impossible since  $\text{dist}_C(S_\ell, u'_1) = k$ , showing that such an  $S_\ell$  does not exist in sub-case (b). Determining the sub-case and  $S_\ell$  requires  $O(1)$  time, proving Case  $r = 3$ .

### 3.6 Case $r = 4$ (Four Ears)

We showed that there can be at most three of  $S_j$  infeasible to  $C + P_1 + P_2 + P_3$ . Now we want to show that adding a fourth ear  $P_4 = u_4 - w_4 - u'_4$  can make at most one more infeasible. As a conclusion, this is a false proposition. Nevertheless, we show how to overcome this difficulty by careful argument. Again, we only need to consider the sub-cases in which every  $P_i$  is an ear of  $P_{i-1}$ , as shown in Fig. 7.

Assume that  $w_4$  is not distance- $k$  dominated by some  $S_q$  that distance- $k$  dominates  $C + P_1 + P_2 + P_3$ . This can happen only if  $\text{dist}_{C+P_1+\dots+P_4}(S_q, u_4) = \text{dist}_{C+P_1+\dots+P_4}(S_q, u'_4) = k$ . Then we can calculate feasible labels  $\text{dist}_C(S_q, u_1)$  and  $\text{dist}_C(S_q, u'_1)$ . Easy calculation shows that it is impossible for sub-cases (a-1) and (b-1). For sub-case (a-2), the unique feasible distance labeling is  $\text{dist}_C(S_q, u_1) = k - 2$  and  $\text{dist}_C(S_q, u'_1) = k - 1$ . For sub-case (b-2), however, there are two feasible labelings:

- $\text{dist}_C(S_q, u_1) = k - 2$  and  $\text{dist}_C(S_q, u'_1) = k - 1$ ,
- $\text{dist}_C(S_q, u_1) = k$  and  $\text{dist}_C(S_q, u'_1) = k - 1$ .

Nevertheless, recall that sub-case (b-2) is derived from sub-case (b) (see Section 3.5), for which all sets  $S_i$  feasible to  $C + P_1 + P_2$  are feasible to  $C + P_1 + P_2 + P_3$  too. Therefore the total number of sets  $S_j$  infeasible to  $C + P_1 + \dots + P_4$  for sub-case (b-2) is still no more than  $2 + 0 + 2 = 4$ . On the other hand, since it is clear that the total running time is  $O(n)$ , we finished proving Case  $r = 4$ . For  $k = 2$ , since  $r = n \bmod (2k + 1) \leq 4$ , the proof for  $\gamma_2(G) \leq \max\{1, \lfloor n/5 \rfloor\}$  finishes here.  $\square$

### 3.7 Cases $r = 5, 6$ (Hence $k \geq 3$ )


Again, we only need to consider the sub-cases in which every  $P_i$  is an ear of  $P_{i-1}$ . For each sub-case, define

$$k_i = \text{the number of } S_j \text{ that cannot distance-}k \text{ dominate } C + P_1 + \dots + P_i.$$


All we want to show (see Section 3.2) is that for all  $r$ ,  $k_r \leq r$ . So far we have shown it for  $r \leq 4$ . Now we prove it for  $r = 5, 6$ . See Figs. 8, 9, in which we start from  $r = 4$  and have marked  $k_r$  for each sub-case. The detail of the distance labels are omitted since it is much technical and not interesting.

## 4. Conclusion and Future Work

This paper showed  $\gamma_k(n) = \max\left\{\left\lfloor \frac{n}{2k+1} \right\rfloor, 1\right\}$  for  $k \leq 3$  and MOGs with a linear-time construction algorithm. In fact, letting  $r = n \bmod (2k + 1)$ , it shows a stronger result that at least  $2k + 1 - r$  distinct distance- $k$  dominating sets of size at most  $\left\lfloor \frac{n}{2k+1} \right\rfloor$  can be found in linear time for all  $n \geq 2k + 1$  and  $r \leq 6$ . Currently we are working on a simple proof for  $n \bmod (2k + 1) \geq 7$  cases (or to develop a counterexample), and trying to improve the results for guarding numbers and vertex cover numbers as considered in Ref. [1].



**Fig. 7** Sub-cases for  $r = 4$ . (a-\*) and (b-\*) are derived from sub-cases (a) and (b) in Fig. 6, respectively. In all sub-cases,  $P_i$  is an ear of  $P_{i-1}$  for all  $i$ .



**Fig. 8** Sub-cases for  $r = 5, 6$  (part 1 of 2).

**Acknowledgments** This work was supported by JSPS KAKENHI Grant Numbers 23700018, 25330026 and John-man Program of Kyoto University. The author would like to thank Prof. Dr. Dorothea Wagner for her kind support. Most of the work was done when the author was visiting her group at KIT, Germany.

Finally the author would like to thank the anonymous referees for their valuable comments.




Fig. 9 Sub-cases for  $r = 5, 6$  (part 2 of 2).

## References

- [1] Canales, S., Hernandez, G., Martins, M. and Matos, I.: Distance domination, guarding and vertex cover for maximal outerplanar graph, arXiv:1307.2043 [cs.CG] (2013).
- [2] Campos, C.N. and Wakabayashi, Y.: On dominating sets of maximal outerplanar graphs, *Discrete Applied Mathematics*, Vol.161, No.3, pp.330–335 (2013).
- [3] Diestel, R.: *Graph Theory*, p.107, Springer-Verlag (2000).
- [4] ElGindy, H.A.: *Hierarchical decomposition of polygons with applications*, Ph.D. thesis, School of Computer Science, University of McGill (1985).
- [5] Haynes, T.W., Hedetniemi, S.T. and Slater, P.J.: *Fundamentals of Domination in Graphs*, Marcel Dekker, New York (1998).
- [6] Haynes, T.W., Hedetniemi, S.T. and Slater, P.J.: *Domination in Graphs: Advanced Topics*, Marcel Dekker, New York (1998).
- [7] Hansberg, A., Meierling, D. and Volkmann, L.: Distance Domination and Distance Irredundance in Graphs, *The Electronic J. Combinatorics*, Vol.14, No.R35 (2007).
- [8] Mitchell, S., Beyer, T. and Jones, W.: Linear Algorithms for Isomorphism of Maximal Outerplanar Graphs, *J. ACM*, Vol.26, No.4, pp.603–610 (1979).
- [9] Meir, A. and Moon, J.W.: Relations between packing and covering number of a tree, *Pacific J. Math.*, Vol.61, pp.225–233 (1975).
- [10] Matheson, L.R. and Tarjan, R.E.: Dominating Sets in Planar Graphs, *European Journal of Combinatorics*, Vol.17, No.6, pp.565–568 (1996).
- [11] Tokunaga, S.: Dominating sets of maximal outerplanar graphs, *Discrete Applied Mathematics*, Vol.161, No.18, pp.3097–3099 (2013).
- [12] Tian, F. and Xu, J.M.: A note on distance domination numbers of graphs, *Australasian J. Combinatorics*, Vol.43, pp.181–190 (2009).



**Liang Zhao** received his B.S. and B.E. degrees from Tsinghua University (Beijing) in 1995, M.E. and Ph.D. degrees from Kyoto University in 1999 and 2002 respectively. Then he worked for Utsunomiya University for four years and since 2006, he has been engaged in Kyoto University. His research interests include

algorithms and applications. He is a member of IEEE, ACM, IEICE and ORSJ.