

# Quasi-periodic solutions of the 2D Euler equation

Nicolas Crouseilles and Erwan Faou

July 12, 2021

## Abstract

We consider the two-dimensional Euler equation with periodic boundary conditions. We construct time quasi-periodic solutions of this equation made of localized travelling profiles with compact support propagating over a stationary state depending on only one variable. The direction of propagation is orthogonal to this variable, and the support is concentrated on flat strips of the stationary state. The frequencies of the solution are given by the locally constant velocities associated with the stationary state.

## 1 Introduction

We consider the two-dimensional Euler equation written in terms of vorticity

$$\partial_t \omega + u \cdot \nabla \omega = 0,$$

where  $\omega(t, x, y) \in \mathbb{R}$ ,  $\nabla = (\partial_x, \partial_y)^T$  with  $(x, y) \in \mathbb{T}^2$  the two-dimensional torus  $(\mathbb{R}/2\pi\mathbb{Z})^2$ . The divergence free velocity field  $u$  is given by the formula

$$u = J \nabla \psi \quad \text{with} \quad \psi = \Delta^{-1} \omega, \quad \text{where} \quad J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad (1.1)$$

$J$  is the canonical symplectic matrix. Here  $\Delta^{-1}$  is the inverse of the Laplace operator on functions with average 0 on  $\mathbb{T}^2$ . We can rewrite this equation as

$$\begin{cases} \partial_t \omega + \{\psi, \omega\} = 0, \\ \Delta \psi = \omega, \end{cases} \quad (1.2)$$

with the 2D Poisson bracket for functions on  $\mathbb{T}^2$ :

$$\{f, g\} = (\partial_x f)(\partial_y g) - (\partial_y f)(\partial_x g).$$

The equation (1.2) possesses many stationary states. For all functions  $F : \mathbb{R} \mapsto \mathbb{R}$  and  $\psi^0 : \mathbb{T}^2 \rightarrow \mathbb{R}$  satisfying  $\Delta\psi^0 = F(\psi^0)$ , then the couple of functions  $\omega(t, x, y) = F(\psi^0(x, y))$  and  $\psi(t, x, y) = \psi^0(x, y)$  solve (1.2). We refer to [1, 4, 5, 8] for further analysis of these particular solutions.

Another class of stationary states are given by functions depending only on one variable (shear flows): for any smooth  $V(x)$  periodic in  $x$ , the couple  $\omega(t, x, y) = V''(x)$  and  $\psi(t, x, y) = V(x)$  is solution of the 2D Euler equation. Note that  $u(x) = (0, V'(x))^T$ .

The goal of this paper is to construct solutions of (1.2) of the form

$$\omega(t, x, y) = V''(x) + \sum_{k=1}^K \Omega_k(x, y - v_k t), \quad (1.3)$$

where the functions  $\Omega_k$  are localized around points  $(x_k, y_k) \in \mathbb{T}^2$  such that  $V''(x) = 0$  in a neighborhood of  $x_k$  which correspond to flat strips of the stationary state  $V''(x)$ . The points  $y_k$  are arbitrary points in  $[0, 2\pi]$ , and  $v_k = V'(x_k)$  is the locally constant velocity associated with  $V''(x)$ . The profiles  $\Omega_k$  are constructed as stationary states of the 2D Euler on  $\mathbb{R}^2$  with radial symmetry around  $(x_k, y_k)$ , compact support and zero average.

This very simple and explicit construction allows to construct quasi-periodic solutions to the 2D Euler equation corresponding to invariant tori of any given dimension in the dynamics. Let us stress that this is in surprising contrast with the traditional situation in nonlinear Hamiltonian PDEs such as Schrödinger and wave equations for which the construction is much more difficult and requires in general the use of Nash-Moser or KAM iterations, see for instance [6, 3, 9, 7, 2].

## 2 Construction

Let  $V(x)$  be a periodic function with zero average in  $x \in [0, 2\pi]$ . We make the following assumptions on  $V$ :

**Hypothesis 2.1** *Let  $K \in \mathbb{N}$ . For all  $k = 1, \dots, K$ , there exist  $a_k < b_k$  in  $[0, 2\pi]$  such that for all  $j, k \in \{1, \dots, K\}$ , we have  $[a_j, b_j] \cap [a_k, b_k] = \emptyset$ , and such that  $V''(x) = 0$  for  $x \in [a_k, b_k]$ .*

This hypothesis implies that for  $x \in [a_k, b_k]$ ,  $V'(x) =: v_k$  is constant. Let us seek a solution  $\omega(t, x, y)$  under the form

$$\omega(t, x, y) = V''(x) + \sum_{k=1}^K \chi_k(t, x, y),$$

where for all  $k$ ,  $\chi_k(t, x, y)$  is of zero average, and the support of  $\chi_k$  and  $\Delta^{-1}\chi_k$  is included in  $]a_k, b_k[$ . This implies in particular that for all  $j$  and  $k$ ,

$$\{\Delta^{-1}\chi_j, \chi_k\} = 0 \quad \text{for } j \neq k.$$

Inserting this decomposition into (1.2), we thus obtain

$$\sum_{k=1}^K \left( \partial_t \chi_k + \{\Delta^{-1}V'', \chi_k\} + \{\Delta^{-1}\chi_k, V''\} + \{\Delta^{-1}\chi_k, \chi_k\} \right) = 0. \quad (2.1)$$

We seek for travelling wave solutions  $\chi_k(t, x, y) = \Omega_k(x, y - v_k t)$ . Equation (2.1) then becomes

$$\sum_{k=1}^K \left( (V'(x) - v_k) \partial_y \Omega_k - V'''(x) \partial_y \Delta^{-1} \Omega_k + \{\Delta^{-1} \Omega_k, \Omega_k\} \right) = 0,$$

where  $\Omega_k(x, y)$  and  $\Psi_k(x, y) = \Delta^{-1}\Omega_k$  have compact support in  $]a_k, b_k[ \times \mathbb{T}$ . Using Hypothesis 2.1, we have:  $V'''(x) = 0$  and  $V'(x) = v_k$  for  $x \in [a_k, b_k] \subset [0, 2\pi]$ . As the intervals  $[a_k, b_k]$  are pairwise disjoint, the system of equations to solve is hence: For all  $k = 1, \dots, K$ ,

$$\{\Psi_k, \Omega_k\}(x, y) = 0 \quad \text{and} \quad \Delta \Psi_k(x, y) = \Omega_k(x, y), \quad (x, y) \in [a_k, b_k] \times \mathbb{T}. \quad (2.2)$$

In other words, the couple  $(\Omega_k, \Psi_k)$  is a smooth stationary state of the Euler equation with support on the flat strip  $]a_k, b_k[ \times \mathbb{T} \subset \mathbb{T}^2$ .

To prove the existence of such functions  $(\Omega_k, \Psi_k)$ , take  $k \in \{1, \dots, K\}$ , and fix  $(x_k, y_k) \in ]a_k, b_k[ \times [0, 2\pi]$ . Let us perform the local action-angle change of coordinates  $x - x_k = \sqrt{2r} \cos \theta$ ,  $y - y_k = \sqrt{2r} \sin \theta$  for  $(x, y)$  close enough to  $(x_k, y_k)$ . The jacobian matrix is

$$M = \begin{pmatrix} \frac{1}{\sqrt{2r}} \cos \theta & \frac{1}{\sqrt{2r}} \sin \theta \\ -\sqrt{2r} \sin \theta & \sqrt{2r} \cos \theta \end{pmatrix},$$

and we verify that this matrix satisfies  $M^T J M = J$ , which means that the change of coordinate is symplectic. Hence this transformation preserves the Poisson bracket and if we define  $\tilde{\Psi}_k(r, \theta) = \Psi_k(x, y)$  and  $\tilde{\Omega}_k(r, \theta) = \Omega_k(x, y)$  we have

$$\begin{aligned} \{\Psi_k, \Omega_k\} &= (\partial_x \Psi_k)(\partial_y \Omega_k) - (\partial_y \Psi_k)(\partial_x \Omega_k) \\ &= (\partial_r \tilde{\Psi}_k)(\partial_\theta \tilde{\Omega}_k) - (\partial_\theta \tilde{\Psi}_k)(\partial_r \tilde{\Omega}_k) = \{\tilde{\Psi}_k, \tilde{\Omega}_k\}. \end{aligned}$$

The Laplace operator in coordinate  $(r, \theta)$  is

$$\Delta f = 2(\partial_r f + r \partial_{rr} f) + \frac{1}{2r} \partial_{\theta\theta} f.$$

Now take a smooth function  $\tilde{\Psi}_k(r)$  on  $[0, +\infty]$  such that  $\int_0^{+\infty} \tilde{\Psi}_k(r) dr = 0$  and with compact support in  $r < \min(|x_k - a_k|, |x_k - b_k|)$ . Set

$$\tilde{\Omega}_k(r) = \Delta \tilde{\Psi}_k(r) = 2(\tilde{\Psi}'_k(r) + r\tilde{\Psi}''_k(r)).$$

We verify that  $\tilde{\Omega}_k(r)$  has compact support, that  $\int_0^\infty \tilde{\Omega}_k(r) dr = 0$ , and that  $\{\tilde{\Psi}_k, \tilde{\Omega}_k\} = 0$  as  $\tilde{\Psi}_k$  and  $\tilde{\Omega}_k$  only depend on  $r$ . Going back to the variables  $(x, y)$ , we verify that the function  $(\Omega_k(x, y), \Psi_k(x, y))$  extended by 0 outside the strip  $]a_k, b_k[ \times \mathbb{T}$  satisfies (2.2) and are of zero average on  $\mathbb{T}^2$ .

This proves the following result:

**Theorem 2.2** *Assume that  $V = V(x)$  is a periodic function with zero average satisfying Hypothesis 2.1. Then for all  $k = 1, \dots, K$ , there exists points  $(x_k, y_k) \in ]a_k, b_k[ \times \mathbb{T}$  and zero average functions  $(\Omega_k, \Psi_k)$  with compact support in  $]a_k, b_k[ \times \mathbb{T}$  and radial symmetry around  $(x_k, y_k)$  such that the couple*

$$\omega(t, x, y) = V''(x) + \sum_{k=1}^K \Omega_k(x, y - v_k t),$$

and

$$\psi(t, x, y) = V(x) + \sum_{k=1}^K \Psi_k(x, y - v_k t),$$

with  $v_k = V'(x_k)$  is a quasi-periodic solution of (1.2).

**Acknowledgements:** It is a great pleasure to thank Sergei Kuksin and Nicolas Depauw for many helpful discussions.

## References

- [1] A. Ambrosetti and M. Struwe, *Existence of Steady Vortex Rings in an Ideal Fluid*, (1989).
- [2] M. Berti and P. Bolle, *Quasi-periodic solutions of NLS on  $T^d$* , Rend. Mat. Acc. Naz. Lincei, 22, (2011) 223–236.
- [3] J. Bourgain, *Construction of approximative and almost-periodic solutions of perturbed linear Schrödinger and wave equations*. Geom. Funct. Anal. 6 (1996) 201–230.
- [4] E. Caglioti, P.-L. Lions, C. Marchioro and M. Pulvirenti, *A Special Class of Stationary Flows for Two-Dimensional Euler Equations: A Statistical Mechanics Description*, Commun. Math. Phys. 143 (1992) 501–525.

- [5] E. Caglioti, M. Pulvirenti and F. Rousset, *On a constrained 2-D Navier-Stokes equation*, Commun. Math. Phys. 290 (2009) 651–677.
- [6] W. Craig and C. E. Wayne, *Periodic solutions of nonlinear Schrödinger equations and the Nash-Moser method*, Hamiltonian mechanics (Toruń, 1993) 103-122.
- [7] H. L. Eliasson, S. B. Kuksin, *KAM for non-linear Schroedinger equation*, Annals of Math. 172 (2010) 371–435)
- [8] M. J. M. Hill, *On a spherical vortex*, Phil. Trans. Roy. Soc. London 185 (1894) 213–245.
- [9] J. Pöschel, *A KAM-theorem for some nonlinear partial differential equations*. Ann. Sc. Norm. Sup. Pisa 23 (1996) 119–148

**Authors address:**

N. Crouseilles and E. Faou, INRIA and ENS Cachan Bretagne, Avenue Robert Schumann, 35170 Bruz, France.

Nicolas.Crouseilles@inria.fr, Erwan.Faou@inria.fr