On the existence of a connected component of a graph

Kirill Gura*

Jeffry L. Hirst[†]

Carl Mummert[‡]

February 7, 2015

Abstract

We study the reverse mathematics and computability of countable graph theory, obtaining the following results. The principle that every countable graph has a connected component is equivalent to ACA_0 over RCA_0 . The problem of decomposing a countable graph into connected components is strongly Weihrauch equivalent to the problem of finding a single component, and each is equivalent to its infinite parallelization. For graphs with finitely many connected components, the existence of a connected component is either provable in RCA_0 or is equivalent to induction for Σ_2^0 formulas, depending on the formulation of the bound on the number of components.

Keywords: Reverse mathematics, Weihrauch reducibility, component, graph, connected, partition, parallelization

MSC Subject Class (2000): 03B30, 03D30, 03F35, 03D45

^{*}Department of Mathematics, Marshall University, 1 John Marshall Drive, Huntington, WV 25755. *Email:* kirill@gura.email

[†]Department of Mathematical Sciences, Appalachian State University, Walker Hall, Boone, NC 28608. *Email:* jlh@math.appstate.edu

[‡]Department of Mathematics, Marshall University, 1 John Marshall Drive, Huntington, WV 25755. *Email:* mummertc@marshall.edu

1 Introduction

The study of countable graph theory in reverse mathematics can be traced back to the founding reverse mathematics works of Friedman [7,8], which include the graph theoretic principles König's Lemma and Weak König's Lemma. We focus here on principles that postulate the existence of a connected component of a countable graph.

Motivated by a suggestion of K. Hatzikiriakou, Hirst [9] proved that the principle that a countable graph may be decomposed into its connected components is equivalent to ACA_0 over RCA_0 . In Theorem 2.1, we show that the principle postulating the existence of a single connected component of each countable graph is already equivalent to ACA_0 . In the subsequent section, we show that, when there is a finite bound on the number of connected components, the strength of statements pertaining to the decomposition depends on the formulation of the finite bound. In some cases, a connected component may be obtained in RCA_0 (Theorem 3.1), while in other cases additional induction is required (Theorem 3.2).

When all connected components of the graph are finite, it is natural to ask whether one can find an infinite set of vertices such that no two are in the same component. Theorem 4.4 shows that ACA_0 is required to find solutions to this problem. Moreover, there is a computable graph such that there is no infinite c.e. set of vertices with no two in the same component. In Section 5, we isolate a combinatorial principle that gives another view of Theorem 2.1.

In the final section, we study the problem of finding a decomposition of a graph into components and the problem of finding a single component of a graph from the viewpoint of Weihrauch reducibility. We show that these principles are strongly Weihrauch equivalent to each other and to their parallelizations, and their strong Weihrauch degree is that of \widehat{LPO} , which is the infinite "parallel" product of the limited principle of omniscience with itself.

We follow Simpson [15] for the basic definitions and background results of reverse mathematics. This paper relies on two standard subsystems of second order arithmetic. RCA_0 includes the basic axioms PA^- , the Δ_1^0 comprehension scheme, and the Σ_1^0 induction scheme. ACA_0 extends RCA_0 with the comprehension and induction schemes for arithmetical formulas. We follow Hirst [9] for definitions of graph theory in reverse mathematics.

2 Existence of a connected component

In RCA₀, a *countable graph* G is a pair of sets (V, E) in which the vertex set V is a nonempty infinite subset of \mathbb{N} and the edge set E is a set of unordered pairs of elements of V. A (*connected*) *component* of a countable graph G is a set $C \subseteq V$ such that:

- (1) for all x and y in C, there is a (finite) path in G from x to y, and
- (2) for all x and y in V, if $x \in C$ and there is a path from x to y then $y \in C$.

RCA₀ proves that a subset of a countable graph is a connected component in this sense if and only if it is a maximal path connected subset of the graph. Our definition of a component has the advantage, compared to "maximal path connected subset," of being stated as a Π_2^0 formula. A set of vertices of a graph is *totally disconnected* if no two vertices in the set are in the same connected component of the graph. As usual, a countable graph (V, E)is *computable* if $V = \mathbb{N}$ and E is a computable relation, and a connected component is computable if its characteristic function is computable.

Our first theorem shows that the problem of constructing even a single component of a countable graph exceeds the power of RCA_0 . As usual, $\mathbb{N}^{<\mathbb{N}}$ denotes the set of finite sequences of natural numbers, including the empty sequence. For $\sigma, \tau \in \mathbb{N}^{<\mathbb{N}}$, $\sigma^{\gamma}\tau$ denotes the concatenation of σ and τ , and $\sigma \subset \tau$ indicates that σ is a proper initial segment of τ .

Theorem 2.1. The principle that every countable graph has a connected component is equivalent to ACA_0 over RCA_0 .

Proof. First, working in ACA₀, let G = (V, E) be a countable graph. By Theorem 2.5 of Hirst [9], there is a function $f: V \to \mathbb{N}$ such that for all vand v' in V, f(v) = f(v') if and only if v and v' lie in the same connected component of G. Let v_0 be an element of V. Then Δ_1^0 comprehension proves the existence of the set $C = \{v \in V : f(v) = f(v_0)\}$, which is a connected component of G.

For the reversal, we work in RCA_0 and assume that every countable graph has a connected component. By Lemma III.1.3 of Simpson [15], to establish ACA_0 it suffices to prove the existence of the range of an arbitrary injection $f: \mathbb{N} \to \mathbb{N}$.

Construction. We construct a countable graph G. For each $\sigma \in \mathbb{N}^{<\mathbb{N}}$ and each $n \in \mathbb{N}$, G has a vertex labeled v_n^{σ} , and these are all the vertices. For

each $\sigma \in \mathbb{N}^{<\mathbb{N}}$ and each $i \in \mathbb{N}$, there is an edge from v_i^{σ} to v_{i+1}^{σ} . If f(i) = j, then for each $\sigma \in \mathbb{N}^{<\mathbb{N}}$ there is an edge from v_i^{σ} to $v_0^{\sigma^{\frown}\langle j \rangle}$. An illustration of a typical piece of the construction is shown in Figure 1.

Verification. Δ_1^0 comprehension suffices to form this set of edges, using f as a parameter, and thus the graph G can be constructed in RCA₀. By assumption, G has a connected component C. Let v_k^{σ} be a vertex of C; then v_0^{σ} is also in C.

Claim: Each $j \in \mathbb{N}$ is in the range of f if and only if $v_0^{\sigma^{\frown}(j)}$ is in C. The forward direction is immediate; if f(i) = j then the construction ensures that there is a path from v_0^{σ} to $v_0^{\sigma^{\frown}(j)}$, and so $v_0^{\sigma^{\frown}(j)} \in C$. For the converse, suppose that $v_0^{\sigma^{\frown}(j)}$ is in C. We can prove from the construction, using Π_1^0 induction on the length of the path, that for any path beginning at $v_0^{\sigma^{\frown}(j)}$ which contains no edge of the form $(v_0^{\sigma^{\frown}(j)}, v_i^{\sigma})$, the final vertex of the path is of the form $v_k^{\sigma^{\frown}(j)^{\frown}\tau}$ for some k and some sequence τ which might be empty. In particular, the final vertex of the path is not v_0^{σ} . Thus, because there is a path from $v_0^{\sigma^{\frown}(j)}$ to v_0^{σ} in C, there must be an edge in G of the form $(v_0^{\sigma^{\frown}(j)}, v_i^{\sigma})$. But then f(i) = j, so j is in the range of f, as desired. This proves the claim.

By the claim, the range of f consists of exactly those j such that $v_0^{\sigma^{\wedge}\langle j \rangle}$ is in C. Thus the range of f can be formed using Δ_1^0 comprehension relative to C.

Substituting a computable enumeration of \emptyset' for f, the construction in the proof yields a purely computability theoretic corollary.

Corollary 2.2. There is a computable graph G such that every connected component of G computes (and is thus Turing equivalent to) \emptyset' .

The graph in the proof of Theorem 2.1 can be constructed with several additional properties, allowing our results to be stated in a sharper form. Provided that the function f used in the construction of Theorem 2.1 is an injection, the graph that is constructed is acyclic. Therefore, sharper versions of Theorem 2.1 and Corollary 2.2 hold, in which "graph" is replaced with "acyclic graph."

We can also characterize the connectivity of the graph. A graph is bounded if there is a function $h: V \to V$ such that whenever (v_1, v_2) is an edge, it follows that $v_2 < h(v_1)$. A computable graph with a computable




Figure 1: Illustration of a typical piece of the construction in Theorem 2.1. The diagonal edges indicate that f(1) = 2 and f(3) = 0.

bounding function is called *highly computable* (or *highly recursive*). If the construction of Theorem 2.1 is modified by replacing each edge from v_i^{σ} to $v_0^{\sigma^{-}\langle j \rangle}$ by an edge from v_i^{σ} to $v_i^{\sigma^{-}\langle j \rangle}$, RCA₀ proves that the resulting graph is bounded. Therefore, a variant Theorem 2.1 holds for countable bounded acyclic graphs, and a variant of Corollary 2.2 holds for highly computable acyclic graphs.

3 Graphs with finitely many components

The graph constructed in Theorem 2.1 has infinitely many connected components. In this section, we prove two results addressing graphs with finitely many connected components. These results show that, although decompositions of these graphs into components can be formed computably, there are subtleties with the manner in which the bound on the number of components is stated.

The first result applies to graphs for which there is a finite set of vertices V_0 such that each vertex of G is either in V_0 or path connected to at least one element of V_0 . Given such a set, RCA_0 can prove that G can be decomposed into its connected components, that is, there is an $f: V \to \mathbb{N}$ such that for all distinct vertices v_1 and v_2 , $f(v_1) = f(v_2)$ if and only if v_1 is path connected to v_2 . Note that if v_0 is a vertex and f is such a decomposition,

then RCA_0 proves the existence of the set $\{v \in V : f(v) = f(v_0)\}$, which is exactly the connected component containing v_0 .

Theorem 3.1. The following is provable in RCA_0 . If G = (V, E) is a countable graph and there is a finite set $V_0 = \{v_0, v_1, \ldots, v_{n-1}\}$ of vertices of G such that each vertex of G is either in V_0 or path connected to at least one element of V_0 , then G can be decomposed into its connected components, and consequently G has a connected component.

Proof. We work in RCA₀. Let G and V_0 be as described above. Form an enumeration $S_0, S_1, \ldots, S_{2^n-1}$ of the subsets of V_0 , ordered so that larger sets come first, that is, if $S_i \subseteq S_j$ then $j \leq i$. Let $\varphi(k)$ denote the formula that asserts that for every finite path p in G, p is not a path between distinct elements of S_k . If S_k is a singleton, then $\varphi(k)$ holds. Note that $\varphi(k)$ is a Π_1^0 formula. By an easy extension of Theorem A of Paris and Kirby [12] to second order arithmetic, Σ_1^0 induction implies the Π_1^0 least element principle. Apply this principle to find the least k_0 such that $\varphi(k_0)$ holds. Because S_{k_0} is totally disconnected and k_0 is minimal with this property, S_{k_0} is a maximal totally disconnected subset of V_0 , and so every vertex not in S_{k_0} is path connected to exactly one vertex in S_{k_0} . Define the function $f: V \to \mathbb{N}$ by f(v) = v if $v \in S_{k_0}$, and f(v) is the unique element of S_{k_0} that is path connected to v otherwise. The function f exists by Δ_1^0 comprehension and is a decomposition of G into connected components. By the comments preceding the statement of the theorem, G has a connected component. \Box

Our second result on graphs with finitely many connected components uses the weaker hypothesis that there is some n such that every collection of n vertices must include at least one path connected pair. Under this weaker assumption, the Σ_2^0 induction scheme is required to prove the existence of a decomposition into connected components, or even the existence of a single component.

Theorem 3.2. RCA_0 proves that the following are equivalent:

- (1) $I\Sigma_2^0$, the induction scheme for Σ_2^0 formulas with set parameters.
- (2) If G = (V, E) is a countable graph and there is some n such that every collection of n vertices includes at least one path connected pair, then G can be decomposed into its connected components.
- (3) If G = (V, E) is a countable graph and there is some n such that every collection of n vertices includes at least one path connected pair, then G has a connected component.

Proof. We will work in RCA_0 throughout. The equivalence of (1) and (2) is Theorem 4.5 of Hirst [9]. We can capitalize on our work above to provide a succinct alternative proof that (1) implies (2). Suppose G is as in (2). By the Π_2^0 least element principle (which can be deduced from (1) in RCA_0 as in Theorem A of Paris and Kirby [12]) there is a least n_0 such that every collection of n_0 vertices contains a path connected pair. By the minimality of n_0 , there is a set V_0 of $n_0 - 1$ vertices no two of which are path connected. For any vertex v, if $v \notin V_0$, then $V_0 \cup \{v\}$ has cardinality n_0 , so v is path connected to an element of V_0 . By Theorem 3.1, G can be decomposed into its connected components.

The proof that (2) implies (3) follows immediately from the comments in the paragraph preceding Theorem 3.1. It remains to show that (3) implies (1). By the second order analog of Theorem A of Paris and Kirby [12], it suffices to use (3) to deduce the Σ_2^0 least element principle. Fix a Σ_0^0 formula $\theta(k, q, s)$, which may have set parameters. Suppose that there is a k for which $(\exists q)(\forall s)\theta(k, q, s)$ holds. We will show that there is a least integer m such that $(\exists q)(\forall s)\theta(m, q, s)$ holds by constructing a graph and finding a connected component.

For each j < k and each n, let b(n, j) be the least number b (if any such number exists) such that:

- 1. if n > 0, then b(n 1, j) is defined and b > b(n 1, j), and
- 2. there is some $s_0 < b$ such that $\neg \theta(j, n, s_0)$ holds.

Informally, b(n, j) is a bound for a witness that $(\forall s)\theta(j, n, s)$ fails. For each j, $(\exists q)(\forall s)\theta(j, q, s)$ holds if and only if only there are only finitely many n for which b(n, j) is defined. The characteristic function for the predicate t = b(n, j) is definable by primitive recursion, so RCA_0 proves its existence. (See Theorem II.3.4 of Simpson [15].) Whenever t = b(n, j) holds, we have $n \leq t$, so $(\exists n)[t = b(n, j)]$ is equivalent to $(\exists n \leq t)[t = b(n, j)]$, a bounded quantifier applied to a primitive recursive predicate. We will use the formula $(\exists n)[t = b(n, j)]$ in the construction of our graph and the bounded quantifier form in verifying that RCA_0 proves the existence of the graph.

Construction. The graph G (see Figure 2) will consist of a root vertex ρ and subgraphs G_{τ} for each nonempty strictly decreasing sequence τ of numbers less than k. There will be $2^k - 1$ such sequences. The vertices of G_{τ} will be $\{u_n^{\tau} : n \in \mathbb{N}\}$ and $\{v_n^{\tau} : n \in \mathbb{N}\}$. Next, we will define E, the set of edges of G. For a sequence consisting of a single number j < k, add the following edges associated with the subgraph $G_{\langle j \rangle}$:

- $(\rho, u_0^{\langle j \rangle}) \in E$. (This edge connects a corner of $G_{\langle j \rangle}$ to the rest of G.)
- $(u_i^{\langle j \rangle}, u_{i+1}^{\langle j \rangle}) \in E.$
- If $\neg(\exists n)[t = b(n, j)]$ then $(v_t^{\langle j \rangle}, v_{t+1}^{\langle j \rangle}) \in E$.
- If $(\exists n)[t = b(n, j)]$ then $(u_t^{\langle j \rangle}, v_t^{\langle j \rangle}) \in E$.

If τ is a strictly descending sequence and j is less than the last element of τ , then add the following edges for $G_{\tau \cap \langle j \rangle}$:

- $(v_0^{\tau}, u_0^{\tau^{-}\langle j \rangle}) \in E$. (This edge connects a corner of $G_{\tau^{-}\langle j \rangle}$ to G_{τ}).
- If $(v_t^{\tau}, v_{t+1}^{\tau}) \in E$, then put edge $(u_t^{\tau \land \langle j \rangle}, u_{t+1}^{\tau \land \langle j \rangle})$ into E. Also, copy the structure of $G_{\langle j \rangle}$, that is:
 - if $\neg(\exists n)[t = b(n, j)]$ then $(v_t^{\tau \land \langle j \rangle}, v_{t+1}^{\tau \land \langle j \rangle}) \in E$, and

• if
$$(\exists n)[t = b(n, j)]$$
 then $(v_t^{\tau \land \langle j \rangle}, u_t^{\tau \land \langle j \rangle}) \in E$.

• If $(v_t^{\tau}, v_{t+1}^{\tau}) \notin E$, then "cap" the construction of $G_{\tau^{\frown}\langle j \rangle}$ by adding $(u_t^{\tau^{\frown}\langle j \rangle}, v_t^{\tau^{\frown}\langle j \rangle}) \in E$, and "link" the continuation of $G_{\tau^{\frown}\langle j \rangle}$ to G_{τ} by adding $(v_{t+1}^{\tau}, u_{t+1}^{\tau^{\frown}\langle j \rangle}) \in E$.

This completes the definition of G.

Figure 2 illustrates a typical construction for G. In the figure, the vertical edge from $u_1^{\langle 0 \rangle}$ to $v_1^{\langle 0 \rangle}$ in the subgraph $G_{\langle 0 \rangle}$ shows that 0 does not witness that $(\exists q)(\forall s)\theta(0,q,s)$, and the similar vertical edge from $u_3^{\langle 1 \rangle}$ to $v_3^{\langle 1 \rangle}$ in $G_{\langle 1 \rangle}$ shows that 0 does not witness $(\exists q)(\forall s)\theta(1,q,s)$. The subgraph $G_{\langle 1,0 \rangle}$ is a copy of $G_{\langle 0 \rangle}$, except at locations where $G_{\langle 1 \rangle}$ has a vertical edge, in which case a cap and a link are added. If $(\forall s)\theta(1,1,s)$ and $\neg(\exists n)(\forall s)\theta(0,n,s)$, then the component of G containing the vertex $v_4^{\langle 1 \rangle}$ would be disconnected from the component containing ρ , and would contain an isomorphic copy of the portion of $G_{\langle 0 \rangle}$ to the right of $u_4^{\langle 0 \rangle}$. Based on the assumption that $\neg(\exists q)(\forall s)\theta(0,q,s)$, this copy of the right portion of $G_{\langle 0 \rangle}$ would lie entirely in the same component as $v_4^{\langle 1 \rangle}$.




Figure 2: Illustration of a typical piece in the construction in Theorem 3.2.

Verification. We claim that RCA_0 proves that G exists. The set of codes for vertices is Δ_1^0 definable. In the definition of the set of edges, the quantified formulas involving t = b(n, j) are equivalent to formulas with bounded quantifiers. Finally, inclusion of any edge with a vertex in $G_{\tau^{\frown}\langle j \rangle}$ depends only on a bounded initial segment of G_{τ} , so RCA_0 proves the existence of the set of edges.

To verify that G has finitely many connected components, we will show that any collection of $2^k(k+3) + 1$ vertices includes at least one path connected pair. (This is an inexact but convenient upper bound.) First note that for each j, $G_{\langle j \rangle}$ has one connected component if $\neg(\exists q)(\forall s)\theta(j,q,s)$ and two components if $(\exists q)(\forall s)\theta(j,q,s)$. Thus, for each j, $G_{\langle j \rangle}$ has at most two connected components. Suppose $W_d = \{w_0^d, w_1^d, \dots, w_n^d\}$ is a sequence of vertices in $G_{\langle j_0, j_1, \dots, j_d \rangle}$, for some n, such that no pair is connected by a path in G. Because at most one connected component of $G_{(j_0,j_1,\ldots,j_d)}$ is not connected to $G_{(j_0,j_1,\dots,j_{d-1})}$, by successively examining finite paths in G we can discover the first n-1 vertices of $G_{\langle j_0, j_1, \dots, j_{d-1} \rangle}$ that are connected to distinct elements of W_d . Let $W_{d-1} = \{w_0^{d-1}, \dots, w_{n-1}^{d-1}\}$ be the sequence of these vertices. Because no two vertices in W_d are connected by a path, no two vertices in W_{d-1} are connected by a path. RCA₀ can prove the existence of the sequence of codes for the finite sets $W_d, W_{d-1}, \ldots, W_0$. The Σ_1^0 least element principle suffices to prove that in each of these sets, no two vertices are path connected. Because W_0 contains at most two vertices, W_d contains at most d+2 vertices. Generalizing, for any strictly descending sequence τ of values less than k, any totally disconnected set of vertices in G_{τ} must be of size at most k + 2. Now suppose V_0 is a subset of the vertices of G of size $2^k(k+3) + 1$. One of these vertices may be ρ , but at least $2^k(k+3)$ of them lie in the $2^k - 1$ subgraphs of the form G_{τ} . Because RCA₀ proves that the sum of less than 2^k numbers each less than k+3 is less than $2^k(k+3)$, some single subgraph G_{τ} must contain at least k+3 vertices. Because no such collection of k+3 vertices can be totally disconnected, some pair must be connected by a path in G. Thus every collection of $2^{k}(k+3) + 1$ vertices of G contains at least one path connected pair.

Apply item (3) to find a connected component of G. Let C denote the set of vertices in this component. We will use C to find the least m such that $(\exists q)(\forall s)\theta(m,q,s)$. In the following argument, each selection of a least element is a consequence of the Σ_1^0 least element principle. If $\rho \in C$ and there is a j such that $v_n^{\langle j \rangle} \notin C$ for some n, then let m be the least such j. If no such j exists, let m = k. If $\rho \notin C$, then there is a shortest sequence τ and a least n_0 such that $v_{n_0}^{\tau} \in C$. Because τ is decreasing, $\min(\tau)$ is the last element of τ . If there is a $j < \min(\tau)$ such that $v_n^{\tau^{\frown}\langle j \rangle} \notin C$ for some $n > n_0$, then let m be the least such j. If no such j exists, let $m = \min \tau$. In each of these cases, m is the least integer such that $(\exists q)(\forall s)\theta(m,q,s)$, as desired. This completes the proof that (3) implies (1) and completes the proof of the theorem.

The previous theorem extends Theorem 4.5 of Hirst [9]. In the context of classical (nonuniform) computability theory, however, there is no issue with induction, and Corollary 4.3 of Hirst [9] shows that each computable graph with finitely many connected components has a computable decomposition into its connected components, using either formulation of the bound on the number of components. Theorem 6.6 shows that situation is more complicated in the context of Weihrauch reducibility.

4 Graphs in which every component is finite

In this section, we turn to a problem that is, in a certain sense, dual to the problem of the previous section. Graphs in which there are only finitely many components must have at least one infinite component. We now consider graphs in which every component is finite. For such graphs, the existence of a single component is trivial, so we turn to principles that imply the existence of infinitely many connected components. There are several seemingly natural ways to state such principles. We show that three are equivalent to each other, and to ACA₀, over RCA₀. Recall that the *canonical index* for a finite set $\{a_0 < a_1 < \cdots < a_k\}$ is the number $\prod_{i \leq k} p_i^{a_i}$, where p_i is the *i*th prime number.

Definition 4.1. The following principles are defined in RCA_0 .

- FC-1: Every infinite graph in which every component is finite has a sequence ⟨d₁, d₂,...⟩ of canonical indices of different components.
- FC-2: Every infinite graph in which every component is finite has a family (C_i : i ∈ N) of distinct connected components. The family is given as a function C: N² → {0,1}.
- FC-3: Every infinite graph in which every component is finite has an infinite totally disconnected set.

Principle FC-3 can be stated equivalently "every infinite graph has either an infinite component or an infinite totally disconnected set." **Lemma 4.2.** The following implications hold over RCA_0 :

$$\mathsf{ACA}_0 \Rightarrow \mathsf{FC}\text{-}1 \Rightarrow \mathsf{FC}\text{-}2 \Rightarrow \mathsf{FC}\text{-}3.$$

Proof. Working in ACA_0 , it is straightforward to prove FC-1 by first decomposing the graph into components and then forming the sequence of canonical indices of the components.

The implication of FC-1 to FC-2 in RCA_0 is straightforward, because we may uniformly construct the characteristic functions for a sequence of sets defined by a sequence of canonical indices.

Finally, the implication from FC-2 to FC-3 proceeds by choosing, for each i, the least v_i such that $C(i, v_i) = 1$. This gives a function f with $f(i) = v_i$, with the range of f being an infinite enumerated set. By a standard argument, every infinite enumerated set has an infinite (decidable) subset. In this situation, any such subset here will be an infinite totally disconnected set in the original graph.

The proof of the next theorem will require a particular technical lemma. Intuitively, this lemma formalizes a standard fact about approximations to \emptyset' that is often required to verify priority arguments: for each *e* there is a *t* such that every program with index less than *e* that halts must halt in no more than *t* steps. This fact is most directly formalized as a Σ_2^0 property of *e*, so some care is necessary to prove the claim in RCA₀.

Lemma 4.3. The following is provable in RCA_0 . Let $f: \mathbb{N} \to \mathbb{N}$ be an injection, and $h_s(e) = \max\{n < s : f(n) \le e\}$ for each e and s. Then for each e there is a t such that $h_t(e') = h_s(e')$ for all $e' \le e$ and all s > t.

Proof. We may construct h, as a function of s and e, using Δ_1^0 comprehension. Fix e. By bounded Σ_1^0 comprehension (Theorem II.3.9 of Simpson [15]), there is a set X such that, for all $i, i \in X$ if and only if $i \leq e$ and $(\exists s)[f(s) = i]$. Define a function $g \colon \mathbb{N} \to \mathbb{N}$ such that g(i) = 0 when $i \notin X$ and $g(i) = (\mu s)[f(s) = i]$ for each $i \in X$. By the strong Σ_1^0 bounding scheme (a consequence of RCA₀ by Exercise II.3.14 of Simpson [15]), the image of the interval [0, e] under g is bounded by some number t. Any such t satisfies the claim.

The proof of the next theorem can be viewed as a priority argument that has been formalized in RCA_0 .

Theorem 4.4. The following are equivalent over RCA_0 :

- ACA_0 .
- FC-*i* for $i \in \{1, 2, 3\}$.

Proof. We work in RCA₀. By Lemma 4.2, it is sufficient to assume FC-3 and prove that the range of an arbitrary injection $f: \mathbb{N} \to \mathbb{N}$ exists. As in Lemma 4.3, Δ_1^0 comprehension is sufficient to define the function $h_s(e) = \max\{n < s : f(n) \le e\}$.

Construction. We form a countable graph G with vertex set $V = \{v_i : i \in \mathbb{N}\}$. We say that i is the *index* of vertex v_i . We construct a sequence $\langle G_s : s \in \mathbb{N} \rangle$ of finite graphs such that the vertices of G_s are $\{v_i : i \leq s\}$ and such that if t > s and v_i, v_j are in G_s then edge (v_i, v_j) is in G_t if and only if it is in G_s . Then $G = \bigcup_{s \in \mathbb{N}} G_s$ can be constructed in RCA₀. To begin, let G_0 be a graph with the single vertex v_0 and no edges.

We say that a vertex v_e of G_s requires attention at stage s + 1 if there is no v_i in the same component of G_s as v_e with i < e (so v_e has the smallest index of all vertices in its component in G_s), and also $h_s(e) > h_t(e)$, where t is maximal such that v_t is in the same component as v_e in G_s . This is a Σ_0^0 property of e and s. If no vertex in G_s requires attention at stage s + 1, let G_{s+1} be G_s with one additional vertex v_{s+1} and no additional edges. Otherwise, there is a minimum e such that v_e requires attention at stage s + 1. In this case, let G_{s+1} be G_s with one new vertex v_{s+1} and an edge between v_{s+1} and v_i when $e \leq i \leq s$.

Verification. It is straightforward to prove by induction that the vertices of G_s are exactly $\{v_0, \ldots, v_s\}$, and that for all v_i, v_j and all s with i < j < s, v_i and v_j are adjacent in G_s if and only if they are adjacent in G_j . We will show that each component of G is of the form $\{v_i \in V : m \leq i < n\}$ for some m < n.

We first show that each component of G is finite. For this, it is enough to prove that for each component C of G there is an r such that C is a component of G_r (in particular, $C \subseteq G_r$). Given a component C of G, choose e to be minimal such that $v_e \in C$. Choose t as in Lemma 4.3, so that $h_t(e') = h_s(e')$ for all $e' \leq e$ and all s > t. No vertex $v_{e'}$ with e' < e can require attention at or after stage e, because otherwise v_e would be in the component of $v_{e'}$ in G. Furthermore, vertex v_e does not require attention at any stage s > t, by the choice of t. Thus no new vertices can be added to the component of v_e after stage $r = \max\{e, t\}$. To show that each connected component of each G_s is of the form $\{v_i \in V : m \leq i < n\}$ for some m < n, note that all components of G_0 are of this form, and the procedure in the construction preserves this property of the graph at each step. Thus, the components of G are either of the desired form or are unbounded. But, by the previous paragraph, every component of G is bounded.

Now assume that A is any infinite totally disconnected set. Arrange A into a sequence $a_0 < a_1 < a_2 < \cdots$ in which the indices of the vertices are strictly increasing. For each i, the component of a_i is an interval $\{v_{e(i)}, v_{e(i)+1}, \ldots, v_{s(i)}\}$. The argument in the previous paragraph shows that the component of each vertex a_i in G is the same as the component of a_i in the induced subgraph consisting of the finite set of vertices with index less than that of a_{i+1} . Therefore, the functions e(i) and s(i) can be formed by Δ_1^0 comprehension using A as a parameter.

Claim: For all i and s, $h_s(e(i)) \leq h_{s(i)}(e(i))$. If not, by the Σ_1^0 least element principle, there is a least i such that, for some s, $h_s(e(i)) > h_{s(i)}(e(i))$. Choose this least i and then choose a minimal witness s. Because s > s(i), v_s is not in the same component of G as $v_{e(i)}$. Thus, by the construction of s, vertex $v_{e(i)}$ requires attention at stage s + 1, and so the construction causes v_s to be in the same component as $v_{e(i)}$, which is impossible. This proves the claim.

Therefore, using the claim, we see that each n is in the range of f if and only if n is in the range of $f \upharpoonright \{0, \ldots, s(i)\}$ for the least i such that the index of a_i is greater than n. This gives a Δ_1^0 definition of the range of f, and so RCA₀ proves that the range exists.

We obtain two computability-theoretic corollaries from the construction of Theorem 4.4.

Corollary 4.5. There is a computable graph in which every component is finite such that every infinite totally disconnected set computes \emptyset' .

Corollary 4.6. There is a computable graph in which every component is finite such that there is no infinite c.e. totally disconnected set of vertices.

Proof. Construct the graph from the proof of Theorem 4.4. Suppose there is an infinite c.e totally disconnected set of vertices, A. Then A has an infinite computable subset, B, which is also totally disconnected. Thus, by Corollary 4.5, B computes \emptyset' , which is impossible.

5 Weak partitions

In this section, we isolate a purely combinatorial principle that underlies the reversal of Theorem 2.1. The principle uses families of sets in the style of Dzhafarov and Mummert [6]. In RCA₀, we define an *enumerated sequence* of enumerated sets to be a function $D: \mathbb{N} \to \mathbb{N} \times \mathbb{N}$. We abuse notation by writing D as a sequence of sets $\langle D_i : i \in \mathbb{N} \rangle$, and by writing $j \in D_i$ if there is an s with $D(s) = \langle i, j \rangle$. We write $D_i = D_j$ if $(\forall n)[n \in D_i \Leftrightarrow n \in D_j]$, which in turn is an abbreviation for

$$(\forall n)[(\exists a)(D(a) = \langle i, n \rangle) \Leftrightarrow (\exists b)(D(b) = \langle j, n \rangle)].$$

We may similarly express $D_i \cap D_j = \emptyset$ as a formula with parameter D. In general, each component is only an enumerated set. We say that D_i exists if there is a set containing exactly those $j \in \mathbb{N}$ for which there is a $k \in \mathbb{N}$ with $D(k) = \langle i, j \rangle$.

Definition 5.1. A weak partition of \mathbb{N} is an enumerated sequence of enumerated sets $\langle D_i : i \in \mathbb{N} \rangle$ that satisfies the following conditions:

- (1) for every $n \in \mathbb{N}$ there is an $i \in \mathbb{N}$ with $n \in D_i$, and
- (2) for all $i, j \in \mathbb{N}$, either $D_i = D_j$ or $D_i \cap D_j = \emptyset$.

A weak partition of \mathbb{N} is a strong partition if it furthermore satisfies the property that $D_i \cap D_j = \emptyset$ when $i \neq j$. Each enumerated set D_i a component of the weak partition.

If $\langle D_i : i \in \mathbb{N} \rangle$ is a strong partition of \mathbb{N} then RCA_0 proves that the set D_j exists for each $j \in \mathbb{N}$. This is because RCA_0 is able to form a function $f: \mathbb{N} \to \mathbb{N}$ such that f(m) is the unique j such that there exists an s with $D(s) = \langle j, m \rangle$; then $D_j = f^{-1}\{j\}$ exists by Δ_1^0 comprehension. The situation is different for weak partitions, however. Rather than providing a direct reversal for the problem of constructing a set in a weak partition of \mathbb{N} , we show that this problem is intricately tied to the problem of finding a component of a countable graph.

Theorem 5.2. The following are equivalent over RCA_0 .

- (1) Every countable graph has a connected component.
- (2) For each weak partition $\langle D_i : i \in \mathbb{N} \rangle$ of \mathbb{N} , there is an *i* such that D_i exists.




Figure 3: Illustration of a typical piece of the construction in Theorem 5.2. The edges beginning at vertices of the form n_i indicate that $0 \in D_0$, $1 \in D_0$, $1 \in D_3$, and $2 \in D_1$. Because $1 \in D_0 \cap D_3$, it must be that $D_0 = D_3$, so the construction will eventually add an edge from n_0 to some vertex of the form $a_{3,i}$.

Proof. Working in RCA₀, we first prove that (1) implies (2). Let $\langle D_i : i \in \mathbb{N} \rangle$ be a weak partition of \mathbb{N} . We build a countable graph G whose vertices are the disjoint union of three infinite sets: $\{d_i : i \in \mathbb{N}\}, \{n_i : i \in \mathbb{N}\}, \text{ and } \{a_{i,s} : i, s \in \mathbb{N}\}$. For all i, j, there is no edge between d_i and d_j and no edge between n_i and n_j . For each i, there is an edge between d_i and $a_{i,0}$, and no edge between d_i and $a_{i,s+1}$ for all i and s, and no other edges among vertices of the form $a_{i,s}$. There is an edge between $a_{i,s}$ and n_j if and only if $D(s) = \langle i, j \rangle$. A diagram of this construction is shown in Figure 3.

Assuming (1), we may choose a connected component C of the graph G. We first claim there is a vertex of the form d_k in C. Let x be any vertex in C. If x is of the form $a_{i,s}$ then d_i is in C. If x is of the form n_j then, by assumption, there is a D_i such that $j \in D_i$. Let s be such that $D(s) = \langle i, j \rangle$. Then there will be an edge in G from n_j to $a_{i,s}$, and thus $a_{i,s}$ and d_i are in C. This proves the claim.

Now choose $d_p \in C$ and let $X = \{j : n_j \in C\}$. We claim that $X = D_p$, which is sufficient to finish the claim that (1) implies (2). As in the previous paragraph, if $j \in D_p$ then $n_j \in C$. For the converse, assume that $n_j \in C$. Then there is a path τ_0 from d_p to n_j in G. Let $\langle n_{i_0}, n_{i_1}, \ldots, n_{i_r} = n_j \rangle$ list the vertices of the form n_i that are visited by τ_0 , in the order they are visited; we will call this sequence the *pattern* of τ_0 . If r = 0 then it is immediate that $j \in D_p$, because the path must be of the form

$$d_p \to a_{p,0} \to a_{p,1} \to \dots \to a_{p,s} \to n_j$$

for some s, and then $D(s) = \langle p, j \rangle$. Otherwise, if r > 0, the matter is more complicated, because we need to verify that a path "directly" from d_p to n_j can be constructed within RCA₀. We will produce a sequence τ_1, \ldots, τ_r of paths from d_p to n_j such that the pattern of τ_m is $\langle n_{i_m}, n_{i_{m+1}}, \ldots, n_{i_r} \rangle$. Then the previous argument will apply to τ_r , completing the argument.

Suppose a path σ is given which begins at a vertex d_p and ends at a vertex n_j with pattern $\langle n_{i_0}, n_{i_1}, \ldots, n_{i_r} \rangle$. If $n_{i_0} = n_{i_1}$, then the vertices visited between the first and second visits to n_{i_0} must all be of the form $a_{k,s}$ for some fixed k. We can remove n_{i_0} and these vertices from the path, and the result will be a shorter path with the same initial and final vertices as σ , but with pattern $\langle n_{i_1}, \ldots, n_{i_r} \rangle$. If $n_{i_0} \neq n_{i_1}$, we look at the vertices visited just before and after n_{i_0} ; the path will be of the form

$$d_p \to \dots \to a_{p,s_0} \to n_{i_0} \to a_{z,s_1} \to \dots \to n_{i_1} \to \dots \to n_{i_r}.$$

Moreover, all the vertices between n_{i_0} and n_{i_1} will be of the form $a_{z,s}$ or d_z . If z = p then we can again remove n_{i_0} by making the first part of the new path proceed directly from d_p to a_{p,s_1} to n_{i_1} and letting the rest be as in σ . If $z \neq p$ then we see that $D(s_0) = \langle p, n_{i_0} \rangle$ and $D(s_1) = \langle z, n_{i_0} \rangle$, so $n_{i_0} \in D_p \cap D_z$. Then, because $n_{i_1} \in D_z$, by assumption n_{i_1} is also in D_p . We may choose any s' with $D(s') = \langle p, n_{i_1} \rangle$ and create a path of the form

$$d_p \to a_{p,0} \to \dots \to a_{p,s'} \to n_{i_1} \to \dots \to n_{i_r}$$

where the part of the path after n_{i_1} is as in σ .

An analysis of the argument of the previous paragraph shows that RCA_0 proves that, given a path σ from a vertex d_p to a vertex n_j with pattern $\langle n_{i_0}, n_{i_1}, \ldots, n_{i_r} \rangle$, with r > 0, there is a path σ' which has the same beginning and ending vertices and has pattern $\langle n_{i_1}, \ldots, n_{i_r} \rangle$. This relationship between σ and σ' can be defined by a Σ_0^0 formula $\phi(\sigma, \sigma')$, so RCA_0 can form the set Z consisting of all such pairs $\langle \sigma, \sigma' \rangle$. Then, by applying Σ_1^0 induction with Z as a parameter, RCA_0 proves that, given a path τ_0 as above, there is a sequence $\langle \tau_0, \ldots, \tau_r \rangle$ with $\phi(\tau_i, \tau_{i+1})$ for all i < r. Then RCA_0 proves that τ_r is a path from d_p to n_j with pattern $\langle n_j \rangle$, and thus $j \in D_p$, as desired.

Now, still working in RCA_0 , we prove that (2) implies (1). Let G be a countable graph. For each vertex v_i , we enumerate a sequence $\langle u_{i,j} : j \in$

 \mathbb{N} by searching (with dovetailing) all possible paths beginning at v_i , and putting $u_{i,j} = w$ if w is the *j*th vertex that is discovered to be path connected to v_i in this way. In particular, we may assume that $u_{i,0} = v_i$ for each *i*. This construction is uniform in *i*, and so there is an enumerated sequence of enumerated sets *D* such that $D_i = \{u_{i,j} : j \in \mathbb{N}\}$ for each *i* (of course D_i may not exist as a set).

We need to verify that D is a weak partition of \mathbb{N} . This follows from the facts that $v_i \in D_i$ for each i and that RCA_0 can concatenate paths in the graph. Thus, if two vertices v and w are both path connected to a vertex u, and v is also path connected to a vertex u', then w is also path connected to u'.

Now we may apply (2) to obtain a set X and index i with $X = D_i$. It is immediate that X is the connected component of vertex v_i in G, as desired.

Applying Theorem 2.1, we obtain a corollary on the reverse mathematics of weak partitions. We also obtain a purely computability theoretic corollary by re-examining the proof of Theorem 5.2 in light of Corollary 2.2.

Corollary 5.3. The principle that if $\langle D_i : i \in \mathbb{N} \rangle$ is a weak partition of \mathbb{N} then D_i exists for some $i \in \mathbb{N}$ is equivalent to ACA₀ over RCA₀.

Corollary 5.4. There is a computable weak partition $\langle D_i : i \in \mathbb{N} \rangle$ of \mathbb{N} (that is, the function $D : \mathbb{N} \to \mathbb{N} \times \mathbb{N}$ is computable) such that for each $i \in \mathbb{N}$ the set D_i computes (and is thus Turing equivalent to) \emptyset' .

6 Parallelization and relationships with LPO

In the previous sections, we have explored the problem of finding a component of a graph using the methods of reverse mathematics. In this section, we study the problem from the viewpoint of Weihrauch reducibility, as described by Brattka and Gherardi [4, 3] and by Dorais, Dzhafarov, Hirst, Mileti, and Shafer [5]. From this viewpoint, each set $C \subseteq 2^{\omega} \times 2^{\omega}$ can be viewed a *Weihrauch problem* in the following way. Each $A \in 2^{\omega}$ is viewed as an *instance* of the problem. Given an instance A, a corresponding *solution* is a $B \in 2^{\omega}$ such that $(A, B) \in C$. The problems studied in this paper are given by Π_2^1 sets, and therefore every instance has at least one solution.

Following Definition 1.5(2) of Dorais *et al.*, we say that a Weihrauch problem $Q \subseteq 2^{\omega} \times 2^{\omega}$ is *strongly Weihrauch reducible* to a Weihrauch problem $R \subseteq 2^{\omega} \times 2^{\omega}$ if there are Turing reductions Φ and Ψ such that whenever

A is an instance of Q, $B = \Phi(A)$ is an instance of R such that whenever T is a solution to B, $S = \Psi(T)$ is a solution to A. This relation is denoted $Q \leq_{sW} R$. Two problems are strongly Weihrauch equivalent if each is strongly Weihrauch reducible to the other.

We have claimed that the proof of Theorem 5.2 provides a tight link between the problem of computing a connected component of a countable graph and the problem of computing a component of a weak partition of \mathbb{N} . This link cannot be established directly through methods of reverse mathematics, however, because the relation of proof theoretic equivalence in reverse mathematics is too coarse. Strong Weihrauch equivalence can be used to establish a tighter link between principles that emphasizes the uniformity of the relationship between the principles. For example, the following corollary follows from an analysis of the proof of Theorem 5.2.

Corollary 6.1. The problem of finding a connected component of a countable graph is strongly Weihrauch equivalent to the problem of finding a component of a weak partition of \mathbb{N} .

In this section, we study two particular Weihrauch problems.

- P: the Weihrauch problem of finding a connected component of a countable graph.
- D: the Weihrauch problem of decomposing a countable graph into connected components. As in Section 3, a decomposition is a map from vertices to \mathbb{N} such that $f(v_0) = f(v_1)$ if and only if v_0 and v_1 are in the same connected component.

We show that P and D are equivalent to their "parallelized" forms, and we obtain results that locate these principles relative to better-known Weihrauch degrees (Theorems 6.4 and 6.5). Many results in this section capitalize on the uniformity of the constructions in the previous sections. The authors thank one of the referees for suggesting the results of Theorems 6.5 and 6.6 and the inclusion of LPO in Theorem 6.4.

We begin by considering the binary parallelization $\langle \mathsf{P}, \mathsf{P} \rangle$. An instance of this problem consists of a pair of countable graphs (G_1, G_2) ; a solution is a pair of sets (C_1, C_2) such that C_1 is a connected component of G_1 and C_2 is a connected component of G_2 . In the terminology more common in the literature on Weihrauch reducibility, the binary parallelization is simply the product of the problem P with itself. Additional information on parallelization and the product operation is given by Brattka and Gherardi [4,3], by Dorais *et al.* [5, Section 2], and by Pauly [13]. **Theorem 6.2.** P is strongly Weihrauch equivalent to its binary parallelization.

Proof. It is immediate that P , like every other problem, is Weihrauch reducible to its parallelization $\langle \mathsf{P}, \mathsf{P} \rangle$. We may effectively convert any computable graph G into an instance of $\langle \mathsf{P}, \mathsf{P} \rangle$ by setting $G_1 = G_2 = G$. Then, given a solution (C_1, C_2) , we simply take the first element of the pair, C_1 .

The proof of the converse is conceptually similar to the proof that the computability problem corresponding to weak König's lemma is Weihrauch equivalent to its binary parallelization. Suppose that $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ are countable graphs. Form a new graph $G_1 \times G_2$ in the following way. The vertices of $G_1 \times G_2$ consist of all pairs (u, v) where $u \in V_1$ and $v \in V_2$. There is an edge between (u, v) and (x, y) if and only if either u = x and $(v, y) \in E_2$, or v = y and $(u, x) \in E_1$. Note that if $(u, x) \in E_1$ and $(v, y) \in E_2$ then there is a path from (u, v) to (x, y) in $G_1 \times G_2$.

Suppose that C is a connected component of $G_1 \times G_2$ and (u, v) is a vertex in C. We claim that a vertex (u, y) is in C if and only if v and y are in the same component of G_2 . For the forward direction, given a path from (u, v) to (u, y) in $G_1 \times G_2$, we can ignore the first components of the vertices, remove any duplicates, and get a path from v to y in G_2 . Conversely, given a path from v to y in G_2 , we can make an ordered pair with u and each vertex of the path to form a path from (u, v) to (u, y) in $G_1 \times G_2$. This proves the claim. Thus we may compute a connected component of G_2 from any connected component of $G_1 \times G_2$. The proof that we may compute a component of G_1 is similar.

According to Definition 2.2 of Dorais *et al.* [5], a Π_2^1 problem R has *finite* tolerance if there is a Turing functional Θ such that whenever B_1 and B_2 are instances of R with $B_1(n) = B_2(n)$ for all $n \ge m$, and S is a solution of B_1 , then $\Theta(S, m)$ is a solution to B_2 . If the graph component problem P had finite tolerance, we could use our Theorem 6.2 above and Theorem 2.5 of Dorais *et al.* [5] to show that the infinite parallelization $\widehat{\mathsf{P}}$ (defined below) is strongly Weihrauch reducible to P. The next theorem rules out this proof strategy.

Theorem 6.3. The problem of finding a connected component of a countable graph does not have finite tolerance.

Proof. We construct a computable graph G with vertex set $V = \{v_i : i \in \mathbb{N}\}$. Such a graph may be presented as the characteristic function of its set of edges, using an indexing $\{e_i : i \in \mathbb{N}\}$ of the edges of the complete graph on V. Without loss of generality, we may assume that the first potential edge e_0 would connect v_0 to v_1 . Let B_1 be the graph consisting of a complete graph the on vertices $\{v_{2i} : i \in \mathbb{N}\}$ and a complete graph on the vertices $\{v_{2i+1} : i \in \mathbb{N}\}$. Let B_2 be B_1 with the edge e_0 added. Let S be the solution to B_1 consisting of the even vertices. Suppose by way of contradiction that Θ is a Turing functional witnessing that P has finite tolerance. Because $B_1(n) = B_1(n)$ for all $n \ge 1$, $\Theta(S, 1)$ is a solution of B_1 , that is, a connected component of B_1 . Because $B_1(n) = B_2(n)$ for all $n \ge 1$, $\Theta(S, 1)$ is also a connected component of B_2 , contradicting the fact that B_1 and B_2 have no common connected components.

Despite the inapplicability of Theorem 2.5 of Dorais *et al.* [5], we can directly prove that the *infinite parallelization* $\widehat{\mathsf{P}}$ is strongly Weihrauch equivalent to P . An instance of the parallelization $\widehat{\mathsf{P}}$ consists of an infinite sequence $\langle G_i \rangle_{i \in \mathbb{N}}$ of countable graphs; a solution is a sequence $\langle C_i \rangle_{i \in \mathbb{N}}$ of sets such that C_i is a component of G_i for each $i \in \mathbb{N}$. An infinite parallelization can also be seen as an infinite product of the problem P with itself.

Our approach will capitalize on established results from the theory of Weihrauch degrees. The following two facts are included in Proposition 4.2 of Brattka and Gherardi [3]. First, Weihrauch reducibility is preserved by parallelization, so $Q \leq_{sW} R$ implies $\widehat{Q} \leq_{sW} \widehat{R}$. Second, parallelization is idempotent, so $\widehat{Q} \equiv_{sW} \widehat{Q}$.

We will make use of the *limited principle of omniscience*, denoted LPO, which asserts that if $p: \mathbb{N} \to \mathbb{N}$ then either there exists an $n \in \mathbb{N}$ such that p(n) = 0, or $p(n) \neq 0$ for all $n \in \mathbb{N}$. Although this principle is trivially provable in classical logic, it has nontrivial Weihrauch degree, as the nonuniform classical proof suggests. The infinite parallelization $\widehat{\mathsf{LPO}}$ is much stronger in reverse mathematics. When stated as a Π_2^1 principle, $\widehat{\mathsf{LPO}}$ is equivalent to ACA_0 over RCA_0 .

Theorem 6.4. Each of the following is strongly Weihrauch equivalent to the others: $P, \hat{P}, D, \hat{D}, and \hat{LPO}$.

Proof. The proof of the theorem will follow easily once we establish the following strong Weihrauch reductions:

$$\widehat{\mathsf{LPO}} \leq_{\mathrm{sW}} \mathsf{P} \leq_{\mathrm{sW}} \mathsf{D} \leq_{\mathrm{sW}} \widehat{\mathsf{LPO}}.$$

Let $\langle p_i \rangle_{i \in \mathbb{N}}$ be a sequence of functions viewed as an instance of $\widehat{\mathsf{LPO}}$. We will conflate ordered pairs with their integer codes and assume the pairing

function is bijective. Define

$$f((i,j)) = \begin{cases} 2i+1 & \text{if } p_i(j) = 0 \text{ and } (\forall t < j)[p_i(t) \neq 0], \\ 2(i,j) & \text{otherwise.} \end{cases}$$

Then f is an injection, and each odd number 2i+1 is in the range of f if and only if $(\exists n)[p_i(n) = 0]$. Use the construction from the proof of Theorem 2.1 to build a graph G; it can be seen that this construction is uniform. As in that theorem, each connected component of G computes (in a uniform manner) the range of f, so $\{i : (\exists n)[p_i(n) = 0]\} = \{i : (\exists n)[f(n) = 2i+1]\}$ is a solution to the instance of $\widehat{\mathsf{LPO}}$ computable from the connected component. Thus $\widehat{\mathsf{LPO}} \leq_{sW} \mathsf{P}$.

If G is a graph with vertex v, each decomposition of G into connected components computes the connected component containing v. Thus $P \leq_{sW} D$. It remains to show that $D \leq_{sW} \widehat{LPO}$. Suppose G is a graph with vertices $\{v_i : i \in \mathbb{N}\}$. For each pair (i, j), let $p_{(i,j)}(n) = 0$ if n is a code for a path from v_i to v_j and let $p_{(i,j)}(n) = 1$ otherwise. Applying \widehat{LPO} yields the set $S = \{(i, j) : \exists n(p_{(i,j)}(n) = 0)\}$, which is the set of codes for all pairs of path connected vertices of G. The function $d(v_n) = (\mu i < n)[(i, n) \in S]$ (where the function returns n if the finite search fails) is a decomposition of G into connected components that is computable from S. Thus $D \leq_{sW} \widehat{LPO}$, completing our initial chain of reductions.

By transitivity of Weihrauch reducibility, we may conclude that $P \equiv_{sW} D \equiv_{sW} \widehat{LPO}$. Because parallelization is idempotent and preserves Weihrauch reducibility, we have $\widehat{P} \equiv_{sW} \widehat{D} \equiv_{sW} \widehat{LPO} \equiv_{sW} \widehat{LPO}$, completing the proof of the theorem.

The techniques of Theorem 6.4 can also be applied to the properties FC -*i* defined in Section 4.

Theorem 6.5. For $i \in \{1, 2, 3\}$, the principles FC-*i* are strongly Weihrauch equivalent to \widehat{LPO} , as are their infinite parallelizations.

Proof. As in the proof of the preceding theorem, we first prove a chain of Weihrauch reductions:

$$\widehat{\mathsf{LPO}} \leq_{sW} \mathsf{FC-3} \leq_{sW} \mathsf{FC-2} \leq_{sW} \mathsf{FC-1} \leq_{sW} \widehat{\mathsf{LPO}}.$$

To prove that $\widehat{\mathsf{LPO}} \leq_{\mathrm{sW}} \mathsf{FC-3}$, fix an instance of $\widehat{\mathsf{LPO}}$ and construct an associated injection f as in the proof of Theorem 6.4. Apply the (uniform) construction from the proof of Theorem 4.4 to f to obtain a graph in which

every component is finite. As in that proof, any infinite totally disconnected set computes the range of f, and so computes a solution to the instance of $\widehat{\mathsf{LPO}}$.

The proof of Lemma 4.2 shows that $\mathsf{FC}\text{-3} \leq_{\mathrm{sW}} \mathsf{FC}\text{-2} \leq_{\mathrm{sW}} \mathsf{FC}\text{-1}$. It remains to prove that $\mathsf{FC}\text{-1} \leq_{\mathrm{sW}} \widehat{\mathsf{LPO}}$. Let G be an infinite graph in which every connected component is finite. Construct an instance of $\widehat{\mathsf{LPO}}$ consisting of a sequence of functions $p_i \colon \mathbb{N} \to \mathbb{N}$, defined as follows. Set $p_i(n) = 1$ if (1) iis a canonical index for F, a finite collection of vertices, (2) G restricted to F is a connected subgraph, and (3) if $v_n \notin F$, then there is no edge between v_n and any vertex listed by F. Otherwise, set $p_i(n) = 0$. Note that i is the canonical index of a connected component of G if and only if $p_i(n) \neq 0$ for all n. Consequently, any solution of the instance of $\widehat{\mathsf{LPO}}$ computes the sequence of codes for connected components satisfying $\mathsf{FC}\text{-1}$.

Because parallelization is idempotent and preserves Weihrauch reducibility, the parallelizations of FC-1, FC-2, and FC-3 are also Weihrauch equivalent to \widehat{LPO} .

The Weihrauch degree of $\widehat{\mathsf{LPO}}$ is the same as that of lim, which has been widely studied. For example, it is Weihrauch complete for effective Σ_2^0 measurable functions [1], and equivalent to computing the Radon-Nikodym derivative [10], constructing a measure with a given closed set as support [14], and computing a subgame perfect equilibrium of an infinite sequential game with open/closed payoffs [11].

Weihrauch reducibility can also be used to analyze statements about connected components and decompositions of graphs with finitely many connected components. We let P_k be the problem of finding a connected component of a countable graph that has exactly k connected components, and let D_k be the problem of decomposing a graph with exactly k connected components into its connected components. These graphs do not come with any witness for the number of components; for example D_5 is simply the restriction of D to graphs with exactly 5 components.

The next theorem relates D_k and P_k to $\mathsf{C}_{\mathbb{N}}$, the Weihrauch principle for closed choice on the natural numbers. The principle $\mathsf{C}_{\mathbb{N}}$ asserts that, given a non-surjective map $p: \mathbb{N} \to \mathbb{N}$, we can find an $n \in \mathbb{N}$ which is not in the range of p. The motivation for the name is that the complement of the range of p can be viewed as a closed subset of \mathbb{N} . For more information about closed choice, see Brattka, de Brecht, and Pauly [2]. **Theorem 6.6.** For each $k \ge 2$, $\mathsf{P}_k \equiv_{sW} \mathsf{D}_k \equiv_{sW} \mathsf{C}_{\mathbb{N}}$.

Proof. First we will show that $C_{\mathbb{N}} \leq_{sW} P_2$. Let $p: \mathbb{N} \to \mathbb{N}$ enumerate the complement of a nonempty subset of \mathbb{N} . Construct a related graph G as follows. Let $V = \{u_i : i \in \mathbb{N}\} \cup \{v_i : i \in \mathbb{N}\}$ be the vertices of G and include all edges of the form (u_i, u_{i+1}) in G. Consider the finite (possibly empty) increasing sequence b_i defined as follows. Let $b_0 = (\mu t)[p(t) = 0]$ if such a t exists. Let $b_{i+1} = (\mu t)[t > b_i \land (\exists s \leq t)(p(s) = i + 1)]$ if such a t exists. Note that there is a computable procedure that determines for each $j \in \mathbb{N}$ whether j is equal to some b_i . For each n, if $n = b_i$, then add the edge (u_n, v_n) to G; otherwise, add (v_n, v_{n+1}) to G. The resulting graph will be similar to the subgraph $G_{\langle 0 \rangle}$ shown in Figure 2, with the number of caps equal to the least n not in the range of p. In particular, the resulting graph G will have exactly two connected components, one of which will contain all vertices of the form u_i .

Suppose C is a connected component of G. If $u_0 \in C$, some (least) v_n must not be in C. Locate it and call it v_{n_0} . If $u_0 \notin C$, some (least) v_n must be in C. Locate it and call it v_{n_0} . In either case, the number of edges of the form (u_j, v_j) with $j \leq n_0$ is the least integer not in the range of p. Thus $C_{\mathbb{N}} \leq_{\mathrm{sW}} \mathsf{P}_2$.

To see that $C_{\mathbb{N}} \leq_{sW} \mathsf{P}_k$ for k > 2, construct k - 1 copies of the graph in the preceding paragraph, using vertices $V_j = \{u_i^j : i \in \mathbb{N}\} \cup \{v_i^j : i \in \mathbb{N}\}$ for each $j \leq k - 1$. For each positive $j \leq k - 1$, add the edge (u_0^0, u_0^j) . By an argument similar to the preceding paragraph, any connected component of the graph can be used to locate an integer not in the range of p.

If G is a graph with vertex v, then from any decomposition of G into connected components we can compute the component containing v. Thus, for each k, $P_k \leq_{sW} D_k$. It remains to show that $D_k \leq_{sW} C_N$. Let G be a graph with exactly k connected components. Let $\{s_i : i \in \mathbb{N}\}$ be a bijective enumeration of the size k subsets of the vertices of G. Let $\{t_j : j \in \mathbb{N}\}$ be an enumeration of all the finite paths in G. We will identify pairs (i, j) with their integer codes, assuming a bijective encoding. Define p(i, j) = i + 1 if t_j includes a path between two vertices of s_i , and p(i, j) = 0 otherwise. Since G has exactly k components, some s_i has no connected vertices, and consequently i+1 is not in the range of p. Thus p is non-surjective. If i+1 is any integer not in the range of p, then every vertex of G is either an element of s_i or connected by a path to exactly one element of s_i . Using s_i , we can compute a decomposition of G into connected components. Thus $D_k \leq_{sW} C_N$. The theorem follows by transitivity of Weihrauch reducibility.

References

- Vasco Brattka, Effective Borel measurability and reducibility of functions, Mathematical Logic Quarterly 51 (2005), no. 1, 19–44.
- [2] Vasco Brattka, Matthew de Brecht, and Arno Pauly, Closed choice and a uniform low basis theorem, Annals of Pure and Applied Logic 163 (2012), no. 8, 968–1008.
- [3] Vasco Brattka and Guido Gherardi, Effective choice and boundedness principles in computable analysis, Bulletin of Symbolic Logic 1 (2011), 73-117. arXiv:0905.4685.
- [4] _____, Weihrauch degrees, omniscience principles and weak computability, J. Symbolic Logic 76 (2011), no. 1, 143–176. MR2791341 (2012c:03186)
- [5] François G. Dorais, Damir D. Dzhafarov, Jeffry L. Hirst, Joseph R. Mileti, and Paul Shafer, On uniform relationships between combinatorial problems, Trans. AMS (2015). To appear.
- [6] Damir D. Dzhafarov and Carl Mummert, On the strength of the finite intersection principle, Israel J. Math. 196 (2013), no. 1, 345–361. MR3096595
- [7] Harvey Friedman, Some systems of second order arithmetic and their use, Proceedings of the International Congress of Mathematicians (Vancouver, B. C., 1974), Vol. 1, 1975, pp. 235–242. MR0429508 (55 #2521)
- [8] _____, Abstracts: Systems of second order arithmetic with restricted induction, I and II, J. Symbolic Logic 41 (1976), no. 2, 557–559.
- Jeffry L. Hirst, Connected components of graphs and reverse mathematics, Arch. Math. Logic 31 (1992), no. 3, 183–192. MR1147740
- [10] Mathieu Hoyrup, Cristóbal Rojas, and Klaus Weihrauch, Computability of the Radon-Nikodym derivative, Computability 1 (2012), no. 1, 3–13.
- [11] Stéphane Le Roux and Arno Pauly, Weihrauch degrees of finding equilibria in sequential games, 2014.
- [12] J. B. Paris and L. A. S. Kirby, Σ_n -collection schemas in arithmetic, Logic Colloquium '77 (Proc. Conf., Wrocław, 1977), 1978, pp. 199–209. MR519815 (81e:03056)
- [13] Arno Pauly, On the (semi)lattices induced by continuous reducibilities, Mathematical Logic Quarterly 56 (2010), no. 5, 488–502.
- [14] Arno Pauly and Willem Fouché, How constructive is constructing measures?, 2014.
- [15] Stephen G. Simpson, Subsystems of second order arithmetic, 2nd ed., Perspectives in Logic, Cambridge University Press, Cambridge; Association for Symbolic Logic, Poughkeepsie, NY, 2009. MR2517689 (2010e:03073)